

THE SHARP LIFESPAN FOR A SYSTEM OF MULTIPLE SPEED WAVE EQUATIONS: RADIAL CASE

MARVIN KOONCE AND JASON METCALFE

Dedicated to Professor Thomas C. Sideris on the occasion of his 70th birthday

ABSTRACT. Ohta examined a system of multiple speed wave equations with small initial data and demonstrated a finite time blowup. We show, in the radial case, that the same system exists almost globally with the same lifespan as a lower bound. To do this, we use integrated local energy estimate, r^p weighted local energy estimates, the Morawetz estimate that results from using the scaling vector field as a multiplier, and mixed speed ghost weights.

1. INTRODUCTION

This article focuses on establishing the sharp lifespan, in the radial case, for a multiple speed system of wave equations with small initial data introduced in [23]. Based on [11], we know that there exists a constant c so that quasilinear wave equations of the form

$$(1.1) \quad \begin{cases} \square u = Q(u, \partial u, \partial^2 u), & (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \\ u(0, \cdot) = \varepsilon f, \quad \partial_t u(0, \cdot) = \varepsilon g, \end{cases}$$

where $f, g \in C_c^\infty(\mathbb{R}^3)$ and Q vanishes to second order at the origin, have solutions on $[0, T_*]$ with $T_* \geq c/\varepsilon^2$ if ε is sufficiently small. With the additional assumption that $(\partial_u^2 Q)(0, 0, 0) = 0$, which in essence rules out u^2 terms and leaves $u\partial u$ nonlinearities at the lowest order, almost global existence $T_* \geq \exp(c/\varepsilon)$ was proved. The latter result was partly extended to systems of equations in [14].

In the case of multiple speed systems of wave equations

$$\square_{c_I} u^I = Q^I(\partial u, \partial^2 u), \quad (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3,$$

almost global existence was established in [10]. Here

$$\square_c = \partial_t^2 - c^2 \Delta$$

denotes the d'Alembertian at speed c . Like in the single speed case [8], [3], if the nonlinearity satisfies a null condition, then global existence can be recovered. In the semilinear case, if

$$Q^I(\partial u) = B_{JK}^{I, \alpha\beta} \partial_\alpha u^J \partial_\beta u^K + C(\partial u),$$

where C vanishes to third order and repeated variables are implicitly summed using the Einstein convention, the null condition is only necessary when $c_I = c_J = c_K$. See, e.g., [26], [24], [27], [12, 13]. The reason that it suffices to only have an assumption on the same speed interactions is that solutions to the wave equation enjoy additional decay off of the light cone, and when there are differing speeds one of the factors will be away from its light cone thus contributing more rapid decay.

For multiple speed analogs of (1.1) with $(\partial_u^2 Q)(0, 0, 0) = 0$, one may wonder if global existence can be recovered provided no quadratic nonlinear term in the equation for u^I has factors that are both at the same speed c_I . More precisely, if we truncate to quadratic level for semilinear equations

$$\square_{c_I} u^I = A_{JK}^{I, \alpha} u^J \partial_\alpha u^K + B_{JK}^{I, \alpha\beta} \partial_\alpha u^J \partial_\beta u^K,$$

Key words and phrases. wave equations, local energy estimates, multiple speed, almost global existence.

The authors gratefully acknowledge the partial support from NSF grants DMS-2054910 and DMS-2135998. The results contained herein were developed as a part of the Undergraduate Honors Thesis of MK. JM thanks Yizhou Gu and Xiao-Ming Porter who completed some calculations related to this material as a part of their undergraduate research.

it may be reasonable to expect global existence provided that

$$(1.2) \quad A_{JK}^{I,\alpha} = 0 = B_{JK}^{I,\alpha\beta}, \quad \text{whenever } c_I = c_J = c_K.$$

In a somewhat surprising result, [23] demonstrated that (1.2) is not a sufficient condition for global existence for sufficiently small initial data. Indeed, for $c > 1$, the following system was considered.

$$(1.3) \quad \begin{cases} \square v = w \partial_t v, & (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3, \\ \square_c w = (\partial_t v)^2, \\ v(0, \cdot) = \varepsilon v_{(0)}, & \partial_t v(0, \cdot) = \varepsilon v_{(1)}, \\ w(0, \cdot) = \varepsilon w_{(0)}, & \partial_t w(0, \cdot) = \varepsilon w_{(1)}. \end{cases}$$

It was established that there are smooth, compactly supported initial data $v_{(j)}, w_{(j)}$ and a constant c_0 so that the lifespan T_* satisfies $T_* \leq \exp(c_0/\varepsilon^2)$.

The current study seeks to show the reverse inequality. That is, for any data $v_{(0)}, v_{(1)}, w_{(0)}, w_{(1)}$ that are smooth and compactly supported, we seek to show that there is a constant \tilde{c} so that $T_\varepsilon \geq \exp(\tilde{c}/\varepsilon^2)$. Without loss of generality, we take the supports of the data to be contained within the unit ball $\{|x| \leq 1\}$. We shall also use time translation symmetry and take the initial data on the time slice $t = 4$. In the current article we consider only the radial case: $v_{(j)}(x) = v_{(j)}(|x|)$, $w_{(j)}(x) = w_{(j)}(|x|)$.

Using the assumption of radial symmetry, we can reduce the question at hand to a problem in $(1+1)$ -dimensions. Indeed, by conjugating, we have

$$\square_c w(t, r) = r^{-1}(\partial_t^2 - c^2 \partial_r^2)(rw).$$

If we set $V(t, r) = rv(t, r)$ and $W(t, r) = rw(t, r)$, we can instead seek sufficiently regular solutions to the $(1+1)$ -dimensional initial-value boundary-value problem

$$(1.4) \quad \begin{cases} \square V = r^{-1}W \partial_t V, & (t, r) \in \mathbb{R}_+ \times \mathbb{R}_+, \\ \square_c W = r^{-1}(\partial_t V)^2, \\ W(t, 0) = V(t, 0) = 0, & \text{for all } t, \\ V(4, \cdot) = \varepsilon V_{(0)}, & \partial_t V(4, \cdot) = \varepsilon V_{(1)}, \\ W(4, \cdot) = \varepsilon W_{(0)}, & \partial_t W(4, \cdot) = \varepsilon W_{(1)}. \end{cases}$$

Here (when applied to V, W) we understand $\square_c = \partial_t^2 - c^2 \partial_r^2$ to be the $(1+1)$ -dimensional d'Alembertian.

If we extend $V_{(j)}, W_{(j)}$ in an odd fashion, $V_{(j)}(-r) = -V_{(j)}(r)$, $W_{(j)}(-r) = -W_{(j)}(r)$, then it is straightforward to check that V, W extend oddly, and we can instead seek to solve

$$(1.5) \quad \begin{cases} \square V = x^{-1}W \partial_t V, & (t, x) \in \mathbb{R}_+ \times \mathbb{R}, \\ \square_c W = x^{-1}(\partial_t V)^2, \\ V(4, \cdot) = \varepsilon V_{(0)}, & \partial_t V(4, \cdot) = \varepsilon V_{(1)}, \\ W(4, \cdot) = \varepsilon W_{(0)}, & \partial_t W(4, \cdot) = \varepsilon W_{(1)}. \end{cases}$$

The main theorem is a statement of almost global existence for (1.5).

Theorem 1.1. *Suppose that $V_{(j)}, W_{(j)} \in C_c^\infty(\mathbb{R})$ and $V_{(j)}(x) = -V_{(j)}(-x)$, $W_{(j)}(x) = -W_{(j)}(-x)$ for $j = 0, 1$. Then there exist constants $\tilde{c}, \varepsilon_0 > 0$ such that when $0 < \varepsilon < \varepsilon_0$, (1.5) has a unique solution $(V, W) \in C^\infty([4, T_\varepsilon] \times \mathbb{R})$, where*

$$(1.6) \quad T_\varepsilon = \exp(\tilde{c}/\varepsilon^2).$$

As indicated above, Theorem 1.1 leads to the following corollary, showing that, in the radial case, the upper bound on the lifespan in [23] is sharp.

Corollary 1.2. *Suppose that $v_{(j)}, w_{(j)} \in C_c^\infty(\mathbb{R}^3)$, and that $v_{(j)}(x) = v_{(j)}(|x|)$, $w_{(j)}(x) = w_{(j)}(|x|)$ for $j = 0, 1$. Then there exist constants $\tilde{c}, \varepsilon_0 > 0$ such that when $0 < \varepsilon < \varepsilon_0$, (1.3) has a unique solution $(v, w) \in C^\infty([0, T_\varepsilon] \times \mathbb{R}^3)$ with T_ε as in (1.6).*

Our proof relies on the method of invariant vector fields with adaptations to a restricted set of vector fields, which is necessitated as the Lorentz boosts have an associated speed and only commute with the d'Alembertian of the same speed. This was pioneered in, e.g., [10], [6, 7], [15, 16, 17], [12, 13]. The method is further simplified in the radial case as the only relevant vector fields are derivatives and the scaling vector field. And in the study at hand, we will rely solely on the scaling vector field to avoid issues with commuting derivatives with the singular weight introduced in the one dimensional reduction.

Like [6] and [16], we will call upon a class of integrated local energy estimates in order to prove long-time existence. These estimates go back to the seminal work [22]. In the current setting, a number of variants are used. These include the r^p -weighted estimates of [4], ghost weighted estimates from [1], and integrated estimate variants of [21] where the scaling vector field is used as a multiplier.

To obtain the requisite decay, rather than using the classical Klainerman-Sobolev estimate [9], which introduces Lorentz boosts, or the weighted Sobolev estimate [9], which provides decay in $|x|$ but fails to capture the added decay off of the light cone, as was done in [6] and [16], we will instead use a class of space-time Klainerman-Sobolev estimates of [20]. The space-time nature of these decay bounds meshes well with the integrated local energy estimates.

In the estimates that we prove, we allow for ghost weights that are associated to a different speed than the equation. This works in the non-radial case (see the forthcoming work [2]) provided that the speed within the ghost weight exceeds that of the speed of the equation. In the radial case, however, no restriction on the speeds is needed. This is the primary place where we rely upon the radially assumption. We anticipate examining the general case in a future study.

2. NOTATION AND DECAY ESTIMATES

For $x \in \mathbb{R}$, we denote $r = |x|$. We will use the standard null coordinates

$$u = t - r, \quad \underline{u} = t + r, \quad \partial_u = \frac{1}{2}(\partial_t - \partial_r), \quad \partial_{\underline{u}} = \frac{1}{2}(\partial_t + \partial_r).$$

The above correspond to speed 1, but at wave speed c , we will instead have

$$u_c = ct - r, \quad \underline{u}_c = ct + r, \quad \partial_{u_c} = \frac{1}{2c}(\partial_t - c\partial_r), \quad \partial_{\underline{u}_c} = \frac{1}{2c}(\partial_t + c\partial_r).$$

We denote the scaling vector field by

$$(2.1) \quad S = t\partial_t + r\partial_r = u\partial_u + \underline{u}\partial_{\underline{u}} = u_c\partial_{u_c} + \underline{u}_c\partial_{\underline{u}_c}.$$

It will be important for later purposes to note

$$[\partial_{u_c}, S] = \partial_{u_c}, \quad [\partial_{\underline{u}_c}, S] = \partial_{\underline{u}_c}, \quad [\square, S] = 2\square.$$

For $k \in \mathbb{N}$, we use the notation

$$|S^{\leq k} w| = \sum_0^k |S^j w|.$$

The space-time Klainerman-Sobolev-type estimates of [20] will be the principal source of decay. See also [19]. The only difference herein is allowing for $c \neq 1$ and simplifications that result from the $(1+1)$ -dimensional regime. As such, we will be brief in the presentation.

If at speed c , the compactly supported data (at $t = 4$) are taken to be supported in the unit ball, the components of the solution will be supported in $C^c = \{(t, x) : t \in [4, T_\varepsilon], r \leq ct - (4c - 1)\}$. We now dyadically decompose C^c in both t and in either r or u_c depending on the proximity to the light cone. We first decompose in t and set

$$C_\tau^c = \{(t, r) \in \mathbb{R}_+ \times \mathbb{R}_+ : t \in [4, T_\varepsilon] \cap [\tau, 2\tau], r \leq ct - (4c - 1)\}.$$

We further break into

$$C_\tau^{c, R=1} = C_\tau^c \cap \{r \leq 2\}, \quad C_\tau^{c, R} = C_\tau^c \cap \{R \leq r \leq 2R\} \text{ when } 1 < R,$$

and

$$C_\tau^{c, U_c} = C_\tau^c \cap \{U_c \leq ct - r \leq 2U_c\} \text{ when } 1 < U_c.$$

We finally set

$$C_\tau^{c,c\tau/2} = C_\tau^c \cap \{ct - r \geq c\tau/2\} \cap \{r \geq c\tau/2\}.$$

Throughout $\tau, R, U_c, U_1 := U$ will be understood to range over dyadic values. This gives

$$(2.2) \quad C_\tau^c = \left(\bigcup_{1 \leq R \leq c\tau/4} C_\tau^{c,R} \right) \cup \left(\bigcup_{2 \leq U_c \leq c\tau/4} C_\tau^{c,U_c} \right) \cup C_\tau^{c,c\tau/2}.$$

We will let $\tilde{C}_\tau^{c,R}$ and \tilde{C}_τ^{c,U_c} denote slight enlargements (in both scales) to allow for tails of cutoff functions. On the components of the decomposition (and their enlargements) we have:

$$\begin{aligned} \langle r \rangle &\approx R, \quad t \approx \tau, \quad u_c \approx \tau \quad \text{on } C_\tau^{c,R}, \quad 1 \leq R \leq c\tau/4, \\ r &\approx \tau, \quad t \approx \tau, \quad \langle u_c \rangle \approx U_c \quad \text{on } C_\tau^{c,U_c}, \quad 2 \leq U_c \leq c\tau/4. \end{aligned}$$

The remaining region $C_\tau^{c,c\tau/2}$ may be thought of as either $R = c\tau/2$ or $U_c = c\tau/2$.

The following are space-time analogs of the Klainerman-Sobolev estimates:

Lemma 2.1 ([20]). *Suppose $W \in C^2([4, T_\varepsilon] \times \mathbb{R})$ is an odd function. If $c > 0$, $\tau \geq 4$, $1 \leq R \leq c\tau/2$, $2 \leq U_c \leq c\tau/4$, then*

$$(2.3) \quad \|r^{-\frac{1}{2}}W\|_{L_t^\infty L_r^\infty(C_\tau^{c,R})} \lesssim \frac{1}{\tau^{\frac{1}{2}}R} \|S^{\leq 1}W\|_{L_t^2 L_r^2(\tilde{C}_\tau^{c,R})} + \frac{1}{\tau^{\frac{1}{2}}} \|\partial_r S^{\leq 1}W\|_{L_t^2 L_r^2(\tilde{C}_\tau^{c,R})},$$

$$(2.4) \quad \|W\|_{L_t^\infty L_r^\infty(C_\tau^{c,U_c})} \lesssim \frac{1}{\tau^{\frac{1}{2}}U_c^{\frac{1}{2}}} \|S^{\leq 1}W\|_{L_t^2 L_r^2(\tilde{C}_\tau^{c,U_c})} + \frac{U_c^{\frac{1}{2}}}{\tau^{\frac{1}{2}}} \|\partial_r S^{\leq 1}W\|_{L_t^2 L_r^2(\tilde{C}_\tau^{c,U_c})}.$$

Here and throughout, L_r^2 is simply the 1-dimensional norm:

$$\|f\|_{L_r^2}^2 = \int_0^\infty |f(r)|^2 dr.$$

Proof. On C_τ^{c,U_c} regions and on $C_\tau^{c,R}$ regions for $R > 1$, the result follows from [20] in one spatial dimension. On the $C_\tau^{c,R=1}$ regions, however, a different change of variables is required to avoid picking up vector fields other than S . Some care is also taken to assist with the singular behavior at $r = 0$ that was introduced in the reduction to one dimension.

Let $\beta : [0, \infty) \rightarrow [0, 1]$ be a smooth cutoff function such that $\beta(y) = 1$ for $y \in [1, 2]$ and $\beta(y) = 0$ for $y \in [0, 1 - \delta] \cup [2 + \delta, \infty)$ where $0 < \delta \ll 1$. We examine $\beta(t/\tau)W(t, r)$. We first change variables to $t = e^s$ and $r = \rho e^s$. Applying the Fundamental Theorem of Calculus in s and ρ (relying up on the fact that $S^{\leq 1}W$ is odd and hence vanishes at $r = 0$), we have

$$|\beta(e^s/\tau)W(e^s, \rho e^s)| \lesssim \int_0^\rho \int_{-\infty}^{\log t} |\partial_\zeta \partial_s (\beta(e^s/\tau)W(e^s, \zeta e^s))| ds d\zeta.$$

Since $\partial_s(W(e^s, \zeta e^s)) = (SW)(e^s, \zeta e^s)$, upon converting back to (t, r) -coordinates, we have

$$(2.5) \quad |\beta(t/\tau)W(t, r)| \lesssim \tau^{-2} \int \int_{[\tau(1-\delta), t] \times [0, r]} |S^{\leq 1}W(t, z)| dz dt \\ + \tau^{-1} \int \int_{[\tau(1-\delta), t] \times [0, r]} |(\partial_r S^{\leq 1}W)(t, z)| dz dt.$$

Applying the Schwarz inequality to the right side and multiplying through by $r^{-\frac{1}{2}}$ then yields

$$\|r^{-\frac{1}{2}}W\|_{L_t^\infty L_r^\infty(C_\tau^{c,R=1})} \lesssim \frac{1}{\tau^{\frac{3}{2}}} \|S^{\leq 1}W\|_{L_t^2 L_r^2(\tilde{C}_\tau^{c,R=1})} + \frac{1}{\tau^{\frac{1}{2}}} \|\partial_r S^{\leq 1}W\|_{L_t^2 L_r^2(\tilde{C}_\tau^{c,R=1})},$$

which is stronger than (2.3) when $R = 1$. □

By writing $\partial_r = \partial_{\underline{u}_c} - \partial_{u_c}$ and using (2.1), we can obtain the following corollary. See, e.g., [19].

Corollary 2.2. *Suppose $W \in C^2([4, T_\varepsilon] \times \mathbb{R})$ is an odd function. If $c > 0$, $\tau \geq 4$, $1 \leq R \leq c\tau/2$, $2 \leq U_c \leq c\tau/4$, then*

$$(2.6) \quad \|r^{-\frac{1}{2}}W\|_{L_t^\infty L_r^\infty(C_\tau^c, R)} \lesssim \frac{1}{\tau^{\frac{1}{2}}R} \|S^{\leq 2}W\|_{L_t^2 L_r^2(\tilde{C}_\tau^c, R)} + \frac{1}{\tau^{\frac{1}{2}}} \|\partial_{\underline{u}_c} S^{\leq 1}W\|_{L_t^2 L_r^2(\tilde{C}_\tau^c, R)},$$

$$(2.7) \quad \|W\|_{L_t^\infty L_r^\infty(C_\tau^c, U_c)} \lesssim \frac{1}{\tau^{\frac{1}{2}}U_c^{\frac{1}{2}}} \|S^{\leq 2}W\|_{L_t^2 L_r^2(\tilde{C}_\tau^c, U_c)} + \frac{\tau^{\frac{1}{2}}}{U_c^{\frac{1}{2}}} \|\partial_{\underline{u}_c} S^{\leq 1}W\|_{L_t^2 L_r^2(\tilde{C}_\tau^c, U_c)}.$$

Using (2.1), we also obtain the following bounds on $\partial V := (\partial_t, \partial_r)V$, which appeared previously in [19]. They are space-time variants of estimates originally from [10] and [25].

Lemma 2.3. *Suppose $V \in C^3([4, T_\varepsilon] \times \mathbb{R})$ is an odd function. If $\tau \geq 4$, $1 \leq R \leq \tau/2$, and $2 \leq U \leq \tau/4$, then*

$$(2.8) \quad \|V\|_{L_t^\infty L_r^\infty(C_\tau^1, R)} \lesssim \frac{1}{\tau^{\frac{1}{2}}R^{\frac{1}{2}}} \|S^{\leq 2}V\|_{L_t^2 L_r^2(\tilde{C}_\tau^1, R)} + \frac{R^{\frac{1}{4}}}{\tau^{\frac{1}{2}}} \|r^{\frac{1}{4}}\partial_{\underline{u}} S^{\leq 1}V\|_{L_t^2 L_r^2(\tilde{C}_\tau^1, R)},$$

$$(2.9) \quad \|V\|_{L_t^\infty L_r^\infty(C_\tau^1, U)} \lesssim \frac{1}{\tau^{\frac{1}{2}}U^{\frac{1}{2}}} \|S^{\leq 2}V\|_{L_t^2 L_r^2(\tilde{C}_\tau^1, U)} + \frac{\tau^{\frac{1}{2}}}{U^{\frac{1}{2}}} \|\partial_{\underline{u}} S^{\leq 1}V\|_{L_t^2 L_r^2(\tilde{C}_\tau^1, U)},$$

$$(2.10) \quad \|\partial V\|_{L_t^\infty L_r^\infty(C_\tau^1, R)} \lesssim \frac{1}{\tau^{\frac{1}{2}}R^{\frac{1}{2}}} \|\partial S^{\leq 2}V\|_{L_t^2 L_r^2(\tilde{C}_\tau^1, R)} + \frac{R^{\frac{1}{4}}}{\tau^{\frac{1}{2}}} \|r^{\frac{1}{4}}\square S^{\leq 1}V\|_{L_t^2 L_r^2(\tilde{C}_\tau^1, R)},$$

$$(2.11) \quad \|\partial V\|_{L_t^\infty L_r^\infty(C_\tau^1, U)} \lesssim \frac{1}{\tau^{\frac{1}{2}}U^{\frac{1}{2}}} \|\partial S^{\leq 2}V\|_{L_t^2 L_r^2(\tilde{C}_\tau^1, U)} + \frac{\tau^{\frac{1}{2}}}{U^{\frac{1}{2}}} \|\square S^{\leq 1}V\|_{L_t^2 L_r^2(\tilde{C}_\tau^1, U)}.$$

Proof. Outside of the $R = 1$ case, the above follow from [20] and [19]. When $R = 1$, we argue as in (2.5). At this point, a different application of the Schwarz inequality yields

$$\beta(t/\tau)V(t, r) \lesssim \frac{1}{\tau^{\frac{3}{2}}} \|S^{\leq 1}V\|_{L_t^2 L_r^2(\tilde{C}_\tau^1, R=1)} + \frac{r^{\frac{1}{4}}}{\tau^{\frac{1}{2}}} \|r^{\frac{1}{4}}\partial_r S^{\leq 1}V\|_{L_t^2 L_r^2(\tilde{C}_\tau^1, R=1)}.$$

Using (2.1) then allows us to recover (2.8). A subsequent application of (2.1), as in [19], yields (2.10). \square

It will be of utmost importance to track the availability to sum over the dyadic ranges R, τ, U, U_c . To this end, we shall use notation such as

$$\|W\|_{\ell_\tau^2 \ell_U^2 \leq \tau/4, L_t^2 L_r^2(C_\tau^1, U)}^2 = \sum_\tau \sum_{U \leq \tau/4} \|W\|_{L_t^2 L_r^2(C_\tau^1, U)}^2, \quad \|W\|_{\ell_U^\infty \ell_\tau^2 \leq \tau/4, L_t^2 L_r^2(C_\tau^1, U)}^2 = \sup_{U \geq 1} \sum_\tau \|W\|_{L_t^2 L_r^2(C_\tau^1, U)}^2.$$

Other variants, such as over $C_\tau^{1, R}, C_\tau^c, U_c$, etc., are similarly defined.

3. LOCAL ENERGY ESTIMATES

In this section we shall gather the energy estimates that will be used in the sequel. These are variants of the integrated local energy estimates, which originated in [22]. See also [6], [28], [16], [18] for subsequent generalizations and applications to nonlinear equations. These ideas are combined with an integrated variant of the classical estimate obtained by using the scaling vector field as a multiplier as was done in [21]. We also utilize a ghost weight as introduced in [1]. As in [2], we use ghost weights where the speed does not necessarily coincide with the given equation. For the speed c component, our principal estimate is an r^p -weighted (with $p = 1$) integrated local energy estimate from [4]. It is combined with a ghost weight as was done in [14]. We pair this with a Hardy inequality that takes advantage of the multiple speeds.

For weighting functions, we shall use, for a parameter $\theta \geq 1$,

$$(3.1) \quad \sigma_\theta(y) = \frac{y}{|y| + \theta}, \quad \sigma'_\theta(y) = \frac{\theta}{(|y| + \theta)^2},$$

which is bounded and C^1 . Moreover, we note that

$$(3.2) \quad \sigma'_\theta(y) \gtrsim \theta^{-1} \quad \text{on } \{\langle y \rangle \approx \theta\}.$$

In order to help control the singularity at $r = 0$ that results from the reduction to one-dimension, we also note

$$(3.3) \quad \frac{d}{dr}(\sigma_R(r))^{\frac{1}{2}} \gtrsim r^{-\frac{1}{2}}R^{-\frac{1}{2}} \quad \text{on } \{\langle r \rangle \approx R\}.$$

Using $(\sigma_R(r))^\delta$ with $0 < \delta < 1$ as a weight in order to gain added control at $r = 0$ has appeared previously in, e.g., [5].

We shall use the following proposition when $p = 0$ and $p = 1$. In the former case, this corresponds to the classical integrated local energy estimates and variants that are available using the ghost weight. For $p = 1$, these are instead variants of [21] where $S = t\partial_t + r\partial_r$ is used as a multiplier. The fact that no upper bound on p is necessary is a consequence of the radially assumption. The angular terms necessitate $p \leq 2$ in more general cases. We start with a lemma, which indicates that the ‘‘good’’ derivative portion of the calculation works independently.

Lemma 3.1. *Suppose that $p \geq 0$, $V \in C^2([4, T] \times \mathbb{R})$, and that there exists $\tilde{R} > 0$ so that $V(t, x) \equiv 0$ whenever $|x| > \tilde{R}$. Then*

$$(3.4) \quad \begin{aligned} & \|r^{-\frac{1}{4}}\langle r \rangle^{-\frac{1}{4}}\langle \underline{u} \rangle^{\frac{p}{2}}\partial_{\underline{u}}V\|_{\ell_r^\infty \ell_r^2 L_t^2 L_r^2(C_r^{1,R})}^2 + \|\langle u \rangle^{-\frac{1}{2}}\langle \underline{u} \rangle^{\frac{p}{2}}\partial_{\underline{u}}V\|_{\ell_{U^c}^\infty \ell_r^2 L_t^2 L_r^2(C_r^{1,U})}^2 + \|\langle u_c \rangle^{-\frac{1}{2}}\langle \underline{u} \rangle^{\frac{p}{2}}\partial_{\underline{u}}V\|_{\ell_{U^c}^\infty \ell_r^2 L_t^2 L_r^2(C_r^{c,U_c})}^2 \\ & \lesssim \|\langle r \rangle^{\frac{p}{2}}\partial_{\underline{u}}V(4, \cdot)\|_{L_r^2}^2 + \int_4^T \int_0^\infty \langle \underline{u} \rangle^p |\square V| |\partial_{\underline{u}}V| dr dt. \end{aligned}$$

Proof. We consider

$$(3.5) \quad \begin{aligned} & \int_4^T \int_0^\infty e^{(\sigma_R(r))^{\frac{1}{2}}} e^{-\sigma_U(t-r)} e^{-\sigma_{U_c}(ct-r)} (1+t+r)^p \square V(t, r) (\partial_t + \partial_r)V(t, r) dr dt \\ & = \frac{1}{2} \int_4^T \int_0^\infty e^{\sigma_R(r)} e^{-\sigma_U(t-r)} e^{-\sigma_{U_c}(ct-r)} (1+t+r)^p (\partial_t - \partial_r) \left((\partial_t + \partial_r)V \right)^2 dr dt. \end{aligned}$$

Integrating by parts gives that this is equivalent to

$$(3.6) \quad \begin{aligned} & \frac{1}{2} \int_0^\infty e^{(\sigma_R(r))^{\frac{1}{2}}} e^{-\sigma_U(T-r)} e^{-\sigma_{U_c}(cT-r)} (1+T+r)^p \left((\partial_t + \partial_r)V(T, r) \right)^2 dr \\ & - \frac{1}{2} \int_0^\infty e^{(\sigma_R(r))^{\frac{1}{2}}} e^{-\sigma_U(4-r)} e^{-\sigma_{U_c}(4c-r)} (5+r)^p \left((\partial_t + \partial_r)V(4, r) \right)^2 dr \\ & + \frac{1}{2} \int_4^T e^{-\sigma_U(t)} e^{-\sigma_{U_c}(ct)} (1+t)^p (\partial_r V(t, 0))^2 dt \\ & + \frac{1}{2} \int_4^T \int_0^\infty \frac{d}{dr}(\sigma_R(r))^{\frac{1}{2}} e^{(\sigma_R(r))^{\frac{1}{2}}} e^{-\sigma_U(t-r)} e^{-\sigma_{U_c}(ct-r)} (1+t+r)^p \left((\partial_t + \partial_r)V \right)^2 dr dt \\ & + \int_4^T \int_0^\infty e^{(\sigma_R(r))^{\frac{1}{2}}} \sigma'_U(t-r) e^{-\sigma_U(t-r)} e^{-\sigma_{U_c}(ct-r)} (1+t+r)^p \left((\partial_t + \partial_r)V \right)^2 dr dt \\ & + \frac{c+1}{2} \int_4^T \int_0^\infty e^{(\sigma_R(r))^{\frac{1}{2}}} e^{-\sigma_U(t-r)} \sigma'_{U_c}(ct-r) e^{-\sigma_{U_c}(ct-r)} (1+t+r)^p \left((\partial_t + \partial_r)V \right)^2 dr dt. \end{aligned}$$

We drop the non-negative first and third terms. We subsequently can restrict the range of the fourth, fifth, and sixth terms so that (3.2) can be applied. We also use that σ_θ is bounded uniformly in θ . For example,

$$\begin{aligned} & \int_4^T \int_0^\infty e^{(\sigma_R(r))^{\frac{1}{2}}} \sigma'_U(t-r) e^{-\sigma_U(t-r)} e^{-\sigma_{U_c}(ct-r)} (1+t+r)^p \left((\partial_t + \partial_r)V \right)^2 dr dt \\ & \gtrsim U^{-1} \int \int_{\{t-r \approx U\}} (1+t+r)^p \left((\partial_t + \partial_r)V \right)^2 dr dt. \end{aligned}$$

Taking appropriate supremums in R, U using the boundedness of σ_θ then yields (3.4). \square

While the previous lemma worked independently, to get similar control on the remaining derivatives, we must examine both the “good” derivative $\partial_t + \partial_r$ and the “bad” derivative $\partial_t - \partial_r$ in unison. In the radial case, this results from the $r = 0$ boundary behavior. In more general situations, there is interaction amongst the angular behavior as well.

Proposition 3.2. *Suppose that $p \geq 0$, $V \in C^2([4, T] \times \mathbb{R})$, and that there exists $\tilde{R} > 0$ so that $V(t, x) \equiv 0$ whenever $|x| > \tilde{R}$. Then*

$$(3.7) \quad \begin{aligned} & \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} \langle \underline{u} \rangle^{\frac{p}{2}} \partial_{\underline{u}} V\|_{\ell_{\tilde{R}}^{\infty} \ell_r^2 L_t^2 L_r^2(C_{\tau}^1, R)}^2 + \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} \langle u \rangle^{\frac{p}{2}} \partial_u V\|_{\ell_{\tilde{R}}^{\infty} \ell_r^2 L_t^2 L_r^2(C_{\tau}^1, R)}^2 \\ & + \|\langle u_c \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{p}{2}} \partial_{\underline{u}} V\|_{\ell_{\tilde{U}_c}^{\infty} \ell_r^2 L_t^2 L_r^2(C_{\tau}^c, U_c)}^2 + \|\langle u_c \rangle^{-\frac{1}{2}} \langle u \rangle^{\frac{p}{2}} \partial_u V\|_{\ell_{\tilde{U}_c}^{\infty} \ell_r^2 L_t^2 L_r^2(C_{\tau}^c, U_c)}^2 \\ & + \|\langle u \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{p}{2}} \partial_{\underline{u}} V\|_{\ell_{\tilde{U}}^{\infty} \ell_r^2 L_t^2 L_r^2(C_{\tau}^1, U)}^2 \lesssim \|\langle r \rangle^{\frac{p}{2}} \partial V(4, \cdot)\|_{L^2}^2 \\ & + \int_4^T \int_0^{\infty} \langle \underline{u} \rangle^p |\square V| |\partial_{\underline{u}} V| dr dt + \int_4^T \int_0^{\infty} \langle u \rangle^p |\square V| |\partial_u V| dr dt. \end{aligned}$$

Proof. We argue much like in the preceding proof but instead examine

$$(3.8) \quad \begin{aligned} & \int_4^T \int_0^{\infty} e^{-(\sigma_R(r))\frac{1}{2}} e^{-\sigma_{U_c}(ct-r)} (1 + |t-r|)^p \square V(t, r) (\partial_t - \partial_r) V(t, r) dr dt \\ & = \frac{1}{2} \int_4^T \int_0^{\infty} e^{-(\sigma_R(r))\frac{1}{2}} e^{-\sigma_{U_c}(ct-r)} (1 + |t-r|)^p (\partial_t + \partial_r) \left((\partial_t - \partial_r) V \right)^2 dr dt. \end{aligned}$$

Integrating by parts shows that this is equal to

$$(3.9) \quad \begin{aligned} & \frac{1}{2} \int_0^{\infty} e^{-(\sigma_R(r))\frac{1}{2}} e^{-\sigma_{U_c}(cT-r)} (1 + |T-r|)^p \left((\partial_t - \partial_r) V(T, r) \right)^2 dr \\ & - \frac{1}{2} \int_0^{\infty} e^{-(\sigma_R(r))\frac{1}{2}} e^{-\sigma_{U_c}(4c-r)} (1 + |4-r|)^p \left((\partial_t - \partial_r) V(4, r) \right)^2 dr - \frac{1}{2} \int_4^T e^{-\sigma_{U_c}(ct)} (1+t)^p (\partial_r V(t, 0))^2 dt \\ & + \frac{1}{2} \int_4^T \int_0^{\infty} \frac{d}{dr} (\sigma_R(r))\frac{1}{2} e^{-(\sigma_R(r))\frac{1}{2}} e^{-\sigma_{U_c}(ct-r)} (1 + |t-r|)^p \left((\partial_t - \partial_r) V \right)^2 dr dt \\ & + \frac{c-1}{2} \int_4^T \int_0^{\infty} e^{-(\sigma_R(r))\frac{1}{2}} \sigma'_{U_c}(ct-r) e^{-\sigma_{U_c}(ct-r)} (1 + |t-r|)^p \left((\partial_t - \partial_r) V \right)^2 dr dt. \end{aligned}$$

If it were not for the $r = 0$ boundary term, we could argue as above to obtain our estimate. Here we instead must sum (3.6) and (3.9). The former provides the desired control in the $r = 0$ boundary term. Then using (3.1) and (3.2) as in the preceding lemma concludes the proof. \square

We next proceed with the speed c estimates. This first estimate combines the r^p weighting of [4] with the ghost weighting of [1] as was done in [14]. Here, however, we use a ghost weight at speed 1 despite working with a solution to a speed c wave equation.

Proposition 3.3. *Suppose that $c > 0$, $W \in C^2([4, T] \times \mathbb{R})$, and that there exists $\tilde{R} > 0$ so that $W(t, x) \equiv 0$ whenever $|x| > \tilde{R}$. Then*

$$(3.10) \quad \begin{aligned} & \|\partial_{\underline{u}_c} W\|_{\ell_{\tilde{R}}^2 \ell_r^2 L_t^2 L_r^2(C_{\tau}^1, R)}^2 + \|\langle u \rangle^{-\frac{1}{2}} r^{\frac{1}{2}} \partial_{\underline{u}_c} W\|_{\ell_{\tilde{U}}^{\infty} \ell_r^2 L_t^2 L_r^2(C_{\tau}^1, U)}^2 \\ & \lesssim \|r^{\frac{1}{2}} \partial_{\underline{u}_c} W(4, \cdot)\|_{L^2}^2 + \int_4^T \int_0^{\infty} r |\square_c W| |\partial_{\underline{u}_c} W| dr dt. \end{aligned}$$

Proof. We now start with

$$\int_4^T \int_0^{\infty} r e^{-\sigma_V(t-r)} \square_c W(t, r) (\partial_t + c\partial_r) W(t, r) dr dt = \frac{1}{2} \int_4^T \int_0^{\infty} r e^{-\sigma_V(t-r)} (\partial_t - c\partial_r) \left((\partial_t + c\partial_r) W \right)^2 dr dt,$$

which upon integrating by parts is seen to be the same as

$$\begin{aligned} & \frac{1}{2} \int_0^\infty r e^{-\sigma_U(T-r)} \left((\partial_t + c\partial_r)W(T, r) \right)^2 dr - \frac{1}{2} \int_0^\infty r e^{-\sigma_U(4-r)} \left((\partial_t + c\partial_r)W(4, r) \right)^2 dr \\ & + \frac{c}{2} \int_4^T \int_0^\infty e^{-\sigma_U(t-r)} \left((\partial_t + c\partial_r)W \right)^2 dr dt + \frac{1+c}{2} \int_4^T \int_0^\infty r \sigma'_U(t-r) e^{-\sigma_U(t-r)} \left((\partial_t + c\partial_r)W \right)^2 dr dt. \end{aligned}$$

After noting the first term is non-negative, we may drop it. The proof is then completed by restricting the range of the last term so that (3.2) may be applied and using the boundedness of σ_θ . \square

The preceding lemma will be paired with two Hardy inequalities as W appears in our nonlinearity without a derivative. The first is a space-time variant of the standard Hardy inequality. See, e.g., [14].

Lemma 3.4. *Suppose that $c > 0$, $W \in C^1([4, T] \times \mathbb{R})$ is an odd function, and that there exists $\tilde{R} > 0$ so that $W(t, x) \equiv 0$ whenever $|x| > \tilde{R}$. Then*

$$(3.11) \quad \|r^{-1}W\|_{L_t^2 L_x^2([4, T] \times \mathbb{R})} \lesssim \|r^{-\frac{1}{2}}W(4, \cdot)\|_{L_x^2} + \|\partial_{\underline{u}_c} W\|_{L_t^2 L_x^2([4, T] \times \mathbb{R})}.$$

Proof. We write

$$\int_4^T \int r^{-2} W^2 dx dt = -\frac{2}{c} \int_4^T \int_0^\infty (\partial_t + c\partial_r)r^{-1} \cdot W^2 dr dt.$$

As W is C^1 and odd, the Mean Value Theorem gives that $W = \mathcal{O}(r)$ near $r = 0$. Thus, these integrals are well-defined and $r^{-1}W^2(t, r) \rightarrow 0$ as $r \rightarrow 0$. Integrating by parts reveals

$$(3.12) \quad \begin{aligned} \int_4^T \int r^{-2} W^2 dx dt + \frac{2}{c} \int_0^\infty r^{-1} (W(T, r))^2 dr \\ = \frac{2}{c} \int_0^\infty r^{-1} (W(4, r))^2 dr + \frac{4}{c} \int_4^T \int_0^\infty r^{-1} W (\partial_t + c\partial_r)W dr dt. \end{aligned}$$

The Schwarz inequality shows that

$$\begin{aligned} \frac{4}{c} \int_4^T \int_0^\infty r^{-1} W (\partial_t + c\partial_r)W dr dt & \lesssim \|r^{-1}W\|_{L_t^2 L_x^2([4, T] \times \mathbb{R})} \|\partial_{\underline{u}_c} W\|_{L_t^2 L_x^2([4, T] \times \mathbb{R})} \\ & \leq \frac{1}{2} \|r^{-1}W\|_{L_t^2 L_x^2([4, T] \times \mathbb{R})}^2 + C \|\partial_{\underline{u}_c} W\|_{L_t^2 L_x^2([4, T] \times \mathbb{R})}^2. \end{aligned}$$

Plugging this into (3.12) and absorbing the $\|r^{-1}W\|_{L_t^2 L_x^2}$ term back into the left side yields the desired estimate. \square

The second Hardy inequality that we rely upon takes advantage of the multiple speed structure.

Lemma 3.5. *Suppose $c > 1$, $W \in C^1([4, T] \times \mathbb{R})$ is an odd function, and that there exists $\tilde{R} > 0$ so that $W(t, x) \equiv 0$ when $|x| > \tilde{R}$. Then*

$$(3.13) \quad \|\langle u \rangle^{-\frac{1}{2}} r^{-\frac{1}{2}} W\|_{\ell_V^\infty \ell_r^2 L_t^2 L_x^2(C_r^1, U)} \lesssim \|r^{-\frac{1}{2}}W(4, \cdot)\|_{L_x^2} + \|\partial_{\underline{u}_c} W\|_{L_t^2 L_x^2([4, T] \times \mathbb{R})}.$$

Proof. The argument here is similar to the preceding, but we take advantage of the difference in speeds between the weight $\langle u \rangle^{-1}$ and the speed c of the equation. To this end, using (3.2), we observe

$$\iint_{\{\langle t-r \rangle \approx U\}} \langle u \rangle^{-1} r^{-1} W^2 dx dt \leq \frac{2}{1-c} \int_4^T \int_0^\infty r^{-1} (\partial_t + c\partial_r)(e^{\sigma_U(t-r)}) \cdot W^2 dr dt$$

and use integration by parts to see that the right side is equivalent to

$$(3.14) \quad \begin{aligned} \frac{2}{1-c} \int_0^\infty r^{-1} e^{\sigma_U(T-r)} (W(T, r))^2 dr + \frac{2}{c-1} \int_0^\infty r^{-1} e^{\sigma_U(4-r)} (W(4, r))^2 dr \\ - \frac{2c}{c-1} \int_4^T \int_0^\infty r^{-2} e^{\sigma_U(t-r)} W^2 dr dt + \frac{4}{c-1} \int_4^T \int_0^\infty r^{-1} e^{\sigma_U(t-r)} W (\partial_t + c\partial_r)W dr dt. \end{aligned}$$

Using the Cauchy-Schwarz inequality to bound

$$(3.15) \quad \frac{4}{c-1} \int_4^T \int_0^\infty r^{-1} e^{\sigma v(t-r)} W (\partial_t + c \partial_r) W \, dr \, dt$$

$$\leq \frac{c}{2(c-1)} \int_4^T \int_0^\infty r^{-2} e^{\sigma v(t-r)} W^2 \, dr \, dt + C \int_4^T \int_0^\infty e^{\sigma v(t-r)} (\partial_{\underline{u}_c} W)^2 \, dr \, dt,$$

plugging this into (3.14), and omitting nonpositive terms from the right produces the desired result. \square

4. PROOF OF THEOREM 1.1

To solve (1.5), we set up an iteration. Let $W_0 \equiv V_0 \equiv 0$, and let V_j, W_j solve

$$(4.1) \quad \begin{cases} \square V_j = x^{-1} W_{j-1} \partial_t V_{j-1}, \\ \square_c W_j = x^{-1} (\partial_t V_{j-1})^2, \\ V_j(4, \cdot) = \varepsilon V_{(0)}, \quad \partial_t V_j(4, \cdot) = \varepsilon V_{(1)}, \\ W_j(4, \cdot) = \varepsilon W_{(0)}, \quad \partial_t W_j(4, \cdot) = \varepsilon W_{(1)}. \end{cases}$$

We shall show that the sequence $((V_j, W_j))_{j=0}^\infty$ is Cauchy on $[4, T_\varepsilon] \times \mathbb{R}$ in an appropriate sense, and by standard results the limit is the desired solution.

4.1. Boundedness. We first show a uniform boundedness of the sequence $((V_j, W_j))$. Let

$$(4.2) \quad M_j = \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} \langle u \rangle^{\frac{1}{2}} S^{\leq 7} \partial_u V_j\|_{\ell_R^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{1,R})} + \|\langle u_c \rangle^{-\frac{1}{2}} \langle u \rangle^{\frac{1}{2}} S^{\leq 7} \partial_u V_j\|_{\ell_{U_c}^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{c,U_c})}$$

$$+ \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_R^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{1,R})} + \|\langle u_c \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_{U_c}^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{c,U_c})}$$

$$+ \|\langle u \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_U^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{1,U})} + \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} S^{\leq 10} \partial V_j\|_{\ell_R^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{1,R})}$$

$$+ \|\langle u_c \rangle^{-\frac{1}{2}} S^{\leq 10} \partial V_j\|_{\ell_{U_c}^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{c,U_c})} + \|\langle u \rangle^{-\frac{1}{2}} S^{\leq 10} \partial_{\underline{u}} V_j\|_{\ell_U^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{1,U})}$$

$$+ \|r^{\frac{1}{2}} \langle u \rangle^{-\frac{1}{2}} S^{\leq 10} \partial_{\underline{u}_c} W_j\|_{\ell_U^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{1,U})} + \|r^{-\frac{1}{2}} \langle u \rangle^{-\frac{1}{2}} S^{\leq 10} W_j\|_{\ell_U^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{1,U})}$$

$$+ \|S^{\leq 10} \partial_{\underline{u}_c} W_j\|_{\ell_R^2 \ell_\tau^2 L_t^2 L_r^2(C_\tau^{1,R})} + \|r^{-1} S^{\leq 10} W_j\|_{\ell_R^2 \ell_\tau^2 L_t^2 L_r^2(C_\tau^{c,R})}.$$

We will label these terms $(I)_j, (II)_j, \dots, (XII)_j$.

Since $\square V_1 \equiv \square_c W_1 \equiv 0$, the smallness of the initial data along with (3.7), (3.10), (3.11), and (3.13) imply that there exists C_0 so that

$$M_1 \leq C_0 \varepsilon.$$

Moreover, V_1 is supported within C^1 . Under the assumption that

$$(4.3) \quad M_k \leq 2C_0 \varepsilon, \quad \text{for } 0 \leq k \leq j-1,$$

and that $\text{supp } V_{j-1} \subset C^1$, we shall show that there exists a fixed constant C such that

$$(4.4) \quad M_j^2 \leq (C_0 \varepsilon)^2 + C(2C_0 \varepsilon)^2 (\log(2 + T_\varepsilon))^{\frac{1}{2}} M_j$$

for $(t, r) \in [4, T_\varepsilon] \times \mathbb{R}_+$. It will also follow immediately that $\text{supp } V_j \subset C^1$. Using (1.6) and absorbing the M_j back into the left side, this implies

$$M_j^2 \leq 2(C_0 \varepsilon)^2 + \tilde{C} \tilde{c} \varepsilon^2.$$

Provided the \tilde{c} in (1.6) is chosen to be sufficiently small, this shows that

$$M_j \leq 2C_0 \varepsilon$$

and completes the inductive proof of boundedness.

4.1.1. *Decay bounds.* To aid in the proof of (4.4), we shall first show some auxiliary decay bounds provided (4.3) holds. In particular, we start with proofs that, for $k \leq j - 1$:

$$(4.5) \quad \|\langle \underline{u}_c \rangle^{\frac{1}{2}} r^{-\frac{1}{2}} S^{\leq 7} W_k\|_{\ell_\tau^2 \ell_R^2 \leq c\tau/2 L_t^\infty L_r^\infty (C_\tau^{c,R})} \lesssim \varepsilon,$$

$$(4.6) \quad \|\langle u_c \rangle^{\frac{1}{2}} \langle r \rangle^{-\frac{1}{2}} S^{\leq 7} W_k\|_{\ell_\tau^2 \ell_{U_c \leq c\tau/4}^2 L_t^\infty L_r^\infty (C_\tau^{c,U_c})} \lesssim \varepsilon,$$

$$(4.7) \quad \|\langle \underline{u} \rangle S^{\leq 5} \partial V_k\|_{\ell_R^\infty \ell_{\tau \geq 2R}^2 L_t^\infty L_r^\infty (C_\tau^{1,R})} \lesssim \varepsilon,$$

$$(4.8) \quad \|\langle u \rangle S^{\leq 5} \partial V_k\|_{\ell_U^\infty \ell_{\tau \geq 4U}^2 L_t^\infty L_r^\infty (C_\tau^{1,U})} \lesssim \varepsilon.$$

We shall provide the proofs of these in order. The first two are easy corollaries of Corollary 2.2.

Proof of (4.5). This is an immediate consequence of (2.6) and (4.3). We apply (2.6) to $S^{\leq 7} W_k$ and note that if W_k is odd then so is SW_k . Then, indeed,

$$\begin{aligned} & \|\langle \underline{u}_c \rangle^{\frac{1}{2}} r^{-\frac{1}{2}} S^{\leq 7} W_k\|_{\ell_\tau^2 \ell_{R \leq c\tau/2}^2 L_t^\infty L_r^\infty (C_\tau^{c,R})} \\ & \lesssim \|\langle r \rangle^{-1} S^{\leq 9} W_k\|_{\ell_\tau^2 \ell_{R \leq c\tau/2}^2 L_t^2 L_r^2 (\tilde{C}_\tau^{c,R})} + \|S^{\leq 8} \partial_{\underline{u}_c} W_k\|_{\ell_\tau^2 \ell_{R \leq c\tau/2}^2 L_t^2 L_r^2 (\tilde{C}_\tau^{c,R})} \\ & \lesssim (XII)_k + (XI)_k, \end{aligned}$$

and the bound follows from (4.3). \square

Proof of (4.6). Applying (2.7), we have

$$\begin{aligned} & \|\langle u_c \rangle^{\frac{1}{2}} \langle r \rangle^{-\frac{1}{2}} S^{\leq 7} W_k\|_{\ell_\tau^2 \ell_{U_c \leq c\tau/4}^2 L_t^\infty L_r^\infty (C_\tau^{c,U_c})} \\ & \lesssim \|\langle r \rangle^{-1} S^{\leq 9} W_k\|_{\ell_\tau^2 \ell_{U_c \leq c\tau/4}^2 L_t^2 L_r^2 (\tilde{C}_\tau^{c,U_c})} + \|S^{\leq 8} \partial_{\underline{u}_c} W_k\|_{\ell_\tau^2 \ell_{U_c \leq c\tau/4}^2 L_t^2 L_r^2 (\tilde{C}_\tau^{c,U_c})} \\ & \lesssim (XII)_k + (XI)_k. \end{aligned}$$

And (4.6) is then a direct consequence of (4.3). \square

We now proceed to the pointwise bounds for V_k with $k \leq j - 1$ using Lemma 2.3. We first consider the bound away from the light cone.

Proof of (4.7). Applying (2.10), and commuting S with \square and ∂ , we have

$$\begin{aligned} & \|\langle \underline{u} \rangle S^{\leq 5} \partial V_k\|_{\ell_R^\infty \ell_{\tau \geq 2R}^2 L_t^\infty L_r^\infty (C_\tau^{1,R})} \\ & \lesssim \|\langle r \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial V_k\|_{\ell_R^\infty \ell_{\tau \geq 2R}^2 L_t^2 L_r^2 (\tilde{C}_\tau^{1,R})} + \|\langle r \rangle^{\frac{1}{4}} r^{\frac{1}{4}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 6} \square V_k\|_{\ell_R^\infty \ell_{\tau \geq 2R}^2 L_t^2 L_r^2 (\tilde{C}_\tau^{1,R})}. \end{aligned}$$

Splitting ∂ into $(\partial_u, \partial_{\underline{u}})$, we bound the first term by

$$\|\langle r \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_k\|_{\ell_R^\infty \ell_{\tau \geq 2R}^2 L_t^2 L_r^2 (\tilde{C}_\tau^{1,R})} + \|\langle r \rangle^{-\frac{1}{2}} \langle u \rangle^{\frac{1}{2}} S^{\leq 7} \partial_u V_k\|_{\ell_R^\infty \ell_{\tau \geq 2R}^2 L_t^2 L_r^2 (\tilde{C}_\tau^{1,R})} \lesssim (III)_k + (I)_k,$$

where we exploit that $u \approx \underline{u}$ on $\tilde{C}_\tau^{1,R}$ regions with $R \leq \tau/2$. This term is $\mathcal{O}(\varepsilon)$ by (4.3).

Moving to the second term, we crudely apply the product rule to $S^{\leq 6} \square V_k$, giving

$$(4.9) \quad |S^{\leq 6} \square V_k| \leq r^{-1} |S^{\leq 6} W_{k-1}| |S^{\leq 6} \partial_t V_{k-1}|,$$

where we note that $|Sr^{-1}| = r^{-1}$. On $\tilde{C}_\tau^{1,R}$ with $R \leq \tau/2$, we have

$$\langle r \rangle \lesssim \langle \underline{u}_c \rangle \approx \langle \underline{u} \rangle \approx \langle u \rangle \approx \tau.$$

This allows us to bound:

$$\begin{aligned} & \|\langle r \rangle^{\frac{1}{4}} r^{\frac{1}{4}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 6} \square V_k\|_{\ell_R^\infty \ell_{\tau \geq 2R}^2 L_t^2 L_r^2 (\tilde{C}_\tau^{1,R})} \lesssim \|r^{-\frac{1}{2}} \langle \underline{u}_c \rangle^{\frac{1}{2}} S^{\leq 6} W_{k-1}\|_{\ell_\tau^\infty \ell_{R \leq \tau/2}^2 L_t^\infty L_r^\infty (\tilde{C}_\tau^{1,R})} \\ & \times \left(\|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 6} \partial_{\underline{u}} V_{k-1}\|_{\ell_R^\infty \ell_{\tau \geq 2R}^2 L_t^2 L_r^2 (\tilde{C}_\tau^{1,R})} + \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} \langle u \rangle^{\frac{1}{2}} S^{\leq 6} \partial_u V_{k-1}\|_{\ell_R^\infty \ell_{\tau \geq 2R}^2 L_t^2 L_r^2 (\tilde{C}_\tau^{1,R})} \right). \end{aligned}$$

We note that since $c > 1$, each $\tilde{C}_\tau^{1,R}$ with $R \leq \tau/2$ is contained in a finite number of regions $C_{\tilde{\tau}}^{c,\tilde{R}}$ with $\tilde{R} \leq c\tilde{\tau}/2$. This allows us to apply (4.5) to the first factor in the right side, which shows that this is $\lesssim \varepsilon[(III)_{k-1} + (I)_{k-1}]$. The desired bound is then a consequence of (4.3). \square

We finally consider the pointwise bound for ∂V_k away from the light cone.

Proof of (4.8). Fixing U and τ with $U \leq \tau/4$, we start with an application of (2.11), which gives that

$$(4.10) \quad \|\langle u \rangle S^{\leq 5} \partial V_k\|_{L_t^\infty L_r^\infty(\tilde{C}_\tau^{1,U})} \lesssim \|\langle u \rangle^{\frac{1}{2}} \langle \underline{u} \rangle^{-\frac{1}{2}} S^{\leq 7} \partial V_k\|_{L_t^2 L_r^2(\tilde{C}_\tau^{1,U})} + \|\langle u \rangle^{\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 6} \square V_k\|_{L_t^2 L_r^2(\tilde{C}_\tau^{1,U})}.$$

For the first term, we have

$$\begin{aligned} \|\langle u \rangle^{\frac{1}{2}} \langle \underline{u} \rangle^{-\frac{1}{2}} S^{\leq 7} \partial V_k\|_{L_t^2 L_r^2(\tilde{C}_\tau^{1,U})} &\lesssim \|\langle r \rangle^{-\frac{1}{2}} \langle u \rangle^{\frac{1}{2}} S^{\leq 7} \partial_u V_k\|_{L_t^2 L_r^2(\tilde{C}_\tau^{1,U})} + \|\langle r \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_k\|_{L_t^2 L_r^2(\tilde{C}_\tau^{1,U})} \\ &\leq (I)_k + (III)_k, \end{aligned}$$

and the bound is an immediate consequence of (4.3).

For the second term in (4.10), we use (4.9). In order to use (4.5) and (4.6), we will consider separately when $\tilde{C}_\tau^{1,U}$ intersects $\bigcup_{R \leq c\tilde{\tau}/2} C_{\tilde{\tau}}^{c,R}$ and $\bigcup_{U_c \leq c\tilde{\tau}/4} C_{\tilde{\tau}}^{c,U_c}$. Away from the speed c light cone, we have

$$\begin{aligned} \|\langle u \rangle^{\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 6} \square V_k\|_{L_t^2 L_r^2(\tilde{C}_\tau^{1,U} \cap (\bigcup_{\tilde{r}} \bigcup_{R \leq c\tilde{\tau}/2} C_{\tilde{\tau}}^{c,R}))} &\lesssim \|\langle \underline{u} \rangle^{\frac{1}{2}} r^{-\frac{1}{2}} S^{\leq 6} W_{k-1}\|_{\ell_{\tilde{r}}^\infty \ell_{R \leq c\tilde{\tau}/2}^\infty L_t^\infty L_r^\infty(C_{\tilde{\tau}}^{c,R})} \\ &\times \left(\|\langle r \rangle^{-\frac{1}{2}} \langle u \rangle^{\frac{1}{2}} S^{\leq 6} \partial_u V_{k-1}\|_{L_t^2 L_r^2(\tilde{C}_\tau^{1,U})} + \|\langle r \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 6} \partial_{\underline{u}} V_{k-1}\|_{L_t^2 L_r^2(\tilde{C}_\tau^{1,U})} \right), \end{aligned}$$

as $\tilde{C}_\tau^{1,U}$ only intersects $\bigcup_{R \leq c\tilde{\tau}/2} C_{\tilde{\tau}}^{c,R}$ for a finite number of $\tilde{\tau}$. Since $\tilde{C}_\tau^{1,U}$ (for fixed U and τ) is contained in a finite number of dyadic regions $C_{\tilde{\tau}}^{1,R}$, it follows, using (4.5) that this is

$$\lesssim \varepsilon[(I)_{k-1} + (III)_{k-1}].$$

After applying (4.3), supremums can be taken over U and τ to obtain the desired result.

Near the speed c light cone, we instead see

$$\begin{aligned} \|\langle u \rangle^{\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 6} \square V_k\|_{L_t^2 L_r^2(\tilde{C}_\tau^{1,U} \cap (\bigcup_{\tilde{r}} \bigcup_{U_c \leq c\tilde{\tau}/4} C_{\tilde{\tau}}^{c,U_c}))} &\lesssim \|\langle u_c \rangle^{\frac{1}{2}} r^{-\frac{1}{2}} S^{\leq 6} W_{k-1}\|_{\ell_{\tilde{r}}^2 \ell_{U_c \leq c\tilde{\tau}/4}^2 L_t^\infty L_r^\infty(C_{\tilde{\tau}}^{c,U_c})} \\ &\times \left(\|\langle u_c \rangle^{-\frac{1}{2}} \langle u \rangle^{\frac{1}{2}} S^{\leq 6} \partial_u V_{k-1}\|_{\ell_{\tilde{r}}^\infty \ell_{U_c}^\infty L_t^2 L_r^2(C_{\tilde{\tau}}^{c,U_c})} + \|\langle u_c \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 6} \partial_{\underline{u}} V_{k-1}\|_{\ell_{\tilde{r}}^\infty \ell_{U_c}^\infty L_t^2 L_r^2(\tilde{C}_\tau^{c,U_c})} \right). \end{aligned}$$

By (4.6), this is bounded by

$$\varepsilon[(II)_{k-1} + (IV)_{k-1}].$$

Thus, an application of (4.3) and taking supremums over U , τ yields the result. \square

4.1.2. *Bound on terms $(I)_j, \dots, (V)_j$.* By applying (3.7) (with $p = 1$) to $S^{\leq 7} V_j$, we see that

$$(4.11) \quad (I)_j^2 + \dots + (V)_j^2 \leq (C_0 \varepsilon)^2 + C \int_4^{T_\varepsilon} \int_0^\infty \langle \underline{u} \rangle |S^{\leq 7} \square V_j| |\partial_{\underline{u}} S^{\leq 7} V_j| dr dt \\ + C \int_4^{T_\varepsilon} \int_0^\infty \langle u \rangle |S^{\leq 7} \square V_j| |\partial_u S^{\leq 7} V_j| dr dt.$$

We need to show that the latter two terms are bounded by

$$(4.12) \quad C \varepsilon^2 (\log(2 + T_\varepsilon))^{\frac{1}{2}} M_j.$$

Due to finite speed of propagation, we note $\partial_t V_{j-1}$, and hence $\square V_j$, vanishes for $r \geq t - 3$.

We shall first decompose these integrals using (2.2) at speed c and note that $C^1 \subset C^c$. For the first of the integrals in (4.11), we have

$$(4.13) \quad \|\langle \underline{u} \rangle S^{\leq 7} \square V_j \cdot S^{\leq 7} \partial_{\underline{u}} V_j\|_{L_t^1 L_r^1(C^1)} \lesssim \|\langle \underline{u} \rangle S^{\leq 7} \square V_j \cdot S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_r^1 L_t^1 L_r^1(C_{\tilde{\tau}}^{c,R=1})} \\ + \|\langle \underline{u} \rangle S^{\leq 7} \square V_j \cdot S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_r^1 \ell_{1 < R \leq c\tilde{\tau}/2}^1 L_t^1 L_r^1(C_{\tilde{\tau}}^{c,R})} + \|\langle \underline{u} \rangle S^{\leq 7} \square V_j \cdot S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_r^1 \ell_{U_c \leq c\tilde{\tau}/4}^1 L_t^1 L_r^1(C_{\tilde{\tau}}^{c,U_c})}.$$

A naive application of the product rule gives that

$$|S^{\leq 7} \square V_j| \leq r^{-1} |S^{\leq 7} W_{j-1}| \left(|S^{\leq 7} \partial_{\underline{u}} V_{j-1}| + |S^{\leq 7} \partial_u V_{j-1}| \right),$$

which we will apply in each instance.

For the first term in the right side of (4.13), since $\langle \underline{u} \rangle \approx \langle u \rangle$ on $C_\tau^{c,R=1}$, we have

$$\begin{aligned} & \|\langle \underline{u} \rangle S^{\leq 7} \square V_j \cdot S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_t^1 L_t^1 L_r^1(C_\tau^{c,R=1})} \lesssim \|\langle \underline{u}_c \rangle^{\frac{1}{2}} r^{-\frac{1}{2}} S^{\leq 7} W_{j-1}\|_{\ell_t^2 L_t^\infty L_r^\infty(C_\tau^{c,R=1})} \\ & \quad \times \left(\|\langle r \rangle^{-\frac{1}{4}} r^{-\frac{1}{4}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_{j-1}\|_{\ell_t^\infty L_t^2 L_r^2(C_\tau^{c,R=1})} + \|\langle r \rangle^{-\frac{1}{4}} r^{-\frac{1}{4}} \langle u \rangle^{\frac{1}{2}} S^{\leq 7} \partial_u V_{j-1}\|_{\ell_t^\infty L_t^2 L_r^2(C_\tau^{c,R=1})} \right) \\ & \quad \times \|\langle r \rangle^{-\frac{1}{4}} r^{-\frac{1}{4}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_t^2 L_t^2 L_r^2(C_\tau^{c,R=1})}, \end{aligned}$$

which, using (4.5), is bounded by $C\varepsilon[(III)_{j-1} + (I)_{j-1}](III)_j$. Since we have (4.3), this term is controlled by (4.12) as desired.

For the next term in (4.13), we use the Schwarz inequality and the facts that $\langle \underline{u} \rangle \approx \tau$ and $\langle r \rangle \lesssim \tau$ on $C_\tau^{c,R}$ with $R \leq c\tau/2$ to see that

$$\begin{aligned} (4.14) \quad & \|\langle \underline{u} \rangle S^{\leq 7} \square V_j \cdot S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_t^1 \ell_{1 < R \leq c\tau/2}^1 L_t^1 L_r^1(C_\tau^{c,R})} \\ & \lesssim \|\langle \underline{u}_c \rangle^{\frac{1}{2}} r^{-\frac{1}{2}} S^{\leq 7} W_{j-1}\|_{\ell_t^2 \ell_{1 < R \leq c\tau/2}^2 L_t^\infty L_r^\infty(C_\tau^{c,R})} \|\langle r \rangle^{-\frac{1}{4}} r^{-\frac{1}{4}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_{j-1}\|_{\ell_t^\infty \ell_R^\infty L_t^2 L_r^2(C_\tau^{c,R})} \\ & \quad \times \|\langle r \rangle^{-\frac{1}{4}} r^{-\frac{1}{4}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_t^2 \ell_{1 < R \leq c\tau/2}^2 L_t^2 L_r^2(C_\tau^{c,R})} \\ & \quad + \|\langle \underline{u}_c \rangle^{\frac{1}{2}} r^{-\frac{1}{2}} S^{\leq 7} W_{j-1}\|_{\ell_t^2 \ell_{1 < R \leq c\tau/2}^2 L_t^\infty L_r^\infty(C_\tau^{c,R})} \|\langle r \rangle^{-\frac{1}{2}} \langle u \rangle^{\frac{1}{2}} S^{\leq 7} \partial_u V_{j-1}\|_{\ell_t^\infty \ell_R^\infty L_t^2 L_r^2(C_\tau^{c,R})} \\ & \quad \times \|\langle u \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_t^2 \ell_{1 < R \leq c\tau/2}^2 L_t^2 L_r^2(C_\tau^{c,R})}. \end{aligned}$$

For the last factor in each term, we sum back up and re-decompose in terms of speed 1 regions to see

$$\begin{aligned} (4.15) \quad & \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_t^2 \ell_{1 < R \leq c\tau/2}^2 L_t^2 L_r^2(C_\tau^{c,R})} + \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_t^2 \ell_{U_c \leq c\tau/4}^2 L_t^2 L_r^2(C_\tau^{c,U_c})} \\ & \lesssim \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_j\|_{L_t^2 L_r^2(C^1)} \\ & \lesssim (\log(2 + T_\varepsilon))^{\frac{1}{2}} \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_t^\infty \ell_R^2 L_t^2 L_r^2(C_\tau^{c,R})} \end{aligned}$$

and similarly

$$\begin{aligned} (4.16) \quad & \|\langle u \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_t^2 \ell_{1 < R \leq c\tau/2}^2 L_t^2 L_r^2(C_\tau^{c,R})} + \|\langle u \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_t^2 \ell_{U_c \leq c\tau/4}^2 L_t^2 L_r^2(C_\tau^{c,U_c})} \\ & \lesssim (\log(2 + T_\varepsilon))^{\frac{1}{2}} \|\langle u \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_t^\infty \ell_R^2 L_t^2 L_r^2(C_\tau^{c,U})}. \end{aligned}$$

Applying (4.5), it then follows that the left side of (4.14) is bounded by

$$C\varepsilon(III)_{j-1} (\log(2 + T_\varepsilon))^{\frac{1}{2}} (III)_j + C\varepsilon(I)_{j-1} (\log(2 + T_\varepsilon))^{\frac{1}{2}} (V)_j,$$

which, owing to (4.3), is controlled by (4.12).

The last term in (4.13) is handled similarly, but we now must rely upon (4.6). Here we use the fact that $r \approx \tau \approx \langle \underline{u} \rangle$ on C_τ^{c,U_c} with $1 \leq U_c \leq c\tau/4$. Indeed,

$$\begin{aligned} (4.17) \quad & \|\langle \underline{u} \rangle S^{\leq 7} \square V_j \cdot S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_t^1 \ell_{U_c \leq c\tau/4}^1 L_t^1 L_r^1(C_\tau^{c,U_c})} \\ & \lesssim \|\langle u_c \rangle^{\frac{1}{2}} r^{-\frac{1}{2}} S^{\leq 7} W_{j-1}\|_{\ell_t^2 \ell_{U_c \leq c\tau/4}^2 L_t^\infty L_r^\infty(C_\tau^{c,U_c})} \|\langle u_c \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_{j-1}\|_{\ell_t^\infty \ell_{U_c}^\infty L_t^2 L_r^2(C_\tau^{c,U_c})} \\ & \quad \times \|\langle r \rangle^{-\frac{1}{4}} r^{-\frac{1}{4}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_t^2 \ell_{U_c \leq c\tau/4}^2 L_t^2 L_r^2(C_\tau^{c,U_c})} \\ & \quad + \|\langle u_c \rangle^{\frac{1}{2}} r^{-\frac{1}{2}} S^{\leq 7} W_{j-1}\|_{\ell_t^2 \ell_{U_c \leq c\tau/4}^2 L_t^\infty L_r^\infty(C_\tau^{c,U_c})} \|\langle u_c \rangle^{-\frac{1}{2}} \langle u \rangle^{\frac{1}{2}} S^{\leq 7} \partial_u V_{j-1}\|_{\ell_t^\infty \ell_{U_c}^\infty L_t^2 L_r^2(C_\tau^{c,U_c})} \\ & \quad \times \|\langle u \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_j\|_{\ell_t^2 \ell_{U_c \leq c\tau/4}^2 L_t^2 L_r^2(C_\tau^{c,U_c})}. \end{aligned}$$

Using (4.6), (4.15), and (4.16), it follows that this is

$$\lesssim \varepsilon(IV)_{j-1} (\log(2 + T_\varepsilon))^{\frac{1}{2}} (III)_j + \varepsilon(IV)_{j-1} (\log(2 + T_\varepsilon))^{\frac{1}{2}} (V)_j.$$

The inductive hypothesis (4.3) then gives that this is bounded by (4.12) as desired.

The same strategy bounds the second integral in (4.11). In fact, since $\langle u \rangle \lesssim \langle \underline{u} \rangle$ in all regions, the argument can be simplified. Indeed, we have

$$(4.18) \quad \begin{aligned} & \|\langle u \rangle S^{\leq 7} \square V_j \cdot S^{\leq 7} \partial_u V_j\|_{\ell_r^1 \ell_R^1 \leq c\tau/2 L_t^1 L_r^1(C_\tau^{c,R})} \lesssim \|\langle \tau \rangle^{\frac{1}{2}} r^{-\frac{1}{2}} S^{\leq 7} W_{j-1}\|_{\ell_r^2 \ell_R^2 \leq c\tau/2 L_t^\infty L_r^\infty(C_\tau^{c,R})} \\ & \times \left(\|\langle r \rangle^{-\frac{1}{4}} r^{-\frac{1}{4}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_{j-1}\|_{\ell_r^\infty \ell_R^\infty L_t^2 L_r^2(C_\tau^{c,R})} + \|\langle r \rangle^{-\frac{1}{4}} r^{-\frac{1}{4}} \langle u \rangle^{\frac{1}{2}} S^{\leq 7} \partial_u V_{j-1}\|_{\ell_r^\infty \ell_R^\infty L_t^2 L_r^2(C_\tau^{c,R})} \right) \\ & \times \|\langle r \rangle^{-\frac{1}{4}} r^{-\frac{1}{4}} \langle u \rangle^{\frac{1}{2}} S^{\leq 7} \partial_u V_j\|_{\ell_r^2 \ell_R^2 \leq c\tau/2 L_t^2 L_r^2(C_\tau^{c,R})}. \end{aligned}$$

Using a direct analog of (4.15) and (4.5), this is

$$\lesssim \varepsilon((III)_{j-1} + (I)_{j-1})(\log(2 + T_\varepsilon))^{\frac{1}{2}}(I)_j,$$

which is in turn controlled by (4.12).

And

$$(4.19) \quad \begin{aligned} & \|\langle u \rangle S^{\leq 7} \square V_j \cdot S^{\leq 7} \partial_u V_j\|_{\ell_r^1 \ell_{U_c}^1 \leq c\tau/4 L_t^1 L_r^1(C_\tau^{c,U_c})} \lesssim \|\langle u_c \rangle^{\frac{1}{2}} r^{-\frac{1}{2}} S^{\leq 7} W_{j-1}\|_{\ell_r^2 \ell_{U_c}^2 \leq c\tau/4 L_t^\infty L_r^\infty(C_\tau^{c,U_c})} \\ & \times \left(\|\langle u_c \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} V_{j-1}\|_{\ell_r^\infty \ell_{U_c}^\infty L_t^2 L_r^2(C_\tau^{c,U_c})} + \|\langle u_c \rangle^{-\frac{1}{2}} \langle u \rangle^{\frac{1}{2}} S^{\leq 7} \partial_u V_{j-1}\|_{\ell_r^\infty \ell_{U_c}^\infty L_t^2 L_r^2(C_\tau^{c,U_c})} \right) \\ & \times \|\langle r \rangle^{-\frac{1}{4}} r^{-\frac{1}{4}} \langle u \rangle^{\frac{1}{2}} S^{\leq 7} \partial_u V_j\|_{\ell_r^2 \ell_{U_c}^2 \leq c\tau/4 L_t^2 L_r^2(C_\tau^{c,U_c})}. \end{aligned}$$

Using the analog of (4.16) and (4.6), this is

$$\lesssim \varepsilon((IV)_{j-1} + (II)_{j-1})(\log(2 + T_\varepsilon))^{\frac{1}{2}}(I)_j,$$

which is in turn controlled by (4.12) after applying (4.3). This completes the proof of the boundedness of terms $(I)_j, \dots, (V)_j$.

4.1.3. Bound on terms $(VI)_j, \dots, (VIII)_j$. We now proceed to considering the high order energy bounds for V_j . We begin by applying (3.7) (with $p = 0$) to $S^{\leq 10} V_j$ to obtain

$$(VI)_j^2 + (VII)_j^2 + (VIII)_j^2 \leq (C_0 \varepsilon)^2 + C \int_4^{T_\varepsilon} \int |\square S^{\leq 10} V_j| |\partial S^{\leq 10} V_j| dr dt.$$

Again, we need to show that the last term is bounded by (4.12). This time, however, we need to apply the product rule more carefully. We note that on each term, there will be one factor with no more than half of the 10 total vector fields. As such, we have

$$|S^{\leq 10} \square V_j| \leq r^{-1} |S^{\leq 5} W_{j-1}| |S^{\leq 10} \partial V_{j-1}| + r^{-1} |S^{\leq 10} W_{j-1}| |S^{\leq 5} \partial V_{j-1}|.$$

When W is lower order, we shall initially decompose in speed c regions. On $C_\tau^{c,R}$ regions, since $\langle r \rangle \lesssim \tau$, we bound

$$(4.20) \quad \begin{aligned} & \|r^{-1} S^{\leq 5} W_{j-1} S^{\leq 10} \partial V_{j-1} S^{\leq 10} \partial V_j\|_{\ell_r^1 \ell_R^1 \leq c\tau/2 L_t^1 L_r^1(C_\tau^{c,R})} \lesssim \|\langle \underline{u}_c \rangle^{\frac{1}{2}} r^{-\frac{1}{2}} S^{\leq 5} W_{j-1}\|_{\ell_r^2 \ell_R^2 \leq c\tau/2 L_t^\infty L_r^\infty(C_\tau^{c,R})} \\ & \times \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} S^{\leq 10} \partial V_{j-1}\|_{\ell_r^\infty \ell_R^\infty L_t^2 L_r^2(C_\tau^{c,R})} \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} S^{\leq 10} \partial V_j\|_{\ell_r^2 \ell_R^2 \leq c\tau/2 L_t^2 L_r^2(C_\tau^{c,R})} \end{aligned}$$

Arguing as in (4.15), the last factor satisfies

$$(4.21) \quad \begin{aligned} & \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} S^{\leq 10} \partial V_j\|_{\ell_r^2 \ell_R^2 \leq c\tau/2 L_t^2 L_r^2(C_\tau^{c,R})} + \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} S^{\leq 10} \partial V_j\|_{\ell_r^2 \ell_{U_c}^2 \leq c\tau/4 L_t^2 L_r^2(C_\tau^{c,U_c})} \\ & \lesssim (\log(2 + T_\varepsilon))^{\frac{1}{2}} \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} S^{\leq 10} \partial V_j\|_{\ell_R^\infty \ell_r^2 L_t^2 L_r^2(C_\tau^{1,R})}. \end{aligned}$$

Using this and (4.5), we see that the left side of (4.20) is

$$\lesssim \varepsilon(VI)_{j-1} (\log(2 + T_\varepsilon))^{\frac{1}{2}} (VI)_j,$$

and thus due to (4.3) is controlled by (4.12). On the C_τ^{c,U_c} regions, we instead have

$$(4.22) \quad \begin{aligned} & \|r^{-1}S^{\leq 5}W_{j-1}S^{\leq 10}\partial V_{j-1}S^{\leq 10}\partial V_j\|_{\ell_\tau^1 \ell_{U_c \leq c\tau/4}^1 L_t^1 L_r^1(C_\tau^{c,U_c})} \lesssim \|\langle u_c \rangle^{\frac{1}{2}} \langle r \rangle^{-\frac{1}{2}} S^{\leq 5}W_{j-1}\|_{\ell_\tau^2 \ell_{U_c \leq c\tau/4}^2 L_t^\infty L_r^\infty(C_\tau^{c,U_c})} \\ & \quad \times \|\langle u_c \rangle^{-\frac{1}{2}} S^{\leq 10}\partial V_{j-1}\|_{\ell_\tau^\infty \ell_{U_c}^\infty L_t^2 L_r^2(C_\tau^{c,U_c})} \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} S^{\leq 10}\partial V_j\|_{\ell_\tau^2 \ell_{U_c \leq c\tau/4}^2 L_t^2 L_r^2(C_\tau^{c,U_c})}. \end{aligned}$$

By (4.6) and (4.21), this is

$$\lesssim \varepsilon(VII)_{j-1}(\log(2+T_\varepsilon))^{\frac{1}{2}}(VI)_j,$$

which is in turn bounded by (4.12) upon applying (4.3).

When ∂V is lower order, we instead use the speed 1 decomposition. Away from the light cone, since $\langle r \rangle \leq \tau$ on $C_\tau^{1,R}$ with $R \leq \tau/2$, we obtain

$$\begin{aligned} & \|r^{-1}S^{\leq 5}\partial V_{j-1}S^{\leq 10}W_{j-1}S^{\leq 10}\partial V_j\|_{\ell_R^1 \ell_{\tau \geq 2R}^1 L_t^1 L_r^1(C_\tau^{1,R})} \\ & \lesssim \|\langle \underline{u} \rangle S^{\leq 5}\partial V_{j-1}\|_{\ell_R^\infty \ell_{\tau \geq 2R}^\infty L_t^\infty L_r^\infty(C_\tau^{1,R})} \|r^{-1}S^{\leq 10}W_{j-1}\|_{\ell_R^2 \ell_{\tau \geq 2R}^2 L_t^2 L_r^2(C_\tau^{1,R})} \\ & \quad \times \|\langle r \rangle^{-\frac{1}{2}} S^{\leq 10}\partial V_j\|_{\ell_R^\infty \ell_{\tau \geq 2R}^\infty L_t^2 L_r^2(C_\tau^{1,R})}. \end{aligned}$$

Here we have used the additional power $\langle \underline{u} \rangle^{-\frac{1}{2}} \leq \langle r \rangle^{-\frac{1}{2}}$ to control the remaining dyadic sum over R . We may apply (4.7) to then see that this is $\lesssim \varepsilon(XI)_{j-1}(VI)_j$, which by (4.3) is better than the required bound (4.12).

When ∂V is lower order and we are near the speed 1 light cone, we instead have

$$\begin{aligned} & \|r^{-1}S^{\leq 5}\partial V_{j-1}S^{\leq 10}W_{j-1}S^{\leq 10}\partial V_j\|_{\ell_U^1 \ell_{\tau \geq 4U}^1 L_t^1 L_r^1(C_\tau^{1,U})} \\ & \lesssim \|\langle u \rangle S^{\leq 5}\partial V_{j-1}\|_{\ell_\tau^\infty \ell_{U \geq \tau/4}^\infty L_t^\infty L_r^\infty(C_\tau^{1,U})} \|\langle u \rangle^{-\frac{1}{2}} r^{-\frac{1}{2}} S^{\leq 10}W_{j-1}\|_{\ell_U^\infty \ell_{\tau \geq 4U}^\infty L_t^2 L_r^2(C_\tau^{1,U})} \\ & \quad \times \|\langle r \rangle^{-\frac{1}{2}} S^{\leq 10}\partial V_j\|_{\ell_\tau^2 \ell_{U \leq \tau/4}^2 L_t^2 L_r^2(C_\tau^{1,U})}. \end{aligned}$$

Here we have again gained ℓ_U^2 summability from the extra factor of $\langle u \rangle^{-\frac{1}{2}}$. We note that the last factor satisfies

$$(4.23) \quad \begin{aligned} & \|\langle r \rangle^{-\frac{1}{2}} S^{\leq 10}\partial V_j\|_{\ell_\tau^2 \ell_{U \leq \tau/4}^2 L_t^2 L_r^2(C_\tau^{1,U})} \lesssim \|\langle r \rangle^{-\frac{1}{2}} S^{\leq 10}\partial V_j\|_{\ell_R^2 \ell_{\tau \geq 4R}^2 L_t^2 L_r^2(C_\tau^{1,R})} \\ & \lesssim (\log(2+T_\varepsilon))^{\frac{1}{2}} \|\langle r \rangle^{-\frac{1}{2}} S^{\leq 10}\partial V_j\|_{\ell_R^\infty \ell_{\tau \geq 4R}^\infty L_t^2 L_r^2(C_\tau^{1,R})}. \end{aligned}$$

It follows from (4.8) that

$$\|r^{-1}S^{\leq 5}\partial V_{j-1}S^{\leq 10}W_{j-1}S^{\leq 10}\partial V_j\|_{\ell_U^1 \ell_{\tau \geq 4U}^1 L_t^1 L_r^1(C_\tau^{1,U})} \lesssim \varepsilon(X)_{j-1}(\log(2+T_\varepsilon))^{\frac{1}{2}}(VI)_j.$$

This gives the bound by (4.12) upon using (4.3).

4.1.4. *Bound on terms $(IX)_j, \dots, (XII)_j$.* Applying (3.10), (3.11), and (3.13) to $S^{\leq 10}W_j$ gives

$$(IX)_j^2 + \dots + (XII)_j^2 \leq (C_0\varepsilon)^2 + C \int_4^{T_\varepsilon} \int r |\square_c S^{\leq 10}W_j| |\partial_{\underline{u}_c} S^{\leq 10}W_j| dr dt.$$

We again seek to show that the last term is bounded by (4.12). The decomposition of the integral will be at speed 1 throughout. The product rule yields

$$|S^{\leq 10}\square_c W_j| \leq r^{-1} |S^{\leq 5}\partial V_{j-1}| |S^{\leq 10}\partial V_{j-1}|.$$

Away from the speed 1 light cone, we have

$$\begin{aligned} & \|rS^{\leq 10}\square_c W_j \cdot S^{\leq 10}\partial_{\underline{u}_c} W_j\|_{\ell_\tau^1 \ell_{R \leq \tau/2}^1 L_t^1 L_r^1(C_\tau^{1,R})} \lesssim \|\langle \underline{u} \rangle S^{\leq 5}\partial V_{j-1}\|_{\ell_R^\infty \ell_{\tau \geq 2R}^\infty L_t^\infty L_r^\infty(C_\tau^{1,R})} \\ & \quad \times \|\langle r \rangle^{-\frac{1}{2}} S^{\leq 10}\partial V_{j-1}\|_{\ell_R^\infty \ell_{\tau \geq 2R}^\infty L_t^2 L_r^2(C_\tau^{1,R})} \|S^{\leq 10}\partial_{\underline{u}_c} W_j\|_{\ell_R^2 \ell_{\tau \geq 2R}^2 L_t^2 L_r^2(C_\tau^{1,R})}. \end{aligned}$$

Here, we have used that $\langle r \rangle \lesssim \langle \underline{u} \rangle$. Moreover, the factor of $\langle \underline{u} \rangle^{-\frac{1}{2}}$ is used to control the remaining ℓ_R^2 summation. As (4.7) gives that this is $\lesssim \varepsilon(VI)_{j-1}(XI)_j$, (4.3) shows that these terms are controlled by (4.12) (without the logarithmic factor, in fact).

Near the speed 1 light cone, we use the Schwarz inequality to bound

$$\begin{aligned} \|rS^{\leq 10} \square_c W_j \cdot S^{\leq 10} \partial_{\underline{u}_c} W_j\|_{\ell_U^1 \ell_{U \leq \tau/4}^1 L_t^1 L_r^1(C_\tau^{1,U})} &\lesssim \|\langle u \rangle S^{\leq 5} \partial V_{j-1}\|_{\ell_{\underline{U}}^\infty \ell_{\tau \geq 4U}^\infty L_t^\infty L_r^\infty(C_\tau^{1,U})} \\ &\times \|\langle r \rangle^{-\frac{1}{2}} S^{\leq 10} \partial V_{j-1}\|_{\ell_{\underline{U}}^2 \ell_t^2 L_r^2(C_\tau^{1,U})} \|r^{\frac{1}{2}} \langle u \rangle^{-\frac{1}{2}} S^{\leq 10} \partial_{\underline{u}_c} W_j\|_{\ell_{\underline{U}}^\infty \ell_t^2 L_r^2(C_\tau^{1,U})}. \end{aligned}$$

The additional $\langle u \rangle^{-\frac{1}{2}}$ allowed us to absorb a square summation in U . Upon applying (4.23) and (4.8), these final terms are

$$\lesssim \varepsilon (\log(2 + T_\varepsilon))^{\frac{1}{2}} (VI)_{j-1}(X)_j.$$

Thus, (4.3) gives that they are controlled by (4.12). This completes the proof of (4.4).

4.2. Convergence. We now show that the sequence $((V_j, W_j))$ converges by showing that it is Cauchy in an appropriate norm. With this aim, we set

(4.24)

$$\begin{aligned} A_j &= \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} \langle u \rangle^{\frac{1}{2}} S^{\leq 7} \partial_u (V_j - V_{j-1})\|_{\ell_R^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{1,R})} + \|\langle u_c \rangle^{-\frac{1}{2}} \langle u \rangle^{\frac{1}{2}} S^{\leq 7} \partial_u (V_j - V_{j-1})\|_{\ell_{\underline{U}_c}^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{c,U_c})} \\ &+ \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} (V_j - V_{j-1})\|_{\ell_R^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{1,R})} + \|\langle u_c \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} (V_j - V_{j-1})\|_{\ell_{\underline{U}_c}^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{c,U_c})} \\ &+ \|\langle u \rangle^{-\frac{1}{2}} \langle \underline{u} \rangle^{\frac{1}{2}} S^{\leq 7} \partial_{\underline{u}} (V_j - V_{j-1})\|_{\ell_{\underline{U}}^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{1,U})} + \|r^{-\frac{1}{4}} \langle r \rangle^{-\frac{1}{4}} S^{\leq 10} \partial (V_j - V_{j-1})\|_{\ell_R^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{1,R})} \\ &+ \|\langle u_c \rangle^{-\frac{1}{2}} S^{\leq 10} \partial (V_j - V_{j-1})\|_{\ell_{\underline{U}_c}^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{c,U_c})} + \|\langle u \rangle^{-\frac{1}{2}} S^{\leq 10} \partial_{\underline{u}} (V_j - V_{j-1})\|_{\ell_{\underline{U}}^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{1,U})} \\ &+ \|r^{\frac{1}{2}} \langle u \rangle^{-\frac{1}{2}} S^{\leq 10} \partial_{\underline{u}_c} (W_j - W_{j-1})\|_{\ell_{\underline{U}}^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{1,U})} + \|r^{-\frac{1}{2}} \langle u \rangle^{-\frac{1}{2}} S^{\leq 10} (W_j - W_{j-1})\|_{\ell_{\underline{U}}^\infty \ell_\tau^2 L_t^2 L_r^2(C_\tau^{1,U})} \\ &+ \|S^{\leq 10} \partial_{\underline{u}_c} (W_j - W_{j-1})\|_{\ell_R^2 \ell_\tau^2 L_t^2 L_r^2(C_\tau^{1,R})} + \|r^{-1} S^{\leq 10} (W_j - W_{j-1})\|_{\ell_R^2 \ell_\tau^2 L_t^2 L_r^2(C_\tau^{c,R})}, \end{aligned}$$

and we will show that, for each j ,

$$(4.25) \quad A_j \leq \frac{1}{2} A_{j-1}.$$

We note that

$$\square(V_j - V_{j-1}) = r^{-1} \left((W_{j-1} - W_{j-2}) \partial_t V_{j-1} + W_{j-2} \partial_t (V_{j-1} - V_{j-2}) \right)$$

and

$$\square_c(W_j - W_{j-1}) = r^{-1} \partial_t (V_{j-1} - V_{j-2}) (\partial_t V_{j-1} + \partial_t V_{j-2}).$$

For this system, we call upon the same arguments as in the proof of (4.4). Doing so yields

$$A_j \lesssim A_{j-1} (M_{j-1} + M_{j-2}),$$

and applying (4.4) with ε sufficiently small completes the proof.

REFERENCES

- [1] S. Alinhac. The null condition for quasilinear wave equations in two space dimensions I. *Invent. Math.*, 145(3):597–618, 2001.
- [2] Benjamin Bechtold and Jason Metcalfe. On a weakly null multiple speed system of wave equations. Forthcoming.
- [3] Demetrios Christodoulou. Global solutions of nonlinear hyperbolic equations for small initial data. *Comm. Pure Appl. Math.*, 39(2):267–282, 1986.
- [4] Mihalis Dafermos and Igor Rodnianski. A new physical-space approach to decay for the wave equation with applications to black hole spacetimes. In *XVIIth International Congress on Mathematical Physics*, pages 421–432. World Sci. Publ., Hackensack, NJ, 2010.
- [5] Kunio Hidano, Chengbo Wang, and Kazuyoshi Yokoyama. On almost global existence and local well posedness for some 3-D quasi-linear wave equations. *Adv. Differential Equations*, 17(3-4):267–306, 2012.
- [6] Markus Keel, Hart F. Smith, and Christopher D. Sogge. Almost global existence for some semilinear wave equations. *J. Anal. Math.*, 87:265–279, 2002. Dedicated to the memory of Thomas H. Wolff.
- [7] Markus Keel, Hart F. Smith, and Christopher D. Sogge. Almost global existence for quasilinear wave equations in three space dimensions. *J. Amer. Math. Soc.*, 17(1):109–153, 2004.
- [8] S. Klainerman. The null condition and global existence to nonlinear wave equations. In *Nonlinear systems of partial differential equations in applied mathematics, Part 1 (Santa Fe, N.M., 1984)*, volume 23 of *Lectures in Appl. Math.*, pages 293–326. Amer. Math. Soc., Providence, RI, 1986.
- [9] Sergiu Klainerman. Uniform decay estimates and the Lorentz invariance of the classical wave equation. *Comm. Pure Appl. Math.*, 38(3):321–332, 1985.

- [10] Sergiu Klainerman and Thomas C. Sideris. On almost global existence for nonrelativistic wave equations in 3D. *Comm. Pure Appl. Math.*, 49(3):307–321, 1996.
- [11] Hans Lindblad. On the lifespan of solutions of nonlinear wave equations with small initial data. *Comm. Pure Appl. Math.*, 43(4):445–472, 1990.
- [12] Jason Metcalfe, Makoto Nakamura, and Christopher D. Sogge. Global existence of quasilinear, nonrelativistic wave equations satisfying the null condition. *Japan. J. Math. (N.S.)*, 31(2):391–472, 2005.
- [13] Jason Metcalfe, Makoto Nakamura, and Christopher D. Sogge. Global existence of solutions to multiple speed systems of quasilinear wave equations in exterior domains. *Forum Math.*, 17(1):133–168, 2005.
- [14] Jason Metcalfe and Taylor Rhoads. Long-time existence for systems of quasilinear wave equations. *Matematica*, 2(1):37–84, 2023.
- [15] Jason Metcalfe and Christopher D. Sogge. Hyperbolic trapped rays and global existence of quasilinear wave equations. *Invent. Math.*, 159(1):75–117, 2005.
- [16] Jason Metcalfe and Christopher D. Sogge. Long-time existence of quasilinear wave equations exterior to star-shaped obstacles via energy methods. *SIAM J. Math. Anal.*, 38(1):188–209, 2006.
- [17] Jason Metcalfe and Christopher D. Sogge. Global existence of null-form wave equations in exterior domains. *Math. Z.*, 256(3):521–549, 2007.
- [18] Jason Metcalfe, Jacob Sterbenz, and Daniel Tataru. Local energy decay for scalar fields on time dependent non-trapping backgrounds. *Amer. J. Math.*, 142(3):821–883, 2020.
- [19] Jason Metcalfe and Alexander Stewart. On a system of weakly null semilinear wave equations. *Anal. Math. Phys.*, 12(5):Paper No. 125, 23, 2022.
- [20] Jason Metcalfe, Daniel Tataru, and Mihai Tohaneanu. Price’s law on nonstationary space-times. *Adv. Math.*, 230(3):995–1028, 2012.
- [21] Cathleen S. Morawetz. The decay of solutions of the exterior initial-boundary value problem for the wave equation. *Comm. Pure Appl. Math.*, 14:561–568, 1961.
- [22] Cathleen S. Morawetz. Time decay for the nonlinear Klein-Gordon equations. *Proc. Roy. Soc. London Ser. A*, 306:291–296, 1968.
- [23] Masahito Ohta. Counterexample to global existence for systems of nonlinear wave equations with different propagation speeds. *Funkcial. Ekvac.*, 46(3):471–477, 2003.
- [24] Thomas C. Sideris. Nonresonance and global existence of prestressed nonlinear elastic waves. *Ann. of Math. (2)*, 151(2):849–874, 2000.
- [25] Thomas C. Sideris and Becca Thomases. Local energy decay for solutions of multi-dimensional isotropic symmetric hyperbolic systems. *J. Hyperbolic Differ. Equ.*, 3(4):673–690, 2006.
- [26] Thomas C. Sideris and Shu-Yi Tu. Global existence for systems of nonlinear wave equations in 3D with multiple speeds. *SIAM J. Math. Anal.*, 33(2):477–488, 2001.
- [27] Christopher D. Sogge. Global existence for nonlinear wave equations with multiple speeds. In *Harmonic analysis at Mount Holyoke (South Hadley, MA, 2001)*, volume 320 of *Contemp. Math.*, pages 353–366. Amer. Math. Soc., Providence, RI, 2003.
- [28] Jacob Sterbenz. Angular regularity and Strichartz estimates for the wave equation. *Int. Math. Res. Not.*, (4):187–231, 2005. With an appendix by Igor Rodnianski.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA, CHAPEL HILL
 Email address: mkoonce@unc.edu, metcalfe@email.unc.edu