An asymptotic on the logarithms of the relative class numbers of imaginary abelian number fields of prime conductor and large degree

Stéphane R. LOUBOUTIN Aix Marseille Université, CNRS, I2M, Marseille, France stephane.louboutin@univ-amu.fr

February 3, 2025

Abstract

An asymptotic on the logarithms of the relative class numbers of the cyclotomic number fields of prime conductors p is known. Here we give an asymptotic on the logarithms of the relative class numbers of the imaginary abelian number fields of prime conductors p and large degrees m = (p-1)/d with $\phi(d) = o(\log p)$. We also show that this asymptotic does not hold true anymore under the only slightly weaker restriction $\phi(d) = O(\log p)$.

1 Introduction

We refer the reader to [Was, Chapters 3, 4 and 11] for more background details. Let \mathbb{K} be an imaginary abelian number field of degree m and prime conductor $p \geq 3$, i.e. let \mathbb{K} be an imaginary subfield of a cyclotomic number field $\mathbb{Q}(\zeta_p)$ (Kronecker-Weber's theorem). Hence mis even and d = (p-1)/m is odd. Its Hasse unit index $Q_{\mathbb{K}}$ is equal to 1 (see e.g. [Lem, Example 5, page 352]). Let $w_{\mathbb{K}}$ be its number of complex roots of unity. Then $w_{\mathbb{K}} = 2p$ if $\mathbb{K} = \mathbb{Q}(\zeta_p)$ but $w_{\mathbb{K}} = 2$ if $\mathbb{K} \subsetneq \mathbb{Q}(\zeta_p)$ (see [Was, Exercise 2.3]). Let \mathbb{K}^+ be the maximal real subfield of \mathbb{K} of degree m/2 fixed by the complex conjugation. The class number $h_{\mathbb{K}^+}$ of \mathbb{K}^+ divides the class number $h_{\mathbb{K}}$ of \mathbb{K} . The relative class number of \mathbb{K} is defined by $h_{\mathbb{K}^+}^- = h_{\mathbb{K}}/h_{\mathbb{K}^+}$. For gcd(t, p) = 1, let σ_t be the \mathbb{Q} -automorphism of $\mathbb{Q}(\zeta_p)$ defined by $\sigma_t(\zeta_p) = \zeta_p^t$. Then $t \mapsto \sigma_t$ a is canonical isomorphic from the multiplicative group $(\mathbb{Z}/p\mathbb{Z})^*$ to the Galois group $Gal(\mathbb{Q}(\zeta_p)/\mathbb{Q})$. Set

$$H := \operatorname{Gal}(\mathbb{Q}(\zeta_p)/\mathbb{K}) = \{ t \in (\mathbb{Z}/p\mathbb{Z})^*; \ \alpha \in \mathbb{K} \Rightarrow \sigma_t(\alpha) = \alpha \},\$$

a subgroup of index m and odd order d = (p-1)/m of the cyclic group $(\mathbb{Z}/p\mathbb{Z})^*$. Let X_p denote the cyclic group of order p-1 of the Dirichlet characters modulo p. Let $X_p^- = \{\chi \in X_p; \chi(-1) = -1\}$ be the set of the (p-1)/2 odd Dirichlet characters modulo p. Now, $-1 \notin H$ and we set

$$X_p^-(H) = \{ \chi \in X_p^-; \ \chi_{/H} = 1 \}.$$

Then $\#X_p^-(H) = m/2$ and we have the relative class number formula

$$h_{\mathbb{K}}^{-} = w_{\mathbb{K}} \left(\frac{p}{4\pi^2}\right)^{m/4} \prod_{\chi \in X_p^-(H)} L(1,\chi).$$

$$\tag{1}$$

⁰2010 Mathematics Subject Classification. Primary. 11R20, 11R29, 11R42.

Key words and phrases. Dirichlet character. L-function. Relative class number. Cyclotomic field.

Using (1) and the arithmetic-geometric mean inequality, we obtain

$$h_{\mathbb{K}}^{-} \le w_{\mathbb{K}} \left(\frac{pM(p,H)}{4\pi^2}\right)^{m/4},\tag{2}$$

where M(p, H) is the mean square value

$$M(p,H) := \frac{1}{\# X_p^-(H)} \sum_{\chi \in X_p^-(H)} |L(1,\chi)|^2.$$

These mean square values have been studied by using Dedekind sums $s(h, p) \in \mathbb{Q}$:

Proposition 1. (See [Lou19, Proposition 2 and Theorem 6]). Let $d \ge 1$ be an odd divisor of p-1, where $p \ge 3$ is an odd prime number. Let H_d be the only subgroup of order d of the multiplicative cyclic group $(\mathbb{Z}/p\mathbb{Z})^*$. Then

$$M(p, H_d) = \frac{\pi^2}{6} \left(1 + \frac{N(H_d, p)}{p} \right),$$

where

$$N(H_d, p) := 12 \sum_{h \in H_d} s(h, p) - p \in \mathbb{Q}.$$

Moreover, for d > 1 the rational number $N(H_d, p)$ is an odd rational integer.

Using Proposition 1 with d = 1 and recalling that $s(1, p) = \frac{(p-1)(p-2)}{12p}$, we recover [Wal]:

$$M(p,\{1\}) := \frac{2}{p-1} \sum_{\chi \in X_p^-} |L(1,\chi)|^2 = \frac{\pi^2}{6} \left(1 - \frac{1}{p}\right) \left(1 - \frac{2}{p}\right) \le \frac{\pi^2}{6} \qquad (p \ge 3).$$
(3)

Using (2) we obtain the upper bound

$$h_{\mathbb{Q}(\zeta_p)}^- \le 2p \left(\frac{pM(p,\{1\})}{4\pi^2}\right)^{(p-1)/4} \le 2p \left(\frac{p}{24}\right)^{(p-1)/4}.$$
(4)

The only other value of d for which an explicit closed formula for $M(p, H_d)$ is known is d = 3:

Proposition 2. (See [Lou16, Theorem 1]). Let $p \equiv 1 \pmod{6}$ be a prime number. Let H_3 be the only subgroup of order 3 of the multiplicative cyclic group $(\mathbb{Z}/p\mathbb{Z})^*$. Let \mathbb{K} be the imaginary subfield of degree (p-1)/3 of the cyclotomic number field $\mathbb{Q}(\zeta_p)$. Then

$$M(p, H_3) = \frac{\pi^2}{6} \left(1 - \frac{1}{p} \right) \le \frac{\pi^2}{6} \text{ and } h_{\mathbb{K}}^- \le 2 \left(\frac{p}{24} \right)^{(p-1)/12}$$
(5)

(compare with (3) and (4), and note the misprint in the exponent in [Lou16, (8)]).

By [LM21, Theorem 1.1], we have the following generalization of (4) and (5):

$$N(p, H_d) = o(1), \ M(p, H_d) = \frac{\pi^2}{6} + o(1) \ \text{and} \ h_{\mathbb{K}}^- \le 2\left(\frac{(1+o(1))p}{24}\right)^{m/4} \tag{6}$$

as p tends to infinity and K runs over the imaginary subfields of $\mathbb{Q}(\zeta_p)$ of large degrees m = (p-1)/d with

$$d \le \frac{\log p}{3\log\log p}.\tag{7}$$

In [MS, Theorem 2.1] it is proved that (6) still holds true under the weaker restriction

$$\phi(d) = o(\log p). \tag{8}$$

Restriction (7) is almost optimal. Indeed, if p runs over the Mersenne primes $p = 2^d - 1$, $d \ge 3$ odd, and H_d is the subgroup of order d of $(\mathbb{Z}/p\mathbb{Z})^*$ generated by 2, then d is prime and

$$N(p, H_d) = 2p - (6d - 3), \ M(p, H_d) = \frac{\pi}{2} \left(1 - \frac{2d - 1}{p}\right) \ \text{and} \ h_{\mathbb{K}}^- \le 2 \left(\frac{p}{8}\right)^{m/4}$$

by [LM23, Theorem 5.4].

Finally, for d_0 a square-free integer, let χ' denote the non-primitive Dirichlet modulo d_0p induced by $\chi \in X_p^-$. By using estimation of the mean square value $M_{d_0}(p, H_d)$ of $|L(1, \chi')|^2$ for $\chi \in X_p^-(H)$, we improved on (4) in [Lou94, Théorème 6] and on (6) in [LM23, Theorem 1.1]: for any given positive constant $C < 4\pi^2$ and for $p \ge p_C$ large enough we have

$$k_{\mathbb{K}}^{-} \le w_{\mathbb{K}} \left(\frac{p}{C}\right)^{m/4} \tag{9}$$

as K runs over the imaginary subfields of $\mathbb{Q}(\zeta_p)$ of large degrees m = (p-1)/d with d as in (7).

In the present paper, we not simply improve on the upper bounds (5), (6) and (9). With restrictions slightly lighter than (7), we prove an asymptotic behavior instead of an upper bound:

Theorem 3. As p tends to infinity and K runs over the imaginary abelian number fields of conductor p and large degrees m = (p-1)/d with d as in (8) we have an explicit asymptotic

$$h_{\mathbb{K}}^{-} = w_{\mathbb{K}} \left(\frac{(1+o(1))p}{4\pi^2} \right)^{m/4}.$$
 (10)

Moreover, this asymptotic is fully explicit under the following stricter restriction

$$d \le \frac{\log p}{\log \log p} \tag{11}$$

Conjecturally, for a given even degree m there are infinitely many imaginary number fields \mathbb{K} of conductor p and relative class number $h_{\mathbb{K}}^- \gg (\sqrt{p} \log \log p)^{m/2}$, see [BCE] and [Jo] for proofs of this conjecture for m = 2. Hence, we do need some restriction on d for (10) to hold true. The following asymptotic (12) shows that (10) does not hold true under the restriction $\phi(d) = O(\log p)$ only slightly weaker than restriction (8):

Theorem 4. Let $p = 2^d - 1$ range over the (conjecturally infinitely many) Mersenne primes, which implies that d is prime. Let \mathbb{K} be the imaginary field of conductor p and large degree $m = (p-1)/d = 2 \cdot \frac{2^{d-1}-1}{d}$. We have an explicit asymptotic

$$h_{\mathbb{K}}^{-} = w_{\mathbb{K}} \left(\frac{(1+o(1))p}{\pi^2} \right)^{m/4}.$$
 (12)

Whereas (10) is an asymptotic on the logarithm of the relative class numbers, it seems hopeless to have an asymptotic on the relative class numbers themselves. Indeed, even in the much studied case of the cyclotomic fields of prime conductors, Kummer's conjecture asserts that

$$h_{\mathbb{Q}(\zeta_p)}^- = 2p(1+o(1)) \left(\frac{p}{4\pi^2}\right)^{(p-1)/4}$$

However A. Granville showed in [Gran] that this conjecture is false if the two well-known and widely believed Hardy-Littlewood's and Elliott-Halberstam's conjectures are true. See also [MP] for more recent results on Kummer's conjecture.

A lot of results have been obtained for relative class number of cyclotomic fields, e.g. see [Deb], [KLMES], [Lep], [LZ], [Mas] and [MM].

However, we could not find results in the spirit of (10) and (12) in the literature.

2 Proof of Theorem 3

To prove (10) we adapt the method developed in [MM, Proof of Theorem 1]. Let us give some details of their method. Since $L(1,\chi) \neq 0$ for $\chi \in X_p^-$, by (1), there exists $r_p > 0$ such that $L(s,\chi) \neq 0$ for $|s-1| < r_p$ and $\chi \in X_p^-$. As in [MM, Section 2] we take $r_p = 2p^{-3}$. Hence $L(s,\chi)$ does not vanish in the union Ω_p of this disc and the half-plane $\sigma = \Re(s) > 1$, by the infinite products of these *L*-functions. Therefore, in this simply connected open set Ω_p there exists an holomorphic determination of

$$f_H(s) := \sum_{\chi \in X_p^-(H)} \log L(s,\chi)$$

such that

$$\log L(s,\chi) = \sum_{n \ge 2} \frac{\Lambda(n)}{\log n} \chi(n) n^{-s} \text{ for } \Re(s) > 1,$$

where $\Lambda(n) = \log q$ if $n = q^k$ is a power of a prime q and $\Lambda(n) = 0$ otherwise (the von Mangoldt's function). The orthogonality relation

$$\varepsilon_H(n) := \frac{1}{\# X_p^-(H)} \sum_{\chi \in X_p^-(H)} \chi(n) = \begin{cases} 1 & \text{if } n \in H, \\ -1 & \text{if } -n \in H \\ 0 & \text{otherwise} \end{cases}$$
(13)

gives $f_H(s) = f_{H,+1}(s) - f_{H,-1}(s)$, where

$$f_{H,\varepsilon}(s) = \frac{m}{2} \sum_{h \in H} \sum_{\substack{n \ge 2\\n \equiv \varepsilon h \pmod{p}}} \frac{\Lambda(n)}{\log n} n^{-s} \quad \text{(for } \Re(s) > 1 \text{ and } \varepsilon \in \{-1, +1\}\text{)}.$$
(14)

Now, by (1) we have

$$h_{\mathbb{K}}^{-} = w_{\mathbb{K}} \left(\frac{p}{4\pi^2} \exp\left(\frac{4}{m} f_H(1)\right) \right)^{m/4}.$$
 (15)

2.1 Bound on $|f_H(1)|$ and proof of (10) from (15)

1. First, for $\sigma > 1$ and $\xi \in (1, \sigma)$, we write

$$f_H(1) = f_H(\sigma) - f'_H(\xi)(\sigma - 1).$$
(16)

2. Second, bearing on [MM, Proof of Lemma 1] and using Montgomery and Vaughan's explicit bound on primes in arithmetic progressions, we will obtain in Lemma 6 a bound of the type

$$|f_H(\sigma)| \le C_{p,d} - \log(\sigma - 1)$$
 (for $1 < \sigma \le 1 + 1/(5\log 2)$). (17)

3. Third, set $\sigma_0 = 1 + p^{-3}$. The closed disc $|s - \sigma_0| \leq 2p^{-3}$ is included in Ω_p and by [MM, Proof of Lemma 2] we have

$$\Re(f_H(s)) = \log \prod_{\chi \in X_p^-(H)} |L(s,\chi)| \le \frac{1}{2}m\log p \quad \text{(for } |s - \sigma_0| \le 2p^{-3}\text{)}.$$

4. Fourth, as in [MM, Lemma 3] we apply the Borel-Carathéodory lemma to $g(s) := f_H(s) - f_H(\sigma_0)$ which satisfies $g(\sigma_0) = 0$ and $\Re(g(s)) \le M = m \log p$ for $|s - \sigma_0| \le R = 2p^{-3}$ to deduce that

 $|f'_{H}(\xi)| = |g'(\xi)| \le 8M/R = 4mp^{3}\log p \quad \text{(for } |\xi - \sigma_{0}| \le R/2 = p^{-3}\text{)}.$ (18)

5. Using (16), (17) and (18), we end with the upper bound

$$|f_H(1)| \le C_{p,d} - \log(\sigma - 1) + 8M(\sigma - 1)/R$$
 (for $1 < \sigma \le \sigma_0 + R/2 = 1 + 2p^{-3}$),

which for the optimal choice $\sigma = 1 + \frac{R}{8M}$ yields

$$|f_H(1)| \le C_{p,d} + \log(8eM/R) = C_{p,d} + \log(4emp^3\log p) \le C_{p,d} + \log(4ep^4\log p).$$

We have thus proved:

Lemma 5. With $C_{p,d}$ as defined in Lemma 6 below, we have

$$|f_H(1)| \le C_{p,d} + \log\left(4ep^4\log p\right).$$
 (19)

Now, first assume that d is as in (8). Using (21) we get that

$$\frac{4}{m}|f_H(1)| = \frac{4d}{p-1}|f_H(1)| = o(1)$$

as p tends to infinity. The first assertion of Theorem 3 follows, by (15).

Second, assume that d is as in (11). Then $p^{1/d} \ge \log p$ and $C_{p,d} \le \frac{p}{2\log p} + \frac{3}{2} + \frac{3}{8}\log p$, by (20). Hence, we also get

$$\frac{4}{m}|f_H(1)| = \frac{4d}{p-1}|f_H(1)| = o(1)$$

as p tends to infinity. The second assertion of Theorem 3 follows, by (15).

2.2 Proof of the bound on $|f_H(\sigma)|$ for $\sigma > 1$ given in (17)

In [MM] the authors dealt only with cyclotomic fields, i.e. with the case m = p-1 and $H = \{1\}$. In the second step of the strategy above they prove the bound $|f_{\{1\}}(\sigma)| \leq 3 + \log(p/(\sigma - 1))$ for $1 < \sigma \leq 5/2$. To obtain this bound they first prove that for m = p - 1 and $H = \{1\}$, we have

$$\frac{m}{2} \sum_{h \in H} \sum_{\substack{q \ge 2 \\ (\text{mod } p)}} q^{-\sigma} \le e^{-1} + \log \frac{3}{2(\sigma - 1)} \quad \text{ (for } 1 < \sigma \le 5/2 \text{ and } \varepsilon \in \{-1, +1\})$$

(sums over primes q). The key point is that in this sum the primality of q implies $q \ge 2p - 1$. This is no longer true for $d \ge 3$. For example, let p be a prime of the form $p = p_0^2 + p_0 + 1$, with p_0 a prime. Take d = 3 and $H = H_3 = \{1, p_0, p_0^2\}$. For $q = p_0$ we get the term $\frac{p-1}{6}p_0^{-\sigma} \ge \frac{p-1}{6}p^{-\sigma/2}$ in the sum above and this term is $\gg p^{1/2}$ in the range $1 < \sigma \le 1 + p^{-5}$. This ruins the hope of having something as good as [MM, Lemma 1] for $d \ge 3$. Instead, we have: An asymptotic on the logarithms of relative class numbers

Lemma 6. For $p \ge 29$ a prime, $1 < \sigma \le 1 + 1/(5 \log 2)$ and H a subgroup of odd order d of the multiplicative group $(\mathbb{Z}/p\mathbb{Z})^*$ we have the following explicit bounds

$$0 \le f_{H,-1}(\sigma), \ f_{H,+1}(\sigma), \ |f_H(\sigma)| \le C_{p,d} - \log(\sigma - 1),$$

where

$$C_{p,d} = \frac{1}{2}p^{1-1/d} + \frac{3}{2} + \frac{3}{8}\log p.$$
 (20)

Moreover, as p tends to infinity and $\phi(d) = o(\log p)$ we can take the following non-explicit expression

$$C_{p,d} = \frac{3}{2} + \frac{3}{8}\log p + o(m), \text{ where } m = (p-1)/d.$$
(21)

Since $f_H(s) = f_{H,+1}(s) - f_{H,-1}(s)$, it suffices to prove the bound for the $f_{H,\varepsilon}(\sigma)$'s with $\varepsilon \in \{\pm 1\}$. To begin with we notice that (14) and $\Lambda(n) \leq \log n$ give

$$0 < f_{H,\varepsilon}(\sigma) \le \Sigma := \frac{m}{2} \sum_{h \in H} \sum_{\substack{n \ge 2\\n \equiv \varepsilon h \pmod{p}}} \frac{1}{n^{\sigma}} \quad \text{(for } \sigma > 1 \text{ and } \varepsilon \in \{-1, +1\}).$$

Now, to bound this double sum Σ we decompose it into 3 sub-sums $\Sigma = \Sigma_1 + \Sigma_2 + \Sigma_3$ by splitting the domain of variation of its summation index n into three parts, Σ_1 for $2 \le n < p$, Σ_2 for $p \le n < 2p$ and Σ_3 for $n \ge 2p$. We bound Σ_1 in (22) and (23), Σ_2 in (24) and Σ_3 in (27).

2.2.1 The range $2 \le n < p$

If $n \equiv \varepsilon h \pmod{p}$ for some $h \in H$ and $n \geq 2$, then $n^d \equiv \varepsilon^d h^d \equiv \varepsilon^d \equiv \pm 1 \pmod{p}$ and $n \geq (p-1)^{1/d}$. Noticing that #H = d and md = p - 1, it follows that

$$\Sigma_{1} := \frac{m}{2} \sum_{h \in H} \sum_{\substack{2 \le n 1 \text{ and } \varepsilon \in \{-1, +1\}).$$
(22)

Using some of the results of the recent preprint [MS], we can improve on (22). This improvement shows that Theorem 3 still holds true with the restriction (8) weaker than the restriction (11).

Lemma 7. Take $\varepsilon \in \{-1, +1\}$. As p tends to infinity and $\phi(d) = o(\log p)$ we have

$$\Sigma_1 = o(m). \tag{23}$$

Proof. We keep the notation of [MS]. Let H' be the subgroup of order 2d of $(\mathbb{Z}/p\mathbb{Z})^*$ generated by -1 and H. Then

$$\Sigma_1 \le \frac{m}{2} \sum_{1 \ne \lambda \in H'} \rho'(\lambda, p)^{-1} \le \frac{m}{2} \sum_{1 \ne \lambda \in H'} \rho(\lambda, p)^{-1},$$

where for $\lambda \in (\mathbb{Z}/p\mathbb{Z})^*$ we set

$$\begin{split} \rho(\lambda,p) &:= \min\{rs \ : \ (r,s) \in \mathbb{N} \setminus \{(0,0)\}, \ r \equiv \lambda s \pmod{p}\} \\ &\leq \rho'(\lambda,p) := \min\{r \geq 1 \ : \ r \equiv \lambda \pmod{p}\}. \end{split}$$

7

Setting $\vartheta(H',p) = \min_{1 \neq \lambda \in H'} \rho(\lambda,p)$ and noticing that $\phi(2d) = \phi(d)$, we have $\vartheta(H',p) \gg p^{1/\phi(d)}$, by [MS, Lemma 4.1]. Consequently,

$$\sum_{1 \neq \lambda \in H'} \rho(\lambda, p)^{-1} \ll \vartheta(H', p)^{-1/10} + dp^{-1/4} \ll p^{-1/(10\phi(d))} + dp^{-1/4} = \exp\left(-\frac{\log p}{10\phi(d)}\right) + dp^{-1/4},$$

by [MS, Lemma 4.3] applied with $\alpha = 1$, $\beta = 1/5$ and $\kappa = 1/10$. The desired result follows.

2.2.2 The range $p \le n < 2p$

Noticing that #H = d and md = p - 1, we clearly have

$$\Sigma_2 := \frac{m}{2} \sum_{h \in H} \sum_{\substack{p \le n < 2p \\ n \equiv \varepsilon h \pmod{p}}} \frac{1}{n^{\sigma}} \le \frac{p-1}{2p} \le \frac{1}{2} \quad \text{(for } \sigma > 1 \text{ and } \varepsilon \in \{-1, +1\}).$$
(24)

2.2.3 The range $n \ge 2p$

To make our proof somewhat clearer than [MM, Proof of Lemma 1], we state and prove two technical Lemmas:

Lemma 8. For $\alpha > 1$, set $X_{\alpha} = 2(\alpha + 1)/(\alpha - 1)$. For $\beta > 0$, $x \ge \exp(2/\beta)$, and $X \ge 2/\beta$, set

$$S_{\beta}(x) = \frac{1}{\sqrt{x}} \sum_{2 \le k \le \beta \log x} x^{1/k} \text{ and } G_{\alpha}(X) = \left(\int_{1/\beta}^{X/2} e^{t} \frac{\mathrm{d}t}{t^{2}}\right) - \frac{\alpha - 1}{X} e^{X/2}$$

If $(\alpha - 1)/(\alpha + 1) \leq \beta$ and $G_{\alpha}(X_{\alpha}) \leq 0$, then $S_{\beta}(x) \leq \alpha$ for $x \geq \exp(2/\beta)$. This holds true for $\alpha = 11/4$ and $\beta = 1/\log(2)$.

Proof. If $\exp(2/\beta) \leq x < \exp(3/\beta)$, then $S_{\beta}(x) = 1 \leq \alpha$. Now assume that $x \geq \exp(3/\beta)$. Take $X = \log x \geq 2/\beta$. Since

$$S_{\beta}(x) - 1 = \frac{1}{\sqrt{x}} \sum_{3 \le k \le \beta \log x} x^{1/k} \le \frac{1}{\sqrt{x}} \int_{2}^{\beta \log x} x^{1/t} \mathrm{d}t = e^{-X/2} \int_{2}^{\beta X} e^{X/t} \mathrm{d}t = X e^{-X/2} \int_{1/\beta}^{X/2} e^{t} \frac{\mathrm{d}t}{t^2},$$

by the change of variable $t \mapsto X/t$, we have

$$(S_{\beta}(x) - \alpha)\frac{e^{X/2}}{X} = (S_{\beta}(x) - 1)\frac{e^{X/2}}{X} - \frac{\alpha - 1}{X}e^{X/2} \le G_{\alpha}(X).$$

Since $G'_{\alpha}(X) = (X_{\alpha} - X) \frac{e^{X/2}}{2(\alpha - 1)X^2}$ and $X_{\alpha} \ge 2/\beta$, we have $G_{\alpha}(X) \le G_{\alpha}(X_{\alpha})$ for $X \ge 2/\beta$. \Box

Lemma 9. Let $N \ge 1$ be a given integer. Let $a_n \in \mathbb{C}$, $n \ge N$. For $x \ge N$, set $A(x) = \sum_{N \le n \le x} a_n$. Let $b \in \mathcal{C}^1([N, +\infty), \mathbb{C})$ be such that $\lim_{n \to \infty} A(x)b(x) = 0$ and $\int_N^{\infty} A(t)b'(t)dt$ is convergent. Then $\sum_{n\ge N} a_n b(n)$ is convergent and its sum is equal to $-\int_N^{\infty} A(t)b'(t)dt$.

Proof. Setting $M = \lfloor x \rfloor$ and noticing that A(t) = A(n) for $n \leq t < n+1$ and $A_N = a_N$, we have

$$\int_{N}^{x} A(t)b'(t)dt = \left(\sum_{n=N}^{M-1} \int_{n}^{n+1} A(n)b'(t)dt\right) + \int_{M}^{x} A(M)b'(t)dt = -\left(\sum_{n=N}^{M} a_{n}b(n)\right) + A(x)b(x).$$

An asymptotic on the logarithms of relative class numbers

Now, set

$$\pi(x;a,p) = \sum_{\substack{2 \le q \le x \\ q \equiv a \pmod{p}}} 1 \text{ and } \Pi(x;a,p) = \sum_{\substack{2 \le n \le x \\ n \equiv a \pmod{p}}} \frac{\Lambda(n)}{\log n} = \sum_{\substack{2 \le q^k \le x \\ q^k \equiv a \pmod{p}}} \frac{1}{k}$$

(sums over primes $q \ge 2$, over integers n and over integers $k \ge 1$ and primes $q \ge 2$). As in [MM, Proof of Lemma 1] we will use Montgomery and Vaughan's bound

$$\pi(x; p, a) < \frac{2x}{(p-1)\log(x/p)} \quad \text{(for } x > p\text{)}.$$
(25)

The multiplicative group $(\mathbb{Z}/p\mathbb{Z})^*$ being cyclic, the set $\{n \in \mathbb{Z}; n^k \equiv a \pmod{p}\}$ is either empty or the disjoint union of gcd(k, p-1) arithmetic progressions of common difference p. Hence, for $X \ge 1$ we have $\#\{n; 1 \le n \le X, n^k \equiv a \pmod{p}\} \le (X/p+1) gcd(k, p-1) \le k(X/p+1)$. Therefore, for $x \ge 4$ we have

$$0 \le \Pi(x; a, p) - \pi(x; a, p) = \sum_{\substack{2 \le k \le \frac{\log x}{\log 2}}} \frac{1}{k} \sum_{\substack{q \le x^{1/k} \\ q^k \equiv a \pmod{p}}} 1 \le \sum_{\substack{2 \le k \le \frac{\log x}{\log 2}}} \left(\frac{x^{1/k}}{p} + 1\right)$$

and for $2 \le x < 4$ we have $\Pi(x; a, p) = \pi(x; a, p)$. Therefore, by Lemma 8 we have

$$0 \le \Pi(x; a, p) - \pi(x; a, p) \le \frac{11\sqrt{x}}{4p} + \frac{3\log x}{2} \qquad (x \ge 2).$$
(26)

For a given $a \in \mathbb{Z}$, taking N = 2p and $b(t) = t^{-\sigma}$ and recalling that md = p - 1, we obtain

$$\frac{md}{2} \sum_{\substack{q \ge 2p \\ q \equiv a \pmod{p}}} \frac{1}{q^{\sigma}} \le \frac{md\sigma}{2} \int_{2p}^{\infty} \frac{\pi(x; p, a)}{x^{\sigma+1}} \mathrm{d}x \le \sigma \int_{2p}^{\infty} \frac{1}{x^{\sigma} \log(x/p)} \mathrm{d}x = \sigma p^{1-\sigma} \int_{(\sigma-1)\log 2}^{\infty} e^{-x} \frac{\mathrm{d}x}{x},$$

by Lemma 9, (25) and by the change of variables $x \mapsto \exp(x/(\sigma - 1))$. For $\varepsilon \in \{-1, +1\}$ and $1 < \sigma \leq 1 + 1/(5 \log 2)$, noticing that #H = d and $\sigma p^{1-\sigma} \leq \sigma e^{1-\sigma} \leq 1$ for $p \geq 3$ and $\sigma > 1$, we get

$$\frac{m}{2} \sum_{h \in H} \sum_{\substack{q \ge 2p \\ q \equiv \varepsilon h \pmod{p}}} \frac{1}{q^{\sigma}} \le \int_{(\sigma-1)\log 2}^{\infty} e^{-x} \frac{\mathrm{d}x}{x} = -\log(\sigma) + I((\sigma-1)\log 2) \le -\log(\sigma-1),$$

since $a \mapsto I(a)$ increases with $a \in (0,1)$ and $I(1/5) \leq 0$, where

$$I(a) = -\log\log 2 + \int_{a}^{1} (e^{-x} - 1)\frac{\mathrm{d}x}{x} + \int_{1}^{\infty} e^{-x}\frac{\mathrm{d}x}{x}.$$

Now, by Lemma 9 and (26), for $\sigma > 1$ we have

$$\frac{md}{2} \sum_{\substack{n \ge 2p \\ n \equiv a \pmod{p}}} \frac{1}{n^{\sigma}} - \frac{md}{2} \sum_{\substack{q \ge 2p \\ q \equiv a \pmod{p}}} \frac{1}{q^{\sigma}} \le \frac{md\sigma}{2} \int_{2p}^{\infty} \frac{\Pi(x; p, a) - \pi(x; p, a)}{x^{\sigma+1}} \mathrm{d}x$$
$$\le \frac{p}{2} \int_{2p}^{\infty} \frac{\frac{11\sqrt{x}}{4p} + \frac{3\log x}{2}}{x^2} \mathrm{d}x = \frac{11}{4\sqrt{2p}} + \frac{3\log(2p)}{8} + \frac{3}{8}$$

where we used and md = p-1 and $\sigma/x^{\sigma-1} \leq 1$ for $\sigma > 1$ and $x \geq \exp(1)$. Noticing that #H = d and that $\frac{11}{4\sqrt{2p}} + \frac{\log 8}{8} + \frac{3}{8} \leq 1$ for $p \geq 29$, for $\varepsilon \in \{-1, +1\}, p \geq 29$ and $1 < \sigma \leq 1 + 1/(5 \log 2)$ we end up with the bond

$$\Sigma_3 := \frac{m}{2} \sum_{h \in H} \sum_{\substack{n \ge 2p \\ n \equiv \varepsilon h \pmod{p}}} \frac{1}{n^{\sigma}} \le 1 + \frac{3}{8} \log p - \log(\sigma - 1).$$
(27)

An asymptotic on the logarithms of relative class numbers

3 Proof of Theorem 4

Here $\operatorname{Gal}(\mathbb{Q}(\zeta_p)/\mathbb{K})$ is the subgroup $H = \{2^k; 0 \le k \le d-1\}$ of order d of the multiplicative group $(\mathbb{Z}/p\mathbb{Z})^*$. For $\chi \in X_p^-(H)$ let χ' denote the character modulo 2p induced by χ , i.e. $\chi'(n) = 0$ is n is even and $\chi'(n) = \chi(n)$ if n is odd. In Ω_p we can still define an holomorphic determination of

$$g_H(s) = \sum_{\chi \in X_p^-(H)} \log L(s, \chi')$$

such that

$$\log L(s,\chi') = \sum_{\substack{n \ge 3\\ n \text{ odd}}} \frac{\Lambda(n)}{\log n} \chi(n) n^{-s} \text{ for } \Re(s) > 1.$$

Hence, here we have $g_H(s) = g_{H,+1}(s) - g_{H,-1}(s)$, where

$$g_{H,\varepsilon}(s) = \frac{m}{2} \sum_{h \in H} \sum_{\substack{n \ge 3 \text{ odd} \\ n \equiv \varepsilon h \pmod{p}}} \frac{\Lambda(n)}{\log n} n^{-s} \quad \text{(for } \Re(s) > 1 \text{ and } \varepsilon \in \{-1, +1\}\text{)}.$$

Since $\chi(2) = +1$, we have $L(s, \chi) = (1 - 2^{-s})^{-1}L(s, \chi')$. Therefore, $L(1, \chi) = 2L(1, \chi')$ and

$$h_{\mathbb{K}}^{-} = 2\left(\frac{p}{\pi^2}\right)^{m/4} \prod_{\chi \in X_p^{-}(H)} L(1,\chi') = 2\left(\frac{p}{\pi^2} \exp\left(\frac{4}{m}g_H(1)\right)\right)^{m/4},$$

by (1). Now we proceed with $g_H(s)$ as we did with $f_H(s)$ with the following modifications. Since the characters χ' are modulo 2p the bound in Point 3 of subsection 2.1 must be changed to $\Re(g_H(s)) \leq \frac{1}{2}m\log(2p)$ and we end up with the bound $|g_H(1)| \leq C_p + \log(4ep^4\log(2p))$, where C_p is obtained by modifying the proof of Lemma 6 in the following way:

In the range $3 \le n < p$ and n odd we notice that $n \equiv h \pmod{p}$ for $h \in H = \{2^k; 0 \le k \le d-1\}$ has no solution whereas $n \equiv -h \pmod{p}$ if and only if $n = p - 2^k$ for some $k \in \{1, \dots, d-1\}$. Hence (22) must be replaced by the much better bound

$$0 \le \frac{m}{2} \sum_{h \in H} \sum_{\substack{2 \le n < p, \ n \text{ odd} \\ n \equiv \varepsilon h \pmod{p}}} \frac{1}{n^{\sigma}} \le \frac{m}{2} \sum_{k=1}^{d-1} \frac{1}{p - 2^k} \le \frac{md}{2(p - 2^{d-1})} = 1 \quad \text{(for } \sigma > 1 \text{ and } \varepsilon \in \{-1, +1\}).$$

In the range $p \leq n < 2p$ and n odd we have

$$\sum_{\substack{h \in H \\ n \equiv \varepsilon h \pmod{p}}} \sum_{\substack{n \sigma \\ (\text{mod } p)}} \frac{1}{n^{\sigma}} \leq \sum_{\substack{h \in H \\ n \equiv \varepsilon h \pmod{p}}} \sum_{\substack{p \leq n < 2p \\ n \equiv \varepsilon h \pmod{p}}} \frac{1}{n^{\sigma}} \quad \text{(for } \sigma > 1 \text{ and } \varepsilon \in \{-1, +1\}).$$

and we can use the bound (24). In the same way, in the range $n \ge 2p$ and n odd we can use the bound (27). Hence, we end up with the bound

$$0 \le g_{H,-1}(\sigma), \ g_{H,+1}(\sigma), \ |g_H(\sigma)| \le C_p - \log(\sigma - 1), \ \text{where } C_p = \frac{5}{2} + \frac{3}{8}\log p.$$

4 Remarks on Theorem 3

An anonymous referee of a previous version of this paper allowed us to include here her/his proof of our Theorem 3 under the restriction (11) base on [GS01] and [Mon]. Her/his following proof is somewhat shorter, but uses more sophisticated tools and gives less explicit results. Noticing that $\#X_n^-(H) = m/2$, by (1) to obtain (10) it suffices to prove that

$$\frac{1}{\#X_p^-(H)} \sum_{\chi \in X_p^-(H)} \log L(1,\chi) = o(1).$$
(28)

Now, by [Mon] and [GS01], there exists a set $\mathcal{E}(p)$ of non-principal characters modulo p of cardinal $\#\mathcal{E}(p) \ll p^{1/4}$ such that for all characters χ modulo p such that for $\chi \notin \mathcal{E}(p)$ we have

$$\log L(1,\chi) = \sum_{n \le (\log p)^{100}} \frac{\Lambda(n)\chi(n)}{n\log n} + O\left(\frac{1}{\log p}\right).$$
⁽²⁹⁾

Indeed, on applying [Mon, Theorem 1] with Q = p, T = p and $\sigma = 1 - 1/40$, there are fewer than $(Q^2T)^{3(1-\sigma)}(\log Q)^{13} = p^{9/40}(\log p)^{13} \ll p^{1/4}$ primitive characters χ with conductor below p for which the Dirichlet *L*-function $L(s,\chi)$ has a zero in the rectangle $1 - 1/40 = \sigma \leq \Re(s) \leq 1$ and $|\Im(s)| \leq p$. We then apply [GS01, Lemma 8.2], to the characters $\chi \in X_p^-(H)$ not in this exceptional set, with the choice $s = \sigma = 1$, t = 0, q = p, $y = (\log p)^{100}$ and $\sigma_0 = \alpha = 1 - 1/40$ (notice that the rectangle $\sigma_0 < \Re(z) \leq 1$ and $|\Im(z) - t| \leq y + 3$ is included in the previous rectangle $1 - 1/40 = \alpha \leq \Re(s) \leq 1$ and $|\Im(s)| \leq p$ p large enough to have $y + 3 \leq p$). We have

$$\sigma_1 = \min\left(\sigma_0 + \frac{1}{\log y}, \frac{\sigma + \sigma_0}{2}\right) = \min\left(1 - \frac{1}{40} + \frac{1}{100\log_2 p}, 1 - \frac{1}{80}\right) = 1 - \frac{1}{40} + \frac{1}{100\log_2 p}$$

(for p large enough). Hence

$$\frac{\log q}{(\sigma_1 - \sigma_0)^2} y^{\sigma_1 - \sigma} = 10^4 \frac{(\log p)(\log_2 p)^2}{(\log p)^{100(1 - \sigma_1)}} = 10^4 \frac{(\log p)(\log_2 p)^2}{(\log p)^{5/2 - \log_2 p}} = 10^4 \frac{(\log p)(\log_2 p)^2}{(\log p)^{5/2}} \exp(1) \ll \frac{1}{\log p}$$

and (29) holds true for $\chi \in X_p^-(H)$ not in the previous exceptional set of cardinality $\ll p^{1/4}$.

Now, for $\chi \in X_p^-(H)$ in this exceptional set we have explicit bounds $\frac{1}{\sqrt{p}} \ll L(1,\chi) \ll \log p$ if χ is quadratic (by the class number formula (1) applied to the imaginary quadratic field $\mathbb{Q}(\sqrt{-p})$ and [Lou96]) and $\frac{1}{\log p} \ll |L(1,\chi)| \ll \log p$ otherwise (e.g. see [Lou92] and [Lou96]). Hence, noticing that for $x \ge 2$ we have

$$\left|\sum_{n \le x} \frac{\Lambda(n)\chi(n)}{n \log n}\right| \le \sum_{n \le x} \frac{1}{n} \ll \log x$$

we deduce that

$$\log L(1,\chi) = \sum_{n \le (\log p)^{100}} \frac{\Lambda(n)\chi(n)}{n\log n} + \begin{cases} O(\log p) & \text{if } \chi \in X_p^-(H) \text{ is exceptional and quadratic} \\ O(\log_2 p) & \text{if } \chi \in X_p^-(H) \text{ is exceptional and not quadratic} \end{cases}$$
(30)

Using (29), (30) and the orthogonality relation (13), and recalling that $\#X_p^-(H) = m/2 = (p-1)/(2d)$ it follows that

$$\frac{1}{\#X_p^-(H)}\sum_{\chi\in X_p^-(H)}\log L(1,\chi) = \sum_{n\leq (\log p)^{100}}\frac{\Lambda(n)}{n\log n}\varepsilon_H(n) + O\left(\frac{dp^{1/4}\log_2 p}{p}\right) + O\left(\frac{1}{\log p}\right).$$

Now we saw in subsection 2.2.1 that the absolute value of the sum in this latter right hand side is less than or equal to $\#H/(p-1)^{1/d} = d/(p-1)^{1/d}$. Therefore (28) and hence (10) hold true under the restriction $d \leq \frac{\log p}{\log \log p}$ assumed in (11).

References

- [BCE] P. T. Bateman, S. Chowla and P. Erdös. Remarks on the size of $L(1, \chi)$. Publ. Math. Debrecen 1 (1950), 165–182.
- [BMS] B. Borda, M. Munsch and I. E. Shparlinski. Pointwise and correlation bounds on Dedekind sums over small subgroups. *Research in Number Theory*, to appear.
- [Deb] K. Debaene. The first factor of the class number of the *p*-th cyclotomic field. Arch. Math. (Basel) **102** (2014), 237–244.
- [Gran] A. Granville. On the size of the first factor of the class number of a cyclotomic field. *Invent.* math. **100** (1990), 321–338.
- [GS01] A. Granville and S. Soundararajan. Large character sums. J. Amer. Math. Soc. 14 (2001), 365–397.
- [Jo] P. T. Joshi. The size of $L(1, \chi)$ for real nonprincipal residue characters χ with prime modulus. J. Number Theory 2 (1970), 58–73.
- [KLMES] N. Kandhil, A. Languasco, P. Moree, S. S. Eddin and A. Sedunova. The Kummer ratio of the relative class number for prime cyclotomic fields. J. Math. Anal. Appl. 538 (2024), 128368.
- [Lem] F. Lemmermeyer. Ideal class groups of cyclotomic number fields. I. Acta Arith. **72** (1995), 347–359.
- [Lep] T. Lepistö. On the growth of the first factor of the class number of the prime cyclotomic field. Ann. Acad. Sci. Fenn. Ser. A. I. 1974, no. 577, 21 pp.
- [LM21] S. Louboutin and M. Munsch. Second moment of Dirichlet L-functions, character sums over subgroups and upper bounds on relative class numbers. Quart. J. Math. 72 (2021), 1379–1399.
- [LM23] S. Louboutin and M. Munsch. Mean square values of L-functions over subgroups for non primitive characters, Dedekind sums and bounds on relative class numbers. Canad. J. Math. 75 (2023), 1711–1743.
- [Lou92] S. Louboutin. Minoration au point 1 des fonctions L et détermination des corps sextiques abéliens totalement imaginaires principaux. Acta Arith. **62** (1992), 109–124.
- [Lou94] S. Louboutin. Quelques formules exactes pour des moyennes de fonctions L de Dirichlet. Canad. Math. Bull. 36 (1993), 190–196. Addendum. Canad. Math. Bull. 37 (1994), p. 89.
- [Lou96] S. Louboutin. Majorations explicites de $|L(1,\chi)|$ (Suite). C. R. Acad. Sci. Paris Sér. I Math. **323** (1996), 443–446.
- [Lou16] S. Louboutin. Dedekind sums, mean square value of L-functions at s = 1 and upper bounds on relative class numbers. Bull. Pol. Acad. Sci. Math. 64 (2016), 165–174.
- [Lou19] S. Louboutin. On the denominator of Dedekind sums. Bull. Korean Math. Soc. 56 (2019), 815–827.
- [LZ] Y. Lu and W. Zhang. On the Kummer conjecture. Acta Arith. 131 (2008), 87–102.
- [Mas] J. M. Masley. On the first factor of the class number of prime cyclotomic fields. J. Number Theory 10 (1978), 273–290.
- [MM] J. M. Masley and H. L. Montgomery. Cyclotomic fields with unique factorization. J. Reine Angew. Math. 286/287 (1976), 248–256.
- [Mon] H. L. Montgomery Zeros of L-functions. Invent. Math 8 (1969), 346–354.
- [MS] M. Munsch and I. E. Shparlinski. Moments and non-vanishing of *L*-functions over thin subgroups. Submitted.
- [MP] M. R. Murty and Y. N. Petridis. On Kummer's conjecture. J. Number Theory **90** (2001), 294–303.
- [Wal] H. Walum. An exact formula for an average of L-series. Illinois J. Math. 26 (1982), 1–3.
- [Was] L. C. Washington. Introduction to Cyclotomic Fields. Second Edition. Graduate Texts in Mathematics 83. Springer-Verlag, New York, 1997.