Modelling Infodemics on a Global Scale: A 30 Countries Study using Epidemiological and Social Listening Data

Edoardo Loru¹, Marco Delmastro², Francesco Gesualdo³, Matteo Cinelli^{1*}

^{1*}Department of Computer Science, Sapienza University of Rome, Via Regina Elena 295, Rome, 00161, Italy.

²Department of Environmental Sciences, Informatics and Statistics, Ca' Foscari University of Venice, Dorsoduro 3246, Venice, 30123, Italy.

³Centro Ricerche Enrico Fermi, Via Panisperna 89 A, Rome, 00184, Italy. ⁴Department of Translational Research and New Technologies in

Medicine and Surgery, University of Pisa, Via Savi 10, Pisa, 56126, Italy.

*Corresponding author(s). E-mail(s): matteo.cinelli@uniroma1.it; Contributing authors: edoardo.loru@uniroma1.it; marco.delmastro@cref.it; francesco.gesualdo@med.unipi.it;

Abstract

Infodemics are a threat to public health, arising from multiple interacting phenomena occurring both online and offline. The continuous feedback loops between the digital information ecosystem and offline contingencies make infodemics particularly challenging to define operationally, measure, and eventually model in quantitative terms. In this study, we present evidence of the effect of various epidemic-related variables on the dynamics of infodemics, using a robust modelling framework applied to data from 30 countries across diverse income groups. We use WHO COVID-19 surveillance data on new cases and deaths, vaccination data from the Oxford COVID-19 Government Response Tracker, infodemic data (volume of public conversations and social media content) from the WHO EARS platform, and Google Trends data to represent information demand. Our findings show that new deaths are the strongest predictor of the infodemic, measured as new document production including social media content and public conversations, and that the epidemic burden in neighbouring countries appears to have a greater impact on document production than the domestic one. Building on these results, we propose a taxonomy that highlights country-specific discrepancies

between the evolution of the infodemic and the epidemic. Further, an analysis of the temporal evolution of the relationship between the two phenomena quantifies how much the discussions around vaccine rollouts may have shaped the development of the infodemic. The insights from our quantitative model contribute to advancing infodemic research, highlighting the importance of a holistic approach integrating both online and offline dimensions.

Keywords: keyword1, Keyword2, Keyword3, Keyword4

1 Introduction

The continuously growing amount of data and information accessible online have generated both opportunities and threats [1-4], one of which is the risk of infodemics [5, 6]. An infodemic is defined as "an excess information of varying quality, including false/misleading information or ambiguous information or both, that spreads in digital and physical environments during a health emergency" [7]. The COVID-19 infodemic was one of the most relevant phenomena that took place in the online ecosystem during the last decade [8]. According to experimental and real-world evidence, the infodemic had a number of offline consequences, such as the use of autocures [9, 10], reduced vaccine acceptance [11] and weakened social cohesion, trust and order [12-14]. Past the COVID-19 health emergency, the scientific community is still actively discussing the infodemic phenomenon, focusing on: the chance of an overflow of the infodemic beyond the boundaries of public health [15, 16]; the role that new technologies, such as generative AI, will play in future infodemics [17]; the systematization of infodemic management frameworks [18-20], often borrowed from epidemiology [21]. While it is reasonable to assume that epidemics and infodemics are interconnected [6] and metaphorically similar [22], simply adapting epidemic management frameworks to infodemic management might not be sufficient.

The infodemic, as highlighted by WHO, extends beyond the spread of information to include its interplay with behavioural, societal, and health system dynamics [7]. Public health outcomes depend on factors such as information avoidance, communitylevel contextualization, and systemic challenges in health communication. Although these broader dimensions are critical to fully understanding the infodemic, most quantitative studies have focused on narrower aspects, often reducing the phenomenon to single measures, such as the circulation of misinformation [8, 23] or information voids [24]. Similarly, in this study, we narrow our focus on document production as a measurable proxy for the infodemic. However, unlike prior studies, we take an additional step by developing an explanatory model to systematically analyse the factors driving infodemic dynamics related to information production and their relationship with epidemic variables. While explanatory models are commonly used in epidemiology to study disease spread [25, 26], their application to infodemics remains largely unexplored. To effectively tackle, manage, and prepare for future infodemics, it is crucial to address this gap by systematically analysing the factors driving their evolution.

To address this need, following the recent recommendation for the use of a holistic approach in the analysis of information ecosystems related to public health [27], the present study proposes a methodological framework for infodemic modelling, applied to the COVID-19 epi-infodemic. We investigate the relationship between epidemic variables, namely new cases and new deaths, and the volume of social media content and public conversations, which we use as a proxy for the infodemic. We develop a global model to identify the key drivers of the infodemic, incorporating the effects of both epidemiological factors (new cases and deaths) and non-epidemiological factors across different areas. Our analysis includes data from 30 countries representing diverse income groups. To ensure robustness, we replicate the model using information demand, a relevant quantity in infodemic studies [24, 28] that we measure through Google Trends data, as an alternative to information production.

2 Data

In this section, we describe the publicly available datasets that we employ in our analyses. A complete breakdown of the collected data is reported in Tab 1.

Data	Source	N. Countries	Time Window (yy-mm-dd)	Time Scale
New Cases	WHO	236	2020-01-03 - 2023-11-09	Daily
New Deaths	WHO	236	2020-01-03 - 2023-11-09	Daily
New Documents	WHO-EARS	30	$2020 ext{-} 12 ext{-} 15 - 2023 ext{-} 11 ext{-} 09$	Daily
Vaccinations (%)	OxCGRT	183	2020-01-01 - 2023-02-28	Daily
Stringency Index	OxCGRT	183	2020-01-01 - 2023-02-28	Daily
Google Trends	Google	30	2020-01-05 - 2023-11-05	Weekly

Table 1 Summary of variables used for data analysis.

2.1 Epidemiological Data

We employ the WHO Coronavirus (COVID-19) surveillance data that report new COVID-19 cases and deaths on a daily scale. The dataset provides information for 236 countries, starting from 2020-01-03. The data, displayed in a dashboard¹, are freely available². Concerning this study, the version of the dataset we use tracks the number of new cases and new deaths until 2023-11-09 and can be retrieved from the Wayback Machine³.

2.2 Infodemics Data

Although the infodemic encompasses a broad conceptualization, which includes behavioural, social, and health system dynamics as highlighted by the WHO [7], in this

¹https://data.who.int/dashboards/covid19/cases?n=c

²https://covid19.who.int/data ³https://web.archive.org/web/20231111122217/https://covid19.who.int/ WHO-COVID-19-global-data.csv

³

study we focus on information production to develop explanatory models. Documents related to COVID-19 are collected on a daily basis by the WHO Early AI-supported Response with Social Listening Platform (EARS). Documents include public conversations and public social media content related to COVID-19, collected using the Twitter API, and data aggregators of public sources such as forums, message boards, blogs, and comments in news. The dataset provides information for 30 countries starting from 2020-12-15 and can be found in the official WHO EARS repository⁴. Additionally, we collect Google Trends data related to COVID-19 for the same period and set of countries encompassed by the previously described epidemiological data, to serve as a representative metric of information demand, rather than production.

Vaccine Data

We use the Oxford Covid-19 Government Response Tracker (OxCGRT) [29], a publicly available⁵ dataset collecting information on the measures enacted by a government in response to the pandemic, to retrieve the status of COVID-19 vaccination for each country. More in detail, the dataset keeps track of the daily proportion of the vaccinated population in 183 countries from 2020-01-01 to 2023-02-28.

Non Pharmaceutical Interventions Data

The aforementioned OxCGRT also includes the Stringency Index (SI) over time for all 183 countries it monitors, from 2020-01-01 to 2023-02-28, using a daily scale. The SI is a composite measure that quantifies the extent of containment and closure measures adopted by a government based on nine indicators, including school and workspace closings, cancellation of public events, restrictions on gatherings, travel controls, and stay-at-home orders. The SI ranges between 0 and 100, where 100 indicates the maximum stringency of non-pharmaceutical interventions, e.g. a full lockdown.

3 Results

Epidemic and infodemic curves

In order to understand the potential association between the infodemic phenomenon, represented by the curve of new documents, and the epidemic phenomenon, represented by the curves of new cases and new deaths, we start our analysis with a comparison of these curves. Fig. 1 shows the cumulative proportion of new cases, new deaths and new documents over time. Overall, we note an association between the two epidemiological variables, which are only partially correlated to the cumulative production of documents. In some cases, we note coupling between the epidemiological and documents curves, as they tend to have peaks and plateaus (i.e. sudden raises or steady behaviours of the variables) at nearly the same time (see, for instance, India, Nicaragua and Philippines). In some other instances, new cases and new deaths are only partially coupled with new documents, showing a single common jump as in the case of Indonesia. Otherwise, as in the cases of UK, South Africa and France, the

⁴https://github.com/citibeats-labs/who-ears

⁵https://github.com/OxCGRT/covid-policy-dataset

⁴

curves tend to show a stronger decoupling, as the production of new documents is steadier than the evolution of the two epidemiological variables. For what concerns the case of France, we note, at the beginning of 2022, a marked peak in cumulative cases while the documents curve maintains a steady trend; this can be interpreted as a signal of saturation in document production or even as a potential signal of information fatigue, a component of epidemic fatigue [30], according to which people and other actors refrained from posting about COVID-19, regardless of epidemic events. Another interesting instance happens when a peak in the curve of documents is not associated with peaks in the epidemiological curves, as in the case of Malta. This situation may either indicate the enforcement of NPIs, which might fuel document production despite a steadiness of the epidemic curve, or a shift of focus in the national debate about the pandemic towards other countries, as observed at a regional to national scale in [31].

Modelling infodemic at a global scale

Given the non-trivial nature of the coupling between document production and epidemiological variables displayed in Fig. 1, and for which we provide a detailed correlation analysis in Appendix A, we employ a fixed effects regression model to understand and quantify the drivers behind the production of documents. Specifically, we utilise a panel regression framework with fixed effects (i.e. separate intercept values that encode country-specific effects) where the daily number of new documents is treated as the dependent variable y. The panel data is balanced (i.e. each panel contains the same number of observations) and includes a total of N = 30 countries with T = 1054 observations each (from 2020-12-21 to 2023-11-09). We specify model (1a) as follows:

$$y_{it} = \alpha_i + \sum_{j=1}^{6} \gamma_j d_{j,t}^{\text{day}} + \sum_{j=1}^{3} \delta_j d_{j,t}^{\text{season}} + \beta_1 C_{it} + \beta_2 \overline{C}_{it} + \epsilon_{it}$$
(1a)

where α_i is the time-invariant fixed effect for country i, $d_{j,t}^{day}$ are dummy variables for the weekdays, and $d_{j,t}^{season}$ for the seasons of the year. The number of new cases in country i at time t is denoted with the variable C_{it} , whereas we denote with \overline{C}_{it} the total number of new cases at time t in all other countries in the same WHO region of country i. We take a right-aligned rolling average of 7 days for both the independent and the dependent variables, in addition to log-transforming them. Similarly to model (1a), we also define a second specification (1b) where we use the number of deaths D_{it} as the epidemiological independent variable

$$y_{it} = \alpha_i + \sum_{j=1}^{6} \gamma_j d_{j,t}^{day} + \sum_{j=1}^{3} \delta_j d_{j,t}^{season} + \beta_1 D_{it} + \beta_2 \overline{D}_{it} + \epsilon_{it}$$
(1b)




Fig. 1 Normalised cumulative curves of new cases, new deaths, and new documents. Some countries exhibit coupling between the epidemiological curves and the production of new documents (e.g. Philippines, Malaysia), but in most cases the coupling is either partial (e.g. India) or absent (e.g. France).

and a third specification (1c) where both cases and deaths are used as independent variables

$$y_{it} = \alpha_i + \sum_{j=1}^{6} \gamma_j d_{j,t}^{\text{day}} + \sum_{j=1}^{3} \delta_j d_{j,t}^{\text{season}} + \beta_1 C_{it} + \beta_2 \overline{C}_{it} + \beta_3 D_{it} + \beta_4 \overline{D}_{it} + \epsilon_{it} \,.$$
(1c)

In Tab. 2, we report the coefficients obtained for all three models and their standard errors, which are robust to heteroskedasticity.

$\mathbf{6}$

	Ν	lew documen	ts
	(1a)	(1b)	(1c)
New cases	0.110^{***} (0.019)		$0.008 \\ (0.021)$
New cases (neighbours)	0.159^{***} (0.016)		$0.007 \\ (0.022)$
New deaths		$\begin{array}{c} 0.159^{***} \\ (0.022) \end{array}$	$\begin{array}{c} 0.151^{***} \\ (0.034) \end{array}$
New deaths (neighbours)		$\begin{array}{c} 0.258^{***} \\ (0.022) \end{array}$	$\begin{array}{c} 0.248^{***} \\ (0.030) \end{array}$
$\begin{array}{c} \text{Observations} \\ \text{R}^2 \\ \text{Adjusted } \text{R}^2 \end{array}$	$31,620 \\ 0.400 \\ 0.399$	$31,620 \\ 0.493 \\ 0.493$	$31,620 \\ 0.493 \\ 0.493$

 Table 2 Fixed-effect panel regression with robust standard errors. The model includes dummy variables for the day of the week and season of the year.

*p<0.1; **p<0.05; ***p<0.01

From models (1a) and (1b) we obtain positive and significant coefficients associated with the independent variables, which indicate an effect of epidemiological variables on new documents. When both variables are employed (model (1c)), only new deaths are significant, likely due to the positive correlation between new cases and new deaths. Overall, the coefficients associated with such variables can be interpreted as the elasticity of the independent variable with respect to the dependent variable, thus providing a quantitative description of the extent to which a unit change in the evolution of new cases or new deaths affects the infodemic process in percentage points. For instance, the elasticity of new deaths being $\simeq 0.16$ implies that a 1% increase in new deaths would yield a 0.16% increase in document production. Furthermore, the property of regression coefficients being elasticity indexes allows for a comparison between models, as the values are independent of both the unit of measurement and the scale of the dependent and independent variables. With this in mind, the most interesting result concerns the relationship between the coefficients related to epidemiological variables in countries belonging to the same WHO region (β_2 and β_4) versus the domestic ones (β_1 and β_3): when the coefficients are significant, the foreign epidemic burden (measured either with new cases or new deaths) has a greater effect on document production than the domestic one. This can be interpreted as quantitative evidence for the fact that COVID-19 was indeed an infodemic, driven more by foreign epidemic burden than by the domestic one.

As a robustness check, we validate the overall effect of the epidemic variables on the information ecosystem by implementing a different model in which we substitute the dependent variable (i.e. new documents gathered by the WHO-EARS platform) with Google Trends data regarding COVID-19 (see Appendix A, Tab.A1). This model

specification is particularly relevant in that we move from an information production (WHO-EARS) to an information demand (Google Trends) perspective, and obtain consistent results.

Finally, to quantify the extent to which other epidemiological variables affect the infodemic phenomenon, we define a regression model that also includes data regarding vaccine coverage and non-pharmaceutical interventions as regressors. In particular, we add to model specification (1b) a new variable V_{it} that represents the change from time t - 1 to time t of the percentage of the vaccinated population in country i, and a variable S_{it} that is the Stringency Index at time t for country i. As for all previous models, we take the right-aligned rolling average of 7 days of V_{it} and S_{it} and then we log-transform them. This regression model is thus specified as follows:

$$y_{it} = \alpha_i + \sum_{j=1}^{6} \gamma_j d_{j,t}^{day} + \sum_{j=1}^{3} \delta_j d_{j,t}^{season} + \beta_1 D_{it} + \beta_2 \overline{D}_{it} + \beta_3 V_{it} + \beta_4 S_{it} + \epsilon_{it}$$
(1d)

Interestingly, the model achieves an adjusted R^2 of 0.43, which is slightly lower than that of the simpler model (1b). However, consistently with our previous results, all included variables have a significant positive effect on the production of new documents, especially the percentage change in the vaccinated population (see Appendix A, Tab A2). To make sure that the potential multicollinearity of the epidemiological variables with the newly added regressors had no impact on the results, we compute the Variance Inflation Factor (VIF) for model (1d), obtaining a VIF ≤ 2 (i.e. non-concerning multicollinearity) for all independent variables.

Temporal sensitivity of the model

As it is reasonable to assume that the epidemic fuels the infodemic consistently throughout its development, we now aim to investigate whether our model reflects the possibly weaker relationship between the phenomena towards the end of the health crisis. From a technical point of view, such variations in the association between the dependent and independent variables over time would be highlighted by changes in the elasticity indexes. Indeed, while the coefficient values for the general model are fixed, we expect them to change when fitting the same model over different time windows over the time frame under study. To this end, we evaluate the temporal sensitivity of the model by studying how the elasticity changes over time with model (1b), as it is the specification that better explains the data in terms of R^2 .

To substantiate this reasoning, in Fig 2 panel A, we report the evolution of the internal elasticity β_1 , which encodes the responsiveness of the infodemic to domestic changes in the new death counts. In detail, we create rolling windows of 6 months and employ model specification (1b). The resulting curve is characterised by a decreasing trend: we observe a peak during the first half of 2021, when the roll-out of the vaccine likely strengthened the impact of the epidemic burden on the infodemic, followed by oscillations around zero starting from the end of 2021.

Finally, as the WHO-EARS data collection only began at the end of 2020, we perform a robustness check including year 2020 but changing the dependent variable. The

new dependent variable shifts the focus towards information demand and is represented by a collection of COVID-19 data from Google Trends. Our findings highlight not only the same decreasing trend of the elasticity coefficient from the first half of 2021 onward, but also an increasing trend during 2020 that reaches the same peak in the first 6 months of 2021 (see Appendix A, Fig A6). Thus, results obtained with Google Trends data and their consistency with those obtained with EARS data along the overlapping time window further indicate the potential role of the production and demand of documents about vaccines on the observed elasticity.

Country-level differences in infodemic modelling

Once modelled and examined the relationship between the epidemic and the infodemic at a global scale, a natural step is to study the infodemic response of single countries. This kind of analysis provides additional insights into how national infodemics may deviate from the general trend.

We perform a regression analysis for each country, and subsequently study the internal β_1 and external β_2 elasticity, where the former represents the elasticity index obtained from domestic epidemiological variables and the latter represents the elasticity index obtained from epidemiological variables of countries in the same WHO region. Fig 3 panel A displays the relationship between internal and external elasticity indexes across countries, showing heterogeneous results. Using the mean values of external and internal elasticity indexes as references, we divide the graph into four quadrants that provide a taxonomy of countries based on the relationship between internal and external elasticity. While most countries cluster near the junction between the axes marked by the average values, we note some exceptions. For instance, Canada (CA), located in the upper left quadrant, exhibits a relatively large external elasticity but an internal elasticity close to 0, suggesting that the evolution of new deaths in neighbouring countries (likely, the US) had a larger impact on document production than the domestic ones. Conversely, Iraq (IQ), located in the lower right quadrant, is characterised by a large internal elasticity and an external elasticity of approximately 0, showing a lower impact of epidemic data from neighbouring countries on the national infodemic. Fig 3 panel B displays the difference between internal and external elasticity indexes for each country. In countries that show a difference above 0, the domestic epidemic burden has a higher impact on the infodemic compared to the burden from neighbouring countries. In, countries with a difference value below 0, the infodemic is mostly fuelled by the foreign epidemiological situation (proportionally to the magnitude of the difference), while the impact of the domestic epidemic burden on document production is limited.

All estimated values of β_1 and β_2 are fully reported in Appendix A Fig A5, including the two resulting from the panel regression with all countries.

4 Discussion and Conclusions

In this paper, we quantitatively analyse how epidemic variables drive the infodemic, using data-driven models based on diverse official sources, including the WHO, the University of Oxford, and Google. The study incorporates data on COVID-19 cases,




Fig. 2 Temporal evolution of the internal elasticity associated with the number of new deaths. We fit the panel regression model specified in (1b) over 6-month sliding windows, and display how the internal elasticity β_1 and its robust standard error evolve. Left, the evolution of β_1 using the number of new documents as the dependent variable, with the inset reporting the total number of new deaths across all modelled countries. Right, the evolution of β_1 using Google Trends data as the dependent variable, as a proxy for information demand rather than production. Both curves show similar behaviour and comparable values, peaking in the first half of 2021 and stabilizing around values close to 0 from the end of 2021 onward.

related deaths, vaccine coverage and non-pharmaceutical interventions, as well as information production and demand. This approach leads to key insights regarding the infodemic phenomenon that can inform public health measures, infodemic management strategies, and, potentially, infodemic prediction models. Our models show a clear relationship between document production and epidemiological variables (i.e., the infodemic and the epidemic), with the number of new cases and new deaths being among the main drivers for the infodemic. Moreover, we show that new deaths outperform other epidemic-related variables as infodemic predictors. The relevance of this variable resonates with recent findings from social psychology, according to which the salience of death brought on by COVID-19 played a central role in driving the attitudes and behaviour of the population [32]. Regarding the epidemic variables, one of the most significant findings is that the epidemiological situation in neighbouring countries emerges as a strong predictor of the infodemic at the national level. If we interpret the foreign epidemic situation as a measure of pandemic proximity, we can speculate that the fear of impending domestic contagions was a significant driver of the infodemic phenomenon. This highlights the critical importance of coordinated, crossborder efforts to mitigate the spread of misinformation and disinformation during health crises.

From a technical perspective, the coefficients in our model represent elasticity factors, providing a robust and scale-independent measure of how percentage changes in independent variables affect the dependent variable (i.e. the intensity of the infodemic). Furthermore, the model displays temporal sensitivity with respect to the epidemic burden and thus has more explanatory power when fitted on the early phases of the




Fig. 3 Country-specific internal and external elasticities associated with the number of new deaths. For each country, we fit a regression model as specified in (1b) and obtain an estimate of the internal elasticity β_1 and of the external elasticity β_2 . (A) Comparison of β_1 and β_2 for all countries, with the vertical and horizontal dashed lines reporting the average β_1 and β_2 , respectively, and the triangle indicating the values resulting from the panel regression. (B) Difference between internal elasticity $\beta_1 - \beta_2$ for all countries. Nicaragua was omitted from this figure because its estimated β_1 was much higher than that of all other countries.

epidemic/pandemic. This aspect of the model highlights its versatility and accuracy in providing robust results that can be used in scenarios where the independent variable is subject to sudden changes. Furthermore, a model characterised by such fine temporal sensitivity enables us to develop reasonable scenarios of infodemic trends by accounting for variations in the effects of different independent variables across different time windows, aligning with common practices in epidemic management [33]. The results are robust to a stress test performed using Google Trends data as an alternative dependent variable [34, 35], therefore considering information demand as proxy for the infodemic. This dual perspective aligns closely with the broader definition of infodemics, which encompasses both the generation and consumption of information [36].

Our study has a number of limitations. First, data on information production are available only after December 2020, and this might result in a partial image of the whole information dynamic. Secondly, despite considering 30 countries, for certain areas of the world, such as Eastern Europe, Russia, China and Australia data are not

available, and therefore a geographical bias could have been introduced in the model, limiting its accuracy and generalisability. Lastly, we considered only document production and information needs as proxies for the infodemic, therefore excluding some other crucial measures of the infodemic, such as information voids and disinformation.

The findings from this study hold significant implications for the management of future infodemics. The influence of neighbouring countries' epidemiological situations underscores the necessity of international cooperation to address the inherently global nature of infodemics. Information spread transcends borders, requiring transnational strategies to counteract misinformation effectively. Our results demonstrate that the evolution of infodemics and their relationship with epidemic variables can be closely monitored, even over short time windows. The proposed modelling framework offers potential for nowcasting and short-term predictions of infodemic trends, making it a valuable tool in critical situations of rapid information spreading. Just as epidemic models allow for the evaluation of different epidemic management strategies, our model enables the estimation and potential prediction of the effects of various infodemic management measures on the infodemic burden. The model's robustness in capturing both information demand and production creates an opportunity for developing early-warning systems to detect and mitigate emerging infodemics promptly. Future specifications of the model could better capture the complexity of the infodemic, enhancing both its accuracy and explanatory power. Assuming some degree of similarity between an infodemic and a syndemic [37, 38], where a multitude of interacting factors such as biological, social, and geopolitical ones play a significant role, additional variables should be considered. In the considered case, variables such as education level, economic indicators, and internet access, are more readily available and could provide valuable insights if included in future analyses. Others, such as health and digital literacy, trust in media and institutions, and characteristics of the media landscape, are potentially harder to measure but could be explored through regional investigations, especially with greater resource allocation. Additionally, combining measures of document production and information demand into a single variable could offer a more integrated perspective on the interplay between these two dimensions of the infodemic. Furthermore, variables like the estimation of information voids and the prevalence of circulating misinformation and disinformation, while requiring advanced computational capabilities, hold significant potential for broadening the scope of infodemic modelling. Including these factors would provide a more comprehensive framework for understanding and addressing the multifaceted dynamics of infodemics and enable policymakers and health organizations to identify and respond to emerging infodemics more effectively, preventing escalation. Overall, this study provides a comprehensive framework for understanding the dynamics of infodemics and offers actionable insights for developing more effective containment strategies in the future. While the model already provides valuable information, it also has the potential for further refinement by incorporating additional variables and perspectives, to enhance its explanatory power and broaden its applicability.

References

- Vosoughi, S., Roy, D., Aral, S.: The spread of true and false news online. science 359(6380), 1146–1151 (2018)
- [2] Shao, C., Ciampaglia, G.L., Varol, O., Yang, K.-C., Flammini, A., Menczer, F.: The spread of low-credibility content by social bots. Nature communications 9(1), 1–9 (2018)
- [3] Cinelli, M., De Francisci Morales, G., Galeazzi, A., Quattrociocchi, W., Starnini, M.: The echo chamber effect on social media. Proceedings of the National Academy of Sciences 118(9), 2023301118 (2021)
- [4] Lorenz-Spreen, P., Oswald, L., Lewandowsky, S., Hertwig, R.: A systematic review of worldwide causal and correlational evidence on digital media and democracy. Nature human behaviour 7(1), 74–101 (2023)
- [5] Zarocostas, J.: How to fight an infodemic. The lancet **395**(10225), 676 (2020)
- [6] Briand, S.C., Cinelli, M., Nguyen, T., Lewis, R., Prybylski, D., Valensise, C.M., Colizza, V., Tozzi, A.E., Perra, N., Baronchelli, A., *et al.*: Infodemics: A new challenge for public health. Cell **184**(25), 6010–6014 (2021)
- [7] Wilhelm, E., Ballalai, I., Belanger, M.-E., Benjamin, P., Bertrand-Ferrandis, C., Bezbaruah, S., Briand, S., Brooks, I., Bruns, R., Bucci, L.M., *et al.*: Measuring the burden of infodemics: Summary of the methods and results of the fifth who infodemic management conference. JMIR infodemiology 3(1), 44207 (2023)
- [8] Cinelli, M., Quattrociocchi, W., Galeazzi, A., Valensise, C.M., Brugnoli, E., Schmidt, A.L., Zola, P., Zollo, F., Scala, A.: The covid-19 social media infodemic. Scientific reports 10(1), 1–10 (2020)
- [9] Aghababaeian, H., Hamdanieh, L., Ostadtaghizadeh, A.: Alcohol intake in an attempt to fight covid-19: A medical myth in iran. Alcohol 88, 29–32 (2020)
- [10] Temple, C., Hoang, R., Hendrickson, R.G.: Toxic effects from ivermectin use associated with prevention and treatment of covid-19. New England Journal of Medicine 385(23), 2197–2198 (2021)
- [11] Loomba, S., De Figueiredo, A., Piatek, S.J., De Graaf, K., Larson, H.J.: Measuring the impact of covid-19 vaccine misinformation on vaccination intent in the uk and usa. Nature human behaviour 5(3), 337–348 (2021)
- [12] Betsch, C., Korn, L., Sprengholz, P., Felgendreff, L., Eitze, S., Schmid, P., Böhm, R.: Social and behavioral consequences of mask policies during the covid-19 pandemic. Proceedings of the National Academy of Sciences 117(36), 21851–21853 (2020)

- [13] Jewett, R.L., Mah, S.M., Howell, N., Larsen, M.M.: Social cohesion and community resilience during covid-19 and pandemics: A rapid scoping review to inform the united nations research roadmap for covid-19 recovery. International Journal of Health Services 51(3), 325–336 (2021)
- [14] Van Der Linden, S.: Misinformation: susceptibility, spread, and interventions to immunize the public. Nature medicine 28(3), 460–467 (2022)
- [15] Pagoto, S.L., Palmer, L., Horwitz-Willis, N.: The next infodemic: abortion misinformation. Journal of Medical Internet Research 25, 42582 (2023)
- [16] Alipour, S., Galeazzi, A., Sangiorgio, E., Avalle, M., Bojic, L., Cinelli, M., Quattrociocchi, W.: Cross-platform social dynamics: an analysis of chatgpt and covid-19 vaccine conversations. Scientific Reports 14(1), 2789 (2024)
- [17] De Angelis, L., Baglivo, F., Arzilli, G., Privitera, G.P., Ferragina, P., Tozzi, A.E., Rizzo, C.: Chatgpt and the rise of large language models: the new ai-driven infodemic threat in public health. Frontiers in Public Health 11, 1166120 (2023)
- [18] Eysenbach, G.: Infodemiology: The epidemiology of (mis) information. The American journal of medicine 113(9), 763–765 (2002)
- [19] Eysenbach, G., et al.: How to fight an infodemic: the four pillars of infodemic management. Journal of medical Internet research 22(6), 21820 (2020)
- [20] Ishizumi, A., Kolis, J., Abad, N., Prybylski, D., Brookmeyer, K.A., Voegeli, C., Wardle, C., Chiou, H.: Beyond misinformation: developing a public health prevention framework for managing information ecosystems. The Lancet Public Health (2024)
- [21] Scales, D., Gorman, J., Jamieson, K.H.: The covid-19 infodemic—applying the epidemiologic model to counter misinformation. New England Journal of Medicine 385(8), 678–681 (2021)
- [22] Simon, F.M., Camargo, C.Q.: Autopsy of a metaphor: The origins, use and blind spots of the 'infodemic'. new media & society 25(8), 2219–2240 (2023)
- [23] Gallotti, R., Valle, F., Castaldo, N., Sacco, P., De Domenico, M.: Assessing the risks of 'infodemics' in response to covid-19 epidemics. Nature human behaviour 4(12), 1285–1293 (2020)
- [24] Purnat, T.D., Vacca, P., Czerniak, C., Ball, S., Burzo, S., Zecchin, T., Wright, A., Bezbaruah, S., Tanggol, F., Dubé, E., *et al.*: Infodemic signal detection during the covid-19 pandemic: development of a methodology for identifying potential information voids in online conversations. JMIR infodemiology 1(1), 30971 (2021)
- [25] Bonaccorsi, G., Pierri, F., Cinelli, M., Flori, A., Galeazzi, A., Porcelli, F., Schmidt,

A.L., Valensise, C.M., Scala, A., Quattrociocchi, W., *et al.*: Economic and social consequences of human mobility restrictions under covid-19. Proceedings of the national academy of sciences **117**(27), 15530–15535 (2020)

- [26] Vandentorren, S., Smaïli, S., Chatignoux, E., Maurel, M., Alleaume, C., Neufcourt, L., Kelly-Irving, M., Delpierre, C.: The effect of social deprivation on the dynamic of sars-cov-2 infection in france: a population-based analysis. The Lancet public health 7(3), 240–249 (2022)
- [27] Jin, S.L., Kolis, J., Parker, J., Proctor, D.A., Prybylski, D., Wardle, C., Abad, N., Brookmeyer, K.A., Voegeli, C., Chiou, H.: Social histories of public health misinformation and infodemics: case studies of four pandemics. The Lancet Infectious Diseases (2024)
- [28] Mavragani, A., Ochoa, G., et al.: Google trends in infodemiology and infoveillance: methodology framework. JMIR public health and surveillance 5(2), 13439 (2019)
- [29] Hale, T., Angrist, N., Goldszmidt, R., Kira, B., Petherick, A., Phillips, T., Webster, S., Cameron-Blake, E., Hallas, L., Majumdar, S., *et al.*: A global panel database of pandemic policies (oxford covid-19 government response tracker). Nature human behaviour 5(4), 529–538 (2021)
- [30] Lilleholt, L., Zettler, I., Betsch, C., Böhm, R.: Development and validation of the pandemic fatigue scale. Nature Communications 14(1), 6352 (2023)
- [31] Castriota, S., Delmastro, M., Tonin, M.: National or local infodemic? the demand for news in italy during covid-19. International Journal of Health Economics and Management 23(4), 507–536 (2023) https://doi.org/10.1007/s10754-023-09350-3
- [32] Pyszczynski, T., Lockett, M., Greenberg, J., Solomon, S.: Terror management theory and the covid-19 pandemic. Journal of Humanistic Psychology 61(2), 173– 189 (2021)
- [33] Jewell, N.P., Lewnard, J.A., Jewell, B.L.: Predictive mathematical models of the covid-19 pandemic: underlying principles and value of projections. Jama 323(19), 1893–1894 (2020)
- [34] Nghiem, L.T.P., Papworth, S.K., Lim, F.K.S., Carrasco, L.R.: Analysis of the capacity of google trends to measure interest in conservation topics and the role of online news. PLOS ONE 11(3), 0152802 (2016) https://doi.org/10.1371/journal. pone.0152802
- [35] Gravino, P., Prevedello, G., Galletti, M., Loreto, V.: The supply and demand of news during covid-19 and assessment of questionable sources production. Nature Human Behaviour 6(8), 1069–1078 (2022) https://doi.org/10.1038/ s41562-022-01353-3

- [36] Health, T.L.P.: Rethinking information ecosystems and infodemics (2024)
- [37] Singer, M., Bulled, N., Ostrach, B., Mendenhall, E.: Syndemics and the biosocial conception of health. The lancet 389(10072), 941–950 (2017)
- [38] Horton, R.: Offline: Covid-19 is not a pandemic. The lancet **396**(10255), 874 (2020)
- [39] Rodríguez-Arias, M.A., Rodó, X.: A primer on the study of transitory dynamics in ecological series using the scale-dependent correlation analysis. Oecologia 138, 485–504 (2004)
- [40] Rodó, X., Rodríguez-Arias, M.-À.: A new method to detect transitory signatures and local time/space variability structures in the climate system: the scale-dependent correlation analysis. Climate Dynamics 27, 441–458 (2006)
- [41] Fontal, A., Bouma, M.J., San-José, A., López, L., Pascual, M., Rodó, X.: Climatic signatures in the different covid-19 pandemic waves across both hemispheres. Nature Computational Science 1(10), 655–665 (2021)

Appendix A

Coupling between epidemic and infodemic using cross-correlation

In order to quantitatively estimate the level of agreement between the number of new documents and the number of new cases, that is the infodemic and epidemic phenomena, we compute the cross-correlation function (CCF) of the two time series, by displacing the latter of $h = 0, \pm 1, \pm 2, \dots, \pm 25$ days. In this analysis, correlation for negative (positive) values of h indicates that the number of documents precedes (follows) the number of cases, whereas a high correlation for h = 0 suggests that the two variables are strongly coupled. In Fig A1, we observe that almost all countries exhibit a positive correlation across all the considered lags, suggesting that the number of new documents and the number of new cases are positively coupled overall. However, we note that not all CCF's display the same profile. For instance, the curves for India, Indonesia, and Senegal are symmetrical around h = 0, meaning that the highest correlation is found when the two time series are not displaced with respect to each other. On the other hand, countries such as Angola, the UK, and Spain show a generally decreasing CCF: correlation is highest when the number of cases is shifted back in time, meaning that the number of documents appears to lead the number of cases. Note that the behaviour observed in the case of countries such as France and Switzerland can be interpreted as a specification of this decreasing trend towards negative values. Although not as common, the opposite behaviour (increasing CCF) can also be observed (e.g., for Nicaragua and Yemen). For a final visual inspection, in Fig A2 we highlight how the cross-correlation function for some countries relates to their respective number of documents and number of cases over time.




Fig. A1 Cross-correlation between number of new cases and number of new documents. On the *x*-axis, how many days the number of documents has been shifted by, with positive (negative) values indicating that the documents time series has been moved backwards (forwards); on the *y*-axis, the sample correlation between the two time series at the given lag.

Transient dynamics of coupling between epidemic and infodemic

The positive cross-correlation observed in the data represents summary information that is useful to understand the general relationship between the variables over the complete time window. Indeed, cross-correlation does not take into account local variations that are present in the data (see for instance the time series in Fig A2). These variations, if present, may represent a localised (e.g., weekly/bi-weekly) decoupling between the epidemic and infodemic curves. To capture these local variations, we




Fig. A2 Cross-correlation between the number of new cases and the number of new documents. Four paradigmatic cases related to Senegal, India, UK and Peru are displayed.

employ the scale-dependent correlation analysis (SDC) [39, 40], a method applied to the time series of both epidemiological and infodemiological variables [41]. The SDC plots can be read as follows: each cell of the grid is coloured according to the Spearman correlation coefficient, with the x and y coordinates corresponding to the position of the two respective time windows of size s = 70 days along each of the time series. The main diagonal shows the correlation between paired segments (h = 0), while other points refer to the correlation between unpaired segments ($h \neq 0$): in the area above the diagonal the first segment of the new cases time series lags the second segment of the new documents time series, while in the area below the diagonal the opposite happens. More in general, the lag h between the cells' position corresponds to the distance

from the diagonal $(h \in \{-21, \ldots, +21\} \text{ days})$, thus the value of the shift between two fragments is the distance in pixels from any point to the main diagonal: positive lags $h \in [1, 21]$ of the new cases time series with respect to the new documents one are visualised as shifts on the *x*-axis, whereas negative lags $h \in [-21, -1]$ (that is positive lags of the new documents time series with respect to the new cases one) are visualised as shifts on the *y*-axis. For instance, the first line of pixels in Fig A3 compares new documents in the interval [1, 70] with new cases in the same interval, while the first pixel on the right of the diagonal compares new documents in the interval [1, 70]with new cases in the interval [2, 71].

The main message deriving from Fig A3, in which only correlations found to be significant in a non-parametric randomised test ($\alpha = 0.01$) are shown and coloured, is systematic evidence for transient dynamics of correlation towards both positive and negative values witnessed by curves oscillating between blue-coloured and red-coloured dots. Nonetheless, not all countries behave the same: some of them display a relatively negative correlation between the two phenomena (e.g., Spain), whereas others (e.g., the UK) switch between positive, negative or non-significant correlation regimes, thus indicating a fluctuating relevance of one variable with respect to the other. While the peculiarity of these coupling dynamics is substantiated by the correlation values themselves, the interplay between new documents and new cases substantially differs from that displayed by comparing epidemiological variables only. In fact, the SDC plot of the number of new cases and new deaths displays, as expected, largely positive correlations (see Appendix A, Fig A4).

Table A1 Fixed-effect panel regression with robust standard		
errors. As the data provided by Google Trends are weekly, unlike the		
models in Tab 2 this regression only takes into account dummy		
variables for the season of the year.		

		Google Tren	ds
	(2a)	(2b)	(2c)
New cases	0.150^{***} (0.025)		0.055^{*} (0.024)
New cases (neighbours)	$\begin{array}{c} 0.132^{***} \\ (0.029) \end{array}$		-0.123^{***} (0.032)
New deaths		$\begin{array}{c} 0.161^{***} \\ (0.027) \end{array}$	0.101^{***} (0.028)
New deaths (neighbours)		$\begin{array}{c} 0.287^{***} \\ (0.025) \end{array}$	$\begin{array}{c} 0.405^{***} \\ (0.033) \end{array}$
Observations	6,000	6,000	6,000
R^2 Adjusted R^2	$\begin{array}{c} 0.440 \\ 0.437 \end{array}$	$0.568 \\ 0.566$	$0.582 \\ 0.580$

*p<0.1; **p<0.05; ***p<0.01




Fig. A3 Scale Dependent Correlation between the number of new daily cases and of new daily documents. For all time windows of s = 70 days of the two time series, the Spearman correlation is computed and reported on the figure if significant (p < 0.01); on the main diagonal, the correlation between paired windows is reported, whereas off-diagonal the windows are shifted with respect to each other. Trinidad and Tobago (TT) and South Africa (ZA) are labelled with their country code rather than their name for visualization purposes.




Fig. A4 Scale Dependent Correlation between the number of new daily cases and of new daily deaths. For all time windows of s = 70 days of the two time series, the Spearman correlation is computed and reported on the figure if significant (p < 0.01); on the main diagonal, the correlation between paired windows is reported, whereas on the off-diagonal the windows are shifted with respect to each other. Trinidad and Tobago (TT) and South Africa (ZA) are labelled with their country code rather than their name for visualisation purposes.



Table A2 Fixed-effect panel regression (1d) with robust standard errors. The model includes dummy variables for the day of the week and season of the year. All independent variables have $VIF \leq 2$, suggesting no multicollinearity.

	New documents
New deaths New deaths (neighbours) Stringency Index Vaccinated population % change	$\begin{array}{c} 0.100^{***} \ (0.030) \\ 0.190^{***} \ (0.040) \\ 0.35^{**} \ (0.11) \\ 1.41^{***} \ (0.28) \end{array}$
Observations R^2 Adjusted R^2	$22,230 \\ 0.432 \\ 0.431$

*p<0.1; **p<0.05; ***p<0.01




Fig. A5 Internal and external elasticity associated with the number of new deaths in model (1b). The exact internal elasticity for Nicaragua is omitted from the visualization because much larger than the rest ($\beta_1 \approx 8$).




Fig. A6 Internal elasticity associated with the number of new deaths in a panel regression framework. 'New documents' refers to the data collected by the WHO-EARS platform, whereas 'Google Trends' to the COVID-19 search interest reported by Google.