Testing quantum gravity with dilute dipolar Bose gases

Asma Tahar Taiba ¹ and Abdelâali Boudjemâa^{2,3}

¹ LPTHIRM, Department of Physics, Faculty of Sciences,

University of Blida 1, P.O. Box. 270, 09000, Blida, Algeria

² Department of Physics, Faculty of Exact Sciences and Informatics, and ³ Laboratory of Mechanics and Energy,

Hassiba Benbouali University of Chlef, P.O. Box 78, 02000, Chlef, Algeria.*

(Dated: February 3, 2025)

We systematically investigate the effects of quantum gravity on the ground-state properties of dilute homogeneous dipolar Bose gases using the Hartree-Fock-Bogoliubov theory based on the generalized uncertainty principle. We calculate quantum gravity corrections to the condensed fraction, the equation of state, the critical temperature and the superfluid fraction. Improved upper bounds on the generalized uncertainty principle parameters are found. We compare our predictions with previous experimental and theoretical results.

I. INTRODUCTION

Quantum theory and general relativity are the fundamental pillars of our current understanding of physics. They successfully describe phenomena at microscopic and macroscopic scales, respectively [1]. Quantum gravity (QG), which aims to unify these incompatible theories, poses significant challenges in modern physics. One of the intriguing manifestation of QG is the so-called generalized uncertainty principle (GUP), which predicts a minimal Planck length in quantum spacetime. The GUP is crucial in diverse QG theories and has been extensively used to explore the Planck-scale phenomenon. The idea of introducing a minimal length into quantum theory has a long history. It was suggested that if one takes into account the gravitational interaction in high energy scatterings, one needs to go to higher center of mass energies (see e.g. [2–6] and references therein). This is essentially a Heisenberg microscope argument [7]. Therefore, it turns out that the usual uncertainty relation is modified yielding a minimal length which is one way to address the infinities that occur in certain solutions in general relativity.

The typical scales associated with QG effects is the Planck scale : 10^{16} TeV or 10^{-35} m (~14 orders of magnitude higher than the scales accessible at the LHC), which renders experiments to test the quantum nature of gravity prohibitive [8]. However, due to rapid advances in quantum technologies tabletop tests of QG become possible suggesting that the low energy signature of QG may be detectable. Recently, many experimental attempts aim at detecting non-classical features of the gravitational interaction have been proposed in [9–13] using principles of quantum mechanics including entanglement, superposition, and decoherence. In their recent work, Westphal et al. [14] have measured gravitational coupling between millimetre-sized masses and Brack et al. [15] have shown the dynamical detection of gravitational coupling between resonating beams in the hertz regime. Most recently, Fuchs *et al.* [16] have identified a way to measure gravity at microscopic levels using massive quantum sensors based on levitated mechanical systems, leading to broaden our understanding of the theory of the QG.

The use of cold atoms including Bose-Einstein condensates (BECs) allows new ways of testing the elusive QG and its effect due to their unprecedented degree of control and sensitivity to ultraweak forces [17–25]. Moreover, based on the generalized uncertainty principle (GUP) [26–41], which enables for investigating QG signatures at lower and more accessible energy levels, the statistical properties of ideal Bose gases have been widely investigated (see, e.g., [42–48] and references therein). The GUP approach was also employed in order to test QG in weakly interacting Bose gases [49].

Motivated by the above fascinating experimental and theoretical works, we investigate in this paper low-energy QG effects using ultracold dipolar BECs since attempts to model a full theory have not been successful until now. Testing QG using dipolar BECs is particularly promising due to the significant role played by dipole-dipole interactions (DDIs) (see for review [50–52]). Ultracold atoms with DDIs which have the same form as the quantum gravitational interactions would be interesting to distinguish the QG signal from electromagnetic force [17], and hence offer a precise environment for testing QG effects. The anisotropic and long-range character of DDIs make these systems especially suitable for refining the bounds on QG parameters and improving the constraints set by the GUP.

The goal is to study the effect of QG in dilute homogeneous dipolar Bose gases using the linear and quadratic form of the GUP (LQGUP) model, which implies a minimum measurable length and a maximum measurable momentum [30, 48]. It has been shown that the LQGUP model predicts stronger QG effects compared to the purely QGUP model notably in the regime of low energies/momenta. Furthermore, the LQGUP when applied to Bose systems, modifies the density of states [48], and thus the corresponding dispersion relation, quantum and thermal fluctuations. We calculate the QG corrections to the condensed fraction, the critical tem-

^{*}a.boudjemaa@univ-chlef.dz

perature, the Lee-Huang-Yang (LHY) equation of state (EoS) and the superfluid fraction using the Hartree-Fock-Bogoliubov (HFB) theory based on a minimal length framework. Our results reveal that the interplay of DDIs and QG corrections tends to improve bounds on the GUP parameters. At higher temperatures, our results reproduce those obtained for an ideal Bose gas, as reported in [48].

The paper is organized as follows. In Sec. II, we introduce the fundamental concepts of dipolar Bose gases. Section III discusses the ground-state properties of homogeneous dipolar Bose gases under the LQGUP. We compute in particular QG corrections to the condensed fraction, the anomalous density, the LHY corrected EoS, and to the superfluid fraction. Section IV delves into the experimental tests of quantum gravity. Finally, we present our conclusions in Sec. V.

II. THEORY OF DIPOLAR BOSE GASES

We consider a three-dimensional (3D) dilute dipolar Bose gas involving quantum and thermal fluctuations at temperature T. Assuming that the dipoles are oriented perpendicularly to the plane. The total interaction between two atoms of mass m and dipole d at \mathbf{r} and \mathbf{r}' , can be represented as:

$$V(\mathbf{r} - \mathbf{r}') = g\delta(\mathbf{r} - \mathbf{r}') + \frac{C_{\rm dd}}{4\pi} \frac{1 - 3\cos^2\theta}{|\mathbf{r}^3 - \mathbf{r}'^3|}.$$
 (1)

The first term accounts for contact interactions component related to the s-wave scattering length a through a coupling strength $g = (4\pi\hbar^2 a/m)$. The second term is the DDI, where the coupling constant $C_{\rm dd}$ is $M_0 M^2$ for particles having a permanent magnetic dipole moment M $(M_0$ is the magnetic permeability in vacuum) and d^2/ϵ_0 for particles having a permanent electric dipole d (ϵ_0 is the permittivity of vacuum), and θ is the angle between the relative position of the particles and the direction of the dipole. The characteristic dipole-dipole distance can be defined as $r_* = mC_{\rm dd}/4\pi\hbar^2$ [53].

The Hamiltonian of the system reads:

$$\hat{H} = \int d\mathbf{r} \,\hat{\psi}^{\dagger}(\mathbf{r}) \left(\frac{-\hbar^2}{2m} \nabla^2 + U(\mathbf{r}) - \mu\right) \hat{\psi}(\mathbf{r}) \qquad (2)$$
$$+ \frac{1}{2} \int d\mathbf{r} \int d\mathbf{r}' \,\hat{\psi}^{\dagger}(\mathbf{r}) \hat{\psi}^{\dagger}(\mathbf{r}') V(\mathbf{r} - \mathbf{r}') \hat{\psi}(\mathbf{r}') \hat{\psi}(\mathbf{r}),$$

where $U(\mathbf{r})$ is the trapping potential, μ is the chemical potential, $\hat{\psi}^{\dagger}$ and $\hat{\psi}$ denote, respectively the usual creation and annihilation field operators, satisfying the usual canonical commutation relations: $[\hat{\psi}(\mathbf{r}), \hat{\psi}^{\dagger}(\mathbf{r}')] = \delta(\mathbf{r} - \mathbf{r}')$.

The dynamics of dipolar BECs including the normal and anomalous fluctuations is govened by the nonlocal generalized Gross-Pitaevskii equation (GGPE) which can be derived through $i\hbar\dot{\Phi} = \partial\langle H \rangle / \partial\Phi^*$, where $\Phi(\mathbf{r}) =$ $\hat{\psi}(\mathbf{r}) - \bar{\psi}(\mathbf{r})$, is the condensate wavefunction and $\bar{\psi}(\mathbf{r})$ is the noncondensed part of the field operator. This yields [54, 55]:

$$i\hbar\dot{\Phi} = \left[\frac{-\hbar^2}{2m}\nabla^2 + U(\mathbf{r}) - \mu + \delta\mu_{\rm LHY} + \int d\mathbf{r}' V(\mathbf{r} - \mathbf{r}')n(\mathbf{r}')\right]\Phi,$$
(3)

where

$$\delta\mu_{\rm LHY}(\mathbf{r})\Phi(\mathbf{r}) = \int d\mathbf{r}' V(\mathbf{r} - \mathbf{r}') \bigg[\tilde{n}(\mathbf{r}, \mathbf{r}')\Phi(\mathbf{r}') \qquad (4)$$
$$+ \tilde{m}(\mathbf{r}, \mathbf{r}')\Phi^*(\mathbf{r}') \bigg],$$

accounts for the LHY quantum corrections to the EoS [54, 56]. In Eq. (4), $\tilde{n}(\mathbf{r}, \mathbf{r}') = \langle \hat{\psi}^{\dagger}(\mathbf{r}) \hat{\psi}(\mathbf{r}') \rangle$ and $\tilde{m}(\mathbf{r}, \mathbf{r}') = \langle \hat{\psi}(\mathbf{r}) \hat{\psi}(\mathbf{r}') \rangle$ are respectively, the normal and anomalous correlation functions represent the dipole exchange interaction between the condensed and noncondensed atoms. In the local limit they reduce, respectively to the noncondensed $\tilde{n}(\mathbf{r})$ and anomalous $\tilde{m}(\mathbf{r})$ densities. The total density is defined as: $n(\mathbf{r}) = n_c(\mathbf{r}) + \tilde{n}(\mathbf{r})$, where $n_c(\mathbf{r}) = |\Phi(\mathbf{r})|^2$ is the condensed density.

The low-lying collective excitations of dipolar BECs can be obtained within the so-called Bogoliubov-de-Gennes equations (BdGE) by considering the small oscillations of the order parameter: $\Phi(\mathbf{r},t) = [\Phi_0(\mathbf{r}) + \delta \Phi(\mathbf{r},t)] \exp(-i\mu t/\hbar)$, where $\delta \Phi(\mathbf{r},t) = u_{\nu}(\mathbf{r}) \exp(-i\varepsilon_{\nu}t/\hbar) + v_{\nu}(\mathbf{r}) \exp(i\varepsilon_{\nu}t/\hbar)$ are small quantum fluctuations with ε_{ν} being the Bogoliubov excitations energy of mode ν . The quasi-particle amplitudes $u_{\nu}(\mathbf{r})$ and $v_{\nu}(\mathbf{r})$ satisfy the following nonlocal BdGEs:

$$\varepsilon_{\nu}u_{\nu}(\mathbf{r}) = \hat{\mathcal{L}}u_{\nu}(\mathbf{r}) + \int d\mathbf{r}' V(\mathbf{r} - \mathbf{r}')n(\mathbf{r}, \mathbf{r}')u_{\nu}(\mathbf{r}') + \int d\mathbf{r}' V(\mathbf{r} - \mathbf{r}')\bar{m}(\mathbf{r}, \mathbf{r}')v_{\nu}(\mathbf{r}'), \quad (5)$$

$$-\varepsilon_{\nu}v_{\nu}(\mathbf{r}) = \hat{\mathcal{L}}v_{\nu}(\mathbf{r}) + \int d\mathbf{r}' V(\mathbf{r} - \mathbf{r}')n(\mathbf{r}, \mathbf{r}')v_{\nu}(\mathbf{r}') + \int d\mathbf{r}' V(\mathbf{r} - \mathbf{r}')\bar{m}(\mathbf{r}, \mathbf{r}')u_{\nu}(\mathbf{r}'), \qquad (6)$$

where $\hat{\mathcal{L}} = (-\hbar^2/2m)\nabla^2 + U(\mathbf{r}) - \mu + \int d\mathbf{r}' V(\mathbf{r} - \mathbf{r}')n(\mathbf{r}')$. In the next sections, we will solve these coupled BdGE analytically for the case of a homogeneous dipolar Bose gas in the presence of QG effects. They enable us to look at how the intriguing interplay of DDIs and QG corrections may improve bounds on the GUP parameters.

III. HOMOGENENOUS DIPOLAR BOSE GASES UNDER THE GUP

The LQGUP taking quantum gravity effects into account is derived from the following commutation relation [30]:

$$[r_i, p_j] = i\hbar \left[\delta_{ij} - \alpha \left(p\delta_{ij} + \frac{p_i p_j}{p} \right) + \beta \left(p^2 \delta_{ij} + 3p_i p_j \right) \right],$$
(7)

where x_i and p_j are the position and momentum operators, respectively and $p = \sqrt{p_i p_j}$, $\alpha = \alpha_0 l_p / \hbar =$ $\alpha_0/(M_pc)$, and $\beta = \beta_0 l_p^2/\hbar^2 = \beta_0/(M_pc)^2$, where α_0 and β_0 are the linear and quadratic deformation GUP parameters which are related to the Planck length, l_p , and the Planck mass $M_p = \sqrt{\hbar c/G}$ with G being the gravitational constant and c denoting the speed of light in vacuum. According to the authors of Ref. [30], the linear correction is suggested by doubly special relativity theories. Current experiments can set upper bounds on the GUP parameter. For instance, the standard model of high-energy physics implies that $\beta_0 < 10^{34}$ [19]. According to the same reference [19], the scanning tunneling microscope delivers the best one $\beta_0 < 10^{21}$ [19]. Other upper bounds have been provided by different approaches, namely the Lamb shift and Landau levels [19], optical systems [39], the light deflection and perihelion precession [57], cold atoms [58], and gravitational systems [59– 61]. However, in the present work we will address QG effects for arbitrary β_0 . For $\alpha = \beta = 0$, one can reproduce the standard Heisenberg uncertainty principle.

One should stress that Eq. (7) implies that standard operator for momentum cannot be used, $p_i \neq -i\hbar\nabla$. However, we introduce a set of canonical operators x_{0i} and p_{0i} , which satisfy a standard commutation relation $[r_{0i}, p_{0i}] = i\hbar\delta_{ij}$ [30]. Doing so, we can write

$$r_i = r_{0i}, \qquad p_i = p_{0i}(1 - \alpha p_0 + 2\beta p_0^2), \qquad (8)$$

where $p = \sqrt{p_{0i}p_{0i}}$.

Furthermore, according to Eq. (7), the deformed density of states, g(E), can be given by [48]

$$\sum_{\mathbf{n}} = \frac{V}{(2\pi\hbar)^3} \int_0^\infty d^3p \qquad (9)$$
$$= \frac{1}{2} \int_0^\pi \sin\theta d\theta \int_0^\infty g(E) \, dE,$$

where

$$g(E) = \frac{(2m)^{3/2}}{4\pi^2\hbar^3} E^{1/2} (1 + 16\alpha\sqrt{m}E^{1/2} - 25\beta mE),$$
(10)

with $E = p^2/2m$ being the energy of free particle [48]. Equations (8) and (9) show that the LQGUP can alter not only the excitations energy but also the statistical and the thermodynamic properties of a weakly interacting dipolar Bose gas.

For concreteness, let us consider a homogeneous 3D Bose gas $(U(\mathbf{r}) = 0)$ with DDI enclosed in a volume V. These unifrom systems constitute a prototype for the experimentally relevant trapped ultracold gases and often lead to the correct physical intuition with respect to their properties. In such a case, the densities n_c , \tilde{n} , and \tilde{m} are



FIG. 1. Bogoliubov spectrum for different values of the GUP parameters, $\overline{\alpha}$, with $\theta = \pi/2$, $\epsilon_{\rm dd} = 0.16$. Here we set $\overline{\alpha} = \alpha m c_{s0} = \alpha_0 (m c_{s0}/M_p c)$ and $\beta_0 = \alpha_0^2$ [48].

constant and the momentum-space interaction is independent of the magnitude of \mathbf{p} and instead depends only on its direction.

From now on we will replace p_0 by p for simplicity. Taking into account effects of QG implied by the LQGUP given in Eq. (8), the GGPE (3) turns out to be given as:

$$i\hbar\dot{\Phi} = \left[\mathcal{E}_{\mathbf{p}} + V(|\mathbf{p}| = 0)n + \delta\mu_{\rm LHY}(\mathbf{p}) - \mu\right]\Phi, \quad (11)$$

where $\mathcal{E}_p = (p^2/2m) \left(1 - 2\alpha p + 5\beta p^2\right)$, and

$$V(\mathbf{p}) = g[1 + \epsilon_{\rm dd}(3\cos^2_{\mathbf{p}}\theta - 1)], \qquad (12)$$

is the Fourier transform of the interaction potential (1), $\epsilon_{\rm dd} = C_{\rm dd}/3g$ is the dimensionless relative strength describing the interplay between the DDI and contact interaction, and θ being the angle between the vector **p** and the polarization direction. Note that the relation (12) is valid only in the ultracold limit where the particle momenta satisfy the inequality $pr_*/\hbar \ll 1$ [49].

The chemical potential is obtained via Eq. (11) [55]

$$\mu = V(|\mathbf{p}| = 0)n + \delta\mu_{\text{LHY}}(\mathbf{p})$$
(13)
$$= V(|\mathbf{p}| = 0)n + \frac{1}{V} \sum_{\mathbf{p}\neq\mathbf{0}} V(\mathbf{p}) \big(\tilde{n}_p + \tilde{m}_p\big),$$

where \tilde{n}_p and \tilde{m}_p stand for the normal and anomalous distributions (see below).

A. Excitations energy

The weakly interacting regime suggests $\tilde{n}/n \ll 1$ and $\tilde{m}/n \ll 1$ [64, 65]. Then, the Bogoliubov excitations

energy can be obtained via the BdGEs (5) and (6), where the Bogoliubov quasiparticle amplitudes take the form $u_p, v_p = (\sqrt{\varepsilon_p/\mathcal{E}_p} \pm \sqrt{\mathcal{E}_p/\varepsilon_p})/2$ [62]. After some algebra we find the QG corrected Bogoliubov dispersion relation:

$$\varepsilon_p = \sqrt{\mathcal{E}_p^2 + 2V(\mathbf{p})\mathcal{E}_p}.$$
 (14)

For small momenta $p \to 0$, the Bogoliubov dispersion relation is phonon-like $\varepsilon_p = c_s(\theta)p$ (quanta of sound waves), where $c_s(\theta) = c_{s0}\sqrt{1 + \epsilon_{dd}(3\cos^2\theta - 1)}$ is the sound velocity which is anisotropic owing to the DDI, with $c_{s0} = \sqrt{gn/m}$ being the sound velocity of a nondipolar BEC. In the high momenta limit $p \to \infty$, the excitations spectrum (14) reduces to $\varepsilon_p = \mathcal{E}_P$. For $\alpha = \beta = 0$, one recovers the standard Bogoliubov energy [62]. It is important to note that, for $\epsilon_{dd} > 1$, the spectrum (14) becomes imaginary giving rise to the collapse of homogeneous dipolar BEC with dominant DDI. Effects of QG on the Bogoliubov excitations spectrum are singinifcant notably for large momenta, $p/mc_{s0} \geq 1$, as shown in Fig. 1.

B. Condensed fraction and transition temperature

In the realm of the Bogoliubov theory, \tilde{n}_p and \tilde{m}_p are defined as:

$$\tilde{n}_p = v_p^2 + (u_p^2 + v_p^2)N_p, \qquad (15)$$

and

$$\tilde{m}_p = u_p v_p (2N_p + 1), \tag{16}$$

where $N_p = [\exp(\varepsilon_p/T) - 1]^{-1}$ are occupation numbers for the excitations. Therefore, explicit formulas for the noncondensed and the anomalous densities can be obtained using the definitions $\tilde{n} = V^{-1} \sum_{\mathbf{p}} \tilde{n}_p$ and $\tilde{m} = -V^{-1} \sum_{\mathbf{p}} \tilde{m}_p$, and the QG corrected density of states in Eq. (10):

$$\tilde{n} = \frac{1}{4} \int_0^{\pi} \sin\theta d\theta \int_0^{\infty} g(E) dE \left[\frac{\mathcal{E}_E + mc_s^2(\theta)}{\varepsilon_E} \right] \quad (17)$$
$$\times \left[\coth\left(\frac{\varepsilon_E}{2T}\right) - 1 \right],$$

and

$$\tilde{m} = -\frac{1}{4} \int_0^\pi \sin\theta d\theta \int_0^\infty g(E) \, dE \frac{mc_s^2(\theta)}{\varepsilon_E} \coth\left(\frac{\varepsilon_E}{2T}\right).$$
(18)

Leading terms in Eqs.(17) and (18) are the zerotemperature contribution to the noncondensed \tilde{n}_0 and anomalous \tilde{m}_0 densities, respectively. Subleading terms represent the contribution of the so-called thermal fluctuations and we denote them as \tilde{n}_T and \tilde{m}_T , respectively. Equations (17) and (18) allow us to determine in a very useful way the condensed fraction and the critical temperature of Bose quantum liquids (see below).

At zero temperature, integral (17) gives for the quantum depletion: $\tilde{n}_0 = (1/3\pi^2)(mc_{s0}/\hbar)^3 f(\overline{\alpha}, \epsilon_{dd})$, where

$$f(\overline{\alpha}, \epsilon_{\rm dd}) = \frac{3}{4\sqrt{2}} \int_0^{\pi} \sin\theta d\theta \int_0^{\infty} dx \sqrt{x} \left(-25\overline{\beta}x^2 + 16\sqrt{x}\overline{\alpha} + 1\right)$$

$$\times \left(\frac{(x^2/2) \left(5\overline{\beta}x^2 - 2\overline{\alpha}x + 1\right) + \epsilon_{\rm dd} \left(3\cos^2\theta - 1\right) + 1}{\sqrt{x^2 \left(5\overline{\beta}x^2 - 2\overline{\alpha}x + 1\right) \left(\epsilon_{\rm dd} \left(3\cos^2\theta - 1\right) + 1\right) + \left(x^4/4\right) \left(5\overline{\beta}x^2 - 2\overline{\alpha}x + 1\right)^2} - 1\right),$$

$$(19)$$

where $\overline{\alpha} = \alpha m c_{s0} = \alpha_0 (m c_{s0}/M_p c)$, and $\overline{\beta} = \beta (m c_{s0})^2 = \beta_0 (m c_{s0}/M_p c)^2$. The result of the numerical integration of Eq. (19) is shown in Fig. 2. We see that the function f increases for $\overline{\alpha} \leq 0.5$, reaches its maximum at $\overline{\alpha} \sim 0.5$, then it decreases for larger $\overline{\alpha}$ regardless of the value of the DDI strength, ϵ_{dd} (see Fig. 2.a). This clearly reveals that the QG effects may significantly modify the quantum depletion and the condensed fraction of the condensate. Surprisingly, the deformation function f is lowering with ϵ_{dd} which is in contrast to dipolar BEC without GUP (see Fig. 2.b). For $\overline{\beta} = \overline{\alpha} = 0$, we reproduce the results of the depletion for a dipolar BEC without GUP, $\tilde{n}_0 = (1/3\pi^2)(m c_{s0}/\hbar)^3 Q_3(\epsilon_{dd})$ [63–65],

where the DDI contribution is described by the function $Q_3(\epsilon_{\rm dd})$ (see Fig. 3 (right panel)), which is special case j = 3 of $Q_j(\epsilon_{\rm dd}) = (1 - \epsilon_{\rm dd})^{j/2} {}_2F_1\left(-\frac{j}{2}, \frac{1}{2}; \frac{3}{2}; \frac{3\epsilon_{\rm dd}}{\epsilon_{\rm dd}-1}\right)$, where ${}_2F_1$ is the hypergeometric function. Note that functions $Q_j(\epsilon_{\rm dd})$ attain their maximal values for $\epsilon_{\rm dd} \approx 1$ and become imaginary for $\epsilon_{\rm dd} > 1$.

For vanishing quadratic QG correction ($\overline{\alpha} = 0$) and for $\epsilon_{\rm dd} = 0$, one can expect that the depletion (19) reduces to our recent result [53].

At low temperatures $T \ll mc_{s0}^2$, the main contribution to integral (17) comes from the low energy branch where $\mathcal{E} \approx E$ and $\varepsilon \approx c_E(\theta)\sqrt{2mE}$. A straightforward calcula-



FIG. 2. Deformation function $f(\overline{\alpha}, \epsilon_{\rm dd})$ which governs the dependence of the condensate depletion as a function of the deformation parameter α in units of mc_{s0} .



FIG. 3. Dipolar functions Q_j vs. the dipolar interaction parameter ϵ_{dd} .

tion gives for the condensed fraction, $n_c/n = 1 - \tilde{n}/n$:

$$\frac{n_c}{n} = 1 - \left(\frac{T}{T_c^0}\right)^2 \left[\frac{\pi^2 \mathcal{Q}_{-1}(\epsilon_{\rm dd})}{3\zeta(3/2)^{4/3}} \left(\xi n^{1/3}\right) + \overline{\alpha} \frac{32\pi\zeta(3)\mathcal{Q}_{-2}(\epsilon_{\rm dd})}{\sqrt{2}\zeta(3/2)^{2/3}} \left(\xi n^{1/3}\right)^3 \left(\frac{T}{T_c^0}\right) - \overline{\beta} \frac{10\pi^6 \mathcal{Q}_{-3}(\epsilon_{\rm dd})}{3\zeta(3/2)^{8/3}} \left(\xi n^{1/3}\right)^5 \left(\frac{T}{T_c^0}\right)^2 \right],$$
(20)

where $\xi = \hbar/(mc_{s0})$ is the healing length and $T_c^0 = (2\pi\hbar^2/\zeta(3/2)^{2/3}m)n^{2/3}$ is the ideal gas transition temperature. In Eq. (20) we utilized the identity $\int_0^\infty x^j dx/(e^x - 1) = \Gamma(j + 1)\zeta(j + 1)$, where $\Gamma(x)$ is the gamma function and $\zeta(x)$ is the Riemann zeta function. For $\overline{\beta} = \overline{\alpha} = 0$, one recovers the results of the depletion for a dipolar BEC without GUP, $\tilde{n}_T = (mT^2/12\hbar^3c_{s0})Q_{-1}(\epsilon_{dd})$ [64, 65]. The behavior of the functions $Q_{-1}(\epsilon_{dd})$, $Q_{-2}(\epsilon_{dd})$, and $Q_{-3}(\epsilon_{dd})$ is displayed in Fig. 3. Equation (20) shows also that the QG corrections (second and last terms in r.h.s) increase with increasing the density n and temperature, T/T_c^0 , and with decreasing boson mass m.

To illustrate our results, we consider two cases namely: 52 Cr atoms with $\epsilon_{dd} = 0.16$ [50] and 168 Er atoms with $\epsilon_{dd} = 0.38$ [66].

Figures 4 (a) and (b) depict that the condensed fraction n_c/n decreases with the reduced temperature, T/T_c^0 ,



FIG. 4. (a) Condensed fraction of Cr BEC, n_c/n , as a function of the reduced temperature, T/T_c^0 , for different values of $\overline{\alpha}$ with $\epsilon_{\rm dd} = 0.16$ and $\xi n^{1/3} = 0.93$ [50]. (b) The same as Fig. 4.a but for Er BEC with $\epsilon_{\rm dd} = 0.38$ and $\xi n^{1/3} = 0.7$ [66]. (c) Condensed fraction as a function of the GUP parameter, $\overline{\alpha}$, for different values of reduced temperature, T/T_c^0 , with $\epsilon_{\rm dd} = 0.16$ and $\xi n^{1/3} = 0.93$. (d) Reduced temperature T/T_c^0 as a function of the condensed fraction, n_c/n , for different values of $\overline{\alpha}$ with $\xi n^{1/3} = 0.93$ and $\epsilon_{\rm dd} = 0.16$.

regardless of the DDI relative strength, $\epsilon_{\rm dd}$. It decays also with the GUP parameter $\overline{\alpha}$ notably for large T as shown in Fig. 4 (c) indicating that QG effects may reduce the condensed density and thus enhance the thermal cloud. For instance, at $T = 0.2T_c^0$, one has $n_c/n \simeq 98\%$ for $\overline{\alpha} = 0$ then it reduces to $n_c/n \simeq 94\%$ for $\overline{\alpha} = 0.1$. Whereas at $T = 0.6T_c^0$, the condensed fraction is about $n_c/n \simeq 70\%$ while it decays to $n_c/n \simeq 10\%$ for $\overline{\alpha} = 0.1$. In Fig. 4 (d) we plot the reduced temperature T/T_c^0 as a function of the condensed fraction, n_c/n , for different values of $\overline{\alpha}$. We see that the GQ corrections lower T/T_c^0 only for relatively small n_c/n . For large n_c/n , the reduced temperature is almost insensitive to the GQ effects. This downshift is a clear indication of the significance of the GQ effects in particular at higher temperature when thermal fraction becomes important.

However, at higher temperatures, $T \gg mc_{s0}^2$, where the main contribution to integral (17) comes from the high energy branch, our results coincide with those obtained in [30] for ideal Bose gases.



FIG. 5. (a) Thermal contribution to the LHY corrected EoS of Cr BEC, $\mu_{\rm LHY}/(ng)$, as a function of the reduced temperature, T/T_c^0 , for different values of $\overline{\alpha}$ with $\epsilon_{\rm dd} = 0.16$ and $\xi n^{1/3} = 0.93$ [50]. (b) The same as Fig. (5.a) but for Er BEC with $\epsilon_{\rm dd} = 0.38$ and $\xi n^{1/3} = 0.7$ [66].

C. Equation of state

Corrections to the chemical potential due to the LHY quantum fluctuations can be derived by combining Eqs. (15), (16) and (13). This yields:

$$\mu_{\rm LHY} = \frac{1}{4} \int_0^{\pi} \sin\theta d\theta \int_0^{\infty} g(E) \, dE \, \tilde{V}(E) \qquad (21)$$
$$\times \left[\frac{\mathcal{E}_E}{\varepsilon_E} \coth\left(\frac{\varepsilon_E}{2T}\right) - 1 \right].$$

This equation is appealing since it enables us to calculate the LHY corrections to all thermodynamic quantities. At low temperatures, the LHY corrected EoS can be written as:

$$\mu_{\rm LHY} = \frac{\pi^2 g T^4}{60m\hbar^3 c_{s0}^5} \left[\frac{\mathcal{Q}_{-3}(\epsilon_{\rm dd})}{8} + \overline{\alpha} \frac{720\zeta(5)\mathcal{Q}_{-4}(\epsilon_{\rm dd})}{\sqrt{2}} \left(\frac{T}{c_{s0}} \right)^2 - \overline{\beta} \frac{125\pi^6 \mathcal{Q}_{-5}(\epsilon_{\rm dd})}{42} \left(\frac{T}{c_{s0}} \right)^2 \right].$$
(22)

In terms of the reduced temperature, $\mu_{\rm LHY}$ reads

$$\frac{\mu_{\rm LHY}}{ng} = \left(\frac{T}{T_c^0}\right)^4 \left[\frac{\pi^6 \mathcal{Q}_{-3}(\epsilon_{\rm dd})}{30\zeta(3/2)^{8/3}} (\xi n^{1/3})^5 + \overline{\alpha} \frac{384\pi^7 \zeta(5) \mathcal{Q}_{-4}(\epsilon_{\rm dd})}{\sqrt{2}\zeta(3/2)^{10/3}} (\xi n^{1/3})^7 \left(\frac{T}{T_c^0}\right) - \overline{\beta} \frac{200\pi^{10} \mathcal{Q}_{-5}(\epsilon_{\rm dd})}{63\zeta(3/2)^4} (\xi n^{1/3})^9 \left(\frac{T}{T_c^0}\right)^2\right].$$
(23)

Again for $\overline{\beta} = \overline{\alpha} = 0$, μ_{LHY} of Eq. (23) simplifies to that for a dipolar BEC without GUP, $g\pi^2 T^4 \mathcal{Q}_{-3}(\epsilon_{\text{dd}})/(60m\hbar^3 c_{s0}^5)$ [64, 65]. The behavior of the functions $Q_{-5}(\epsilon_{\text{dd}})$, and $Q_{-7}(\epsilon_{\text{dd}})$ is shown in Fig. 3.

Figure 5 shows that for $\overline{\alpha} \gtrsim 2 \times 10^{-2}$, the thermal contribution to the LHY corrected EoS, $\mu_{\rm LHY}/(ng)$, deviates from that of a dipolar BEC without GUP at higher temperatures owing to QG effects. Remarkably, this discrepancy becomes pronounced for small $\epsilon_{\rm dd}$ (see Fig. 5 (a)).

D. Superfluidity

Now we focus on the behavior of the superfluid fraction of a dipolar BEC under GUP. In a 3D dipolar BEC the superfluid density n_s is a tensor quantity with components n_s^{ij} due to the peculiar anisotropy property of the DDI [67, 68]. This means that n_s depends on the direction of the superfluid motion with respect to the orientation of the dipoles. It can be found by applying a Galilean boost with the total momentum of the moving system $\mathbf{P} = mV(n\mathbf{v_s} + n_n\mathbf{v_n})$, where $\mathbf{v_s}$ denotes the superfluid velocity and $\mathbf{v_n} = \mathbf{u} - \mathbf{v_s}$ is normal fluid velocity with \mathbf{u} being a boost velocity [67, 68]. Keeping only linear term in P, we get

$$\frac{n_s^{ij}}{n} = \delta_{ij} - \frac{1}{Tnm} \int \frac{d^3p}{(2\pi\hbar)^3} \left[\frac{p_i p_j}{4\sinh^2(\varepsilon_k/2T)} \right]. \quad (24)$$

Applying the QG corrected density of states in Eq. (10), the superfluid fraction (24) turns out to be given:

$$\frac{n_s^{ij}}{n} = \delta_{ij} - \frac{1}{Tnm} \int_0^\pi \sin\theta d\theta \int_0^\infty g(E) \, dE \left[\frac{\sqrt{E_i E_j}}{4 \sinh^2(\varepsilon_k/2T)} \right]$$
(25)

At low temperatures $T \ll ng$, the parallel direction of the superfluid fraction reads

$$\frac{n_s^{\parallel}}{n} = 1 - \left(\frac{T}{T_c^0}\right)^4 \left[\frac{\pi^7 \mathcal{Q}_{-5}^{\parallel}(\epsilon_{\rm dd})}{30\zeta(3/2)^{8/3}} (\xi n^{1/3})^5 + \overline{\alpha} \frac{480\pi^4 \zeta(5) \mathcal{Q}_{-6}^{\parallel}(\epsilon_{\rm dd})}{\sqrt{2}\zeta(3/2)^{10/3}} (\xi n^{1/3})^6 \left(\frac{T}{T_c^0}\right) - \overline{\beta} \frac{25\pi^{11} \mathcal{Q}_{-7}^{\parallel}(\epsilon_{\rm dd})}{8\zeta(3/2)^4} (\xi n^{1/3})^9 \left(\frac{T}{T_c^0}\right)^2\right]. \tag{26}$$

where the functions

$$\mathcal{Q}_{j}^{\parallel}(\epsilon_{\rm dd}) = \frac{2(1-\epsilon_{\rm dd})^{j/2}}{9(j+1)(j+3)\epsilon_{\rm dd}^{2}} \left[(\epsilon_{\rm dd}-1)^{2} {}_{2}F_{1}\left(-\frac{1}{2},-\frac{j}{2};\frac{1}{2};\frac{3\epsilon_{\rm dd}}{\epsilon_{\rm dd}-1}\right) + (2\epsilon_{\rm dd}+1)((3j+4)\epsilon_{\rm dd}-1)\left(1-\frac{3\epsilon_{\rm dd}}{\epsilon_{\rm dd}-1}\right)^{j/2} \right],$$

behave as $Q_j^{\parallel}(\epsilon_{dd} = 0) = 2/3$ and imaginary for $\epsilon_{dd} > 1$ (see also Fig. 6.a). In the perpendicular direction, one has

$$\frac{n_s^{\perp}}{n} = 1 - \left(\frac{T}{T_c^0}\right)^4 \left[\frac{\pi^7 \mathcal{Q}_{-5}^{\perp}(\epsilon_{\rm dd})}{30\zeta(3/2)^{8/3}} (\xi n^{1/3})^5 + \overline{\alpha} \frac{480\pi^4 \zeta(5) \mathcal{Q}_{-6}^{\perp}(\epsilon_{\rm dd})}{\sqrt{2}\zeta(3/2)^{10/3}} (\xi n^{1/3})^6 \left(\frac{T}{T_c^0}\right) - \overline{\beta} \frac{25\pi^{11} \mathcal{Q}_{-7}^{\perp}(\epsilon_{\rm dd})}{8\zeta(3/2)^4} (\xi n^{1/3})^9 \left(\frac{T}{T_c^0}\right)^2\right]. \tag{27}$$

where $Q_j^{\perp}(\epsilon_{dd}) = Q_j(\epsilon_{dd}) - Q_j^{\parallel}(\epsilon_{dd})$ which increases monotonically with ϵ_{dd} and becomes complex for $\epsilon_{dd} > 1$ (see Fig. 6.b). For $\overline{\beta} = \overline{\alpha} = 0$, the superfluid fraction in both directions reduces to that obtained for a dipolar BEC without GUP [67, 68].



FIG. 6. Parallel $\mathcal{Q}_{j}^{\parallel}(\epsilon_{dd})$ (a) and perpendicular $\mathcal{Q}_{j}^{\perp}(\epsilon_{dd})$ (b) dipolar functions and vs. the dipolar interaction parameter ϵ_{dd} .

Figure 7 depicts that the superfluid fraction is decreasing with the reduced temperature in both directions, regardless of the value of the GUP parameter and of DDI strength. A direct comparison between both components shows that n_s^{\parallel} and n_s^{\perp} coincide for $\overline{\alpha} = 0$ and for $\epsilon_{dd} = 0$ and hence, well reproduce the standard two-body contact interaction result. We see also that for relatively high temperatures $T \gtrsim 0.45 T_c^0$ and for $\epsilon_{\rm dd} = 0.16, n_s^{\perp}$ is slightly larger than n_s^{\parallel} while the situation is inverted in the case of Er BEC with $\epsilon_{\rm dd} = 0.38$ where $n_s^{\parallel} > n_s^{\perp}$. This reveals that QG effects on the superfluidity could be pronounced in perpendicular direction rather than in the parallel direction depending on the relative strength of the DDI. Another important remark is that the superfluid fraction increases with the GUP parameter in contrast to the condensed fraction. For sufficiently large values of $\overline{\alpha}$, it diverges from that of the ordinary dipolar BEC in both components. Our results would provide unique new insight into the physics of superfluidity in neutron stars [69] and thus will furnish a probe of the neutron star interior.



FIG. 7. Superfluid fractions n_s^{\parallel}/n (a) and n_s^{\perp}/n (b) of Cr BEC, as a function of the reduced temperature, T/T_c^0 , for different values of $\overline{\alpha}$ with $\epsilon_{\rm dd} = 0.16$ and $\xi n^{1/3} = 0.93$ [50]. (c) -(d) The same as Figs. 4.(a) and (b) but for Er BEC with $\epsilon_{\rm dd} = 0.38$ and $\xi n^{1/3} = 0.7$ [66].

IV. EXPERIMENTAL TEST OF QUANTUM GRAVITY

In order to constrain parameters in the previous GUP proposal, it is necessary to relate it to two observables namely: the condensed fraction (20) and the superfluid fraction obtained from Eqs. (26) and (27). In our analysis, two kinds of dipolar atomic systems are being treated. The first consists of ⁵²Cr atoms with *s*-wave scattering length $a = 100 a_0$ (a_0 is the Bohr radius) and relative

DDI strength $\epsilon_{\rm dd} = 0.16$ [50]. Next we consider ¹⁶⁸Er BEC with $a = 175 a_0$, and $\epsilon_{\rm dd} = 0.38$ [66]. The average density of both species is $n = 5 \times 10^{20} {\rm m}^{-3}$.

	$\epsilon_{ m dd}$	T/T_c^0	n_c/n	$lpha_0$	eta_0
^{52}Cr	0.16	0.20	95%	3.42×10^{22}	1.17×10^{45}
	0.16	0.95	15%	1.03×10^{25}	1.05×10^{50}
¹⁶⁸ Er	0.38	0.20	97%	2.60×10^{22}	6.70×10^{44}
	0.38	0.95	35%	7.76×10^{24}	6.03×10^{49}

TABLE I. Typical values of α_0 and β_0 extracted from the superfluid fraction of ⁵²Cr atoms [50] and ¹⁶⁸Er atoms [66].

$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		T/T_c^0	n_s^{\parallel}/n	n_s^{\perp}/n	$lpha_0$	eta_0
168 Er 0.35 08% 08% 2.60 × 10 ²⁴ 6.70×1	$^{52}\mathrm{Cr}$	0.35	95%	95% 45%	3.42×10^{24}	1.17×10^{49} 4.70×10^{51}
EI 0.35 3670 3670 2.00×10 0.70×1	¹⁶⁸ Er	0.35	98%	98%	2.60×10^{24}	6.70×10^{48}

TABLE II. Typical values of α_0 and β_0 extracted from the superfluid fraction of ⁵²Cr atoms [50] and ¹⁶⁸Er atoms [66].

Table I shows that at sufficiently low temperature, $T/T_c^0 \simeq 0.2$, where the ground-state population n_c/n is large, our model predicts for the GUP parameters

 $\alpha_0 \sim 10^{22}$ and $\beta_0 \sim 10^{44}$. Clearly, these values improve the bounds set by the model of an ideal Bose gas [48] and by the model of BEC with a pure contact interaction [49]. The reason is that β_0 strongly depends on $\epsilon_{\rm dd}$. For instance, β_0 (¹⁶⁸Er) is one order of magnitude higher than β_0 (⁵²Cr).

However, the bounds on QG parameters obtained from measuring the superfluid fraction in both parallel and perpendicular directions are $\alpha_0 \sim 10^{24}$ and $\beta_0 \sim 10^{48}$ as shown in Table II. Although these bounds are better than the results obtained for weakly interacting Bose gases [49], they are weaker than those set by high-energy physics [70] and measurements of an ideal Bose gas [48]. Therefore, they are somehow not interesting compared to previous ones.

V. CONCLUSIONS

In this paper we studied effects of QG due to GUP on the ground-state properties of dipolar BECs under the aim to constrain the GUP parameters. Using the HFB-LQGUP approach we calculate corrections to the condensed fraction, the critical temperature, the EoS, and the superfluid fraction. We showed that the intriguing interplay of QG and DDIs may significantly affect these quantities. We also discussed the possible experimental tests of our theoretical predictions. Our theory predicted that better bounds require a strong relative DDI strength and a large condensed fraction (i.e. low temperatures). Compared to bounds set from high-energy physics and other experiments and theories [70], our bounds on the GUP obtained from the condensed fraction are better while those obtained from the superfluid fraction are worse. Our findings can be readily probed in current experiments, and might bring us closer to understanding whether gravity can be reconciled with quantum mechanics.

- [1] R. Penrose, On the gravitization of quantum mechanics 1: Quantum state reduction, Found. Phys. 44, 557 (2014).
- [2] D.Amati, M.Ciafaloni, G.Veneziano, Phys. Lett. B 197, 81 (1987).
- [3] F.Scardigli, Phys. Lett. B **452**, 39 (1999).
- [4] R.J.Adler, D.I.Santiago, Mod. Phys. Lett. A 14, 1371 (1999).
- [5] L. N. Chang, Z. Lewis, Dj. Minic, T. Takeuchi, Advances in High Energy Physics 2011, 493514 (2011).
- [6] M. Bishop, J. Contreras, and D. Singleton, Universe, 8, 192 (2022).
- [7] W. Heisenberg, The Physical Principles of the Quantum Theory, University of Chicago Press, Dover Publications (1930).
- [8] M. Bronstein Gen. Relativ.and Gravit. 44, 267,(2012).
- [9] S. Bose, A. Mazumdar, G. W. Morley, H. Ulbricht, M. Toros, M. Paternostro, A. A. Geraci, P. F. Barker, M. S. Kim, and G. Milburn, Phys. Rev. Lett. **119**, 240401

(2017).

- [10] A. Belenchia, R. M. Wald, F. Giacomini, E. Castro-Ruiz, C. Brukner, and M. Aspelmeyer, Phys.Rev.D 98, 126009 (2018).
- [11] M. Zych, F.Costa, I. Pikovski, and C. Brukner. Nat. Commun 10, 3772 (2019).
- [12] R. Penrose, Gen.Relativ.Gravit. 28, 581 (1996).
- [13] R. Penrose, Phil. Trans.R.Soc. A **356**, 1927 (1998).
- [14] T. Westphal, H. Hepach, J. Pfaff, M. Aspelmeyer, Nature 591, 225–228 (2021).
- [15] T. Brack, B. Zybach, F. Balabdaoui, S. Kaufmann, F. Palmegiano, J.-C. Tomasina, S. Blunier, D. Scheiwiller, J. Fankhauser, J. Dual, Nat. Phys. 18, 952–957 (2022).
- [16] Fuchs et al., Sci. Adv. 10, eadk 2949 (2024).
- [17] R. Howl, V. Vedral, D. Naik, M. Christodoulou, C. Rovelli and A. Iyer, Phys. Rev. X Quantum, 2, 010325 (2021).
- [18] K. Shiraishi, Prog. Theor. Phys. 77 975 (1987).

- [19] S. Das, and E.C. Vagenas, Phys. Rev. Lett. 101, 221301 (2008).
- [20] F. Briscese, M. Grether and M. de Llano, Euro. Phys. Lett., 98, 6 (2012).
- [21] F. Briscese, Phys. Lett. B, **718**, 214 (2012).
- [22] M. M. Dos Santos, T. Oniga, A. S. Mcleman, M. Caldwell and C. H. - T. Wang, J. Plasma Physics , **79**, 437–442 (2013).
- [23] J. Hansson, S. Francois, International Journal of Modern Physics D, 26, 1743003 (2017).
- [24] M. Jaffe, P. Haslinger, V. Xu, P. Hamilton, A. Upadhye, B. Elder, J. Khoury and H. Müller, Nat. Phys 13, 938 (2017).
- [25] S. A. Haine, New J. Phys., 23, 033020, (2021).
- [26] M. Maggiore, Phys. Lett. B 304, 65 (1993); Phys. Lett.
 B 319, 83 (1993); Phys. Rev. D 49, 5182 (1994).
- [27] A. Kempf, G. Mangano and R. B. Mann, Phys. Rev. D 52, 1108 (1995).
- [28] F. Scardigli, Phys. Lett. B 452, 39 (1999).
- [29] L. N. Chang, D. Minic, N. Okamura and T. Takeuchi, Phys. Rev. D 65, 125028 (2002).
- [30] A. F. Ali, S. Das, E. C. Vagenas, Phys.Rev.D 84, 044013 (2011).
- [31] M. Sprenger, P. Nicolini, M. Bleicher, Class. Quantum Gravity, 28, 235019 (2011).
- [32] I. Pikovski, M;R. Vanner, M. Aspelmeyer, M. Kim, C. Brukner, Nat. Phys. 8, 393 (2012).
- [33] V. Husain, S. Seahra, S. Webster, Phys. Rev. D 88, 024014 (2013).
- [34] P.Pedram, Phys. Rev. D 91, 063517 (2015).
- [35] Z. Feng, H.L. Li, X .T. Zu, and S. Z. Yang, Eur. Phys. J. C 76, 1 (2016).
- [36] H. Shababi and W. S. Chung, Phys. Lett. B 770, 445 (2017).
- [37] G. Gecim, and Y. Sucu, Phys. Lett. B 773, 391 (2017).
- [38] F. Scardigli, G. Lambiase and E. C. Vagenas, Phys. Lett. B 767, 242 (2017).
- [39] M. C. Braidotti, Z. H. Musslimani, C. Conti, Physica D 338, 34 (2017).
- [40] P. Bosso, S. Das, I. Pikovski, M. R. Vanner, Phys. Rev. A 96, 023849 (2017).
- [41] R.Casadio and F.Scardigli, Phys. Lett. B 807, 135558 (2020).
- [42] T. Fityo, Phys. Lett. A 372, 5872 (2008).
- [43] B. Vakili, M. A. Gorji, J. Stat. Mech. P10013 (2012).

- [44] E. Castellanos and C. Laemmerzahl, Phys. Lett. B 731, 1 (2014).
- [45] X. Zhang and C. Tian, Chinese. Phys. Lett. **32**, 010303 (2015).
- [46] H. L. Li, J. X. Ren, W. W. Wang, B. Yang and H. J. Shen, J. Stat. Mech. 023106 (2018).
- [47] S. Dey, V. Hussin, International Journal of Theoretical Physics 58, 3138 (2019).
- [48] S. Das and M. Fridman, Phys. Rev. D 104, 026014 (2021).
- [49] A. Boudjemâa, Eur. Phys. J. Plus 137, 256 (2022).
- [50] T. Lahaye et al., Rep. Prog. Phys. 72, 126401 (2009)
- [51] L.D. Carr, D. DeMille, R.V. Krems, and J. Ye, New. J. Phys 11, 055049 (2009).
- [52] M. A. Baranov, Physics Reports 464, 71 (2008).
- [53] A. Boudjemaa and G.V. Shlyapnikov, Phys. Rev. A 87, 025601 (2013).
- [54] A. Boudjemâa, and N. Guebli, Phys. Rev. A 102, 023302 (2020).
- [55] A. Boudjemâa, Phys. Lett. A 465, 128712 (2023).
- [56] T. D. Lee, K. Huang and C. N. Yang, Phys. Rev 106, 1135 (1957).
- [57] F. Scardigli, R. Casadio, Eur. Phys. J. C 75, 425 (2015).
- [58] D. Gao, M. Zhan, Phys. Rev. A 94, 013607 (2016).
- [59] Z. W. Feng, S. Z. Yang b, H. L. Li, X. T. Zu, Phys. Lett.B 768, 81 (2017).
- [60] J. C. S. Neves, Eur. Phys. J. C 80, 343 (2020).
- [61] A. Das, S. Das, N. R. Mansour, E. C. Vagenas, Phys. Lett. B 819, 136429 (2021).
- [62] N. N. Bogolubov, J. Phys. (Moscow) **11** 23 (1947).
- [63] Aristeu R. P. Lima and Axel Pelster, Phys. Rev. A 84, 041604 (R) (2011); Phys. Rev. A 86, 063609 (2012).
- [64] A. Boudjemâa, J. Phys. B: At. Mol. Opt. Phys. 48, 035302 (2015).
- [65] A. Boudjemâa, J. Phys. A: Math. Theor. 49, 285005 (2016).
- [66] K. Aikawa et al., Phys. Rev. Lett. 108, 210401 (2012).
- [67] B. Nikolic, A. Balaz, and A. Pelster, Phys. Rev. A 88, 013624 (2013).
- [68] A. Boudjemâa, Annals of Physics, **381**, 68 (2017).
- [69] G. Baym, C. Pethick and D. Pines, Nature 224, 673 (1969).
- [70] F. Scardigli, J. Phys.: Conf. Ser. 1275, 012004 (2019).