

# A NOTE ON THE LIOUVILLE THEOREM OF FULLY NONLINEAR ELLIPTIC EQUATIONS

DONGSHENG LI AND LICHUN LIANG

ABSTRACT. In this paper, a new method is presented to investigate the asymptotic behavior of solutions to the fully nonlinear uniformly elliptic equation  $F(D^2u) = 0$  in exterior domains. This method does not depend on the  $C^2$  regularity of  $F$  and the dimension  $n$ .

## 1. INTRODUCTION

In this paper, we are concerned with the Liouville theorem of fully nonlinear elliptic equations of the form

$$F(D^2u) = 0 \tag{1.1}$$

in the whole space and exterior domains. We assume that  $F$  is uniformly elliptic, i.e., there are two constants  $0 < \lambda \leq \Lambda < \infty$  such that

$$\lambda\|N\| \leq F(M + N) - F(M) \leq \Lambda\|N\|$$

for any  $M, N \in \mathcal{S}^{n \times n}$  with  $N \geq 0$ , where  $\mathcal{S}^{n \times n}$  is the space of all real  $n \times n$  symmetric matrices.

There have been a number of works on rigidity results for fully nonlinear elliptic equations. In view of the Evans-Krylov estimate, it is easily seen that any entire quadratic growth solution  $u$  of (1.1) must be a quadratic polynomial if  $F$  is convex or concave. In 2020, Li, Li and Yuan [5] made the following extension to the result: for  $n \geq 3$  and  $F \in C^2$ , if the solution  $u$  of (1.1) in  $\mathbb{R}^n \setminus \overline{B_1}$  has quadratic growth, then  $u$  must be close to a quadratic polynomial at infinity with the error  $|x|^{2-n}$ . Furthermore, the error term  $O(|x|^{2-n})$  was refined to  $d|x|^{2-n} + O(|x|^{1-n})$  for some constant  $d$  by Hong employing the Kelvin transformation in [4]. Recently, for  $n = 2$ , the asymptotic result was established with a logarithm term by Li and Liu [6] using the quasiconformal mapping. These asymptotic results can be applied to some degenerate elliptic equations such as Monge-Ampère equations, special Lagrangian equations, quadratic Hessian equations and inverse harmonic Hessian equations.

In this paper, our purpose is to present a different proof for the asymptotic result. What's more important is that this proof does not depend on the  $C^2$  regularity assumption on  $F$  and the dimension  $n$ .

We can now formulate our main result.

**Theorem 1.1.** *Let  $F \in C^{1, \frac{2}{n}}(\mathcal{S}^{n \times n})$  be uniformly elliptic and convex. Assume that  $u$  is a viscosity solution of (1.1) and satisfies*

$$|u(x)| \leq C|x|^2, \quad x \in \mathbb{R}^n \setminus \overline{B_1} \tag{1.2}$$

---

*Key words and phrases.* Fully Nonlinear Elliptic Equation; Exterior domain; Asymptotic behavior.

for some constant  $C > 0$ . Then there exist some  $A \in \mathcal{S}^{n \times n}$ ,  $b, e \in \mathbb{R}^n$  and  $c, d \in \mathbb{R}$  such that for any  $0 < \alpha < 1$ ,

$$u(x) = \frac{1}{2}x^T Ax + b \cdot x + d\Gamma(x) + c + \frac{e \cdot x}{|x^T(DF(A))^{-1}x|^{\frac{n}{2}}} + O(|x|^{1-n-\alpha}) \quad \text{as } |x| \rightarrow \infty,$$

where

$$\Gamma(x) = \begin{cases} |x^T(DF(A))^{-1}x|^{\frac{2-n}{2}} & n \geq 3, \\ \frac{1}{2} \log |x^T(DF(A))^{-1}x| & n = 2. \end{cases}$$

## 2. THE PROOF OF THEOREM 1.1

Now let us prove the following lemma.

**Lemma 2.1.** *Let  $u$  be a positive solution of  $a_{ij}D_{ij}u = 0$  in  $\mathbb{R}^2 \setminus \overline{B_1}$ , where  $a_{ij}$  is uniformly elliptic with the ellipticity constants  $0 < \lambda \leq \Lambda < \infty$  and  $a_{ij} \rightarrow \delta_{ij}$  as  $|x| \rightarrow \infty$ . Then for any  $0 < \alpha < 1$ , there exists a constant  $R_\alpha > 1$  depending only on  $\alpha$  and  $a_{ij}$  such that*

$$0 \leq u \leq C|x|^\alpha, \quad x \in \mathbb{R}^n \setminus \overline{B_{R_\alpha}}$$

for some constant  $C > 0$ .

*Proof. Step 1:* We will show that  $v := |x|^\alpha$  is a subsolution of  $a_{ij}D_{ij}u = 0$  in  $\mathbb{R}^2 \setminus \overline{B_{R_\alpha}}$  for some  $R_\alpha > 1$ .

A simple computation yields

$$D_{ij}v = \alpha|x|^{\alpha-2}\delta_{ij} + \alpha(\alpha-2)|x|^{\alpha-4}x_i x_j$$

and

$$\Delta v = \alpha^2|x|^{\alpha-2}.$$

Then we have

$$a_{ij}D_{ij}v = (a_{ij} - \delta_{ij})D_{ij}v + \Delta v \geq (\alpha^2 - \epsilon(|x|)\alpha(3-\alpha))|x|^{\alpha-2},$$

where  $\epsilon(|x|) \rightarrow 0$  as  $|x| \rightarrow \infty$ . Hence letting  $R_\alpha$  be chosen large enough such that

$$\alpha^2 - \epsilon(|x|)\alpha(3-\alpha) \geq 0, \quad |x| \geq R_\alpha,$$

we obtain the desired result.

*Step 2:* We will prove that

$$a := \sup_{x \in \mathbb{R}^2 \setminus \overline{B_{R_\alpha}}} \frac{u(x)}{v(x)} < +\infty.$$

To obtain a contradiction, suppose  $a = +\infty$ . Then for any constant  $\varepsilon > 0$ , there exist sequences of  $\{R_i\}_{i=1}^\infty \subset \mathbb{R}^+$  increasing and  $\{x_i\}_{i=1}^\infty$  with  $R_1 \geq R_\alpha$  and  $x_i \in \partial B_{R_i}$  such that

$$\varepsilon u(x_i) \geq v(x_i).$$

Applying the Harnack inequality to  $u$  in  $\partial B_{R_i}$ , we have

$$\varepsilon u(x) \geq C\varepsilon u(x_i) \geq Cv(x_i) = Cv(x), \quad x \in \partial B_{R_i},$$

---

THE LIOUVILLE THEOREM

---

where  $C$  depends only on  $\lambda$ ,  $\Lambda$  and  $n$ . Next, for any  $x \in \mathbb{R}^2 \setminus \overline{B_{R_\alpha}}$  with  $R_\alpha \leq |x| \leq R_i$  for some  $i \geq 4$ , by the comparison principle in  $B_{R_i} \setminus \overline{B_{R_\alpha}}$ , we obtain

$$v(x) \leq \max_{\partial B_{R_\alpha}} v + \frac{\varepsilon}{C} u(x), \quad x \in \mathbb{R}^2 \setminus \overline{B_{R_\alpha}}.$$

Then letting  $\varepsilon \rightarrow 0$  yields

$$|x|^\alpha = v(x) \leq \max_{\partial B_{R_\alpha}} v = |R_\alpha|^\alpha, \quad x \in \mathbb{R}^2 \setminus \overline{B_{R_\alpha}},$$

which leads to a contradiction.

*Step 3:* We complete this proof.

By the definition of  $a$ , there exist some  $R \geq R_\alpha$  and  $x_0 \in \partial B_R$  such that

$$u(x_0) \geq \frac{a}{2} v(x_0).$$

If  $R_\alpha \leq R \leq R_\alpha + 1$ , then we have

$$a \leq \frac{2u(x_0)}{v(x_0)} = \frac{2u(x_0)}{|R|^\alpha} \leq 2|R_\alpha|^{-\alpha} \|u\|_{L^\infty(B_{R_\alpha+1} \setminus B_{R_\alpha})}.$$

If  $R \geq R_\alpha + 1$ , by the Harnack inequality in  $\partial B_R$ , then we have

$$u(x) \geq Cu(x_0) \geq C \frac{a}{2} v(x_0) = C \frac{a}{2} v(x), \quad x \in \partial B_R,$$

where  $C$  depends only on  $\lambda$ ,  $\Lambda$  and  $n$ . Furthermore, applying the comparison principle yields

$$C \frac{a}{2} v(x) \leq C \frac{a}{2} |R_\alpha|^\alpha + u(x), \quad x \in B_R \setminus \overline{B_{R_\alpha}}.$$

Taking  $x \in \partial B_{R_\alpha+1}$ , we obtain

$$a \leq Cu(x) \leq C \|u\|_{L^\infty(B_{R_\alpha+1} \setminus B_{R_\alpha})}.$$

□

*Proof of Theorem 1.1.* By the Evans-Krylov theorem,  $u \in C^2$ . We assume  $F(0) = 0$ . The proof will be divided into five steps.

*Step 1:* We will show that there exists some  $A \in \mathcal{S}^{n \times n}$  such that

$$D^2 u(x) \rightarrow A \quad \text{as } |x| \rightarrow \infty.$$

For any  $e \in \mathbb{R}^n$  with  $|e| = 1$ , the pure second derivative  $u_{ee}$  belongs to the subsolution set  $\underline{S}(\lambda/n, \Lambda)$  (see [2] for the definition). Therefore, by the aid of the weak Harnack inequality and the Evans-Krylov estimate, we can complete this step (see [5, Lemma 2.1] for the detailed proof).

*Step 2:* We will find some quadratic polynomial  $Q(x) = \frac{1}{2} x^T \tilde{A} x + \tilde{b} \cdot x + \tilde{c}$  for some  $\tilde{A} \in \mathcal{S}^{n \times n}$  with  $F(\tilde{A}) = 0$ ,  $\tilde{b} \in \mathbb{R}^n$  and  $\tilde{c} \in \mathbb{R}$  such that  $Q \leq u$  in  $\mathbb{R}^n \setminus \overline{B_1}$ .

For  $k > 1$ , let  $u_k$  be a viscosity solution of

$$\begin{cases} F(D^2 u_k) = 0 & \text{in } B_k, \\ u_k = u & \text{on } \partial B_k. \end{cases}$$

Clearly, there exists infinite  $k > 1$  such that we have either

$$\max_{x \in \partial B_1} (u_k(x) - u(x)) \geq 0$$

or

$$\min_{x \in \partial B_1} (u_k(x) - u(x)) \leq 0.$$

Without loss of generality, we assume there exists infinite  $k > 1$  such that  $M_k := \max_{x \in \partial B_1} (u_k(x) - u(x)) = u_k(x_k) - u(x_k) > 0$  for  $x_k \in \partial B_1$  and set

$$\bar{u}_k(x) = u_k(x) - M_k$$

for all  $x \in B_k$ . Clearly, observing

$$\bar{u}_k \leq u \quad \text{on} \quad \partial B_1 \cup \partial B_k,$$

we apply the comparison principle to obtain

$$\bar{u}_k \leq u \quad \text{in} \quad B_k \setminus \overline{B_1}. \quad (2.1)$$

For any  $r > 1$  and  $k \geq 2r$ , applying the Evans-Krylov estimate and the Alexandroff-Bakelman-Pucci estimate to  $F(D^2 \bar{u}_k) = 0$  in  $B_k$ , we have

$$\|D^2 \bar{u}_k\|_{L^\infty(B_r)} \leq \|D^2 u_k\|_{L^\infty(B_{k/2})} \leq \frac{C}{k^2} \|u_k\|_{L^\infty(B_k)} \leq \frac{C}{k^2} \|u_k\|_{L^\infty(\partial B_k)},$$

where  $C > 0$  depends only on  $n$ ,  $\lambda$  and  $\Lambda$ . Combining this with the quadratic growth condition (1.2), we know that  $D^2 \bar{u}_k$  is bounded in  $B_r$  for  $k \geq 2r$ . Now we consider the function

$$w_k(x) = \bar{u}_k(x) - D\bar{u}_k(x_k) \cdot (x - x_k)$$

and then  $w_k(x_k) = |Dw_k(x_k)| = 0$ . Therefore, it is easily seen that

$$\|w_k\|_{L^\infty(B_r)} \leq Cr^2$$

for some constant  $C > 0$  independent of  $k$ . Furthermore, from the Hölder estimate applied to  $F(D^2 w_k) = 0$  in  $B_k$ , it follows that there exists a subsequence of  $\{w_k\}_{k \geq 2}$  that converges uniformly to a function  $w \in C(\mathbb{R}^n)$  in compact sets of  $\mathbb{R}^n$ . Moreover, we have

$$F(D^2 w) = 0 \quad \text{in} \quad \mathbb{R}^n$$

in the viscosity sense with

$$|w(x)| \leq C(1 + |x|^2), \quad x \in \mathbb{R}^n$$

for some constant  $C > 0$ . Therefore, by the Evans-Krylov estimate, we see that  $w$  is a quadratic polynomial  $P$ .

Recalling (2.1), we have

$$w_k(x) \leq u(x) - D\bar{u}_k(x_k) \cdot (x - x_k), \quad x \in B_k \setminus \overline{B_1}. \quad (2.2)$$

By taking some point  $x_0 \in \partial B_3(x_k)$  such that  $x_0 - x_k$  along the direction of  $D\bar{u}_k(x_k)$ , we obtain

$$3|D\bar{u}_k(x_k)| \leq u(x_0) - w_k(x_0) \leq C$$

for some constant  $C > 0$  independent of  $k$ . Consequently, there exist some  $b_\infty \in \mathbb{R}^n$  and  $x_\infty \in \partial B_1$  such that, up to a subsequence,

$$D\bar{u}_k(x_k) \rightarrow b_\infty \quad \text{and} \quad x_k \rightarrow x_\infty.$$

Finally, letting  $k \rightarrow \infty$  in (2.2), we have

$$P = w \leq u(x) - b_\infty \cdot (x - x_\infty), \quad x \in B_k \setminus \overline{B_1}.$$

Hence  $Q = P + b_\infty \cdot (x - x_\infty)$  is the desired quadratic polynomial.

*Step 3:* We claim that  $A = \tilde{A}$  and

$$|D^2u(x) - A| = O(|x|^{-n+\delta}) \quad \text{as } |x| \rightarrow \infty \quad (2.3)$$

for all  $\delta > 0$ .

Clearly,  $u - Q$  satisfies the linear elliptic equation

$$a_{ij}(x)D_{ij}(u - Q)(x) = 0, \quad x \in \mathbb{R}^n \setminus \overline{B_1}, \quad (2.4)$$

where  $a_{ij}(x) = \int_0^1 F_{ij}(tD^2u(x) + (1-t)D^2Q(x)) dt$ . Since  $F \in C^1$ , we combine this with Step 1 to conclude that

$$a_{ij}(x) \rightarrow a_{ij}^\infty \quad \text{as } |x| \rightarrow \infty,$$

where  $a_{ij}^\infty = \int_0^1 F_{ij}(tA + (1-t)\tilde{A}) dt$ .

*Case  $n \geq 3$ .* By the asymptotic theorem established by Li, Li and Yuan [5, Theorem 2.2] for positive solutions of non-divergence linear elliptic equations, there exists a constant  $c$  such that

$$u(x) = Q(x) + c + o(|x|^{2-n+\delta}) \quad \text{as } |x| \rightarrow \infty \quad (2.5)$$

for all  $\delta > 0$ .

For any  $x \in \mathbb{R}^n \setminus \overline{B_1}$  with  $|x| > 2$ , let

$$v(y) = \left(\frac{2}{|x|}\right)^2 (u - Q - c) \left(x + \frac{|x|}{2}y\right), \quad y \in B_1.$$

Then we have

$$F(D^2v(y) + \tilde{A}) = 0, \quad y \in B_1.$$

Applying the Evans-Krylov estimate, we obtain

$$|D^2u(x) - \tilde{A}| = |D^2v(0)| \leq C|x|^{-n+\delta}, \quad x \in \mathbb{R}^n \setminus \overline{B_1}$$

for some constant  $C > 0$ , which easily yields  $A = \tilde{A}$  and (2.3).

*Case  $n = 2$ .* By Lemma 2.1, we have

$$u(x) = Q(x) + O(|x|^\delta) \quad \text{as } |x| \rightarrow \infty \quad (2.6)$$

for all  $\delta > 0$ . Following the same arguments to the case  $n \geq 3$ , we also obtain  $A = \tilde{A}$  and (2.3).

*Step 4:* The aim of this step is to refine the error to  $O(|x|^{2-n})$  ( $n \geq 3$ ) and  $O(|x|^{-1})$  ( $n = 2$ ) for the asymptotic results (2.5) and (2.6).

Without loss of generality, we can assume  $a_{ij}^\infty = \delta_{ij}$ . We take  $0 < \delta < \frac{1}{2}$  small enough. Since  $F \in C^{1, \frac{2}{n}}$ , we have

$$|a_{ij}(x) - \delta_{ij}| \leq C|x|^{\frac{2}{n}(-n+\delta)}, \quad x \in \mathbb{R}^n \setminus \overline{B_1}$$

for some constant  $C > 0$ .

*Case  $n \geq 3$ .* Once again using the asymptotic theorem ([5, Theorem 2.2]), we obtain

$$u(x) = Q(x) + c + O(|x|^{2-n}) \quad \text{as } |x| \rightarrow \infty.$$

*Case  $n = 2$ .* Clearly, we have

$$\Delta(u - Q)(x) = (\delta_{ij} - a_{ij}(x))D_{ij}(u - Q)(x) = O(|x|^{-2+\delta}|x|^{-2+\delta}) \quad \text{as } |x| \rightarrow \infty.$$

By [6, Lemma 3.4], there exists some function

$$w(x) = O(|x|^{-2+2\delta+\epsilon})$$

for any  $0 < \epsilon < 1$  such that

$$\Delta(u - Q - w)(x) = 0$$

and

$$(u - Q - w)(x) = O(|x|^\delta) \quad \text{as } |x| \rightarrow \infty.$$

Then applying [6, Lemma 3.5] leads to

$$(u - Q - w)(x) = d \log |x| + c + O(|x|^{-1})$$

for some  $c, d \in \mathbb{R}$ . Hence

$$u(x) = Q(x) + d \log |x| + c + O(|x|^{-1})$$

*Step 5:* We will determine the term  $\frac{x}{|x|^n}$ .

*Case  $n \geq 3$ .* We consider the function

$$E(x) = u(x) - Q(x) - c = O(|x|^{2-n}) \quad \text{as } |x| \rightarrow \infty.$$

By the Evans-Krylov estimate, we have

$$D^2 E(x) = O(|x|^{-n}) \quad \text{as } |x| \rightarrow \infty.$$

Since  $E$  satisfies the linear equation (2.4) and  $F \in C^{1, \frac{2}{n}}$ , we obtain

$$\Delta E(x) = (\delta_{ij} - a_{ij}(x)) D_{ij} E(x) := g(x) = O(|x|^{-2} |x|^{-n}), \quad x \in \mathbb{R}^n \setminus \overline{B_R}$$

for some constant  $R > 1$ . Hence the Kelvin transformation of  $E$ , defined by

$$K[E](x) = |x|^{2-n} E\left(\frac{x}{|x|^2}\right) \quad \text{for } x \in B_{\frac{1}{R}} \setminus \{0\},$$

satisfies

$$\Delta K[E](x) = |x|^{-n-2} g\left(\frac{x}{|x|^2}\right) := \tilde{g}(x) \quad \text{in } B_{\frac{1}{R}} \setminus \{0\},$$

where  $\tilde{g}$  satisfies

$$|\tilde{g}(x)| \leq C \quad \text{in } B_{\frac{1}{R}} \setminus \{0\}$$

for some constant  $C > 0$ .

Clearly,  $\tilde{g} \in L^\infty(B_{1/R})$ . By the  $L^p$  regularity, we have  $K[E] \in W^{2,p}(B_{1/R})$  for any  $1 < p < \infty$ . Hence  $K[E] \in C^{1,\alpha}(B_{1/R})$  for any  $0 < \alpha \leq 1 - \frac{n}{p}$  with  $p \in (n, \infty)$ , which follows from the Sobolev imbedding theorem. More precisely, there exist some constant  $\tilde{C} > 0$  and linear function  $\tilde{b} \cdot x + \tilde{c}$  such that

$$\left| K[E](x) - \tilde{b} \cdot x - \tilde{c} \right| \leq \tilde{C} |x|^{1+\alpha} \quad \text{in } B_{\frac{1}{R}}.$$

Therefore, going back to  $E$ , we have

$$\left| E(x) - \tilde{c} |x|^{2-n} - \frac{\tilde{b} \cdot x}{|x|^n} \right| \leq \frac{\tilde{C}}{|x|^{n-1+\alpha}} \quad \text{in } \mathbb{R}^n \setminus \overline{B_R}.$$

*Case  $n = 2$ .* We let

$$E(x) = u(x) - Q(x) - d \log |x| - c = O(|x|^{-1}) \quad \text{as } |x| \rightarrow \infty.$$

The rest of the proof runs as before (or see [6, pp. 10-11] for the detailed proof).  $\square$

---

## THE LIOUVILLE THEOREM

---

*Remark 2.2.* For  $n \geq 3$ , if  $F \in C^1$ , we can obtain the error term  $o(|x|^{2-n+\delta})$  for any  $\delta > 0$ ; if  $F \in C^{1,\gamma}$  for some  $0 < \gamma < 1$ , the error term is  $O(|x|^{2-n})$ . When  $F$  is Lipschitz continuous, Lian and Zhang [7] obtained an asymptotic result with the error  $O(|x|^{1-(n-1)\frac{\lambda}{\lambda}})$  that is the fundamental solution of the Pucci's operator (see [1, 3] for more details).

### ACKNOWLEDGEMENT

This work is supported by NSFC 12071365.

### REFERENCES

- [1] S. N. Armstrong, B. Sirakov, and C. K. Smart, Fundamental solutions of homogeneous fully nonlinear elliptic equations, *Comm. Pure Appl. Math.*, 64 (2011), no. 6, 737-777.
- [2] Caffarelli, Luis A.; Cabré, Xavier Fully nonlinear elliptic equations. American Mathematical Society Colloquium Publications, 43. American Mathematical Society, Providence, RI, 1995. vi+104 pp. ISBN: 0-8218-0437-5
- [3] D. A. Labutin, Isolated singularities for fully nonlinear elliptic equations, *J. Differential Equations*, 177 (2001), no. 1, 49-76.
- [4] G.H. Hong, A Remark on Monge-Ampère equation over exterior domains, 2020, arXiv:2007.12479.
- [5] D. Li, Z. Li, and Y. Yuan, A Bernstein problem for special Lagrangian equations in exterior domains, *Adv. Math.* 361 (106927) (2020) 29 pp.
- [6] D. Li and R. Liu, Quasiconformal mappings and a Bernstein type theorem over exterior domains in  $\mathbb{R}^2$ , *Calc. Var. Partial Differential Equations* 63 (2024), no. 8, Paper No. 209, 13 pp.
- [7] Y. Y. Lian and K. Zhang, Asymptotic behavior for fully nonlinear elliptic equations in exterior domains, arXiv:2401.05829.

SCHOOL OF MATHEMATICS AND STATISTICS, XI'AN JIAOTONG UNIVERSITY, XI'AN, P.R.CHINA 710049.

SCHOOL OF MATHEMATICS AND STATISTICS, XI'AN JIAOTONG UNIVERSITY, XI'AN, P.R.CHINA 710049.

*Email address:* lidsh@mail.xjtu.edu.cn

*Email address:* lianglichun126@stu.xjtu.edu.cn