On the minimal components of substitution subshifts

Raphael Henry Aix Marseille Univ, CNRS, I2M, Marseille, France raphael.henry@univ-amu.fr

February 3, 2025

Abstract

In this paper we study substitutions on $A^{\mathbb{Z}}$ where *A* is a finite alphabet. We precisely characterize the minimal components of substitution subshifts, give an optimal bound for their number and describe their dynamics. The explicitness of these results provides a method to algorithmically compute and count the minimal components of a given substitution subshift.

1 Introduction

In symbolic dynamics, substitutions on a finite alphabet A are a rich way to generate infinite words, for a general background see [1] or the upcoming book [2]. In particular, substitution subshifts on $A^{\mathbb{Z}}$ have been extensively studied, for a general overview see [14] and for various computability results see [3]. A major interest regarding subshifts is to understand when they are *minimal*, meaning that the orbit of every point is dense. For example it is well-known that a primitive substitution generates a minimal subshift (see for example [14]), also Damanik and Lenz [6] proved that a substitution subshift is minimal if and only if it is linearly repetitive. In a recent interest towards non-minimal substitution subshifts, Maloney and Rust [11] identified that tameness, a property of the substitution, is essential for minimality. This led to a characterization of minimal substitution subshifts by Shimomura [15]: the class of minimal substitution subshifts is equal to the class of subshifts generated by tame and l-primitive (i.e., primitive when removing the bounded letters) substitutions. Naturally, the next step is to focus on non-minimal substitution subshifts. In [3], Béal, Perrin and Restivo show that any substitution subshift is quasi-minimal, meaning that it has a finite number of sub-subshifts. In particular it has a finite number of minimal components, which raises the question of characterizing and counting them.

For growing substitutions, a first type of minimal components is known: Durand [8] introduces *main sub-subtitutions*, that are primitive restrictions of a specific power of the original substitution. Main sub-substitutions always exist when the original substitution is growing, and they are primitive so each of them generates a minimal component. An analog of these minimal components is introduced by Cortez and Solomyak [5] in the framework of tiling substitutions on \mathbb{R}^d , and they show that they are precisely the minimal components of the tiling dynamical system. In particular, it is remarkable that every minimal component is itself a substitution system.

For non-growing substitutions, another type of minimal component comes from the bounded letters. This idea originates from the closely related framework of D0L-systems, where Ehrenfeucht and Rozenberg [9] begin the investigation of infinite repetitions in their language and Klouda and Starosta [10] provide effective results. In particular, they show that if a substitution is *wild* (i.e., not tame), its language contains infinite repetitions over bounded words. Notably, this idea was already used by Pansiot [12] to characterize purely morphic words with factor complexity $\Theta(n^2)$. This transposes to the framework of substitution subshifts where Maloney and Rust [11] and Shimomura [15] show that a wild substitution produces periodic words over bounded letters, and in particular their orbit is a minimal component.

The main contribution of this paper is the characterization of the minimal components of any substitution subshift: they are either the subshifts generated by a generalization of main sub-substitutions, defined with a new construction to identify the tame and l-primitive sub-substitutions instead of the primitive ones; or single periodic orbits produced by wild substitutions, defined by introducing specific computable words. One key step of this result is to show that the substitution induces a map on the subshifts of the substitution subshift, and that it is a permutation. In a second time, we show that this map acts specifically on the minimal components and that its action is described by the directed graphs introduced in our proof. We also aim to make our result as effective as possible, which allows us in the last sections to count and bound the number of minimal components for all substitutions on two letters. We provide a Python implementation of the computation of minimal components at https://codeberg.org/RaphaelHENRY/MinimalComponents.git.

2 Preliminaries

2.1 Notations

Words

Given a finite alphabet *A*, *A*^{*} denotes the set of all words on *A*, in particular ε is the empty word and we define $A^+ := A^* \setminus \{\varepsilon\}$. For $u = u_0 u_1 \dots u_{n-1} \in A^*$, we write |u| := n the *length* of *u*. To simplify some expressions, we will use the following

notation: if $(u_i)_{0 \le i \le n-1} \in (A^*)^n$, we write the concatenation from left to right as

$$\bigsqcup_{i=0}^{n-1} u_i := u_0 u_1 \dots u_{n-2} u_{n-1} \qquad \text{with the convention that } \bigsqcup_{i=0}^{-1} u_i := \varepsilon.$$

If words $u, u', v, w \in A^*$ are such that w = uvu', we say that v is a *factor* of w and we write $v \sqsubset w$. If $u = \varepsilon$, v is a *prefix* of w, and if $u' = \varepsilon$, v is a *suffix* of w. If $u, v \in A^*$ and there exists $w, w' \in A^*$ such that u = ww' and v = w'w, we say that u is a *cyclic shift* of v. A word $u \in A^+$ is primitive if $u = v^k$ implies that k = 1. The shortest word v such that $u = v^k$ for some k is the *primitive root* of u.

Let $A^{\mathbb{Z}}$ be the set of all two-sided sequences of letters of A. For $x \in A^{\mathbb{Z}}$ and $i - 1 \leq j \in \mathbb{Z}$, we write the word $x_{[i,j]} := x_i...x_j$, with the convention that $x_{[i,i-1]} = \varepsilon$. The word $x_{[i,j]}$ is a *factor* of x and we write $x_{[i,j]} \sqsubset x$. If $u \in A^+$, we write the infinite word to the right $u^{\omega} := uuu...$ and the infinite word to the left ${}^{\omega}u := ...uuu$. By writing a dot right to the left of the letter of index 0, we define the bi-infinite words ${}^{\omega}u^{\omega} := {}^{\omega}u.u^{\omega}$ and ${}^{\omega}uvw^{\omega} := {}^{\omega}u.vw^{\omega}$.

Substitutions

A map $\varphi : A^* \to A^*$ is a *morphism* if for all $u, v \in A^*$, $\varphi(uv) = \varphi(u)\varphi(v)$. In this case, we write $\varphi : A \to A^*$ and the language of φ is the set

$$\mathscr{L}(\varphi) := \{ u \in A^* \mid \exists a \in A, \exists n \ge 0, u \sqsubset \varphi^n(a) \}.$$

If in addition φ is *non-erasing*, i.e., for all $a \in A$, $\varphi(a) \neq \varepsilon$, then φ is a *substitution* and we write $\varphi : A \to A^+$. We also define $|\varphi| := \max_{a \in A} |\varphi(a)|$.

If $A = \{a_1, ..., a_n\}$ and $\varphi : A \to A^+$ is a substitution, the substitution matrix M is the matrix of size $n \times n$ such that, for all $i, j \in \{1, ..., n\}, M_{i,j} = |\varphi(a_j)|_{a_i}$ the number of a_i in $\varphi(a_j)$. We say that φ is *primitive* if there exists $m \ge 1$ such that, for every $a, b \in A, a \sqsubset \varphi^m(b)$. If $D \subset A$ is a subalphabet such that $\varphi(D) \subset D^+$, we have the restriction $\varphi|_D : D \to D^+$.

Growth rate of letters

Let $\varphi : A \to A^+$ be a substitution. If $a \in A$ is such that the sequence $(|\varphi^n(a)|)_{n\geq 0}$ is bounded, we say that *a* is *bounded* and we define *B* the set of bounded letters. Otherwise, we say that *a* is *growing* and we define *C* the set of growing letters. If $B = \emptyset$, we say that φ is *growing*. If $u \in A^*$, we define $alph_C(u) := \{c \in C \mid c \sqsubset u\}$. If $X \subset A^{\mathbb{Z}}$, we define the alphabet of growing letters in X

$$alph_C(X) := \{c \in C \mid \exists x \in X, c \sqsubset x\}.$$

Subshifts

We define the right shift $T : A^{\mathbb{Z}} \to A^{\mathbb{Z}}$ where $T(x)_i = x_{i-1}$. A non-empty set $X \subset A^{\mathbb{Z}}$ is a *subshift* if it is closed for the discrete topology on $A^{\mathbb{Z}}$ and satisfies

T(X) = X. If $x \in A^{\mathbb{Z}}$, the smallest subshift containing *x* is the set

$$X(x) := \overline{\{T^k(x) \mid k \in \mathbb{Z}\}}$$

In particular, if $u \in A^+$, $X({}^{\omega}u^{\omega})$ is a single periodic orbit. The language of the subshift *X* is the set

$$\mathscr{L}(X) := \{ u \in A^* \mid \exists x \in X, u \sqsubset x \}.$$

A word *u* is said to occur with *bounded gaps* in X if there exists $L \ge 1$ such that, for all $v \in \mathcal{L}(X)$, if $|v| \ge L$ then $u \sqsubset v$.

Given a substitution $\varphi : A \to A^+$ where $C \neq \emptyset$, the associated *substitution subshift* is the set

$$X_{\varphi} := \left\{ x \in A^{\mathbb{Z}} \mid \forall u \sqsubset x, u \in \mathscr{L}(\varphi) \right\}.$$

In particular, $\mathscr{L}(X_{\varphi}) \subset \mathscr{L}(\varphi)$. If $A \subset \mathscr{L}(X_{\varphi})$, φ is admissible.

Example 2.1. The substitution $\varphi : 0 \mapsto 12, 1 \mapsto 22, 2 \mapsto 11$ is not admissible because $X_{\varphi} = X(^{\omega}12^{\omega}) \cup X(^{\omega}21^{\omega})$ and $0 \notin \mathscr{L}(X_{\varphi})$.

Minimality

A subshift X is *minimal* if it does not contain any subshift other than itself.

Remark 2.2. If X is a subshift, then the three following properties are equivalent:

- (i) X is minimal.
- (ii) For every $x \in X$, X(x) = X.
- (iii) Every word of $\mathscr{L}(X)$ occurs with bounded gaps in *X*.
- A proof can be found in [13] for example.

If *X* is a subshift, a minimal subshift $Y \subset X$ is called a *minimal component* of *X*. If *X* has a unique minimal component, we say it is *essentially minimal*, which is not equivalent to being minimal as illustrated with Example 2.17. More generally, if *X* contains a finite number of subshifts, we say it is *quasi-minimal*. In particular, every substitution subshift is quasi-minimal.

Proposition 2.3 ([3, Proposition 10.8]). Let $\varphi : A \to A^+$ be a substitution. Then X_{φ} is quasi-minimal.

D0L-systems

A D0L-system is a triplet $G = (A, \varphi, w)$ where A is a finite alphabet, $\varphi : A \to A^*$ is a morphism and w is a word in A^+ . The associated language is the set

$$\mathscr{L}(G) := \{ u \in A^* \mid \exists n \ge 0, u \sqsubset \varphi^n(w) \}.$$

2.2 Minimality and beyond

2.2.1 Characterization of minimality

The minimality of substitution subshifts relies on two properties of the substitution.

The first property is about the production of bounded letters. If $\varphi : A \to A^+$ is a substitution, we say that a letter $c \in C$ is *left-isolated* (resp. *right-isolated*) if there exist $n \ge 1$, $u \in B^+$ and $v \in A^*$ such that $\varphi^n(c) = ucv$ (resp. $\varphi^n(c) = vcu$). We define C_{liso} (resp. C_{riso}) the set of left-isolated (resp. right-isolated) letters, and C_{niso} the set of growing letters that are not left or right-isolated.

Definition 2.4. We say that a substitution φ is *tame* if $C_{liso} = C_{riso} = \emptyset$, otherwise we say it is *wild*.

Example 2.5. Consider the substitution $\varphi : 0 \mapsto 001, 1 \mapsto 1$. We have $C = \{0\}$, $B = \{1\}$ and $0 \notin C_{liso}$ but $0 \in C_{riso}$, therefore φ is wild.

Note that growing substitutions are always tame. The second property is a softer version of primitivity.

Definition 2.6. We say that a substitution φ is *l*-primitive if there exists $n \ge 1$ such that, for all $a, b \in C$, $a \sqsubset \varphi^n(b)$.

Note that for growing substitutions, l-primitivity is the same as primitivity. By combining the two properties, Shimomura gives a characterization of minimal substitution subshifts:

Theorem 2.7 ([15, Theorem A]). Let $\varphi : A \to A^+$ be a substitution.

(*i*) Suppose that φ is tame and *l*-primitive. Then, X_{φ} is minimal.

(ii) Suppose that X_{φ} is minimal and is not a single periodic orbit. Then, φ is tame, and there exists a unique subalphabet $D \subset A$ and a restriction $\varphi|_D : D \to D^+$ such that $\varphi|_D$ is tame, *l*-primitive, and $X_{\varphi} = X_{\varphi|_D}$.

As single periodic orbits are minimal and can be expressed as a substitution subshift, this result gives an equivalent class for minimal substitution subshifts.

Corollary 2.8 ([15, Corollary B]). Let \mathscr{M} be the class of all minimal substitution subshifts. Let \mathscr{M}' be the class of all X_{φ} such that φ is a tame and *l*-primitive substitution. Then, it follows that $\mathscr{M} = \mathscr{M}'$.

2.2.2 Minimal components for growing substitutions

When $\varphi : A \to A^+$ is a growing substitution, Durand [8] defines *main subsubstitutions*. As one goal of this paper is to generalize them to any substitution, we will describe them only briefly.

By raising the matrix of φ to a sufficient power p, Durand introduces *principal primitive components* of A, that are the disjoint alphabets $A_i \subset A$ such that $\varphi^p(A_i) \subset A_i^+$ and the restriction $\varphi^p|_{A_i}$ is primitive. The $\varphi^p|_{A_i}$ are called *main subsubstitutions*, they always exist and the substitution subshifts $X_{\varphi^p|_{A_i}}$ are minimal components of X_{φ} .

Example 2.9. Consider the growing substitution $\varphi : 0 \mapsto 11, 1 \mapsto 00$. Looking at φ^2 , its main sub-substitutions are $\varphi^2|_{\{0\}}$ and $\varphi^2|_{\{1\}}$, so $X_{\varphi^2|_{\{0\}}} = \{{}^{\omega}0{}^{\omega}\}$ and $X_{\varphi^2|_{\{1\}}} = \{{}^{\omega}1{}^{\omega}\}$ are minimal components of X_{φ} . Note that, in this example, they are its only minimal components.

Let us draw a parallel with a result from Cortez and Solomyak [5] in the framework of admissible substitution tiling spaces on \mathbb{R}^d .

Lemma 2.10 ([5, Lemma 2.9]). Let ω be an admissible tile substitution. For all $k \ge 2$, $X_{\omega^k} = X_{\omega}$.

This allows to replace ω by a sufficient power ω^p that satisfies property (2.2), i.e., its matrix provides alphabets A_i and the primitive substitutions $\varphi^p|_{A_i}$ like Durand. What is remarkable in this framework is that this provides all the minimal components:

Proposition 2.11 ([5, Lemma 2.10 (i)]). Let ω be an admissible tile substitution satisfying (2.2). Then the minimal components of X_{ω} are the $X_{\omega|_{A}}$.

As the A_i are pairwise disjoint, it follows an upper bound on the number of minimal components.

Corollary 2.12 ([5, Corollary 2.11]). Let ω be an admissible tile substitution on a set of tiles A. Then X_{ω} has at most |A| minimal components.

2.2.3 Minimal components for non-growing substitutions

When a substitution is not growing, the production of bounded letters is closely related to tameness:

Proposition 2.13 ([11, Theorem 2.9], [15, Proposition 3.17]). Let $\varphi : A \to A^+$ be a substitution. Then the words in $\mathscr{L}(X_{\varphi}) \cap B^*$ have bounded length if and only φ is tame.

Remark 2.14. A similar result was proven earlier for the language of D0L-systems in [9, proof of Lemma 2.1] with a different vocabulary: tameness is called the *edge condition*, and the fact that $\mathscr{L}(G) \cap B^*$ is infinite is called *pushyness*.

In particular, if a substitution φ is wild, then there exists words in $X_{\varphi} \cap B^{\mathbb{Z}}$ and they are shown to have a strong periodic structure. Firstly, Proposition 2.13 relies in part on the following result:

Lemma 2.15 ([11, Lemma 2.8],[15, Lemma 3.8]). Let $\varphi : A \to A^+$ be a substitution. If φ is wild, then X_{φ} contains a periodic word in $B^{\mathbb{Z}}$.

Proof. Let *c* ∈ *C*_{*liso*} (resp. *c* ∈ *C*_{*riso*}), so that there exist *n* ≥ 1, *u* ∈ *B*⁺ and *v* ∈ *A*^{*} such that $\varphi^n(c) = ucv$ (resp. $\varphi^n(c) = vcu$). Then, for all $l \ge 1$, there exists $v_l \in A^*$ such that $\varphi^{nl}(c) = \varphi^{l-1}(u)...\varphi(u)ucv_l$ (resp. $\varphi^{nl}(c) = v_lcu\varphi(u)...\varphi^{l-1}(u)$). As $u \in B^+$, the sequence $(\varphi^l(u))_{l\ge 0}$ is eventually periodic so, as *l* grows, the $\varphi^{nl}(c)$ produce a periodic word in $X_{\varphi} \cap B^{\mathbb{Z}}$.

Remark 2.16. The proof of Lemma 2.15 is in fact more precise than its statement: it shows that every letter of C_{liso} (resp. of C_{riso}) generates a periodic word $x \in X_{\varphi} \cap B^{\mathbb{Z}}$, therefore it generates the minimal component $X(x) \subset X_{\varphi} \cap B^{\mathbb{Z}}$.

Example 2.17. Consider the substitution $\varphi : 0 \mapsto 101, 1 \mapsto 1$, for which $C = \{0\}$ and $B = \{1\}$. We have $0 \in C_{liso} \cap C_{riso}$ and ${}^{\omega}1^{\omega} \in X_{\varphi} \cap B^{\mathbb{Z}}$ so $\{{}^{\omega}1^{\omega}\} \subset X_{\varphi} \cap B^{\mathbb{Z}}$ is a minimal component. Note that $X_{\varphi} = X({}^{\omega}101^{\omega})$ so $\{{}^{\omega}1^{\omega}\}$ is its unique minimal component, thus X_{φ} is essentially minimal but not minimal.

More generally, every word in $X_{\varphi} \cap B^{\mathbb{Z}}$ has a periodic structure.

Proposition 2.18 ([3, Proposition 4.3]). Let $\varphi : A \to A^+$ be a substitution. If $x \in X_{\varphi} \cap B^{\mathbb{Z}}$, then there exists $u, v, w \in B^+$ and $k \in \mathbb{Z}$ such that $x = T^k(^{\omega}uwv^{\omega})$, where the lengths of u, v, w are bounded by a computable integer depending only on φ .

This provides a description of the minimal components of X_{φ} in $B^{\mathbb{Z}}$.

Corollary 2.19. Let $\varphi : A \to A^+$ be a substitution and let $X \subset B^{\mathbb{Z}}$ be a minimal component of X_{φ} , then there exists a word $u \in B^+$ such that $X = X(^{\omega}u^{\omega})$. In other words, every minimal component of $X_{\varphi} \cap B^{\mathbb{Z}}$ is a single periodic orbit.

Proof. Let $x \in X$. Then, by Proposition 2.18, there exist $u, v, w \in B^+$ and $k \in \mathbb{Z}$ such that $x = T^k({}^{\omega}uwv{}^{\omega})$. By closeness of subshifts, ${}^{\omega}u^{\omega} \in X$ so, as X is minimal, $X = X({}^{\omega}u^{\omega})$.

2.3 Results

In this paper, we give a complete and effective characterization of the minimal components of X_{φ} where φ is any substitution. To achieve that, we distinguish two types of minimal components.

Definition 2.20. Let $\varphi : A \to A^+$ be a substitution and let *X* be a minimal component of X_{φ} . If $X \subset B^{\mathbb{Z}}$, we say that *X* is *wild*, otherwise we say that *X* is *tame*.

This definition is inspired by the two types of minimal components previously described. On the one hand, main sub-substitutions provide tame minimal components in the growing case, our goal is to generalize them to the general case and to show that they are precisely the minimal components, as done in [5] for admissible tile substitutions. On the other hand, Corollary 2.19 states that all the wild minimal components are single periodic orbits, our goal is to give an explicit characterization that is suited for describing the dynamics, computing, and counting the minimal components. This is the main result of this paper:

Theorem 2.21. Let $\varphi : A \to A^+$ be a substitution.

(*i*) The tame minimal components of X_{φ} are the $X_{\varphi^k|_{D \cup B}}$ where $D \subset C_{niso}$ is a minimal alphabet of period k.

(ii) The wild minimal components of X_{φ} are the $X(^{\omega}LP(c)^{\omega})$ where $c \in C_{liso}$ and LP(c) is a computable word in B^+ , and the $X(^{\omega}RP(c)^{\omega})$ where $c \in C_{riso}$ and RP(c) is a computable word in B^+ .

Remark 2.22. In the growing case, the definition of main sub-substitutions and Proposition 2.11 rely on raising the substitution to a sufficient power. This has the benefit of simplifying the cyclic behaviour of subalphabets, but we do not do this here for several reasons:

(i) Lemma 2.10 also holds for admissible substitutions on $A^{\mathbb{Z}}$ but not for non-admissible ones: in Example 2.1, we have $X_{\varphi} = X(^{\omega}12^{\omega}) \cup X(^{\omega}21^{\omega})$ and $X_{\varphi^2} = X(^{\omega}21^{\omega})$,

(ii) It would hide all the dynamical aspects we focus on in Sections 3.1.3 and 5,

(iii) It would make computation significantly longer when we want our result to be as efficient as possible.

In Section 3 we prove Theorem 2.21 (i) by generalizing main sub-substitutions to the non-growing case. We first introduce minimal alphabets (Definition 3.5), not with matrices but with an oriented graph, and we show that the restrictions to these alphabets provide tame minimal components. To prove that all tame components have this form, we show that the substitution φ induces a map $\tilde{\varphi}$ on the subshifts of X_{φ} and that it is a permutation.

In Section 4 we prove Theorem 2.21 (ii) by constructing the computable words LP(c) and RP(c) (Equations (1) and (2)) that depend only on φ and the letter *c*. By construction, ${}^{\omega}LP(c){}^{\omega}$ and ${}^{\omega}RP(c){}^{\omega}$ are the periodic words exhibited in Remark 2.16, and we take inspiration from a result in D0L-systems to show that they are precisely the periodic words in Corollary 2.19.

In Section 5 we show that $\tilde{\varphi}$ acts on the tame (resp. wild) components of X_{φ} as a permutation, and that its action is described by the directed graphs we built to prove our theorem.

Example 2.23. The different constructions and results throughout Sections 3 to 5 will be illustrated with the substitution

 $\varphi: 0 \mapsto 141, 1 \mapsto 00, 2 \mapsto 242, 3 \mapsto 5435, 4 \mapsto 5, 5 \mapsto 6, 6 \mapsto 5, 5 \mapsto 6, 5 \mapsto 5, 5 \mapsto 6, 5 \mapsto 5, 5 \mapsto 6, 5 \mapsto 6, 5 \mapsto 5, 5 \mapsto 6, 5 \mapsto 6, 5 \mapsto 5, 5 \mapsto 6, 5 \mapsto 5, 5 \mapsto 6, 5 \mapsto 6, 5 \mapsto 5, 5 \mapsto 6, 5 \mapsto 5, 5 \mapsto 6, 5 \mapsto 5, 5 \mapsto 6, 5 \mapsto 6, 5 \mapsto 5, 5 \mapsto 6, 5 \mapsto 5, 5 \mapsto 6, 5 \mapsto 5, 5 \mapsto 5$

for which $C = \{0, 1, 2, 3\}, B = \{4, 5, 6\}$ and $C_{liso} = C_{riso} = \{3\}$.

Remark 2.24. In practice, we display minimal components in their reduced form: for every tame component $X = X_{\varphi^k|_{D\cup B}}$, there is a unique alphabet *E* such that $D \subset E \subset D \cup B$, $\varphi^k|_E$ is defined and admissible and $X = X_{\varphi^k|_E}$; for every wild component $X = X({}^{\omega}u^{\omega})$, there is a primitive word *v* such that $X = X({}^{\omega}v^{\omega})$.

In the last two sections we emphasize the effectiveness of our characterization. In Section 6 we begin by showing a known computability result for which we could not find a proper proof:

Proposition 2.25. Let $\varphi : A \to A^+$ be a substitution. Then B and C are computable.

As an application of our theorem, we are able to compute and count the minimal components of a given substitution subshift. If $\varphi : A \to A^+$ is a substitution, the number of minimal components denoted by $MC(\varphi)$.

Corollary 2.26. Let $\varphi : A \to A^+$ be a substitution. Then $MC(\varphi)$ is computable.

In particular, we can decide if X_{φ} is essentially minimal. A Python implementation of the computation of *B* and *C*, of the minimal components of a given substitution subshift as well as their number can be found at https://codeberg.org/RaphaelHENRY/MinimalComponents.git. According to Remark 2.24, we output the tame components of the form $X_{\varphi^k|_E}$ as the couple (E, k) and the wild components as the primitive word v.

We also bound $MC(\varphi)$ by the size of the alphabet, as in Corollary 2.12.

Corollary 2.27. Let $\varphi : A \rightarrow A^+$ be a substitution. (i) If |B| = 0, then $MC(\varphi) \le |C| = |A|$. (ii) If |B| = 1, then $MC(\varphi) \le |C| = |A| - 1$. (iii) If $|B| \ge 2$, then $MC(\varphi) \le 2|C| \le 2|A| - 4$.

We provide examples to show that these upper bounds - in fact every number between 1 and the upper bound - can be reached.

Finally, in Section 7 we detail the computation of the minimal components for all substitutions on two letters, which provides numerous examples to illustrate the tools developed in this paper.

Remark 2.28. The way we characterize, compute and count the wild components can be directly used to characterize, compute and count the infinite repetitions over bounded letters in a D0L-system.

3 Tame minimal components

3.1 Dynamics of alphabets

3.1.1 Minimal alphabets

In this section we identify the subalphabets $D \subset A$ for which there exists $k \ge 1$ such that $\varphi^k(D) \subset D^+$ in order to study the restrictions $\varphi^k|_D : D \to D^+$. Our goal is to determine when such a restriction is l-primitive and tame (so that it generates a minimal substitution subshift) so we study the action of φ on the subalphabets of *C* rather than on the subalphabets of *A*, and we later add the bounded letters.

Definition 3.1. In order to represent how φ acts on the subalphabets of *C*, we define the directed graph *G* := (*V*, *E*) by

•
$$V := \mathscr{P}(C) \setminus \{\emptyset\},\$$

•
$$E := \left\{ \left(D, \bigcup_{a \in D} alph_{\mathbb{C}}(\varphi(a)) \right) \middle| D \in V \right\} \subset V^2.$$

If $(D, D') \in E$, we write $D \to D'$. If $k \ge 1$ and $D \xrightarrow{\longrightarrow} \dots \xrightarrow{\longrightarrow} D'$, we write $D \xrightarrow{\longrightarrow} D'$, k times

and in that case $D' = \bigcup_{a \in D} alph_C(\varphi^k(a)).$

Example 3.2. Following Example 2.23, the graph *G* has 15 vertices so let us display the orbit of the singletons only:



The following lemma shows that *G* behaves well with inclusion and union.

Lemma 3.3. Let $k \ge 1$ and $D_1, D_2, D_3, D_4 \in V$. (*i*) If $D_1 \subset D_2, D_1 \xrightarrow{k} D_3$ and $D_2 \xrightarrow{k} D_4$, then $D_3 \subset D_4$. (*ii*) If $D_1 \rightarrow D_3$ and $D_2 \rightarrow D_4$, then $D_1 \cup D_2 \rightarrow D_3 \cup D_4$.

Proof. The proof is left to the reader.

Remark 3.4. Lemma 3.3 (ii) implies that the graph *G* is entirely determined by the alphabets *D* such that $\{a\} \rightarrow D$. We call these alphabets the *generators* of *G*, they are particularly relevant when we do computations.

New let us identify the cyclic behaviors in *G*.

Definition 3.5. We say that $D \in V$ is a *k-periodic* alphabet if *k* is the smallest positive integer such that $D \xrightarrow{k} D$. We say that $D \in V$ is a *minimal* alphabet if there exists $k \ge 1$ such that D is *k*-periodic and has no proper periodic subalphabet.

Note that every periodic alphabet contains at least one minimal alphabet.

Lemma 3.6. Let *D* be a minimal alphabet of period *k* and let *E* such that $D \rightarrow E$. Then *E* is a minimal alphabet of period *k*.

Proof. First, *E* is a *k*-periodic alphabet. Let $E' \subset E$ be a *k'*-periodic subalphabet. Also let *D'* be the *k'*-periodic alphabet such that $E' \xrightarrow[k-1]{k-1} D'$. We have $E \xrightarrow[k-1]{k-1} D$ so, by Lemma 3.3 (i), $D' \subset D$. Then, as *D* is minimal, we have D' = D and k' = k. Finally, $E' \xrightarrow[k]{k} E'$ and $E' \xrightarrow[k-1]{k-1} D \to E$ so E' = E.

We now show an equivalent property to the minimality of alphabets, which will be easier to handle in the proofs.

Lemma 3.7. Let *D* be a *k*-periodic alphabet. Then *D* is minimal if and only if for all $a \in D$, there exists $l_a \ge 1$ such that $\{a\} \xrightarrow[k]{} D$.

Proof. Suppose that *D* is minimal. Let $a \in D$, and for all $l \ge 1$, let D_l be the subalphabet such that $\{a\} \xrightarrow{kl} D_l$. Then, as $\{a\} \subset D$ and $D \xrightarrow{kl} D$, by Lemma 3.3 (i), $D_l \subset D$. As *V* is finite, the sequence $(D_l)_{l\ge 1}$ is eventually periodic, which means that there exists $l_a \ge 1$ such that D_{l_a} is a periodic alphabet. Then, by minimality of D, $D_{l_a} = D$.

Suppose that for all $a \in D$, there exists $l_a \ge 1$ such that $\{a\} \xrightarrow{kl_a} D$. Let $D' \subset D$ be a minimal k'-periodic alphabet and let $a \in D'$. By supposition, there exists l_a such that $\{a\} \xrightarrow{kl_a} D$. Moreover, by the previous implication, there exists $l'_a \ge 1$ such that $\{a\} \xrightarrow{kl_a} D'$. Setting $l = max(l_a, l'_a)$, we get $\{a\} \xrightarrow{kk'l} D$ and $\{a\} \xrightarrow{kk'l} D'$, so D = D'. Hence D is minimal.

Remark 3.8. Lemma 3.7 implies that the minimal alphabets are in the orbit of the singletons $\{a\}$ for $a \in C$. Therefore, when searching for minimal alphabets we only need to compute these orbits instead of the entire graph *G*.

Example 3.9. Following Example 3.2, the minimal alphabets are the 2-periodic alphabets {0} and {1} and the 1-periodic alphabets {2} and {3}.

We also prove a natural fact that will be useful when we count minimal components in Section 6.

Proposition 3.10. *The minimal alphabets are pairwise disjoint.*

Proof. Let *D* and *D'* be two minimal alphabets of respective period *k* and *k'* such that $D \cap D' \neq \emptyset$. Let $a \in D \cap D'$. By Lemma 3.7, there exists $l_a \ge 1$ such that $\{a\} \xrightarrow{kl_a} D$ and $l'_a \ge 1$ such that $\{a\} \xrightarrow{kl'_a} D'$. By setting $l = \max(l_a, l'_a)$, we get $\{a\} \xrightarrow{kk'l} D$ and $\{a\} \xrightarrow{kk'l} D'$, hence D = D'.

3.1.2 l-primitivity on subalphabets

If *D* is a *k*-periodic alphabet, we have the restriction $\varphi^k|_{D\cup B} : D \cup B \to (D \cup B)^+$. We show here that the restrictions to minimal alphabets are precisely the l-primitive sub-substitutions of φ .

Lemma 3.11. Let D be a k-periodic alphabet. Then $\varphi^k|_{D\cup B}$ is l-primitive if and only if for all $a \in D$, there exists $l_a \ge 1$ such that $\{a\} \xrightarrow[k]{} D$.

Proof. Suppose that $\varphi^k|_{D\cup B}$ is l-primitive, i.e., there exists $n \ge 1$ such that, for all $a, b \in D$, $a \sqsubset \varphi^{kn}(b)$. So for all $a \in D$, $alph_C(\varphi^{kn}(a)) = D$, which means that $\{a\} \xrightarrow{k_n} D$.

Suppose that for all $a \in D$, there exists $l_a \ge 1$ such that $\{a\} \xrightarrow{kl_a} D$. By setting $n = \max_{a \in D} l_a$, we get $\{a\} \xrightarrow{kn} D$ for all $a \in D$, i.e., $alph_C(\varphi^{kn}(a)) = D$, which means that $\varphi^k|_{D \cup B}$ is l-primitive.

Lemmas 3.7 and 3.11 bring the following corollary:

Corollary 3.12. Let D be a k-periodic alphabet. Then D is minimal if and only if $\varphi^k|_{D\cup B}$ is l-primitive.

3.1.3 Dynamics of subshifts

A substitution $\varphi : A \to A^+$ induces a map $\tilde{\varphi}$ on the subshifts of X_{φ} : if $X \subset X_{\varphi}$ is a subshift, we define the subshift

$$\tilde{\varphi}(X) := \left\{ T^k(\varphi(x)) \mid x \in X, k \in \mathbb{Z} \right\} \subset X_{\varphi}.$$

Lemma 3.13. Let X be a subshift of X_{φ} . Then, for all $x \in X$, there exist $y \in X_{\varphi}$ and $k \in \mathbb{Z}$ such that $x = T^{k}(\varphi(y))$.

Proof. As $x \in X_{\varphi}$, for all $n \ge 0$, there exist $a_n \in A$ and $m_n \ge 1$ such that $x_{[-n,n]} \sqsubset \varphi^{m_n}(a_n) = \varphi(\varphi^{m_n-1}(a_n))$. Then there exist $u_n, v_n \in A^*$ and $k_n \in \llbracket 0, |\varphi| \rrbracket$ such that $u_n v_n = \varphi^{m_n-1}(a_n)$, $x_{[-n,k_n-1]}$ is a suffix of u_n and $x_{[k_n,n]}$ is a prefix if v_n . In particular, $|u_n v_n| \to_{n\to\infty} \infty$. By compacity, the couple $({}^{\omega}u_n v_n^{\omega}, k_n)$ has an adherent point (y,k), and $y \in X_{\varphi}$. By construction, for all $n \ge 0$, $x_{[-n,n]} = T^{k_n}({}^{\omega}\varphi(u_n)\varphi(v_n)^{\omega})_{[-n,n]} = T^{k_n}(\varphi({}^{\omega}u_n v_n^{\omega}))_{[-n,n]}$ so $T^k(\varphi(y)) = x$, therefore $y \in \varphi^{-1}(X)$.

This means that the reciprocal of $\tilde{\varphi}$ is defined, it is the subshift

$$\tilde{\varphi}^{-1}(X) := \left\{ x \in X_{\varphi} \mid \varphi(x) \in X \right\} \subset X_{\varphi}.$$

such that, for all subshift $X \subset X_{\varphi}$, $\tilde{\varphi}(\tilde{\varphi}^{-1}(X)) = X$. More generally, for all subshift $X \subset X_{\varphi}$ and all $n \ge 1$, $\tilde{\varphi}^n(\tilde{\varphi}^{-n}(X)) = X$.

Remark 3.14. If $X \subset X_{\varphi}$ is a subshift such that $alph_{C}(X) \neq \emptyset$, then $alph_{C}(\tilde{\varphi}(X)) \neq \emptyset$ and $alph_{C}(X) \rightarrow alph_{C}(\tilde{\varphi}(X))$. More generally, for $n \ge 1$, $alph_{C}(X) \rightarrow alph_{C}(\tilde{\varphi}^{n}(X))$.

We can now prove that the subshifts of X_{φ} have a strong cyclic behavior.

Proposition 3.15. Let $X \subset X_{\varphi}$ be a subshift. There exists $k \ge 1$ such that $\tilde{\varphi}^{k}(X) = X$, and if in addition $alph_{C}(X) \ne \emptyset$, then $alph_{C}(X)$ is a k-periodic alphabet and $X_{\varphi^{k}|alph_{C}(X)\cup B} \subset X$.

Proof. Consider the sequence of subshifts $(\tilde{\varphi}^{-n}(X))_{n\geq 0}$. By Proposition 2.3, X_{φ} has a finite number of subshifts so this sequence is ultimately periodic. By setting $l \geq 0$ and $k \geq 1$ the smallest integers such that $\tilde{\varphi}^{-l}(X) = \tilde{\varphi}^{-(l+k)}(X)$, we obtain $\tilde{\varphi}^{k}(X) = \tilde{\varphi}^{l+k}(\tilde{\varphi}^{-l}(X)) = \tilde{\varphi}^{l+k}(\tilde{\varphi}^{-(l+k)}(X)) = X$.

If $alph_{C}(X) \neq \emptyset$, with Remark 3.14 we have $alph_{C}(X) \xrightarrow{k} alph_{C}(\tilde{\varphi}^{k}(X)) = alph_{C}(X)$, which means that $alph_{C}(X)$ is *k*-periodic. Let $u \in \mathscr{L}(X_{\varphi^{k}|_{alph_{C}(X)\cup B}})$, then there exist $c \in alph_{C}(X)$ and $l \geq 1$ such that $u \sqsubset \varphi^{kl}(c)$. As $c \in alph_{C}(X)$, there exists $x \in X$ such that $c \sqsubset x$, then $u \sqsubset \varphi^{kl}(c) \sqsubset \varphi^{kl}(X) = X$ so $u \in \mathscr{L}(X)$. Therefore $\mathscr{L}(X_{\varphi^{k}|_{alph_{C}(X)\cup B}}) \subset \mathscr{L}(X)$, which means that $X_{\varphi^{k}|_{alph_{C}(X)\cup B}} \subset X$.

Remark 3.16. This means that $\tilde{\varphi}$ is a permutation on the subshifts of X_{φ} .

3.2 Proof of Theorem 2.21 (i)

With Corollary 3.12 we identified l-primitive sub-substitutions, now let us add tameness to obtain minimal components.

Proposition 3.17. Let $D \subset C_{niso}$ be a minimal alphabet of period k. Then $X_{\varphi^k|_{D\cup B}}$ is a tame minimal component of X_{φ} .

Proof. By Corollary 3.12, $\varphi^k|_{D\cup B}$ is 1-primitive. $\varphi^k|_{D\cup B}$ is also tame because $D \subset C_{niso}$, so Theorem 2.7 provides the minimality. The inclusion in X_{φ} follows directly from the definition of substitution subshifts.

We proved that $\varphi^{k}|_{D\cup B}$ is tame so, by Proposition 2.13, the words in $\mathscr{L}(X_{\varphi^{k}|_{D\cup B}}) \cap B^{*}$ have bounded length. This implies that $X_{\varphi^{k}|_{D\cup B}}$ contains growing letters, meaning that it is a tame minimal component of X_{φ} .

The converse relies on Proposition 3.15.

Proposition 3.18. Let X be a tame minimal component of X_{φ} . Then $alph_{C}(X) \subset C_{niso}$ and there exists $k \ge 1$ such that $alph_{C}(X)$ is a minimal alphabet of period k and $X = X_{\varphi^{k}|_{alph_{C}(X) \cup B}}$.

Proof. Proposition 3.15 provides $k \ge 1$ such that $alph_C(X)$ is *k*-periodic and $X_{\varphi^k|_{alph_C(X)\cup B}} \subset X$. Then, by minimality of *X*, we have equality.

As $alph_{\mathbb{C}}(X) \neq \emptyset$ and X is minimal, growing letters occur with bounded gaps in X. In other words, the words in $\mathscr{L}(X_{\varphi^k|_{alph_{\mathbb{C}}(X)\cup B}}) \cap B^*$ have bounded length, so, by Proposition 2.13, $\varphi^k|_{alph_{\mathbb{C}}(X)\cup B}$ is tame, i.e., $alph_{\mathbb{C}}(X) \subset C_{niso}$.

Proposition 3.15 also states that $\tilde{\varphi}^k(X) = X$, so, for all $d \in alph_C(X)$, $\varphi^{kl}(d) \in \mathscr{L}(X)$. As X is minimal, for l large enough, $\varphi^{kl}(d)$ contains every growing letter of $alph_C(X)$. This means that $\varphi^k|_{alph_C(X)\cup B}$ is l-primitive, and by Corollary 3.12 $alph_C(X)$ is minimal.

Propositions 3.17 and 3.18 complete the proof of Theorem 2.21 (i).

Example 3.19. Following Example 3.9, we recall that $C_{niso} = \{0, 1, 2\}$ so the minimal alphabets included in C_{niso} are $\{0\}$ of period 2, $\{1\}$ of period 2 and $\{2\}$ of period 1. We then add the bounded letters that appear in the associated subshift, so the tame minimal components of X_{φ} are $X_{\varphi^2|_{[0,5]}}$, $X_{\varphi^2|_{[1,4,6]}}$ and $X_{\varphi|_{[2,4,5_6]}}$.

4 Wild minimal components

4.1 Maximal bounded factors

The purpose of this section is to prove Proposition 4.35, which is an analog of a result proved for D0L-systems:

Proposition 4.1 ([10, Theorem 12]). Let $G = (A, \varphi, w)$ be a pushy D0L-system (i.e., φ is wild). Then there exists $L \ge 1$ and a finite subset $U \subset B^+$ such that any factor from $\mathscr{L}(G) \cap B^*$ has one of the following forms:

• w_1 • $w_1 u_1^{k_1} w_2$ • $w_1 u_1^{k_1} w_2 u_2^{k_2} w_3$ where $u_1, u_2 \in U$, $|w_j| < L$ for all $j \in \{1, 2, 3\}$, and $k_1, k_2 \ge 1$.

A similar result can also be found for purely morphic words in [4, Proposition 4.7.62]. We are going to be more precise by explicitly defining the words u_1 and u_2 as the computable words LP(c) and RP(c). To achieve that, we define the following: if $u \in A^*$, we say that v is a *maximal bounded factor* of u if $v \in \mathcal{L}(u) \cap B^*$ and it is not a factor of any other word of $\mathcal{L}(u) \cap B^*$. In particular, every word of $\mathcal{L}(u) \cap B^*$ is a factor of a maximal bounded factor of u.

As only growing letters generate images of arbitrarily large length, for all $u \in \mathscr{L}(\varphi) \cap B^*$, there exists $c \in C$ and $k \ge 0$ such that $u \sqsubset \varphi^k(c)$. In particular, u is a factor of a maximal bounded factor of $\varphi^k(c)$. We say that a word is a maximal bounded factor of a $\varphi^k(c)$.

4.1.1 1-blocks

We adapt here the notion of 1-blocks, introduced by Devyatov in [7] for morphic words, to substitution subshifts.

Definition 4.2. If $u \in A^*$ contains a growing letter, we denote the first growing letter in u by LC(u) and the prefix before LC(u) by $LB(u) \in B^*$. Symmetrically, we denote the last growing letter in u by RC(u) and the suffix after RC(u) by $RB(u) \in B^*$.

Example 4.3. If $C = \{0, 1\}$ and $B = \{2\}$, the words 22012001 and 212 have the following decomposition:

	22	0	1200	1	ε	1	2	0	2	
1	$ \rightarrow $	$ \frown$		\frown	\frown	and	\frown		$ \rightarrow $	•
	LB	LC		RC	RB		LB	LC=RC	RB	

Definition 4.4. If $c \in C$ and $k \ge 1$, a 1-block of $\varphi^k(c)$ is a triplet $(a, u, b) \in C \times B^* \times C$ such that $aub \sqsubset \varphi^k(c)$. In general, we say that (a, u, b) is a 1-block for φ if it is a 1-block of a $\varphi^k(c)$.

We now notice a most important fact:

Remark 4.5. Let $c \in C$ and $k \ge 1$. The maximal bounded factors of $\varphi^k(c)$ have one of the following forms:

- $LB(\varphi^k(c)),$
- *u* where (a, u, b) is a 1-block of $\varphi^k(c)$,
- $RB(\varphi^k(c))$.

Note that a 1-block for φ can be a 1-block of several $\varphi^k(c)$, and a 1-block of a $\varphi(c)$ is called an *origin*. In particular, φ has a finite number of origins.

Definition 4.6. Let $\mathfrak{u} = (a, u, b)$ be a 1-block of $\varphi^k(c)$. As $aub \sqsubset \varphi^k(c)$, we have $\varphi(a) \varphi(u) \varphi(b) \sqsubset \varphi^{k+1}(c)$, and in particular

$$v = (RC(\varphi(a)), RB(\varphi(a)), \varphi(u) LB(\varphi(b)), LC(\varphi(b)))$$

is a 1-block of $\varphi^{k+1}(c)$. We call v the *descendant* of u and we write v = Dc(u).

Lemma 4.7. Let $c \in C$. For all $k \ge 1$ and all 1-block \mathfrak{u} of $\varphi^k(c)$, there exist an origin \mathfrak{v} and $0 \le l < k$ such that $\mathfrak{u} = Dc^l(\mathfrak{v})$.

Proof. Let us prove it by induction on *k*. If k = 1, u is itself an origin and $u = Dc^0(u)$. Let $k \ge 1$ such that for every $c \in C$ and every 1-block u of $\varphi^k(c)$, there exist an origin v and $0 \le l < k$ such that $u = Dc^l(v)$. Let $c \in C$ and let u = (a, u, b) be a 1-block of $\varphi^{k+1}(c)$. When desubstituting once, the growing letters *a* and *b* come from growing letters in $\varphi^k(c)$, and we have two possibilities:

• Either *a* and *b* come from the same growing letter $a' \sqsubset \varphi^k(c)$.



Figure 1: $aub \sqsubset \varphi(a') \sqsubset \varphi^{k+1}(c)$.

Then $aub \sqsubset \varphi(a')$, which means that u is itself an origin. • Or *a* and *b* come from two different growing letters $a', b' \sqsubseteq \varphi^k(c)$.



Figure 2: The word $a'u'b' \sqsubset \varphi^k(c)$ generates $aub \sqsubset \varphi^{k+1}(c)$.

Then *a*' and *b*' are separated by the word *u*' in $\varphi^k(c)$, and $\varphi(u') \sqsubset u \in B^*$ so $u' \in B^*$. In particular, $\mathfrak{u}' = (a', u', b')$ is a 1-block of $\varphi^k(c)$ so, by hypothesis, there exists an origin \mathfrak{v} and $0 \le l < k$ such that $\mathfrak{u}' = Dc^l(\mathfrak{v})$. We have $a = RC(\varphi(a'))$ and $b = LC(\varphi(b'))$ so $u = RB(\varphi(a')) \varphi(u') LB(\varphi(b'))$, therefore $\mathfrak{u} = Dc(\mathfrak{u}') = Dc^{l+1}(\mathfrak{v})$.

This leads to the following definition.

Definition 4.8. An *evolution of 1-blocks* is a sequence \mathscr{E} where \mathscr{E}_0 is an origin for φ and, for all $l \ge 1$, $\mathscr{E}_l = Dc^l(\mathscr{E}_0)$. In particular, each evolution is determined by its first term and φ has a finite number of evolutions of 1-blocks.

Remark 4.9. Lemma 4.7 means that every 1-block for φ belongs to an evolution.

4.1.2 Decomposition of 1-blocks

In order to study the structure of 1-blocks, let us start with a useful lemma.

Lemma 4.10. *Let u be a word containing a growing letter. Then, for all* $k \ge 0$ *,*

$$LB(\varphi^{k}(u)) = \varphi^{k}(LB(u)) LB(\varphi^{k}(LC(u))),$$

$$RB(\varphi^{k}(u)) = RB(\varphi^{k}(RC(u))) \varphi^{k}(RB(u)).$$

Proof. We have $LB(\varphi^k(u)) = LB(\varphi^k(LB(u)) \varphi^k(LC(u))...) = \varphi^k(LB(u)) LB(\varphi^k(LC(u)))$. The right case is symmetric.

We deduce a decomposition of 1-blocks.

Lemma 4.11. Let \mathscr{E} be an evolution of 1-blocks for φ and set $\mathscr{E}_0 = (a, u, b)$. For all $l \ge 0$, we have

$$\mathscr{E}_l = \left(RC(\varphi^l(a)) , RB(\varphi^l(a)) \varphi^l(u) LB(\varphi^l(b)) , LC(\varphi^l(b)) \right).$$

Proof. Let us prove it by induction on *l*. If l = 0, this is directly true. Let $l \ge 0$ such that $\mathscr{E}_l = RB(\varphi^l(a)) \varphi^l(u) LB(\varphi^l(b))$. If $\mathscr{E}_{l+1} = (a', u', b')$, we have $\mathscr{E}_{l+1} = Dc(\mathscr{E}_l)$ so $a' = RC(\varphi(RC(\varphi^l(a)))) = RC(\varphi^{l+1}(a))$,

$$u' = RB(\varphi(RC(\varphi^{l}(a)))) \varphi(RB(\varphi^{l}(a))) \varphi^{l+1}(u) \varphi(LB(\varphi^{l}(b))) LB(\varphi(LC(\varphi^{l}(b))))$$
$$= RB(\varphi^{l+1}(a)) \varphi^{l+1}(u) LB(\varphi^{l+1}(b)) \text{ with Lemma 4.10,}$$

and
$$b' = LC(\varphi(LC(\varphi^{l}(b)))) = LC(\varphi^{l+1}(b)).$$

This lemma allows us to refine Remark 4.5:

Remark 4.12. Let $c \in C$ and $k \ge 1$. The maximal bounded factors of $\varphi^k(c)$ have one of the following forms:

(i)
$$LB(\varphi^{k}(c))$$
,
(ii) $RB(\varphi^{l}(a)) \varphi^{l}(u) LB(\varphi^{l}(b))$ where $0 \le l < k$ and (a, u, b) is an origin for φ ,
(iii) $RB(\varphi^{k}(c))$.

Now let us decompose the words $LB(\varphi^k(c))$ and $RB(\varphi^k(c))$.

4.1.3 Decomposition of $LB(\varphi^k(c))$

The following lemma provides a decomposition of $LB(\varphi^k(c))$ into k parts. Lemma 4.13. Let $c \in C$. Then, for all $k \ge 1$,

$$LB\left(\varphi^{k}(c)\right) = \bigsqcup_{j=0}^{k-1} \varphi^{k-j-1}\left(LB(\varphi(LC(\varphi^{j}(c))))\right).$$

Proof. Let us prove it by induction on k. If k = 1, this is directly true.

Let
$$k \ge 1$$
 such that $LB(\varphi^k(c)) = \bigsqcup_{j=0}^{k-1} \varphi^{k-j-1} (LB(\varphi(LC(\varphi^j(c))))))$. Then

$$LB(\varphi^{k+1}(c)) = \varphi(LB(\varphi^{k}(c))) LB(\varphi(LC(\varphi^{k}(c)))) \text{ with Lemma 4.10}$$
$$= \bigsqcup_{j=0}^{k-1} \varphi^{k-j}(LB(\varphi(LC(\varphi^{j}(c))))) . LB(\varphi(LC(\varphi^{k}(c)))))$$
$$= \bigsqcup_{j=0}^{k} \varphi^{k-j}(LB(\varphi(LC(\varphi^{j}(c))))).$$

In the rest of this section, we are going to identify the periodic structure of this decomposition. First, let us introduce the directed graph G_L , which corresponds to the directed graph UL_G in [10, Section 3.3].

Definition 4.14. In order to represent how φ acts on the leftmost letter of $\varphi^k(c)$, we define the directed graph $G_L := (V_L, E_L)$ by

- $V_L = C$, • $E_L := \{(c, LC(\varphi(c))) \mid c \in C\}.$
- If $(a, b) \in \mathscr{E}_L$, we write $a \xrightarrow{L} b$.

Definition 4.15. We say that a collection $\mathscr{C} = (c_i)_{0 \le i \le p-1} \in C^p$ is a *p*-cycle of G_L if the c_i are all distinct, for all $i \le p-2$, $c_i \xrightarrow{L} c_{i+1}$ and $c_{p-1} \xrightarrow{L} c_0$. If $c \in C$ belongs to a *p*-cycle of G_L , we say that *c* is *left-p-periodic*.

Example 4.16. Following Example 2.23, the graph *G*_{*L*} is the following:



The letters 0 and 1 are left-2-periodic, and the letters 2 and 3 are left-1-periodic.

Definition 4.17. If $\mathscr{C} = (c_i)_{0 \le i \le p-1}$ is a *p*-cycle of G_L , for all $i \in [[0, p-1]]$, we define the word

$$L(c_i) := \bigsqcup_{j=0}^{p-1} \varphi^{p-j-1} \left(LB(\varphi(c_{i+j[p]})) \right).$$

Remark 4.18. Let \mathscr{C} be a *p*-cycle of G_L . For all $c \in \mathscr{C}$, Lemma 4.13 provides $L(c) = LB(\varphi^p(c))$. This means that L(c) is the smallest bounded word generated at the left of $c: \varphi^p(c) = L(c) cv$. In particular, $c \in C_{liso}$ if and only if $L(c) \neq \varepsilon$.

Also, by construction, $L(c) \neq \varepsilon$ if and only if there exists $c \in \mathscr{C}$ such that $LB(\varphi(c)) \neq \varepsilon$. This means that, in a cycle of G_L , either every letter is left-isolated or no letter is left-isolated.

Remark 4.19. If $c \in C$ is left-*p*-periodic, then, for all $l \ge 0$, Lemma 4.13 also provides

$$LB\left(\varphi^{pl}(c)\right) = \bigsqcup_{j=0}^{l} \varphi^{p(l-j-1)}(L(c)).$$

Definition 4.20. Let $\mathscr{C} = \{c_i\}_{0 \le i \le p-1}$ be a *p*-cycle of G_L . For all $i \in [\![0, p-1]\!]$, $L(c_i) \in B^*$ so the sequence $(\varphi^{pi}(L(c)))_{i\ge 0}$ is eventually periodic. Set $q_i \ge 0$ and $p_i \ge 1$ the smallest integers such that $\varphi^{q_i+p_i}(LB(\varphi(c_i))) = \varphi^{q_i}(LB(\varphi(c_i)))$. Also set $q \ll max_{0\le i\le p-1} q_i$ and $p \ll lcm(c_0, ..., c_{p-1})$. Then, for all $i \in [\![0, p-1]\!]$, $q \ll$ and $p \ll$ are the smallest integers such that $\varphi^{q \ll +p_{\mathscr{C}}}(L(c_i)) = \varphi^{q \ll}(L(c_i))$. We then define the part before the left period

$$LQ(c) := \bigsqcup_{j=0}^{q_{\mathscr{C}}-1} \varphi^{p(q_{\mathscr{C}}-j-1)}(L(c)),$$

and the left period

$$LP(c) := \bigsqcup_{j=0}^{p_{\mathscr{C}}-1} \varphi^{p(q_{\mathscr{C}}+p_{\mathscr{C}}-j-1)}(L(c)),$$
(1)

so that $\varphi^{pp_{\mathscr{C}}}(LP(c)) = LP(c)$. In particular, $c \in C_{liso}$ if and only if $LP(c) \neq \varepsilon$.

Example 4.21. Following Example 4.16, we compute *LP* of the left-periodic letters.

• In the 2-cycle of G_L {0,1}, $\varphi(LB(\varphi(0))) = \varphi(LB(\varphi(1))) = \varepsilon$ so, by Remark 4.18, we have $L(0) = L(1) = \varepsilon$. This means that 0 and 1 are not left-isolated, and $LP(0) = LP(1) = \varepsilon$.

• In the 1-cycle of G_L {2}, $\varphi(LB(\varphi(2))) = \varepsilon$ so, similarly, $2 \notin C_{liso}$ and $LP(2) = \varepsilon$.

• In the 1-cycle of $G_L \mathcal{C} = \{3\}, \varphi(LB(\varphi(3))) = 54$. We have $L(3) = \varphi(LB(\varphi(3))) = 54$, $q_{\mathcal{C}} = 1$ and $p_{\mathcal{C}} = 2$. Then $3 \in C_{liso}$ and $LP(3) = \varphi^2(54) \varphi(54) = 5665$.

Remark 4.22. Let $\mathscr{C} = \{c_i\}_{0 \le i \le p-1}$ be a *p*-cycle of G_L . Then, for all $i \in \llbracket 0, p-1 \rrbracket$,

$$LP(c_i) = \bigsqcup_{j=0}^{pp_{\mathscr{C}}-1} \varphi^{p(q_{\mathscr{C}}+p_{\mathscr{C}})-j-1} \left(LB(\varphi(c_{i+j[p]})) \right).$$

Definition 4.23. Let \mathscr{C} be a *p*-cycle of G_L . If $l \ge q_{\mathscr{C}}$, there exist unique integers $l' \ge 0$ and $0 \le l'' \le p_{\mathscr{C}} - 1$ such that $l = q_{\mathscr{C}} + l'p_{\mathscr{C}} + l''$. Then, for all $c \in \mathscr{C}$, we define the suffix of LP(c)

$$LP_{l''}(c) := \bigsqcup_{j=0}^{l''-1} \varphi^{p(q_{\mathscr{C}}+l''-j-1)}(L(c)).$$

Remark 4.24. Let \mathscr{C} is a *p*-cycle of G_L . With Remark 4.19 and the definitions of LQ, LP and $LP_{l''}$, for all $c \in \mathscr{C}$ and for all $l = q_{\mathscr{C}} + l'p_{\mathscr{C}} + l'' \ge q_{\mathscr{C}}$ we have

$$LB\left(\varphi^{pl}(c)\right) = LP_{l''}(c) LP(c)^{l'} LQ(c).$$

Our goal is to decompose $LB(\varphi^k(c))$ when *c* is any growing letter, not necessarily left-periodic, and *k* is any integer.

Definition 4.25. If $c \in C$, set r_c the first integer such that $LC(\varphi^{r_c}(c))$ is a left-*p*-periodic letter. Let $\mathscr{C} = \{c_i\}_{0 \le i \le p-1}$ be the *p*-cycle of G_L such that $c_0 = LC(\varphi^{r_c}(c))$. For $k \ge r_c + pq_{\mathscr{C}}$, there exists unique integers $l \ge q_{\mathscr{C}}$ and $0 \le i \le p-1$ such that $k = r_c + i + pl = r_c + i + p(q_{\mathscr{C}} + l'p_{\mathscr{C}} + l'')$. We then define the word

$$LE_k(c) := \varphi^{p(q_{\mathscr{C}}+l'p_{\mathscr{C}}+l'')} \left(LB(\varphi^{r_c+i}(c)) \right) LP_{l''}(c_i).$$

Note that the words $LE_k(c)$ have bounded length because $LB(\varphi^{r_c+i}(c)) \in B^*$ and the $LP_{l''}(c_i)$ have bounded length.

We finally obtain the desired decomposition.

Proposition 4.26. Let $c \in C$ and let $\{c_i\}_{0 \le i \le p-1}$ be the *p*-cycle of G_L such that $c_0 = LC(\varphi^{r_c}(c))$. For all $k = r_c + i + p(q_{\mathscr{C}} + l'p_{\mathscr{C}} + l'') \ge r_c + pq_{\mathscr{C}}$, we have

$$LB(\varphi^{k}(c)) = LE_{k}(c) LP(c_{i})^{l'} LQ(c_{i}).$$

Proof. By construction, $LC(\varphi^{r_c+i}(c)) = c_i$, so

$$LB(\varphi^{k}(c)) = \varphi^{p(q_{\mathscr{C}}+l'p_{\mathscr{C}}+l'')} (LB(\varphi^{r_{c}+i}(c))) LB(\varphi^{p(q_{\mathscr{C}}+l'p_{\mathscr{C}}+l'')}(LC(\varphi^{r_{c}+i}(c))))$$
with Lemma 4.10
$$= \varphi^{p(q_{\mathscr{C}}+l'p_{\mathscr{C}}+l'')} (LB(\varphi^{r_{c}+i}(c))) LB(\varphi^{p(q_{\mathscr{C}}+l'p_{\mathscr{C}}+l'')}(c_{i}))$$
$$= \varphi^{p(q_{\mathscr{C}}+l'p_{\mathscr{C}}+l'')} (LB(\varphi^{r_{c}+i}(c))) LP_{l''}(c_{i}) LP(c_{i})^{l'} LQ(c_{i})$$
with Remark 4.24
$$= LE_{k}(c) LP(c_{i})^{l'} LQ(c_{i}).$$

Corollary 4.27. Let $\varphi : A \to A^+$ be a substitution. There exists a finite subset $Q_L \subset B^* \times B^*$ for which, for every $c \in C$ and every $k \ge 0$, there exist a left-periodic letter $a \in C$, $l \ge 0$ and $(u_1, u_2) \in Q_L$ such that

$$LB\left(\varphi^{k}(c)\right) = u_{1} LP(a)^{l} u_{2}.$$

Proof. Let $c \in C$ and let $\mathscr{C} = \{c_i\}_{0 \le i \le p-1}$ be the *p*-cycle of G_L such that $c_0 = LC(\varphi^{r_c}(c))$. If $k \ge r_c + pq_{\mathscr{C}}$, with Proposition 4.26 we can write $LB(\varphi^k(c)) = u_1 LP(c_i)^l u_2$ where $u_1 = LE_k(c)$ and $u_2 = LQ(c_i)$ for some *l* and *i*, and in that case there is a finite number of such u_1 and u_2 . If $k < r_c + pq_{\mathscr{C}}$, we can write $LB(\varphi^k(c)) = u_2$ and there is a finite number of such u_2 .

4.1.4 Decomposition of $RB(\varphi^k(c))$

This is the exact symmetric of the decomposition of $LB(\varphi^k(c))$, so we only give the key definitions and results.

Definition 4.28. In order to represent how φ acts on the rightmost letter of $\varphi^k(c)$, we define the directed graph $G_R := (V_R, E_R)$ by

• $V_R = C$, • $E_R := \{(c, RC(\varphi(c))) \mid c \in C\}.$

We have the same notions of *p*-cycle of G_R and *right-p*-periodic letters.

Example 4.29. Following Example 2.23, the graph *G*_{*R*} is the following:



The letters 0 and 1 are right-2-periodic, and the letters 2 and 3 are right-1-periodic.

Definition 4.30. If $\mathscr{C} = (c_i)_{0 \le i \le p-1}$ is a *p*-cycle of G_R , for all $i \in [[0, p-1]]$, we define the word

$$R(c_i) := \bigsqcup_{j=0}^{p-1} \varphi^j \left(RB(\varphi(c_{i+p-j-1[p]})) \right).$$

Definition 4.31. Let $\mathscr{C} = \{c_i\}_{0 \le i \le p-1}$ be a *p*-cycle of G_R . For all $i \in [\![0, p-1]\!]$, $R(c_i) \in B^*$ so the sequence $(\varphi^{pi}(R(c)))_{i\ge 0}$ is eventually periodic. Set $q_i \ge 0$ and $p_i \ge 1$ the smallest integers such that $\varphi^{q_i+p_i}(RB(\varphi(c_i))) = \varphi^{q_i}(RB(\varphi(c_i)))$. Also set $q_{\mathscr{C}} = \max_{0 \le i \le p-1} q_i$ and $p_{\mathscr{C}} = lcm(c_0, ..., c_{p-1})$. Then, for all $i \in [\![0, p-1]\!]$, $q_{\mathscr{C}}$ and $p_{\mathscr{C}}$ are the smallest integers such that $\varphi^{q_{\mathscr{C}}+p_{\mathscr{C}}}(R(c_i)) = \varphi^{q_{\mathscr{C}}}(R(c_i))$. We then define the part before the right period

$$RQ(c) := \bigsqcup_{j=0}^{q_{\mathscr{C}}-1} \varphi^{pj}(R(c)),$$

and the right period

$$RP(c) := \bigsqcup_{j=0}^{p_{\mathscr{C}}-1} \varphi^{p(q_{\mathscr{C}}+j)}(R(c))$$
(2)

so that $\varphi^{pp_{\mathscr{C}}}(RP(c)) = RP(c)$. In particular, $c \in C_{riso}$ if and only if $RP(c) \neq \varepsilon$.

Example 4.32. Following Example 4.29, we compute *RP* of the right-periodic letters.

• Similarly to the left side, $RP(0) = RP(1) = RP(2) = \varepsilon$.

• In the 1-cycle of $G_R \mathscr{C} = \{3\}, \varphi(RB(\varphi(3))) = 5$. We have $R(3) = \varphi(LB(\varphi(3))) = 5$, $q_{\mathscr{C}} = 0$ and $p_{\mathscr{C}} = 2$. Then $3 \in C_{riso}$ and $RP(3) = 5 \varphi(5) = 56$.

If $c \in \mathscr{C}$ a *p*-cycle of G_R , for $k \ge r_c + pq_{\mathscr{C}}$, there exists unique integers $l \ge q_{\mathscr{C}}$ and $0 \le i \le p - 1$ such that $k = r_c + i + pl = r_c + i + p(q_{\mathscr{C}} + l'p_{\mathscr{C}} + l'')$. We then define $RP_{l''}(c)$ as the prefix of RP(c), and $RE_k(c)$, both have bounded length and we obtain the same decomposition.

Proposition 4.33. Let $c \in C$ and let $\{c_i\}_{0 \le i \le p-1}$ be the p-cycle of G_R such that $c_0 = RC(\varphi^{r_c}(c))$. For all $k = r_c + i + p(q_{\mathscr{C}} + l'p_{\mathscr{C}} + l'') \ge r_c + pq_{\mathscr{C}}$, we have

$$RB(\varphi^{k}(c)) = RQ(c_{i}) RP(c_{i})^{l'} RE_{k}(c).$$

Corollary 4.34. Let $\varphi : A \to A^+$ be a substitution. There exists a finite subset $Q_R \subset B^* \times B^*$ for which, for every $c \in C$ and every $k \ge 0$, there exist a right-periodic letter $a \in C$, $l \ge 0$ and $(u_1, u_2) \in Q_R$ such that

$$RB\left(\varphi^{k}(c)\right) = u_1 RP(a)^l u_2.$$

4.1.5 Decomposition of maximal bounded factors

To recapitulate, Remark 4.12 states that every maximal bounded factor for φ can be decomposed into some $LB(\varphi^k(c))$ where $c \in C$, some $\varphi^k(u)$ where u is part of an origin, and some $RB(\varphi^k(c))$ where $c \in C$. The $\varphi^k(u)$ are easy to understand, and with Corollaries 4.27 and 4.34 we decomposed $LB(\varphi^k(c))$ and $RB(\varphi^k(c))$. We now deduce a precise decomposition of the maximal bounded factors.

Proposition 4.35. Let $\varphi : A \to A^+$ be a substitution. There exists a finite subset $Q \subset B^* \times B^* \times B^*$ such that every maximal bounded factor for φ has the form

$$u_1 RP(a)^p u_2 LP(b)^q u_3$$

where $(u_1, u_2, u_3) \in Q$, $a \in C$ is left-periodic, $b \in C$ is right-periodic and $p, q \ge 0$.

Proof. If *u* is a maximal factor of a $\varphi^k(c)$, Remark 4.12 gives three cases to consider:

(i) $u = LB(\varphi^k(c))$. By Corollary 4.27, there exists $(u_2, u_3) \in Q_L$, $b \in C$ leftperiodic and $q \ge 0$ such that $u = u_2 LP(b)^q u_3$, and there is a finite number of such u_2 and u_3 . (ii) $u = RB(\varphi^{l}(a)) \varphi^{l}(v) LB(\varphi^{l}(b))$ where (a, v, b) is an origin for φ . By Corollary 4.34, there exists $(u_1, v_1) \in Q_R$, $a \in C$ right-periodic and $p \ge 0$ such that $RB(\varphi^{l}(a)) = u_1 RP(a)^p v_1$. By Corollary 4.27, there exists $(v_2, u_3) \in Q_L$, $b \in C$ left-periodic and $q \ge 0$ such that $LB(\varphi^{l}(b)) = v_2 LP(b)^q u_3$. By setting $u_2 = v_1 \varphi^{l}(v)v_2$, we can write $u = u_1 RP(a)^p u_2 LP(b)^q u_3$, and there is a finite number of such u_1, u_2 and u_3 because $v \in B^*$.

(iii) $u = RB(\varphi^k(c))$. Similarly to the first case, we can write $u = u_1 RP(a)^p u_2$ with $a \in C$ right-periodic, and there is a finite number of such u_1 and u_2 .

Remark 4.36. This proposition decomposes the maximal bounded factor for φ , which will naturally provide a decomposition of every factor of $\mathscr{L}(\varphi) \cap B^*$. It makes two improvements from Proposition 4.1:

(i) It holds not only when φ is wild, but in the general case. In particular, this allows to prove Proposition 2.13: if φ is tame, then every LP(c) and RP(c) is empty, so the maximal bounded factors have bounded length; if φ is wild, then one LP(c) (resp. RP(c)) is non-empty so, by Proposition 4.26 (resp. Proposition 4.33), the words of $\mathscr{L}(X_{\varphi}) \cap B^*$ have unbounded length.

(ii) It explicitly defines the words that occur periodically in the decomposition.

4.2 **Proof of Theorem 2.21 (ii)**

We first show that the repetitions arising in Propositions 4.26 and 4.33 provide the wild minimal components exhibited in Remark 2.16.

Proposition 4.37. Let $c \in C_{liso}$ (resp. $c \in C_{riso}$). Then the subshift $X(^{\omega}LP(c)^{\omega})$ (resp. $X(^{\omega}RP(c)^{\omega})$) is a wild minimal component of X_{ω} .

Proof. Let $c \in C_{liso}$ be a left-*p*-periodic letter. We have $r_c = 0$ so, with Proposition 4.26, for all $l' \ge 0$, $LP(c)^{l'} \sqsubset LB\left(\varphi^{p(q_{\mathscr{C}}+l'p_{\mathscr{C}})}(c)\right)$. Therefore $X(^{\omega}LP(c)^{\omega}) \subset X_{\varphi}$. Symmetrically, if $c \in C_{riso}$, $X(^{\omega}RP(c)^{\omega}) \subset X_{\varphi}$.

The converse relies on Corollaries 4.27 and 4.34, this is a more precise version of Corollary 2.19.

Proposition 4.38. Let X be a wild minimal component of X_{φ} . Then X satisfies at least one of the following properties:

- There exists $c \in C_{liso}$ such that $X = X(^{\omega}LP(c)^{\omega})$.
- There exists $c \in C_{riso}$ such that $X = X(^{\omega}RP(c)^{\omega})$.

Proof. Let $x \in X$. Suppose that for every $c \in C_{liso}$ (resp. $c \in C_{riso}$), there exists $k_c \ge 1$ such that $LP(c)^{k_c} \not\subset x$ (resp. $RP(c)^{k_c} \not\subset x$). Set K to be the maximum of all k_c , and set $l_P := \max_{\substack{c \text{ left-periodic} \\ (u_1,u_2,u_3) \in Q}} |u_1| + |u_2| + |u_3|$ where Q is the finite set from Proposition 4.35. Let $u \sqsubset x$ such that $|u| = l_P + l_Q + 1$. As $v \in \mathscr{L}(X_{\varphi}) \cap B^*$, there exists a maximal bounded factor for φu such that $v \sqsubset u$, and, by Proposition 4.35, there

exist (u_1, u_2, u_3) , $a \in C$ right-periodic, $b \in C$ left-periodic and $p, q \ge 0$ such that $u = u_1 RP(a)^p u_2 LP(b)^q u_3$. Let us write $v = v_1 v_2 v_3 v_4 v_5$ where $v_1 \sqsubset u_1, v_2 \sqsubset RP(a)^p$, $v_3 \sqsubset u_2, v_4 \sqsubset LP(b)^q$ and $v_5 \sqsubset u_3$. We then have $|v_1| + |v_3| + |v_5| \le |u_1| + |u_2| + |u_3| \le l_Q$. If $a \in C_{riso}$, by definition of K we have $RP(a)^K \not \sqsubset v_2$ so $|v_2| < |RP(a)^{K+2}|$, otherwise $|v_2| = 0$. Similarly, if $b \in C_{liso}$, we have $|v_4| < |LP(b)^{K+2}|$, otherwise $|v_4| = 0$. Finally, we have $|v_2| + |v_4| \le l_P$, so $|v| \le l_Q + l_P$, which is a contradiction.

We just proved that either there exists $c \in C_{liso}$ such that for all $k \ge 1$, $LP(c)^k \sqsubset x$, which means that $X({}^{\omega}LP(c)^{\omega}) \subset X$, or there exists $c \in C_{riso}$ such that for all $k \ge 1$, $RP(c)^k \sqsubset x$, which means that $X({}^{\omega}RP(c)^{\omega}) \subset X$. As X is minimal, we get the equality. \Box

Propositions 4.37 and 4.38 complete the proof of Theorem 2.21 (ii).

Example 4.39. Following Examples 4.21 and 4.32, the wild minimal components of X_{φ} are $X(^{\omega}(5665)^{\omega})$ and $X(^{\omega}(56)^{\omega})$.

5 Dynamics of minimal components

In this section, we follow Remark 3.16 by showing that $\tilde{\varphi}$ acts on the tame (resp. wild) minimal components of X_{φ} as a permutation.

5.1 Dynamics of tame minimal components

Proposition 5.1. Let $D \subset C_{niso}$ be a minimal alphabet of period k and let E such that $D \to E$. Then $E \subset C_{niso}$ is a minimal alphabet of period k and $\tilde{\varphi}(X_{\varphi^k|_{D\cup B}}) = X_{\varphi^k|_{F\cup B}}$.

Proof. By Lemma 3.6, *E* is a minimal alphabet of period *k*. In particular, we have $E \xrightarrow[k-1]{k-1} D$. Let $e \in E$ be a left-periodic letter, and let \mathscr{C} be the cycle of G_L

such that $e \in \mathscr{C}$. Let $d \in \mathscr{C}$ such that $e \xrightarrow{L}_{k-1} d$, then $d = LC(\varphi^{k-1}(e)) \sqsubset \varphi^{k-1}(e)$ so $d \in D$ and $d \notin C_{liso}$. Then, by Remark 4.18, $e \notin C_{liso}$. We just proved that $E \cap C_{liso} = \emptyset$, and with the same argument $E \cap C_{riso} = \emptyset$ so $E \subset C_{niso}$.

Let $u \in \mathscr{L}(\tilde{\varphi}(X_{\varphi^{k}|_{D\cup B}}))$. This provides $x \in X_{\varphi^{k}|_{D\cup B}}$ and $n \in \mathbb{Z}$ such that $u \sqsubset T^{n}(\varphi(x))$. Then $u \sqsubset \varphi(x)$, so there exists $u' \sqsubset x$ such that $u \sqsubset \varphi(u')$. We have $u' \in \mathscr{L}(X_{\varphi^{k}|_{D\cup B}})$ so there exists $d \in D$ and $l \ge 1$ such that $u' \sqsubset \varphi^{kl}(d)$. We also have $E \xrightarrow[k-1]{} D$ so there exists $e \in E$ such that $d \sqsubset \varphi^{k-1}(e)$. Then $u \sqsubset \varphi(u') \sqsubset \varphi^{kl+1}(d) \sqsubset \varphi^{k(l+1)}(e)$, which means that $u \in \mathscr{L}(X_{\varphi^{k}|_{E\cup B}})$. We just proved that $\mathscr{L}(\tilde{\varphi}(X_{\varphi^{k}|_{D\cup B}})) \subset \mathscr{L}(X_{\varphi^{k}|_{E\cup B}})$, which means that $\tilde{\varphi}(X_{\varphi^{k}|_{D\cup B}}) \subset X_{\varphi^{k}|_{E\cup B}}$. Finally, by Proposition 3.17, $X_{\varphi^{k}|_{E\cup B}}$ is minimal so we have equality.

This allows us to define the following graph.

Definition 5.2. We define the directed graph $G_t = (V_t, E_t)$ by

- V_t the set of tame minimal components of X_{φ} ,
- $E_t := \{ (X, Y) \mid X, Y \in V_t, \tilde{\varphi}(X) = Y \}.$

Then Proposition 5.1 means that G_t is in correspondence with the subgraph of *G* restricted to the minimal alphabets included in C_{niso} .

Example 5.3. Following Examples 3.2 and 3.19, the graph *G*_t is the following:



5.2 Dynamics of wild minimal components

Let us give another way to show that two words are cyclic shifts.

Lemma 5.4. Let $u, v \in A^*$. If there exists $w \in A^*$ such that $w \sqsubset u$, $w \sqsubset v$ and wu = vw, then u is a cyclic shift of v.

Proof. $w \sqsubset u$ and $w \sqsubset v$ provide $u', v' \in A^*$ such that u = u'w and v = wv'. Then we have wu'w = wv'w, so u' = v'.

We deduce the dynamics of the wild minimal components.

Proposition 5.5. Let $a, b \in C_{liso}$ such that $a \xrightarrow{L} b$. Then $\tilde{\varphi}(X(^{\omega}LP(a)^{\omega})) = X(^{\omega}LP(b)^{\omega})$. *Proof.* Let $\mathscr{C} = \{c_i\}_{0 \le i \le p-1}$ be the *p*-cycle of G_L such that $a = c_0$ and $b = c_{1[p]}$ (*b* has index 1[*p*] in \mathscr{C} because *p* can be equal to 1). With Remark 4.22, we have

$$\begin{split} \varphi\left(LP(c_{0})\right) \ \varphi^{pq_{\mathscr{C}}}\left(LB(\varphi(c_{0}))\right) &= \bigsqcup_{j=0}^{pp_{\mathscr{C}}-1} \varphi^{p(q_{\mathscr{C}}+p_{\mathscr{C}})-j}\left(LB(\varphi(c_{j[p]}))\right) \ . \ \varphi^{pq_{\mathscr{C}}}\left(LB(\varphi(c_{0}))\right) \\ &= \varphi^{p(q_{\mathscr{C}}+p_{\mathscr{C}})}\left(LB(\varphi(c_{0}))\right) \bigsqcup_{j=1}^{pp_{\mathscr{C}}} \varphi^{p(q_{\mathscr{C}}+p_{\mathscr{C}})-j}\left(LB(\varphi(c_{j[p]}))\right) \\ &= \varphi^{pq_{\mathscr{C}}}\left(LB(\varphi(c_{0}))\right) \bigsqcup_{j=0}^{pp_{\mathscr{C}}-1} \varphi^{p(q_{\mathscr{C}}+p_{\mathscr{C}})-j-1}\left(LB(\varphi(c_{j+1[p]}))\right) \\ &= \varphi^{pq_{\mathscr{C}}}\left(LB(\varphi(c_{0}))\right) \ LP(c_{1[p]}). \end{split}$$

Then, by Lemma 5.4, φ (*LP*(*a*)) is a cyclic shift of *LP*(*b*).

With the same arguments, we have the symmetric result.

Proposition 5.6. Let $a, b \in C_{riso}$ such that $a \xrightarrow{R} b$. Then $\tilde{\varphi}(X({}^{\omega}RP(a){}^{\omega})) = X({}^{\omega}RP(b){}^{\omega})$.

This allows us to define the following graph.

Definition 5.7. We define the directed graph $G_w := (V_w, E_w)$ by

- V_w the set of wild minimal components of X_{φ} ,
- $E_w := \{(X, Y) \mid X, Y \in V_w, \tilde{\varphi}(X) = Y\}.$

Then Propositions 5.5 and 5.6 mean that G_w is determined by G_L and G_R .

Remark 5.8. Note that different letters $a, b \in B$ can satisfy $X(^{\omega}LP(a)^{\omega}) = X(^{\omega}LP(b)^{\omega})$, $X(^{\omega}LP(a)^{\omega}) = X(^{\omega}RP(b)^{\omega})$ or $X(^{\omega}RP(a)^{\omega}) = X(^{\omega}RP(b)^{\omega})$. Therefore, G_w is in correspondence with the union of the subgraph of G_L restricted to C_{liso} and the subgraph of G_R restricted to C_{riso} .

Example 5.9. Following Examples 4.16, 4.29 and 4.39, the graph G_w is the following:



6 Counting minimal components

6.1 Computing *B* and *C*

We take inspiration in [7, Lemmas 3.1 and 3.2] to define the set of *periodic* letters $P \subset B$ which is essential to characterize *B*.

Definition 6.1. If $a \in B$, either *a* occurs in a $\varphi^n(a)$ and we say that *a* is *periodic*, or *a* never occurs in $\varphi^n(a)$ and we say that *a* is *pre-periodic*. We write *P* the set of periodic letters and *PP* the set of pre-periodic letters.

Lemma 6.2. Let $a \in A$. Then the following holds.

(*i*) $a \in P$ if and only if there exists $n \ge 1$ such that $\varphi^n(a) = a$. (*ii*) If $a \in P$, then, for all $m \ge 1$, $\varphi^m(a) \in P$. (*iii*) $a \in B$ if and only if $\varphi^{|A|}(a) \in P^+$.

Proof. (i) If *a* is periodic, there exist $n \ge 1$ and $u, v \in A^*$ such that $\varphi^n(a) = uav$. If $(u, v) \ne (\varepsilon, \varepsilon)$, then, by iterating the substitution φ^n on *a*, we would get $a \in C$, therefore $\varphi^n(a) = a$.

If there exists $n \ge 1$ such that $\varphi^n(a) = a$, then, for all $m \ge 1$, $\varphi^{mn}(a) = a$ so $1 = |\varphi^{mn}(a)| \ge |\varphi^m(a)| \ge 1$ so $|\varphi^m(a)| = 1$. This means that $a \in B$, and it occurs in $\varphi^n(a)$ so $a \in P$.

(ii) If $a \in P$, we already proved that, for all $m \ge 1$, $|\varphi^m(a)| = 1$. (i) provides $n \ge 1$ such that $\varphi^n(a) = a$, so $\varphi^n(\varphi^m(a)) = \varphi^m(\varphi^n(a)) = \varphi^m(a)$) and, by (i), $\varphi^m(a) \in P$.

(iii) If $a \in B$, suppose that $\varphi^{|A|}(a) \notin P^+$. In particular, there exists a letter $a_{|A|} \in PP$ such that $a_{|A|} \sqsubset \varphi^{|A|}(a)$. This provides $a_{|A|-1} \sqsubset \varphi^{|A|-1}(a)$ such that $a_{|A|} \sqsubset \varphi(a_{|A|-1})$, and, by (ii), $a_{|A|-1} \in PP$. By iterating this construction, we obtain a finite sequence $(a_i)_{0 \le i \le |A|} \in PP^{|A|+1}$ such that $a_{i+1} \sqsubset \varphi(a_i)$. We have |A| + 1 letters so there exists i < j such that $a_i = a_j$, and then $a_i \sqsubset \varphi^{j-i}(a_i)$ so $a_i \notin PP$, contradiction.

If $\varphi^{|A|}(a) \in P^+$, in particular $\varphi^{|A|}(a) \in B^+$ so $a \in B$.

We now have a method to compute *B*, as stated in Proposition 2.25.

Proof of Proposition 2.25. We first compute the alphabet *P*: for $a \in A$, we compute the $\varphi^n(a)$ and either we reach *n* such that $|\varphi^n(a)| \ge 2$, and by Lemma 6.2 (ii) $a \notin P$, or we reach *n* for which there exists m < n such that $\varphi^n(a) = \varphi^m(a) \in A$, and if m = 0 then, by Lemma 6.2 (ii), $a \in P$, or if m > 0 then $a \notin P$.

We can now compute *B*: for $a \in A$, we compute $\varphi^{|A|}(a)$ and we check if it belongs to P^+ to use Lemma 6.2 (iii). The remaining letters are in *C*.

6.2 Computing $MC(\varphi)$

Essentially, computing the minimal components is the same as counting them. We recall that the number of minimal components of X_{φ} is $MC(\varphi)$, and we also define $TMC(\varphi)$ the number of tame minimal components and $WMC(\varphi)$ the number of wild minimal components.

Proposition 6.3. Let $\varphi : A \to A^+$ be a substitution. Then TMC(φ) is computable.

Proof. Once Proposition 2.25 provides *C*, we can compute *G*, the minimal alphabets and their period. By checking if the minimal alphabets contain a left or right-isolated letter, we obtain $TMC(\varphi)$.

Proposition 6.4. Let $\varphi : A \to A^+$ be a substitution. Then WMC(φ) is computable.

Proof. Once Proposition 2.25 provides *B*, we can compute G_L , G_R , LP(c) and RP(c) for every left or right-periodic letter. In order to differenciate the $X(^{\omega}LP(c)^{\omega})$ and $X(^{\omega}RP(c)^{\omega})$, we compute the primitive root of the words LP(c) and RP(c) and we check if one is a cyclic shift of another to remove duplicates.

Propositions 6.3 and 6.4 complete the proof of Corollary 2.26. For details about the implementation of these algorithms, the reader can take a look at the associated Python code.

6.3 Bounding $MC(\varphi)$

Let us first bound *TMC*(φ) and *WMC*(φ) separately.

Lemma 6.5. Let $\varphi : A \to A^+$ be a substitution. Then (i) $TMC(\varphi) \le |C_{niso}|$, (ii) $WMC(\varphi) \le |C_{liso}| + |C_{riso}|$.

Proof. (i) Set *n* to be the number of minimal alphabets included in C_{niso} . By Theorem 2.21, $TMC(\varphi) = n$, and with Proposition 3.10, $n \le |C_{niso}|$.

(ii) $WMC(\varphi) \le |C_{liso}| + |C_{riso}|$ comes directly from Theorem 2.21 (ii).

We can then bound $MC(\varphi)$ using the size of the alphabet, as stated in Corollary 2.27.

Proof of Corollary 2.27. By Lemma 6.5, we have $MC(\varphi) = TMC(\varphi) + WMC(\varphi) \le |C_{niso}| + |C_{liso}| + |C_{riso}|$.

(i) If $B = \emptyset$, $C_{niso} = C = A$ and $C_{liso} = C_{riso} = \emptyset$ so $MC(\varphi) \le |C_{niso}| = |C| = |A|$.

(ii) If $B = \{b\}$, $\{{}^{\omega}b{}^{\omega}\}$ is the only possible wild minimal component, so $WMC(\varphi) \leq 1$. If $C_{liso} = C_{riso} = \emptyset$, then $MC(\varphi) \leq |C_{niso}| \leq |C| = |A| - 1$. If $C_{liso} \neq \emptyset$ or $C_{riso} \neq \emptyset$, then $MC(\varphi) \leq |C_{niso}| + 1 \leq |C| = |A| - 1$.

(iii) If $|B| \ge 2$, $|C_{liso}| \le |C| - |C_{niso}|$ and $|C_{riso}| \le |C| - |C_{niso}|$ so $MC(\varphi) \le 2|C| - |C_{niso}| \le 2|C| \le 2|A| - 4$.

The following examples show that these bounds are optimal.

(i) This bound is reached every singleton in *G* is a minimal alphabet.

Example 6.6. Let $k \ge 1$, $A = \{a_1, ..., a_k\}$ and for all $i \in \llbracket 1, k \rrbracket$, $\varphi : a_i \mapsto a_i a_i$. We have $B = \emptyset$, the minimal alphabets are the 1-periodic alphabets $\{a_i\}$ and they generate the tame minimal components $\{{}^{\omega}a_i^{\omega}\}$, therefore $MC(\varphi) = |A|$.

(ii) This bound is reached similarly to (i), but we have two possibilities: • $TMC(\varphi) = |A| - 1$ and $WMC(\varphi) = 0$

Example 6.7. Let $k \ge 2$, $A = \{b, a_1, ..., a_{k-1}\}$ and $\varphi : a_i \mapsto a_i a_i$ for all $i \in \llbracket 1, k-2 \rrbracket$, $a_{k-1} \mapsto a_{k-1} ba_{k-1}, b \mapsto b$. We have $B = \{b\}$, the minimal alphabets included in C_{niso} are the $\{a_i\}$ for $i \in \llbracket 1, k-1 \rrbracket$ and they generate the tame minimal components $X_{\varphi|_{[a_i,b]}} = \{{}^{\omega}(a_i b){}^{\omega}\}$ for $i \in \llbracket 1, k-1 \rrbracket$. Therefore $MC(\varphi) = |A| - 1$.

• $TMC(\varphi) = |A| - 2$ and $WMC(\varphi) = 1$

Example 6.8. Let $k \ge 2$, $A = \{b, a_1, ..., a_{k-1}\}$ and $\varphi : a_i \mapsto a_i a_i$ for all $i \in [\![1, k-2]\!]$, $a_{k-1} \mapsto a_{k-1}b, b \mapsto b$. We have $B = \{b\}$ and $a_{k-1} \in C_{riso}$ so the minimal alphabets included in C_{niso} are the 1-periodic alphabets $\{a_i\}$ for $i \in [\![1, k-2]\!]$ and they generate the tame minimal components $X_{\varphi|_{[a_i]}} = \{{}^{\omega}a_i^{\omega}\}$ for $i \in [\![1, k-2]\!]$. Moreover, $\{{}^{\omega}b{}^{\omega}\}$ is the unique wild component, therefore $MC(\varphi) = |A| - 1$.

(iii) This bound is reached when |B| = 2, $C_{liso} = C_{riso} = C$ and all $X(^{\omega}LP(c)^{\omega})$ and $X(^{\omega}RP(c)^{\omega})$ are distinct.

Example 6.9. Let $k \ge 3$, $A = \{a, b, c_1, ..., c_{k-2}\}$ and $\varphi : a \mapsto a, b \mapsto b, c_i \mapsto a^{2i-2}bc_ia^{2i-1}b$. We have $B = \{a, b\}$, $C = \{c_i \mid 1 \le i \le k-2\}$ and every growing letter is left-isolated with period 1 and right-isolated with period 1. Then the minimal components of X_{φ} are the $X(^{\omega}(a^ib)^{\omega})$ for all $0 \le i \le 2k-5$, therefore $MC(\varphi) = 2k - 4 = 2|A| - 4$.

Remark 6.10. By doing any combination of $LP(c_i)$ and $RP(c_i)$ in Example 6.9, $MC(\varphi)$ can reach every number between 1 and 2|A| - 4.

7 Minimal components on two letters

Substitutions on one letter are trivial, but substitutions on the binary alphabet $\{0, 1\}$ already generate interesting subshifts. We have a reasonable number of

cases to analyse, so let us compute and display the minimal components of X_{φ} for all of them. We recall that, in order to generate a subshift, substitutions must have a growing letter so, without loss of generality, we assume that the letter 0 as always growing. Then the first distinction is wether 1 is growing or bounded.

7.1 $C = \{0, 1\}$

Theorem 2.21 tells us that X_{φ} only has tame minimal components, and Corollary 2.27 (i) tells us that it has at most two. They are characterized by the minimal alphabets, and we recall from Remark 3.4 that *G* is uniquely determined by its generators *D* and *D'* such that $\{0\} \rightarrow D$ and $\{1\} \rightarrow D'$. The following table displays the minimal components in each case, white cells mean that X_{φ} is minimal, light gray cells mean that X_{φ} is sometimes minimal and sometimes not, and gray cells mean that X_{φ} is not minimal.

D' D	{1}	{0}	{0,1}
{0}	$\{{}^{\omega}0{}^{\omega}\},\{{}^{\omega}1{}^{\omega}\}$	$\{\omega 0^{\omega}\}$	$\{\omega 0^{\omega}\}$
{1}	$\{\omega 1^{\omega}\}$	$\{{}^{\omega}0{}^{\omega}\},\{{}^{\omega}1{}^{\omega}\}$	X_{φ}
{0,1}	$\{^{\omega}1^{\omega}\}$	X_{φ}	X_{φ}

Let us detail the computations. Note that switching 0 and 1 brings the analysis down to six cases.

7.1.1 $D = \{0\}, D' = \{1\}$

This is one of the two cases where X_{φ} has two minimal components. The orbits in *G* are



The minimal alphabets are the 1-periodic alphabets {0} and {1} so the minimal components of X_{φ} are $X_{\varphi|_{[0]}} = {}^{\omega}0^{\omega}$ } and $X_{\varphi|_{[1]}} = {}^{\omega}1^{\omega}$ }.

Example 7.1. Consider the substitution $\varphi : 0 \mapsto 00, 1 \mapsto 11$.

7.1.2 $D = \{1\}, D' = \{1\}$

The orbits in *G* are



The unique minimal alphabet is the 1-periodic alphabet {1} so the unique minimal component of X_{φ} is $X_{\varphi|_{[1]}} = \{^{\omega}1^{\omega}\}$. In fact this is the pathological case where $\varphi(\{0,1\}) \in \{1\}^+$, so $X_{\varphi} = \{^{\omega}1^{\omega}\}$ and it is minimal.

Example 7.2. Consider the substitution $\varphi : 0 \mapsto 1, 1 \mapsto 11$.

7.1.3 $D = \{1\}, D' = \{0\}$

This is the second case when X_{φ} has two minimal components. The orbits in *G* are



The minimal alphabets are the 2-periodic alphabets {0} and {1} so the minimal components of X_{φ} are $X_{\varphi^2|_{[0]}} = \{{}^{\omega}0^{\omega}\}$ and $X_{\varphi^2|_{[1]}} = \{{}^{\omega}1^{\omega}\}$.

Example 7.3. Consider the substitution $\varphi : 0 \mapsto 11, 1 \mapsto 00$.

7.1.4 $D = \{0, 1\}, D' = \{1\}$

The orbits in *G* are



The unique minimal alphabet is the 1-periodic alphabet {1} so the unique minimal component of X_{φ} is $X_{\varphi|_{(1)}} = \{\omega 1^{\omega}\}$. In this case, there are examples where X_{φ} is not minimal and others where it is minimal.

Example 7.4. Consider the substitution $\varphi : 0 \mapsto 101, 1 \mapsto 11$. We have $X_{\varphi} = {}^{\omega}101^{\omega}$ } so it is not minimal.

Example 7.5. Consider the substitution $\varphi : 0 \mapsto 01, 1 \mapsto 11$. Every word of $\mathscr{L}(\varphi)$ contains at most one 0, so $X_{\varphi} = \{{}^{\omega}1^{\omega}\}$ and it is minimal.

7.1.5 $D = \{0, 1\}, D' = \{0\}$

The orbits in *G* are



The only minimal alphabet is the 1-periodic alphabet $\{0, 1\}$ so the unique minimal component is X_{φ} itself. A more direct method is to apply Theorem 2.7 (i), which tells us that X_{φ} is minimal.

Example 7.6. Consider the Fibonacci substitution $\varphi : 0 \mapsto 01, 1 \mapsto 0$.

7.1.6 $D = \{0, 1\}, D' = \{0, 1\}$

This is the primitive case. The orbits in *G* are



Similarly to the previous case, X_{φ} is minimal.

Example 7.7. Consider the Thue-Morse substitution $\varphi : 0 \mapsto 01, 1 \mapsto 10$.

7.2 $C = \{0\}, B = \{1\}$

In this case, φ is l-primitive and the unique minimal alphabet is {0}. Also, Corollary 2.27 (ii) tells us that X_{φ} has a unique minimal component, that will depend on wether φ is tame or wild.

7.2.1 φ is tame

The alphabet $\{0\} \subset C_{niso}$ is minimal so, by Theorem 2.21 (ii), the unique minimal com- ponent of X_{φ} is itself. A more direct method is to apply Theorem 2.7 (i), which tells us that X_{φ} is minimal.

Example 7.8. Consider Example 6.7 where k = 2, that is $\varphi : 0 \mapsto 010, 1 \mapsto 1$. Then $X_{\varphi} = X(^{\omega}(01)^{\omega})$ is a single periodic orbit.

7.2.2 φ is wild

Necessarily, $LP(0) \in \{1\}^+$ or $RP(0) \in \{1\}^+$, so, by Theorem 2.21 (ii), the minimal component of X_{φ} is $\{{}^{\omega}1{}^{\omega}\}$. There are examples where X_{φ} is not minimal and others where it is minimal.

Example 7.9. In Example 2.17, $X_{\varphi} = X(^{\omega}101^{\omega})$ and its unique minimal component is $\{^{\omega}1^{\omega}\}$, so X_{φ} is not minimal.

Example 7.10. Consider Example 6.8 where k = 2, that is $\varphi : 0 \mapsto 01, 1 \mapsto 1$. We have $X_{\varphi} = {}^{\omega}1^{\omega}$ so it is minimal.

7.3 Larger alphabets

In general the number of cases explodes. Just looking at the graph *G*, there are $(2^{|C|} - 1)^{|C|}$ possible graphs, and finding how many there are up to permutation of letters seems really hard.

Given the graph *G*, set MA(G) to be the number of minimal alphabets in *G*. By Theorem 2.21 (i), we have $TMC(\varphi) \le MA(G) \le |C|$, and $TMC(\varphi) = MA(G)$ when φ is growing. The following table displays the number of graphs *G* on |C| letters who have MA(G) minimal alphabets. The computations were done with the code shared with this paper.

C MA(G)	1	2	3	4	5
1	1	0	0	0	0
2	7	2	0	0	0
3	292	45	6	0	0
4	46739	3490	372	24	0
5	27714556	848005	62690	3780	120

This strongly suggests that almost all graphs *G* have a unique minimal alphabet.

Remark 7.11. We would also like to do statistics on the substitutions instead of the graphs *G*, for example: computing the distribution of $MC(\varphi)$ when |A| and $|\varphi|$ are fixed.

Acknowledgements

The author thanks F. Durand and S. Starosta for kindly detailing their work. He also thanks J. Cassaigne for his precious insights.

References

- [1] J.P. Allouche and J. Shallit. *Automatic Sequences: Theory, Applications, Generalizations*. Cambridge University Press, 2003.
- [2] M.P. Béal, F. Durand, and D. Perrin. *Symbolic dynamics and substitutions*. 2024. Preprint.
- [3] M.P. Béal, D. Perrin, and A. Restivo. Decidable problems in substitution shifts. *Journal of Computer and System Sciences*, 143:103529, 2024.
- [4] J. Cassaigne and F. Nicolas. Factor complexity. In V. Berthé, editor, Combinatorics, Automata and Number Theory, Encyclopedia of Mathematics and its Applications 135, pages 163–247. Cambridge University Press, 2010.
- [5] M.I. Cortez and B. Solomyak. Invariant measures for non-primitive tiling substitutions. *Journal d'Analyse Mathématique*, 115:293–342, 2010.
- [6] D. Damanik and D. Lenz. Substitution dynamical systems: characterization of linear repetitivity and applications. *Journal of Mathematical Analysis* and Applications, 321(2):766–780, 2006.
- [7] R. Devyatov. On factor complexity of morphic sequences. *Moscow Mathematical Journal*, 18:211–303, 04 2018.

- [8] F. Durand. A theorem of Cobham for non-primitive substitutions. *Acta Arithmetica*, 104(3):225–241, June 2002.
- [9] A. Ehrenfeucht and G. Rozenberg. Repetition of subwords in D0L languages. *Information and Control*, 59(1):13–35, 1983.
- [10] K. Klouda and Š. Starosta. An algorithm for enumerating all infinite repetitions in a D0L-system. *Journal of Discrete Algorithms*, 33:130–138, July 2015.
- [11] G. Maloney and D. Rust. Beyond primitivity for one-dimensional substitution subshifts and tiling spaces. *Ergodic Theory and Dynamical Systems*, 38(3):1086–1117, 2018.
- [12] J.J. Pansiot. Complexité des facteurs des mots infinis engendrés par morphismes itérés. In J. Paredaens, editor, Automata, Languages and Programming, pages 380–389. Springer Berlin Heidelberg, 1984.
- [13] N. Pythéas Fogg, V. Berthé, S. Ferenczi, C. Mauduit, and A. Siegel. Substitutions in dynamics, arithmetics and combinatorics. Lecture Notes in Mathematics 1794. Springer Berlin Heidelberg, 2002.
- [14] M. Queffélec. Substitution dynamical systems spectral analysis, volume 1294 of Lect. Notes Math. Springer, Cham, 1987.
- [15] T. Shimomura. A simple approach to minimal substitution subshifts. *Topology and its Applications*, 260:203–214, 2019.