

What is causal about causal models and representations?

Frederik Hytting Jørgensen, Luigi Gresele, and Sebastian Weichwald

Copenhagen Causality Lab, Department of Mathematical Sciences, and Pioneer Centre for AI,
University of Copenhagen, Denmark

Abstract

Causal Bayesian networks are ‘causal’ models since they make predictions about interventional distributions. To connect such causal model predictions to real-world outcomes, we must determine which actions in the world correspond to which interventions in the model. For example, to interpret an action as an intervention on a treatment variable, the action will presumably have to a) change the distribution of treatment in a way that corresponds to the intervention, and b) not change other aspects, such as how the outcome depends on the treatment; while the marginal distributions of some variables may change as an effect. We introduce a formal framework to make such requirements for different interpretations of actions as interventions precise. We prove that the seemingly natural interpretation of actions as interventions is circular: Under this interpretation, every causal Bayesian network that correctly models the observational distribution is trivially also interventionally valid, and no action yields empirical data that could possibly falsify such a model. We prove an impossibility result: No interpretation exists that is non-circular and simultaneously satisfies a set of natural desiderata. Instead, we examine non-circular interpretations that may violate some desiderata and show how this may in turn enable the falsification of causal models. By rigorously examining how a causal Bayesian network could be a ‘causal’ model of the world instead of merely a mathematical object, our formal framework contributes to the conceptual foundations of causal representation learning, causal discovery, and causal abstraction, while also highlighting some limitations of existing approaches.

1 Introduction

Causal Bayesian networks are mathematical models that induce multiple distributions over some random variables [Spirtes et al., 2001, Pearl, 2009, Peters et al., 2017]. A causal Bayesian network describes one reference distribution, called the observational distribution, and a procedure to derive interventional distributions. As such, a causal Bayesian network is a concise mathematical model of several distributions indexed by interventions.

Causal reasoning using causal models is seemingly intuitive once we assign names to the variables in the model based on the real-world quantities they aim to represent. The term ‘intervention’ is suggestive and one might use interventions on model variables to reason about actions that perturb the corresponding real-world quantities. Yet, without making the correspondence between model interventions and actions explicit, we blur the line between mathematical model and real-world substantiation. It is then unclear

what predictions the model makes about the effects of actions in the world and how interventional predictions may be used to falsify a causal model.

1.1 Dialogue – What is an intervention?

The following dialogue illustrates the conflict one runs into when using a causal Bayesian network, a mathematical model, to reason about actions and observations of some real-world quantities while using the word ‘intervention’ ambiguously.

Omar: I have two quantities that I model as random variables A and B . I have measured them and the data seems to perfectly match the joint normal distribution

$$\mathcal{L}^{\mathcal{O}}(A, B) = \mathcal{N}\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}\right).$$

I am sure that there is no unobserved confounding,¹ but I am not sure if A causes B or B causes A . Do you think $A \rightarrow B$ or $A \leftarrow B$ is correct?

Sofia: I am sure that A causes B .

Omar: How do you know?

Sofia: Try to intervene on B . If A causes B , then we would expect that intervening on B changes the conditional distribution of B given A but does not change the marginal distribution of A . For example, if you intervene to set B equal to 5, you will observe that (A, B) follows joint distribution

$$\mathcal{L}^{\text{do}(B=5)}(A, B) = \mathcal{N}(0, 1) \otimes \delta_5,$$

where δ_5 is the Dirac distribution with support $\{5\}$.

Omar: Okay, I tried. I did something and now I observe that A and B follow joint normal distribution

$$\mathcal{N}\left(\begin{pmatrix} 2 \\ 0 \end{pmatrix}, \begin{pmatrix} 4 & 3 \\ 3 & 7 \end{pmatrix}\right).$$

Sofia: Now B follows distribution $\mathcal{N}(0, 7)$. I proposed that you intervene to make B have point mass in 5. Could you try again?

Omar: Ah, sorry. I thought I implemented the intervention you suggested, but I can see now that I did not. I will try something different such that B has point mass in 5: Now, A and B have distribution

$$\delta_1 \otimes \delta_5.$$

So B has distribution δ_5 , but I did not get the interventional distribution that you said I would get. Does that mean that A does not cause B ?

Sofia: Haha, you also intervened on A and set it equal to 1! When you intervene on B , you only intervene on B . Make sure to only change the conditional distribution of B given A . Do not change the marginal distribution of A .

¹This assumption eases the presentation of the example but is not necessary to arrive at the problem this example illustrates.

Omar: Okay, now I think I did the right thing and indeed, just like you predicted, I now get

$$\mathcal{N}(0, 1) \otimes \delta_5.$$

Sofia: See, as I predicted, A causes B and not vice-versa.

1.2 What went wrong?

Let us assume that Sofia is right that $A \rightarrow B$. Based on her correct model (and the observational distribution), she makes the following prediction:

- (P) If you intervene $\text{do}(B = 5)$, then you will observe the distribution $\mathcal{N}(0, 1) \otimes \delta_5$ over (A, B) .

Proposition (P) is ambiguous: It is clear what distribution Sofia’s model implies under the intervention $\text{do}(B = 5)$, however, it is unclear when the antecedent is satisfied in the world. Here is an explanation provided by Pearl, Glymour, and Jewell:

The difference between intervening on a variable and conditioning on that variable should, hopefully, be obvious. When we intervene on a variable in a model, we fix its value. We *change* the system, and the values of other variables often change as a result. [Pearl et al., 2016, page 54]

One possible way to understand this in the context of (P) is as follows:

- (P1) If you do something to change the system such that you observe $B = 5$ with probability 1, then you will observe the distribution $\mathcal{N}(0, 1) \otimes \delta_5$ over (A, B) .

Even though Sofia believes that $A \rightarrow B$, she apparently thinks that (P1) is false. When Omar does something such that (A, B) has distribution $\delta_1 \otimes \delta_5$ and the antecedent of (P1) is thus satisfied, she objects that Omar intervened on both nodes. Instead, she proposes (P2) as an analysis of (P):

- (P2) If you do something to change the system such that you observe $B = 5$ with probability 1 while not changing the marginal distribution of A , then you will observe the distribution $\mathcal{N}(0, 1) \otimes \delta_5$ over (A, B) .

Interpreting (P) as (P2) cannot be correct because if $A \leftarrow B$, then (P) is false and (P2) is true. (P2) is a mathematical truth that holds irrespectively of whether $A \rightarrow B$ or $A \leftarrow B$, while (P) has different truth values depending on the causal structure. In [Corollary 3.3](#), we show that interpreting (P) as (P2) means that every causal Bayesian network that correctly models the observational distribution is interventionaly valid (see [Definition 2.7](#)).

1.3 Contribution

We introduce a mathematical framework that explicitly links causal models and the real-world data-generating processes they are models of. This enables a transparent and formal argument showing that the seemingly natural interpretation of actions as interventions is circular, suggesting that interventional (layer 2 [Ibeling and Icard, 2020, Bareinboim et al., 2022]) predictions are not inherently free from the philosophical intricacies of falsifiability that have been debated for counterfactual (layer 3) predictions [Dawid, 2000, Shpitser

and Pearl, 2007, Raghavan and Bareinboim, 2024]. Our work thus challenges the common assumption that interventional predictions allow for falsification of causal models.² Our formalism enables us to work out interpretations that do allow for falsification of causal models via interventional predictions and to transparently discuss their pros and cons, see Section 5. This is necessary to make sense of, for example, causal discovery and causal representation learning, where it is presupposed that not all causal Bayesian networks that induce the correct observational distribution are also interventionally valid.

Our contributions also highlight that intuition and commonsense may not be the best arbiters for determining if and how causal models make falsifiable predictions about the effects of certain actions. Therefore, we present many examples throughout the paper to illustrate our theoretical results and their formal implications. Some examples may appear contrived and abstract, but this is intentional: They are designed to explain and highlight the principles and technical subtleties involved in establishing a formal relationship between causal models and real-world data-generating processes.

1.4 Article outline

Section 2 – Framework: We introduce the framework used in this article. Instead of assuming that the underlying data-generating process is a causal model, we have a generic set of distributions indexed by actions. We draw an important distinction between 1) a model emulating the distributions of a representation and 2) a model being an interventionally valid model of a representation.

Section 3 – A circular interpretation: We formalize the seemingly natural interpretation of actions as interventions and show that it is circular, rendering every CBN that correctly models the observational distribution interventionally valid.

Section 4 – An impossibility result for interpretations: We discuss different intuitive properties of interpretations of actions as interventions and show that an interpretation that satisfies four intuitive desiderata is necessarily circular.

Section 5 – Non-circular interpretations: We define and discuss five non-circular interpretations of actions as interventions, making it possible for a causal model to be falsified.

Section 6 – Implications for related research: We discuss implications of our work for causal representation learning, causal discovery, and causal abstraction. We discuss connections to the philosophical literature on the logic of conditionals.

Section 7 – Conclusion

²For example, Loftus [2024, Position 4] posits that “Causal models can be falsified in more ways than predictive models. This is usually good.”; Peters et al. [2017, Section 6.8] that “if an interventional model predicts the observational distribution correctly but does not predict what happens in a randomized experiment, the model is still considered to be falsified”; and Raghavan and Bareinboim [2024] that “It is commonly believed that, in a real-world environment, samples can only be drawn from observational and interventional distributions [... whereas sampling from] counterfactual distributions, is believed to be inaccessible almost by definition.”

2 Framework

We choose to formalize causal models as causal Bayesian networks [Spirtes et al., 2001, Pearl, 2009, Peters et al., 2017] instead of, for example, structural causal models (SCMs). This choice eases the mathematical presentation in the present manuscript, while we think that the considerations in this paper apply equally to SCMs.

Definition 2.1. Causal Bayesian network. A *causal Bayesian network* (CBN) \mathfrak{C} , also called a causal graphical model, over real-valued random variables $\mathbf{Z} = (Z_1, \dots, Z_n)$ is a directed acyclic graph (DAG) \mathcal{G}^3 over nodes $[n] = \{1, \dots, n\}^4$ and a collection of Markov kernels $\{\mathbf{pa}_i \mapsto p_i^{\mathfrak{C}}(\cdot \mid \mathbf{pa}_i) \cdot \nu_i \mid i \in [n]\}^5$. These Markov kernels induce a joint distribution, called the observational distribution, denoted by $\mathcal{L}^{\mathfrak{C}}(\mathbf{Z})$, with density

$$p_{\mathbf{Z}}^{\mathfrak{C}}(z_1, \dots, z_n) = \prod_{i=1}^n p_i^{\mathfrak{C}}(z_i \mid \mathbf{pa}_i).$$

Given some nonempty subset $J \subseteq [n]$, interventional distributions are obtained by, for each $j \in J$, replacing the kernel of j with some new kernel $\mathbf{pa}_j \mapsto q_j(\cdot \mid \mathbf{pa}_j) \cdot \mu_j$. We denote this intervention by $\text{do}(j \leftarrow q_j, j \in J)$.⁶ The interventional distribution under intervention $\text{do}(j \leftarrow q_j, j \in J)$ is denoted by $\mathcal{L}^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in J)}(\mathbf{Z})$ and has density given by

$$\begin{aligned} & p_{\mathbf{Z}}^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in J)}(\mathbf{z}) \\ &= \prod_{i=1}^n p_i^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in J)}(z_i \mid \mathbf{pa}_i) \\ &= \prod_{i \notin J} p_i^{\mathfrak{C}}(z_i \mid \mathbf{pa}_i) \prod_{i \in J} q_i(z_i \mid \mathbf{pa}_i). \end{aligned}$$

$p_i^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in J)}$ denotes the i 'th kernel given by \mathfrak{C} and intervention $d = \text{do}(j \leftarrow q_j, j \in J)$, that is, $p_i^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in J)} = p_i^{\mathfrak{C}}$ for $i \notin J$, and $p_i^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in J)} = q_i$ for $i \in J$. In this work, we do not consider interventions that change the DAG \mathcal{G} . We assume that interventions induce distributions different from the observational distribution, that is, they satisfy

$$\mathcal{L}^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in J)}(\mathbf{Z}) \neq \mathcal{L}^{\mathfrak{C}}(\mathbf{Z}).$$

This, for example, rules out intervening only on source nodes⁷ without changing at least some of their marginal distributions.

³ Throughout, we assume acyclicity to ensure that the collection of Markov kernels induces a well-defined distribution under every intervention.

⁴ The nodes of a graph \mathcal{G} are technically the natural numbers $[n]$, but to improve readability we also consider the corresponding coordinates of $\mathbf{Z} = (Z_1, \dots, Z_n)$ to be the nodes of \mathcal{G} .

⁵ By \mathbf{PA}_i we denote the variables of \mathbf{Z} whose indices corresponds to the parents of i in \mathcal{G} , and by $\mathbf{pa}_i \in \mathbb{R}^{|\mathbf{PA}_i|}$ we denote some value of these variables. The conditional distribution $\mathcal{L}^{\mathfrak{C}}(Z_i \mid \mathbf{PA}_i = \mathbf{pa}_i)$ has a density $p_i^{\mathfrak{C}}(\cdot \mid \mathbf{pa}_i)$ w.r.t. the σ -finite measure ν_i on \mathbb{R} . The measure ν_i is fixed across all \mathbf{pa}_i .

⁶ An intervention $\text{do}(j \leftarrow q_j, j \in J)$ is allowed to change both the densities $p_j(\cdot \mid \cdot)$ and the measures ν_j , even though the change of measure is suppressed in the notation. Sometimes, we write $\text{do}(Z_j = z)$ and take this to mean that the kernel of j is replaced with $\mathbf{pa}_j \mapsto \delta_z$, where δ_z is the Dirac distribution with support $\{z\}$. We also sometimes use notation like $\text{do}(Z_j \leftarrow \mathcal{N}(0, 1))$ and take this to mean that the kernel of j is replaced with $\mathbf{pa}_j \mapsto \mathcal{N}(0, 1)$, that is, Z_j is set to follow a standard normal distribution and is made independent of its parents.

⁷ A node is a *source node* if it has no parents.

We say a DAG \mathcal{G} is *complete* if for every distinct nodes Z_j and Z_i , either $Z_j \rightarrow Z_i$ or $Z_j \leftarrow Z_i$; we say that a CBN is *complete* if its DAG is complete. \circ

Definition 2.2. Single-node, multi-node, minimal, decomposable, and perfect interventions. We say that an intervention $\text{do}(j \leftarrow q_j, j \in J)$ is a *single-node intervention* if $|J| = 1$ and a *multi-node intervention* if $|J| > 1$. We say that an intervention $\text{do}(j \leftarrow q_j, j \in J)$ is *minimal* if for every nonempty proper subset $J^* \subsetneq J$ and kernels $\{q_j^*\}_{j \in J^*}$,

$$\mathcal{L}^{\mathbb{C}; \text{do}(j \leftarrow q_j, j \in J)}(\mathbf{Z}) \neq \mathcal{L}^{\mathbb{C}; \text{do}(j \leftarrow q_j^*, j \in J^*)}(\mathbf{Z}),$$

that is, if no intervention on a proper subset of the nodes induces the same interventional distribution. We say that an intervention $\text{do}(j \leftarrow q_j, j \in J)$ is *decomposable* if $\text{do}(j \leftarrow q_j, j \in J^*)$ is an intervention for every nonempty subset $J^* \subseteq J$, that is, if $\mathcal{L}^{\mathbb{C}; \text{do}(j \leftarrow q_j, j \in J^*)}(\mathbf{Z}) \neq \mathcal{L}^{\mathbb{C}}(\mathbf{Z})$ for every nonempty subset $J^* \subseteq J$. We say that a kernel $\mathbf{pa}_i \mapsto p_i^{\mathbb{C}}(\cdot | \mathbf{pa}_i) \cdot \nu_i$ is *perfect* if the measure $p_i^{\mathbb{C}}(\cdot | \mathbf{pa}_i) \cdot \nu_i$ is identical for all \mathbf{pa}_i . We say that an intervention $\text{do}(j \leftarrow q_j, j \in J)$ is *perfect* if q_j is perfect for all $j \in J$. \circ

One of the novelties of this work is that we do not assume that the data-generating process is a causal Bayesian network or a structural causal model. Instead, we remain agnostic as to how the data is generated and then ask if a representation of the data-generating process can be modeled by a causal Bayesian network.

Definition 2.3. Data-generating process. A *data-generating process* is a tuple $\mathcal{D} = (\mathcal{A}, \{\mathcal{L}^a(\mathbf{X}^*)\}_{a \in \mathcal{A}})$ where

- \mathcal{A} is a set of *actions*,
- one action $\mathcal{O} \in \mathcal{A}$ is called the *observational regime*, and
- $\mathbf{X}^* \in \mathbb{R}^m$ is a multivariate random variable that we call *low-level features* with distributions given by $\mathcal{L}^a(\mathbf{X}^*)$ for each action $a \in \mathcal{A}$. \circ

In general, we assume that observed data consists of low-level features, such as image pixels, which may not be directly suited for causal modeling (see, for example, Schölkopf et al. [2021]). Instead, we may aim to causally model a high-level representation given by functions of these pixels, rather than the individual pixels themselves.

Unlike most works on causal representation learning, we do not focus on the challenges of learning or identifying such representations. Instead, we investigate the implications of hypothesizing that a representation is modeled by a causal model.

Definition 2.4. Representation. A *representation* of a data-generating process $\mathcal{D} = (\mathcal{A}, \{\mathcal{L}^a(\mathbf{X}^*)\}_{a \in \mathcal{A}})$ is a multivariate random variable $\mathbf{Z}^* = (Z_1^*, \dots, Z_n^*) = h(\mathbf{X}^*) = (h_1(\mathbf{X}^*), \dots, h_n(\mathbf{X}^*))$ for measurable functions h_1, \dots, h_n . \circ

See Definition 6.3 for a definition of ‘causal representation’.

We will consider a correspondence between variables $\mathbf{Z} = (Z_1, \dots, Z_n)$ in a causal Bayesian network and a representation $\mathbf{Z}^* = (Z_1^*, \dots, Z_n^*)$ of a data-generating process.⁸ The observational distribution of the representation \mathbf{Z}^* is given by the push-forward measure

⁸ We mark variables whose distributions are derived from the data-generating process with a superscript $*$ and the corresponding variables in a CBN without the superscript.




Figure 1: The framework presented in this article has three main components. 1) The observed low-level features \mathbf{X}^* , 2) a representation $\mathbf{Z}^* := h(\mathbf{X}^*)$, and 3) a hypothesized causal model \mathcal{C} with variables \mathbf{Z} .

$\mathcal{L}^{\mathcal{O}}(\mathbf{Z}^*) = h(\mathcal{L}^{\mathcal{O}}(\mathbf{X}^*))$. We say that a CBN \mathcal{C} over nodes \mathbf{Z} is *compatible* with representation \mathbf{Z}^* if $\mathcal{L}^{\mathcal{C}}(\mathbf{Z}) = \mathcal{L}^{\mathcal{O}}(\mathbf{Z}^*)$, which in particular implies that \mathbf{Z} and \mathbf{Z}^* have the same dimension. In addition to the observational distribution, for each action $a \in \mathcal{A}$, we have a distribution $\mathcal{L}^a(\mathbf{Z}^*) = h(\mathcal{L}^a(\mathbf{X}^*))$. Likewise, the causal Bayesian network \mathcal{C} induces different interventional distributions $\mathcal{L}^{\mathcal{C}; \text{do}(j \leftarrow q_j, j \in J)}(\mathbf{Z})$ for different interventions $\text{do}(j \leftarrow q_j, j \in J)$. In [Definition 2.7](#), we make precise what it means for the CBN \mathcal{C} to be an interventionally valid model of a representation. See [Figure 1](#) for a visual summary of the setting considered in this work.

Throughout this work, we assume that the distribution of \mathbf{Z}^* under action $a \in \mathcal{A}$, denoted by $\mathcal{L}^a(\mathbf{Z}^*)$, has a density w.r.t. some product of σ -finite measures. We write $\mathcal{L}^a(Z_j^* | \mathbf{Y}^*) \sim p$ to imply that the kernel $\mathbf{y}^* \mapsto p(\cdot | \mathbf{y}^*) \cdot \nu$ is a regular conditional probability distribution of Z_j^* given $\mathbf{Y}^* \subseteq \mathbf{Z}^*$ under distribution $\mathcal{L}^a(\mathbf{Z}^*)$.

In [Appendix A](#), we provide a notation overview.

2.1 Emulation and interventional validity

Since CBNs are convenient to describe multiple distributions, we often use a CBN to describe the distributions of a representation.

Definition 2.5. Representation emulated by a CBN. Let a data-generating process $\mathcal{D} = (\mathcal{A}, \{\mathcal{L}^a(\mathbf{X}^*)\}_{a \in \mathcal{A}})$ and a CBN \mathcal{A} over nodes $\mathbf{Z} = (Z_1, \dots, Z_n)$ be given. We say that a representation $\mathbf{Z}^* = (Z_1^*, \dots, Z_n^*)$ of \mathcal{D} is *emulated by \mathcal{A} and interventions \mathcal{I}^** if \mathcal{I}^* is a set of interventions in \mathcal{A} and there is a surjective function $g : \mathcal{A} \setminus \{\mathcal{O}\} \rightarrow \mathcal{I}^*$ such that

1. $\mathcal{L}^{\mathcal{O}}(\mathbf{Z}^*) = \mathcal{L}^{\mathcal{A}}(\mathbf{Z})$, and

2. $\mathcal{L}^a(\mathbf{Z}^*) = \mathcal{L}^{\mathcal{A};g(a)}(\mathbf{Z})$ for all $a \in \mathcal{A} \setminus \{\mathcal{O}\}$.

We refer to g as the *link*. ◦

Importantly, stating that a representation is emulated by a CBN \mathcal{A} and interventions \mathcal{I}^* implies no other claims about the data-generating process than the distributions it induces for the set of actions. Moreover, for every data-generating process $\mathcal{D} = (\mathcal{A}, \{\mathcal{L}^a(\mathbf{X}^*)\}_{a \in \mathcal{A}})$ and representation \mathbf{Z}^* , there exists a CBN \mathcal{A} and a set of interventions \mathcal{I}^* in \mathcal{A} such that \mathbf{Z}^* is emulated by \mathcal{A} and \mathcal{I}^* . To see this, pick a complete CBN \mathcal{A} such that $\mathcal{L}^{\mathcal{A}}(\mathbf{Z}) = \mathcal{L}^{\mathcal{O}}(\mathbf{Z}^*)$, and for each $a \in \mathcal{A} \setminus \{\mathcal{O}\}$, let d^a be an intervention such that $\mathcal{L}^a(Z_i^* | \mathbf{PA}_i^*) \sim p_i^{\mathcal{A};d^a}$ for all $i \in [n]$ (this is possible since $\mathcal{L}^a(\mathbf{Z}^*)$ has density w.r.t. a product of σ -finite measures). Now $\mathcal{L}^a(\mathbf{Z}^*) = \mathcal{L}^{\mathcal{A};d^a}(\mathbf{Z})$ for every $a \in \mathcal{A} \setminus \{\mathcal{O}\}$ since \mathcal{A} is complete, so \mathbf{Z}^* is emulated by \mathcal{A} and $\mathcal{I}^* = \{d^a \mid a \in \mathcal{A} \setminus \{\mathcal{O}\}\}$. Thus, while we may imagine that the data-generating process can be any physical mechanism, every representation of such a mechanism can be emulated by a CBN (and is in fact emulated by several different CBNs if there is more than one node).⁹ We rely on [Definition 2.5](#) to simplify our presentation (avoiding other mathematical descriptions of data-generating processes in terms of, for example, stochastic differential equations or exhaustive enumerations of the distributions for each action).

The variables in causal discovery and the latent variables in causal representation learning are high-level features and it may be unclear what constitutes an intervention on those variables. In particular, interventions are usually ambiguous [[Spirtes and Scheines, 2004](#), [Rubenstein et al., 2017](#)]. Consider performing a phacoemulsification cataract surgery. Assume that we have a high-level feature that indicates whether the surgery was performed. Interventions on this variable do not correspond to unique physical processes. The surgery may be performed at different times of the day, in different locations, by various different doctors using slightly different types of equipment, etc.

In particular, if we are given a representation \mathbf{Z}^* and a CBN compatible with variables \mathbf{Z} , whether or not the model correctly predicts the effects of actions depends on which actions in the data-generating process correspond to which interventions in the CBN. Interpretations of actions formalize this correspondence by assigning subsets of the modeled interventions \mathcal{I} to each action $a \in \mathcal{A}$. This allows us to discuss the implications of different interpretations.

Definition 2.6. Interpretations of actions. Let a data-generating process \mathcal{D} , representation \mathbf{Z}^* , and a compatible CBN \mathcal{C} be given. An *interpretation* \mathbf{Int} is a mapping that takes an action $a \in \mathcal{A}$ and a set of modeled interventions \mathcal{I} in \mathcal{C} as input and outputs a subset of \mathcal{I} . For a given set of modeled interventions \mathcal{I} , an interpretation induces a function $\mathbf{Int}^{\mathcal{I}} : \mathcal{A} \rightarrow \mathcal{P}(\mathcal{I})$, where $\mathcal{P}(\mathcal{I})$ is the power set of \mathcal{I} . ◦

Notice that we do not necessarily map every action to an intervention, that is, $\mathbf{Int}^{\mathcal{I}}(a)$ can be the empty set. Likewise, for an intervention in $d \in \mathcal{I}$, there may be no action $a \in \mathcal{A}$ such that $d \in \mathbf{Int}^{\mathcal{I}}(a)$.

Without committing to a specific interpretation, it is unclear whether and how a CBN correctly predicts the distribution of a data-generating process’s representation for a given action. Consequently, it is then unclear what is causal about a causal model and

⁹ This argument generalizes one of the main points made in [Eberhardt \[2016\]](#), namely that the same system can seemingly be described equally well by different causal models.

what the criteria for falsifying it as a valid model of a representation should be. Once given an interpretation \mathbf{Int} , we can ask whether a CBN makes a correct prediction about the distribution of \mathbf{Z}^* under a given action. In particular, if we interpret action a as intervention d , that is, if $d \in \mathbf{Int}^{\mathcal{I}}(a)$, we can ask if $\mathcal{L}^a(\mathbf{Z}^*) = \mathcal{L}^{\mathfrak{C};d}(\mathbf{Z})$.

Definition 2.7. Interventional validity. Let a data-generating process \mathcal{D} , representation \mathbf{Z}^* , compatible CBN \mathfrak{C} , set of interventions \mathcal{I} in \mathfrak{C} , and an interpretation \mathbf{Int} be given. If for all $a \in \mathcal{A}$ and $d \in \mathcal{I}$

$$d \in \mathbf{Int}^{\mathcal{I}}(a) \Rightarrow \mathcal{L}^a(\mathbf{Z}^*) = \mathcal{L}^{\mathfrak{C};d}(\mathbf{Z}),$$

we say that \mathfrak{C} is an $\mathcal{I} - \mathbf{Int}$ valid model of \mathbf{Z}^* . ◦

3 \mathbf{Int}_C : The seemingly natural interpretation of actions as interventions is circular

In a CBN, an intervention $\text{do}(j \leftarrow q_j, j \in J)$ modifies the kernels of Z_j given \mathbf{PA}_j for $j \in J$, while keeping the kernels fixed for $j \notin J$. Since we do not consider interventions that change the graph, we have, for all interventions d , that $\mathcal{L}^{\mathfrak{C};d}(\mathbf{Z})$ is Markov w.r.t. the DAG \mathcal{G} of the CBN \mathfrak{C} . These considerations might compel us to consider the following interpretation.

Definition 3.1. \mathbf{Int}_C . The seemingly natural interpretation. Let a data-generating process \mathcal{D} , representation \mathbf{Z}^* , compatible CBN \mathfrak{C} , and set of interventions \mathcal{I} in \mathfrak{C} be given. We define interpretation \mathbf{Int}_C by the following rule: An intervention $\text{do}(j \leftarrow q_j, j \in J) \in \mathcal{I}$ is in $\mathbf{Int}_C^{\mathcal{I}}(a)$ if and only if the following 3 conditions hold:

- 1) $\mathcal{L}^a(Z_i^* \mid \mathbf{PA}_i^*) \sim q_i$ for all $i \in J$. That is, the action sets the conditionals of intervened nodes correctly. For example, if we interpret an action as intervention $\text{do}(Z_i = 4)$, then Z_i^* must have Dirac distribution with support $\{4\}$ under that action.
- 2) $\mathcal{L}^a(Z_i^* \mid \mathbf{PA}_i^*) \sim p_i^{\mathfrak{C}}$ for all $i \notin J$. Intuitively, we do not intervene on nodes not in J .
- 3) $\mathcal{L}^a(\mathbf{Z}^*)$ is Markov w.r.t. the DAG of \mathfrak{C} . That is, we do not introduce dependencies. ◦

The C in \mathbf{Int}_C is for ‘circular’. This interpretation interprets an action as an intervention d in \mathfrak{C} if and only if the action induces the interventional distribution given by \mathfrak{C} and d .

Proposition 3.2. Let a data-generating process \mathcal{D} , representation \mathbf{Z}^* , compatible CBN \mathfrak{C} , and set of interventions \mathcal{I} in \mathfrak{C} be given. Then $d \in \mathcal{I}$ is in $\mathbf{Int}_C^{\mathcal{I}}(a)$ if and only if $\mathcal{L}^a(\mathbf{Z}^*) = \mathcal{L}^{\mathfrak{C};d}(\mathbf{Z})$.

Proof. Let $d \in \mathcal{I}$ and $a \in \mathcal{A}$ be given. If $\mathcal{L}^a(\mathbf{Z}^*) = \mathcal{L}^{\mathfrak{C};d}(\mathbf{Z})$, this immediately implies 1)–3) in [Definition 3.1](#) and hence that $d \in \mathbf{Int}_C^{\mathcal{I}}(a)$.

Assume that $d = \text{do}(j \leftarrow q_j, j \in J) \in \mathbf{Int}_C^{\mathcal{I}}(a)$. Since $\mathcal{L}^a(\mathbf{Z}^*)$ is Markov w.r.t. the DAG of \mathfrak{C} , there exists some CBN \mathfrak{A} that has the same DAG as \mathfrak{C} such that $\mathcal{L}^a(\mathbf{Z}^*) = \mathcal{L}^{\mathfrak{A}}(\mathbf{Z})$.

We then have that

$$\begin{aligned}\mathcal{L}^a(\mathbf{Z}^*) &= \mathcal{L}^{\mathcal{A}}(\mathbf{Z}) \\ &= \mathcal{L}^{\mathcal{A}; \text{do}(j \leftarrow q_j, j \in J; j \leftarrow p_j^{\mathcal{C}}, j \notin J)}(\mathbf{Z}) \\ &= \mathcal{L}^{\mathcal{C}; \text{do}(j \leftarrow q_j, j \in J)}(\mathbf{Z}),\end{aligned}$$

where the second equality follows by using 1) and 2) of [Definition 3.1](#). \square

The following is a formalization of a circularity that has been hinted at in previous works [[Baumgartner, 2009](#), [Woodward, 2023](#), [Janzing and Meija, 2024](#)].¹⁰

Corollary 3.3. Int_C is circular. Let a data-generating process \mathcal{D} and representation \mathbf{Z}^* be given. Then every compatible CBN \mathcal{C} is an $\mathcal{I} - \text{Int}_C$ valid model of \mathbf{Z}^* for every set of interventions \mathcal{I} in \mathcal{C} . \circ

[Corollary 3.3](#) highlights the need for alternative interpretations of which actions constitute interventions since otherwise there is nothing ‘causal’ about a causal Bayesian network: Observational and interventional validity are equivalent under interpretation Int_C and interventional predictions do not help with falsification (in contrast to common assumptions, see also [Footnote 2](#)). If we dropped condition 3) of [Definition 3.1](#), then every compatible CBN with a complete DAG would still be interventionally valid.

4 Impossibility result for non-circular interpretations

We now present five desiderata **D0–D4** for interpretations of actions as interventions. Since each desideratum appears intuitively reasonable, one might expect that a reasonable interpretation should satisfy all of them. We show in [Proposition 4.1](#) that if an interpretation satisfies **D1–D4**, then it is the circular interpretation Int_C ([Definition 3.1](#)), which renders all compatible models interventionally valid.

Desideratum D0: Correct conditionals on intervened nodes. If we interpret action a as an intervention $\text{do}(j \leftarrow q_j, j \in J)$, then that action must set the conditional distribution of intervened nodes given their parents correctly. Formally, an interpretation Int satisfies desideratum **D0** if

- For every set of modeled interventions \mathcal{I} and every action $a \in \mathcal{A}$,
if $\text{do}(j \leftarrow q_j, j \in J) \in \text{Int}^{\mathcal{I}}(a)$, then $\mathcal{L}^a(\mathbf{Z}_i^* \mid \mathbf{PA}_i^*) \sim q_i$ for all $i \in J$.

We believe that any reasonable interpretation satisfies **D0** and therefore do not consider interpretations that may violate **D0**. In the context of hard interventions, **D0** is sometimes referred to as ‘effectiveness’ [[Galles and Pearl, 1998](#), [Bareinboim et al., 2022](#), [Ibeling and Icard, 2023](#)]. Effectiveness is similarly considered an axiom in [Park et al. \[2023\]](#).

¹⁰ For example: “If, as Pearl apparently intends, we understand this [the notion of intervention] to include the requirement that an intervention on X_i must leave intact the causal mechanism if any, that connects X_i to its possible effects Y , then an obvious worry about circularity arises [...]” [[Woodward, 2023](#)].

Desideratum D1: If it behaves like an intervention, it is that intervention.

If an action a induces a distribution that equals a distribution induced by the model under an intervention in the intervention set, then we should interpret that action as that intervention. Formally, an interpretation \mathbf{Int} satisfies desideratum **D1** if

- For every set of modeled interventions \mathcal{I} , every action $a \in \mathcal{A}$, and every intervention $d \in \mathcal{I}$, if $\mathcal{L}^a(\mathbf{Z}^*) = \mathcal{L}^{\mathbb{C};d}(\mathbf{Z})$, then $d \in \mathbf{Int}^{\mathcal{I}}(a)$.¹¹

Desideratum D2: An action should not be interpreted as distinct interventions.

If the action a is interpreted as two distinct interventions, then these two interventions should induce the same interventional distribution. Formally, an interpretation \mathbf{Int} satisfies desideratum **D2** if

- For every set of modeled interventions \mathcal{I} , every action $a \in \mathcal{A}$, and every interventions $b, d \in \mathcal{I}$, if $d \in \mathbf{Int}^{\mathcal{I}}(a)$ and $b \in \mathbf{Int}^{\mathcal{I}}(a)$, then $\mathcal{L}^{\mathbb{C};d}(\mathbf{Z}) = \mathcal{L}^{\mathbb{C};b}(\mathbf{Z})$.

Desideratum D3: Interpretations should not depend on the intervention set \mathcal{I} .

Whether we interpret an action a as an intervention $d \in \mathcal{I}$ should not depend on which other interventions are in \mathcal{I} . Formally, an interpretation \mathbf{Int} satisfies desideratum **D3** if

- For every sets of modeled interventions \mathcal{I} and \mathcal{I}' , every action $a \in \mathcal{A}$, and every intervention $d \in \mathcal{I} \cap \mathcal{I}'$, $d \in \mathbf{Int}^{\mathcal{I}}(a) \Leftrightarrow d \in \mathbf{Int}^{\mathcal{I}'}(a)$.

Desideratum D4: An intervention does not create new dependencies. If an action a does not induce a distribution that is Markov w.r.t. the DAG, then we should not interpret a as an intervention (in this work, as is common, we only consider interventions that do not introduce dependencies between variables). Formally, an interpretation \mathbf{Int} satisfies desideratum **D4** if

- For every set of modeled interventions \mathcal{I} and every action $a \in \mathcal{A}$, if $\mathcal{L}^a(\mathbf{Z}^*)$ is not Markov w.r.t. \mathcal{G} , then $\mathbf{Int}^{\mathcal{I}}(a) = \emptyset$.

Proposition 4.1. Impossibility result. Let a data-generating process \mathcal{D} , representation \mathbf{Z}^* , and a compatible CBN \mathbb{C} be given. Let \mathbf{Int} be an interpretation that satisfies desiderata **D1–D4**. Then, for every set of modeled interventions \mathcal{I} in \mathbb{C} and for all actions $a \in \mathcal{A}$, $\mathbf{Int}^{\mathcal{I}}(a) = \mathbf{Int}_{\mathbb{C}}^{\mathcal{I}}(a)$.

Proof. Let $a \in \mathcal{A}$ and \mathcal{I} be given. From [Proposition 3.2](#) and **D1**, it follows that $\mathbf{Int}_{\mathbb{C}}^{\mathcal{I}}(a) \subseteq \mathbf{Int}^{\mathcal{I}}(a)$. Assume that $d \in \mathbf{Int}^{\mathcal{I}}(a)$. By **D4** we have that $\mathcal{L}^a(\mathbf{Z}^*)$ is Markov w.r.t. the DAG of \mathbb{C} . Since $\mathcal{L}^a(\mathbf{Z}^*)$ is Markov w.r.t. the DAG of \mathbb{C} , we can find an intervention b such that $\mathcal{L}^{\mathbb{C};b}(\mathbf{Z}) = \mathcal{L}^a(\mathbf{Z}^*)$, namely an intervention b such that $\mathcal{L}^a(\mathbf{Z}_i^* \mid \mathbf{PA}_i^*) \sim p_i^{\mathbb{C};b}$ for all $i \in [n]$. Consider $\tilde{\mathcal{I}} = \mathcal{I} \cup \{b\}$. By condition **D1**, $b \in \mathbf{Int}^{\tilde{\mathcal{I}}}(a)$, and by condition **D3**,

¹¹ If an interpretation \mathbf{Int} additionally satisfies the reverse implication of **D1**, such that “For every set of modeled interventions \mathcal{I} , actions $a \in \mathcal{A}$, and $d \in \mathcal{I}$, $\mathcal{L}^a(\mathbf{Z}^*) = \mathcal{L}^{\mathbb{C};d}(\mathbf{Z})$ if and only if $d \in \mathbf{Int}^{\mathcal{I}}(a)$ ”, then $\mathbf{Int}^{\mathcal{I}}(a) = \mathbf{Int}_{\mathbb{C}}^{\mathcal{I}}(a)$ for every action $a \in \mathcal{A}$ and every set of interventions \mathcal{I} , rendering any compatible CBN $\mathcal{I} - \mathbf{Int}$ valid for every set of interventions \mathcal{I} in that CBN.

$d \in \mathbf{Int}^{\tilde{\mathcal{I}}}(a)$. By condition **D2**, this implies that $\mathcal{L}^{\mathbb{C};d}(\mathcal{Z}) = \mathcal{L}^{\mathbb{C};b}(\mathcal{Z})$. [Proposition 3.2](#) then gives us that $d \in \mathbf{Int}_C^{\mathcal{I}}(a)$ since $\mathcal{L}^a(\mathcal{Z}^*) = \mathcal{L}^{\mathbb{C};d}(\mathcal{Z})$. \square

\mathbf{Int}_C satisfies **D0**, so desiderata **D1–D4** together imply **D0**.

5 Non-circular interpretations

We now consider, in turn, possible interpretations that may violate either one of the desiderata **D1** and **D2** to avoid the circularity of [Corollary 3.3](#) implied by satisfying all desiderata. In [Appendix D.1](#) and [Appendix D.2](#), we consider interpretations that may violate **D3** and **D4**, respectively. Taken together, this shows that no proper subset of the desiderata **D1–D4** implies any other of the desiderata **D1–D4**, so the desiderata can be considered separately. In [Section 5.3](#), we consider an interpretation that takes action complexity into account. See [Table 1](#) for an overview of all interpretations considered in this paper.

Table 1: Overview of interpretations presented in this work.

Interpretation	Definition: Interpret action a as intervention d if..	Violated desiderata	Can falsify a model?
\mathbf{Int}_C Definition 3.1 (Circular)	<ul style="list-style-type: none"> .. a sets the conditionals of intervened nodes correctly, \wedge a changes no other conditionals, \wedge a does not introduce dependencies. 	None.	No. (\mathbf{Int}_C is circular, Corollary 3.3 .)
\mathbf{Int}_P Definition 5.1 (Perfect)	<ul style="list-style-type: none"> .. a sets the conditionals of intervened nodes correctly, \wedge non-intervened nodes not independent of parents, \wedge a does not introduce dependencies. 	D1 (If it behaves like an intervention, it is that intervention.)	Yes. (Falsified if an action behaves like an imperfect intervention, Proposition 5.2 .)
\mathbf{Int}_S Definition 5.4 (Single-node)	<ul style="list-style-type: none"> .. a sets the conditionals of intervened nodes correctly, \wedge a changes the distribution of intervened nodes \wedge a does not introduce dependencies. 	D2 (An action should not be interpreted as distinct interventions.)	Yes. (Falsified is an action behaves like a multi-node intervention, Proposition 5.5 .)
$\mathbf{Int}_{\tilde{\mathcal{I}},f}$ Definition D.1 (f -least in $\tilde{\mathcal{I}}$)	<ul style="list-style-type: none"> .. $d \in \mathbf{Int}_C^{\mathcal{I}}(a)$, \vee d is least element in $\tilde{\mathcal{I}} \cap \mathbf{Int}_S^{\mathcal{I}}(a)$, in the strict total order on $\tilde{\mathcal{I}}$ induced by f. 	D3 (Interpretations should not depend on the intervention set \mathcal{I} .)	Yes. (See Example D.2 .)
\mathbf{Int}_M Definition D.3 (Markov)	<ul style="list-style-type: none"> .. a sets the conditionals of intervened nodes correctly, \wedge a changes no other conditionals. 	D4 (An intervention does not create new dependencies.)	Yes, but can only falsify CBNs with non-complete DAGs. (Falsified if an action introduces dependencies between variables, Proposition D.4 .)
\mathbf{Int}_K Definition 5.7 (Complexity K)	.. $a \in \arg \min_{a \in \mathcal{A}: d \in \mathbf{Int}_S^{\mathcal{I}}(a)} K(a)$.	D1 and D2 .	Yes. (See Example 5.9 .)

5.1 \mathbf{Int}_P : Letting imperfect interventions falsify a model is one way out of circularity

Informal overview of Section 5.1. If we insist that all actions correspond to perfect interventions, then it becomes possible to falsify a causal model: If we perform an action and the resulting observation cannot be explained by a perfect intervention, the causal model can be rejected. Explaining these observations by imperfect soft interventions may be a slippery slope leading to a circular interpretation where every observed distribution can be explained by some complex intervention in the model. In [Example 5.3](#), we show how to falsify a causal model under \mathbf{Int}_P (to be defined in [Definition 5.1](#)).

Consider the following interpretation that may violate **D1**, but satisfies **D0** and **D2–D4**.

Definition 5.1. \mathbf{Int}_P . An interpretation violating only D1. Let a data-generating process \mathcal{D} , representation \mathbf{Z}^* , compatible CBN \mathfrak{C} , and set of interventions \mathcal{I} in \mathfrak{C} be given. We define the interpretation \mathbf{Int}_A by the following rule: An intervention $d = \text{do}(j \leftarrow q_j, j \in J) \in \mathcal{I}$ is in $\mathbf{Int}_P^{\mathcal{I}}(a)$ if and only if the following four conditions hold:

- 1) d is a perfect intervention.
- 2) For all $i \in J$,

$$\mathcal{L}^a(Z_i^* \mid \mathbf{PA}_i^*) \sim q_i.$$

That is, \mathbf{Int}_P satisfies **D0** (correct conditionals on intervened nodes).

- 3) For all $i \notin J$,

$$\begin{aligned} &\mathbf{PA}_i^* \text{ is empty and } \mathcal{L}^a(Z_i^*) = \mathcal{L}^{\mathfrak{C}}(Z_i), \text{ or} \\ &\mathbf{PA}_i^* \text{ is nonempty and } Z_i^* \not\perp\!\!\!\perp \mathbf{PA}_i^* \text{ in } \mathcal{L}^a(\mathbf{Z}^*). \end{aligned}$$

That is, nodes not intervened on are either source nodes with unchanged distributions or not independent of their parents.

- 4) $\mathcal{L}^a(\mathbf{Z}^*)$ is Markov w.r.t. the DAG of \mathfrak{C} . That is, \mathbf{Int}_P satisfies **D4** (an intervention does not create new dependencies).¹²

◦

The P in \mathbf{Int}_P is for ‘perfect’. Condition 3) ensures that \mathbf{Int}_P satisfies **D2** (an action should not be interpreted as distinct interventions), see [Appendix B](#). It is straightforward to verify that \mathbf{Int}_P satisfies **D3** (interpretations should not depend on the intervention set \mathcal{I}). \mathbf{Int}_P may violate **D1** (if it behaves like an intervention, it is that intervention) because it may be that $\mathcal{L}^a(\mathbf{Z}^*) = \mathcal{L}^{\mathfrak{C};d}(\mathbf{Z})$ for some $d \in \mathcal{I}$ that is not a perfect intervention and some $a \in \mathcal{A}$, and thus $d \notin \mathbf{Int}_P^{\mathcal{I}}(a)$. Under a non-circular interpretation like \mathbf{Int}_P , a CBN \mathfrak{C} can be an invalid model of a representation even though the representation is emulated by \mathfrak{C} . We now provide a partial characterization for when this happens under interpretation \mathbf{Int}_P .

Proposition 5.2. Let a data-generating process \mathcal{D} be given. Assume that \mathbf{Z}^* is emulated by CBN \mathfrak{C} and interventions \mathcal{I}^* .

¹²[Proposition 5.2](#) would still hold if we omitted condition 4) of [Definition 5.1](#).

(1) If \mathcal{I}^* only contains perfect interventions, then \mathfrak{C} is $\mathcal{I} - \mathbf{Int}_P$ valid model of \mathbf{Z}^* for every set of interventions \mathcal{I} in \mathfrak{C} .

(2) On the other hand, if \mathcal{I}^* contains a minimal and decomposable intervention $\text{do}(j \leftarrow q_j, j \in J)$ for which there exist $s, t \in J$ such that $Z_s \perp\!\!\!\perp \mathbf{PA}_s$ and $Z_t \not\perp\!\!\!\perp \mathbf{PA}_t$ in $\mathcal{L}^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in J)}(\mathbf{Z})$, then there exists a set of interventions \mathcal{I} in \mathfrak{C} such that \mathfrak{C} is not an $\mathcal{I} - \mathbf{Int}_P$ valid model of \mathbf{Z}^* .

Proof. (1) Assume that \mathcal{I}^* contains only perfect interventions. Consider some $a \in \mathcal{A}$ and some intervention $d = \text{do}(j \leftarrow q_j, j \in J) \in \mathbf{Int}_P^{\mathcal{I}}(a)$. We need to show that $\mathcal{L}^{\mathfrak{C}; d}(\mathbf{Z}) = \mathcal{L}^a(\mathbf{Z}^*)$. By assumption, \mathbf{Z}^* is emulated by \mathfrak{C} and \mathcal{I}^* , so there is a perfect intervention $d^* = g(a) = \text{do}(j \leftarrow \tilde{q}_j, j \in \tilde{J}) \in \mathcal{I}^*$ such that $\mathcal{L}^a(\mathbf{Z}^*) = \mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z})$. We have

$$\begin{aligned} \mathcal{L}^a(\mathbf{Z}^*) &= \mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z}) \\ &= \mathcal{L}^{\mathfrak{C}; \text{do}(j \leftarrow \tilde{q}_j, j \in \tilde{J})}(\mathbf{Z}) \\ &= \mathcal{L}^{\mathfrak{C}; \text{do}(j \leftarrow \tilde{q}_j, j \in \tilde{J} \setminus J; j \leftarrow q_j, j \in J)}(\mathbf{Z}) \\ &= \mathcal{L}^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in J)}(\mathbf{Z}) \\ &= \mathcal{L}^{\mathfrak{C}; d}(\mathbf{Z}) \end{aligned}$$

The third equality follows from condition 2) of [Definition 5.1](#). The fourth equality follows from condition 3) and the fact that d^* is perfect: For every node $i \notin J$, either (1) \mathbf{PA}_i is empty and $\mathcal{L}^{\mathfrak{C}; d^*}(Z_i) = \mathcal{L}^{\mathfrak{C}}(Z_i)$, or (2) \mathbf{PA}_i is nonempty and $Z_i \perp\!\!\!\perp \mathbf{PA}_i$ in $\mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z})$, which implies that $i \notin \tilde{J}$ since d^* is perfect.

(2) We want to find an action a , a set of interventions \mathcal{I} , and an intervention $d \in \mathbf{Int}_P^{\mathcal{I}}(a)$ such that $\mathcal{L}^a(\mathbf{Z}^*) \neq \mathcal{L}^{\mathfrak{C}; d}(\mathbf{Z})$. By assumption, \mathcal{I}^* contains a minimal and decomposable intervention $d^* = \text{do}(j \leftarrow \tilde{q}_j, j \in \tilde{J})$ for which there exist $s, t \in \tilde{J}$ such that $Z_s \perp\!\!\!\perp \mathbf{PA}_s$ and $Z_t \not\perp\!\!\!\perp \mathbf{PA}_t$ in $\mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z})$. Let $\tilde{J} := \{j \in \tilde{J} \mid Z_j \perp\!\!\!\perp \mathbf{PA}_j \text{ in } \mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z})\} \subseteq \tilde{J} \setminus \{t\} \subsetneq \tilde{J}$, and let $C := \{j \in [n] \setminus \tilde{J} \mid Z_j \perp\!\!\!\perp \mathbf{PA}_j \text{ in } \mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z}) \text{ and } \mathbf{PA}_j \neq \emptyset\}$. Let $\text{do}(j \leftarrow q_j, j \in \tilde{J} \cup C)$ be a perfect intervention such that $\mathcal{L}^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in \tilde{J} \cup C)}(Z_i) = \mathcal{L}^{\mathfrak{C}; d^*}(Z_i)$ for all $i \in \tilde{J} \cup C$. For every $i \in \tilde{J} \cup C$, $\mathcal{L}^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in \tilde{J} \cup C)}(Z_i) = \mathcal{L}^{\mathfrak{C}; d^*}(Z_i)$ implies that $\mathcal{L}^{\mathfrak{C}; d^*}(Z_i \mid \mathbf{PA}_i) \sim p_i^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in \tilde{J} \cup C)}$ since $Z_i \perp\!\!\!\perp \mathbf{PA}_i$ in $\mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z})$ and $p_i^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in \tilde{J} \cup C)}$ is perfect. Let the set of modeled interventions consist of this intervention, that is, $\mathcal{I} = \{\text{do}(j \leftarrow q_j, j \in \tilde{J} \cup C)\}$. We have that $\text{do}(j \leftarrow q_j, j \in \tilde{J} \cup C) \in \mathbf{Int}_P^{\mathcal{I}}(a)$ since 1) it is perfect, 2) sets the conditionals of intervened nodes correctly, 3) every node i not in $\tilde{J} \cup C$ is either a source node not in \tilde{J} , or $Z_i \perp\!\!\!\perp \mathbf{PA}_i$ in $\mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z})$,¹³ and condition 4) of [Definition 5.1](#) trivially holds.

If $\mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z}) = \mathcal{L}^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in \tilde{J} \cup C)}(\mathbf{Z})$, then $\mathcal{L}^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in \tilde{J} \cup C)}(Z_i \mid \mathbf{PA}_i) \sim p_i^{\mathfrak{C}}$ for all $i \in C$ since nodes in C are not intervened upon by d^* . Therefore, if $\mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z}) = \mathcal{L}^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in \tilde{J} \cup C)}(\mathbf{Z})$, we would also have that $\mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z}) = \mathcal{L}^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in \tilde{J})}(\mathbf{Z})$, but this is a contradiction since $\tilde{J} \subsetneq \tilde{J} \cup C$ and d^* is minimal. Therefore, we can conclude that $\mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z}) \neq \mathcal{L}^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in \tilde{J} \cup C)}(\mathbf{Z})$ and that \mathfrak{C} is not an $\mathcal{I} - \mathbf{Int}_P$ valid model of \mathbf{Z}^* . \square

¹³ Let i be a node not in $\tilde{J} \cup C$. Since i is not in \tilde{J} , either $Z_i \not\perp\!\!\!\perp \mathbf{PA}_i$ in $\mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z})$ or i is not in \tilde{J} . If $Z_i \perp\!\!\!\perp \mathbf{PA}_i$ in $\mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z})$ and i is not in \tilde{J} , i must be a source node, otherwise i would be in C .




Figure 2: In [Example 5.3](#), we assume that the low-level representation (LDL^*, HDL^*, HD^*) is emulated by a CBN \mathcal{A} , with graph given in [Figure 2a](#), and single-node interventions in \mathcal{A} . Under \mathbf{Int}_P , we falsify the $(TC^*, HD^*) := (LDL^* + HDL^*, HD^*)$ -compatible CBN \mathcal{C} with graph given in [Figure 2b](#).

If there exists an action that induces a distribution like an intervention in [Proposition 5.2](#) (2), that is, a perfect intervention on some nodes and an imperfect intervention on others, then a model can be falsified. Since for a given representation Z^* , it may be difficult to rule out such actions, [Proposition 5.2](#) (2) highlights a limitation of causal modeling using \mathbf{Int}_P .

We now show an example where we falsify a CBN under interpretation \mathbf{Int}_P .

Example 5.3. Falsifying a total cholesterol model under interpretation \mathbf{Int}_P . We consider the causal effect of total cholesterol on heart disease [[Spirtes and Scheines, 2004](#)]. Throughout this example, we use the shorthands LDL for low-density lipoprotein, HDL for high-density lipoprotein, HD for heart disease, and TC for total cholesterol. Let \mathcal{A} be a CBN given by the graph depicted in [Figure 2a](#) and kernels

$$\begin{aligned}\mathcal{L}^{\mathcal{A}}(\text{LDL}) &= \mathcal{N}(0, 1) \\ \mathcal{L}^{\mathcal{A}}(\text{HDL}) &= \mathcal{N}(0, 1) \\ \mathcal{L}^{\mathcal{A}}(\text{HD} \mid \text{LDL} = x, \text{HDL} = y) &= \mathcal{N}(2x - y, 1).\end{aligned}$$

Assume that $X^* = (\text{LDL}^*, \text{HDL}^*, \text{HD}^*)$ is emulated by \mathcal{A} and perfect interventions $\mathcal{I}^* = \{\text{do}(\text{LDL} \leftarrow \mathcal{N}(y, 1), \text{HDL} \leftarrow \mathcal{N}(x, 1)) \mid x, y \in \mathbb{R}\}$. Consider the representation given by $Z^* = (\text{TC}^*, \text{HD}^*) = (\text{LDL}^* + \text{HDL}^*, \text{HD}^*)$. Let \mathcal{C} be the Z^* -compatible CBN with graph given in [Figure 2b](#) and kernels given by

$$\begin{aligned}\mathcal{L}^{\mathcal{C}}(\text{TC}) &= \mathcal{N}(0, 2) \\ \mathcal{L}^{\mathcal{C}}(\text{HD} \mid \text{TC} = t) &= \mathcal{N}\left(\frac{t}{2}, \frac{11}{2}\right).\end{aligned}$$

We consider perfect shift interventions that change the mean of total cholesterol $\mathcal{I} = \{\text{do}(\text{TC} \leftarrow \mathcal{N}(t, 2)) \mid t \in \mathbb{R}\}$. \mathcal{C} is not an $\mathcal{I} - \mathbf{Int}_P$ valid model of Z^* . To see this, consider an action a' with $\mathcal{L}^{a'}(\text{TC}^*, \text{HD}^*) = \mathcal{L}^{\mathcal{A}; \text{do}(\text{LDL} \leftarrow \mathcal{N}(1, 1), \text{HDL} \leftarrow \mathcal{N}(0, 1))}(\text{LDL} + \text{HDL}, \text{HD})$ (such an action exists by definition of \mathcal{I}^*). We have that $\text{do}(\text{TC} \leftarrow \mathcal{N}(1, 2)) \in \mathbf{Int}_P^{\mathcal{I}}(a')$ since $\mathcal{L}^{a'}(\text{TC}^*) = \mathcal{N}(1, 2)$ and $\text{HD}^* \not\perp\!\!\!\perp \text{TC}^*$ in $\mathcal{L}^{a'}(\text{TC}^*, \text{HD}^*)$. But $\mathcal{L}^{a'}(\text{TC}^*, \text{HD}^*) \neq \mathcal{L}^{\mathcal{C}; \text{do}(\text{TC} \leftarrow \mathcal{N}(1, 2))}(\text{TC}, \text{HD})$, for example, because the expected value $\mathbb{E}^{a'}(\text{HD}^*) = 2$ is not

equal to $\mathbb{E}^{\mathfrak{C}; \text{do}(\text{TC} \leftarrow \mathcal{N}(1,2))}(\text{HD}) = \frac{1}{2}$. This falsifies CBN \mathfrak{C} as an $\mathcal{I} - \mathbf{Int}_P$ valid model of $(\text{TC}^*, \text{HD}^*)$, and a' can be thought of as the falsifying experiment. Note that to falsify \mathfrak{C} , one need not have any knowledge about how total cholesterol is constituted of low-density lipoprotein and high-density lipoprotein. The model is falsified purely based on the distribution $(\text{TC}^*, \text{HD}^*)$ under action a' .

Under \mathbf{Int}_C : On the other hand, we cannot falsify \mathfrak{C} as an $\mathcal{I} - \mathbf{Int}_C$ valid model of $(\text{TC}^*, \text{HD}^*)$. $\text{do}(\text{TC} \leftarrow \mathcal{N}(1,2)) \notin \mathbf{Int}_C^{\mathcal{I}}(a')$ since it is not the case that $\mathcal{L}^{a'}(\text{HD}^* \mid \text{TC}^*) \sim p_{\text{HD}^* \mid \text{TC}^*}^{\mathfrak{C}; \text{do}(\text{TC} \leftarrow \mathcal{N}(1,2))}$. Under action a' , the conditional distribution of heart disease given total cholesterol is different than in the observational regime. Therefore, under interpretation \mathbf{Int}_C , we could only interpret a' as an imperfect multi-node intervention on total cholesterol and heart disease. Indeed, \mathfrak{C} is an $\mathcal{I} - \mathbf{Int}_C$ valid model of \mathbf{Z}^* , where $\text{do}(\text{TC} \leftarrow \mathcal{N}(t,2)) \in \mathbf{Int}_C^{\mathcal{I}}(a)$ if $\mathcal{L}^a(\text{TC}^*, \text{HD}^*) = \mathcal{L}^{\mathcal{A}; \text{do}(\text{LDL} \leftarrow \mathcal{N}(\frac{1}{2},1), \text{HDL} \leftarrow \mathcal{N}(\frac{1}{2},1))}(\text{LDL} + \text{HDL}, \text{HD})$, that is, an action is only interpreted as a single-node intervention on total cholesterol if the conditional distribution of heart disease given total cholesterol is the same as in the observational regime. \circ

5.2 \mathbf{Int}_S : Letting multi-node interventions falsify a model is another way out of circularity

In this section, we discuss another option for avoiding circularity: letting multi-node interventions falsify a causal model. We structure the arguments analogous to those in [Section 5.1](#).

Informal overview of [Section 5.2](#). If we insist that all actions correspond to single-node interventions, then it becomes possible to falsify a causal model: If we perform an action and the resulting distribution cannot be explained by a single-node intervention, the causal model can be rejected. Explaining these observations by multi-node interventions may be a slippery slope leading to a circular interpretation where every observed distribution can be explained by some complex intervention in the model. In [Example 5.6](#), we show how to falsify a causal model under \mathbf{Int}_S (to be defined in [Definition 5.4](#)).

Consider the following interpretation that may violate **D2**, but satisfies **D0**, **D1**, **D3**, and **D4**.

Definition 5.4. \mathbf{Int}_S . An interpretation violating only **D2.** Let a data-generating process \mathcal{D} , representation \mathbf{Z}^* , compatible CBN \mathfrak{C} , and set of interventions \mathcal{I} in \mathfrak{C} be given. We define the interpretation \mathbf{Int}_S by the following rule: An intervention $\text{do}(j \leftarrow q_j, j \in J) \in \mathcal{I}$ is in $\mathbf{Int}_S^{\mathcal{I}}(a)$ if and only if the following three conditions hold:

- 1) $\mathcal{L}^a(Z_i^* \mid \mathbf{PA}_i^*) \sim q_i$ for all $i \in J$. That is, \mathbf{Int}_S satisfies **D0** (correct conditionals on intervened nodes).
- 2) $\mathcal{L}^a(Z_i^* \mid \mathbf{PA}_i^*) \approx p_i^{\mathfrak{C}}$ for all $i \in J$. That is, the kernels of the observational distribution are incompatible with the conditionals of intervened nodes under action a .¹⁴

¹⁴ [Proposition 5.5](#) (1) would not hold if we omitted condition 2) of [Definition 5.4](#), see [Appendix C](#) for a counterexample. [Proposition 5.5](#) (2) would still hold if we omitted condition 2).

- 3) $\mathcal{L}^a(\mathbf{Z}^*)$ is Markov w.r.t. the DAG of \mathfrak{C} . That is, \mathbf{Int}_S satisfies **D4** (an intervention does not create new dependencies).¹⁵

◦

The S in \mathbf{Int}_S is for ‘single-node’. \mathbf{Int}_S may violate **D2**, that is, an action can be interpreted as distinct interventions, because if $\text{do}(j \leftarrow q_j, j \in J) \in \mathbf{Int}_S^{\mathcal{I}}(a)$ is minimal and decomposable, then $\text{do}(j \leftarrow q_j, j \in J^*) \in \mathbf{Int}_S^{\mathcal{I}}(a)$ for every nonempty subset $J^* \subsetneq J$, but $\mathcal{L}^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in J)}(\mathbf{Z}) \neq \mathcal{L}^{\mathfrak{C}; \text{do}(j \leftarrow q_j, j \in J^*)}(\mathbf{Z})$.

Proposition 5.5. Let a data-generating process \mathcal{D} be given. Assume that \mathbf{Z}^* is emulated by CBN \mathfrak{C} and interventions \mathcal{I}^* .

- (1) If \mathcal{I}^* only contains single-node interventions, then \mathfrak{C} is an $\mathcal{I} - \mathbf{Int}_S$ valid model of \mathbf{Z}^* for every set of interventions \mathcal{I} in \mathfrak{C} .
- (2) On the other hand, if \mathcal{I}^* contains a minimal and decomposable multi-node intervention, then there exists a set of interventions \mathcal{I} in \mathfrak{C} such that \mathfrak{C} is not an $\mathcal{I} - \mathbf{Int}_S$ valid model of \mathbf{Z}^* .

Proof. (1) Assume that \mathcal{I}^* contains only single-node interventions. Consider some action a . Since \mathbf{Z}^* is emulated by \mathfrak{C} and \mathcal{I}^* , there is a single-node intervention $d^* \in \mathcal{I}^*$ such that $\mathcal{L}^a(\mathbf{Z}^*) = \mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z})$. Consider some intervention $d = \text{do}(j \leftarrow q_j, j \in J) \in \mathbf{Int}_S^{\mathcal{I}}(a)$. We want to show that $\mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z}) = \mathcal{L}^{\mathfrak{C}; d}(\mathbf{Z})$ and do this by arguing that $\mathcal{L}^{\mathfrak{C}; d^*}(Z_i | \mathbf{PA}_i) \sim p_i^{\mathfrak{C}; d}$ for all $i \in [n]$.

Per condition 1) of [Definition 5.4](#), we have for all $i \in J$ that

$$\mathcal{L}^{\mathfrak{C}; d^*}(Z_i | \mathbf{PA}_i) \sim q_i = p_i^{\mathfrak{C}; d}.$$

By condition 2), $\mathcal{L}^{\mathfrak{C}; d^*}(Z_i | \mathbf{PA}_i) \sim p_i^{\mathfrak{C}}$ for all $i \in J$, so the single-node intervention d^* must be a single-node intervention on a node in J . Therefore, d^* does not intervene on nodes outside of J , and thus we have for all $i \notin J$ that

$$\mathcal{L}^{\mathfrak{C}; d^*}(Z_i | \mathbf{PA}_i) \sim p_i^{\mathfrak{C}} = p_i^{\mathfrak{C}; d}.$$

In summary, we have established that $\mathcal{L}^{\mathfrak{C}; d^*}(Z_i | \mathbf{PA}_i) \sim p_i^{\mathfrak{C}; d}$ for all $i \in [n]$, which implies that $\mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z}) = \mathcal{L}^{\mathfrak{C}; d}(\mathbf{Z})$.

- (2) Assume that there is a minimal and decomposable multi-node intervention $d^* = \text{do}(j \leftarrow q_j, j \in J) \in \mathcal{I}^*$ and let a be such that $\mathcal{L}^a(\mathbf{Z}^*) = \mathcal{L}^{\mathfrak{C}; d^*}(\mathbf{Z})$. Fix some $j' \in J$ and let $\mathcal{I} = \{\text{do}(j' \leftarrow q_{j'})\}$ (which is well-defined since d^* is decomposable and thus $\mathcal{L}^{\mathfrak{C}; \text{do}(j' \leftarrow q_{j'})}(\mathbf{Z}) \neq \mathcal{L}^{\mathfrak{C}}(\mathbf{Z})$). $\text{do}(j' \leftarrow q_{j'}) \in \mathbf{Int}_S^{\mathcal{I}}(a)$ as 1) $\mathcal{L}^a(Z_{j'}^* | \mathbf{PA}_{j'}^*) \sim q_{j'}$, 2) $\mathcal{L}^a(Z_{j'}^* | \mathbf{PA}_{j'}^*) \sim p_{j'}^{\mathfrak{C}}$ (since d^* is minimal), and condition 3) trivially holds. But $\mathcal{L}^a(\mathbf{Z}^*) \neq \mathcal{L}^{\mathfrak{C}; \text{do}(j' \leftarrow q_{j'})}(\mathbf{Z})$ since d^* is minimal, and therefore, \mathfrak{C} is not an $\mathcal{I} - \mathbf{Int}_S$ valid model of \mathbf{Z}^* . ◻

¹⁵ [Proposition 5.5](#) would still hold if we omitted condition 3) of [Definition 5.4](#).

If there exists an action that induces a distribution like a multi-node intervention, then the model can be falsified. Since for a given representation \mathbf{Z}^* , it may be difficult to rule out such actions, [Proposition 5.5](#) (2) highlights a limitation of causal modeling using \mathbf{Int}_S .

In [Example 5.3](#), we could have falsified the model \mathbf{C} under interpretation \mathbf{Int}_S , like we did under interpretation \mathbf{Int}_P .

The following example is more extreme: \mathbf{Z}^* is emulated by \mathbf{A} and multi-node interventions \mathcal{I}^* in a way that is fine-tuned to make the opposite causal direction interventionally valid.

Example 5.6. a) Falsify a model under \mathbf{Int}_S , and b) reverse the causal direction.

Assume that $\mathbf{Z}^* = (Z_1^*, Z_2^*)$ is emulated by CBN \mathbf{A} and interventions \mathcal{I}^* . Let \mathbf{A} be given by DAG $Z_1 \rightarrow Z_2$ and kernels

$$\begin{aligned}\mathcal{L}^{\mathbf{A}}(Z_1) &= \text{Ber}(0.5) \\ \mathcal{L}^{\mathbf{A}}(Z_2 \mid Z_1 = 1) &= \text{Ber}(0.6) \\ \mathcal{L}^{\mathbf{A}}(Z_2 \mid Z_1 = 0) &= \text{Ber}(0.4),\end{aligned}$$

and let

$$\begin{aligned}\mathcal{I}^* = \{ & \text{do}(Z_1 \leftarrow \text{Ber}(0.6), Z_2 = 1), \\ & \text{do}(Z_1 \leftarrow \text{Ber}(0.4), Z_2 = 0), \\ & \text{do}(Z_1 = 1, Z_2 \leftarrow \text{Ber}(0.5)), \\ & \text{do}(Z_1 = 0, Z_2 \leftarrow \text{Ber}(0.5))\}.\end{aligned}$$

a) Falsify \mathbf{A} . Let $\mathcal{I} = \{\text{do}(Z_1 = 1), \text{do}(Z_1 = 0), \text{do}(Z_2 = 1), \text{do}(Z_2 = 0)\}$ be the set of modeled interventions. Let a be an action such that $\mathcal{L}^a(\mathbf{Z}^*) = \mathcal{L}^{\mathbf{A}; \text{do}(Z_1 \leftarrow \text{Ber}(0.6), Z_2 = 1)}(\mathbf{Z})$ (which exists since $\text{do}(Z_1 \leftarrow \text{Ber}(0.6), Z_2 = 1) \in \mathcal{I}^*$). Now \mathbf{A} is falsified as an $\mathcal{I} - \mathbf{Int}_S$ valid model of \mathbf{Z}^* , for example, because $\text{do}(Z_2 = 1) \in \mathbf{Int}_S^{\mathcal{I}}(a)$, but $\mathcal{L}^a(Z_1^*) = \text{Ber}(0.6) \neq \text{Ber}(0.5) = \mathcal{L}^{\mathbf{A}; \text{do}(Z_2 = 1)}(Z_1)$.

b) Reversing the causal direction. Instead, consider CBN \mathbf{C} with DAG $Z_1 \leftarrow Z_2$ and kernels

$$\begin{aligned}\mathcal{L}^{\mathbf{C}}(Z_2) &= \text{Ber}(0.5) \\ \mathcal{L}^{\mathbf{C}}(Z_1 \mid Z_2 = 1) &= \text{Ber}(0.6) \\ \mathcal{L}^{\mathbf{C}}(Z_1 \mid Z_2 = 0) &= \text{Ber}(0.4).\end{aligned}$$

\mathbf{C} is an $\mathcal{I} - \mathbf{Int}_S$ valid model of \mathbf{Z}^* for every set of interventions \mathcal{I} in \mathbf{C} , even though it has opposite causal direction than \mathbf{A} . This is true per [Proposition 5.5](#) (1) because \mathbf{Z}^* is emulated by \mathbf{C} and a set of single-node interventions, namely:

$$\begin{aligned}\mathcal{L}^{\mathbf{A}; \text{do}(Z_1 \leftarrow \text{Ber}(0.6), Z_2 = 1)}(Z_1, Z_2) &= \mathcal{L}^{\mathbf{C}; \text{do}(Z_2 = 1)}(Z_1, Z_2) \\ \mathcal{L}^{\mathbf{A}; \text{do}(Z_1 \leftarrow \text{Ber}(0.4), Z_2 = 0)}(Z_1, Z_2) &= \mathcal{L}^{\mathbf{C}; \text{do}(Z_2 = 0)}(Z_1, Z_2) \\ \mathcal{L}^{\mathbf{A}; \text{do}(Z_1 = 1, Z_2 \leftarrow \text{Ber}(0.5))}(Z_1, Z_2) &= \mathcal{L}^{\mathbf{C}; \text{do}(Z_1 = 1)}(Z_1, Z_2) \\ \mathcal{L}^{\mathbf{A}; \text{do}(Z_1 = 0, Z_2 \leftarrow \text{Ber}(0.5))}(Z_1, Z_2) &= \mathcal{L}^{\mathbf{C}; \text{do}(Z_1 = 0)}(Z_1, Z_2).\end{aligned}$$

In particular, \mathcal{Z}^* is emulated by both \mathcal{A} and \mathcal{C} , but only \mathcal{C} is a $\mathcal{I} - \mathbf{Int}_S$ valid model of \mathcal{Z}^* for $\mathcal{I} = \{\text{do}(Z_1 = 1), \text{do}(Z_1 = 0), \text{do}(Z_2 = 1), \text{do}(Z_2 = 0)\}$. The interventions \mathcal{I}^* in \mathcal{A} are fine-tuned to mimic single-node interventions in \mathcal{C} . The lack of fine-tuning between interventions has previously been suggested as a possible fundamental property of causal models [Janzing and Schölkopf, 2010, Janzing et al., 2016]. \circ

5.3 \mathbf{Int}_K : Interventions as simple actions

The larger the set of actions the more conditions must be satisfied for a causal model to be interventionally valid. More formally, there exists interpretations \mathbf{Int} such that if \mathcal{C} is an $\mathcal{I} - \mathbf{Int}$ valid model of \mathcal{Z}^* with set of actions \mathcal{A} , then \mathcal{C} may not be an $\mathcal{I} - \mathbf{Int}$ valid model of \mathcal{Z}^* with set of actions $\mathcal{A}' \supsetneq \mathcal{A}$. For example, under \mathbf{Int}_P , the model \mathcal{C} in Example 5.3 is not interventionally valid but it would be for a sufficiently small subset of \mathcal{A} ; and every compatible model is interventionally valid if $\mathcal{A} = \{\mathcal{O}\}$. More generally, Proposition 5.5 (2) and Proposition 5.2 (2) show that while \mathbf{Int}_S and \mathbf{Int}_P avoid the circularity of interpretation \mathbf{Int}_C (Corollary 3.3 and Proposition 4.1), these interpretations may prevent interventionally valid causal modeling of sensible representations if the set of actions is large.

Informal overview of Section 5.3. In this section, we present an interpretation \mathbf{Int}_K which has intermediate restrictiveness between \mathbf{Int}_S and \mathbf{Int}_C : $\mathcal{I} - \mathbf{Int}_S$ validity implies $\mathcal{I} - \mathbf{Int}_K$ validity and, since every compatible CBN is $\mathcal{I} - \mathbf{Int}_C$ valid, $\mathcal{I} - \mathbf{Int}_K$ validity implies $\mathcal{I} - \mathbf{Int}_C$ validity (while the reverse implications do not hold). We accomplish this by considering the complexity of actions and disqualifying actions that are not the most simple implementations of an intervention. The following thought experiment motivates why the complexity of actions is relevant to deciding which actions should be considered as which interventions.

Thought experiment. Suppose I am leading a sedentary lifestyle and am considering taking up smoking during my vacation. I have the following question: ‘How would smoking one pack of cigarettes over a week affect my heart health?’ What kind of experiment could be relevant to answer this question? Following Dawid [2021] and setting ethical issues aside, one approach might be the following study: Pay participants who are similar to me (including having a sedentary lifestyle) to smoke a pack of cigarettes over a week and then measure an indicator of heart health before and after. However, if participants started a rigorous exercise routine to offset the negative effects of smoking, the study would no longer capture how smoking alone impacts heart health; rather, the before-after measurements would reflect the combined effects of smoking and exercising on heart health. Naively, we want the participants to take up smoking while keeping everything else fixed. This is both a) impossible and b) undesirable. a) It is impossible because taking up smoking will necessarily affect other things as well. As Lewis [1973] explains “If we try too hard for exact similarity to the actual world in one respect, we will get excessive differences in some other respect.”¹⁶ In our case, smoking will inevitably impact other

¹⁶ Lewis [1973, page 9] puts it like this, considering what would happen if kangaroos had no tails: “We might think it best to confine our attention to worlds where kangaroos have no tails and everything else is as it actually is; but there are no such worlds. Are we to suppose that kangaroos have no tails but that their tracks in the sand are as they actually are? Then we shall have to suppose that these tracks

aspects of participants’ lives such as taking smoking breaks or carrying a lighter. b) It is undesirable because we do not want to keep heart health fixed. The problem seems to be that taking up smoking and exercise is not the most simple modification of the participants’ daily routine that involves taking up smoking, and this might disqualify the participants’ behavior as being an intervention on smoking.

Another way to understand the problem above is that the study participants are implementing interventions in a way that is counter to a commonsense consensus on how interventions should be implemented. As pointed out in Peters et al. [2017, page 121], the notion of falsification “includes the assumption that there is an agreement about what a randomized experiment should look like”. In this work, we attempt to move beyond human intuition toward mathematical precision. We do not think that human intuition is never good enough to do useful causal modeling. However, we believe that in many circumstances, such as those encountered in causal representation learning and causal abstraction, it is entirely unclear how human intuition could serve as a secure foundation, and we doubt that there is an implicit commonsense consensus on what constitutes a valid action to implement some intervention.

We now provide an interpretation that is based on a given notion of complexity of actions. For this, assume that there is some complexity measure $K : \mathcal{A} \rightarrow \mathbb{R}_+$ that assigns a positive real number to each action.

Definition 5.7. Int_K . An interpretation violating D1 and D2. Let a data-generating process \mathcal{D} , representation \mathcal{Z}^* , compatible CBN \mathcal{C} , set of interventions \mathcal{I} in \mathcal{C} , and complexity measure $K : \mathcal{A} \rightarrow \mathbb{R}_+$ be given. We define the interpretation Int_K by the following rule: An intervention $d \in \mathcal{I}$ is in $\text{Int}_K^{\mathcal{I}}(a)$ if and only if $a \in \arg \min_{a \in \mathcal{A}: d \in \text{Int}_K^{\mathcal{I}}(a)} K(a)$. \circ

Under this interpretation, the complexity measure of actions affects which actions are interpreted as implementations of which interventions and thereby whether a model is interventionaly valid. If a representation \mathcal{Z}^* is emulated by \mathcal{C} and \mathcal{I}^* with link $g : \mathcal{A} \setminus \{\mathcal{O}\} \rightarrow \mathcal{I}^*$, we show that \mathcal{C} is an $\mathcal{I}^* - \text{Int}_K$ valid model of \mathcal{Z}^* if the complexity measure K is strictly increasing in the number of nodes intervened upon by $g(a)$ (and some regularity conditions are met). We use $|g(a)|$ to denote the number of nodes intervened upon by $g(a)$ in \mathcal{C} .¹⁷ The complexity of actions ought to be given by some consideration that is external to the model; defining the complexity of actions in terms of $|g(a)|$ would be circular since $|g(a)|$ depends on the CBN \mathcal{C} .

Proposition 5.8. Let a data-generating process \mathcal{D} be given. Assume that \mathcal{Z}^* is emulated by CBN \mathcal{C} and interventions \mathcal{I}^* with link $g : \mathcal{A} \setminus \{\mathcal{O}\} \rightarrow \mathcal{I}^*$. Assume that

1. $K(a) = t(|g(a)|)$ for some strictly increasing $t : \mathbb{R} \rightarrow \mathbb{R}$.
2. $\mathcal{L}^a(\mathcal{Z}^*)$ has the same finite support for every $a \in \mathcal{A}$.

are produced in a way quite different from the actual way. Are we to suppose that kangaroos have no tails but that their genetic makeup is as it actually is? Then we shall have to suppose that genes control growth in a way quite different from the actual way (or else that there is something, unlike anything there actually is, that removes the tails). And so it goes; respects of similarity and difference trade-off. If we try too hard for exact similarity to the actual world in one respect, we will get excessive differences in some other respect.”

¹⁷ Formally, if \mathcal{Z}^* is emulated by \mathcal{C} and \mathcal{I}^* with link $g : \mathcal{A} \setminus \{\mathcal{O}\} \rightarrow \mathcal{I}^*$, we define $|g(a)| = |J|$ for $g(a) = \text{do}(j \leftarrow q_j, j \in J)$.




Figure 3: Depiction of the situation in [Example 5.9](#). There are 6 different locations, three on each the left and right side S^* , with associated rewards R^* (+1 or -1). Observationally, the stick figure picks a position P^* using a uniform distribution over the 6 locations. Whether the causal model $S \rightarrow R$ makes correct predictions about interventions on left/right, depends on which specific actions are interpreted as interventions.

3. Every $d \in \mathcal{I}$ is minimal and $\mathcal{I} \subseteq \mathcal{I}^*$.

Then \mathfrak{C} is an $\mathcal{I} - \mathbf{Int}_K$ valid model of \mathbf{Z}^* .

Proof. Let $a' \in \mathcal{A}$, $\mathcal{I} \subseteq \mathcal{I}^*$, and $d = \text{do}(j \leftarrow q_j, j \in J) \in \mathbf{Int}_K^{\mathcal{I}}(a')$ be given. We want to show that $\mathcal{L}^{a'}(\mathbf{Z}^*) = \mathcal{L}^{\mathfrak{C};d}(\mathbf{Z})$.

Since $d \in \mathbf{Int}_S^{\mathcal{I}}(a')$ (by definition of \mathbf{Int}_K), and since $\mathcal{L}^a(\mathbf{Z}^*)$ has the same finite support for every a , we have for every $i \in J$ that $p^{\mathfrak{C};g(a')}(z_i \mid \mathbf{pa}_i) = p^{\mathfrak{C};d}(z_i \mid \mathbf{pa}_i)$ for every z_i and \mathbf{pa}_i in the support.

Since $g^{-1}(\{d\})$ is nonempty (because $g : \mathcal{A} \setminus \{\mathcal{O}\} \rightarrow \mathcal{I}^*$ is surjective and $d \in \mathcal{I} \subseteq \mathcal{I}^*$) and $d \in \mathbf{Int}_S^{\mathcal{I}}(a)$ for every $a \in g^{-1}(\{d\})$, we have that $K(a') \leq t(|J|)$, which implies that $|g(a')| \leq |J|$.

Since every intervention in \mathcal{I} is minimal, d is minimal, and we have for every $i \in J$ that $p^{\mathfrak{C};g(a')}(z_i \mid \mathbf{pa}_i) \neq p^{\mathfrak{C}}(z_i \mid \mathbf{pa}_i)$ for some z_i and \mathbf{pa}_i in the support. Since $|g(a')| \leq |J|$, this means that $g(a')$ is an intervention on the nodes in J , and only those. Therefore, for $i \notin J$, we must have $p^{\mathfrak{C};g(a')}(z_i \mid \mathbf{pa}_i) = p^{\mathfrak{C}}(z_i \mid \mathbf{pa}_i)$ for every z_i and \mathbf{pa}_i in the support as these nodes are not intervened upon by $g(a')$. In summary, we have that $p^{\mathfrak{C};g(a')}(z) = p^{\mathfrak{C};d}(z)$ for all z in the support. \square

We now provide a simple example with two different complexity measures and investigate how they affect the interventional validity of a model.

Example 5.9. Making an \mathbf{Int}_S invalid model valid by specifying a complexity measure. Consider CBN \mathfrak{A} with graph $P \rightarrow R$ and kernels

$$P \sim \text{Unif}([6])$$

$$R := \begin{cases} +1 & P \in \{1, 3, 5\} \\ -1 & P \in \{2, 4, 6\} \end{cases}.$$

Assume that (P^*, R^*) is emulated by CBN \mathcal{A} and intervention set

$$\mathcal{I}^* = \{d \mid d \text{ is a single-node intervention on } P \text{ such that } \mathbb{P}^{\mathcal{A};d}(P \in [6]) = 1 \}.$$

P^* stands for position and R^* for reward, see [Figure 3](#). Consider the feature left/right given by $S^* = \mathbb{1}(P^* \in \{4, 5, 6\})$, which is 1 if the position is on the right and 0 if the position is on the left. We now investigate if \mathfrak{C} , given by $S \rightarrow R$ and kernels

$$\begin{aligned} \mathbb{P}^{\mathfrak{C}}(S = 1) &= \frac{1}{2} \\ \mathbb{P}^{\mathfrak{C}}(S = 0) &= \frac{1}{2} \\ \mathbb{P}^{\mathfrak{C}}(R = +1 \mid S = 0) &= \frac{2}{3} \\ \mathbb{P}^{\mathfrak{C}}(R = -1 \mid S = 0) &= \frac{1}{3} \\ \mathbb{P}^{\mathfrak{C}}(R = +1 \mid S = 1) &= \frac{1}{3} \\ \mathbb{P}^{\mathfrak{C}}(R = -1 \mid S = 1) &= \frac{2}{3}, \end{aligned}$$

is an interventionally valid model of (S^*, R^*) under interventions $\mathcal{I} = \{\text{do}(S = 0), \text{do}(S = 1)\}$. Notice that we have defined the kernels so that \mathfrak{C} is compatible with (S^*, R^*) .

\mathfrak{C} is not an $\mathcal{I} - \mathbf{Int}_S$ valid model of (S^*, R^*) . To see this, consider action a such that $\mathbb{P}^a(P^* = 1, R^* = 1) = 1$ (such an action exists since it corresponds to an intervention in \mathcal{I}^*). Now $\text{do}(S = 0) \in \mathbf{Int}_S^{\mathcal{I}}(a)$ as $\mathcal{L}^a(S^*) = \delta_0$ (correct conditionals on intervened nodes) and $\mathcal{L}^a(S^*) \approx p_S^{\mathfrak{C}}$. However, $\mathbb{P}^a(R^* = 1) = 1 \neq \frac{2}{3} = \mathbb{P}^{\mathfrak{C};\text{do}(S=0)}(R = 1)$.

Making \mathfrak{C} interventionally valid by using reverse entropy as complexity. If we define the complexity of actions a by the inverse entropy of P^* in $\mathcal{L}^a(P^*)$, $K(a) = \frac{1}{H^a(P^*)}$, then \mathfrak{C} is an $\mathcal{I} - \mathbf{Int}_K$ valid model of (S^*, R^*) . Using the reverse entropy as complexity may be reasonable since higher entropy of P^* intuitively means less specificity and thus intuitively corresponds to a less complex action. Using this measure of complexity, for example, implies that the least complex action that goes left with probability 1, is an action a such that $\mathcal{L}^a(P^*) = \text{Unif}(\{1, 2, 3\})$, which implies that $\mathcal{L}^a(R^* \mid S^* = 0) = \mathcal{L}^{\mathfrak{C}}(R \mid S = 0)$. Therefore, for all a , $\text{do}(S = 0) \in \mathbf{Int}_K^{\mathcal{I}}(a)$ only if $\mathcal{L}^a(R^* \mid S^* = 0) = \mathcal{L}^{\mathfrak{C}}(R \mid S = 0)$ and $\mathbb{P}^a(S^* = 0) = 1$; and likewise $\text{do}(S = 1) \in \mathbf{Int}_K^{\mathcal{I}}(a)$ only if $\mathcal{L}^a(R^* \mid S^* = 1) = \mathcal{L}^{\mathfrak{C}}(R^* \mid S^* = 1)$ and $\mathbb{P}^a(S^* = 1) = 1$. This implies that for all a , $\text{do}(S = 0) \in \mathbf{Int}_K^{\mathcal{I}}(a)$ only if $\mathcal{L}^a(S^*, R^*) = \mathcal{L}^{\mathfrak{C};\text{do}(S=0)}(S, R)$ and $\text{do}(S = 1) \in \mathbf{Int}_K^{\mathcal{I}}(a)$ only if $\mathcal{L}^a(S^*, R^*) = \mathcal{L}^{\mathfrak{C};\text{do}(S=1)}(S, R)$, so \mathfrak{C} is an $\mathcal{I} - \mathbf{Int}_K$ valid model of (S^*, R^*) .

Falsifying \mathfrak{C} if distance is used as complexity measure. If we instead assume that complexity is given by the number of steps required to get from the starting position to the respective field, that is,

$$K(a) = \mathbb{E}^a(3 \cdot \mathbb{1}(P^* \in \{1, 3, 4, 6\}) + 2 \cdot \mathbb{1}(P^* \in \{2, 5\})),$$

then \mathfrak{C} is not an $\mathcal{I} - \mathbf{Int}_K$ valid model of (S^*, R^*) . To see this, consider an action a such that $\mathbb{P}^a(P^* = 2, R^* = -1) = 1$ (such an action exists since it corresponds to

an intervention in \mathcal{I}^*). Then $\text{do}(S = 0) \in \mathbf{Int}_K^{\mathcal{I}}(a)$ since the least complex action that goes left with probability 1 is the one that always goes to position $P^* = 2$, but $\mathbb{P}^a(R^* = -1) = 1 \neq \frac{1}{3} = \mathbb{P}^{\mathfrak{C}; \text{do}(S=0)}(R = -1)$.

That there exist actions such that $\mathcal{L}^a(R^* | S^*) \approx p_{R|S}^{\mathfrak{C}}$ even though intuitively all actions seem to be modifying side rather than the reward mechanism, is an example of what [Zhu et al. \[2024, Definition 6\]](#) call ‘macro-confounding’, here between R^* and S^* . \circ

6 Discussion of implications for related research

In this section, we will examine causal representation learning, causal discovery, and causal abstraction in light of the previous sections. Most notably, we argue that identifiability is not sufficient for interventional validity and that causal abstraction rests on an infinite regress. We also discuss connections to the philosophical literature on the logic of conditionals and, in [Appendix F](#), related work by [Janzing and Meija \[2024\]](#).

6.1 Causal representation learning

6.1.1 Identifiability & interventional validity in causal representation learning

Works on interventional causal representation learning [[Squires et al., 2023](#), [Buchholz et al., 2024](#), [Jin and Syrkanis, 2024](#), [Varici et al., 2024](#), [von Kügelgen et al., 2024](#), [Zhang et al., 2024](#)] often consider the following setting: There are some latent variables $\mathbf{Z}^* = (Z_1^*, \dots, Z_n^*)$ emulated by a causal Bayesian network \mathfrak{A} (with DAG \mathcal{G}) and interventions \mathcal{I}^* ,¹⁸ the observed data $\mathbf{X} = (X_1^*, \dots, X_m^*)$ is given by some mixing function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, $\mathbf{X}^* = f(\mathbf{Z}^*)$, where f is commonly assumed (at least) to be a diffeomorphism onto its image (e.g., [von Kügelgen et al. \[2024\]](#), [Varici et al. \[2024\]](#)). In this setting, the goal is to recover \mathbf{Z}^* and \mathcal{G} from the distribution of \mathbf{X}^* in different environments $E \subseteq \mathcal{A}$, that is, from $\{\mathcal{L}^a(\mathbf{X}^*)\}_{a \in E}$.¹⁹ Even under strong assumptions, this can usually only be done up to certain ambiguities. At most, we can identify the unmixing function $f^{-1} : \text{Im}(f) \rightarrow \mathbb{R}^n$ up to an equivalence class. Definition 2.6 of [von Kügelgen et al. \[2024\]](#) introduces one such equivalence class.

Definition 6.1. \sim_{CRL} [[von Kügelgen et al., 2024, Definition 2.6](#)]. Let \mathcal{H} be a set of unmixing functions $h : \text{Im}(f) \rightarrow \mathbb{R}^n$, and let \mathfrak{G} be the set of DAGs over n nodes. Let \sim_{CRL} be the equivalence relation over $\mathcal{H} \times \mathfrak{G}$ defined as

$$(h_1, \mathcal{G}_1) \sim_{\text{CRL}} (h_2, \mathcal{G}_2) \iff (h_2, \mathcal{G}_2) = (\mathbf{P}_{\pi^{-1}} \circ \phi \circ h_1, \pi(\mathcal{G}_1))$$

for some element-wise diffeomorphism $\phi(\mathbf{v}) = (\phi_1(v_1), \dots, \phi_n(v_n))$ of \mathbb{R}^n and graph isomorphism $\pi : \mathcal{G}_1 \mapsto \mathcal{G}_2$, where \mathbf{P}_{π} is the corresponding permutation matrix over $[n]$. \circ

¹⁸ The data-generating process is often described in terms of a latent or augmented structural causal model — see, for example, [von Kügelgen et al. \[2024, Section 2.1\]](#), [Buchholz et al. \[2024, Assumption 2\]](#), and [Li et al. \[2024, Definition 2.1\]](#).

¹⁹ Some works assume a known graph and focus only on learning the unmixing function [[Wendong et al., 2024](#), [Li et al., 2024](#)]. Other works consider additional assumptions on the mixing and latent distribution [[Ahuja et al., 2023](#)]. In this section, we focus on works that address the problem of learning causal representations based on multiple interventions or domains, rather than on counterfactual and multiview data [[Brehmer et al., 2022](#), [von Kügelgen et al., 2021](#), [Daunhawer et al., 2023](#), [Xu et al., 2024](#), [Yao et al., 2024](#)], or purely observational data [[Xie et al., 2020](#), [Kivva et al., 2021](#), [Welch et al., 2024](#)], or settings based on temporal structure [[Ahuja et al., 2022](#), [Lachapelle et al., 2022](#), [Lippe et al., 2022](#)].

\sim_{CRL} **preserves causal validity if the latent representation is emulated by single-node interventions.** The \sim_{CRL} equivalence class is quite special: If $\mathbf{Z}^* = f^{-1}(\mathbf{X}^*)$ is emulated by a CBN $\tilde{\mathcal{A}}$ and single-node interventions in $\tilde{\mathcal{A}}$ and $(h, \tilde{\mathcal{G}}) \sim_{\text{CRL}} (f^{-1}, \mathcal{G})$, then there exists a CBN $\tilde{\tilde{\mathcal{A}}}$ such that $h(\mathbf{X}^*)$ is emulated by single-node interventions in $\tilde{\tilde{\mathcal{A}}}$.²⁰ This implies that $\tilde{\tilde{\mathcal{A}}}$ is an $\mathcal{I} - \mathbf{Int}_S$ valid model of $h(\mathbf{X}^*)$ for every set of interventions \mathcal{I} in $\tilde{\tilde{\mathcal{A}}}$, see [Proposition 5.5](#) (1). From this point of view, there is no sense in which $f^{-1}(\mathbf{X}^*)$ is more a ‘causal representation’ or the ‘ground truth’ than $h(\mathbf{X}^*)$, and thus the choice of one representation as the assumed latent ground-truth is arbitrary.²¹

\sim_{CRL} -**identifiability does not imply interventional validity.** Suppose that we have assumptions in place such that f^{-1} is identifiable up to \sim_{CRL} based on $\{\mathcal{L}^a(\mathbf{X}^*)\}_{a \in E}$, and assume that $(h, \tilde{\mathcal{G}}) \sim_{\text{CRL}} (f^{-1}, \mathcal{G})$. Let $\tilde{\tilde{\mathcal{A}}}$ (with graph $\tilde{\tilde{\mathcal{G}}}$) and $\tilde{\mathcal{I}}^*$ be such that $h(\mathbf{X}^*)$ is emulated by $\tilde{\tilde{\mathcal{A}}}$ and $\tilde{\mathcal{I}}^*$ with link $\tilde{g} : \mathcal{A} \setminus \{\mathcal{O}\} \rightarrow \tilde{\mathcal{I}}^*$. Then, it may be the case that $\tilde{\tilde{\mathcal{A}}}$ is not a $\{d \in \tilde{\mathcal{I}}^* \mid d \text{ is single-node intervention}\} - \mathbf{Int}_S$ valid model of $h(\mathbf{X}^*)$; in particular, [Proposition 5.5](#) (1) only establishes validity if all interventions in $\tilde{\mathcal{I}}^*$ are single-node. In fact, as we will illustrate in [Example 6.2](#), $\tilde{g}(E) - \mathbf{Int}_S$ can be an invalid model of $h(\mathbf{X}^*)$ even if $\tilde{g}(a)$ were a single-node intervention for all $a \in E$. Therefore, to ensure interventional validity we need to not only make assumptions about the observed environments (often necessary for identifiability), but also about the entire set of possible actions (this type of extrapolation to hypothetical (future) environments is discussed in [Bühlmann \[2020\]](#)). Conversely, even if the identified latent model is interventionally valid, other models outside the equivalence class can also be interventionally valid (see [Section 6.1.2](#) and [Example 6.6](#)).

We now present an example that illustrates the distinction between identifiability and interventional validity.

Example 6.2. Identifiability does not imply interventional validity. Assume a data-generating process where $\mathcal{A} = \{\mathcal{O}, a_1, a_2, a_3\}$ and that the latent variables (A^*, B^*) have the following distributions

$$\begin{aligned}\mathcal{L}^{\mathcal{O}}(A^*, B^*) &= \mathcal{N}\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & \frac{1}{2} \\ \frac{1}{2} & 1 \end{pmatrix}\right) \\ \mathcal{L}^{a_1}(A^*, B^*) &= \mathcal{N}\left(\begin{pmatrix} 1 \\ \frac{1}{2} \end{pmatrix}, \begin{pmatrix} 1 & \frac{1}{2} \\ \frac{1}{2} & 1 \end{pmatrix}\right) \\ \mathcal{L}^{a_2}(A^*, B^*) &= \mathcal{N}\left(\begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}\right) \\ \mathcal{L}^{a_3}(A^*, B^*) &= \mathcal{N}\left(\begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 & \frac{1}{2} \\ \frac{1}{2} & 1 \end{pmatrix}\right)\end{aligned}$$

²⁰ This follows from Proposition B.1 in [von Kügelgen et al. \[2024\]](#).

²¹ [von Kügelgen et al. \[2024\]](#) posit that “since the scale of the variables is arbitrary, we clearly cannot predict the exact outcomes of interventions.” This statement may be too strong if we adopt the perspective that there is no unique ‘true’ causal representation, see [Definition 6.3](#). When do we interpret an action a as corresponding to intervention $\text{do}(H_1 := 3)$? Under interpretation \mathbf{Int}_S , for example, this would simply be an action such that $\mathcal{L}^a(h(\mathbf{X}^*)_1) = \delta_3$ (assuming that condition 2) and 3) of [Definition 5.4](#) are also satisfied). It is no more difficult to predict the outcome of interventions using the representation $h(\mathbf{X}^*)$ than the representation $f^{-1}(\mathbf{X}^*)$.

Let \mathcal{A} be a CBN with graph $A \rightarrow B$ and $\mathcal{I}_{\mathcal{A}}^*$, consisting only of minimal interventions, be such that (A^*, B^*) is emulated by \mathcal{A} and $\mathcal{I}_{\mathcal{A}}^*$ with link $g_{\mathcal{A}} : \mathcal{A} \setminus \{\mathcal{O}\} \rightarrow \mathcal{I}_{\mathcal{A}}^*$. From $\mathcal{L}^{a_1}(A^*, B^*)$, $\mathcal{L}^{a_2}(A^*, B^*)$, and the fact that $\mathcal{I}_{\mathcal{A}}^*$ only contains minimal interventions, we can deduce that $g_{\mathcal{A}}(a_1) = \text{do}(A \leftarrow \mathcal{N}(1, 1))$ and $g_{\mathcal{A}}(a_2) = \text{do}(B \leftarrow \mathcal{N}(1, 1))$ are perfect single-node interventions, and that $g_{\mathcal{A}}(a_3)$ is a multi-node intervention. Assume that we observe $\mathbf{X}^* = f(A^*, B^*)$ for a linear injective mixing function f in environments $E = \{\mathcal{O}, a_1, a_2\}$. Following Squires et al. [2023], we assume that the non-observational environments, $\{a_1, a_2\}$, correspond to perfect single-node interventions.²² Since this assumption is met here, we can identify \mathcal{A} up to permutation and scaling [Squires et al., 2023]. But the CBN \mathcal{A} with graph $A \rightarrow B$ is neither $\{g_{\mathcal{A}}(a_1), g_{\mathcal{A}}(a_2)\} - \mathbf{Int}_S$ nor $\{g_{\mathcal{A}}(a_1), g_{\mathcal{A}}(a_2)\} - \mathbf{Int}_P$ valid.²³ Therefore, to ensure that identifiability implies interventional validity, we need additional assumptions. \circ

6.1.2 A definition of causal representations

Using the notion of interventional validity (Definition 2.7), we now provide a formalization of what it means for a representation to be a causal representation. To the best of our knowledge, this is the first definition in the literature that defines a causal representation based on its properties rather than stipulating that a certain representation is the ‘ground truth’ causal one.²⁴

Definition 6.3. Causal Representation. Let a data-generating process \mathcal{D} be given. We say that a representation \mathbf{Z}^* of \mathcal{D} is an $\mathcal{I} - \mathbf{Int}$ causal representation of \mathcal{D} if there exists a CBN \mathcal{C} and a set of interventions \mathcal{I} in \mathcal{C} such that \mathcal{C} is an $\mathcal{I} - \mathbf{Int}$ valid model of \mathbf{Z}^* . \circ

The larger \mathcal{I} is, the more interventions the model purports to make predictions about, and the interpretation \mathbf{Int} specifies which actions are interpreted as which interventions. The interpretation \mathbf{Int}_C is special as every representation \mathbf{Z}^* of a data-generating process is an $\mathcal{I} - \mathbf{Int}_C$ causal representation if \mathcal{I} is a set of interventions in a \mathbf{Z}^* -compatible CBN; that is, without committing to certain non-circular interpretations, there is apparently nothing distinctively causal about causal representations.

A consequence of Definition 6.3 is that there is no unique causal representation, which clarifies how we may model a data-generating process at different levels of abstraction, see Section 6.3, and, as suggested by Sadeghi and Soo [2024], that “[t]here is no need to take the true causal graph as the primitive object”. It also suggests an approach to causal representation learning that focuses on learning representations that satisfy interventional validity rather than recovering some ‘true’ latent representation emulated by a CBN. In fact, a key insight of the present work is that a representation \mathbf{Z}^* can be emulated by a CBN \mathcal{C} without \mathcal{C} being an $\mathcal{I} - \mathbf{Int}$ valid model of \mathbf{Z}^* (for non-circular interpretation \mathbf{Int} and nonempty set of interventions \mathcal{I} , see, for example, Propositions 5.2, 5.5, and 5.8).

²² Specifically $\mathcal{L}^{a_1}(A^*, B^*) = \mathcal{L}^{\mathcal{A}; \text{do}(A \leftarrow \mathcal{N}(1, 1))}(A, B)$ and $\mathcal{L}^{a_2}(A^*, B^*) = \mathcal{L}^{\mathcal{A}; \text{do}(B \leftarrow \mathcal{N}(1, 1))}(A, B)$.

²³ To see this, notice, for example, that $g_{\mathcal{A}}(a_1) = \text{do}(A \leftarrow \mathcal{N}(1, 1)) \in \mathbf{Int}_S^{\{g_{\mathcal{A}}(a_1), g_{\mathcal{A}}(a_2)\}}(a_3)$ and $g_{\mathcal{A}}(a_1) = \text{do}(A \leftarrow \mathcal{N}(1, 1)) \in \mathbf{Int}_P^{\{g_{\mathcal{A}}(a_1), g_{\mathcal{A}}(a_2)\}}(a_3)$, that is, a_3 is interpreted as the intervention $g_{\mathcal{A}}(a_1)$, while the distributions $\mathcal{L}^{a_3}(A^*, B^*) \neq \mathcal{L}^{\mathcal{C}; g(a_1)}(A, B)$ do not match.

²⁴ Cohen [2022] raised the related point that existing frameworks for causal modeling “give no guidance regarding variable [...] representation, and [...] no indication as to which behaviour policies or physical transformations of state space shall count as interventions.”

Furthermore, the assumptions that render a latent CBN representation identifiable (up to some equivalence class) need not ensure it is a causal representation with desirable properties such as interventional validity, see [Section 6.1.1](#).

Toward a new approach to causal representation learning. We can formulate the task as follows: Let a data-generating process \mathcal{D} and observed low-level features \mathbf{X}^* be given. Find a non-circular interpretation \mathbf{Int} and a transformation h such that $h(\mathbf{X}^*)$ is an $\mathcal{I} - \mathbf{Int}$ causal representation for a suitable set of interventions \mathcal{I} .

Requiring that $h(\mathbf{X}^*)$ is a causal representation is not sufficient to get an interesting representation. For example, the trivial representation $Z^* = h(\mathbf{X}^*) := 0$ is a causal representation, according to [Definition 6.3](#), since the CBN with one node Z and distribution $\mathbb{P}(Z = 0) = 1$ is an $\mathcal{I} - \mathbf{Int}$ valid model of $Z^* = 0$ for every set of interventions \mathcal{I} in \mathfrak{C} if \mathbf{Int} is an interpretation that satisfies **D0**. Therefore in addition to requiring interventional validity, we probably want our representations to satisfy further criteria (some of which have, sometimes implicitly, motivated representation learning approaches); for example:

- We may want interventions in \mathcal{I} to be implementable, that is, that for every intervention $d \in \mathcal{I}$ there exists an action a such that $d \in \mathbf{Int}^{\mathcal{I}}(a)$.
This criterion is reminiscent of the idea that causes must be manipulable [[Cartwright, 2007](#), [Glymour and Glymour, 2014](#), [Pearl, 2018, 2019](#)].
- We may want that for each variable there exists a single-node intervention in \mathcal{I} that intervenes on it.
This criterion is reminiscent of the idea of autonomy [[Aldrich, 1989](#)]. See [Janzing and Meija \[2024\]](#) for a related requirement that we discuss in [Appendix F](#).
- We may want to model only some interventions on specific nodes, that is, \mathcal{I} should be restricted in some way.
[[Dawid, 2021](#)]
- We may want the transformation h to disregard little or no information.
See [Bengio et al. \[2013\]](#) for a review of representation learning where this is discussed as a desirable criterion.

- We may want the representation to have a prespecified number of nodes. [Kekić et al., 2024]
- We may want the representation to contain a prespecified aspect of the data-generating process, that is, parts of h may be prespecified. [Chalupka et al., 2015, Weichwald et al., 2016]
- We may want the resulting representation to be useful for (computationally or statistically efficiently) solving a prespecified set of downstream tasks. See, for example, Kinney and Watson [2020], Gultchin et al. [2021], Lachapelle et al. [2023], Dyer et al. [2024], Saengkyongam et al. [2024], Christgau and Hansen [2024].

What precise properties to require of a causal representation and what it may be used for, is in our opinion a neglected question; see [Woodward, 2016, Bing et al., 2024, Cadei et al., 2024] for discussions. While some existing representation learning approaches strive for representations that satisfy criteria similar to the ones listed above, the resulting causal model does not make precise empirical predictions when there is no explicit choice of how to interpret actions as interventions. Since many transformations of the observed data may yield causal representations, how to choose between representations and which properties to impose is a pressing question for future research.

Learning objectives and identifiability. Once the requirements for a causal representation are well specified, an important question for future research is the precise formulation of a learning objective for causal representations in the sense of Definition 6.3. Existing learning objectives in causal representation learning may yield causal representations but will require an explicit choice of a suitable interpretation and additional assumptions. Furthermore, our work suggests a different perspective on identifiability theory: Rather than focusing on identifiability as a guarantee for the learned representation to be disentangled with respect to some ‘true’ latent variables, an identifiability analysis could assess whether all maximizers of a given learning objective share properties we deem desirable (similar in spirit to, for example, the analysis by Marconato et al. [2024]). For example, in Section 6.1.1, we show that the \sim_{CRL} equivalence class preserves interventional validity under the interpretation \mathbf{Int}_S (Definition 5.4). If one is willing to make assumptions on the set of all possible actions and not only those that induce the distributions used for learning the representation, then an identifiability result proving that maximizers of a given objective are in the \sim_{CRL} equivalence class would thus provide a sufficient (but not necessary) condition for all (or none) of the corresponding representations to be causal representations. Since coarser equivalence classes than \sim_{CRL} may preserve interventional validity (depending on the chosen interpretation and assumptions), it may be possible to simplify the learning problem.

6.2 Causal discovery

What do we assume when we assume that observed variables are described by a CBN? The starting point of causal discovery is, in the words of Dawid [2010, Assumption 6.1], the assumption that “[t]here exists some DAG $[\mathcal{G}]$ that is a causal DAG representation of the system.” Dawid [2010] considers this a “strong [a]ssumption”, suggesting that the assumption does not amount to mere emulation, which is always possible (see Definition 2.5 and discussion below it). Our work makes precise what this

assumption may amount to other than emulation, namely the existence of a CBN \mathfrak{C} and a set of interventions \mathcal{I} in \mathfrak{C} such that \mathfrak{C} is an \mathcal{I} – **Int** valid model of the given representation \mathbf{Z}^* (see [Definition 6.3](#)). Whether this assumption holds depends both on the interpretation **Int** and the modeled interventions \mathcal{I} . We clarify that this assumption can fail to be true, namely for a given representation \mathbf{Z}^* , non-circular interpretation **Int**, and set of interventions \mathcal{I} , there may not exist an \mathcal{I} – **Int** valid model of \mathbf{Z}^* . We demonstrate this by the classical example of the effect of total cholesterol on heart disease. Here, our framework enables us to make the claim formally precise that this representation does not admit a causal model.

Example 6.4. Total cholesterol [Example 5.3](#) continued. Maybe no causal graph is adequate. Assume again that (LDL^*, HDL^*, HD^*) is emulated by \mathfrak{A} and interventions $\mathcal{I}^* = \{\text{do}(LDL \leftarrow \mathcal{N}(y, 1), HDL \leftarrow \mathcal{N}(x, 1)) \mid x, y \in \mathbb{R}\}$, and let $(TC^*, HD^*) = (LDL^* + HDL^*, HD^*)$. In this example, we argue that the representation (TC^*, HD^*) is not a causal representation given interpretation **Int**_S and set of perfect interventions $\mathcal{I} = \{\text{do}(TC \leftarrow \mathcal{N}(1, 2)), \text{do}(HD \leftarrow \mathcal{N}(1, 6))\}$ that change the mean of total cholesterol or heart disease, respectively. Observationally, total cholesterol and heart disease are correlated. Therefore, every compatible CBN must have an edge between those two nodes, that is, either $TC \rightarrow HD$ or $TC \leftarrow HD$.

Let \mathfrak{C} be a compatible CBN with graph $TC \rightarrow HD$. Then, similar to [Example 5.3](#), we have an action a such that $\text{do}(TC \leftarrow \mathcal{N}(1, 2)) \in \mathbf{Int}_{\mathcal{I}}^{\mathfrak{C}}(a)$, but $\mathbb{E}^a(HD^*) = 2 \neq \frac{1}{2} = \mathbb{E}^{\mathfrak{C}; \text{do}(TC \leftarrow \mathcal{N}(1, 2))}(HD)$.

We now argue that having heart disease cause total cholesterol also results in an invalid model. Let \mathfrak{h} be a compatible CBN with graph $TC \leftarrow HD$. Let a be an action such that $\mathcal{L}^a(LDL^*, HDL^*, HD^*) = \mathcal{L}^{\mathfrak{A}; \text{do}(LDL \leftarrow \mathcal{N}(\frac{1}{2}, 1), HDL \leftarrow \mathcal{N}(0, 1))}(LDL, HDL, HD)$. Now, $\text{do}(HD \leftarrow \mathcal{N}(1, 6)) \in \mathbf{Int}_{\mathcal{I}}^{\mathfrak{h}}(a)$, but the distributions do not match, for example, because $\mathbb{E}^a(TC^*) = \frac{1}{2}$, while

$$\begin{aligned} & \mathbb{E}^{\mathfrak{h}; \text{do}(HD \leftarrow \mathcal{N}(1, 6))}(TC) \\ &= \mathbb{E}^{\mathfrak{h}; \text{do}(HD \leftarrow \mathcal{N}(1, 6))} \left(\mathbb{E}^{\mathfrak{h}; \text{do}(HD \leftarrow \mathcal{N}(1, 6))}(TC \mid HD) \right) \\ &\stackrel{*}{=} \mathbb{E}^{\mathfrak{h}; \text{do}(HD \leftarrow \mathcal{N}(1, 6))} \left(\frac{HD}{6} \right) \\ &= \frac{1}{6}, \end{aligned}$$

where $\stackrel{*}{=}$ follows by calculating the conditional mean in the joint normal distribution $\mathcal{L}^{\mathfrak{h}; \text{do}(HD \leftarrow \mathcal{N}(1, 6))}(TC, HD) = \mathcal{N} \left(\begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 & 1 \\ 1 & 6 \end{pmatrix} \right)$.

In summary, there does not exist an \mathcal{I} – **Int**_S valid model of (TC^*, HD^*) . ◦

Identifiability does not imply interventional validity in causal discovery. In [Example 6.4](#), we argue that the representation (TC^*, HD^*) does not admit an interventionally valid causal model (for a given interpretation and set of interventions). In this example, it is also unclear which assumptions one would use to identify the causal direction. Unfortunately, identifiability does not ensure interventional validity. For example, if the

representation \mathbf{Z}^* is emulated by \mathfrak{C} and \mathcal{I}^* with link g and the observed environments $\{\mathcal{L}^{\mathfrak{C};g(a)}(\mathbf{Z}^*)\}_{a \in E}$, $E \subseteq \mathcal{A}$, correspond to single-node interventions in \mathfrak{C} , then we can identify \mathfrak{C} [Eberhardt et al., 2006]. But this does not guarantee that \mathfrak{C} is a $\{g(E)\} - \mathbf{Int}$ valid model of \mathbf{Z}^* for the non-circular interpretations \mathbf{Int}_S and \mathbf{Int}_P , see Example 6.2. Work on causal discovery focuses on identifiability of an emulating CBN \mathfrak{C} since it is implicitly assumed that \mathfrak{C} is a valid model of \mathbf{Z}^* if \mathbf{Z}^* is emulated by \mathfrak{C} and interventions in \mathfrak{C} . One of the main contributions of this paper is to clearly distinguish the notions of ‘emulated by’ (Definition 2.5) and ‘interventional validity’ (Definition 2.7); see also Propositions 5.2, 5.5, and 5.8.

6.3 Causal abstraction

Causal abstraction is about transforming one causal model into another causal model. Which constraints such transformations ought to satisfy has been up for debate [Rubenstein et al., 2017, Beckers and Halpern, 2019, Otsuka and Saigo, 2022, Massidda et al., 2023, Otsuka and Saigo, 2024]. The argumentation has been rooted in intuitions about which models can intuitively be considered abstractions of other models. The aim then has been to find mathematical formalizations that capture these intuitions. In this work, we take a different approach: Instead of considering when one model is a ‘valid abstraction’ of another model, we ask when a model is an interventional valid model of a representation. Rather than relying on intuitions about what models ought to count as abstractions of other models, our approach suggests that model transformations and abstractions should preserve or induce interventional validity. In this section, we argue that existing notions of abstraction do not necessarily align with the goal of preserving or inducing interventional validity, and that we need interpretations of actions as interventions to avoid an infinite regress.

Existing notions of abstractions may not preserve interventional validity. The following example shows that τ -abstractions [Beckers and Halpern, 2019] do not preserve interventional validity. We focus on τ -abstractions since this is the strictest notion among those in the literature that considers a restricted set of low-level interventions. It follows that exact transformations, as presented in Rubenstein et al. [2017], also do not preserve interventional validity.

Example 6.5. Transforming an interventional valid model into an invalid model by a τ -abstraction. Consider the SCM $M_{\mathbf{X}}$ given by

$$X_1 := U_1 \sim \text{Unif}([4]).$$

Let \mathfrak{A} be a single-node CBN with observational distribution induced by $M_{\mathbf{X}}$. Assume that \mathbf{X}^* is emulated by \mathfrak{A} and interventions $\mathcal{I}^* = \{\text{do}(X_1 = x) \mid x \in [4]\}$. Since \mathcal{I}^* only has single-node interventions, \mathfrak{A} is an $\mathcal{I} - \mathbf{Int}_S$ valid model of \mathbf{X}^* for every set of interventions \mathcal{I} in \mathfrak{A} , see Proposition 5.5 (1). We now present a τ -abstraction of $(M_{\mathbf{X}}, \mathcal{I}^*)$ that does not preserve interventional validity.

Consider SCM $M_{\mathbf{Y}}$ given by

$$\begin{aligned} Y_1 &:= N_1 \\ Y_2 &:= N_2 \end{aligned}$$

with $N_1, N_2 \stackrel{\text{iid}}{\sim} \text{Unif}(\{0, 1\})$, and let $\tau : [4] \rightarrow \{0, 1\}^2$ be given by

$$x \mapsto (\mathbb{1}(x = 3) + \mathbb{1}(x = 4), \mathbb{1}(x = 2) + \mathbb{1}(x = 4)).$$

τ can be viewed as mapping from the integers $[4]$ to a binary representation of those integers. We can verify that $M_{\mathbf{Y}}$ is a τ -abstraction of $(M_{\mathbf{X}}, \mathcal{I}^*)$ as defined by [Beckers and Halpern \[2019\]](#), see [Appendix E.1](#). Let \mathfrak{C} be a CBN with graph and observational distribution induced by $M_{\mathbf{Y}}$, and let \mathcal{I} be a set of interventions in \mathfrak{C} such that $\text{do}(Y_1 = 0) \in \mathcal{I}$. \mathfrak{C} is compatible with $\mathbf{Y}^* := \tau(X_1)$ but is not an \mathcal{I} - \mathbf{Int}_S valid model of \mathbf{Y}^* . To see this, consider a such that $\mathcal{L}^a(X_1^*) = \mathcal{L}^{\mathfrak{A}; \text{do}(X_1=1)}(\mathbf{X})$ (such an action exists since $\text{do}(X_1 = 1) \in \mathcal{I}^*$). Now, $\text{do}(Y_2 = 0) \in \mathbf{Int}_S^{\mathcal{I}}(a)$ (since $\mathbb{P}^a(Y_2^* = 0) = 1$) but $\mathcal{L}^a(\mathbf{Y}^*) \neq \mathcal{L}^{\mathfrak{C}; \text{do}(Y_2=0)}(\mathbf{Y})$, for example, because $\mathbb{P}^a(Y_1^* = 0) = 1$ while $\mathcal{L}^{\mathfrak{C}; \text{do}(Y_2=0)}(Y_1) = \text{Unif}\{0, 1\}$. \circ

Similarly, in [Appendix E.2](#) we show that interventional validity is not preserved by constructive soft abstractions [[Massidda et al., 2023](#)] under interpretation \mathbf{Int}_P . While it is possible that some notions of abstractions preserve validity for some interpretations, this would be by coincidence rather than per definition. In [Example 6.6](#), we show that we can preserve or induce interventional validity by a transformation that is not a valid τ -abstraction.

Existing notions of abstraction may disallow transforming invalid models into valid models. It may seem puzzling why it would ever be useful to have a high-level model if a low-level model is known. While one motivation may be interpretability, our framework highlights another reason: Maybe we can transform an interventional invalid model into an interventional valid model. We show an example of this below, providing a new formal argument for why high-level models may be preferable to low-level models (for other motivations for high-level causal models see [Hoel et al. \[2013\]](#), [Hoel \[2017\]](#), [Anand et al. \[2023\]](#), [Zennaro et al. \[2024\]](#)).

Example 6.6. Transforming an interventional invalid model into an interventional valid model. Consider the SCM $M_{\mathbf{X}}$ given by

$$\begin{aligned} X_1 &:= U_1 \\ X_2 &:= U_2 \\ X_3 &:= X_1 + X_2 + U_3, \end{aligned}$$

where $U_1, U_2, U_3 \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1)$. Let \mathfrak{A} be a CBN with graph and observational distribution induced by $M_{\mathbf{X}}$. Assume that \mathbf{X}^* is emulated by \mathfrak{A} and interventions $\mathcal{I}^* = \{\text{do}(X_1 = x_1, X_2 = x_2), \text{do}(X_1 = x_1), \text{do}(X_2 = x_2) \mid x_1, x_2 \in \mathbb{R}\}$. \mathfrak{A} is not \mathcal{I}^* - \mathbf{Int}_S valid model of \mathbf{X}^* . To see this, fix $x_1, x_2 \in \mathbb{R}$ and consider an action a such that $\mathcal{L}^a(\mathbf{X}^*) = \mathcal{L}^{\mathfrak{C}; \text{do}(X_1=x_1, X_2=x_2)}(\mathbf{X})$ (which exists since $\text{do}(X_1 = x_1, X_2 = x_2) \in \mathcal{I}^*$). Now $\text{do}(X_1 = x_1) \in \mathbf{Int}_S^{\mathcal{I}^*}(a)$, but $\mathcal{L}^a(\mathbf{X}^*) \neq \mathcal{L}^{\mathfrak{C}; \text{do}(X_1=x_1)}(\mathbf{X})$, for example, because $\mathcal{L}^a(X_2^*)$ has point mass while $\mathcal{L}^{\mathfrak{C}; \text{do}(X_1=x_1)}(X_2)$ is a normal distribution.

Instead, consider now the transformation $\tau : \mathbb{R}^3 \rightarrow \mathbb{R}^2, (x_1, x_2, x_3) \mapsto (x_1 + x_2, x_3)$, and the SCM $M_{\mathbf{Y}}$ given by

$$\begin{aligned} Y_1 &:= \sqrt{2}N_1 \\ Y_2 &:= Y_1 + N_2, \end{aligned}$$

where $N_1, N_2 \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1)$. Let \mathfrak{C} be a CBN with graph and observational distribution induced by $M_{\mathbf{Y}}$. \mathfrak{C} is an $\mathcal{I} - \mathbf{Int}_S$ (and $\mathcal{I} - \mathbf{Int}_P$) valid model of $\tau(\mathbf{X}^*)$ for every set of interventions \mathcal{I} in \mathfrak{C} . But $M_{\mathbf{Y}}$ is not a τ -abstraction of $(M_{\mathbf{X}}, \mathcal{I}^*)$ as defined by [Beckers and Halpern \[2019\]](#), see [Appendix E.3](#). This shows that sometimes a transformation can induce interventional validity without being a τ -abstraction.

If instead $\mathcal{I}^* = \{\text{do}(X_1 = x_1), \text{do}(X_2 = x_2) \mid x_1, x_2 \in \mathbb{R}\}$, then \mathfrak{A} would be an $\mathcal{I} - \mathbf{Int}_S$ (and $\mathcal{I} - \mathbf{Int}_P$) valid model of \mathbf{X}^* for every set of interventions \mathcal{I} in \mathfrak{A} . But $M_{\mathbf{Y}}$ would still not be a τ abstraction of $(M_{\mathbf{X}}, \mathcal{I}^*)$. This shows that sometimes a transformation can preserve interventional validity without being a τ -abstraction. \circ

We think that the perspective of preserving or inducing interventional validity is useful to rigorously ground the notion of valid model transformations. We now argue that the foundation of causal abstraction is dubious without an explicit interpretation.

Causal abstraction rests on an infinite regress. In existing works on abstraction [[Rubenstein et al., 2017](#), [Beckers and Halpern, 2019](#), [Beckers et al., 2020](#), [Rischel and Weichwald, 2021](#), [Massidda et al., 2023](#), [Xia and Bareinboim, 2024](#)] there is a map $\omega : \mathcal{I}_L \rightarrow \mathcal{I}_H$ between interventions in the low-level model and the high-level model. Implicitly, this suggests an interpretation \mathbf{Int} such that for all $a \in \mathcal{A}$, $d_H \in \mathbf{Int}^{\mathcal{I}_H}(a)$ if and only if $d_L \in \mathbf{Int}^{\mathcal{I}_L}(a)$ for some $d_L \in \omega^{-1}(\{d_H\})$. But to determine if $d_L \in \mathbf{Int}^{\mathcal{I}_L}(a)$ we would presumably need yet another model on an even lower level, leading to an infinite regress. The definition of interventions by [Woodward \[2005\]](#) suffers from an analogous problem as explained by [Baumgartner \[2009\]](#). In some concrete applications, this potentially infinite regress may come to a halt at a level of abstraction where there is no ambiguity about which actions constitute interventions. One example of this is the work on causal abstraction of artificial neural networks, where there seems to be no ambiguity about what constitutes an intervention on the level of neuron activations [[Geiger et al., 2021, 2022, 2024a,b](#)]. We think this is the exception rather than the rule; in most applications, there is no level of abstraction where interventions are non-ambiguous. The case of neural networks is peculiar because the network is implemented to literally be a causal model and the structure is given by the network topology. This means that on the level of neurons, we can adopt the circular interpretation \mathbf{Int}_C ([Definition 3.1](#)); it does not matter that other causal models of the neural network are also interventionally valid under interpretation \mathbf{Int}_C because we have prior justification to regard one of them as the causal model. Since this case is an exception, we usually need an interpretation that does not lead to a (potentially infinite) regress. The interpretations presented in this work avoid the regress by depending only on the distribution of the variables on the one given modeling level (and potentially taking the complexity of actions into account).

6.4 Logic of conditionals

In this section, we clarify the connections between causal models and the logic of conditionals. The connection between causal models and (counterfactual) conditionals has received attention from researchers questioning the use of causal models in algorithmic fairness [[Hu and Kohler-Hausmann, 2020](#), [Kasirzadeh and Smart, 2021](#)]. Our work is, as far as we know, the first to spell out a precise connection between interpretations of causal models and different analyses of conditionals. It turns out that considerations from the

philosophical literature on conditionals are relevant for how we interpret causal models, contrary to what is suggested, for example, in Pearl [2009] (see below).

The material condition. In Section 1.2 we considered the proposition

(P) If you intervene $\text{do}(B = 5)$, then you will observe the distribution $\mathcal{N}(0, 1) \otimes \delta_5$ over (A, B) .

Since we assume that Sofia is right that A causes B , the CBN implies $\mathcal{L}^{\text{do}(B=5)}(A, B) = \mathcal{N}(0, 1) \otimes \delta_5$, so (P) should be a true proposition. The proposition has the form of a conditional, that is, a proposition of the form ‘If p , then q ’ [Egré and Rott, 2021]. In mathematics, it is common to interpret conditionals as the material condition ‘ $p \Rightarrow q$ ’, where $p \Rightarrow q \equiv \neg p \vee q$.²⁵ The material condition does not provide the correct analysis of propositions like (P); this can be seen by considering the analogous proposition

(P’) If you intervene $\text{do}(B = 6)$, then you will observe the distribution $\mathcal{N}(0, 1) \otimes \delta_5$ over (A, B) .

Since $\mathcal{L}^{\text{do}(B=6)}(B) = \delta_6 \neq \delta_5$, (P’) should be a false proposition. Furthermore, (P’) should be false regardless of whether anyone intervened $\text{do}(B = 6)$. For example, if someone intervened $\text{do}(B = 7)$ that would not make (P’) true. However, if we analyze (P’) as a material condition, then (P’) is true if you do not perform the intervention $\text{do}(B = 6)$, that is, $p \Rightarrow q$ is true if p is false.

Another way to see that (P’) is not a material condition is that both (P’) and the reverse

(P’’) If you observe the distribution $\mathcal{N}(0, 1) \otimes \delta_5$ over (A, B) , then you had intervened $\text{do}(B = 6)$.

should be false. But this is not possible if (P’) is a material condition since it is a tautology that $(p \Rightarrow q) \vee (q \Rightarrow p)$. These are some of the ‘paradoxes of material conditions’. Since (P’) is apparently not a material condition, by analogy, (P) also is not a material condition.²⁶

Strict implication. An alternative interpretation of propositions like (P) is that they express strict implication [Lewis, 1912, Zach, 2019]. Using the notation of modal logic, strict implication \rightarrow is defined by $p \rightarrow q \equiv \Box(p \Rightarrow q)$ which means that the material condition $p \Rightarrow q$ is true in every ‘accessible world’ [Egré and Rott, 2021]. Consider the interpretation of (P) as (P1):

(P1) If you do something to change the system such that you observe $B = 5$ with probability 1, then you will observe the distribution $\mathcal{N}(0, 1) \otimes \delta_5$ over (A, B) .

We regard the ‘accessible worlds’ as the distributions $\{\mathcal{L}^a(A, B)\}_{a \in \mathcal{A}}$ induced by the available actions \mathcal{A} . If we interpret (P1) as a strict implication with this set of accessible worlds, then (P1) means that

(P-strict) For every $a \in \mathcal{A}$, $(\mathcal{L}^a(B) = \delta_5) \Rightarrow (\mathcal{L}^a(A, B) = \mathcal{N}(0, 1) \otimes \delta_5)$.

²⁵ We use ‘ \Rightarrow ’ rather than the more commonly used ‘ \rightarrow ’ to distinguish from graph notation.

²⁶ That causal statements are not to be analyzed as material conditions has, in the words of Shoham [1990], “been taken into account by all philosophers interested in the subject.”

In the introductory example, see [Section 1.1](#), this turned out to be false since there apparently was an action a such that $\mathcal{L}^a(A, B) = \delta_0 \otimes \delta_5$. Interpreting (P1) as (P-strict) is analogous to interpretation \mathbf{Int}_S . Someone might object that (P1) should not be interpreted as a strict implication and below we provide arguments against interpreting (P1) as (P-strict).

Why (P-strict) is probably not the correct analysis of (P). Assume for the sake of argument that (P-strict) provides the correct analysis of (P). Then, by analogy,

(D-strict) For every $a \in \mathcal{A}$, $(\mathcal{L}^a(A, B) = \delta_0 \otimes \delta_5) \Rightarrow (\mathcal{L}^a(A, B) = \mathcal{N}(0, 1) \otimes \delta_5)$.

provides the correct analysis of

(D) If you intervene $\text{do}(A = 0, B = 5)$, then you will observe the distribution $\mathcal{N}(0, 1) \otimes \delta_5$ over (A, B) .

Since we assume that $A \rightarrow B$, (P) is true. Since we assume that (P-strict) provides the correct analysis of (P), (P-strict) must also be true. (D) is false and since (D-strict) is supposed to provide an analysis of (D), (D-strict) must also be false. But now we have a contradiction since (P-strict) implies (D-strict). This follows from monotonicity of strict implication: $p \rightarrow q \models (p \wedge d) \rightarrow q$. Here, (P-strict) implies (D-strict) because $\mathbb{P}(A = 0, B = 5) = 1$ if and only if $\mathbb{P}(A = 0) = 1 \wedge \mathbb{P}(B = 5) = 1$. Therefore, if (P1) is to provide the correct analysis of (P), we must interpret (P1) as a conditional that does not satisfy monotonicity.

Minimal change semantics. [Stalnaker \[1968\]](#) and [Lewis \[1973\]](#) used this type of argument to show that subjunctive conditionals are generally not strict implications (though this is not undisputed [[von Fintel, 2001](#), [Gillies, 2007](#), [Williamson, 2020](#)]). Here is an example from [Lewis \[1973\]](#):

If Otto had come, it would have been a lively party; but if both Otto and Anna had come, it would have been a dreary party; but if Waldo had come as well, it would have been lively; but ... [[Lewis, 1973](#), Page 10]

Since these sentences are felicitous, they cannot express strict implications. As an alternative [Stalnaker \[1968\]](#) and [Lewis \[1973\]](#) proposed an analysis based on minimal change semantics. In essence, this approach considers a subjunctive conditional to be true if the consequent is true in the closest possible world where the antecedent is true, see [Zach \[2019\]](#) for a simple exposition. If we measure ‘closeness of possible worlds’ by the complexity K of the action, then this approach is analogous to using interpretation \mathbf{Int}_K . In contrast to \mathbf{Int}_S , \mathbf{Int}_K is a non-monotonic interpretation: Consider two interventions $\text{do}(j \leftarrow q_j, j \in J), \text{do}(j \leftarrow q_j, j \in J') \in \mathcal{I}$ with $J \supseteq J'$. Then $\text{do}(j \leftarrow q_j, j \in J) \in \mathbf{Int}_S^{\mathcal{I}}(a)$ implies that $\text{do}(j \leftarrow q_j, j \in J') \in \mathbf{Int}_S^{\mathcal{I}}(a)$. This inference is not valid for \mathbf{Int}_K since the actions that change the conditional distributions for all nodes in J may be more complex than actions that change the conditional distribution only for nodes in J' . The drawback of using an interpretation like \mathbf{Int}_K is that we need a theory of complexity of actions, or, alternatively, of similarity among worlds. This approach is rife with difficulties [[Fine, 1975](#), [Lewis, 1979](#), [Hajek, 2014](#)].²⁷ According to Pearl

²⁷ One seemingly simple way to define similarity of worlds is using the causal model. This is suggested

Such difficulties do not enter the structural account. In contrast with Lewis’s theory, counterfactuals [including interventional claims] are not based on an abstract notion of similarity among hypothetical worlds; instead, they rest directly on the mechanisms [...] that produce those worlds and on the invariant properties of those mechanisms. [...] [S]imilarities and priorities – if they are ever needed – may be read into the $do(\cdot)$ operator as an afterthought [...], but they are not basic to the analysis. [Pearl, 2009, page 239-240]

If one relies on an interpretation like \mathbf{Int}_K these notions are in fact basic to the analysis. Using the $do(\cdot)$ operator to refer both to operations in a mathematical model and to actions in the world, risks obfuscating these fundamental issues, rather than solving them.²⁸

7 Conclusion

Without specifying which real-world actions correspond to which interventions in the mathematical model, it is unclear what it means for a causal Bayesian network to be a valid causal model of a representation. We develop a formal framework for reasoning about the interventional validity of a causal model, which depends on the chosen interpretation of actions as interventions. We discuss different interpretations by considering five desiderata, **D0–D4**, some of which must be violated to escape circularity. Only when the interpretation is made precise can causal models make testable predictions about future observations of a real-world system, which is crucial to enable falsification. We submit that rigorous thinking about the relationship between real-world systems and causal models is critical to facilitate the use of causal models in practice. Otherwise, it is unclear how to use causal models to predict the effect of an action unless the system has already been observed under that very action, echoing the conclusion in “Use and Abuse of Regression” [Box, 1966] that “[t]o find out what happens to a system when you interfere with it you have to interfere with it (not just passively observe it).”

Acknowledgments. LG was supported by the Danish Data Science Academy, which is funded by the Novo Nordisk Foundation (NNF21SA0069429).

in Galles and Pearl [1998], where they write “In essence, causal models define an obvious distance measure among worlds $d(w, w')$, given by the minimal number of local interventions needed for transforming w into w' .” This idea is problematic: If the distances among worlds are given by the causal model, then it amounts to using the circular interpretation \mathbf{Int}_C .

²⁸ One example of this conflation appears in Galles and Pearl [1998]: “[D]efine the action $do(X = x)$ as the minimal change in [the causal model] M required to make $X = x$ [...]”

References

- Kartik Ahuja, Jason Hartford, and Yoshua Bengio. Properties from mechanisms: An equivariance perspective on identifiable representation learning. In *International Conference on Learning Representations*, 2022. (Cited on page 23.)
- Kartik Ahuja, Divyat Mahajan, Yixin Wang, and Yoshua Bengio. Interventional causal representation learning. In *International Conference on Machine Learning*, 2023. (Cited on page 23.)
- John Aldrich. Autonomy. *Oxford Economic Papers*, 41(1):15–34, 1989. (Cited on page 26.)
- Tara V. Anand, Adele H. Ribeiro, Jin Tian, and Elias Bareinboim. Causal effect identification in cluster dags. In *AAAI Conference on Artificial Intelligence*, 2023. (Cited on page 30.)
- Elias Bareinboim, Juan Correa, Duligur Ibeling, and Thomas Icard. On Pearl’s hierarchy and the foundations of causal inference. In *Probabilistic and Causal Inference: the Works of Judea Pearl*, pages 507–556. ACM Books, 2022. (Cited on pages 3 and 10.)
- Michael Baumgartner. Interdefining causation and intervention. *Dialectica*, 63(2):175–194, 2009. (Cited on pages 10 and 31.)
- Sander Beckers and Joseph Y. Halpern. Abstracting causal models. In *AAAI Conference on Artificial Intelligence*, 2019. (Cited on pages 29, 30, 31, 47, and 48.)
- Sander Beckers, Frederick Eberhardt, and Joseph Y. Halpern. Approximate causal abstractions. In *Uncertainty in Artificial Intelligence*, 2020. (Cited on page 31.)
- Yoshua Bengio, Aaron Courville, and Pascal Vincent. Representation learning: A review and new perspectives. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 35(8):1798–1828, 2013. (Cited on page 26.)
- Simon Bing, Tom Hochsprung, Jonas Wahl, Urmi Ninad, and Jakob Runge. Invariance & causal representation learning: Prospects and limitations. *Transactions on Machine Learning Research*, 2024. (Cited on page 27.)
- George Box. Use and abuse of regression. *Technometrics*, 8(4):625–629, 1966. (Cited on page 34.)
- Johann Brehmer, Pim De Haan, Phillip Lippe, and Taco S. Cohen. Weakly supervised causal representation learning. In *Advances in Neural Information Processing Systems*, 2022. (Cited on page 23.)
- Simon Buchholz, Goutham Rajendran, Elan Rosenfeld, Bryon Aragam, Bernhard Schölkopf, and Pradeep Ravikumar. Learning linear causal representations from interventions under general nonlinear mixing. In *Advances in Neural Information Processing Systems*, 2024. (Cited on page 23.)
- Peter Bühlmann. Invariance, causality and robustness. *Statistical Science*, 35(3):404–426, 2020. (Cited on page 24.)
- Riccardo Cadei, Lukas Lindorfer, Sylvia Cremer, Cordelia Schmid, and Francesco Locatello.

- Smoke and mirrors in causal downstream tasks. *Preprint arXiv:2402.12980*, 2024. (Cited on page 27.)
- Nancy Cartwright. *Hunting causes and using them: Approaches in philosophy and economics*. Cambridge University Press, 2007. (Cited on page 26.)
- Krzysztof Chalupka, Pietro Perona, and Frederick Eberhardt. Visual causal feature learning. In *Uncertainty in Artificial Intelligence*, 2015. (Cited on page 27.)
- Alexander Mangulad Christgau and Niels Richard Hansen. Efficient adjustment for complex covariates: Gaining efficiency with DOPE. *Preprint arXiv:2402.12980*, 2024. (Cited on page 27.)
- Taco Cohen. Towards a grounded theory of causation for embodied AI. *Preprint arXiv:2206.13973*, 2022. (Cited on page 25.)
- Imant Daunhawer, Alice Bizeul, Emanuele Palumbo, Alexander Marx, and Julia E. Vogt. Identifiability results for multimodal contrastive learning. In *International Conference on Learning Representations*, 2023. (Cited on page 23.)
- Philip Dawid. Causal inference without counterfactuals. *Journal of the American statistical Association*, 95(450):407–424, 2000. (Cited on page 3.)
- Philip Dawid. Beware of the DAG! In *Causality: Objectives and Assessment*, pages 59–86, 2010. (Cited on page 27.)
- Philip Dawid. Decision-theoretic foundations for statistical causality. *Journal of Causal Inference*, 9(1), 2021. (Cited on pages 19 and 26.)
- Joel Dyer, Nicholas George Bishop, Yorgos Felekis, Fabio Massimo Zennaro, Ani Calinescu, Theodoros Damoulas, and Michael J. Wooldridge. Interventionally consistent surrogates for complex simulation models. In *Advances in Neural Information Processing Systems*, 2024. (Cited on page 27.)
- Frederick Eberhardt. Green and grue causal variables. *Synthese*, 193:1029–1046, 2016. (Cited on page 8.)
- Frederick Eberhardt, Clark Glymour, and Richard Scheines. N-1 experiments suffice to determine the causal relations among n variables. *Innovations in Machine Learning: Theory and Applications*, pages 97–112, 2006. (Cited on page 29.)
- Paul Egré and Hans Rott. The Logic of Conditionals. In *The Stanford Encyclopedia of Philosophy*. Metaphysics Research Lab, Stanford University, 2021. (Cited on page 32.)
- Kit Fine. Critical notice of lewis’ counterfactuals. *Mind*, 84:451–458, 1975. (Cited on page 33.)
- David Galles and Judea Pearl. An axiomatic characterization of causal counterfactuals. *Foundations of Science*, 3:151–182, 1998. (Cited on pages 10 and 34.)
- Atticus Geiger, Hanson Lu, Thomas Icard, and Christopher Potts. Causal abstractions of neural networks. In *Advances in Neural Information Processing Systems*, 2021. (Cited on page 31.)

- Atticus Geiger, Zhengxuan Wu, Hanson Lu, Josh Rozner, Elisa Kreiss, Thomas Icard, Noah Goodman, and Christopher Potts. Inducing causal structure for interpretable neural networks. In *International Conference on Machine Learning*, 2022. (Cited on page 31.)
- Atticus Geiger, Duligur Ibeling, Amir Zur, Maheep Chaudhary, Sonakshi Chauhan, Jing Huang, Aryaman Arora, Zhengxuan Wu, Noah Goodman, Christopher Potts, et al. Causal abstraction: A theoretical foundation for mechanistic interpretability. *Preprint arXiv:2301.04709*, 2024a. (Cited on page 31.)
- Atticus Geiger, Zhengxuan Wu, Christopher Potts, Thomas Icard, and Noah Goodman. Finding alignments between interpretable causal variables and distributed neural representations. In *Causal Learning and Reasoning*, 2024b. (Cited on page 31.)
- Anthony S. Gillies. Counterfactual scorekeeping. *Linguistics and Philosophy*, 30:329–360, 2007. (Cited on page 33.)
- Clark Glymour and Madelyn R Glymour. Commentary: race and sex are causes. *Epidemiology*, 2014. (Cited on page 26.)
- Limor Gultchin, David Watson, Matt Kusner, and Ricardo Silva. Operationalizing complex causes: A pragmatic view of mediation. In *International Conference on Machine Learning*, 2021. (Cited on page 27.)
- Alan Hajek. *Most Counterfactuals Are False*. 2014. URL <https://philarchive.org/rec/HJEMCA>. (Cited on page 33.)
- Erik P. Hoel. When the map is better than the territory. *Entropy*, 19(5):188, 2017. (Cited on page 30.)
- Erik P. Hoel, Larissa Albantakis, and Giulio Tononi. Quantifying causal emergence shows that macro can beat micro. *Proceedings of the National Academy of Sciences*, 110(49):19790–19795, 2013. (Cited on page 30.)
- Lily Hu and Issa Kohler-Hausmann. What's sex got to do with machine learning? In *Conference on Fairness, Accountability, and Transparency*, 2020. (Cited on page 31.)
- Duligur Ibeling and Thomas Icard. Probabilistic reasoning across the causal hierarchy. In *AAAI Conference on Artificial Intelligence*, 2020. (Cited on page 3.)
- Duligur Ibeling and Thomas Icard. Comparing causal frameworks: Potential outcomes, structural models, graphs, and abstractions. In *Advances in Neural Information Processing Systems*, 2023. (Cited on page 10.)
- Dominik Janzing and Sergio H. G. Mejia. A phenomenological account for causality in terms of elementary actions. *Journal of Causal Inference*, 12(1), 2024. (Cited on pages 10, 23, 26, 49, and 50.)
- Dominik Janzing and Bernhard Schölkopf. Causal inference using the algorithmic Markov condition. *IEEE Transactions on Information Theory*, 56(10):5168–5194, 2010. (Cited on page 19.)
- Dominik Janzing, Rafael Chaves, and Bernhard Schölkopf. Algorithmic independence

- of initial condition and dynamical law in thermodynamics and causal inference. *New Journal of Physics*, 18(9):093052, 2016. (Cited on page 19.)
- Jikai Jin and Vasilis Syrgkanis. Learning causal representations from general environments: Identifiability and intrinsic ambiguity. In *Advances in Neural Information Processing Systems*, 2024. (Cited on page 23.)
- Atoosa Kasirzadeh and Andrew Smart. The use and misuse of counterfactuals in ethical machine learning. In *Conference on Fairness, Accountability, and Transparency*, 2021. (Cited on page 31.)
- Armin Kekić, Bernhard Schölkopf, and Michel Besserve. Targeted reduction of causal models. In *Uncertainty in Artificial Intelligence*, 2024. (Cited on page 27.)
- David Kinney and David Watson. Causal feature learning for utility-maximizing agents. In *International Conference on Probabilistic Graphical Models*, pages 257–268, 2020. (Cited on page 27.)
- Bohdan Kivva, Goutham Rajendran, Pradeep Ravikumar, and Bryon Aragam. Learning latent causal graphs via mixture oracles. In *Advances in Neural Information Processing Systems*, 2021. (Cited on page 23.)
- Sébastien Lachapelle, Pau Rodriguez, Yash Sharma, Katie E Everett, Rémi Le Priol, Alexandre Lacoste, and Simon Lacoste-Julien. Disentanglement via mechanism sparsity regularization: A new principle for nonlinear ICA. In *Causal Learning and Reasoning*, 2022. (Cited on page 23.)
- Sébastien Lachapelle, Tristan Deleu, Divyat Mahajan, Ioannis Mitliagkas, Yoshua Bengio, Simon Lacoste-Julien, and Quentin Bertrand. Synergies between disentanglement and sparsity: Generalization and identifiability in multi-task learning. In *International Conference on Machine Learning*, 2023. (Cited on page 27.)
- Steffen Lauritzen. *Lectures on Graphical Models, 3rd edition*. Department of Mathematical Sciences, Faculty of Science, University of Copenhagen, 2019. ISBN 978-87-70787-53-6. (Cited on page 50.)
- Clarence Irving Lewis. Implication and the algebra of logic. *Mind*, 21(84):522–531, 1912. (Cited on page 32.)
- David Lewis. *Counterfactuals*. Blackwell, 1973. (Cited on pages 19 and 33.)
- David Lewis. Counterfactual dependence and time’s arrow. *Noûs*, 13(4):455–476, 1979. (Cited on page 33.)
- Adam Li, Yushu Pan, and Elias Bareinboim. Disentangled representation learning in non-Markovian causal systems. In *Advances in Neural Information Processing Systems*, 2024. (Cited on page 23.)
- Phillip Lippe, Sara Magliacane, Sindy Löwe, Yuki M Asano, Taco Cohen, and Stratis Gavves. Citris: Causal identifiability from temporal intervened sequences. In *International Conference on Machine Learning*, 2022. (Cited on page 23.)
- Joshua Loftus. Position: The causal revolution needs scientific pragmatism. In *International Conference on Machine Learning*, 2024. (Cited on page 4.)

- Emanuele Marconato, Sébastien Lachapelle, Sebastian Weichwald, and Luigi Gresele. All or none: Identifiable linear properties of next-token predictors in language modeling. *Preprint arXiv:2410.23501*, 2024. (Cited on page 27.)
- Riccardo Massidda, Atticus Geiger, Thomas Icard, and Davide Bacciu. Causal abstraction with soft interventions. In *Causal Learning and Reasoning*, 2023. (Cited on pages 29, 30, 31, 47, and 48.)
- Jun Otsuka and Hayato Saigo. On the equivalence of causal models: A category-theoretic approach. In *Causal Learning and Reasoning*, pages 634–646, 2022. (Cited on page 29.)
- Jun Otsuka and Hayato Saigo. Process theory of causality: a category-theoretic perspective. *Behaviormetrika*, 51(1):21–36, 2024. (Cited on page 29.)
- Junhyung Park, Simon Buchholz, Bernhard Schölkopf, and Krikamol Muandet. A measure-theoretic axiomatisation of causality. In *Advances in Neural Information Processing Systems*, 2023. (Cited on page 10.)
- Judea Pearl. *Causality*. Cambridge University Press, second edition, 2009. (Cited on pages 1, 5, 32, and 34.)
- Judea Pearl. Does obesity shorten life? Or is it the soda? On non-manipulable causes. *Journal of Causal Inference*, 6(2), 2018. (Cited on page 26.)
- Judea Pearl. On the interpretation of do(x). *Journal of Causal Inference*, 2019. (Cited on page 26.)
- Judea Pearl, Madelyn Glymour, and Nicholas P Jewell. *Causal inference in statistics: A primer*. John Wiley & Sons, 2016. (Cited on page 3.)
- Jonas Peters, Dominik Janzing, and Bernhard Schölkopf. *Elements of Causal Inference: Foundations and Learning Algorithms*. Adaptive Computation and Machine Learning. MIT Press, 2017. (Cited on pages 1, 4, 5, and 20.)
- Arvind Raghavan and Elias Bareinboim. Counterfactual realizability. Technical Report R-113, Causal Artificial Intelligence Lab, Columbia University, 2024. (Cited on page 4.)
- Eigil F. Rischel and Sebastian Weichwald. Compositional abstraction error and a category of causal models. In *Uncertainty in Artificial Intelligence*, 2021. (Cited on page 31.)
- Paul K. Rubenstein, Sebastian Weichwald, Stephan Bongers, Joris M. Mooij, Dominik Janzing, Moritz Grosse-Wentrup, and Bernhard Schölkopf. Causal consistency of structural equation models. In *Uncertainty in Artificial Intelligence*, 2017. (Cited on pages 8, 29, and 31.)
- Kayvan Sadeghi and Terry Soo. Axiomatization of interventional probability distributions. *Biometrika*, 2024. (Cited on page 25.)
- Sorawit Saengkyongam, Elan Rosenfeld, Pradeep Kumar Ravikumar, Niklas Pfister, and Jonas Peters. Identifying representations for intervention extrapolation. In *International Conference on Learning Representations*, 2024. (Cited on page 27.)

- Bernhard Schölkopf, Francesco Locatello, Stefan Bauer, Nan Rosemary Ke, Nal Kalchbrenner, Anirudh Goyal, and Yoshua Bengio. Toward causal representation learning. *Proceedings of the IEEE*, 109(5):612–634, 2021. (Cited on page 6.)
- Yoav Shoham. Nonmonotonic reasoning and causation. *Cognitive Science*, 14(2):213–252, 1990. (Cited on page 32.)
- Ilya Shpitser and Judea Pearl. What counterfactuals can be tested. In *Uncertainty in Artificial Intelligence*, 2007. (Cited on page 3.)
- Peter Spirtes and Richard Scheines. Causal inference of ambiguous manipulations. *Philosophy of Science*, 71:833 – 845, 2004. (Cited on pages 8 and 15.)
- Peter Spirtes, Clark Glymour, and Richard Scheines. *Causation, prediction, and search*. MIT press, 2001. (Cited on pages 1 and 5.)
- Chandler Squires, Anna Seigal, Salil S Bhate, and Caroline Uhler. Linear causal disentanglement via interventions. In *International Conference on Machine Learning*, 2023. (Cited on pages 23 and 25.)
- Robert Stalnaker. A theory of conditionals. In *Studies in Logical Theory*, pages 98–112. Blackwell, 1968. (Cited on page 33.)
- Burak Varici, Emre Acartürk, Karthikeyan Shanmugam, and Ali Tajer. General identifiability and achievability for causal representation learning. In *International Conference on Artificial Intelligence and Statistics*, 2024. (Cited on page 23.)
- Kai von Fintel. Counterfactuals in a Dynamic Context. In *Ken Hale: A Life in Language*. The MIT Press, 2001. (Cited on page 33.)
- Julius von Kügelgen, Yash Sharma, Luigi Gresele, Wieland Brendel, Bernhard Schölkopf, Michel Besserve, and Francesco Locatello. Self-supervised learning with data augmentations provably isolates content from style. In *Advances in Neural Information Processing Systems*, 2021. (Cited on page 23.)
- Julius von Kügelgen, Michel Besserve, Liang Wendong, Luigi Gresele, Armin Kekić, Elias Bareinboim, David Blei, and Bernhard Schölkopf. Nonparametric identifiability of causal representations from unknown interventions. In *Advances in Neural Information Processing Systems*, 2024. (Cited on pages 23 and 24.)
- Sebastian Weichwald, Moritz Grosse-Wentrup, and Arthur Gretton. MERLiN: Mixture Effect Recovery in Linear Networks. *IEEE Journal of Selected Topics in Signal Processing*, 10(7):1254–1266, 2016. (Cited on page 27.)
- Ryan Welch, Jiaqi Zhang, and Caroline Uhler. Identifiability guarantees for causal disentanglement from purely observational data. In *Advances in Neural Information Processing Systems*, 2024. (Cited on page 23.)
- Liang Wendong, Armin Kekić, Julius von Kügelgen, Simon Buchholz, Michel Besserve, Luigi Gresele, and Bernhard Schölkopf. Causal component analysis. In *Advances in Neural Information Processing Systems*, 2024. (Cited on page 23.)
- Timothy Williamson. *Suppose and Tell: The Semantics and Heuristics of Conditionals*. Oxford University Press, 2020. (Cited on page 33.)

- James Woodward. *Making things happen: A theory of causal explanation*. Oxford university press, 2005. (Cited on page 31.)
- James Woodward. The problem of variable choice. *Synthese*, 193:1047–1072, 2016. (Cited on page 27.)
- James Woodward. Causation and Manipulability. In *The Stanford Encyclopedia of Philosophy*. Metaphysics Research Lab, Stanford University, 2023. (Cited on page 10.)
- Kevin Xia and Elias Bareinboim. Neural causal abstractions. In *AAAI Conference on Artificial Intelligence*, 2024. (Cited on page 31.)
- Feng Xie, Ruichu Cai, Biwei Huang, Clark Glymour, Zhifeng Hao, and Kun Zhang. Generalized independent noise condition for estimating latent variable causal graphs. In *Advances in Neural Information Processing Systems*, 2020. (Cited on page 23.)
- Danru Xu, Dingling Yao, Sebastien Lachapelle, Perouz Taslakian, Julius von Kügelgen, Francesco Locatello, and Sara Magliacane. A sparsity principle for partially observable causal representation learning. In *International Conference on Machine Learning*, 2024. (Cited on page 23.)
- Dingling Yao, Danru Xu, Sebastien Lachapelle, Sara Magliacane, Perouz Taslakian, Georg Martius, Julius von Kügelgen, and Francesco Locatello. Multi-view causal representation learning with partial observability. In *International Conference on Learning Representations*, 2024. (Cited on page 23.)
- Richard Zach. *Boxes and Diamonds: An Open Introduction to Modal Logic*. Open Logic Project, 2019. (Cited on pages 32 and 33.)
- Fabio Massimo Zennaro, Nicholas George Bishop, Joel Dyer, Yorgos Felekis, Ani Calinescu, Michael J. Wooldridge, and Theodoros Damoulas. Causally abstracted multi-armed bandits. In *Uncertainty in Artificial Intelligence*, 2024. (Cited on page 30.)
- Jiaqi Zhang, Kristjan Greenewald, Chandler Squires, Akash Srivastava, Karthikeyan Shanmugam, and Caroline Uhler. Identifiability guarantees for causal disentanglement from soft interventions. In *Advances in Neural Information Processing Systems*, 2024. (Cited on page 23.)
- Yuchen Zhu, Kailash Budhathoki, Jonas M Kübler, and Dominik Janzing. Meaningful causal aggregation and paradoxical confounding. In *Causal Learning and Reasoning*, 2024. (Cited on page 23.)

A Notation

Symbol	Description
\mathcal{A}	A set of actions, see Definition 2.3 .
\mathcal{O}	$\mathcal{O} \in \mathcal{A}$ denotes the action corresponding to the observational regime, see Definition 2.3 .
\mathcal{G}	Directed acyclic graph over nodes $\{1, \dots, n\}$.
$[n]$	The set $\{1, \dots, n\}$.
$\mathfrak{C}, \mathfrak{A}, \mathfrak{h}$	Causal Bayesian networks (CBNs), see Definition 2.1 .
h	Function $\mathbb{R}^m \rightarrow \mathbb{R}^n$ transforming low-level features, see Definition 2.4 .
\mathbf{Z}	Multivariate random variable of CBN variables $\mathbf{Z} \in \mathbb{R}^n$.
\mathbf{X}^*	Multivariate random variable of low-level features, $\mathbf{X}^* \in \mathbb{R}^m$, see Definition 2.3 .
\mathbf{Z}^*	Multivariate representation of a data-generating process, $\mathbf{Z}^* = h(\mathbf{X}^*) \in \mathbb{R}^n$, see Definition 2.4 .
\mathcal{L}	Denotes a distribution. For example, $\mathcal{L}^a(h(\mathbf{X}^*))$ is the distribution induced by action $a \in \mathcal{A}$ over $h(\mathbf{X}^*)$, see Definition 2.4 , and $\mathcal{L}^{\mathfrak{C};d}(\mathbf{Z})$ denotes the distribution induced by CBN \mathfrak{C} over variables \mathbf{Z} and intervention d , see Definition 2.1 .
$\mathcal{L}^a(Z_j^* \mathbf{Y}^*) \sim p$	Denotes the claim that the kernel $\mathbf{y}^* \mapsto p(\cdot \mathbf{y}^*) \cdot \nu$ is a regular conditional probability distribution of Z_j^* given $\mathbf{Y}^* \subseteq \mathbf{Z}^*$ under distribution $\mathcal{L}^a(\mathbf{Z}^*)$ (for some fixed σ -finite measure ν , which is suppressed in the notation).
\mathcal{I}	Set of interventions in a CBN.
\mathcal{I}^*	Set of interventions in a CBN. Used to denote interventions in a CBN that emulate a representation, see Definition 2.5 .
$\text{do}(j \leftarrow q_j, j \in J)$	An intervention on nodes J , see Definition 2.1 .
d, d^*	Denotes interventions, $d \in \mathcal{I}$, $d^* \in \mathcal{I}^*$.
$p_i^{\mathfrak{C}}, p_i^{\mathfrak{C};d}$	$p_i^{\mathfrak{C}}$ denotes the i 'th kernel given by CBN \mathfrak{C} . $p_i^{\mathfrak{C};d}$ denotes i 'th kernel given by CBN \mathfrak{C} and intervention $d = \text{do}(j \leftarrow q_j, j \in J)$, that is, $p_i^{\mathfrak{C};d} = p_i^{\mathfrak{C}}$ for $i \notin J$, and $p_i^{\mathfrak{C};d} = q_i$ for $i \in J$.
\mathbf{Int}	An interpretation that takes a set of interventions \mathcal{I} in a CBN and an action $a \in \mathcal{A}$ and outputs a subset $\mathbf{Int}^{\mathcal{I}}(a)$ of \mathcal{I} , see Definition 2.6 and Table 1 .
$\mathbf{Int}^{\mathcal{I}}$	Given a set of interventions \mathcal{I} in a CBN, an interpretation induces a mapping $\mathbf{Int}^{\mathcal{I}} : \mathcal{A} \rightarrow \mathcal{P}(\mathcal{I})$, see Definition 2.6 and Table 1 .
$\mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$	Joint normal distribution with mean $\boldsymbol{\mu}$ and covariance $\boldsymbol{\Sigma}$.
δ_x	Dirac distribution with support $\{x\}$ ($x \in \mathbb{R}$).
$\text{Unif}(S)$	Uniform distribution over finite set S .
$\text{Ber}(p)$	Bernoulli distribution with mean p .
\otimes	$\nu_1 \otimes \nu_2$ denotes the product measure of two σ -finite measures ν_1 and ν_2 .

B Interpretation Int_P satisfies D2

Proposition B.1. Int_P satisfies desideratum D2.

Proof. Let a data-generating process \mathcal{D} , representation \mathbf{Z}^* , compatible CBN \mathfrak{C} , and set of interventions \mathcal{I} in \mathfrak{C} be given.

Let action $a \in \mathcal{A}$ be arbitrary. We want to show that every intervention in $\text{Int}_P^{\mathcal{I}}(a)$ induces the same distribution.

Let $\tilde{J} = \{i \in [n] \mid Z_i^* \perp\!\!\!\perp \mathbf{PA}_i^* \text{ in } \mathcal{L}^a(\mathbf{Z}^*)\}$, and let $b = \text{do}(j \leftarrow \tilde{q}_j, j \in \tilde{J})$, be a perfect intervention where each \tilde{q}_j is given such that $\mathcal{L}^a(Z_i^* \mid \mathbf{PA}_i^*) \sim \tilde{q}_i$ for every $i \in \tilde{J}$.

Let $d = \text{do}(j \leftarrow q_j, j \in J) \in \text{Int}_P^{\mathcal{I}}(a)$ be given arbitrarily (if $\text{Int}_P^{\mathcal{I}}(a)$ is empty, there is nothing to show). We now argue that $\mathcal{L}^{\mathfrak{C};d}(\mathbf{Z}) = \mathcal{L}^{\mathfrak{C};b}(\mathbf{Z})$. For every $i \in J$, since d is perfect and sets correct conditionals on intervened nodes (**D0**), we have that $Z_i^* \perp\!\!\!\perp \mathbf{PA}_i^*$ in $\mathcal{L}^a(\mathbf{Z}^*)$ (so $i \in \tilde{J}$), and thus $p_i^{\mathfrak{C};b}$ is also perfect. So for $i \in J$, both $p_i^{\mathfrak{C};b}$ and $p_i^{\mathfrak{C};d}$ are the constant kernel $\mathbf{pa}_i \mapsto \mathcal{L}^a(Z_i)$, and therefore $\mathcal{L}^{\mathfrak{C};d}(Z_i \mid \mathbf{PA}_i) \sim p_i^{\mathfrak{C};b}$.

Since d satisfies condition 3) of [Definition 5.1](#) (and by the definition of \tilde{J}), every $i \in \tilde{J} \setminus J$ is a source node and $\mathcal{L}^{\mathfrak{C};b}(Z_i) = \mathcal{L}^{\mathfrak{C}}(Z_i)$ which, since d is not intervening on $i \in \tilde{J} \setminus J$, implies $\mathcal{L}^{\mathfrak{C};d}(Z_i) = \mathcal{L}^{\mathfrak{C};b}(Z_i)$, so $\mathcal{L}^{\mathfrak{C};d}(Z_i \mid \mathbf{PA}_i) \sim p_i^{\mathfrak{C};b}$ for every $i \in \tilde{J} \setminus J$.

Finally, for every $i \notin \tilde{J}$, we also have $\mathcal{L}^{\mathfrak{C};d}(Z_i \mid \mathbf{PA}_i) \sim p_i^{\mathfrak{C};b}$ since neither d nor b can be interventions on nodes outside of \tilde{J} as they are perfect interventions. Therefore, $\mathcal{L}^{\mathfrak{C};d}(\mathbf{Z}) = \mathcal{L}^{\mathfrak{C};b}(\mathbf{Z})$, and since $d \in \text{Int}_P^{\mathcal{I}}(a)$ was arbitrary, we conclude that every intervention in $\text{Int}_P^{\mathcal{I}}(a)$ must induce the same distribution, namely $\mathcal{L}^{\mathfrak{C};b}(\mathbf{Z})$. \square

C Counterexamples regarding [Footnote 14](#)

Consider interpretation $\text{Int}_{\tilde{S}}$ defined exactly as Int_S in [Definition 5.4](#), except that we drop condition 2). We now argue that [Proposition 5.5](#) (1) is invalid for $\text{Int}_{\tilde{S}}$.

Example C.1. $\text{Int}_{\tilde{S}}$ violates [Proposition 5.5](#) (1). Let \mathfrak{C} be given by graph $X \rightarrow Y$ and kernels:

$$\begin{aligned}\mathcal{L}^{\mathfrak{C}}(X) &= \text{Ber}(0.5) \\ \mathcal{L}^{\mathfrak{C}}(Y \mid X = 0) &= \text{Ber}(0.4) \\ \mathcal{L}^{\mathfrak{C}}(Y \mid X = 1) &= \text{Ber}(0.6).\end{aligned}$$

Assume that (X^*, Y^*) is emulated by \mathfrak{C} and interventions $\mathcal{I}^* = \{d^*\}$, where d^* is a single-node intervention on X given by

$$\mathcal{L}^{\mathfrak{C};d^*}(X) = \delta_0.$$

Now, let $\mathcal{I} = \{d\}$, where d is a single-node intervention on Y given by

$$\begin{aligned}\mathcal{L}^{\mathfrak{C};d}(Y \mid X = 0) &= \text{Ber}(0.4) \\ \mathcal{L}^{\mathfrak{C};d}(Y \mid X = 1) &= \text{Ber}(0.7).\end{aligned}$$

Let a be an action such that $\mathcal{L}^a(X^*, Y^*) = \mathcal{L}^{\mathfrak{C}; d^*}(X, Y)$. $d \in \mathbf{Int}_{\tilde{\mathcal{S}}}^{\mathcal{I}}(a)$ since $\mathcal{L}^a(Y^* | X^*) \sim p_{Y^*|X^*}^{\mathfrak{C}; d}$, but $\mathcal{L}^a(X^*, Y^*) \neq \mathcal{L}^{\mathfrak{C}; d}(X, Y)$, contradicting [Proposition 5.5](#) (1). Notice that $d \notin \mathbf{Int}_{\tilde{\mathcal{S}}}^{\mathcal{I}}(a)$ since $\mathcal{L}^a(Y^* | X^*) \sim p_{Y^*|X^*}^{\mathfrak{C}}$, violating condition 2) of [Definition 5.4](#). \circ

D Further non-circular interpretations ([Section 5](#))

D.1 $\mathbf{Int}_{\tilde{\mathcal{I}}, f}^{\mathcal{I}}$: Violating only **D3**

We provide an interpretation that satisfies **D0–D4**, except **D3**. We do not expect that this interpretation will be useful in itself; rather, we provide it to show that **D0**, **D1**, **D2**, and **D4** do not imply **D3**. We leave it for future work to investigate if there exist interesting interpretations that may violate **D3**.

Definition D.1. $\mathbf{Int}_{\tilde{\mathcal{I}}, f}^{\mathcal{I}}$. **An interpretation violating only **D3**.** Let a data-generating process \mathcal{D} , representation \mathbf{Z}^* , compatible CBN \mathfrak{C} , countable set of interventions $\tilde{\mathcal{I}}$ in \mathfrak{C} , and an injective function $f : \tilde{\mathcal{I}} \rightarrow \mathbb{N}$ be given. In addition, let a set of modeled interventions \mathcal{I} in \mathfrak{C} be given. We define the interpretation $\mathbf{Int}_{\tilde{\mathcal{I}}, f}^{\mathcal{I}}$ by the following rule: An intervention $d \in \mathcal{I}$ is in $\mathbf{Int}_{\tilde{\mathcal{I}}, f}^{\mathcal{I}}(a)$ if and only if

- a) $d \in \mathbf{Int}_{\mathcal{C}}^{\mathcal{I}}(a)$,
- or
- b) the following two conditions hold:
 - 1) $d \in \tilde{\mathcal{I}} \cap \mathbf{Int}_{\tilde{\mathcal{S}}}^{\mathcal{I}}(a)$, see [Definition 5.4](#).
 - 2) For every $b \in \mathcal{I} \setminus \{d\}$, if $b \in \tilde{\mathcal{I}} \cap \mathbf{Int}_{\tilde{\mathcal{S}}}^{\mathcal{I}}(a)$, then $f(b) > f(d)$.

\circ

$\mathbf{Int}_{\mathcal{C}}^{\mathcal{I}}(a) \subseteq \mathbf{Int}_{\tilde{\mathcal{I}}, f}^{\mathcal{I}}(a)$ such that **D1** (if it behaves like an intervention, it is that intervention) is satisfied. To ensure that $\mathbf{Int}_{\tilde{\mathcal{I}}, f}^{\mathcal{I}}(a)$ is not generally a subset of $\mathbf{Int}_{\mathcal{C}}^{\mathcal{I}}(a)$, we also let $d \in \mathcal{I}$ be in $\mathbf{Int}_{\tilde{\mathcal{I}}, f}^{\mathcal{I}}(a)$ if two conditions are satisfied. Informally, this works as follows: For an action a , we check which interventions are in $\tilde{\mathcal{I}} \cap \mathbf{Int}_{\tilde{\mathcal{S}}}^{\mathcal{I}}(a) \subseteq \mathcal{I}$. If $\tilde{\mathcal{I}} \cap \mathbf{Int}_{\tilde{\mathcal{S}}}^{\mathcal{I}}(a)$ is nonempty, we interpret a as the least element in $\tilde{\mathcal{I}} \cap \mathbf{Int}_{\tilde{\mathcal{S}}}^{\mathcal{I}}(a)$, using f as the ordering. This ensures that $\mathbf{Int}_{\tilde{\mathcal{I}}, f}^{\mathcal{I}}$ does not violate **D2** (an action should not be interpreted as distinct interventions). $\mathbf{Int}_{\tilde{\mathcal{I}}, f}^{\mathcal{I}}(a)$ satisfies **D0** (correct conditionals on intervened nodes) and **D4** (an intervention does not create new dependencies) since $\mathbf{Int}_{\tilde{\mathcal{S}}}$ and $\mathbf{Int}_{\mathcal{C}}$ satisfy these desiderata. On the other hand, $\mathbf{Int}_{\tilde{\mathcal{I}}, f}^{\mathcal{I}}$ may violate **D3** (interpretations should not depend on the intervention set \mathcal{I}), as shown in the following example.

Example D.2. **Interpretation $\mathbf{Int}_{\tilde{\mathcal{I}}, f}^{\mathcal{I}}$ may violate **D3**.** Let \mathfrak{C} be a CBN whose DAG has no edges and with kernels given by

$$\begin{aligned} Z_1 &\sim \mathcal{N}(0, 1) \\ Z_2 &\sim \mathcal{N}(0, 1). \end{aligned}$$

Let $\mathcal{I}^* = \{\text{do}(Z_1 = 0, Z_2 = 0)\}$, $\tilde{\mathcal{I}} = \{\text{do}(Z_1 = 0), \text{do}(Z_1 = 0, Z_2 = 0)\}$, and define $f : \tilde{\mathcal{I}} \rightarrow \mathbb{N}$ by $f(\text{do}(Z_1 = 0, Z_2 = 0)) = 1$ and $f(\text{do}(Z_1 = 0)) = 2$. Assume that (Z_1^*, Z_2^*) is emulated by \mathfrak{C} and \mathcal{I}^* . Consider an action $a \in \mathcal{A} \setminus \{O\}$; since \mathcal{I}^* has only one element, $\mathcal{L}^a(\mathbf{Z}^*) = \mathcal{L}^{\mathfrak{C}; \text{do}(Z_1=0, Z_2=0)}(\mathbf{Z})$. Now $\text{do}(Z_1 = 0) \in \mathbf{Int}_{\tilde{\mathcal{I}}, f}^{\{\text{do}(Z_1=0)\}}(a)$, but $\text{do}(Z_1 = 0) \notin \mathbf{Int}_{\tilde{\mathcal{I}}, f}^{\{\text{do}(Z_1=0), \text{do}(Z_1=0, Z_2=0)\}}(a) = \{\text{do}(Z_1 = 0, Z_2 = 0)\}$, contradicting **D3**. This example also shows that it is possible to falsify a compatible model under $\mathbf{Int}_{\tilde{\mathcal{I}}, f}$ since $\text{do}(Z_1 = 0) \in \mathbf{Int}_{\tilde{\mathcal{I}}, f}^{\{\text{do}(Z_1=0)\}}(a)$ but $\mathcal{L}^a(Z_2^*) = \delta_0 \neq \mathcal{N}(0, 1) = \mathcal{L}^{\mathfrak{C}; \text{do}(Z_1=0)}(Z_2)$. \circ

D.2 \mathbf{Int}_M : Violating only D4 renders all CBNs with complete DAGs interventionally valid

Assume that we have a representation \mathbf{Z}^* and a compatible and complete CBN \mathfrak{C} . If \mathbf{Int} is an interpretation that satisfies **D1–D3**, then \mathfrak{C} is an $\mathcal{I} - \mathbf{Int}$ valid model of \mathbf{Z}^* for every set of interventions \mathcal{I} in \mathfrak{C} . The proof of this is the same as the proof of [Proposition 4.1](#), except that we do not need **D4** to have that $\mathcal{L}^a(\mathbf{Z}^*)$ is Markov w.r.t. to the graph of \mathfrak{C} since this is a complete DAG (and Markovianity trivially holds). Therefore, violating only **D4** is not a viable strategy to avoid circularity if one believes that not all complete and compatible CBNs should be considered interventionally valid. Consider the following interpretation that may violate **D4**, but satisfies **D0–D3**.

Definition D.3. \mathbf{Int}_M . An interpretation violating only D4. Let a data-generating process \mathcal{D} , representation \mathbf{Z}^* , compatible CBN \mathfrak{C} , and set of interventions \mathcal{I} in \mathfrak{C} be given. We define the interpretation \mathbf{Int}_M by the following rule: An intervention $\text{do}(j \leftarrow q_j, j \in J) \in \mathcal{I}$ is in $\mathbf{Int}_M^{\mathcal{I}}(a)$ if and only if the following two conditions hold:

- 1) $\mathcal{L}^a(Z_i^* \mid \mathbf{PA}_i^*) \sim q_i$ for all $i \in J$. That is, \mathbf{Int}_M satisfies **D0** (correct conditionals on intervened nodes).
- 2) $\mathcal{L}^a(Z_i^* \mid \mathbf{PA}_i^*) \sim p_i^{\mathfrak{C}}$ for all $i \notin J$. Intuitively, we do not intervene on nodes not in J .

\circ

The M in \mathbf{Int}_M is for ‘Markov’. Interpretation \mathbf{Int}_M is the same as \mathbf{Int}_C except that we do not require that $\mathcal{L}^a(\mathbf{Z}^*)$ is Markov w.r.t. the graph of \mathfrak{C} for a to be interpreted as an intervention. Since \mathbf{Int}_M satisfies **D0–D3** but not **D4**, interventional validity requires actions to not introduce new dependencies, as shown by the following proposition.

Proposition D.4. Let a data-generating process \mathcal{D} , representation \mathbf{Z}^* , and compatible CBN \mathfrak{C} with graph \mathcal{G} be given.

- (1) If $\mathcal{L}^a(\mathbf{Z})$ is Markov w.r.t. \mathcal{G} for all a , then \mathfrak{C} is an $\mathcal{I} - \mathbf{Int}_M$ valid model of \mathbf{Z}^* for every set of interventions \mathcal{I} in \mathfrak{C} .
- (2) If there exists an action a such that $\mathcal{L}^a(\mathbf{Z})$ is not Markov w.r.t. \mathcal{G} , then there exists a set of interventions \mathcal{I} in \mathfrak{C} such that \mathfrak{C} is not an $\mathcal{I} - \mathbf{Int}_M$ valid model of \mathbf{Z}^*

Proof. (1) Let action a and set of interventions \mathcal{I} in \mathfrak{C} be given. Assume that $d \in \mathbf{Int}_M^{\mathcal{I}}(a)$. We want to show $\mathcal{L}^a(\mathbf{Z}^*) = \mathcal{L}^{\mathfrak{C}; d}(\mathbf{Z})$. The proof is the same as for [Proposition 3.2](#).

(2) Let a be an action such that $\mathcal{L}^a(\mathbf{Z}^*)$ is not Markov w.r.t. \mathcal{G} . Consider intervention $d = \text{do}(j \leftarrow q_j, j \in [n])$ such that $\mathcal{L}^a(Z_j^* | \mathbf{PA}_j^*) \sim q_j$ for all $j \in [n]$, and let $\mathcal{I} = \{d\}$. Now $d \in \mathbf{Int}_M^{\mathcal{I}}(a)$ but $\mathcal{L}^a(\mathbf{Z}^*) \neq \mathcal{L}^{\mathfrak{C};d}(\mathbf{Z})$ since $\mathcal{L}^a(\mathbf{Z}^*)$ is not Markov w.r.t. \mathcal{G} , while $\mathcal{L}^{\mathfrak{C};d}(\mathbf{Z})$ is Markov w.r.t. \mathcal{G} . \square

We now provide an example of how \mathbf{Int}_M might falsify a model.

Example D.5. Falsifying a model under interpretation \mathbf{Int}_M .


Let the set of actions be $\mathcal{A} = \{\mathcal{O}, a\}$. Assume that $\mathcal{L}^{\mathcal{O}}(X^*, Y^*, Z^*)$ is given by the observational distribution induced by the SCM

$$\begin{aligned} X^* &:= \mathcal{E}_1 \\ Y^* &:= X^* + \mathcal{E}_2 \\ Z^* &:= \mathcal{E}_3 \end{aligned}$$

with $\mathcal{E}_1, \mathcal{E}_2, \mathcal{E}_3 \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1)$, and assume that $\mathcal{L}^a(X^*, Y^*, Z^*)$ is given by the observational distribution induced by the SCM

$$\begin{aligned} X^* &:= \mathcal{E}_1 \\ Y^* &:= \frac{X^*}{2} + \frac{Z^*}{\sqrt{2}} + \frac{\mathcal{E}_2}{\sqrt{2}} \\ Z^* &:= \mathcal{E}_3 \end{aligned}$$

with $\mathcal{E}_1, \mathcal{E}_2, \mathcal{E}_3 \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1)$. Consider the compatible CBN \mathfrak{C} given by the graph



and kernels

$$\begin{aligned} X &\sim \mathcal{N}(0, 1) \\ Y | X = x &\sim \mathcal{N}(x, 1) \\ Z &\sim \mathcal{N}(0, 1). \end{aligned}$$

Let $\mathcal{I} = \{\text{do}(Y \leftarrow \mathcal{N}(\frac{x}{2}, 1))\}$.²⁹ Now, $\text{do}(Y \leftarrow \mathcal{N}(\frac{x}{2}, 1)) \in \mathbf{Int}_M^{\mathcal{I}}(a)$ since $\mathcal{L}^a(X^*) = \mathcal{N}(0, 1)$, $\mathcal{L}^a(Y^* | X^*) \sim \mathcal{N}(\frac{x^*}{2}, 1)$, and $\mathcal{L}^a(Z^*) = \mathcal{N}(0, 1)$, in line with the corresponding conditionals in $\mathcal{L}^{\mathfrak{C};\text{do}(Y \leftarrow \mathcal{N}(\frac{x}{2}, 1))}(X, Y, Z)$. But $\mathcal{L}^a(X^*, Y^*, Z^*) \neq \mathcal{L}^{\mathfrak{C};\text{do}(Y \leftarrow \mathcal{N}(\frac{x}{2}, 1))}(X, Y, Z)$, for example, because $Y^* \not\perp\!\!\!\perp Z^*$ in $\mathcal{L}^a(X^*, Y^*, Z^*)$ while $Y \perp\!\!\!\perp Z$ in $\mathcal{L}^{\mathfrak{C};\text{do}(Y \leftarrow \mathcal{N}(\frac{x}{2}, 1))}(X, Y, Z)$. Intuitively, action a behaves like an intervention that introduces probabilistic dependence between variables Y and Z . \circ

²⁹ By $\mathcal{N}(\frac{x}{2}, 1)$ we denote the kernel $x \mapsto \mathcal{N}(\frac{x}{2}, 1)$, resulting in an imperfect intervention.

E Details on Causal Abstraction (Section 6.3)

E.1 Further details on Example 6.5

To check that $(M_{\mathbf{Y}}, \mathcal{I}_{\mathbf{Y}})$ for $\mathcal{I}_{\mathbf{Y}} = \omega_{\tau}(\mathcal{I}^*)$ is a τ -abstraction of $(M_{\mathbf{X}}, \mathcal{I}^*)$ [Definition 3.13 in Beckers and Halpern, 2019], we must check that

1. τ is surjective.
2. There exists a surjective function $\tau_U : [4] \rightarrow \{0, 1\}^2$ such that $\tau(M_{\mathbf{X}}^d(u_1)) = M_{\mathbf{Y}}^{\omega_{\tau}(d)}(\tau_U(u_1))$ for every $d \in \mathcal{I}^*$ and $u_1 \in [4]$.
3. The third condition of Definition 3.13 in Beckers and Halpern [2019] holds since ω_{τ} is defined for all interventions in \mathcal{I}^* and we choose $\mathcal{I}_{\mathbf{Y}} = \omega_{\tau}(\mathcal{I}^*)$.

$\tau : [4] \rightarrow \{0, 1\}^2$ given by $x \mapsto (\mathbb{1}(x = 3) + \mathbb{1}(x = 4), \mathbb{1}(x = 2) + \mathbb{1}(x = 4))$ is clearly surjective. For each $d \in \mathcal{I}^*$, we compute $\omega_{\tau}(d)$ using Definition 3.12 of Beckers and Halpern [2019]:

- $\omega_{\tau}(\text{do}(X_1 = 1)) = \text{do}(Y_1 = 0, Y_2 = 0)$,
- $\omega_{\tau}(\text{do}(X_1 = 2)) = \text{do}(Y_1 = 0, Y_2 = 1)$,
- $\omega_{\tau}(\text{do}(X_1 = 3)) = \text{do}(Y_1 = 1, Y_2 = 0)$,
- $\omega_{\tau}(\text{do}(X_1 = 4)) = \text{do}(Y_1 = 1, Y_2 = 1)$.

Let $\tau_U := \tau$. Now, it is straightforward to check the second requirement. For example,

$$\tau\left(M_{\mathbf{X}}^{\text{do}(X_1=1)}(4)\right) = \tau(1) = (0, 0) = M_{\mathbf{Y}}^{\text{do}(Y_1=0, Y_2=0)}(\tau_U(4)),$$

and

$$\tau\left(M_{\mathbf{X}}^{\text{do}(X_1=3)}(4)\right) = \tau(3) = (1, 0) = M_{\mathbf{Y}}^{\text{do}(Y_1=1, Y_2=0)}(\tau_U(4)).$$

(Note that ω_{τ} would map the empty intervention to the empty intervention and $\tau(M_{\mathbf{X}}(u_1)) = M_{\mathbf{Y}}(\tau_U(u_1))$ for every $u_1 \in [4]$, that is, $M_{\mathbf{Y}}$ is also a τ -abstraction of $M_{\mathbf{X}}$ if we additionally included the empty intervention in \mathcal{I}^* (this is analogous to the CBN \mathfrak{C} being compatible with $\mathbf{Y}^* = \tau(X_1)$).

E.2 Interventional validity is not preserved by constructive soft abstractions under interpretation Int_P

The following example shows that constructive soft abstractions [Massidda et al., 2023] do not preserve interventional validity; the example is a slight modification of Example 4 in Massidda et al. [2023], adding nodes X_0 and Y_0 .

Example E.1. Transforming an interventionally valid model into an invalid model by a constructive soft abstraction. Let SCM $M_{\mathbf{X}}$ be given by

$$\begin{aligned} X_0 &:= U_0 \\ X_1 &:= U_1 \\ X_2 &:= U_2 \\ X_3 &:= \min(X_1, X_2) \\ X_4 &:= (X_1 - X_2)^2 \end{aligned}$$

with $U_0, U_1, U_2 \stackrel{\text{iid}}{\sim} \text{Unif}(\{0, 1\})$. and let SCM $M_{\mathbf{Y}}$ be given by

$$\begin{aligned} Y_0 &:= N_0 \\ Y_1 &:= N_1 \\ Y_2 &:= N_2 \\ Y_3 &:= Y_1 + Y_2 \end{aligned}$$

with $N_0, N_1, N_2 \stackrel{\text{iid}}{\sim} \text{Unif}(\{0, 1\})$. Let $\tau : \{0, 1\}^5 \rightarrow \{0, 1\}^3 \times \{0, 1, 2, 3\}$ be given by $(x_0, x_1, x_2, x_3, x_4) \mapsto (x_0, x_1, x_2, 2x_3 + x_4)$. $M_{\mathbf{Y}}$ is a constructive soft τ -abstraction of $M_{\mathbf{X}}$ [Massidda et al., 2023]. Let \mathfrak{A} be a CBN with graph and observational distribution induced by $M_{\mathbf{X}}$. Assume that \mathbf{X}^* is emulated by \mathfrak{A} and $\mathcal{I}^* = \{\text{do}(X_0 = 0, X_4 = 1)\}$. Since \mathcal{I}^* only contains perfect interventions, \mathfrak{A} is an $\mathcal{I} - \text{Int}_P$ valid model of \mathbf{X}^* for every set of interventions \mathcal{I} in \mathfrak{A} , see Proposition 5.2 (1). Let \mathfrak{C} be a CBN with graph and observational distribution induced by $M_{\mathbf{Y}}$, and let \mathcal{I} be a set of interventions in \mathfrak{C} such that $\text{do}(Y_0 = 0) \in \mathcal{I}$. Then, \mathfrak{C} is not an $\mathcal{I} - \text{Int}_P$ valid model of $\mathbf{Y}^* := \tau(\mathbf{X}^*)$. To see this let a be an action such that $\mathcal{L}^a(\mathbf{X}^*) = \mathcal{L}^{\mathfrak{A}; \text{do}(X_0=0, X_4=1)}(\mathbf{X})$ (such an action exists since $\text{do}(X_0 = 0, X_4 = 1) \in \mathcal{I}^*$). Now $\text{do}(Y_0 = 0) \in \text{Int}_P^{\mathcal{I}}(a)$ but $\mathcal{L}^a(\mathbf{Y}^*) \neq \mathcal{L}^{\mathfrak{C}; \text{do}(Y_0=0)}(\mathbf{Y})$, for example, because $\mathbb{E}^a(Y_3^*) = \mathbb{E}^{\mathfrak{A}; \text{do}(X_0=0, X_4=1)}(2X_3 + X_4) = \frac{3}{2} \neq 1 = \mathbb{E}^{\mathfrak{C}; \text{do}(Y_0=0)}(Y_3)$. \circ

E.3 Further details on Example 6.6

To show that $(M_{\mathbf{Y}}, \mathcal{I}_{\mathbf{Y}})$ is not a τ -abstraction of $(M_{\mathbf{X}}, \mathcal{I}^*)$ [Definition 3.13 in Beckers and Halpern, 2019], we must show a violation of at least one of the following criteria

1. τ is surjective.
2. There exists a surjective function $\tau_U : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ such that $\tau(M_{\mathbf{X}}^d(u_1, u_2, u_3)) = M_{\mathbf{Y}}^{\omega_{\tau}(d)}(\tau_U(u_1, u_2, u_3))$ for every $d \in \mathcal{I}^*$ and $(u_1, u_2, u_3) \in \mathbb{R}^3$.
3. The third condition of Definition 3.13 in Beckers and Halpern [2019] would be trivially violated if we chose $\mathcal{I}_{\mathbf{Y}} \neq \omega_{\tau}(\mathcal{I}^*)$. Therefore, we choose $\mathcal{I}_{\mathbf{Y}} = \omega_{\tau}(\mathcal{I}^*)$ (here, ω_{τ} is defined for all interventions in \mathcal{I}^*).

Assume for contradiction that $(M_{\mathbf{Y}}, \omega_{\tau}(\mathcal{I}^*))$ is a τ -abstraction of $(M_{\mathbf{X}}, \mathcal{I}^*)$. Notice that $\omega_{\tau}(\text{do}(X_1 = x_1)) = \emptyset$ for all $x_1 \in \mathbb{R}$ [Beckers and Halpern, 2019, Definition 3.12]. The second condition then implies that

$$\tau(M_{\mathbf{X}}^{\text{do}(X_1=x_1)}(u_1, u_2, u_3)) = (x_1 + u_2, x_1 + u_2 + u_3) = M_{\mathbf{Y}}(\tau_U(u_1, u_2, u_3)),$$

for every $x_1, u_1, u_2, u_3 \in \mathbb{R}$. But this is impossible since no function $\tau_U : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ exists that simultaneously satisfies all the above equations, for example, since this would imply that the first coordinate $M_Y(\tau_U(1, 1, 1))_1 = \sqrt{2}\tau_U(1, 1, 1)_1 = x + 1$ for every $x \in \mathbb{R}$.


F Relation to Phenomenological causality

Janzing and Mejia [2024] present a related approach to grounding causality in actions. Translated into the notation of the present work, Definition 3 in Janzing and Mejia [2024] states that a graph \mathcal{G} is a valid causal graph over variables $\mathbf{Z}^* = (Z_1^*, \dots, Z_n^*)$ relative to a set of actions \mathcal{A} if (1) $\mathcal{L}^{\mathcal{O}}(\mathbf{Z}^*)$ is Markov w.r.t. \mathcal{G} , and (2) $\mathcal{A} \setminus \{\mathcal{O}\} = \cup_{j=1}^n \mathcal{A}_j$, where each \mathcal{A}_j is nonempty, and for every $a \in \mathcal{A}_j$,

$$\begin{aligned} \mathcal{L}^a(Z_j^* | \mathbf{PA}_j^*) &\neq \mathcal{L}^{\mathcal{O}}(Z_j^* | \mathbf{PA}_j^*), \text{ and} \\ \mathcal{L}^a(Z_i^* | \mathbf{PA}_i^*) &= \mathcal{L}^{\mathcal{O}}(Z_i^* | \mathbf{PA}_i^*) \text{ for } i \neq j. \end{aligned}$$

Interpreting this definition within our framework of interventional validity, it is similar to \mathbf{Int}_S in Definition 5.4.³⁰ Indeed, analogous to Proposition 5.5 (2), if \mathbf{Z}^* is emulated by CBN \mathfrak{C} (with graph \mathcal{G}) and interventions \mathcal{I}^* , then (using the terminology of Janzing and Mejia [2024]) \mathcal{G} is not a valid causal graph over variables \mathbf{Z}^* if \mathcal{I}^* contains a minimal decomposable multi-node intervention. There are two differences between \mathbf{Int}_S and Definition 3 of Janzing and Mejia [2024] to take notice of.

1) Janzing and Mejia [2024] require that each \mathcal{A}_j is nonempty, that is, intuitively, that it is possible to intervene on every node. This requirement may be overly restrictive as the following example illustrates. Assume we are interested in the causal effect of a treatment T^* on an outcome Y^* , where W^* is a confounder. Say we model the situation using CBN \mathfrak{C} over nodes (T, Y, W) with graph \mathcal{G} :



If, for every $a \in \mathcal{A}$,

$$\mathcal{L}^a(Y^* | T^*, W^*) \sim p_{Y|T,W}^{\mathfrak{C}}, \text{ and} \tag{1}$$

$$\mathcal{L}^a(W^*) \sim p_W^{\mathfrak{C}}, \tag{2}$$

then \mathcal{G} is not a valid causal graph (using the terminology of Janzing and Mejia [2024]) because no action changes the conditional distribution of outcome given treatment and confounding nor the marginal distribution of the confounding. Yet, this invariance is what enables \mathfrak{C} to be useful for making predictions about the actions that affect the conditional distribution of T^* given W^* . Indeed, our formal framework correctly captures

³⁰ Since Janzing and Mejia [2024] also require that \mathcal{A} only consists of ‘elementary actions’, Definition 3 of Janzing and Mejia [2024] also bears resemblance to \mathbf{Int}_K (Definition 5.7). In our work, we do not require that \mathcal{A} only consists of ‘elementary actions’, and instead draw a formal distinction between \mathbf{Int}_K and \mathbf{Int}_S .

the usefulness of such model: If [Equation \(1\)](#) and [Equation \(2\)](#) hold for every $a \in \mathcal{A}$, then \mathfrak{C} is an $\mathcal{I} - \mathbf{Int}$ valid model of (T^*, Y^*, W^*) for every set of interventions \mathcal{I} in \mathfrak{C} if \mathbf{Int} is an interpretation that satisfies **DO**, which includes all interpretations considered in this paper.

In addition, there is a subtle technical issue in [Janzing and Meija \[2024\]](#) that our framework handles correctly.

2) Under \mathbf{Int}_S , [Definition 5.4](#), we interpret an action a as an intervention $d = \text{do}(j \leftarrow q_j)$ on node j if $\mathcal{L}^a(Z_j^* | \mathbf{PA}_j^*) \sim q_j$, $\mathcal{L}^a(Z_j^* | \mathbf{PA}_j^*) \approx p_j^{\mathfrak{C}}$, and $\mathcal{L}^a(\mathbf{Z}^*)$ is Markov w.r.t. the DAG. Therefore, whether $d \in \mathbf{Int}_S^{\mathcal{I}}(a)$ depends on both the graph and the specific kernels in the CBN model. In the approach by [Janzing and Meija \[2024\]](#), to decide whether a causal graph is valid, one has to determine for all i whether the conditional distributions $\mathcal{L}^a(Z_i^* | \mathbf{PA}_i^*)$ are the same as $\mathcal{L}^{\mathcal{O}}(Z_i^* | \mathbf{PA}_i^*)$. However, whether or not $\mathcal{L}^a(Z_i^* | \mathbf{PA}_i^*) = \mathcal{L}^{\mathcal{O}}(Z_i^* | \mathbf{PA}_i^*)$ cannot in general be determined based on $\mathcal{L}^a(\mathbf{Z}^*)$ and $\mathcal{L}^{\mathcal{O}}(\mathbf{Z}^*)$, and thus it is unclear what it means. To illustrate this point, consider that $\mathcal{L}^{\mathcal{O}}(Z_1^*, Z_2^*) = \mathcal{N}\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}\right)$ and $\mathcal{L}^a(Z_1^*, Z_2^*) = \delta_0 \otimes \delta_5$ and ask whether $\mathcal{L}^a(Z_2^* | Z_1^*) = \mathcal{L}^{\mathcal{O}}(Z_2^* | Z_1^*)$? One might argue that this equality does not hold because $p_{Z_2^* | Z_1^*}^{\mathcal{O}}(\cdot | 0)$ is a normal density with mean 0 and therefore does not have point mass in 5. This, however, does not follow from the observational distribution $\mathcal{L}^{\mathcal{O}}(Z_1^*, Z_2^*)$ since conditional distributions of Z_2^* given Z_1^* can always be changed on null sets w.r.t. $\mathcal{L}(Z_1^*)$ without altering the joint distribution of (Z_1^*, Z_2^*) [[Lauritzen, 2019](#)]. In particular, there exists a CBN \mathfrak{C} such that $\mathcal{L}^{\mathcal{O}}(Z_1^*, Z_2^*) = \mathcal{L}^{\mathfrak{C}}(Z_1^*, Z_2^*)$ and $\mathcal{L}^a(Z_1^*, Z_2^*) = \mathcal{L}^{\mathfrak{C}; \text{do}(Z_1^*=0)}(Z_1^*, Z_2^*)$. We avoid this problem by comparing $\mathcal{L}^a(Z_2^* | Z_1^*)$ with the specific kernels given by the CBN and write, for example, $\mathcal{L}^a(Z_2^* | Z_1^*) \sim p_{Z_2^* | Z_1^*}^{\mathfrak{C}}$ if $p_{Z_2^* | Z_1^*}^{\mathfrak{C}}$ is a valid Markov kernel for conditional distribution of Z_2^* given Z_1^* in $\mathcal{L}^a(Z_1^*, Z_2^*)$.

This subtlety occurs even for discrete variables. Let CBN \mathfrak{A} be given by graph $X \rightarrow Y$ and kernels

$$\begin{aligned} \mathcal{L}^{\mathfrak{A}}(X) &= \delta_0 \\ \mathcal{L}^{\mathfrak{A}}(Y | X = 0) &= \text{Ber}(0.4) \\ \mathcal{L}^{\mathfrak{A}}(Y | X = 1) &= \text{Ber}(0.6). \end{aligned}$$

Assume that (X^*, Y^*) is emulated by \mathfrak{A} and interventions $\mathcal{I}^* = \{\text{do}(X = 1)\}$. Given action $a \in \mathcal{A} \setminus \{\mathcal{O}\}$, there is no way to determine whether $\mathcal{L}^{\mathcal{O}}(Y^* | X^*) = \mathcal{L}^a(Y^* | X^*)$ since the support for X^* under the two distributions is not overlapping. But we can, for example, say that $\mathcal{L}^a(Y^* | X^*) \sim p_{Y^* | X^*}^{\mathfrak{C}; \text{do}(X=1)} = p_{Y^* | X^*}^{\mathfrak{C}}$ and $\mathcal{L}^{\mathcal{O}}(Y^* | X^*) \sim p_{Y^* | X^*}^{\mathfrak{C}; \text{do}(X=1)} = p_{Y^* | X^*}^{\mathfrak{C}}$.