On the algebraic independence of G-functions and E-functions

DANIEL VARGAS-MONTOYA

ABSTRACT. Let $f_1(z), \ldots, f_m(z)$ be power series in $\mathbb{Q}_p[[z]]$ such that, for every $1 \leq i \leq m$, $f_i(z)$ is solution of a differential operator $\mathcal{L}_i \in E_p[d/dz]$, where E_p is the field of analytic elements. We prove that if, for every $1 \leq i \leq m$, \mathcal{L}_i has a strong Frobenius structure and has maximal order multiplicity at zero (MOM) then $f_1(z), \ldots, f_m(z)$ are algebraically dependent over E_p if and only if there are integers a_1, \ldots, a_m not all zero, such that $f_1^{a_1}(z) \cdots f_m^{a_m}(z) \in E_p$. The main consequence of this result is that it allows us to study the algebraic independence of a large class of *G*-functions and certain *E*-functions.

1. Introduction

The class of *E*-functions and *G*-functions were introduced by Siegel in his seminal 1929 paper [26]. The main purpose of Siegel in introducing such classes was to generalise the classical theorems of *Her*mite and *Lindemann-Weierstrass*. Let us recall that a power series $f(z) = \sum_{n \ge 0} a_n z^n$ is a *G*-function if the coefficients a_n are algebraic numbers and there is a real number C > 0 such that:

1. the power series f(z) is solution of a nonzero differential operator with coefficients in $\overline{\mathbb{Q}}(z)$;

2. the absolute values of all Galois conjugates of a_n are at most C^{n+1} for all $n \ge 0$;

3. there is a sequence of positive integers D_m such that, for all integers $m \ge 0$, $|D_m| < C^{m+1}$ and, for all $n \le m$, $D_m a_n$ is an algebraic integer.

Among the G-functions, we have the hypergeometric series ${}_{n}F_{n-1}$ with rational parameters, and diagonal of rational functions.

Furthermore, a power series $F(z) = \sum_{n \ge 0} \frac{a_n}{n!} z^n$ is a *E*-function if and only if $f(z) = \sum_{n \ge 0} a_n z^n$ is a *G*-function. In particular, that implies that if F(z) is a *E*-function then F(z) is solution of a nonzero differential operator with coefficients in $\overline{\mathbb{Q}}(z)$. Classical examples of *E*-functions are the exponential function, sine, cosine, the hypergeometric series ${}_{p}F_{p}$ with rational parameters and the Bessel function.

The study of algebraic independence of E and G-functions finds its motivation in Siegel's work [26], where he developed a method to prove the algebraic independence of values of E-functions at algebraic points. The turning point of this method is given by Siegel-Shidlovskii's Theorem. Recall that this theorem says that if $F_1(z), \ldots, F_n(z)$ are E-functions and Y'(z) = A(z)Y(z) where Y(z) = $(F_1(z), \ldots, F_n(z))^t$ and A(z) is a matrix with coefficients in $\overline{\mathbb{Q}}(z)$ then, for almost every algebraic number α , we have the equality deg $\operatorname{tr}_{\overline{\mathbb{Q}}(z)}(\overline{\mathbb{Q}}(z)(F_1(z), \ldots, F_n(z))) = \operatorname{deg} \operatorname{tr}_{\overline{\mathbb{Q}}}(\overline{\mathbb{Q}}(F_1(\alpha), \ldots, F_n(\alpha)))$. Siegel also mentioned that his method should also work for *G*-functions. But, it follows from [5] that Siegel-Shidlovskii's Theorem does not hold for *G*-functions. Nevertheless, over the past years, several results on transcendence [4] and algebraic independence of *G*-functions [1, 29] have been obtained. Concerning the transcendence and algebraic independence of values of *G*-functions at algebraic points not much is known.⁽¹⁾

Since E and G-functions are solutions of differential operators with coefficients in $\overline{\mathbb{Q}}(z)$, differential Galois theory provides a natural framework to study the algebraic independence of these power series. For instance, by using this theory, some criteria determining the algebraic independence of E-functions are given in [3] and [27]. Further, Beukers and Heckman [4] also used this theory to give a characterization of hypergeometric series ${}_{n}F_{n-1}$ with rational parameters that are algebraic over $\mathbb{Q}(z)$. In order to study the algebraic independence of power series $f_1(z), \ldots, f_m(z)$, we must compute the differential Galois group associated with a differential operator annihilating these power series. In general, it is not easy to determine this Galois group. Nevertheless, in the present work we overcome this problem for a class of E and G-functions by showing that any power series f(z) in such class is a solution of a differential operator of order 1 with coefficients in the field of analytic elements E_p . More precisely, we focus on the set of power series $f(z) \in 1 + z\mathbb{Q}_p[[z]]$ such that f(z) is solution of a differential operator $\mathcal{L} \in E_p[d/dz]$ such that \mathcal{L} has maximal order multiplicity at zero (MOM) and has strong Frobenius structure. It turns out that there are E and G-functions that are annihilated by differential operators satisfying these two conditions. When \mathcal{L} belongs to $\mathbb{Q}(z)[d/dz]$, the MOM property is equivalent to saying that \mathcal{L} has maximal unipotent monodromy at zero (MUM). The notion of strong Frobenius structure was introduced by Dwork [15] and it corresponds to a Frobenius action on the set of solutions of \mathcal{L} . The reader can find the precise definition of both notions in Section 2. So, our main result says

Theorem 1.1 (Theorems 2.3 and 2.5). — Let $f_1(z), \ldots, f_m(z)$ be power series in $1 + z\mathbb{Q}_p[[z]]$ and let us suppose that, for every $i \in \{1, \ldots, m\}$, $f_i(z)$ is solution of a differential operator \mathcal{L}_i with coefficients E_p such that \mathcal{L}_i has a strong Frobenius structure and is MOM at zero. Then:

(i) $f_1(z), \ldots, f_m(z)$ are algebraically dependent over E_p if and only if there are $a_1, \ldots, a_m \in \mathbb{Z}$ not all zero such that

 $f_1(z)^{a_1}\cdots f_m(z)^{a_m}\in E_p;$

(ii) $f_1(z), \ldots, f_m(z)$ are algebraically dependent over E_p if and only if, for any $(r_1, \ldots, r_m) \in \mathbb{N}^m$, $f_1^{(r_1)}, \ldots, f_m^{(r_m)}$ are algebraically dependent over E_p .

The proof of Theorem 1.1 combines differential Galois theory and the theory of p-adic differential equations. Using the latter, we prove the following result:

Theorem 1.2 (Theorem 3.1). — Let f(z) be power series in $1 + z\mathbb{Q}_p[[z]]$. Suppose that f(z) is solution of a differential operator \mathcal{L} with coefficients E_p . If \mathcal{L} is MOM at zero and has a strong Frobenius structure then:

(i) $f(z) \in 1 + z\mathbb{Z}_p[[z]];$ (ii) f'(z)/f(z) belongs to E_p .

 $^{^{(1)}}$ For more details on what is known about the transcendence of values of G-functions at algebraic points, we refer the reader to [25, Sec 7].

Then, by combining the previous result with a fundamental result of differential Galois theory due to Kolchin [22], we are able to prove Theorem 1.1.

We make Theorem 1.1 effective by giving a criterion that determines when $f_1(z)^{a_1} \cdots f_m(z)^{a_m} \notin E_p$ for any $(a_1, \ldots, a_m) \in \mathbb{Z}^m \setminus \{(0, \ldots, 0)\}$. This criterion, given by Theorem 6.12, relies on the study of the poles and residues of the power series $f'_1(z)/f_1(z), \ldots, f'_m(z)/f_m(z)$. As an illustration of this criterion, we prove the following result:

Theorem 1.3 (Theorem 2.10). — Let π_3 be in $\overline{\mathbb{Q}}$ such that $\pi_3^2 = -3$. Let us consider the following power series

$$J_0(z) = \sum_{n \ge 0} \frac{(-1)^n}{4^n (n!)^2} z^{2n}, \quad \mathfrak{f}(z) = \sum_{n \ge 0} = \frac{-1}{(2n-1)64^n} \binom{2n}{n}^3 z^n,$$

and

$$\mathfrak{A}(z) = \sum_{n \ge 0} \left(\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 \right) z^n.$$

Then, for all integers $r, s, k \ge 0$, the power series $J_0^{(r)}(\pi_3 z)$, $\mathfrak{A}^{(s)}(z)$, and $\mathfrak{f}^{(k)}(z)$ are algebraically independent over E_3 .

This result implies that, for all integers $r, s, k \ge 0$, the power series $J_0^{(r)}(\pi_3 z)$, $\mathfrak{A}^{(s)}(z)$, and $\mathfrak{f}^{(k)}(z)$ are algebraically independent over $\mathbb{Q}(z)$ because, for all prime numbers p, $\mathbb{Q}(z) \subset E_p$. In this case, the power series $J_0(\pi_3 z)$ is a *E*-function and the power series $\mathfrak{f}(z)$ and $\mathfrak{A}(z)$ are *G*-functions.

A similar statement to Theorem 1.1 was proven by Adamczeswki, Bell and Delaygue [1] for the set of power series that are *p*-Lucas for infinitely many prime numbers *p*. We recall that a power series $f(z) = \sum_{n \ge 0} a_n z^n \in \mathbb{Q}[[z]]$ is *p*-Lucas if $a_0 = 1$, f(z) belongs to $\mathbb{Z}_{(p)}[[z]]^{(2)}$, and

$$f(z) = \left(\sum_{n=0}^{p-1} a_n z^n\right) f(z)^p \mod p\mathbb{Z}_{(p)}[[z]].$$

Let S be an infinite set of prime numbers. In [1], the authors introduced the set of power series $\mathcal{L}(S)$. The main example of this set is given by the power series that are *p*-Lucas for all $p \in S$. The main result of [1] states that if $f_1(z), \ldots, f_m(z)$ belong to $\mathcal{L}(S)$ then the power series $f_1(z), \ldots, f_m(z)$ are algebraically dependent over $\mathbb{Q}(z)$ if and only if there are $a_1, \ldots, a_m \in \mathbb{Z}$ not all zero such that $f_1(z)^{a_1} \cdots f_m(z)^{a_m} \in \mathbb{Q}(z)$. In the approach established in [1] it is crucial to assume that S is infinite. In the same work, the authors showed that a large class of G-functions are p-Lucas for infinitely many primes p. Further, in [29] we introduced the set $\mathcal{L}^2(S)$. This set contains $\mathcal{L}(S)$, and a criterion for the algebraic independence of the power series in $\mathcal{L}^2(S)$ has been given. In [29] this criterion is illustrated by showing the algebraic independence of many G-functions (see e.g [29, Theorems 9.1, 9.2]). Again, in the approach used in [29] it is essential that the S to be infinite.

The main difference between the method developed in the present work and the previous ones lies in the fact that the assumption that S is infinite is removed. Our method focuses on a single prime number p. One of the benefits of working with a single prime number p is that we can now study the algebraic independence of E-functions. Indeed, the sets $\mathcal{L}(S)$ and $\mathcal{L}^2(S)$ do not contain any E-function. It is also important to highlight that the methods used in [1, 29] do not allow to study the algebraic independence of derivatives.

⁽²⁾ Recall that $\mathbb{Z}_{(p)}$ is the ring of rational numbers a/b such that p does not divide b and a is coprime to b.

DANIEL VARGAS-MONTOYA

The outline of the paper is as follows. In Section 2 we recall the definition of the field of analytic elements, strong Frobenius structure and we state our main results, Theorems 2.3 and 2.5. Actually, Theorem 1.1 is a weak version of these two theorems. In section 3, we prove Theorems 2.3 and Theorem 2.5. The proof of Theorem 2.3 relies on Theorem 1.2 and Kolchin's Theorem. Section 4 is devoted to proving (i) of Theorem 1.2 and Section 5 is dedicated to proving (ii) of Theorem 1.2. Finally, in Section 6 we provide a necessary condition for determining the algebraic independence over E_p . This condition is given by Theorem 6.12 and we illustrate this criterion by proving the algebraic independence of some G-functions and E-functions.

Acknowledgements. We warmly thank Charlotte Hardouin for the valuable discussions regarding this project. The second part of Theorem 1.1 was motivated by a question from Tanguy Rivoal about whether our method could be used to study the algebraic independence of derivatives; we also thank him.

2. Main results

We first recall the definitions of the field of analytic elements, strong Frobenius structure, and MOM operators.

2.1. Analytic elements.

Let p be a prime number and let \mathbb{Q}_p be the field of p-adic numbers. It is well-know that the field \mathbb{Q}_p is equipped with the non-Archimedean norm $|\cdot|$ such that |p| = 1/p. This norm will be called the p-adic norm. Let $\overline{\mathbb{Q}_p}$ be an algebraic closure of \mathbb{Q}_p . It is also well-known that this norm extends to a unique way to the field $\overline{\mathbb{Q}_p}$ and it is still called the p-adic norm. Finally, \mathbb{C}_p will be the completion of $\overline{\mathbb{Q}_p}$ with respect to the p-adic norm.

From now on, $K \subset \mathbb{C}_p$ designates a complete extension of \mathbb{Q}_p with respect to the *p*-adic norm.

We remind the reader that the Amice ring with coefficient in K is the following ring

$$\mathcal{A}_{K} = \left\{ \sum_{n \in \mathbb{Z}} a_{n} z^{n} : a_{n} \in K, \lim_{n \to -\infty} |a_{n}| = 0 \text{ and } \sup_{n \in \mathbb{Z}} |a_{n}| < \infty \right\}.$$

This ring is equipped with the Gauss norm, which is defined as follows:

$$\left|\sum_{n\in\mathbb{Z}}a_nz^n\right|_{\mathcal{G}}=\sup_{n\in\mathbb{Z}}|a_n|.$$

Since K is complete, by Proposition 1.1 of [10], the ring \mathcal{A}_K is also complete for the Gauss norm. Furthermore, it follows from Proposition 1.2 of [10] that $h = \sum_{n \in \mathbb{Z}} h_n z^n \in \mathcal{A}_K$ is a unit element if only if, there is $n_0 \in \mathbb{Z}$ such that $|h| = |h_{n_0}| \neq 0$. In particular, a non-zero polynomial with coefficients in K is a unit element in \mathcal{A}_K and therefore, $K(z) \subset \mathcal{A}_K$. For this reason, the field K(z) is equipped with the Gauss. Actually, for any polynomial $P(z) = \sum_{i=0}^n a_i z^i$ and $Q(z) = \sum_{j=0}^m b_j z^j$, we have

$$\left|\frac{P(z)}{Q(z)}\right|_{\mathcal{G}} = \frac{\max\{|a_i|\}_{1 \le i \le n}}{\max\{|b_j|\}_{1 \le j \le n}}$$

The field of *analytic elements* with coefficients in K, denoted by E_K , is the completion of K(z) with respect to the Gauss norm. As \mathcal{A}_K is complete and $K(z) \subset \mathcal{A}_K$ then $E_K \subset \mathcal{A}_K$. When $K = \mathbb{C}_p$, the field $E_{\mathbb{C}_p}$ will be denoted by E_p and the Amice ring $\mathcal{A}_{\mathbb{C}_p}$ will be denoted by \mathcal{A}_p .

Remark 2.1. We have $\mathbb{Q}(z) \subset \mathbb{Q}_p(z) \subset E_K$ because $\mathbb{Q} \subset \mathbb{Q}_p \subset K$ and it is clear that $K(z) \subset E_K$. Similarly, we also have $E_K \subset E'_K$ if $K \subset K'$.

Now, let $D_0 = \{x \in \mathbb{C}_p : |x| < 1\}$. We then denote by $K_0(z)$ the ring of rational functions $A(z)/B(z) \in K(z)$ such that $B(x) \neq 0$ for all $x \in D_0$. We denote by $E_{0,K}$ the completion of $K_0(z)$ with respect to the Gauss norm. It is clear that $E_{0,K} \subset E_K$ and that $E_{0,K} \subset K[[z]]$. When $K = \mathbb{C}_p$, the ring E_{0,\mathbb{C}_p} is denoted $E_{0,p}$.

2.2. Strong Frobenius structure and MOM operators

We recall that two matrices $B_1, B_2 \in M_n(E_K)$ are E_p -equivalent if there exists $T \in GL_n(E_p)$ such that $\delta T = B_1T - TB_2$, where $\delta = z \frac{d}{dz}$. We also recall that an endomorphism $\sigma : K \to K$ is a Frobenius endomorphism if σ is continuous and, for all $x \in K$ such that $|x| \leq 1$, we have $|\sigma(x) - x^p| < 1$. Let us assume that K is equipped with a Frobenius endomorphism σ . We put $\boldsymbol{\sigma} = K(z) \to E_K$ given by $\boldsymbol{\sigma}(\sum_{i=0}^l a_i z^i / \sum_{j=0}^s b_j z^j) = \sum_{i=0}^l \sigma(a_i) z^{ip} / \sum_{j=0}^s \sigma(b_j) z^{jp}$. Since σ is continuous, $\boldsymbol{\sigma}$ is continuous. So, $\boldsymbol{\sigma}$ extends to a continuous endomorphism of E_K which is denoted again by $\boldsymbol{\sigma}$.

Let $L = \delta^n + a_1(z)\delta^{n-1} + \cdots + a_{n-1}(z)\delta + a_n(z)$ be a differential operator in $E_K[\delta]$. Then the companion of L is the matrix

$$A = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & 1 \\ -a_n(z) & -a_{n-1}(z) & -a_{n-2}(z) & \dots & -a_2(z) & -a_1(z) \end{pmatrix}$$

We recall We say that L has a strong Frobenius structure of period h if the matrices A and $p^h \sigma^h(A)$ are E_p -equivalent, where $\sigma^h(A)$ is the matrix obtained after applying σ^h to each entry of A. Notice that if K is totally ramified over \mathbb{Q}_p (that is, its residue field is \mathbb{F}_p) then the identity $i = K \to K$ is a Frobenius endomorphism. From now on, will say that K is a Frobenius field if the identity map $i: K \to K$ is a Frobenius endomorphism. Therefore, if K is a Frobenius field then $L \in E_K[\delta]$ has a strong Frobenius structure of period h if the matrices A and $p^h A(z^{p^h})$ are E_p -equivalent. Here is two important examples of Frobenius field.

• The field \mathbb{Q}_p is a Frobenius field because the residue field of \mathbb{Q}_p is \mathbb{F}_p .

• Any root of $X^{p-1} + p$ is called the *Dwork's contant* and it will be denoted by π_p . The field $\mathbb{Q}_p(\pi_p)$ is a Frobenius field. Indeed, since π_p is a root of $X^{p-1} + p$ and this polynomial is an Eisenstein polynomial, we get that $\mathbb{Q}_p(\pi_p)$ is totally ramified over \mathbb{Q}_p and therefore, $\mathbb{Q}_p(\pi_p)$ is a Frobenius field.

Let L be in $\mathbb{Q}(z)[\delta]$ and let A be its companion matrix. Remark 2.1 implies $L \in E_{\mathbb{Q}_p}[\delta]$. Then, we will say that L has a *strong Frobenius structure* of period h for p if L seen as element of $E_{\mathbb{Q}_p}[\delta]$ has strong Frobenius structure of period h. Since \mathbb{Q}_p is a Frobenius field, that is equivalent to saying that A and $p^h A(z^{p^h})$ are E_p -equivalent.

Finally, we say that $L = \delta^n + a_1(z)\delta^{n-1} + \cdots + a_{n-1}(z)\delta + a_n(z) \in E_K[\delta]$ is MOM (maximal order of multiplicity) at zero if, for all $i \in \{1, \ldots, n\}$, $a_i(z) \in K[[z]]$ and $a_i(0) = 0$. Actually, when $L \in \mathbb{Q}(z)[\delta]$ being MOM at zero is equivalent to saying that L is MUM (maximal unipotent monodromy) at zero.

2.3. Main results

We first introduce the set of power series $\mathcal{MF}(K)$.

Definition 2.2. — Let $K \subset \mathbb{C}_p$ be a complete extension of \mathbb{Q}_p . We let $\mathcal{MF}(K)$ denote the set of power series $f(z) \in 1 + zK[[z]]$ such that f(z) is a solution of a monic differential operator $L = \delta^n + a_1(z)\delta^{n-1} + \cdots + a_{n-1}(z)\delta + a_n(z) \in E_{0,K}[\delta]$ with the following properties:

- (1) L is MOM at zero;
- (2) L has a strong Frobenius structure;
- (3) $|a_i(z)|_{\mathcal{G}} \leq 1$ for all $1 \leq i \leq n$.

We are now ready to state our main results.

Theorem 2.3. — Let K be a finite extension of \mathbb{Q}_p and $f_1(z), \ldots, f_m(z)$ be power series in $\mathcal{MF}(K)$. If K is a Frobenius field then $f_1(z), \ldots, f_m(z)$ are algebraically dependent over E_K if and only if there exist $a_1, \ldots, a_m \in \mathbb{Z}$ not all zero, such that

$$f_1(z)^{a_1}\cdots f_m(z)^{a_m}\in E_{0,K}.$$

The main ingredients in the proof of Theorem 2.3 are Theorem 3.1 and Kolchin's Theorem (see Theorem 3.2). Actually, Theorem 3.1 says that if f(z) belongs to $\mathcal{MF}(K)$ with K a Frobenius field then f'(z)/f(z), the logarithmic derivative of f(z), belongs to $E_{0,K}$. Finally, by Kolchin's Theorem, our Theorem 2.3 follows.

Remark 2.4. — We would like to point out that the strategy used to prove Theorem 2.3 also allows us to study the algebraic independence of power series that are not necessarily in $\mathcal{MF}(K)$. For example, let us take f(z) and g(z) in $\mathcal{MF}(K)$.

- (1) Not necessarily f(z)g(z) is in $\mathcal{MF}(K)$. Nevertheless, we are able to study the algebraic independence of f(z)g(z) and $h(z) \in K[[z]]$ such that $h'(z)/h(z) \in E_{0,K}$. In fact, since f(z) and $g(z) \in \mathcal{MF}(K)$, Theorem 3.1 implies that f'(z)/(f) and g'(z)/g(z) are in $E_{0,K}$ and thus $f'(z)/(f) + g'(z)/g(z) \in E_{0,K}$. But, it is easy to see that f(z)g(z) is solution of $\frac{d}{dz} - f'(z)/(f) - g'(z)/g(z)$. So, from Kolchin's Theorem, we get that f(z)g(z) and h(z) are algebraically dependent over $E_{0,K}$ if and only if there are $a, b \in \mathbb{Z}$ not all zero such that $(f(z)g(z))^ah(z)^b \in E_{0,K}$.
- (2) Not necessarily f'(z) is in $\mathcal{MF}(K)$. Nevertheless, thanks to Theorem 2.5, we can study the algebraic independence of f'(z) and g'(z) over E_K .

Given a power series f(z) and an integer r > 0, $f^{(r)}(z)$ is the r-th derivative of f(z).

Theorem 2.5. — Let K be a finite extension of \mathbb{Q}_p and $f_1(z), \ldots, f_m(z)$ be in $\mathcal{MF}(K)$ such that, for all $1 \leq i \leq m$, $f_i^{(r)}(z)$ is not zero for any integer $r \geq 0$. If K is a Frobenius field then the following statements are equivalent:

ALGEBRAIC INDEPENDENCE

- (i) $f_1(z), \ldots, f_m(z)$ are algebraically dependent over E_K ;
- (ii) for all $(r_1, \ldots, r_m) \in \mathbb{N}^m$, $f_1^{(r_1)}, \ldots, f_m^{(r_m)}$ are algebraically dependent over E_K ; (iii) for some $(s_1, \ldots, s_m) \in \mathbb{N}^m$, $f_1^{(s_1)}, \ldots, f_m^{(s_m)}$ are algebraically dependent over E_K .

2.4. Algebraic independence of G-functions and E-functions over the field of analytic elements

Theorems 2.3 and 2.5 put us in a position to study the algebraic independence of G-functions and E-functions over the field E_p because numerous G-functions belong to $\mathcal{MF}(\mathbb{Q}_p)$ and some E-functions belong to $\mathcal{MF}(K)$ with $K = \mathbb{Q}_p(\pi_p)$, where π_p is the Dwork's constant. Let us see that in more detail.

• G-functions

A very important class of G-functions is given by diagonals of rational functions. We recall that $f(z) = \sum_{n \ge 0} a_n z^n \in \mathbb{Q}[[z]]$ is a diagonal of a rational function if there are an integer m > 0 and $F(z_1, \ldots, z_m) = \sum_{(i_1, \ldots, i_m) \in \mathbb{N}^m} a_{(i_m, \ldots, i_m)} z_1^{i_1} \cdots z_m^{i_m} \in \mathbb{Q}[[z_1, \ldots, z_m]] \cap \mathbb{Q}(z_1, \ldots, z_m)$ such that, for all integers $n \ge 0$, $a_{(n,\dots,n)} = a_n$. An interesting example of diagonals of rational functions is given by the hypergeometric series of the shape

$$f_{\alpha} = \sum_{j \ge 0} \frac{(\alpha_1)_j \cdots (\alpha_n)_j}{j!^n} z^j,$$

where $\boldsymbol{\alpha} = (\alpha_1, \ldots, \alpha_n) \in \mathbb{Q}^n$ and $(x)_i$ denotes the Pochhammer symbol, that is, $(x)_0 = 1$ and $(x)_{j} = x(x+1)\cdots(x+j-1)$ for j > 0.

Notice that f_{α} is the Hadamard product ⁽³⁾ of $(1-z)^{-\alpha_1}, \ldots, (1-z)^{-\alpha_n}$. Further, for every $i \in \{1, \ldots, n\}, (1-z)^{-\alpha_i}$ is algebraic over $\mathbb{Q}(z)$. Then, by Furstenberg [20], $(1-z)^{-\alpha_i}$ is a diagonal of a rational function. Consequently, f_{α} is a diagonal of a rational function because it is well-known that the Hadamard product of diagonals of rational functions is again a diagonal of a rational function.

Let d_{α} be the least common multiple of the denominators of α_i 's. If for every $1 \leq i \leq n, \alpha_i - 1 \notin \mathbb{Z}$ and p does not divide d_{α} then f_{α} belongs to $\mathcal{MF}(\mathbb{Q}_p)$. Indeed f_{α} is solution of the hypergeometric operator $\mathcal{H}_{\alpha} = \delta^n - z \prod_{i=1}^n (\delta + \alpha_i)$. This differential operator is MOM at zero and, by [28, Theorem 6.2], the assumption $\alpha_i - 1 \notin \mathbb{Z}$ for every $1 \leq i \leq n$ implies that \mathcal{H}_{α} has a strong Frobenius structure for p.

More in general, we have the following situation. Let f(z) be a diagonal of rational function. Then, thanks to Christol [11], we know that f(z) is solution of a Picard-Fuchs equation. It turns out that a lot diagonal of rational functions are solutions of MOM operators. So, let us show that if f(z) is solution of a MOM differential operator $\mathcal{D} \in \mathbb{Q}(z)[\delta]$ then f(z) belongs $\mathcal{MF}(\mathbb{Q}_p)$ for almost every prime number p. In fact, let $\mathcal{R} \in \mathbb{Q}(z)[\delta]$ be the minimal differential operator of f(z) and let N be the differential module over E_p associated to R. We are going to prove that \mathcal{R} is MOM at zero and has a strong Frobenius structure for almost every prime number p. Because of the minimality of \mathcal{R} , we get $\mathcal{D} \in \mathbb{Q}(z)[\delta]\mathcal{R}$ and as \mathcal{D} is MOM at zero then \mathcal{R} is MOM at zero. Now, we will see that N has a strong Frobenius structure for almost every prime number p. By Christol [11], we know that f(z) is solution of a Picard-Fuchs equation $\mathcal{L} \in \mathbb{Q}(z)[\delta]$. Let M be the differential module over E_p associated to \mathcal{L} . Then, according to [13, Theorem 4.2.6] or [10, Proposition 7.1], M is semi-simple and since N

⁽³⁾ Given two power series $f(z) = \sum_{n \ge 0} z^n$ and $g(z) = \sum_{n \ge 0} z^n$, the Hadamard product of f(z) and g(z) is the power series $\sum_{n \ge 0} a_n b_n z^n$

is a submodule of M, N is also semi-simple. In particular, $N = \bigoplus_{i=1}^{r} N_i$, where the N_i 's are simple modules. Furthermore, according to Theorem 22.2.1 of [21], M has a strong Frobenius structure of period 1 for almost every prime number p. Let p be one of these prime numbers and denote by ϕ the Frobenius functor. Thanks to Krull-Schmidt's Theorem (see e.g [12, p. 83]), we conclude that there exist two different natural numbers m and m' such that $\phi^m(N_i)$ and $\phi^{m'}(N_i)$ are isomorphic as differential modules over E_p . Whence, N_i has a strong Frobenius structure for p. Finally, N has also strong Frobenius structure for p since ϕ respects the direct sum.

For example the generating power series of Apéry's numbers

$$\mathfrak{A}(z) = \sum_{n \ge 0} \left(\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 \right) z^n$$

belongs to $\mathcal{MF}(\mathbb{Q}_p)$ for almost every prime number p because it is solution of the MOM differential operator $\delta^3 - z(2\delta + 1)(17\delta^2 + 17\delta + 5) - z^2(\delta + 1)^3$ and, by Theorem 3.5 of [6], $\mathfrak{A}(z)$ is a diagonal of a rational function.

As an illustration of Theorems 2.3 and 2.5, we show the algebraic independence over $E_{\mathbb{Q}_3}$ of some hypergeometric series f_{α} and $\mathfrak{A}(z)$.

Theorem 2.6. Let $\alpha = (-1/2, 1/2, 1/2)$, $\beta = (1/2, 1/2, 1/2)$, and $\gamma = (1/2, 1/2)$ and let $\mathfrak{f}(z) = f_{\alpha}$, $\mathfrak{h}(z) = f_{\beta}$, and $\mathfrak{g}(z) = f_{\gamma}$ and let K be a finite extension of \mathbb{Q}_3 . If K is a Frobenius field then, for all integers $r, s \ge 0$, we have

- (i) $\mathfrak{f}^{(s)}(z)$ and $\mathfrak{h}^{(r)}(z)$ are algebraically dependent over E_K . More precisely, $\mathfrak{f}^{(s)}(z)/\mathfrak{h}^{(r)}(z)$ belongs to E_K ;
- (ii) $\mathfrak{h}^{(r)}(z)$ and $\mathfrak{g}^{(s)}(z)$ are algebraically independent over E_K ;
- (iii) $f^{(r)}(z)$ and $g^{(s)}(z)$ are algebraically independent over E_K .

We also have

Theorem 2.7. Let $\alpha = (-1/2, 1/2, 1/2)$, $\beta = (1/2, 1/2, 1/2)$ and let $\mathfrak{f}(z) = f_{\alpha}$, $\mathfrak{h}(z) = f_{\beta}$ and let K be a finite extension of \mathbb{Q}_3 . If K is a a Frobenius field then, for all integers $r, s \ge 0$, we have

- (i) $\mathfrak{h}^{(r)}(z)$ and $\mathfrak{A}^{(s)}(z)$ are algebraically independent over E_K ;
- (ii) $\mathfrak{f}^{(r)}(z)$ and $\mathfrak{A}^{(s)}(z)$ are algebraically independent over E_K .

Remark 2.8. — By using the main result of [1] we can prove that $\mathfrak{h}(z)$ and $\mathfrak{g}(z)$ are algebraically independent over $\mathbb{Q}(z)$ and the main results of [29] imply that $\mathfrak{f}(z)$ and $\mathfrak{g}(z)$ are algebraically independent over $\mathbb{Q}(z)$. We recall that the approach used in the works [1, 29] is based on the fact the fact the power series $\mathfrak{f}(z)$, $\mathfrak{h}(z)$ and $\mathfrak{A}(z)$ satisfy "*p*-Lucas" congruences for almost every prime number *p*. It turns out that the derivatives of these power series do not satisfy "*p*-Lucas" congruences for any prime number *p* and thus, we are not able to use the results of [1, 29] in order to study the algebraic independence of the derivatives. Nevertheless, as Theorems 2.6 and 2.7 show our approach allows to study the algebraic independence of the derivatives. Furthermore, as we have already said in the introduction, in the approach used in [1, 29] it is necessary to work with infinitely many primes *p* whereas our approach works for a single prime number *p*.

• E-functions

Here we give some examples of E-functions satisfying the assumptions of Theorem 2.3. Let π_p be the Dwork's constant. Then, for every p, the exponential

$$\exp(\pi_p z) = 1 + \frac{\pi_p}{1} z + \frac{\pi_p^2}{2!} z^2 + \dots + \frac{\pi_p^n}{n!} z^n + \dotsb$$

is a E-function and it is solution of the differential operator $\delta - z\pi_p$. According to Dwork [15], for any prime number $p, \delta - z\pi_p$ has a strong Frobenius structure for p. So, $\exp(\pi_p z)$ belongs to $\mathcal{MF}(\mathbb{Q}_p(\pi_p))$. We recall that $\mathbb{Q}_p(\pi_p)$ is a Frobenius field for any p. Another example is given by the Bessel function

$$J_0(\pi_p z) = \sum_{n \ge 0} \frac{(-1)^n \pi_p^{2n}}{4^n (n!)^2} z^{2n}$$

The Bessel function is solution of $\delta^2 - \pi_p^2 z^2$. It is clear that this differential operator is MOM at zero and, by Dwork [17], if $p \neq 2$ then it has strong Frobenius structure. So, $J_0(\pi_p z) \in \mathcal{MF}(\mathbb{Q}_p(\pi_p))$ for any $p \neq 2$. Another examples are also given by Dwork in his work [16]. More precisely, Theorem 3.1 of [16] implies that the logarithmic derivative of some hypegeometric E-functions belong to $E_{0,K}$ with $K = \mathbb{Q}_p(\pi_p)$ and thus, these hypergeometric *E*-functions are in $\mathcal{MF}(\mathbb{Q}_p(\pi_p))$.

As an illustration of Theorems 2.3 and 2.5, we show

Theorem 2.9. — Let p be an odd prime number and K be a finite extension of $\mathbb{Q}_p(\pi_p)$ and let $\mathfrak{n}(z)$ be the E-function

$$\sum_{n \ge 0} \frac{1}{n!} \left(\sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} \right) z^n.$$

If K is a Frobenius field then:

- (i) the power series $\exp(\pi_p z)$ is algebraic over E_K ;
- (ii) $J_0(\pi_p z)$ is transcendental over E_p and, for all $s \ge 0$, $J_0^{(s)}(\pi_p z)$ is transcendental over the ring $E_K[\exp(\pi_p z)];$
- (iii) For all $s \ge 0$, $\mathfrak{y}^{(s)}(\pi_p z)$ is transcendental over the ring $E_K[\exp(\pi_p z)]$.

E-functions and G-functions

Theorem 2.10. — Let K be a finite extension of $\mathbb{Q}_3(\pi_3)$. If K is a Frobenius field then, for all integers $r, s, k \ge 0$,

- (i) the power series $J_0^{(r)}(\pi_3 z)$, $\mathfrak{A}^{(s)}(z)$, and $\mathfrak{h}^{(k)}(z)$ are algebraically independent over E_K , (ii) the power series $J_0^{(r)}(\pi_3 z)$, $\mathfrak{A}^{(s)}(z)$, and $\mathfrak{f}^{(k)}(z)$ are algebraically independent over E_K .

Theorems 2.6, 2.7, 2.9, and 2.10 are proven in Section 6.

3. Proof of Theorems 2.3 and 2.5

As we have already said, the proof of Theorem 2.3 relies on the Theorem 3.1 and Kolchin's Theorem. We recall that for the field K, the set $\mathcal{O}_K = \{x \in K : |x| \leq 1\}$ is a ring and it is usually called the ring of integers of K.

Theorem 3.1. — Let K be a finite extension of \mathbb{Q}_p and f(z) be in $\mathcal{MF}(K)$. If K is a Frobenius field then:

(i) $f(z) \in 1 + z\mathcal{O}_K[[z]];$ (ii) $f(z)/f(z^{p^h}) \in E_{0,K}$, where h is the period of the strong Frobenius structure; (iii) $f'(z)/f(z) \in E_{0,K}.$

Theorem 3.2 (Kolchin's Theorem [22]). — Let R be a differential ring, B be a differential field such that $B \subset R$, and $f_1(z), \ldots, f_m(z)$ be in R. Suppose that $f'_1(z)/f_1(z), \ldots, f'_m(z)/f_m(z) \in B$. Then $f_1(z), \ldots, f_m(z)$ are algebraically dependent over B if and only if there exist $a_1, \ldots, a_m \in \mathbb{Z}$ not all zero, such that

$$f_1(z)^{a_1}\cdots f_m(z)^{a_m}\in B.$$

We are going to apply Kolchi's Theorem by taking R as \mathcal{A}_K and B as E_K .

Proof of Theorem 2.3. —

Let $f_1(z), \ldots, f_m(z) \in \mathcal{MF}(K)$. According to (i) of Theorem 3.1, $f_1(z), \ldots, f_m(z)$ belong to $1 + z\mathcal{O}_K[[z]]$ and thus, $f_1(z), \ldots, f_m(z) \in \mathcal{A}_K$. It also follows from (iii) of Theorem 3.1 that $f'_1(z)/f_1(z), \ldots, f'_m(z)/f_m(z)$ belong to $E_{0,K}$. Since $E_{0,K} \subset E_K$, by Kolchin's Theorem, we conclude that $f_1(z), \ldots, f_m(z)$ are algebraically dependent over E_K if and only if there exist $a_1, \ldots, a_m \in \mathbb{Z}$ not all zero, such that

$$f_1(z)^{a_1}\cdots f_m(z)^{a_m}\in E_K.$$

In order to complete the proof, we need to show that $h(z) = f_1(z)^{a_1} \cdots f_m(z)^{a_m} \in E_{0,K}$. By (i) of Theorem 3.1 again, we deduce that, for every $1 \leq i \leq m$, $f_i(z)$ converges in $D_0 = \{x \in \mathbb{C}_p : |x| < 1\}$. Moreover, $|f_i(x)| = 1$ for all $x \in D_0$ because $f_i(0) = 1$ and |x| < 1. In particular, |h(x)| = 1 for all $x \in D_0$ and consequently, $h(z) \in E_{0,K}$.

In oder to prove Theorem 2.5, we first prove the following lemma.

Lemma 3.3. — Let K be a finite extension of \mathbb{Q}_p and f(z) be in $\mathcal{MF}(K)$ such that $f^{(r)}(z)$ is not zero for any integer $r \ge 0$. If K is a Frobenius field then, , for all $r \in \mathbb{N}$, $f(z)/f^{(r)}(z)$ belongs to $E_K \setminus \{0\}$.

Proof. — We proceed by induction on $r \ge 1$. As K is a Frobenius field, from Theorem 3.1, we deduce that $f(z)/f'(z) \in E_K \setminus \{0\}$. Now, we assume that $f(z)/f^{(r)}(z) \in E_K \setminus \{0\}$. Thus, $\frac{d}{dz}(f(z)/f^{(r)}(z))$ belongs to E_K and it is clear that

$$\frac{d}{dz}\left(\frac{f(z)}{f^{(r)}(z)}\right) = \frac{f'(z)}{f^{(r)}(z)} - \frac{f(z)f^{(r+1)}(z)}{(f^{(r)}(z))^2}.$$

Since, by hypothesis induction, $f^{(r)}(z)/f(z) \in E_K$, we have

$$\frac{f^{(r)}(z)}{f(z)} \cdot \frac{d}{dz} \left(\frac{f(z)}{f^{(r)}(z)} \right) = \frac{f'(z)}{f(z)} - \frac{f^{(r+1)}(z)}{f^{(r)}(z)} \in E_K.$$

But, from Theorem 3.1 again, we have $f'(z)/f(z) \in E_K$ and thus, $f^{(r+1)}(z)/f^{(r)}(z) \in E_K$. But, by hypothesis induction again, we know that $f^{(r)}(z)/f(z) \in E_K$ and therefore,

$$\frac{f^{(r)}(z)}{f(z)} \cdot \frac{f^{(r+1)}(z)}{f^{(r)}(z)} = \frac{f^{(r+1)}(z)}{f(z)} \in E_K \setminus \{0\}.$$

Proof of Theorem 2.5. —

 $(1) \Rightarrow (2)$. Let $(r_1, \ldots, r_m) \in \mathbb{N}^m$. Since $f_1(z), \ldots, f_m(z)$ are algebraically dependent over E_K , by Theorem 2.3, there are $a_1, \ldots, a_m \in \mathbb{Z}$ not all zero such that

$$f_1(z)^{a_1}\cdots f_m(z)^{a_m} \in E_{0,K} \setminus \{0\}$$

Further, it is clear that

$$f_1(z)^{a_1} \cdots f_m(z)^{a_m} = \left(\frac{f_1(z)}{f_1^{(r_1)}(z)}\right)^{a_1} \left(f_1^{(r_1)}(z)\right)^{a_1} \cdots \left(\frac{f_m(z)}{f_m^{(r_m)}(z)}\right)^{a_m} \left(f_m^{(r_m)}(z)\right)^{a_m}$$
$$= \left(\frac{f_1(z)}{f_1^{(r_1)}(z)}\right)^{a_1} \cdots \left(\frac{f_m(z)}{f_m^{(r_m)}(z)}\right)^{a_m} \left(f_1^{(r_1)}(z)\right)^{a_1} \cdots \left(f_m^{(r_m)}(z)\right)^{a_m}$$

By Lemma 3.3, for all $1 \leq i \leq m$, $f_i(z)/f_i^{(r_i)}(z) \in E_K \setminus \{0\}$ and thus

$$\left(\frac{f_1(z)}{f_1^{(r_1)}(z)}\right)^{a_1}\cdots\left(\frac{f_m(z)}{f_m^{(r_m)}(z)}\right)^{a_m}\in E_K\setminus\{0\}.$$

Hence

$$\left(f_1^{(r_1)}(z)\right)^{a_1}\cdots\left(f_m^{(r_m)}(z)\right)^{a_m} = f_1(z)^{a_1}\cdots f_m(z)^{a_m} \left(\frac{f_1^{(r_1)}(z)}{f_1(z)}\right)^{a_1}\cdots\left(\frac{f_m^{(r_m)}(z)}{f_m(z)}\right)^{a_m} \in E_K \setminus \{0\}.$$

Consequently, $f_1^{(r_1)}, \ldots, f_m^{(r_m)}$ are algebraically dependent over E_K .

 $(2) \Rightarrow (3)$ It is clear.

(3) \Rightarrow (1) Suppose that there exists $(s_1, \ldots, s_m) \in \mathbb{N}^m$ such that $f_1^{(s_1)}, \ldots, f_m^{(s_m)}$ are algebraically dependent over E_K . So

$$P(f_1^{(s_1)},\ldots,f_m^{(s_m)}) = \sum_{(i_1,\ldots,i_m)\in\mathbb{N}^m} b(i_1,\ldots,i_m) \left(f_1^{(s_1)}\right)^{i_1}\cdots\left(f_m^{(s_m)}\right)^{i_m} = 0,$$

where $P(X_1, ..., X_m) = \sum_{(i_1, ..., i_m) \in \mathbb{N}^m} b(i_1, ..., i_m) X_1^{i_1} \cdots X_m^{i_m} \in E_K[X_1, ..., X_m] \setminus \{0\}$. Therefore, $Q(f_1, ..., f_m) = 0$, where

$$Q(X_1, \dots, X_m) = \sum_{(i_1, \dots, i_m) \in \mathbb{N}^m} b(i_1, \dots, i_m) \left(\frac{f_1^{(s_1)}}{f_1}\right)^{i_1} \cdots \left(\frac{f_m^{(s_m)}}{f_m}\right)^{i_m} X_1^{i_1} \cdots X_m^{i_m}.$$

But $Q(X_1, \ldots, X_m) \in E_K[X_1, \ldots, X_m] \setminus \{0\}$ because by Lemma 3.3, for all $1 \leq i \leq m$, $f_i^{(s_i)}/f_i \in E_K \setminus \{0\}$. For this reason, $f_1(z), \ldots, f_m(z)$ are algebraically dependent over E_K .

4. Constructing Frobenius antecedents and Cartier operators

The aim of this section is to prove (i) of Theorem 3.1. To this end, we provide an explicit construction of the *Frobenius antecedent* for MOM differential operators $\mathcal{L} \in E_p[\delta]$. Our construction is based on the Christol's work [9]. We say that a matrix $A \in M_n(E_p)$ has a Frobenius antecedent if there exists a matrix $B \in M_n(E_p)$ such that A and $pB(z^p)$ are E_p -equivalent. So, the goal of this section is to prove that if $\mathcal{L} \in E_p[\delta]$ is a MOM differential operator with a strong Frobenius structure and A is the companion matrix of \mathcal{L} , then there is a MOM differential operator $\mathcal{L}_1 \in E_p[\delta]$ with a strong Frobenius structure such that A and $pA_1(z^p)$ are E_p -equivalent, where A_1 is the companion matrix of \mathcal{L}_1 . By definition, this means that there is a matrix $H_1 \in GL_n(E_p)$ such that $\delta H = AH_1 - pH_1A(z^p)$. The matrix H_1 is called the *matrix of passage*. Using the *Cartier operator*, we provide in Proposition 4.4 an explicit construction of H_1 and prove some *p*-adic properties of the matrix H_1 that are fundamental in the proof of Theorem 3.1. For this purpose, we first introduce the set \mathcal{M}_K .

Definition 4.1. — Let K be a complete extension of \mathbb{Q}_p . The set \mathcal{M}_K is the set of square matrices A with coefficients in E_K satisfying the following conditions:

- (i) A is a matrix with coefficients in $E_{0,K}$.
- (ii) The eigenvalues of A(0) are all equal to zero.
- (iii) There is a square matrix U invertible with coefficients in the generic disc $D(t, 1^{-})$ such that $\delta U = AU$.

The relation between the sets $\mathcal{MF}(K)$ and \mathcal{M}_K is given by the fact that if $\mathcal{L} \in E_{0,K}[\delta]$ is MOM at zero and has a strong Frobenius structure then the companion matrix of \mathcal{L} is in \mathcal{M}_K . Indeed, if A is the companion matrix of \mathcal{L} then it is clear that A is a matrix with coefficients in $E_{0,K}$ and the MOM condition is equivalent to the fact that the eigenvalues of A(0) are all equal to zero. Finally, since \mathcal{L} has a strong Frobenius structure, it follows from Propositions 4.1.2, 4.6.4, and 4.7.2 of [8] that there exists a matrix U invertible with coefficients in the generic disc $D(t, 1^-)$ such that $\delta U = AU$.

Remark 4.2. — Let K be a complete extension of \mathbb{Q}_p . Suppose that $A \in M_n(E_{0,K})$ and that the eigenvalues of A(0) are all equal to zero. In particular, the eigenvalues of A(0) are prepared⁽⁴⁾. Moreover, $A \in M_n(K[[z]])$ because $E_{0,K} \subset K[[z]]$. Therefore, it follows from [18, Chap. III, Proposition 8.5] that the differential system $\delta X = AX$ has a fundamental matrix of solutions of the shape $Y_A X^{A(0)}$, where $Y_A \in GL_n(K[[z]])$, $Y_A(0) = I$ and $X^{A(0)} = \sum_{j \ge 0} A(0)^j \frac{(Logz)^j}{j!}$. Since the eigenvalues of A(0) are all equal to zero, we have $A(0)^n = 0$. For this reason, $X^{A(0)} = \sum_{j=0}^{n-1} A(0)^j \frac{(Logz)^j}{j!}$. The matrix Y_A will be called the *uniform part* of the system $\delta X = AX$. Now, let L be in $E_{0,K}[\delta]$ and let A be its companion matrix. It is clear that if L is MOM at zero then the eigenvalues of A(0) are all equal to zero. Then, from the above argument, we deduce that L has a solution $f(z) \in 1 + zK[[z]]$. Moreover, since Logz is transcendental over K[[z]], it follows that if $g(z) \in K[[z]]$ is a solution of L then there exists a constant $c \in K$ such that g(z) = cf(z).

4.1. Constructing Frobenius antecedents

An important ingredient in the construction of the Frobenius antecedent for the elements in \mathcal{M}_K is the *Cartier operator*. We recall that the Cartier operator is the \mathbb{C}_p -linear map $\Lambda_p : \mathbb{C}_p[[z]] \to \mathbb{C}_p[[z]]$ given by $\Lambda_p(\sum_{n \ge 0} a_n z^n) = \sum_{n \ge 0} a_{np} z^n$. If A is a matrix with coefficients in $\mathbb{C}_p[[z]]$, the matrix $\Lambda_p(A)$ is the matrix obtained after applying Λ_p to each entry of A.

In [9], Christol gave a construction of Frobenenius antecedent for a more general set than the set \mathcal{M}_K . Nevertheless, our main observation is that when A belongs to \mathcal{M}_K we can construct the matrix of passage explicitly. This is done in Proposition 4.4. To prove this proposition, we need the following

⁽⁴⁾ Given a matrix $C \in M_n(\mathbb{C}_p)$, we say that their eigenvalues are prepared if the following condition holds: given two eigenvalues λ and α then, $\lambda - \alpha$ is an integer if only if $\lambda = \alpha$.

ALGEBRAIC INDEPENDENCE

lemma whose proof follows the same line as the proof of Lemma 5.1 of [9]. Finally, given a matrix $A = (a_{i,j})_{1 \leq i,j \leq n}$ with coefficients in E_K , we set $||A|| = \max\{|a_{i,j}|_{\mathcal{G}}\}_{1 \leq i,j \leq n}$.

Lemma 4.3. — Let K be a finite extension of \mathbb{Q}_p . Let \mathcal{L} be a MOM differential operator in $E_{0,K}[\delta]$ of order n, let A be the companion matrix of \mathcal{L} , and let $Y_A X^{A(0)}$ be a fundamental matrix of solutions of $\delta X = AX$ with Y_A the uniform part. If A belongs to \mathcal{M}_K and $||A|| \leq 1$ then there exists $F \in M_n(E_{0,K})$ such that the matrices A and $pF(z^p)$ are E_K -equivalent. Moreover, the matrix $H_0 = Y_A(\Lambda_p(Y_A)(z^p))^{-1}$ belongs to $GL_n(E_{0,K})$, $||H_0|| = 1 = ||H_0^{-1}||$,

$$\delta H_0 = pF(z^p)H_0 - H_0A,$$

and a fundamental matrix of solutions of $\delta X = FX$ is given by $\Lambda_p(Y_A) X^{\frac{1}{p}A(0)}$.

Proof. — We first construct the matrix H_0 . For this purpose, let us consider the sequence $\{A_j(z)\}_{j \ge 0}$, where $A_0(z) = I$ is the identity matrix and $A_{j+1}(z) = \delta A_j(z) + A_j(z)(A(z) - jI)$. By assumption, $A \in M_n(E_{0,K})$. Then, $A_j(z) \in M_n(E_{0,K})$ for all $j \ge 0$. Since $X^p - 1 = (X - 1)((X - 1)^{p-1} + pt(X))$ with $t(X) \in \mathbb{Z}[X]$, it follows that if $\xi^p = 1$ then $|\xi - 1| \le |p|^{1/(p-1)}$. Furthermore, for all integers $j \ge 1$, we have $|p|^{j/(p-1)} < |j!|$. Thus, for all $j \ge 1$, $|(\xi - 1)^j/j!| < 1$. Since the norm is non-Archimedean and $\sum_{\xi^p=1} (\xi - 1)^j/j$ is a rational number, we conclude that, for all $j \ge 0$, $\left|\sum_{\xi^p=1} \frac{(\xi - 1)^j}{j!}\right| \le 1/p$. Thus, $\left|\sum_{\xi^p=1} \frac{(\xi - 1)^j}{pj!}\right| \le 1$. That is, for all $j \ge 0$, $\sum_{\xi^p=1} \frac{(\xi - 1)^j}{pj!} \in \mathbb{Z}_p$. So, for all $j \ge 0$, the matrix

$$A_j\left(\sum_{\xi^p=1}\frac{(\xi-1)^j}{pj!}\right)$$

belongs to $M_n(E_{0,K})$. We set

$$H_0 = \sum_{j \ge 0} A_j(z) \left(\sum_{\xi^p = 1} \frac{(\xi - 1)^j}{pj!} \right).$$

Let us show that $H_0 \in GL_n(E_{0,K})$. We first prove that $H_0 \in M_n(E_{0,K})$. That is equivalent to saying that

$$\lim_{j \to \infty} ||A_j|| \left| \sum_{\xi^p = 1} \frac{(\xi - 1)^j}{pj!} \right| = 0$$
(4.1)

because $E_{0,K}$ is complete with respect to the Gauss norm and it is non-Archimedean. Indeed, as there is a square matrix U invertible in the generic disc $D(t, 1^-)$ such that $\delta U = AU$, it follows from Remark b) of [9, p.155], $\lim_{j\to\infty} ||A_j|| = 0$. Therefore, (4.1) follows immediately since, for all $j \ge 0$, $\left|\sum_{\xi^p=1} \frac{(\xi-1)^j}{pj!}\right| \le 1$.

Further, we have
$$||H_0|| \leq 1$$
. Indeed, by assumption, $||A|| \leq 1$ and consequently, $||A_j|| \leq 1$ for all $j \geq 0$. We also have proven that, for all $j \geq 0$, $\sum_{\xi^{p=1}} \frac{(\xi-1)^j}{pj!} \in \mathbb{Z}_p$ and therefore, for all $j \geq 0$, $||A_j|| \left| \sum_{\xi^{p=1}} \frac{(\xi-1)^j}{pj!} \right| \leq 1$. As the Gauss norm is non-Archimedean, we conclude that $||H_0|| \leq 1$.

Now, we are going to prove that $H_0 = \Lambda_p(Y_A)(z^p)Y_A^{-1}$. Let us write $Y_A = \sum_{j \ge 0} Y_j z^j$. We have $A(0)^n = 0$ because L is MOM and thus, it follows from [9, p.165] that $H_0Y_A = \sum_{j \ge 0} Y_{jp}z^{jp}$. So, $H_0Y_A = \Lambda_p(Y_A)(z^p)$. Therefore, $H_0 = \Lambda_p(Y_A)(z^p)Y_A^{-1}$. Finally, $H_0 \in GL_n(E_{0,K})$ because $H_0(0)$ is the identity matrix. In particular, $||H_0|| = 1 = ||H_0^{-1}||$.

Now, we proceed to construct the matrix F. For that, we set

$$F(z) = [\delta(\Lambda_p(Y_A)) + \frac{1}{p}\Lambda_p(Y_A)A(0)](\Lambda_p(Y_A))^{-1}.$$
(4.2)

As $H_0Y_A = \Lambda_p(Y_A)(z^p)$ then, from Equation (4.2) we obtain,

$$pF(z^p) = [p(\delta(\Lambda_p(Y_A)))(z^p) + H_0Y_AA(0)][Y_A^{-1}H_0^{-1}].$$

We also have

$$\delta(H_0)Y_A + H_0(\delta Y_A) = \delta(H_0Y_A) = \delta(\Lambda_p(Y_A)(z^p)) = p(\delta(\Lambda_p(Y_A)))(z^p).$$

Since $\delta(Y_A z^{A(0)}) = A Y_A z^{A(0)}$, it follows that $\delta Y_A = A Y_A - Y_A A(0)$. Thus,

$$p(\delta(\Lambda_p(Y_A)))(z^p) = \delta(H_0)Y_A + H_0[AY_A - Y_AA(0)]$$

Then,

$$pF(z^{p}) = [p(\delta(\Lambda_{p}(Y_{A})))(z^{p}) + H_{0}Y_{A}A(0)][Y_{A}^{-1}H_{0}^{-1}]$$

= $[(\delta H_{0})Y_{A} + H_{0}AY_{A}][Y_{A}^{-1}H_{0}^{-1}]$
= $(\delta H_{0})H_{0}^{-1} + H_{0}AH_{0}^{-1}.$

Consequently,

$$\delta H_0 = pF(z^p)H_0 - H_0A.$$

As $pF(z^p) = (\delta H_0 + H_0 A) H_0^{-1}$ and $H_0 \in GL_n(E_{0,K})$, and $A \in M_n(E_{0,K})$ then, we obtain $F(z) \in M_n(E_{0,K})$.

Finally, we show that the matrix $\Lambda_p(Y_A)X^{\frac{1}{p}A(0)}$ is the fundamental matrix of solutions of $\delta X = FX$. It is clear that $\Lambda_p(Y_A)(0) = I$ and from Equality (4.2), we obtain $F(z)\Lambda_p(Y_A) = \delta(\Lambda_p(Y_A)) + \Lambda_p(Y_A)\frac{1}{p}A(0)$. Thus,

$$\begin{split} \delta(\Lambda_p(Y_L)X^{\frac{1}{p}N}) &= \delta(\Lambda_p(Y_A))X^{\frac{1}{p}A(0)} + \Lambda_p(Y_A)\frac{1}{p}A(0)X^{\frac{1}{p}A(0)} \\ &= (\delta(\Lambda_p(Y_A)) + \Lambda_p(Y_A)\frac{1}{p}A(0))X^{\frac{1}{p}A(0)} \\ &= F(z)\Lambda_p(Y_A)X^{\frac{1}{p}A(0)}. \end{split}$$

This completes the proof.

The new ingredient in our construction of the Frobenius antecedent is given by the following proposition. Actually, it follows from this proposition that if $\mathcal{L} \in E_{0,K}[\delta]$ is a MOM differential operator with a strong Frobenius structure and A is its companion matrix then there is a MOM differential operator $\mathcal{L}_1 \in E_{0,K}[\delta]$ with a strong Frobenius structure such that A and $pA_1(z^p)$ are E_p -equivalent, where A_1 is the companion matrix of \mathcal{L}_1 . The construction of \mathcal{L}_1 relies on the construction of the matrix F of the previous lemma.

Proposition 4.4. — Let the assumptions be as in Lemma 4.3 and let $f(z) \in 1 + zK[[z]]$ be a solution of \mathcal{L} . Then there exists a MOM differential operator $\mathcal{L}_1 \in E_{0,K}[\delta]$ of order n such that:

(a) $\Lambda_p(f)$ is a solution of \mathcal{L}_1 ,

(b) if A_1 is the companion matrix of \mathcal{L}_1 then a fundamental matrix of solutions of $\delta X = A_1 X$ is given by

$$\begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & 1/p & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1/p^{n-2} & 0 \\ 0 & 0 & \dots & 0 & 1/p^{n-1} \end{pmatrix} \Lambda_p(Y_A) \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & p & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & p^{n-2} & 0 \\ 0 & 0 & \dots & 0 & p^{n-1} \end{pmatrix} X^{A(0)},$$

- (c) A_1 belongs to \mathcal{M}_K and $||A_1|| \leq 1$,
- (d) the matrices A and $pA_1(z^p)$ are E_K -equivalent. Moreover,

$$\delta(\widetilde{H}_1) = A\widetilde{H}_1 - p\widetilde{H}_1 A_1(z^p)$$

with

$$\widetilde{H}_{1} = Y_{A}(\Lambda_{p}(Y_{A})(z^{p}))^{-1} \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & p & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & p^{n-2} & 0 \\ 0 & 0 & \dots & 0 & p^{n-1} \end{pmatrix} \in GL_{n}(E_{0,K})$$

and $||\tilde{H}_1|| = 1$.

Proof. —

We start by constructing the MOM differential operator $\mathcal{L}_1 \in E_{0,K}[\delta]$ of order *n*. According to Lemma 4.3, there is a matrix $F \in M_n(E_{0,K})$ such that *A* and $pF(z^p)$ are E_K -equivalent. Let us write $F = (b_{i,j})_{1 \leq i,j \leq n}$. We set

$$\mathcal{L}_1 := \delta^n - b_{n,n} \delta^{n-1} - \frac{1}{p} b_{n,n-1} \delta^{n-2} - \dots + \frac{1}{p^{n-i}} b_{n,i} \delta^{i-1} - \dots - \frac{1}{p^{n-2}} b_{n,2} \delta - \frac{1}{p^{n-1}} b_{n,1}.$$

 \mathcal{L}_1 belongs to $E_{0,K}[\delta]$ because F is a matrix with coefficients in $E_{0,K}$ and it is clear that the order of \mathcal{L}_1 is n. In addition, \mathcal{L}_1 is MOM because, from Lemma 4.3, we know that

$$\delta H_0 = pF(z^p)H_0 - H_0A, \tag{4.3}$$

with $H_0 = Y_A(\Lambda_p(Y_A)(z^p))^{-1} \in GL_n(E_{0,K})$. By hypotheses, $Y_A(0) = I$. Thus, $H_0(0) = I$ and therefore, by substituting z = 0 into Equation (4.3), we get $F(0) = \frac{1}{p}A(0)$. In particular, for all $i \in \{1, \ldots, n\}$, $b_{n,i}(0) = \frac{1}{p}a_{n,i}(0)$. But, by assumption $A \in \mathcal{M}_K$ and thus, all eigenvalues of A(0) are equal to zero. Since A is the companion matrix of \mathcal{L} , that is equivalent to saying that, $a_{n,i}(0) = 0$ for all $1 \leq i \leq n$. So, for every $1 \leq i \leq n$, $b_{n,i}(0) = 0$.

We now proceed to prove the statements (a)-(d).

Proof of (a). We are going to see that $\Lambda_p(f)$ is a solution of \mathcal{L}_1 . By Lemma 4.3, we know that a fundamental matrix of solutions of $\delta X = FX$ is given by $\Lambda_p(Y_A)X^{\frac{1}{p}A(0)}$. Let us write $Y_A = (f_{i,j})_{1 \leq i,j \leq n}$. As Y_A is the uniform part of $\delta X = AX$ and $f(z) \in 1 + zK[[z]]$ is a solution of L_1 then, for all $i \in \{1, \ldots, n\}$, $f_{i,1} = \delta^{i-1}f$. Thus, the vector $(\Lambda_p(f_{1,1}), \ldots, \Lambda_p(f_{n,1}))^t$ is a solution of $\delta \vec{y} = F\vec{y}$. So, we have

$$F\begin{pmatrix}\Lambda_p(f)\\\Lambda_p(\delta f)\\\vdots\\\Lambda_p(\delta^{n-1}f)\end{pmatrix} = F\begin{pmatrix}\Lambda_p(f_{1,1})\\\Lambda_p(f_{2,1})\\\vdots\\\Lambda_p(f_{n,1})\end{pmatrix} = \begin{pmatrix}\delta(\Lambda_p(f_{1,1}))\\\delta(\Lambda_p(f_{2,1}))\\\vdots\\\delta(\Lambda_p(f_{2,1}))\end{pmatrix} = \begin{pmatrix}\delta(\Lambda_p(f))\\\delta(\Lambda_p(\delta f))\\\vdots\\\delta(\Lambda_p(\delta^{n-1}f))\end{pmatrix}.$$

Then, from the previous equality, we get

$$b_{n,1}\Lambda_p(f) + b_{n,2}\Lambda_p(\delta f) + \dots + b_{n,i}\Lambda_p(\delta^{i-1}f) + \dots + b_{n,n}\Lambda_p(\delta^{n-1}f) = \delta(\Lambda_p(\delta^{n-1}f)).$$

But, $\Lambda_p \circ \delta = p\delta \circ \Lambda_p$ and thus, we obtain

 $b_{n,1}\Lambda_p(f) + pb_{n,2}\delta(\Lambda_p(f)) + \dots + p^{i-1}b_{n,i}\delta^{i-1}(\Lambda_p(f)) + \dots + p^{n-1}b_{n,n}\delta^{n-1}(\Lambda_p(f)) = p^{n-1}\delta^n(\Lambda_p(f)).$ Consequently, $\Lambda_p(f)$ is a solution of \mathcal{L}_1 .

Proof of (b). Recall that $Y_A = (f_{i,j})_{1 \leq i,j \leq n}$. We first prove that, for all integers $i \in \{1, \ldots, n-1\}$ and all $k \in \{1, \ldots, n\}$,

$$f_{i,k-1} + \delta f_{i,k} = f_{i+1,k}.$$

For every $i, j \in \{1, \ldots, n\}$, we set $\phi_{i,j} = \sum_{k=1}^{j} f_{i,k} \frac{(Logz)^{j-k}}{(j-k)!}$. Notice that $Y_A X^{A(0)} = (\phi_{i,j})_{1 \leq i,j \leq n}$. Since, $Y_A X^{A(0)}$ is a fundamental matrix of $\delta X = AX$ and A is the companion matrix of L, it follows that, for every $i \in \{2, \ldots, n\}$ and $j \in \{1, \ldots, n\}$, $\phi_{i,j} = \delta \phi_{i-1,j}$. Therefore,

$$\sum_{k=1}^{j} f_{i+1,k} \frac{(Logz)^{j-k}}{(j-k)!} = \phi_{i+1,j} = \delta\phi_{i,j} = \delta(f_{i,1}) \frac{(Logz)^{j-k}}{(j-k)!} + \sum_{k=2}^{j} (f_{i,k-1} + \delta(f_{i,k})) \frac{(Logz)^{j-k}}{(j-k)!}.$$

As Logz is transcendental over K[[z]] then, it follows from the previous equality that, for all $k \in \{1, \ldots, n\}, f_{i,k-1} + \delta f_{i,k} = f_{i+1,k}$.

Now, we proceed to prove that

$$T = \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & 1/p & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1/p^{n-2} & 0 \\ 0 & 0 & \dots & 0 & 1/p^{n-1} \end{pmatrix} \Lambda_p(Y_A) \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & p & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & p^{n-2} & 0 \\ 0 & 0 & \dots & 0 & p^{n-1} \end{pmatrix} X^{A(0)}$$

is a fundamental matrix of solutions of $\delta X = A_1 X$, where A_1 is the companion matrix of \mathcal{L}_1 . To this end, we will first prove that, for every $j \in \{1, \ldots, n\}$,

$$\theta_j = \sum_{k=1}^{j} p^{k-1} \Lambda_p(f_{1,k}) \frac{(Logz)^{j-k}}{(j-k)!}$$

is a solution of \mathcal{L}_1 . Notice that, the vector $(\theta_1, \theta_2, \ldots, \theta_n)$ corresponds to the first row of T. Now, if we put $\Lambda_p(Y_A)X^{\frac{1}{p}A(0)} = (\eta_{i,j})_{1 \leq i,j \leq n}$ then, for every $i, j \in \{1, \ldots, n\}$,

$$\eta_{i,j} = \sum_{k=1}^{j} \Lambda_p(f_{i,k}) \frac{(Logz)^{j-k}}{p^{j-k}(j-k)!}.$$

We are going to see that, for every $j, l \in \{1, ..., n\}$, $\frac{1}{p^{n-l}} \delta^{l-1} \theta_j = \frac{1}{p^{n-j}} \eta_{l,j}$. To prove this equality, we are going to proceed by induction on $l \in \{1, ..., n\}$. For l = 1, it is clear that $\frac{1}{p^{n-1}} \theta_j = \frac{1}{p^{n-j}} \eta_{1,j}$. Now, we suppose that for some $l \in \{1, ..., n-1\}$, we have $\frac{1}{p^{n-l}} \delta^{l-1} \theta_j = \frac{1}{p^{n-j}} \eta_{l,j}$. So, $\frac{1}{p^{n-l}} \delta^l \theta_j = \frac{1}{p^{n-j}} \delta(\eta_{l,j})$.

But, $\delta(\eta_{l,j}) = \frac{1}{p} \eta_{l+1,j}$. In fact, as $\Lambda_p \circ \delta = p\delta \circ \Lambda_p$ and $f_{l,k-1} + \delta(f_{l,k}) = f_{l+1,k}$ for all $k \in \{1, \ldots, n\}$ then

$$\begin{split} \delta(\eta_{l,j}) &= \sum_{k=1}^{j} \delta(\Lambda_{p}(f_{l,k})) \frac{(Logz)^{j-k}}{p^{j-k}(j-k)!} + \Lambda_{p}(f_{l,k}) \frac{(Logz)^{j-k-1}}{p^{j-k}(j-k-1)!} \\ &= \sum_{k=1}^{j} \frac{1}{p} \Lambda_{p}(\delta(f_{l,k})) \frac{(Logz)^{j-k}}{p^{j-k}(j-k)!} + \frac{1}{p} \Lambda_{p}(f_{l,k}) \frac{(Logz)^{j-k-1}}{p^{j-k-1}(j-k-1)!} \\ &= \frac{1}{p} \left[\Lambda_{p}(\delta(f_{l,1})) \frac{(Logz)^{j-1}}{p^{j-1}(j-1)!} + \sum_{k=2}^{j} \Lambda_{p}(f_{l,k-1} + \delta f_{l,k})) \frac{(Logz)^{j-k}}{p^{j-k}(j-k)!} \right] \\ &= \frac{1}{p} \left[\Lambda_{p}(f_{l+1,1}) \frac{(Logz)^{j-1}}{p^{j-1}(j-1)!} + \sum_{k=2}^{j} \Lambda_{p}(f_{l+1,k}) \frac{(Logz)^{j-k}}{p^{j-k}(j-k)!} \right] \\ &= \frac{1}{p} \eta_{l+1,j}. \end{split}$$

Thus, $\frac{1}{p^{n-l}}\delta^l\theta_j = \frac{1}{p^{n-j}}\delta(\eta_{l,j}) = \frac{1}{p^{n-j}}(\frac{1}{p}\eta_{l+1,j})$. Whence, $\frac{1}{p^{n-l-1}}\delta^l\theta_j = \frac{1}{p^{n-j}}\eta_{l+1,j}$. So that, we conclude that, for every $j, l \in \{1, \dots, n\}, \frac{1}{p^{n-l}}\delta^{l-1}\theta_j = \frac{1}{p^{n-j}}\eta_{l,j}$.

As $\Lambda_p(Y_A)X^{\frac{1}{p}A(0)}$ is a fundamental matrix of $\delta X = FX$, it follows that, for every $j \in \{1, \ldots, n\}$,

$$F\begin{pmatrix}\eta_{1,j}\\\eta_{2,j}\\\vdots\\\eta_{n,j}\end{pmatrix} = \begin{pmatrix}\delta(\eta_{1,j})\\\delta(\eta_{2,j})\\\vdots\\\delta(\eta_{2,j})\end{pmatrix}.$$

Thus, for every $j \in \{1, \ldots, n\}$,

$$b_{n,1}\eta_{1,j} + b_{n,2}\eta_{2,j} + \dots + b_{n,k}\eta_{k,j} + \dots + b_{n,n}\eta_{n,j} = \delta(\eta_{n,j}).$$

So,

$$b_{n,1}\frac{1}{p^{n-j}}\eta_{1,j} + b_{n,2}\frac{1}{p^{n-j}}\eta_{2,j} + \dots + b_{n,k}\frac{1}{p^{n-j}}\eta_{k,j} + \dots + b_{n,n}\frac{1}{p^{n-j}}\eta_{n,j} = \frac{1}{p^{n-j}}\delta(\eta_{n,j}).$$

But, we know that, for every $l \in \{1, \ldots, n\}$, $\frac{1}{p^{n-l}}\delta^{l-1}\theta_j = \frac{1}{p^{n-j}}\eta_{l,j}$. Thus,

$$\frac{1}{p^{n-1}}b_{n,1}\theta_j + \frac{1}{p^{n-2}}b_{n,2}\delta\theta_j + \dots + \frac{1}{p^{n-k}}b_{n,k}\delta^{k-1}\theta_j + \dots + b_{n,n}\delta^{n-1}\theta_j = \delta^n\theta_j.$$

Therefore, θ_i is a solution of \mathcal{L}_1 .

Moreover, $\theta_1, \ldots, \theta_n$ are linearly independent over K because Log(z) is transcendental over K[[z]]. Since A_1 is the companion matrix of \mathcal{L}_1 and $\theta_1, \ldots, \theta_n$ are linearly independent over K, it follows that the matrix $(\delta^{i-1}\theta_j)_{1 \leq i,j \leq n}$ is a fundamental matrix of solutions of $\delta X = A_1 X$. We are going to see that $T = (\delta^{i-1}\theta_j)_{1 \leq i,j \leq n}$. As we have already seen, for every $i, j \in \{1, \ldots, n\}, \frac{1}{p^{n-i}}\delta^{i-1}\theta_j = \frac{1}{p^{n-j}}\eta_{i,j}$. Hence, for every $i, j \in \{1, \ldots, n\}$

$$\delta^{i-1}\theta_j = \frac{p^j}{p^i}\eta_{i,j} = \frac{p^j}{p^i}\sum_{k=1}^j \Lambda_p(f_{i,k}) \frac{(Logz)^{j-k}}{p^{j-k}(j-k)!} = \sum_{k=1}^j \frac{\Lambda_p(f_{i,k})}{p^{i-k}} \frac{(Logz)^{j-k}}{(j-k)!}.$$

But,

$$\begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & 1/p & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1/p^{n-2} & 0 \\ 0 & 0 & \dots & 0 & 1/p^{n-1} \end{pmatrix} \Lambda_p(Y_A) \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & p & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & p^{n-2} & 0 \\ 0 & 0 & \dots & 0 & p^{n-1} \end{pmatrix} = \left(\frac{\Lambda_p(f_{i,k})}{p^{i-k}}\right)_{1 \le i,k \le n}$$

Thus $T = (\delta^{i-1}\theta_j)_{1 \leq i,j \leq n}$. Consequently, T is a fundamental matrix of $\delta X = A_1 X$.

Proof of (c) Since \mathcal{L}_1 belongs to $E_{0,K}[\delta]$, A_1 is a matrix with coefficients in $E_{0,K}$. Further, we have already observed that $b_{n,i}(0) = 0$ for all $1 \leq i \leq n$. Thus, the eigenvalues of $A_1(0)$ are all equal to zero.

Now, we prove that there exists U_1 invertible with coefficients in $D(t, 1^-)$ such that $\delta U_1 = A_1 U_1$. Recall that $Y_A = (f_{i,j})_{1 \leq i,j \leq n}$ is the uniform part of $\delta X = AX$. Since there exists a matrix U invertible with coefficients in $D(t, 1^-)$ such that $\delta U = AU$, it follows from Theorem 2 of [9] that, for all $i, j \in \{1, \ldots, n\}$, the radius of convergence of $f_{i,j}$ is greater than or equal to 1. So, for all $i, j \in \{1, \ldots, n\}$, the radius of convergence of $\Lambda_p(f_{i,j})$ is greater than or equal to 1 because the *p*-adic norm is non-Archimedean. Further, from (b) we know that

$$\begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & 1/p & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1/p^{n-2} & 0 \\ 0 & 0 & \dots & 0 & 1/p^{n-1} \end{pmatrix} \Lambda_p(Y_A) \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & p & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & p^{n-2} & 0 \\ 0 & 0 & \dots & 0 & p^{n-1} \end{pmatrix}$$

is the uniform part of $\delta X = A_1 X$. In particular, each entry of the uniform part of $\delta X = A_1 X$ has radius of convergence greater than or equal to 1. Therefore, following Theorem 2 of [9], we conclude that there exists a matrix U_1 invertible with coefficients in $D(t, 1^-)$ such that $\delta U_1 = A_1 U_1$. Consequently, the matrix A_1 belongs to \mathcal{M}_K .

Finally, since U_1 is a invertible matrix with coefficients in $D(t, 1^-)$ such that $\delta U_1 = A_1 U_1$, the Frobenius-Dwork's Theorem [7, Proposition 8.1] implies $||A_1|| \leq 1$.

Proof of (d). Let \widetilde{H}_1 be the matrix

$$Y_A(\Lambda_p(Y_A)(z^p))^{-1} \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & p & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & p^{n-2} & 0 \\ 0 & 0 & \dots & 0 & p^{n-1} \end{pmatrix}$$

From Lemma 4.3, we know that the matrix $H_0 = Y_A(\Lambda_p(Y_A)(z^p))^{-1}$ belongs to $GL_n(E_{0,K})$. Thus, \tilde{H}_1 belongs to $GL_n(E_{0,K})$. By again invoking Lemma 4.3, we have $||H_0|| = 1$. So that, $||\tilde{H}_1|| \leq 1$. But, $\tilde{H}_1(0) = diag(1, p, \ldots, p^{n-1})$ and thus, $||\tilde{H}_1|| = 1$. Now, we show that

$$\delta(\tilde{H}_1) = A\tilde{H}_1 - p\tilde{H}_1A_1(z^p).$$

We infer from (b) that

$$T(z^{p}) = \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & 1/p & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1/p^{n-2} & 0 \\ 0 & 0 & \dots & 0 & 1/p^{n-1} \end{pmatrix} \Lambda_{p}(Y_{A})(z^{p}) \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & p & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & p^{n-2} & 0 \\ 0 & 0 & \dots & 0 & p^{n-1} \end{pmatrix} X^{pA(0)},$$

is a fundamental matrix of solutions of $\delta X = pA_1(z^p)X$. Further, it is clear that

$$\widetilde{H}_1 T(z^p) = Y_A \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & p & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & p^{n-2} & 0 \\ 0 & 0 & \dots & 0 & p^{n-1} \end{pmatrix} X^{pA(0)}.$$

But,

$$\begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & p & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & p^{n-2} & 0 \\ 0 & 0 & \dots & 0 & p^{n-1} \end{pmatrix} X^{pA(0)} = X^{A(0)} \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & p & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & p^{n-2} & 0 \\ 0 & 0 & \dots & 0 & p^{n-1} \end{pmatrix}.$$

Consequently,

$$\widetilde{H}_1 T(z^p) = Y_A X^{A(0)} \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & p & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & p^{n-2} & 0 \\ 0 & 0 & \dots & 0 & p^{n-1} \end{pmatrix}.$$

Thus, $\widetilde{H}_1T(z^p)$ is a fundamental matrix of $\delta X = AX$. In addition, we have proven in (b) that $\delta T = A_1T$ and thus, $\delta(T(z^p)) = pA_1(z^p)T(z^p)$. So, we get

$$\begin{aligned} A\widetilde{H}_1T(z^p) &= \delta(\widetilde{H}_1T(z^p)) \\ &= \delta(\widetilde{H}_1)T(z^p) + \widetilde{H}_1\delta(T(z^p)) \\ &= \delta(\widetilde{H}_1)T(z^p) + \widetilde{H}_1(pA_1(z^p))T(z^p). \end{aligned}$$

Hence, $A\widetilde{H}_1 = \delta(\widetilde{H}_1) + \widetilde{H}_1 p A_1(z^p)$. For this reason, we have

$$\delta(H_1) = AH_1 - pH_1A_1(z^p).$$

We have the following corollary

Corollary 4.5. — Let the assumptions be as in Lemma 4.3 and let $f(z) \in 1+zK[[z]]$ be a solution of \mathcal{L} . Then, for every integer m > 0, there exists a MOM differential operator $\mathcal{L}_m \in E_{0,K}[\delta]$ of order n such that:

(a) $\Lambda_p^m(f)$ is a solution of \mathcal{L}_m ,

(b) if A_m denotes the companion matrix of \mathcal{L}_m then a fundamental matrix of solutions of $\delta X = A_m X$ is given by

$$\begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & 1/p^m & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1/p^{m(n-2)} & 0 \\ 0 & 0 & \dots & 0 & 1/p^{m(n-1)} \end{pmatrix} \Lambda_p^m(Y_A) \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & p^m & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & p^{m(n-2)} & 0 \\ 0 & 0 & \dots & 0 & p^{m(n-1)} \end{pmatrix} X^{A(0)},$$

- (c) A_m belongs to \mathcal{M}_K and $||A_m|| \leq 1$,
- (d) the matrices A and $p^m A_m(z^{p^m})$ are E_K -equivalent. Moreover,

$$\delta(\tilde{H}_m) = A\tilde{H}_m - p^m\tilde{H}_m A_m(z^{p^m})$$

with

$$\widetilde{H}_m = Y_A(\Lambda_p^m(Y_A)(z^{p^m}))^{-1} \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & p^m & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & p^{m(n-2)} & 0 \\ 0 & 0 & \dots & 0 & p^{m(n-1)} \end{pmatrix} \in GL_n(E_K) \cap M_n(E_{0,K})$$

and $||\tilde{H}_m|| = 1$.

Proof. —

We proceed by induction on $m \in \mathbb{Z}_{>0}$. For m = 1 we are in a position to apply Proposition 4.4 and thus, there is a monic differential operator $\mathcal{L}_1 \in E_{0,K}[\delta]$ of order n such that the conditions (a)-(d) are satisfied. Now, suppose that for some integer m > 0 the conditions (a)-(d) are satisfied. We are going to see that for m + 1 the conditions (a)-(d) also hold. So, by induction hypothesis, there is MOM differential operator $\mathcal{L}_m \in E_{0,K}[\delta]$ of order n such that: $\Lambda_p^m(f)$ is solution of \mathcal{L}_m , the companion matrix of \mathcal{L}_m , noted by A_m , belongs to \mathcal{M}_K , $||A_m|| \leq 1$, and a fundamental matrix of solutions of the system $\delta X = A_m X$ is given by $Y_{A_m} X^{A(0)}$, where

$$Y_{A_m} = diag(1, 1/p^m, \dots, 1/p^{m(n-2)}, 1/p^{m(n-1)}) \cdot \Lambda_p(Y_A) \cdot diag(1, p^m, \dots, p^{m(n-2)}, p^{m(n-1)}).$$

We also know, by the induction hypothesis, that the matrix

$$\widetilde{H}_m = Y_A(\Lambda_p^m(Y_A)(z^{p^m}))^{-1} \cdot diag(1, p^m, \dots, p^{m(n-2)}, p^{m(n-1)})$$

belongs to $GL_n(E_K) \cap M_n(E_{0,K}), ||\widetilde{H}_m|| = 1$, and

$$\delta(\widetilde{H}_m) = A\widetilde{H}_m - p^m \widetilde{H}_m A_m(z^{p^m}).$$
(4.4)

Now, by applying Proposition 4.4 to the differential operator \mathcal{L}_m , it follows that there exists a MOM differential operator $\mathcal{L}_{m+1} \in E_{0,K}[\delta]$ of oder n such that: $\Lambda_p^{m+1}(f) = \Lambda_p(\Lambda_p^m(f))$ is a solution of \mathcal{L}_{m+1} , the companion matrix of \mathcal{L}_{m+1} , noted by A_{m+1} , belongs to \mathcal{M}_K , $||A_{m+1}|| \leq 1$, and a fundamental matrix of solutions of $\delta X = A_{m+1}X$ is given by

$$diag(1, 1/p, \dots, 1/p^{n-2}, 1/p^{n-1}) \cdot \Lambda_p(Y_{A_m}) \cdot diag(1, p, \dots, p^{n-2}, p^{n-1}) X^{A(0)}.$$

Since $Y_{A_m} = diag(1, 1/p^m, \dots, 1/p^{m(n-2)}, 1/p^{m(n-1)}) \cdot \Lambda_p(Y_A) \cdot diag(1, p^m, \dots, p^{m(n-2)}, p^{m(n-1)})$, we get that a fundamental of solutions of $\delta X = A_{m+1}X$ is the matrix

$$diag(1, 1/p^{m+1}, \dots, 1/p^{(m+1)(n-1)}) \cdot \Lambda_p^{m+1}(Y_A) \cdot diag(1, p^{m+1}, \dots, p^{(m+1)(n-1)}) X^{A(0)}.$$

By again invoking Proposition 4.4, we know that the matrix

$$\widetilde{G}_1 = Y_{A_m}(\Lambda_p(Y_{A_m})(z^p))^{-1} \cdot diag(1, p, \dots, p^{n-2}, p^{n-1})$$

belongs to $GL_n(E_K) \cap M_n(E_{0,K}), ||\widetilde{G}_1|| \leq 1$, and $\delta(\widetilde{G}_1) = A_m \widetilde{G}_1 - p \widetilde{G}_1 A_{m+1}(z^p).$

Thus,

$$\delta(\widetilde{G}_{1}(z^{p^{m}})) = p^{m}A_{m}(z^{p^{m}})\widetilde{G}_{1}(z^{p^{m}}) - \widetilde{G}_{1}(z^{p^{m}})p^{m+1}A_{m+1}(z^{p^{m+1}}).$$
(4.5)
= $\widetilde{H}_{1} \widetilde{C}_{1}(z^{p^{m}})$ Then

We put $\widetilde{H}_{m+1} = \widetilde{H}_m \widetilde{G}_1(z^{p^m})$. Then,

$$\widetilde{H}_{m+1} = Y_A(\Lambda_p^{m+1}(Y_A)(z^{p^{m+1}}))^{-1} \cdot diag(1, p^{m+1}, \dots, p^{(m+1)(n-1)})$$

belongs to $GL_n(E_K) \cap M_n(E_{0,K})$.

Furthermore, from Equations (4.4) and (4.5), we obtain

$$\delta(\tilde{H}_{m+1}) = A\tilde{H}_{m+1} - p^{m+1}\tilde{H}_{m+1}A_{m+1}(z^{p^{m+1}}).$$

Finally, $||\widetilde{H}_{m+1}|| = 1$ given that $||\widetilde{G}_1|| = 1 = ||\widetilde{H}_m||$ and

$$\widetilde{H}_{m+1}(0) = diag(1, p^{m+1}, \dots, p^{(m+1)(n-1)})$$

Therefore, the conditions (a)-(d) also hold for m + 1.

Before proving (i) of Theorem 3.1, we have the following remark.

Remark 4.6. Let \mathcal{L} be in $E_{0,K}[\delta]$ of order n, let A be the companion matrix of \mathcal{L} , and let $H = (h_{i,j})_{1, \leq i,j \leq n} \in M_n(E_{0,K}[[z]])$ such that $\delta H = AH - HB$, where $B \in M_n(E_{0,K})$. If the vector $(y_1, \ldots, y_n)^t \in K[[z]]^n$ is a solution of the system $\delta \vec{y} = B\vec{y}$ then $h_{1,1}y_1 + h_{1,2}y_2 + \cdots + h_{1,n}y_n$ is a solution of \mathcal{L} . Indeed, the equality $\delta H = AH - HB$ implies that $H(y_1, \ldots, y_n)^t$ is a solution of the system $\delta \vec{y} = A\vec{y}$ and given that A is the companion matrix of \mathcal{L} then $h_{1,1}y_1 + h_{1,2}y_2 + \cdots + h_{1,n}y_n$ is a solution of \mathcal{L} .

4.2. Proof of (i) of Theorem 3.1

Let us write $f(z) = \sum_{j \ge 0} f_j z^j$. In order to prove that $f(z) \in 1 + z\mathcal{O}_K[[z]]$, it is sufficient to show that, for all integers m > 0, $f_0, f_1, \ldots, f_{p^m-1} \in \mathcal{O}_K$. Since $f(z) \in \mathcal{MF}(K)$, there is a MOM differential operator $\mathcal{L} \in E_p[\delta]$ such that \mathcal{L} has a strong Frobenius structure and the coefficients of \mathcal{L} have Gauss norm less than or equal to 1. Let us say that the order of \mathcal{L} is n. Let m > 0 be an integer and let A be the companion matrix of \mathcal{L} . Since \mathcal{L} has a strong Frobenius structure, it follows from Propositions 4.1.2, 4.6.4, and 4.7.2 of [8] that there exists a matrix U invertible with coefficients in the generic dis $D(t, 1^-)$ such that $\delta U = AU$. Now, the eigenvalues of A(0) are all equal to zero because, by hypotheses, \mathcal{L} is MOM at zero. Hence, A belongs to \mathcal{M}_K . Further, we also have $||A|| \leq 1$ given that the coefficients of \mathcal{L} have norm less than or equal to 1. So, by Corollary 4.5, there exists $A_m \in \mathcal{M}_K$ and $\widetilde{H}_m \in GL_n(E_K) \cap M_n(E_{0,K})$ such that $||A_m|| \leq 1, ||\widetilde{H}_m|| = 1$, and

$$\delta(\widetilde{H}_m) = A\widetilde{H}_m - p^m \widetilde{H}_m A_m(z^{p^m}).$$
(4.6)

Again by Corollary 4.5, we know that $\Lambda_p^m(f)$ is a solution of \mathcal{L}_m . As A_m is the companion matrix of \mathcal{L}_m then the vector $(\Lambda_p^m(f), \delta(\Lambda_p^m(f)), \ldots, \delta^{n-1}(\Lambda_p^m(f)))^t$ is a solution of the system $\delta \vec{y} = A_m \vec{y}$. So, the vector $(\Lambda_p^m(f)(z^{p^m}), \delta(\Lambda_p^m(f))(z^{p^m}), \ldots, \delta^{n-1}(\Lambda_p^m(f))(z^{p^m}))^t$ is a solution of the system $\delta \vec{y} = p^m A_m(z^{p^m})\vec{y}$.

If we put $\tilde{H}_m = (h_{i,j})_{1 \leq i,j \leq n}$ then, thanks to Equation (4.6) we can apply Remark 4.6 and we deduce that

$$h_{1,1}\Lambda_p^m(f)(z^{p^m}) + h_{1,2}(\delta(\Lambda_p^m(f)))(z^{p^m}) + \dots + h_{1,n}(\delta^{n-1}(\Lambda_p^m(f)))(z^{p^m})$$

is a solution of \mathcal{L} . Further, according to Remark 4.2, there exists $c \in \mathbb{C}_p$ such that

$$h_{1,1}\Lambda_p^m(f)(z^{p^m}) + h_{1,2}(\delta(\Lambda_p^m(f)))(z^{p^m}) + \dots + h_{1,n}(\delta^{n-1}(\Lambda_p^m(f)))(z^{p^m}) = cf(z).$$
(4.7)

As f(0) = 1 and, for all integers j > 0, $(\delta^j(\Lambda_p^m(f)))(z^{p^m})(0) = 0$ then, from the previous equality, we have $h_{1,1}(0) = c$. But, by (d) of Corollary 4.5, we conclude that $h_{1,1}(0) = 1$. So that, c = 1. Again by (d) of Corollary 4.5, we infer that, for all $j \in \{2, \ldots, n\}$, $h_{1,j}(0) = 0$ and it is clear that, for all integers j > 0, $(\delta^j(\Lambda_p^m(f)))(z^{p^m}) \in z^{p^m} K[[z]]$. Hence, from Equation (4.7), we obtain

$$h_{1,1}\Lambda_p^m(f)(z^{p^m}) \equiv f \mod z^{p^m+1}K[[z]].$$
 (4.8)

Let us write $h_{1,1} = \sum_{j \ge 0} h_j z^j$. Thus, by Equation (4.8), we conclude that, for every integer $k \in \{1, \ldots, p^m - 1\}, h_k = f_k$. But, from (d) of Corollary 4.5, we know that $||\tilde{H}_m|| = 1$ and thus, $|h_{1,1}|_{\mathcal{G}} \le 1$. So that, $|f_k| \le 1$ for every integer $k \in \{1, \ldots, p^m - 1\}$. Whence, for every integer $k \in \{1, \ldots, p^m - 1\}, f_k \in \mathcal{O}_K$.

5. Constructing analytic elements and congruences modulo p^m

The aim of this section is to prove (ii) and (iii) of Theorem 3.1. In this direction, the key result is Proposition 5.5. To prove this proposition, we first establish some results concerning Cartier operator and congruences modulo p^m .

First of all, recall that if K is a finite extension of \mathbb{Q}_p then the maximal ideal $\mathfrak{m}_{\mathcal{O}_K}$ of \mathcal{O}_K is a principal ideal. An element $\pi \in K$ will be called a uniformizer of \mathcal{O}_K if $\mathfrak{m}_{\mathcal{O}_K} = (\pi)$. We also recall that $K_0(z)$ is the ring of rational functions A(z)/B(z) with $A(z), B(z) \in K[z]$ such that, for all $x \in D_0$, $B(x) \neq 0$.

5.1. Cartier operator and congruences modulo p^m .

The main result of this section is Lemma 5.3, where we prove that $\frac{f(z)}{\Lambda_p(f)(z^p)}$ belongs to E_p . We have the following two auxiliary lemmas.

Lemma 5.1. — Let the assumptions be as in Lemma 4.3. Then, for all integers m > 0, there exists a matrix $H_m \in GL_n(E_{0,K})$ such that $||H_m|| = 1 = |H_m^{-1}||$ and

$$\Lambda_p^m(Y_A)(\Lambda^{m+1}(Y_A)(z^p))^{-1} = diag(1, p^m, \dots, p^{m(n-1)}) \cdot H_m \cdot diag(1, 1/p^m, \dots, 1/p^{m(n-1)}).$$

Moreover, $H_m(0) = I = H_m^{-1}(0)$ and the first entry of the matrices H_m and H_m^{-1} has Gauss norm 1.

Proof. — Let m > 0 be an integer. From Corollary 4.5, it follows that there exits a MOM differential operator \mathcal{L}_m in $E_{0,K}[\delta]$ of order n such that $\Lambda_p^m(f)$ is a solution of \mathcal{L}_m . Let A_m be the companion

matrix of \mathcal{L}_m . Again by Corollary 4.5, we know that A_m belongs to \mathcal{M}_K , $||A_m|| \leq 1$ and that a fundamental matrix of solutions of the system $\delta X = A_m X$ is given by $Y_{A_m} X^{A(0)}$, where

$$Y_{A_m} = diag(1, 1/p^m, \dots, 1/p^{m(n-1)}) \cdot \Lambda_p^m(Y_A) \cdot diag(1, p^m, \dots, p^{m(n-1)}).$$

Now, by applying Lemma 4.3 to A_m , we conclude that the matrix $H_m = Y_{A_m}(\Lambda_p(Y_{A_m})(z^p))^{-1}$ belongs to $GL_n(E_{0,K})$ and that $||H_m|| = 1 = ||H_m^{-1}||$. It is clear that $H_m(0) = I = H_m^{-1}(0)$ and that

$$H_m = diag(1, 1/p^m, \dots, 1/p^{m(n-1)}) \cdot \Lambda_p^m(Y_A)(\Lambda_p^{m+1}(z^p))^{-1} \cdot diag(1, p^m, \dots, p^{m(n-1)})$$

As $||H_m|| = 1 = |H_m^{-1}||$ and $H_m(0) = I = H_m(0)^{-1}$ then, the first entry of the matrices H_m and H_m^{-1} has norm 1.

The following the lemma is technical but, it gives congruences modulo p^m .

Lemma 5.2. — For every integer $l \ge 0$, let H_l be the matrix defined in Lemma 5.1 and for every integer $m \ge 2$, we define

$$\prod_{l=1}^{m-1} \left[H_l(z^{p^l}) \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & p & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & p^{n-1} \end{pmatrix} \right] = (s_{\mu,\nu})_{1 \le \mu, \nu \le n}.$$

Then:

(i) $|s_{1,1}|_{\mathcal{G}} = 1 = |s_{1,1}(0)|$ and $s_{1,1}$ belongs to $1 + z\mathcal{O}_K[[z^p]]$.

(ii) For all $\nu \in \{2, \ldots, n\}$ and all $\mu \in \{2, \ldots, n\}$,

$$s_{\mu,1}s_{1,\nu} - s_{1,1}s_{\mu,\nu} \in p^{m-1}\mathcal{O}_K[[z]].$$

Proof. — Notice that $s_{\mu,\nu}$ belongs to $E_{0,K} \cap \mathcal{O}_K[[z^p]]$ because, by Lemma 5.1, we deduce that, for all l > 0, $||H_l(z^{p^l})|| \leq 1$ and $H_l(z^{p^l}) \in GL_n(E_{0,K})$.

Let us write $H_l(z^{p^l}) = (r_{i,j}^{(l)})_{1 \leq i,j \leq n}$. Thus,

$$s_{\mu,\nu} = \sum_{j_1=1}^n \sum_{j_2=1}^n \cdots \sum_{j_{m-2}=1}^n p^{j_1+\dots+j_{m-2}+\nu-m+1} r^{(1)}_{\mu,j_{m-2}} r^{(2)}_{j_{m-2},j_{m-3}} r^{(3)}_{j_{m-3},j_{m-4}} \cdots r^{(m-1)}_{j_1,\nu}.$$
 (5.1)

(i) It is clear that

$$s_{1,1} = r_{1,1}^{(1)} r_{1,1}^{(2)} r_{1,1}^{(3)} \cdots r_{1,1}^{(m-1)} + \sum_{\substack{1 \le j_1, \dots, j_{m-2} \le n, \\ (j_1, \dots, j_{m-2}) \ne (1, \dots, 1)}} p^{j_1 + \dots + j_{m-2} + 1 - m + 1} r_{1, j_{m-2}}^{(1)} r_{j_{m-2}, j_{m-3}}^{(2)} \cdots r_{j_1, 1}^{(m-1)}.$$

Let $\mathbf{J} = (j_1, \ldots, j_{m-2})$ be in $\{1, \ldots, n\}^{m-2}$ such that $\mathbf{J} \neq (1, \ldots, 1)$. Therefore, there exists $\kappa \in \{1, \ldots, m-2\}$ such that $j_{\kappa} \ge 2$. So that,

$$j_1 + \dots + j_{m-2} + 1 - m + 1 = j_1 + \dots + j_{m-2} - m + 2 \ge m - 3 + 2 - m + 2 = 1 > 0.$$

From Lemma 5.1, we know that $||H_l|| = 1$ for all l > 0. So, for all $i, j \in \{1, \ldots, n\}$ and, for all integers $l > 0, |r_{i,j}^{(l)}| \leq 1$. Consequently,

$$\left| \sum_{\substack{1 \leq j_1, \dots, j_{m-2} \leq n, \\ (j_1, \dots, j_{m-2}) \neq (1, \dots, 1)}} p^{j_1 + \dots + j_{m-2} + 1 - m + 1} r_{1, j_{m-2}}^{(1)} r_{j_{m-2}, j_{m-3}}^{(2)} \cdots r_{j_1, 1}^{(m-1)} \right|_{\mathcal{G}} < 1$$

But, from Lemma 5.1, we know that, for all integers l > 0, $|r_{1,1}^{(l)}| = 1$. Therefore, $|s_{1,1}|_{\mathcal{G}} = 1$. Finally, we know that, for all integers l > 0, $H_l(0) = I$. Hence, if $(j_1, \ldots, j_{m-2}) \neq (1, \ldots, 1)$ then $r_{1,j_{m-2}}^{(1)}r_{j_{m-2},j_{m-3}}^{(2)}\cdots r_{j_{1,1}}^{(m-1)}(0) = 0$ and it is clear that $r_{1,1}^{(1)}r_{1,1}^{(2)}r_{1,1}^{(3)}\cdots r_{1,1}^{(m-1)}(0) = 1$. Thus, $|s_{1,1}(0)| = 1$. So, we have $|s_{1,1}|_{\mathcal{G}} = 1 = |s_{1,1}(0)|$. Consequently, we obtain $\frac{1}{s_{1,1}}$ belongs to $1 + \mathcal{O}_K[[z]] \cap E_K$. Even, we have $\frac{1}{s_{1,1}} \in 1 + \mathcal{O}_K[[z^p]]$ because, by construction, it is clear that $s_{1,1} \in \mathcal{O}_K[[z^p]]$ and we have already seen that $s_{1,1}(0) = 1$.

(ii) From Equation (5.1), we have

$$s_{\mu,1}s_{1,\nu} - s_{1,1}s_{\mu,\nu} = \sum_{\substack{1 \le j_1, \dots, j_{m-2} \le n, \\ 1 \le t_1, \dots, t_{m-2} \le n \\ 1 \le j_1, \dots, j_{m-2} \le n \\ -}} \sum_{\substack{p^{\gamma=1}}} p^{m-2} (j_{\gamma} + t_{\gamma}) + \nu - 2m + 3 \\ - \sum_{\substack{1 \le j_1, \dots, j_{m-2} \le n \\ 1 \le j_1, \dots, j_{m-2} \le n \\ -}} p^{m-2} (j_{\gamma} + t_{\gamma}) + \nu - 2m + 3 \\ p^{\gamma=1} r_{1,j_{m-2}}^{(1)} r_{j_{m-2},j_{m-3}}^{(2)} \cdots r_{j_{1,1}}^{(m-1)} r_{\mu,t_{m-2}}^{(2)} r_{t_{m-2},t_{m-3}}^{(2)} \cdots r_{t_{1,\nu}}^{(m-1)}$$

Let $\mathbf{J} = (j_1, \ldots, j_{m-2})$ and $\mathbf{T} = (t_1, \ldots, t_{m-2})$ be in $\{1, \ldots, n\}^{m-2}$ such that $j_{\kappa} = t_{\kappa}$ for some $\kappa \in \{1, \ldots, m-2\}$. Let us consider $\mathbf{J}' = (j'_1, \ldots, j'_{m-2})$ and $\mathbf{T}' = (t'_1, \ldots, t'_{m-2})$ defined as follows

$$(j'_1, \dots, j'_{\kappa-1}, j'_{\kappa}, j'_{\kappa+1}, j'_{\kappa+2}, \dots, j'_{m-2}) = (j_1, \dots, j_{\kappa-1}, j_{\kappa}, t_{\kappa+1}, t_{\kappa+2}, \dots, t_{m-2})$$

and

$$(t'_1, \dots, t'_{\kappa-1}, t'_{\kappa}, t'_{\kappa+1}, t'_{\kappa+2}, \dots, t'_{m-2}) = (t_1, \dots, t_{\kappa-1}, t_{\kappa}, j_{\kappa+1}, \dots, j_{m-2})$$

So that, we have

$$\begin{bmatrix} r_{\mu,j_{m-2}}^{(1)} r_{j_{m-2},j_{m-3}}^{(2)} \cdots r_{j_{1,1}}^{(m-1)} \end{bmatrix} \begin{bmatrix} r_{1,t_{m-2}}^{(1)} r_{t_{m-2},t_{m-3}}^{(2)} \cdots r_{t_{1,\nu}}^{(m-1)} \end{bmatrix} = \\ \begin{bmatrix} r_{\mu,j_{m-2}}^{(1)} \cdots r_{j_{\kappa+1,j_{\kappa}}}^{(m-\kappa-1)} r_{j_{\kappa},j_{\kappa-1}}^{(m-\kappa)} r_{j_{\kappa-1,j_{\kappa-2}}}^{(m-\kappa+1)} \cdots r_{j_{1,1}}^{(m-1)} \end{bmatrix} \begin{bmatrix} r_{1,t_{m-2}}^{(1)} \cdots r_{t_{\kappa+1,t_{\kappa}}}^{(m-\kappa-1)} r_{t_{\kappa,t-1}}^{(m-\kappa)} r_{t_{\kappa-1,t_{\kappa-2}}}^{(m-\kappa+1)} \cdots r_{t_{1,\nu}}^{(m-1)} \end{bmatrix} = \\ \begin{bmatrix} r_{1,t_{m-2}}^{(1)} \cdots r_{t_{\kappa+1,t_{\kappa}}}^{(m-\kappa-1)} r_{j_{\kappa},j_{\kappa-1}}^{(m-\kappa)} r_{j_{\kappa-1,j_{\kappa-2}}}^{(m-\kappa+1)} \cdots r_{j_{1,1}}^{(m-1)} \end{bmatrix} \begin{bmatrix} r_{\mu,j_{m-2}}^{(1)} \cdots r_{j_{\kappa+1,j_{\kappa}}}^{(m-\kappa-1)} r_{t_{\kappa-1,t_{\kappa-2}}}^{(m-\kappa+1)} \cdots r_{t_{1,\nu}}^{(m-1)} \end{bmatrix} = \\ \begin{bmatrix} r_{1,t_{m-2}}^{(1)} \cdots r_{t_{\kappa+1,t_{\kappa}}}^{(m-\kappa-1)} r_{j_{\kappa,j_{\kappa-1}}}^{(m-\kappa)} r_{j_{\kappa-1,t_{\kappa-2}}}^{(m-\kappa+1)} \cdots r_{j_{1,1}}^{(m-1)} \end{bmatrix} \begin{bmatrix} r_{\mu,j_{m-2}}^{(1)} \cdots r_{j_{\kappa+1,j_{\kappa}}}^{(m-\kappa-1)} r_{t_{\kappa-1,t_{\kappa-2}}}^{(m-\kappa+1)} \cdots r_{t_{1,\nu}}^{(m-1)} \end{bmatrix} = \\ \begin{bmatrix} r_{1,j_{m-2}}^{(1)} \cdots r_{j_{\kappa+1,j_{\kappa}}}^{(m-\kappa-1)} r_{j_{\kappa},j_{\kappa-1}}^{(m-\kappa)} r_{j_{\kappa-1,j_{\kappa-2}}}^{(m-\kappa+1)} \cdots r_{j_{1,1}}^{(m-1)} \end{bmatrix} \begin{bmatrix} r_{\mu,t_{m-2}}^{(1)} \cdots r_{t_{\kappa+1,t_{\kappa}}}^{(m-\kappa-1)} r_{t_{\kappa-1,t_{\kappa-2}}}^{(m-\kappa+1)} \cdots r_{t_{1,\nu}}^{(m-1)} \end{bmatrix} \end{bmatrix} = \\ \end{bmatrix}$$

Moreover, we also have $\sum_{\gamma=1}^{m-2} (j_{\gamma} + t_{\gamma}) = \sum_{\gamma=1}^{m-2} (j'_{\gamma} + t'_{\gamma}).$ Thus, we obtain $p^{\sum(j_{\gamma}+t_{\gamma})+\nu-2m+3} r^{(1)}_{\mu,j_{m-2}} r^{(2)}_{j_{m-2},j_{m-3}} \cdots r^{(m-1)}_{j_{1,1}} r^{(1)}_{1,t_{m-2}} r^{(2)}_{t_{m-2},t_{m-3}} \cdots r^{(m-1)}_{t_{1,\nu}} - p^{\sum(j'_{\gamma}+t'_{\gamma})+\nu-2m+3} r^{(1)}_{1,j'_{m-2}} r^{(2)}_{j'_{m-2},j'_{m-3}} \cdots r^{(m-1)}_{j'_{1,1}} r^{(1)}_{\mu,t'_{m-2}} r^{(2)}_{t'_{m-2},t'_{m-3}} \cdots r^{(m-1)}_{t'_{1,\nu}} - 0$

Consequently,

$$\begin{split} s_{\mu,1}s_{1,\nu} &- s_{1,1}s_{\mu,\nu} = \\ &\sum_{\substack{1 \leq j_1, \dots, j_{m-2} \leq n \\ j_i \neq t_i \forall i}} p^{m-2}_{\gamma=1}(j_{\gamma} + t_{\gamma}) + \nu - 2m + 3 \\ r_{\mu,j_{m-2}}^{(1)}r_{j_{m-2},j_{m-3}}^{(2)} \cdots r_{j_{1,1}}^{(m-1)}r_{1,t_{m-2}}^{(2)}r_{t_{m-2},t_{m-3}}^{(2)} \cdots r_{t_{1,\nu}}^{(m-1)} \\ - \\ &\sum_{\substack{m=2 \\ j_i \neq t_i \forall i}} p^{m-2}_{\gamma=1}(j_{\gamma} + t_{\gamma}) + \nu - 2m + 3 \\ p^{\gamma=1} & r_{1,j_{m-2}}^{(1)}r_{j_{m-2},j_{m-3}}^{(2)} \cdots r_{j_{1,1}}^{(m-1)}r_{\mu,t_{m-2}}^{(1)}r_{t_{m-2},t_{m-3}}^{(2)} \cdots r_{t_{1,\nu}}^{(m-1)} \\ & r_{1,j_{m-2}}^{(1)}r_{j_{m-2},j_{m-3}}^{(2)} \cdots r_{j_{1,1}}^{(m-1)}r_{\mu,t_{m-2}}^{(2)}r_{t_{m-2},t_{m-3}}^{(2)} \cdots r_{t_{1,\nu}}^{(m-1)} \end{split}$$

Let $\mathbf{J} = (j_1, \ldots, j_{m-2})$ and $\mathbf{T} = (t_1, \ldots, t_{m-2})$ be in $\{1, \ldots, n\}^{m-2}$ such that, for all $\kappa \in \{1, \ldots, m-2\}$, $j_{\kappa} \neq t_{\kappa}$. Hence, for all $\kappa \in \{1, \ldots, m-2\}$, $j_{\kappa} + t_{\kappa} \ge 3$. Therefore,

$$\sum_{\gamma=1}^{m-2} j_{\gamma} + t_{\gamma} \ge 3m - 6.$$

As $\nu \ge 2$ then

$$\sum_{\gamma=1}^{n-2} (j_{\gamma} + t_{\gamma}) + \nu - 2m + 3 + \ge m - 1.$$

Therefore,

$$s_{\mu,1}s_{1,\nu} - s_{1,1}s_{\mu,\nu} \in p^{m-1}\mathcal{O}_K[[z]]$$

We are now ready to prove that $\frac{f(z)}{\Lambda_p(f)(z^p)}$ belongs to E_p .

1

Lemma 5.3. — Let the assumptions be as in Lemma 4.3, $f(z) \in 1 + zK[[z]]$ be a solution of \mathcal{L} , and π be a uniformizer of K. Then, for all integers m > 0, there exists $D_m(z) \in K_0(z)$ with norm 1 such that

$$D_m(z)(\Lambda_p(f))(z^p) \equiv f(z) \mod \pi^m \mathcal{O}_K[[z]].$$

Proof. — We split the proof into two cases.

Case m = 1. According to Proposition 4.4, there exists a MOM differential operator $\mathcal{L}_1 \in E_{0,K}[\delta]$ such that $\Lambda_p(f)$ is a solution of \mathcal{L}_1 . Let A_1 be the companion matrix of \mathcal{L}_1 and let $B_0 =$

 $Y_A(\Lambda(Y_A)(z^p))^{-1}$, where Y_A is the uniform part of $\delta X = AX$. Again by Proposition 4.4, the matrix $\widetilde{H}_1 = B_0 diag(1, p, \dots, p^{n-1})$ belongs to $GL_n(E_K) \cap M_n(E_{0,K})$ and

$$\delta \widetilde{H}_1 = A \widetilde{H}_1 - p \widetilde{H}_1 A_1(z^p).$$
(5.2)

Since $\Lambda_p(f)$ is solution of \mathcal{L}_1 , the vector $((\Lambda_p(f))(z^p), (\delta\Lambda_p(f))(z^p), \dots, (\delta^{n-1}\Lambda_p(f))(z^p))^t$ is a solution of $\delta \vec{y} = pA_1(z^p)\vec{y}$. Thus, from Equation (5.2), a solution of $\delta \vec{y} = A(z)\vec{y}$ is given by the vector $\widetilde{H}_1((\Lambda_p(f))(z^p), (\delta\Lambda_p(f))(z^p), \dots, (\delta^{n-1}\Lambda_p(f))(z^p))^t$. Let us write $B_0 = (b_{i,j})_{1 \leq i, \leq n}$. As A is the companion matrix of \mathcal{L} then

$$b_{1,1}(\Lambda_p(f))(z^p) + pb_{1,2}(\delta\Lambda_p(f))(z^p) + \dots + p^{n-1}b_{1,n}(\delta^{n-1}\Lambda_p(f))(z^p)$$

is a solution of \mathcal{L} . Following Remark 4.2, there exists $c \in K$ such that

$$b_{1,1}(\Lambda_p(f))(z^p) + pb_{1,2}(\delta\Lambda_p(f))(z^p) + \dots + p^{n-1}b_{1,n}(\delta^{n-1}\Lambda_p(f))(z^p) = cf(z).$$
(5.3)

But c = 1 because $B_0(0) = I$, f(0) = 1 and, for all integers j > 0, $(\delta^j \Lambda_p(f))(0) = 0$. It is clear that $p \in \pi \mathcal{O}_K$ and, by Lemma 4.3, we know that $||B_0|| = 1$ and that $B_0 \in GL_n(E_{0,K})$. Therefore, we can reduce Equation (5.3) modulo π and, we obtain

$$b_{1,1}(\Lambda_p(f))(z^p) \equiv f(z) \mod \pi \mathcal{O}_K[[z]].$$

We have $|b_{1,1}| = 1$ because $||B_0|| = 1$ and $b_{1,1}(0) = 1$. Thus, since $b_{1,1} \in E_{0,K}$, there exists $D_1 \in K_0(z)$ with norm equal to 1 such that $b_{1,1} \equiv D_1 \mod \pi \mathcal{O}_K[[z]]$. Consequently,

$$D_1(\Lambda_p(f))(z^p) \equiv f(z) \mod \pi \mathcal{O}_K[[z]].$$

Case $m \ge 2$. By Corollary 4.5, there exists a MOM differential operator $\mathcal{L}_m \in E_{0,K}[\delta]$ of order n such that $\Lambda_p^m(f)$ is a solution of \mathcal{L}_m . Let A_m be the companion matrix of \mathcal{L}_m . Then, by invoking Corollary 4.5 again, A_m belongs to \mathcal{M}_K and $||A_m|| \le 1$. Let us consider

$$\widetilde{H}_m = Y_A(\Lambda_p^m(Y_A)(z^{p^m}))^{-1} \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & p^m & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & p^{m(n-2)} & 0 \\ 0 & 0 & \dots & 0 & p^{m(n-1)} \end{pmatrix}$$

By (d) of Corollary 4.5, \widetilde{H}_m belongs to $GL_n(E_K) \cap M_n(E_{0,K})$, $||\widetilde{H}_m|| = 1$ and

$$\delta(\widetilde{H}_m) = A\widetilde{H}_m - p^m \widetilde{H}_m A_m(z^{p^m}).$$
(5.4)

Let us write $\widetilde{H}_m = (\widetilde{h}_{i,j}^{(m)})_{1 \leq i,j \leq n}$. Hence $|\widetilde{h}_{1,1}^{(m)}|_{\mathcal{G}} \leq 1$. But, $\widetilde{h}_{1,1}^{(m)}(0) = 1$ given that $\widetilde{H}_m(0) = diag(1, p^m, \dots, p^{m(n-1)})$. Thus, $|\widetilde{h}_{1,1}^{(m)}|_{\mathcal{G}} = 1$.

As $\Lambda_p^m(f)$ is a solution of \mathcal{L}_m and A_m is the companion matrix of \mathcal{L}_m then the vector given by $(\Lambda_p^m(f), \delta(\Lambda_p^m(f)), \dots, \delta^{n-1}(\Lambda_p^m(f)))^t$ is a solution of $\delta \vec{y} = A_m \vec{y}$. Thus, it follows that the vector $(\Lambda_p^m(f)(z^{p^m}), \delta(\Lambda_p^m(f))(z^{p^m}), \dots, \delta^{n-1}(\Lambda_p^m(f))(z^{p^m}))^t$ is a solution of $\delta \vec{y} = p^m A_m(z^{p^m})\vec{y}$. Therefore, thanks to Equation (5.4) we can apply Remark 4.6 and we deduce that

$$\tilde{h}_{1,1}^{(m)}\Lambda_p^m(f)(z^{p^m}) + \tilde{h}_{1,2}^{(m)}\delta(\Lambda_p^m(f))(z^{p^m}) + \dots + \tilde{h}_{1,n}^{(m)}\delta^{n-1}(\Lambda_p^m(f))(z^{p^m})$$

is a solution of \mathcal{L} .

By hypotheses A belongs to \mathcal{M}_K . In particular, the eigenvalues of A(0) are all equal to zero and therefore, according to Remark 4.2, there exists $c \in K$ such that

$$\widetilde{h}_{1,1}^{(m)}\Lambda_p^m(f)(z^{p^m}) + \widetilde{h}_{1,2}^{(m)}\delta(\Lambda_p^m(f))(z^{p^m}) + \dots + \widetilde{h}_{1,n}^{(m)}\delta^{n-1}(\Lambda_p^m(f))(z^{p^m}) = cf(z).$$

As, f(0) = 1 and, for all integers j > 0, $\delta^j(\Lambda_p^m(f))(0) = 0$, then $\tilde{h}_{1,1}^{(m)}(0) = c$. But, we know that $\tilde{h}_{1,1}^{(m)}(0) = 1$. So that, c = 1. Consequently,

$$\widetilde{h}_{1,1}^{(m)}\Lambda_p^m(f)(z^{p^m}) + \widetilde{h}_{1,2}^{(m)}\delta(\Lambda_p^m(f))(z^{p^m}) + \dots + \widetilde{h}_{1,n}^{(m)}\delta^{n-1}(\Lambda_p^m(f))(z^{p^m}) = f(z).$$
(5.5)

The rest of the proof is devoted to showing that, for all $\nu \in \{2, \ldots, n\}$, there exists $g_{\nu}(z) \in E_{0,K} \cap \mathcal{O}_K[[z]]$ such that

$$\widetilde{h}_{1,\nu}^{(m)} \equiv \widetilde{h}_{1,1}^{(m)} g_{\nu}(z^p) \mod \pi^m \mathcal{O}_K[[z]].$$
(5.6)

For each integer $l \ge 0$, we set $B_l = \Lambda_p^l(Y_A)(\Lambda_p^{l+1}(Y_A)(z^p))^{-1}$. Thus, we have

$$Y_A(\Lambda_p^m(Y_A)(z^{p^m}))^{-1} = \prod_{l=0}^{m-1} B_l(z^{p^l}).$$

Furthermore, we infer from Lemma 5.1 that, for every integer l > 0,

$$B_{l}(z^{p^{l}}) = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & p^{l} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & p^{l(n-1)} \end{pmatrix} H_{l}(z^{p^{l}}) \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1/p^{l} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1/p^{l(n-1)} \end{pmatrix}$$

where $H_l \in GL_n(E_{0,K})$, $H_l(0) = I = H_l(0)^{-1}$, and $||H_l|| = 1 = ||H_l^{-1}||$. Therefore,

$$\widetilde{H}_{m} = B_{0} \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & p & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & p^{n-1} \end{pmatrix} \prod_{l=1}^{m-1} \begin{bmatrix} H_{l}(z^{p^{l}}) \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & p & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & p^{n-1} \end{pmatrix} \end{bmatrix}.$$
(5.7)

Let us write $H_l(z^{p^l}) = (r_{i,j}^{(l)})_{1 \leq i,j \leq n}$. We also write

$$\prod_{l=1}^{m-1} \left[H_l(z^{p^l}) \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & p & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & p^{n-1} \end{pmatrix} \right] = (s_{\mu,\nu})_{1 \le \mu, \nu \le n}.$$

We recall that $B_0 = (b_{i,j})_{1 \leq i,j \leq n}$. Then, from Equation (5.7), we get that, for all $\nu \in \{1, \ldots, n\}$, we have

$$\widetilde{h}_{1,\nu}^{(m)} = b_{1,1}s_{1,\nu} + pb_{1,2}s_{2,\nu} + \dots + p^{n-1}b_{1,n}s_{n,\nu}.$$

We are going to see that, for all $\nu \in \{2, \ldots, n\}$,

$$\widetilde{h}_{1,1}^{(m)} \frac{s_{1,\nu}}{s_{1,1}} \equiv \widetilde{h}_{1,\nu}^{(m)} \mod \pi^m \mathcal{O}_K[[z]].$$
(5.8)

It is clear that

$$\widetilde{h}_{1,1}^{(m)} \frac{s_{1,\nu}}{s_{1,1}} - \widetilde{h}_{1,\nu}^{(m)} = \sum_{\mu=2}^{n} p^{\mu-1} b_{1,\mu} \left(s_{\mu,1} \frac{s_{1,\nu}}{s_{1,1}} - s_{\mu,\nu} \right).$$

Since, by definition, $B_0 = Y_A(\Lambda_p(Y_A)(z^p))^{-1}$, it follows from Lemma 4.3 that $B_0 \in GL_n(E_{0,K})$ and that $||B_0|| = 1$. In particular, for all $\mu \in \{2, \ldots, n\}$, $b_{1,\mu} \in \mathcal{O}_K[[z]]$. Further, $s_{\mu,\nu} \in E_{0,K} \cap \mathcal{O}_K[[z]]$ for all $1 \leq \mu, \nu \leq n$ because, according to Lemma 5.1, $H_l \in GL_n(E_{0,K})$ and $||H_l|| = 1$ for all integers l > 0. Finally, from (i) of Lemma 5.2, we deduce that $1/s_{1,1} \in 1 + \mathcal{O}_K[[z]]$. Therefore, for all $\mu \in \{2, \ldots, n\}$,

$$p^{\mu-1}b_{1,\mu}\left(s_{\mu,1}\frac{s_{1,\nu}}{s_{1,1}}-s_{\mu,\nu}\right) \in \mathcal{O}_K[[z]].$$

Hence, to prove the equality described by Equation (5.8) it is sufficient to show that, for all $\mu \in \{2, \ldots, n\}$, $p^{\mu-1}b_{1,\mu}\left(s_{\mu,1}\frac{s_{1,\nu}}{s_{1,1}} - s_{\mu,\nu}\right) \in \pi^m \mathcal{O}_K[[z]]$. As $s_{1,1}$ belongs to $1 + z\mathcal{O}_K[[z]]$ then, it follows that $p^{\mu-1}b_{1,\mu}\left(s_{\mu,1}\frac{s_{1,\nu}}{s_{1,1}} - s_{\mu,\nu}\right) \in \pi^m \mathcal{O}_K[[z]]$ if and only if $p^{\mu-1}b_{1,\mu}\left(s_{\mu,1}s_{1,\nu} - s_{1,1}s_{\mu,\nu}\right) \in \pi^m \mathcal{O}_K[[z]]$. But, according to (ii) of Lemma 5.2, we know that, for all $\mu \in \{2, \ldots, n\}$ and for all $\nu \in \{2, \ldots, n\}$,

$$(s_{\mu,1}s_{1,\nu} - s_{1,1}s_{\mu,\nu}) \in p^{m-1}\mathcal{O}_K[[z]].$$

Notice that $m - 1 + \mu - 1 \ge m$ because $\mu \ge 2$. Hence,

$$p^{\mu-1}b_{1,\mu}(s_{\mu,1}s_{1,\nu}-s_{1,1}s_{\mu,\nu}) \in p^m \mathcal{O}_K[[z]].$$

Since $p \in \pi \mathcal{O}_K$,

$$p^{\mu-1}b_{1,\mu}\left(s_{\mu,1}s_{1,\nu}-s_{1,1}s_{\mu,\nu}\right)\in\pi^{m}\mathcal{O}_{K}[[z]].$$

Therefore, for all $\nu \in \{2, \ldots, n\}$,

$$\widetilde{h}_{1,1}^{(m)} \frac{s_{1,\nu}}{s_{1,1}} \equiv \widetilde{h}_{1,\nu}^{(m)} \mod \pi^m \mathcal{O}_K[[z]].$$

Then, it follows from Equation (5.5) that

$$\widetilde{h}_{1,1}^{(m)}\left(\Lambda_p^m(f)(z^{p^m}) + \frac{s_{1,2}}{s_{1,1}}\delta(\Lambda_p^m(f))(z^{p^m}) + \dots + \frac{s_{1,n}}{s_{1,1}}\delta^{n-1}(\Lambda_p^m(f))(z^{p^m})\right) \equiv f \mod \pi^m \mathcal{O}_K[[z]].$$

But, $\Lambda_p^m(f)(z^{p^m}) + \frac{s_{1,2}}{s_{1,1}}\delta(\Lambda_p^m(f))(z^{p^m}) + \dots + \frac{s_{1,n}}{s_{1,1}}\delta^{n-1}(\Lambda_p^m(f))(z^{p^m}) \in \mathcal{O}_K[[z^p]]$ because, by construction, for all $\mu, \nu \in \{1, \dots, n\}$, $s_{\mu,\nu} \in \mathcal{O}_K[[z^p]]$ and we have already seen that, $1/s_{1,1} \in 1 + z\mathcal{O}_K[[z^p]]$. In addition, from (i) of Theorem 3.1, we know that $f(z) \in 1 + z\mathcal{O}_K[[z]]$ and thus, $\delta^j(\Lambda_p^m(f)) \in \mathcal{O}_K[[z]]$ for all integers $j \ge 0$. So, let d(z) be in $\mathcal{O}_K[[z]]$ such that

$$d(z^{p}) = \Lambda_{p}^{m}(f)(z^{p^{m}}) + \frac{s_{1,2}}{s_{1,1}}\delta(\Lambda_{p}^{m}(f))(z^{p^{m}}) + \dots + \frac{s_{1,n}}{s_{1,1}}\delta^{n-1}(\Lambda_{p}^{m}(f))(z^{p^{m}}).$$

Therefore,

$$\widetilde{h}_{1,1}^{(m)}d(z^p) \equiv f \mod \pi^m \mathcal{O}_K[[z]].$$

By applying Λ_p to the previous equality, we get

$$\Lambda_p\left(\widetilde{h}_{1,1}^{(m)}\right)d(z) \equiv \Lambda_p(f) \bmod \pi^m \mathcal{O}_K[[z]].$$

As $\widetilde{h}_{1,1}^{(m)} \in 1 + z\mathcal{O}_K[[z]]$ then $\Lambda_p\left(\widetilde{h}_{1,1}^{(m)}\right) \in 1 + z\mathcal{O}_K[[z]]$ and thus $\Lambda_p\left(\widetilde{h}_{1,1}^{(m)}\right) \mod \pi^m \mathcal{O}_K$ is a unit element of the ring $(\mathcal{O}_K/\pi^m)[[z]]$. So

$$d(z^p) \equiv \frac{(\Lambda_p(f))(z^p)}{(\Lambda_p(\widetilde{h}_{1,1}^{(m)}))(z^p)} \mod \pi^m \mathcal{O}_K[[z]].$$

Consequently,

$$\frac{\widetilde{h}_{1,1}^{(m)}}{\Lambda_p((\widetilde{h}_{1,1}^{(m)}))(z^p)}(\Lambda_p(f))(z^p) \equiv f \mod \pi^m \mathcal{O}_K[[z]]$$

We know that $\tilde{h}_{1,1}^{(m)} \in E_{0,K}$ and, from Proposition 5.1 of [10], we deduce that $\Lambda_p(\tilde{h}_{1,1}^{(m)}) \in E_{0,K}$. Since $|\tilde{h}_{1,1}^{(m)}|_{\mathcal{G}} = 1 = |\tilde{h}_{1,1}^{(m)}(0)|$, we have $|\Lambda_p(\tilde{h}_{1,1}^{(m)})|_{\mathcal{G}} = 1 = |\Lambda_p(\tilde{h}_{1,1}^{(m)})(0)|$. Thus, $\tilde{h}_{1,1}^{(m)}/\Lambda_p((\tilde{h}_{1,1}^{(m)}))(z^p)$ is an element of $E_{0,K}$ with norm 1. For this reason, there exists $D_m \in K_0(z)$ with norm 1 such that $\tilde{h}_{1,1}^{(m)}/\Lambda_p((\tilde{h}_{1,1}^{(m)}))(z^p) \equiv D_m \mod \pi^m \mathcal{O}_K[[z]]$. So that,

$$D_m(\Lambda_p(f))(z^p) \equiv f \mod \pi^m \mathcal{O}_K[[z]].$$

As a consequence of Lemma 5.3 we have the following result

Corollary 5.4. — Let the assumptions be as in Lemma 4.3, $f(z) \in 1 + zK[[z]]$ be a solution of \mathcal{L} , and π be a uniformizer of K. Then, for all integers $k \ge 1$, there is $P_k(z) \in K_0(z)$ with norm 1 such that

$$P_k(z)(\Lambda_p^k(f))(z^{p^k}) \equiv f(z) \bmod \pi^k \mathcal{O}_K[[z]].$$

Proof. — Let $m \ge 1$ be an integer. It follows from Corollary 4.5 that $\Lambda_p^m(f)$ is a solution of a MOM differential operator $\mathcal{L}_m \in E_{0,K}[\delta]$ such that its companion matrix A_m belongs to \mathcal{M}_K and $||A_m|| \le 1$. Then, by Lemma 5.3, we conclude that $\frac{\Lambda_p^m(f(z))}{(\Lambda_p^{m+1}(f))(z^p)}$ belongs to $E_{0,K}$. In particular, we deduce that, for all integers $m \ge 1$, $\frac{(\Lambda_p^m(f))(z^{p^m})}{\Lambda_p^{m+1}(f)(z^{p^{m+1}})}$ belongs to $E_{0,K}$. Now, it is clear that for any integer $k \ge 1$,

$$\frac{f}{(\Lambda_p^k(f))(z^{p^k})} = \frac{f(z)}{(\Lambda_p(f))(z^p)} \cdot \frac{(\Lambda^p(f))(z^p)}{(\Lambda_p^2(f))(z^{p^2})} \cdot \frac{(\Lambda_p^2(f))(z^{p^2})}{(\Lambda_p^3(f))(z^{p^3})} \cdots \frac{(\Lambda_p^{k-2}(f))(z^{p^{k-2}})}{(\Lambda_p^{k-1}(f))(z^{p^{k-1}})} \cdot \frac{(\Lambda_p^{k-1}(f))(z^{p^{k-1}})}{(\Lambda_p^k(f))(z^{p^k})}.$$

Therefore, $\frac{f}{(\Lambda_p^k(f))(z^{p^k})}$ belongs to $E_{0,K}$. Whence, there is $P_k(z) \in K_0(z)$ with norm 1 such that

$$P_k(z)(\Lambda_p^k(f))(z^{p^k}) \equiv f(z) \bmod \pi^k \mathcal{O}_K[[z]].$$

5.2. Key Proposition

The following proposition is fundamental in the proof of (ii) of Theorem 3.1.

Proposition 5.5. — Let K be a finite extension of \mathbb{Q}_p and let π be a uniformizer of \mathcal{O}_K . Let \mathcal{L} be a MOM differential operator in $E_{0,K}[\delta]$, let f(z) be a solution of \mathcal{L} in 1 + zK[[z]], and let A be the companion matrix of \mathcal{L} . Suppose that $||A|| \leq 1$ and that K is a Frobenius field. If \mathcal{L} has a strong Frobenius structure of period h then, for all integers $k \geq 0$, there is $Q_k(z) \in K_0(z)$ with norm 1 such that

$$Q_k(z) \equiv \frac{\Lambda_p^{kh}(f(z))}{f(z)} \mod \pi^{kh} \mathcal{O}_K[[z]].$$

Before proving Proposition 5.5, we need the following remark which will be also useful in Section 6.

Remark 5.6. Let K be an extension of \mathbb{Q}_p and let P(z) be $K_0(z)$. If $|P(z)|_{\mathcal{G}} \leq 1$ then $P(z) \in \mathcal{O}_K[[z]]$. Since 0 is not a pole of P(z), we have $P(z) = \sum_{n \geq 0} a_n z^n$ with $a_n \in K$ for all $n \geq 0$. But, $|a_n| \leq 1$ for all $n \geq 0$ because $|P(z)|_{\mathcal{G}} \leq 1$ and thus, $1 \geq \sup\{|a_n|\}_{n \geq 0}$. Proof of Proposition 5.5. —

Let A be the companion matrix of \mathcal{L} . We assume that the order of \mathcal{L} is equal to n. By hypotheses, we know that \mathcal{L} has a strong Frobenius structure of period h. Since K is a Frobenius field, the identity $i: K \to K$ map is a Frobenius endomorphism. Thus, there is a matrix $H \in GL_n(E_p)$ such that $\delta H = AH - p^h HA(z^{p^h})$. Since, by hypotheses \mathcal{L} is MOM at zero, it follows from Remark 4.2 that a fundamental matrix of solutions of $\delta X = AX$ is given by $Y_A X^{A(0)}$ with $Y_A \in GL_n(K[[z]])$ and $Y_A(0) = I$. So, a fundamental matrix of solutions of $\delta X = p^h A(z^{p^h})$ is given by $Y_A(z^{p^h})X^{p^hA(0)}$. Therefore, the equality $\delta H = AH - p^h HA(z^{p^h})$ implies that $HY_A(z^{p^h})X^{p^hA(0)}$ is a fundamental matrix of solution of $\delta X = AX$. So that, there exists $C \in GL_n(K)$ such that $HY_A(z^{p^h})X^{p^hA(0)} =$ $Y_A X^{A(0)}C$. Since $Y_A(0) = I$, $H \in M_n(E_p)$, and since Log(z) is transcendental over $\mathbb{C}_p[[z, 1/z]]$, we have $HY_A(z^{p^h}) = Y_AC$. Therefore, $H \in M_n(K[[z]])$. So, there exists $d \in K$ such that ||H|| = |d|. Notice that $d \neq 0$ given that $H \in GL_n(E_p)$. Let us set $T = \frac{1}{d}H$. Then, $T \in GL_n(E_p) \cap M_n(K[[z]])$, ||T|| = 1, and $\delta T = AT - p^h TA(z^{p^h})$.

Let $k \ge 1$ be an integer. We set $T_k = T(z)T(z^{p^h})\cdots T(z^{p^{(k-1)h}})$. Thus, $T_k \in GL_n(E_p) \cap M_n(K[[z]])$, $||T_k|| \le 1$, and

$$\delta T_k = AT_k - p^{kh} T_k A(z^{p^{kh}}).$$
(5.9)

So, \mathcal{L} has a strong Frobenius structure of period kh. Since f(z) is a solution of \mathcal{L} and A is the companion matrix of \mathcal{L} , the vector $(f, \delta f, \ldots, \delta^{n-1}f)^t$ is a solution of the system $\delta \vec{y} = A\vec{y}$. Thus, the vector $(f(z^{p^{kh}}), (\delta f)(z^{p^{kh}}), \ldots, (\delta^{n-1}f)(z^{p^{kh}}))^t$ is a solution of the system $\delta \vec{y} = p^{kh}A(z^{p^{kh}})\vec{y}$. Let us write $T_k = (t_{i,j}^{(k)})_{1 \leq i,j \leq n}$. Then, thanks to Equation (5.9), we deduce from Remark 4.6 that

$$t_{1,1}^{(k)}f(z^{p^{kh}}) + t_{1,2}^{(k)}(\delta f)(z^{p^{kh}}) + \dots + t_{1,n}^{(k)}(\delta^{n-1}f)(z^{p^{kh}})$$

is a solution of \mathcal{L} .

This solution is different from zero because $(f(z^{p^h}), (\delta f)(z^{p^h}), \ldots, (\delta^{n-1}f)(z^{p^h}))^t$ is not the zero vector and T_k belongs to $GL_n(E_p)$. By hypotheses, we know that \mathcal{L} is MOM at zero. Hence, according to Remark 4.2, there is $c \in K$ such that

$$t_{1,1}^{(k)}f(z^{p^{kh}}) + t_{1,2}^{(k)}(\delta f)(z^{p^{kh}}) + \dots + t_{1,n}^{(k)}(\delta^{n-1}f)(z^{p^{kh}}) = cf.$$

By applying Λ_p^{kh} to the previous equality, we have

$$\Lambda_p^{kh}(t_{1,1}^{(k)})f + \Lambda_p^{kh}(t_{1,2}^{(k)})\delta f + \dots + \Lambda_p^{kh}(t_{1,n}^{(k)})\delta^{n-1}f = c\Lambda_p^{kh}(f).$$
(5.10)

We have $c \neq 0$ because we have already seen that $t_{1,1}^{(k)}f(z^{p^{kh}}) + t_{1,2}^{(k)}(\delta f)(z^{p^{kh}}) + \cdots + t_{1,n}^{(k)}(\delta^{n-1}f)(z^{p^{kh}})$ is a solution of \mathcal{L} different from zero. As, for all integers j > 0, $(\delta^j f)(0) = 0$ and f(0) = 1, then $t_{1,1}^{(k)}(0) = c$. But, we know that $||T_k|| \leq 1$. Hence, $|c| \leq 1$ and $c = \pi^r u$, where u is a unit of \mathcal{O}_K and $r \in \mathbb{N}_{\geq 0}$. Therefore, from Equation (5.10), we get

$$\frac{\Lambda_p^{kh}(t_{1,1}^{(k)})f + \Lambda_p^{kh}(t_{1,2}^{(k)})\delta f + \dots + \Lambda_p^{kh}(t_{1,n}^{(k)})\delta^{n-1}f}{\pi^r} = u\Lambda_p^{kh}(f).$$
(5.11)

According to Corollary 5.4, there exists $P_{r+kh}(z) \in K_0(z)$ with norm 1 such that

$$P_{r+kh}[(\Lambda_p^{r+kh}f)(z^{p^{r+kh}})] \equiv f \mod \pi^{r+kh}\mathcal{O}_K[[z]].$$
(5.12)

Thus, for all integers i > 0,

$$(\delta^i P_{r+kh})[(\Lambda_p^{r+kh}f)(z^{p^{r+kh}})] \equiv \delta^i f \mod \pi^{r+kh} \mathcal{O}_K[[z]].$$

ALGEBRAIC INDEPENDENCE

In particular, for all integers $i \ge 0$, there exists $d_i \in \mathcal{O}_K[[z]]$ such that

$$(\delta^i P_{r+kh})[(\Lambda_p^{r+kh}f)(z^{p^{r+kh}})] - \delta^i f = \pi^{r+kh} d_i$$

So, by using Equation (5.11), we obtain the following equalities

$$\begin{split} &\left[\frac{\sum_{i=0}^{n-1}\Lambda_p^{kh}(t_{1,i+1}^{(k)})\delta^i P_{r+kh}}{\pi^r}\right](\Lambda_p^{r+kh}f)(z^{p^{r+kh}}) - u\Lambda_p^{kh}(f) \\ &= \frac{\sum_{i=0}^{n-1}\Lambda_p^{kh}(t_{1,i+1}^{(k)})\left[\delta^i P_{r+kh}(\Lambda_p^{r+kh}f)(z^{p^{r+kh}}) - \delta^i f\right]}{\pi^r} \\ &= \frac{\sum_{i=0}^{n-1}\Lambda_p^{kh}(t_{1,i+1}^{(k)})(\pi^{r+kh}d_i)}{\pi^r} \\ &= \pi^{kh}\left(\sum_{i=0}^{n-1}\Lambda_p^{kh}(t_{1,i+1}^{(k)})d_i\right). \end{split}$$

 So

$$\frac{\sum_{i=0}^{n-1} \Lambda_p^{kh}(t_{1,i+1}^{(k)}) \delta^i P_{r+kh}}{\pi^r} = \frac{\pi^{kh} \left(\sum_{i=0}^{n-1} \Lambda_p^{kh}(t_{1,i+1}^{(k)}) d_i \right) + u \Lambda_p^{kh}(f)}{(\Lambda_p^{r+kh} f)(z^{p^{r+kh}})}$$

From (i) of Theorem 3.1, $f(z) \in 1 + z\mathcal{O}_K[[z]]$. Thus, $(\Lambda_p^{r+kh}f)(z^{p^{r+kh}})$ is a unit of $\mathcal{O}_K[[z]]$ and from the previous equality we then have

$$\frac{\sum_{i=0}^{n-1} \Lambda_p^{kh}(t_{1,i+1}^{(k)}) \delta^i P_{r+kh}}{\pi^r} \in \mathcal{O}_K[[z]]$$

So, by reducing the last equality modulo π^{kh} and using the fact that $(\Lambda_p^{r+kh}f)(z^{p^{r+kh}})$ is a unit of $\mathcal{O}_K[[z]]$, we get that

$$\frac{\sum_{i=0}^{n-1} \Lambda_p^{kh}(t_{1,i+1}^{(k)}) \delta^i P_{r+kh}}{\pi^r} (\Lambda_p^{r+kh} f)(z^{p^{r+kh}}) \equiv u \Lambda_p^{kh}(f) \mod \pi^{kh} \mathcal{O}_K[[z]].$$
(5.13)

It follows from Equation (5.12) that,

$$P_{r+kh}[(\Lambda_p^{r+kh}f)(z^{p^{r+kh}})] \equiv f \mod \pi^{kh} \mathcal{O}_K[[z]].$$

Since $f(0) = 1 = \Lambda_p^{r+kh}(f)(0)$, we obtain $P_{r+kh}(0) \equiv 1 \mod \pi^{kh}$. Further, we know that $|P_{r+kh}|_{\mathcal{G}} = 1$ and, by Remark 5.6, we obtain $P_{r+kh} \in \mathcal{O}_K[[z]]$ and thus, P_{r+kh} is a unit of $\mathcal{O}_K[[z]]$. Whence, $P_{r+kh} \mod \pi^{kh} \mathcal{O}_K[[z]]$ belongs to $1 + z(\mathcal{O}_K/\pi^{kh})[[z]]$. So that, $P_{r+kh} \mod \pi^{kh} \mathcal{O}_K[[z]]$ is a unit element of $(\mathcal{O}_K/\pi^{kh})[[z]]$. Consequently,

$$(\Lambda_p^{r+kh}f)(z^{p^{r+kh}}) = \frac{f}{P_{r+kh}} \mod \pi^{kh} \mathcal{O}_k[[z]].$$

By substituting the above equality into Equation (5.13), we get

$$\frac{\sum_{i=0}^{n-1} \Lambda_p^{kh}(t_{1,i+1}^{(k)}) \delta^i P_{r+kh}}{\pi^r} \frac{f}{P_{r+kh}} \equiv u \Lambda_p^{kh}(f) \bmod \pi^{kh} \mathcal{O}_K[[z]].$$

Since $f(z) \in 1 + z\mathcal{O}_K[[z]]$, $f \mod \pi^{rh}\mathcal{O}_K[[z]]$ is a unit element of $(\mathcal{O}_K/\pi^{kh})[[z]]$. Then, from the previous equality, we have

$$\frac{\sum_{i=0}^{n-1} \Lambda_p^{kh}(t_{1,i+1}^{(k)}) \delta^i P_{r+kh}}{\pi^r} \frac{u^{-1}}{P_{r+kh}} \equiv \frac{\Lambda_p^{kh}(f)}{f} \mod \pi^{kh} \mathcal{O}_K[[z]].$$

Now, we are going to see that there exists $Q_k \in K_0(z)$ with norm 1 such that

$$\frac{\sum_{i=0}^{n-1} \Lambda_p^{kh}(t_{1,i+1}^{(k)}) \delta^i P_{r+kh}}{\pi^r} \frac{u^{-1}}{P_{r+kh}} \equiv Q_k \mod \pi^{kh} \mathcal{O}_K[[z]].$$

We know that P_{r+kh} has norm equal to 1 and that $P_{r+kh}(0) \equiv 1 \mod \pi^{kh}$. For this reason $P_{r+kh}(0)$ is a unit element of \mathcal{O}_K . But, P_{r+kh} belongs to $\mathcal{O}_K[[z]]$ because $P_{r+kh} \in K_0(z)$ and $|P_{r+kh}|_{\mathcal{G}} = 1$. Thus, P_{r+kh} is a unit element of $\mathcal{O}_K[[z]]$. So that, $1/P_{r+kh} \in K_0(z)$. Now, as $t_{1,1}^{(k)}, t_{1,2}^{(k)}, \dots, t_{1,n}^{(k)}$ belong to $E_p \cap K[[z]]$ then, by Proposition 5.1 of [10], we deduce that $\Lambda_p^{kh}(t_{1,1}^{(k)}), \Lambda_p^{kh}(t_{1,2}^{(k)}), \dots, \Lambda_p^{kh}(t_{1,n}^{(k)})$ belong to $E_p \cap K[[z]]$. Further, for all integers i > 0, $\delta^i P_{r+kh}$ belongs to $E_K \cap K[[z]]$ given that P_{r+kh} belongs to $K_0(z) \subset E_p \cap K[[z]]$. Therefore,

$$\frac{\sum_{i=0}^{n-1} \Lambda_p^{kh}(t_{1,i+1}^{(k)}) \delta^i P_{r+kh}}{\pi^r} \frac{u^{-1}}{P_{r+kh}}$$

belongs to $E_p \cap K[[z]]$. In particular, there exists $Q_k \in K(z) \cap K[[z]]$ such that

$$\frac{\sum_{i=0}^{n-1} \Lambda_p^{kh}(t_{1,i+1}^{(k)}) \delta^i P_{r+kh}}{\pi^r} \frac{u^{-1}}{P_{r+kh}} \equiv Q_k \mod \pi^{kh} \mathcal{O}_K[[z]].$$

Thus,

$$Q_k \equiv \frac{\Lambda_p^{kh}(f)}{f} \mod \pi^{kh} \mathcal{O}_K[[z]].$$
(5.14)

Since $|\Lambda_p^{kh}(f)/f|_{\mathcal{G}} = 1$, from Equation (5.14), we have $|Q_k|_{\mathcal{G}} = 1$ Thus, $Q_k \in \mathcal{O}_K[[z]]$. But, from Equation (5.14), we also have $Q_k(0) \equiv 1 \mod \pi^k$. Hence, for all xD_0 , $|Q_k(x)| = 1$ and consequently, $Q_k \in K_0(z)$.

That completes de proof.

5.3. Proof of (ii) and (iii) of Theorem 3.1

Let A be the companion matrix of \mathcal{L} . By hypotheses, we know that \mathcal{L} has a strong Frobenius structure and let h be the period.

(ii) We first prove that $f(z)/f(z^{p^h}) \in E_{0,K}$. Since \mathcal{L} has a strong Frobenius structure, it follows from Propositions 4.1.2, 4.6.4, and 4.7.2 of [8] that there exists a matrix U invertible with coefficients in the generic disc $D(t, 1^-)$ such that $\delta U = AU$. Now, the eigenvalues of A(0) are all equal to zero because, by hypotheses, \mathcal{L} is MOM at zero. It is clear that $A \in M_n(E_{0,K})$ because, by hypotheses, \mathcal{L} is a differential operator with coefficients in $E_{0,K}$. Hence, A belongs to \mathcal{M}_K . Further, by hypotheses, we know that the coefficients of \mathcal{L} have norm less than or equal to 1. So, $||A|| \leq 1$. This puts us in a position to apply Corollary 5.4. Thus, for any integer $k \ge 1$, there is $P_{kh}(z) \in K_0(z)$ with norm 1 such that

$$P_{kh}(z)(\Lambda_p^{kh}(f))(z^{p^{kh}}) \equiv f(z) \mod \pi^{kh} \mathcal{O}_K[[z]].$$

According to (i) of Theorem 3.1, $f(z) \in 1 + z\mathcal{O}_K[[z]]$. Hence, we can reduce f(z) modulo π^{kh} . So, $f(z) \mod \pi^{kh} \in 1 + z \left(\mathcal{O}_K/\pi^{kh}\right)[[z]]$. Hence, $f(z) \mod \pi^{kh}$ is a unit element of the ring $\left(\mathcal{O}_K/\pi^{kh}\right)[[z]]$. Thus,

$$P_{kh}(z)\frac{(\Lambda_p^{kh}(f))(z^{p^{kh}})}{f(z^{p^{kh}})}f(z^{p^{kh}}) \equiv f(z) \bmod \pi^{kh}\mathcal{O}_K[[z]].$$

But, according to Proposition 5.5, there is $Q_k(z) \in K_0(z)$ with norm 1 such that

$$Q_k(z) \equiv \frac{\Lambda_p^{kh}(f(z))}{f(z)} \mod \pi^{kh} \mathcal{O}_K[[z]].$$

Therefore,

$$P_{kh}(z)Q_k(z^{p^{kh}})f(z^{p^{kh}}) \equiv f(z) \mod \pi^{kh}\mathcal{O}_K[[z]].$$

We put $B_{kh}(z) = P_{kh}(z)Q_k(z^{p^{kh}})$. It is clear that $B_{kh}(z) \in K_0(z)$ and its norm is equal to 1.

Since k is an arbitrary element in $\mathbb{N}_{>0}$ we conclude that, for all integers $k \ge 1$,

$$f(z) \equiv B_{kh}(z)f(z^{p^{kh}}) \mod \pi^{kh}\mathcal{O}_K[[z]].$$
(5.15)

We now prove that, for all integers $k \ge 1$,

$$\frac{f}{f(z^{p^h})} \equiv \frac{B_{(k+1)h}(z)}{B_{kh}(z^{p^h})} \mod \pi^{kh} \mathcal{O}_K[[z]].$$

It follows from (5.15) that $f(z) \equiv B_{kh}(z)f(z^{p^{kh}}) \mod \pi^{kh}\mathcal{O}_K[[z]]$. Hence,

$$f(z^{p^{h}}) \equiv B_{kh}(z^{p^{h}})f(z^{p^{(k+1)h}}) \mod \pi^{kh} \mathcal{O}_{K}[[z]].$$
(5.16)

Again, it follows from (5.15) that $f(z) \equiv B_{(k+1)h}(z)f(z^{p^{(k+1)h}}) \mod \pi^{(k+1)h}\mathcal{O}_K[[z]]$. As $\pi^{(k+1)h}\mathcal{O}_K \subset \pi^{kh}\mathcal{O}_K$, we have

$$f(z) \equiv B_{(k+1)h}(z) f(z^{p^{(k+1)h}}) \mod \pi^{kh} \mathcal{O}_K[[z]].$$
(5.17)

Thus, from Equations (5.16) and (5.17), we obtain

$$\frac{f}{f(z^{p^h})} \equiv \frac{B_{(k+1)h}(z)}{B_{kh}(z^{p^h})} \mod \pi^{kh} \mathcal{O}_K[[z]].$$

Consequently, $f/f(z^{p^h})$ is an analytic element. In order to prove that $f/f(z^{p^h})$ belongs to $E_{0,K}$, we have to prove that, for all integers $k \ge 1$, the rational function $\frac{B_{(k+1)h}(z)}{B_{kh}(z^{p^h})}$ belongs to $K_0(z)$. Let k be a positive integer. It follows from Equation (5.15) that

$$f(z) - B_{kh}(z)f(z^{p^{kh}}) \in \pi^{kh}\mathcal{O}_K[[z]].$$

Thus, $1 - B_{kh}(0) \in \pi^{kh} \mathcal{O}_K$. Since the norm is non-Archimedean and $|\pi| < 1$, $|B_{kh}(0)| = 1$. Further, we know that $|B_{kh}(z)|_{\mathcal{G}} = 1$ and thus, for all $x \in D_0$, $|B_{kh}(x)| = |B_{kh}(0)| = 1$. In particular, for all $x \in D_0$, $B_{kh}(x) \neq 0$. Therefore, $1/B_{kh}(z) \in K_0(z)$. Finally, as $B_{(k+1)h}(z)$ belongs to $K_0(z)$ then $\frac{B_{(k+1)h}(z)}{B_{kh}(z)^{ph}}$ belongs to $K_0(z)$.

(iii) We now prove that $f'(z)/f(z) \in E_{0,K}$. We put $H(z) = f(z)/f(z^{p^h})$. According to (ii), H belongs to $E_{0,K}$. Let us show that H is a unit of $E_{0,K}$. For this purpose, we need to show that, for all $x \in D_0$, $H(x) \neq 0$. We know from (i) of Theorem 3.1 that $f(z) \in 1 + z\mathcal{O}_K[[z]]$. Since the norm is non-Archimedean, we get that, for all $x \in D_0$, |f(x)| = 1. Whence, for all $x \in D_0$, |H(x)| = 1 and therefore, $H(x) \neq 0$ for all $x \in D_0$. Now, let us show that H'(z) also belongs to $E_{0,K}$. Since $H(z) \in E_K$, $H'(z) \in E_K$. So, it is sufficient to show that, for all $x \in D_0$, H'(x) converges. It is not hard to see that

$$H'(z) = H(z) \left[\frac{f'(z)}{f(z)} - p^h z^{p^h - 1} \frac{f'(z^{p^h})}{f(z^{p^h})} \right].$$

As $f(z) \in 1 + z\mathcal{O}_K[[z]]$ then, for all $x \in D_0$, f(x) converges. So, from the previous equality, we conclude that, for all $x \in D_0$, H'(x) converges. For every $m \ge 0$, we consider $H_m = H(z)H(z^{p^h})\cdots H(z^{p^{m^h}})$. So, $H_m \in E_{0,K}$ and is a unit of $E_{0,K}$. Further, we also have

$$H'_{m}(z) = H_{m}(z) \left[\frac{f'(z)}{f(z)} - p^{(m+1)h} z^{p^{(m+1)h} - 1} \frac{f'(z^{p^{(m+1)h}})}{f(z^{p^{(m+1)h}})} \right].$$
(5.18)

Since $f(z) \in 1 + z\mathcal{O}_K[[z]]$, we get that, for every integer $m \ge 0$, $\frac{f'(z^{p^{(m+1)h}})}{f(z^{p^{(m+1)h}})} \in \mathcal{O}_K[[z]]$ and $H_m \in 1 + z\mathcal{O}_K[[z]]$. As $p \in (\pi)\mathcal{O}_K$ then, it is derived from Equation (5.18) that, for every integer $m \ge 0$,

$$\frac{H'_m(z)}{H_m(z)} \equiv \frac{f'(z)}{f(z)} \mod \pi^{(m+1)h} \mathcal{O}_K[[z]].$$

But, for every integer $m \ge 0$, $\frac{H'_m(z)}{H_m(z)} \in E_{0,K}$. The ring $E_{0,K}$ is complete because K is complete and thus, we have $f'(z)/f(z) \in E_{0,K}$. This completes the proof.

Remark 5.7. — It is clear from the previous proof that condition (ii) of Theorem 3.1 implies condition (iii) of Theorem 3.1.

6. A criterion for algebraic independence

The purpose of this section is to prove Theorem 6.12, which provides a necessary condition for determining when power series in $\mathcal{MF}(K)$ are algebraically independent over $E_{0,K}$. As an illustration of this criterion, we prove at the end of this section Theorems 2.6, 2.7, 2.9, and 2.10. We first recall the notions of pole and residue for an analytic element, as well as the notions of an exponent and regular singular point for a differential operator with coefficients in a ring of characteristic p > 0.

We recall that $\mathcal{O}_{\mathbb{C}_p}$ is the set of elements in \mathbb{C}_p with norm less than or equal to 1. The maximal ideal of $\mathcal{O}_{\mathbb{C}_p}$ is denoted by \mathfrak{m} . For any $\alpha \in \mathcal{O}_{\mathbb{C}_p}$, we put $D_{\alpha} = \{x \in \mathcal{O}_{\mathbb{C}_p} : |x - \alpha| < 1\}$. We also put $D_{\infty} = \{x \in \mathbb{C}_p : |x| \ge 1\}$. Note that $\mathfrak{m} = D_0$.

Remark 6.1. Let α, β be in $\mathcal{O}_{\mathbb{C}_p}$. As the norm is non-Archimedean then, $D_{\alpha} \cap D_{\beta} = \emptyset$ if and only if $|\alpha - \beta| = 1$ and $D_{\alpha} = D_{\beta}$ if and only if $|\alpha - \beta| < 1$.

6.0.1. Poles and residues

We recall the notions of *pole* and *residue* for an analytic element and prove some properties related to them. Let α be in $\mathcal{O}_{\mathbb{C}_p}$ and let $\mathbb{C}_p[z]_{\mathfrak{m}_\alpha}$ be the set of rational functions R(z) in $\mathbb{C}_p(z)$ such that every pole of R(z) belongs to D_α . Then, we let E^α denote the completion of $\mathbb{C}_p[z]_{\mathfrak{m}_\alpha}$ for the Gauss norm. For any set $A \subset \mathbb{C}_p$, E(A) is the completion of ring of the rational functions in $\mathbb{C}_p(z)$ that do not have poles in A. Note that $E(D_0) = E_{0,p}$.

Let f be in $E_{0,K}$ bounded⁽⁵⁾. According to Mittag-Lefflet's Theorem (see [19, Theorem 15.1]), there exists a unique sequence $\{\alpha_n\}_{n\geq 1}$ in $\mathcal{O}_{\mathbb{C}_p}$ such that, for every n, $|\alpha_n| = 1$, $|\alpha_n - \alpha_m| = 1$ for

⁽⁵⁾ We say that $f \in E_{0,K}$ is bounded if there exits a constant C > 0 such that for all $x \in D_0$, $|f(x)| \leq C$.

every $n \neq m$ and there exists a unique sequence $\{f_n\}_{n \geq 0}$ in $E_{0,K}$ such that f_0 belongs to $E(\mathcal{O}_{\mathbb{C}_p})$ and, for every n > 0, f_n belongs to E^{α_n} , and

$$f = \sum_{n=0}^{\infty} f_n. \tag{6.1}$$

Definition 6.2 (Pole). — Let f be in $E_{0,K}$ bounded and let α be in $\mathcal{O}_{\mathbb{C}_p}$. We say that D_{α} is a pole of f if there exists $n \ge 1$ such that $\alpha_n \in D_{\alpha}$ and $f_n \ne 0$. We denote the set of poles of f by Pol(f).

In order to define the residue for an analytic element in $E_{0,K}$, we need the following result.

Lemma 6.3. — Let α and β be in \mathbb{C}_p .

(i) If $|\alpha - \beta| < 1$ then

$$\frac{1}{z-\beta} = \frac{1}{z-\alpha} \sum_{l \ge 0} \left(\frac{\beta-\alpha}{z-\alpha}\right)^l.$$

(ii) If $|\alpha - \beta| \ge 1$ then $\frac{1}{z-\beta} = \sum_{s \ge j} a_s (z-\alpha)^s$ with $a_s \in \mathcal{O}_{\mathbb{C}_p}$ for all $s \ge 0$.

 $\begin{array}{lll} Proof. & -(\mathrm{i}) & \mathrm{Since} \ |\alpha - \beta| < 1, \ \mathrm{we \ have \ } \lim_{l \to \infty} |\alpha - \beta|^l = 0. \ \mathrm{If} \ |\alpha| \leqslant 1 \ \mathrm{then} \ \left| \frac{\beta - \alpha}{z - \alpha} \right|_{\mathcal{G}} = \\ |\beta - \alpha| < 1 \ \mathrm{and \ if} \ |\alpha| > 1 \ \mathrm{then} \ \left| \frac{\beta - \alpha}{z - \alpha} \right|_{\mathcal{G}} = \frac{|\beta - \alpha|}{|\alpha|} < 1 \ \mathrm{because} \ |\beta - \alpha| < 1 < |\alpha|. \ \mathrm{In \ both} \\ \mathrm{cases \ we \ obtain \ } \lim_{l \to \infty} \left| \left(\frac{\beta - \alpha}{z - \alpha} \right)^l \right|_{\mathcal{G}} = 0 \ \mathrm{and \ thus, \ the \ sum \ } \sum_{l \ge 0} \left(\frac{\beta - \alpha}{z - \alpha} \right)^l \ \mathrm{exists \ in \ } E_p. \ \mathrm{Let \ us \ put} \\ x_0 = \frac{1}{z - \alpha} \sum_{l \ge 0} \left(\frac{\beta - \alpha}{z - \alpha} \right)^l \ \mathrm{and} \ S_1 = \frac{1}{z - \alpha} + \frac{\beta - \alpha}{(z - \alpha)(z - \beta)}. \ \mathrm{So, \ } S_1 - x_0 = \left(\frac{\beta - \alpha}{z - \alpha} \right) \left(\frac{1}{z - \beta} - x_0 \right) \ \mathrm{and \ for \ this} \\ \mathrm{reason, \ } S_1 - \left(\frac{\beta - \alpha}{z - \beta} \right) \frac{1}{z - \beta} = x_0 \left(1 - \frac{\beta - \alpha}{z - \beta} \right). \ \mathrm{But, \ it \ is \ clear \ that \ } S_1 = \frac{1}{z - \beta}. \ \mathrm{Whence, \ } S_1 - \left(\frac{\beta - \alpha}{z - \beta} \right) \frac{1}{z - \beta} = \\ \frac{1}{z - \beta} \left(1 - \frac{\beta - \alpha}{z - \beta} \right). \ \mathrm{Therefore, \ } x_0 = \frac{1}{z - \beta}. \end{array}$

(ii) It is clear that $\frac{1}{z-\beta} = \sum_{s \ge 0} \frac{(-1)^s}{(\alpha-\beta)^{s+1}} (z-\alpha)^s$.

Remark 6.4. Let R(z) be in $\mathbb{C}_p(z)$ and let $\alpha_1, \ldots, \alpha_r$ be the poles of R(z). Suppose that $|\alpha_i| = 1$ for all $1 \leq i \leq r$. Since \mathbb{C}_p is algebraically closed, by partial fraction decomposition, we get that $R(z) = T(z) + F_1(z) + \cdots + F_r(z)$, where T(z) is a polynomial and $F_i(z) = \sum_{k=1}^{n_i} \frac{a_{i,k}}{(z-\alpha_i)^k}$, where $a_{i,k} \in \mathbb{C}_p$ and n_i is the order of R(z) at α_i . In particular, for all $i \in \{1, \ldots, r\}$, $res(R(z), \alpha_i) = a_{i,1}$.

(1) Let α be in $\{\alpha_1, \ldots, \alpha_r\}$. We are going to prove that

$$R(z) = \sum_{s \in \mathbb{Z}} a_s (z - \alpha)^s \text{ with } a_s \in \mathbb{C}_p \text{ for all } s \in \mathbb{Z} \text{ and } a_{-1} = \sum_{\alpha_i \in S} a_{i,1}.$$

where S is the set of poles of R(z) in D_{α} .

Without loss of generality we can assume that $S = \{\alpha_1, \ldots, \alpha_r\} \cap D_{\alpha} = \{\alpha_1, \ldots, \alpha_q\}$. It is clear that $R(z) = T(z) + \sum_{i=1}^{q} F_i(z) + \sum_{i=q+1}^{r} F_i(z)$. So, by (ii) of Lemma 6.3, we have $\sum_{i=q+1}^{r} F_i(z) \in \mathbb{C}_p[[z - \alpha]]$ and thus, $T(z) + \sum_{i=q+1}^{r} F_i(z) \in \mathbb{C}_p[[z - \alpha]]$. In addition, by (i) of Lemma 6.3, we deduce that, for all $1 \leq i \leq q$,

$$\frac{1}{z-\alpha_i} = \frac{1}{z-\alpha} \sum_{l \ge 0} \left(\frac{\alpha_i - \alpha}{z-\alpha}\right)^l.$$

So,

$$\sum_{i=1}^{q} F_i(z) = \sum_{i=1}^{q} \left(\sum_{k=1}^{n_i} a_{i,k} \left(\frac{1}{z - \alpha} \sum_{l \ge 0} \left(\frac{\alpha_i - \alpha}{z - \alpha} \right)^l \right)^k \right)$$

Thus, $R(z) = \sum_{s \in \mathbb{Z}} a_s (z - \alpha)^s$ with $a_s \in \mathbb{C}_p$ for all $s \in \mathbb{Z}$ and $a_{-1} = \sum_{i=1}^q a_{i,1}$. (2) Let α be in $\{\alpha_1, \ldots, \alpha_r\}$ and let $\tau \in D_\alpha$. We are going to see that

$$R(z) = \sum_{s \in \mathbb{Z}} b_s (z - \tau)^s \text{ with } b_s \in \mathbb{C}_p \text{ for all } s \in \mathbb{Z} \text{ and } b_{-1} = \sum_{\alpha_i \in S} a_{i,1},$$

where S is the set of poles of R(z) in D_{α} .

By (1), we know that $R(z) = \sum_{s \in \mathbb{Z}} a_s(z - \alpha)^s$ with $a_s \in \mathbb{C}_p$ for all $s \in \mathbb{Z}$ and $a_{-1} = \sum_{\alpha_i \in S} a_{i,1}$. So $R(z) = \sum_{s \leqslant -1} a_s(z - \alpha)^s + \sum_{s \geqslant 0} a_s(z - \alpha)^s$. Notice that $z - \alpha = z - \tau + \tau - \alpha$ and hence, for all integers $s \geqslant 0$, $(z - \alpha)^s = \sum_{k=0}^s {s \choose k} (\tau - \alpha)^{n-k} (z - \tau)^k$. Since $|\tau - \alpha| < 1$, for all integers $m \geqslant 1$, $t_m := \sum_{k=m}^\infty {k \choose m} (\tau - \alpha)^{k-m}$ exists in \mathbb{C}_p and therefore

$$\sum_{s \ge 0} a_s (z - \alpha)^s = \sum_{s \ge 0} a_s t_s (z - \tau)^s \in \mathbb{C}_p[[z - \tau]].$$

Since $|\tau - \alpha| < 1$, Lemma 6.3 gives that

$$\frac{1}{z-\alpha} = \frac{1}{z-\tau} \sum_{l \ge 0} \left(\frac{\alpha-\tau}{z-\tau}\right)^l.$$

Consequently,

$$\sum_{s \leqslant -1} a_s (z - \tau)^s = \sum_{s \geqslant 1} a_s \left(\frac{1}{z - \tau} \sum_{l \geqslant 0} \left(\frac{\alpha - \tau}{z - \tau} \right)^l \right)^s$$

Thus, $R(z) = \sum_{s \in \mathbb{Z}} b_s (z - \tau)^s$ with $b_s \in \mathbb{C}_p$ for all $s \in \mathbb{Z}$ and $b_{-1} = a_{-1}$ and, by (1) we know that $a_{-1} = \sum_{\alpha_i \in S} a_{i,1}$.

Definition 6.5 (Residue). — Let f be in $E_{0,K}$ bounded and let α be in $\mathcal{O}_{\mathbb{C}_p}$ such that D_{α} is a pole of f. Following the decomposition given by Equation (6.1), that means that there exits $n \ge 1$ such that $\alpha_n \in D_{\alpha}$ and $f_n \ne 0$. Since $f_n \in E^{\alpha_n}$, f_n is the limit of rational functions $R_j(z) \in \mathbb{C}_p(z)$ such that every pole of $R_j(z)$ belongs to D_{α_n} . Let $\tau \in D_{\alpha_n}$. Then, according to (2) of Remark 6.4, for every j, $R_j(z) = \sum_{s \in \mathbb{Z}} a_{s,j}(z - \tau)^s$ with $a_{s,j} \in \mathbb{C}_p$ for all $s \in \mathbb{Z}$. So, the residue of f at D_{α} is $res(f, D_{\alpha}) = \lim_{j \to \infty} a_{-1,j}$.

This definition does not depend on the choice of τ . In fact, let τ' be in D_{α_n} . Then, by (2) of Remark 6.4 again, we get that, for every j, $R_j(z) = \sum_{s \in \mathbb{N}} b_{s,j}(z - \tau')^s$ with $b_{-1,j} = a_{-1,j}$.

Remark 6.6. — Let R(z) be in $\mathbb{C}_p(z)$ and let $\alpha_1, \ldots, \alpha_r$ be the poles of R(z). Suppose that $|\alpha_i| = 1$ for all $1 \leq i \leq r$. Let α be in $\{\alpha_1, \ldots, \alpha_r\}$. It follows from the definition of residue and Remark 6.4 that $res(R, D_\alpha) = \sum_{\alpha_i \in D_\alpha} res(R, \alpha_i)$.

Lemma 6.7. Let K be a finite extension of \mathbb{Q}_p and let f(z) be in K[[z]] such that f(x) converges for all $x \in D_0$ and $f'(z)/f(z) \in E_{0,p}$.

(i) If D_{α} is a pole of f'(z)/f(z) then $res(f'(z)/f(z), D_{\alpha})$ belongs to \mathbb{Z}_p .

(ii) Suppose that $f(z) \in K[[z]]$ where K is a finite extension of \mathbb{Q}_p and let $R(z) = \prod_{i=1}^s (z - \beta_i)^{n_i}$ where, for each $1 \leq i \leq s$, $n_i \in \mathbb{Z} \setminus \{0\}$ and $|\beta_i| = 1$. If $\left|\frac{f'(z)}{f(z)} - \frac{R'(z)}{R(z)}\right|_{\mathcal{G}} \leq 1/p^n$ and D_α is a pole of f'(z)/f(z) then

$$res(f'(z)/f(z), D_{\alpha}) = \left(\sum_{\beta_i \in D_{\alpha}} n_i\right) \mod p^n \mathbb{Z}_p.$$

Proof. —

(i) We first show by induction on $n \ge 1$ that there exists a sequence $\{R_n(z)\}_{n\ge 1}$ of rational functions with coefficients in $\mathbb{C}_p(z)$ such that, for all $n, R_n(x) \ne 0$ for all $x \in D_0$ and

$$\left| \frac{f'(z)}{f(z)} - \sum_{i=1}^{n} \frac{R'_i(z)}{R_i(z)} \right|_{\mathcal{G}} < 1/p^n.$$
(6.2)

Since f(x) converges for all $x \in D_0$ and $f'(z)/f(z) \in E_{0,p}$, it follows from Theorem 3.1 of [24] that there exists $R_1(z) \in \mathbb{C}_p(z) \cap E_{0,p}$ such that, $R_1(x) \neq 0$ for all $x \in D_0$ and $\left| \frac{f'(z)}{f(z)} - \frac{R'_1(z)}{R_1(z)} \right|_{\mathcal{G}} < 1/p$. Now, let us suppose that there are $R_1(z), \ldots, R_n(z) \in \mathbb{C}_p(z)$ such that, for all $1 \leq j \leq n, R_j(x) \neq 0$ for all $x \in D_0$ and that Equation (6.2) is satisfied. Now, we put $g(z) = \frac{f(z)}{R_1(z)R_2(z)\cdots R_n(z)}$. Then

$$\frac{g'(z)}{g(z)} = \frac{f'(z)}{f(z)} - \sum_{i=1}^{n} \frac{R'_i(z)}{R_i(z)}$$

Notice that g(x) converges for all $x \in D_0$ and that g'(z)/g(z) belongs to $E_{0,p}$. Then, by applying Theorem 3.1 of [24] to g(z), there exists $R_{n+1}(z) \in \mathbb{C}_p(z) \cap E_{0,p}$ such that, $R_{n+1}(x) \neq 0$ for all $x \in D_0$ and $\left| \frac{g'(z)}{g(z)} - \frac{R'_{n+1}(z)}{R_{n+1}(z)} \right|_{\mathcal{G}} < 1/p^{n+1}$. In particular, we obtain

$$\left| \frac{f'(z)}{f(z)} - \sum_{i=1}^{n+1} \frac{R'_i(z)}{R_i(z)} \right|_{\mathcal{G}} < 1/p^{n+1}.$$

So, by induction we conclude that there is a sequence $\{R_n(z)\}_{n \ge 1}$ of rational functions with coefficients in \mathbb{C}_p such that, for all n, Equation (6.2) is satisfied. We now put $Q_j(z) = \sum_{i=1}^j \frac{R'_i(z)}{R_i(z)}$. Then, it follows from (6.2) that $f'(z)/f(z) = \lim_{j \to \infty} Q_j(z)$. Further, $Q_j(z)$ is a rational function of the shape $\sum \frac{n_i}{z - \alpha_i}$, where the n_i 's are integers. Further, it is clear that

$$Q_j(z) = \sum_{\alpha_i \in D_\alpha} \frac{n_i}{z - \alpha_i} + \sum_{\alpha_i \notin D_\alpha} \frac{n_i}{z - \alpha_i}.$$

By (ii) of Lemma 6.3, we deduce that $\sum_{\alpha_i \notin D_\alpha} \frac{n_i}{z - \alpha_i} \in \mathbb{C}_p[[z - \alpha]]$. Thus, from Remark 6.6, we deduce that $res(Q_j(z), D_\alpha) = \sum_{\alpha_i \in D_\alpha} n_i \in \mathbb{Z}$. So, $res(f'/f, D_\alpha) = \lim_{j \to \infty} res(Q_j(z), D_\alpha)$ and since, for every j, $res(Q_j(z), D_\alpha)$ is an integer, we get that $res(f'/f, D_\alpha)$ belongs to \mathbb{Z}_p .

(ii) Since $\left|\frac{f'(z)}{f(z)} - \frac{R'(z)}{R(z)}\right|_{\mathcal{G}} \leq 1/p^n$, it follows that $\frac{f'(z)}{f(z)} = \frac{R'(z)}{R(z)} \mod \pi_L^n \mathcal{O}_L[[z]]$, where $L = K(\beta_1, \dots, \beta_s)$

and π_L is a uniformizer of L. It is clear that $\frac{R'(z)}{R(z)} = \sum_{i=1}^s \frac{n_i}{z-\beta_i}$. So, Remark 6.6, we deduce that $res(R'(z)/R(z), D_\alpha) = \sum_{\beta_i \in D_\alpha} n_i$. Thus, we have $res(f'(z)/f(z), D_\alpha) = res(R'(z)/R(z), D_\alpha) \mod p^n$. So,

$$res(f'(z)/f(z), D_{\alpha}) = \left(\sum_{\beta_i \in D_{\alpha}} n_i\right) \mod p^n.$$

Let $f(z) = \sum_{j \ge 0} a_j (z - \alpha)^j \in \mathbb{C}_p[[z - \alpha]], \alpha \in \mathbb{C}_p$, and be a > 0 an integer different from. In this section, we define an *a*-th root of f(z). Let us suppose that $a_0 \neq 0$. Since \mathbb{C}_p is algebraically closed, the polynomial $X^a - a_0$ has all its roots in \mathbb{C}_p . We denote by $a_0^{1/a}$ one of these roots. So, the roots of $X^a - a_0$ are $a_0^{1/a} \exp(\frac{2\pi i k}{a})$ with $k = 0, 1, \ldots, a - 1$. We then define an *a*-th root of f(z) as follows

$$f(z)^{1/a} := a_0^{1/a} \left(\sum_{s \ge 0} \binom{1/a}{s} \frac{1}{a_0^s} T^s \right), \text{ where } \binom{1/a}{s} = \frac{(1/a)(1/a-1)\cdots(1/a-s+1)}{s!}$$

and $T(z) = \sum_{j \ge 1} a_j (z - \alpha)^j$. Then $f(z)^{1/a} = \sum_{j \ge 0} b_j (z - \alpha)^j$, where $b_0 = a_0^{1/a}$ and for any j > 0, $b_j \in \mathbb{Z} \left[a_1, \dots, a_j, \binom{1/a}{1}, \dots, \binom{1/a}{j}, \frac{1}{a_0}, a_0^{1/a} \right]$.

Recall that, for any set $A \subset \mathbb{C}_p \cup \{\infty\}$, E(A) is the the completion of the ring of the rational functions in $\mathbb{C}_p(z)$ that do not have poles in $A^{(6)}$.

Lemma 6.8. (i) Let $f(z) = \sum_{n \ge 0} a_n (z - \alpha)^n$ where, for all $n \ge 0$, $|a_n| \le 1$ and $|a_0| = 1$. If a is an integer such that p does not divide a then

$$f(z)^{1/a} = \sum_{n \ge 0} b_n (z - \alpha)^n \in \mathcal{O}_{\mathbb{C}_p}[[z - \alpha]] \quad and \ |b_0| = 1.$$

(ii) Let $P(z) \in \mathbb{C}_p(z)$ such that $P(z) \in E(D_\alpha)$ with $\alpha \in \mathcal{O}_{\mathbb{C}_p}$. Suppose that the poles and zeros of P(z) belong to $\mathcal{O}_{\mathbb{C}_p}$ and that $P(\alpha) \neq 0$. Then $P(z) = \sum_{\geq 0} a_n(z-\alpha)^n \in \mathcal{O}_L[[z-\alpha]]$ and $|a_0| = 1$, where L is a finite extension of \mathbb{Q}_p . Moreover, if p does not divide a then

$$P(z)^{1/a} = \sum_{n \ge 0} b_n (z - \alpha)^n \in \mathcal{O}_F[[z - \alpha]] \quad and \quad |b_0| = 1,$$

where F is the finite extension of L containing all the roots of $X^a - P(\alpha)$.

Proof. — (i) By definition,

$$f(z)^{1/a} = a_0^{1/a} \left(\sum_{s \ge 0} {\binom{1/a}{s} \frac{1}{a_0^s} T^s} \right) = \sum_{j \ge 0} b_j (z - \alpha)^j,$$

where $b_0 = a_0^{1/a}$ and for j > 0,

$$b_j \in \mathbb{Z}\left[a_1, \dots, a_j, \binom{1/a}{1}, \dots, \binom{1/a}{j}, \frac{1}{a_0}, a_0^{1/a}\right].$$

Since p does not divide a, $\binom{1/a}{s} \in \mathbb{Z}_p$ and we know that $|a_n| \leq 1$ for any $n \geq 0$. Thus, for all integers $n, |b_n| \leq 1$. Finally, as $b_0 = a_0^{1/a}$ then $|b_0| = 1$.

(ii) Let us write $P(z) = \frac{\prod_{i=1}^{n} (z-\alpha_i)^{\mu_i}}{\prod_{j=1}^{m} (z-\beta_j)^{\nu_j}}$, where $\mu_1, \ldots, \mu_n, \nu_1, \ldots, \nu_m$ belong to $\mathbb{N}_{>0}$. Let $L = \mathbb{Q}_p(\alpha, \alpha_1, \ldots, \alpha_n, \beta_1, \ldots, \beta_m)$. By assumption, $\beta_1, \ldots, \beta_m \in \mathcal{O}_{\mathbb{C}_p}$ and $P(z) \in E(D_\alpha)$. Then, Remark 6.1

⁽⁶⁾ We say that $R(z) = P(z)/Q(z) \in \mathbb{C}_p(z)$ does not have pole at ∞ if $deg(P) \leq deg(Q)$.

implies that, for every $1 \leq j \leq m$, $|\beta_j - \alpha| = 1$. Similarly, by assumption, $\alpha_1, \ldots, \alpha_n \in \mathcal{O}_{\mathbb{C}_p}$ and $P(\alpha) \neq 0$ and thus, Remark 6.1 implies that, for every $1 \leq i \leq n$, $|\alpha_i - \alpha| = 1$. Consequently, $|P(\alpha)| = 1$.

Now, it is clear that, for all $(i, j) \in \{1, \dots, n\} \times \{1, \dots, m\}$,

$$(z - \alpha_i)^{\mu_i} = \sum_{s=0}^{\mu_i} {\mu_i \choose s} (\alpha - \alpha_i)^s (z - \alpha)^{\mu_i - s} \in \mathcal{O}_L[z - \alpha]$$
$$(z - \beta_j)^{\nu_j} = \sum_{s=0}^{\nu_j} {\nu_j \choose s} (\alpha - \beta_j)^s (z - \alpha)^{\nu_j - s} \in \mathcal{O}_L[z - \alpha].$$

Since $|\alpha - \beta_j| = 1$, we have $|\alpha - \beta_j|^{\nu_j} = 1$. So, for all $1 \leq j \leq m$, $(z - \beta_j)^{\nu_j}$ is a unit of the ring $\mathcal{O}_L[[z - \alpha]]$ and thus, $\frac{1}{(z - \beta_j)^{\nu_j}} \in \mathcal{O}_L[[z - \alpha]]$. Consequently, $P(z) = \sum_{n \geq 0} a_n (z - \alpha)^n \in \mathcal{O}_L[[z - \alpha]]$ and $|a_0| = 1$ because $P(\alpha) = a_0$ and we have already seen that $|P(\alpha)| = 1$.

Finally, from (i) we know that $P(z)^{1/a} = \sum_{n \ge 0} b_n (z - \alpha)^n$, where, $b_0 = a_0^{1/a}$ and for j > 0, $b_j \in \mathbb{Z}\left[a_1, \ldots, a_j, \binom{1/a}{1}, \ldots, \binom{1/a}{j}, \frac{1}{a_0}, a_0^{1/a}\right]$. Thus, $P(z)^{1/a} \in \mathcal{O}_F[[z - \alpha]]$ because $\binom{1/a}{s} \in \mathbb{Z}_p$ for all $s \ge 0$ (recall that p does not divide a) and we know that $a_n \in \mathcal{O}_L$ for any $n \ge 1$. Finally, $|b_0| = 1$ because $b_0^a = a_0 = P(\alpha)$ and we have already seen that $|P(\alpha)| = 1$.

Remark 6.9. Let $f(z) = \sum_{n \ge 0} a_n z^n$ be in $\mathcal{O}_K[[z]]$, K an extension of \mathbb{Q}_p , such that $|f|_{\mathcal{G}} = 1 = |a_0|$. It follows from the proof of previous lemma that if p does not divide a then $f(z)^{1/a} = \sum_{n \ge 0} b_n z^n \in \mathcal{O}_L[[z]]$ and $|f^{1/a}|_{\mathcal{G}} = 1 = |b_0|$, where L is a finite extension of K containing all the roots of $X^a - a_0$.

6.0.3. Exponents and regular singular points

Let K be a finite extension of \mathbb{Q}_p and let $\mathcal{O}_{K(z)}$ be the set of rational functions $R(z) \in K(z)$ with Gauss norm less than or equal to 1 and let \mathcal{M}_K be the set of rational functions $R(z) \in K(z)$ with Gauss norm less than 1. Note that \mathcal{M}_K is the maximal ideal of $\mathcal{O}_{K(z)}$. Now, let us take \mathcal{L} in $\mathcal{O}_{K(z)}[d/dz]$. Then, we can reduce \mathcal{L} modulo \mathcal{M}_K^t for any integer $t \ge 1$. The reduction of \mathcal{L} modulo \mathcal{M}_K^t is denoted by $\mathcal{L}_{|\mathcal{M}_K^t}$.

Definition 6.10. — Let $\mathcal{L} = \frac{d}{dz^n} + b_1(z)\frac{d}{dz^{n-1}} + \cdots + b_{n-1}(z)\frac{d}{dz} + b_n(z)$ be in $\mathcal{O}_{K(z)}[d/dz]$ with K a finite extension of \mathbb{Q}_p , let α be an element of \mathbb{C}_p of norm 1, let $L = K(\alpha)$ and let $t \ge 0$ be an integer. We say that α is a regular singular point of $\mathcal{L}_{|\mathcal{M}_K^t|}$ if, for every $1 \le j \le n$, $(z - \alpha)^j b_j(z) = F_j(z) \mod \mathfrak{M}_L^t \mathcal{O}_{L(z)}$ with $F_j \in L(z) \cap \mathcal{O}_L[[z - \alpha]]$. Further, any solution of the polynomial

 $P_{\alpha,t}(X) = X(X-1)\cdots(X-n+1)+s_1X(X-1)\cdots(X-n+2)+\cdots+s_{n-1}X+s_n, \ s_j = F_j(\alpha) \ \text{mod} \ \pi_L^t \mathcal{O}_L$ is called an exponent of $L_{|\mathcal{M}_K^t}$ at α . This polynomial will be called the indicial polynomial of $\mathcal{L}_{|\mathcal{M}_K^t}$ at α . Notice that $P_{\alpha,t}(X)$ is a polynomial with coefficients in \mathcal{O}_L/π_L^t .

Lemma 6.11. — Let $\mathcal{L} = \frac{d}{dz^n} + b_1(z)\frac{d}{dz^{n-1}} + \cdots + b_{n-1}(z)\frac{d}{dz} + b_n(z)$ be in $\mathcal{O}_{K(z)}[d/dz]$ with K a finite extension of \mathbb{Q}_p , let α be an element of \mathbb{C}_p with norm 1, $L = K(\alpha)$ and let $h(z) = (z - \alpha)^{\mu} \left(\sum_{j \ge 0} a_j(z - \alpha)^j \right) \in (z - \alpha)^{\mu} \mathcal{O}_K[[z - \alpha]]$ with $\mu \in \mathbb{Z}_{(p)}$ and $|a_0| = 1$. Suppose that α is a regular singular point of $\mathcal{L}_{|\mathcal{M}_K^t}$ for some integer $t \ge 0$. If $h(z) \mod \pi_L^t \mathcal{O}_L[[z - \alpha]]$ is solution of $\mathcal{L}_{|\mathcal{M}_L^t}$ then $\mu \mod \pi_L^t$ is an exponent of $\mathcal{L}_{|\mathcal{M}_K^t}$ at α , where π_L is a uniformizer of L.

Proof. — By assumption, for all $1 \leq j \leq n$, $(z-\alpha)^j b_j(z) = F_j(z) \mod \mathcal{M}_L^t \mathcal{O}_{L(z)}$, where $F_j \in L(z) \cap$ $\mathcal{O}_L[[z-\alpha]]$. Notice that $(z-\alpha)$ is a unit in $\mathcal{O}_{L(z)}$ because $|\alpha| = 1$. Then, $b_j(z) = \frac{F_j(z)}{(z-\alpha)^j} \mod \mathfrak{M}_L^t \mathcal{O}_{L(z)}$. By assumption again, $h(z) \mod \pi_L^t$ is solution of $\mathcal{L}_{|\mathcal{M}_L^t}$. Then,

$$h^{(n)} + \frac{F_1}{(z-\alpha)}h^{(n-1)} + \dots + \frac{F_n}{(z-\alpha)^n}h(z) = 0 \mod \pi_L^t \mathcal{O}_L\left[\left[z-\alpha, (z-\alpha)^{\mu}, \frac{1}{z-\alpha}\right]\right].$$
 (6.3)

It is clear that $h(z) \mod \pi_L^t = \overline{a_0}(z - \overline{\alpha})^\mu + \overline{a_1}(z - \overline{\alpha})^{\mu+1} + \dots + \overline{a_j}(z - \overline{\alpha})^{\mu+j} + \dots +$, where for any $x \in \mathcal{O}_L, \overline{x}$ denotes the reduction of x modulo π_L^t . It follows from Equation (6.3) that

$$a_0[\mu(\mu-1)\cdots(\mu-n+1)+s_1\mu(\mu-1)\cdots(\mu-n+2)+\cdots+s_{n-1}\mu+s_n]=0 \mod \pi_L^t.$$

But $a_0 \mod \pi_L^t \mathcal{O}_L$ is a unit of the ring \mathcal{O}_L/π_L^t because $|a_0| = 1$. So $\mu \mod \pi_L^t$ is a root of $P_{\alpha,t}(X)$.

6.1. A criterion for algebraic independence

We can now state our criterion for algebraic independence.

Theorem 6.12. — Let K be a finite extension of \mathbb{Q}_p . Suppose that K is a Frobenius field and let $f_1(z), \ldots, f_r(z)$ be in $\mathcal{MF}(K)$. Let $\mathcal{L}_1, \ldots, \mathcal{L}_r$ be in $\mathcal{O}_{K(z)}[d/dz]$ such that $\mathcal{L}_i(f_i) = 0$ for every $1 \leq i \leq r.$

- (1) Suppose that, for every $i \in \{1, \ldots, r\}$, there exists $D_{\alpha_{c_i}} \in \mathbf{Pol}(f'_i(z)/f_i(z))$ such that $D_{\alpha_{c_i}} \notin \mathcal{O}(f'_i(z))$ $\operatorname{Pol}(f'_j(z)/f_j(z)) \text{ for all } j \in \{1, \ldots, r\} \setminus \{i\} \text{ and that } \operatorname{res}(\frac{f'_i(z)}{f_i(z)}, D_{\alpha_{c_i}}) \neq 0.$ (2) Suppose that, for each $i \in \{1, \ldots, r\}, \alpha_{c_i}$ is a regular singular point of $\mathcal{L}_{i|\mathcal{M}_{K}^2}$.

If $f_1(z), \ldots, f_r(z)$ are algebraically dependent over E_K then there exists $s \in \{1, \ldots, r\}$ such that $res\left(\frac{f'_{s}(z)}{f_{s}(z)}, D_{\alpha_{c_{s}}}\right)$ belongs to \mathbb{Z}_{p} and its reduction modulo p^{2} is an exponent of $\mathcal{L}_{s|\mathcal{M}_{K}^{2}}$ at $\alpha_{c_{s}}$.

The proof of this theorem is based on Theorem 6.13, which is a more explicit version of Theorem 2.3.

Theorem 6.13. — Let K be a finite extension of \mathbb{Q}_p and $f_1(z), \ldots, f_m(z)$ be in $\mathcal{MF}(K)$. Let us suppose that, for every $1 \leq i \leq m$, there exists $D_{\alpha_{c_i}} \in \mathbf{Pol}(f'_i(z)/f_i(z))$ such that $D_{\alpha_{c_i}} \notin \mathbf{Pol}(f'_i(z)/f_i(z))$ $Pol(f'_j(z)/f_j(z))$ for all $j \in \{1, \ldots, m\} \setminus \{i\}$ and that $res\left(\frac{f'_i(z)}{f_i(z)}, D_{\alpha_{c_i}}\right) \neq 0$. If K is a Frobenius field and $f_1(z), \ldots, f_m(z)$ are algebraically dependent over E_K then there exist $d_1, \ldots, d_m \in \mathbb{Z}$ not all zero, $\mathfrak{a}_1(z) \in L(z)$ and $\mathfrak{j}_1(z) \in \mathcal{O}_L[[z]]$, L a finite extension of K, such that

(i)

$$f_1^{d_1}(z) \cdots f_m^{d_m}(z) = \prod_{i=1}^m (z - \alpha_{c_i})^{\Gamma_i} \mathfrak{a}_1(z) \mathfrak{j}_1(z), \tag{6.4}$$

where $\Gamma_i = d_i \cdot res\left(\frac{f_i'(z)}{f_i(z)}, D_{\alpha_{c_i}}\right) \in \mathbb{Z}.$

Moreover,

- (ii) there exists $d_s \in \{d_1, \ldots, d_m\}$ such that p does not divide d_s ,
- (iii) $|\mathfrak{a}_1(z)|_{\mathcal{G}} = 1$ and, for every $1 \leq i \leq m$, $\mathfrak{a}_1(z) \in E(D_{\alpha_{c_i}})$ and $\mathfrak{a}_1(\alpha_{c_i}) \neq 0$, and all the poles and zeros of $\mathfrak{a}_1(z)$ have norm 1.

- (iv) $|\mathfrak{j}_1(z)| = 1$, $\mathfrak{j}_1(z) = \mathfrak{j}_1(0) \mod \pi_L \mathcal{O}_L[[z]]$, and $\mathfrak{j}'(z)$ belongs to $\pi_L^2 \mathcal{O}_L[[z]]$,
- (v) there is an integer N > 0 such that $(f_1^{d_1}(z) \cdots f_m^{d_m}(z))^N \in E_{0,K}$.

Before proving Theorem 6.12, we need the following remark.

Remark 6.14. Let f(z) be in $\mathcal{MF}(K)$ with K a finite extension of \mathbb{Q}_p . If K is a Frobenius field then $f(z) \equiv P(z)t(z^{p^2}) \mod \mathcal{O}_K[[z]]$, where $P(z) \in K_0(z), |P(z)|_{\mathcal{G}} = 1$, and $t(z) \in C_0(z)$ $1 + z\mathcal{O}_K[[z]]$. Indeed, thanks to Theorem 3.1, $f(z)/f(z^{p^h}) \in E_{0,p}$, where h is the period of the Frobenius structure of the differential operator annihilating by f(z). Thus, there is $B(z) \in K_0(z)$ such that $f(z)/f(z^{p^h}) \equiv B \mod \pi_K^2 \mathcal{O}_K[[z]]$. By Theorem 3.1 again, we also know that $f(z) \in 1 + z\mathcal{O}_K[[z]]$ and thus, $|f(z)/f(z^{p^h})|_{\mathcal{G}} = 1$. Given that the norm is non-Archimedean, the equality $f(z)/f(z^{p^h}) \equiv B \mod B$ $\pi_K^2 \mathcal{O}_K[[z]]$ implies that $|B(z)|_{\mathcal{G}} = 1$. In addition, $f(z)/f(z^{p^h}) \equiv B \mod \pi_K^2 \mathcal{O}_K[[z]]$ also implies that $f(z)/f(z^{p^{2h}}) \equiv B(z)B(z^{p^{h}}) \mod \pi_{K}^{2}\mathcal{O}_{K}[[z]].$ Whence, $f(z) \equiv B(z)B(z^{p^{h}})f(z^{p^{2h}}) \mod \pi_{K}^{2}\mathcal{O}_{K}[[z]].$ So, we take $P(z) = B(z)B(z^{p^{h}})$ and $t(z) = f(z^{p^{2h-2}}).$

Proof of Theorem 6.12. — Let us write $\mathbf{Pol}(f'_i(z)/f_i(z)) = \{D_{\alpha_{1,i}}, \dots, D_{\alpha_{g_i,i}}\}$. By Theorem 3.1, we know that $f'_i(z)/f_i(z) \in E_{0,K}$ and thus, by using Equation (6.1), we conclude that $|\alpha_{l,i}| = 1$ for every $1 \leq l \leq g_i$.

Since $f_1(z), \ldots, f_r(z)$ are algebraically dependent over E_K , according to Theorem 6.13, there are $d_1, \ldots, d_r \in \mathbb{Z}$ not all zero such that

$$f_1^{d_1}\cdots f_r^{d_r} = \mathfrak{q}_1(z)\mathfrak{a}_1(z)\mathfrak{j}_1(z) \text{ where } \mathfrak{q}_1(z) = \prod_{i=1}^r (z-\alpha_{c_i})^{\Gamma_i}, \ \mathfrak{a}_1(z) \in \boldsymbol{L}(z), \text{ and } \mathfrak{j}_1(z) \in \mathcal{O}_{\boldsymbol{L}}[[z]],$$

where \boldsymbol{L} is a finite extension of K.

We also know from Theorem 6.13 that $\mathfrak{j}_1'(z) \in \pi_L^2 \mathcal{O}_L[[z]], |\mathfrak{j}_1(0)| = 1, \text{ and } \Gamma_i = d_i \cdot res\left(\frac{f_i'(z)}{f_i(z)}, D_{\alpha_{c_i}}\right) \in \mathcal{O}_L[[z]]$ \mathbb{Z} for all $1 \leq i \leq r$, and that there is $d_s \in \{d_1, \ldots, d_r\}$ such that p does not divide d_s . Further, Theorem 6.13 also says that, for all $1 \leq i \leq r$, $\mathfrak{a}_1(z) \in E(D_{\alpha_{c_i}})$ and $\mathfrak{a}_1(\alpha_{c_i}) \neq 0$, and all the poles and zeros of $\mathfrak{a}_1(z)$ have norm 1.

It follows from Remark 6.14 that, for every, $j \in \{1, \ldots, r\}$, $f_j(z) = P_j(z)t_j(z^{p^2}) \mod \pi_K^2 \mathcal{O}_K[[z]]$, where $P_j(z) \in K_0(z)$ with $|P_j(z)|_{\mathcal{G}} = 1$ and $t_j(z) \in 1 + z\mathcal{O}_K[[z]]$. So, for every $1 \leq j \leq r$, we have $\left|\frac{f_j'(z)}{f_j(z)} - \frac{P_i'(z)}{P_i(z)}\right|_{\mathcal{G}} \leqslant 1/p^2$. Therefore, the poles and zeros of $P_j(z)$ belong to $D_{\alpha_{1,j}} \cup \cdots \cup D_{\alpha_{g_j,j}}$. In particular, $P_j(z) \in E(D_0)$ for all $1 \leq j \leq r$. Further, for all $1 \leq j \leq r$ and, for all $1 \leq l \leq g_j, 0 \notin D_{\alpha_{l,j}}$ because $|\alpha_{l,j}| = 1$ and consequently $P_j(0) \neq 0$ for all $1 \leq i \leq r$.

Note that, for all $j \in \{1, \ldots, r\} \setminus \{s\}, P_j(z) \in E(D_{\alpha_{c_s}})$ and $P_j(\alpha_{c_s}) \neq 0$. Indeed, since $j \neq s$, we get, by assumption, that $D_{\alpha_{c_s}} \notin \mathbf{Pol}(f'_j(z)/f_j(z))$. But, recall that $\mathbf{Pol}(f'_j(z)/f_j(z)) = \{D_{\alpha_{1,j}}, \ldots, D_{\alpha_{g_j,j}}\}$ and we have already seen that the poles and zeros of $P_j(z)$ belong to $D_{\alpha_{1,j}} \cup \cdots \cup D_{\alpha_{g_j,j}}$. Whence, for all $j \in \{1, \ldots, r\} \setminus \{s\}, P_j(z) \in E(D_{\alpha_{c_s}})$ and $P_j(\alpha_{c_s}) \neq 0$.

Now, let F be the finite extension of L such that every root of the following polynomials is in F.

- (1) $X^{d_s} P_j(\alpha_{c_s})$, with $j \in \{1, \dots, r\} \setminus \{s\}$, (2) $X^{d_s} - \alpha_{c_i}$ for all $1 \leq i \leq r$, (3) $X^{d_s} - P_j(0)$ for all $1 \leq j \leq r$,
- (4) $X^{d_s} \mathfrak{a}_1(\alpha_{c_s}), X^{d_s} \mathfrak{j}_1(0), \text{ and } X^{d_s} \mathfrak{a}_1(0).$

DANIEL VARGAS-MONTOYA

We know that $|\alpha_{c_i}| = 1$ for every $1 \leq i \leq r$. So, it follows from (ii) of Lemma 6.8 that, for every $1 \leq i \leq r$, $(z - \alpha_{c_i})^{\frac{\Gamma_i}{d_s}} \in \mathcal{O}_{\mathbf{F}}[[z]]$. We also know that all the poles and zeros of $\mathfrak{a}_1(z)$ have norm 1. Hence, $\mathfrak{a}_1(z) \in E(D_0)$ and $\mathfrak{a}_1(0) \neq 0$. Then, by (ii) of Lemma 6.8, we get that $\mathfrak{a}_1(z)^{1/d_s}$ belongs to $\mathcal{O}_{\mathbf{F}}[[z]]$.

We have already seen that, for all $1 \leq j \leq r$, $P_j(z) \in E(D_0)$ and that $P_j(0) \neq 0$. So, from (ii) of Lemma 6.8, we deduce that, $j \in \{1, \ldots, r\}$, $P_j(z)^{\frac{-d_j}{d_s}} \in \mathcal{O}_{\mathbf{F}}[[z]]$. In addition, from Lemma 6.8, we also know that $|P_j(0)| = 1$ for all $1 \leq j \leq r$.

Since, for every, $i \in \{1, ..., r\}$, $f_i(z) = P_i(z)t_i(z^{p^2}) \mod \pi_K^2 \mathcal{O}_K[[z]]$, there exists $g_i \in \mathcal{O}_K[[z]]$ such that $f_i(z) = P_i(z)t_i(z^{p^2}) + \pi_K^2 g_i(z)$. So, it is clear that $f_i(z) = P_i(z)t_i(z^{p^2}) \left(1 + \pi_K^2 \frac{g_i(z)}{P_i(z)t_i(z^{p^2})}\right)$. Whence

$$f_i(z)^{-\frac{d_i}{d_s}} = (P_i(z)t_i(z^{p^2}))^{-\frac{d_i}{d_s}} \left(1 + \pi_L^2 \frac{g_i(z)}{P_i(z)t_i(z^{p^2})}\right)^{-\frac{a_i}{d_s}}.$$

We have

$$\left(1 + \pi_K^2 \frac{g_i(z)}{P_i(z)t_i(z^{p^2})}\right)^{-\frac{d_i}{d_s}} = \sum_{j \ge 0} \binom{-d_i/d_s}{j} \left(\pi^2 \frac{g_i(z)}{P_i(z)t_i(z^{p^2})}\right)^j$$

Given that $|P_i(z)t_i(z^{p^2})|_{\mathcal{G}} = 1$ and that $g_i \in \mathcal{O}_K[[z]]$, we have $\left|\frac{g_i(z)}{P_i(z)t_i(z^{p^2})}\right|_{\mathcal{G}} \leq 1$. In addition, for all $j \ge 0$, $\binom{-d_i/d_s}{j} \in \mathbb{Z}_p$ because p does not divide d_s . Therefore, $\left(1 + \pi_K^2 \frac{g_i(z)}{P_i(z)t_i(z^{p^2})}\right)^{-\frac{d_i}{d_s}} = 1 \mod \pi_F^2$.

Thus, we deduce that $f_i(z)^{-\frac{d_i}{d_s}} = (P_i(z)t_i(z^{p^2}))^{-\frac{d_i}{d_s}} \mod \pi_F^2 \mathcal{O}_F[[z]].$

We know that, for every $1 \leq i \leq r$, $t_i(z) \in 1 + z\mathcal{O}_K[[z]]$ and thus, by Remark 6.9, we deduce that, for every $i \in \{1, \ldots, r\}$,

$$t_i(z)^{-\frac{a_i}{d_s}} = \sum_{j \ge 0} \rho_{i,j} z^j \in \mathcal{O}_F[[z]] \text{ with } |\rho_{i,0}| = 1.$$

In addition, we know that $j_1(z) \in \mathcal{O}_L[[z]]$, with $|j_1(0)| = 1$ and thus, from Remark 6.9, we deduce that $j_1(z)^{1/d_s} \in \mathcal{O}_F[[z]]$.

Now, it is clear that

$$f_s(z) = \prod_{i=1}^r (z - \alpha_{c_i})^{\frac{\Gamma_i}{d_s}} \mathfrak{a}_1(z)^{1/d_s} \mathfrak{j}_1(z)^{1/d_s} \prod_{i=1, i \neq s}^r f_i(z)^{-\frac{d_i}{d_s}}.$$

So, by reducing the previous equality modulo π_F^2 , we get

$$f_s(z) = \mathfrak{r}(z)\mathfrak{c}(z) \mod \pi_F^2 \mathcal{O}_F[[z]]$$

where

$$\mathfrak{r}(z) = \prod_{i=1}^{r} (z - \alpha_i)^{\frac{\Gamma_i}{d_s}} \mathfrak{a}_1(z)^{1/d_s} \left(\prod_{i=1, i \neq s}^{r} P_i(z)^{-\frac{d_i}{d_s}} \right) \text{ and } \mathfrak{c}(z) = \left(\prod_{i=1, i \neq s}^{r} \left(\sum_{j \ge 0} \rho_{i,j} z^{p^2 j} \right) \right) \mathfrak{j}_1(z)^{1/d_s}.$$

By assumption, $\mathcal{L}_s(f_s) = 0$. So $\mathcal{L}_{s|\mathcal{M}_F^2}(f_s \mod \pi_F^2 \mathcal{O}_F) = 0$. Thus, $\mathcal{L}_{s|\mathcal{M}_F^2}(\mathfrak{r}(z)\mathfrak{c}(z)) = 0$. Now, we know that and $\mathfrak{j}'_1(z) \in \pi_L^2 \mathcal{O}_L[[z]]$ and give that $\pi_L^2 \in \pi_F^2 \mathcal{O}_F$ then $\mathfrak{j}'_1(z) \in \pi_F^2 \mathcal{O}_F[[z]]$. Further, we also have $p^2 \in \pi_F^2 \mathcal{O}_F$. Therefore, $\mathfrak{c}'(z) = 0 \mod \pi_F^2 \mathcal{O}_F[[z]]$. Whence

$$\mathcal{L}_{s|\mathcal{M}_{F}^{2}}(\mathfrak{r}(z)\mathfrak{c}(z)) = \mathfrak{c}(z)\mathcal{L}_{s|\mathcal{M}_{F}^{2}}(\mathfrak{r}(z)) = 0$$

Note that $\mathfrak{c}(z) \mod \pi_F^2 \mathcal{O}_F[[z]]$ is a unit of the ring $\frac{\mathcal{O}_F}{\pi_F^2}[[z]]$ because $|\rho_{i,0}| = 1$ for all $i \in \{1, \ldots, r\}$ and $|\mathfrak{j}_1(z)| = 1$. Therefore, $\mathcal{L}_{s|\mathcal{M}_F^2}(\mathfrak{r}(z)) = 0$

By assumption, $D_{\alpha_{c_s}} \notin \mathbf{Pol}(f'_i(z)/f_i(z))$ for all $i \in \{1, \ldots, r\} \setminus \{s\}$. So, $D_{\alpha_{c_i}} \neq D_{\alpha_{c_s}}$ for all $i \in \{1, \ldots, r\} \setminus \{s\}$ because $D_{\alpha_{c_i}} \in \mathbf{Pol}(f'_i(z)/f_i(z))$. Then, Remark 6.1 implies that if $1 \leq i \leq r$ with $i \neq s$ then $|\alpha_{c_s} - \alpha_{c_i}| = 1$. Since p does not divide d_s and Γ_i is an integer, from Lemma 6.8, we get that, for all $1 \leq i \leq r$ with $i \neq s$,

$$(z - \alpha_{c_i})^{\frac{\Gamma_i}{d_s}} = \sum_{j \ge 0} \zeta_{j,i} (z - \alpha_{c_s})^j \in \mathcal{O}_{\boldsymbol{F}}[[z - \alpha_{c_s}]]$$

with $|\zeta_{0,i}| = 1$.

Consequently,

$$\left(\prod_{i=1,i\neq s}^{r} (z-\alpha_{c_i})^{\Gamma_i/d_s}\right) = \left(\prod_{i=1,i\neq s}^{r} \left(\sum_{j\geqslant 0} \zeta_{j,i}(z-\alpha_{c_s})^j\right)\right) = \sum_{j\geqslant 0} \xi_j(z-\alpha_{c_s})^j \in \mathcal{O}_{\boldsymbol{F}}[[z-\alpha_{c_s}]]$$

with $|\xi_0| = 1$.

We know that $\mathfrak{a}_1(z) \in E(D_{\alpha_{c_s}})$, $\mathfrak{a}_1(\alpha_{c_s}) \neq 0$ and all the poles and zeros of $\mathfrak{a}_1(z)$ have norm 1. So, by (ii) of Lemma 6.8, we get that

$$\mathfrak{a}_1(z)^{1/d_s} = \sum_{j \ge 0} \epsilon_j (z - \alpha_{c_s})^j \in \mathcal{O}_{\mathbf{F}}[[z - \alpha_{c_s}]] \text{ with } |\epsilon_0| = 1.$$

We have seen that if $i \in \{1, ..., r\} \setminus \{s\}$ then $P_i(z) \in E(D_{\alpha_{c_s}})$ and $P_i(\alpha_{c_s}) \neq 0$. Consequently, by Lemma 6.8, we deduce that, for any $i \neq s$,

$$P_i(z)^{-\frac{d_i}{d_s}} = \sum_{j \ge 0} \gamma_{i,j} (z - \alpha_{c_s})^j \in \mathcal{O}_F[[z - \alpha_{c_s}]] \text{ with } |\gamma_{i,0}| = 1.$$

Then

$$\left(\sum_{j\geq 0}\xi_j(z-\alpha_{c_s})^j\right)\left(\sum_{j\geq 0}\epsilon_j(z-\alpha_{c_s})^j\right)\prod_{i=1,i\neq s}^r\left(\sum_{j\geq 0}\gamma_{i,j}(z-\alpha_{c_s})^j\right)\in\mathcal{O}_{\boldsymbol{F}}[[z-\alpha_{c_s}]]$$

and $\left|\xi_0\epsilon_0\prod_{i=1,i\neq s}^r\gamma_{i,0}\right|=1.$

Further, it is clear that

$$\mathfrak{r}(z) = (z - \alpha_{c_s})^{\frac{\Gamma_s}{d_s}} \left(\sum_{j \ge 0} \xi_j (z - \alpha_{c_s})^j \right) \left(\sum_{j \ge 0} \epsilon_j (z - \alpha_{c_s})^j \right) \prod_{i=1, i \ne s}^r \left(\sum_{j \ge 0} \gamma_{i,j} (z - \alpha_{c_s})^j \right).$$

Note that Γ_s/d_s belongs to $\mathbb{Z}_{(p)}$ because Γ_s is an integer and p does not divide d_s . So, we are able to apply Lemma 6.11 and we deduce that $\Gamma_s/d_s \mod p^2$ is an exponent of $L_{s|\mathcal{M}_F^2}$ at α_{c_s} . Finally, we already know that

$$\frac{\Gamma_s}{d_s} = res\left(\frac{f'_s(z)}{f_s(z)}, D_{\alpha_{c_s}}\right).$$

6.2. Proof of Theorem 6.13

The proof of this Theorem 6.13 is derived from Theorems 2.3 and 6.16. Let us say that Theorem 6.16 is a direct consequence of Theorems 59.8 and 59.11 of [19]. In order to state this theorem, we need to recall the notion of a *strongly copiercing sequence associated to* D_0 . Following [19, p. 160], we say that an infinite sequence $(a_n, b_n)_{n \ge 0}$ is a strongly copiercing sequence associated to D_0 if, for all $n \ge 0$, $a_n, b_n \in \mathbb{C}_p \setminus D_0$, $|a_n - b_n| < \delta(a_n, D_0)^{(7)}$ and $\lim_{n \to \infty} \frac{|a_n - b_n|}{\delta(a_n, D_0)} = 0$.

Remark 6.15. If $(a_n, b_n)_{n \ge 0}$ is a strongly copiercing sequence associated to D_0 then $\mathfrak{t}(z) = \prod_{i=0}^{\infty} \left(\frac{z-a_n}{z-b_n}\right)$ belongs to $E_{0,p}$, $|\mathfrak{t}(z)-1|_{\mathcal{G}} < 1$, and $|\mathfrak{t}(0)| = 1$. In order to prove that, let us consider $\mathfrak{t}_r(z) = \prod_{n=1}^r \left(\frac{z-a_n}{z-b_n}\right)$ for every $r \ge 1$. Then

$$\mathfrak{t}_r(z) - \mathfrak{t}_{r+1}(z) = \mathfrak{t}_r(z) \left(1 - \frac{z - a_{r+1}}{z - b_{r+1}} \right).$$

Since, for all $n \ge 1$, $|a_n| \ge 1$ and the norm is non-Archimedean, we get that $\delta(a_n, D_0) = |a_n|$ and for this reason $|a_n - b_n| < |a_n|$ for all $n \ge 1$. So, using the fact that norm is non-Archimedean again, we get that $|b_n| = |a_n|$ for all $n \ge 1$. Hence, for every $r \ge 1$,

$$\left|1 - \frac{z - a_r}{z - b_r}\right|_{\mathcal{G}} = \frac{|a_r - b_r|}{|b_r|} = \frac{|a_r - b_r|}{\delta(a_r, D_0)} < 1.$$
(6.5)

As $\lim_{n\to\infty} \frac{|a_n-b_n|}{\delta(a_n,D_0)} = 0$ then $\lim_{r\to\infty} \left| 1 - \frac{z-a_r}{z-b_r} \right|_{\mathcal{G}} = 0$. Whence, $\mathfrak{t}(z)$ is the limit of the rational functions $\mathfrak{t}_r(z)$. Note that this rational function does have poles in the D_0 because $b_n \notin D_0$ for all $n \ge 1$. Further, it follows from Equation (6.5) that, for all $n \ge 1$, $\frac{z-a_n}{z-b_n} = 1 \mod \mathcal{MO}_{\mathbb{C}_p(z)}$, where \mathcal{M} is the maximal ideal of $\mathcal{O}_{\mathbb{C}_p(z)}$. Thus, for every $r \ge 1$, $\mathfrak{t}_r(z) = 1 \mod \mathcal{MO}_{\mathbb{C}_p(z)}$. Whence, $\mathfrak{t}(z) = 1 \mod \mathcal{MO}_{\mathbb{C}_p(z)}$ and consequently, $|\mathfrak{t}(z) - 1|_{\mathcal{G}} < 1$. Finally, by Equation (6.5) again, we get that, for ever $r \ge 1$, $\frac{a_r}{b_r} = 1 \mod \mathfrak{mO}_{\mathbb{C}_p}$ and hence $\mathfrak{t}_r(0) = 1 \mod \mathfrak{mO}_{\mathbb{C}_p}$ for every $r \ge 1$. So $\mathfrak{t}(0) = 1 \mod \mathfrak{mO}_{\mathbb{C}_p}$. Whence, $|\mathfrak{t}(0)| = 1$.

Theorem 6.16 (Theorems 59.8 and 59.11 [19]). — Let K be a finite extension of \mathbb{Q}_p and f(z) be in $1 + z\mathcal{O}_K[[z]]$ such that $f'(z)/f(z) \in E_{0,K}$. Let $D_{\alpha_1}, \ldots, D_{\alpha_q}$ be the poles of f'(z)/f(z) such that $\mu_i = res\left(\frac{f'(z)}{f(z)}, D_{\alpha_i}\right) \neq 0$ for all $1 \leq i \leq q$. If $f(z) \in E_{0,K}$ then $\mu_i \in \mathbb{Z}$ and, for every $i \in \{1, \ldots, q\}$, there is $\beta_i \in D_{\alpha_i}$ such that

$$f(z) = \prod_{i=1}^{q} (z - \beta_i)^{\mu_i} \mathfrak{t}(z),$$
(6.6)

where $\mathfrak{t}(z) = \prod_{i=1}^{\infty} \left(\frac{z-a_n}{z-b_n}\right)$ with $(a_n, b_n)_{n \ge 1}$ a strongly copiercing sequence associated to D_0 . Moreover, $\frac{f'(z)}{f(z)} - \sum_{i=1}^{q} \frac{\mu_i}{z-\beta_i}$ is integrable in $E_{0,p}$. That is, there exists $F \in E_{0,p}$ such that $F'(z) = \frac{f'(z)}{f(z)} - \sum_{i=1}^{q} \frac{\mu_i}{z-\beta_i}$.

Remark 6.17. We would like to say that Theorem 59.8 of [19] is stated in a slightly different way. Let us be more precise. In [19], the author considers $f'/f \in E(D)$, where D is a well pierced set⁽⁸⁾. Then, he shows that f(z) is invertible in E(D) if and only if f(z) satisfies a decomposition

 $^{^{(7)}\}delta(a, D_0) = \inf\{|a - x| : x \in D_0\}.$

⁽⁸⁾ Following [19, p.106], we say that a set $D \subset \mathbb{C}_p$ is well pierced if $\delta(\overline{D}, \mathbb{C}_p \setminus D) > 0$, where \overline{D} is the topological closure of D and $\delta(\overline{D}, \mathbb{C}_p \setminus D) := \inf\{|b-a| : b \in \mathbb{C}_p \setminus D, a \in \overline{D}\}$. For example, D_0 is well pierced because $\overline{D_0} = D_0$ and $\delta(D_0, \mathbb{C}_p \setminus D_0) = 1$.

of the shape (6.6). So, Theorem 6.16 is a version of this theorem for $D = D_0$ because D_0 is a well pierced set and if $f(z) \in 1 + z\mathcal{O}_K[[z]] \cap E_{0,K}$ then f(z) is an invertible element of $E_{0,K}$ and thus f(z)is invertible element of $E(D_0)$ because $E(D_0) = E_{0,p}$. Finally, the fact that $f'(z)/f(z) - \sum_{i=1}^{q} \frac{\mu_i}{z - \beta_i}$ is integrable in $E_{0,p}$ is a direct consequence of Theorem 59.8 and Theorem 59.11 of [19].

We are now ready to prove Theorem 6.13.

Proof of Theorem 6.13. — Let K be a finite extension of \mathbb{Q}_p and $f_1(z), \ldots, f_m(z)$ be in $\mathcal{MF}(K)$ with K a Frobenius field.

Suppose that $f_1(z), \ldots, f_m(z)$ are algebraically dependent over E_K . By Theorem 2.3, there exist $a_1, \ldots, a_m \in \mathbb{Z}$ not all zero, such that

$$f_1(z)^{a_1}\cdots f_m(z)^{a_m}\in E_{0,K}.$$

We put $h(z) = f_1(z)^{a_1} \cdots f_m(z)^{a_m}$. Notice that $h(z) \in 1 + z\mathcal{O}_K[[z]]$ because, by (i) of Theorem 3.1, for all $1 \leq i \leq m$, $f_i(z) \in 1 + z\mathcal{O}_K[[z]]$. Further, by (iii) of Theorem 3.1, we know that, for all $1 \leq i \leq m$, $f'_i(z)/f_i(z) \in E_{0,K}$ and thus,

$$\frac{h'(z)}{h(z)} = \sum_{i=1}^{m} a_i \frac{f'_i(z)}{f_i(z)} \in E_{0,K}.$$

Further, h'(z)/h(z) is bounded because, by Theorem 3.1, for all $1 \leq i \leq m$ $f_i(z) \in 1 + z\mathcal{O}_K[[z]]$. Let $D_{\alpha_1}, \ldots, D_{\alpha_q}$ be the poles of h'(z)/h(z) such that $res\left(\frac{h'(z)}{h(z)}, D_{\alpha_i}\right) \neq 0$ for all $1 \leq i \leq q$. Notice D_0 is not a pole of h'(z)/h(z) because $h'(z)/h(z) \in E_{0,K}$. Then, by Equation (6.1), we conclude that, for all $1 \leq i \leq q$, $|\alpha_i| = 1$.

Given that $h(z) \in E_{0,K}$, from Theorem 6.16 we deduce that, there are β_1, \ldots, β_q such that $\beta_i \in D_{\alpha_i}$ and

$$h(z) = \prod_{i=1}^{q} (z - \beta_i)^{\mu_i} \mathfrak{t}(z) \text{ with } \mu_i = res\left(\frac{h'}{h}, D_{\alpha_i}\right) \in \mathbb{Z},$$
(6.7)

 $\mathfrak{t}(z) = \prod_{i=0}^{\infty} \left(\frac{z-a_n}{z-b_n}\right) \text{ with } (a_n, b_n)_{n \ge 0} \text{ a strongly copiercing sequence associated to } D_0. \text{ In addition, by Remark 6.15, we know that } \mathfrak{t}(z) \text{ belongs to } E_{0,p}, |\mathfrak{t}(z) - 1|_{\mathcal{G}} < 1, \text{ and } |\mathfrak{t}(0)| = 1.$

Let us show that, for all $1 \leq i \leq m$, $D_{\alpha_{c_i}}$ belongs to $\mathbf{Pol}(h'(z)/h(z))$ and that $res\left(\frac{h'(z)}{h(z)}, D_{\alpha_{c_i}}\right) = a_i \cdot res\left(\frac{f'_i(z)}{f_i(z)}, D_{\alpha_{c_i}}\right)$. By assumption, we know that, for every $1 \leq i \leq m$, $D_{\alpha_{c_i}} \in \mathbf{Pol}(f'_i(z)/f_i(z))$ and that $D_{\alpha_{c_i}} \notin \mathbf{Pol}(f'_j(z)/f_j(z))$ for every $j \neq i$. Thus, for every $1 \leq i \leq m$, $D_{\alpha_{c_i}}$ is a pole of h'(z)/h(z) and $res\left(\frac{h'(z)}{h(z)}, D_{\alpha_{c_i}}\right) = a_i \cdot res\left(\frac{f'_i(z)}{f_i(z)}, D_{\alpha_{c_i}}\right)$.

By assumption, we know that, for all $i \in \{1, \ldots, m\}$, $res\left(\frac{f'_i(z)}{f_i(z)}, D_{\alpha_{c_i}}\right) \neq 0$. Further, we know that $a_j \neq 0$ for some $j \in \{1, \ldots, m\}$. So, $res\left(\frac{h'(z)}{h(z)}, D_{\alpha_{c_j}}\right) \neq 0$. Thus, we can then suppose without loss of generality that $res\left(\frac{h'(z)}{h(z)}, D_{\alpha_{c_i}}\right) \neq 0$ for all $1 \leq i \leq m$. We can also suppose that $D_{\alpha_1} = D_{\alpha_{c,1}}, \ldots, D_{\alpha_m} = D_{\alpha_{c,m}}$.

Theorem 6.16 aslo implies that $\frac{h'(z)}{h(z)} - \sum_{i=1}^{q} \frac{\mu_i}{z-\beta_i}$ is integrable in $E_{0,p}$. Consequently, according to [19, Lemma 59.2], $\frac{h'(z)}{h(z)} - \sum_{i=1}^{m} \frac{\mu_i}{z-\alpha_{c_i}} - \sum_{i=m+1}^{q} \frac{\mu_i}{z-\beta_i}$ is also integrable in $E_{0,p}$ and thus, according to

Theorem 59.11 of [19], there exits an integer $e \ge 0$ such that

$$\frac{h'(z)}{h(z)} - \sum_{i=1}^{m} \frac{\mu_i}{z - \alpha_{c_i}} - \sum_{i=m+1}^{q} \frac{\mu_i}{z - \beta_i} = \frac{1}{p^e} \left(\sum_{i=1}^{\infty} \frac{1}{z - c_n} - \frac{1}{z - d_n} \right),\tag{6.8}$$

where $(c_n, d_n)_{n \ge 1}$ is a strongly copiercing sequence associated to D_0 . So, by Remark 6.15, we have $\mathfrak{o}(z) = \prod_{n=1}^{\infty} \left(\frac{z-c_n}{z-d_n}\right) \in E_{0,p}$ and $|\mathfrak{o}(0)| = 1$.

Consequently, Equation (6.8) implies that there exists $\lambda \in \mathbb{C}_p$ such that

$$h(z) = \lambda \mathfrak{q}(z)\mathfrak{a}(z)\mathfrak{j}(z) \text{ with } \mathfrak{q}(z) = \prod_{i=1}^{m} (z - \alpha_{c_i})^{\mu_i}, \ \mathfrak{a}(z) = \prod_{i=m+1}^{q} (z - \beta_i)^{\mu_i} \text{ and } \mathfrak{j}(z) = \mathfrak{o}(z)^{1/p^e}.$$
(6.9)

We proceed to show that $|\lambda| = 1$. We know that, for all $1 \leq i \leq q$, $|\alpha_i| = 1$ and given that $D_{\alpha_1} = D_{\alpha_{c,1}}, \ldots, D_{\alpha_m} = D_{\alpha_{c,m}}$ and $\beta_{m+1} \in D_{\alpha_m}, \ldots, \beta_q \in D_{\alpha_q}$, Remark 6.1 implies that, for all $1 \leq i \leq m$, $|\alpha_{c_i}| = 1$ and that, for all $m+1 \leq i \leq q$, $|\beta_i| = 1$. Thus, $|\mathfrak{q}(0)| = 1$ and $|\mathfrak{a}(0)| = 1$. As $h(z) \in 1 + z\mathcal{O}_K[[z]]$ then Equation (6.9) gives $1 = \lambda \mathfrak{q}(0)\mathfrak{a}(0)\mathfrak{o}(0)^{1/p^e}$ and $|\mathfrak{o}(0)^{1/p^e}| = 1$ because $|\mathfrak{o}(0)| = 1$. As a consequence, we then have $|\lambda| = 1$.

We put $s = \min\{v_p(a_1), \ldots, v_p(a_m)\}$. Let L_1 be the field $K(\zeta_1, \ldots, \zeta_{p^s}, \lambda, \alpha_{c_1}, \ldots, \alpha_{c_m}, \beta_1, \ldots, \beta_q)$, where $\zeta_1, \ldots, \zeta_{p^s}$ are the p^s -th roots of unity. We proceed to show that j(z) belongs to $\mathcal{O}_{L_1}[[z]]$ and that it is a unit of this ring. Indeed, we know that $h(z) \in 1 + z\mathcal{O}_K[[z]]$ and it is clear that $\lambda \mathfrak{q}(z)\mathfrak{a}(z) \in L_1(z)$. However, we know that $|\alpha_{c_i}| = 1$ for any $1 \leq i \leq m$ and $|\beta_i| = 1$ for any $m + 1 \leq i \leq q$, and that $|\lambda| = 1$. Thus, we get that $|\lambda \mathfrak{q}(z)\mathfrak{a}(z)|_{\mathcal{G}} = 1$ and that $\lambda \mathfrak{q}(z)\mathfrak{a}(z) \in E(D_0)$. Hence, by Remark 5.6, $\mathfrak{q}(z)\mathfrak{a}(z) \in \mathcal{O}_{L_1}[[z]]$ and since $|\lambda \mathfrak{q}(0)\mathfrak{a}(0)| = |\lambda| |\prod_{i=1}^m (-\alpha_{c_i})^{\mu_i}| |\prod_{i=m+1}^q (-\beta_i)^{\mu_i}| = 1$, we get that $\lambda \mathfrak{q}(z)\mathfrak{a}(z)$ is a unit of $\mathcal{O}_{L_1}[[z]]$. As consequence, it follows from (6.9) that j(z) belongs to $\mathcal{O}_{L_1}[[z]]$ and it is a unit because |j(0)| = 1.

We now show that $\lambda \mathbf{j}(z) = 1 \mod \pi_{L_1} \mathcal{O}_{L_1}[[z]]$. We prove, by using Equation (6.7), in a similar way as above that $\mathfrak{t}(z)$ belongs to $\mathcal{O}_{L_1}[[z]]$. Since $|\mathfrak{t}(z) - 1|_{\mathcal{G}} < 1$, we have $\mathfrak{t}(z) = 1 \mod \pi_{L_1} \mathcal{O}_{L_1}[[z]]$. So, by reducing Equation (6.7) modulo π_{L_1} we get $h(z) = \prod_{i=1}^q (z - \beta_i)^{\mu_i} \mod \pi_{L_1} \mathcal{O}_{L_1}[[z]]$. In addition, for all $1 \leq i \leq m, |\beta_i - \alpha_{c_i}| < 1$ and thus, $\prod_{i=1}^q (z - \beta_i)^{\mu_i} = \prod_{i=1}^m (z - \alpha_{c_i})^{\mu_i} \prod_{i=m+1}^q (z - \beta_i)^{\mu_i} \mod \pi_{L_1} \mathcal{O}_{L_1}[[z]]$. So, by reducing Equation (6.9) modulo π_{L_1} we conclude that $\lambda \mathbf{j}(z) = 1 \mod \pi_{L_1} \mathcal{O}_{L_1}[[z]]$.

We now proceed to prove the statements (i)-(iv).

(i) For every $i \in \{1, \ldots, m\}$, we put $d_i = \frac{a_i}{p^s}$. Then $d_i \in \mathbb{Z}$ for all $i \in \{1, \ldots, m\}$ because, by definition, $s = \min\{v_p(a_1), \ldots, v_p(a_m)\}$. So, Equation (6.9) gives us

$$h_1(z) = \zeta \mathfrak{q}_1(z) \mathfrak{a}_1(z) \mathfrak{j}_1(z) \text{ with } \mathfrak{q}_1(z) = \prod_{i=1}^m (z - \alpha_{c_i})^{\Gamma_i}, \ \mathfrak{a}_1(z) = \prod_{i=m+1}^q (z - \beta_i)^{\Gamma_i} \text{ and } \mathfrak{j}_1(z) = (\lambda \mathfrak{j}(z))^{1/p^s},$$
(6.10)

where, $h_1(z) = f_1(z)^{d_1} \cdots f_m(z)^{d_r}$, $\Gamma_i = \frac{\mu_i}{p^s}$ for every $1 \le i \le q$ and ζ is a p^s -th root of unity.

Notice that, for all $1 \leq i \leq q$,

$$\Gamma_i = \sum_{j=1}^m \frac{a_j}{p^s} \cdot res\left(\frac{f_i'(z)}{f_i(z)}, D_{\alpha_i}\right) = \sum_{j=1}^m d_j \cdot res\left(\frac{f_j'(z)}{f_j(z)}, D_{\alpha_i}\right) = res\left(\frac{h_1'(z)}{h_1(z)}, D_{\alpha_i}\right).$$

By assumption, we know that, for all $1 \leq i \leq m$, $D_{\alpha_{c_i}} \in \mathbf{Pol}(f'_i(z)/f_i(z))$ and that $D_{\alpha_{c_i}} \notin \mathbf{Pol}(f'_j(z)/f_j(z))$ for all $j \in \{1, \ldots, m\} \setminus \{i\}$. Since, for all $1 \leq i \leq m$, $D_{\alpha_i} = D_{\alpha_{c_i}}$, we get that,

for all $1 \leq i \leq m$,

$$\Gamma_i = d_i \cdot res\left(\frac{f_i'(z)}{f_i(z)}, D_{\alpha_{c_i}}\right).$$

We now prove that, for any $1 \leq i \leq q$, Γ_i belongs to \mathbb{Z} . By Theorem 3.1, we know that $f_1(z), \ldots, f_m(z)$ belong to $1 + z\mathcal{O}_K[[z]]$ and, by Theorem 3.1 again, we also know that $f'_1(z)/f_1(z), \ldots, f'_m(z)/f_m(z)$ belong to $E_{0,K}$. So, we are able to apply Lemma 6.7 and thus, we conclude that $res\left(\frac{f'_i(z)}{f_j(z)}, D_{\alpha_i}\right)$ belongs to \mathbb{Z}_p for all $(i, j) \in \{1, \ldots, q\} \times \{1, \ldots, m\}$. Then $\Gamma_i \in \mathbb{Z}_p$ because $d_j \in \mathbb{Z}$ for all $1 \leq j \leq m$. As $\mu_i/p^s = \Gamma_i$ and $\mu_i \in \mathbb{Z}$, and $\Gamma_i \in \mathbb{Z}_p$ then p^s divides μ_i and thus, we get that $\Gamma_i \in \mathbb{Z}$.

Therefore $\mathfrak{q}_1(z)$ and $\mathfrak{a}_1(z)$ belong to $L_1(z)$. Now, we prove that $\mathfrak{q}_1(z)$ belongs to $\mathcal{O}_{L_1}[[z]]$ and that it is a unit of this ring. Indeed, we have already seen that $|\alpha_{c_i}| = 1$ for any $1 \leq i \leq m$ and therefore, $|\mathfrak{q}_1(z)|_{\mathcal{G}} = 1 = |\mathfrak{q}_1(0)|$. Further, we have $\mathfrak{q}_1(z) \in E(D_0)$ because $|\alpha_{c_i}| = 1$ for all $1 \leq i \leq m$. So, Remark 5.6 implies that $\mathfrak{q}_1(z) \in \mathcal{O}_{L_1}[[z]]$ and given that $|\mathfrak{q}_1(0)| = 1$ then $\mathfrak{q}_1(z)$ is a unit of $\mathcal{O}_{L_1}[[z]]$. In a similar fashion, we prove that $\mathfrak{a}_1(z)$ belongs to $\mathcal{O}_{L_1}[[z]]$ and that it is a unit of this ring. We also have $h_1(z) \in 1 + z\mathcal{O}_{L_1}[[z]]$ because $f_1(z), \ldots, f_m(z) \in 1 + z\mathcal{O}_K[[z]]$ and for all $1 \leq i \leq m, d_i \in \mathbb{Z}$ and hence, for every $1 \leq i \leq m, f_i(z)^{d_i} \in 1 + z\mathcal{O}_K[[z]]$. Therefore, from Equation (6.10), we obtain

$$\mathfrak{j}_1(z) = \frac{h_1(z)}{\zeta \mathfrak{q}_1(z)\mathfrak{a}_1(z)} \in \mathcal{O}_{L_1}[[z]] \text{ and } |\mathfrak{j}_1(z)|_{\mathcal{G}} = 1 = |\mathfrak{j}_1(0)|.$$

Whence, $\mathfrak{j}_1(z)$ is a unit of $\mathcal{O}_{L_1}[[z]]$.

(ii) It is clear, by construction, that p does not divide d_s .

(iii) Notice that $|\mathfrak{a}_1(z)|_{\mathcal{G}} = 1$ because for all $m + 1 \leq i \leq q$, $|\beta_i| = 1$. Since the poles and zeros of $\mathfrak{a}_1(z)$ are in $\{\beta_{m+1}, \ldots, \beta_q\}$, we get that all the poles and zeros of $\mathfrak{a}_1(z)$ have norm 1. Recall that we have assumed that, for all $1 \leq i \leq m$, $D_{\alpha_i} = D_{\alpha_{c_i}}$. Further, for all $1 \leq i, j \leq q$, $D_{\alpha_i} \neq D_{\alpha_j}$ for $j \neq i$ because $D_{\alpha_1}, \ldots, D_{\alpha_q}$ are the poles of h'(z)/h(z). In particular, Remark 6.1 implies that, for all $1 \leq i \leq m$, $\beta_{m+1}, \ldots, \beta_q$ do not belong to $D_{\alpha_{c_i}}$. So, for every $1 \leq i \leq m$, $\mathfrak{a}_1(z) \in E(D_{\alpha_{c_i}})$ and $\mathfrak{a}_1(\alpha_{c_i}) \neq 0$.

(iv) We have already seen that $|\mathfrak{j}_1(0)| = 1$. We now prove that $\mathfrak{j}_1(z) = \mathfrak{j}_1(0) \mod \pi_{L_1}\mathcal{O}_{L_1}[[z]]$. Let us write $\mathfrak{j}_1(z) = \sum_{j \ge 0} \kappa_j z^j$. It is clear that $\lambda \mathfrak{j}(z) = \mathfrak{j}_1(z)^{p^s}$ and that $\mathfrak{j}_1(z)^{p^s} = \sum_{j \ge 0} \kappa_j^{p^s} z^{jp^s} \mod \pi_{L_1}\mathcal{O}_{L_1}[[z]]$. But we have already proven that $\lambda \mathfrak{j}(z) = 1 \mod \pi_{L_1}\mathcal{O}_{L_1}[[z]]$ and thus $\kappa_j \in \pi_{L_1}\mathcal{O}_{L_1}$ for all $j \ge 1$.

Finally, we proceed to prove that $j'_1(z) \in \pi_L^2 \mathcal{O}_L[[z]]$, where L is a finite extension of L_1 . Since $j_1(z) = j_1(0) \mod \pi_{L_1} \mathcal{O}_{L_1}[[z]]$, by reducing Equation (6.10) modulo π_{L_1} , we have

$$h_1(z) = \zeta \mathfrak{j}_1(0)\mathfrak{q}_1(z)\mathfrak{a}_1(z) \mod \pi_{L_1}\mathcal{O}_{L_1}[[z]].$$

Notice that $j_1(0)$ is a unit of \mathcal{O}_{L_1} because $|j_1(0)| = 1$. In addition, we have already seen that $\mathfrak{q}_1(z)$ and $\mathfrak{a}_1(z)$ are units of $\mathcal{O}_{L_1}[[z]]$ but $h_1(z)$ is also a unit of this ring because $h_1(z) \in 1 + z\mathcal{O}_K[[z]]$. So, from the equality $h_1(z) = \zeta j_1(0)\mathfrak{q}_1(z)\mathfrak{a}_1(z) \mod \pi_{L_1}\mathcal{O}_{L_1}[[z]]$, we obtain

$$\frac{h_1'(z)}{h_1(z)} = \frac{\mathfrak{q}_1'(z)}{\mathfrak{q}_1(z)} + \frac{\mathfrak{a}_1'(z)}{\mathfrak{a}_1(z)} \mod \pi_{L_1} \mathcal{O}_{L_1}[[z]]$$
(6.11)

Let us show that $\frac{j'_1(z)}{j_1(z)}$ belongs to $E_{0,p}$. From Equation (6.10), we deduce that

$$\frac{\mathfrak{j}_1'(z)}{\mathfrak{j}_1(z)} = \frac{h_1'(z)}{h_1(z)} - \frac{\mathfrak{q}_1'(z)}{\mathfrak{q}_1(z)} - \frac{\mathfrak{a}_1'(z)}{\mathfrak{a}_1(z)}.$$

Given that $\frac{h'_1(z)}{h_1(z)} = \sum_{i=1}^m d_i \frac{f'_i(z)}{f_i(z)}$ and $\frac{f'_i(z)}{f_i(z)} \in E_{0,K}$ for all $1 \leq i \leq m$, we have $\frac{h'_1(z)}{h_1(z)} \in E_{0,K}$. Further, $\frac{\mathfrak{q}'_1(z)}{\mathfrak{q}_1(z)} = \sum_{i=1}^m \frac{\Gamma_i}{z - \alpha_{c_i}}$ and since $|\alpha_{c_i}| = 1$ for all $1 \leq i \leq m$, we get $\frac{\mathfrak{q}'_1(z)}{\mathfrak{q}_1(z)} \in E_{0,p}$. In a similar way, we show that $\frac{\mathfrak{a}'_1(z)}{\mathfrak{a}_1(z)} \in E_{0,p}$. Whence, $\mathfrak{j}'_1(z)/\mathfrak{j}_1(z)$ belongs to $E_{0,p}$.

Now, as $\mathfrak{j}_1(z) \in \mathcal{O}_{L_1}[[z]]$ and $\mathfrak{j}'_1(z)/\mathfrak{j}_1(z) \in E_{0,p}$, it follows from Theorem 3.1 of [24] that there exits $R_2(z) \in \mathbb{C}_p(z)$ such that $R_2(z)$ has no poles or zeros in D_0 and

$$\left|\frac{j_1'(z)}{j_1(z)} - \frac{R_2'(z)}{R_2(z)}\right|_{\mathcal{G}} < 1/p^2.$$
(6.12)

In consequence

$$\frac{h_1'(z)}{h_1(z)} = \frac{\mathfrak{q}_1'(z)}{\mathfrak{q}_1(z)} + \frac{\mathfrak{a}_1'(z)}{\mathfrak{a}_1(z)} + \frac{R_2'(z)}{R_2(z)} + g_2$$
(6.13)

with $g_2 \in E_{0,p}$ and $|g_2|_{\mathcal{G}} < 1/p^2$.

Let L be the extension of L_1 obtained by adding the zeros and poles of $R_2(z)$. Let us show that $R'_2(z)/R_2(z)$ belongs to $\mathcal{O}_L[[z]]$. It is clear that $R'_2(z)/R_2(z) \in L(z)$. Further, $j'_1(z)/j_1(z) \in \mathcal{O}_L[[z]]$ because we know that $j_1(z)$ is a unit of the ring $\mathcal{O}_{L_1}[[z]]$. So, we deduce, from Equation (6.12), that

$$\left|\frac{R_{2}'(z)}{R_{2}(z)}\right|_{\mathcal{G}} = \left|\frac{R_{2}'(z)}{R_{2}(z)} - \frac{j_{1}'(z)}{j_{1}(z)} + \frac{j_{1}'(z)}{j_{1}(z)}\right|_{\mathcal{G}} \le \max\left\{\left|\frac{j_{1}'(z)}{j_{1}(z)} - \frac{R_{2}'(z)}{R_{2}(z)}\right|_{\mathcal{G}}, \left|\frac{j_{1}'(z)}{j_{1}(z)}\right|_{\mathcal{G}}\right\} \le 1.$$

Further, $R'_2(z)/R_2(z)$ belongs to $E(D_0)$ because D_0 is not a pole of $h'_1(z)/h_1(z)$ (recall that $h'_1(z)/h_1(z) \in E_{0,K}$). Then, by Remark 5.6, $R'_2(z)/R_2(z)$ belongs $\mathcal{O}_L[[z]]$.

Consequently, it follows from Equation (6.13) that $g_2 \in \mathcal{O}_L[[z]]$ and since $|g_2|_{\mathcal{G}} < 1/p^2$, $g_2 \in \pi_L^2 \mathcal{O}_L[[z]]$. So, reducing Equation (6.13) modulo π_L^2 , we have

$$\frac{h_1'(z)}{h_1(z)} = \frac{\mathfrak{q}_1'(z)}{\mathfrak{q}_1(z)} + \frac{\mathfrak{a}_1'(z)}{\mathfrak{a}_1(z)} + \frac{R_2'(z)}{R_2(z)} \mod \pi_L^2 \mathcal{O}_L[[z]].$$
(6.14)

From Equation (6.11), we know that $\frac{h'_1(z)}{h_1(z)} = \frac{\mathfrak{q}'_1(z)}{\mathfrak{q}_1(z)} + \frac{\mathfrak{a}'_1(z)}{\mathfrak{a}_1(z)} \mod \pi_L \mathcal{O}_L[[z]]$ and therefore, from Equation (6.14), we have $\frac{R'_2(z)}{R_2(z)} \in \pi_L \mathcal{O}_L[[z]]$.

Let us write

$$\frac{R_2'(z)}{R_2(z)} = \sum_{\tau \in S_1} \frac{c_\tau}{z - \tau},$$

where S_1 is the set of poles and zero of $R_2(z)$ and $c_{\tau} \in \mathbb{Z} \setminus \{0\}$.

So, for all $\tau \in S_1$, $D_{\tau} \in \{D_{\alpha_1}, \ldots, D_{\alpha_q}\}$. Given that, $\frac{\mathfrak{q}'_1(z)}{\mathfrak{q}_1(z)} = \sum_{i=1}^m \frac{\Gamma_i}{z - \alpha_{c_i}}$ and $\frac{\mathfrak{a}'_1(z)}{\mathfrak{a}_1(z)} = \sum_{i=m+1}^q \frac{\Gamma_i}{z - \beta_i}$ then, by using (ii) of Lemma 6.7, we deduce from Equation (6.14) that, for every $1 \leq i \leq q$,

$$res\left(\frac{h_1'(z)}{h_1(z)}, D_{\alpha_i}\right) = \left(\Gamma_i + \sum_{\tau \in S_1 \cap D_{\alpha_i}} c_{\tau}\right) \mod p^2.$$
(6.15)

But, $\Gamma_i = res\left(\frac{h_1'(z)}{h_1(z)}, D_{\alpha_i}\right)$. Whence,

$$\sum_{\tau \in S_1 \cap D_{\alpha_i}} c_\tau = 0 \bmod p^2 \mathbb{Z}.$$
(6.16)

Further, if $\tau \in D_{\alpha_i} \cap S_1$, by Lemma 6.3, we conclude that

$$\frac{1}{z-\tau} = \frac{1}{z-\alpha_i} \sum_{l \ge 0} \left(\frac{\tau-\alpha_i}{z-\alpha_i}\right)^l$$

Since $|\alpha_i - \tau| < 1$, we have $\alpha_i - \tau \in \pi_L \mathcal{O}_L$. So, from the previous equality, we get

$$\frac{1}{z-\tau} = \frac{1}{z-\alpha_i} + \frac{\tau-\alpha_i}{(z-\alpha_i)^2} \mod \pi_L^2 \mathcal{O}_L[[z]].$$

But $c_{\tau} \in \pi_L \mathcal{O}_L$ because $\frac{R'_2(z)}{R_2(z)} \in \pi_L \mathcal{O}_L[[z]]$. So, from the previous equality, we get

$$\frac{c_{\tau}}{z-\tau} = \frac{c_{\tau}}{z-\alpha_i} \mod \pi_L^2 \mathcal{O}_L[[z]].$$
(6.17)

Therefore, combining Equations (6.14), (6.16), and (6.17), we deduce that

$$\frac{h_1'(z)}{h_1(z)} = \sum_{i=0}^m \frac{\Gamma_i + \sum_{\tau \in S \cap D_{\alpha_i}} c_\tau}{z - \alpha_{c_i}} + \sum_{i=m+1}^q \frac{\Gamma_i + \sum_{\tau \in S \cap D_{\alpha_i}} c_\tau}{z - \beta_i} = \sum_{i=0}^m \frac{\Gamma_i}{z - \alpha_{c_i}} + \sum_{i=m+1}^q \frac{\Gamma_i}{z - \beta_i} \mod \pi_L^2 \mathcal{O}_L[[z]]$$

It follows from Equation (6.10) that

$$\frac{h_1'(z)}{h_1(z)} = \sum_{i=0}^m \frac{\Gamma_i}{z - \alpha_{c_i}} + \sum_{i=m+1}^q \frac{\Gamma_i}{z - \beta_i} + \frac{\mathfrak{j}_1'(z)}{\mathfrak{j}_1(z)}$$

So, from the previous two equalities we get

$$\frac{\mathfrak{j}_1'(z)}{\mathfrak{j}_1(z)} \in \pi_L^2 \mathcal{O}_L[[z]].$$

We have already seen that $\mathfrak{j}_1(z)$ is a unit of $\mathcal{O}_L[[z]]$ and therefore, $\mathfrak{j}'_1(z) \in \pi_L^2 \mathcal{O}_L[[z]]$.

(v) Since, for all $1 \leq i \leq r$, $p^s d_i = a_i$, we have $(f_1^{d_1}(z) \cdots f_r^{d_r}(z))^{p^s} = f_1^{a_1}(z) \cdots f_r^{a_r}(z)$. But, we know that $f_1^{a_1}(z) \cdots f_r^{a_r}(z) \in E_{0,K}$.

6.3. Proof of Theorems 2.6 and 2.7

As illustration of Theorem 6.12, we prove Theorems 2.6 and 2.7. The proof of both theorems runs along the same lines, we only need to verify that the assumptions of Theorem 6.12 are satisfied.

Proof of Theorem 2.6. — The power series $\mathfrak{h}(z)$, $\mathfrak{f}(z)$ and $\mathfrak{g}(z)$ are respectively solution of the following hypergeometric operators

$$\mathcal{L} := \delta^3 - z(\delta + 1/2)^3, \quad \delta^3 - z(\delta - 1/2)(\delta + 1/2)^2, \quad \mathcal{D} := \delta^2 - z(\delta + 1/2)^2.$$

It is clear that these differential operators are MOM at zero and, it follows from [28, Theorem 6.2] that they have strong Frobenius structure for p = 3. For this reason $\mathfrak{h}(z)$, $\mathfrak{g}(z)$, $\mathfrak{g}(z)$ belong to $\mathcal{MF}(\mathbb{Q}_3)$. Recall that \mathbb{Q}_3 is a Frobenius field. In addition, since K is an extension of \mathbb{Q}_3 , we have $\mathcal{MF}(\mathbb{Q}_3) \subset \mathcal{MF}(K)$. So $\mathfrak{h}(z)$, $\mathfrak{f}(z)$, $\mathfrak{g}(z)$ belong to $\mathcal{MF}(K)$.

(i). We first show that $\mathfrak{f}(z)/\mathfrak{h}(z)$ belongs to $E_{\mathbb{Q}_3}$. By [28, Theorem 6.2] again, we know that $\delta^3 - z(\delta + 1/2)^3$ has a strong Frobenius structure for p = 3 of period 1. Thus, according to (ii) of Theorem 3.1, we get that $\mathfrak{h}(z)/\mathfrak{h}(z^3) \in E_{\mathbb{Q}_3}$. But, according to Corollary 2 of Section 1 of [14] and Theorem 2 of [14], $\mathfrak{f}(z)/\mathfrak{h}(z^3) \in E_{\mathbb{Q}_3}$. Given that $E_{\mathbb{Q}_3}$ is a field we obtain $\mathfrak{f}(z)/\mathfrak{h}(z) \in E_{\mathbb{Q}_3}$.

Finally, as $\mathfrak{h}(z)$ and $\mathfrak{f}(z)$ belong to $\mathcal{MF}(\mathbb{Q}_3)$ and \mathbb{Q}_3 is a Frobenius field then, from Lemma 3.3, we deduce that $\mathfrak{f}^{(s)}(z)/\mathfrak{f}(z)$ and $\mathfrak{h}(z)/\mathfrak{h}^{(r)}(z)$ belong to $E_{\mathbb{Q}_3}$ for all integers $r, s \ge 0$. Consequently,

$$\frac{\mathfrak{f}^{(s)}(z)}{\mathfrak{h}^{(r)}(z)} = \frac{\mathfrak{f}^{(s)}(z)}{\mathfrak{f}(z)} \cdot \frac{\mathfrak{f}(z)}{\mathfrak{h}(z)} \cdot \frac{\mathfrak{h}(z)}{\mathfrak{h}^{(r)}(z)}$$

belongs to $E_{\mathbb{Q}_3}$ for all integers $r, s \ge 0$.

Given that K is an extension of \mathbb{Q}_3 , we get $E_{\mathbb{Q}_3} \subset E_K$ and thus, for all integers $r, s \ge 0$, $\frac{\mathfrak{f}^{(s)}(z)}{\mathfrak{h}^{(r)}(z)}$ belongs to E_K .

(ii). In order to prove that, for all integers $r, s \ge 0$, $\mathfrak{h}^{(r)}(z)$ and $\mathfrak{g}^{(s)}(z)$ are algebraically independent over E_K , it sufficient, according to Theorem 2.5, to show that $\mathfrak{h}(z)$ and $\mathfrak{g}(z)$ are algebraically independent over E_K . Our strategy is to suppose that $\mathfrak{h}(z)$ and $\mathfrak{g}(z)$ are algebraically dependent over E_K and then, by applying Theorem 6.12, we get a contradiction. For this purpose, we split our argument into several steps.

First Step We show that $\mathfrak{h}'(z)/\mathfrak{h}(z) \in E(\mathcal{O}_{\mathbb{C}_3} \setminus D_{\alpha_1})$ and that $\mathfrak{g}'(z)/\mathfrak{g} \in E(\mathcal{O}_{\mathbb{C}_3} \setminus D_{\alpha_2})$ with $\alpha_1 = 1$ and $\alpha_2 = -1$.

It follows from Corollary 1 of Section 1 of [14] and Theorem 2 of [14] that, for all integers $r \ge 1$,

$$\frac{\mathfrak{h}(z)}{\mathfrak{h}(z^3)} = \frac{\mathfrak{h}_r(z)}{\mathfrak{h}_{r-1}(z^3)} \mod 3^r \mathbb{Z}[[z]], \tag{6.18}$$

where, for $r \ge 0$, $\mathfrak{h}_r(z)$ is the $3^r - 1$ -th truncation of $\mathfrak{h}(z)$. Thus, according to Lemma 2 of [2], $\mathfrak{h}(z)/\mathfrak{h}(z^3)$ extends to $\Omega = \{x \in \mathcal{O}_{\mathbb{C}_3} : |\mathfrak{h}_1(x)| = 1\}$. As $\mathfrak{h}_1(z) = 1 + \frac{1}{8}z + \frac{27}{512}z^2$ then, it is not hard to see that, $|\mathfrak{h}_1(x)| < 1$ if and only if $x \in D_{\alpha_1}$. So, $\Omega = \mathcal{O}_{\mathbb{C}_3} \setminus D_{\alpha_1}$. We put $H(z) = \mathfrak{h}(z)/\mathfrak{h}(z^3)$. Then $H \in E(\Omega)$. Moreover, from (iii) of Lemma 2 of [2], we also have |H(x)| = 1 for all $x \in \Omega$. Hence, H is a unit of the ring $E(\Omega)$. It is not hard to see that

$$H'(z) = H(z) \left[\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)} - 3z^2 \frac{\mathfrak{h}'(z^3)}{\mathfrak{h}(z^3)} \right].$$

Therefore, if for every $m \ge 0$, we consider $H_m = H(z)H(z^p)\cdots H(z^{p^m})$ then $H_m \in E(\Omega)$ and $|H_m(x)| = 1$ for all $x \in \Omega$. So H_m is a unit of $E(\Omega)$ and we also have the equality

$$H'_{m}(z) = H_{m}(z) \left[\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)} - 3^{m+1} z^{3^{m+1}-1} \frac{\mathfrak{h}'(z^{3^{m+1}})}{\mathfrak{h}(z^{3^{m+1}})} \right]$$

Therefore, for all integers $m \ge 0$,

$$\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)} \equiv \frac{H'_m(z)}{H_m(z)} \bmod 3^{m+1}$$

Since, for all $m \ge 0$, H_m is a unit of $E(\Omega)$, from the previous congruences, we conclude that $\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)}$ belongs to $E(\Omega)$.

We have a similar situation for \mathfrak{g} . It follows from Corollary 1 of Section 1 of [14] and Theorem 2 of [14] that, for all integers $r \ge 1$,

$$\frac{\mathfrak{g}(z)}{\mathfrak{g}(z^3)} = \frac{\mathfrak{g}_r(z)}{\mathfrak{g}_{r-1}(z^3)} \mod 3^r \mathbb{Z}[[z]], \tag{6.19}$$

where, for $r \ge 0$, $\mathfrak{g}_r(z)$ is the $3^r - 1$ -th truncation of $\mathfrak{h}(z)$. Thus, according to Lemma 2 of [2], $\mathfrak{g}(z)/\mathfrak{g}(z^3)$ extends to $\Delta = \{x \in \mathcal{O}_{\mathbb{C}_3} : |\mathfrak{g}_1(x)| = 1\}$. As $\mathfrak{g}_1(z) = 1 + \frac{1}{4}z + \frac{9}{64}z^2$ then, it is not hard to see that,

 $|\mathfrak{g}_1(x)| < 1$ if and only if $x \in D_{\alpha_2}$. So, $\Delta = \mathcal{O}_{\mathbb{C}_3} \setminus D_{\alpha_2}$. We put $G(z) = \mathfrak{g}(z)/\mathfrak{g}(z^3)$. Then $G \in E(\Delta)$. Moreover, from (iii) of Lemma 2 of [2], we also have |G(x)| = 1 for all $x \in \Delta$. Hence, G is a unit of the ring $E(\Delta)$. So, $G_m = G(z)G(z^p)\cdots G(z^{p^m})$ is a unit of $G_m \in E(\Delta)$ and $|G_m(x)| = 1$ for all $x \in \Delta$. By proceeding as before, we conclude that, for all integers $m \ge 0$,

$$\frac{\mathfrak{g}'(z)}{\mathfrak{g}(z)} \equiv \frac{G'_m(z)}{G_m(z)} \mod 3^{m+1}$$

Since, for all $m \ge 0$, G_m is a unit of $E(\Delta)$, from the previous congruences, we conclude that $\frac{\mathfrak{g}'(z)}{\mathfrak{g}(z)}$ belongs to $E(\Delta)$.

Second Step We show that D_{α_1} is a pole of o of $\mathfrak{h}'/\mathfrak{h}$ and that D_{α_2} is a pole of $\mathfrak{g}'/\mathfrak{g}$. It follows from Equations (6.18) and (6.19) that

$$\mathfrak{h}(z) = \mathfrak{h}_1(z)\mathfrak{h}(z^3) \mod 3, \quad \mathfrak{g}(z) = \mathfrak{g}_1(z)\mathfrak{g}(z^3).$$

For this reason

$$\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)} = \frac{1}{z-1} \mod 3, \quad \frac{\mathfrak{g}'(z)}{\mathfrak{g}(z)} = \frac{1}{1+z} \mod 3.$$
(6.20)

So D_{α_1} is a pole of $\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)}$ and D_{α_2} is a pole of $\frac{\mathfrak{g}'(z)}{\mathfrak{g}(z)}$.

We now put

$$\lambda_1 = res\left(\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)}, D_{\alpha_1}\right) \text{ and } \lambda_2 = res\left(\frac{\mathfrak{g}'(z)}{\mathfrak{g}(z)}, D_{\alpha_2}\right).$$

Third Step We now show that $\lambda_1 \mod 9 = 4 = \lambda_2 \mod 9$.

We deduce from Equation (6.18) that

$$\frac{\mathfrak{h}(z)}{\mathfrak{h}(z^9)} = \frac{\mathfrak{h}_2(z)}{\mathfrak{h}_1(z^3)} \frac{\mathfrak{h}_2(z^3)}{\mathfrak{h}_1(z^9)} \bmod 9.$$

Therefore,

$$\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)} = \frac{8+8z+5z^2}{1-z^3} = \frac{2z+1}{z^2+z+1} - \frac{7}{z-1} \mod 9$$

Let $z_0, z_1 \in \overline{\mathbb{Q}_3}$ be the roots of $1 + z + z^2$. Then $|z_0| = 1 = |z_1|, z_0 \neq z_1$, and $z_0, z_1 \in D_{\alpha_1}$ because $1 + z + z^2 = (1 - z)^2 \mod 3$. Furthermore, it is clear that

$$\frac{2z+1}{z^2+z+1} - \frac{7}{z-1} = \frac{1}{z-z_0} + \frac{1}{z-z_1} - \frac{7}{z-1}.$$

Since $1, z_0, z_1 \in D_{\alpha_1}$, it follows from (1) of Remark 6.4 and (ii) of Lemma 6.7 that

$$\lambda_1 \mod 9 = (1+1-7) \mod 9 = -5 \mod 9 = 4.$$

Now, from Equation (6.19), we deduce that

$$\frac{\mathfrak{g}(z)}{\mathfrak{g}(z^9)} = \frac{\mathfrak{g}_2(z)}{\mathfrak{g}_1(z^3)} \frac{\mathfrak{g}_2(z^3)}{\mathfrak{g}_1(z^9)} \bmod 9.$$

Therefore,

$$\frac{\mathfrak{g}'(z)}{\mathfrak{g}(z)} = \frac{-7 - 5z + 3z^2 + 4z^3 + 8x^4 - 7z^6 - 5z^7}{(z-1)(z+1)(z^6 - z^3 + 1)} \mod 9$$

Note that

$$\frac{-7-5z+3z^2+4z^3+8x^4-7z^6-5z^7}{(z-1)(z+1)(z^6-z^3+1)} = \frac{3-3z^4}{z^6-z^3+1} - \frac{1}{2(z+1)} \mod 9.$$

Let s be the set of the roots of $z^6 - z^3 + 1$. Since $z^6 - z^3 + 1 = (z+1)^3 \mod 3$, we have $s \subset D_{\alpha_2}$. Therefore, by (1) of Remark 6.4 and (ii) of Lemma 6.7, we deduce that

$$\lambda_2 \mod 9 = \left(\sum_{\tau \in s} res\left(\frac{3-3z^4}{z^6-z^3+1}, \tau\right) - \frac{1}{2}\right) \mod 9.$$

But $\sum_{\tau \in S} res\left(\frac{3-3z^4}{z^6-z^3+1}, \tau\right) = 0$ and consequently $\lambda_2 = 4 \mod 9$.

Fourth Step. We now proceed to see that the conditions (1)-(2) of Theorem 6.12 are fulfilled. In First Step we proved that $\mathfrak{h}'(z)/\mathfrak{h}(z) \in E(\mathcal{O}_{\mathbb{C}_3} \setminus D_{\alpha_1})$ with $\alpha_1 = 1$ and that $\mathfrak{g}'(z)/\mathfrak{g}(z) \in E(\mathcal{O}_{\mathbb{C}_3} \setminus D_{\alpha_2})$ with $\alpha_2 = -1$. We also proved in Second Step that D_{α_1} is a pole of $\mathfrak{h}'(z)/\mathfrak{h}(z)$ and that D_{α_2} is a pole of $\mathfrak{g}'(z)/\mathfrak{g}(z)$. Thus, $\operatorname{Pol}(\mathfrak{h}'(z)/\mathfrak{h}(z)) = \{D_{\alpha_1}\}$ and $\operatorname{Pol}(\mathfrak{g}'(z)/\mathfrak{g}(z)) = \{D_{\alpha_2}\}$. In addition, notice that $D_{\alpha_1} \neq D_{\alpha_2}$ because $|\alpha_1 - \alpha_2| = 1$.

Finally, we have $\mathcal{L}(\mathfrak{h}) = 0$ and $\mathcal{D}(\mathfrak{g}) = 0$. Let us write \mathcal{L} and \mathcal{D} in terms of d/dz. Then,

$$\mathcal{L}_1 = \frac{d}{dz^3} + \left(\frac{3}{z} - \frac{3}{2(1-z)}\right) \frac{d}{dz^2} + \left(\frac{1}{z^2} - \frac{3}{2z(1-z)} - \frac{1}{z(1-z)}\right) \frac{d}{dz} - \frac{1}{8z^2(z-1)}$$
$$\mathcal{D}_1 = \frac{d}{dz^2} + \frac{1-2z}{z(z-1)} \frac{d}{dz} - \frac{1}{4z(z-1)}$$

So, it is clear that α_1 is a regular singular point of $\mathcal{L}_{1|\mathcal{M}^2}$ and α_2 is also a regular singular point of $\mathcal{D}_{1|\mathcal{M}^2}$. The indicial polynomial of $\mathcal{L}_{1|\mathcal{M}^2}$ at α_1 is the $P = X(X-1)(X-2) + \frac{3}{2}X(X-1) \in \frac{\mathbb{Z}}{9\mathbb{Z}}[X]$ and the indicial polynomial of $\mathcal{D}_{1|\mathcal{M}^2}$ at α_2 is $Q = X(X-1)(X-2) \in \frac{\mathbb{Z}}{9\mathbb{Z}}[X]$.

So, the conditions (1)-(2) of Theorem 6.12 are satisfied and, in oder to apply it, we suppose, towards a contradiction, that $\mathfrak{h}(z)$ and $\mathfrak{g}(z)$ are algebraically dependent over E_K and thus we have two cases.

case A: $res\left(\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)}, D_{\alpha_1}\right)$ belongs to \mathbb{Z}_3 and its reduction modulo p^2 is an exponent of $\mathcal{L}_{1|\mathcal{M}^2}$ at α_1 . Or

case B: es $\left(\frac{\mathfrak{g}'(z)}{\mathfrak{g}(z)}, D_{\alpha_2}\right)$ belongs to \mathbb{Z}_3 and its reduction modulo p^2 is an exponent of $\mathcal{D}_{1|\mathcal{M}^2}$ at α_2 .

We are going to see that case A and B lead to a contradiction. According to Third Step,

$$res\left(\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)}, D_{\alpha_1}\right) \mod 9 = \lambda_1 \mod 9 = 4.$$

So case A implies that 4 is a solution of P. That is a contradiction because P(4) = 6. We now consider case B. According to Third Step,

$$res\left(\frac{\mathfrak{g}'(z)}{\mathfrak{g}(z)}, D_{\alpha_2}\right) \mod 9 = \lambda_2 \mod 9 = 4.$$

Therefore, 4 is a solution of Q. Again, that is a contradiction because Q(4) = 6.

(iii). Suppose that $\mathfrak{f}(z)$ and $\mathfrak{g}(z)$ are algebraically dependent over E_K . Then, by Theorem 2.3, there are integers a, b not all zero such that $\mathfrak{f}^a(z)\mathfrak{g}^b(z) \in E_K$. According to (i), $\mathfrak{f}/\mathfrak{h} \in E_K$. Thus,

$$\mathfrak{h}^{a}(z)\mathfrak{g}^{b}(z) = rac{\mathfrak{h}^{a}(z)}{\mathfrak{f}^{a}(z)}\mathfrak{f}^{a}(z)\mathfrak{g}^{b}(z) \in E_{K}.$$

Whence, $\mathfrak{g}(z)$ and $\mathfrak{h}(z)$ are algebraically dependent over E_K . That is a contradiction because, according to (ii), $\mathfrak{g}(z)$ and $\mathfrak{h}(z)$ are algebraically independent over E_K . Therefore, $\mathfrak{f}(z)$ and $\mathfrak{g}(z)$ are algebraically

independent over E_K . Consequently, Theorem 2.5 implies that, for all integers $r, s \ge 0$, $\mathfrak{f}^{(r)}(z)$ and $\mathfrak{g}^{(s)}(z)$ are algebraically independent over E_K .

Proof of Theorem 2.7. — As we have already observed at the very beginning of the proof of Theorem 2.6 that $\mathfrak{h}(z)$ belongs to $\mathcal{MF}(K)$. We now prove that $\mathfrak{A}(z) \in \mathcal{MF}(K)$. Since $\mathcal{MF}(\mathbb{Q}_3) \subset \mathcal{MF}(K)$, it is sufficient to prove that $\mathfrak{A}(z) \in \mathcal{MF}(\mathbb{Q}_3)$. Indeed, by [2, Theorem 1], we get that $\mathfrak{A}(z)/\mathfrak{A}(z^3)$ belongs to E_{0,\mathbb{Q}_3} . Consequently, from Remark 5.7, we obtain $\mathfrak{A}'(z)/\mathfrak{A}(z)$ belongs to E_{0,\mathbb{Q}_3} . Further, it is clear that $\mathfrak{A}(z)$ is solution of the differential operator $\delta - \frac{\delta \mathfrak{A}(z)}{\mathfrak{A}(z)}$. Notice that this differential operator is MOM at zero. In addition, this differential operator belongs to $E_{\mathbb{Q}_3}[\delta]$ because $\mathfrak{A}'(z)/\mathfrak{A}(z) \in E_{\mathbb{Q}_3}$ and has a Frobenius structure because $\mathfrak{A}(z)/\mathfrak{A}(z^3) \in E_{\mathbb{Q}_3}$. Finally $\left|\frac{\delta \mathfrak{A}(z)}{\mathfrak{A}(z)}\right|_{\mathcal{G}} \leq 1$ because $\mathfrak{A}(z) \in 1 + z\mathbb{Z}[[z]]$. So, we conclude that $\mathfrak{A}(z) \in \mathcal{MF}(\mathbb{Q}_3)$. Further, we also know that $\mathfrak{A}(z)$ is solution of the differential operator

$$\mathcal{H} = \frac{d}{dz^3} + \frac{3 - 153z + 6z^2}{(1 - 34z + z^2)z} \frac{d}{dz^2} + \frac{1 - 112z + 7z^2}{(1 - 34z + z^2)z^2} \frac{d}{dz} + \frac{z - 5}{(1 - 34z + z^2)z^2}.$$

(i) According to Theorem 2.5, in oder to prove that, for all integers $r, s \ge 0$, $\mathfrak{h}^{(r)}(z)$ and $\mathfrak{A}^{(s)}(z)$ are algebraically independent over E_K , it is sufficient to prove that $\mathfrak{h}(z)$ and $\mathfrak{A}(z)$ are algebraically independent over E_K . For this purpose, we suppose that $\mathfrak{h}(z)$ and $\mathfrak{A}(z)$ are algebraically dependent over E_K and then, by applying Theorem 6.12, we get a contradiction. For this purpose, we split our argument into several steps.

First Step We show that $\mathfrak{A}'(z)/\mathfrak{A}(z) \in E(\mathcal{O}_{\mathbb{C}_3} \setminus D_{\alpha_2})$ with $\alpha_2 = -1$. In fact, it follows from [2, Theorem 1] that, for all integers $r \ge 1$,

$$\frac{\mathfrak{A}(z)}{\mathfrak{A}(z^3)} = \frac{\mathfrak{A}_r(z)}{\mathfrak{A}_{r-1}(z^3)} \mod 3^r \mathbb{Z}[[z]], \tag{6.21}$$

where for all $r \ge 0$, $\mathfrak{A}_r(z)$ is the $3^r - 1$ -th truncation of $\mathfrak{A}(z)$. Thus, according to Lemma 2 of [2], $\mathfrak{A}(z)/\mathfrak{A}(z^3)$ extends to $\Lambda = \{x \in \mathcal{O}_{\mathbb{C}_3} : |\mathfrak{A}_1(x)| = 1\}$. As $\mathfrak{A}_1(z) = 1 + 5z + 73z^2$ then, it is not hard to see that, $|\mathfrak{A}_1(x)| < 1$ if and only if $x \in D_{\alpha_2}$. So, $\Lambda = \mathcal{O}_{\mathbb{C}_p} \setminus D_{\alpha_2}$. So, proceeding in a similar fashion as in the First Step of the proof of Theorem 2.6, we conclude that $\mathfrak{A}'(z)/\mathfrak{A}(z)$ belongs to $E(\Lambda)$.

Second Step We now show that D_{α_1} , $\alpha_1 = 1$, is a pole of $\mathfrak{h}'/\mathfrak{h}$ and that D_{α_2} is a pole of $\mathfrak{A}'/\mathfrak{A}$ with $\alpha_2 = -1$. From Second Step of the proof of Theorem 2.6, we know that D_{α_1} is a pole of $\mathfrak{h}'(z)/\mathfrak{h}(z)$. Now, we deduce from Equation (6.21) that

$$\frac{\mathfrak{A}(z)}{\mathfrak{A}(z^9)} = \frac{\mathfrak{A}_2(z)}{\mathfrak{A}_1(z^3)} \frac{\mathfrak{A}_2(z^3)}{\mathfrak{A}_1(z^9)} \bmod 9\mathbb{Z}[[z]]$$

Therefore,

$$\frac{\mathfrak{A}'(z)}{\mathfrak{A}(z)} = \frac{5+4z+8z^2+4z^3+5z^4+z^5}{1-z^6} \mod 9\mathbb{Z}.$$

Note that

$$\frac{5+4z+8z^2+4z^3+5z^4+z^5}{1-z^6} = \frac{2z-1}{z^2-z+1} + \frac{3}{2(z+1)} \mod 9\mathbb{Z}.$$
(6.22)

Let s be the roots of $z^2 - z + 1$ in \mathbb{C}_3 . We have $s \subset D_{\alpha_2}$ because $(z^2 - z + 1) = (z + 1)^2 \mod 3$. So, D_{α_2} is a pole of $\mathfrak{A}'(z)/\mathfrak{A}(z)$.

We now put

$$\epsilon_1 = res\left(\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)}, D_{\alpha_1}\right) \text{ and } \epsilon_2 = res\left(\frac{\mathfrak{A}'(z)}{\mathfrak{A}(z)}, D_{\alpha_2}\right).$$

Third Step We now prove that $\epsilon_1 \mod 9 = 4$ and that $\epsilon_2 \mod 9 = 8$. According to Third Step of the proof of Theorem 2.6, $\epsilon_1 \mod 9 = 4$.

Now, regarding ϵ_2 , we have $\sum_{\tau \in s} \left(\frac{2z-1}{z^2-z+1}, \tau\right) = 2$. Thus, by (1) of Remark 6.4 and (ii) of Lemma 6.7, we deduce from Equation (6.22) that

$$\epsilon_2 \mod 9 = (2 + 3/2) \mod 9 = 8.$$

Fourth Step We now proceed to see that the conditions (1)-(2) of Theorem 6.12 are fulfilled. It follows from First Step and Second Step that $\mathbf{Pol}(\mathfrak{h}'(z)/\mathfrak{h}(z)) = \{D_{\alpha_1}\}$ and $\mathbf{Pol}(\mathfrak{A}'(z)/\mathfrak{A}(z)) = \{D_{\alpha_2}\}$. In addition, $D_{\alpha_1} \neq D_{\alpha_2}$ because $|\alpha_1 - \alpha_2| = 1$.

Further, we have seen in the proof of Theorem 2.6 that $\mathcal{L}_1(\mathfrak{h}) = 0$, α_1 is regular singular point of $\mathcal{L}_{1|\mathcal{M}^2}$ and that the indicial polynomial of $\mathcal{L}_{1|\mathcal{M}^2}$ at α_1 is $P = X(X-1)(X-2) + \frac{3}{2}X(X-1) \in \frac{\mathbb{Z}}{9\mathbb{Z}}[X]$. Now $\mathcal{H}(\mathfrak{A}) = 0$. We are going to see that α_2 is a regular singular point of $\mathcal{H}_{|\mathcal{M}^2}$. We first observe that $(1 - 34z + z^2) = (1 + z)^2 \mod 9\mathbb{Z}[[z]]$ and that $3 - 153z + 6z^2 = (1 + z)(3 + 6z) \mod 9\mathbb{Z}[[z]]$. Whence,

$$(z+1)\frac{3-153z+6z^2}{z(1-34z+z^2)} = \frac{3+6z}{z} \mod \mathcal{M}^2 \mathcal{O}_{\mathbb{C}_p(z)},$$
$$(1+z)^2 \frac{1-112z+7z^2}{z^2(1-34z+z^2)} = \frac{1+5z+7z^2}{z^2} \mod \mathcal{M}^2 \mathcal{O}_{\mathbb{C}_p(z)},$$
$$(z+1)^3 \frac{z-5}{z^2(1-34z+z^2)} = \frac{(z+1)(z-5)}{z^2} \mod \mathcal{M}^2 \mathcal{O}_{\mathbb{C}_p(z)}.$$

Further, it is clear that $\frac{3+6z}{z}$, $\frac{1+5z+7z^2}{z^2}$ and that $\frac{(z+1)(z-5)}{z^2}$ belong to $\mathbb{Z}_3[[z+1]]$. So α_2 is a regular singular point of $\mathcal{H}_{|\mathcal{M}^2}$. In addition, the indicial polynomial of $\mathcal{H}_{|\mathcal{M}^2}$ at α_2 is

$$R = X(X-1)(X-2) + 3X(X-1) + 3X = X(X^2+2) \in \frac{\mathbb{Z}}{9\mathbb{Z}}[[z]].$$

So, the conditions (1)-(2) of Theorem 6.12 are satisfied and, in oder to apply it, we suppose, towards a contradiction, that $\mathfrak{h}(z)$ and $\mathfrak{A}(z)$ are algebraically dependent over E_K and thus we have two cases

case A: $res\left(\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)}, D_{\alpha_1}\right)$ belongs to \mathbb{Z}_3 and its reduction modulo p^2 is an exponent of $\mathcal{L}_{1|\mathcal{M}^2}$ at α_1 . Or

case B: $res\left(\frac{\mathfrak{A}'(z)}{\mathfrak{A}(z)}, D_{\alpha_2}\right)$ belongs to \mathbb{Z}_3 and its reduction modulo p^2 is an exponent of $\mathcal{H}_{1|\mathcal{M}^2}$ at α_2 . Finally, let us see that cases A and B lead to a contradiction. From Third Step, we have

$$res\left(\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)}, D_{\alpha_1}\right) \mod 9 = \lambda_1 \mod 9 = 4.$$

So case A implies that 4 is a solution of P. That is a contradiction because P(4) = 6. Again, by Third Step,

$$res\left(rac{\mathfrak{A}'(z)}{\mathfrak{A}(z)}, D_{\beta_1}
ight) \mod 9 = \lambda_2 \mod 9 = 8.$$

Therefore, 8 is a solution of R. Again, that is a contradiction because R(8) = 6.

(ii) Suppose that $\mathfrak{f}(z)$ and $\mathfrak{A}(z)$ are algebraically dependent over E_K . Then, there are integers a, b not all zero such that $\mathfrak{f}^a(z)\mathfrak{A}^b(z) \in E_K$. According to (i) of Theorem 2.6, $\mathfrak{f}/\mathfrak{h} \in E_K$. Thus,

$$\mathfrak{h}^{a}(z)\mathfrak{A}^{b}(z) = \frac{\mathfrak{h}^{a}(z)}{\mathfrak{f}^{a}(z)}\mathfrak{f}^{a}(z)\mathfrak{A}^{b}(z) \in E_{K}.$$

Whence, $\mathfrak{h}(z)$ and $\mathfrak{A}(z)$ are algebraically dependent over E_K . That contradicts (ii). Therefore, $\mathfrak{f}(z)$ and $\mathfrak{A}(z)$ are algebraically independent over E_K . Consequently, Theorem 2.5 implies that, for all integers $r, s \ge 0$, $\mathfrak{f}^{(r)}(z)$ and $\mathfrak{A}^{(s)}(z)$ are algebraically independent over E_K .

6.4. Proof of Theorem 2.9

In order to prove Theorem 2.9, we need the following lemma.

Lemma 6.18. — Let p be an odd prime, π_p be in $\overline{\mathbb{Q}_p}$ such that $\pi_p^{p-1} = -p$ and let N > 0 be an integer.

(i) For an integer $n \ge 1$, let $n = n_0 + n_1 p + \dots + n_l p^l$ be its p-adic expansion and let $\Sigma_n = n_0 + n_1 + \dots + n_l$. Then

$$\frac{\pi_p^{2n}}{4^n (n!)^2} \bigg|_p = (1/p)^{\frac{2\Sigma_n}{p-1}}.$$
(6.23)

(ii) For every $1 < s \leq N$,

$$\left| \binom{N}{s} f_1(z)^s \right|_{\mathcal{G}} < |Nf_1(z)|_{\mathcal{G}}, \tag{6.24}$$

where
$$f_1(z) = \sum_{n \ge 1} \frac{\pi_p^{2n}}{4^n (n!)^2} z^{2n}$$

Proof. —

(i) It is well-known that, for every integer $n \ge 1$, $|n!|_p = (1/p)^{(n-\Sigma_n)/(p-1)}$. Whence, we immediately obtain the equality (6.23) for every integer $n \ge 1$.

(ii) It follows from (i) that $|f_1(z)|_{\mathcal{G}} = (1/p)^{2/p-1}$. Thus, the inequality (6.24) is equivalent to proving that, for every $1 < s \leq N$,

$$\left|\binom{N}{s}\right|_{p} < \left|N\right|_{p} p^{\frac{2(s-1)}{p-1}}.$$
(6.25)

Let us suppose that p does not divide N. Then $|N|_p = 1$ and since s > 1, we have $p^{2(s-1)/(p-1)} > 1$. Further, the p-adic norm of $\binom{N}{s}$ is less than or equal to 1 given that $\binom{N}{s}$ is an integer. Hence, the inequality (6.25) holds.

Let us now suppose that p divides N and let r be the p-adic valuation of N. Then, we have $|N|_p p^{2(s-1)/(p-1)} = p^{\frac{2(s-1)}{p-1}-r}$. If $\frac{2(s-1)}{p-1} > r$ then it is clear that the inequality (6.25) is true. Now, we suppose that $\frac{1}{p-1} \leq \frac{2(s-1)}{p-1} \leq r$. We first observe that $(N-1)(N-2)(N-3)(N-4)\cdots(N-s+1) = (-1)^{s-1}(s-1)! + Nt$ for some integer t > 0. Whence,

$$(N-1)(N-2)(N-3)(N-4)\cdots(N-s+1) = (-1)^{s-1}p^{\frac{s-1-\Sigma_{s-1}}{p-1}}a + p^rbt,$$

where a, b are integers not divisible by p. Since $\frac{1}{p-1} \leq \frac{2(s-1)}{p-1} \leq r$ and s > 1, we have $\frac{1}{p-1} \leq \frac{s-1}{p-1} < \frac{2(s-1)}{p-1} \leq r$. In addition, it is clear that $s - 1 - \sum_{s-1} < s - 1$ because s > 1. So $\frac{s-1-\sum_{s-1}}{p-1} < \frac{s-1}{p-1} < r$. Thus

$$|(N-1)(N-2)(N-3)(N-4)\cdots(N-s+1)|_p = (1/p)^{\frac{s-1-\Sigma_{s-1}}{p-1}}.$$

Consequently,

$$|N(N-1)(N-2)\cdots(N-s+1)|_p = (1/p)^{r+\frac{s-1-\Sigma_{s-1}}{p-1}}.$$

So, by using the fact that $v_p(s!) = (s - \Sigma_s)/(p-1)$, we get

$$\left|\binom{N}{s}\right|_{p} = \left(\frac{1}{p}\right)^{r + \frac{\Sigma_{s} - \Sigma_{s-1} - 1}{p-1}}$$

It is not hard to see that, for every $s \ge 2$, $s - \Sigma_{s-1} > 1$ and that $\Sigma_s \ge 1$. Thus,

$$\frac{\sum_{s} - \sum_{s-1} - 1}{p-1} + \frac{s-1}{p-1} = \frac{s - \sum_{s-1} - 1 + \sum_{s} - 1}{p-1} > 0.$$

So

$$\frac{\sum_s - \sum_{s-1} - 1}{p-1} > -\frac{s-1}{p-1} > \frac{-2(s-1)}{p-1}$$

Whence,

$$r + \frac{\sum_s - \sum_{s-1} - 1}{p - 1} > r - \frac{2(s - 1)}{p - 1}$$

Notice that $0 \leq r - 2(s-1)/(p-1)$ because we have assumed that $2(s-1)/(p-1) \leq r$. Then

$$\left|\binom{N}{s}\right|_{p} = (1/p)^{r + \frac{\Sigma_{s} - \Sigma_{s-1} - 1}{p-1}} \leq (1/p)^{r - \frac{2(s-1)}{p-1}} = |N|_{p} p^{\frac{2(s-1)}{p-1}}$$

That completes the proof of the inequality (6.25).

Proof of Theorem 2.9. —

(i) We are going to see that $\exp(\pi_p z)^p$ belongs to E_p . We have

$$\exp(\pi_p z)^p = \exp(p\pi_p z) = \sum_{n \ge 0} \frac{p^n \pi_p^n}{n!} z^n.$$

It follows from (i) of Lemma 6.18 that

$$\left|\frac{p^n \pi_p^n}{n!}\right|_p = (1/p)^{n + \frac{\Sigma_n}{p-1}}.$$

So $\lim_{n\to\infty} \left| \frac{p^n \pi_p^n}{n!} \right|_p = 0$. Consequently, $\exp(\pi_p z)^p$ belongs to E_K .

(ii) We put $\mathfrak{B}(z) = J_0(\pi_p z)$. We first show that $\mathfrak{B}(z)$ is transcendental over E_p . For this purpose, aiming for a contradiction, suppose that $\mathfrak{B}(z)$ is algebraic over E_p . Given that $J_0(z)$ is solution of $\delta^2 + z^2$ then $\mathfrak{B}(z)$ is solution of $\delta^2 - \pi_p^2 z^2$. It is clear that this differential operator is MOM at zero and given that by assumption $p \neq 2$ then, according to Dwork [17, §4], this differential operator has a strong Frobenius structure. Hence, $\mathfrak{B}(z) \in \mathcal{MF}(K)$ and, by Theorem 3.1, $\mathfrak{B}'(z)/\mathfrak{B}(z)$ belongs to

_	
Г	٦
	1

 $E_{0,K}$. Since $E_{0,K} \subset E_p$ and $\mathfrak{B}(z)$ is algebraic over E_p , if follows from Theorem 3.2 that there exists an integer N > 0 such that $\mathfrak{B}(z)^N \in E_p$. It is clear that

$$\mathfrak{B}(z) = \sum_{n \ge 0} \frac{(-1)^n \pi_p^{2n}}{4^n (n!)^2} z^{2n}.$$

We set $f_1(z) = \sum_{n \ge 1} \frac{(-1)^n \pi_p^{2n}}{4^n (n!)^2} z^{2n}$. Then $\mathfrak{B}(z)^N = (1 + f_1(z))^N = \sum_{s=0}^N {N \choose s} f_1(z)^s$.

According to (ii) of Lemma 6.18 we know that, for every $1 < s \leq N$,

$$\left|\binom{N}{s}f_1(z)^s\right|_{\mathcal{G}} < |Nf_1(z)|_{\mathcal{G}}.$$

From (i) of Lemma 6.18, we deduce that $|f_1(z)|_{\mathcal{G}} = (1/p)^{2/p-1}$. Thus, for all $1 < s \leq N$,

$$\left| \binom{N}{s} f_1(z)^s \right|_{\mathcal{G}} < (1/p)^{v_p(N) + \frac{2}{p-1}}.$$
(6.26)

By definition, π_p is solution of the Eisenstein polynomial $X^{p-1} + p$ and thus, π_p is a uniformizer of $\mathbb{Q}_p(\pi_p)$. Given that, for any $s \ge 1$, $\binom{N}{s} f_1(z)^s \in \mathbb{Q}_p(\pi_p)[[z]]$ so Inequality (6.26) implies that, for every $1 < s \le N$, $\binom{N}{s} f_1(z)^s \in (\pi_p)^{r(p-1)+3} \mathcal{O}_{\mathbb{Q}_p(\pi_p)}[[z]]$. Consequently, for every $1 < s \le N$,

$$\binom{N}{s} f_1(z)^s = 0 \mod \pi_p^{(p-1)v_p(N)+3} \mathcal{O}_{\mathbb{Q}_p(\pi_p)}[[z]].$$

Therefore, we obtain

$$\mathfrak{B}(z)^N = 1 + N f_1(z) \mod \pi_p^{(p-1)v_p(N)+3} \mathcal{O}_{\mathbb{Q}_p(\pi_p)}[[z]].$$
(6.27)

Again, from (i) of Lemma 6.18, we conclude that, for all $n \ge 0$,

$$\left| \frac{\pi_p^{2p^n}}{4^{p^n} (p^n)!^2} \right|_p = (1/p)^{2/p-1}.$$
(6.28)

Consequently, for all $n \ge 0$, $N \frac{\pi_p^{2p^n}}{4^{p^n}(p^n)!^2} \notin \pi_p^{(p-1)v_p(N)+3} \mathcal{O}_{\mathbb{Q}_p(\pi_p)}$.

Now, it is clear that if $m \notin \{p^n\}_{n \ge 0}$ then $\Sigma_m > 1$ and thus, by using (i) of Lemma 6.18 again, we obtain

$$\left| \frac{\pi_p^{2m}}{4^m(m)!^2} \right|_p = (1/p)^{\frac{2\Sigma_m}{p-1}} < (1/p)^{\frac{2}{p-1}}.$$

In particular, if $m \notin \{p^n\}_{n \ge 0}$ then $N \frac{\pi_p^{2m}}{4^m (m)!^2} \in \pi_p^{(p-1)v_p(N)+3} \mathcal{O}_{\mathbb{Q}_p(\pi_p)}$. Hence, we get

$$1 + Nf_1(z) = 1 - N \sum_{n \ge 1} \frac{\pi_p^{2p^n}}{4^{p^n}(p^n)!^2} z^{2p^n} \mod \pi_p^{(p-1)v_p(N)+3} \mathcal{O}_{\mathbb{Q}_p(\pi_p)}[[z]].$$
(6.29)

Since $\mathfrak{B}(z)^N$ belongs to E_p and $\mathfrak{B}(z)^N \in \mathbb{Q}_p(\pi_p)[[z]]$ there are a rational function $R(z) \in \mathbb{Q}_p(\pi_p)(z)$ such that

$$\mathfrak{B}(z)^N = R(z) \mod \pi_p^{(p-1)v_p(N)+3} \mathcal{O}_{\mathbb{Q}_p(\pi_p)}[[z]].$$

Let us write $R(z) = \sum_{n \ge -M} a_n z^n$ with $M \in \mathbb{N}$. Then, it follows from the previous equality and Equations (6.27) and (6.29) that

$$\sum_{n \ge -M} a_n z^n = 1 - N \sum_{n \ge 1} \frac{\pi_p^{2p^n}}{4^{p^n} (p^n)!^2} z^{2p^n} \mod \pi_p^{(p-1)v_p(N)+3} \mathcal{O}_{\mathbb{Q}_p(\pi_p)}[[z]].$$
(6.30)

DANIEL VARGAS-MONTOYA

On the one hand, since R(z) is a rational function, the sequence $\{a_n\}_{n \ge 0}$ is almost periodic. That is, for any $\epsilon > 0$, there are an integer n_{ϵ} such that, for all $n \ge n_{\epsilon}$, $|a_{n+n_{\epsilon}} - a_n| < \epsilon$. On the other hand, it follows from Equation (6.30) that, for all $m \notin \{2p^k : k \ge 1\}$, $a_m \in \pi_p^{(p-1)v_p(N)+3}\mathcal{O}_{\mathbb{Q}_p(\pi_p)}$ and thus, $|a_m| \le (1/p)^{v_p(N)+\frac{3}{p-1}}$. Now, if we take $\epsilon' = (1/p)^{v_p(N)+\frac{3}{p-1}}$ there an integer $n_{\epsilon'}$ such that, for all $n \ge n_{\epsilon'}$, $|a_{n+n_{\epsilon'}} - a_n| < \epsilon'$. In addition, it is clear that exists $n_0 \in \{2p^k : k \ge 1\}$ such that $n_0 \ge n_{\epsilon'}$ and $n_0 + n_{\epsilon'} \notin \{2p^k : k \ge 1\}$. If $n_0 = 2p^{k_0}$, it follows from (6.30) that

$$a_{n_0} = \lambda \mod \pi_p^{(p-1)v_p(N)+3} \mathcal{O}_{\mathbb{Q}_p(\pi_p)} \text{ with } \lambda = -N \frac{\pi_p^{2p^{k_0}}}{4p^{k_0}(p^{k_0})!^2}.$$

Thus $|a_{n_0} - \lambda| \leq \epsilon'$. Further, from Equation (6.28), we have $|\lambda| = (1/p)^{v_p(N) + \frac{2}{p-1}} > \epsilon'$. Since $|a_{n_0}| = |a_{n_0} - \lambda + \lambda|$ and $|a_{n_0} - \lambda| \leq \epsilon' < |\lambda|$, and since moreover the norm is non-Archimedean, we get that $|a_{n_0} - \lambda + \lambda| = |\lambda|$ and hence, $|a_{n_0}| = |\lambda|$. Given that $n_0 + n_{\epsilon'} \notin \{2p^k : k \geq 1\}$ we know that $|a_{n_0+n_{\epsilon'}}| \leq \epsilon'$. Therefore, $|a_{n_0}| > |a_{n_0+n_{\epsilon'}}|$. Again, given that the norm is non-Archimedean, we get that $|a_{n_0+n_{\epsilon'}} - a_{n_0}| = |\lambda| > \epsilon'$. But, that is a contradiction because $|a_{n_0+n_{\epsilon'}} - a_{n_0}| < \epsilon'$. So $\mathfrak{B}(z)$ is transcendental over E_p and since $E_K \subset E_p$, $\mathfrak{B}(z)$ is transcendental over E_K .

Now, by Theorem 2.5, we get that, for all $s \ge 0$, $\mathfrak{B}^{(s)}(z)$ is transcendental over E_K . Finally, for all $s \ge 0$, $\mathfrak{B}^{(s)}(z)$ is transcendental over $E_K[\exp(\pi_p z)]$ because $\mathfrak{B}^{(s)}(z)$ is transcendental over E_K and, according to (i), $\exp(\pi_p z)$ is algebraic over E_K .

(iii) We put $\mathfrak{B}_1(z) = J_0(2i\sqrt{2\pi_p}z)$ and $\mathfrak{B}_2(z) = \mathfrak{g}(\pi_p z)$. We first show that $\mathfrak{B}_2(z)$ is transcendental over are E_K . It turns out that $\mathfrak{g}(z) = \exp(3z)J_0(2i\sqrt{2}z)$. So $\mathfrak{B}_2(z) = \exp(3\pi_p z)\mathfrak{B}_1(z)$ and

$$\frac{\mathfrak{B}_{2}'(z)}{\mathfrak{B}_{2}(z)} = \frac{3\pi_{p}\exp(3\pi_{p}z)}{\exp(3\pi_{p}z)} + \frac{2i\sqrt{2}\mathfrak{B}'(2i\sqrt{2}z)}{\mathfrak{B}(2i\sqrt{2}z)} = 3\pi_{p} + \frac{2i\sqrt{2}\mathfrak{B}'(2i\sqrt{2}z)}{\mathfrak{B}(2i\sqrt{2}z)} = 3\pi_{p} + \frac{\mathfrak{B}_{1}'(z)}{\mathfrak{B}_{1}(z)}$$

As we have already said at the beginning of (ii), $\mathfrak{B}(z) \in \mathcal{MF}(K)$. Then, according to (iii) of Theorem 3.1, $\mathfrak{B}'(z)/\mathfrak{B}(z) \in E_{0,K}$. In particular, $\mathfrak{B}'(z)/\mathfrak{B}(z) \in E_{0,p}$. Since $p \neq 2$, the *p*-adic norm of $2i\sqrt{2}$ is 1. Thus, $\mathfrak{B}'(2i\sqrt{2}z)/\mathfrak{B}(2i\sqrt{2}z)$ is still an element of $E_{0,p}$. Whence, $\mathfrak{B}'_1(z)/\mathfrak{B}_1(z)$ belongs to $E_{0,p}$ and hence, $\mathfrak{B}'_2(z)/\mathfrak{B}_2(z)$ is also in $E_{0,p}$. Suppose, towards a contradiction, that $\mathfrak{B}_2(z)$ is transcendental over E_K . So, by Theorem 3.2, we deduce that there there is a nonzero integer *a* such that $\mathfrak{B}_2(z)^a \in E_p$. That is, $\exp(3\pi_p z)^a \mathfrak{B}_1(z)^a \in E_p$. According to (i), $\exp(\pi_p z)^p \in E_K$ and since $\exp(3\pi_p z) = \exp(\pi_p z)^3$, we have $\exp(\pi_p z)^{3p} \in E_K$. Consequently, $\mathfrak{B}_1(z)^{pa} \in E_p$. As the *p*-adic norm of $2i\sqrt{2}$ is 1, we get $\mathfrak{B}(z)^{pa} \in E_p$. But, according to (ii), $\mathfrak{B}(z)$ is transcendental over E_p and thus a = 0. Which is a contradiction. So $\mathfrak{B}_2(z)$ is transcendental over E_K . In addition, for all integers $s \ge 0$, $\mathfrak{B}_2^{(s)}(z)/\mathfrak{B}_2(z) \in E_p$ because $\mathfrak{B}'_2(z)/\mathfrak{B}_2(z) \in E_p$. Consequently, Theorem 3.2 and the fact that $\mathfrak{B}_2(z)$ is transcendental over E_p imply that $\mathfrak{B}_2^{(s)}(z)$ is transcendental over E_p . Finally, for all integers $s \ge 0$, $\mathfrak{B}_2^{(s)}(z)$ is transcendental over the ring $E_K[\exp(\pi_p z)]$ because, according to (i), $\exp(\pi_p z)$ is algebraic over E_K and we have already seen that $\mathfrak{B}_2^{(s)}(z)$ is transcendental over E_K .

6.5. Proof of Theorem 2.10

Proof of Theorem 2.10. — Let $\mathfrak{B}(z) = J_0(\pi_3 z)$ and K be an finite extension of $\mathbb{Q}_3(\pi_3)$ such that K is a Frobenius field. At the beginning of the proof of Theorem 2.6 we proved that $\mathfrak{h}(z) \in \mathcal{MF}(K)$ and we have also seen at the beginning of proof of Theorem 2.7 that $\mathfrak{A}(z) \in \mathcal{MF}(K)$. The fact that $\mathfrak{B}(z) \in \mathcal{MF}(K)$ has been shown at the beginning of (ii) of the proof of Theorem 2.9.

(i) We have already seen in the proof of Theorems 2.6 and 2.7 that $\mathfrak{h}'(z)/\mathfrak{h}(z)$ belongs to $E(\mathcal{O}_{\mathbb{C}_3} \setminus D_1)$ and that $\mathfrak{A}'(z)/\mathfrak{A}(z)$ belongs to $E(\mathcal{O}_{\mathbb{C}_3} \setminus D_{-1})$. We have also shown there that D_1 is the only pole of $\mathfrak{h}'(z)/\mathfrak{h}(z)$ and that D_{-1} is the only pole of $\mathfrak{A}'(z)/\mathfrak{A}(z)$.

Notice that π_3 is a uniformizer of $\mathbb{Q}_3(\pi_3)$. As we already said $\mathfrak{B}(z)$ is solution of the MOM differential operator $\mathcal{B} = \delta^2 + \pi_3^2 z$ and, by [17, §4], this differential operator has strong Frobenius structure with period 1. So, according to Theorem 3.1, $\mathfrak{B}(z) \in 1 + \pi_3 \mathcal{O}_{\mathbb{Q}_3(\pi_3)}[[z]], T := \mathfrak{B}(z)/\mathfrak{B}(z^3) \in E_{0,\mathbb{Q}_3(\pi_3)}$ and $\mathfrak{B}'(z)/\mathfrak{B}(z) \in E_{0,\mathbb{Q}_3(\pi_3)}$. We are going to show that if $\alpha \in \mathcal{O}_{\mathbb{C}_3}$ then D_α is not a pole of $\mathfrak{B}'(z)/\mathfrak{B}(z)$. It follows from [17, § 6] that T converges in $D(0, 1^+) = \{x \in \mathbb{C}_3 : |x| \leq 1\}$. Moreover, according to Remark 5.7, we have

$$\frac{\mathfrak{B}'(z)}{\mathfrak{B}(z)} \equiv \frac{T'_m(z)}{T_m(z)} \mod \pi_3^{m+1} \mathcal{O}_{\mathbb{Q}_3(\pi_3)}[[z]] \text{ with } T_m = T(z)T(z^3)\cdots T(z^{3^m}).$$
(6.31)

Notice that $H \in 1 + \pi_3 \mathcal{O}_{\mathbb{Q}_3(\pi_3)}[[z]]$ because $\mathfrak{B}(z) \in 1 + \pi_3 \mathcal{O}_{\mathbb{Q}_3(\pi_3)}[[z]]$. Thus, if $\alpha \in \mathcal{O}_{\mathbb{C}_3}$, $|H(\alpha)| = 1$ and consequently, for all $m \ge 1$, $|H_m(\alpha)| = 1$. So, from Equation (6.31), D_α is not a pole of $\mathfrak{B}'(z)/\mathfrak{B}(z)$.

Aiming for a contradiction, suppose that $\mathfrak{h}(z), \mathfrak{A}(z)$, and $\mathfrak{B}(z)$ are algebraically dependent over E_K . Then, according to Theorem 2.3, there are integers a, b, and c not all zero such that $\mathfrak{h}(z)^a \mathfrak{A}(z)^b \mathfrak{B}(z)^c \in E_{0,K}$ and, further, by Theorem 6.13, we also have

$$\mathfrak{h}(z)^{a}\mathfrak{A}(z)^{b}\mathfrak{B}(z)^{c} = \prod_{i=1}^{2} (z - \beta_{i})^{\mu_{i}} \mathfrak{t}(z), \qquad (6.32)$$

where $\mu_i = res\left(a\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)} + b\frac{\mathfrak{A}'(z)}{\mathfrak{A}(z)} + c\frac{\mathfrak{B}'(z)}{\mathfrak{B}(z)}, D_{\beta_i}\right) \in \mathbb{Z}, \beta_1 \in D_1 \text{ and } \beta_2 \in D_{-1}, \text{ and } \mathfrak{t}(z) = \prod_{n=1}^{\infty} \left(\frac{z-a_n}{z-b_n}\right),$ where $(a_n, b_n)_{n \ge 1}$ is a strongly copiercing sequence associated to D_0 .

Notice that Remark 6.1 implies $D_{\beta_1} = D_1$ and $D_{\beta_2} = D_{-1}$.

As K is a Frobenius then, by Theorem 2.7, the power series $\mathfrak{h}(z)$ and $\mathfrak{A}(z)$ are algebraically independent over E_K and consequently, $c \neq 0$. Now, from Equation (6.32), we deduce that

$$c\frac{\mathfrak{B}'(z)}{\mathfrak{B}(z)} = \sum_{i=1}^{2} \frac{\mu_i}{z - \beta_i} - a\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)} - b\frac{\mathfrak{A}'(z)}{\mathfrak{A}(z)} + \frac{\mathfrak{t}'(z)}{\mathfrak{t}(z)}.$$
(6.33)

Now, we have the following factorization $\mathfrak{t}(z) = \prod_{i=1}^{4} \mathfrak{t}_i(z)$, where for $i \in \{1, 2\}$, $\mathfrak{t}_i(z) = \prod_{b_n \in D_{\beta_i}} \left(\frac{z-a_n}{z-b_n}\right)$, $\mathfrak{t}_3(z) = \prod_{b_n \in \mathcal{O}_{\mathbb{C}_3} \setminus D_{\beta_1} \cup D_{\beta_2}} \left(\frac{z-a_n}{z-b_n}\right)$ and $\mathfrak{t}_4(z) = \prod_{b_n \notin \mathcal{O}_{\mathbb{C}_3}} \left(\frac{z-a_n}{z-b_n}\right)$. As the sequence $(a_n, b_n)_{n \ge 1}$ is a strongly copiercing sequence associated to D_0 then the sequences used to define $\mathfrak{t}_1(z), \ldots, \mathfrak{t}_4(z)$ are also strongly copiercing sequence associated to D_0 . Thus, by Remark 6.15, $\mathfrak{t}_1(z), \ldots, \mathfrak{t}_4(z)$ belong to E_p . Further, it follows from Equation (6.33) that

$$c\frac{\mathfrak{B}'(z)}{\mathfrak{B}(z)} = \frac{\mu_1}{z - \beta_1} - a\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)} + \frac{\mathfrak{t}'_1(z)}{\mathfrak{t}_1(z)} + \frac{\mu_2}{z - \beta_2} - b\frac{\mathfrak{A}'(z)}{\mathfrak{A}(z)} + \frac{\mathfrak{t}'_2(z)}{\mathfrak{t}_2(z)} + \sum_{i=3}^4 \frac{\mathfrak{t}'_i(z)}{\mathfrak{t}_i(z)}.$$
(6.34)

On the one hand, by using Mittag-Leffet's Theorem (see Equation (6.1)), we know that there exits a unique sequence $\{\alpha_n\}_{n\geq 1}$ in $\mathcal{O}_{\mathbb{C}_3}$ such that, for every, $|\alpha_n| = 1$, $|\alpha_n - \alpha_m| = 1$ for every $n \neq m$ and there exists a unique sequence $\{f_n\}_{n\geq 0}$ in $E_{0,K}$ such that $f_0 \in E(\mathcal{O}_{\mathbb{C}_3})$, for n > 0, $f_n \in E^{\alpha_n}$, and

$$c\frac{\mathfrak{B}'(z)}{\mathfrak{B}(z)} = \sum_{n \ge 0} f_n.$$

But, we have already seen that for any $\alpha \in \mathcal{O}_{\mathbb{C}_3}$, D_α is not a pole of $\mathfrak{B}'(z)/\mathfrak{B}(z)$ and thus, $f_n = 0$ for all n > 0.

On the other hand, it is clear D_{β_1} is the only of pole of $\frac{\mu_1}{z-\beta_1} - a\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)} + \frac{\mathfrak{t}'_1(z)}{\mathfrak{t}_1(z)}$ and thus it belongs to E^{β_1} . Similarly, D_{β_2} is the only pole of $\frac{\mu_2}{z-\beta_2} - b\frac{\mathfrak{A}'(z)}{\mathfrak{A}(z)} + \frac{\mathfrak{t}'_2(z)}{\mathfrak{t}_2(z)}$ and thus it belongs to E^{β_2} .

Since $f_n = 0$ for all n > 0 it follows from Equation (6.34) that

$$\frac{\mu_1}{z-\beta_1} - a\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)} + \frac{\mathfrak{t}'_1(z)}{\mathfrak{t}_1(z)} = 0 \text{ and } \frac{\mu_2}{z-\beta_2} - b\frac{\mathfrak{A}'(z)}{\mathfrak{A}(z)} + \frac{\mathfrak{t}'_2(z)}{\mathfrak{t}_2(z)} = 0.$$

Given that $\frac{\mathfrak{t}'_1(z)}{\mathfrak{t}_1(z)} = \sum_{b_n \in D_{\beta_1}} \left(\frac{1}{z-a_n} - \frac{1}{z-b_n} \right)$ and $(a_n, b_n)_{b_n \in D_{\beta_1}}$ is a strongly copiercing sequence associated to D_0 , from Theorem 59.1 of [19], we deduce that $\frac{\mu_1}{z-\beta_1} - a\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)}$ is integrable in $E_{0,p}$. Since D_{β_1} is the only pole of $\mathfrak{h}'(z)/\mathfrak{h}(z)$ and $\mu_1 = res(a\mathfrak{h}'(z)/\mathfrak{h}(z), D_{\beta_1})$, we get, from Lemma 59.2 of [19], that $\frac{\mu_1}{z-1} - a\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)}$ is also integrable in $E_{0,p}$. Further, Theorem 59.11 of [19] implies that there exists an integer $e \ge 0$ such that

$$\frac{\mu_1}{z-1} - a\frac{\mathfrak{h}'(z)}{\mathfrak{h}(z)} = \frac{1}{p^e} \sum_{n \ge 1} \left(\frac{1}{z-c_n} - \frac{1}{z-d_n} \right),$$

where $(c_n, d_n)_{n \ge 1}$ is a strongly copiercing sequence associated to D_0 . So, by Remark 6.15, $\mathfrak{j}(z) = \prod_{n=1}^{\infty} \left(\frac{z-c_n}{z-d_n}\right)$ belongs to $E_{0,p}$. Thus, there exists $\lambda \in \mathbb{C}_3 \setminus \{0\}$ such that $\mathfrak{h}^a(z) = \lambda(z-1)^{\mu_1}\mathfrak{j}(z)^{1/p^e}$. Whence, $\mathfrak{h}^{ap^e}(z) = (z-1)^{\mu_1 p^e}\mathfrak{j}_1(z)$, with $\mathfrak{j}_1(z) = \lambda^{p^e}\mathfrak{j}(z)$. We know that $\mathfrak{h}(z) \in 1 + z\mathbb{Z}_3[[z]]$ and thus $\mathfrak{h}(z)/(z-1) \in -1 + z\mathbb{Z}_3[[z]]$. So, we conclude that $\mathfrak{j}_1(z) \in E_{0,\mathbb{Q}_3}$

Similarly, we can prove that there exists an integer $m \ge 0$ such that $\mathfrak{A}^{bp^m}(z) = (z+1)^{\mu_2 p^m} \mathfrak{j}_2(z)$, with $\mathfrak{j}_2(z) \in E_{0,\mathbb{Q}_3}$.

Thus, $\mathfrak{h}^{ap^e}(z)\mathfrak{A}^{bp^m}(z) \in E_{0,\mathbb{Q}_3}$. But, we know that $\mathfrak{h}(z)$ and $\mathfrak{A}(z)$ are algebraically independent over $E_{\mathbb{Q}_3}$ and therefore, $ap^e = 0 = bp^m$. Whence a = 0 = b. But, we know that $\mathfrak{h}(z)^a\mathfrak{A}(z)^b\mathfrak{B}(z)^c \in E_{0,K}$ and thus $\mathfrak{B}(z)^c \in E_{0,K}$, which is a contradiction because $c \neq 0$ and $\mathfrak{B}(z)$ is transcendental over E_K .

Therefore, $\mathfrak{h}(z)$, $\mathfrak{A}(z)$, and $\mathfrak{B}(z)$ are algebraically independent over E_K . So, from Theorem 2.5 we conclude that, for all integers $r, s, k \ge 0$, $\mathfrak{B}^{(r)}(z)$, the power series $\mathfrak{A}^{(s)}(z)$, and $\mathfrak{h}^{(k)}(z)$ are algebraically independent over E_K .

(ii) Aiming for a contradiction, suppose that $\mathfrak{f}(z), \mathfrak{A}(z)$ and $\mathfrak{B}(z)$ are algebraically dependent over E_K . At the beginning of the proof of Theorem 2.6 we showed that $\mathfrak{f}(z) \in \mathcal{MF}(K)$ and we have already seen that $\mathfrak{A}(z)$ and $\mathfrak{B}(z)$ belong to $\mathcal{MF}(K)$. Thus, according to Theorem 2.3, there are integers a, b, and c not all zero such that $\mathfrak{f}(z)^a \mathfrak{A}(z)^b \mathfrak{B}(z)^c \in E_{0,K}$. But, according to (i) of Theorem 2.6, $\mathfrak{h}/\mathfrak{f}$ belongs to $E_{\mathbb{Q}_3}$. Thus

$$\mathfrak{h}(z)^{a}\mathfrak{A}(z)^{b}\mathfrak{B}(z)^{c} = \left(\frac{\mathfrak{h}(z)}{\mathfrak{f}(z)}\right)^{a}\mathfrak{f}(z)^{a}\mathfrak{A}(z)^{b}\mathfrak{B}(z)^{c} \in E_{K}.$$

That is a contradiction because, according to (i), $\mathfrak{h}(z)$, $\mathfrak{A}(z)$ and $\mathfrak{B}(z)$ are algebraically independent over E_K . Therefore, $\mathfrak{f}(z)$, $\mathfrak{A}(z)$ and $\mathfrak{B}(z)$ are algebraically independent over E_K . Finally, according to Theorem 2.5, for all integers r, s, $k \ge 0$, $\mathfrak{B}^{(r)}(z)$, $\mathfrak{A}^{(s)}(z)$, and $\mathfrak{f}^{(k)}(z)$ are algebraically independent over E_K .

Bibliography

- B. ADAMCZEWSKI, J. P. BELL, AND E. DELAYGUE, Algebraic independence of G-functions and congruences "à la Lucas", Ann. Sci. Éc. Norm. Supér 52 (2019), 515–559.
- M. ANTON AND M. VLASENKO Dwork's congruences for the constant terms of powers of a Laurent polynomial, Int. J. Number Theory 12 (2016), 313–321.
- [3] F. BEUKERS, W.D. BROWNAWELL, G. HECKMAN, Siegel normality, Annals of Math 127 (1988), 279-308.
- [4] F. BEUKERS, AND G. HECKMAN, Monodromy for the hypergeometric function ${}_{n}F_{n-1}$, Invent. Math 95 (1989), 325–354.
- [5] F. BEUKERS AND J. WOLFART, Algebraic values of hypergeometric functions, New advances in transcendence theory (Durham, 1986), 68-81, Cambridge Univ. Press, Cambridge, ; MR0971994.
- [6] A. BOSTAN, P. LAIREZ, B. SALVY, Mutiple binomial sums, J. Symbolic Comput 80 (2017), 351–386.
- [7] G. CHRISTOL, Systèmes différentiels linéaires p-adiques, structures de Frobenius faible, Bull. Soc. Math. France 109 (1981), 83–122.
- [8] G. CHRISTOL, Modules différentiels et équations différentielles p-adiques, Queen's papers in pure and applied mathematics 66, Queen's University, Kingston, 1983.
- [9] G. CHRISTOL, Un théorème de transfert pour les disques singuliers réguliers, Astérisque 119-120 (1984), 151-168.
- [10] G. CHRISTOL, Fonctions et éléments algébriques, Pacific J. Math 125 (1986), 1–37.
- G. CHRISTOL, Diagonales de fractions rationnelles et équations de Picard-Fuchs, Group de travail d'analyse ultramétrique 12 (1984/85), Exp No 13, 12 pp.
- [12] C. CURTIS AND I. REINER, Representation theory of finite groups and associative algebras American Mathematical Soc. Vol 366.
- [13] P. DELIGNE, Théorie de Hodge II, Inst. Hautes Études Sci. Publ. Math 40 (1971) 5-57.
- [14] B.M. DWORK, *p-adic cycles*, Inst. Hautes Études Sci. Publ. Math **37** (1969), 27–115.
- [15] B. DWORK, On p-adic differential equations I. The Frobenius structure of differential equations, Bull. Soc. Math. France 39-40 (1974), 27–37.
- [16] B. DWORK, On p-adic differential equations. IV. Generalized hypergeometric functions as p-adic analytic functions in one variable, Ann. Sci. École Norm. Sup. 6 (1973), 295–315.
- [17] B. DWORK, Bessel functions as p-adic functions of the argument, Duke Math. 41 (1974), 711-738.
- [18] B. DWORK, G. GEROTTO, AND F. SULLIVAN, An introduction to G-functions, Annals of Mathematics Studies 133, Princeton University Press, 1994.
- [19] A. ESCASSUT, Analytic Elements in p-adic Analysis, World Scientific Publishing Co., Inc., River Edge, NJ, 1995. xii+390 pp.
- [20] H. FURSTENBERG, Algebraic functions over finite fields, J. Algebra 7 (1967) 271–277.
- [21] K. KEDLAYA, p-adic differential equations, Cambridge studies in advanced mathematics 125, Cambridge University Press, 2010.
- [22] E. R. KOLCHIN, Algebraic groups and algebraic dependence, Amer. J. Math 90 (1968), 1151-1164.
- [23] M. KONTSEVICH, D. ZAGIER, Periods, in Mathematics unlimited 2001 and beyond Springer, 2001, 771–808.
- [24] P. ROBBA, Caractérisation des dérivées logarithmiques, Groupe d'Étude d'Analyse Ultramétrique, (1974/75), Exp. No 12, 6 pp.
- [25] T. RIVOAL, Les E-fonctions et G-fonctions de Siegel, Périodes et transcendance, Éditions de l'École polytechnique (2024), 189-284.
- [26] C. SIEGEL, Über einige Anwendungen diophantischer Approximationen, Abhandlungen Akad. Berlin 1929.
- [27] A. SHIDLOVSKII, Transcendental Numbers, W.de Gruyter, Berlin, New York, 1989.
- [28] D. VARGAS-MONTOYA, Algébricité modulo p, séries hypergéométriques et structure de Frobenius forte. Bull. Soc. Math. France 149 (2021), 439–477.
- [29] D. VARGAS-MONTOYA, Monodromie unipotente maximale, congruences "à la Lucas" et indépendance algébrique, Trans. Amer. Math. Soc. 377 (2024), 167–202.