## $\sqrt{-3}$ -SELMER GROUPS, IDEAL CLASS GROUPS AND LARGE 3-SELMER RANKS

SOMNATH JHA, DIPRAMIT MAJUMDAR AND PRATIKSHA SHINGAVEKAR

ABSTRACT. We consider the family of elliptic curves  $E_{a,b}: y^2 = x^3 + a(x-b)^2$  with  $a, b \in \mathbb{Z}$ . These elliptic curves have a rational 3-isogeny, say  $\varphi$ . We give an upper and a lower bound on the rank of the  $\varphi$ -Selmer group of  $E_{a,b}$  over  $K := \mathbb{Q}(\zeta_3)$  in terms of the 3-part of the ideal class group of certain quadratic extension of K. Using our bounds on the Selmer groups, we construct infinitely many curves in this family with arbitrary large 3-Selmer rank over K and no non-trivial K-rational point of order 3. We also show that for a positive proportion of natural numbers n, the curve  $E_{n,n}/\mathbb{Q}$  has root number -1 and 3-Selmer rank = 1.

#### INTRODUCTION

The *n*-Selmer group of an elliptic curve E defined over a number field F encapsulates various important arithmetic properties of E. Indeed, via the *n*-descent exact sequence

$$0 \longrightarrow E(F)/nE(F) \longrightarrow \operatorname{Sel}^{n}(E/F) \longrightarrow \operatorname{III}(E/F)[n] \longrightarrow 0,$$
(1)

it is closely related to the Mordell-Weil and the Tate-Shafarevich groups of the elliptic curve. A similar consideration holds, more generally, for an isogeny induced Selmer group of an elliptic curve (see (4)). Several important results on *n*-Selmer group of elliptic curves have been established recently by Bhargava et.al, Mazur-Rubin and many others. Soon after Cassels defined the Selmer group of an elliptic curve, he considered [Ca1] the relation between a (degree 3) isogeny induced Selmer group and the Tate-Shafarevich group of an elliptic curve over  $\mathbb{Q}$  with the 3-part of the ideal class group of certain quadratic extension of  $\mathbb{Q}$ . There are many results comparing the Selmer group of an elliptic curve with the ideal class group of a suitable number field including [BK, Če, Li, SS].

For a number field F and a finite set S of its finite primes, we let  $\mathcal{O}_S$  be the ring of S-integers in F and  $Cl_S(F)$  be its S-ideal class group. We denote by  $h_S^3$  the 3-rank of the  $Cl_S(F)$  i.e.  $h_S^3 := \dim_{\mathbb{F}_3} Cl_S(F) \otimes \mathbb{F}_3 = \dim_{\mathbb{F}_3} Cl_S(F)[3]$ . Throughout this paper, we set  $K := \mathbb{Q}(\zeta_3)$ , where  $\zeta_3 = \frac{-1+\sqrt{-3}}{2}$  is a primitive cube root of unity.

Now, let E be an elliptic curve over  $\mathbb{Q}$  with a rational isogeny  $\varphi: E \to \widehat{E}$  of degree 3. Then (up to a change of coordinates), the equation of E can be written either in the form (i)  $E_a: y^2 = x^3 + a$  with  $a \in \mathbb{Z}$ ,  $a \neq 0$  or (ii)  $E_{a,b}: y^2 = x^3 + a(x-b)^2$  with  $a, b \in \mathbb{Z}$ ,  $ab(4a + 27b) \neq 0$  (see [JMS] for details). In [JMS], we studied the  $\varphi$ -Selmer groups  $\operatorname{Sel}^{\varphi}(E_a/K)$  of the curves  $E_a: y^2 = x^3 + a$  over K. In fact, the  $\varphi$ -Selmer groups of  $E_a$  have also been studied by Bandini, [BES], Nekovář etc., whereas the study of the  $\varphi$ -Selmer groups of the curves  $E_{a,b}$  remains somewhat unexplored. In the first part of this article, we will complement the study on  $\varphi$ -Selmer rank bounds of  $E_a$  in [JMS] by exploring the  $\varphi$ - and 3-Selmer groups of the curves  $E_{a,b}: y^2 = x^3 + a(x-b)^2$  over K, denoted by  $\operatorname{Sel}^{\varphi}(E_{a,b}/K)$  and  $\operatorname{Sel}^3(E_{a,b}/K)$  respectively. The main goal of this article is to give an upper and a lower bound on the  $\mathbb{F}_3$ -ranks of  $\operatorname{Sel}^{\varphi}(E_{a,b}/K)$ , in terms of  $h_S^3$  for certain S-class groups of  $K(\sqrt{a})$ . Here the sets S are certain subsets of the set of primes of bad reduction of  $E_{a,b}$  over K (Definition 3.2). We then use these bounds for the Selmer groups induced by  $\varphi$  and by it's dual isogeny  $\widehat{\varphi}$  to get the upper and lower bounds for the 3-Selmer group  $\operatorname{Sel}^3(E_{a,b}/K)$ . The main result of this article is as stated below (cf. Theorem 3.8 for details).

**Theorem 0.1.** Let  $E_{a,b}: y^2 = x^3 + a(x-b)^2$  with  $a, b \in \mathbb{Z}$  and  $ab(4a+27b) \neq 0$  and  $\psi_{a,b}: E_{a,b} \rightarrow \widehat{E}_{a,b}$  be the rational isogeny of degree 3, defined in (2). Let  $S_1$ ,  $S_2$  and  $S_3$  (as in the Definition 3.2) be certain subsets of the set of primes of bad reduction of  $E_{a,b}$  over K. Assume that  $a \notin K^{*2}$  and put  $L = L_a := K[X]/(X^2 - a)$ .

<sup>2020</sup> Mathematics Subject Classification. Primary 11G05, 11R29, 11R34; Secondary 11G40, 11S25.

Key words and phrases. Elliptic curves, 3-Selmer groups, Ideal class groups, Arbitrary large 3-Selmer ranks.

Denote by  $S_{i,j}(L)$  the set of primes of L lying above  $S_i \cup S_j$ . If  $3 \nmid a$ , then

$$\max \left\{ h_{S_{1,2}(L)}^3, \ h_{S_{1,3}(L)}^3 + |S_3| - |S_2| - 1 \right\} \le \dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \\ \le \min \left\{ h_{S_{1,2}(L)}^3 + |S_{1,2}(L)| + |S_3| - |S_2| + 1, \ h_{S_{1,3}(L)}^3 + |S_{1,3}(L)| + 2 \right\}.$$

Further, the global root number  $\omega(E_{a,b}/K)$  is given by  $\omega(E_{a,b}/K) = (-1)^{|S_2|+|S_3|+1}$ .

In particular, if  $S_1$ ,  $S_2$  and  $S_3$  are all empty, then  $\dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \in \{h_L^3, h_L^3 + 1\}$  and the global root number  $\omega(E_{a,b}/K) = -1$ .

The bounds obtained here are indeed sharp and we have explicit numerical examples (see Example (2),  $\S$ 3) where the upper and the lower bound differ by 1.

We consider a subfamily  $E_{n,n}$ , where  $n \in \mathbb{N}$  and make suitable choices of n, such that  $\operatorname{Sel}^{\psi_{n,n}}(E_{n,n}/\mathbb{Q})$  vanishes. Using this, we prove in Proposition 3.10 that for at least 6% of square-free natural numbers n, the 3-rank of the 3-Selmer group of  $E_{n,n}$  over  $\mathbb{Q}$  i.e.  $\dim_{\mathbb{F}_3} \operatorname{Sel}^3(E_{n,n}/\mathbb{Q})$  is equal to 1.

We now briefly explain the key ideas behind the proof of Theorem 0.1. The existence of a rational 3isogeny  $\psi_{a,b}: E_{a,b} \to \widehat{E}_{a,b}$  implies that the residual representation  $E_{a,b}(\overline{\mathbb{Q}})[3]$  is a reducible  $G_{\mathbb{Q}}$ -module and gives rise to a  $G_{\mathbb{Q}}$ -stable order 3 subgroup  $(0, \pm \sqrt{a}b)$  which naturally corresponds to the quadratic étale algebra  $L_a := K[X]/(X^2 - a)$ . The Galois cohomology group  $H^1(G_K, E_{a,b}[\psi_{a,b}])$  can be identified with the kernel of the norm maps  $L_a^*/L_a^{*3} \longrightarrow K^*/K^{*3}$  (Prop. 1.2) and one obtains an explicit description of the isogeny induced Selmer group  $\operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K)$  in (6). We compute the images of the local Kummer maps and use some results in algebraic number theory (see §1.2, Props. 1.12 and 1.13) to first arrive at some preliminary bounds for  $\operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K)$  (Theorem 3.5) via certain S-ideal class groups of  $L_a$ . Further, we obtain sharper bounds for  $\operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K)$  in Theorem 3.8. This is achieved by using the formula of Cassels (13), that involves the study of local Tamagawa numbers to compare the sizes of  $\psi_{a,b}$ - and  $\widehat{\psi}_{a,b}$ -Selmer groups of  $E_{a,b}$  and  $\widehat{E}_{a,b}$ , respectively.

We then use these upper and lower bounds on  $\operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K)$  in Theorem 3.8 to exhibit a sub-family of the curves  $E_{a,b}$  that has arbitrary large 3-Selmer ranks in Theorem 4.1. Producing families of elliptic curves with large Selmer rank has been of considerable interest in the literature. Cassels [Ca1] produced an infinite family of elliptic curves  $E/\mathbb{Q}$  satisfying j(E) = 0,  $E(\mathbb{Q})[3] \neq 0$  and such that  $\dim_{\mathbb{F}_3} \operatorname{Sel}^3(E/\mathbb{Q})$  is unbounded as E varies. On the other hand, [KS] have produced families of elliptic curves with large *n*-Selmer groups over  $\mathbb{Q}$ , for  $n \geq 5$ . There is a recent important work of Bhargava et. al. [BKOS] on the arbitrarily large growth of 3-torsion of the Tate-Shafarevich groups of elliptic curves over number fields with cyclic 9-isogenies. See Remark 4.2 for details on related works. We prove the following result in this direction.

**Theorem 0.2.** Given any  $n \ge 0$ , there exist infinitely many elliptic curves  $E_{a,1}: y^2 = x^3 + a(x-1)^2$ , with  $a \in \mathbb{Z}$  such that  $\dim_{\mathbb{F}_3} \operatorname{Sel}^3(E_{a,1}/K) \ge \dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,1}}(E_{a,1}/K) \ge 2n$ . These curves have a rational 3-isogeny and no non-trivial 3-torsion points over K.

The article consists of 4 sections and is structured as follows: In §1, we state some preliminary definitions and recall some algebraic results from [JMS] which will be used later. In §2, we will compute the images of local Kummer maps at each prime of K. Our main results on the  $\mathbb{F}_3$ -rank bounds on the Selmer groups (and the proof of Proposition 3.10) is contained in §3. The result on arbitrary large 3-Selmer ranks is discussed in §4. Table 2 contains examples of the curves  $E_{a,b}$  for which we compute the bounds on their  $\psi_{a,b}$ - and 3-Selmer groups over K using our results from §3.

### 1. Preliminaries

1.1. The curves  $E_{a,b}$  and the 3-isogenies. The rational degree 3-isogeny corresponds to a  $G_{\mathbb{Q}}$ -invariant subgroup C of  $E_{a,b}(\overline{\mathbb{Q}})$  of order 3 and given C, the rational 3-isogeny  $\varphi$  is obtained as  $\varphi : E_{a,b} \to E_{a,b}/C$ . Via a change of variables  $(x, y) \mapsto (\frac{x}{9} - \frac{4a}{3}, \frac{y}{27})$ , we can identify  $E_{a,b}/C$  with  $\hat{E}_{a,b} := E_{-27a,4a+27b}$ . Thus, we obtain a rational 3-isogeny  $\psi_{a,b} : E_{a,b} \to \hat{E}_{a,b}$  given explicitly using [Vé] as

$$\psi_{a,b}(x,y) = \left(\frac{9(x^3 + \frac{4}{3}ax^2 - 4abx + 4ab^2)}{x^2}, \frac{27y(x^3 + 4abx - 8ab^2)}{x^3}\right).$$
(2)

Putting d = 4a + 27b, we obtain the corresponding dual isogeny  $\widehat{\psi}_{a,b} : \widehat{E}_{a,b} \to E_{a,b}$  given by

$$\widehat{\psi}_{a,b}(x,y) = \left(\frac{x^3 - 36ax^2 + 108adx - 108ad^2}{3^4x^2}, \frac{y(x^3 - 108adx + 216ad^2)}{3^6x^3}\right).$$
(3)

Set  $\zeta_3 := \frac{-1+\sqrt{-3}}{2}$  and  $K := \mathbb{Q}(\zeta_3)$ , and note that there is an isomorphism  $\lambda_c : E_{ac^2, bc^2} \xrightarrow{\sim} E_{a,b}$  defined by  $(x,y) \mapsto (\bar{c}^{-2}x, c^{-3}y)$  over K for any  $c \in K^*$ . Thus, without any loss of generality, we can assume that gcd(a,b) is square-free in K. Further, observe that  $\widehat{\psi}_{a,b} = \lambda_{27} \circ \psi_{-27a,d}$  and hence one has an isomorphism of Selmer groups  $\operatorname{Sel}^{\widehat{\psi}_{a,b}}(\widehat{E}_{a,b}/K) \cong \operatorname{Sel}^{\psi_{-27a,d}}(E_{-27a,d}/K).$ 

**Notation:** For any finite set X, |X| denotes its cardinality.

- Given a multiplicative group G, define the subgroup  $G^3 := \{x^3 \mid x \in G\}$ .
- For abelian groups A, A' and a homomorphism  $\eta: A \to A'$ , let  $A[\eta]$  denote the kernel of  $\eta$ .
- F denotes an arbitrary number field and let  $\Sigma_F$  be the set of all its finite places. For a finite subset S of  $\Sigma_F$ ,  $\mathcal{O}_S$  denotes the ring of S-integers in F. For  $\varpi \in \Sigma_F$ ,  $F_{\varpi}$  will denote the completion of F at  $\varpi$  and its ring of integers will be denoted by  $\mathcal{O}_{F_{\pi}}$ .
- $\zeta_3 = \frac{-1+\sqrt{-3}}{2}$  is the primitive third root of unity,  $K = \mathbb{Q}(\zeta_3)$  and  $\mathcal{O}_K = \mathbb{Z}[\zeta_3]$  the ring of integers of K. We will denote a finite prime of  $\mathcal{O}_K$  by  $\mathfrak{q}$ .
- $(\mathfrak{p}) = (1 \zeta_3)$  is the unique prime of K above 3 with uniformizer  $\mathfrak{p} = 1 \zeta_3$  in  $\mathcal{O}_{K_{\mathfrak{p}}}$ . By abuse of notation, we simply write  $\mathfrak{p}$  for the ideal ( $\mathfrak{p}$ ) of  $\mathcal{O}_K$ .
- We set  $L = L_a = \frac{K[X]}{(X^2 a)}$ . Thus L/K is a quadratic field extension if  $a \notin K^{*2}$ . Otherwise,  $L \cong K \times K$ if  $a \in K^{*2}$ .
- Define A := O<sub>L</sub>, the ring of integers of L if a ∉ K<sup>\*2</sup>. If a ∈ K<sup>\*2</sup>, then set A := O<sub>K</sub> × O<sub>K</sub>.
  Next, we define the quadratic étale K<sub>q</sub>-algebras: L<sub>q</sub> = K<sub>q</sub>[X]/(X<sup>2</sup>-a). Thus L<sub>q</sub>/K<sub>q</sub> is a quadratic field extension, if a ∉ K<sup>\*2</sup><sub>q</sub>; otherwise, L<sub>q</sub> ≅ K<sub>q</sub> × K<sub>q</sub>, if a ∈ K<sup>\*2</sup><sub>q</sub>.
- Define A<sub>q</sub> := { the ring of integers O<sub>Lq</sub> of the field L<sub>q</sub>, if a ∉ K<sub>q</sub><sup>\*2</sup>, O<sub>Kq</sub> × O<sub>Kq</sub>, if a ∈ K<sub>q</sub><sup>\*2</sup>.
  N<sub>L/K</sub> : L → K denotes the field norm if L is field and the multiplication of co-ordinates if L ≅ K×K.
- A similar notation is followed for  $L_{\mathfrak{g}}$  and  $K_{\mathfrak{g}}$ .

For  $F \in \{K, K_{\mathfrak{q}}\}$ , let  $\delta_{\psi_{a,b},F} : \widehat{E}_{a,b}(F) \longrightarrow \widehat{E}_{a,b}(F)/\psi_{a,b}(E(F)) \xrightarrow{\overline{\delta}_{\psi_{a,b},F}} H^1(G_F, E[\psi_{a,b}])$  be the Kummer map. Then we have the following commutative diagram:

**Definition 1.1.** The  $\psi_{a,b}$ -Selmer group of  $E_{a,b}$  over K, denoted  $\operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K)$ , is defined as

$$\operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) := \left\{ \xi \in H^1(G_K, E_{a,b}[\psi_{a,b}]) \mid \operatorname{res}_{\mathfrak{q}}(\xi) \in \operatorname{Im} \delta_{\psi_{a,b}, K_{\mathfrak{q}}} \text{ for every } \mathfrak{q} \in \Sigma_K \right\}.$$

Setting  $\operatorname{III}(E_{a,b}/K) := \operatorname{Ker}(H^1(G_K, E_{a,b}) \to \prod_{\mathfrak{q} \in \Sigma_K} H^1(G_{K_{\mathfrak{q}}}, E_{a,b}))$  to be the Tate-Shafarevich group of  $E_{a,b}$ over K, we get the fundamental exact sequence:

$$0 \longrightarrow \widehat{E}_{a,b}(K)/\psi_{a,b}(E_{a,b}(K)) \longrightarrow \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \longrightarrow \operatorname{III}(E_{a,b}/K)[\psi_{a,b}] \longrightarrow 0$$
(4)

For the isogeny given by the multiplication-by-3 map on  $E_{a,b}$ , the 3-Selmer group Sel<sup>3</sup> $(E_{a,b}/K)$  is defined in a similar manner.

The following result can be deduced from [BES, Prop. 24].

**Proposition 1.2.** There is an isomorphism of group schemes  $E_{a,b}[\psi_{a,b}] \cong \operatorname{Ker}(\operatorname{Res}_{K}^{L_{a}} \mu_{3} \to \mu_{3})$  and an induced isomorphism

 $H^1(G_K, E_{a,b}[\psi_{a,b}]) \cong (L_a^*/L_a^{*3})_{N=1},$ 

where  $(L_a^*/L_a^{*3})_{N=1}$  denotes the kernel of the norm map  $\overline{N}_{L_a/K}: L_a^*/L_a^{*3} \to K^*/K^{*3}$ . The same statement holds if we replace  $(E_{a,b}, \psi_{a,b})$  by  $(\widehat{E}_{a,b}, \widehat{\psi}_{a,b})$ .

We denote the composite map  $\widehat{E}_{a,b}(K) \longrightarrow \widehat{E}_{a,b}(K)/\psi_{a,b}(E_{a,b}(K)) \hookrightarrow H^1(G_K, E_{a,b}[\psi_{a,b}]) \xrightarrow{\sim} (L^*/L^{*3})_{N=1}$ again by  $\delta_{\psi_{a,b},K}$ , in view of Prop. 1.2. Similarly, we will view  $\delta_{\psi_{a,b},K_{\mathfrak{q}}}(\widehat{E}_{a,b}(K_{\mathfrak{q}}))$  as a subset of  $(L_{\mathfrak{q}}^*/L_{\mathfrak{q}}^{*3})_{N=1}$ .

We have the commutative diagram,

To ease the notation, via the canonical embedding  $\iota_{\mathfrak{q}}: K \to K_{\mathfrak{q}}$ , we identify  $\iota_{\mathfrak{q}}(x)$  with  $x \in K$ ; similarly for L. Hence from (5), if  $\overline{x} \in (L^*/L^{*3})_{N=1}$ , then  $\overline{x} \in (L^*_{\mathfrak{q}}/L^{*3}_{\mathfrak{q}})_{N=1}$ .

We now have the following alternative description of  $\operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K)$ :

$$\operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) = \left\{ \overline{x} \in (L^*/L^{*3})_{N=1} \mid \overline{x} \in \operatorname{Im} \delta_{\psi_{a,b},K_{\mathfrak{q}}} \text{ for all } \mathfrak{q} \in \Sigma_K \right\}.$$
(6)

Similarly, if  $a \in K^{*2}$ , then the definition in (6) can be written more explicitly:

$$\operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) = \left\{ (\overline{x}_1, \overline{x}_2) \in (K^*/K^{*3} \times K^*/K^{*3})_{N=1} \mid (\overline{x}_1, \overline{x}_2) \in \operatorname{Im} \delta_{\psi_{a,b}, K_{\mathfrak{q}}} \text{ for all } \mathfrak{q} \in \Sigma_K \right\}.$$
(7)

These will be our working definitions of the Selmer groups for the rest of the article.

1.2. Some algebraic number theory. Here, we will recall some basic algebraic results from [JMS] which will be used later. The proofs can be found in [JMS, §2].

**Definition 1.3.** We define  $(A^*/A^{*3})_{N=1}$  to be the kernel of the norm map  $\overline{N}_{L/K} : A^*/A^{*3} \to \mathcal{O}_K^*/\mathcal{O}_K^{*3}$ induced from the norm map  $\overline{N}_{L/K} : L^*/L^{*3} \to K^*/K^{*3}$ . The group  $(A_{\mathfrak{q}}^*/A_{\mathfrak{q}}^{*3})_{N=1}$  is defined similarly by replacing A and  $\mathcal{O}_K$  with  $A_{\mathfrak{q}}$  and  $\mathcal{O}_{K_{\mathfrak{q}}}$ , respectively, in this definition.

**Proposition 1.4.** For 
$$\mathfrak{q} \nmid 3$$
, we have  $\left| \begin{pmatrix} A_{\mathfrak{q}}^* \\ A_{\mathfrak{q}}^{*3} \end{pmatrix}_{N=1} \right| = \begin{cases} 1, & \text{if } a \notin K_{\mathfrak{q}}^{*2}, \\ 3, & \text{if } a \in K_{\mathfrak{q}}^{*2}. \end{cases}$   
On the other hand, for  $\mathfrak{p} \mid 3$ , we have  $\left| \begin{pmatrix} A_{\mathfrak{p}}^* \\ A_{\mathfrak{p}}^{*3} \end{pmatrix}_{N=1} \right| = \begin{cases} 9, & \text{if } a \notin K_{\mathfrak{p}}^{*2}, \\ 27, & \text{if } a \in K_{\mathfrak{p}}^{*2}. \end{cases}$ 

For a local field F with the ring of integers  $\mathcal{O}_F$  and uniformizer  $\pi$ , we let  $U_F^n := 1 + \pi^n \mathcal{O}_F$ . Given an element  $u \in \mathcal{O}_F^*$ , we let  $\overline{u}$  denote its image in  $\mathcal{O}_F^*/U_F^n \cong (\mathcal{O}_F/\pi^n \mathcal{O}_F)^*$ .

**Definition 1.5.** For a local field F, define

$$V_F = \left\{ u \in \mathcal{O}_F^* \mid \overline{u} \in \mathcal{O}_F^* / U_F^3 \text{ satisfies } \overline{u} = \alpha^3 \text{ for some } \alpha \in \mathcal{O}_F^* / U_F^3 \right\}$$

Next, we define a subset  $V_3$  of  $A^*_{\mathfrak{p}}$ :

$$V_{3} = \begin{cases} \left\{ u \in V_{L_{\mathfrak{p}}} \mid N_{L_{\mathfrak{p}}/K_{\mathfrak{p}}}(u) \in \mathcal{O}_{K_{\mathfrak{p}}}^{*3} \right\}, & \text{if } a \notin K_{\mathfrak{p}}^{*2}, \\ \left\{ (u_{1}, u_{2}) \in V_{K_{\mathfrak{p}}} \times V_{K_{\mathfrak{p}}} \mid u_{1}u_{2} \in \mathcal{O}_{K_{\mathfrak{p}}}^{*3} \right\}, & \text{if } a \in K_{\mathfrak{p}}^{*2}. \end{cases}$$

Note that  $A_{\mathfrak{p}}^{*3} \subseteq V_3$  and  $V_3/A_{\mathfrak{p}}^{*3} \subseteq (A_{\mathfrak{p}}^*/A_{\mathfrak{p}}^{*3})_{N=1}$ , whether or not *a* is a square in  $K_{\mathfrak{p}}^*$ .

**Lemma 1.6.** Let 
$$a \in K_{\mathfrak{p}}^{*2}$$
, then  $\left(\frac{A_{\mathfrak{p}}^{*}}{A_{\mathfrak{p}}^{*3}}\right)_{N=1} \cong \frac{U_{K_{\mathfrak{p}}}^{1}}{U_{K_{\mathfrak{p}}}^{4}}$  and  $\frac{V_{3}}{A_{\mathfrak{p}}^{*3}} \cong \frac{U_{K_{\mathfrak{p}}}^{3}}{U_{K_{\mathfrak{p}}}^{4}}$ .

**Proposition 1.7.** We have 
$$\left|\frac{V_3}{A_{\mathfrak{p}}^{*3}}\right| = 1$$
 if  $a \notin K_{\mathfrak{p}}^{*2}$  and  $\left|\frac{V_3}{A_{\mathfrak{p}}^{*3}}\right| = 3$  if  $a \in K_{\mathfrak{p}}^{*2}$ .

**Notation**: For a non-archimedean local field F with discrete valuation v and  $\overline{x} \in F^*/F^{*3}$ , define  $v(\overline{x}) \in \mathbb{Z}/3\mathbb{Z}$  by  $v(\overline{x}) := v(x) \pmod{3}$ , for any lift  $x \in F^*$ . Similarly, for an element  $(\overline{y}, \overline{z})$  of  $F^*/F^{*3} \times F^*/F^{*3}$ , we define  $v(\overline{y}, \overline{z}) := (v(y) \pmod{3}, v(z) \pmod{3}) \in \mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$ . If both  $v(y), v(z) \equiv 0 \pmod{3}$ , then by abuse of notation, we write  $v(\overline{y}, \overline{z}) \equiv 0 \pmod{3}$ .

For a prime  $q \in \Sigma_K$ , we denote the induced valuation on  $L_q$  simply by  $v_q$ .

**Lemma 1.8.** For an element 
$$\overline{x}$$
 of  $\left(\frac{L_{\mathfrak{q}}^*}{L_{\mathfrak{q}}^{*3}}\right)_{N=1}$ , one has  $\overline{x} \in \left(\frac{A_{\mathfrak{q}}^*}{A_{\mathfrak{q}}^{*3}}\right)_{N=1}$  if and only if  $v_{\mathfrak{q}}(\overline{x}) \equiv 0 \pmod{3}$ .  $\Box$ 

**Definition 1.9.** Let S be a finite set of finite primes of K and recall that  $\mathcal{O}_S$  denotes the ring of S-integers in K. If  $a \notin K^{*2}$ , then  $L = K(\sqrt{a})$ . Let  $S(L) = \{\mathfrak{Q} \in \Sigma_L \mid \mathfrak{Q} \cap K \in S\}$ . Further,  $\mathcal{O}_{S(L)}$  will denote the ring of S(L)-integers in L. Let  $x \in L^*$  be any lift of  $\overline{x} \in L^*/L^{*3}$ . We define the following  $\mathbb{F}_3$ -vector spaces:

$$M(S,a) = \left\{ \overline{x} \in L^*/L^{*3} : L(\sqrt[3]{x})/L \text{ is unramified and } x \in L^{*3}_{\mathfrak{q}} \text{ for all } \mathfrak{q} \in S \right\},$$

$$N(S,a) = \left\{ \overline{x} \in L^*/L^{*3} : (x) = I^3 \text{ for some fractional ideal } I \text{ of } \mathcal{O}_{S(L)} \right\}.$$

Note that if  $\overline{x} \in M(S, a)$ , then any lift  $x \in L^*$  of  $\overline{x}$  satisfies  $(x) = I^3$  for some fractional ideal I of  $\mathcal{O}_{S(L)}$  ([Wa, Ch. 9]). This implies that  $N_{L/K}(x) \in K^{*3}$ . So,  $M(S, a) \subset N(S, a) \subset (L^*/L^{*3})_{N=1}$ .

**Definition 1.10.** Let S be a finite set of finite primes of K. When  $a \in K^{*2}$ , we define

 $N'(S,a) := \left\{ (\overline{x}_1, \overline{x}_2) \in \left( K^* / K^{*3} \times K^* / K^{*3} \right)_{N=1} \mid v_{\mathfrak{q}}(\overline{x}_1, \overline{x}_2) \equiv 0 \pmod{3} \text{ for all } \mathfrak{q} \notin S \right\}.$ 

**Definition 1.11.** We assume that  $a \notin K^{*2}$  and denote by  $Cl_{S(L)}(L)$  the S(L)-ideal class group of L i.e. the class group of  $\mathcal{O}_{S(L)}$ . Further, let  $h_{S(L)}^3$  denote the 3-rank of  $Cl_{S(L)}(L)$  i.e.  $h_{S(L)}^3 := \dim_{\mathbb{F}_3} Cl_{S(L)}(L) \otimes_{\mathbb{Z}} \mathbb{F}_3 = \dim_{\mathbb{F}_3} Cl_{S(L)}(L)[3]$ . In particular, if S(L) is empty, then  $Cl_{\emptyset}(L) = Cl(L)$  is the ideal class group of L and  $h_L^3 = \dim_{\mathbb{F}_3} Cl(L) \otimes_{\mathbb{Z}} \mathbb{F}_3$ .

**Proposition 1.12.** If  $a \notin K^{*2}$ , we have  $\dim_{\mathbb{F}_3} M(S, a) = h^3_{S(L)}$  and  $\dim_{\mathbb{F}_3} N(S, a) = h^3_{S(L)} + |S(L)| + 2$ . On the other hand, one has that  $\dim_{\mathbb{F}_3} N'(S, a) = |S| + 1$  when  $a \in K^{*2}$ .

We have an equivalently description of the  $\mathbb{F}_3$ -modules M(S, a), N(S, a) and N'(S, a), which are compatible with the definition of Selmer groups given in (6) and (7).

**Proposition 1.13.** When  $a \notin K^{*2}$ , we have

$$\begin{split} M(S,a) &= \Big\{ \overline{x} \in \Big(\frac{L^*}{L^{*3}}\Big)_{N=1} \Big| \ \overline{x} \in \Big(\frac{A_{\mathfrak{q}}^*}{A_{\mathfrak{q}}^{*3}}\Big)_{N=1} \ \text{if } \mathfrak{q} \notin S \cup \{\mathfrak{p}\}, \ \overline{x} = \overline{1} \ \text{if } \mathfrak{q} \in S \ \text{and } \overline{x} \in \frac{V_3}{A_{\mathfrak{p}}^{*3}} \ \text{if } \mathfrak{p} \notin S \Big\} \\ \text{and } N(S,a) &= \Big\{ \overline{x} \in \Big(\frac{L^*}{L^{*3}}\Big)_{N=1} \Big| \ \overline{x} \in \Big(\frac{A_{\mathfrak{q}}^*}{A_{\mathfrak{q}}^{*3}}\Big)_{N=1} \ \text{for all } \mathfrak{q} \notin S \Big\}. \\ \text{Whereas for } a \in K^{*2}, \ N'(S,a) &= \Big\{ (\overline{x}_1, \overline{x}_2) \in \Big(\frac{K^*}{K^{*3}} \times \frac{K^*}{K^{*3}}\Big)_{N=1} \Big| \ (\overline{x}_1, \overline{x}_2) \in \Big(\frac{A_{\mathfrak{q}}^*}{A_{\mathfrak{q}}^{*3}}\Big)_{N=1} \ \text{for all } \mathfrak{q} \notin S \Big\}. \\ \Box \end{split}$$

2. Local theory for the curves  $E_{a,b}$ 

Let F be a local field of characteristic 0 and  $\omega$  be a uniformizer of  $\mathcal{O}_F$ . Suppose  $\varphi: E \to \widehat{E}$  be an isogeny of elliptic curves and  $c_{\omega}(E)$  and  $c_{\omega}(\widehat{E})$  denote the Tamagawa numbers of E and  $\widehat{E}$  at  $\omega$ . By a theorem of Schaefer [Sc, Lemma 3.8], we know that

$$|\widehat{E}(F)/\varphi(E(F))| = \frac{||\varphi'(0)||_{\omega}^{-1} \times |E(F)[\varphi]| \times c_{\omega}(\widehat{E})}{c_{\omega}(E)},$$
(8)

where  $||.||_{\omega}$  is the norm in F and  $\varphi'(0)$  is the leading coefficient of the power series representation of  $\varphi$  on the formal group of  $\hat{E}$  [Si1, §4]. Using (8), we will compute  $|\hat{E}(K_{\mathfrak{q}})/\varphi(E(K_{\mathfrak{q}}))|$  for all  $\mathfrak{q}$ .

Consider the curves  $E_{a,b}: y^2 = x^3 + a(x-b)^2$ , with  $ab(4a+27b) \neq 0$ ,  $a,b \in \mathbb{Z}$  such that gcd(a,b) is square-free in K and the rational 3-isogeny  $\psi_{a,b}: E_{a,b} \to \widehat{E}_{a,b}$ , where  $\widehat{E}_{a,b}: y^2 = x^3 - 27a(x-d)^2$  with d = 4a+27b. In this subsection, we give an explicit description of the Kummer maps  $\delta_{\psi_{a,b},K_{\mathfrak{q}}}$  for all  $\mathfrak{q} \in \Sigma_K$ . But first, we state a general result (cf. [JMS, Lemma 3.1] for the proof).

**Lemma 2.1.** Let  $\varphi : E \to \widehat{E}$  be a 3-isogeny defined over K. Recall from §1 that  $\delta_{\varphi,K_{\mathfrak{q}}}(E(K_{\mathfrak{q}})) \subset (L_{\mathfrak{q}}^*/L_{\mathfrak{q}}^{*3})_{N=1}$ . If E (and hence  $\widehat{E}$ ) has good reduction at a prime  $\mathfrak{q} \nmid 3$  in K, then for any  $P \in \widehat{E}(K_{\mathfrak{q}})$ ,  $v_{\mathfrak{q}}(\delta_{\varphi,K_{\mathfrak{q}}}(P)) \equiv 0 \pmod{3}$ . In particular,  $\delta_{\varphi,K_{\mathfrak{q}}}(\widehat{E}(K_{\mathfrak{q}})) \subset (A_{\mathfrak{q}}^*/A_{\mathfrak{q}}^{*3})_{N=1}$ .

For  $P \in \widehat{E}(K_{\mathfrak{q}})$ ,  $\delta_{\varphi,K_{\mathfrak{q}}}(P) = \overline{t} \in L_{\mathfrak{q}}^*/L_{\mathfrak{q}}^{*3}$ , if  $L_{\mathfrak{q}}$  is a field and  $\delta_{\varphi,K_{\mathfrak{q}}}(P) = (\overline{t_1},\overline{t_2})$ , if  $L_{\mathfrak{q}} \cong K_{\mathfrak{q}} \times K_{\mathfrak{q}}$ . To ease the notation, we write  $\delta_{\varphi,K_{\mathfrak{q}}}(P) = t$  or  $(t_1,t_2)$ , respectively.

**Proposition 2.2.** [Ca3, §14, §15] Let  $\psi_{a,b} : E_{a,b} \to \widehat{E}_{a,b}$  be the K-isogeny defined in (2). Recall that d = 4a + 27b. We have the Kummer map  $\delta_{\psi_{a,b},K_{\mathfrak{q}}} : \widehat{E}_{a,b}(K_{\mathfrak{q}}) \to (L_{\mathfrak{q}}^*/L_{\mathfrak{q}}^{*3})_{N=1}$  with the kernel  $\operatorname{Ker}(\delta_{\psi_{a,b},K_{\mathfrak{q}}}) = \psi_{a,b}(E_{a,b}(K_{\mathfrak{q}})).$ 

If  $L_{\mathfrak{q}}$  is a field, then  $\delta_{\psi_{a,b},K_{\mathfrak{q}}}$  is given by  $\delta_{\psi_{a,b},K_{\mathfrak{q}}}(P) = \begin{cases} 1, & \text{if } P = O, \\ y(P) - \mathfrak{p}^3 \sqrt{a} (x(P) - d), & \text{otherwise.} \end{cases}$ If  $L_{\mathfrak{q}} \cong K_{\mathfrak{q}} \times K_{\mathfrak{q}}$ , then  $\delta_{\psi_{a,b},K_{\mathfrak{q}}}$  is given by

$$\delta_{\psi_{a,b},K_{\mathfrak{q}}}(P) = \begin{cases} (1,1), & \text{if } P = O, \\ (2\mathfrak{p}^{3}\sqrt{a}d, \frac{1}{2\mathfrak{p}^{3}\sqrt{a}d}), & \text{if } P = (0,\mathfrak{p}^{3}\sqrt{a}d), \\ (-\frac{1}{2\mathfrak{p}^{3}\sqrt{a}d}, -2\mathfrak{p}^{3}\sqrt{a}d), & \text{if } P = (0,-\mathfrak{p}^{3}\sqrt{a}d), \\ (y(P)-\mathfrak{p}^{3}\sqrt{a}(x(P)-d), y(P)+\mathfrak{p}^{3}\sqrt{a}(x(P)-d)), & \text{if } P \notin \widehat{E}_{a,b}(K_{\mathfrak{q}})[\widehat{\psi}_{a,b}]. & \Box \end{cases}$$

**Lemma 2.3.** [Si1, § 7] Let  $\ell \in \mathbb{Z}$  be a prime. If F is a local field,  $\omega$  is a prime in F and  $\varphi : E \to \widehat{E}$  is an  $\ell$ -isogeny over F, then

- (1) E has good reduction at  $\omega$  if and only if  $\hat{E}$  has good reduction at  $\omega$ .
- (2) E has split multiplicative reduction at  $\omega$  if and only if  $\hat{E}$  has split multiplicative reduction at  $\omega$ . In this situation, one of E and  $\hat{E}$  is of Kodaira type  $I_n$ , while the other is of the type  $I_{\ell n}$  with  $n = -v_{\omega}(j(E))$ .
- 2.1. Local theory at primes  $\mathfrak{q} \nmid 3$ . Let  $\mathfrak{q}$  be a prime of  $K = \mathbb{Q}(\zeta_3)$  not dividing 3.

**Proposition 2.4.** If  $a \notin K_{\mathfrak{q}}^{*2}$ , then we have  $\frac{\widehat{E}_{a,b}(K_{\mathfrak{q}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{q}}))} = \left(\frac{A_{\mathfrak{q}}^*}{A_{\mathfrak{q}}^{*3}}\right)_{N=1} = \{1\}.$ 

Proof. Since  $a \notin K_{\mathfrak{q}}^{*2}$ , we have  $E_{a,b}(K_{\mathfrak{q}})[\psi_{a,b}] = \{O\}$ . Using (8), we get  $\left|\frac{\widehat{E}_{a,b}(K_{\mathfrak{q}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{q}}))}\right| = \frac{c_{\mathfrak{q}}(\widehat{E}_{a,b})}{c_{\mathfrak{q}}(E_{a,b})}$  and  $\left|\frac{E_{a,b}(K_{\mathfrak{q}})}{\widehat{\psi}_{a,b}(\widehat{E}_{a,b}(K_{\mathfrak{q}}))}\right| = \frac{c_{\mathfrak{q}}(\widehat{E}_{a,b})}{c_{\mathfrak{q}}(\widehat{E}_{a,b})}$ . Thus,  $\frac{\widehat{E}_{a,b}(K_{\mathfrak{q}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{q}}))}$  is trivial, also  $\left(A_{\mathfrak{q}}^*/A_{\mathfrak{q}}^{*3}\right)_{N=1} = \{1\}$  by Prop. 1.4.

**Proposition 2.5.** Suppose  $a \in K_{\mathfrak{q}}^{*2}$  and  $E_{a,b}$  (or equivalently  $\widehat{E}_{a,b}$ ) has good reduction at  $\mathfrak{q}$ . Then we have that  $\left|\frac{\widehat{E}_{a,b}(K_{\mathfrak{q}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{q}}))}\right| = 3$  and hence  $\frac{\widehat{E}_{a,b}(K_{\mathfrak{q}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{q}}))} \cong \left(\frac{A_{\mathfrak{q}}^{*}}{A_{\mathfrak{q}}^{*3}}\right)_{N=1}$ .

Proof. As  $a \in K_{\mathfrak{q}}^{*2}$ ,  $|E_{a,b}[\psi_{a,b}](K_{\mathfrak{q}})| = 3$  and again  $||\psi'_{a,b}(0)||_{\mathfrak{q}}^{-1} = 1$  by [Sc]. Since  $\mathfrak{q}$  is a good prime for  $E_{a,b}$  (and  $\widehat{E}_{a,b}$ ), we have  $|\frac{\widehat{E}_{a,b}(K_{\mathfrak{q}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{q}}))}| = 3$  by (8). The result follows from Prop. 1.4 and Lemma 2.1.

Next, let  $\mathfrak{q} \nmid 3$  be a prime of K such that  $a \in K_{\mathfrak{q}}^{*2}$  and  $\mathfrak{q} \mid \Delta_{E_{a,b}} = -2^4 a^2 b^3 d$ . The local data obtained for each such  $\mathfrak{q}$  using Tate's algorithm [Si2, Ch. 4, Algo. 9.4], the Schaefer's formula (8) and Prop. 1.4 is consolidated in Table 1 below.

$a \in K_{\mathfrak{q}}^{*2}$ and $\mathfrak{q}$ divides $\Delta_{E_{a,b}}$	Tamagawa Numbers	
$\begin{array}{c c} \mathfrak{q} \mid a \text{ and} \\ \upsilon_{\mathfrak{q}}(4ab^2) \equiv 0 \pmod{6} \end{array}$	$c_{\mathfrak{q}}(E_{a,b}) = c_{\mathfrak{q}}(\widehat{E}_{a,b}) = 1$	$\frac{\widehat{E}_{a,b}(K_{\mathfrak{q}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{q}}))} = \left(\frac{A_{\mathfrak{q}}^*}{A_{\mathfrak{q}}^{*3}}\right)_{N=1}$
$\begin{array}{c c} \mathfrak{q} \mid a \text{ and} \\ \upsilon_{\mathfrak{q}}(4ab^2) \equiv 2 \pmod{6} \end{array}$	$c_{\mathfrak{q}}(E_{a,b}) = c_{\mathfrak{q}}(\widehat{E}_{a,b}) = 3$	$\frac{\widehat{E}_{a,b}(K_{\mathfrak{q}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{q}}))} \cap \left(\frac{A_{\mathfrak{q}}^*}{A_{\mathfrak{q}}^{*3}}\right)_{N=1} = \{1\}$
$\begin{array}{c c} \mathfrak{q} \mid a \text{ and} \\ \upsilon_{\mathfrak{q}}(4ab^2) \equiv 4 \pmod{6} \end{array}$	$c_{\mathfrak{q}}(E_{a,b}) = c_{\mathfrak{q}}(\widehat{E}_{a,b}) = 3$	$\frac{\widehat{E}_{a,b}(K_{\mathfrak{q}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{q}}))} \cap \left(\frac{A_{\mathfrak{q}}^*}{A_{\mathfrak{q}}^{*3}}\right)_{N=1} = \{1\}$
$\mathfrak{q} \mid b \text{ and } \mathfrak{q} \nmid 2a$	$c_{\mathfrak{q}}(E_{a,b}) = 3\upsilon_{\mathfrak{q}}(b)$ $c_{\mathfrak{q}}(\widehat{E}_{a,b}) = \upsilon_{\mathfrak{q}}(b)$	$\{1\} = \frac{\widehat{E}_{a,b}(K_{\mathfrak{q}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{q}}))} \subsetneq \left(\frac{A_{\mathfrak{q}}^*}{A_{\mathfrak{q}}^{*3}}\right)_{N=1}$
$\mathfrak{q} \mid d \text{ and } \mathfrak{q} \nmid 2a$	$c_{\mathfrak{q}}(E_{a,b}) = v_{\mathfrak{q}}(d)$ $c_{\mathfrak{q}}(\widehat{E}_{a,b}) = 3v_{\mathfrak{q}}(d)$	$\left(\frac{A_{\mathfrak{q}}^*}{A_{\mathfrak{q}}^{*3}}\right)_{N=1} \subsetneq \frac{\widehat{E}_{a,b}(K_{\mathfrak{q}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{q}}))}$
$\begin{array}{c} \mathfrak{q} = 2\\ 2 \nmid a \text{ and } 2 \nmid b \end{array}$	$c_{\mathfrak{q}}(E_{a,b}) = c_{\mathfrak{q}}(\widehat{E}_{a,b}) = 3$	$\frac{\widehat{E}_{a,b}(K_{\mathfrak{q}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{q}}))} \cap \left(\frac{A_{\mathfrak{q}}^*}{A_{\mathfrak{q}}^{*3}}\right)_{N=1} = \{1\}$
$\begin{aligned} \mathbf{q} &= 2\\ 2 \nmid a \text{ and } 2    b \end{aligned}$	$c_{\mathfrak{q}}(E_{a,b}) = c_{\mathfrak{q}}(\widehat{E}_{a,b}) = 3$	$\frac{\widehat{E}_{a,b}(K_{\mathfrak{q}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{q}}))} \cap \left(\frac{A_{\mathfrak{q}}^*}{A_{\mathfrak{q}}^{*3}}\right)_{N=1} = \{1\}$
q = 2, 2 \ a and 4   b	$c_{\mathfrak{q}}(\widehat{E}_{a,b}) = v_{\mathfrak{q}}(d) - 2$ $c_{\mathfrak{q}}(\widehat{E}_{a,b}) = 3(v_{\mathfrak{q}}(d) - 2)$	$\left(\frac{A_{\mathfrak{q}}^*}{A_{\mathfrak{q}}^{*3}}\right)_{N=1} \subsetneq \frac{\widehat{E}_{a,b}(K_{\mathfrak{q}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{q}}))}$
q = 2, 2 \not a and 8 \not b	$c_{\mathfrak{q}}(\overline{E}_{a,b}) = 3(\upsilon_{\mathfrak{q}}(b) - 2)$ $c_{\mathfrak{q}}(\widehat{E}_{a,b}) = \upsilon_{\mathfrak{q}}(b) - 2$	$\{1\} = \frac{\widehat{E}_{a,b}(K_{\mathfrak{q}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{q}}))} \subsetneq \left(\frac{A_{\mathfrak{q}}^*}{A_{\mathfrak{q}}^{*3}}\right)_{N=1}$

TABLE 1. Local data at a prime  $\mathfrak{q} \nmid 3$ 

2.2. Local theory at the prime  $\mathfrak{p} \mid 3$ . Recall that  $\mathfrak{p}$  is the unique prime dividing 3 in K. Proposition 2.6. For  $a \notin K_{\mathfrak{p}}^{*2}$ , one has  $\frac{\widehat{E}_{a,b}(K_{\mathfrak{p}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))} \subset \left(\frac{A_{\mathfrak{p}}^{*}}{A_{\mathfrak{p}}^{*3}}\right)_{N=1}$ .

*Proof.* The proof is similar to [JMS, Lemma 3.9].

We divide the study of image of local Kummer maps  $\delta_{\psi_{a,b},K_{\mathfrak{p}}}$  at  $\mathfrak{p}$  into three subcases  $3 \nmid ab, 3 \mid b$  and  $3 \mid a$ .

2.2.1. **Case-1:**  $3 \nmid ab$ . In this case, we find a p-minimal model for  $\widehat{E}_{a,b}$  and use it to compute the image of the Kummer map  $\delta_{\psi_{a,b},K_{\mathfrak{p}}}$ . Note that  $E_{a,b}: y^2 = x^3 + a(x-b)^2$  is p-minimal and has good reduction at p. By Tate's algorithm [Si2],

Note that  $E_{a,b}: y^2 = x^3 + a(x-b)^2$  is p-minimal and has good reduction at p. By Tate's algorithm [Si2], the p-minimal model for  $\widehat{E}_{a,b}$  is given by  $E' := E'_{a,b}$ 

$$E': Y^2 = X^3 + a_2 X^2 + b_2 X + c_2, (9)$$

where  $a_2 = -a(3+5\zeta_3)$ ,  $b_2 = a(a(-5+2\zeta_3)+18b\zeta_3^2)$  and  $c_2 = a(a^2(2+3\zeta_3)-2ab(1-9\zeta_3)-27b^2)$ . The following map  $\theta: E' \to E_{-27a,d}$  is then a  $G_K$ -isomorphism

$$\theta(x(P), y(P)) = (x'(P), y'(P)) := (\mathfrak{p}^2(\mathfrak{p}^2 x(P) - a - a\mathfrak{p}), \mathfrak{p}^6 y(P)).$$
(10)

The isogeny  $\Psi_{a,b} := \theta^{-1} \circ \psi_{a,b} : E_{a,b} \to E'$  is given by

$$\Psi_{a,b}(x,y) = (X,Y) := \left(\frac{\zeta_3(x^3 - a(3\mathfrak{p} + 5/3)x^2 - 4abx + 4ab^2)}{x^2}, \frac{-y(x^3 + 4abx - 8ab^2)}{x^3}\right).$$
(11)

Then, using formal groups of elliptic curves as in [Sc, pg. 92], we compute

$$Z := -\frac{X}{Y} = \zeta_3 \cdot \frac{x}{y} \left( 1 - \frac{a(3\mathfrak{p} + 5/3)x^2 + 8abx - 12ab^2}{x^3 + 4abx - 8ab^2} \right) = -\zeta_3 \cdot z + O(z^2).$$

Hence,  $\Psi'_{a,b}(0) = -\zeta_3$  and  $||\Psi'_{a,b}(0)||_{\mathfrak{p}}^{-1} = 1.$ 

**Proposition 2.7.** Assume that  $3 \nmid ab$ . We have

$$\begin{array}{ll} \text{(1)} & \text{If } a \notin K_{\mathfrak{p}}^{*2}, \text{ then } \left| \frac{\widehat{E}_{a,b}(K_{\mathfrak{p}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))} \right| = 1 \text{ and } \frac{\widehat{E}_{a,b}(K_{\mathfrak{p}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))} \subsetneq \left( \frac{A_{\mathfrak{p}}^{*}}{A_{\mathfrak{p}}^{*3}} \right)_{N=1}. \\ \text{(2)} & \text{If } a \in K_{\mathfrak{p}}^{*2}, \text{ then } \left| \frac{\widehat{E}_{a,b}(K_{\mathfrak{p}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))} \right| = 3 \text{ and } \frac{\widehat{E}_{a,b}(K_{\mathfrak{p}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))} \subsetneq \left( \frac{A_{\mathfrak{p}}^{*}}{A_{\mathfrak{p}}^{*3}} \right)_{N=1}. \end{array}$$

Proof. We know that  $c_{\mathfrak{p}}(E_{a,b}) = 1$ , as  $E_{a,b}$  has good reduction at  $\mathfrak{p}$ ; thus  $c_{\mathfrak{p}}(E') = 1$ . Also, we have  $||\Psi'_{a,b}(0)||_{\mathfrak{p}}^{-1} = 1$ . Using the Schaefer's formula (8), we obtain  $\left|\frac{E'(K_{\mathfrak{p}})}{\Psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))}\right| = \begin{cases} 1, & \text{if } a \notin K_{\mathfrak{p}}^{*2}, \\ 3, & \text{if } a \in K_{\mathfrak{p}}^{*2}, \end{cases}$  Now,  $\left|\frac{\widehat{E}_{a,b}(K_{\mathfrak{p}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))}\right| = \left|\frac{\theta(E'(K_{\mathfrak{p}}))}{\theta \circ \Psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))}\right| = \left|\frac{E'(K_{\mathfrak{p}})}{\Psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))}\right| = \left|\frac{1}{\theta \circ \Psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))}\right| = \left|\frac{1}{\Phi \circ \Psi_{a,b}(E_{a,b}(K_{\mathfrak{p}))}}\right| = \left|\frac{1}{\Phi \circ \Psi_{a,b}(E_{a,b}(K_{\mathfrak{p}))}\right| = \left$ 

When  $a \notin K_{\mathfrak{p}}^{*2}$ , then  $\left| \left( \frac{A_{\mathfrak{p}}^{*}}{A_{\mathfrak{p}}^{*3}} \right)_{N=1} \right| = 9$  by Prop. 1.4 and the strict inequality in (1) follows. When  $a \in K_{\mathfrak{p}}^{*2}$ , then  $\left| \left( \frac{A_{\mathfrak{p}}^{*}}{A_{\mathfrak{p}}^{*3}} \right)_{N=1} \right| = 27$  by Prop. 1.4. Thus, to complete the proof, it suffices to show, by Lemma 1.8, that  $v_{\mathfrak{p}}(\delta_{\psi_{a,b},K_{\mathfrak{p}}}(P')) \equiv 0 \pmod{3}$  for all  $P' \in \widehat{E}_{a,b}(K_{\mathfrak{p}})$ . We now prove this by showing that for any  $P \in E'(K_{\mathfrak{p}})$  we have  $v_{\mathfrak{p}}(\delta_{\psi_{a,b},K_{\mathfrak{p}}}(\theta(P))) \equiv 0 \pmod{3}$ .

Consider a point  $P \in E'(K_{\mathfrak{p}})[\widehat{\Psi}_{a,b}]$  and recall that  $3 \nmid ab$  and  $\mathfrak{p} \mid 3$ . Hence,  $v_{\mathfrak{p}}(a) = v_{\mathfrak{p}}(d) = 0$  and  $v_{\mathfrak{p}}(2\mathfrak{p}^{3}d\sqrt{a}) = 3$ . Now, note here that  $\theta(P) = (0, \pm \mathfrak{p}^{3}d\sqrt{a}) \in \widehat{E}_{a,b}(K_{\mathfrak{p}})[\widehat{\psi}_{a,b}]$  and we have  $v_{\mathfrak{p}}(\delta_{\psi_{a,b},K_{\mathfrak{p}}}(\theta(P))) \equiv 0 \pmod{3}$ .

Next, take  $P \notin E'(K_{\mathfrak{p}})[\widehat{\Psi}_{a,b}]$  and  $\theta(P) = (x'(P), y'(P))$ . If  $v_{\mathfrak{p}}(y'(P)) \neq v_{\mathfrak{p}}(\mathfrak{p}^{3}\sqrt{a}(x'(P) - d))$ , then as in the proof of [JMS, Prop. 3.7(1)], we get  $v_{\mathfrak{p}}(\delta_{\psi_{a,b},K_{\mathfrak{p}}}((x'(P), y'(P)))) \equiv 0 \pmod{3}$ . So, we only consider the case  $v_{\mathfrak{p}}(y'(P)) = v_{\mathfrak{p}}(\mathfrak{p}^{3}\sqrt{a}(x'(P) - d)) = n$ , say.

Further, if  $v_{\mathfrak{p}}(x'(P)) = 0$ , then  $v_{\mathfrak{p}}(y'(P)) = 0$  but  $v_{\mathfrak{p}}(\mathfrak{p}^3\sqrt{a}(x'(P)-d)) \ge 3$ , a contradiction. Now if  $v_{\mathfrak{p}}(x'(P)) < 0$ , then  $x'(P) = \mathfrak{p}^2(\mathfrak{p}^2x(P) - a - \mathfrak{p}a)$  implies that  $v_{\mathfrak{p}}(x(P)) < 0$ . This gives  $v_{\mathfrak{p}}(x(P)) = -2r$  for some  $r \ge 3$ . Thus,  $v_{\mathfrak{p}}(\mathfrak{p}^3\sqrt{a}(x'(P)-d)) = -2r + 7 \ne -3r + 6 = v_{\mathfrak{p}}(y'(P))$ . This is a contradiction.

Finally, the case  $x'(P) \neq 0$  and  $v_{\mathfrak{p}}(x'(P)) > 0$  and  $v_{\mathfrak{p}}(y'(P)) = v_{\mathfrak{p}}(\mathfrak{p}^3\sqrt{a}(x'(P)-d))$  can be proved in a similar manner as in [JMS, Lemma 3.9].

**Proposition 2.8.** For  $3 \nmid ab$  and  $a \in K_{\mathfrak{p}}^{*2}$ , we have  $\frac{\widehat{E}_{a,b}(K_{\mathfrak{p}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))} = \frac{V_3}{A_{\mathfrak{p}}^{*3}}$ .

Proof. Note that  $|\text{Im} (\delta_{\psi_{a,b},K_{\mathfrak{p}}} \circ \theta)| = 3$  by Prop. 2.7. Let  $P = (x(P), y(P)) \in E'(K_{\mathfrak{p}}), \theta(P) = (x'(P), y'(P))$ and  $(C, D) := \delta_{\psi_{a,b},K_{\mathfrak{p}}} \circ \theta(P) = \delta_{\psi_{a,b},K_{\mathfrak{p}}}(x'(P), y'(P))$ . Then  $(C, D) \in (A_{\mathfrak{p}}^*/A_{\mathfrak{p}}^{*3})_{N=1}$  by Prop. 2.7. Recall from Definition 1.5 that  $V_3 = \{(u_1, u_2) \in V_{K_{\mathfrak{p}}} \times V_{K_{\mathfrak{p}}} \mid u_1 u_2 \in \mathcal{O}_{K_{\mathfrak{p}}}^{*3}\}$  and  $|V_3/A_{\mathfrak{p}}^{*3}| = 3$  from Prop. 1.7. Thus, it reduces to show that Im  $(\delta_{\psi_{a,b},K_{\mathfrak{p}}} \circ \theta) \subset V_3/A_{\mathfrak{p}}^{*3}$ . By Lemma 1.8,  $(v_{\mathfrak{p}}(C), v_{\mathfrak{p}}(D)) \equiv 0 \pmod{3}$ . Also, as elements of  $K^*/K^{*3} \times K^*/K^{*3}$ ,

$$(\overline{C},\overline{D}) = (\overline{\mathfrak{p}^{-\upsilon_{\mathfrak{p}}(C)}u_1^3C}, \ \overline{\mathfrak{p}^{-\upsilon_{\mathfrak{p}}(D)}u_2^3D}) \ \text{ for any } \ u_1, u_2 \in \mathcal{O}_{K_{\mathfrak{p}}}^*.$$

This gives  $\mathfrak{p}^{-v_{\mathfrak{p}}(C)}u_1^3C \cdot \mathfrak{p}^{-v_{\mathfrak{p}}(D)}u_2^3D \in \mathcal{O}_{K_{\mathfrak{p}}}^{*3}$ . Hence, it is enough to show that there exists some  $u \in \mathcal{O}_{K_{\mathfrak{p}}}^*$  such that either  $\overline{\mathfrak{p}^{-v_{\mathfrak{p}}(C)}u^3C} \in (\mathcal{O}_{K_{\mathfrak{p}}}^*/U_{K_{\mathfrak{p}}}^3)^3$  or  $\overline{\mathfrak{p}^{-v_{\mathfrak{p}}(D)}u^3D} \in (\mathcal{O}_{K_{\mathfrak{p}}}^*/U_{K_{\mathfrak{p}}}^3)^3$ .

For  $(x'(P), y'(P)) \notin \widehat{E}_{a,b}(K_{\mathfrak{p}})[\widehat{\psi}_{a,b}]$ , recall that  $C = y'(P) - \mathfrak{p}^3 \sqrt{a}(x'(P) - d)$  and  $D = y'(P) + \mathfrak{p}^3 \sqrt{a}(x'(P) - d)$ . If  $v_{\mathfrak{p}}(x(P)) \geq 0$ , then  $v_{\mathfrak{p}}(y(P)) \geq 0$ . This implies,  $v_{\mathfrak{p}}(y'(P)) \geq 6$  and  $v_{\mathfrak{p}}(x'(P)) = 2$ , which shows that  $v_{\mathfrak{p}}(C) = v_{\mathfrak{p}}(D) = 3$ . Then

$$\mathfrak{p}^{-3}C = \mathfrak{p}^3 y(P) - \sqrt{a}(\mathfrak{p}^4 x(P) - \mathfrak{p}^2 a - \mathfrak{p}^3 a - d) \equiv a\sqrt{a}\mathfrak{p}^2 + \sqrt{a}d \equiv a\sqrt{a}(\mathfrak{p}^2 + 4) \equiv a\sqrt{a} \pmod{\mathfrak{p}^3},$$

where the last two congruences use the facts that  $d \equiv 4a \pmod{\mathfrak{p}^3}$  and  $\mathfrak{p}^2 + 4 \equiv 1 \pmod{\mathfrak{p}^3}$ . Hence, we obtain that  $\overline{\mathfrak{p}^{-\upsilon_{\mathfrak{p}}(C)}(\sqrt{a})^{-3}C} = \overline{1} \in \left(\mathcal{O}_{K_{\mathfrak{p}}}^*/U_{K_{\mathfrak{p}}}^3\right)^3$ .

For the rest of the proof, we deal with the case  $v_{\mathfrak{p}}(x(P)) < 0$ . Thus,  $v_{\mathfrak{p}}(x(P)) = -2r$  and  $v_{\mathfrak{p}}(y(P)) = -3r$ for some  $r \in \mathbb{N}$ . By the theory of formal groups, there exists  $0 \neq z = \mathfrak{p}^r w$  with  $w \in \mathcal{O}_{K_{\mathfrak{p}}}^*$  such that  $P(z) = P = (x(P), y(P)) := (x(P(z)), y(P(z))) \in E'(K_{\mathfrak{p}})$ . We divide this in two cases :  $r \ge 2$  and r = 1.

(1) Assume that  $v_{\mathfrak{p}}(x(P(z))) = -2r$ , where  $0 \neq z = \mathfrak{p}^r w$  with  $r \geq 2$ . Therefore, one has  $v_{\mathfrak{p}}(x'(P(z))) = -2r + 4$  and  $v_{\mathfrak{p}}(y'(P(z))) = -3r + 6$ , which in turn implies that  $v_{\mathfrak{p}}(C) = v_{\mathfrak{p}}(D) = -3r + 6 \le 0.$  Now,  $\mathfrak{p}^{3r-6}w^{3}C = z^{3}y(P(z)) - w\sqrt{a}\mathfrak{p}^{r+1}z^{2}x(P(z)) + w^{3}a\sqrt{a}\mathfrak{p}^{3r-1} + w^{3}a\sqrt{a}\mathfrak{p}^{3r} + dw^{3}\sqrt{a}\mathfrak{p}^{3r-3}$  $\equiv z^3 y(P(z)) \pmod{\mathfrak{p}^3} \equiv -1 \pmod{\mathfrak{p}^3}.$ 

Thus we have  $\overline{\mathfrak{p}^{-\upsilon_{\mathfrak{p}}(C)}w^{-3}C} = -\overline{1} \in \left(\mathcal{O}_{K_{\mathfrak{p}}}^*/U_{K_{\mathfrak{p}}}^3\right)^3$  in this case. (2) Now assume that r = 1 i.e.  $\upsilon_{\mathfrak{p}}(x(P(z))) = -2$ .

(a) First consider the case  $x(P(z)) \neq \frac{1}{p^2}(a + pa)$ . In this case,  $v_{\mathfrak{p}}(y'(P(z))) = 3$  and  $v_{\mathfrak{p}}(\mathfrak{p}^3\sqrt{a}(x'(P(z)) - d)) = 3$ . Thus, at least one of  $v_{\mathfrak{p}}(C)$  and  $v_{\mathfrak{p}}(D)$  is 3. Further,  $w\sqrt{a} \equiv \pm 1 \pmod{\mathfrak{p}}$  and we only prove the case  $w\sqrt{a} \equiv -1 \pmod{\mathfrak{p}}$  here. The case  $w\sqrt{a} \equiv 1 \pmod{\mathfrak{p}}$  can be proved in a similar manner. Let  $z = \mathfrak{p}w$  and  $w\sqrt{a} \equiv -1 \pmod{\mathfrak{p}}$ . Then

$$\begin{aligned} \mathfrak{p}^{-3}w^3C &\equiv z^3y(P(z)) - w\sqrt{a}\mathfrak{p}^2 z^2 x(P(z)) + w^3 a\sqrt{a}\mathfrak{p}^2 + dw^3\sqrt{a} \pmod{\mathfrak{p}^3} \\ &\equiv -1 + a_2\mathfrak{p}^2w^2 - w\sqrt{a}\mathfrak{p}^2 + w^3 a\sqrt{a}\mathfrak{p}^2 + 4w^3 a\sqrt{a} \pmod{\mathfrak{p}^3} \\ &\equiv -1 + 4w^3 a\sqrt{a} + \mathfrak{p}^2(a_2w^2 + w^3 a\sqrt{a} - w\sqrt{a}) \pmod{\mathfrak{p}^3} \\ &\equiv -5 + a_2\mathfrak{p}^2 \pmod{\mathfrak{p}^3} \equiv -5 + a\mathfrak{p}^2 \pmod{\mathfrak{p}^3} \equiv -5 + \mathfrak{p}^2 \pmod{\mathfrak{p}^3} \equiv 1 \pmod{\mathfrak{p}^3}. \end{aligned}$$

Thus, we obtain  $\overline{\mathfrak{p}^{-\upsilon_{\pi}(C)}w^3C} = \overline{1} \in \left(\mathcal{O}_{K_{\mathfrak{p}}}^*/U_{K_{\mathfrak{p}}}^3\right)^3$  in this case. (b) Finally, consider the case  $x(P(z)) = \frac{1}{p^2}(a + pa)$ .

In this situation,  $\theta(P(z)) = (x'(P(z), y'(P(z)) \text{ satisfies } x'(P(z)) = 0 \text{ and therefore we have } \delta_{\psi_{a,b},K_{\mathfrak{p}}} \circ \theta(P(z)) = (\pm 2\mathfrak{p}^{3}d\sqrt{a}, \pm \frac{1}{2\mathfrak{p}^{3}d\sqrt{a}}).$  Now note that

$$\mathfrak{p}^{-3}(\pm 2\mathfrak{p}^3 d\sqrt{a}) = \pm 2d\sqrt{a} \equiv \pm 8a\sqrt{a} \pmod{\mathfrak{p}^3} = (\pm 2\sqrt{a})^3 \pmod{\mathfrak{p}^3}.$$

Therefore, in this case, we obtain  $\mathfrak{p}^{-\upsilon_{\mathfrak{p}}(C)}(\pm 2\sqrt{a})^{-3}C = 1 \in (\mathcal{O}_{K_{\mathfrak{p}}}^*/U_{K_{\mathfrak{p}}}^s)^{\sim}$ . 

2.2.2. Case-2:  $3 \mid b$ . In this case, we assume that  $3 \mid b$  (this implies  $3 \nmid a$ ) and study the relationship

between the groups  $\frac{\widehat{E}_{a,b}(K_{\mathfrak{p}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))}$ ,  $\frac{V_3}{A_{\mathfrak{p}}^{*3}}$  and  $\left(\frac{A_{\mathfrak{p}}^*}{A_{\mathfrak{p}}^{*3}}\right)_{N=1}^{N=1}$ . By Tate's algorithm,  $E_{a,b}: y^2 = x^3 + a(x-b)^2$  is  $\mathfrak{p}$ -minimal and has multiplicative reduction of type  $I_n$  at  $\mathfrak{p}$ , where  $n = 3v_{\mathfrak{p}}(b)$ . Thus, the Tamagawa number  $c_{\mathfrak{p}}(E_{a,b}) = \begin{cases} 2, & \text{if } a \notin K_{\mathfrak{p}}^{*2}, \\ 3 v_{\mathfrak{p}}(b), & \text{if } a \in K_{\mathfrak{p}}^{*2}. \end{cases}$ 

Observe that the equation of  $\widehat{E}_{a,b}: y^2 = x^3 - 27a(x-d)^2$  is not p-minimal and by Tate's algorithm, the  $\mathfrak{p}$ -minimal model for  $\widehat{E}_{a,b}$  is given by  $E': Y^2 = X^3 + a_2X^2 + b_2X + c_2$ , where  $a_2, b_2, c_2$  are as defined in (9). As in the previous case,  $\theta: E' \to \widehat{E}_{a,b}$  as in (10) defines a K-isomorphism and the isogeny  $\Psi_{a,b}: E_{a,b} \to E'$ is again given by (11).

Moreover, E' has multiplicative reduction of type  $I_n$  at  $\mathfrak{p}$ , where  $n = v_{\mathfrak{p}}(b)$ . Thus,  $c_{\mathfrak{p}}(E') = 2$ , if  $a \notin K_{\mathfrak{p}}^{*2}$  and  $c_{\mathfrak{p}}(E') = v_{\mathfrak{p}}(b)$ , if  $a \in K_{\mathfrak{p}}^{*2}$ . In particular, note that  $c_{\mathfrak{p}}(E_{a,b}) = 3c_{\mathfrak{p}}(\widehat{E}_{a,b})$ , when  $a \in K_{\mathfrak{p}}^{*2}$ . Using formal groups, we have that  $\Psi'_{a,b}(0) = -\zeta_3$  and hence,  $||\Psi'_{a,b}(0)||_{\mathfrak{p}}^{-1} = 1$ .

**Lemma 2.9.** For  $3 \mid b$ , we have (whether or not  $a \in K_{\mathfrak{p}}^{*2}$ ) that  $\left|\frac{\widehat{E}_{a,b}(K_{\mathfrak{p}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))}\right| = 1$  and thus trivially,  $\frac{\widehat{E}_{a,b}(K_{\mathfrak{p}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))} \subsetneq \left(\frac{A_{\mathfrak{p}}^{*}}{A_{\mathfrak{p}}^{*3}}\right)_{N=1}$ .

*Proof.* This follows easily from the above discussion and using (8).

2.2.3. **Case-3:**  $3 \mid a$ . We finally consider the case  $3 \mid a$ ; this implies  $3 \nmid b$ . We study the bounds for  $\frac{\widehat{E}_{a,b}(K_{\mathfrak{p}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))}$  using  $\frac{V_3}{A_{\mathfrak{p}}^{*3}}$  and  $\left(\frac{A_{\mathfrak{p}}^*}{A_{\mathfrak{p}}^{*3}}\right)_{N=1}$ .

If  $a \notin K_{\mathfrak{p}}^{*2}$ , then we know that  $\{1\} = \frac{V_3}{A_{\mathfrak{p}}^{*3}} \subset \frac{\hat{E}_{a,b}(K_{\mathfrak{p}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))} \subset \left(\frac{A_{\mathfrak{p}}^*}{A_{\mathfrak{p}}^{*3}}\right)_{N=1}$  from Props. 2.6 and 1.7. From now on, we take  $a \in K_{\mathfrak{p}}^{*2}$ . First, we consider the special case when  $3^6 \mid d = 4a + 27b$ ; then 27  $\mid\mid a$ .

From now on, we take  $a \in K_{\mathfrak{p}}^{-2}$ . First, we consider the special case when  $3^{\circ} \mid d = 4a + 27b$ ; then  $27 \mid a$ . Setting  $a_1 = -\frac{a}{27}$  and  $b_1 = \frac{d}{3^{\circ}}$ , we see that  $\hat{E}_{a,b} \cong E_{a_1,b_1}$  and  $E_{a,b} = \hat{E}_{a_1,b_1}$ . Note here that  $3 \nmid a_1$  and  $a_1 \in K_{\mathfrak{p}}^{*2}$ . Then following a similar proof as in Case 1 §2.2.1 and Case 2 §2.2.2, it follows that:

(1) If 3<sup>6</sup> || *d* or equivalently, 
$$3 \nmid b_1$$
, we see that  $\frac{V_3}{A_{\mathfrak{p}^3}^*} \cong \frac{E_{a,b}(K_{\mathfrak{p}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))} \subset \left(\frac{A_{\mathfrak{p}}}{A_{\mathfrak{p}^3}^*}\right)_{N=1}$ .  
(2) If 3<sup>7</sup> | *d* or equivalently 3 | *b*<sub>1</sub>, then  $\{1\} = \frac{\hat{E}_{a,b}(K_{\mathfrak{p}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{p}}))} \subsetneq \left(\frac{A_{\mathfrak{p}}^*}{A_{\mathfrak{p}^3}^*}\right)_{N=1}$ .

We summarize this in the following remark:

**Remark 2.10.** For  $a \notin K_{\mathfrak{p}}^{*2}$ , we have  $|V_3/A_{\mathfrak{p}}^{*3}| = 1$  by Prop. 1.7. Then

(1) If 
$$3 \nmid a$$
, then  $\frac{V_3}{A_{\mathfrak{p}}^{*3}} = \delta_{\psi_{a,b},K_{\mathfrak{p}}}(\widehat{E}_{a,b}(K_{\mathfrak{p}})) \subsetneq \left(\frac{A_{\mathfrak{p}}}{A_{\mathfrak{p}}^{*3}}\right)_{N=1}$  by Props. 2.7(1) and 2.9(1).  
(2) If  $3 \mid a$ , then  $\frac{V_3}{A_{\mathfrak{p}}^{*3}} \subset \delta_{\psi_{a,b},K_{\mathfrak{p}}}(\widehat{E}_{a,b}(K_{\mathfrak{p}})) \subset \left(\frac{A_{\mathfrak{p}}}{A_{\mathfrak{p}}^{*3}}\right)_{N=1}$  by Prop. 2.6.

For  $a \in K_{\mathfrak{p}}^{*2}$ , we have  $|V_3/A_{\mathfrak{p}}^{*3}| = 3$  by Prop. 1.7. Then

(1) If 
$$3 \nmid ab$$
, then  $\frac{V_3}{A_{\mathfrak{p}}^{*3}} = \delta_{\psi_{a,b},K_{\mathfrak{p}}}(\widehat{E}_{a,b}(K_{\mathfrak{p}})) \subsetneq \left(\frac{A_{\mathfrak{p}}^*}{A_{\mathfrak{p}}^{*3}}\right)_{N=1}$  by Prop. 2.8.

(2) If 
$$3 \mid b$$
, then  $\{1\} = \delta_{\psi_{a,b},K_{\mathfrak{p}}}(\widehat{E}_{a,b}(K_{\mathfrak{p}})) \subsetneq \left(\frac{A_{\mathfrak{p}}}{A_{\mathfrak{p}}^{*3}}\right)_{N=1}$  by Prop. 2.9(2).

(3) If 27 || a and 3<sup>6</sup> || d, then 
$$\frac{V_3}{A_{\mathfrak{p}}^{*3}} \cong \delta_{\psi_{a,b},K_{\mathfrak{p}}}(E_{a,b}(K_{\mathfrak{p}})) \subsetneq \left(\frac{A_{\mathfrak{p}}}{A_{\mathfrak{p}}^{*3}}\right)_{N=1}$$
 by Props. 2.7(2) and 2.8.

(4) If 27 || a and 3<sup>7</sup> | d, then 
$$\{1\} = \delta_{\psi_{a,b},K_{\mathfrak{p}}}(\widehat{E}_{a,b}(K_{\mathfrak{p}})) \subsetneq \left(\frac{A_{\mathfrak{p}}^*}{A_{\mathfrak{p}}^*}\right)_{N=1}$$
 by Lemma 2.9(2).

3. Bounds for the 
$$\psi_{a,b}$$
- and 3-Selmer Groups of  $E_{a,b}/K$ 

In this subsection, we give bounds on  $\dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K)$  and  $\dim_{\mathbb{F}_3} \operatorname{Sel}^3(E_{a,b}/K)$  in terms of  $\dim_{\mathbb{F}_3} Cl_S(L)[3]$  for certain sets S that we define shortly.

**Remark 3.1.** As explained in §1, we have  $\operatorname{Sel}^{\widehat{\psi}_{a,b}}(\widehat{E}_{a,b}/K) \cong \operatorname{Sel}^{\psi_{-27a,d}}(E_{-27a,d}/K)$  using  $\widehat{\psi}_{a,b} = \lambda_{27} \circ \psi_{-27a,d}$ . Further, writing  $gcd(-27a,d) = t^2g$ , where g is square-free in K,  $\widehat{E}_{a,b}$  can be identified with  $E_{-27a/t^2,d/t^2}$  up to an isomorphism defined over K. Thus to compute bounds of  $\dim_{\mathbb{F}_3} \operatorname{Sel}^{\widehat{\psi}_{a,b}}(\widehat{E}_{a,b}/K)$ , it suffices to compute the same for  $\operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K)$  for all  $a, b \in \mathbb{Z}$  with gcd(a, b) square-free in K.

Recall that  $a, b \in \mathbb{Z}$  with gcd(a, b) square-free in K; In particular,  $3 \nmid gcd(a, b)$ . Set

$$T_{2,1} := \begin{cases} \{2\mathcal{O}_K\}, & \text{if } a \in K_2^{*2}, 2 \nmid a \text{ and } 4 \nmid b, \\ \emptyset, & \text{otherwise.} \end{cases} \quad T_{3,1} := \begin{cases} \{\mathfrak{p}\}, & \text{if } a \in K_\mathfrak{p}^{*2}, \mathfrak{p} \mid a \text{ and } v_\mathfrak{p}(a) \neq 6, \\ \{\mathfrak{p}\}, & \text{if } a \in K_\mathfrak{p}^{*2}, v_\mathfrak{p}(a) = 6 \text{ and } v_\mathfrak{p}(d) < 12, \\ \emptyset, & \text{otherwise.} \end{cases}$$

$$T_{2,2} := \begin{cases} \{2\mathcal{O}_K\}, & \text{if } a \in K_2^{*2}, 2 \nmid a \text{ and } 8 \mid b, \\ \emptyset, & \text{otherwise.} \end{cases} \qquad T_{3,2} := \begin{cases} \{\mathfrak{p}\}, & \text{if } a \in K_\mathfrak{p}^{*2}, \mathfrak{p} \mid b \text{ and } \mathfrak{p} \nmid a, \\ \emptyset, & \text{otherwise.} \end{cases}$$
$$T_{2,3} := \begin{cases} \{2\mathcal{O}_K\}, & \text{if } a \in K_2^{*2}, 2 \nmid a \text{ and } 4 \mid \mid b, \\ \emptyset, & \text{otherwise.} \end{cases} \qquad T_{3,3} := \begin{cases} \{\mathfrak{p}\}, & \text{if } a \in K_\mathfrak{p}^{*2} \text{ and } v_\mathfrak{p}(d) > 12, \\ \emptyset, & \text{otherwise.} \end{cases}$$

**Definition 3.2.** We define the following subsets of  $\Sigma_K$ , the set of all finite primes of K:

$$S_{1} = \left\{ \mathfrak{q} \in \Sigma_{K} \setminus \{\mathfrak{p}\} \mid a \in K_{\mathfrak{q}}^{*2}, \ \mathfrak{q} \mid a \text{ and } v_{\mathfrak{q}}(4ab^{2}) \not\equiv 0 \pmod{6} \right\} \cup T_{2,1} \cup T_{3,1},$$

$$S_{2} = \left\{ \mathfrak{q} \in \Sigma_{K} \setminus \{\mathfrak{p}\} \mid a \in K_{\mathfrak{q}}^{*2}, \ \mathfrak{q} \mid b \text{ and } \mathfrak{q} \nmid 2a \right\} \cup T_{2,2} \cup T_{3,2} \text{ and}$$

$$S_{3} = \left\{ \mathfrak{q} \in \Sigma_{K} \setminus \{\mathfrak{p}\} \mid a \in K_{\mathfrak{q}}^{*2}, \ \mathfrak{q} \mid d \text{ and } \mathfrak{q} \nmid 2a \right\} \cup T_{2,3} \cup T_{3,3}.$$

Observe that  $S_1$ ,  $S_2$  and  $S_3$  are pairwise disjoint finite sets of primes of K.

**Definition 3.3.** Let L be a field. Define the following subsets of  $\Sigma_L$ :

 $S_{1,2}(L) := \{ \mathfrak{Q} \in \Sigma_L \mid \mathfrak{Q} \cap \mathcal{O}_K \in S_1 \cup S_2 \} \text{ and } S_{1,3}(L) := \{ \mathfrak{Q} \in \Sigma_L \mid \mathfrak{Q} \cap \mathcal{O}_K \in S_1 \cup S_3 \}.$ 

**Remark 3.4.** Using Tate's algorithm, we observe that if  $q \in S_1 \cup S_2$  (resp.  $q \in S_1 \cup S_3$ ) including q = p, then  $E_{a,b}$  and  $\hat{E}_{a,b}$  have either split multiplicative reduction or additive reduction of Kodaira type IV or IV\* at  $\mathfrak{q}$ .

The  $\mathbb{F}_3$ -subspaces  $M(S_1 \cup S_2, a)$ ,  $N(S_1 \cup S_3, a)$  and  $N'(S_1 \cup S_3, a)$  of  $(L^*/L^{*3})_{N-1}$  are defined in Prop. 1.13.

**Theorem 3.5.** We have the inclusion of  $\mathbb{F}_3$ -modules and the inequalities of their 3-ranks:

(1) If  $a \notin K^{*2}$ , then  $M(S_1 \cup S_2, a) \subset \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \subset N(S_1 \cup S_3, a)$ . Hence, we get that

$$h_{S_{1,2}(L)}^3 \le \dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \le h_{S_{1,3}(L)}^3 + |S_{1,3}(L)| + 2.$$

(2) If 
$$a \in K^{*2}$$
, then  $\operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \subset N'(S_1 \cup S_3, a)$ . Hence,  $\dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \leq |S_1 \cup S_3| + 1$ .

*Proof.* We start with the case  $a \notin K^{*2}$ . From Props. 2.4, 2.5 and Table 1 (for  $\mathfrak{q} \nmid 3$ ) and Props. 2.6, 2.7, 2.9 and Remark 2.10 (for  $\mathfrak{p} \mid 3$ ), it follows that Im  $\delta_{\psi_{a,b},K_{\mathfrak{q}}} \subset (A^*_{\mathfrak{q}}/A^{*3}_{\mathfrak{q}})_{N=1}$  for all  $\mathfrak{q} \notin S_1 \cup S_3$ . Thus, by Prop. 1.13,  $\operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \subset N(S_1 \cup S_3, a).$ 

On the other hand, it follows from Props. 2.4, 2.5 and Table 1, that  $(A_{\mathfrak{q}}^*/A_{\mathfrak{q}}^{*3})_{N=1} \subset \operatorname{Im} \delta_{\psi_{a,b},K_{\mathfrak{q}}}$  for  $\mathfrak{q} \notin S_1 \cup S_2 \cup \{\mathfrak{p}\}$ . Further, from Props. 2.7, 2.9, Remark 2.10, it follows that  $V_3/A_{\mathfrak{p}}^{*3} \subset \operatorname{Im} \delta_{\psi_{a,b},K_{\mathfrak{p}}}$ , if  $\mathfrak{p} \notin S_1 \cup S_2$ . Thus, from Prop. 1.13, we get  $M(S_1 \cup S_2, a) \subset \mathrm{Sel}^{\psi_{a,b}}(E_{a,b}/K)$ .

Now if  $a \in K^{*2}$ , then from Prop. 2.5 and Table 1 (for  $\mathfrak{q} \nmid 3$ ) and Props. 2.7, 2.9 and Remark 2.10 (for  $\mathfrak{p} \mid 3$ ), it follows that Im  $(\delta_{\psi_{a,b},K_{\mathfrak{q}}}) \subset (A_{\mathfrak{q}}^*/A_{\mathfrak{q}}^{*3})_{N=1}$  for all  $\mathfrak{q} \notin S_1 \cup S_3$ . Hence, by Prop. 1.13,  $\operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \subset N'(S_1 \cup S_3, a).$ 

In both cases (1) and (2), the statements about ranks follow directly from Prop. 1.12.

**Remark 3.6.** Following Remark 3.1, the bounds for  $\operatorname{Sel}^{\widehat{\psi}_{a,b}}(\widehat{E}_{a,b}/K)$  are computed similarly:

- (1) If  $a \notin K^{*2}$ , then  $h^3_{S_{1,3}(L)} \leq \dim_{\mathbb{F}_3} \operatorname{Sel}^{\widehat{\psi}_{a,b}}(\widehat{E}_{a,b}/K) \leq h^3_{S_{1,2}(L)} + |S_{1,2}(L)| + 2.$
- (2) If  $a \in K^{*2}$ , then  $\dim_{\mathbb{F}_3} \operatorname{Sel}^{\widehat{\psi}_{a,b}}(\widehat{E}_{a,b}/K) \leq |S_1 \cup S_2| + 1$ .

Theorem 3.5 and Remark 3.6 have the following consequence on  $\text{Sel}^3(E_{a,b}/K)$ :

**Corollary 3.7.** If  $a \notin K^{*2}$ , then from [SS, Lemma 6.1] we have the exact sequence

$$0 \to \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \to \operatorname{Sel}^{3}(E_{a,b}/K) \to \operatorname{Sel}^{\widehat{\psi}_{a,b}}(\widehat{E}_{a,b}/K) \to \frac{\operatorname{III}(\widehat{E}_{a,b}/K)[\widehat{\psi}_{a,b}]}{\psi_{a,b}(\operatorname{III}(E_{a,b}/K)[3]} \to 0$$
(12)

and it follows that

$$\max \{h_{S_{1,2}(L)}^3, rk \; E_{a,b}(K)\} \leq \dim_{\mathbb{F}_3} \operatorname{Sel}^3(E_{a,b}/K) \leq h_{S_{1,2}(L)}^3 + h_{S_{1,3}(L)}^3 + |S_{1,2}(L)| + |S_{1,3}(L)| + 4.$$
  
If  $a \in K^{*2}$ , then  $rk \; E_{a,b}(K) \leq \dim_{\mathbb{F}_3} \operatorname{Sel}^3(E_{a,b}/K) \leq |S_1 \cup S_2| + |S_1 \cup S_3| + 2.$ 

**Theorem 3.8** (Refined Bounds). If  $a \notin K^{*2}$  and  $3 \nmid a$ , then we have that

$$\max \{h_{S_{1,2}(L)}^3, h_{S_{1,3}(L)}^3 + |S_3| - |S_2| - 1\} \le \dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \\ \le \min \{h_{S_{1,2}(L)}^3 + |S_{1,2}(L)| + |S_3| - |S_2| + 1, h_{S_{1,3}(L)}^3 + |S_{1,3}(L)| + 2\}.$$

In particular, if  $S_1 \cup S_2 \cup S_3$  is empty, then  $\dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \in \{h_L^3, h_L^3 + 1\}.$ 

*Proof.* Using [Ca2, Eq. (1.22), (1.26) and (3.4)], we get that

$$\frac{|\operatorname{Sel}^{\psi_{a,b}}(\widehat{E}_{a,b}/K)|}{|\operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K)|} = \frac{|\widehat{E}_{a,b}(K)[\widehat{\psi}_{a,b}]|}{|E_{a,b}(K)[\psi_{a,b}]|} \cdot \frac{|E_{a,b}(\mathbb{C})[\psi_{a,b}]|}{|\widehat{E}_{a,b}(\mathbb{C})/\psi_{a,b}(E_{a,b}(\mathbb{C}))|} \cdot \prod_{\mathfrak{q}\in\Sigma_K} \frac{|E_{a,b}(K_{\mathfrak{q}})[\psi_{a,b}]|}{|\widehat{E}_{a,b}(K_{\mathfrak{q}})/\psi_{a,b}(E_{a,b}(K_{\mathfrak{q}}))|}.$$
 (13)

Also  $\left|\frac{\widehat{E}_{a,b}(K_{\mathfrak{q}})}{\psi_{a,b}(E_{a,b}(K_{\mathfrak{q}}))}\right| = \frac{||\psi'_{a,b}(0)||_{\mathfrak{q}}^{-1} \cdot |E_{a,b}(K_{\mathfrak{q}})[\psi_{a,b}]| \cdot c_{\mathfrak{q}}(\widehat{E}_{a,b})}{c_{\mathfrak{q}}(E_{a,b})}$ , where  $||\psi'_{a,b}(0)||_{\mathfrak{q}}^{-1} = 1$  for any  $\mathfrak{q} \in \Sigma_K$  by (8). Now, note that  $|E_{a,b}(\mathbb{C})[\psi_{a,b}]| = 3$  and  $\psi_{a,b} : E_{a,b}(\mathbb{C}) \to \widehat{E}_{a,b}(\mathbb{C})$  is surjective. Thus, (13) becomes  $\tfrac{|\mathrm{Sel}^{\hat{\psi}_{a,b}}(\hat{E}_{a,b}/K)|}{|\mathrm{Sel}^{\psi_{a,b}}(E_{a,b}/K)|} = 3 \cdot \prod_{\mathfrak{q} \in \Sigma_K} \tfrac{c_\mathfrak{q}(E_{a,b})}{c_\mathfrak{q}(\hat{E}_{a,b})}$ 

Define  $S' := \{ \mathfrak{q} \in \Sigma_K \mid c_\mathfrak{q}(E_{a,b}) = 3 \ c_\mathfrak{q}(\widehat{E}_{a,b}) \}$  and  $S'' := \{ \mathfrak{q} \in \Sigma_K \mid c_\mathfrak{q}(\widehat{E}_{a,b}) = 3 \ c_\mathfrak{q}(E_{a,b}) \}$ . Then we have  $\frac{|\operatorname{Sel}^{\hat{\psi}_{a,b}}(\hat{E}_{a,b}/K)|}{|\operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K)|} = 3^{|S'|-|S''|+1}$ . Under the assumption  $3 \nmid a$ , it is easy to see from the local theory of the curves  $E_{a,b}$  (Table 1, §2.2.2) that  $S' = S_2$  and  $S'' = S_3$ . This implies  $\frac{|\operatorname{Sel}^{\psi_{a,b}}(\widehat{E}_{a,b}/K)|}{|\operatorname{Sel}^{\psi_{a,b}}(\widehat{E}_{a,b}/K)|} = 3^{|S_2| - |S_3| + 1}$ .

Rearranging the terms and using Remark 3.6 we get that

$$\dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) = \dim_{\mathbb{F}_3} \operatorname{Sel}^{\widehat{\psi}_{a,b}}(\widehat{E}_{a,b}/K) + |S_3| - |S_2| - 1 \ge h_{S_{1,3}(L)}^3 + |S_3| - |S_2| - 1.$$

This along with Theorem 3.5 gives that  $\dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \ge \max\{h_{S_{1,2}(L)}^3, h_{S_{1,3}(L)}^3 + |S_3| - |S_2| - 1\}.$ Similarly by Remark 3.6, we have that

$$\dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(\widehat{E}_{a,b}/K) + |S_3| - |S_2| - 1 = \dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \le h_{S_{1,2}(L)}^3 + |S_{1,2}(L)| + |S_3| - |S_2| + 1.$$

Together with Theorem 3.5, this gives that

$$\dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \le \min\{h_{S_{1,2}(L)}^3 + |S_{1,2}(L)| + |S_3| - |S_2| + 1, \ h_{S_{1,3}(L)}^3 + |S_{1,3}(L)| + 2\}.$$

This completes the proof.

**Example (1):** Take a = 29 and b = 76. Then  $S_1 = S_2 = \emptyset$  and  $S_3 = \{2\mathcal{O}_K\}$ ; also,  $h_{S_{1,2}(L)}^3 = h_L^3 = 1$  and  $h^3_{S_{1,3}(L)} = 0.$  So, by Theorem 3.8,  $1 \leq \dim_{\mathbb{F}_3} \mathrm{Sel}^{\psi_{a,b}}(E_{a,b}/K) \leq 3.$ 

**Example (2):** For a = 79 and b = 79, observe that  $S_1 = S_2 = S_3 = \emptyset$  and  $h^3_{S_{1,2}(L)} = h^3_{S_{1,3}(L)} = h^3_L = 2$ . So, by Theorem 3.8,  $2 \leq \dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \leq 3$ .

**Example (3):** Taking a = 2263 and b = 72, we have that  $S_1 = S_2 = S_3 = \emptyset$  and  $h_L^3 = 3$ . Thus, we get that  $3 \leq \dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \leq 4$  using Theorem 3.8.

**Example (4):** If a = 113 and b = 22, then  $S_1 = \{2\mathcal{O}_K\}, S_2 = \{11\mathcal{O}_K\}$  and  $S_3 = \emptyset$ . We get that  $Cl_{S_{1,2}(L)(L)} = \mathbb{Z}/6$  and  $Cl_{S_{1,3}(L)(L)} = \mathbb{Z}/3$ . Thus, Theorem 3.8 gives that  $1 \leq \dim_{\mathbb{F}_3} \text{Sel}^{\psi_{a,b}}(E_{a,b}/K) \leq 5$ .

**Corollary 3.9.** Assume that  $a \notin K^{*2}$  and  $3 \nmid a$ . Then the global root number  $\omega(E_{a,b}/K)$  is determined by the parity of  $|S_2| + |S_3|$ . More precisely,

$$\omega(E_{a,b}/K) = (-1)^{|S_2| + |S_3| + 1} = \begin{cases} -1, & \text{if } |S_2| \equiv |S_3| \pmod{2}, \\ 1, & \text{if } |S_2| \not\equiv |S_3| \pmod{2}. \end{cases}$$
(14)

*Proof.* From the exact sequence (12), we obtain

$$\dim_{\mathbb{F}_3} \operatorname{Sel}^3(E_{a,b}/K) \equiv \dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) + \dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(\widehat{E}_{a,b}/K) \pmod{2} \tag{15}$$

since  $\dim_{\mathbb{F}_3} \frac{\operatorname{III}(\hat{E}_{a,b}/K)[\psi_{a,b}]}{\psi_{a,b}(\operatorname{III}(E_{a,b}/K)[3])}$  is even [BES, Prop. 49]. Also, from the proof of Theorem 3.8, we get

$$\dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) = \dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(\widehat{E}_{a,b}/K) + |S_3| - |S_2| - 1$$

and hence  $\dim_{\mathbb{F}_3} \operatorname{Sel}^3(E_{a,b}/K) \equiv |S_3| - |S_2| - 1 \pmod{2}$ . Now  $E_{a,b}(K)[3] = 0$  since  $a \notin K^{*2}$  by assumption. It follows that  $r = \operatorname{corank} \operatorname{Sel}^{3^{\infty}}(E_{a,b}/K) \equiv \dim_{\mathbb{F}_3} \operatorname{Sel}^3(E_{a,b}/K) \pmod{2}$ . The result follows from 3-parity conjecture of  $E_{a,b}$  over K which is known due to Nekovář and Dokchitser-Dokchitser.  $\square$ 

By studying the image of the Kummer maps over  $\mathbb{Q}_{\ell}$ , following a method similar to that in Section 2, one can obtain an upper bound of  $\dim_{\mathbb{F}_3} \mathrm{Sel}^{\psi_{a,b}}(E_{a,b}/\mathbb{Q})$ . In particular, if this upper bound is small, it is possible to determine the dimension of the 3-Selmer group of  $E_{a,b}$  over  $\mathbb{Q}$ . Using this approach, we provide an infinite family of elliptic curves  $E_{a,b}$  with 3-Selmer rank equal to 1.

**Proposition 3.10.** For a positive proportion of natural numbers n, one has that

$$\omega(E_{n,n}/\mathbb{Q}) = -1, \ \omega(E_{-3n,-3n}/\mathbb{Q}) = +1 \ and \ \dim_{\mathbb{F}_3} \operatorname{Sel}^3(E_{n,n}/\mathbb{Q}) = 1$$

Proof. Let n be a square-free natural number satisfying  $n \equiv 2$  or 3 (mod 4),  $n \equiv 2 \pmod{3}$  and n is a quadratic non-residue modulo 31. Let  $K_n := \mathbb{Q}(\sqrt{-3n})$  and  $\mathcal{O}_{K_n}$  be its ring of integers. Consider the elliptic curve  $E_{n,n}$ . Then for the above choice of n, for every prime  $\ell$ , using (8) one can show that  $\operatorname{Im} \delta_{\psi_{n,n},\mathbb{Q}_{\ell}} \subset (A_{\ell}^*/A_{\ell}^{*3})_{N=1}$ , where  $A_{\ell} := \frac{\mathbb{Z}_{\ell}[X]}{(X^2+3n)}$ . Consequently, we have that  $\dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{n,n}}(E_{n,n}/\mathbb{Q})$  is bounded by  $h_{K_n}^3 + \dim_{\mathbb{F}_3} \frac{\mathcal{O}_{K_n}^*}{\mathcal{O}_{K_n}^*} = h_{K_n}^3$ . As  $|E_{n,n}(\mathbb{R})[\psi_{n,n}]| = 3$ , from Cassels' formula (13) and Schaefer's formula (8), we obtain that  $\frac{|\operatorname{Sel}^{\psi_{n,n}}(\widehat{E}_{n,n}/\mathbb{Q})|}{|\operatorname{Sel}^{\psi_{n,n}}(E_{n,n}/\mathbb{Q})|} = 3$ . Using this, we deduce from [SS, Lemma 6.1] (see also (15)) that  $\dim_{\mathbb{F}_3} \operatorname{Sel}^3(E_{n,n}/\mathbb{Q}) \equiv 1 \pmod{2}$  and as  $E_{n,n}(\mathbb{Q})[3] = 0$ , we have that corank  $\operatorname{Sel}^{3^{\infty}}(E_{n,n}/\mathbb{Q}) \equiv$  $\dim_{\mathbb{F}_3} \operatorname{Sel}^3(E_{n,n}/\mathbb{Q}) \equiv 1 \pmod{2}$ . It follows that  $\omega(E_{n,n}/\mathbb{Q}) = -1$ , by the 3-parity conjecture. Further, observe that  $E_{n,n}$  is isomorphic to  $E_{-3n,-3n}$  over K and  $\omega(E_{n,n}/K) = -1$  by Corollary 3.9. Hence, we get that  $\omega(E_{-3n,-3n}/\mathbb{Q}) = +1$ .

By a result of Nakagawa-Horie (which can be conveniently found in [By, Lemma 2.2]), we see that in each of the 30 equivalent classes,  $n \equiv a \pmod{372}$  determined by the conditions  $n \equiv 2 \text{ or } 3 \pmod{4}$ ,  $n \equiv 2 \pmod{$ 

Let  $m, M \in \mathbb{N}$  be such that  $d := \operatorname{gcd}(m, M)$  is square free. Then we have

$$\sum_{\substack{0 < n \le X \\ \equiv m \pmod{M}}} \mu^2(n) = \frac{X}{M\zeta(2)} \Big(\prod_{p \mid \frac{M}{d}} \frac{1}{1 - p^{-2}}\Big) + o(X).$$

It follows that the number of square-free integers  $0 < n \le X$ , with  $n \equiv 2, 11 \pmod{12}$  and n is not a square modulo 31 is of the order of  $\frac{30X}{372\zeta(2)} \cdot \frac{2^2}{2^2-1} \cdot \frac{3^2}{3^2-1} \cdot \frac{31^2}{31^2-1} = \frac{31X}{2^8\zeta(2)}$ .

Putting all of it together, we see that for  $X \gg 0$ , the number of square-free natural numbers  $n \leq X$ , such that  $\omega(E_{n,n}/\mathbb{Q}) = -1$ ,  $\omega(E_{-3n,-3n}/\mathbb{Q}) = +1$  and  $\dim_{\mathbb{F}_3} \operatorname{Sel}^3(E_{n,n}/\mathbb{Q}) = 1$  is at least  $\frac{31X}{2^9\zeta(2)}$ . This completes the proof.

### 4. Arbitrary large rank of 3-Selmer groups of elliptic curves over K

We construct infinite families of elliptic curves with arbitrary large 3-Selmer group over K.

**Theorem 4.1.** Given any  $n \ge 0$ , there exist infinitely many elliptic curves  $E_{a,1}: y^2 = x^3 + a(x-1)^2$  with  $a \in \mathbb{Z}$  such that  $\dim_{\mathbb{F}_3} \operatorname{Sel}^3(E_{a,1}/K) \ge \dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,1}}(E_{a,1}/K) \ge 2n$ . These curves have a rational 3-isogeny and no non-trivial 3-torsion points over K.

Proof. Take  $a = \frac{p_1 p_2 \cdots p_{2n+1} - 27}{4}$ , where  $p_i$ 's are all distinct rational primes and each  $p_i \equiv -1 \pmod{12}$ . Then  $p_1 p_2 \cdots p_{2n+1} = 12s - 1$  for some s and  $a = \frac{12s-28}{4} = 3s - 7 \equiv -1 \pmod{3}$ . Hence,  $a \notin K^{*2}$ ; moreover,  $3 \nmid a$  and  $a \notin K^{*2}_{\mathfrak{p}}$ . Consider the sets  $S_2$ ,  $S_3$  from the Definition 3.2 and observe that  $S_2 = \emptyset$ . Further,  $d := 4a + 27 = p_1 p_2 \cdots p_{2n+1}$ , so each  $p_i \mid d$  and  $p_i \nmid 2a$ . Also, since each  $p_i \equiv -1 \pmod{12}$ , we see that each  $p_i$  is inert in K and  $\left(\frac{-3}{p_i}\right) = -1$  for all i. Hence, either  $\left(\frac{a}{p_i}\right) = 1$  or  $\left(\frac{-3a}{p_i}\right) = 1$ , which implies  $a \in K^{*2}_{p_i}$  for all  $p_i$ . This in turn implies that each  $p_i \in S_3$ ; in fact,  $S_3 = \{p_1, \cdots, p_{2n+1}\}$ . Indeed, if  $\mathfrak{q} \neq p_i$  for any i, then  $\mathfrak{q} \nmid d$ , so  $\mathfrak{q} \notin S_3$ . As  $a \notin K^{*2}_{\mathfrak{p}}$ ,  $\mathfrak{p} \notin S_3$  and  $2\mathcal{O}_K \notin S_3$ . Thus, we have  $|S_3| = 2n + 1$ . By Theorem 3.8, we get that  $\dim_{\mathbb{F}_3} \mathrm{Sel}^{\psi_{a,1}}(E_{a,1}/K) \geq 2n$ .

Clearly, there are infinitely many choices of a for the given n, hence the statement holds.

**Remark 4.2** (Related works). In [Ca1], Cassels showed that  $\dim_{\mathbb{F}_3} \operatorname{Sel}^3(E/\mathbb{Q})$  for a class of elliptic curves E, each having a rational 3-torsion point and j-invariant 0, can be arbitrarily large. For p = 2, [Bö, Kr] prove that for any given n > 0, there exists an elliptic curve E such that  $\dim_{\mathbb{F}_2} \operatorname{Sel}^2(E/\mathbb{Q}) \ge n$ , assuming that  $E[2] \subset E(\mathbb{Q})$ . On the other hand, for certain  $n \ge 5$ , there are several results (see [KS, Fi]) proving that the n-Selmer groups of elliptic curves over  $\mathbb{Q}$  can be arbitrarily large.

In contrast, in our result on  $\dim_{\mathbb{F}_3} \operatorname{Sel}^3(E_{a,b}/K)$  in Theorem 4.1, the elliptic curves  $E_{a,b}/\mathbb{Q}$  have a  $\mathbb{Q}$ rational 3-isogeny, do not have a non-trivial 3-torsion point over K and the *j*-invariant  $j(E_{a,b}) \neq 0$ .

In their recent article [BKOS, Theorem 1.3], the authors show that for a fixed number field F and  $r \ge 1$ , almost all elliptic curves E/F with a cyclic F-rational 9-isogeny have a positive proportion of quadratic twists  $E_s$  with  $\operatorname{III}(E_s/F)[3] \ge 9^r$  (see [BKOS, Theorem 1.3] for a precise statement). In a different direction, given a prime p and an elliptic curve E defined over a number field F, it was shown by [Če] that  $\dim_{\mathbb{F}_p} \operatorname{Sel}^p(E/F')$ is unbounded where F' varies over  $\mathbb{Z}/p\mathbb{Z}$ -extensions of F. There are important works of Mazur-Rubin on large corank of  $p^{\infty}$ -Selmer groups.

We have the following interesting application of Theorem 3.5.

**Corollary 4.3.** Given a biquadratic number field L' containing  $\zeta_3$ , there exist infinitely many elliptic curves  $E_{a,b}$  and  $\widehat{E}_{a,b}$  having

 $\dim_{\mathbb{F}_3} \operatorname{Sel}^{\widehat{\psi}_{a,b}}(\widehat{E}_{a,b}/K) - 2 \leq \dim_{\mathbb{F}_3} Cl(L')[3] \leq \dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K) \leq \dim_{\mathbb{F}_3} \operatorname{Sel}^3(E_{a,b}/K).$ 

Proof. Note that  $L' \cong \mathbb{Q}(\sqrt{-3}, \sqrt{a'})$  for some square-free integer a'. Since  $-3a' \in L'$ , we may assume  $3 \nmid a'$ . Set  $T_{a'} := \{\ell \in \Sigma_{\mathbb{Q}} : \ell \nmid 3a', \ell \equiv 1 \pmod{3} \text{ and } \left(\frac{a'}{\ell}\right) = -1\}$ . It is easy to see that  $T_{a'}$  is an infinite set. Take a := 16a' if  $a' \equiv 1 \pmod{4}$  and a := a' otherwise. We can verify that for such a and for every  $\ell \in T_{a'}$ , the sets  $S_1$  and  $S_2$  in the Definition 3.2 are both empty.

For a fixed *a* as above, consider the elliptic curves  $E_{a,\ell_i}$  with  $\ell_i \in T_{a'}$ . We claim that  $\{E_{a,\ell_i} \mid \ell_i \in T_{a'}\}$  is an infinite set of non-isomorphic elliptic curves over *K*. To see this, we compute the *j*-invariant  $j(E_{a,\ell_i}) = -\frac{2^8 a(a+6\ell_i)^3}{\ell_i^3(4a+27\ell_i)}$  and observe that there are at most four choices of  $\ell_i$  such that  $j(E_{a,\ell_i})$  are equal. For each of  $\{E_{a,\ell_i} \mid \ell_i \in T_{a'}\}$ , we get that  $S_1 \cup S_2 = \emptyset$  and the result is immediate from Theorem 3.5.

# EXAMPLES

Table 2 contains the bounds on  $\dim_{\mathbb{F}_3} \operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K)$  and  $\dim_{\mathbb{F}_3} \operatorname{Sel}^3(E_{a,b}/K)$  of elliptic curves  $E_{a,b}$  computed using SageMath/Magma. We define  $s_l^{\psi}$  and  $s_u^{\psi}$  to be the lower and upper bounds on the  $\operatorname{Sel}^{\psi_{a,b}}(E_{a,b}/K)$ , respectively, and  $s_l^3$  and  $s_u^3$  to be the lower and upper bounds on the  $\operatorname{Sel}^3(E_{a,b}/K)$ , respectively, following Theorem 3.5, Corollary 3.7 and Theorem 3.8. Also, r denotes  $\operatorname{rk} E_{a,b}(K)$ , which is computed using Magma under the assumption of GRH. In the cases when Magma can not compute r precisely, [-, -] will denote the bounds on r given by the software.

a, b	$S_1$	$S_2$	$S_3$	$h^3_{S_{1,2}(L)}$	$h^3_{S_{1,3}(L)}$	r	$s_l^{\psi_{a,b}}$	$s_u^{\psi_{a,b}}$	$s_l^3$	$s_u^3$
79,131	Ø	$\{131\mathcal{O}_K\}$	Ø	2	2	[0, 2]	2	4	2	10
137, 127	$\{2\mathcal{O}_K\}$	Ø	$\{41\mathcal{O}_K\}$	1	1	[1, 2]	1	5	1	12
137, 137	$\{2\mathcal{O}_K\}$	Ø	Ø	1	1	1	1	4	1	9
137, 143	$\{2\mathcal{O}_K\}$	$\{11\mathcal{O}_K\}$	$\{4409\mathcal{O}_K\}$	1	1	1	1	6	1	13
142,83	Ø	$\{83\mathcal{O}_K\}$	$\{53\mathcal{O}_K\}$	2	2	3	2	5	3	11
142,194	Ø	Ø	$\{2903\mathcal{O}_K\}$	2	2	[0, 2]	2	4	2	8
1714, 20	Ø	$\{5\mathcal{O}_K\}$	Ø	3	3	2	3	5	3	12
2230, 48	Ø	$\{\mathfrak{p}\}$	$\{1277\mathcal{O}_K\}$	2	3	1	2	5	2	11
2230,533	Ø	$\{41\mathcal{O}_K\}$	Ø	3	3	2	3	5	3	12
2263,587	Ø	$\{587\mathcal{O}_K\}$	Ø	3	3	2	3	5	3	12
2659, 29	Ø	$\{29\mathcal{O}_K\}$	Ø	3	3	2	3	5	3	12
2659,98	Ø	Ø	$\{29\mathcal{O}_K\}$	3	3	2	3	5	3	10
3023, 273	Ø	Ø	$\{19463\mathcal{O}_K\}$	3	3	2	3	5	3	10
3023,678	Ø	$\{113\mathcal{O}_K\}$	Ø	3	3	0	3	5	3	12
79,193	Ø	Ø	Ø	2	2	[2,3]	2	3	2	7
142, 12	Ø	$\{\mathfrak{p}\}$	Ø	1	2	0	1	3	1	7
142,67	Ø	Ø	Ø	2	2	1	2	3	2	7
223, 3	Ø	$\{\mathfrak{p}\}$	Ø	1	2	0	1	3	1	7
223, 63	Ø	$\{\mathfrak{p}\}$	Ø	1	2	0	1	3	1	7
2659, 24	Ø	$\{\mathfrak{p}\}$	Ø	2	3	0	2	4	2	9
2659, 39	Ø	$\{\mathfrak{p}\}$	Ø	2	3	0	2	4	2	9
3023, 18	Ø	Ø	Ø	3	3	[1,3]	3	4	3	9
3391, 12	Ø	$\{\mathfrak{p}\}$	Ø	2	3	0	2	4	2	9
3391,67	Ø	Ø	Ø	3	3	[0, 1]	3	4	3	9
3667, 3	Ø	{ <b>p</b> }	Ø	2	3	0	2	4	2	9
3667, 63	Ø	{ <b>p</b> }	Ø	2	3	[0,2]	2	4	2	9
4094, 23	Ø	Ø	Ø	3	3	[0,1]	3	4	3	9
4094,74	Ø	Ø	Ø	3	3	[0,1]	3	4	3	9
4151, 14	Ø	Ø	Ø	3	3	[0,3]	3	4	3	9
4279, 39	Ø	{¤}	Ø	2	3	[0, 2]	2	4	2	9
4279,66	Ø	{p}	Ø	2	3	0	2	4	2	9
4778,48	Ø	Ø	Ø	3	3	[0,1]	3	4	3	9
4910,14	Ø	Ø	Ø	3	3	[0,1]	3	4	3	9
4910,38	Ø	Ø	Ø	3	3	[0,1]	3	4	3	9
43063, 7	Ø	Ø	Ø	4	4	[0, 1]	3	4	3	9
43063,96	Ø	{p}	Ø	3	4	0	3	5	3	11
51694, 13	Ø	Ø	Ø	4	4	[1, 3]	4	5	4	11
51694, 21	Ø	{p}	Ø	3	4	0	3	5	3	11
53507, 18	Ø	Ø	Ø	4	4	1	4	5	4	11
53507,74	Ø	Ø	Ø	4	4	[0,1]	4	5	4	11
529987, 32	Ø	Ø	Ø	4	4	[0, 1]	4	5	4	11
529987, 108	Ø	{\$}	Ø	3	4	0	3	5	3	11

TABLE 2. Bounds on the  $\psi_{a,b}$ - and 3-Selmer groups of the curves  $E_{a,b}: y^2 = x^3 + a(x-b)^2$ 

# References

[BES]	M. Bhargava, N. Elkies & A. Shnidman, The average size of the 3-isogeny Selmer groups of elliptic curves $y^2 = x^3 + k$ ,
[BK] [BKOS]	<ul> <li>A. Brumer &amp; K. Kramer, The rank of elliptic curves, Duke Math J. 44(4) (1972), 715-743.</li> <li>M. Bhargava, Z. Klagsbrun, R. Oliver &amp; A. Shnidman, Elements of given order in Tate–Shafarevich groups of abelian varieties in quadratic twist families. Algebra Number Theory 15(3) (2021), 627-655</li> </ul>
[Bö]	R. Bölling, Die Ordnung der Schaftrewitsch-Tate Gruppe kann beliebig groß werden, Math. Nachr. 67 (1975), 157–179.
[Bv]	D. Byeon, Class numbers of quadratic fields $\mathbb{Q}(\sqrt{D})$ and $\mathbb{Q}(\sqrt{tD})$ , Proc. Amer. Math. Soc. <b>132</b> (11) (2004), 3137–3140.
[Ca1]	J. W. S. Cassels, Arithmetic on curves of genus 1. VI. The Tate-Šafarevič group can be arbitrarily large, J. Reine Angw. Math. 214/215 (1964), 65-70.
[Ca2]	J. W. S. Cassels, Arithmetic on curves of genus 1. VIII. On conjectures of Birch and Swinnerton-Dyer, J. Reine Angw.
	Math. 217 (1965), 180-199.
[Ca3]	J. W. S. Cassels, Lectures on elliptic curves, LMS Student Texts 24, Cambridge Uni. Press (1991).
[Če]	K. Česnavičius, Selmer groups and class groups, Compositio Math. 151 (2015), 416-434.
[Fi]	T. Fisher, Some examples of 5 and 7 descent for elliptic curves over Q, J. Eur. Math. Soc. 3 (2001), 169–201.
[JMS]	Somnath Jha, Dipramit Majumdar and Pratiksha Shingavekar, 3-Selmer group, ideal class groups and cube sum problem, https://arxiv.org/abs/2207.12487.
[Kr]	K. Kramer, A family of semistable elliptic curves with large Tate-Shafarevitch groups, <i>Proc. Amer. Math. Soc.</i> 89 (1983), 379–386.
[KS]	R. Kloosterman & E. Schaefer, Selmer groups of elliptic curves that can be arbitrarily large, J. Number Theory 99 (2003), 148-163.
[Li]	C. Li, 2-Selmer groups, 2-class groups and rational points on elliptic curves, Trans. AMS 371 (2019), 4631-4653.
[Sc]	E. Schaefer, Class groups and Selmer groups, J. Number Theory 56(1) (1996), 79-114.
[Si1]	J. Silverman, The arithmetic of elliptic curves, GTM 106, Springer-Verlag, New York (1986).
[Si2]	J. Silverman, Advanced topics in the arithmetic of elliptic curves, GTM 151, Springer-Verlag, 1994.
[SS]	E. Schaefer & M. Stoll, How to do a <i>p</i> -descent on an elliptic curve, <i>Trans. AMS</i> <b>356</b> (3) (2003), 1209–1231.
[Vé]	J. Vélu, Isogénies entre courbes elliptiques, C. R. Acad. Sci. Paris 273 (1971), 238-241.
[Wa]	L. C. Washington, Introduction to cyclotomic fields, GTM 83 Springer-Verlag, New York (2012).

Somnath Jha, jhasom@iitk.ac.in, Indian Institute of Technology Kanpur, India

DIPRAMIT MAJUMDAR, DIPRAMIT@IITM.AC.IN, INDIAN INSTITUTE OF TECHNOLOGY MADRAS, INDIA

Pratiksha Shingavekar, pshingavekar@gmail.com, Chennai Mathematical Institute, India