# ON FINITE CONVERGENCE AND MINIMIZER EXTRACTION IN MOMENT RELAXATIONS WITH CORRELATIVE SPARSITY

GIOVANNI FANTUZZI AND FEDERICO FUENTES

ABSTRACT. We provide a new sufficient condition to detect the finite convergence of moment relaxations of polynomial optimization problems with correlative sparsity. The condition requires that certain moment matrices in the relaxation admit a flat extension, and that the variable cliques used to construct the relaxation satisfy a 'running intersection' property. The proof also reveals an algorithm to extract at least as many minimizers for the original polynomial optimization problem as the smallest rank of the moment matrices in its relaxation. The necessity of the running intersection property is demonstrated with an illustrative example.

## 1. INTRODUCTION

Moment-sum-of-squares (moment-SOS) relaxations are an established approach to bound from below the global minimum of a polynomial optimization problem (POP) by solving semidefinite programs [9, 12, 19]. Raising the relaxation order to improve the lower bound rapidly increases the size of these semidefinite programs, often resulting in high or outright prohibitive computational costs and memory footprint. To avoid this, one can modify standard relaxations to exploit properties of the POP such as symmetries [2, 5, 13, 20] and sparsity [10, 14, 21–26]. A key question is then whether these modified relaxations enjoy the same good theoretical properties as the standard ones. In particular, one would like to have asymptotic convergence results for the lower bounds, sufficient conditions to detect finite convergence, and techniques to extract one or more POP minimizers when these conditions hold.

As we explain next, this work provides new rank conditions that, combined with a well-known 'running intersection' property, imply the finite convergence of moment relaxations of POPs with *correlative sparsity* [10, 21]. The proof also reveals an algorithm to recover at least as many optimizers for the POP as the smallest rank of the moment matrices in its relaxation.

1.1. Main result. Fix  $n \in \mathbb{N}$  and cover the set  $[n] := \{1, \ldots, n\}$  by subsets  $\Delta_1, \ldots, \Delta_m \subset [n]$ . We slightly abuse established terminology and call these subsets *cliques*. We also assume without loss of generality that no clique is a subset of another clique. Given a 'global' vector  $\mathbf{x} = (x_1, \ldots, x_n)$ , let  $\mathbf{x}_{\Delta_i}$  be the 'clique' subvector of  $\mathbf{x}$  indexed by the elements of  $\Delta_i$ . Given positive integers  $N_1, \ldots, N_m$  and polynomials  $f_i : \mathbb{R}^{|\Delta_i|} \to \mathbb{R}$  and  $\mathbf{g}_i : \mathbb{R}^{|\Delta_i|} \to \mathbb{R}^{N_i}$ , consider the POP

(POP) 
$$f^* := \min_{\mathbf{x} \in \mathbb{R}^n} \sum_{i=1}^m f_i(\mathbf{x}_{\Delta_i}) \text{ s.t. } \mathbf{g}_i(\mathbf{x}_{\Delta_i}) \ge 0 \quad \forall i \in [m].$$

Dept. of Mathematics, Friedrich-Alexander Universität Erlangen-Nürnberg Institute for Mathematical and Computational Engineering (IMC), School of Engineering and Faculty of Mathematics, Pontificia Universidad Católica de Chile

E-mail addresses: giovanni.fantuzzi@fau.de, federico.fuentes@uc.cl.

Date: February 4, 2025.

Its sparse moment relaxation of order  $\omega \in \mathbb{N}$  is

(MOM) 
$$f_{\omega}^* := \min_{\mathbf{y} \in \mathbb{R}^{|\mathbb{N}_{2\omega}^n|}} \langle \mathbf{f}, \mathbf{y} \rangle \quad \text{s.t.} \quad \begin{cases} \mathbb{M}_{\Delta_i}^{\omega}(\mathbf{y}) \succeq 0 \quad \forall i \in [m], \\ \mathbb{M}_{\Delta_i}^{\omega}(\mathbf{g}_i \mathbf{y}) \succeq 0 \quad \forall i \in [m], \\ \mathcal{L}_{\mathbf{y}}(1) = 1, \end{cases}$$

where  $|\mathbb{N}_{2\omega}^n| = \binom{n+2\omega}{n}$ , **f** is the vector of coefficients of  $f(x) = \sum_{i=1}^m f_i(\mathbf{x}_{\Delta_i})$  with respect to the monomial basis, angled brackets indicate an inner product,  $\mathbb{M}_{\Delta_i}^{\omega}(\mathbf{y})$ and  $\mathbb{M}_{\Delta_i}^{\omega}(\mathbf{g}_i \mathbf{y})$  are the *moment* and *localizing matrices* of order  $\omega$  associated to the clique  $\Delta_i$ , and  $\mathcal{L}_{\mathbf{y}}$  is the Riesz functional associated to the vector  $\mathbf{y}$ . Definitions are given in section 2.

Set  $d_i = \lceil \frac{1}{2} \deg \mathbf{g}_i \rceil$  and let the 'clique overlap' matrix  $\mathbb{M}_{\Delta_i \cap \Delta_j}^{\omega}(\mathbf{y})$  denote the largest principal submatrix of the moment matrix  $\mathbb{M}_{\Delta_i}^{\omega}(\mathbf{y})$  that, up to a symmetric permutation, is also a principal submatrix of  $\mathbb{M}_{\Delta_j}^{\omega}(\mathbf{y})$ . Our main result, proven in section 3, is the following.

**Theorem 1.1.** Suppose that the cliques  $\Delta_1, \ldots, \Delta_m$  satisfy the following running intersection property:

(RIP)  $\forall i \geq 2, \quad \exists j \in [i-1] \quad such \ that \quad \Delta_i \cap (\Delta_1 \cup \cdots \cup \Delta_{i-1}) \subset \Delta_j.$ 

If  $\mathbf{y}$  is optimal for (MOM) and

(1.1a) 
$$\operatorname{rank} \mathbb{M}_{\Delta_i}^{\omega}(\mathbf{y}) = \operatorname{rank} \mathbb{M}_{\Delta_i}^{\omega-d_i}(\mathbf{y})$$

(1.1b) 
$$\operatorname{rank} \mathbb{M}_{\Delta_i \cap \Delta_j}^{\omega}(\mathbf{y}) = \operatorname{rank} \mathbb{M}_{\Delta_i \cap \Delta_j}^{\omega - 1}(\mathbf{y})$$

for all  $i \in [m]$  and at least one  $j \in [i-1]$  satisfying (RIP), then  $f_{\omega}^* = f^*$ . Moreover, one can extract from **y** at least  $r_{\min}$  distinct global minimizers for (POP), where

(1.2) 
$$r_{\min} := \min_{i \in [m]} \operatorname{rank} \mathbb{M}_{\Delta_i}^{\omega}(\mathbf{y})$$

Remark 1.1. In some cases when (RIP) does not hold, the cliques can be reordered so that it does. Our notation  $\Delta_1, \ldots, \Delta_m$  always refers to the reordered cliques. If we say that (RIP) fails, we mean there is no clique reordering for which it holds.

Remark 1.2. Rank conditions like (1.1) are typically called *flat extension* or *flat truncation* conditions because they imply that the moment matrix  $\mathbb{M}_{\Delta_i}^{\omega}(\mathbf{y})$  can be truncated to the submatrix  $\mathbb{M}_{\Delta_i}^{\omega-d_i}(\mathbf{y})$  without changing its rank.

Further remarks on Theorem 1.1 are given after its proof (see section 3.2).

1.2. Relation to other conditions implying finite convergence. Rank-based sufficient conditions to detect the finite convergence of moment relaxations for correlatively sparse POPs have already been proposed in the literature. For example, Lasserre [10, Theorem 3.7] proves that  $f_{\omega}^* = f^*$  if (1.1a) holds and the clique overlap matrices  $\mathbb{M}_{\Delta_i \cap \Delta_j}^{\omega}(\mathbf{y})$  have rank one. Theorem 1.1 weakens this rank-one condition at the expense of requiring (RIP). We believe this trade-off to be very useful in practice because Lasserre's rank-one condition can only hold if (POP) has a unique minimizer, or if the entries of multiple minimizers  $\mathbf{x}_1^*, \ldots, \mathbf{x}_r^*$  whose index appears in at least two cliques have the same value. The latter is a rather special situation. In contrast, Theorem 1.1 can be applied to POPs with generic multiple minimizers. Moreover, one can always choose to enforce (RIP) through a so-called *chordal ex*tension process [21]. In fact, this is typically (but not always) done because (RIP) ensures the asymptotic convergence of  $f^*_\omega$  to  $f^*$  as  $\omega \to \infty$  when the quadratic module of weighted sum-of-squares polynomials generated by each of the polynomial vectors  $\mathbf{g}_i$  in (POP) is Archimedean (see [10, Theorem 3.6] and [8, Theorem 1], as well as [6, Theorem 5] for a simple proof).

Theorem 1.1 is also very similar to a claim by Nie *et al.* [18, Theorem 3.4] stating that, irrespective of (RIP), it is possible to extract optimizers of (POP) (and thus detect finite convergence) if there is a positive integer r such that

(1.3a) 
$$\operatorname{rank} \mathbb{M}_{\Delta_i}^{\omega}(\mathbf{y}) = \operatorname{rank} \mathbb{M}_{\Delta_i}^{\omega-d_i}(\mathbf{y}) = r$$

(1.3b) 
$$\operatorname{rank} \mathbb{M}_{\Delta_i \cap \Delta_i}^{\omega}(\mathbf{y}) = \operatorname{rank} \mathbb{M}_{\Delta_i \cap \Delta_i}^{\omega-1}(\mathbf{y}) = r$$

for all  $i \in [m]$  and all  $j \in [m]$  such that  $\Delta_i \cap \Delta_j \neq \emptyset$ . In section 4 we give a simple counterexample showing that this claim is, in fact, *false*. Therefore, within the proof strategy of Theorem 1.1, the RIP cannot be dropped without either imposing much stronger assumptions on the ranks of the clique overlap matrices  $\mathbb{M}^{\omega}_{\Delta_i \cap \Delta_j}(\mathbf{y})$ , or resorting to sufficient conditions for finite convergence not based solely on the rank of the moment matrices (see, for instance, [18, Theorem 3.3], which is a key ingredient in the proof of Theorem 1.1).

Finally, we remark that many of the sufficient conditions to detect the finite convergence of (MOM) from [10, 18] were recently generalized to sparse polynomial matrix optimization problems [15]. We wonder if a version of Theorem 1.1 holds in this setting, too, but leave answering this question to future work.

## 2. NOTATION AND BASIC DEFINITIONS

2.1. Notation. Let  $\mathbb{N}$  be the non-negative integers. We write  $\mathbf{x}^{\alpha} = x_1^{\alpha_1} \cdots x_n^{\alpha_n}$  for the *n*-variate monomial with independent variables  $\mathbf{x} = (x_1, \ldots, x_n)$  and exponent  $\alpha = (\alpha_1, \ldots, \alpha_n) \in \mathbb{N}^n$ ,  $\mathbb{R}[\mathbf{x}]_d$  for the space of polynomials with variable  $\mathbf{x}$  of total degree up to d, and  $\mathbb{N}_d^n$  for the set of exponents  $\alpha \in \mathbb{N}^n$  with  $|\alpha_1| + \cdots + |\alpha_n| \leq d$ . Given a polynomial

(2.1) 
$$f(x) = \sum_{\alpha \in \mathbb{N}_d^n} f_\alpha \mathbf{x}^\alpha$$

in  $\mathbb{R}[\mathbf{x}]_d$ , we collect its  $|\mathbb{N}_d^n| = \binom{n+d}{n}$  coefficients and monomials in the vectors

(2.2a) 
$$\mathbf{f} := (f_{\alpha})_{\alpha \in \mathbb{N}^n_d},$$

(2.2b) 
$$[\mathbf{x}]_d := (\mathbf{x}^{\alpha})_{\alpha \in \mathbb{N}^n}.$$

The *Riesz functional* associated to a vector  $\mathbf{y} = (y_{\alpha})_{\alpha \in \mathbb{N}_{d}^{n}}$  is the unique countinuous linear functional  $\mathcal{L}_{\mathbf{y}}$  on  $\mathbb{R}[\mathbf{x}]_{d}$  satisfying  $\mathcal{L}_{\mathbf{y}}(\mathbf{x}^{\alpha}) = y_{\alpha}$  for every exponent  $\alpha \in \mathbb{N}_{d}^{n}$ . The definition implies that  $\mathcal{L}_{\mathbf{y}}(f) = \langle \mathbf{f}, \mathbf{y} \rangle$  for every  $f \in \mathbb{R}[\mathbf{x}]_{d}$ .

Lastly, given an index set  $\Delta \subset [n]$  and a 'global' vector  $\mathbf{x} = (x_1, \ldots, x_n)$ , we denote by  $\mathbf{x}_{\Delta}$  the subvector of  $\mathbf{x}$  indexed by the elements of  $\Delta$ . Let  $P_{\Delta} : \mathbb{R}^n \to \mathbb{R}^{|\Delta|}$  be the projection operator satisfying  $P_{\Delta}\mathbf{x} = \mathbf{x}_{\Delta}$  and denote its pseudoinverse by  $P_{\Delta}^{\dagger} : \mathbb{R}^{|\Delta|} \to \mathbb{R}^n$ . We identify  $P_{\Delta}$  with the  $|\Delta| \times n$  matrix listing the rows of the  $n \times n$  identity matrix indexed by  $\Delta$ , and  $P_{\Delta}^{\dagger}$  with its transpose. Note that  $P_{\Delta}^{\dagger}\mathbf{z}$  simply 'lifts' a vector  $\mathbf{z} \in \mathbb{R}^{|\Delta|}$  to a vector in  $\mathbb{R}^n$  by placing  $\mathbf{z}$  in the entries indexed by  $\Delta$ , and padding the rest with zeros. With this interpretation, it is clear that

(2.3a) 
$$P_{\Delta}P_{\Delta}^{\dagger}\mathbf{z} = \mathbf{z} \quad \forall \mathbf{z} \in \mathbb{R}^{|\Delta|}$$

(2.3b)  $P_{\Delta_1} P_{\Delta_2}^{\dagger} \mathbf{z} = \mathbf{0} \quad \forall \mathbf{z} \in \mathbb{R}^{|\Delta_2|}, \ \Delta_1 \cap \Delta_2 = \emptyset.$ 

2.2. Moment and localizing matrices. Consider problem (POP) with cliques  $\Delta_1, \ldots, \Delta_m$  that cover [n]. Set

(2.4) 
$$d_0 = \left\lceil \frac{1}{2} \deg f \right\rceil$$
 and  $d_i = \left\lceil \frac{1}{2} \deg \mathbf{g}_i \right\rceil$   $\forall i \in [m]$ .

Fix a positive integer  $\omega \ge \max\{d_0, \ldots, d_m\}$  and a vector  $\mathbf{y} = (y_\alpha)_{\alpha \in \mathbb{N}_{2\omega}^n}$ . For every  $i \in [m]$ , the moment matrix of order  $d \le \omega$  associated to clique  $\Delta_i$  is defined as

(2.5) 
$$\mathbb{M}^{d}_{\Delta_{i}}(\mathbf{y}) := \mathcal{L}_{\mathbf{y}}\left([\mathbf{x}_{\Delta_{i}}]_{d} \otimes [\mathbf{x}_{\Delta_{i}}]_{d}\right),$$

where  $\otimes$  denotes the usual tensor product. The localizing matrix of order  $d \in [d_i, \omega]$ associated to the clique  $\Delta_i$  and the entry  $g_{i,j}$  of the polynomial vector  $\mathbf{g}_i : \mathbb{R}^{|\Delta_i|} \to \mathbb{R}^{N_i}$  is instead defined as

(2.6) 
$$\mathbb{M}^{d}_{\Delta_{i}}(g_{i,j}\mathbf{y}) := \mathcal{L}_{\mathbf{y}}\left(g_{i,j}(\mathbf{x}_{\Delta_{i}})\left([\mathbf{x}_{\Delta_{i}}]_{d-d_{i}}\otimes[\mathbf{x}_{\Delta_{i}}]_{d-d_{i}}\right)\right).$$

The localizing matrix associated to the full polynomial vector  $\mathbf{g}_i$  is then simply the block-diagonal matrix

(2.7) 
$$\mathbb{M}^{d}_{\Delta_{i}}(\mathbf{g}_{i}\mathbf{y}) = \begin{bmatrix} \mathbb{M}^{d}_{\Delta_{i}}(g_{i,1}\mathbf{y}) & & \\ & \ddots & \\ & & \mathbb{M}^{d}_{\Delta_{i}}(g_{i,N_{i}}\mathbf{y}) \end{bmatrix}$$

Note that, for fixed  $i \in [m]$ , the moment and localizing matrices depend not on the full vector **y**, but only on (a subset of) the entries of the subvector

(2.8) 
$$\mathbf{y}_{\Delta_i} := \mathcal{L}_{\mathbf{y}}\left([\mathbf{x}_{\Delta_i}]_{2\omega}\right).$$

These vectors play a key role in the rest of the paper.

## 3. Proof of Theorem 1.1

We now prove Theorem 1.1. The argument, detailed in section 3.1, follows ideas from [18] and relies on classical results on the existence of atomic representing measures for truncated moment sequences (see [3, 4, 7, 11, 16, 17] for details). More comments on the proof and some of its implications are offered in section 3.2.

3.1. The proof. Let y be an optimal solution of (MOM). Fix  $i \in [m]$  and set

(3.1) 
$$K_i := \left\{ \mathbf{z} \in \mathbb{R}^{|\Delta_i|} : \ \mathbf{g}_i(\mathbf{z}) \ge 0 \right\}$$

It is well known (see, e.g., [3, Theorem 1.6] and [11, Theorem 1.6]) that if the rank conditions in (1.1a) hold, then the vector  $\mathbf{y}_{\Delta_i}$  defined in (2.8) is the moment vector of an atomic probability measure supported on  $r_i = \operatorname{rank} \mathbb{M}_{\Delta_i}^{\omega}(\mathbf{y})$  points in  $K_i$ . More precisely, for every  $i \in [m]$  there exist points  $\mathbf{z}^{i,1}, \ldots, \mathbf{z}^{i,r_i} \in K_i$  such that  $\mathbf{y}_{\Delta_i}$  lies in the interior of the convex hull of  $\{[\mathbf{z}^{i,1}]_{2\omega}, \ldots, [\mathbf{z}^{i,r_i}]_{2\omega}\}$ , i.e.,

(3.2a) 
$$\mathbf{y}_{\Delta_i} = \lambda_{i,1} [\mathbf{z}^{i,1}]_{2\omega} + \dots + \lambda_{i,r_i} [\mathbf{z}^{i,r_i}]_{2\omega},$$

$$\mathbf{z}^{i,1},\ldots,\mathbf{z}^{i,r_i}\in K_i$$

(3.2c) 
$$\lambda_{i,1} + \dots + \lambda_{i,r_i} = 1, \quad \lambda_{i,1}, \dots, \lambda_{i,r_i} \in (0,1).$$

To prove Theorem 1.1 we will apply the following theorem of Nie et al. [18].

**Theorem 3.1** (Adapted from [18, Theorem 3.3]). Let  $\mathbf{y}$  be optimal for (MOM) and assume the vectors  $\mathbf{y}_{\Delta_i}$  in (2.8) satisfy (3.2). Suppose  $\mathbf{x}^* \in \mathbb{R}^n$  satisfies

$$(3.3) P_{\Delta_i} \mathbf{x}^* \in \left\{ \mathbf{z}^{i,1}, \dots, \mathbf{z}^{i,r_i} \right\} \quad \forall i \in [m]$$

Then,  $f_{\omega}^* = f^*$  and  $\mathbf{x}^*$  is a minimizer of (POP).

The strategy to construct  $\mathbf{x}^* \in \mathbb{R}^n$  satisfying (3.3) is as follows. By assumption (see also Remark 1.1) we have ordered the cliques  $\Delta_1, \ldots, \Delta_m$  to ensure (RIP). Given this ordering, we proceed sequentially in a clique-by-clique fashion, concatenating vectors  $\mathbf{z}^{(1)}, \ldots, \mathbf{z}^{(m)}$  with  $\mathbf{z}^{(i)} \in {\mathbf{z}^{i,1}, \ldots, \mathbf{z}^{i,r_i}}$  obtained from (3.2) in such a way that the entries indexed by the intersection of cliques match. Specifically, starting with i = 1, for each clique index  $i \in [m]$  we first carefully select  $\mathbf{z}^{(i)}$  to match with the vectors  $\mathbf{z}^{(1)}, \ldots, \mathbf{z}^{(i-1)}$  already selected. Then, we construct a corresponding vector  $\mathbf{x}^{(i)} \in \mathbb{R}^n$  that satisfies

(3.4a) 
$$P_{\Delta_k} \mathbf{x}^{(i)} = \mathbf{z}^{(k)} \quad \forall k \le i,$$

(3.4b) 
$$P_{\Delta} \mathbf{x}^{(i)} = \mathbf{0} \qquad \forall \Delta \subset (\Delta_1 \cup \cdots \cup \Delta_i)^c$$

where  $\Delta^c$  denotes the complement of  $\Delta$  in [n]. After m steps we obtain the point  $\mathbf{x}^* = \mathbf{x}^{(m)}$ . We now make this construction rigorous.

For i = 1, we fix  $\mathbf{z}^{(1)} \in {\{\mathbf{z}^{1,1}, \ldots, \mathbf{z}^{1,r_1}\}}$  arbitrarily and set  $\mathbf{x}^{(1)} = P_{\Delta_1}^{\dagger} \mathbf{z}^{(1)}$ . One easily checks that the two conditions in (3.4) hold using the identities in (2.3).

Next, we proceed by induction. Fix  $i \in [m]$  with  $i \ge 2$  and assume we have already found points  $\mathbf{z}^{(1)}, \ldots, \mathbf{z}^{(i-1)}$  and a point  $\mathbf{x}^{(i-1)}$  satisfying the conditions in (3.4) with i - 1 in place of i. To obtain  $\mathbf{x}^{(i)}$ , we distinguish two cases.

**Case 1:**  $\Delta_i \cap (\Delta_1 \cup \cdots \cup \Delta_{i-1}) = \emptyset$ . In this case, clique  $\Delta_i$  does not overlap with any of the previous ones. Then, it suffices to pick any  $\mathbf{z}^{(i)} \in {\mathbf{z}^{i,1}, \ldots, \mathbf{z}^{i,r_i}}$  and set (3.5)  $\mathbf{x}^{(i)} = \mathbf{x}^{(i-1)} + P^{\dagger}_{\Delta_i} \mathbf{z}^{(i)}$ .

Using the identities in (2.3), it is straightforward to verify that condition (3.4b) holds. To check that (3.4a) also holds, note that for every k < i we have  $\Delta_i \cap \Delta_k = \emptyset$  by assumption and, therefore,

(3.6) 
$$P_{\Delta_k} \mathbf{x}^{(i)} = P_{\Delta_k} \mathbf{x}^{(i-1)} + \underbrace{P_{\Delta_k} P_{\Delta_i}^{\dagger} \mathbf{z}^{(i)}}_{=\mathbf{0}} = P_{\Delta_k} \mathbf{x}^{(i-1)} = \mathbf{z}^{(k)}.$$

Similarly, since  $\Delta_i \cap (\Delta_1 \cup \cdots \cup \Delta_{i-1})$  is empty by assumption, and since  $\mathbf{x}^{(i-1)}$  satisfies (3.4b) with i-1 in place of i by the induction hypothesis, we find that

(3.7) 
$$P_{\Delta_i} \mathbf{x}^{(i)} = \underbrace{P_{\Delta_i} \mathbf{x}^{(i-1)}}_{=\mathbf{0}} + P_{\Delta_i} P_{\Delta_i}^{\dagger} \mathbf{z}^{(i)} = P_{\Delta_i} P_{\Delta_i}^{\dagger} \mathbf{z}^{(i)} = \mathbf{z}^{(i)}.$$

We have therefore verified (3.4a), as desired.

**Case 2:**  $\Delta_i \cap (\Delta_1 \cup \cdots \cup \Delta_{i-1}) \neq \emptyset$ . In this case, clique  $\Delta_i$  overlaps with at least one previous clique. To take care of these intersections, recall that the assumptions of Theorem 1.1 ensure that there exists  $j \in [i-1]$  such that

$$(3.8) \qquad \qquad \Delta_i \cap (\Delta_1 \cup \dots \cup \Delta_{i-1}) \subset \Delta_j$$

and such that (1.1b) holds. For this particular index j, we have from (3.2) that

(3.9) 
$$\mathbf{y}_{\Delta_{i}\cap\Delta_{j}} = \lambda_{j,1} \left[ P_{\Delta_{i}\cap\Delta_{j}} P_{\Delta_{j}}^{\dagger} \mathbf{z}^{j,1} \right]_{2\omega} + \dots + \lambda_{j,r_{j}} \left[ P_{\Delta_{i}\cap\Delta_{j}} P_{\Delta_{j}}^{\dagger} \mathbf{z}^{j,r_{j}} \right]_{2\omega} \\ = \lambda_{i,1} \left[ P_{\Delta_{i}\cap\Delta_{j}} P_{\Delta_{i}}^{\dagger} \mathbf{z}^{i,1} \right]_{2\omega} + \dots + \lambda_{i,r_{i}} \left[ P_{\Delta_{i}\cap\Delta_{j}} P_{\Delta_{i}}^{\dagger} \mathbf{z}^{i,r_{i}} \right]_{2\omega}.$$

Now, set  $s_{ij} = \operatorname{rank} \mathbb{M}_{\Delta_i \cap \Delta_j}^{\omega}(\mathbf{y})$  and note that  $s_{ij} \leq \min\{r_i, r_j\}$  because the rank of a matrix cannot be smaller than that of a submatrix. By classical results on moment matrices (see, e.g., [11, Corollary 1.4]), the rank conditions in (1.1b) ensure that there is a unique choice of points  $\mathbf{v}^1, \ldots, \mathbf{v}^{s_{ij}} \in \mathbb{R}^{|\Delta_i \cap \Delta_j|}$  and of scalars  $\mu_1, \ldots, \mu_{s_{ij}} \in (0, 1)$  with unit sum such that the vector  $\mathbf{y}_{\Delta_i \cap \Delta_j} = \mathcal{L}_{\mathbf{y}}([\mathbf{x}_{\Delta_i \cap \Delta_j}]_{2\omega})$  can be represented as

(3.10) 
$$\mathbf{y}_{\Delta_i \cap \Delta_j} = \mu_1 [\mathbf{v}^1]_{2\omega} + \dots + \mu_{s_{ij}} [\mathbf{v}^{s_{ij}}]_{2\omega}.$$

The uniqueness of this representation and the inequality  $s_{ij} \leq \min\{r_i, r_j\}$  allow us to make two key observations. First, since  $P_{\Delta_j} \mathbf{x}^{(i-1)} = \mathbf{z}^{(j)} \in \{\mathbf{z}^{j,1}, \ldots, \mathbf{z}^{j,r_j}\}$  there exists an index  $q \in [s_{ij}]$  such that

(3.11) 
$$\mathbf{v}^{q} = P_{\Delta_{i} \cap \Delta_{j}} P_{\Delta_{j}}^{\dagger} \mathbf{z}^{(j)} = P_{\Delta_{i} \cap \Delta_{j}} P_{\Delta_{j}}^{\dagger} P_{\Delta_{j}} \mathbf{x}^{(i-1)}.$$

Second, there must exist an index  $\ell \in [r_i]$  such that

$$(3.12) P_{\Delta_i \cap \Delta_j} P_{\Delta_i}^{\dagger} \mathbf{z}^{i,\ell} = \mathbf{v}^q.$$

We now choose  $\mathbf{z}^{(i)} = \mathbf{z}^{i,\ell}$  and define

(3.13) 
$$\mathbf{x}^{(i)} = \mathbf{x}^{(i-1)} + P^{\dagger}_{\Delta_i \setminus \Delta_j} P_{\Delta_i \setminus \Delta_j} P^{\dagger}_{\Delta_i} \mathbf{z}^{(i)}.$$

To conclude the proof, we need to verify that  $\mathbf{x}^{(i)}$  satisfies the conditions in (3.4).

Condition (3.4b) holds because  $P_{\Delta} \mathbf{x}^{(i-1)} = \mathbf{0}$  when  $\Delta \cap (\Delta_1 \cup \cdots \cup \Delta_i) = \emptyset$  by the induction hypothesis, while  $P_{\Delta}P_{\Delta_i \setminus \Delta_j}^{\dagger}$  is the zero matrix since  $\Delta \cap (\Delta_i \setminus \Delta_j) = \emptyset$ . Next, we check (3.4a) for arbitrary clique index k < i. (The case k = i is

considered separately below). The running intersection property implies that

$$(3.14) \qquad \Delta_i \setminus (\Delta_1 \cup \cdots \cup \Delta_{i-1}) = \Delta_i \setminus [\Delta_i \cap (\Delta_1 \cup \cdots \cup \Delta_{i-1})] = \Delta_i \setminus \Delta_j,$$

so  $\Delta_k \cap (\Delta_i \setminus \Delta_j) = \emptyset$ . This, in turn, means that  $P_{\Delta_k} P_{\Delta_i \setminus \Delta_j}^{\dagger}$  is the zero matrix. Consequently, since  $\mathbf{x}^{(i-1)}$  satisfies (3.4a) with i-1 in place of i by the induction assumption,

$$(3.15) \qquad P_{\Delta_k} \mathbf{x}^{(i)} = P_{\Delta_k} \mathbf{x}^{(i-1)} + P_{\Delta_k} P_{\Delta_i \setminus \Delta_j}^{\dagger} P_{\Delta_i \setminus \Delta_j} P_{\Delta_i}^{\dagger} \mathbf{z}^{(i)} = P_{\Delta_k} \mathbf{x}^{(i-1)} = \mathbf{z}^{(k)}.$$

There remains only to verify (3.4a) for k = i. For this, observe that by (3.14),  $(\Delta_i \setminus \Delta_j) \subset (\Delta_1 \cup \cdots \cup \Delta_{i-1})^c$  and recall that  $\mathbf{x}^{(i-1)}$  satisfies (3.4b) with i-1 in place of i by the induction hypothesis. Then,

$$P_{\Delta_i \setminus \Delta_j} \mathbf{x}^{(i-1)} = \mathbf{0},$$

(3.16b) 
$$P_{\Delta_i \cap \Delta_j} \mathbf{x}^{(i-1)} = P_{\Delta_i \cap \Delta_j} P_{\Delta_j}^{\dagger} P_{\Delta_j} \mathbf{x}^{(i-1)}.$$

Using these identities, and recalling that  $\mathbf{z}^{(i)} = \mathbf{z}^{i,\ell}$  where  $\mathbf{z}^{i,\ell}$  satisfies (3.12), we obtain

$$(3.17) \qquad P_{\Delta_{i}}\mathbf{x}^{(i-1)} = P_{\Delta_{i}}P_{\Delta_{i}\backslash\Delta_{j}}^{\dagger}P_{\Delta_{i}\backslash\Delta_{j}}\mathbf{x}^{(i-1)} + P_{\Delta_{i}}P_{\Delta_{i}\cap\Delta_{j}}^{\dagger}P_{\Delta_{i}\cap\Delta_{j}}\mathbf{x}^{(i-1)}$$

$$\stackrel{(3.16a)}{=} P_{\Delta_{i}}P_{\Delta_{i}\cap\Delta_{j}}^{\dagger}P_{\Delta_{i}\cap\Delta_{j}}\mathbf{x}^{(i-1)}$$

$$\stackrel{(3.16b)}{=} P_{\Delta_{i}}P_{\Delta_{i}\cap\Delta_{j}}^{\dagger}P_{\Delta_{i}\cap\Delta_{j}}P_{\Delta_{j}}^{\dagger}P_{\Delta_{j}}\mathbf{x}^{(i-1)}$$

$$\stackrel{(3.11)}{=} P_{\Delta_{i}}P_{\Delta_{i}\cap\Delta_{j}}^{\dagger}\mathbf{v}^{q}$$

$$\stackrel{(3.12)}{=} P_{\Delta_{i}}P_{\Delta_{i}\cap\Delta_{j}}^{\dagger}P_{\Delta_{i}\cap\Delta_{j}}P_{\Delta_{i}}^{\dagger}\mathbf{z}^{(i)}.$$

We can therefore finally verify that

$$(3.18) \qquad P_{\Delta_{i}}\mathbf{x}^{(i)} = P_{\Delta_{i}}\mathbf{x}^{(i-1)} + P_{\Delta_{i}}P_{\Delta_{i}\backslash\Delta_{j}}^{\dagger}P_{\Delta_{i}\backslash\Delta_{j}}P_{\Delta_{i}}^{\dagger}\mathbf{z}^{(i)}$$

$$\stackrel{(3.17)}{=} P_{\Delta_{i}}P_{\Delta_{i}\cap\Delta_{j}}^{\dagger}P_{\Delta_{i}\cap\Delta_{j}}P_{\Delta_{i}}^{\dagger}\mathbf{z}^{(i)} + P_{\Delta_{i}}P_{\Delta_{i}\backslash\Delta_{j}}^{\dagger}P_{\Delta_{i}\backslash\Delta_{j}}P_{\Delta_{i}}^{\dagger}\mathbf{z}^{(i)}$$

$$= P_{\Delta_{i}}\left(P_{\Delta_{i}\cap\Delta_{j}}^{\dagger}P_{\Delta_{i}\cap\Delta_{j}} + P_{\Delta_{i}\backslash\Delta_{j}}^{\dagger}P_{\Delta_{i}\backslash\Delta_{j}}\right)P_{\Delta_{i}}^{\dagger}\mathbf{z}^{(i)}$$

$$= P_{\Delta_{i}}\left(P_{\Delta_{i}}^{\dagger}P_{\Delta_{i}}\right)P_{\Delta_{i}}^{\dagger}\mathbf{z}^{(i)} \stackrel{(2.3a)}{=} \mathbf{z}^{(i)}.$$

This completes the construction in Case 2.

To conclude, we iterate the above steps for  $i \in [m]$  and set  $\mathbf{x}^* = \mathbf{x}^{(m)}$  to obtain a vector that, by Theorem 3.1, is a minimizer for (POP). In fact, by varying the choice of  $\mathbf{z}^{(1)} \in {\{\mathbf{z}^{1,1}, \dots, \mathbf{z}^{1,r_1}\}}$  in the first step of the construction, we can recover at least  $r_1 = \operatorname{rank} \mathbb{M}^{\omega}_{\Delta_1}(\mathbf{y}) \geq r_{\min}$  distinct minimizers. Theorem 1.1 is proved.  $\Box$ 

#### 3.2. Additional remarks on Theorem 1.1. A few remarks are in order.

First, Theorem 1.1 and Theorem 3.1 hold also if one adds equality constraints to (POP) because each equality can always be replaced with two inequalities. This observation suffices to extend the theorems, but we refer to [18] for a description of sparse moment relaxations that explicitly handles equalities, which should be preferred for practical computations. Note also that the proof of Theorem 1.1 requires neither the feasible set of (POP) nor the sets  $K_i$  in (3.1) to be compact. In particular, it does not require the polynomial vectors  $\mathbf{g}_i$  in (POP) to define Archimedean quadratic modules of weighted sum-of-squares polynomials, which is usually assumed to ensure the asymptotic convergence of  $f_{\omega}^*$  to  $f^*$  [6, Theorem 5].

Second, the proof shows that we can extract at least  $r_1 = \operatorname{rank} \mathbb{M}_{\Delta_1}^{\omega}(\mathbf{y})$  minimizers of (POP). Since this number depends on the particular ordering of the cliques  $\Delta_1, \ldots, \Delta_m$ , we wonder if computational techniques from graph theory can be used to determine a clique ordering that satisfies (RIP) and maximizes  $r_1$ . In fact, one would like to find the clique ordering that maximizes the actual number of recoverable minimizers of (POP). This problem, however, has a combinatorial nature because one must count all possible ways to combine the 'clique' vectors  $\mathbf{z}^{i,j}$  into a 'global' vector  $\mathbf{x}^*$ , and thus appears to be challenging.

On the other hand, no optimization of the clique ordering is needed if an optimal solution  $\mathbf{y}$  of (MOM) satisfies the stronger rank conditions in (1.3): in this case, the construction in our proof produces exactly r minimizers of (POP) for any clique ordering satisfying (RIP). This is because the representations of  $\mathbf{y}_{\Delta_i}$  in (3.9) and (3.10) have  $r_i = s_{ij} = r$  and must therefore be the same, making the choice of  $\mathbf{z}^{(i)}$  unique in all but the first iteration, where one can choose from r distinct vectors.

Finally, to the best of our knowledge, all rank-based sufficient conditions for the finite convergence of (MOM) have been proved by explicitly constructing optimizers of (POP) that satisfy the assumptions of Theorem 3.1. What differs are the details of the construction and, more importantly, their underlying assumptions. In Theorem 1.1, we assume (RIP) and flat extension conditions on the moment matrices for each clique and for certain clique overlaps. Lasserre [10, Theorem 3.7] replaces (RIP) with a rank-one condition on all overlap moment matrices, and is able to recover exactly  $\prod_{i \in [m]} r_i$  minimizers. We wonder if different assumptions on the cliques and the ranks of the moment matrices can be used. We also wonder if finite convergence can be detected without extracting optimizers and applying Theorem 3.1. We discuss this point further in section 4.2 after presenting an example where  $f_{\omega}^* = f^*$ , but no minimizers can be extracted because both (RIP) and the rank-one conditions in Lasserre's theorem fail.

## 4. Necessity of (RIP)

We now demonstrate that, if one drops (RIP) from Theorem 1.1, then it may be impossible to construct a minimizer of (POP) from the representations (3.2) of the vectors  $\mathbf{y}_{\Delta_i}$ , even if the rank conditions in (1.1) are replaced by their stronger version (1.3). This observation disproves the claims in [18, Theorem 3.4].

4.1. The counterexample. Fix n = 3 and consider the cliques  $\Delta_1 = \{1, 2\}$ ,  $\Delta_2 = \{2, 3\}$  and  $\Delta_3 = \{1, 3\}$ , which do *not* satisfy (RIP). The trivariate POP

(4.1)  

$$f^* = \min_{\mathbf{x} \in \mathbb{R}^3} \quad (x_1 - x_2)^2 + (x_2^2 - 1)^2 + (x_3^2 - 1)$$

$$\text{s.t.} \quad g_1(\mathbf{x}_{\Delta_1}) := 3 - x_1^2 - x_2^2 \ge 0$$

$$g_2(\mathbf{x}_{\Delta_2}) := 3 - x_2^2 - x_3^2 \ge 0$$

$$g_3(\mathbf{x}_{\Delta_3}) := 3 - x_3^2 - x_1^2 \ge 0$$

has the form (POP) for our choice of cliques. It is clear that the global minimum  $f^* = 0$  is achieved by the four minimizers

(4.2) 
$$\mathbf{x}_1^* = (1, 1, 1), \qquad \mathbf{x}_2^* = (1, 1, -1), \\ \mathbf{x}_3^* = (-1, -1, 1), \qquad \mathbf{x}_4^* = (-1, -1, -1).$$

We construct the sparse moment relaxation (MOM) of (4.1) of order  $\omega = 2$ . Once the constraint  $\mathcal{L}_{\mathbf{y}}(1) = 1$  is enforced, the resulting optimization problem depends on a vector  $\mathbf{y}$  with 30 entries denoted  $y_{\alpha_1\alpha_2\alpha_3}$  and indexed by the exponents  $\alpha = (\alpha_1, \alpha_2, \alpha_3) \in \mathbb{N}_4^3$  supported on one of the three cliques. The objective function to be minimized is

$$(4.3) 2 + y_{200} - 2y_{110} - y_{020} + y_{040} - 2y_{002} + y_{004}.$$

Expressions for the moment and localizing matrices to be made positive semidefinite are given in appendix A.

The optimal value of this moment relaxation cannot be smaller than zero. This is because it is bounded below by the best lower bound on  $f^*$  certifiable by correlatively sparse sum-of-squares polynomials (see, e.g., [10]), which is zero because the polynomial being minimized in (4.1) is already such a sum of squares. With this observation, one can easily verify that a particular optimal solution of our moment relaxation is the vector  $\mathbf{y}$  with the following entries:

$$\begin{array}{rll} y_{100}=0, & y_{010}=0, & y_{001}=0, & y_{200}=1, & y_{110}=1, & y_{020}=1, \\ y_{101}=-1, & y_{011}=1, & y_{002}=1, & y_{300}=0, & y_{210}=0, & y_{120}=0, \\ (4.4) & y_{030}=0, & y_{201}=0, & y_{021}=0, & y_{102}=0, & y_{012}=0, & y_{003}=0, \\ y_{400}=1, & y_{310}=1, & y_{220}=1, & y_{130}=1, & y_{040}=1, & y_{301}=-1, \\ y_{031}=1, & y_{202}=1, & y_{022}=1, & y_{103}=-1, & y_{013}=1, & y_{004}=1. \end{array}$$

Similarly, using the expressions for the matrices  $\mathbb{M}^1_{\Delta_i}(\mathbf{y})$ ,  $\mathbb{M}^2_{\Delta_i \cap \Delta_j}$  and  $\mathbb{M}^1_{\Delta_i \cap \Delta_j}(\mathbf{y})$  given in appendix A, one verifies (with  $\omega = 2$  and  $d_1 = d_2 = d_3 = 1$ ) that

(4.5a) 
$$\operatorname{rank} \mathbb{M}_{\Delta_i}^{\omega}(\mathbf{y}) = \operatorname{rank} \mathbb{M}_{\Delta_i}^{\omega-d_i}(\mathbf{y}) = 2 \quad \forall i \in [3],$$

(4.5b) 
$$\operatorname{rank} \mathbb{M}_{\Delta_i \cap \Delta_j}^{\omega}(\mathbf{y}) = \operatorname{rank} \mathbb{M}_{\Delta_i \cap \Delta_j}^{\omega - 1}(\mathbf{y}) = 2 \quad \forall i, j \in [3].$$

Thus, the rank conditions in (1.3) hold with r = 2.

Now, as in the proof of Theorem 1.1, these rank conditions imply that the vectors  $\mathbf{y}_{\Delta_1}$ ,  $\mathbf{y}_{\Delta_2}$  and  $\mathbf{y}_{\Delta_3}$  (not stated for brevity) can be expressed as in (3.2) for  $r_1 = r_2 = r_3 = 2$ . Indeed, one finds that

(4.6a) 
$$\mathbf{y}_{\Delta_1} = \frac{1}{2} [(1,1)]_4 + \frac{1}{2} [(-1,-1)]_4$$

(4.6b) 
$$\mathbf{y}_{\Delta_2} = \frac{1}{2} [(1,1)]_4 + \frac{1}{2} [(-1,-1)]_4$$

(4.6c) 
$$\mathbf{y}_{\Delta_3} = \frac{1}{2} [(1,-1)]_4 + \frac{1}{2} [(-1,1)]_4$$

However, no vector  $\mathbf{x}^* \in \mathbb{R}^3$  simultaneously satisfies

(4.7a) 
$$P_{\Delta_1} \mathbf{x}^* \in \{(1,1), (-1,-1)\},\$$

(4.7b) 
$$P_{\Delta_2} \mathbf{x}^* \in \{(1,1), (-1,-1)\},\$$

(4.7c)  $P_{\Delta_3} \mathbf{x}^* \in \{(1, -1), (-1, 1)\}.$ 

In particular, none of the POP minimizers in (4.2) satisfy (4.7). It is then impossible to recover a minimizer of (4.1) from the representations in (4.6).

4.2. Additional comments. We conclude with two comments about our example. First, it is common to select the cliques for a correlative sparse POP as the maximal cliques of its sparsity graph [21], which has the integers  $\{1, \ldots, n\}$  as its vertices and an edge (i, j) if two variables  $x_i$  and  $x_j$  are either multiplied together in the objective, or appear together in at least one constraint. When applied to (4.1), this selection strategy gives a single clique  $\Delta = \{1, 2, 3\}$ , meaning that the POP is not correlatively sparse in the usual sense. However, there are examples with  $n \ge 4$  variables that are correlatively sparse in the usual sense and such that (MOM) admits optimal solutions satisfying the rank conditions in (1.3), but from which no minimizer can be recovered due the failure of (RIP). One such example is

(4.8)  

$$f^* = \min_{\mathbf{x} \in \mathbb{R}^4} (x_1 - x_2)^2 + (x_2 - x_3)^2 + (x_3^2 - 1)^2 + (x_4^2 - 1)^2$$

$$g_{\mathbf{x}, \mathbf{x}, \mathbf{$$

whose sparsity graph has maximal cliques  $\Delta_1 = \{1, 2\}, \Delta_2 = \{2, 3\}, \Delta_3 = \{3, 4\}$ and  $\Delta_4 = \{1, 4\}$  that do not satisfy (RIP). We omit the details for brevity. We note only that the sparsity graphs of this and other similar examples are necessarily not chordal, otherwise their maximal cliques would satisfy (RIP) [1].

Second, but perhaps more important, we have shown that, if (RIP) fails, then the rank conditions in (1.3) may not be sufficient to enable the extraction of minimizers for (POP). However, since  $f_{\omega}^* = f^*$  in our example, we cannot conclude that they do not imply the finite convergence of the sparse moment relaxation (MOM). Consequently, proving or disproving the following claim remains an open problem.

**Claim 4.1.** If **y** is an optimal solution of (MOM) and the rank conditions in (1.3) hold for some integer r > 1, then  $f_{\omega}^* = f^*$ .

We currently do not know if this claim is true or false. If true, however, a proof will require new techniques that avoid extracting minimizers for (POP) and thus, as mentioned in section 3.2, considerably depart from those available in the literature.

## APPENDIX A.

This appendix gives the expressions for various moment matrices used in section 4. They arise when constructing the sparse moment relaxation (MOM) of the POP (4.1) using the cliques  $\Delta_1 = \{1, 2\}, \Delta_2 = \{2, 3\}$  and  $\Delta_3 = \{1, 3\}$ . The expressions we report already take into account the constraint  $\mathcal{L}_{\mathbf{y}}(1) = 1$  in (MOM). Throughout this section, given a square matrix  $\mathbf{A}$ , we denote by  $\mathbf{A}_{[i_1,...,i_K]}$  its principal submatrix comprised of the  $i_1, \ldots, i_K$ -th rows and columns.

The moment matrices for the relaxation order  $\omega = 2$  are

$$\mathbb{M}^2_{\Delta_1}(\mathbf{y}) = egin{bmatrix} 1 & y_{100} & y_{010} & y_{200} & y_{110} & y_{020} \ y_{100} & y_{200} & y_{110} & y_{300} & y_{210} & y_{120} \ y_{010} & y_{110} & y_{020} & y_{210} & y_{120} & y_{030} \ y_{200} & y_{300} & y_{210} & y_{400} & y_{310} & y_{220} \ y_{110} & y_{210} & y_{120} & y_{310} & y_{220} & y_{130} \ y_{020} & y_{120} & y_{030} & y_{220} & y_{130} \ y_{020} & y_{120} & y_{030} & y_{220} & y_{130} \ y_{040} \end{bmatrix},$$

$$\mathbb{M}^{2}_{\Delta_{2}}(\mathbf{y}) = \begin{bmatrix} 1 & y_{010} & y_{001} & y_{020} & y_{011} & y_{002} \\ y_{010} & y_{020} & y_{011} & y_{030} & y_{021} & y_{012} \\ y_{001} & y_{011} & y_{002} & y_{021} & y_{012} & y_{003} \\ y_{020} & y_{030} & y_{021} & y_{040} & y_{031} & y_{022} \\ y_{011} & y_{021} & y_{012} & y_{031} & y_{022} & y_{013} \\ y_{002} & y_{012} & y_{300} & y_{022} & y_{013} & y_{004} \end{bmatrix},$$
$$\mathbb{M}^{2}_{\Delta_{3}}(\mathbf{y}) = \begin{bmatrix} 1 & y_{100} & y_{001} & y_{200} & y_{101} & y_{002} \\ y_{100} & y_{200} & y_{101} & y_{300} & y_{201} & y_{102} \\ y_{001} & y_{101} & y_{002} & y_{201} & y_{102} & y_{003} \\ y_{200} & y_{300} & y_{201} & y_{400} & y_{301} & y_{202} \\ y_{101} & y_{201} & y_{102} & y_{301} & y_{202} & y_{103} \\ y_{002} & y_{102} & y_{003} & y_{202} & y_{103} & y_{004} \end{bmatrix}.$$

The moment matrices for the relaxation order  $\omega = 1$  are then obtain from these expressions by taking only the leading  $3 \times 3$  submatrices, i.e.,

$$\begin{split} \mathbb{M}^{1}_{\Delta_{1}}(\mathbf{y}) &= \mathbb{M}^{2}_{\Delta_{1}}(\mathbf{y})_{[1,2,3]}, \\ \mathbb{M}^{1}_{\Delta_{2}}(\mathbf{y}) &= \mathbb{M}^{2}_{\Delta_{2}}(\mathbf{y})_{[1,2,3]}, \\ \mathbb{M}^{1}_{\Delta_{3}}(\mathbf{y}) &= \mathbb{M}^{2}_{\Delta_{3}}(\mathbf{y})_{[1,2,3]}. \end{split}$$

The clique overlap matrices for the relaxation order  $\omega = 2$  are

$$\begin{split} \mathbb{M}^{2}_{\Delta_{1}\cap\Delta_{2}}(\mathbf{y}) &= \mathbb{M}^{2}_{\Delta_{1}}(\mathbf{y})_{[1,3,6]} = \mathbb{M}^{2}_{\Delta_{2}}(\mathbf{y})_{[1,2,4]},\\ \mathbb{M}^{2}_{\Delta_{2}\cap\Delta_{3}}(\mathbf{y}) &= \mathbb{M}^{2}_{\Delta_{2}}(\mathbf{y})_{[1,3,6]} = \mathbb{M}^{2}_{\Delta_{3}}(\mathbf{y})_{[1,3,6]},\\ \mathbb{M}^{2}_{\Delta_{1}\cap\Delta_{3}}(\mathbf{y}) &= \mathbb{M}^{2}_{\Delta_{1}}(\mathbf{y})_{[1,2,4]} = \mathbb{M}^{2}_{\Delta_{3}}(\mathbf{y})_{[1,2,4]}. \end{split}$$

Again, their analogue for the relaxation order  $\omega = 1$  is obtained simply by taking the leading  $2 \times 2$  submatrices, i.e.,

$$\begin{split} \mathbb{M}^{1}_{\Delta_{1}\cap\Delta_{2}}(\mathbf{y}) &= \mathbb{M}^{2}_{\Delta_{1}\cap\Delta_{2}}(\mathbf{y})_{[1,2]},\\ \mathbb{M}^{1}_{\Delta_{2}\cap\Delta_{3}}(\mathbf{y}) &= \mathbb{M}^{2}_{\Delta_{2}\cap\Delta_{3}}(\mathbf{y})_{[1,2]},\\ \mathbb{M}^{1}_{\Delta_{1}\cap\Delta_{3}}(\mathbf{y}) &= \mathbb{M}^{2}_{\Delta_{1}\cap\Delta_{3}}(\mathbf{y})_{[1,2]}. \end{split}$$

Lastly, the localizing matrices associated to the inequality constraints of (4.1) for the relaxation order  $\omega = 2$  are given explicitly by

$$\mathbb{M}^{2}_{\Delta_{1}}(g_{1}\mathbf{y}) = \begin{bmatrix} 3 - y_{200} - y_{020} & 3y_{100} - y_{300} - y_{120} & 3y_{010} - y_{210} - y_{030} \\ 3y_{100} - y_{300} - y_{120} & 3y_{200} - y_{400} - y_{220} & 3y_{110} - y_{310} - y_{130} \\ 3y_{010} - y_{210} - y_{030} & 3y_{110} - y_{310} - y_{130} & 3y_{020} - y_{040} - y_{220} \end{bmatrix},$$
$$\mathbb{M}^{2}_{\Delta_{2}}(g_{2}\mathbf{y}) = \begin{bmatrix} 3 - y_{020} - y_{002} & 3y_{010} - y_{030} - y_{012} & 3y_{001} - y_{021} - y_{003} \\ 3y_{010} - y_{030} - y_{012} & 3y_{020} - y_{040} - y_{022} & 3y_{011} - y_{031} - y_{013} \\ 3y_{001} - y_{021} - y_{003} & 3y_{011} - y_{031} - y_{013} & 3y_{002} - y_{004} - y_{022} \end{bmatrix},$$
$$\mathbb{M}^{2}_{\Delta_{3}}(g_{3}\mathbf{y}) = \begin{bmatrix} 3 - y_{200} - y_{002} & 3y_{100} - y_{300} - y_{102} & 3y_{001} - y_{201} - y_{003} \\ 3y_{100} - y_{300} - y_{102} & 3y_{200} - y_{400} - y_{202} & 3y_{101} - y_{201} - y_{003} \\ 3y_{001} - y_{201} - y_{003} & 3y_{101} - y_{301} - y_{103} & 3y_{002} - y_{004} - y_{202} \end{bmatrix}.$$

## References

- J. R. S. Blair and B. Peyton. An introduction to chordal graphs and clique trees. In Graph theory and sparse matrix computation, volume 56 of IMA Vol. Math. Appl., pages 1–29. Springer, New York, 1993.
- [2] G. Blekherman and C. Riener. Symmetric non-negative forms and sums of squares. Discrete Comput. Geom., 65(3):764-799, 2021.
- [3] R. E. Curto and L. A. Fialkow. The truncated complex K-moment problem. Trans. Amer. Math. Soc., 352(6):2825–2855, 2000.

- [4] L. A. Fialkow. The truncated K-moment problem: A survey. In Operator theory: the state of the art, volume 18 of Theta Ser. Adv. Math., pages 25–51. Theta, Bucharest, 2016.
- [5] K. Gatermann and P. A. Parrilo. Symmetry groups, semidefinite programs, and sums of squares. J. Pure Appl. Algebra, 192(1-3):95–128, 2004.
- [6] D. Grimm, T. Netzer, and M. Schweighofer. A note on the representation of positive polynomials with structured sparsity. Arch. Math. (Basel), 89(5):399–403, 2007.
- [7] D. Henrion and J.-B. Lasserre. Detecting global optimality and extracting solutions in GloptiPoly. In *Positive polynomials in control*, volume 312 of *Lect. Notes Control Inf. Sci.*, pages 293–310. Springer, Berlin, 2005.
- [8] M. Kojima and M. Muramatsu. A note on sparse SOS and SDP relaxations for polynomial optimization problems over symmetric cones. *Comput. Optim. Appl.*, 42 (1):31–41, 2009.
- J. B. Lasserre. Global optimization with polynomials and the problem of moments. SIAM J. Optim., 11(3):796–817, 2001.
- [10] J. B. Lasserre. Convergent SDP-relaxations in polynomial optimization with sparsity. SIAM J. Optim., 17(3):822–843, 2006.
- [11] M. Laurent. Revisiting two theorems of Curto and Fialkow on moment matrices. Proc. Amer. Math. Soc., 133(10):2965–2976, 2005.
- [12] M. Laurent. Sums of squares, moment matrices and optimization over polynomials, volume 149 of IMA Vol. Math. Appl., pages 157–270. Springer, New York, 2009.
- [13] J. Löfberg. Pre-and post-processing sum-of-squares programs in practice. IEEE Trans. Automatic Control, 54:1007–1011, 2009.
- [14] V. Magron and J. Wang. Sparse polynomial optimization—theory and practice, volume 5 of Series on Optimization and its Applications. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2023.
- [15] J. Miller, J. Wang, and F. Guo. Sparse polynomial matrix optimization. arXiv:2411.15479 [math.OC], 2024.
- [16] J. Nie. Certifying convergence of Lasserre's hierarchy via flat truncation. Math. Program., 142(1-2, Ser. A):485–510, 2013.
- [17] J. Nie. Optimality conditions and finite convergence of Lasserre's hierarchy. Math. Program., 146(1-2, Ser. A):97–121, 2014.
- [18] J. Nie, Z. Qu, X. Tang, and L. Zhang. A characterization for tightness of the sparse moment-SOS hierarchy. arXiv:2406.06882 [math.OC], 2024.
- [19] P. A. Parrilo. Semidefinite programming relaxations for semialgebraic problems. Math. Program., 96(2, Ser. B):293–320, 2003.
- [20] C. Riener, T. Theobald, L. J. Andrén, and J. B. Lasserre. Exploiting symmetries in SDP-relaxations for polynomial optimization. *Math. Oper. Res.*, 38(1):122–141, 2013.
- [21] H. Waki, S. Kim, M. Kojima, and M. Muramatsu. Sums of squares and semidefinite program relaxations for polynomial optimization problems with structured sparsity. *SIAM J. Optim.*, 17(1):218–242, 2006.
- [22] J. Wang, H. Li, and B. Xia. A new sparse SOS decomposition algorithm based on term sparsity. In ISSAC'19—Proceedings of the 2019 ACM International Symposium on Symbolic and Algebraic Computation, pages 347–354. ACM, New York, 2019.
- [23] J. Wang, V. Magron, and J.-B. Lasserre. TSSOS: a moment-SOS hierarchy that exploits term sparsity. SIAM J. Optim., 31(1):30–58, 2021.
- [24] J. Wang, V. Magron, and J.-B. Lasserre. Chordal-TSSOS: a moment-SOS hierarchy that exploits term sparsity with chordal extension. SIAM J. Optim., 31(1):114–141, 2021.
- [25] J. Wang, V. Magron, J. B. Lasserre, and N. H. A. Mai. CS-TSSOS: correlative and term sparsity for large-scale polynomial optimization. ACM Trans. Math. Software, 48(4):Art. 42, 26, 2022.
- [26] Y. Zheng, G. Fantuzzi, and A. Papachristodoulou. Chordal and factor-width decompositions for scalable semidefinite and polynomial optimization. *Annu. Rev. Control*, 52:243–279, 2021.