

Local Quantum Mechanical Prediction of the Singlet State

Carl F. Diether, III*

Einstein Centre for Local-Realistic Physics, N. California, USA Branch

(Dated: 2-1-2025)

We deduce the quantum mechanical prediction of $-\mathbf{a} \cdot \mathbf{b}$ for the singlet spin state employing local measurement functions following Bell's approach. Our derivation is corroborated through a computational simulation conducted via the Mathematica programming environment.

In this manuscript, we elucidate the quantum mechanical (QM) prediction for the singlet spin state in the context of the EPR-Bohm thought experiment (EPRB) [1, 4], employing local measurement functions in accordance with the methodologies of Bell [2] and Christian [3]. Within the framework of the EPRB experiment, we consider two spin one-half quantum particles (an electron and a positron) propagating in opposite directions subsequent to their generation from a singlet state particle. Subsequently, at Stations A and B—situated at a spacelike separation—measurements of their spins are conducted along independently chosen unit vectors \mathbf{a} and \mathbf{b} . We posit that the progenitor particle of the pair possesses zero spin, thereby endowing the pair with the following singlet wavefunction:

$$|\Psi_{\mathbf{n}}\rangle = \frac{1}{\sqrt{2}} \left\{ |\mathbf{n}, +\rangle_1 \otimes |\mathbf{n}, -\rangle_2 - |\mathbf{n}, -\rangle_1 \otimes |\mathbf{n}, +\rangle_2 \right\}, \quad (1)$$

where

$$\boldsymbol{\sigma} \cdot \mathbf{n} |\mathbf{n}, \pm\rangle = \pm |\mathbf{n}, \pm\rangle, \quad (2)$$

characterizes the quantum mechanical eigenstates where the particles exhibit spin in the "up" or "down" orientation measured in units of $\hbar = 2$. Here, $\boldsymbol{\sigma}$ denotes the Pauli spin "vector" ($\sigma_x, \sigma_y, \sigma_z$), while \mathbf{n} represents the particle's spin vector oriented in an arbitrary direction.

The rotational symmetry inherent in the singlet state $\Psi_{\mathbf{n}}$ guarantees that the expectation values for the individual spin observables $\boldsymbol{\sigma} \cdot \mathbf{a}$ and $\boldsymbol{\sigma} \cdot \mathbf{b}$ are

$$\begin{aligned} \mathcal{E}_{q.m.}(\mathbf{a}) &= \langle \Psi_{\mathbf{n}} | \boldsymbol{\sigma} \cdot \mathbf{a} \otimes \mathbb{1} | \Psi_{\mathbf{n}} \rangle \\ &= \langle \Psi_{\mathbf{n}} | \boldsymbol{\sigma} \cdot \mathbf{a} | \Psi_{\mathbf{n}} \rangle = 0 \end{aligned} \quad (3)$$

$$\begin{aligned} \text{and } \mathcal{E}_{q.m.}(\mathbf{b}) &= \langle \Psi_{\mathbf{n}} | \mathbb{1} \otimes \boldsymbol{\sigma} \cdot \mathbf{b} | \Psi_{\mathbf{n}} \rangle \\ &= \langle \Psi_{\mathbf{n}} | \boldsymbol{\sigma} \cdot \mathbf{b} | \Psi_{\mathbf{n}} \rangle = 0, \end{aligned} \quad (4)$$

where $\mathbb{1}$ is the identity matrix. The expectation value of the joint observable $\boldsymbol{\sigma} \cdot \mathbf{a} \otimes \boldsymbol{\sigma} \cdot \mathbf{b}$ is [5]

$$\mathcal{E}_{q.m.}(\mathbf{a}, \mathbf{b}) = \langle \Psi_{\mathbf{n}} | \boldsymbol{\sigma} \cdot \mathbf{a} \otimes \boldsymbol{\sigma} \cdot \mathbf{b} | \Psi_{\mathbf{n}} \rangle = -\mathbf{a} \cdot \mathbf{b}, \quad (5)$$

regardless of the relative distance between the two remote locations represented by the unit vectors \mathbf{a} and \mathbf{b} .

We shall now formulate certain explicitly local measurement functions, akin to those in Bell's framework [2]. This approach yields the aforementioned result of $-\mathbf{a} \cdot \mathbf{b}$ and aligns with the eigenvalues of observable operators that entail spins being detected by detectors, with the division of the singlet vector governed by the conservation of spin angular momentum, as discussed in $\mathbf{s}_1 + \mathbf{s}_2 = 0$, so that:

$$\mathbf{n} = \mathbf{s} = \mathbf{s}_1 = -\mathbf{s}_2. \quad (6)$$

In the interaction of two vectors, the resultant is composed of a scalar, $(\mathbf{u} \cdot \mathbf{v})$, in addition to a vector via a cross-product, $(\mathbf{u} \times \mathbf{v}) = \mathbf{r}$, which may collectively be represented as a quaternion, $\mathbf{q}(\mathbf{u} \cdot \mathbf{v}, \mathbf{r})$, as a single entity. Employing this notion in conjunction with eq.(6), we define

$$\mathbf{r}_a = \frac{\mathbf{a} \times \mathbf{s}_1}{\|\mathbf{a} \times \mathbf{s}_1\|} \quad \text{and} \quad \mathbf{r}_b = \frac{\mathbf{s}_2 \times \mathbf{b}}{\|\mathbf{s}_2 \times \mathbf{b}\|}, \quad (7)$$

*Electronic address: fred.diether@einstein-physics.org

$$\mu_{\mathbf{a}} = \text{sgn}(\mathbf{a} \cdot \mathbf{s}_1) \mathbf{a}, \quad (8)$$

$$\mu_{\mathbf{b}} = \text{sgn}(\mathbf{b} \cdot \mathbf{s}_2) \mathbf{b}, \quad (9)$$

$$\begin{aligned} A(\mathbf{a}, \mathbf{s}_1) &:= \lim_{\mathbf{s}_1 \rightarrow \mu_{\mathbf{a}}} \left[\langle \phi_{\mathbf{n}} | (\boldsymbol{\sigma} \cdot \mathbf{a}) (\boldsymbol{\sigma} \cdot \mathbf{s}_1) | \phi_{\mathbf{n}} \rangle + i \mathbf{r}_a \right], \\ &= \lim_{\mathbf{s}_1 \rightarrow \mu_{\mathbf{a}}} \left[\mathbf{a} \cdot \mathbf{s}_1 + i \mathbf{r}_a \right], \\ &= \lim_{\mathbf{s}_1 \rightarrow \mu_{\mathbf{a}}} \left[\mathbf{q}_a(\mathbf{a} \cdot \mathbf{s}_1, \mathbf{r}_a) \right], \\ &= \text{sgn}(\mathbf{a} \cdot \mathbf{s}_1) = \pm 1, \end{aligned} \quad (10)$$

$$\begin{aligned} B(\mathbf{b}, \mathbf{s}_2) &:= \lim_{\mathbf{s}_2 \rightarrow \mu_{\mathbf{b}}} \left[\langle \chi_{\mathbf{n}} | (\boldsymbol{\sigma} \cdot \mathbf{s}_2) (\boldsymbol{\sigma} \cdot \mathbf{b}) | \chi_{\mathbf{n}} \rangle + i \mathbf{r}_b \right], \\ &= \lim_{\mathbf{s}_2 \rightarrow \mu_{\mathbf{b}}} \left[\mathbf{s}_2 \cdot \mathbf{b} + i \mathbf{r}_b \right], \\ &= \lim_{\mathbf{s}_2 \rightarrow \mu_{\mathbf{b}}} \left[\mathbf{q}_b(\mathbf{s}_2 \cdot \mathbf{b}, \mathbf{r}_b) \right], \\ &= \text{sgn}(\mathbf{s}_2 \cdot \mathbf{b}) = \mp 1, \end{aligned} \quad (11)$$

where

$$|\phi_{\mathbf{n}}\rangle = \frac{1}{\sqrt{2}} \left\{ |\mathbf{n}, +\rangle_1 |\mathbf{n}, -\rangle_1 \right\} \quad (12)$$

$$\text{and } |\chi_{\mathbf{n}}\rangle = \frac{1}{\sqrt{2}} \left\{ |\mathbf{n}, +\rangle_2 |\mathbf{n}, -\rangle_2 \right\}. \quad (13)$$

The derivation of the expectation value of the matrix calculation pertaining to the measurement functions A and B is detailed in the Appendix. In this context, $\boldsymbol{\sigma} \cdot \mathbf{a}$ and $\boldsymbol{\sigma} \cdot \mathbf{b}$ are designated as the detectors utilized by Alice and Bob, respectively, each possessing no angular momentum at the instant of detection. $\boldsymbol{\sigma} \cdot \mathbf{s}_1 = -\boldsymbol{\sigma} \cdot \mathbf{s}_2$ denotes the spin of the fermions these detectors receive, which constitutes the basis for conducting the EPRB experiment. The replacement limit functions simulate the functioning of the polarizers at the detection stations, while $|\phi_{\mathbf{n}}\rangle$ and $|\chi_{\mathbf{n}}\rangle$ are elementary products now symbolizing the wavefunction of the individual particles. The original singlet state is thus partitioned between two distinct simple product bra-kets. In the subsequent stage of the A and B functions, it becomes apparent that quaternions are delineated from the preceding stage. Although the measurement functions exhibit a deterministic appearance, quantum mechanics precludes the prediction of outcomes on an event-by-event basis. An analytical depiction of these outcomes, following proper evaluation, does not yield the negative cosine curve. The probability remains constant at fifty percent for either +1 or -1 outcomes (up or down). The cross products contained within the functions neither influence this probability nor alter the $-\mathbf{a} \cdot \mathbf{b}$ expectation value through the product of the functions. As the replacement limit functions mimic the role of detection polarizers, over numerous events, the mean action should converge to approximately fifty percent.

The measurement functions delineate the detection processes occurring at two potentially spacelike-separated observation stations belonging to Alice and Bob. Despite possibly transpiring concurrently, $A(\mathbf{a}, \mathbf{s}_1)$ and $B(\mathbf{b}, \mathbf{s}_2)$ constitute independent physical processes that are *not* constrained by the conservation of the initial zero spin angular momentum. In advancing the AB product calculation, the k indices are omitted subsequent to the initial step. Employing the "product of limits equal to limits of product" principle results in the computation of the expectation value as follows [8]:

$$\mathcal{E}(\mathbf{a}, \mathbf{b}) = \lim_{n \gg 1} \left[\frac{1}{n} \sum_{k=1}^n A(\mathbf{a}^k, \mathbf{s}_1^k) B(\mathbf{b}^k, \mathbf{s}_2^k) \right], \quad (14)$$

$$= \lim_{n \gg 1} \left\{ \frac{1}{n} \sum_{k=1}^n \left[\lim_{\mathbf{s}_1 \rightarrow \mu_{\mathbf{a}}} \{ \mathbf{q}_a(\mathbf{a} \cdot \mathbf{s}_1, \mathbf{r}_a) \} \right] \left[\lim_{\mathbf{s}_2 \rightarrow \mu_{\mathbf{b}}} \{ \mathbf{q}_b(\mathbf{s}_2 \cdot \mathbf{b}, \mathbf{r}_b) \} \right] \right\}, \quad (15)$$

$$= \lim_{n \gg 1} \left[\frac{1}{n} \sum_{k=1}^n \lim_{\substack{\mathbf{s}_1 \rightarrow \mu_{\mathbf{a}} \\ \mathbf{s}_2 \rightarrow \mu_{\mathbf{b}}}} \left\{ \mathbf{q}_a(\mathbf{a} \cdot \mathbf{s}_1, \mathbf{r}_a) \mathbf{q}_b(\mathbf{s}_2 \cdot \mathbf{b}, \mathbf{r}_b) \right\} \right], \quad (16)$$

$$= \lim_{n \gg 1} \left[\frac{1}{n} \sum_{k=1}^n \lim_{\substack{\mathbf{s}_1 \rightarrow \mu_a \\ \mathbf{s}_2 \rightarrow \mu_b}} \left\{ \text{Re}(\mathbf{q}_a) \text{Re}(\mathbf{q}_b) - \text{Im}(\mathbf{q}_a) \cdot \text{Im}(\mathbf{q}_b) + \mathbf{r}_0 \right\} \right], \quad (17)$$

$$= -\cos(\theta_{\mathbf{ab}}) + \lim_{n \gg 1} \left[\frac{1}{n} \sum_{k=1}^n \lim_{\substack{\mathbf{s}_1 \rightarrow \mu_a \\ \mathbf{s}_2 \rightarrow \mu_b}} \left\{ \mathbf{r}_0 \right\} \right], \quad (18)$$

$$= -\cos(\theta_{\mathbf{ab}}) + \vec{\mathbf{0}}, \quad (19)$$

$$= -\mathbf{a} \cdot \mathbf{b}, \quad (20)$$

$$\text{where } \mathbf{r}_0(\mathbf{s}_1, \mathbf{s}_2) = \frac{(\mathbf{a} \cdot \mathbf{s}_1)(\mathbf{s}_2 \times \mathbf{b}) + (\mathbf{s}_2 \cdot \mathbf{b})(\mathbf{a} \times \mathbf{s}_1) - (\mathbf{a} \times \mathbf{s}_1) \times (\mathbf{s}_2 \times \mathbf{b})}{\sin(\theta_{\mathbf{ab}})}. \quad (21)$$

In step (17), we have employed an identity pertinent to the multiplication of two quaternions, as referenced in [7, 8]. Consequently, the product of two quaternions results in another quaternion, and it is noted that \mathbf{r}_0 encompasses all cross products in the numerator which become zero upon taking the limits. In step (18), the limit replacement functions are applied exclusively to the cross-products of \mathbf{r}_0 , resulting in a null vector. Consequently, the desired result is obtained through a wholly local process. It is apparent that quantum mechanics is perhaps completed by the 3-sphere topology discussed in [7, 8] through the utilization of quaternions, and that the singlet exhibits a parallelized 3-sphere topology that is transmitted to the particle pair [3, 6–8]. This demonstration can be readily achieved through the application of Pauli algebra.

We have verified the above analytical calculation through a computer simulation using Mathematica programming language [9] and the result can be seen in Figure 1. The Mathematica notebook file is also available at [9].




FIG. 1: Plot of product calculation from the simulation. Blue is the correlation data, magenta is the negative cosine curve for an exact match.

As demonstrated, it is feasible to arrive at a solution for a local QM result. The solution is achieved through the application of 3-sphere topology using quaternions. This discourse addresses the presumptions made by John Bell regarding the non-local nature of quantum mechanics in the EPRB scenario. Nonetheless, Bell provided valuable insights into the formulation of local measurement functions. Subsequently, Joy Christian refined the formulation by incorporating limit replacement functions, which facilitated accurate product calculations [3, 6–8]. It is probable that previous assertions claiming "ordinary quantum mechanics is not locally causal" are incorrect [10].

Acknowledgments

The author thanks Joy Christian and Jay Yablon, who contributed much to this paper via many discussions.

Appendix: Matrix Expectation Value Calculation for Functions A and B

We demonstrate here the mathematical steps for the calculation of the matrix expectation value in the measurement functions that produce cosine with $\mathbf{s}_1 = \mathbf{s}$.

$$\mathcal{E}(\cos(\theta_{\mathbf{as}_1})) = \langle \phi_{\mathbf{n}} | (\boldsymbol{\sigma} \cdot \mathbf{a}) (\boldsymbol{\sigma} \cdot \mathbf{s}_1) | \phi_{\mathbf{n}} \rangle \quad (\text{A.1})$$

$$= \frac{1}{2} \begin{pmatrix} 1 & 0 \end{pmatrix} \cdot \begin{pmatrix} a_x s_x + a_y s_y + a_z s_z + i a_x s_y - i a_y s_x & a_z s_x - a_x s_z - i a_z s_y + i a_y s_z \\ -a_z s_x + a_x s_z - i a_z s_y + i a_y s_z & a_x s_x + a_y s_y + a_z s_z - i a_x s_y + i a_y s_x \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (\text{A.2})$$

$$+ \begin{pmatrix} 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} a_x s_x + a_y s_y + a_z s_z + i a_x s_y - i a_y s_x & a_z s_x - a_x s_z - i a_z s_y + i a_y s_z \\ -a_z s_x + a_x s_z - i a_z s_y + i a_y s_z & a_x s_x + a_y s_y + a_z s_z - i a_x s_y + i a_y s_x \end{pmatrix} \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (\text{A.3})$$

$$= \frac{1}{2} ((a_x s_x + a_y s_y + a_z s_z + i a_x s_y - i a_y s_x) + (a_x s_x + a_y s_y + a_z s_z - i a_x s_y + i a_y s_x)) \quad (\text{A.4})$$

$$= a_x s_x + a_y s_y + a_z s_z = \mathbf{a} \cdot \mathbf{s}_1 \quad (\text{A.5})$$

Note that the calculation for Bob's station is similar giving $\mathbf{s}_2 \cdot \mathbf{b}$. Since there is only one particle per detection station, it is either "up" or "down" upon detection.

-
- [1] D. Bohm and Y. Aharonov, *Discussions of Experimental Proof for the Paradox of Einstein, Rosen, and Podolsky*, Phys. Rev., **108**, 1079 (1957).
- [2] J. S. Bell, *On the Einstein-Podolsky-Rosen paradox*, Physics, vol. 1, pp. 195–200 (1964).
- [3] J. Christian, *Quantum correlations are weaved by the spinors of the Euclidean primitives*. R. Soc. Open Sci., **5**, 180526; doi:10.1098/rsos.180526 (2018), arXiv:1806.02392; See also: *Eight-dimensional Octonion-like but Associative Normed Division Algebra*, arXiv:1908.06172 (2019).
- [4] A. Einstein, B. Podolsky and N. Rosen. *Can Quantum-Mechanical Description of Physical Reality be Considered Complete?*. Phys. Rev., **1935**, 47 (10), 777–780.
- [5] J. Yablon, *Why Quantum Mechanics is Local and Realistic with Uncertainty-Hidden Variables, without Instantaneous Action at a Distance, and is Not Incomplete*, [Online] Available: <https://jayryablon.files.wordpress.com/2019/10/lrhvcqm-1.1.pdf>, (2019).
- [6] J. Christian, *Bell's Theorem Versus Local Realism in a Quaternionic Model of Physical Space*, IEEE Access, **7**, 133388; doi:10.1109/ACCESS.2019.2941275 (2019); see also arXiv:1405.2355 (2014).
- [7] J. Christian, *Dr. Bertlmann's Socks in a Quaternionic World of Ambidextral Reality*, IEEE Access, **8**, 191028; doi:10.1109/ACCESS.2020.3031734 (2020).
- [8] J. Christian, *Local origins of quantum correlations rooted in geometric algebra*, doi:10.13140/RG.2.2.35755.72480 (2022).
- [9] C. F. Diether III, *Simulation of the 3-Sphere Model for the singlet correlations*, [Online] Available: <https://notebookarchive.org/local-quantum-mechanical-prediction-of-the-singlet-state--2022-04-1eu4roy> or <https://sciphysicsfoundations.com/viewtopic.php?p=886#p886>, accessed Jul., 10 (2022).
- [10] J.S. Bell, *Speakable and Unsayable in Quantum Mechanics*. Cambridge University Press, **2004**.