MATRIX SYSTEMS, ALGEBRAS, AND OPEN MAPS

STEPHAN WEIS

ABSTRACT. Every state on the algebra M_n of complex $n \times n$ matrices restricts to a state on any matrix system. Whereas the restriction to a matrix system is generally not open, we prove that the restriction to every *-subalgebra of M_n is open. This simplifies topology problems in matrix theory and quantum information theory.

In honor of Ilya Matveevich Spitkovsky, for his 70th birthday.

1. INTRODUCTION

In the work of Choi and Effros [15], a matrix system on \mathbb{C}^n is a complex linear subspace \mathcal{R} of the full matrix algebra M_n that is self-adjoint (the conjugate transpose A^* of every $A \in \mathcal{R}$ lies in \mathcal{R}) and contains the $n \times n$ identity matrix $\mathbb{1}_n$, see also [5,42]. Let \mathcal{R} be a matrix system on \mathbb{C}^n . If \mathcal{R} is closed under matrix multiplication we call it a *-subalgebra of M_n . The dual space to \mathcal{R} is denoted by $\mathcal{R}^* := \{\ell : \mathcal{R} \to \mathbb{C} \mid \ell \text{ is } \mathbb{C}\text{-linear}\}$ and the cone of positive semidefinite matrices in \mathcal{R} by $\mathcal{C}(\mathcal{R})$. The state space of \mathcal{R} is

$$\mathcal{S}(\mathcal{R}) := \left\{ \ell \in \mathcal{R}^* \mid \forall A \in \mathcal{C}(\mathcal{R}) : \ell(A) \ge 0, \ell(\mathbb{1}_n) = 1 \right\}.$$

The restriction $\mathcal{S}(\mathcal{M}_n) \to \mathcal{S}(\mathcal{R})$, $\ell \mapsto \ell|_{\mathcal{R}}$ of states to \mathcal{R} is continuous and affine. Its analytic properties would be perfectly clear if it were not for the openness that fails in Example 1.1. Let K, L be subsets of some Euclidean spaces endowed with their relative topologies [27]. A map $f : K \to L$ is open at $x \in K$ if the image of every neighborhood of x in K is a neighborhood of f(x) in L. The map f is open if it is open at every point in K.

It is helpful to represent states as matrices. The antilinear isomorphism $\mathcal{R} \to \mathcal{R}^*, A \mapsto \langle A, \cdot \rangle$ restricts by Lemma 2.2 to the affine isomorphism

$$r_{\mathcal{R}}: \mathcal{D}(\mathcal{R}) \to \mathcal{S}(\mathcal{R}), \quad \rho \mapsto \langle \rho, \cdot \rangle,$$

where $\langle A, B \rangle := \operatorname{tr}(A^*B)$ is the Frobenius inner product of $A, B \in M_n$,

$$\begin{split} \mathcal{H}(\mathcal{R}) &:= \left\{ A \in \mathcal{R} \mid A^* = A \right\}, \\ \mathcal{C}(\mathcal{R})^{\vee} &:= \left\{ A \in \mathcal{H}(\mathcal{R}) \mid \forall B \in \mathcal{C}(\mathcal{R}) : \langle A, B \rangle \geq 0 \right\}, \\ \text{nd} \quad \mathcal{D}(\mathcal{R}) &:= \left\{ \rho \in \mathcal{C}(\mathcal{R})^{\vee} \mid \operatorname{tr}(\rho) = 1 \right\}. \end{split}$$

Generalizing a term of von Neumann algebras [10], we refer to the elements of $\mathcal{D}(\mathcal{R})$ as *density matrices*. The inclusion $\mathcal{C}(\mathcal{R}) \subset \mathcal{C}(\mathcal{R})^{\vee}$ can be strict. For example, the density matrix diag(-1, 5, 2)/6 of the matrix system spanned by $\mathbb{1}_3$ and diag(1, -1, 0) is indefinite. It is well known, see Rem. 2.8, that the identity $\mathcal{C}(\mathcal{A}) = \mathcal{C}(\mathcal{A})^{\vee}$ holds for every *-subalgebra \mathcal{A} of M_n .

a

Date: Draft of February 5, 2025.

²⁰²⁰ Mathematics Subject Classification. 46A30, 15A60, 47A12, 81P16.

Key words and phrases. open map, numerical range, matrix system, quantum state.

Example 1.1. We write block diagonal matrices as direct sums, for instance

$$\begin{pmatrix} A & 0 \\ 0 & c \end{pmatrix} = A \oplus c \in \mathcal{M}_3 \quad \text{for every} \quad A \in \mathcal{M}_2 \,, \quad c \in \mathbb{C} \cong \mathcal{M}_1 \,.$$

Denoting the imaginary unit by $i \in \mathbb{C}$ and the *Pauli matrices* by

$$X := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad Y := \begin{pmatrix} 0 & -\mathbf{i} \\ \mathbf{i} & 0 \end{pmatrix}, \quad Z := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

we define the matrix system $\mathcal{R} := \operatorname{span}_{\mathbb{C}}(\mathbb{1}_3, X \oplus 1, Z \oplus 0)$. The orthogonal projection of \mathbb{C}^2 onto the line spanned by $|+\rangle := (1,1)^T/\sqrt{2}$ is written $|+\rangle\langle+|$ in *Dirac's notation* [7,11]. The open set $\mathcal{O} := \{\ell \in \mathcal{S}(M_3) \mid \ell(0 \oplus 1) > 0\}$ contains

$$\omega_{\lambda} := r_{\mathrm{M}_{3}} \left[(1 - \lambda) \left| + \right\rangle \! \left| + \right\rangle \! \left| \oplus \lambda \right], \quad \lambda \in (0, 1].$$

So $\mathcal{O}|_{\mathcal{R}} := \{\ell|_{\mathcal{R}} : \ell \in \mathcal{O}\}$ contains $\omega_{\lambda}|_{\mathcal{R}}$ but none of the restriction $\ell_{\theta}|_{\mathcal{R}}$ of

$$\ell_{\theta} := r_{\mathrm{M}_{3}} \left[\frac{1}{2} \left(\mathbb{1}_{2} + \cos(\theta) X + \sin(\theta) Z \right) \oplus 0 \right], \quad \theta \in (0, 2\pi).$$

Specifically, ℓ_{θ} has the value $\ell_{\theta}(A_{\theta}) = 1$ at $A_{\theta} := \cos(\theta)(X \oplus 1) + \sin(\theta)Z \oplus 0$ and $\ell(A_{\theta}) \leq (\cos(\theta) - 1)\ell(0 \oplus 1) + 1$ holds for every $\ell \in \mathcal{S}(M_3)$. Hence

$$\ell|_{\mathcal{R}}(A_{\theta}) - \ell_{\theta}|_{\mathcal{R}}(A_{\theta}) \le (\cos(\theta) - 1)\ell(0 \oplus 1) < 0, \quad \ell \in \mathcal{O}.$$

This shows that $\mathcal{O}|_{\mathcal{R}}$ is not a neighborhood of $\omega_{\lambda}|_{\mathcal{R}}$ as $\lim_{\theta \to 0} \ell_{\theta}|_{\mathcal{R}} = \omega_{\lambda}|_{\mathcal{R}}$. In conclusion, $\mathcal{S}(M_3) \to \mathcal{S}(\mathcal{R}), \ \ell \mapsto \ell|_{\mathcal{R}}$ is not open at ω_{λ} for any $\lambda \in (0, 1]$.

Asking where $\mathcal{S}(M_n) \to \mathcal{S}(\mathcal{R}), \ell \mapsto \ell|_{\mathcal{R}}$ is open is the same, by Coro. 2.7 b), as inquiring at which density matrices the orthogonal projection

$$\mathcal{D}(\mathcal{M}_n) \to \mathcal{D}(\mathcal{R})$$
 (1.1)

is open. The map (1.1) is defined in (2.2) and (2.3) below. For now suffice it to say that it is a restriction of the orthogonal projection of M_n onto \mathcal{R} .

Corey et al. [18] and Leake et al. [29, 30] first studied a problem of numerical ranges closely related to (1.1). The problem (1.1) was studied by Weis [52, 53] and Rodman et al. [46] when $\mathcal{D}(\mathcal{R})$ is replaced with the affinely isomorphic joint numerical range. Numerical ranges are the topic of Sec. 4.

Theorem 1.2. If \mathcal{A} is a *-subalgebra of M_n , then the orthogonal projection $\mathcal{D}(M_n) \to \mathcal{D}(\mathcal{A})$ is open.

Thm. 1.2 can simplify the problem (1.1), and related continuity problems, if \mathcal{R} is included in a *-subalgebra of M_n smaller than M_n , as we show in Sec. 8. Examples from quantum information theory are presented in Sec. 9.

Thm. 1.2 is proved in Sec. 6. The main ideas are that \mathcal{A} is a direct sum of full matrix algebras and that $\mathcal{D}(M_n)$ is stable and highly symmetric. Thereby, a convex set K is *stable* if the *midpoint map*

$$K \times K \to K$$
, $(x, y) \mapsto \frac{1}{2}(x+y)$

is open. A convex set is always (except Rem. 1.3) understood to be included in a Euclidean space. Debs [20] proved the stability of $\mathcal{D}(\mathcal{A})$ for $\mathcal{A} = M_n$, Papadopoulou [41] achieved it for *-subalgebras \mathcal{A} of M_n , and we do it for *real* *-subalgebras \mathcal{A} of M_n in Sec. 5 using Clausing's work on retractions [16]. The analogue of Thm. 1.2 is established for the algebra $\mathcal{A} = M_n(\mathbb{R})$ of real $n \times n$ matrices in Sec. 7. The Secs. 2 and 3 collect preliminaries.

- **Remark 1.3.** a) Vesterstrøm [51] proved that the restriction of states to the center of a von Neumann algebra is open. Thm. 1.2 is a noncommutative analogue in the finite-dimensional setting.
- b) Stability is a meaningful concept in optimal control [41,57] and quantum information theory [48] because of the following "CE-property". A compact convex set K is stable if and only if for every continuous function $f: K \to \mathbb{R}$, the envelope $f^{\vee}(x) := \sup\{h(x) : h \leq f\}, x \in K$ is continuous, see [39,51] and [48]. Here, the supremum is taken over all continuous affine functions $h: K \to \mathbb{R}$ whose graphs lie below the graph of f.

2. STATES AND DENSITY MATRICES

The aim of this section is to translate openness questions from states to density matrices. In the sequel, we refer to matrix systems and *-subalgebras synonymously as *complex matrix systems* and *complex *-subalgebras*, respectively. We introduce their real counterparts because real *-subalgebras have a greater variety of state spaces than the complex ones.

A real matrix system on \mathbb{C}^n is a real linear subspace \mathcal{R} of M_n that is self-adjoint and contains $\mathbb{1}_n$. Let \mathcal{R} denote a real matrix system on \mathbb{C}^n . We endow \mathcal{R} with the Euclidean scalar product

$$\mathcal{R} \times \mathcal{R} \to \mathbb{R}, \quad (A, B) \mapsto \operatorname{Re}\langle A, B \rangle,$$

$$(2.1)$$

where $\langle A, B \rangle = \text{tr } A^*B$ and Re(a + ib) = a is the real part of a complex number, $A, B \in M_n$, $a, b \in \mathbb{R}$. The positive cone, space of hermitian matrices, dual cone, and space of density matrices are defined verbatim to their respective complex counterparts defined in Sec. 1, and are denoted by

$$\mathcal{C}(\mathcal{R}), \quad \mathcal{H}(\mathcal{R}), \quad \mathcal{C}(\mathcal{R})^{\vee}, \quad \text{and} \quad \mathcal{D}(\mathcal{R}).$$

We call \mathcal{R} a *real* *-*subalgebra* of M_n if \mathcal{R} is closed under matrix multiplication.

Generalizing a definition from real algebras [32, Sec. 4.5], we define the real state space of \mathcal{R} as

$$\mathcal{S}_{\mathbb{R}}(\mathcal{R}) := \left\{ \ell \in \mathcal{R}^*_{\mathbb{R},0} \mid \forall A \in \mathcal{C}(\mathcal{R}) : \ell(A) \ge 0, \ell(\mathbb{1}_n) = 1 \right\},\$$

where

$$\mathcal{R}^*_{\mathbb{R},0} := \{\ell : \mathcal{R} \to \mathbb{R} \mid \ell \text{ is } \mathbb{R}\text{-linear}, \forall A \in \mathcal{H}^-(\mathcal{R}) : \ell(A) = 0\}$$

is the space of real functionals vanishing on the skew-hermitian matrices

$$\mathcal{H}^{-}(\mathcal{R}) := \{A \in \mathcal{R} : A^* = -A\}.$$

Lemma 2.1. If \mathcal{R} is a real matrix system on \mathbb{C}^n , then the map

$$r_{\mathbb{R},\mathcal{R}}: \mathcal{D}(\mathcal{R}) \to \mathcal{S}_{\mathbb{R}}(\mathcal{R}), \quad \rho \mapsto \operatorname{Re}\langle \rho, \cdot \rangle$$

is a real affine isomorphism between compact convex sets.

Proof. As \mathcal{R} is the orthogonal direct sum $\mathcal{R} = \mathcal{H}(\mathcal{R}) \oplus \mathcal{H}^{-}(\mathcal{R})$, the real linear isomorphism [23, Sec. 67]

$$r_{\mathbb{R},\mathcal{R}}: \mathcal{R} \to \{\ell: \mathcal{R} \to \mathbb{R} \mid \ell \text{ is } \mathbb{R}\text{-linear}\}, \quad A \mapsto \operatorname{Re}\langle A, \cdot \rangle$$

restricts to the real linear isomorphism $\mathcal{H}(\mathcal{R}) \to \mathcal{R}^*_{\mathbb{R},0}$. Restricting $r_{\mathbb{R},\mathcal{R}}$ further, we obtain an injective map whose domain is $\mathcal{D}(\mathcal{R})$. So, it suffices to prove $r_{\mathbb{R},\mathcal{R}}(\rho) \in \mathcal{S}_{\mathbb{R}}(\mathcal{R})$ for $\rho \in \mathcal{D}(\mathcal{R})$, and that $r_{\mathbb{R},\mathcal{R}} : \mathcal{D}(\mathcal{R}) \to \mathcal{S}_{\mathbb{R}}(\mathcal{R})$

is surjective. Both assertions are straightforward to verify. The convex set $\mathcal{D}(\mathcal{R})$ is compact, see [44, Lemma 3.3], because $\mathbb{1}_n$ lies in the interior of $\mathcal{C}(\mathcal{R})$ in the topology of $\mathcal{H}(\mathcal{R})$. The convex set $\mathcal{S}_{\mathbb{R}}(\mathcal{R})$ is compact as it is the image of a compact set under a continuous map.

Returning to complex functionals, we consider the real vector space

$$\mathcal{R}^*_{her} := \{\ell : \mathcal{R} \to \mathbb{C} \mid \ell \text{ is } \mathbb{C}\text{-linear}, \forall A \in \mathcal{H}(\mathcal{R}) : \ell(A) \in \mathbb{R}\}$$

of complex functionals taking real values on hermitian matrices.

Lemma 2.2. If \mathcal{R} is a complex matrix system on \mathbb{C}^n , then the map

$$r_{\mathcal{R}}: \mathcal{D}(\mathcal{R}) \to \mathcal{S}(\mathcal{R}), \quad \rho \mapsto \langle \rho, \cdot \rangle$$

is a real affine isomorphism between compact convex sets.

Proof. The complex antilinear isomorphism $r_{\mathcal{R}} : \mathcal{R} \to \mathcal{R}^*, A \mapsto \langle A, \cdot \rangle$ restricts to a real linear isomorphism $\mathcal{H}(\mathcal{R}) \to \mathcal{R}^*_{her}$. Furthermore,

$$\alpha: \mathcal{R}^*_{\mathbb{R},0} \to \mathcal{R}^*_{\mathrm{her}}\,, \quad \alpha(\ell)[A + \mathrm{i}B] := \ell(A) + \mathrm{i}\ell(B)\,, \quad A, B \in \mathcal{H}(\mathcal{R})\,,$$

is a real linear isomorphism, whose inverse is given by $\ell \mapsto \operatorname{Re} \circ \ell$. So, the following diagram commutes. (Note that two arrows in opposite directions denote a bijection.)



Moreover, if $\ell_1 \in \mathcal{R}^*_{\mathbb{R},0}$ and $\ell_2 \in \mathcal{R}^*_{her}$ satisfy $\ell_2 = \alpha(\ell_1)$, then $\ell_1 \in \mathcal{S}_{\mathbb{R}}(\mathcal{R})$ holds if and only if $\ell_2 \in \mathcal{S}(\mathcal{R})$. Hence, the claim follows from Lemma 2.1. \Box

Example 2.3. Real *-subalgebras of M_n have a richer class of state spaces than the complex ones. The *Bloch ball* [7, Sec. 5.2]

$$\mathcal{D}(M_2) = \left\{ \frac{1}{2} (\mathbb{1}_2 + c_X X + c_Y Y + c_Z Z) \colon c_X, c_Y, c_Z \in \mathbb{R}, c_X^2 + c_Y^2 + c_Z^2 \le 1 \right\}$$

is a three-dimensional Euclidean ball of radius $1/\sqrt{2}$. The set of density matrices of the algebra $M_2(\mathbb{R})$ of real 2×2 matrices is the great disk

$$\mathcal{D}(\mathcal{M}_2(\mathbb{R})) = \left\{ \frac{1}{2} (\mathbb{1}_2 + c_X X + c_Z Z) : c_X, c_Z \in \mathbb{R}, c_X^2 + c_Z^2 \le 1 \right\}$$

of the Bloch ball $\mathcal{D}(M_2)$. There is no complex *-subalgebra of M_n in any dimension n whose state space is a disk.

Remark 2.4. Real and complex matrix systems have the same families of state spaces. If \mathcal{R} is a real or complex matrix system on \mathbb{C}^n , then the real matrix system of hermitian matrices $\mathcal{H}(\mathcal{R})$ has the same set of hermitian matrices and hence the same set of density matrices as \mathcal{R} . Conversely, any real matrix system \mathcal{R} included in $\mathcal{H}(M_n)$ has the same set of hermitian matrices and the same set of density matrices as the complex matrix system $\mathcal{R} \oplus i\mathcal{R}$. The set of density matrices of a real or complex matrix system \mathcal{R} is affinely isomorphic the state space of \mathcal{R} by Lemma 2.1 or 2.2, respectively.

Let $\mathcal{R}_1, \mathcal{R}_2$ be real matrix systems on \mathbb{C}^n and let $\mathcal{R}_2 \subset \mathcal{R}_1$. We abbreviate $\mathcal{H}_i := \mathcal{H}(\mathcal{R}_i), \ \mathcal{C}_i := \mathcal{C}(\mathcal{R}_i), \ \mathcal{C}_i^{\vee} := \mathcal{C}^{\vee}(\mathcal{R}_i), \ \mathcal{D}_i := \mathcal{D}(\mathcal{R}_i), \ \mathcal{S}_{\mathbb{R},i} := \mathcal{S}_{\mathbb{R}}(\mathcal{R}_i), \ \mathcal{S}_i := \mathcal{S}(\mathcal{R}_i), \ r_{\mathbb{R},i} := r_{\mathbb{R},\mathcal{R}_i}, \ \text{and} \ r_i := r_{\mathcal{R}_i}, \ i = 1, 2.$ Usually, the orthogonal projection of \mathcal{R}_1 onto \mathcal{R}_2 is the idempotent self-adjoint linear map $\mathcal{R}_1 \to \mathcal{R}_1$ whose range is \mathcal{R}_2 . Reducing the codomain to the range, we get a map

$$\pi: \mathcal{R}_1 \to \mathcal{R}_2 \,, \tag{2.2}$$

whose value at $A \in \mathcal{R}_1$ is specified by the equations $\operatorname{Re}\langle A - \pi(A), B \rangle = 0$ for all $B \in \mathcal{R}_2$. We refer to π as the *orthogonal projection* of \mathcal{R}_1 onto \mathcal{R}_2 in this paper. The notation $\mathcal{R}_1 \to \mathcal{R}_2$ conveying domain and range is useful especially in Sec. 6. The adjoint of π is the embedding $\mathcal{R}_2 \to \mathcal{R}_1, A \mapsto A$. If \mathcal{R}_1 and \mathcal{R}_2 are complex matrix systems, then the Frobenius inner product induces the same orthogonal projection as the Euclidean scalar product (2.1).

As \mathcal{R}_i is the orthogonal direct sum of the spaces of its hermitian and skew-hermitian matrices, i = 1, 2, the map (2.2) restricts to $\pi : \mathcal{H}_1 \to \mathcal{H}_2$, which we denote (aware of the notational imprecision) by the same symbol π . The value of π at $A \in \mathcal{H}_1$ is specified by $\langle A - \pi(A), B \rangle = 0$ for all $B \in \mathcal{H}_2$.

A cone in a Euclidean space $(E, \langle\!\langle \cdot, \cdot \rangle\!\rangle)$ is a subset C of E that is closed under multiplication with positive scalars. A base of a cone C is a subset B of C such that $0 \notin \text{aff } B$ and such that for all nonzero $x \in C$ there exist $y \in B$ and s > 0 such that x = sy holds. Note that we have $B = C \cap \text{aff } B$ for every base B of a cone C. The set $M^{\vee} := \{x \in E \mid \forall y \in M : \langle\!\langle x, y \rangle\!\rangle \ge 0\}$ is a closed convex cone for every subset $M \subset E$, called the *dual cone* to M. Regarding duality of convex cones, we refer to [45, Sec. 14].

Lemma 2.5. Let $\mathcal{R}_1, \mathcal{R}_2$ be real matrix systems on \mathbb{C}^n such that $\mathcal{R}_2 \subset \mathcal{R}_1$ and let $\pi : \mathcal{H}_1 \to \mathcal{H}_2$ denote the orthogonal projection. Then $\mathcal{C}_2^{\vee} = \pi(\mathcal{C}_1^{\vee})$ and $\mathcal{D}_2 = \pi(\mathcal{D}_1)$ holds.

Proof. This lemma and its proof are similar to [44, Prop. 5.2]. The inclusions $\pi(\mathcal{C}_1^{\vee}) \subset \mathcal{C}_2^{\vee}$ and $\mathcal{C}_2 \supset (\pi(\mathcal{C}_1^{\vee}))^{\vee}$ are straightforward to verify and imply

$$\pi(\mathcal{C}_1^{\vee}) \subset \mathcal{C}_2^{\vee} \subset [\pi(\mathcal{C}_1^{\vee})]^{\vee \vee}$$

Since $[\pi(\mathcal{C}_1^{\vee})]^{\vee\vee}$ is the closure of the convex cone $\pi(\mathcal{C}_1^{\vee})$, it suffices to show that $\pi(\mathcal{C}_1^{\vee})$ is closed. By [44, Lemma 3.1], this would follow if $\pi(\mathcal{D}_1)$ was a compact base of $\pi(\mathcal{C}_1^{\vee})$. As $\mathbb{1}_n$ lies in the interior of \mathcal{C}_1 in the topology of \mathcal{H}_1 , we know that \mathcal{D}_1 is a compact base of \mathcal{C}_1^{\vee} , see [44, Lemma 3.3]. Hence $\pi(\mathcal{D}_1)$ is a compact base of $\pi(\mathcal{C}_1^{\vee})$ provided we establish that $0 \notin \operatorname{aff} \pi(\mathcal{D}_1)$. But this is clear from π being trace-preserving:

$$\operatorname{tr}(\pi(A)) = \langle \mathbb{1}_n, \pi(A) \rangle = \langle \mathbb{1}_n, A \rangle = \operatorname{tr}(A), \quad A \in \mathcal{H}_1.$$

This completes the proof of $C_2^{\vee} = \pi(C_1^{\vee})$. The identity $C_2^{\vee} = \pi(C_1^{\vee})$ and the fact that π is trace-preserving imply $\mathcal{D}_2 = \pi(\mathcal{D}_1)$.

Lemma 2.5 shows that the orthogonal projection $\pi : \mathcal{R}_1 \to \mathcal{R}_2$ restricts to the map

$$\pi: \mathcal{D}_1 \to \mathcal{D}_2 \,, \tag{2.3}$$

which we call the *orthogonal projection* of \mathcal{D}_1 onto \mathcal{D}_2 . To avoid any possible confusion, the orthogonal projections $\mathcal{R}_1 \to \mathcal{R}_2$ and $\mathcal{H}_1 \to \mathcal{H}_2$ are written without function labels from here on (Lemma 3.7 is an exception).

Proposition 2.6. Let $\mathcal{R}_1, \mathcal{R}_2$ be real matrix systems on \mathbb{C}^n such that $\mathcal{R}_2 \subset \mathcal{R}_1$ and let $\pi : \mathcal{D}(\mathcal{R}_1) \to \mathcal{D}(\mathcal{R}_2)$ denote the orthogonal projection.

- a) The diagram a) below commutes. Both $\pi : \mathcal{D}_1 \to \mathcal{D}_2$ and $\mathcal{S}_{\mathbb{R},1} \to \mathcal{S}_{\mathbb{R},2}$, $\ell \mapsto \ell|_{\mathcal{R}_2}$ are surjective maps.
- b) If $\mathcal{R}_1, \mathcal{R}_2$ are complex matrix systems on \mathbb{C}^n , then the diagram b) below commutes and the maps $\pi : \mathcal{D}_1 \to \mathcal{D}_2$ and $\mathcal{S}_1 \to \mathcal{S}_2$, $\ell \mapsto \ell|_{\mathcal{R}_2}$ are onto.

Proof. a) The horizontal arrows of diagram a) and b) are obtained in Lemma 2.1 and Lemma 2.2, respectively, and $\pi : \mathcal{D}_1 \to \mathcal{D}_2$ is onto by Lemma 2.5. The diagram a) commutes because for all $\rho \in \mathcal{D}_1$ and $A \in \mathcal{R}_2$ we have

$$[r_{\mathbb{R},2} \circ \pi(\rho)](A) = \operatorname{Re}\langle \pi(\rho), A \rangle = \operatorname{Re}\langle \rho, A \rangle = [r_{\mathbb{R},1}(\rho)](A) = r_{\mathbb{R},1}(\rho)|_{\mathcal{R}_2}(A).$$

Therefore and since $\pi : \mathcal{D}_1 \to \mathcal{D}_2$ is surjective, the map $\mathcal{S}_{\mathbb{R},1} \to \mathcal{S}_{\mathbb{R},2}$ is surjective as well. The proof of b) is similar.

Corollary 2.7. Let $\mathcal{R}_1, \mathcal{R}_2$ be real matrix systems on \mathbb{C}^n such that $\mathcal{R}_2 \subset \mathcal{R}_1$, let $\pi : \mathcal{D}(\mathcal{R}_1) \to \mathcal{D}(\mathcal{R}_2)$ denote the orthogonal projection, and let $\rho \in \mathcal{D}_1$.

- a) The map $\pi : \mathcal{D}_1 \to \mathcal{D}_2$ is open at ρ if and only if $\mathcal{S}_{\mathbb{R},1} \to \mathcal{S}_{\mathbb{R},2}$, $\ell \mapsto \ell|_{\mathcal{R}_2}$ is open at $r_{\mathbb{R},1}(\rho)$.
- b) If $\mathcal{R}_1, \mathcal{R}_2$ are complex matrix systems on \mathbb{C}^n , then $\pi : \mathcal{D}_1 \to \mathcal{D}_2$ is open at ρ if and only if $\mathcal{S}_1 \to \mathcal{S}_2$, $\ell \mapsto \ell|_{\mathcal{R}_2}$ is open at $r_1(\rho)$.

Proof. This follows directly from Prop. 2.6.

Remark 2.8. Let $\mathcal{R}_1, \mathcal{R}_2$ be real matrix systems on \mathbb{C}^n and let $\mathcal{R}_2 \subset \mathcal{R}_1$. Then $\mathcal{D}_2 \supset \mathcal{D}_1 \cap \mathcal{R}_2$ holds, but the converse inclusion is wrong in general, as Ex. 2.9 shows. However, if \mathcal{R}_2 is a real *-subalgebra of M_n , then we have

$$\mathcal{D}_2 = \mathcal{D}_1 \cap \mathcal{R}_2 \,. \tag{2.4}$$

Indeed, $C_2^{\vee} = C_2$ holds [28, Thm. III.2.1] as the space of hermitian matrices \mathcal{H}_2 is a Euclidean Jordan algebra with Jordan product $A \circ B = \frac{1}{2}(AB + BA)$ and inner product $(A, B) \mapsto \operatorname{Re}\langle A, B \rangle$, $A, B \in \mathcal{H}_2$. So, $C_2^{\vee} = C_2 \subset C_1 \subset C_1^{\vee}$ proves $C_2^{\vee} = C_2$, which implies (2.4). See also the Notes to Chapter 6 in [2].

Example 2.9. Despite $\mathcal{R}_2 \subset \mathcal{R}_1 \subset M_3$, the inclusions $\mathcal{D}_2 \subset \mathcal{D}_1 \subset \mathcal{D}(M_3)$ fail if $\mathcal{R}_2 := \operatorname{span}_{\mathbb{C}}(\mathbb{1}_3, Z \oplus 0)$ and $\mathcal{R}_1 := \operatorname{span}_{\mathbb{C}}(\mathbb{1}_3, X \oplus 1, Z \oplus 0)$. We have

$$A_{\lambda} \in \mathcal{D}_2 \Leftrightarrow |\lambda| \leq \frac{3}{2}, \quad A_{\lambda} \in \mathcal{D}_1 \Leftrightarrow |\lambda| \leq \sqrt{2}, \quad A_{\lambda} \in \mathcal{D}(\mathcal{M}_3) \Leftrightarrow |\lambda| \leq 1$$

for $A_{\lambda} := (\mathbb{1}_3 + \lambda Z \oplus 0)/3 \in \mathcal{R}_2, \ \lambda \in \mathbb{R}$. The second equivalence is obtained by minimizing $\langle A_{\lambda}, A \rangle$ for fixed λ over $A \in \mathcal{C}_1$, that is, by minimizing $\langle A_{\lambda}, \mathbb{1}_3 + c_1(X \oplus 1) + c_2 Z \oplus 0 \rangle = 1 + \frac{1}{3}c_1 + \frac{2}{3}\lambda c_2$ on the unit disk of \mathbb{R}^2 .

3. Direct convex sums

This section addresses affinely independent convex sets, their convex hulls, and maps defined thereon. Let

$$\Delta_m := \{ (s_1, \dots, s_m) \in \mathbb{R}^m \mid \forall i : s_i \ge 0, s_1 + \dots + s_m = 1 \}$$

denote the probability simplex, and

$$\Delta_m(\epsilon, s_1, \dots, s_m) := \Delta_m \cap \bigoplus_{i=1}^m (s_i - \epsilon, s_i + \epsilon)$$

the open hypercube of edge length 2ϵ centered at $(s_1, \ldots, s_m) \in \Delta_m$.

A family of convex subsets K_1, \ldots, K_m of a Euclidean space is *affinely* independent if every point in their convex hull can be expressed by a unique convex combination $s_1x_1 + \cdots + s_mx_m$. This means that $(s_1, \ldots, s_m) \in \Delta_m$ is unique and $x_i \in K_i$ is unique for all *i* for which if $s_i > 0, i = 1, \ldots, m$. The *direct convex sum* [1] of a family K_1, \ldots, K_m of affinely independent convex sets is defined as their convex hull

$$K_1 \oplus_{\mathbf{c}} \cdots \oplus_{\mathbf{c}} K_m := \operatorname{conv}(K_1 \cup \cdots \cup K_m).$$

If K_1, \ldots, K_m is a family of affinely independent compact convex sets, then their direct convex sum is compact [45, Thm. 17.2]. A compactness argument allows us to describe a base of open neighborhoods.

Lemma 3.1. Let K_1, \ldots, K_m be affinely independent compact convex subsets of a Euclidean space, and let $x_i \in K_i$, $i = 1, \ldots, m$, and $(s_1, \ldots, s_m) \in \Delta_m$. Let $I := \{i \in \{1, \ldots, m\} \mid s_i > 0\}, \delta := \min_{i \in I} s_i$, and define

$$O_I(\epsilon, (A_i)_{i \in I}) := \{ t_1 y_1 + \dots + t_m y_m \mid (t_1, \dots, t_m) \in \Delta_m(\epsilon, s_1, \dots, s_m), \\ \forall i : y_i \in K_i \text{ and } (i \in I \Rightarrow y_i \in A_i) \}$$

for every $\epsilon \in (0, \delta]$ and open set A_i in the relative topology of K_i containing x_i , i = 1, ..., m. Then the family $\{O_I(\epsilon, (A_i)_{i \in I})\}$ is a local base of open neighborhoods at $s_1x_1 + \cdots + s_mx_m$ in the relative topology of $K_1 \oplus_c \cdots \oplus_c K_m$.

Proof. As $K := K_1 \oplus_{\mathbf{c}} \cdots \oplus_{\mathbf{c}} K_m$ is a metric space, it suffices to show that there are arbitrary small family members and $O := O_I(\epsilon, (A_i)_{i \in I})$ is open.

Let $B_i := A_i$ if $i \in I$ and $B_i := K_i$ if $i \notin I$, i = 1, ..., m. Then the set

$$U := \Delta_m(\epsilon, s_1, \dots, s_m) \oplus \bigoplus_i B_i$$

is open in the relative topology of the compact set $\tilde{K} := \Delta_m \oplus \bigoplus_i K_i$. The complement

$$U^{\complement} := \widetilde{K} \setminus U = \underbrace{\Delta_m(\epsilon, s_1, \dots, s_m)^{\complement} \oplus \bigoplus_i K_i}_{C :=} \cup \bigcup_{j \in I} \underbrace{\Delta_m \oplus (\bigoplus_{i \neq j} K_i) \oplus A_j^{\complement}}_{C_j :=}$$

is compact. Hence, its image under the continuous surjective map

 $\beta: \widetilde{K} \to K$, $((t_i)_{i=1}^m, (y_i)_{i=1}^m) \mapsto \sum_{i=1}^m t_i y_i$

is compact. We prove that $O = \beta(U)$ is open in K by showing that $\beta(U)$ is disjoint from $\beta(U^{\complement})$. Let

$$u := (t_1, \ldots, t_m, y_1, \ldots, y_m) \in U.$$

Then $\beta(u) \notin \beta(C)$ follows as the vector $(t_i)_{i=1}^m$ is uniquely determined by $\beta(u)$. For every $j \in I$ we have $s_j > 0$, hence $t_j > 0$ by the definition of O. Thus, y_j is uniquely determined by $\beta(u)$, which proves $\beta(u) \notin \beta(C_j)$.

Let $|M| := \sup_{x,y \in M} |y - x|$ denote the diameter of a set M. The distance of $x := s_1 x_1 + \dots + s_m x_m$ from a point $t_1 y_1 + \dots + t_m y_m$ in O is bounded by

$$\sum_{i} \left(|s_i - t_i| |x_i| + t_i |x_i - y_i| \right) \le \epsilon \sum_{i} |x_i| + \sum_{i \in I} t_i |A_i| + \epsilon \sum_{i \notin I} |K_i|.$$

Choosing open sets $(A_i)_{i \in I}$ with diameters at most ϵ , we obtain

$$|O| \le 2\sup_{y \in O} |x - y| \le 2\epsilon \left(\sum_i |x_i| + 1 + \sum_{i \notin I} |K_i| \right),$$

which completes the proof, as the compact sets K_i have finite diameters and as ϵ can be chosen arbitrarily small.

Let E_1, \ldots, E_m be Euclidean spaces. We consider E_i as a subspace of the direct sum $\bigoplus_{i=1}^m E_i$ via the embedding

$$E_i \to \bigoplus_{j=1}^m E_j, \quad x \mapsto (\underbrace{0, \dots, 0}_{i-1 \text{ zeros}}, x, \underbrace{0, \dots, 0}_{m-i \text{ zeros}}), \quad i = 1, \dots, m$$

Proposition 3.2. Let $K_i \subset E_i$ and $L_i \subset F_i$ be compact convex subsets of Euclidean spaces E_i and F_i , i = 1, ..., m, such that $K_1, ..., K_m$ are affinely independent in $\bigoplus_{i=1}^m E_i$ and $L_1, ..., L_m$ are affinely independent in $\bigoplus_{i=1}^m F_i$. If $f_i : K_i \to L_i$ is a map, i = 1, ..., m, then a map

$$f_1 \oplus_{\mathbf{c}} \cdots \oplus_{\mathbf{c}} f_m : K_1 \oplus_{\mathbf{c}} \cdots \oplus_{\mathbf{c}} K_m \to L_1 \oplus_{\mathbf{c}} \cdots \oplus_{\mathbf{c}} L_m$$
(3.1)

is well defined by

$$s_1x_1 + \dots + s_mx_m \mapsto s_1f_1(x_1) + \dots + s_mf_m(x_m)$$

If f_i is open and surjective for i = 1, ..., m, then $f_1 \oplus_c \cdots \oplus_c f_m$ is open.

Proof. The map is well defined as the sets K_1, \ldots, K_m are affinely independent and the sets L_1, \ldots, L_m are affinely independent. Regarding the openness of $f_1 \oplus_{\mathbf{c}} \cdots \oplus_{\mathbf{c}} f_m$, it suffices to show that the images of the members of a base of the relative topology of $K_1 \oplus_{\mathbf{c}} \cdots \oplus_{\mathbf{c}} K_m$ are open. Using the base of Lemma 3.1, we have

$$f_1 \oplus_{\mathbf{c}} \cdots \oplus_{\mathbf{c}} f_m[O_I(\epsilon, (A_i)_{i \in I})] = O_I[\epsilon, (f(A_i))_{i \in I}],$$

which is open in $L_1 \oplus_c \cdots \oplus_c L_m$. The required identity of $f_i(K_i) = L_i$ for every $i \notin I$ is a consequence of f_i being surjective.

We call the map (3.1) the *direct convex sum* of the maps f_1, \ldots, f_m . Next, we recall a sufficient condition for the affine independence of convex sets [17].

Remark 3.3. Let K_i be a convex subset of a Euclidean space E_i such that $0 \notin \text{aff } K_i$ holds for $i = 1, \ldots, m$. Then K_1, \ldots, K_m are affinely independent in the direct sum $\bigoplus_{i=1}^m E_i$.

Returning to matrix systems, we consider a real matrix system \mathcal{R}_i on \mathbb{C}^{n_i} , $i = 1, \ldots, m$. The direct sum $\mathcal{R} := \bigoplus_{i=1}^m \mathcal{R}_i$ is a real matrix system on $\mathbb{C}^{n_1 + \cdots + n_m}$. The Frobenius inner product of $(A_i)_{i=1}^m, (B_i)_{i=1}^m \in \mathcal{R}$ is $\langle (A_i)_{i=1}^m, (B_i)_{i=1}^m \rangle = \sum_{i=1}^m \langle A_i, B_i \rangle$.

Lemma 3.4. Let \mathcal{R}_i be a real matrix system on \mathbb{C}^{n_i} , $i = 1, \ldots, m$. Then

$$\mathcal{C}(\mathcal{R}_1 \oplus \cdots \oplus \mathcal{R}_m) = \mathcal{C}(\mathcal{R}_1) \oplus \cdots \oplus \mathcal{C}(\mathcal{R}_m),$$

$$\mathcal{C}^{\vee}(\mathcal{R}_1 \oplus \cdots \oplus \mathcal{R}_m) = \mathcal{C}^{\vee}(\mathcal{R}_1) \oplus \cdots \oplus \mathcal{C}^{\vee}(\mathcal{R}_m),$$

and $\mathcal{D}(\mathcal{R}_1 \oplus \cdots \oplus \mathcal{R}_m) = \mathcal{D}(\mathcal{R}_1) \oplus_{\mathrm{c}} \cdots \oplus_{\mathrm{c}} \mathcal{D}(\mathcal{R}_m).$

8

Proof. The first identity is clear. The second one follows by induction from m = 2, a case that is easy to verify. By Rem. 3.3 and because $\mathcal{D}(\mathcal{R}_i)$ is included in the hyperplane of trace-one matrices $i = 1, \ldots, m$, the sets $\mathcal{D}(\mathcal{R}_1), \ldots, \mathcal{D}(\mathcal{R}_m)$ are affinely independent in the direct sum $\bigoplus_{i=1}^m \mathcal{H}(\mathcal{R}_i)$. The third identity follows from the second one by enforcing the trace to be one.

Example 3.5. Let $\mathcal{R} := \operatorname{span}_{\mathbb{R}}(\mathbb{1}_3, X \oplus 1, Z \oplus 0).$

a) The set of density matrices $\mathcal{D}(M_2 \oplus M_1)$ of the *-subalgebra $M_2 \oplus M_1$ of M_3 is a symmetric cone that is the direct convex sum of the Bloch ball $\mathcal{D}(M_2)$ and the singleton $\mathcal{D}(M_1) = \{1\}$, see Lemma 3.4 and Ex. 2.3. The closed segment

$$\mathcal{G} := \left[\left| + \right\rangle\!\!\left\langle + \right| \oplus 0, 0 \oplus 1 \right] = \left\{ \left(1 - \lambda \right) \left| + \right\rangle\!\!\left\langle + \right| \oplus \lambda \colon \lambda \in [0, 1] \right\}$$

is a generatrix of this cone. One proves along the lines of Ex. 1.1 that the orthogonal projection

$$\pi: \mathcal{D}(M_2 \oplus M_1) \to \mathcal{D}(\mathcal{R})$$

is not open at any point in the half-open segment

 $\mathcal{G}_0 := \left(\left| + \right\rangle \!\! \left\langle + \right| \oplus 0, 0 \oplus 1 \right] = \mathcal{G} \setminus \left\{ \left| + \right\rangle \!\! \left\langle + \right| \oplus 0 \right\}.$

Thereby, the equivalence of states and density matrices is described in Coro. 2.7. It is important to observe that $\frac{1}{2}(\mathbb{1}_2 + \cos(\theta)X + \sin(\theta)Z) \oplus 0$ and $\cos(\theta)(X \oplus 1) + \sin(\theta)Z \oplus 0$ are matrices in $M_2 \oplus M_1$ for all $\theta \in \mathbb{R}$. The map π is open at every point in the complement of \mathcal{G}_0 because $\mathcal{D}(M_2 \oplus M_1)$ is a cone over a ball, see [52, Lemma 4.17] for a detailed proof.

b) The orthogonal projection $\pi : \mathcal{D}(M_2 \oplus M_1) \to \mathcal{D}(\mathcal{R})$ has an instructive geometry. The set of density matrices $\mathcal{D}(\mathcal{R}) = \pi(\mathcal{D}(M_2) \oplus_c \{1\})$ is the convex hull of the projected ball $\pi(\mathcal{D}(M_2) \oplus \{0\})$ and the singleton $\pi(0 \oplus 1)$ by Lemma 2.5. In turn, $\pi(\mathcal{D}(M_2) \oplus \{0\})$ is the filled ellipse of all points

$$\pi \left(\frac{1}{2} (\mathbb{1}_2 + c_X X + c_Y Y + c_Z Z) \oplus 0 \right)$$

$$= \frac{1}{2} (\mathbb{1}_3 - M + c_X (3M - \mathbb{1}_3) + c_Z Z \oplus 0),$$
(3.2)

where $c_X, c_Y, c_Z \in \mathbb{R}$ satisfy $c_X^2 + c_Y^2 + c_Z^2 = 1$, and $M := \frac{1}{2}(|+\rangle \langle +| \oplus 1)$ is the midpoint of the generatrix \mathcal{G} . Note that $\mathbb{1}_3, (3X - \mathbb{1}_2) \oplus 2, Z \oplus 0$ is an orthogonal basis of \mathcal{R} . The choice of $c_X = 1, c_Y = c_Z = 0$ yields

$$\pi(|+\rangle\langle +|\oplus 0) = M = \pi(0\oplus 1).$$
(3.3)

Thus, \mathcal{G} is perpendicular to \mathcal{R} , and $\mathcal{D}(\mathcal{R}) = \pi(\mathcal{D}(M_2) \oplus \{0\})$ is an ellipse. Moreover, $\pi^{-1}(M) = \mathcal{G}$ holds by equation (3.2) and (3.3).

c) The ellipse $\mathcal{D}(\mathcal{R})$ has the semiaxes $\sqrt{3/8}$ and $1/\sqrt{2}$. This follows from the formula (3.2) when (c_X, c_Y, c_Z) is assigned the values of $(\pm 1, 0, 0)$ and $(0, 0, \pm 1)$. In retrospect to Rem. 2.9, the value $(1/3, 0, \sqrt{8/9})$ confirms that $(\mathbb{1}_3 + \lambda Z \oplus 0)/3$ is contained in $\mathcal{D}(\mathcal{R})$ if and only if $|\lambda| \leq \sqrt{2}$.

Example 3.6. The set of density matrices $\mathcal{D}(M_2(\mathbb{R}) \oplus M_1(\mathbb{R}))$ of the real *-subalgebra $M_2(\mathbb{R}) \oplus M_1(\mathbb{R})$ of M_3 is a symmetric cone, which is the direct convex sum of the great disk $\mathcal{D}(M_2(\mathbb{R}))$ of the Bloch ball and a singleton.

As in Ex. 3.5, the closed segment \mathcal{G} is a generatrix of this cone, which is the fiber of the orthogonal projection

$$\pi: \mathcal{D}(M_2(\mathbb{R}) \oplus M_1(\mathbb{R})) \to \mathcal{D}(\mathcal{R})$$

over M. The map π is not open at any point in the half-open segment \mathcal{G}_0 and open at every point in the complement. The lack of openness can be visualized graphically in three-space by observing that π projects the cone $\mathcal{D}(M_2(\mathbb{R}) \oplus M_1(\mathbb{R}))$ along its generatrix \mathcal{G} to the ellipse $\mathcal{D}(\mathcal{R})$.

Lemma 3.7. Let $\mathcal{R}_i, \mathcal{R}'_i$ be real matrix systems on \mathbb{C}^{n_i} such that $\mathcal{R}_i \subset \mathcal{R}'_i$, let $\pi_i : \mathcal{R}'_i \to \mathcal{R}_i$ denote the orthogonal projection, $i = 1, \ldots, m$, and consider the direct sums $\mathcal{R}' := \bigoplus_{i=1}^m \mathcal{R}'_i$ and $\mathcal{R} := \bigoplus_{i=1}^m \mathcal{R}_i$. The orthogonal projection $\pi : \mathcal{D}(\mathcal{R}') \to \mathcal{D}(\mathcal{R})$ equals

$$\pi_1 \oplus_{\mathbf{c}} \cdots \oplus_{\mathbf{c}} \pi_m : \mathcal{D}(\mathcal{R}'_1) \oplus_{\mathbf{c}} \cdots \oplus_{\mathbf{c}} \mathcal{D}(\mathcal{R}'_m) \to \mathcal{D}(\mathcal{R}_1) \oplus_{\mathbf{c}} \cdots \oplus_{\mathbf{c}} \mathcal{D}(\mathcal{R}_m).$$

Proof. The orthogonal projections $\pi_i : \mathcal{D}(\mathcal{R}'_i) \to \mathcal{D}(\mathcal{R}_i), i = 1, ..., m$, and $\pi : \mathcal{D}(\mathcal{R}') \to \mathcal{D}(\mathcal{R})$ are well defined by Lemma 2.5. A straight-forward computation shows that the orthogonal projection $\pi : \mathcal{R}' \to \mathcal{R}$ is the direct sum $\bigoplus_{i=1}^{m} \pi_i$. The claim then follows from the third identity of Lemma 3.4 and from the definition of the direct convex sum of maps in formula (3.1). \Box

4. Numerical ranges

The orthogonal projection $\mathcal{D}(M_n) \to \mathcal{D}(\mathcal{R})$ may be restricted to the set $\exp \mathcal{D}(M_n)$ of extreme points¹ of $\mathcal{D}(M_n)$. The openness of this restriction was studied in matrix theory [18,29–31,34,35,49,50,53] for two-dimensional state spaces $\mathcal{D}(\mathcal{R})$ represented as numerical ranges (see Rem. 4.3).

Let $A_1, \ldots, A_k \in \mathcal{H}(\mathcal{M}_n)$ and consider the real matrix system

$$\mathcal{R}(A_1,\ldots,A_k) := \operatorname{span}_{\mathbb{R}}(\mathbb{1}_n,A_1,\ldots,A_k).$$

The image of $\mathcal{D}(M_n)$ under the real linear map

 $v: \mathcal{H}(\mathcal{M}_n) \to \mathbb{R}^k, \quad B \mapsto (\langle B, A_1 \rangle, \dots, \langle B, A_k \rangle)^{\mathrm{T}}$

is the joint numerical range $V(A_1, \ldots, A_k) := v(\mathcal{D}(\mathbf{M}_n)) \subset \mathbb{R}^k$, see [9]. Aware of the notational imprecision, we use the same label v also for several restrictions of v, among others for

$$v: \mathcal{D}(\mathbf{M}_n) \to V(A_1, \dots, A_k), \qquad \rho \mapsto (\langle \rho, A_1 \rangle, \dots, \langle \rho, A_k \rangle)^{\mathrm{T}}.$$
 (4.1)

Lemma 4.1. The following diagram commutes.



¹An point in a convex set K is an *extreme point* [45] of K if there is no way to express it as a convex combination (1 - s)x + sy such that $x, y \in K$ and $s \in (0, 1)$, except by taking x = y. We denote the set of extreme points of K by ex K.

Proof. Let $\mathcal{R} := \mathcal{R}(A_1, \ldots, A_k)$. By Lemma 2.5, the orthogonal projection $\pi : \mathcal{D}(\mathcal{M}_n) \to \mathcal{D}(\mathcal{R})$ is surjective. It is straightforward to verify that $v : \mathcal{H}(\mathcal{M}_n) \to \mathbb{R}^k$ factors through \mathcal{R} , in the sense that $v = v \circ \pi$ holds. Hence, the commutativity of the diagram is implied by the injectivity of v restricted to the affine space $\{B \in \mathcal{H}(\mathcal{R}) : \operatorname{tr}(B) = 1\}$. Let B_1, B_2 be contained in this affine space. If $v(B_1) = v(B_2)$, then

$$0 = v(B_1 - B_2) = \langle B_1 - B_2, A_i \rangle_{i=1}^k$$

and
$$0 = \operatorname{tr}(B_1) - \operatorname{tr}(B_2) = \langle B_1 - B_2, \mathbb{1}_n \rangle.$$

This implies $B_1 - B_2 = 0$ as $B_1 - B_2 \in \mathcal{H}(\mathcal{R}) = \operatorname{span}_{\mathbb{R}}(\mathbb{1}_n, A_1, \dots, A_k)$. \Box

In the remainder of this section, let k = 2 and $A := A_1 + iA_2$. The image of the unit sphere $\mathbb{C}S^n := \{ |\varphi\rangle \in \mathbb{C}^n : \langle \varphi | \varphi \rangle = 1 \}$ under the hermitian quadratic form $f_A : \mathbb{C}^n \to \mathbb{C}, |\varphi\rangle \mapsto \langle \varphi | A\varphi \rangle$ is the numerical range

$$W(A) := f_A(\mathbb{C}S^n) \subset \mathbb{C}$$

Here, $\langle \varphi_1 | \varphi_2 \rangle := \overline{x_1} y_1 + \dots + \overline{x_n} y_n$ is the inner product of $|\varphi_1\rangle = (x_1, \dots, x_n)^T$ and $|\varphi_2\rangle = (y_1, \dots, y_n)^T$ in \mathbb{C}^n . We use the same label f_A to denote the map

$$f_A : \mathbb{C}S^n \to W(A), \quad |\varphi\rangle \mapsto \langle \varphi | A\varphi \rangle.$$
 (4.2)

Minkowski's theorem [47] asserts that every compact convex set is the convex hull of its extreme points.

Proposition 4.2. The following diagram commutes.

Proof. The bottom right triangle is the case k = 2 of Lemma 4.1. The map $f_A : \mathbb{C}S^n \to W(A)$ factors through $\exp \mathcal{D}(M_n)$, $\mathcal{D}(M_n)$, and $V(A_1, A_2)$, as $\mathbb{C}S^n \to \mathcal{D}(M_n)$, $|\varphi\rangle \mapsto |\varphi\rangle\langle\varphi|$ parametrizes the extreme points of $\mathcal{D}(M_n)$, see for example [7, Sec. 5.1], and since for all $|\varphi\rangle \in \mathbb{C}S^n$ we have

$$f_A(|\varphi\rangle) = \langle \varphi | A\varphi \rangle = \operatorname{tr}(|\varphi\rangle\langle \varphi | A) = \langle |\varphi\rangle\langle \varphi | , A \rangle$$
$$= \langle |\varphi\rangle\langle \varphi | , A_1 \rangle + i \langle |\varphi\rangle\langle \varphi | , A_2 \rangle .$$

It remains to show that $g: W(A) \to V(A_1, A_2), z \mapsto (\operatorname{Re}(z), \operatorname{Im}(z))^{\mathrm{T}}$ is onto (being the restriction of a bijection, the map g is one-to-one).

First, we show that $\operatorname{ex} V(A_1, A_2)$ is included in the image of g. The preimage of every extreme point x of $V(A_1, A_2)$ under $v : \mathcal{D}(M_n) \to V(A_1, A_2)$ contains an extreme point of $\mathcal{D}(M_n)$. This is true because the preimage of xis a face \mathcal{F} of $\mathcal{D}(M_n)$, which has an extreme point ρ by Minkowski's theorem, since $\mathcal{D}(M_n)$, and hence \mathcal{F} , is compact. Since ρ is also an extreme point of $\mathcal{D}(M_n)$, the claim follows from $\mathbb{C}S^n \to \operatorname{ex} \mathcal{D}(M_n)$ being onto.

Second, the convex hull of $\exp(A_1, A_2)$ is included in the image of g because W(A) is convex by the Toeplitz-Hausdorff theorem [12, 37]. Third, the map g is onto, because $V(A_1, A_2)$ is the convex hull of its extreme points, again by Minkowski's theorem.

The affine isomorphism $W(A) \cong \mathcal{D}(\mathcal{R}(A_1, A_2))$ of Prop. 4.2 has an analogue in the much more general setting of matrix-valued states [22, Thm. 5.1].

Let f_A^{-1} denote the multi-valued inverse of $f_A : \mathbb{C}S^n \to W(A)$. Corey et al. [18] define f_A^{-1} to be strongly continuous at $z \in W(A)$ if the map f_A is open at every point in the fiber $f_A^{-1}(z)$ of f_A over z.

Remark 4.3. Strong continuity can be described in terms of standard results on the numerical range. We refer to Sec. 8 of [29] and the references therein.

There exists a family of orthonormal bases $|\varphi_1(\theta)\rangle, \ldots, |\varphi_n(\theta)\rangle$ of \mathbb{C}^n that is analytically parametrized by a real number $\theta \in \mathbb{R}$; and there are analytic functions $\lambda_i : \mathbb{R} \to \mathbb{R}$, called *eigenfunctions* [30], such that

$$\left(\cos(\theta)A_1 + \sin(\theta)A_2\right) |\varphi_i(\theta)\rangle = \lambda_i(\theta) |\varphi_i(\theta)\rangle, \quad \theta \in \mathbb{R}, \quad i = 1, \dots, n.$$

For every $i = 1, \ldots, n$, the numerical range W(A) includes the image $\text{Img}(z_i)$ of the curve

$$z_i : \mathbb{R} \to \mathbb{C}, \quad \theta \mapsto e^{\mathrm{i}\theta} (\lambda_i(\theta) + \mathrm{i}\lambda'_i(\theta)).$$

Every extreme point of W(A) is contained in $\text{Img}(z_i)$ for some i = 1, ..., n.

If $z \in W(A)$ is not an extreme point of W(A), then f_A^{-1} is strongly continuous at z [18, Thm. 4]. Let z be an extreme point of W(A). Then there are $\theta_0 \in \mathbb{R}$ and $i_0 \in \{1, \ldots, n\}$ such that $z = z_{i_0}(\theta_0)$. Now, f_A^{-1} is strongly continuous at z if and only if $z_i(\theta_0) = z$ implies $z_i = z_{i_0}$ for all $i = 1, \ldots, n$, see [30, Thm. 2.1.1]. Leake at al. [30] state the latter condition by saying that the eigenfunctions corresponding to z at θ_0 do not split.

Remark 4.4. Strong continuity is connected to the openness of a linear map. For all $z \in W(A)$, the multi-valued map f_A^{-1} is strongly continuous at z if and only if the restricted linear map $v : \mathcal{D}(M_n) \to V(A_1, A_2)$, introduced in (4.1), is open at every point in the fiber $v^{-1}[(\operatorname{Re}(z), \operatorname{Im}(z))^{\mathrm{T}}]$, see [53, Coro. 5.2]. Moreover, for every $x \in V(A_1, A_2)$, the map v is open at some point in the relative interior² of $v^{-1}(x)$ if and only if v is open at every point in $v^{-1}(x)$.

Proposition 4.5. Let $z \in W(A)$, let $x := (\operatorname{Re}(z), \operatorname{Im}(z))^{\mathrm{T}} \in \mathbb{R}^2$, let ρ be the unique density matrix of $\mathcal{R}(A_1, A_2)$ that satisfies $v(\rho) = x$, and let $\pi: \mathcal{D}(M_n) \to \mathcal{D}(\mathcal{R}(A_1, A_2))$ denote the orthogonal projection. Then the following assertions are equivalent.

- f_A⁻¹ is strongly continuous at z,
 v: D(M_n) → V(A₁, A₂) is open at every point in v⁻¹(x),
 π: D(M_n) → D(R(A₁, A₂)) is open at every point in π⁻¹(ρ).

We have $z \in ex W(A) \Leftrightarrow x \in ex V(A_1, A_2) \Leftrightarrow \rho \in ex \mathcal{D}(\mathcal{R}(A_1, A_2))$. If $z \in ex W(A)$, then $z = z_{i_0}(\theta_0)$ for some $\theta_0 \in \mathbb{R}$ and $i_0 \in \{1, \ldots, n\}$, and the following assertions are equivalent.

- f_A^{-1} is strongly continuous at z,
- the eigenfunctions corresponding to z at θ_0 do not split,
- π is open at some point in the relative interior of $\pi^{-1}(\rho)$.

Proof. The claims follow directly from Rem. 4.3 and 4.4, and Prop. 4.2.

12

²The relative interior [45] of a convex set K, denoted by ri(K), is the interior of K in the relative topology of the affine hull of K.

Example 4.6. Prop. 4.5 ignores that $\pi : \mathcal{D}(M_n) \to \mathcal{D}(\mathcal{R}(A_1, A_2))$ could be open at some point in the fiber $\pi^{-1}(\rho)$ over an extreme point ρ of $\mathcal{D}(\mathcal{R}(A_1, A_2))$, but not open anywhere in the relative interior of $\pi^{-1}(\rho)$.

This occurs for $A_1 := X \oplus 1$ and $A_2 := Z \oplus 0$. As discussed in Ex. 3.5 a) and b), the orthogonal projection $\pi_{\mathcal{R}} : \mathcal{D}(M_2 \oplus M_1) \to \mathcal{D}(\mathcal{R}(A_1, A_2))$ is open at $|+\rangle\langle+|\oplus 0$ and nowhere else in the fiber $\pi_{\mathcal{R}}^{-1}(M) = [|+\rangle\langle+|\oplus 0, 0\oplus 1]$ over $M = \frac{1}{2}(|+\rangle\langle+|\oplus 1)$. In particular, $\pi_{\mathcal{R}}$ is not open anywhere in the relative interior of $\pi_{\mathcal{R}}^{-1}(M)$. Ex. 8.3 proves the analogue for the larger fiber $\pi^{-1}(M)$.

5. Retractions

Generalizing a known result, we prove that the set of density matrices of every real *-subalgebra of M_n is a stable convex set. The proof relies on retractions of state spaces, a topic that also proved helpful in the foundations of quantum information theory [24].

Remark 5.1 (Stability of state spaces).

- a) The stability problem of a finite-dimensional compact convex set K is completely solved [40]. The *d*-skeleton of K is the union of all faces³ of K of dimension at most d. The convex set K is stable if and only if for every nonnegative integer d, the *d*-skeleton of K is closed.
- b) The set of density matrices $\mathcal{D}(M_n)$ is stable [20] because all its *d*-skeletons are closed. The closedness follows from three arguments: First, the set $\mathcal{D}(M_n)$ is a compact convex set of dimension $n^2 - 1$. Second, every nonempty face of M_n is unitarily similar to $\mathcal{D}(M_l) \oplus \{0\}$ for some positive integer *l*, and third, and the unitary group U(n) is compact.
- c) The state space $\mathcal{S}(\mathcal{A})$ of every *-subalgebra \mathcal{A} of M_n is stable [41] as it is a direct convex sum of state spaces of full matrix algebras M_n . As $\mathcal{S}(\mathcal{A})$ is stable, the set of density matrices $\mathcal{D}(\mathcal{A})$ is stable, too, by Lemma 2.2.

A retraction is an affine map $f: K \to L$ between compact convex sets K, L which is left-inverse to an affine map $g: L \to K$, called a section.

Proposition 5.2. If \mathcal{A} is a real *-subalgebra of M_n , then the orthogonal projection $\mathcal{D}(M_n) \to \mathcal{D}(\mathcal{A})$ is a retraction and $\mathcal{D}(\mathcal{A})$ is stable.

Proof. The orthogonal projection $\pi : \mathcal{D}(M_n) \to \mathcal{D}(\mathcal{A})$ is well defined by Lemma 2.5 and it is a retraction because the inclusion $\mathcal{D}(\mathcal{A}) \subset \mathcal{D}(M_n)$ of equation (2.4) provides a section for π . Coro. 1.3 in [16] asserts that the image of a stable convex set under a retraction is stable. Therefore, and since $\mathcal{D}(M_n)$ is stable by Rem. 5.1 b), the convex set $\mathcal{D}(\mathcal{A})$ is stable. \Box

Example 5.3. Prop. 5.2 does not generalize to arbitrary matrix systems. We consider the hermitian matrices

 $A_1 := X \oplus 1 \oplus 1$, $A_2 := Z \oplus 0 \oplus 0$, $A_3 := 0 \oplus (-1) \oplus 1$,

in the algebra $M_2 \oplus M_1 \oplus M_1$. The joint numerical range $V(A_1, A_2, A_3)$, introduced in Sec. 4, is easier to handle algebraically than the set of density

³A face [45] of a convex set K is a convex subset $F \subset K$ such that $x, y \in F$ is implied by $(1-s)x + sy \in F$ for all $x, y \in K$ and $s \in (0, 1)$.

matrices $\mathcal{D}(\mathcal{R}(A_1, A_2, A_3))$, to which it is affinely isomorphic by Lemma 4.1. Since $A_1, A_2, A_3 \in M_2 \oplus M_1 \oplus M_1$, we have the standard result of

$$V(A_1, A_2, A_3) = \operatorname{conv} \left(V(X, Z, 0) \cup V(1, 0, -1) \cup V(1, 0, 1) \right).$$

Hence, $V(A_1, A_2, A_3) = \operatorname{conv}(S)$ is the convex hull of

$$S := \{ (c_1, c_2, 0)^{\mathrm{T}} \in \mathbb{R}^3 \mid c_1^2 + c_2^2 = 1 \} \cup \{ (1, 0, -1)^{\mathrm{T}}, (1, 0, 1)^{\mathrm{T}} \}.$$

The set of extreme points $S \setminus \{(1, 0, 0)^{\mathrm{T}}\}$ of $V(A_1, A_2, A_3)$ is not closed, hence $V(A_1, A_2, A_3)$ is not stable by Rem. 5.1 a).

6. Proof of Thm. 1.2

The main ideas in establishing Thm. 1.2 are that the set of density matrices $\mathcal{D}(M_n)$ is a stable convex set and that $\mathcal{D}(M_n)$ is highly symmetric. Being stable and symmetric, $\mathcal{D}(M_{p+q})$ projects openly onto $\mathcal{D}(M_p \oplus M_q)$. Loosely speaking, Thm. 1.2 is obtained by combining various such open projections, since every *-subalgebra of M_n is a direct sum of full matrix algebras.

If K is a stable convex set, then the *arithmetic mean map*

$$K^{\times k} \to K$$
, $(x_1, \dots, x_k) \mapsto \frac{1}{k} \sum_{i=1}^k x_i$

the k-fold cartesian product of K is open for all positive integers k. More generally, for every tuple (s_1, \ldots, s_k) of nonnegative real numbers adding up to one, the map

$$K^{\times k} \to K$$
, $(x_1, \dots, x_k) \mapsto \sum_{i=1}^k s_i x_i$

is open. The latter assertion is proved for k = 2 in Prop. 1.1 in [17]. By induction, it is true for every k > 2 as well.

Lemma 6.1. Let \mathcal{R} be a real matrix system on \mathbb{C}^n and let $\mathcal{D}(\mathcal{R})$ be stable and invariant under an orthogonal transformation $\gamma : \mathcal{R} \to \mathcal{R}$ that generates a finite cyclic group. Let $\mathcal{A} := \{A \in \mathcal{R} : \gamma(A) = A\}$ be a real *-subalgebra of M_n . Then the orthogonal projection $\mathcal{D}(\mathcal{R}) \to \mathcal{D}(\mathcal{A})$ is open.

Proof. Let k denote the order of the cyclic group generated by γ . The main idea is to write the orthogonal projection $\pi : \mathcal{D}(\mathcal{R}) \to \mathcal{D}(\mathcal{A})$ in terms of the arithmetic mean map

$$a: \mathcal{D}(\mathcal{R})^{\times k} \to \mathcal{D}(\mathcal{R}), \quad (\rho_1, \dots, \rho_k) \mapsto \frac{1}{k} \sum_{i=1}^k \rho_i,$$

which, as discussed above, is open because $\mathcal{D}(\mathcal{R})$ is stable. The proof is done in two steps. First, we prove that

$$a(\gamma(\mathcal{O}) \times \dots \times \gamma^k(\mathcal{O})) \cap \mathcal{A}$$
 (6.1)

is an open subset of $\mathcal{D}(\mathcal{A})$ for all open subsets \mathcal{O} of $\mathcal{D}(\mathcal{R})$. Secondly, we prove that

$$a(\gamma(\mathcal{K}) \times \dots \times \gamma^k(\mathcal{K})) \cap \mathcal{A} = \pi(\mathcal{K})$$
(6.2)

holds for all convex subsets \mathcal{K} of $\mathcal{D}(\mathcal{R})$. The assertions (6.1) and (6.2) together show that $\pi(\mathcal{O})$ is an open subset of $\mathcal{D}(\mathcal{A})$ for all convex open subsets \mathcal{O} of $\mathcal{D}(\mathcal{R})$, and hence for all open subsets.

First, we prove that the set in (6.1) is open. As $\mathcal{D}(\mathcal{R})$ is invariant under the orthogonal transformation γ , the map γ restricts to a homeomorphism $\mathcal{D}(\mathcal{R}) \to \mathcal{D}(\mathcal{R})$. It follows that $\gamma(\mathcal{O}) \times \cdots \times \gamma^k(\mathcal{O})$ is an open subset of the *k*-fold cartesian product $\mathcal{D}(\mathcal{R})^{\times k}$. Then

$$\widetilde{\mathcal{O}} := a(\gamma(\mathcal{O}) \times \cdots \times \gamma^k(\mathcal{O}))$$

is an open subset of $\mathcal{D}(\mathcal{R})$, because $\mathcal{D}(\mathcal{R})$ is stable. Finally, $\widetilde{\mathcal{O}} \cap \mathcal{A}$ is an open subset of $\mathcal{D}(\mathcal{A})$ by equation (2.4), which shows that $\widetilde{\mathcal{O}} \cap \mathcal{A}$ equals $\widetilde{\mathcal{O}} \cap \mathcal{D}(\mathcal{A})$.

Secondly, we prove the formula (6.2), beginning with the inclusion " \supset ". Let $\rho \in \mathcal{K} \subset \mathcal{D}(\mathcal{R})$. The density matrix $\sigma := a(\gamma(\rho), \ldots, \gamma^k(\rho))$ is invariant under γ . This implies $\sigma \in \mathcal{A}$, hence $\sigma \in \mathcal{D}(\mathcal{A})$ by (2.4). Since γ is selfadjoint, for all $A \in \mathcal{H}(\mathcal{A})$ we have

$$\operatorname{Re}\langle\rho,A\rangle = \operatorname{Re}\langle a(\rho,\ldots,\rho),A\rangle = \operatorname{Re}\langle a(\gamma(\rho),\ldots,\gamma^k(\rho)),A\rangle = \operatorname{Re}\langle\sigma,A\rangle,$$

hence $\pi(\rho) = \sigma$. Regarding the inclusion " \subset ", let $\rho_i \in \mathcal{K}$, $i = 1, \ldots, k$. Let $\sigma := a(\gamma(\rho_1), \ldots, \gamma^k(\rho_k))$ and $\tau := a(\rho_1, \ldots, \rho_k)$. Then for all $A \in \mathcal{H}(\mathcal{A})$

$$\operatorname{Re}\langle\sigma,A\rangle = \operatorname{Re}\langle a(\gamma(\rho_1),\ldots,\gamma^k(\rho_k)),A\rangle = \operatorname{Re}\langle a(\rho_1,\ldots,\rho_k),A\rangle = \operatorname{Re}\langle\tau,A\rangle$$

holds. If $\sigma \in \mathcal{A}$, then $\sigma = \pi(\tau)$ follows. As \mathcal{K} is convex, $\tau \in \mathcal{K}$ holds and we obtain $\sigma \in \pi(\mathcal{K})$.

Remark 6.2. A hermitian matrix $M \in \mathcal{H}(M_{p+q})$ in the block form

$$M = \begin{pmatrix} A & B \\ B^* & C \end{pmatrix}, \quad A \in \mathcal{H}(M_p), \quad B \in \mathbb{C}^{p \times q}, \quad C \in \mathcal{H}(M_q)$$

is positive semidefinite if and only if the top left block A is positive semidefinite, the range of B is included in the range of A, and the generalized Schur complement $M/A = C - B^*A^-B$ is positive semidefinite, where A^- is a generalized inverse of A, that is to say, $A^- \in M_p$ and $AA^-A = A$ holds [25].

Proposition 6.3. The orthogonal projection $\mathcal{D}(M_{p+q}) \to \mathcal{D}(M_p \oplus M_q)$ is open.

Proof. Lemma 6.1 proves the claim when $\mathcal{R} := M_{p+q}$ and $\mathcal{A} := M_p \oplus M_q$. The set of density matrices $\mathcal{D}(M_{p+q})$ is stable by Rem. 5.1 b). The reflection $\gamma : M_{p+q} \to M_{p+q}$ at the subspace $M_p \oplus M_q$ generates a group of order two. We show that $\mathcal{D}(M_{p+q})$ is invariant under γ . In block form, the reflection reads

$$\gamma: \begin{pmatrix} A & B \\ D & C \end{pmatrix} \mapsto \begin{pmatrix} A & -B \\ -D & C \end{pmatrix} +$$

where $A \in \mathcal{M}_p$, $B \in \mathbb{C}^{p \times q}$, $C \in \mathcal{M}_q$, and $D \in \mathbb{C}^{q \times p}$. The space of hermitian matrices is invariant under γ , which restricts to

$$\mathcal{H}(\mathcal{M}_{p+q}) \to \mathcal{H}(\mathcal{M}_{p+q}), \quad M = \begin{pmatrix} A & B \\ B^* & C \end{pmatrix} \mapsto M' = \begin{pmatrix} A & -B \\ -B^* & C \end{pmatrix},$$

where $A \in \mathcal{H}(M_p)$, $B \in \mathbb{C}^{p \times q}$, and $C \in \mathcal{H}(M_q)$. By Rem. 6.2, the map $M \mapsto M'$ preserves the positive semidefiniteness since M and M' have the same diagonal blocks and both off-diagonal blocks differ in a sign, so that M'/A = M/A holds. The map γ preserves the trace. The set of fixed points $M_p \oplus M_q = \{A \in M_{p+q} : \gamma(A) = A\}$ is a *-subalgebra of M_{p+q} . \Box

Corollary 6.4. The orthogonal projection $\mathcal{D}(M_n) \to \mathcal{D}(M_{n_1} \oplus \cdots \oplus M_{n_k})$ is open, where $n = n_1 + \cdots + n_k$.

Proof. Proceeding by induction, we observe that the orthogonal projection $\mathcal{D}(M_{n_1}) \to \mathcal{D}(M_{n_1})$ is of course open. Let $k \geq 1$ and assume that the orthogonal projection

$$\pi_k: \mathcal{D}(\mathcal{M}_{n_1+\cdots+n_k}) \to \mathcal{D}(\mathcal{M}_{n_1} \oplus \cdots \oplus \mathcal{M}_{n_k})$$

is open. The orthogonal projection $M_{n_1+\dots+n_{k+1}} \to M_{n_1} \oplus \dots \oplus M_{n_{k+1}}$ factors into the orthogonal projections

 $\mathcal{M}_{n_1+\dots+n_{k+1}} \to \mathcal{M}_{n_1+\dots+n_k} \oplus \mathcal{M}_{n_{k+1}}$

and $M_{n_1+\dots+n_k} \oplus M_{n_{k+1}} \to M_{n_1} \oplus \dots \oplus M_{n_k} \oplus M_{n_{k+1}}$.

Hence, by Lemma 2.5, the map π_{k+1} factors into the orthogonal projections

$$\mathcal{D}(\mathcal{M}_{n_1+\dots+n_{k+1}}) \to \mathcal{D}(\mathcal{M}_{n_1+\dots+n_k} \oplus \mathcal{M}_{n_{k+1}})$$
(6.3)

and
$$\mathcal{D}(\mathcal{M}_{n_1+\dots+n_k} \oplus \mathcal{M}_{n_{k+1}}) \to \mathcal{D}(\mathcal{M}_{n_1} \oplus \dots \oplus \mathcal{M}_{n_k} \oplus \mathcal{M}_{n_{k+1}}).$$
 (6.4)

The map (6.3) is open by Prop. 6.3. Lemma 3.7 shows that (6.4) equals

$$\mathcal{D}(\mathcal{M}_{n_1+\dots+n_k})\oplus_{\mathbf{c}}\mathcal{D}(\mathcal{M}_{n_{k+1}})\to\mathcal{D}(\mathcal{M}_{n_1}\oplus\dots\oplus\mathcal{M}_{n_k})\oplus_{\mathbf{c}}\mathcal{D}(\mathcal{M}_{n_{k+1}})$$

which is open by Prop. 3.2 (for m = 2) and by the induction hypothesis. Being the composition of two open maps, π_{k+1} is open.

Proposition 6.5. The orthogonal projection $\mathcal{D}(\bigoplus_{i=1}^k M_q) \to \mathcal{D}(M_q \otimes \mathbb{1}_k)$ is open.

Proof. Lemma 6.1 proves the claim when $\mathcal{R} := \bigoplus_{i=1}^{k} M_q$ and $\mathcal{A} := M_q \otimes \mathbb{1}_k$. The convex set $\mathcal{D}(\mathcal{R})$ is stable by Rem. 5.1 c) and equals the k-fold direct convex sum $\mathcal{D}(\mathcal{R}) = \mathcal{D}(M_q) \oplus_c \cdots \oplus_c \mathcal{D}(M_q)$ of $\mathcal{D}(M_q)$ by Lemma 3.4. Hence, the cyclic permutation $\gamma = (1, \ldots, k)$ defines the orthogonal transformation

 $\gamma: \mathcal{D}(\mathcal{R}) \to \mathcal{D}(\mathcal{R}), \quad (\sigma_1, \dots, \sigma_k) \mapsto (\sigma_{\gamma^{-1}(1)}, \dots, \sigma_{\gamma^{-1}(k)}),$

which generates a group of order k. Clearly, $\mathcal{D}(\mathcal{R})$ is invariant under γ and $\mathcal{A} = \{A \in \mathcal{R} : \gamma(A) = A\}$ is a *-subalgebra of M_{kq} .

Proof of Thm. 1.2. Since \mathcal{A} is a *-subalgebra of M_n , there exists a unitary $n \times n$ matrix U such that $U\mathcal{A}U^* = \bigoplus_{i=1}^m \mathcal{A}_i$, where $\mathcal{A}_i := M_{q_i} \otimes \mathbb{1}_{k_i}$ for every $i = 1, \ldots, m$, and $q_1k_1 + \cdots + q_mk_m = n$, see [21, Thm. 5.6].

As $\mathcal{D}(\mathbf{M}_n) \to \mathcal{D}(\mathbf{M}_n)$, $\rho \mapsto U\rho U^*$ is a homeomorphism, it suffices to prove that the orthogonal projection $\pi : \mathcal{D}(\mathbf{M}_n) \to \mathcal{D}(\bigoplus_{i=1}^m \mathcal{A}_i)$ is open. The orthogonal projection $\mathbf{M}_n \to \bigoplus_{i=1}^m \mathcal{A}_i$ factors into the orthogonal projections

and
$$\mathcal{B}_1 \oplus \cdots \oplus \mathcal{B}_m \to \mathcal{A}_1 \oplus \cdots \oplus \mathcal{A}_m$$
,

where $\mathcal{B}_i := \bigoplus_{j=1}^{k_i} M_{q_i}$, i = 1, ..., m. By Lemma 2.5, the map π factors into

$$\mathcal{D}(\mathcal{M}_n) \to \mathcal{D}(\mathcal{B}_1 \oplus \cdots \oplus \mathcal{B}_m),$$
 (6.5)

and
$$\mathcal{D}(\mathcal{B}_1 \oplus \cdots \oplus \mathcal{B}_m) \to \mathcal{D}(\mathcal{A}_1 \oplus \cdots \oplus \mathcal{A}_m).$$
 (6.6)

The map (6.5) is open by Coro. 6.4 (for $k = k_1 + \cdots + k_m$). Lemma 3.7 shows that the map (6.6) equals

$$\mathcal{D}(\mathcal{B}_1) \oplus_{\mathrm{c}} \cdots \oplus_{\mathrm{c}} \mathcal{D}(\mathcal{B}_m) \to \mathcal{D}(\mathcal{A}_1) \oplus_{\mathrm{c}} \cdots \oplus_{\mathrm{c}} \mathcal{D}(\mathcal{A}_m),$$

which is open by Prop. 3.2 and Prop. 6.5. In conclusion, π is open as it is a composition of two open maps.

7. Real *-subalgebras of M_n

Every real *-subalgebra of M_n is *-isomorphic [21, Thm. 5.22] to a direct sum of algebras of real, complex, and quaternionic q-by-q-matrices of various sizes q. We are here interested in the algebra $M_n(\mathbb{R})$ of real $n \times n$ matrices.

Proposition 7.1. The orthogonal projection $\mathcal{D}(M_n) \to \mathcal{D}(M_n(\mathbb{R}))$ is open.

Proof. Lemma 6.1 proves the claim when $\mathcal{R} := M_n$ and $\mathcal{A} := M_n(\mathbb{R})$. The set of density matrices $\mathcal{D}(M_n)$ is stable by Rem. 5.1 b). The reflection $\gamma : M_n \to M_n$ at the real subspace $M_n(\mathbb{R})$ generates a group of order two.

The algebra M_n is the orthogonal direct sum $M_n = M_n(\mathbb{R}) \oplus i M_n(\mathbb{R})$ and the reflection at the real subspace $M_n(\mathbb{R})$ reads

$$\gamma: \mathbf{M}_n \to \mathbf{M}_n, \quad A + \mathbf{i}B \mapsto A - \mathbf{i}B, \quad A, B \in \mathbf{M}_n(\mathbb{R}).$$

The orthogonal transformation γ preserves the space of hermitian matrices, which is the orthogonal direct sum

$$\mathcal{H}(\mathcal{M}_n) = \operatorname{Sym}_n(\mathbb{R}) \oplus \operatorname{i} \operatorname{Skew}_n(\mathbb{R})$$

of the space of real symmetric matrices

$$\operatorname{Sym}_{n}(\mathbb{R}) := \{A \in \operatorname{M}_{n}(\mathbb{R}) \colon A^{\mathrm{T}} = A\} = \mathcal{H}(\operatorname{M}_{n}(\mathbb{R}))$$

and the space of skew-symmetric matrices

$$\operatorname{Skew}_n(\mathbb{R}) := \{A \in \operatorname{M}_n(\mathbb{R}) \colon A^{\mathrm{T}} = -A\}.$$

We prove that γ preserves the trace and the positive semidefiniteness on the space of hermitian matrices. Let $A, C \in \operatorname{Sym}_n(\mathbb{R})$ and $B, D \in \operatorname{Skew}_n(\mathbb{R})$. Then $\operatorname{tr}(A + iB) = \operatorname{tr}(A)$ shows that the trace is preserved. It is well known that a matrix is positive semidefinite if and only if its inner product with the square of every hermitian matrix is nonnegative. Thus

$$\langle A - iB, (C + iD)^2 \rangle = \langle A + iB, (C - iD)^2 \rangle$$

shows that A - iB is positive semidefinite if A + iB is positive semidefinite. Clearly, $M_n(\mathbb{R}) = \{A \in M_n : \gamma(A) = A\}$ is a real *-subalgebra of M_n . \Box

As the orthogonal projection $M_n \to M_n(\mathbb{R})$ is the entrywise real part, we denote by Re : $\mathcal{D}(M_n) \to \mathcal{D}(M_n(\mathbb{R}))$ the orthogonal projection of $\mathcal{D}(M_n)$ onto $\mathcal{D}(M_n(\mathbb{R}))$.

Example 7.2. We consider the chain $M_3 \supset M_2 \oplus M_1 \supset M_2(\mathbb{R}) \oplus M_1(\mathbb{R})$ of real *-subalgebras of M_3 . The orthogonal projections

$$M_3 \longrightarrow M_2 \oplus M_1 \longrightarrow M_2(\mathbb{R}) \oplus M_1(\mathbb{R})$$

restrict by Lemma 2.5 to

 $\mathcal{D}(M_3) \xrightarrow{\pi_1} \mathcal{D}(M_2 \oplus M_1) \xrightarrow{\pi_2} \mathcal{D}(M_2(\mathbb{R}) \oplus M_1(\mathbb{R})) \,.$

The map π_1 is open by Prop. 6.3. By Lemma 3.7, the map π_2 is the direct convex sum

$$\operatorname{Re} \oplus_{\operatorname{c}} \operatorname{id} : \mathcal{D}(\operatorname{M}_2) \oplus_{\operatorname{c}} \{1\} \to \mathcal{D}(\operatorname{M}_2(\mathbb{R})) \oplus_{\operatorname{c}} \{1\}$$

of Re : $\mathcal{D}(M_2) \to \mathcal{D}(M_2(\mathbb{R}))$ and the identity map id : $\{1\} \to \{1\}$. The map Re is open by Prop. 7.1, hence π_2 is open by Prop. 3.2. The orthogonal projection $\mathcal{D}(M_3) \to \mathcal{D}(M_2(\mathbb{R}) \oplus M_1(\mathbb{R}))$ is open, as it is a composition of two open maps.

8. TOPOLOGY SIMPLIFIED BY ALGEBRA

Thm. 1.2 can simplify topology problems. Given topological spaces K, L, a map $f : K \to L$ is *continuous* [27] at $x \in K$ if the preimage of every neighborhood of f(x) in L is a neighborhood of x in K.

Lemma 8.1. Let $\mathcal{R}_1, \mathcal{R}_2$ be real matrix systems on \mathbb{C}^n such that $\mathcal{R}_2 \subset \mathcal{R}_1$. Let $\pi_1 : \mathcal{D}(\mathcal{M}_n) \to \mathcal{D}(\mathcal{R}_1)$ and $\pi_2 : \mathcal{D}(\mathcal{R}_1) \to \mathcal{D}(\mathcal{R}_2)$ denote the orthogonal projections and assume the orthogonal projection $\pi_2 \circ \pi_1 : \mathcal{D}(\mathcal{M}_n) \to \mathcal{D}(\mathcal{R}_2)$ is open. Let $f : \mathcal{D}(\mathcal{R}_2) \to T$ be a map to a topological space T. Let $\rho \in \mathcal{D}(\mathcal{R}_1)$.

- a) The map $f \circ \pi_2 : \mathcal{D}(\mathcal{R}_1) \to T$ is open at ρ if and only if $f : \mathcal{D}(\mathcal{R}_2) \to T$ is open at $\pi_2(\rho)$.
- b) The map $f \circ \pi_2 : \mathcal{D}(\mathcal{R}_1) \to T$ is continuous at ρ if and only if the map $f : \mathcal{D}(\mathcal{R}_2) \to T$ is continuous at $\pi_2(\rho)$.

Proof. The orthogonal projection $\mathcal{D}(\mathcal{M}_n) \to \mathcal{D}(\mathcal{R}_2)$ equals indeed $\pi_2 \circ \pi_1$ by Lemma 2.5.

We begin with the implication " \Rightarrow " of a). If $\mathcal{N}_2 \subset \mathcal{D}(\mathcal{R}_2)$ is a neighborhood of $\pi_2(\rho)$, then

$$f(\mathcal{N}_2) = (f \circ \pi_2) \circ \pi_2^{-1}(\mathcal{N}_2)$$

is a neighborhood of $f(\pi_2(\rho))$ because π_2 is continuous and $f \circ \pi_2$ is open at ρ . Regarding the implication " \Leftarrow ", we choose a neighborhood $\mathcal{N}_1 \subset \mathcal{D}(\mathcal{R}_1)$ of ρ . Then

$$f \circ \pi_2(\mathcal{N}_1) = f \circ (\pi_2 \circ \pi_1) \circ \pi_1^{-1}(\mathcal{N}_1)$$

is a neighborhood of $f(\pi_2(\rho))$ because π_1 is continuous, $\pi_2 \circ \pi_1$ is open, and f is open at $\pi_2(\rho)$.

To prove b) we choose a neighborhood $\mathcal{N}_T \subset T$ of $f(\pi_2(\rho))$. Regarding the implication " \Rightarrow ", the preimage

$$f^{-1}(\mathcal{N}_T) = (\pi_2 \circ \pi_1) \circ \pi_1^{-1} \circ (f \circ \pi_2)^{-1}(\mathcal{N}_T)$$

is a neighborhood of $\pi_2(\rho)$, because $f \circ \pi_2$ is continuous at ρ , the map π_1 is continuous, and $\pi_2 \circ \pi_1$ is open. Regarding the implication " \Leftarrow ", the preimage

$$(f \circ \pi_2)^{-1}(\mathcal{N}_T) = \pi_2^{-1} \circ f^{-1}(\mathcal{N}_T)$$

is a neighborhood of ρ , as f is continuous at $\pi_2(\rho)$, and π_2 is continuous. \Box

Remark 8.2 (Simplifying openness problems). Let $\pi : \mathcal{D}(M_n) \to \mathcal{D}(\mathcal{R})$ be the orthogonal projection to a real matrix system \mathcal{R} on \mathbb{C}^n and let \mathcal{A} be a *-subalgebra of M_n such that $\mathcal{R} \subset \mathcal{A}$. Then $\pi = \pi_{\mathcal{R}} \circ \pi_{\mathcal{A}}$ factors into the orthogonal projections $\pi_{\mathcal{A}} : \mathcal{D}(M_n) \to \mathcal{D}(\mathcal{A})$ and $\pi_{\mathcal{R}} : \mathcal{D}(\mathcal{A}) \to \mathcal{D}(\mathcal{R})$. For every $\rho \in \mathcal{D}(M_n)$ the map π is open at ρ if and only if $\pi_{\mathcal{R}}$ is open at $\pi_{\mathcal{A}}(\rho)$.

As a word of explication, the map π factors by Lemma 2.5. The claim follows from Lemma 8.1 a), by letting $\mathcal{R}_1 := M_n$ and $\mathcal{R}_2 := \mathcal{A}$, and by taking $\pi_{\mathcal{R}}$ as the map $f : \mathcal{D}(\mathcal{R}_2) \to T$. The assumptions of the lemma are met since $\pi_{\mathcal{A}}$ is open by Thm. 1.2.

The following two examples are concerned with openness. Continuity is a topic of Sec. 9 below.

Example 8.3. Let $\mathcal{R} := \operatorname{span}_{\mathbb{R}}(\mathbb{1}_3, X \oplus 1, Z \oplus 0).$

- a) Rem. 8.2 solves the openness problem of the orthogonal projection π : $\mathcal{D}(M_3) \to \mathcal{D}(\mathcal{R})$ to \mathcal{R} , based on the solution of the openness problem of $\pi_{\mathcal{R}}: \mathcal{D}(\mathcal{A}) \to \mathcal{D}(\mathcal{R})$, where $\mathcal{A} := M_2 \oplus M_1$. By Ex. 3.5 a), the map $\pi_{\mathcal{R}}$ is not open at any point in the half-open segment $\mathcal{G}_0 = (|+\rangle \langle +| \oplus 0, 0 \oplus 1] \subset \mathcal{D}(\mathcal{A})$ and open at every point in the complement. By Rem. 8.2, the map $\pi: \mathcal{D}(M_3) \to \mathcal{D}(\mathcal{R})$ is not open at any point in $\pi_{\mathcal{A}}^{-1}(\mathcal{G}_0)$ and open at every point in the complement.
- b) To describe $\pi_{\mathcal{A}}^{-1}(\mathcal{G}_0)$, we study the fibers of the orthogonal projection

$$\mathcal{H}(\mathcal{M}_3) \to \mathcal{H}(\mathcal{A}), \quad \begin{pmatrix} A & |\varphi\rangle\\ \langle \varphi| & c \end{pmatrix} \mapsto \begin{pmatrix} A & 0\\ 0 & c \end{pmatrix},$$

where $A \in \mathcal{H}(M_2)$, $|\varphi\rangle \in \mathbb{C}^2 \cong \mathbb{C}^{2\times 1}$, and $c \in \mathbb{R} \cong \mathcal{H}(M_1)$. Every point in $\mathcal{G} = [|+\rangle \langle +| \oplus 0, 0 \oplus 1]$ is of the form $(1-\lambda) |+\rangle \langle +| \oplus \lambda$ for some $\lambda \in [0,1]$. Using the generalized Schur complement (Rem. 6.2), one verifies that the fiber of $\pi_{\mathcal{A}}$ over this point is the set of all matrices

$$\rho(\lambda, z) := \begin{pmatrix} (1-\lambda) \mid + \rangle \langle + \mid & z \mid + \rangle \\ \overline{z} \langle + \mid & \lambda \end{pmatrix}, \quad z \in \mathbb{C}, \quad |z|^2 \le \lambda (1-\lambda), \quad (8.1)$$

where |z| denotes the absolute value of $z \in \mathbb{C}$. In conclusion, the orthogonal projection $\pi : \mathcal{D}(M_3) \to \mathcal{D}(\mathcal{R})$ is not open at any point of

$$\pi_{\mathcal{A}}^{-1}(\mathcal{G}_0) = \{\rho(\lambda, z) \colon z \in \mathbb{C}, |z|^2 \le \lambda(1-\lambda), \lambda \in (0, 1]\}$$

and open at every point in the complement.

τ

c) We verify a claim made in Ex. 4.6. The segment $\mathcal{G} = [|+\rangle\langle +|\oplus 0, 0\oplus 1]$ is the fiber of $\pi_{\mathcal{R}}$ over $M = \frac{1}{2}(|+\rangle\langle +|\oplus 1) \in \mathcal{D}(\mathcal{R})$ by Ex. 3.5 b), so $\pi_{\mathcal{A}}^{-1}(\mathcal{G})$ is the fiber of $\pi = \pi_{\mathcal{R}} \circ \pi_{\mathcal{A}}$ over M. As recalled in part a) above, the map π is open at $|+\rangle\langle +|\oplus 0$ but not open at any point of $\pi_{\mathcal{A}}^{-1}(\mathcal{G}_0)$. We now observe that π is not open at any point in the relative interior of $\pi^{-1}(M)$, as we have the chain of inclusions

$$\pi_{\mathcal{A}}\Big(\operatorname{ri}\left((\pi_{\mathcal{R}}\circ\pi_{\mathcal{A}})^{-1}(M)\right)\Big) = \operatorname{ri}\left(\pi_{\mathcal{A}}\left((\pi_{\mathcal{R}}\circ\pi_{\mathcal{A}})^{-1}(M)\right)\right)$$
$$=\operatorname{ri}(\pi_{\mathcal{R}}^{-1}(M)) = \operatorname{ri}(\mathcal{G}) = \mathcal{G}_0 \setminus \{0 \oplus 1\} \subset \mathcal{G}_0,$$

whose first equality holds by [45, Thm. 6.6].

Example 8.4. The recipe of Rem. 8.2 helps analyze the openness of the orthogonal projection $\mathcal{D}(M_n) \to \mathcal{D}(\mathcal{R}(P,Q))$ to the real matrix system

$$\mathcal{R}(P,Q) = \operatorname{span}_{\mathbb{R}}(\mathbb{1}_n, P, Q)$$

generated by two orthogonal projections $P, Q \in M_n$, that is to say, matrices satisfying $P = P^2 = P^*$ and $Q = Q^2 = Q^*$. It is well known that the matrix system $\mathcal{R}(P,Q)$ is included in a surprisingly small *-subalgebra of M_n , see Coro. 2.2 in the survey [8] by Böttcher and Spitkovsky, and the references therein. More precisely, there exists a unitary $n \times n$ matrix U, nonnegative integers $m_1 \leq 4$ and m_2 , and positive integers k_i , $i = 1, \ldots, m_1$ satisfying $k_1 + \cdots + k_{m_1} + 2m_2 = n$, such that $\mathcal{R} := U\mathcal{R}(P,Q)U^*$ is included in

$$\mathcal{A} := \left(\bigoplus_{i=1}^{m_1} \mathrm{M}_1 \otimes \mathbb{1}_{k_i} \right) \oplus \left(\bigoplus_{j=1}^{m_2} \mathrm{M}_2 \right).$$

Since $\rho \mapsto U\rho U^*$ is a homeomorphism of $\mathcal{D}(M_n)$, the openness problems of the orthogonal projections $\mathcal{D}(M_n) \to \mathcal{D}(\mathcal{R}(P,Q))$ and $\mathcal{D}(M_n) \to \mathcal{D}(\mathcal{R})$

are equivalent. The second one is substantially simplified by the method of Rem. 8.2 as $\mathcal{D}(\mathcal{A})$ has a rather simple shape. It is the direct convex sum of several three-dimensional Euclidean balls and a simplex of dimension at most three by Lemma 3.4 and Ex. 2.3. This observation should also simplify the strong continuity problem for the numerical range W(P + iQ).

9. Continuity in quantum information theory

We discuss continuity problems of entropic inference maps and of measures of correlation. We assume that \mathcal{R} is a real matrix system on \mathbb{C}^n and that, without loss of generality (see Rem. 2.4), we have $\mathcal{R} \subset \mathcal{H}(M_n)$.

Example 9.1 (Maximum entropy inference I). The purpose of the maximum entropy inference method is to update a prior probability distribution if new information becomes available in the form of constraints that specify a set of possible posterior probability distributions. The preferred posterior is that which minimizes the relative entropy from the prior subject to the available constraints, see Chap. 8 in Caticha's book [13], and the references therein. Whereas the maximum entropy method has a sound axiomatic basis, the analogous inference method for quantum states lacks an axiomatic foundation and remains a controversial topic [3].

Analytic properties of quantum mechanical inference are interesting despite unsolved foundational issues. Notably, quantum inference maps have discontinuities [55] that aroused interest in theoretical physics [14,26,36,56].

Linear constraints⁴ on $\mathcal{D}(M_n)$ are defined by the orthogonal projection

$$\pi: \mathcal{D}(\mathcal{M}_n) \to \mathcal{D}(\mathcal{R}).$$

The relative entropy $S : \mathcal{D}(M_n) \times \mathcal{D}(M_n) \to [0, +\infty]$ is an asymmetric distance. It is defined as $S(\rho_1, \rho_2) := \operatorname{tr}[\rho_1(\log(\rho_1) - \log(\rho_2))]$ if the range of ρ_1 is included in the range of ρ_2 and as $S(\rho_1, \rho_2) := +\infty$ otherwise, for all $\rho_1, \rho_2 \in \mathcal{D}(M_n)$. Let $\tau \in \mathcal{D}(M_n)$, the prior, be a density matrix of maximal rank n. Then

$$\phi_{\tau}: \mathcal{D}(\mathbf{M}_n) \to \mathbb{R}, \quad \rho \mapsto -S(\rho, \tau)$$

is continuous and strictly concave. So the maximum entropy inference map

$$\Psi_{\tau} : \mathcal{D}(\mathcal{R}) \to \mathcal{D}(\mathcal{M}_n), \quad \sigma \mapsto \operatorname{argmax}_{\rho \in \pi^{-1}(\sigma)} \phi_{\tau}(\rho) \tag{9.1}$$

is well defined, see [52, Def. 1.1] and the references therein.

We quote [52, Thm. 4.9] under the assumptions of Ex. 9.1.

Theorem 9.2. For any $\sigma \in \mathcal{D}(\mathcal{R})$, the inference map $\Psi_{\tau} : \mathcal{D}(\mathcal{R}) \to \mathcal{D}(\mathcal{M}_n)$ is continuous at σ if and only if the orthogonal projection $\pi : \mathcal{D}(\mathcal{M}_n) \to \mathcal{D}(\mathcal{R})$ is open at $\Psi_{\tau}(\sigma)$.

⁴Linear constraints can be defined in terms of expectation values. Let A_1, \ldots, A_k be hermitian $n \times n$ matrices such that $\mathcal{R} = \operatorname{span}_{\mathbb{R}}(\mathbb{1}_n, A_1, \ldots, A_k)$. The observables represented by A_1, \ldots, A_k have the *expectation values* $v(\rho) = (\langle \rho, A_1 \rangle, \ldots, \langle \rho, A_k \rangle)^{\mathrm{T}}$ if $\rho \in \mathcal{D}(\mathcal{M}_n)$ is the system state [7]. The fiber $\pi^{-1}(\sigma)$ over $\sigma \in \mathcal{D}(\mathcal{R})$ is the set of $\rho \in \mathcal{D}(\mathcal{M}_n)$ whose expectation values are $v(\rho) = v(\sigma)$, see Lemma 4.1.

Example 9.3 (Maximum entropy inference II). We resume Ex. 9.1. Thm. 9.2 translates the continuity problem of the inference map Ψ_{τ} into an openness problem of the orthogonal projection π . Whether Rem. 8.2 can simplify this openness problem depends on the location of the posteriors.

The range of Ψ_{τ} can be described explicitly. The manifold

$$\mathcal{E}_{\tau}(\mathcal{R}) := \left\{ \frac{e^{\log(\tau) + A}}{\operatorname{tr}(e^{\log(\tau) + A})} \mid A \in \mathcal{R} \right\}$$
(9.2)

is known as a Gibbsian family or exponential family, see [38,43,52,55] and the references therein (and recall $\mathcal{R} = \mathcal{H}(\mathcal{R})$). The manifold $\mathcal{E}_{\tau}(\mathcal{R})$ is included in $\Psi_{\tau}(\mathcal{D}(\mathcal{R}))$). But not just that, the set $\Psi_{\tau}(\mathcal{D}(\mathcal{R}))$ of maximum entropy inference states equals

$$\{\rho_1 \in \mathcal{D}(\mathcal{M}_n) : \inf_{\rho_2 \in \mathcal{E}_\tau(\mathcal{R})} S(\rho_1, \rho_2) = 0\}, \qquad (9.3)$$

which is a subset of the Euclidean closure of $\mathcal{E}_{\tau}(\mathcal{R})$, see (D5) in [52]. The set (9.3) is called *reverse information closure* or the *rI-closure* [19] of $\mathcal{E}(\mathcal{R})$.

Let \mathcal{A} denote the smallest *-subalgebra \mathcal{A} of M_n that contains \mathcal{R} .

a) Rem. 8.2 is of no avail if $\tau = \mathbb{1}_n/n$ is the uniform prior. In this case,

$$\phi_{\tau}(\rho) = -S(\rho, \mathbb{1}_n/n) = S(\rho) - \log(n), \quad \rho \in \mathcal{D}(\mathcal{M}_n)$$

is the von Neumann entropy $S(\rho) := -\operatorname{tr}[\rho \log(\rho)]$ up to a constant. By functional calculus, the exponential family $\mathcal{E}_{\tau}(\mathcal{R})$ is included in \mathcal{A} and so is the set of inference states $\Psi_{\tau}(\mathcal{D}(\mathcal{R}))$ as per the formula (9.3), because \mathcal{A} is closed. Rem. 8.2 is of no help since all posteriors lie in \mathcal{A} anyways. The same conclusion is true for every prior τ in \mathcal{A} replacing $\mathbb{1}_n/n$.

- b) If the prior τ lies outside of \mathcal{A} then the exponential family $\mathcal{E}_{\tau}(\mathcal{R})$ is disjoint from $\mathcal{D}(\mathcal{A})$, again by functional calculus. Rem. 8.2 brings an advantage if the orthogonal projection $\pi_{\mathcal{R}} : \mathcal{D}(\mathcal{A}) \to \mathcal{D}(\mathcal{R})$ is easier to analyze than $\pi : \mathcal{D}(M_n) \to \mathcal{D}(\mathcal{R})$. An example is given in Ex. 8.4 where \mathcal{A} has a simple structure and a much smaller dimension than M_n .
- c) The real matrix system $\mathcal{R} := \operatorname{span}_{\mathbb{R}}(\mathbb{1}_3, X \oplus \mathbb{1}, Z \oplus 0)$ in Ex. 8.3 has the property that the orthogonal projection $\pi : \mathcal{D}(M_3) \to \mathcal{D}(\mathcal{R})$ is not open anywhere in the relative interior of the fiber over $M := \frac{1}{2}(|+\rangle\langle +|\oplus 1\rangle)$ and open at every point in the complement of that fiber. For every $\sigma \in \mathcal{D}(\mathcal{R})$ and prior τ , the posterior $\Psi_{\tau}(\sigma)$ is contained in the relative interior of the fiber $\pi^{-1}(\sigma)$ over every σ by Coro. 5.7 and Lemma 5.8 in [52]. Thm. 9.2 then shows that Ψ_{τ} is discontinuous at M and continuous at every point different from M (for all priors τ).

Example 9.4 (Maximum entropy inference III). An important example from physics is the real matrix system of *local Hamiltonians* [14, 58].

Every unit $i \in \Omega := \{1, 2, \ldots, N\}$ of an N-qubit system is associated with a copy \mathcal{A}_i of the algebra M_2 . The subsystem with units in a subset $\nu \subset \Omega$ is associated with the tensor product algebra $\mathcal{A}_{\nu} := \bigotimes_{i \in \nu} \mathcal{A}_i$, whose identity we denote by $\mathbb{1}_{\nu}$. We have $\mathcal{A}_{\Omega} = M_n$ for $n = 2^N$. The algebra \mathcal{A}_{ν} embeds into \mathcal{A}_{Ω} via the map $\mathcal{A}_{\nu} \to \mathcal{A}_{\Omega}$, $A \mapsto A \otimes \mathbb{1}_{\bar{\nu}}$, where $\bar{\nu} = \Omega \setminus \nu$ is the complement of ν . Let \mathfrak{g} be a family of subsets of Ω . A \mathfrak{g} -local Hamiltonian is a hermitian matrix in \mathcal{A}_{Ω} of the form

$$\sum_{\nu \in \mathfrak{g}} A_{\nu} \otimes \mathbb{1}_{\bar{\nu}}, \quad A_{\nu} \in \mathcal{H}(\mathcal{A}_{\nu}), \quad \nu \in \mathfrak{g}.$$

$$(9.4)$$

We denote the real matrix system of all \mathfrak{g} -local Hamiltonians by $\mathcal{R}_{\mathfrak{g}}$ and the orthogonal projection by $\pi_{\mathfrak{g}}: \mathcal{D}(\mathcal{A}_{\Omega}) \to \mathcal{D}(\mathcal{R}_{\mathfrak{g}}).$

The partial trace $\operatorname{Tr}_{\bar{\nu}} : \mathcal{A}_{\Omega} \to \mathcal{A}_{\nu}$ is the adjoint of the embedding $\mathcal{A}_{\nu} \to \mathcal{A}_{\Omega}$ and satisfies $\langle A \otimes \mathbb{1}_{\bar{\nu}}, B \rangle = \langle A, \operatorname{Tr}_{\bar{\nu}}(B) \rangle$ for every $A \in \mathcal{A}_{\nu}, B \in \mathcal{A}_{\Omega}$. The partial trace $\operatorname{Tr}_{\bar{\nu}}(\rho)$ of $\rho \in \mathcal{D}(\mathcal{A}_{\Omega})$ is a density matrix of \mathcal{A}_{ν} called *reduced density matrix*. Let

$$\operatorname{red}_{\mathfrak{g}}: \mathcal{A}_{\Omega} \to \prod_{\nu \in \mathfrak{g}} \mathcal{A}_{\nu}, \quad A \mapsto [\operatorname{Tr}_{\bar{\nu}}(A)]_{\nu \in \mathfrak{g}}$$

$$\tag{9.5}$$

denote the map from \mathcal{A}_{Ω} to the cartesian product of the algebras $(\mathcal{A}_{\nu})_{\nu \in \mathfrak{g}}$ that assigns partial traces to the subsystems specified by the members of \mathfrak{g} .

Linear constraints on $\mathcal{D}(\mathcal{A}_{\Omega})$ have been defined in terms of reduced density matrices, see [14, 38] and [58, Sec. 1.4.2]. This is formalized in the following diagram, which commutes by formula (19) in [54]. (Obvious restrictions of red_g are omitted.)



The commutative diagram shows that the fiber $\pi_{\mathfrak{g}}^{-1}(\sigma)$ over $\sigma \in \mathcal{D}(\mathcal{R}_{\mathfrak{g}})$ is the set of all $\rho \in \mathcal{D}(\mathcal{A}_{\Omega})$ such that $\operatorname{red}_{\mathfrak{g}}(\rho) = \operatorname{red}_{\mathfrak{g}}(\sigma)$. Thm. 9.2 proves that the pullback $\Xi_{\tau} := \Psi_{\tau} \circ \operatorname{red}_{\mathfrak{g}}^{-1}$ of the inference map Ψ_{τ} under $\operatorname{red}_{\mathfrak{g}}^{-1}$ is continuous at $(\rho_{\nu})_{\nu \in \mathfrak{g}} \in \operatorname{red}_{\mathfrak{g}}[\mathcal{D}(\mathcal{A}_{\Omega})]$ if and only if $\operatorname{red}_{\mathfrak{g}}$ is open at $\Xi_{\tau}[(\rho_{\nu})_{\nu \in \mathfrak{g}}] \in \mathcal{D}(\mathcal{A}_{\Omega})$.

A discontinuity of Ξ_{τ} (with uniform prior τ) was reported for N = 3qubits and $\mathfrak{g} := \{\{1,2\},\{2,3\},\{3,1\}\}$ at $(\rho_{\nu})_{\nu \in \mathfrak{g}} \in \operatorname{red}_{\mathfrak{g}}[\mathcal{D}(\mathcal{A}_{\Omega})]$, where

$$\rho_{\nu} := \frac{1}{2} (|00\rangle\langle 00| + |11\rangle\langle 11|), \quad \nu \in \mathfrak{g}.$$

This result by Chen et al. [14, Ex. 4] has an interesting interpretation. The map red_g being open⁵ at $\rho \in \mathcal{D}(\mathcal{A}_{\Omega})$ means that any sufficiently small change of red_g(ρ) within red_g[$\mathcal{D}(\mathcal{A}_{\Omega})$] is matched by an arbitrarily small change of ρ within $\mathcal{D}(\mathcal{A}_{\Omega})$. Conversely, if the openness fails, then there are arbitrarily small changes of red_g(ρ) that are only matched by changes of ρ beyond some positive threshold (in the metric sense). Loosely speaking, a small change of a subsystem abruptly changes the entire system. Such behavior is associated with phase transitions [14]. It motivates every attempt to study the openness problem of $\pi_{\mathfrak{g}}$. This should be done by tuning the interaction pattern \mathfrak{g} to a concrete system. Whether enclosing $\mathcal{R}_{\mathfrak{g}}$ into a *-subalgebra of \mathcal{A}_{Ω} could simplify the problem, as suggested in Rem. 8.2, is not yet clarified.

Focusing on the continuity of correlations measures, we begin with theory.

Example 9.5 (Entropy distance I). Let $\tau := \mathbb{1}_n/n$ and let $\mathcal{E}(\mathcal{R}) := \mathcal{E}_{\tau}(\mathcal{R})$ be an exponential family, see Ex. 9.3. The *entropy distance* from $\mathcal{E}(\mathcal{R})$ is

 $d: \mathcal{D}(\mathcal{M}_n) \to \mathbb{R}, \quad \rho_1 \mapsto \inf_{\rho_2 \in \mathcal{E}(\mathcal{R})} S(\rho_1, \rho_2).$ (9.6)

⁵The openness of $\operatorname{red}_{\mathfrak{g}}$ at $\rho \in \mathcal{D}(\mathcal{A}_{\Omega})$ is a priori weaker than the continuity of Ξ_{τ} at $\operatorname{red}_{\mathfrak{g}}(\rho)$. The continuity means that any sufficiently small change of $\operatorname{red}_{\mathfrak{g}}(\rho)$ can be matched by an arbitrarily small change of ρ inside the range of Ξ_{τ} and not just anywhere within $\mathcal{D}(\mathcal{A}_{\Omega})$. Somewhat surprisingly, the two propositions are equivalent by Thm. 9.2.

Using the von Neumann entropy $S = \phi_{\tau} + \log(n)$, we define

$$\Psi: \mathcal{D}(\mathcal{R}) \to \mathcal{D}(\mathcal{M}_n), \quad \sigma \mapsto \operatorname{argmax}_{\rho \in \pi^{-1}(\sigma)} S(\rho)$$

Clearly, the map Ψ equals the inference map Ψ_{τ} defined earlier in Ex. 9.1. The entropy distance can be written as the difference

$$d(\rho) = S(\Psi \circ \pi(\rho)) - S(\rho), \quad \rho \in \mathcal{D}(\mathcal{M}_n)$$
(9.7)

between the value of the von Neumann entropy at ρ and its maximal value on the fiber of $\pi : \mathcal{D}(M_n) \to \mathcal{D}(\mathcal{R})$ that contains ρ , see p. 1288 in [52].

Formula (9.7) suggests studying the *rI-projection* [19]

$$\Pi: \mathcal{D}(\mathcal{M}_n) \to \mathcal{D}(\mathcal{M}_n), \quad \Pi = \Psi \circ \pi.$$

By (9.3), the range of Π is the rI-closure of $\mathcal{E}(\mathcal{R})$. We also consider the restriction $d|_{\mathcal{D}(\mathcal{A})}$ of d and the following restriction of Π .

$$\Pi|_{\mathcal{D}(\mathcal{A})}: \mathcal{D}(\mathcal{A}) \to \mathcal{D}(\mathcal{A}), \quad \rho \mapsto \Pi(\rho)$$

Here, \mathcal{A} is real *-subalgebra of M_n that includes \mathcal{R} . The map $\Pi|_{\mathcal{D}(\mathcal{A})}$ is well defined because $\Psi(\mathcal{D}(\mathcal{R}))$ is included in $\mathcal{D}(\mathcal{A})$, as observed in Ex. 9.3 a).

Example 9.6 (Entropy distance II). The exponential family $\mathcal{E}(\mathcal{R}_g)$ of local Hamiltonians (Ex. 9.4) is interesting because the entropy distance from it quantifies many-body correlations. Amari [4] and Ay [6] studied this type of correlation measure in probability theory. Linden et al. [33] introduced it as a difference of von Neumann entropies like formula (9.7), see also [58, Sec. 1.4.2]. And Zhou [59] proved the equivalence of the two representations (9.6) and (9.7) for density matrices of maximal rank n, see also [38]; the equivalence is true without rank restrictions as presented in Ex. 9.5.

We quote from [52, Lemma 5.15 and 4.5] under the assumptions of Ex. 9.5.

Lemma 9.7. Le \mathcal{A} be real *-subalgebra of M_n that includes \mathcal{R} .

- a) For every $\rho \in \mathcal{D}(\mathcal{A})$, the map $\Pi|_{\mathcal{D}(\mathcal{A})}$ is continuous at ρ if and only if $d|_{\mathcal{D}(\mathcal{A})}$ is continuous at ρ .
- b) For every $\sigma \in \mathcal{D}(\mathcal{R})$, the inference map Ψ is continuous at σ if and only if d is continuous at every point in the fiber $\pi^{-1}(\sigma)$.

With reference to Ex. 9.5, we simplify the continuity problem of the entropy distance (and of other invariant maps) using *-subalgebras.

Remark 9.8 (Continuity of the rI-projection). Let \mathcal{A} be a *-subalgebra of M_n such that $\mathcal{R} \subset \mathcal{A}$. As in Rem. 8.2, the map π factors into $\pi = \pi_{\mathcal{R}} \circ \pi_{\mathcal{A}}$. Then, for every $\rho \in \mathcal{D}(M_n)$ the rI-projection Π is continuous at ρ if and only if $\Pi|_{\mathcal{D}(\mathcal{A})}$ is continuous at $\pi_{\mathcal{A}}(\rho)$.

This assertions needs a justification. By its definition, the map $\Pi = \Psi \circ \pi$ is invariant under $\pi_{\mathcal{A}}$ in the sense that $\Pi = \Pi \circ \pi_{\mathcal{A}}$ holds. This shows $\Pi = \Pi|_{\mathcal{D}(\mathcal{A})} \circ \pi_{\mathcal{A}}$. The claim then follows from Lemma 8.1 b), by letting $\mathcal{R}_1 := M_n$ and $\mathcal{R}_2 := \mathcal{A}$, and by taking $\Pi|_{\mathcal{D}(\mathcal{A})}$ as the map $f : \mathcal{D}(\mathcal{R}_2) \to T$. The assumptions of the lemma are met since $\pi_{\mathcal{A}}$ is open by Thm. 1.2.

Example 9.9 (Entropy distance III). Rem. 9.8 helps solve the continuity problem of the entropy distance from $\mathcal{E}(\mathcal{R})$ for $\mathcal{R} := \operatorname{span}_{\mathbb{R}}(\mathbb{1}_3, X \oplus 1, Z \oplus 0)$,

based on a solution for $d|_{\mathcal{A}(\mathcal{D})}$ in the algebra $\mathcal{A} := M_2(\mathbb{R}) \oplus M_1(\mathbb{R})$ that was obtained using asymptotic curvature estimates [52].

Real *-subalgebras are excluded from Rem. 9.8 but the conclusion is still true, as Ex. 7.2 can replace Thm. 1.2 in this remark.

We recall from Ex. 3.6 that the generatrix $\mathcal{G} = [|+\rangle\langle +| \oplus 0, 0 \oplus 1]$ of the cone $\mathcal{D}(\mathcal{A})$ is the fiber of the orthogonal projection $\pi_{\mathcal{R}} : \mathcal{D}(\mathcal{A}) \to \mathcal{D}(\mathcal{R})$ over $M := \frac{1}{2}(|+\rangle\langle +| \oplus 1)$. Thm. 5.18 in [52] shows that $d|_{\mathcal{D}(\mathcal{A})}$ is discontinuous at every point in the half-open segment

$$\mathcal{G}_d := \left[|+\rangle \langle +| \oplus 0, M \right] = \left\{ (1-\lambda) |+\rangle \langle +| \oplus \lambda \colon \lambda \in [0, \frac{1}{2}) \right\}$$

and continuous at every point in the complement. (Unitary similarity with respect to $U \oplus 1$ matches \mathcal{R} with the problem considered in [52], as the similarity with respect to $U := \exp(i\frac{\pi}{3\sqrt{3}}(X+Y+Z))$ permutes the Pauli matrices X, Y, Z cyclicly.) Lemma 9.7 then shows that the restricted rIprojection $\Pi|_{\mathcal{D}(\mathcal{A})}$ is discontinuous at every point in \mathcal{G}_d and continuous at every point in the complement. Now Rem. 9.8 proves that the rI-projection Π is discontinuous at every point in $\pi_{\mathcal{A}}^{-1}(\mathcal{G}_d)$ and continuous at every point in the complement. A second application of Lemma 9.7 finally shows that the entropy distance d is discontinuous at every point in $\pi_{\mathcal{A}}^{-1}(\mathcal{G}_d)$ and continuous at every point in the complement.

The points in the set $\pi_{\mathcal{A}}^{-1}(\mathcal{G}_d)$ are explicitly described in Ex. 8.3 b). The result is consistent with the inference Ψ being discontinuous at M and continuous at every point different from M (see Ex. 9.3 c)) as per Lemma 9.7 b).

Acknowledgments. The author is grateful to several people for discussions related to this work, among them are Achim Clausing, Chi-Kwong Li, Tim Netzer, Ivan Todorov, and Ilya M. Spitkovsky. The author gave a lecture on the results of this paper in the special session "Numerical Ranges" of the conference IWOTA 2024 in Canterbury, UK, and wishes to thank the organizers for this opportunity.

References

- [1] E. M. Alfsen, Compact Convex Sets and Boundary Integrals, Berlin: Springer, 1971.
- [2] E. M. Alfsen and F. W. Shultz, Geometry of State Spaces of Operator Algebras, Boston: Birkhäuser, 2003.
- [3] S. A. Ali, C. Cafaro, A. Giffin, C. Lupo, and S. Mancini, On a differential geometric viewpoint of Jaynes' MaxEnt method and its quantum extension, AIP Conf Proc 1443, 120–128 (2012).
- [4] S.-I. Amari, Information geometry on hierarchy of probability distributions, IEEE Transactions on Information Theory 47:5, 1701–1711 (2001).
- [5] W. Arveson, The noncommutative Choquet boundary III: Operator systems in matrix algebras, Mathematica Scandinavica 106:2, 196–210 (2010).
- [6] N. Ay, An information-geometric approach to a theory of pragmatic structuring, Annals of Probability 30:1, 416–436 (2002).
- [7] I. Bengtsson and K. Życzkowski, Geometry of Quantum States: An Introduction to Quantum Entanglement, 2nd ed., Cambridge: Cambridge University Press, 2017.
- [8] A. Böttcher and I. M. Spitkovsky, A gentle guide to the basics of two projections theory, Linear Algebra and its Applications 432:6, 1412–1459 (2010).
- [9] F.F. Bonsall and J. Duncan, Numerical Ranges of Operators on Normed Spaces and of Elements of Normed Algebras, London: Cambridge University Press, 1971.

- [10] O. Bratteli and D. W. Robinson, Operator Algebras and Quantum Statistical Mechanics 1, Berlin: Springer, 1987.
- [11] A. Buckley, The mathematics of quantum information theory: Geometry of quantum states, Master thesis, Lugano, Switzerland, 2021. [Retrieved 2025-01-27]
- [12] C. Davis, The Toeplitz-Hausdorff theorem explained, Canadian Mathematical Bulletin 14:2, 245–246 (1971).
- [13] A. Caticha, *Entropic Physics*, 2022. [Retrieved 2025-01-27]
- [14] J. Chen, Z. Ji, C.-K. Li, Y.-T. Poon, Y. Shen, N. Yu, B. Zeng, and D. Zhou, Discontinuity of maximum entropy inference and quantum phase transitions, New Journal of Physics 17:8, 083019 (2015).
- [15] M.-D. Choi and E. G. Effros, *Injectivity and operator spaces*, Journal of Functional Analysis 24:2, 156–209 (1977).
- [16] A. Clausing, Retractions and open mappings between convex sets, Math Z. 160:3, 263–274 (1978).
- [17] A. Clausing and S. Papadopoulou, Stable convex sets and extremal operators, Math. Ann. 231:3, 193–203 (1978).
- [18] D. Corey, C. R. Johnson, R. Kirk, B. Lins, and I. Spitkovsky, Continuity properties of vectors realizing points in the classical field of values, Linear and Multilinear Algebra 61:10, 1329–1338 (2013).
- [19] I. Csiszár and F. Matúš, Information projections revisited, IEEE Transactions on Information Theory 49:6, 1474–1490 (2003).
- [20] G. Debs, Some general methods for constructing stable convex sets, Mathematische Annalen 241:2, 97–105 (1979).
- [21] D. R. Farenick, Algebras of Linear Transformations, New York: Springer, 2001.
- [22] D. R. Farenick, Pure matrix states on operator systems, Linear Algebra and its Applications 393, 149–173 (2004).
- [23] P. R. Halmos, Finite-Dimensional Vector Spaces, New York: Springer, 1974.
- [24] P. Harremoës, Entropy on spin factors, in: N. Ay, et al. (eds.), Information Geometry and Its Applications, Cham: Springer, 2018.
- [25] R. A. Horn and F. Zhang, *Basic Properties of the Schur Complement*, in: F. Zhang (ed.), The Schur Complement and Its Applications, New York: Springer, 2005.
- [26] F. Huber and O. Gühne, Characterizing ground and thermal states of few-body Hamiltonians, Physical Review Letters 117:1, 010403 (2016).
- [27] J. L. Kelley, *General Topology*, New York: Springer, 1975.
- [28] J. Faraut and A. Korányi, Analysis on Symmetric Cones, Oxford: Oxford University Press, 1994.
- [29] T. Leake, B. Lins, and I. M. Spitkovsky, Pre-images of boundary points of the numerical range, Operators and Matrices 8:3, 699–724 (2014).
- [30] T. Leake, B. Lins, and I. M. Spitkovsky, *Inverse continuity on the boundary of the numerical range*, Linear and Multilinear Algebra 62:10, 1335–1345 (2014).
- [31] T. Leake, B. Lins, and I. M. Spitkovsky, Corrections and additions to 'Inverse continuity on the boundary of the numerical range', Linear and Multilinear Algebra 64:1, 100–104 (2016).
- [32] B. Li, Real Operator Algebras, World Scientific, 2003.
- [33] N. Linden, S. Popescu, and W. Wootters, Almost every pure state of three qubits is completely determined by its two-particle reduced density matrices, Physical Review Letters 89:20, 207901 (2002).
- [34] B. Lins and P. Parihar, Continuous selections of the inverse numerical range map, Linear and Multilinear Algebra 64:1, 87–99 (2016).
- [35] B. Lins and I. M. Spitkovsky, Inverse continuity of the numerical range map for Hilbert space operators, Operators and Matrices 14:1, 77–90 (2020).
- [36] M. Lostaglio, D. Jennings, and T. Rudolph, Thermodynamic resource theories, non-commutativity and maximum entropy principles, New Journal of Physics 19:4, 043008 (2017).
- [37] B. Maier and T. Netzer, A note on the Carathéodory number of the joint numerical range. arXiv:2409.04444 [math.FA].

- [38] S. Niekamp, T. Galla, M. Kleinmann, and O. Gühne, Computing complexity measures for quantum states based on exponential families, Journal of Physics A: Mathematical and Theoretical 46:12, 125301 (2013).
- [39] R. C. O'Brien, On the openness of the barycentre map, Math. Ann. 223:3, 207–212 (1976).
- [40] S. Papadopoulou, On the geometry of stable compact convex sets, Math. Ann. 229:3, 193–200 (1977).
- [41] S. Papadopoulou, Stabile konvexe Mengen, Jahresbericht der Deutschen Mathematiker-Vereinigung 84:2, 92–106 (1982).
 [Retrieved 2025-01-27]
- [42] V.I. Paulsen, Completely Bounded Maps and Operator Algebras, New York: Cambridge University Press, 2002.
- [43] D. Pavlov, B. Sturmfels, and S. Telen, *Gibbs manifolds*, Info Geo 7:S2, 691–717 (2024).
- [44] D. Plaumann, R. Sinn, and S. Weis, Kippenhahn's theorem for joint numerical ranges and quantum states, SIAM J. Appl. Algebra Geometry 5:1, 86–113 (2021).
- [45] R.T. Rockafellar, Convex Analysis, Princeton: Princeton University Press, 1970.
- [46] L. Rodman, I. M. Spitkovsky, A. Szkoła, and S. Weis, Continuity of the maximum-entropy inference: convex geometry and numerical ranges approach, Journal of Mathematical Physics 57:1, 015204 (2016).
- [47] R. Schneider, Convex Bodies: The Brunn-Minkowski Theory, 2nd ed., New York: Cambridge University Press, 2014.
- [48] M. E. Shirokov, Stability of convex sets and applications, Izvestiya: Mathematics 76:4, 840–856 (2012).
- [49] I. M. Spitkovsky and S. Weis, Pre-images of extreme points of the numerical range, and applications, Operators and Matrices 10:4, 1043–1058 (2016).
- [50] I. M. Spitkovsky and S. Weis, Signatures of quantum phase transitions from the boundary of the numerical range, Journal of Mathematical Physics 59:12, 121901 (2018).
- [51] J. Vesterstrøm, On open maps, compact convex sets, and operator algebras, Journal of the London Mathematical Society s2-6:2, 289-297 (1973).
- [52] S. Weis, Continuity of the maximum-entropy inference, Communications in Mathematical Physics 330:3, 1263–1292 (2014).
- [53] S. Weis, Maximum-entropy inference and inverse continuity of the numerical range, Reports on Mathematical Physics 77:2, 251–263 (2016).
- [54] S. Weis and J. Gouveia, The face lattice of the set of reduced density matrices and its coatoms, Info Geo 6:1, 293–326 (2023).
- [55] S. Weis and A. Knauf, Entropy distance: New quantum phenomena, Journal of Mathematical Physics 53:10, 102206 (2012).
- [56] A. Winter, Tight uniform continuity bounds for quantum entropies: Conditional entropy, relative entropy distance and energy constraints, Communications in Mathematical Physics 347:1, 291–313 (2016).
- [57] H. Witsenhausen, A minimax control problem for sampled linear systems, IEEE Trans. Automat. Contr. 13:1, 5–21 (1968).
- [58] B. Zeng, X. Chen, D.-L. Zhou, and X.-G. Wen, Quantum Information Meets Quantum Matter, New York: Springer, 2019.
- [59] D. Zhou, Irreducible multiparty correlations can be created by local operations, Physical Review A 80:2, 022113 (2009).

Stephan Weis Czech Technical University in Prague Faculty of Electrical Engineering Karlovo náměstí 13 12000, Prague 2 Czech Republic e-mail maths@weis-stephan.de

26