

TREE PROPERTIES AT SUCCESSORS OF SINGULARS OF MANY COFINALITIES

WILLIAM ADKISSON

ABSTRACT. From many supercompact cardinals, we show that it is consistent for the tree property to hold at many small successors of singular cardinals, each with a different cofinality. In particular, we construct a model in which the tree property holds at $\aleph_{\omega+\omega+1}$ and at \aleph_{ω_n+1} for all $0 < n < \omega$. We show that this can be done for the strong tree property as well, and extend the technique to large uncountable sequences of desired cofinalities.

1. INTRODUCTION

The tree property is a generalization of König's Lemma to uncountable cardinals, and is closely related to large cardinals. In particular, the tree property holds at an inaccessible cardinal κ if and only if κ is weakly compact. The tree property can consistently hold at small cardinals, but it retains some large cardinal strength: Mitchell and Silver [11] showed that the tree property at \aleph_2 is equiconsistent with a weakly compact.

In the 70s, Magidor asked whether it was consistent for every regular cardinal greater than \aleph_1 to have the tree property. Since then, set theorists have constructed models where the tree property holds at more and more successive cardinals; see for instance [1], [4], [10], and [12].

More recently, there has been a resurgence of interest in strengthenings of the tree property that are linked with more powerful large cardinals. The strong tree property is a generalization of the tree property that holds at an inaccessible cardinal κ if and only if κ is strongly compact [9]. Like the tree property, this property can consistently hold at small cardinals, and is viewed as strong evidence for the presence of a strongly compact cardinal.

Magidor's question can be extended to these stronger properties, and in the past decade many results about the tree property have been generalized. See for instance [5], [14], [8], and [2].

If the tree property and its strengthenings are to hold at every regular cardinal, they will need to hold at many successors of singular cardinals. In particular, they will need to hold at successors of singulars of many different cofinalities. It is easy to arrange this situation at the level of large cardinals: Magidor and Shelah [10] proved that the successor of a limit of supercompacts always has the tree property, so a long enough sequence of supercompacts will give rise to many such limits of different cofinalities. Obtaining this result at small cardinals is much more difficult. In [2], the author forced the strong tree property at finitely many successors of singulars with different cofinalities simultaneously. Unfortunately, the techniques used in that paper are fundamentally restricted to finite sequences of cofinalities, and cannot be easily generalized.

In this paper we present a different technique that can be used to obtain the tree property and the strong tree property at successors of singulars of infinitely many cofinalities simultaneously. In particular, we will obtain the strong tree property at $\aleph_{\omega+\omega+1}$ along with \aleph_{ω_n+1} for all $0 < n < \omega$.

In Section 2, we define the strong tree property and list a number of standard lemmas that we will use. In section 3 we will obtain the tree property at $\aleph_{\omega+\omega+1}$ and each \aleph_{ω_n+1} . In section 4, we prove the same result for the strong tree property. Section 5 extends these results to uncountably many successors of singulars of different cofinalities.

2. PRELIMINARIES AND BRANCH LEMMAS

First, let us define the tree property and strong tree property.

Definition 2.1. A tree T is a κ -tree if it has height κ and levels of size $< \kappa$. A cardinal κ has the *tree property* if every κ -tree has a cofinal branch.

The strong tree property is concerned with a more general class of objects, called $\mathcal{P}_\kappa(\lambda)$ -lists.

Definition 2.2. A sequence $d = \langle d_z \mid z \in \mathcal{P}_\kappa(\lambda) \rangle$ is a $\mathcal{P}_\kappa(\lambda)$ -list if $d_z \subseteq z$ for all $z \in \mathcal{P}_\kappa(\lambda)$.

Definition 2.3. The z -th level of a $\mathcal{P}_\kappa(\lambda)$ -list, written L_z , is defined as

$$L_z := \{d_y \cap z \mid y \supseteq z\}.$$

A $\mathcal{P}_\kappa(\lambda)$ -list is *thin* if $|L_z| < \kappa$ for all $z \in \mathcal{P}_\kappa(\lambda)$.

Definition 2.4. A subset $b \subseteq \lambda$ is a *cofinal branch* through a $\mathcal{P}_\kappa(\lambda)$ -list d if $b \cap z \in L_z$ for all $z \in \mathcal{P}_\kappa(\lambda)$.

Definition 2.5. The *strong tree property* holds at κ if for all $\lambda > \kappa$, every thin $\mathcal{P}_\kappa(\lambda)$ -list has a cofinal branch.

Fact 2.6. [6, Lemma 3.4] Suppose $\lambda < \lambda'$. If every thin $\mathcal{P}_\kappa(\lambda')$ -list has a cofinal branch, then so does every thin $\mathcal{P}_\kappa(\lambda)$ -list.

In the specific case where $\kappa = \lambda$, it is often convenient to restrict to elements of κ rather than of $\mathcal{P}_\kappa(\kappa)$.

Definition 2.7. Let κ be a regular cardinal. A sequence $d = \langle d_\alpha \mid \alpha < \kappa \rangle$ is a κ -list if $d_\alpha \subseteq \alpha$ for all $\alpha < \kappa$. The α -th level, denoted L_α , is the set $\{d_\beta \cap \alpha \mid \alpha < \beta < \kappa\}$. A κ -list is *thin* if every level has size $< \kappa$. A set $b \subseteq \kappa$ is an *cofinal branch* through a κ -list d if $b \cap \alpha \in L_\alpha$ for all $\alpha < \kappa$.

Since κ is unbounded in $\mathcal{P}_\kappa(\kappa)$, we have the following easy facts:

Fact 2.8. If every thin κ -list has a cofinal branch, then every thin $\mathcal{P}_\kappa(\kappa)$ -list has a cofinal branch.

Fact 2.9. Let $d = \langle d_z \mid z \in \mathcal{P}_\kappa(\kappa) \rangle$ be a thin $\mathcal{P}_\kappa(\kappa)$ -list. Let d^* be the restriction of d to elements indexed by ordinals; that is, $d^* = \langle d_\alpha \mid \alpha < \kappa \rangle$. Then d^* is a thin κ -list, and if d^* has a cofinal branch, then so does d .

We will make heavy use of the following general lemma for obtaining cofinal branches through lists at successors of singulars. It follows immediately from [2, Theorem 3.14] and is a slight weakening of it, requiring stronger assumptions and proving a weaker result; we use this form because the hypotheses are straightforward to verify and the conclusion is precisely what is required for our argument.

Lemma 2.10. *Let τ be a regular cardinal. Let $\langle \kappa_\rho \mid \rho < \tau \rangle$ be a continuous increasing sequence of regular cardinals above τ , with supremum ν . Let $I \subseteq \kappa_0$, and fix $\rho' < \tau$. For each $\mu \in I$, let \mathbb{L}_μ be a forcing poset of size $\leq \kappa_{\rho'}$. Suppose the following hold:*

- (1) *For each $\mu \in I$, \mathbb{L}_μ is the product of forcings \mathbb{P}_μ and \mathbb{Q}_μ , where $|\mathbb{P}_\mu| < \mu^+$ and \mathbb{Q}_μ is μ^+ -closed.*
- (2) *κ_0 is ν^+ -supercompact, with a normal measure U_0 on $\mathcal{P}_{\kappa_0}(\nu^+)$ and a corresponding embedding i such that $\nu \in i(I)$.*
- (3) *For all ordinals $\rho < \tau$ there is a generic ν^+ -supercompactness embedding $j_{\rho+2}$ with domain V and critical point $\kappa_{\rho+2}$, added by a poset \mathbb{F} whose full support power \mathbb{F}^{κ_ρ} is $< \kappa_\rho$ -distributive in V .*

Then there exists $\mu \in I$ such that in the extension of V by \mathbb{L}_μ , every thin ν^+ -list has a cofinal branch.

In particular, the conclusion of this lemma implies the tree property at ν^+ .

A key tool used to work with cofinal branches through trees and thin lists is the (thin) κ -approximation property.

Definition 2.11. Let κ be regular, λ be an ordinal, and \mathbb{P} be a forcing notion in a model V . A \mathbb{P} -name \dot{b} for a subset of λ is κ -approximated by \mathbb{P} over V if for all $z \in (\mathcal{P}_\kappa(\lambda))^V$, $\Vdash_{\mathbb{P}} \dot{b} \cap z \in V$.

A \mathbb{P} -name \dot{b} for a subset of λ is *thinly κ -approximated by \mathbb{P} over V* if it is κ -approximated by \mathbb{P} over V , and furthermore for every $z \in (\mathcal{P}_\kappa(\lambda))^V$, $|\{x \in V \mid \exists p \in \mathbb{P} \Vdash_{\mathbb{P}} x = \dot{b} \cap z\}| < \kappa$.

Definition 2.12. Let κ be regular. A forcing \mathbb{P} has the κ -approximation property over a model V if for every ordinal λ and \mathbb{P} -name \dot{b} for a subset of λ , if \dot{b} is κ -approximated by \mathbb{P} over V , then $\Vdash_{\mathbb{P}} \dot{b} \in V$.

A forcing \mathbb{P} has the *thin κ -approximation property* over V if for every ordinal λ and every \mathbb{P} -name \dot{b} for a subset of λ , if \dot{b} is thinly κ -approximated by \mathbb{P} over V , then $\Vdash_{\mathbb{P}} \dot{b} \in V$.

The κ -approximation property implies the thin κ -approximation property.

A cofinal branch through a thin $\mathcal{P}_\kappa(\lambda)$ list is always thinly approximated over V , by any forcing. A similar argument shows that branches through a κ -tree are thinly approximated by any forcing; see [13, Section 2] for details.

Lemma 2.13. *Let d be a thin $\mathcal{P}_\kappa(\lambda)$ list in V , and let \mathbb{P} be a notion of forcing over V . Suppose \dot{b} is a \mathbb{P} -name for a cofinal branch through this list. Then \dot{b} is thinly κ -approximated by \mathbb{P} over V .*

Proof. Let \dot{b} be a name for an cofinal branch, and let $z \in (\mathcal{P}_\kappa(\lambda))^V$. Since \dot{b} is forced to be cofinal, it must meet every level; that is, the empty condition forces that $\dot{b} \cap z \in L_z$, for all $z \in \mathcal{P}_\kappa(\lambda)$. Since L_z is in V , $\dot{b} \cap z$ is likewise forced to be in V . Since the list is thin, $|L_z| < \kappa$, and so there are fewer than κ possibilities for $\dot{b} \cap z$. \square

Many standard branch lemmas for trees can be generalized to use the (thin) approximation property. We will make use of the following:

Lemma 2.14. [13, Lemma 2.4] *Let κ be a regular cardinal. Suppose that \mathbb{P} is a forcing poset such that $\mathbb{P} \times \mathbb{P}$ is κ -cc. Then \mathbb{P} has the κ -approximation property.*

Corollary 2.15. *Let κ be a regular cardinal. Suppose that \mathbb{P} is a forcing poset with $|\mathbb{P}| < \kappa$. Then \mathbb{P} has the κ -approximation property.*

Lemma 2.16. [2, Lemma 2.20] *Suppose ν is a singular strong limit cardinal with cofinality τ . Let \mathbb{Q} be a μ^+ -closed forcing over a model V for some $\mu < \nu$ with $\tau \leq \mu$, and let $\mathbb{P} \in V$ be a forcing poset with $|\mathbb{P}| \leq \mu$. Then \mathbb{Q} has the ν^+ -approximation property in the generic extension of V by \mathbb{P} .*

Finally, our argument will make crucial use of the following standard absorption lemma.

Lemma 2.17. [3, Theorem 14.3] *Let κ be an inaccessible cardinal, and let $\delta < \kappa$ be regular. Let \mathbb{P} be a δ -closed forcing poset with $|\mathbb{P}| < \kappa$. Then there is a forcing projection from $\text{Col}(\delta, < \kappa)$ to \mathbb{P} whose quotient is $\text{Col}(\delta, < \kappa)$.*

3. THE TREE PROPERTY AT COUNTABLY MANY COFINALITIES

First we will show that the tree property can be forced to hold at $\aleph_{\omega+\omega+1}$ and at each \aleph_{ω_n+1} simultaneously for $0 < n < \omega$. To obtain the tree property at each \aleph_{ω_n+1} using Lemma 2.10, we need to choose a singular cardinal μ_n whose successor will become ω_{n+1} in the final model. The cofinality of μ_n needs to be the cardinal that will become ω_n in final model; that is, $\text{cf}(\mu_n) = \mu_{n-1}^+$, and μ_n will be collapsed to μ_{n-1}^+ . Once all of these collapses are complete, \aleph_ω will be the supremum of the sequence $\langle \mu_n^+ \mid n < \omega \rangle$. Since each μ_n^+ is the successor of a singular, there is no reason to expect the tree property to hold at $\aleph_{\omega+1}$ in this construction; we settle for the next best cardinal, $\aleph_{\omega+\omega+1}$.

The primary difficulty is in selecting the sequence $\langle \mu_i \mid i < \omega \rangle$. If we choose them one at a time, we run into difficulties. In particular, when selecting each μ_i we need complete knowledge of the poset (so all μ_j for $j \neq i$) to ensure that we will have the tree property at the appropriate cardinals. If we select a different μ_i for every possible tail of the sequence, we are unable to combine all of our choices at the limit stage, since we will never make a selection that does not depend on other information. Working within these constraints, we can obtain some partial results as in [2], but those techniques only work for finitely many cofinalities.

Our solution is to obtain the tree property in a larger poset that does not depend on any future choices. This poset will project down to our target model, regardless of the selections we make, and the quotient of this projection will not add branches through trees. This approach is loosely inspired by the arguments of Golshani and Hayut in [7], although the details of the construction are quite different.

Theorem 3.1. *Let $\langle \kappa_\alpha \mid \alpha < \kappa_0 \rangle$ be an continuous increasing sequence of cardinals with the following properties:*

- $\kappa_{\alpha+1} = \kappa_\alpha^+$ for all limit ordinals α
- κ_n is indestructibly supercompact for all $n < \omega$
- $\kappa_{\alpha+2}$ is indestructibly supercompact for all $\omega \leq \alpha < \kappa_0$.

Then there is a generic extension in which the tree property holds at $\aleph_{\omega+\omega+1}$ and at \aleph_{ω_n+1} for all $0 < n < \omega$.

Proof. Define the poset \mathbb{H} as follows:

$$\mathbb{H} := \left(\prod_{n < \omega} \text{Col}(\kappa_n, < \kappa_{n+1}) \right) \times \left(\prod_{\omega \leq \rho < \tau} \text{Col}(\kappa_{\rho+1}, < \kappa_{\rho+2}) \right).$$

Let H be generic for \mathbb{H} , and work in $V[H]$. Note that in $V[H]$, requirements (2) and (3) of Lemma 2.10 are satisfied.

Let I be the collection of all strictly increasing sequences $\langle \mu_i \mid i < \omega \rangle$, where $\mu_i < \kappa_0$ is a singular cardinal of cofinality μ_{i-1}^+ . For each sequence $s = \langle \mu_i \mid i < \omega \rangle \in I$, define

$$\mathbb{L}_s := \text{Col}(\omega, \mu_0) \times \left(\prod_{n < \omega} \text{Col}(\mu_n^+, \mu_{n+1}) \right) \times \text{Col} \left(\left(\sup_{n < \omega} \mu_n \right)^+, < \kappa_0 \right).$$

Given such a sequence s , for all i with $0 < i < \omega$, define $\nu_i(s) := \sup_{\rho < \mu_{i-1}^+} \kappa_\rho$. Let $\nu_0(s) = \kappa_0^{+\omega}$. In the extension of $V[H]$ by \mathbb{L}_s , $\nu_i(s)$ will become $\aleph_{\omega+\omega_i}$. Note that $\nu_i(s)$ only depends on the initial segment $\langle \nu_n \mid n < i \rangle$; when these values are fixed, as they will be throughout our argument, we will usually omit the parameter s .

Our goal is to inductively build a sequence s such that in the extension of $V[H]$ by a generic for \mathbb{L}_s , the tree property will hold at every $\nu_i^+(s)$. We will start at μ_0 and work our way up.

Lemma 3.2. *There is $\mu_0^* < \kappa_0$ such that for all $s = \langle \mu_i \mid i < \omega \rangle \in I$ with $\mu_0 = \mu_0^*$, the tree property holds at $\nu_0^+(s)$ in $V[H][\mathbb{L}_s]$.*

Proof. For all $\mu_0 < \kappa_0$, define

$$\mathbb{L}_0^*(\mu_0) = \text{Col}(\omega, \mu_0) \times \text{Col}(\mu_0^+, < \kappa_0) \times \left(\prod_{\mu_0 < \alpha < \kappa_0} \text{Col}(\alpha^+, < \kappa_0) \right),$$

where the final product has μ_0^{++} -support. Let $L_0^*(\mu_0)$ be generic for $\mathbb{L}_0^*(\mu_0)$.

Let us examine the properties of this poset. The first component, $\text{Col}(\omega, \mu_0)$ has size $< \mu_0^+$, while the remainder of the forcing is μ_0^+ -closed. The full poset has size $< \kappa_0^{++} = \kappa_2$. Thus $\mathbb{L}_0^*(\mu_0)$ satisfies the first condition for Lemma 2.10.

It follows that there is some $\mu_0^* < \kappa_0$ such that in $V[H][L_0^*(\mu_0^*)]$, the tree property holds at ν_0^+ .

Now, let $s = \langle \mu_i \mid i < \omega \rangle \in I$ be a sequence such that $\mu_0 = \mu_0^*$, and let \dot{T} be a \mathbb{L}_s -name for a ν_0^+ -tree with no branch. We claim that $\mathbb{L}_0^*(\mu_0^*)$ projects to \mathbb{L}_s . Clearly $\text{Col}(\omega, \mu_0^*)$ projects to itself via the identity. By Lemma 2.17, since $\prod_{n < \omega} \text{Col}(\mu_n^+, \mu_{n+1})$ is $(\mu_0^*)^+$ -closed and has size $< \kappa_0$, it is absorbed by $\text{Col}((\mu_0^*)^+, < \kappa_0)$. Furthermore, the quotient is isomorphic to $\text{Col}((\mu_0^*)^+, < \kappa_0)$. Finally, since $\sup_{i < \omega} (\mu_i)$ is in the interval (μ_0, κ_0) , $\prod_{\mu_0 < \alpha < \kappa_0} \text{Col}(\alpha^+, < \kappa_0)$ projects to $\text{Col}(\left(\sup_{i < \omega} \mu_i\right)^+, < \kappa_0)$. The quotient of this projection is the full product, without whichever coordinate happened to correspond to the supremum of the μ_i 's.

Let L_s be generic for \mathbb{L}_s over $V[H]$, and let $L^*(\mu_0^*)$ be a generic for $\mathbb{L}_0^*(\mu_0^*)$ over $V[H]$ projecting to L_s . Let T be the interpretation of \dot{T} using the generic L_s . Since

T is in $V[H][L_s]$, and $\mathbb{L}_0^*(\mu_0^*)$ projects to \mathbb{L}_s , T must be in $V[H][L_0^*(\mu_0^*)]$. Being a tree is upwards absolute, and ν_0^+ is preserved by this quotient, so T remains a ν_0^+ -tree in $V[H][L_0^*(\mu_0^*)]$. Since the tree property holds in $V[H][L_0^*(\mu_0^*)]$, there is a branch b through T in $V[H][L_0^*(\mu_0^*)]$.

Note that the quotient of the projection from $\mathbb{L}_0^*(\mu_0^*)$ to \mathbb{L}_s has size $< \nu_0^+$, since ν_0 is a strong limit cardinal. By Corollary 2.15 this means that the quotient cannot add branches to ν_0^+ -trees, so b must be in $V[H][L_s]$.

We conclude that the tree property holds at ν_0^+ in $V[H][L_s]$ for all sequences $s \in I$ starting with μ_0^* . \square

Now we prove the inductive step.

Lemma 3.3. *Fix $\langle \mu_i^* \mid i < k \rangle$. There exists μ_k^* such that for all sequences $s = \langle \mu_i \mid i < \omega \rangle$ with $\mu_i = \mu_i^*$ for $i \leq k$, the tree property holds at $\nu_k^+(s)$ in $V[H][L_s]$.*

Proof. For all μ_i with $\mu_i^* < \mu_i < \kappa_0$, define

$$\mathbb{P}_k^*(\mu_k) := \text{Col}(\omega, \mu_0^*) \times \left(\prod_{n < k-1} \text{Col}((\mu_n^*)^+, \mu_{n+1}^*) \right) \times \text{Col}((\mu_{k-1}^*)^+, \mu_k)$$

and

$$\mathbb{Q}_k^*(\mu_k) := \text{Col}(\mu_k^+, < \kappa_0) \times \left(\prod_{\mu_k < \alpha < \kappa_0} \text{Col}(\alpha^+, < \kappa_0) \right),$$

where the product has μ_k^{++} -support. Finally, we define

$$\mathbb{L}_k^*(\mu_k) := \mathbb{P}_k^*(\mu_k) \times \mathbb{Q}_k^*(\mu_k).$$

Note that $|\mathbb{P}_k^*(\mu_k)| = \mu_k$, and $\mathbb{Q}_k^*(\mu_k)$ is μ_k^+ -closed; $|\mathbb{L}_k^*(\mu_k)| < \kappa_0^{++} = \kappa_2$. Thus it meets the first hypothesis of Lemma 2.10.

We apply Lemma 2.10 to conclude that there is some μ_k^* such that in the extension of $V[H]$ by a generic L for $\mathbb{L}_k^*(\mu_k^*)$, the tree property holds at ν_k^+ .

Now, let $s = \langle \mu_i \mid i < \omega \rangle$ with $\mu_n = \mu_n^*$ for $n \leq k$. In $V[H]$, let \hat{T} be a \mathbb{L}_s -name for a ν_k^+ -tree.

As before, we wish to show that $\mathbb{L}_k^*(\mu_k^*)$ projects to \mathbb{L}_s . Note that $\mathbb{L}_s = \mathbb{P}_s \times \mathbb{Q}_s$, where

$$\mathbb{P}_s = \text{Col}(\omega, \mu_0) \times \left(\prod_{n < k} \text{Col}(\mu_n^+, \mu_{n+1}) \right)$$

and

$$\mathbb{Q}_s = \left(\prod_{k \leq n < \omega} \text{Col}(\mu_n^+, \mu_{i+1}) \right) \times \text{Col} \left(\left(\sup_{i < \omega} \mu_i \right)^+, < \kappa_0 \right).$$

Note that since $\mu_n^* = \mu_n$ for $n \leq k$, $\mathbb{P}_s = \mathbb{P}_k^*(\mu_k^*)$.

By Lemma 2.17, noting that $\prod_{k \leq n < \omega} \text{Col}(\mu_n^+, \mu_{i+1})$ is $(\mu_k^*)^+$ -closed and has size less than κ_0 , the first component of $\mathbb{Q}_k^*(\mu_k^*)$ projects onto the first component of \mathbb{Q}_s . The second component of $\mathbb{Q}_k^*(\mu_k^*)$ projects onto the second component of \mathbb{Q}_s by restricting to the coordinate indexed by $\alpha = \sup_{i < \omega} \mu_i$.

Let L_s be generic for \mathbb{L}_s over $V[H]$, and let $L_k^*(\mu_k^*)$ be a generic for $\mathbb{L}_k^*(\mu_k^*)$ over $V[H]$ projecting to L_s . Let T be the interpretation of \hat{T} by this generic. Then $T \in V[H][L_s]$. Being a ν_k^+ -tree is upwards absolute, so T remains a tree in $V[H][L_k^*(\mu_k^*)]$. As before, since the tree property holds in this model, there must

be a cofinal branch through T in $V[H][L_k^*(\mu_k^*)]$. The quotient has size $< \nu_k^+$, so it cannot have added the branch. We conclude that T must have a cofinal branch in $V[H][L_s]$. \square

Let $s = \langle \mu_i^* \mid i < \omega \rangle$. Let L_s be generic for \mathbb{L}_s . Then for each $i < \omega$, by Lemma 3.2 (for $i = 0$) and Lemma 3.3, we conclude that in $V[H][L_s]$, the tree property holds at $\nu_i^+(s)$. In this extension, $\nu_i^+(s) = \aleph_{\omega+\omega_i+1}$, so we have the desired result. \square

4. THE STRONG TREE PROPERTY AT COUNTABLY MANY COFINALITIES.

In this section we extend our construction to force the strong tree property at $\aleph_{\omega+\omega+1}$ and at each \aleph_{ω_n+1} simultaneously for $n < \omega$. When we attempt to apply the techniques of the previous section, we run into a new obstacle: thin $\mathcal{P}_{\nu^+}(\lambda)$ -lists are not necessarily upwards absolute. When we build our larger model where the strong tree property will hold, we add new small subsets of λ . The list we wish to find a branch through may not be a $\mathcal{P}_\kappa(\lambda)$ -list in the larger model, so we can't apply the strong tree property to find the branch. To get around this, we use an auxiliary forcing that collapses λ to ν^+ . This lets us work with a single cardinal parameter, reducing the problem to finding a branch through a ν^+ -list. Since thin ν^+ -lists are upwards absolute, we can use the techniques of the previous section.

Theorem 4.1. *Let $\langle \kappa_\alpha \mid \alpha < \kappa_0 \rangle$ be an continuous increasing sequence of cardinals with the following properties:*

- $\kappa_{\alpha+1} = \kappa_\alpha^+$ for all limit ordinals α
- κ_n is indestructibly supercompact for all $n < \omega$
- $\kappa_{\alpha+2}$ is indestructibly supercompact for all $\omega \leq \alpha < \kappa_0$.

Then there is a generic extension in which the strong tree property holds at $\aleph_{\omega+\omega+1}$ and at \aleph_{ω_n+1} for all $0 < n < \omega$.

Proof. Define the poset \mathbb{H} as before:

$$\mathbb{H} := \left(\prod_{n < \omega} \text{Col}(\kappa_n, < \kappa_{n+1}) \right) \times \left(\prod_{\omega \leq \rho < \kappa_0} \text{Col}(\kappa_{\rho+1}, < \kappa_{\rho+2}) \right).$$

Let H be generic for \mathbb{H} . As before, $V[H]$ meets the requirements given in Lemma 2.10.

Let I be the collection of all strictly increasing sequences $\langle \mu_i \mid i < \omega \rangle$, where $\mu_i < \kappa_0$ is a singular cardinal of cofinality μ_{i-1}^+ . For each sequence $s = \langle \mu_i \mid i < \omega \rangle \in I$, define

$$\mathbb{L}_s := \text{Col}(\omega, \mu_0) \times \left(\prod_{n < \omega} \text{Col}(\mu_n^+, \mu_{n+1}) \right) \times \text{Col} \left(\left(\sup_{n < \omega} \mu_n \right)^+, < \kappa_0 \right).$$

Given such a sequence s , for all i with $0 < i < \omega$, define $\nu_i(s) := \sup_{\rho < \mu_{i-1}^+} \kappa_\rho$. As before, we will freely omit the parameter s . Let $\nu_0 = \kappa_0^{+\omega}$. In the extension of $V[H]$ by \mathbb{L}_s , ν_i will become $\aleph_{\omega+\omega_i}$.

As before, we inductively build a sequence s such that in the extension of $V[H]$ by a generic for \mathbb{L}_s , the strong tree property will hold at every ν_i^+ . To do this, we will need to make use of an auxiliary collapse.

Lemma 4.2. *There is $\mu_0^* < \kappa_0$ such that for all $s = \langle \mu_i \mid i < \omega \rangle \in I$ with $\mu_0 = \mu_0^*$, the strong tree property holds at ν_0^+ in $V[H][\mathbb{L}_s]$.*

Proof. Fix $\lambda \geq \nu_0^+$. Let K_λ be generic for $\text{Col}(\nu_0^+, \lambda)^{V[H]}$. In $V[H][K_\lambda]$, λ is collapsed to an ordinal with cardinality and cofinality ν_0^+ . Note that $V[H][K_\lambda]$ still satisfies the second and third criteria of Lemma 2.10.

For all $\mu_0 < \kappa_0$, working in $V[H]$, define

$$\mathbb{L}_0^*(\mu_0) = \text{Col}(\omega, \mu_0) \times \text{Col}(\mu_0^+, < \kappa_0) \times \left(\prod_{\mu_0 < \alpha < \kappa_0} \text{Col}(\alpha^+, < \kappa_0) \right),$$

where the final product has μ_0^{++} -support.

As before, the first component $\text{Col}(\omega, \mu_0)$ has size $< \mu_0^+$, while the remainder of the forcing is μ_0^+ -closed. The full poset has size $< \kappa_0^{++} = \kappa_2$. Thus $\mathbb{L}_0^*(\mu_0)$ satisfies the conditions for Lemma 2.10, with $V[H][K_\lambda]$ as the ground model. Applying this lemma, we obtain $\mu_0^*(\lambda) < \kappa_0$ such that in $V[H][K_\lambda][\mathbb{L}_0^*(\mu_0^*(\lambda))]$, every thin ν_0^+ -list has a cofinal branch.

Now, returning to $V[H]$ for a moment, let $s = \langle \mu_i \mid i < \omega \rangle \in I$ be a sequence such that $\mu_0 = \mu_0^*(\lambda)$. Let \dot{d} be a \mathbb{L}_s -name for a thin $\mathcal{P}_{\nu_0^+}(\lambda)$ -list with no branch. Let L_s be generic for \mathbb{L}_s . Note that $\text{Col}(\nu_0^+, \lambda)^{V[H]}$ is ν_0^+ -distributive in $V[H][L_s]$ by Easton's Lemma. In particular, $\mathcal{P}_{\nu_0^+}(\lambda)^{V[H][L_s]} = \mathcal{P}_{\nu_0^+}(\lambda)^{V[H][L_s][K_\lambda]}$.

Move to $V[H][L_s][K_\lambda]$, and let d be the interpretation of \dot{d} . By assumption d is a thin $\mathcal{P}_{\nu_0^+}(\lambda)$ -list in $V[H][L_s]$; it remains a thin $\mathcal{P}_{\nu_0^+}(\lambda)$ -list in $V[H][L_s][K_\lambda]$, since $\mathcal{P}_{\nu_0^+}(\lambda)$ is the same in both models. In $V[H][L_s][K_\lambda]$, since $|\lambda| = \nu_0^+$, there is an order isomorphism from $\mathcal{P}_{\nu_0^+}(\lambda)$ to $\mathcal{P}_{\nu_0^+}(\nu_0^+)$. Thus there is a thin $\mathcal{P}_{\nu_0^+}(\nu_0^+)$ -list d' that is isomorphic to d .

By Fact2.9, we can build a thin κ -list d^* such that d' has a cofinal branch if d^* does. Since d' is isomorphic to d , we conclude that if d^* has a cofinal branch, so does d .

Our goal now is to bring d^* up to $V[H][K_\lambda][\mathbb{L}_0^*(\mu_0^*(\lambda))]$, where we can obtain a branch for this list. We claim that $\mathbb{L}_0^*(\mu_0^*(\lambda))$ projects to \mathbb{L}_s . Clearly $\text{Col}(\omega, \mu_0^*)$ projects to itself via the identity. By Lemma 2.17, since $\prod_{n < \omega} \text{Col}(\mu_n^+, \mu_{n+1}^+)$ is $(\mu_0^*(\lambda))^+$ -closed and has size $< \kappa_0$, it is absorbed by $\text{Col}((\mu_0^*(\lambda))^+, < \kappa_0)$, with a quotient isomorphic to $\text{Col}((\mu_0^*(\lambda))^+, < \kappa_0)$. Finally, since $\sup_{i < \omega} (\mu_i)$ is in the interval $(\mu_0^*(\lambda), \kappa_0)$, the final product $\prod_{\mu_0^*(\lambda) < \alpha < \kappa_0} \text{Col}(\alpha^+, < \kappa_0)$ projects to $\text{Col}((\sup_{i < \omega} \mu_i)^+, < \kappa_0)$ via restricting to the appropriate coordinate of the product. The quotient of this projection is the product with that coordinate removed.

Since the list d^* is in $V[H][L_s]$, and $\mathbb{L}_0^*(\mu_0^*(\lambda))$ projects to \mathbb{L}_s , d^* must be in $V[H][\mathbb{L}_0^*(\mu_0^*(\lambda))]$. Since the levels of d^* are indexed by ordinals, and ν^+ is preserved by this quotient, d^* remains a thin ν_0^+ -list in this model. Every thin ν_0^+ -list in $V[H][K_\lambda][\mathbb{L}_0^*(\mu_0^*(\lambda))]$ has cofinal branch, so we obtain a branch b^* through d^* in $V[H][\mathbb{L}_0^*(\mu_0^*(\lambda))]$.

As before, the quotient of the projection from $\mathbb{L}_0^*(\mu_0^*(\lambda))$ to \mathbb{L}_s has size $< \nu_0^+$. By Corollary 2.15 this means that the quotient has the ν_0^+ -approximation property over $V[H][L_s][K_\lambda]$, and thus cannot add cofinal branches through thin ν^+ -lists. We conclude that b^* must be in $V[H][L_s][K_\lambda]$. Since d^* has a cofinal branch in $V[H][L_s][K_\lambda]$, d must also have a cofinal branch b in that model.

Since $\text{Col}(\nu_0^+, \lambda)^{V[H]}$ is ν^+ -closed in $V[H]$, and \mathbb{L}_s has size $< \nu_0^+$, by Lemma 2.16, $\text{Col}(\nu_0^+, \lambda)^{V[H]}$ has the thin ν_0^+ -approximation property over $V[H][L_s]$. It follows that $\text{Col}(\nu_0^+, \lambda)^{V[H]}$ could not have added b , so $b \in V[H][L_s]$. We conclude that for

all sequences $s \in I$ starting with $\mu_0^*(\lambda)$, in $V[H][L_s]$ every thin $\mathcal{P}_{\nu_0^+}(\lambda)$ -list has a cofinal branch.

Since there are only κ_0 options for each $\mu_0^*(\lambda)$, there must be a fixed μ_0^* such that for unboundedly many $\lambda \geq \kappa_0$, $\mu_0^*(\lambda) = \mu_0^*$. By Fact 2.6, to check the strong tree property we only need to verify that it holds for unboundedly many λ . So for all sequences $s \in I$ starting with μ_0^* , the strong tree property holds at ν_0^+ in $V[H][L_s]$. \square

Lemma 4.3. *Fix $\langle \mu_i^* \mid i < k \rangle$. There exists μ_k^* such that for all sequences $s = \langle \mu_i \mid i < \omega \rangle$ with $\mu_i = \mu_i^*$ for $i \leq k$, the strong tree property holds at ν_k^+ in $V[H][L_s]$.*

Proof. Fix $\lambda \geq \nu_k^+$. Let K_λ be generic for $\text{Col}(\nu_k, \lambda)^{V[H]}$. Note that in $V[H][K_\lambda]$, λ is collapsed to an ordinal with cardinality and cofinality ν_k^+ .

For all μ_k with $\mu_{k-1}^* < \mu_k < \kappa_0$, in $V[H]$ define

$$\mathbb{P}_k^*(\mu_k) := \text{Col}(\omega, \mu_0^*) \times \left(\prod_{n < k-1} \text{Col}((\mu_n^*)^+, \mu_{n+1}^*) \right) \times \text{Col}((\mu_{k-1}^*)^+, \mu_k)$$

and

$$\mathbb{Q}_k^*(\mu_k) := \text{Col}(\mu_k^+, < \kappa_0) \times \left(\prod_{\mu_k < \alpha < \kappa_0} \text{Col}(\alpha^+, < \kappa_0) \right),$$

where the product has μ_k^{++} -support. Define

$$\mathbb{L}_k^*(\mu_k) := \mathbb{P}_k^*(\mu_k) \times \mathbb{Q}_k^*(\mu_k).$$

Note that $|\mathbb{P}_k^*(\mu_k)| = \mu_k$, $\mathbb{Q}_k^*(\mu_k)$ is μ_k^+ -closed, and $|\mathbb{L}_k^*(\mu_k)| < \kappa_0^{++} = \kappa_2$. So $\mathbb{L}_k^*(\mu_k)$ meets the hypotheses of Lemma 2.10, applied in $V[H][K_\lambda]$.

We apply Lemma 2.10 to conclude that there is some $\mu_k^*(\lambda)$ such that in the extension of $V[H][K_\lambda]$ by a generic L for $\mathbb{L}_k^*(\mu_k^*(\lambda))$, every thin ν_k^+ -list has a cofinal branch.

In $V[H]$, let $s = \langle \mu_i \mid i < \omega \rangle$ with $\mu_n = \mu_n^*$ for $n < k$ and $\mu_k = \mu_k^*(\lambda)$, and let \dot{d} be a \mathbb{L}_s -name for a thin $\mathcal{P}_{\nu_k^+}(\lambda)$ -list. Let L_s be generic for \mathbb{L}_s . Once again, $\mathcal{P}_{\nu_k^+}(\lambda)^{V[H][K_\lambda]} = \mathcal{P}_{\nu_k^+}(\lambda)^{V[H]}$, and λ is collapsed to ν_k^+ . As before, applying Fact 2.9, we obtain a thin ν_k^+ -list d^* that has a cofinal branch if and only if d has a cofinal branch.

Working in $V[H][K_\lambda]$, we wish to show that $\mathbb{L}_k^*(\mu_k^*(\lambda))$ projects to \mathbb{L}_s . Note that $\mathbb{L}_s = \mathbb{P}_s \times \mathbb{Q}_s$, where

$$\mathbb{P}_s = \text{Col}(\omega, \mu_0) \times \left(\prod_{n < k} \text{Col}(\mu_n^+, \mu_{n+1}) \right)$$

and

$$\mathbb{Q}_s = \left(\prod_{k < n < \omega} \text{Col}(\mu_n^+, \mu_{i+1}) \right) \times \text{Col} \left((\sup_{i < \omega} \mu_i)^+, < \kappa_0 \right).$$

Note that since $\mu_n^* = \mu_n$ for $n < k$ and $\mu_k^*(\lambda) = \mu_k$, $\mathbb{P}_s = \mathbb{P}_k^*(\mu_k^*)$. By Lemma 2.17, noting that $\prod_{k < n < \omega} \text{Col}(\mu_n^+, \mu_{i+1})$ is $(\mu_k^*(\lambda))^+$ -closed and has size less than κ_0 , the first component of $\mathbb{Q}_k^*(\mu_k^*(\lambda))$ projects onto the first component of \mathbb{Q}_s . The second component of $\mathbb{Q}_k^*(\mu_k^*(\lambda))$ projects onto the second component of \mathbb{Q}_s by restricting to the coordinate indexed by $\alpha = \sup_{i < \omega} \mu_i$.

Let $L_k^*(\mu_k^*)$ be a generic for $\mathbb{L}_k^*(\mu_k^*)$ projecting to L_s . Since the levels of d^* are indexed by ordinals and ν_k^+ is preserved, d^* remains a thin ν_k^+ -tree in $V[H][L_k^*(\mu_k^*(\lambda))]$. Every thin ν_k^+ -list has a cofinal branch in this model, so there must be a cofinal branch through d^* in $V[H][L_k^*(\mu_k(\lambda))]$. Since the quotient has size $< \nu_k^+$, it cannot have added the branch by Lemma 2.14, so d^* must have a cofinal branch in $V[H][K_\lambda][L_s]$.

Since we have found such a cofinal branch for d^* , we conclude that d has a cofinal branch b in $V[H][K_\lambda][L_s]$. Since \mathbb{L}_s has size less than ν , and $\text{Col}(\nu^+, \lambda)^{V[H]}$ is ν^+ -closed in $V[H]$, applying Lemma 2.16 we conclude that $\text{Col}(\nu^+, \lambda)^{V[H]}$ could not have added a branch through d over $V[H][L_s]$.

It follows that for any sequence $s \in I$ starting with $\langle \mu_n^* \mid n < k \rangle \frown \mu_k^*(\lambda)$, any thin $\mathcal{P}_{\nu_k^+(s)}(\lambda)$ -list in $V[H][L_s]$ has a cofinal branch.

Since there are at most κ_0 many options for $\mu_k^*(\lambda)$, there is a fixed μ_k^* such that for unboundedly many λ , $\mu_k^*(\lambda) = \mu_k^*$. We conclude (using Fact 2.6) that for any sequence $s \in I$ starting with $\langle \mu_n^* \mid n < k \rangle \frown \mu_k^*(\lambda)$, the strong tree property holds at $\nu_k^+(s)$ in $V[H][L_s]$. \square

Let $s = \langle \mu_i^* \mid i < \omega \rangle$, and let L_s be generic for \mathbb{L}_s . As before, applying Lemma 3.2 (for $i = 0$) and Lemma 3.3 inductively, we conclude that in $V[H][L_s]$, the strong tree property holds at $\nu_i^+ = \aleph_{\omega+\omega_i+1}$ for all $i < \omega$. \square

5. EXTENDING FURTHER

The arguments in the previous two sections are not limited to countable sequences. The primary restriction on the number of cofinalities is that the sequence of μ_i 's needs to stay below κ_0 . In addition, if we are attempting to obtain the tree property at $\aleph_{\omega_\alpha+1}$, our argument requires that $\aleph_{\omega_\alpha} > \omega_\alpha$. Since we need our target cofinalities to be regular, we avoid this problem by restricting to successor cardinals.

Note however that the more cofinalities we wish to include, the larger our starting point will be. If our desired cofinalities are the first τ -many successor cardinals for some fixed $\tau < \kappa_0$, we will select a new $\omega_{\alpha+1}$ for all $\alpha < \tau$, and κ_0 will be collapsed to a finite successor of \aleph_τ . Then we will obtain the strong tree property at cardinals of the form $\aleph_{\tau+\omega_\alpha+1}$.

Theorem 5.1. *Let $\langle \kappa_\alpha \mid \alpha < \kappa_0 \rangle$ be an continuous increasing sequence of cardinals with the following properties:*

- $\kappa_{\alpha+1} = \kappa_\alpha^+$ for all limit ordinals α
- κ_n is indestructibly supercompact for all $n < \omega$
- $\kappa_{\alpha+2}$ is indestructibly supercompact for all $\omega \leq \alpha < \kappa_0$.

Fix $\tau < \kappa_0$. Then there is a generic extension in which the strong tree property holds at $\aleph_{\tau+\omega_\alpha+1}$ for all successor ordinals $\alpha < \tau$.

Proof. Our construction is as before.

Define the poset \mathbb{H} as follows:

$$\mathbb{H} := \left(\prod_{n < \omega} \text{Col}(\kappa_n, < \kappa_{n+1}) \right) \times \left(\prod_{\omega \leq \rho < \kappa_0} \text{Col}(\kappa_{\rho+1}, < \kappa_{\rho+2}) \right).$$

Let I be the collection of all continuous strictly increasing sequences $\langle \mu_\alpha \mid \alpha < \tau \rangle$. For all sequences $s = \langle \mu_\alpha \mid \alpha < \tau \rangle \in I$, define:

$$\mathbb{L}_s := \text{Col}(\omega, \mu_0) \times \left(\prod_{\alpha < \tau} \text{Col}(\mu_\alpha^+, \mu_{\alpha+1}) \right) \times \text{Col}((\sup_{\alpha < \tau} \mu_{\alpha+1})^+, < \kappa_0).$$

Note that after forcing with \mathbb{L}_s , μ_α^+ will become $\aleph_{\omega_{\alpha+1}}$. We inductively select μ_α^* such that for any sequence $s \in I$ beginning with $\langle \mu_\beta^* \mid \beta \leq \alpha \rangle$, the strong tree property will hold at $\aleph_{\tau+\alpha+1}$ in $V[H][L_s]$. The base case is more or less identical to Lemma 4.2, using

$$\mathbb{L}_0^*(\mu_0) = \text{Col}(\omega, \mu_0) \times \text{Col}(\mu_0^+, < \kappa_0) \times \left(\prod_{\mu_0 < \beta < \kappa_0} \text{Col}(\beta^+, < \kappa_0) \right).$$

At limit stages γ , since the sequence must be continuous, we set $\mu_\gamma^* = \sup_{\alpha < \gamma} \mu_\alpha^*$. At successor stages, we follow the argument of Lemma 4.3, using

$$\begin{aligned} \mathbb{P}_{\alpha+1}^*(\mu_{\alpha+1}) &:= \text{Col}(\omega, \mu_0^*) \times \left(\prod_{\beta < \alpha} \text{Col}((\mu_\beta^*)^+, \mu_{\beta+1}^*) \right) \times \text{Col}(\mu_\alpha^*, \mu_{\alpha+1}), \\ \mathbb{Q}_{\alpha+1}^*(\mu_{\alpha+1}) &:= \text{Col}(\mu_{\alpha+1}^+, < \kappa_0) \times \left(\prod_{\mu_{\alpha+1} < \beta < \kappa_0} \text{Col}(\beta^+, < \kappa_0) \right), \end{aligned}$$

and

$$\mathbb{L}_{\alpha+1}^*(\mu_{\alpha+1}) := \mathbb{P}_{\alpha+1}^*(\mu_{\alpha+1}) \times \mathbb{Q}_{\alpha+1}^*(\mu_{\alpha+1}).$$

When this induction concludes, we will have a sequence $s \in I$ such that in $V[H][L_s]$, the strong tree property holds at $\aleph_{\tau+\omega_\alpha+1}$ for all successor ordinals $\alpha < \tau$. \square

REFERENCES

- [1] Uri Abraham. Aronszajn trees on \aleph_2 and \aleph_3 . *Ann. Pure Appl. Logic*, 24(3):213–230, 1983.
- [2] William Adkisson. The strong and super tree properties at successors of singular cardinals. *The Journal of Symbolic Logic*, 89(3):1251–1283, 2024.
- [3] James Cummings. Iterated forcing and elementary embeddings. In *Handbook of set theory. Vols. 1, 2, 3*, pages 775–883. Springer, Dordrecht, 2010.
- [4] James Cummings and Matthew Foreman. The tree property. *Adv. Math.*, 133(1):1–32, 1998.
- [5] Laura Fontanella. Strong tree properties for small cardinals. *J. Symbolic Logic*, 78(1):317–333, 2013.
- [6] Laura Fontanella. The strong tree property at successors of singular cardinals. *J. Symb. Log.*, 79(1):193–207, 2014.
- [7] Mohammad Golshani and Yair Hayut. The tree property on a countable segment of successors of singular cardinals. *Fund. Math.*, 240(2):199–204, 2018.
- [8] Sherwood Hachtman and Dima Sinapova. The super tree property at the successor of a singular. *Israel J. Math.*, 236(1):473–500, 2020.
- [9] Thomas J. Jech. Some combinatorial problems concerning uncountable cardinals. *Ann. Math. Logic*, 5:165–198, 1972/73.
- [10] Menachem Magidor and Saharon Shelah. The tree property at successors of singular cardinals. *Arch. Math. Logic*, 35(5-6):385–404, 1996.
- [11] William Mitchell. Aronszajn trees and the independence of the transfer property. *Ann. Math. Logic*, 5:21–46, 1972/73.
- [12] Itay Neeman. The tree property up to $\aleph_{\omega+1}$. *J. Symbolic Logic*, 79(2):429–459, 2014.
- [13] Spencer Unger. Aronszajn trees and the successors of a singular cardinal. *Arch. Math. Logic*, 52(5-6):483–496, 2013.
- [14] Spencer Unger. A model of Cummings and Foreman revisited. *Ann. Pure Appl. Logic*, 165(12):1813–1831, 2014.