

**VERTEX ALGEBRAS RELATED TO REGULAR REPRESENTATIONS
OF SL_2**

DRAŽEN ADAMOVIĆ AND ANTUN MILAS

ABSTRACT. We construct a family of potentially quasi-lisse (non-rational) vertex algebras, denoted by \mathcal{C}_p , $p \geq 2$, which are closely related to the vertex algebra of chiral differential operators on $SL(2)$ at level $-2 + \frac{1}{p}$. The parameter p also serves as a dilation parameter of the weight lattice of A_1 . We prove that for $p = 3$, we have $\mathcal{C}_3 \cong L_{-5/3}(\mathfrak{g}_2)$. Moreover, we also establish isomorphisms between \mathcal{C}_4 and \mathcal{C}_5 and certain affine W -algebras of types F_4 and E_8 , respectively. In this way, we resolve the problem of decomposing certain conformal embeddings of affine vertex algebras into affine W -algebras. An important feature is that \mathcal{C}_p is $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded with finite-dimensional graded subspaces and convergent characters. Therefore, for all $p \geq 2$, we show that the characters of \mathcal{C}_p exhibit modularity, supporting the conjectural quasi-lisse property.

CONTENTS

1. Introduction	2
1.1. Regular representations of vertex algebras	2
1.2. Minimal and Weil representations	2
1.3. Our paper and main results	2
1.4. Notation	4
2. Inverse Quantum Hamiltonian Reduction	5
3. Construction of the vertex algebra \mathcal{C}_p	6
4. The vertex algebra $L_{-\frac{5}{3}}(G_2)$	9
4.1. Decomposition of $M_{(2)}$	9
4.2. Isomorphism $\mathcal{C}_3 \cong L_{-5/3}(G_2)$	11
5. The vertex algebra \mathcal{C}_4 as inverse QHR of $L_{-3/2}(\mathfrak{sl}_3)$	12
5.1. The decomposition of $L_{-3/2}(\mathfrak{sl}_3)$ as $L_{-6}(\mathfrak{sl}_2) \otimes L_{\mathfrak{c}_{1,4}}^{Vir}$ -module	12
6. Realization of \mathcal{C}_p as affine W -algebras: the cases $p = 4, 5$	14
6.1. Affine W -algebras	14
6.2. Affine W -algebra $\mathcal{W}_k(F_4, A_1 + \tilde{A}_1)$	14
6.3. The vertex algebra $\mathcal{W}_{-30+31/5}(E_8, A_4 + A_2)$	16
7. Proof of some character identities	17
8. MLDE for \mathcal{C}_p	20
9. Modularity of the character of \mathcal{C}_n	21
9.1. Appell-Lerch series	22
10. Future work	23
References	24

2010 *Mathematics Subject Classification*. Primary 17B69; Secondary 17B20, 17B67.
Key words and phrases. vertex algebra, W -algebras, affine Lie algebra, regular representations.

1. INTRODUCTION

1.1. Regular representations of vertex algebras. The study of the regular representation of vertex algebras began with the work of Feigin and Parkhomenko [17] and more comprehensively by Frenkel and Styrkas [19] on their work on modified regular representation. An alternative approach involves the geometric construction of algebras of global chiral differential operators \mathcal{D}_X^{ch} on the cotangent bundle T^*G , which can be traced back to earlier works [1, 23, 24]; see also [31] for connection between two approaches. A more recent perspective on chiral differential operators $\mathcal{D}_{G,k}^{ch}$ was undertaken by Arakawa, who explored them within the context of quasi-lisse vertex algebras and genus zero S -class theories in 4d $N = 2$ SCFT [4].

These works are rooted in classical representation theory, starting with the regular representation $\mathcal{O}(G)$, here G is semisimple, that decomposes by the Peter–Weyl theorem

$$(1.1) \quad \mathcal{O}(G) = \bigoplus_{\lambda \in P_+} V_\lambda \otimes V_{\lambda^*},$$

where V_λ is finite-dimensional irreducible $\mathfrak{g} = \text{Lie}(G)$ -module of highest weight λ and $\lambda^* = -w_0(\lambda)$ ¹. In the affine Lie algebra setting, the algebra of function is replaced with the vertex algebra of chiral differential operators that decomposes as $\hat{\mathfrak{g}} \times \hat{\mathfrak{g}}$ -module

$$(1.2) \quad \mathcal{D}_{G,k}^{ch} \cong \bigoplus_{\lambda \in P_+} L_{-k-h^\vee, \lambda}(\mathfrak{g}) \otimes L_{k-h^\vee, \lambda^*}(\mathfrak{g}),$$

where $k \notin \mathbb{Q}$ is a generic level and h^\vee is the dual Coxeter number.

1.2. Minimal and Weil representations. The oscillator, or Weil representation, of $\mathfrak{sp}(2n)$, also known as the metaplectic representation, is a fundamental tool in harmonic analysis and representation theory. Weil-type representations also exist for exceptional Lie algebras. For example, S. Gelfand [22] studied a representation of the exceptional Lie algebra \mathfrak{g}_2 , realized as differential operators acting on a suitable space of Laurent polynomials in three variables; see also [30]. These representations are minimal because their annihilator J in the universal enveloping algebra $U(\mathfrak{g})$ coincides with the Joseph ideal \mathcal{J}_0 - the unique completely prime ideal satisfying $\text{Ass}(\mathcal{J}_0) = \overline{\mathcal{O}}_{\min}$.

Minimal and Weil representations also appear in the context of vertex algebras. It is known [2] that for every affine vertex operator algebras $L_k(\mathfrak{g})$, with $k = -\frac{h^\vee}{6} - 1$, in the Deligne series

$$A_1 \subset A_2 \subset G_2 \subset D_4 \subset E_6 \subset E_7 \subset E_8,$$

the Zhu algebra $A(L_k(\mathfrak{g}))$ contains $\mathcal{U}(\mathfrak{g})/J_0$ as one of its components, indicating that the minimal representations of \mathfrak{g} also appear as the top component of certain $L_k(\mathfrak{g})$ -modules. Interestingly, this property also arises for vertex algebras outside Deligne's series. For the symplectic Lie algebra \mathfrak{sp}_{2n} , this occurs for the vertex algebra $L_{-1/2}(\mathfrak{sp}_{2n})$ [3], which is also the \mathbb{Z} -graded part of the $\beta\gamma$ -system $M_{(n)}$ of rank n . Vertex algebras in these series are *quasi-lisse*, meaning their associated variety $X_{L_k(\mathfrak{g})} := \text{Specm}(R_{L_k(\mathfrak{g})})$ admits finitely many symplectic leaves.

1.3. Our paper and main results. In this paper, we construct a family of vertex algebras that, on one hand, deform the regular representation of $\hat{\mathfrak{sl}}_2$, and, on the other, exhibit characteristics of the Deligne series of vertex algebras at admissible level discussed above. Consequently, these algebras have finite-dimensional $\mathbb{Z}_{\geq 0}$ -subspaces and are expected to be quasi-lisse. Although \mathcal{C}_p is modeled on $\mathcal{D}_{SL(2), -2 + \frac{1}{p}}^{ch}$, which pairs the levels $-2 + \frac{1}{p}$ and $-2 - \frac{1}{p}$ as in (1.2), there is a crucial difference here: \mathcal{C}_p pairs levels $-2 - p$ and $-2 + \frac{1}{p}$ as in the quantum version of the geometric Langlands correspondence [21].

¹There is also version of this result for $\mathcal{O}(Bw_0B)$ where Bw_0B is the big Bruhat cell [19].

To this end, we consider extensions of the vertex algebra $L_k(\mathfrak{sl}_2) \otimes L_{k'}(\mathfrak{sl}_2)$ such that $k + 2 = \frac{1}{p}$, $k' + 2 = -p$. One of our main results is the following theorem.

Theorem 1.1. *For every $p \geq 1$, there is a simple vertex algebra \mathcal{C}_p such that $L_{-2+\frac{1}{p}}(\mathfrak{sl}_2) \otimes L_{-2-p}(\mathfrak{sl}_2)$ is conformally embedded into \mathcal{C}_p , and we have the following decomposition:*

$$\mathcal{C}_p = \bigoplus_{\ell=0}^{\infty} L_{\widehat{\mathfrak{sl}_2}}(-2+p+p\ell)\Lambda_0 + p\ell\Lambda_1 \otimes L_{\widehat{\mathfrak{sl}_2}}(-2-\frac{1}{p}+\ell)\Lambda_0 + \ell\Lambda_1.$$

To prove this result, we use the following vertex algebras constructed in the literature on chiral differential operators [8] (see [3]):

$$\mathcal{U}_p := \bigoplus_{\ell=0}^{\infty} L_{\widehat{\mathfrak{sl}_2}}(-2+\frac{1}{p}+\ell)\Lambda_0 + \ell\Lambda_1 \otimes L_{\widehat{\mathfrak{sl}_2}}(-2-\frac{1}{p}+\ell)\Lambda_0 + \ell\Lambda_1,$$

and

$$\mathcal{V}_p := H_{DS,f}(\mathcal{U}_p) = \bigoplus_{\ell=0}^{\infty} L^{Vir}(c_{1,-p}, h_{1,\ell+1}^{1,-p}) \otimes L_{\widehat{\mathfrak{sl}_2}}(-2-\frac{1}{p}+\ell)\Lambda_0 + \ell\Lambda_1.$$

We notice that

$$L^{Vir}(c_{1,-p}, h_{1,\ell+1}^{1,-p}) = L^{Vir}(c_{-p,1}, h_{1,np+1}^{-p,1}),$$

which indicates that we can apply the inverse Quantum Hamiltonian Reduction (QHR) from a suitable category of $L_{c_{1,-p}}^{Vir}$ -modules to the category $KL_{-p-2}(\mathfrak{sl}_2)$ (cf. Section 2). As a consequence, we get that inverse QHR acts from the category of $L_{c_{1,-p}}^{Vir} \otimes L_{-2+\frac{1}{p}}(\mathfrak{sl}_2)$ -modules to the category of $L_{-2-p}(\mathfrak{sl}_2) \otimes L_{-2+\frac{1}{p}}(\mathfrak{sl}_2)$ -modules, and we prove that \mathcal{C}_p is exactly the inverse QHR of \mathcal{V}_p .

Theorem 1.2. *There is an action of the Lie algebra $\mathfrak{g}_0 = \mathfrak{sl}_2 \times \mathfrak{sl}_2$ on $\mathcal{V}_p \otimes \Pi(0)^{1/2}$ so that*

$$\mathcal{C}_p = \left(\mathcal{V}_p \otimes \Pi(0)^{1/2} \right)^{\text{int}_{\mathfrak{g}_0}} \cong \bigoplus_{\ell=0}^{\infty} L_{\widehat{\mathfrak{sl}_2}}(-2+p+p\ell)\Lambda_0 + p\ell\Lambda_1 \otimes L_{\widehat{\mathfrak{sl}_2}}(-2-\frac{1}{p}+\ell)\Lambda_0 + \ell\Lambda_1.$$

For small values of p , we have more explicit realizations of \mathcal{C}_p , which does not use chiral differential operators and results from [4].

- For $p = 2$, it was proven in [12, Section 9] that $\mathcal{C}_2 \cong M_{(3)}$. This result was obtained in the context of conformal embeddings $\mathfrak{sl}_2 \times \mathfrak{sl}_2 \hookrightarrow \mathfrak{sp}_6$ at $k = -\frac{1}{2}$. Note that $M_{(3)}^{\mathbb{Z}_2} \cong L_{-1/2}(\mathfrak{sp}_6)$.
- For $p = 3$, we prove that $\mathcal{C}_3 \cong L_{-\frac{5}{3}}(G_2)$. In order to see this, we first prove that the rank two Weyl vertex algebra $M_{(2)}$ is completely reducible as a module for $L_{-5}(\mathfrak{sl}_2) \otimes V_{c_{1,3}}^{Vir}$ (cf. Theorem 4.2). Then we show that the vertex algebra $L_{-\frac{5}{3}}(G_2)$ is obtained by inverse QHR of $M_{(2)}$:

$$L_{-\frac{5}{3}}(G_2) = \text{Ker}_{M_{(2)} \otimes \Pi(0)^{1/2}} S,$$

which implies that $\mathcal{C}_3 \cong L_{-\frac{5}{3}}(G_2)$ (cf. Theorem 4.4).

As a byproduct, we obtain the decomposition of conformal embedding $\mathfrak{sl}_2 \times \mathfrak{sl}_2 \hookrightarrow G_2$ at $k = -\frac{5}{3}$ initiated in [10].

Since $L_{-\frac{5}{3}}(G_2)$ is admissible affine vertex algebra, it is quasi-lisse. So \mathcal{C}_3 is quasi-lisse.

- For $p = 4$, in Theorem 5.3 we prove that \mathcal{C}_4 is realized as the inverse quantum hamiltonian reduction of $L_{-3/2}(\mathfrak{sl}_3)$. The key point for this result is the decomposition of $L_{-3/2}(\mathfrak{sl}_3)$ as $L_{-6}(\mathfrak{sl}_2) \otimes V_{c_{1,4}}^{Vir}$ -modules (cf. Theorem 5.2).
- Next we identify \mathcal{C}_4 with the affine vertex algebra $\mathcal{W}_{-\frac{23}{4}}(F_4, A_1 + \tilde{A}_1)$. Since it is obtained by the QHR of the admissible affine vertex algebra $L_{-\frac{23}{4}}(F_4)$, we conclude that \mathcal{C}_4 is quasi-lisse (cf. Proposition 6.3).

- For $p = 5$, in Theorem 6.9 we established an isomorphism

$$\mathcal{C}_5 \cong \mathcal{W}_{-30+31/5}(E_8, A_4 + A_2).$$

Since $k = -30 + 31/5$ is admissible level for E_8 we also get that \mathcal{C}_5 is quasi lisse; see Proposition 6.8.

- It is interesting to notice that for $p = 2, 3, 4, 5$, the relevant levels of affine vertex algebra needed to define \mathcal{C}_p are given by $-h^\vee + \frac{h+1}{p}$. For $p \geq 6$, we do not expect \mathcal{C}_p to be isomorphic to another affine W -algebra.

By a result of Arakawa and Kawasetsu [6], it is known that the character of a quasi-lisse vertex operator algebra satisfies a particular modular linear differential equation (MLDE). Motivated by their work, we construct explicit MLDEs of minimal order satisfied by $\text{ch}[\mathcal{C}_p]$ for $p \in \{2, 3, 4, 5\}$ (see Section 8). That would also follow from general properties of the associated varieties.

In the last section we obtained a closed formula for $\text{ch}[\mathcal{C}_p]$ and discuss modular transformation properties; see Theorem 9.3. Based on that, we present further evidence that \mathcal{C}_p has a unique ordinary module.

1.4. Notation. Throughout we will be using the following notation:

- \mathfrak{g} will denote a finite-dimensional simple Lie algebra of rank ℓ , $\mathfrak{h} \subset \mathfrak{g}$ its Cartan subalgebra and we choose $\{\alpha_1, \dots, \alpha_\ell\}$ a set of simple roots and ω_i be the corresponding fundamental weights. Subject to this, we denote by Q the root lattice, P the weight lattice, and P_+ the cone of dominant integral weights.
- Given $\lambda \in P_+$, we denote by $V_{\mathfrak{g}}(\lambda)$ the irreducible finite-dimensional \mathfrak{g} -module of highest weight λ .
- $Vir = \text{Span}\{L(n), n \in \mathbb{Z}, c\}$ will denote the Virasoro Lie algebra with the usual bracket relations and central element c .
- V_c^{Vir} denotes the universal Virasoro vertex algebra of central charge c .
- L_c^{Vir} denotes the simple Virasoro vertex algebra of central charge c .
- Let $L^{Vir}(c, h)$ will denote the irreducible highest weight V_c^{Vir} -module with highest weight h .
- We write $\tilde{L}^{Vir}(c, h)$ for the highest weight V_c^{Vir} -module with highest weight h , which is not necessary irreducible.
- $\hat{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}C$ is the affine Lie algebra associated to \mathfrak{g} with central element C .
- $V^k(\mathfrak{g})$ will denote the universal affine vertex algebra associated to \mathfrak{g} of level k .
- $L_k(\mathfrak{g})$ will denote the simple affine vertex algebra associated to \mathfrak{g} of level k .
- $L_{\hat{\mathfrak{g}}}(\Lambda)$ will denote the irreducible $\hat{\mathfrak{g}}$ -module of highest weight module Λ . We denote the fundamental weights by Λ_i , $0 \leq i \leq \ell$. We also use sometimes $L_{\hat{\mathfrak{g}}, k}(\lambda)$, $\lambda \in \mathfrak{h}^*$ to denote the same module.
- We write $\tilde{L}_{\hat{\mathfrak{g}}}(\Lambda)$ for the highest weight $\hat{\mathfrak{g}}$ -module of highest weight module Λ , which is not-necessary irreducible.
- $H_{DS}(\cdot)$ will denote the Drinfeld-Sokolov reduction functor [4].
- $\mathcal{W}^k(\mathfrak{g}, f)$ will denote the universal affine W -algebra of level k and nilpotent element f . We also denote by $\mathcal{W}_k(\mathfrak{g}, f)$ its simple quotient.
- $M_{(n)}$ denotes the $\frac{1}{2}$ -shifted $\beta\gamma$ -system of rank n , also known as the Weyl vertex algebra.
- Given a conformal vertex algebra V -module of central charge c , and a V -module M , then

$$\text{ch}[M] = \text{tr}_M q^{L(0) - \frac{c}{24}},$$

will denote the character of M .

- If a semi-simple Lie algebra \mathfrak{g} acts on a vertex algebra V by derivations, let $V^{\text{int}_{\mathfrak{g}}}$ denote the maximal \mathfrak{g} -integrable part of V , i.e., the direct sum of all finite-dimensional \mathfrak{g} -submodules of V . The space $V^{\text{int}_{\mathfrak{g}}}$ is a vertex subalgebra of V .

- Vertex algebra V is called quasi-lisse if the associative variety $X_{L_k(\mathfrak{g})} = \text{Specm}(R_{L_k(\mathfrak{g})})$ admits finitely many symplectic leaves (cf. [6]).

2. INVERSE QUANTUM HAMILTONIAN REDUCTION

We shall follow realization of $L_k(\mathfrak{sl}_2)$ from [9]. Let $k+2 = \frac{p'}{p}$. For any two co-prime integers p, p' we set

$$c_{p',p} := 1 - 6 \frac{(p-p')^2}{pp'}.$$

For $r, s \in \mathbb{Z}$, we set

$$h_{r,s}^{p',p} = \frac{(sp - rp')^2 - (p-p')^2}{4pp'}.$$

Recall the following well-known fusion rules in the category of $L_{c_{1,p}}^{Vir}$ -modules for $i \geq 2$:

$$(2.1) \quad L^{Vir}(c_{1,p}, h_{1,2}^{1,p}) \times L^{Vir}(c_{1,p}, h_{1,i}^{1,p}) = L^{Vir}(c_{1,p}, h_{1,i-1}^{1,p}) + L^{Vir}(c_{1,p}, h_{1,i+1}^{1,p}).$$

Let ω be the conformal vector in $V_{c_{p',p}}^{Vir}$. Define

$$(2.2) \quad \Pi(0) := M_{c,d}(1) \otimes \mathbb{C}[\mathbb{Z}c] \quad \text{and} \quad \Pi(0)^{\frac{1}{2}} = M_{c,d}(1) \otimes \mathbb{C}[\mathbb{Z}\frac{c}{2}],$$

where $M_{c,d}(1)$ is the Heisenberg vertex algebra generated by fields $c(z)$ and $d(z)$.

We also have the following \mathbb{Z}_2 -gradation on $\Pi(0)^{\frac{1}{2}}$

$$\Pi(0)^{\frac{1}{2}} = \Pi_0(0)^{\frac{1}{2}} + \Pi_1(0)^{\frac{1}{2}},$$

where $\Pi_0(0)^{\frac{1}{2}}$ and $\Pi_1(0)^{\frac{1}{2}} = \Pi(0).e^{\frac{c}{2}}$.

Now [9] gives homomorphism of vertex algebras

$$\Phi : V^k(\mathfrak{sl}_2) \rightarrow V_{c_{p',p}}^{Vir} \otimes \Pi(0)$$

such that

$$(2.3) \quad \begin{aligned} e &\mapsto \mathbf{1} \otimes e^c, \\ h &\mapsto 2[\mathbf{1} \otimes \mu(-1)\mathbf{1}], \\ f &\mapsto (k+2)\omega \otimes e^{-c} - \mathbf{1} \otimes [\nu(-1)^2 - (k+1)\nu(-2)]e^{-c}, \end{aligned}$$

where

$$(2.4) \quad \mu = \frac{1}{2}d + \frac{k}{4}c, \quad \nu = \frac{1}{2}d - \frac{k}{4}c.$$

As in [9] we have screening operator

$$(2.5) \quad S = \text{Res}_z Y(v^{(k)} \otimes e^\nu, z),$$

where $v^{(k)}$ is a singular vector for the Virasoro algebra highest weight module of conformal weight $h_{2,1}^{p',p} = \frac{3}{4}k + 1$.

The following result follows from [9] and [13]:

Proposition 2.1. *Assume that $k \notin \mathbb{Z}_{\geq 0}$, $k \neq -2$ and $n \in \mathbb{Z}_{\geq 0}$. Then we have*

$$(2.6) \quad \begin{aligned} L_{\widehat{\mathfrak{sl}_2}}((k-n)\Lambda_0 + n\Lambda_1) &\cong V^k(\mathfrak{sl}_2).(v_{1,n+1} \otimes e^{\frac{n}{2}c}) \\ &\cong \text{Ker}_{L^{Vir}(c_{p',p}, h_{1,n+1}^{p',p}) \otimes \Pi(0)^{1/2}} S \\ &\cong \left(L^{Vir}(c_{p',p}, h_{1,np+1}^{1,p}) \otimes \Pi(0)^{1/2} \right)^{\text{int}_{\mathfrak{sl}_2}}. \end{aligned}$$

In particular, $L_k(\mathfrak{sl}_2) = \left(L_{c_{p',p}}^{Vir} \otimes \Pi(0)^{1/2} \right)^{\text{int}_{\mathfrak{sl}_2}}$.

This result also holds for negative levels such that $k + 2 < 0$. Let us look the negative level case in more details. We shall consider special cases:

$$k + 2 = -\frac{1}{p} \quad \text{and} \quad k + 2 = -p,$$

where $p \in \mathbb{Z}_{>0}$. We have

$$H_{DS}(L_{\widehat{\mathfrak{sl}_2}}((-2 - \frac{1}{p} - n)\Lambda_0 + n\Lambda_1) = L^{Vir}(c_{1,-p}, h_{1,n+1}^{1,-p}),$$

$$H_{DS}(L_{\widehat{\mathfrak{sl}_2}}((-2 - p - np)\Lambda_0 + np\Lambda_1) = L^{Vir}(c_{-p,1}, h_{1,np+1}^{-p,1}).$$

Since $h_{1,n+1}^{-1,p} = h_{1,np+1}^{-p,1}$, we have

$$L^{Vir}(c_{1,-p}, h_{1,n+1}^{-1,p}) = L^{Vir}(c_{-p,1}, h_{1,np+1}^{-p,1}),$$

which implies

$$H_{DS}(L_{\widehat{\mathfrak{sl}_2}}((-2 - \frac{1}{p} - n)\Lambda_0 + n\Lambda_1) = H_{DS}(L_{\widehat{\mathfrak{sl}_2}}((-2 - p - np)\Lambda_0 + np\Lambda_1).$$

Corollary 2.2. *Let $k + 2 = -p$ and n a positive integer. Then we have*

$$L_{\widehat{\mathfrak{sl}_2}}((k - np)\Lambda_0 + np\Lambda_1) \cong \text{Ker}_{L^{Vir}(c_{-p,1}, h_{1,np+1}^{-p,1}) \otimes \Pi(0)^{1/2}S} \cong \text{Ker}_{L^{Vir}(c_{-p,1}, h_{1,n+1}^{1,-p}) \otimes \Pi(0)^{1/2}S},$$

where S is screening operator given by (2.5).

Moreover, we have

$$L_{\widehat{\mathfrak{sl}_2}}((k - np)\Lambda_0 + np\Lambda_1) = \left(L^{Vir}(c_{-p,1}, h_{1,np+1}^{-p,1}) \otimes \Pi(0)^{1/2} \right)^{\text{int}_{\mathfrak{sl}_2}}.$$

Recall that the fusion product of two subsets A, B of a vertex algebra V is

$$A \cdot B = \text{span}_{\mathbb{C}}\{a_n b \mid n \in \mathbb{Z}, a, b \in V\}.$$

This product is associative due to the Borcherds identity.

3. CONSTRUCTION OF THE VERTEX ALGEBRA \mathcal{C}_p

Next result is a slight variation of the result on generic levels mentioned in the introduction as it only requires some minor modification due to rationality of the level involved.

Theorem 3.1. *For each $p \in \mathbb{Z}_{\geq 1}$, there exists a simple vertex operator algebra structure on the $L_{-2+\frac{1}{p}}(\mathfrak{sl}_2) \otimes L_{-2-\frac{1}{p}}(\mathfrak{sl}_2)$ -module*

$$\mathcal{U}_p := \bigoplus_{\ell=0}^{\infty} L_{\widehat{\mathfrak{sl}_2}}(-2 + \frac{1}{p} + \ell)\Lambda_0 + \ell\Lambda_1 \otimes L_{\widehat{\mathfrak{sl}_2}}(-2 - \frac{1}{p} + \ell)\Lambda_0 + \ell\Lambda_1.$$

\mathcal{U}_p is generated by the generators of $L_{-2-1/p}(\mathfrak{sl}_2) \otimes L_{-2+1/p}(\mathfrak{sl}_2)$ and top component of the module $L_{\widehat{\mathfrak{sl}_2}}(-3 + \frac{1}{p})\Lambda_0 + \Lambda_1 \otimes L_{\widehat{\mathfrak{sl}_2}}(-3 - \frac{1}{p})\Lambda_0 + \Lambda_1$ isomorphic to $V_{\mathfrak{sl}_2}(\omega_1) \otimes V_{\mathfrak{sl}_2}(\omega_1)$.

Proof. It is known that $L_{-2+\frac{1}{p}}(\mathfrak{sl}_2) \otimes L_{-2-\frac{1}{p}}(\mathfrak{sl}_2) \hookrightarrow \mathcal{D}_{SL_2, -2+\frac{1}{p}}^{ch}$ and that [8]

$$\left(\mathcal{D}_{SL_2, -2+\frac{1}{p}}^{ch} \right)^{t\mathfrak{sl}_2[t] \times t\mathfrak{sl}_2[t]} \cong \mathcal{O}(SL_2) = \bigoplus_{\ell \in \mathbb{N}_0} V_{\mathfrak{sl}_2}(\ell\omega_1) \otimes V_{\mathfrak{sl}_2}(\ell\omega_1).$$

Combining this with complete reducibility of the categories $KL^{-2+\frac{1}{p}}(\mathfrak{sl}_2)$ and $KL^{-2-\frac{1}{p}}(\mathfrak{sl}_2)$ we get that $\mathcal{D}_{-2+\frac{1}{p}}^{ch}$ is isomorphic to \mathcal{U}_p (exactly the same argument is used Section 9.1.6 of [3] for the generic level). But it is known that $\mathcal{D}_{-2+\frac{1}{p}}^{ch}$ is simple; see again [3].

Using $\text{gr}(\mathcal{D}_{G,k}^{ch}) \cong \mathcal{O}(J_{\infty}(T^*G))$ (arc algebra of $T^*G \cong \mathfrak{g}^* \times G$), we see that $\mathcal{D}_{G,k}^{ch}$ is generated by the top component $O(G)$, and elements of \mathfrak{g} , embedded in the degree one component. For $G = SL_2$, it is not hard to see using explicit action of $\mathfrak{sl}_2 \times \mathfrak{sl}_2$ on $O(SL_2)$ (see for instance [19]) that $V(1) \otimes V(1)$ generates $O(SL_2)$, and therefore

the algebra $\mathcal{D}_{SL_2, k}^{ch}$ is generated by vectors in $L_{\widehat{\mathfrak{sl}_2}}((-2-k)\Lambda_0) \otimes L_{\widehat{\mathfrak{sl}_2}}((-2+k)\Lambda_0)$ and $L_{\widehat{\mathfrak{sl}_2}}((-3-k)\Lambda_0 + \Lambda_1) \otimes L_{\widehat{\mathfrak{sl}_2}}((-3+k)\Lambda_0 + \Lambda_1)$, for all k . In particular, everything holds for $k = -2 + \frac{1}{p}$. \square

Remark 3.2. When $k \notin \mathbb{Q}$, that is level k is generic, then a version of this result was also obtained more explicitly by I. Frenkel and K. Styrkas [19] using Wakimoto-type construction. In their setup, $\mathcal{D}_{SL_2, k}^{ch}$ appears as a subalgebra of their modified regular representation. Their proof also relies on the use of semi-simplicity of the category $KL^k(\mathfrak{sl}_2)$ when k is generic; see also [28].

Let f (resp. \bar{f}) be principal nilpotent element in first (respond. second) copy of \mathfrak{sl}_2 . Arakawa in [8] (see also comments in [28]) consider the QHR related to f and \bar{f} . Applying his construction on the vertex algebra \mathcal{U}_p , which is simple by the theorem above, we get:

Corollary 3.3. *We have the following simple vertex algebras:*

$$\begin{aligned} \mathcal{V}_p &= H_{DS, f}(\mathcal{U}_p) = \bigoplus_{\ell=0}^{\infty} L^{Vir}(c_{1, -p}, h_{1, \ell+1}^{1, -p}) \otimes L_{\widehat{\mathfrak{sl}_2}}\left(-\left(2 - \frac{1}{p} + \ell\right)\Lambda_0 + \ell\Lambda_1\right), \\ H_{DS, \bar{f}}(\mathcal{U}_p) &= \bigoplus_{\ell=0}^{\infty} L_{\widehat{\mathfrak{sl}_2}}\left(-\left(2 + \frac{1}{p} + \ell\right)\Lambda_0 + \ell\Lambda_1\right) \otimes L^{Vir}(c_{1, p}, h_{1, \ell+1}^{1, p}), \\ I_{SL_2}^{-2-\frac{1}{p}} &= H_{DS, f, \bar{f}}(\mathcal{U}_p) = \bigoplus_{\ell=0}^{\infty} L^{Vir}(c_{1, -p}, h_{1, \ell+1}^{1, -p}) \otimes L^{Vir}(c_{1, p}, h_{1, \ell+1}^{1, p}). \end{aligned}$$

The vertex algebra $I_{SL_2}^k$ is called the *chiral universal centralizer* [4, 19], and $H_{DS, \bar{f}}(\mathcal{U}_p)$ is a certain equivariant \mathcal{W} -algebra [4].

Theorem 3.4. *Let $\mathfrak{g}_0 = \mathfrak{sl}_2 \times \mathfrak{sl}_2$. There exist homomorphism of vertex algebras*

$$(3.1) \quad V(\mathfrak{g}_0) = V^{-p-2}(\mathfrak{sl}_2) \otimes V^{-2+\frac{1}{p}}(\mathfrak{sl}_2) \rightarrow \mathcal{V}_p \otimes \Pi(0)^{1/2}$$

such that

$$\mathcal{C}_p = \left(\mathcal{V}_p \otimes \Pi(0)^{1/2}\right)^{\text{int}_{\mathfrak{g}_0}} \cong \bigoplus_{\ell=0}^{\infty} W_p(\ell),$$

where

$$W_p(\ell) = L_{\widehat{\mathfrak{sl}_2}}(-2+p+p\ell)\Lambda_0 + p\ell\Lambda_1 \otimes L_{\widehat{\mathfrak{sl}_2}}\left(-\left(2 - \frac{1}{p} + \ell\right)\Lambda_0 + \ell\Lambda_1\right),$$

and \mathfrak{g}_0 acts on $\mathcal{V}_p \otimes \Pi(0)^{1/2}$ via the homomorphism (3.1). Moreover, \mathcal{C}_p is a simple vertex algebra generated by $W_p(0) + W_p(1)$.

Proof. The decomposition follows directly by applying Corollary 2.2. It remains to prove the simplicity of \mathcal{C}_p . Let

$$V_p(\ell) = L^{Vir}(c_{1, -p}, h_{1, \ell+1}^{1, -p}) \otimes L_{\widehat{\mathfrak{sl}_2}}\left(-\left(2 - \frac{1}{p} + \ell\right)\Lambda_0 + \ell\Lambda_1\right) \subset \mathcal{V}_p.$$

Since \mathcal{U}_p is weakly generated by $A = \mathfrak{sl}_2 \times \mathfrak{sl}_2 + V_{\mathfrak{sl}_2}(\omega_1) \otimes V_{\mathfrak{sl}_2}(\omega_1) \subset L_{-2-1/p}(\mathfrak{sl}_2) \otimes L_{-2+1/p}(\mathfrak{sl}_2) \oplus L_{\widehat{\mathfrak{sl}_2}}(-3 + \frac{1}{p})\Lambda_0 + \Lambda_1 \otimes L_{\widehat{\mathfrak{sl}_2}}(-3 - \frac{1}{p})\Lambda_0 + \Lambda_1$, one can prove (as in [25]) that the vertex algebra \mathcal{V}_p is generated by

$$A^f = \mathbb{C}f \times \mathfrak{sl}_2 \oplus \mathbb{C}v_{-1} \otimes V_{\mathfrak{sl}_2}(\omega_1),$$

where v_{-1} is the lowest weight vector in $V_{\mathfrak{sl}_2}(\omega_1)$. The space A^f generates $L_{c_{1, -p}}^{Vir} \otimes L_{-2+\frac{1}{p}}(\mathfrak{sl}_2)$ -module $V_p(0) \oplus V_p(1)$.

Since the vertex algebra \mathcal{V}_p is simple, we have that $V_p(1) \cdot V_p(\ell) \neq \{0\}$. We claim that

$$(3.2) \quad V_p(1) \cdot V_p(\ell) = V_p(\ell+1) \oplus V_p(\ell-1) \quad (\ell \in \mathbb{Z}_{\geq 1}).$$

By using the fusion rules for $L_{-2+\frac{1}{p}}(\mathfrak{sl}_2)$ -modules, we have that

$$(3.3) \quad V_p(1) \cdot V_p(\ell) \subseteq V_p(\ell+1) \oplus V_p(\ell-1) \quad (\ell \in \mathbb{Z}_{\geq 1}).$$

Assume that there exists minimal $\ell_0 \in \mathbb{Z}_{\geq 1}$ such that the equation (3.2) does not hold. Then we have two possibilities:

- (i) $V_p(1) \cdot V_p(\ell_0) = V_p(\ell_0+1)$; or
- (ii) $V_p(1) \cdot V_p(\ell_0) = V_p(\ell_0-1)$.

If (i) holds, using (3.3) we easily get that $\bigoplus_{\ell=\ell_0}^{\infty} V_p(\ell)$ is a proper ideal in \mathcal{V}_p . This is not possible, since \mathcal{V}_p is simple.

If (ii) holds, we get that $\bigoplus_{\ell=0}^{\ell_0} V_p(\ell)$ is a proper subalgebra of \mathcal{V}_p , which is generated by $V_p(0) + V_p(1)$. This contradicts the fact that $V_p(0) + V_p(1)$ generates \mathcal{V}_p . Therefore (3.2) holds.

We get that

$$\mathcal{C}_p = \bigoplus_{\ell=0}^{\infty} W_p(\ell),$$

where

$$W_p(\ell) = L_{\widehat{\mathfrak{sl}}_2}(-2+p+p\ell)\Lambda_0 + p\ell\Lambda_1 \otimes L_{\widehat{\mathfrak{sl}}_2}(-2-\frac{1}{p}+\ell)\Lambda_0 + \ell\Lambda_1.$$

Next we shall prove that the following fusion holds

$$(3.4) \quad W_p(1) \cdot W_p(\ell) = W_p(\ell+1) \oplus W_p(\ell-1) \quad (\ell \in \mathbb{Z}_{\geq 1}).$$

The fusion rules for $L_{c_{1,p}}^{Vir}$ -modules (2.1) and the decomposition of \mathcal{C}_p implies that

$$(3.5) \quad W_p(1) \cdot W_p(\ell) \subseteq W_p(\ell-1) \oplus W_p(\ell+1).$$

Using Proposition 2.1 again we get that

$$W_p(\ell) = W_p(0)(v_{1,1+\ell p} \otimes v_{\ell} \otimes e^{\frac{p\ell}{2}c}) \subseteq V_p(\ell) \otimes \Pi(0)^{1/2},$$

where v_{ℓ} is a highest weight vector in $L_{\widehat{\mathfrak{sl}}_2}(-2+\ell-\frac{1}{p})\Lambda_0 + \ell\Lambda_1$ and $v_{1,1+p\ell}$ highest weight vector in $L^{Vir}(c_{1,-p}, h_{1,p\ell+1}^{-p,1})$. Moreover, the top component of $V_p(\ell)$ is embedded into $W_p(\ell)$ in the following way

$$(3.6) \quad V_p(\ell)_{top} \cong \mathbb{C}v_{1,1+p\ell} \otimes V_{\widehat{\mathfrak{sl}}_2}(\ell\omega_1) \otimes \mathbb{C}e^{\frac{p\ell}{2}c} \subseteq W_p(\ell).$$

The fusion product $V_p(1) \cdot V_p(\ell) = V_p(\ell-1) + V_p(\ell+1)$, ($\ell \geq 1$) of $V_p(0)$ -submodules of \mathcal{V}_p , implies that

$$v_{1,1+p(\ell\pm 1)} \otimes v_{\ell\pm 1} \in V_p(1)_{top} \cdot V_p(\ell)_{top}.$$

Applying (3.6) we get $W_p(1) \cdot W_p(\ell)$ contains non-zero vectors

$$v_{1,1+p(\ell+1)} \otimes v_{\ell+1} \otimes e^{\frac{(\ell+1)p}{2}c} \in W_p(\ell+1),$$

$$v_{1,1+p(\ell-1)} \otimes v_{\ell-1} \otimes e^{\frac{p(\ell-1)}{2}c} = ((e_2)_{-1})^p \left(v_{1,1+p(\ell-1)} \otimes v_{\ell-1} \otimes e^{\frac{p(\ell-1)}{2}c} \right) \in W_p(\ell-1),$$

where $e_2 = e^c \in W_p(0)$.

This proves that

$$W_p(1) \cdot W_p(\ell) \supseteq W_p(\ell-1) \oplus W_p(\ell+1),$$

and therefore the fusion product (3.4) holds. Since all components $W_p(\ell)$ appearing in the decomposition of \mathcal{C}_p are simple $W_p(0)$ -modules, we get \mathcal{C}_p is a simple vertex algebra.

The fusion product (3.4) also implies that \mathcal{C}_p is generated by $W_p(0) + W_p(1)$. □

Corollary 3.5. *For $p > 2$, \mathcal{C}_p is strongly generated by six generators of weight 1, generators of $\widehat{\mathfrak{sl}}_2 \times \widehat{\mathfrak{sl}}_2$, and $2p+2$ primaries of weight $\frac{p-1}{2}$. Therefore \mathcal{C}_p is a W -algebra of type $(1^6, (\frac{p-1}{2})^{2p+2})$. For $p = 2$, the algebra is of type $(\frac{1}{2})^6$.*

Proof. For the generators of weight $\frac{p-1}{2}$ we choose a basis of the top component of $W(1)^{top}$ of conformal weight $\frac{p-1}{2}$. To see that these are strong generators, it is enough to observe, using the fusion rules in the theorem, that we have

$$W_p(\ell+1)^{top} \subset W_p(1)_{-1}^{top}(W_p(\ell)^{top}).$$

This inductively gives that all $W_p(\ell)$ are strongly generated by $W_p(1)^{top}$ and $\widehat{\mathfrak{sl}}_2 \times \widehat{\mathfrak{sl}}_2$. For $p=2$, we have $\mathcal{C}_2 = M_{(3)}$ so the type is clearly as stated. \square

For even p , vertex algebras \mathcal{C}_p are all $\frac{1}{2}\mathbb{Z}$ -graded, so we can consider the maximal \mathbb{Z} -graded subalgebra \mathcal{C}_p^0 .

Corollary 3.6. *For p even, \mathcal{C}_p^0 is simple and we have a decomposition*

$$\mathcal{C}_p^0 = \bigoplus_{\ell=0}^{\infty} W_p(2\ell).$$

Remark 3.7. *Another approach for constructing \mathcal{C}_p is through its identification as the inverse QHR of the equivariant W -algebra $H_{DS,f}(\mathcal{D}_{SL_2,k}^{ch})$ at positive integer level $k = p-2$. It can be shown that, for $p=3$, this equivariant W -algebra is isomorphic to the Weyl vertex algebra $M_{(2)}$ (see Section 4.1 below), and for $p=4$, it is isomorphic to the affine vertex algebra $L_{-\frac{3}{2}}(\mathfrak{sl}_3)$. In what follows, we present alternative realizations of \mathcal{C}_p for these two special cases, which may be of independent interest.*

4. THE VERTEX ALGEBRA $L_{-\frac{5}{3}}(G_2)$

In this section we shall see that $\mathcal{C}_3 \cong L_{-\frac{5}{3}}(G_2)$ and identify $L_{-\frac{5}{3}}(G_2)$ as a subalgebra of $M_{(2)} \otimes \Pi(0)^{1/2}$.

4.1. Decomposition of $M_{(2)}$. Recall that the Weyl vertex algebra $M_{(m)}$ is generated by bosonic fields a_i^{\pm} ($i=1, \dots, m$) with λ -products

$$[(a_i^{\pm})_{\lambda}(a_j^{\pm})] = 0, \quad [(a_i^+)_{\lambda}(a_j^-)] = \delta_{i,j}.$$

The vertex algebra $M_{(m)}$ has the following \mathbb{Z}_2 -gradation

$$M_{(m)} = M_{(m)}^0 + M_{(m)}^1,$$

where $M_{(m)}^0 = L_{-1/2}(\mathfrak{sp}(2m))$, $M_{(m)}^1 = L_{\widehat{\mathfrak{sp}(2m)}}(-\frac{3}{2}\Lambda_0 + \Lambda_1)$ (cf. [18]).

In this section we prove the complete reducibility of $M_{(2)}$ as $L_{-5}(\mathfrak{sl}_2) \otimes L_{-7}^{Vir}$ -module and find the explicit decomposition. It is interesting that this decomposition gives a vertex-algebraic interpretation of the q -series identity from [26]. Let $(a)_n = (a; q)_n := \prod_{i=0}^{n-1} (1 - aq^i)$, so that

$$(q; q)_{\infty} = \prod_{i \geq 1} (1 - q^i).$$

We shall first identify characters of $L_k(\mathfrak{sl}_2)$ -modules for $k = -5$ and L_c^{Vir} for $c = c_{1,3} = -7$. Then

$$h_{1,3}^{1,3} = \frac{(3i-1)^2 - 4}{12} = \frac{(i-1)(3i+1)}{4}.$$

Lemma 4.1. *Let $\ell \in \mathbb{Z}_{\geq 0}$, $\ell \geq 2$, we have:*

$$\begin{aligned} \text{ch}[L_{\widehat{\mathfrak{sl}}_2}(-5-3\ell)\Lambda_0 + 3\ell\Lambda_1] &= q^{-5/24} \frac{1}{(q)_{\infty}^3} \left((3\ell+1)q^{-\frac{\ell(3\ell+2)}{4}} - (3\ell-1)q^{-\frac{\ell(3\ell-2)}{4}} \right) \\ &\quad + \text{ch}[L_{\widehat{\mathfrak{sl}}_2}(-5-3(\ell-2))\Lambda_0 + 3(\ell-2)\Lambda_1] \\ \text{ch}[L^{Vir}(c_{1,3}, h_{1,3})] &= q^{7/24} \frac{1}{(q)_{\infty}} \left(q^{h_{1,3}^{1,3}} - q^{h_{1,-3}^{1,3}} \right). \end{aligned}$$

In particular,

$$\begin{aligned} \text{ch}[L_{\widehat{\mathfrak{sl}_2}}(-5-6\ell)\Lambda_0 + 6\ell\Lambda_1] &= q^{-5/24} \frac{1}{(q)_\infty^3} \left(\sum_{i=0}^{\ell} (6i+1)q^{-i(3i+1)} - \sum_{i=1}^{\ell} (6i-1)q^{-i(3i-1)} \right) \\ \text{ch}[L_{\widehat{\mathfrak{sl}_2}}(-2-6\ell)\Lambda_0 + (6\ell-3)\Lambda_1] &= q^{-5/24} \frac{1}{(q)_\infty^3} \sum_{i=1}^{\ell} \left((6i-2)q^{-\frac{(2i-1)(6i-1)}{4}} - (6i-4)q^{-\frac{(2i-1)(6i-5)}{4}} \right) \end{aligned}$$

Proof. Similarly as in [12, Section 9]. \square

Next result shows that $M_{(2)}$ is a completely reducible as $L_{-5}(\mathfrak{sl}_2) \otimes L_{c_{1,3}}^{Vir}$ -module.

Theorem 4.2. $M_{(2)}$ is a completely reducible $L_{-5}(\mathfrak{sl}_2) \otimes L_{c_{1,3}}^{Vir}$ -module and the following decomposition holds

$$(4.1) \quad M_{(2)} = \bigoplus_{\ell=0}^{\infty} V(\ell), \quad \text{where}$$

$$(4.2) \quad V(\ell) = L_{\widehat{\mathfrak{sl}_2}}(-5+3\ell)\Lambda_0 + 3\ell\Lambda_1 \otimes L^{Vir}(c_{1,3}, \frac{\ell(3\ell+4)}{4}).$$

The decomposition satisfies the following fusion product:

$$(4.3) \quad V(1) \cdot V(\ell) = V(\ell-1) + V(\ell+1) \quad (\ell \geq 1).$$

Proof. First we notice that there is a homomorphism $\Phi : V^{-5}(\mathfrak{sl}_2) \rightarrow M_{(2)}$ (cf. [15, Section 3.3]) uniquely determined by

$$e \mapsto a_2^+ a_1^- : + \frac{1}{\sqrt{3}} : (a_2^-)^2 : , \quad h \mapsto 3 : a_1^+ a_1^- : + : a_2^+ a_2^- : , \quad f \mapsto 3 : a_1^+ a_2^- : - \frac{1}{\sqrt{3}} : (a_2^+)^2 : .$$

Let $U = \text{Im}(\Phi)$. U is a VOA of central charge $c_1 = \frac{3 \cdot \frac{-5}{3}}{-\frac{5}{3}+2} = -5$. Then $\text{Com}(U, M_{(2)})$ is a vertex operator algebra of central charge $c = c_{1,3} = -7$. We conclude that there is conformal embedding

$$U \otimes L_{c_{1,3}}^{Vir} \hookrightarrow M_{(2)}.$$

For $\ell \in \mathbb{Z}_{\geq 0}$, we define:

$$w_\ell = : (a_1^-)^\ell : .$$

We directly see that w_ℓ is a singular vector and it generates the following highest weight module:

$$\tilde{V}(\ell) := (L_{-5}(\mathfrak{sl}_2) \otimes L_{c_{1,3}}^{Vir}) \cdot w_\ell = \tilde{L}_{\widehat{\mathfrak{sl}_2}}(-5+3\ell)\Lambda_0 + 3\ell\Lambda_1 \otimes \tilde{L}^{Vir}(c_{1,3}, \frac{\ell(3\ell+4)}{4}),$$

where $\tilde{L}_{\widehat{\mathfrak{sl}_2}}(\Lambda)$ is a certain highest weight $L_{-5}(\mathfrak{sl}_2)$ -module of highest weight Λ , and $\tilde{L}^{Vir}(c_{1,3}, h)$ certain highest weight $L_{c_{1,3}}^{Vir}$ -module. (Note that we still don't know if these modules are irreducible). So

$$\sum_{\ell \in \mathbb{Z}_{\geq 0}} \tilde{V}(\ell) \subset M_{(2)}.$$

In order to prove decomposition (4.1) is suffices to prove the character identity

$$(4.4) \quad \text{ch}[M_{(2)}] = \sum_{\ell=0}^{\infty} \text{ch}[L_{\widehat{\mathfrak{sl}_2}}(-5+3\ell)\Lambda_0 + 3\ell\Lambda_1] \cdot \text{ch}[L^{Vir}(c_{1,3}, \frac{\ell(3\ell+4)}{4})],$$

which would also imply that $\tilde{V}(\ell) = V(\ell)$ (simplicity of each component). Using the fusion rules (2.1) for $p = 3$, we conclude that

$$(4.5) \quad V(1) \cdot V(\ell) \subset V(\ell+1) + V(\ell-1).$$

Since $V(1)_{top} = \text{span}_{\mathbb{C}}\{a_1^+, a_1^-, a_2^+, a_2^-\}$, we get

$$(a_1^+)_0 w_\ell = \ell w_{\ell-1}, \quad (a_1^-)_{-1} w_\ell = w_{\ell+1},$$

which implies that $V(\ell+1) + V(\ell-1) \subset V(1) \cdot V(\ell)$. Using (4.5) we get the fusion product (4.3).

So it remains to prove the character identity (4.4). This is proved in Lemma 7.1 in Section 7. The proof follows. \square

4.2. Isomorphism $\mathcal{C}_3 \cong L_{-5/3}(G_2)$. Now we shall apply inverse quantum Hamiltonian reduction procedure as in Section 2 for the level $k = -\frac{5}{3}$.

The vertex algebra $M_{(m)}$ has \mathbb{Z}_2 -twisted module, which we denote by $M_{(m)}^{tw}$ (cf. [?]) and which is generated by cyclic vector $\mathbf{1}^{tw}$. For $m = 2$, we get that $M_{(2)}^{tw}$ is a module for $L_{-5}(\mathfrak{sl}_2) \otimes L_{c_{1,3}}^{Vir}$. Moreover, by direct calculation we can check that $L^{Vir}(c_{1,3}, -\frac{1}{4}) = L^{Vir}(c_{1,3}, h_{2,1}^{1,3}) \cong L_{c_{1,3}}^{Vir} \cdot \mathbf{1}^{tw}$.

We consider the vertex algebra $M_{(2)} \otimes \Pi(0)^{1/2}$ and its \mathbb{Z}_2 -twisted module $M_{(2)}^{tw} \otimes \Pi(0)^{1/2} \cdot e^\nu$. The Virasoro highest weight vector $v^{(k)}$ is now represented by $\mathbf{1}^{tw} \in M_{(2)}^{tw}$. As in (2.5) we have the screening operator

$$(4.6) \quad S = \text{Res}_z \mathcal{Y}(\mathbf{1}^{tw} \otimes e^\nu, z) : M_{(2)} \otimes \Pi(0)^{1/2} \rightarrow M_{(2)}^{tw} \otimes \Pi(0)^{1/2} \cdot e^\nu$$

where \mathcal{Y} is intertwining operator of type

$$\begin{pmatrix} M_{(2)}^{tw} \otimes \Pi(0)^{1/2} \cdot e^\nu \\ M_{(2)}^{tw} \otimes \Pi(0)^{1/2} \cdot e^\nu \quad M_{(2)} \otimes \Pi(0)^{1/2} \end{pmatrix},$$

which is transpose of the vertex operator Y_{tw} who defines the structure of a twisted module on $M_{(2)}^{tw} \otimes \Pi(0)^{1/2} \cdot e^\nu$. Since S is residuum of the intertwining operator among (twisted) $M_{(2)} \otimes \Pi(0)^{1/2}$ -modules, we get:

Lemma 4.3. $\text{Ker}_{M_{(2)} \otimes \Pi(0)^{1/2}} S$ is a vertex subalgebra of $M_{(2)} \otimes \Pi(0)^{1/2}$.

Here is our main theorem:

Theorem 4.4. Let $\mathfrak{g}_0 = \mathfrak{sl}_2 \times \mathfrak{sl}_2$. There exist homomorphism of vertex algebras

$$(4.7) \quad V(\mathfrak{g}_0) = V^{-5}(\mathfrak{sl}_2) \otimes V^{-\frac{5}{3}}(\mathfrak{sl}_2) \rightarrow M_{(2)} \otimes \Pi(0)^{1/2}$$

such that

(1)

$$W = \text{Ker}_{M_{(2)} \otimes \Pi(0)^{1/2}} S = \left(M_{(2)} \otimes \Pi(0)^{1/2} \right)^{\text{int}_{\mathfrak{g}_0}} = \bigoplus_{\ell \in \mathbb{Z}_{\geq 0}} W(\ell)$$

where

$$\begin{aligned} W(\ell) &= L_{\widehat{\mathfrak{sl}_2}}(- (5 + 3\ell)\Lambda_0 + 3\ell\Lambda_1) \otimes \text{Ker}_{L^{Vir}(c_{1,3}, \frac{\ell(3\ell+4)}{4}) \otimes \Pi(0)^{1/2}} S \\ &= L_{\widehat{\mathfrak{sl}_2}}(- (5 + 3\ell)\Lambda_0 + 3\ell\Lambda_1) \otimes L_{\widehat{\mathfrak{sl}_2}}(- (\frac{5}{3} + \ell)\Lambda_0 + \ell\Lambda_1), \end{aligned}$$

S is the screening operator defined by (4.6), and $\left(M_{(2)} \otimes \Pi(0)^{1/2} \right)^{\text{int}_{\mathfrak{g}_0}}$ is the maximal integrable part of $M_{(2)} \otimes \Pi(0)^{1/2}$ with respect to \mathfrak{g}_0 -action induced from the homomorphism (4.7).

2) The decomposition in (1) satisfies the following fusion product:

$$W(1) \cdot W(\ell) = W(\ell-1) + W(\ell+1) \quad (\ell \geq 1).$$

(3) W is a simple vertex algebra.

(4) $W \cong \mathcal{C}_3 \cong L_{-\frac{5}{3}}(G_2)$.

Proof. The assertion (1) follows directly from Proposition 2.1 and Theorem 4.2. The fusion rules for $L_{c_{1,p}}^{Vir}$ -modules (2.1) now implies that

$$(4.8) \quad W(1) \cdot W(\ell) \subset W(\ell - 1) \oplus W(\ell + 1).$$

Using Proposition 2.1 again we get that

$$W(\ell) = W(0)(v_{3\ell} \otimes v_{1,1+\ell} \otimes e^{\frac{\ell}{2}c}) = W(0)(w_\ell \otimes e^{\frac{\ell}{2}c}) \subset V(\ell) \otimes \Pi(0)^{1/2},$$

where w_ℓ is identified with the vector $v_{3\ell} \otimes v_{1,1+\ell}$, such that $v_{3\ell}$ is a highest weight vector in $L_{\widehat{\mathfrak{sl}}_2}(-5 + 3\ell)\Lambda_0 + 3\ell\Lambda_1$ and $v_{1,1+\ell}$ highest weight vector in $L^{Vir}(c_{1,3}, h_{1,\ell+1}^{1,3})$. Now applying the fusion product $V(1) \cdot V(\ell) = V(\ell - 1) + V(\ell + 1)$ ($\ell \geq 1$) of $V(0)$ -submodules of $M_{(2)}$ and using (4.8) we get that the fusion product $W(1) \cdot W(\ell)$ contains non-zero vectors

$$w_{\ell+1} \otimes e^{\frac{\ell+1}{2}c} \in W(\ell + 1), \quad w_{\ell-1} \otimes e^{\frac{\ell+1}{2}c} = (e_2)_{-1}(w_{\ell-1} \otimes e^{\frac{\ell-1}{2}c}) \in W(\ell - 1),$$

where $e_2 = e^c \in W(0)$.

Indeed, we have

$$(w_1 \otimes e^{\frac{c}{2}})_{-1}(w_\ell \otimes e^{\frac{\ell}{2}c}) = (w_{\ell+1} \otimes e^{\frac{\ell+1}{2}c}) \in W(1) \cdot W(\ell).$$

Next we notice that $a_1^+ = \mu_1 f(0)^3 w_1$, for certain non-zero constant μ_1 , so

$$a_1^+ \otimes e^{\frac{c}{2}} = \mu_1 f(0)^3 w_1 \otimes e^{\frac{c}{2}} \in W(1).$$

This implies

$$(a_1^+ \otimes e^{\frac{c}{2}})_0(w_\ell \otimes e^{\frac{\ell}{2}c}) = \ell(w_{\ell-1} \otimes e^{\frac{\ell+1}{2}c}) \in W(1) \cdot W(\ell).$$

This proves that

$$W(1) \cdot W(\ell) \supset W(\ell - 1) \oplus W(\ell + 1),$$

and therefore the assertion (2) holds. Since all components $W(\ell)$ are simple $W(0)$ -modules, the assertion (2) implies that W is a simple vertex algebra.

Using (2) and (3) we see that W is a simple vertex algebra generated by $W(0) + W(1)$. More precisely, W is generated by the generators $L_{-5/3}(\mathfrak{sl}_2) \otimes L_{-5}(\mathfrak{sl}_2)$ which span $\mathfrak{g}_0 = \mathfrak{sl}_2 \times \mathfrak{sl}_2$ and the top component of $L_{\widehat{\mathfrak{sl}}_2}(-8\Lambda_0 + 3\Lambda_1) \otimes L_{\widehat{\mathfrak{sl}}_2}(-\frac{8}{3}\Lambda_0 + \Lambda_1)$ which is isomorphic to $\mathfrak{g}_1 = V_{\mathfrak{sl}_2}(3\omega_1) \otimes V_{\mathfrak{sl}_2}(\omega_1)$.

This gives that the weight one space of V with bracket $[x, y] = x_{(0)}y$ is the Lie algebra $\mathfrak{g} = \mathfrak{g}_0 + \mathfrak{g}_1 \cong G_2$. \square

Remark 4.5. Note that in [15] the authors also found a homomorphism $\Phi_1 : V^{-5/3}(G_2) \rightarrow M_{(2)} \otimes \Pi(0)^{1/2}$, but they don't provide the proof of simplicity of $\text{Im}(\Phi_1)$ or decomposition of conformal embedding.

5. THE VERTEX ALGEBRA \mathcal{C}_4 AS INVERSE QHR OF $L_{-3/2}(\mathfrak{sl}_3)$

5.1. The decomposition of $L_{-3/2}(\mathfrak{sl}_3)$ as $L_{-6}(\mathfrak{sl}_2) \otimes L_{c_{1,4}}^{Vir}$ -module. We shall identify characters of $L_k(\mathfrak{sl}_2)$ -modules for $k = -6$ and L_c^{Vir} for $c = c_{1,4} = -\frac{25}{2}$.

Then

$$h_{1,i}^{1,4} = \frac{(4i-1)^2 - 9}{16} = \frac{(i-1)(2i+1)}{2}.$$

Similarly as in [12, Section 9], we have

Lemma 5.1. Let $\ell \in \mathbb{Z}_{\geq 0}$, $\ell \geq 2$, we have:

$$\begin{aligned} \text{ch}[L_{\widehat{\mathfrak{sl}}_2}(-6 - 4\ell)\Lambda_0 + 4\ell\Lambda_1] &= q^{-3/16} \frac{1}{(q)_\infty^3} \left((4\ell + 1)q^{-\frac{\ell(2\ell+1)}{2}} - (4\ell - 1)q^{-\frac{\ell(2\ell-1)}{2}} \right) \\ &\quad + \text{ch}[L_{\widehat{\mathfrak{sl}}_2}(-6 - 4(\ell - 2))\Lambda_0 + 4(\ell - 2)\Lambda_1], \\ \text{ch}[L^{Vir}(c_{1,4}, h_{i,1})] &= q^{25/48} \frac{1}{(q)_\infty} \left(q^{h_{1,i}^{1,4}} - q^{h_{1,-i}^{1,4}} \right). \end{aligned}$$

In particular,

$$\begin{aligned} \text{ch}[L_{\widehat{\mathfrak{sl}}_2}(-6-8\ell)\Lambda_0 + 8\ell\Lambda_1] &= q^{-3/16} \frac{1}{(q)_\infty^3} \left(\sum_{i=0}^{\ell} (8i+1)q^{-i(4i+1)} - \sum_{i=1}^{\ell} (8i-1)q^{-i(4i-1)} \right) \\ \text{ch}[L_{\widehat{\mathfrak{sl}}_2}(-2-8\ell)\Lambda_0 + (8\ell-4)\Lambda_1] &= q^{-3/16} \frac{1}{(q)_\infty^3} \sum_{i=1}^{\ell} \left((8i-3)q^{-\frac{(2i-1)(4i-1)}{2}} - (8i-5)q^{-\frac{(2i-1)(4i-3)}{2}} \right) \end{aligned}$$

Consider now Lie algebra \mathfrak{sl}_3 with standard generators

$$e_{\alpha_1}, e_{\alpha_2}, e_{\theta}, f_{\alpha_1}, f_{\alpha_2}, f_{\theta}, h_{\alpha_1}, h_{\alpha_2}$$

and the simple affine vertex algebra $L_{-3/2}(\mathfrak{sl}_3)$ of level $-\frac{3}{2}$.

Theorem 5.2. $L_{-3/2}(\mathfrak{sl}_3)$ is completely reducible as an $L_{-6}(\mathfrak{sl}_2) \otimes L_{c_{1,4}}^{Vir}$ -module and the following decomposition holds

$$(5.1) \quad L_{-3/2}(\mathfrak{sl}_3) = \bigoplus_{\ell=0}^{\infty} \left(L_{\widehat{\mathfrak{sl}}(2)}(-6+4\ell)\Lambda_0 + 4\ell\Lambda_1 \right) \otimes L^{Vir}(c_{1,4}, \frac{\ell(2\ell+3)}{2}).$$

Proof. There is embedding $L_{-6}(\mathfrak{sl}_2) \hookrightarrow L_{-3/2}(\mathfrak{sl}_3)$, given by

$$e \mapsto \sqrt{2}(e_{\alpha_1} + e_{\alpha_2}), \quad f \mapsto \sqrt{2}(f_{\alpha_1} + f_{\alpha_2}), \quad h = 2h_{\alpha_1} + 2h_{\alpha_2}.$$

Since the commutant $\text{Com}(L_{-6}(\mathfrak{sl}_2), L_{-3/2}(\mathfrak{sl}_3))$ has the central charge

$$c = -8 - \frac{9}{2} = -\frac{25}{2} = c_{1,4},$$

we conclude that we have an conformal embedding $L_{-6}(\mathfrak{sl}_2) \otimes L_{c_{1,4}}^{Vir} \hookrightarrow L_{-3/2}(\mathfrak{sl}_3)$, so we can consider $L_{-3/2}(\mathfrak{sl}_3)$ as an $L_{-6}(\mathfrak{sl}_2) \otimes L_{c_{1,4}}^{Vir}$ -module.

The proof now follow from the q -series identity given in Lemma 7.2, using the same arguments as in the proof of Theorem 4.2. \square

The inverse reduction functor gives that for each $\ell \in \mathbb{Z}_{\geq 0}$ we have

$$L_{\widehat{\mathfrak{sl}}_2}(-\frac{7}{4} + \ell)\Lambda_0 + \ell\Lambda_1 = \left(L^{Vir}(c_{1,4}, \frac{\ell(2\ell+3)}{2}) \otimes \Pi(0)^{1/2} \right)^{\mathfrak{sl}_2}.$$

Using the simplicity of $L_{-3/2}(\mathfrak{sl}_3)$ and the same arguments as in the proof of Theorem 4.4 we get:

Theorem 5.3. Let $\mathfrak{g}_0 = \mathfrak{sl}_2 \times \mathfrak{sl}_2$. There exist homomorphism of vertex algebras

$$(5.2) \quad V(\mathfrak{g}_0) = V^{-6}(\mathfrak{sl}_2) \otimes V^{-\frac{7}{4}}(\mathfrak{sl}_2) \rightarrow L_{-3/2}(\mathfrak{sl}_3) \otimes \Pi(0)^{1/2}$$

such that

$$\begin{aligned} \mathcal{C}_4 &= \left(L_{-3/2}(\mathfrak{sl}_3) \otimes \Pi(0)^{1/2} \right)^{\text{int}_{\mathfrak{g}_0}} \\ &= \bigoplus_{\ell=0}^{\infty} \left(L_{\widehat{\mathfrak{sl}}_2}(-6+4\ell)\Lambda_0 + 4\ell\Lambda_1 \right) \otimes L_{\widehat{\mathfrak{sl}}_2}(-\frac{7}{4} + \ell)\Lambda_0 + \ell\Lambda_1, \end{aligned}$$

where the Lie algebra \mathfrak{g}_0 acts on $L_{-3/2}(\mathfrak{sl}_3) \otimes \Pi(0)^{1/2}$ via the homomorphism (5.2). Moreover, \mathcal{C}_4 is a simple vertex algebra.

Remark 5.4. As in Section 4.2, one can see that there exist twisted $L_{-3/2}(\mathfrak{sl}_3)$ -module $L_{-3/2}^{tw}(\mathfrak{sl}_3)$ generated by twisted highest weight vector $\mathbf{1}^{tw}$ such that $L_{c_{1,4}}^{Vir} \mathbf{1}^{tw} = L^{Vir}(c_{1,4}, h_{2,1}^{1,4}) \mathbf{1}^{tw}$ and that the screening operator (2.5) for $k = -7/4$ can be extended to a operator

$$S = \text{Res}_z \mathcal{Y}(\mathbf{1}^{tw} \otimes e^\nu, z) : L_{-3/2}(\mathfrak{sl}_3) \otimes \Pi(0)^{1/2} \rightarrow L_{-3/2}^{tw}(\mathfrak{sl}_3) \otimes \Pi(0)^{1/2} \cdot e^\nu$$

such that $\mathcal{C}_4 = \text{Ker}_{L_{-3/2}(\mathfrak{sl}_3) \otimes \Pi(0)^{1/2}} S$.

6. REALIZATION OF \mathcal{C}_p AS AFFINE W -ALGEBRAS: THE CASES $p = 4, 5$

We identified \mathcal{C}_3 as affine vertex algebra $L_{-\frac{5}{3}}(G_2)$. In this section we shall see that \mathcal{C}_p for $p = 4, 5$ admits realization as simple affine W -algebras. Moreover, we shall see that in these cases \mathcal{C}_p are obtained by QHR from admissible affine vertex algebras, which would imply that they are quasi-lisse.

6.1. Affine W -algebras. Let us briefly recall construction of affine W -algebra associated to a nilpotent element f in a simple Lie algebra \mathfrak{g} . Given nilpotent f , we denote by (e, f, h) an \mathfrak{sl}_2 -triple with a good grading, so that in particular $\mathfrak{g} = \bigoplus_{j \in \frac{1}{2}\mathbb{Z}} \mathfrak{g}_j$, where $\mathfrak{g}_j = \{y : [x, y] = jy\}$ and let $x = \frac{h}{2}$. We denote by $\mathcal{W}^k(\mathfrak{g}, f)$ the universal affine W -algebra associated with \mathfrak{g} and such f , where $\mathcal{W}^k(\mathfrak{g}, f) = H_{DS, f}^0(V^k(\mathfrak{g}))$. This vertex algebra is conformal provided that $k \neq -h^\vee$ and its central charge is given by [25]

$$c = \frac{k \dim(\mathfrak{g})}{k + h^\vee} - 12k|x|^2 - \sum_{\alpha \in S_+} (12m_\alpha^2 - 12m_\alpha + 2) - \frac{1}{2} \mathfrak{g}_{1/2},$$

where S_+ is a basis of $\mathfrak{g}_{>0}$ and m_α is the x -eigenvalue of α , and $(\cdot|\cdot) = \frac{1}{2h^\vee} \kappa(\cdot|\cdot)$ denotes the normalized Killing form. The universal affine W -algebra $\mathcal{W}^k(\mathfrak{g}, f)$ is strongly generated by the fields corresponding to homogeneous basis elements of $\mathfrak{g}_{\leq 0}^f = \mathfrak{g}^f \cap \mathfrak{g}_{\leq 0}$ inside the centralizer \mathfrak{g}^f (cf. [25]).

6.2. Affine W -algebra $\mathcal{W}_k(F_4, A_1 + \tilde{A}_1)$. In this section we shall prove that \mathcal{C}_4 is quasi-lisse and is realized as the affine W -algebra $\mathcal{W}_k(F_4, A_1 + \tilde{A}_1)$ with $k = -\frac{23}{4}$. Here we are using classification of nilpotent orbits and at the same time we slightly abuse notation for W -algebras, indicating the nilpotent orbit $A_1 + \tilde{A}_1$ instead of a nilpotent element $f_{A_1 + \tilde{A}_1}$ that belongs to it.

Let \mathfrak{g} be the exceptional simple Lie algebra of type F_4 . We shall now study vertex algebra $\mathcal{W}_k(\mathfrak{g}, A_1 + \tilde{A}_1)$ at admissible level $k = -\frac{23}{4}$.

Take nilpotent element f in the Lie algebra of type $A_1 + \tilde{A}_1$, and corresponding \mathfrak{sl}_2 -triple (e, f, x) . Then $\text{ad}(x)$ defines the following gradation of \mathfrak{g}^f :

$$\mathfrak{g}^f = \mathfrak{g}_{-3/2}^f \oplus \mathfrak{g}_{-1}^f \oplus \mathfrak{g}_{-1/2}^f \oplus \mathfrak{g}^{\mathfrak{h}}$$

such that $\mathfrak{g}^{\mathfrak{h}} = \mathfrak{sl}_2 \times \mathfrak{sl}_2$ (here $\mathfrak{g}^{\mathfrak{h}} = \mathfrak{g}_0 \cap \mathfrak{g}^f$) and as $\mathfrak{g}^{\mathfrak{h}}$ -modules:

$$\begin{aligned} \mathfrak{g}_{-3/2}^f &= V_{\mathfrak{sl}_2}(\omega_1) \otimes V_{\mathfrak{sl}_2}(0) \\ \mathfrak{g}_{-1}^f &= V_{\mathfrak{sl}_2}(0) \otimes V_{\mathfrak{sl}_2}(4\omega_1) + V_{\mathfrak{sl}_2}(0) \otimes V_{\mathfrak{sl}_2}(0), \\ \mathfrak{g}_{-1/2}^f &= V_{\mathfrak{sl}_2}(\omega_1) \otimes V_{\mathfrak{sl}_2}(4\omega_1). \end{aligned}$$

Moreover, there is a vertex algebra homomorphism

$$\Phi : V^{8k+40}(\mathfrak{sl}_2) \otimes V^{k+4}(\mathfrak{sl}_2) \hookrightarrow \mathcal{W}^k(F_4, A_1 + \tilde{A}_1).$$

Applying [25], we get that $\mathcal{W}^k(F_4, A_1 + \tilde{A}_1)$ is strongly generated by

- the generators of $\text{Im}(\Phi)$, i.e. by $J^{\{x\}}$, $x \in \mathfrak{g}^{\mathfrak{h}}$,
- the Virasoro field L of central charge $c = -\frac{6(3+k)(14+3k)}{9+k}$,
- 10-generators of conformal weight $3/2$, which span $\mathfrak{g}^{\mathfrak{h}}$ -module $V_{\mathfrak{sl}_2}(\omega_1) \otimes V_{\mathfrak{sl}_2}(4\omega_1)$, these generators can be taken to be primary for L and $\text{Im}(\Phi)$, which span $\mathfrak{g}^{\mathfrak{h}}$ -module,
- 6 generators of conformal weight 2 which consists of L and 5 primary vectors for L and $\text{Im}(\Phi)$, which span $\mathfrak{g}^{\mathfrak{h}}$ -module $V_{\mathfrak{sl}_2}(0) \otimes V_{\mathfrak{sl}_2}(4\omega_1) \oplus V_{\mathfrak{sl}_2}(0) \otimes V_{\mathfrak{sl}_2}(0)$. Note that the multiplicity of the trivial representation is one.
- two generators of conformal weight $5/2$, which span $\mathfrak{g}^{\mathfrak{h}}$ -module $V_{\mathfrak{sl}_2}(\omega_1) \otimes V_{\mathfrak{sl}_2}(0)$.

Let $\mathcal{W}_k(F_4, A_1 + \tilde{A}_1)$ be the simple quotient of $\mathcal{W}^k(F_4, A_1 + \tilde{A}_1)$, and let $f : \mathcal{W}^k(F_4, A_1 + \tilde{A}_1) \rightarrow \mathcal{W}_k(F_4, A_1 + \tilde{A}_1)$ be the quotient homomorphism. Then we have a homomorphism

$$\Phi_1 = f \circ \Phi : V^{8k+40}(\mathfrak{sl}_2) \otimes V^{k+4}(\mathfrak{sl}_2) \hookrightarrow \mathcal{W}_k(F_4, A_1 + \tilde{A}_1).$$

Central charge of the vertex algebra $\text{Im}(\Phi_1)$ is given by

$$c_{\text{sug}} = \frac{3(8k+40)}{8k+42} + \frac{3(k+4)}{k+6}.$$

By applying [14, Theorem 3.1] we get:

- The embedding $\text{Im}(\Phi_1) \hookrightarrow \mathcal{W}_k(F_4, A_1 + \tilde{A}_1)$ is conformal if and only if $c = c_{\text{sug}}$, i.e., when $k \in \{-\frac{23}{4}, -4, -5, -\frac{9}{2}\}$.

Remark 6.1. *This result also appears in a much more general setup in [11]. Here we include details because of reader convenience.*

Now we shall consider the case of conformal level $k = -\frac{23}{4}$. Then $\text{Im}(\Phi_1)$ is isomorphic to the simple affine vertex algebra $L_{-6}(\mathfrak{sl}_2) \otimes L_{-\frac{7}{4}}(\mathfrak{sl}_2)$, and we have conformal embeddings

$$L_{-6}(\mathfrak{sl}_2) \otimes L_{-\frac{7}{4}}(\mathfrak{sl}_2) \hookrightarrow \mathcal{W}_{-\frac{23}{4}}(F_4, A_1 + \tilde{A}_1).$$

Proposition 6.2. • $\mathcal{W}_{-\frac{23}{4}}(F_4, A_1 + \tilde{A}_1)$ is strongly generated by the generators of $\text{Im}(\Phi_1)$ and by 10 generators of conformal weight $\frac{3}{2}$. In particular, $k = -\frac{23}{4}$ is not a collapsing level.

Proof. By applying again results from [14] we get that in $\mathcal{W}_{-\frac{23}{4}}(F_4, A_1 + \tilde{A}_1)$ we have:

- $L = L_{\text{sug}}$, where L_{sug} is the Sugawara Virasoro vector of $L_{-6}(\mathfrak{sl}_2) \otimes L_{-\frac{7}{4}}(\mathfrak{sl}_2)$,
- If G is a generator of $\mathcal{W}_{-\frac{23}{4}}(F_4, A_1 + \tilde{A}_1)$ of conformal weight 2 and $\frac{5}{2}$, $G \notin \mathbb{C}L$, when $G = 0$ in $\mathcal{W}_{-\frac{23}{4}}(F_4, A_1 + \tilde{A}_1)$.

Since the conformal weight $3/2$ space is irreducible \mathfrak{g}^{\natural} -module, we have two possibilities

- all generators of conformal weight $3/2$ vanish in $\mathcal{W}_{-\frac{23}{4}}(F_4, A_1 + \tilde{A}_1)$;
- all generators of conformal weight $3/2$ are non-zero in $\mathcal{W}_{-\frac{23}{4}}(F_4, A_1 + \tilde{A}_1)$.

The first possibility leads to the conclusion that $\mathcal{W}_{-\frac{23}{4}}(F_4, A_1 + \tilde{A}_1) = L_{-6}(\mathfrak{sl}_2) \otimes L_{-\frac{7}{4}}(\mathfrak{sl}_2)$, but this contradicts Proposition 6.3 below.

Therefore (ii) holds. The proof follows. \square

Next we have the following result:

Proposition 6.3. *The vertex algebra $\mathcal{W}_{-\frac{23}{4}}(F_4, A_1 + \tilde{A}_1)$ is quasi-lisse.*

Proof. Since $L_{-23/4}(\mathfrak{f}_4)$ is quasi-lisse, using results of Arakawa et al. [5] and Proposition 6.3 in [6], it suffices to check that the condition $f_{A_1 + \tilde{A}_1}$ is inside the closure of the nilpotent orbit of \mathbb{O}_q (in our case $q = 4$). This is equivalent to $f_{A_1 + \tilde{A}_1} \in \{x \in \mathfrak{f}_4 : \pi_{\theta_s}(x)^4 = 0\}$ where π_{θ_s} is the irreducible finite-dimensional representation of \mathfrak{f}_4 with highest weight θ_s where θ_s is the highest short root. We have $\theta_s = 2\alpha_1 + 3\alpha_2 + 4\alpha_3 + 2\alpha_4 = \omega_1$ and thus $\pi_{\theta_s} = \pi_{\omega_1}$ is the fundamental 26-dimensional representation. Explicit computation shows that $\pi_{\omega_1}(f_{A_1 + \tilde{A}_1})^4 = 0$ as required. \square

Next we want to identify $\mathcal{W}_{-\frac{23}{4}}(F_4, A_1 + \tilde{A}_1)$ with the vertex algebra V from previous section. We have that both algebras are simple, and extensions of the vertex algebra $L_{-6}(\mathfrak{sl}_2) \otimes L_{-\frac{7}{4}}(\mathfrak{sl}_2)$ by weight $\frac{3}{2}$ generators.

In [7, Theorem 3.1], the authors considered the problem of uniqueness of minimal W -algebras, which are of the same type as simple W -algebra $\mathcal{W}_{-\frac{23}{4}}(F_4, A_1 + \tilde{A}_1)$. Using the same proof as in [7] we can prove:

Proposition 6.4. *Assume that $\mathcal{A} = \bigoplus_{\ell \in \frac{1}{2}\mathbb{Z}} \mathcal{A}(\ell)$ is another simple, $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded conformal vertex algebra, such that*

- (1) *There is a conformal embeddings $L_{-6}(\mathfrak{sl}_2) \otimes L_{-\frac{7}{4}}(\mathfrak{sl}_2) \hookrightarrow \mathcal{A}$.*
- (2) *$\mathcal{A}(\frac{1}{2}) = \{0\}$, and \mathcal{A} is strongly generated by $\mathcal{A}(1) + \mathcal{A}(3/2)$ and $\mathcal{A}(1) \cong \mathfrak{g}^{\natural}$, $\mathcal{A}(\frac{3}{2}) \cong V_{A_1}(\omega_1) \otimes V_{A_1}(4\omega_1)$ as \mathfrak{g}^{\natural} -modules.*
- (3) *The weight $3/2$ fields in both \mathcal{A} and $\mathcal{W}_{-\frac{23}{4}}(F_4, A_1 + \tilde{A}_1)$ carry the same OPE with the Virasoro field and weight one fields.*

Then $\mathcal{A} \cong \mathcal{W}_{-\frac{23}{4}}(F_4, A_1 + \tilde{A}_1)$.

It is easy to check that our vertex algebra V satisfies the same conditions as required for the vertex algebra \mathcal{A} above. Thus we get:

Theorem 6.5. *We have*

$$V \cong \mathcal{W}_{-\frac{23}{4}}(F_4, A_1 + \tilde{A}_1).$$

6.3. The vertex algebra $\mathcal{W}_{-30+31/5}(E_8, A_4 + A_2)$. Now we shall prove that $\mathcal{C}_p \cong \mathcal{W}_{-30+31/5}(E_8, A_4 + A_2)$ for $p = 5$. As a consequence, we shall get that \mathcal{C}_5 is quasi-lisse. The approach is similar to those in Subsection 6.2 above, so we omit some details.

Let now \mathfrak{g} denoted the exceptional Lie algebra of type E_8 . Take nilpotent element f in the Lie algebra of type $A_4 + A_2$, and corresponding \mathfrak{sl}_2 -triple (e, f, x) .

One can show that there is a vertex algebra homomorphism

$$\Phi : V^{15k+350}(\mathfrak{sl}_2) \otimes V^{k+22}(\mathfrak{sl}_2) \hookrightarrow \mathcal{W}^k(\mathfrak{g}, f).$$

Applying [25] (details will be also presented in [11]) we get that $\mathcal{W}^k(\mathfrak{g}, f)$ is strongly generated by

- the generators of $\text{Im}(\Phi)$, i.e. by $J^{\{x\}}$, $x \in \mathfrak{g}^{\natural}$,
- the Virasoro field L of central charge $c = -\frac{6(12490+1095k+24k^2)}{30+k}$,
- 18-generators of conformal weight 2,
- 14 generators of conformal weight 3,
- 13 generators of conformal weight 4,
- 3 generators of conformal weight 5.

Let $\mathcal{W}_k(\mathfrak{g}, f)$ be the simple quotient of $\mathcal{W}^k(\mathfrak{g}, f)$, and let $\pi : \mathcal{W}^k(\mathfrak{g}, f) \rightarrow \mathcal{W}_k(\mathfrak{g}, f)$ be the quotient homomorphism. Then we have a homomorphism

$$\Phi_1 = \pi \circ \Phi : V^{15k+350}(\mathfrak{sl}_2) \otimes V^{k+22}(\mathfrak{sl}_2) \rightarrow \mathcal{W}_k(\mathfrak{g}, f).$$

By applying [14, Theorem 3.1] we get that the embedding $\text{Im}(\Phi_1) \hookrightarrow \mathcal{W}_k(\mathfrak{g}, f)$ is conformal for $k = -\frac{119}{5}$. Then $\text{Im}(\Phi_1)$ is isomorphic to the simple affine vertex algebra $L_{-7}(\mathfrak{sl}_2) \otimes L_{-\frac{9}{5}}(\mathfrak{sl}_2)$, and we have the conformal embedding

$$L_{-7}(\mathfrak{sl}_2) \otimes L_{-\frac{9}{5}}(\mathfrak{sl}_2) \hookrightarrow \mathcal{W}_k(\mathfrak{g}, f).$$

Remark 6.6. *This result also will appear in [11] in more general context, where all conformal embeddings of affine vertex algebra to affine W -algebras of type E will be classified.*

Proposition 6.7. *Let $k = -30 + \frac{31}{5} = -\frac{119}{5}$. Then $\mathcal{W}_k(\mathfrak{g}, f)$ is strongly generated by the generators of $\text{Im}(\Phi_1)$ and by 12 generators of conformal weight 2.*

Proof. By applying again results from [14] we get that in $\mathcal{W}_k(\mathfrak{g}, f)$ we have:

- $L = L_{\text{Sug}}$, where L_{Sug} is the Sugawara Virasoro vector of $L_{-7}(\mathfrak{sl}_2) \otimes L_{-\frac{9}{5}}(\mathfrak{sl}_2)$.
- If G is a generator of $\mathcal{W}^k(\mathfrak{g}, f)$ of conformal weight 3, 4 or 5 then $G = 0$ in $\mathcal{W}_k(\mathfrak{g}, f)$.

Since the conformal weight 2 consist of $\mathcal{C}L + U$, where U is an irreducible \mathfrak{g}^{\natural} -module, we have two possibilities

- (i) all generators from U vanish in $\mathcal{W}_k(\mathfrak{g}, f)$;

(ii) all generators from U are non-zero in $\mathcal{W}_k(\mathfrak{g}, f)$.

The first possibility leads to the conclusion that $\mathcal{W}_k(\mathfrak{g}, f) = L_{-7}(\mathfrak{sl}_2) \otimes L_{-\frac{9}{5}}(\mathfrak{sl}_2)$, but this contradicts Proposition 6.8 below. Therefore (ii) holds. The proof follows. \square

Next we have the following result:

Proposition 6.8. *The vertex algebra $\mathcal{W}_k(\mathfrak{g}, f)$ is quasi-lisse.*

Proof. Since $L_k(E_8)$ is quasi-lisse, using results of Arakawa et al. [5] and Proposition 6.3 in [6], it suffices to check that the condition f is inside the closure of the nilpotent orbit of \mathbb{O}_q (in our case $q = 5$). This follows using explicit computation as in Proposition 6.3. \square

Finally, we identify $\mathcal{W}_k(E_8, f)$ with the vertex algebra \mathcal{C}_5 . We know that they are both algebras are simple, and are extensions of the vertex algebra $L_{-7}(\mathfrak{sl}_2) \otimes L_{-\frac{9}{5}}(\mathfrak{sl}_2)$ and weight 2 generators. By using similar arguments as in the proof of Theorem 6.5 we get:

Theorem 6.9. *We have an isomorphism of vertex algebras*

$$\mathcal{C}_5 \cong \mathcal{W}_k(\mathfrak{g}, f).$$

7. PROOF OF SOME CHARACTER IDENTITIES

Lemma 7.1. *The following q -series identity holds:*

$$(7.1) \quad \text{ch}[M_{(2)}] = \sum_{\ell=0}^{\infty} \text{ch}[L_{\widehat{\mathfrak{sl}}_2}(-5 + 3\ell)\Lambda_0 + 3\ell\Lambda_1] \cdot \text{ch}[L^{Vir}(c_{1,3}, \frac{\ell(3\ell+4)}{4})].$$

Proof. Basic manipulation with q -series (here $c = -2$) gives

$$\begin{aligned} \text{ch}[M_{(2)}] &= q^{-c/24} \prod_{n=1}^{\infty} \frac{1}{(1 - q^{n-1/2})^4} = q^{1/12} \prod_{n=1}^{\infty} (1 - q^{n-1/2})^{-4} \\ &= q^{1/12} \frac{1}{(q^4_{\infty})} \left(\frac{(q^2_{\infty})}{(q^{1/2}_{\infty})} \right)^4 = q^{1/12} \frac{1}{(q^4_{\infty})} \Delta(q^{1/2})^4, \end{aligned}$$

where

$$\Delta(q) = \sum_{n \in \mathbb{Z}_{\geq 0}} q^{n(n+1)/2} = \frac{(q^2_{\infty})^2}{(q)_{\infty}}.$$

Using Lemma 5.1 we get that the q -character of the right side of (5.1) is given by $\frac{q^{1/12}}{(q^4_{\infty})} \mathcal{B}[q]$ where

$$\begin{aligned} \mathcal{B}[q] &:= \sum_{\ell=0}^{\infty} \sum_{i=0}^{\ell} (6i+1) \left(q^{\ell(3\ell+2)-i(3i+1)} - q^{(\ell+1)(3\ell+1)-i(3i+1)} \right) \\ &+ \sum_{\ell=1}^{\infty} \sum_{i=1}^{\ell} (6i-1) \left(-q^{\ell(3\ell+2)-i(3i-1)} + q^{(\ell+1)(3\ell+1)-i(3i-1)} \right) \\ &+ \sum_{\ell=1}^{\infty} \sum_{i=1}^{\ell} (6i-2) \left(q^{\frac{(2\ell-1)(6\ell+1)-(2i-1)(6i-1)}{4}} - q^{\frac{(2\ell+1)(6\ell-1)-(2i-1)(6i-1)}{4}} \right) \\ &+ \sum_{\ell=1}^{\infty} \sum_{i=1}^{\ell} (6i-4) \left(-q^{\frac{(2\ell-1)(6\ell+1)-(2i-1)(6i-5)}{4}} + q^{\frac{(2\ell+1)(6\ell-1)-(2i-1)(6i-5)}{4}} \right). \end{aligned}$$

Therefore the proof of the theorem is now reduced to the following identity:

$$\frac{(q^2_{\infty})^8}{(q^{1/2}_{\infty})^4} = \Delta(q^{1/2})^4 = \mathcal{B}[q]$$

By [26] we have that

$$\Delta(q)^4 = \sum_{j,k=0}^{\infty} (2k+1)q^{\frac{(2j+1)(2k+1)-1}{2}}.$$

Next we organize pairs $(j, k) \in \mathbb{Z}_{\geq 0}^2$, such that $k \equiv 0, 2 \pmod{3}$, into eight congruent classes. Inspection shows that each such pair satisfies one and only one condition as in the table below.

	Condition	$\ell \in \mathbb{Z}$	$i \in \mathbb{Z}$	Relations
(1)	$\frac{j}{2} - \frac{k}{6} \in \mathbb{Z}_{\geq 0}$	$\frac{k}{6} + \frac{j}{2}$	$\frac{j}{2} - \frac{k}{6}$	$k = 3(\ell - i), j = \ell + i$ $6i + 1 = 3j - k + 1$
(2)	$\frac{j}{2} - \frac{k}{6} \in \mathbb{Z}_{< 0}$	$\frac{k}{6} + \frac{j}{2}$	$\frac{k}{6} - \frac{j}{2}$	$k = 3(\ell + i), j = \ell - i$ $-(6i - 1) = 3j - k + 1$
(3)	$\frac{k}{6} - \frac{j}{2} + \frac{1}{2} \in \mathbb{Z}_{\geq 0}$	$\frac{k}{6} + \frac{j+1}{2}$	$\frac{k}{6} - \frac{j-1}{2}$	$k = 3(\ell + i - 1), j = \ell - i$ $-(6i - 4) = 3j - k + 1$
(4)	$\frac{k}{6} - \frac{j}{2} + \frac{1}{2} \in \mathbb{Z}_{< 0}$	$\frac{k}{6} + \frac{j+1}{2}$	$\frac{j+1}{2} - \frac{k}{6}$	$k = 3(\ell - i), j = \ell + i - 1$ $6i - 2 = 3j - k + 1$
(5)	$\frac{k}{6} - \frac{j}{2} - \frac{1}{3} \in \mathbb{Z}_{\geq 0}$	$\frac{k+1}{6} + \frac{j-1}{2}$	$\frac{k+1}{6} - \frac{j+1}{2}$	$k = 3(\ell + i) + 2, j = \ell - i$ $-(6i + 1) = 3j - k + 1$
(6)	$\frac{k}{6} - \frac{j}{2} - \frac{1}{3} \in \mathbb{Z}_{< 0}$	$\frac{k+1}{6} + \frac{j-1}{2}$	$-\frac{k+1}{6} + \frac{j+1}{2}$	$k = 3(\ell - i) + 2, j = \ell + i$ $6i - 1 = 3j - k + 1$
(7)	$\frac{j}{2} - \frac{k}{6} + \frac{5}{6} \in \mathbb{Z}_{\geq 0}$	$\frac{k+1}{6} + \frac{j}{2}$	$\frac{j}{2} - \frac{k+1}{6} + 1$	$k = 3(\ell - i) + 2, j = \ell + i - 1$ $6i - 4 = 3j - k + 1$
(8)	$\frac{j}{2} - \frac{k}{6} + \frac{5}{6} \in \mathbb{Z}_{< 0}$	$\frac{k+1}{6} + \frac{j}{2}$	$\frac{k+1}{6} - \frac{j}{2}$	$k = 3(\ell + i) - 1, j = \ell - i$ $-(6i - 2) = 3j - k + 1$

For $m = 1, \dots, 8$, denote by $S_m = \{(j, k) \in \mathbb{Z}_{\geq 0}^2 \mid (j, k) \text{ satisfies Condition } (m)\}$. Then,

$$\mathbb{Z}_{\geq 0}^2 \setminus \{(j, 3s + 1) \mid s, j \geq 0\} = \bigcup_{1 \leq m \leq 8} S_m, \quad S_i \cap S_j = \emptyset \text{ for } i \neq j.$$

Condition (1) in the table gives the following q -series relation:

$$\begin{aligned} \sum_{\ell=0}^{\infty} \sum_{i=0}^{\ell} (6i+1)q^{\ell(3\ell+2)-i(3i+1)} &= \sum_{k=3(\ell-i), j=\ell+i}^{\infty} (3j-k+1)q^{\frac{(2j+1)(2k+1)-1}{4}} \\ &= \sum_{(j,k) \in S_1} (3j-k+1)q^{\frac{(2j+1)(2k+1)-1}{4}}. \end{aligned}$$

Similarly, using Conditions (2)-(8), we obtain:

$$\begin{aligned} -\sum_{\ell \geq 1} \sum_{i=1}^{\ell} (6i-1)q^{\ell(3\ell+2)-i(3i-1)} &= \sum_{(j,k) \in S_2} (3k-j+1)q^{\frac{(2j+1)(2k+1)-1}{4}}, \\ -\sum_{\ell \geq 1} \sum_{i=1}^{\ell} (6i-4)q^{\frac{(2\ell-1)(6\ell+1)-(2i-1)(6i-5)}{4}} &= \sum_{(j,k) \in S_3} (3j-k+1)q^{\frac{(2j+1)(2k+1)-1}{4}}, \\ \sum_{\ell \geq 1} \sum_{i=1}^{\ell} (6i-2)q^{\frac{(2\ell-1)(6\ell+1)-(2i-1)(6i-1)}{4}} &= \sum_{(j,k) \in S_4} (3j-k+1)q^{\frac{(2j+1)(2k+1)-1}{4}}, \end{aligned}$$

$$\begin{aligned}
& - \sum_{\ell \geq 0} \sum_{i=0}^{\ell} (6i+1)q^{(\ell+1)(3\ell+1)-i(3i+1)} = \sum_{(j,k) \in S_5} (3j-k+1)q^{\frac{(2j+1)(2k+1)-1}{4}}, \\
& \sum_{\ell \geq 1} \sum_{i=1}^{\ell} (6i-1)q^{(\ell+1)(3\ell+1)-i(3i-1)} = \sum_{(j,k) \in S_6} (3j-k+1)q^{\frac{(2j+1)(2k+1)-1}{4}}, \\
& \sum_{\ell \geq 1} \sum_{i=1}^{\ell} (6i-4)q^{\frac{(2\ell+1)(6\ell-1)-(2i-1)(6i-5)}{4}} = \sum_{(j,k) \in S_7} (3j-k+1)q^{\frac{(2j+1)(2k+1)-1}{4}}, \\
& - \sum_{\ell \geq 1} \sum_{i=1}^{\ell} (6i-2)q^{\frac{(2\ell+1)(6\ell-1)-(2i-1)(6i-1)}{4}} = \sum_{(j,k) \in S_8} (3j-k+1)q^{\frac{(2j+1)(2k+1)-1}{4}}.
\end{aligned}$$

Next we notice that for any symmetric polynomial $\sigma(x, y)$ we have

$$(7.2) \quad \sum_{j,s=0}^{\infty} jq^{\sigma(j,s)} = \sum_{j,s=0}^{\infty} sq^{\sigma(j,s)}.$$

This implies that

$$\begin{aligned}
\mathcal{B}[q] &= \sum_{j,k=0}^{\infty} (3j-k+1)q^{\frac{(2j+1)(2k+1)-1}{4}} - \sum_{j=0}^{\infty} \sum_{s=0}^{\infty} (3j-(3s+1)+1)q^{\frac{(2j+1)(2(3s+1))-1}{4}} \\
&= \sum_{j,k=0}^{\infty} (3j-k+1)q^{\frac{(2j+1)(2k+1)-1}{4}} - \sum_{j,s=0}^{\infty} 3(j-s)q^{\frac{3(2s+1)(2j+1)-1}{4}} \\
&\stackrel{\text{by (7.2)}}{=} \underbrace{\sum_{j,k=0}^{\infty} (2k+1)q^{\frac{(2j+1)(2k+1)-1}{4}}}_{\text{by (7.2)}} = \Delta(q^{1/2})^4.
\end{aligned}$$

□

Lemma 7.2. *The decomposition (5.1) holds at the level of characters.*

Proof. The character of $L_{-3/2}(\mathfrak{sl}_3)$ is

$$\text{ch}[L_{-3/2}(\mathfrak{sl}_3)] = q^{-c/24} \frac{(q^2)_{\infty}^8}{(q)_{\infty}^8} = q^{1/3} \frac{1}{(q)_{\infty}^4} \left(\frac{(q^2)_{\infty}^2}{(q)_{\infty}} \right)^4 = q^{1/3} \frac{1}{(q)_{\infty}^4} \Delta(q)^4,$$

where by a Gauss' identity

$$\Delta(q) := \sum_{n \in \mathbb{Z}_{\geq 0}} q^{n(n+1)/2} = \frac{(q^2)_{\infty}^2}{(q)_{\infty}}.$$

Using Lemma 5.1 we get that the character of the right side of (5.2) is given by $\frac{q^{1/3}}{(q)_{\infty}^4} \mathcal{C}[q]$ where

$$\begin{aligned}
\mathcal{C}[q] &:= \sum_{\ell=0}^{\infty} \sum_{i=0}^{\ell} (8i+1) \left(q^{\ell(4\ell+3)-i(4i+1)} - q^{(\ell+1)(4\ell+1)-i(4i+1)} \right) \\
&+ \sum_{\ell=1}^{\infty} \sum_{i=1}^{\ell} (8i-1) \left(-q^{\ell(4\ell+3)-i(4i-1)} + q^{(\ell+1)(4\ell+1)-i(4i-1)} \right) \\
&+ \sum_{\ell=1}^{\infty} \sum_{i=1}^{\ell} (8i-3) \left(q^{\frac{(2\ell-1)(4\ell+1)-(2i-1)(4i-1)}{2}} - q^{\frac{(2\ell+1)(4\ell-1)-(2i-1)(4i-1)}{2}} \right) \\
&+ \sum_{\ell=1}^{\infty} \sum_{i=1}^{\ell} (8i-5) \left(-q^{\frac{(2\ell-1)(4\ell+1)-(2i-1)(4i-3)}{2}} + q^{\frac{(2\ell+1)(4\ell-1)-(2i-1)(4i-3)}{2}} \right).
\end{aligned}$$

Therefore the proof of the theorem is now reduced to the following q -series identity:

$$\frac{\phi(q^2)^8}{\phi(q)^4} = \Delta(q)^4 = \mathcal{C}[q].$$

Using similar calculation as in previous section, we prove that

$$\mathcal{C}[q] = \sum_{j,k=0}^{\infty} (4j - 2k + 1)q^{\frac{(2j+1)(2k+1)-1}{2}} = \sum_{j,k=0}^{\infty} (2k + 1)q^{\frac{(2j+1)(2k+1)-1}{2}} = \Delta(q)^4.$$

The proof follows. \square

8. MLDE FOR \mathcal{C}_p

We proved that for any $n \geq 2$, we have a simple vertex algebra structure on the module

$$\mathcal{C}_n := \bigoplus_{\ell=0}^{\infty} L_{\widehat{\mathfrak{sl}}_2}(-(-n-2+n\ell)\Lambda_0 + n\ell\Lambda_1) \otimes L_{\widehat{\mathfrak{sl}}_2}\left(-\left(\frac{2n-1}{n} + \ell\right)\Lambda_0 + \ell\Lambda_1\right),$$

of central charge $c_n = \frac{6(1+n-n^2)}{n}$. This vertex algebra is $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded for n even and $\mathbb{Z}_{\geq 0}$ -graded for n odd, so some features will be slightly different.

To compute the characters of modules $L_{\widehat{\mathfrak{sl}}_2}(-(-n-2+n\ell)\Lambda_0 + n\ell\Lambda_1)$, and thus of \mathcal{C}_n , we proceed as earlier in the cases of $n = 3$ and $n = 4$ and obtain the following formula:

Theorem 8.1. *For $n \geq 2$, we have*

$$\begin{aligned} (q)_{\infty}^6 q^{c/24} \text{ch}[\mathcal{C}_n](q) &= \sum_{i \geq 0, k \geq 0} (1 + 2(i+k))(1 + 2in)q^{k(k+1)n+i(2kn+n-1)} \\ &\quad - \sum_{i \geq 0, k \geq 0} \left(3 + 2(i+k)\right)(1 + 2in)q^{k(1+k)n+(i+1)(1+n+2kn)} \\ &\quad + \sum_{i \geq 0, k \geq 0} 2(1+i+k)(1+n+2in)q^{\frac{1}{2}+(k^2-\frac{1}{2})n+(1+i)(-1+n+2kn)} \\ &\quad - \sum_{i \geq 0, k \geq 0} 2(1+i+k)(-1+n+2in)q^{-\frac{1}{2}+(-\frac{1}{2}+k^2)n+(1+i)(1+n+2kn)}. \end{aligned}$$

Observe that we have signature $(1,1)$ -indefinite theta functions appearing above, so it is not totally clear that $\text{ch}[\mathcal{C}_n]$ are modular forms; see the last section for a thorough discussion about modularity of these series.

Recall the the spaces of classical holomorphic modular forms for $\Gamma(1) = SL(2, \mathbb{Z})$ and for the principal congruence subgroup $\Gamma(2)$ are given by (as graded rings):

$$\bigoplus_{k=0}^{\infty} M_k(\Gamma(1), \mathbb{C}) = \mathbb{C}[E_4, E_6],$$

where $E_{2k} = -\frac{B_{2k}}{(2k)!} + \frac{2}{(2k-1)!} \sum_{n \geq 1} \frac{n^{2k-1} q^n}{1-q^n}$, and by

$$\bigoplus_{k=0}^{\infty} M_k(\Gamma(2), \mathbb{C}) = \mathbb{C}[\theta_2(\tau)^4, \theta_3(\tau)^4],$$

where $\theta_2(\tau) = \sum_{n \in \mathbb{Z} + \frac{1}{2}} q^{\frac{1}{2}n^2}$ and $\theta_3(\tau) = \sum_{n \in \mathbb{Z}} q^{\frac{1}{2}n^2}$ are Jacobi's theta functions. Let

$$\Theta_{r,s}(\tau) := \theta_2(\tau)^{4r} \theta_3(\tau)^{4s}$$

so that $\Theta_{r,s}(\tau)$, with $2r + 2s = 2k$ spans $M_{2k}(\Gamma(2))$. Then we use Serre's derivatives:

$$(8.1) \quad \partial_{(k)} f(q) = \left(q \frac{d}{dq} + kE_2(q) \right) f(q),$$

and

$$(8.2) \quad D_q^{(k)} f(q) := \partial_{(2k-2)} \circ \cdots \circ \partial_{(2)} \circ \partial_{(0)} f(q),$$

where we set $D_q^{(0)}f(q) := f(q)$. We are interested in MLDEs of a particular type $\mathcal{D}_q^{(k)}f = 0$, where

$$(8.3) \quad \mathcal{D}_q^{(k)} \equiv D_q^{(k)} + \sum_{r=1}^k f_r(q) D_q^{(k-r)}, \quad f_r(q) \in M_{2r}(\Gamma, \mathbb{C}),$$

where $\Gamma = \Gamma(1)$ or $\Gamma(2)$. This type of equations have been studied in the context of vertex algebra characters for a long time [27], and also in connection with Schur's indices in 4d $N = 2$ SCFTs [16].

Next result would follow if \mathcal{C}_n is quasi-lisse, essentially using an argument from [6].

Conjecture 8.2. *For $n \geq 2$, the character $\text{ch}[\mathcal{C}_n]$ satisfies an MLDE with holomorphic coefficients for $M_*(\Gamma(1))$ for n odd and with $M_*(\Gamma(2))$ for n even.*

In support for this conjecture we present a few examples:

- (1) The lowest order MLDE for $\text{ch}[\mathcal{C}_2]$ is

$$D_q^{(1)}f(q) - \left(\frac{1}{8}\Theta_{1,0} + \frac{1}{8}\Theta_{0,1}\right)f(q) = 0$$

- (2) The lowest order MLDE for $\text{ch}[\mathcal{C}_3]$ is

$$D_q^{(2)}f(q) - 75E_4f(q) = 0$$

- (3) The lowest order MLDE for $\text{ch}[\mathcal{C}_4]$ is

$$D_q^{(3)}f(q) + \left(-\frac{5}{16}\Theta_{1,0} + \frac{5}{16}\Theta_{0,1}\right)D_q^{(2)}f(q) + \left(-\frac{77}{2304}\Theta_{2,0} - \frac{89}{1152}\Theta_{1,1} - \frac{17}{2304}\Theta_{0,2}\right)D_q^{(1)}f(q) \\ + \left(\frac{33}{4096}\Theta_{3,0} + \frac{33}{4096}\Theta_{2,1} - \frac{197}{4096}\Theta_{1,2} + \frac{3}{4096}\Theta_{0,3}\right)f(q) = 0.$$

- (4) The lowest order MLDE for $\text{ch}[\mathcal{C}_5]$ is

$$D_q^{(6)}f(q) - 161E_4D_q^{(4)}f(q) - \frac{28812}{75}E_6D_q^{(3)}f(q) - \frac{8965187}{75}E_8D_q^{(2)}f(q) - \frac{192787364}{125}E_{10}D_q^{(1)}f(q) \\ + \left(-\frac{5599287}{5}E_4^3 - \frac{48993336}{25}E_6^2\right)f(q) = 0.$$

9. MODULARITY OF THE CHARACTER OF \mathcal{C}_n

Here we discuss modular properties of characters of \mathcal{C}_n . To make formulas more transparent, we multiply the character with $\eta(\tau)^6$ so that only the indefinite-theta function remains, i.e. we are interested in modular properties of

$$A_n(\tau) := \eta(\tau)^6 \text{ch}[\mathcal{C}_n].$$

We first start with the formula obtained earlier.

$$(q)_\infty^6 q^{c/24} \text{ch}[\mathcal{C}_n](q) = \sum_{i \geq 0, k \geq 0} (1 + 2(i+k))(1 + 2in)q^{k(k+1)n + i(2kn + n - 1)} \\ - \sum_{i \geq 0, k \geq 0} \left(3 + 2(i+k)\right)(1 + 2in)q^{k(1+k)n + (i+1)(1+n+2kn)} \\ + \sum_{i \geq 0, k \geq 0} 2(1+i+k)(1+n+2in)q^{\frac{1}{2} + (k^2 - \frac{1}{2})n + (1+i)(-1+n+2kn)} \\ - \sum_{i \geq 0, k \geq 0} 2(1+i+k)(-1+n+2in)q^{-\frac{1}{2} + (-\frac{1}{2} + k^2)n + (1+i)(1+n+2kn)}.$$

Then, after manipulating the q -series and organizing terms based on whether i is even or odd, the RHS of the formula yields

$$(q)_\infty^6 q^{c/24} \text{ch}[\mathcal{C}_n](q) = \left(\sum_{i \geq 0, k \geq 0} - \sum_{i < 0, k < 0} \right) (1+i+2k)(1+in) q^{k(k+1)n + \frac{i}{2}(2kn+n-1)}.$$

Using $c = 6(1 + \frac{1}{n} - n)$, we multiply the above equation by $q^{-c/24+1/4} = q^{-\frac{1}{4n} + \frac{n}{4}}$, obtaining

$$A_n(\tau) = \left(\sum_{\ell_1, \ell_2 \geq 0} - \sum_{\ell_1, \ell_2 < 0} \right) (1+2\ell_1+\ell_2)(1+n\ell_2) q^{n(\ell_1 + \frac{n-1}{2n})^2 + n(\ell_1 + \frac{n-1}{2n})(\ell_2 + \frac{1}{n})}.$$

Remark 9.1. Now letting $Q(x_1, x_2) = nx_1^2 + nx_1x_2$, $A = \begin{pmatrix} 2n & n \\ n & 0 \end{pmatrix}$, and $\lambda = (\frac{n-1}{2n}, \frac{n+1}{2n}) \in A^{-1}\mathbb{Z}^2/\mathbb{Z}^2$ (this is true only for n odd but; for n even we have to take first $\tau \rightarrow 2\tau$). With these shifts we can write $A_n(\tau)$ above as

$$A_n(\tau) = \frac{n}{2} \sum_{(\ell_1, \ell_2) \in \mathbb{Z}^2 + (\frac{n-1}{2n}, \frac{1}{n})} (\text{sgn}(\ell_1) + \text{sgn}(\ell_2)) p(\ell_1, \ell_2) q^{n\ell_1^2 + n\ell_1\ell_2}$$

where $p(\ell_1, \ell_2) = \ell_2(2\ell_1 + \ell_2)$. It is easy to verify that the polynomial $\ell_2(2\ell_1 + \ell_2)$ is A -harmonic. There is yet another expression for $A_n(\tau)$ easily proven

$$A_n(\tau) = \sum_{\substack{\ell_1 \equiv \ell_2 \pmod{2} \\ 0 \leq |\ell_2| \leq \ell_1; \ell_1 \geq 0; \ell_2 \in \mathbb{Z}}} (1+\ell_1)(1+n\ell_2) q^{\frac{n}{4}(\ell_1+1)^2 - \frac{n}{4}(\ell_2 + \frac{1}{n})^2},$$

but this will not be used in the rest of the paper.

9.1. Appell-Lerch series. To determine modular properties of $A_n(\tau)$ we need a few facts about generalized Appell-Lerch series $\kappa_\ell(x, y, q)$. Let (here $x = e(z_1)$ and $y = e(z_2)$)

$$(9.1) \quad \kappa_\ell(x, y, q) = \sum_{m \in \mathbb{Z}} \frac{q^{\frac{\ell m^2}{2}} x^{\ell m}}{1 - xyq^m},$$

with a double series representation (in a certain domain):

$$\kappa_\ell(x, y, q) = \left(\sum_{i, j \geq 0} - \sum_{i, j < 0} \right) x^{i+\ell j} y^i q^{\frac{\ell}{2}j^2 + ij}.$$

Consider now $\kappa_2(x, y, q)$ with shifts along x and y variables

$$B_n(x, y, q) := \kappa_2(q^{\frac{n}{2}}x, q^{-\frac{1}{2}}y^n, q^n).$$

Then it is easy to see that

$$q^{-\frac{1}{4n} + \frac{n}{4}} \frac{\partial^2}{\partial x \partial y} (xy B_n(x, y, q)) \Big|_{x=y=1} = A_n(\tau).$$

Some special linear combinations of the Appell-Lerch series can be expressed through theta functions. For κ_2 this formula is quite nice and reads (see for instance [26])

$$(9.2) \quad \kappa_2(x, y, q) - \kappa_2(x^{-1}, y, q) = -\frac{(q; q)_\infty^3 \theta(x^2; q)}{\theta(xy; q) \theta(xy^{-1}; q)}.$$

where $\theta(x; q) = \sum_{n \in \mathbb{Z}} (-1)^n q^{\frac{n^2}{2} + \frac{n}{2}} x^n$. Differentiation of (9.1), and after specializing at $x = y = 1$, yields the following identity

$$\frac{\partial^2}{\partial x \partial y} (xy \kappa_2(q^{-n/2}x^{-1}, q^{-1/2}y^n, q^n)) \Big|_{x=y=1} = -\frac{\partial^2}{\partial x \partial y} (xy B_n(x, y, q)) \Big|_{x=y=1}.$$

Using this relation, together with (9.2), and some manipulations with q -series gives

Proposition 9.2. *We have for $n \geq 2$,*

$$A_n(\tau) = \frac{1}{2} \frac{\partial^2}{\partial x \partial y} \left(\frac{\eta(n\tau)^3 \vartheta_{n,n}(x^2)}{\vartheta_{n,2n-1}(xy^n) \vartheta_{n,2n+1}(xy^{-n})} \right) \Big|_{x=y=1},$$

where $\vartheta_{n,a}(x; q) = \sum_{n \in \mathbb{Z}} (-1)^k q^{\frac{a}{2}(k + \frac{a}{2n})^2} x^{k + \frac{a}{2n}}$, is a Jacobi form of weight $\frac{1}{2}$.

Theorem 9.3. *The series $A_n(\tau)$ is a modular form (on a certain congruence group) of weight 3. Consequently, $\text{ch}[C_n]$ is modular of weight zero.*

Proof. Using Proposition 9.2 together with $\vartheta_{n,n}(1) = 0$, after differentiation we obtain that $A_n(\tau)$ can be expressed using specialized (at $x = y = 1$) theta series and their first derivatives, so $A_n(\tau)$ is clearly modular. Since we started from a Jacobi form of weight one, after two differentiations and specializations we get a modular form of weight 3. Since $\text{ch}[C_n] = \frac{1}{\eta(\tau)^6} A_n(\tau)$, the character has weight 0. \square

Remark 9.4. (a) In [12], a closed formula for $A_2(\tau)$ is obtained as a specialization of the Kac-Wakimoto denominator formula for $\widehat{\text{osp}(3|2)}$ [26] :

$$A_2(\tau) = \frac{\eta(\tau)^{12}}{\eta(\tau/2)^6}.$$

(b) For $n = 3$, in [6], the character of $L_{-\frac{5}{3}}(G_2)$ was computed using the fact that it satisfies the MLDE

$$D_q^{(2)} f(q) - 75E_4 f(q) = 0.$$

This gives

$$A_3(\tau) = \frac{\eta(3\tau)^6 E_1^{(3)}(\tau)}{\eta(\tau)^2}$$

where $E_1^{(3)}$ is an Eisenstein series of weight 1. Both proofs use different techniques compared to Theorem 9.3.

(c) Analysis of character-like solutions of MLDEs in Section 8, for low p , reveals that C_p most likely has a unique ordinary module.

Based on $p = 2, 3, 4, 5$ cases studied previously and Theorem 9.3 we expect

Conjecture 9.5. *Vertex operator algebra C_p is quasi-lisse for all p .*

10. FUTURE WORK

There are several directions that we would like to pursue.

- (1) We plan to generalize some results of this paper by extending them to higher-rank simple Lie algebras at levels $-h^\vee + \frac{1}{p}$ and $-h^\vee - p$, starting with \mathfrak{sl}_3 . Although some results on the structure of such representations at these affine levels are missing (or they are subject to values of Kazhdan-Lusztig polynomials), we believe that higher rank simple vertex algebras $C_{p,\mathfrak{g}}$ generalizing $\mathfrak{g} = \mathfrak{sl}_2$, can be rigorously constructed using methods of this paper for every simple \mathfrak{g} . If \mathfrak{g} is of ADE type, we expect that we have a simple vertex algebra structure on

$$C_{p,\mathfrak{g}} = \bigoplus_{\lambda \in P^+} L_{-p-h^\vee, p\lambda}(\mathfrak{g}) \otimes L_{-h^\vee + \frac{1}{p}, \lambda^*}(\mathfrak{g}).$$

Again, unlike $\mathcal{D}_{G, -h^\vee + \frac{1}{p}}^{ch}$, this space is $\mathbb{Z}_{\geq 0}$ -graded with finite-dimensional graded subspaces. For $p = 2$ and $\mathfrak{g} = \mathfrak{sl}_3$, we expect $C_{2, \mathfrak{sl}_3} \cong L_{-\frac{5}{2}}(F_4)$, from the Deligne series [6].

- (2) As hinted in the introduction, studying non-ordinary modules for C_p would be interesting, as it provides a direct connection to minimal representations. Another direction for exploration is the construction of logarithmic modules, as in [13].

- (3) The modularity of characters discussed in the final section suggests new families of indefinite-type modular theta functions with harmonic coefficients arising from the “numerator” of $\text{ch}[C_{p,g}]$ with harmonic coefficients (see Remark 9.1). Related higher rank q -series with polynomial coefficients have been previously studied by S. Zwegers and others [29, 32].

Acknowledgements. D.A. is partially supported by the Croatian Science Foundation under the project IP-2022-10-9006. We would like to thank Igor Frenkel for bringing [19] to our attention a while ago, and Sander Zwegers for discussion regarding indefinite-theta functions.

REFERENCES

- [1] S. Arkhipov, and D. Gaitsgory. Differential operators on the loop group via chiral algebras, *International Mathematics Research Notices* 2002.4 (2002): 165-210.
- [2] T. Arakawa and A. Moreau. Joseph ideals and lisse minimal-algebras. *Journal of the Institute of Mathematics of Jussieu* 17.2 (2018): 397-417.
- [3] T. Arakawa and A. Moreau, *Arc spaces and vertex algebras*, Springer Monograph, April 17 (2021): 170.
- [4] T. Arakawa, Chiral algebras of class S and Moore-Tachikawa symplectic varieties. arXiv preprint arXiv:1811.01577.
- [5] T. Arakawa, J. van Ekeren, and A. Moreau 2020. On the nilpotent orbits arising from admissible affine vertex algebras, *Proc. Lond. Math. Soc.*, 125 (2022), 681–699.
- [6] T. Arakawa, K. Kawasetsu, Quasi-lisse vertex algebras and modular linear differential equations, *Lie Groups, Geometry, and Representation Theory: A Tribute to the Life and Work of Bertram Kostant* (2018): 41-57.
- [7] T. Arakawa, T. Creutzig, K. Kawasetsu, A. Linshaw, Orbifolds and cosets of minimal W -algebras, *Comm. Math. Phys.* 355 (2017), no. 1, 339–372
- [8] T. Arakawa, Chiral algebras of class S and Moore-Tachikawa symplectic varieties, arXiv:1811.01577.
- [9] D. Adamović Realizations of simple affine vertex algebras and their modules: the cases $\widehat{\mathfrak{sl}}_2$ and $\widehat{osp}(1, 2)$, *Communications in Mathematical Physics*, March 2019, Volume 366, Issue 3, pp 1025-1067; arXiv:1711.11342 [math.QA].
- [10] D. Adamović, V. G. Kac, P. Möseneder Frajria, P. Papi, O. Perše, Finite vs infinite decompositions in conformal embeddings, *Communications in Mathematical Physics* 348 (2016) 445–473.
- [11] D. Adamović, T. Arakawa, T. Creutzig, A. Linshaw, A. Moreau, P. Möseneder Frajria, P. Papi, to appear
- [12] D. Adamović, V. G. Kac, P. Möseneder Frajria, P. Papi, O. Perše, On classification of non-equal rank affine conformal embeddings and applications, *Selecta Mathematica New Series*, July 2018, Volume 24, Issue 3, pp 2455-2498
- [13] D. Adamović, T. Creutzig, N. Genra, J. Yang, The vertex algebras $\mathcal{V}^{(p)}$ and $\mathcal{R}^{(p)}$, *Communications in Mathematical Physics* volume 383, pages1207-1241(2021) arXiv:2001.08048 [math.RT].
- [14] D. Adamović, P. Möseneder Frajria, P. Papi, New approaches for studying conformal embeddings and collapsing levels for W -algebras, *International Mathematics Research Notices*, Volume 2023, Issue 22, November 2023, Pages 19431–19475, rna138, arXiv:2203.08497 [math.RT]
- [15] C. Beem, C. , Meneghelli, C. and L. Rastelli, Free field realizations from the Higgs branch. *J. High Energ. Phys.* 2019, 58 (2019). [https://doi.org/10.1007/JHEP09\(2019\)058](https://doi.org/10.1007/JHEP09(2019)058)
- [16] C. Beem and L. Rastelli. Vertex operator algebras, Higgs branches, and modular differential equations. *Journal of High Energy Physics* 2018, no. 8 (2018): 1-72.
- [17] Feigin, B., and S. Parkhomenko. Regular representation of affine Kac-Moody algebras. *Algebraic and Geometric Methods in Mathematical Physics: Proceedings of the Kaciveli Summer School, Crimea, Ukraine, 1993*. Dordrecht: Springer Netherlands, 1996.
- [18] A. Feingold, and I. Frenkel. Classical affine algebras. *Advances in Mathematics* 56.2 (1985): 117-172.
- [19] I. Frenkel, and K. Styrkas. Modified regular representations of affine and Virasoro algebras, VOA structure and semi-infinite cohomology. *Advances in Mathematics* 206.1 (2006): 57-111.
- [20] The GAP Group, GAP – Groups, Algorithms, and Programming, Version 4.14.0; 2024.
- [21] D. Gaitsgory, Quantum Langlands correspondence. arXiv preprint arXiv:1601.05279.
- [22] S. Gelfand, The Weil representation of a Lie algebra of type G_2 and related representations of SL_3 , *Funktsional. Anal. i Prilozhen.* 14 (1980), no. 1, 51–52.
- [23] V. Gorbounov, F. Malikov, V. Schechtman, Gerbes of chiral differential operators, II, *math.AG/0003170v2*, August 2000
- [24] V. Gorbounov, F. Malikov, and V. Schechtman, On chiral differential operators over homogeneous spaces, *International Journal of Mathematics and Mathematical Sciences*, 2001;26(2):83-106.
- [25] V. Kac, M. Wakimoto, Quantum reduction and representation theory of superconformal algebras, *Adv. Math.*, **185**(2):400–458, 2004.

- [26] V. Kac, M. Wakimoto, Integrable highest weight modules over affine superalgebras and number theory. In: Brylinski, J.-L., Brylinski, R., Guillemin, V., Kac, V. (Eds.) Lie Theory and Geometry, Progress in Mathematics, vol. 123, pp. 415–456. Birkhäuser, Boston (1994)
- [27] A. Milas, On certain automorphic forms associated to rational vertex operator algebras, Proceedings of a Workshop on the Moonshine Conjectures and Vertex Algebras 2004, London Mathematical Society Lecture Note Series, Cambridge University Press.
- [28] R. McRae, A general mirror equivalence theorem for coset vertex operator algebras, Sci. China Math., Vol. 67 (2024), no. 10, 2237-2282
- [29] C. Roehrig and S. Zwegers. Theta series for quadratic forms of signature $(n - 1, 1)$ with (spherical) polynomials. International Journal of Number Theory 18.07 (2022): 1491-1515.
- [30] G. Savin, An analogue of the Weil representation for G_2 , Journal für die reine und angewandte Mathematik 435 (1993): 115-126.
- [31] M. Zhu, Vertex operator algebras associated to modified regular representations of affine Lie algebras. Advances in Mathematics. 2008 Dec 1;219(5):1513-47.
- [32] S. Zwegers, Mock theta functions, arXiv preprint arXiv:0807.4834 (2008).

(Dražen Adamović) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, UNIVERSITY OF ZAGREB, 10000 ZAGREB, CROATIA
Email address, Dražen Adamović: adamovic@math.hr

(Antun Milas) DEPARTMENT OF MATHEMATICS AND STATISTICS, SUNY-ALBANY, ALBANY NY 12222, USA
Email address, Antun Milas: amilas@albany.edu