# LIFESPAN ESTIMATE FOR THE SEMILINEAR REGULAR EULER-POISSON-DARBOUX-TRICOMI EQUATION

YUEQUN LI AND FEI GUO

ABSTRACT. In this paper, we begin by establishing local well-posedness for the semilinear regular Euler-Poisson-Darboux-Tricomi equation. Subsequently, we derive a lifespan estimate with the Strauss index given by  $p = p_S(n + \frac{\mu}{m+1}, m)$  for any  $\delta > 0$ , where  $\delta$  is a parameter to describe the interplay between damping and mass. This is achieved through the construction of a new test function derived from the Gaussian hypergeometric function and a second-order ordinary differential inequality, as proven by Zhou [36]. Additionally, we extend our analysis to prove a blow-up result with the index  $p = \max\{p_S(n + \frac{\mu}{m+1}, m), p_F((m+1)n + \frac{\mu-1-\sqrt{\delta}}{2})\}$  by applying Kato's Lemma (i.e., Lemma 5.1), specifically in the case of  $\delta = 1$ .

*Keywords*: Semilinear Euler-Poisson-Darboux-Tricomi equation; Strauss exponent; Fujita exponent; Gaussian hypergeometric function; Test function.

AMS Subject Classification (2020): 35B44, 35G25

### 1. INTRODUCTION

We consider the following semilinear Euler-Poisson-Darboux-Tricomi equation (EPDT) with a power nonlinearity

$$\begin{cases} \partial_t^2 u - t^{2m} \Delta u + \frac{\mu}{t} \partial_t u + \frac{\nu^2}{t^2} u = |u|^p, & t \ge 1, \\ u(1,x) = \varepsilon u_0(x), \ \partial_t u(1,x) = \varepsilon u_1(x), & x \in \mathbb{R}^n, \end{cases}$$
(1.1)

where  $m \ge 0$ ,  $n \ge 1$  denotes the space dimension and  $\varepsilon > 0$  is a sufficiently small number describing the smallness of the initial value. Palmieri [24] first named (1.1) the "EPDT" while investigating its critical exponent problem, one can see [5], [28] and their references to obtain more results on the generalized Euler-Poisson-Darboux equation. (1.1) is known as scale-invariant since the corresponding linear equation is invariant under the scaling

$$v(t,x) = u(\lambda^{\frac{1}{m+1}}t,\lambda x).$$
(1.2)

(1.1) becomes the generalized Euler-Poisson-Darboux equation when  $m = \nu = 0$ , which is commonly encountered in the fields of gas dynamics, fluid mechanics, elasticity, plasticity, and various physical and material science problems. If the initial values are given at time zero, this problem is referred to as the singular problem, and if the initial values are given at a non-zero time, it is called the regular problem. D'Abbicco, Lucente and Reissig [8] determined the critical exponent for the regular problem and proved blow-up results in the case of  $n \ge 2$  with  $\mu = 2$ , and the global well-posedness was only proved in low-dimensional

Date: Sept. 10, 2024.

cases (n = 2, 3). Subsequently, they [7] demonstrated the existence of global solutions for the regular problem with spherically symmetric initial data. However, concerning the singular problem, to our best knowledge, there are currently no satisfactory results available and some partial results can be referred to [5, 6].

When  $\mu = \nu^2 = 0$ , (1.1) reduces to the semilinear Tricomi equation

$$\partial_t^2 u - t^{2m} \Delta u = |u|^p, \tag{1.3}$$

and it is often used in gas dynamics to describe the conversion between ultrasound and infrasound, one can refer [9] for more details. At the early stage, mathematicians attempted to establish the expression of explicit solutions for the linear Tricomi equation. Barros-Neto and Gelfand [1, 2, 3] found the fundamental solution for 1-D problem  $yu_{xx} + u_{yy} = 0$  and then Yagdjian [33] established the explicit fundamental solution for the generalized Gellerstedt operator  $\partial_t^2 - t^{2m}\Delta$  in  $\mathbb{R}^{n+1}$ . In recent years, the large time behavior of nonlinear Tricomi equations has attracted much attention. Many mathematicians dedicated to the critical exponent problem for (1.3), namely, it is expected that there exists an index  $p_c(n,m)$ such that if  $p > p_c(n,m)$ , (1.3) has small data global solution and when 1 ,the solution to (1.3) will blow up even for small initial values. Note that when <math>m = 0, (1.3) becomes the classical semilinear wave equation

$$\partial_t^2 u - \Delta u = |u|^p \tag{1.4}$$

with the critical exponent  $p_S(n)$ , which is the positive root of the quadratic equation

$$(n-1)p^2 - (n+1)p - 2 = 0$$
 for  $n \ge 2$ , (1.5)

and  $p_S(1) = +\infty$ , one can refer [12, 13, 19, 29, 31, 34, 35, 36] for more details about Strauss conjecture. In the past few years, He, Witt and Yin [16, 17, 18] systematically studied (1.3) and determined the critical exponent  $p_S(n,m)$ , more precisely,  $p_S(n,m)$  is the positive root of the quadratic equation

$$((m+1)n-1)p^2 - ((m+1)(n-2)+3)p - 2(m+1) = 0.$$
(1.6)

It should be pointed out that the critical exponent  $p_S(n,m)$  extends  $p_S(n)$  because (1.6) reduces to (1.5) when m = 0, that is  $p_S(n) = p_S(n,0)$ . Subsequently, Lin and Tu [20] obtained the upper bound estimate of lifespan by the test function method when 1 , and Sun [30] gave the lower bound estimate based on the Strichartz estimate established in [18] when <math>1 .

When m = 0, (1.1) becomes the semilinear wave equation with scale invariant damping and mass

$$\begin{cases} \partial_t^2 u - \Delta u + \frac{\mu}{t} \partial_t u + \frac{\nu^2}{t^2} u = |u|^p, & t \ge 1, \\ u(1,x) = \varepsilon u_0(x), \ \partial_t u(1,x) = \varepsilon u_1(x), & x \in \mathbb{R}^n. \end{cases}$$
(1.7)

It has been found that the scale of the non-negative number  $\delta = (\mu - 1)^2 - 4\nu^2$  reflects an important relationship between the damping term  $\frac{\mu}{t}\partial_t u$  and the mass term  $\frac{\nu^2}{t^2}u$ . Specifically speaking, in [25], it has been proved that the solution to (1.7) blows up when  $1 denotes the Fujita index appearing in the semilinear heat equation (c.f. [11, 15]). For large <math>\delta$  ( $\delta \geq (n+1)^2$ ), (1.7) can be called "parabolic-like" since it has been

proved that  $p_F(n + \frac{\mu - 1 - \sqrt{\delta}}{2})$  is the critical exponent in [21, 23, 26]. Subsequently, it is shown in [27] that the solution blows up when  $1 for all <math>\delta > 0$  and  $p = p_S(n + \mu)$  for  $\delta < n^2$ .

As for (1.1), Palmieri [24] proved that for all  $\delta \ge 0$ , the solution blows up when  $1 and established the upper bound estimate of the lifespan utilizing the iterative argument, where <math>p_S(n + \frac{\mu}{m+1}, m)$  is the positive root to the following quadratic equation

$$((m+1)n - 1 + \mu)p^2 - ((m+1)(n-2) + 3 + \mu)p - 2(m+1) = 0.$$
(1.8)

Note that  $p_S(n + \mu, 0) = p_S(n + \mu)$ , so we can make a reasonable conjecture that, for different values of  $\delta$ ,  $p_S(n + \frac{\mu}{m+1}, m)$ ,  $p_F((m + 1)n + \frac{\mu-1-\sqrt{\delta}}{2})$  or the larger of the two indices represents the critical exponent of (1.1). The primary objective in this paper is to demonstrate that for  $\delta < (m + 1)^2 n^2$ , the solution to (1.1) blows up in finite time and give the upper bound lifespan estimate when  $p = p_S(n + \frac{\mu}{m+1}, m)$ . Due to the appearance of  $t^m$  in the Gellerstedt operator, the test function that works for the wave operator is no longer applicable, this compels us to search for a new one that depends on m. Inspired by the treatment of  $p = p_S(n, 0)$  in [27], we construct a new test function using Gaussian hypergeometric functions, and when m = 0, this function is the same as the one used in [27]. Additionally, when  $\delta = 1$ , (1.1) can be transformed into a Tricomi-type equation without damping and mass terms, and then the second objective of this paper is to give a blow up result for  $p = \max\{p_S(n + \frac{\mu}{m+1}, m), p_F((m + 1)n + \frac{\mu-1-\sqrt{\delta}}{2})\}$  under the condition  $\delta = 1$ . It is a pity that when  $\delta \neq 1$ , we have not found a suitable method to handle the case of  $p = p_F((m + 1)n + \frac{\mu-1-\sqrt{\delta}}{2})$ , however, in our next paper, we demonstrate that  $p_F((m + 1)n + \frac{\mu-1-\sqrt{\delta}}{2})$  is the critical exponent for large  $\delta$ .

This paper is organized as follows. In Section 2, we provide the definition of the energy solution to (1.1) and present the main results. Section 3 is devoted to prove the local well-posedness of (1.1) by deriving  $(L^1 \cap L^2) - L^2$  estimates to the solutions of the corresponding linear equation. In Section 4, we give the new test function, and consequently, obtain the upper bound estimate of the lifespan with the Strauss index given by  $p = p_S(n + \frac{\mu}{m+1}, m)$ . A blow up result is derived in Section 5 in the case of  $\delta = 1$  with the index  $p = \max\{p_S(n + \frac{\mu}{m+1}, m), p_F((m+1)n + \frac{\mu-1-\sqrt{\delta}}{2})\}$ .

### 2. MAIN RESULTS

Before stating our results, we give the definition of energy solution for (1.1). Let us introduce the space

$$D^{\sigma}(\mathbb{R}^{n}) = \begin{cases} \left( L^{1}(\mathbb{R}^{n}) \cap H^{\sigma}(\mathbb{R}^{n}) \right) \times \left( L^{1}(\mathbb{R}^{n}) \cap H^{\sigma-1}(\mathbb{R}^{n}) \right), & \sigma > 1, \\ \left( L^{1}(\mathbb{R}^{n}) \cap H^{\sigma}(\mathbb{R}^{n}) \right) \times \left( L^{1}(\mathbb{R}^{n}) \cap L^{2}(\mathbb{R}^{n}) \right), & \sigma \in [0, 1] \end{cases}$$
(2.1)

with the norm

$$\|(u_0, u_1)\|_{D^{\sigma}} = \begin{cases} \|u_0\|_{L^1} + \|u_0\|_{H^{\sigma}} + \|u_1\|_{L^1} + \|u_1\|_{H^{\sigma-1}}, & \sigma > 1, \\ \|u_0\|_{L^1} + \|u_0\|_{H^{\sigma}} + \|u_1\|_{L^1} + \|u_1\|_{L^2}, & \sigma \in [0, 1], \end{cases}$$
(2.2)

and  $D^1 := D$ .

**Definition 2.1.** Let  $(u_0, u_1) \in D$  be compactly supported in  $\mathbb{R}^n$  and  $u \in L^p_{loc}([1, T] \times \mathbb{R}^n)$ . We say that u is an energy solution of (1.1) if

$$u \in C\left([1,T); H^1(\mathbb{R}^n)\right) \cap C^1\left([1,T); L^2(\mathbb{R}^n)\right)$$

satisfies for any  $\Phi(t,x) \in C_0^{\infty}([1,T) \times \mathbb{R}^n)$  and  $t \in [1,T)$ ,

$$\int_{1}^{t} \int_{\mathbb{R}^{n}} u(s,x) \Big( \partial_{s}^{2} \Phi(s,x) - s^{2m} \Delta \Phi(s,x) - \partial_{t} \Big( \frac{\mu}{s} \Phi(s,x) \Big) + \frac{\nu^{2}}{s^{2}} \Phi(s,x) \Big) dx ds$$

$$+ \int_{\mathbb{R}^{n}} \Big( \partial_{t} u(t,x) \Phi(t,x) - u(t,x) \partial_{t} \Phi(t,x) + \frac{\mu}{t} u(t,x) \Phi(t,x) \Big) dx$$

$$= \int_{1}^{t} \int_{\mathbb{R}^{n}} |u(s,x)|^{p} \Phi(s,x) dx ds$$

$$+ \varepsilon \int_{\mathbb{R}^{n}} \Big( -u_{0}(x) \partial_{t} \Phi(1,x) + \big( \mu u_{0}(x) + u_{1}(x) \big) \Phi(1,x) \big) dx.$$

$$(2.3)$$

To begin with, we establish the local well-posedness of (1.1).

**Proposition 2.1.** Let  $(u_0, u_1) \in D$  compactly supported in  $B_M(0) = \{x : |x| \leq M\}$  with some M > 0. Suppose

$$1 (2.4)$$

and  $\mu$ ,  $\nu \geq 0$  such that

$$\delta := (\mu - 1)^2 - 4\nu^2 > 0, \tag{2.5}$$

then, there exists a T > 1 and a unique solution  $u \in C([1,T); H^1(\mathbb{R}^n)) \cap C^1([1,T); L^2(\mathbb{R}^n))$ to (1.1).

The lifespan estimate of the solutions to (1.1) is as follows.

**Theorem 2.1.** Let  $m, \mu, \nu \ge 0, n \ge 1$  and  $0 < \delta < (m+1)^2 n^2$ . Assume that the initial data  $(u_0, u_1) \in D$  be nonnegative, not identically zero and  $supp(u_0, u_1) \subset \{x : |x| \le M\}$  for some  $0 < M < \frac{1}{m+1}$ . Consider  $p = p_S(n + \frac{\mu}{m+1}, m)$ , we suppose  $p > \frac{2(m+1)}{(m+1)n-\sqrt{\delta}}$ , furthermore, when n = 1, we assume  $\mu \ge m$ , then there exists a  $\varepsilon_0 = \varepsilon_0(u_0, u_1, m, n, p, M, \mu, \nu) > 0$  such that for any  $0 < \varepsilon \le \varepsilon_0$ , the solution u to (1.1) blows up in finite time and we have the upper estimate of the lifespan  $T(\varepsilon)$ 

$$T(\varepsilon) \le e^{C\varepsilon^{-p(p-1)}}$$

**Remark 2.1.** We assess the reasonability for the assumption  $p > \frac{2(m+1)}{(m+1)n-\sqrt{\delta}}$ . Since  $p = p_S(n + \frac{\mu}{m+1}, m)$ , the assumption is actually a restriction on  $\mu$ ,  $\nu$ . We can easily show that n = 3,  $\mu = 3$  and  $\nu = \frac{1}{2}$  are suitable for  $0 < 3 = \delta < 9(m+1)^2$  and this assumption, furthermore, we can check that (2.4) holds in this case.

**Theorem 2.2.** Assume that  $m, \mu, \nu \ge 0, n \ge 1$  and  $\delta = 1$ , the initial data  $(u_0, u_1) \in D$  be nonnegative, not identically zero and compactly supported in  $\{x : |x| \le M\}$  for some M > 0. Then the solution to (1.1) blows up in finite time when  $p = \max\{p_S(n + \frac{\mu}{m+1}, m), p_F((m+1)n + \frac{\mu-1-\sqrt{\delta}}{2})\}$ .

**Remark 2.2.** After a complex algebraic calculation, we discover that  $p_S(n + \frac{\mu}{m+1}, m) > p_F((m+1)n + \frac{\mu-1-\sqrt{\delta}}{2})$  holds when  $n \ge 3$  in the case of  $\delta = 1$  for any  $m \ge 0$ , then in view of Theorem 2.1, we only need to deal with the low dimensions n = 1, 2, and the details can be found in Section 5.

### 3. The Proof Framework of Proposition 2.1

Through two changes of variables, we can transform (1.1) into a damped wave equation without the mass term. By following the approach in [26] and [32], we can establish  $(L^1 \cap L^2) - L^2$  estimates for the corresponding linear equation of (1.1). Subsequently, the existence of local solutions can be obtained by the contraction mapping principle.

Consider the corresponding linear equation of (1.1)

$$\begin{cases} \partial_t^2 u - t^{2m} \Delta u + \frac{\mu}{t} \partial_t u + \frac{\nu^2}{t^2} u = 0, \qquad t \ge \tau \ge 1, \end{cases}$$
(3.1)

$$u(\tau, x) = u_0(x), \ \partial_t u(\tau, x) = u_1(x), \qquad x \in \mathbb{R}^n,$$

let

$$u(t,x) = t^{\frac{\sqrt{\delta} - \mu + 1}{2}} w(t,x), \qquad (3.2)$$

where  $\delta$  is defined by (2.5), (3.1) is transformed into

$$\begin{cases} \partial_t^2 w - t^{2m} \Delta w + \left(1 + \sqrt{\delta}\right) \frac{\partial_t w}{t} = 0, & t \ge \tau \ge 1, \\ w(\tau, x) = w_0(x), \ \partial_t w(\tau, x) = w_1(x), & x \in \mathbb{R}^n, \end{cases}$$
(3.3)

where

$$w_0(x) = \tau^{-\frac{\sqrt{\delta}-\mu+1}{2}} u_0(x),$$
  
$$w_1(x) = \tau^{-\frac{\sqrt{\delta}-\mu+1}{2}} u_1(x) - \frac{\sqrt{\delta}-\mu+1}{2} \tau^{-\frac{\sqrt{\delta}-\mu+3}{2}} u_0(x).$$

Set

$$s = \frac{t^{m+1}}{m+1}, \ \eta = \frac{\tau^{m+1}}{m+1} \text{ and } v(s,x) = w\Big(\Big((m+1)s\Big)^{\frac{1}{m+1}}, x\Big), \tag{3.4}$$

(3.3) can be transformed into

$$\begin{cases} \partial_s^2 v - \Delta v + \left(1 + \frac{\sqrt{\delta}}{m+1}\right) \frac{\partial_s v}{s} = 0, \quad s \ge \eta, \\ v(\eta, x) = v_0(x), \; \partial_s v(\eta, x) = v_1(x), \quad x \in \mathbb{R}^n, \end{cases}$$
(3.5)

where

$$v_0(x) = w_0(x) = \left((m+1)\eta\right)^{-\frac{\sqrt{\delta}-\mu+1}{2(m+1)}} u_0(x),$$
  

$$v_1(x) = \left((m+1)\eta\right)^{-\frac{m}{m+1}} w_1(x)$$
  

$$= \left((m+1)\eta\right)^{-\frac{m}{m+1}-\frac{\sqrt{\delta}-\mu+1}{2(m+1)}} u_1(x) - \frac{\sqrt{\delta}-\mu+1}{2} ((m+1)\eta)^{-\frac{m}{m+1}-\frac{\sqrt{\delta}-\mu+3}{2(m+1)}} u_0(x).$$

We will use the ideas outlined in [26] to establish  $(L^1 \cap L^2) - L^2$  estimates for (3.5), subsequently, the  $(L^1 \cap L^2) - L^2$  estimates for (3.1) can be obtained through (3.2) and (3.4). Here, we only list the results.

**Lemma 3.1.** Assume that  $u_0 = 0$ ,  $u_1 \in L^1 \cap H^k$  for some k > 0 and  $\delta > 0$ . Then for any  $k \in [0, 1]$ , the solution u to (3.1) satisfies the decay estimate

$$\|u(t,\cdot)\|_{\dot{H}^{k}} \le C_{m} \left( \|u_{1}\|_{L^{1}} + \tau^{(m+1)\frac{n}{2}} \|u_{1}\|_{L^{2}} \right)$$

$$\int t^{-\frac{\mu+m}{2}} \tau^{-(m+1)(k+\frac{n}{2})+\frac{\mu+m}{2}+1}, \quad \text{if} \quad k > \frac{\sqrt{\delta}}{2(m+1)} + \frac{1}{2} - \frac{n}{2}, \quad (3.6a)$$

$$\times \left\{ t^{-\frac{\mu+m}{2}} \tau^{-\frac{\sqrt{\delta}-\mu-1}{2}} (1+\log\frac{t}{\tau})^{\frac{1}{2}}, \quad if \quad k = \frac{\sqrt{\delta}}{2(m+1)} + \frac{1}{2} - \frac{n}{2}, \tag{3.6b} \right\}$$

$$t^{-(m+1)(k+\frac{n}{2})+\frac{\sqrt{\delta}-\mu+1}{2}}\tau^{-\frac{\sqrt{\delta}-\mu-1}{2}}, \text{ if } k < \frac{\sqrt{\delta}}{2(m+1)} + \frac{1}{2} - \frac{n}{2}.$$
 (3.6c)

*Moreover, for any*  $k \geq 1$ *,* 

$$\|\partial_t u(t,\cdot)\|_{\dot{H}^{k-1}} \le C_m \left( \|u_1\|_{L^1} + \tau^{(m+1)(\frac{n}{2}+k-1)} \|u_1\|_{\dot{H}^{k-1}} + \tau^{(m+1)(\frac{n}{2}+[k-2]_+)} \|u_1\|_{\dot{H}^{[k-2]_+}} \right)$$

$$\int t^{m-\frac{\mu+m}{2}} \tau^{-(m+1)(k+\frac{n}{2})+\frac{\mu+m}{2}+1}, \qquad \text{if} \quad k > \frac{\sqrt{\delta}}{2(m+1)} + \frac{1}{2} - \frac{n}{2}, \qquad (3.7a)$$

$$\times \left\{ t^{m - \frac{\mu + m}{2}} \tau^{-\frac{\sqrt{\delta} - \mu - 1}{2}} (1 + \log \frac{t}{\tau})^{\frac{1}{2}}, \quad if \quad k = \frac{\sqrt{\delta}}{2(m+1)} + \frac{1}{2} - \frac{n}{2}, \quad (3.7b)$$

$$\left(t^{m-(m+1)(k+\frac{n}{2})+\frac{\sqrt{\delta}-\mu+1}{2}}\tau^{-\frac{\sqrt{\delta}-\mu-1}{2}}, if k < \frac{\sqrt{\delta}}{2(m+1)} + \frac{1}{2} - \frac{n}{2}, \quad (3.7c)\right)$$

where  $[x]_{+} = \max\{x, 0\}.$ 

**Lemma 3.2.** Assume that  $(u_0, u_1) \in D^k$  for some k > 0 and  $\delta > 0$ . Then for any  $k \in [0, 1]$ , the solution u to (3.1) with  $\tau = 1$  satisfies the decay estimates

$$||u(t,\cdot)||_{\dot{H}^k} \le C_m ||(u_0,u_1)||_{D^k}$$

$$\int t^{-\frac{\mu+m}{2}}, \qquad \qquad \text{if} \quad k > \frac{\sqrt{\delta}}{2(m+1)} + \frac{1}{2} - \frac{n}{2}, \qquad (3.8a)$$

$$\times \left\{ t^{-\frac{\mu+m}{2}} (1+\log t)^{\frac{1}{2}}, \quad if \quad k = \frac{\sqrt{\delta}}{2(m+1)} + \frac{1}{2} - \frac{n}{2}, \quad (3.8b) \right\}$$

$$\left(t^{-(m+1)(k+\frac{n}{2})+\frac{\sqrt{\delta}-\mu+1}{2}}, if k < \frac{\sqrt{\delta}}{2(m+1)} + \frac{1}{2} - \frac{n}{2}, \quad (3.8c)\right)$$

and for any  $k \geq 1$ ,

 $\|\partial_t u(t,\cdot)\|_{\dot{H}^{k-1}} \le C_m \|(u_0,u_1)\|_{D^k}$ 

$$t^{m-\frac{\mu+m}{2}},$$
 if  $k > \frac{\sqrt{\delta}}{2(m+1)} + \frac{1}{2} - \frac{n}{2},$  (3.9a)

$$\times \left\{ t^{m - \frac{\mu + m}{2}} (1 + \log t)^{\frac{1}{2}}, \quad if \quad k = \frac{\sqrt{\delta}}{2(m+1)} + \frac{1}{2} - \frac{n}{2}, \quad (3.9b) \right\}$$

$$t^{m-(m+1)(k+\frac{n}{2})+\frac{\sqrt{\delta}-\mu+1}{2}}, \ if \ k < \frac{\sqrt{\delta}}{2(m+1)} + \frac{1}{2} - \frac{n}{2}.$$
 (3.9c)

Denote by  $E_0(t, 1, x)$  and  $E_1(t, 1, x)$  the fundamental solution to (3.1) with initial data  $(u_0, u_1) = (\delta_0, 0)$  and  $(0, \delta_0)$ , respectively, where  $\delta_0$  is the Dirac function. By Duhamel's principle,  $\int_1^t E_1(t, \tau, x) *_{(x)} F(\tau, x) d\tau$  is the solution to the inhomogeneous linear Cauchy problem

$$\begin{cases} \partial_t^2 u - t^{2m} \Delta u + \frac{\mu}{t} \partial_t u + \frac{\nu^2}{t^2} u = F(t, x), & t \ge \tau \ge 1, \\ t = \tau : \ u = 0, \ \partial_t u = 0, & x \in \mathbb{R}^n, \end{cases}$$

where  $*_{(x)}$  denotes the convolution with respect to x. Introduce the Banach space

$$X(T) := \left\{ u \in C([1,T]; H^1) \cap C^1([1,T]; L^2) \\ \text{such that } \operatorname{supp} u(t, \cdot) \subset B_{\phi(t) - \phi(1) + M} \text{ for all } t \in [1,T] \right\}$$
(3.10)

equipped with the norm  $||u||_{X(T)} = \max_{t \in [1,T]} M[u](t)$ , where  $M[u](t) := ||u(t, \cdot)||_{L^2} + ||\partial_t u(t, \cdot)||_{L^2} + ||\nabla u(t, \cdot)||_{L^2}$ . Let

$$X(T,K) := \left\{ u \in X(T) : \|u\|_{X(T)} \le K \right\}$$
(3.11)

for some T > 1, K > 0. We define the operator

$$\mathcal{N}: u \in X(T) \longrightarrow \mathcal{N}(u) = u^l + u^n,$$

where  $u^l = E_0(t, 1, x) *_{(x)} u_0(x) + E_1(t, 1, x) *_{(x)} u_1(x)$  is the solution to (3.1) with  $\tau = 1$ , and  $u^n = \int_1^t E_1(t, \tau, x) *_{(x)} |u(\tau, x)|^p d\tau$ , i.e.,  $\mathcal{N}(u)$  is the solution to

$$\begin{cases} \partial_t^2 u - t^{2m} \Delta u + \frac{\mu}{t} \partial_t u + \frac{\nu^2}{t^2} u = |u|^p, \quad t \ge 1, \\ u(1,x) = u_0(x), \ \partial_t u(1,x) = u_1(x), \quad x \in \mathbb{R}^n. \end{cases}$$

Once we have proved that for any  $u, \tilde{u} \in X(T, K)$ ,

$$\|\mathcal{N}(u)\|_{X(T)} \le C_{1T} \|(u_0, u_1)\|_D + C_{2T} \|u\|_{X(T)}^p, \tag{3.12}$$

$$\|\mathcal{N}(u) - \mathcal{N}(\tilde{u})\|_{X(T)} \le C_{3T} \|u - \tilde{u}\|_{X(T)} \left( \|u\|_{X(T)}^{p-1} + \|\tilde{u}\|_{X(T)}^{p-1} \right), \tag{3.13}$$

where  $C_{1T}$  is bounded,  $C_{2T}, C_{3T} \to 0$  as  $T \to 1^+$ , we can obtain the local solution to (1.1). Indeed, for sufficiently large K, we can choose T sufficiently close to 1, ensuring that  $\mathcal{N}$  maps X(T, K) into itself and  $\mathcal{N}$  is a contraction mapping, therefore, we can get the local existence and uniqueness of the solution to (1.1) in X(T) by the contraction mapping principle.

Our aim now is to prove (3.12) and (3.13) based on Lemmas 3.1-3.2. The Gagliardo-Nirenberg inequality will be used, one can refer to [10].

**Lemma 3.3.** (*Gagliardo–Nirenberg Inequality*) Let  $1 < p, q, r < \infty$  and  $\theta \in [0, 1]$  satisfy

$$\frac{1}{p} = \theta(\frac{1}{r} - \frac{1}{n}) + (1 - \theta)\frac{1}{q}$$

Then there exists a constant C = C(p, q, r, n) > 0 such that for any  $h \in C_0^1(\mathbb{R}^n)$ ,

$$\|h\|_{L^p} \le C \|h\|_{L^q}^{1-\theta} \|\nabla h\|_{L^r}^{\theta}.$$
(3.14)

We proceed with the proof by considering five cases. Case 1:  $\frac{\sqrt{\delta}}{2(m+1)} + \frac{1}{2} - \frac{n}{2} > 1$ , i.e.,  $\delta > (m+1)^2(n+1)^2$ . Taking k = 0, 1 in (3.8c) and k = 1 in (3.9c) respectively, we get

$$\begin{aligned} \|u^{l}(t,\cdot)\|_{L^{2}} &\leq C_{m}t^{-(m+1)\frac{n}{2}+\frac{\sqrt{\delta}-\mu+1}{2}}\|(u_{0},u_{1})\|_{D^{0}} \leq C_{m}t^{\frac{\sqrt{\delta}+1}{2}}\|(u_{0},u_{1})\|_{D}, \\ \|\nabla u(t,\cdot)\|_{L^{2}} &\leq C_{m}t^{-(m+1)(1+\frac{n}{2})+\frac{\sqrt{\delta}-\mu+1}{2}}\|(u_{0},u_{1})\|_{D} \leq C_{m}t^{\frac{\sqrt{\delta}+1}{2}}\|(u_{0},u_{1})\|_{D}, \\ \|\partial_{t}u(t,\cdot)\|_{L^{2}} &\leq C_{m}t^{m-(m+1)(1+\frac{n}{2})+\frac{\sqrt{\delta}-\mu+1}{2}}\|(u_{0},u_{1})\|_{D} \leq C_{m}t^{\frac{\sqrt{\delta}+1}{2}}\|(u_{0},u_{1})\|_{D}, \end{aligned}$$

which yields

$$M[u^{l}](t) \le C_{m} t^{\frac{\sqrt{\delta}+1}{2}} \|(u_{0}, u_{1})\|_{D}.$$
(3.15)

By Lemma 3.3, Hölder's inequality and the definition of X(T), we have

$$\begin{aligned} \||u(\tau,\cdot)|^{p}\|_{L^{2}} &= \|u(\tau,\cdot)\|_{L^{2p}}^{p} \\ &\leq C \|u(\tau,\cdot)\|_{L^{2}}^{\theta(2p)p} \|\nabla u(\tau,\cdot)\|_{L^{2}}^{(1-\theta(2p))p} \leq CM[u]^{p}(\tau) \end{aligned}$$
(3.16)

and

$$\begin{aligned} \||u(\tau,\cdot)|^{p}\|_{L^{1}} &= \|u(\tau,\cdot)\|_{L^{p}}^{p} \leq C \|u(\tau,\cdot)\|_{L^{2p}}^{p} (\phi(\tau) - \phi(1) + M)^{\frac{n}{2}} \\ &\leq C\tau^{\frac{(m+1)n}{2}} \|u(\tau,\cdot)\|_{L^{2}}^{\theta(2p)p} \|\nabla u(\tau,\cdot)\|_{L^{2}}^{(1-\theta(2p))p} \leq C\tau^{\frac{(m+1)n}{2}} M[u]^{p}(\tau), \end{aligned}$$
(3.17)

where

$$\theta(2p) = \frac{n(p-1)}{2p} \tag{3.18}$$

and the condition (2.4) ensures that  $\theta(2p) \in [0,1]$ . Then by taking k = 0 in (3.6c), using Duhamel's principle and (3.16)-(3.17), we can estimate  $||u^n(t,\cdot)||_{L^2}$  as

The estimates of  $\|\nabla u(t,\cdot)\|_{L^2}$  and  $\|\partial_t u(t,\cdot)\|_{L^2}$  can be derived by taking k = 1 in (3.6c) and (3.7c) and we list the results as

$$\|\nabla u^{n}(t,\cdot)\|_{L^{2}} \leq C_{m} t^{\frac{\sqrt{\delta}}{2}+1} \int_{1}^{t} M[u]^{p}(\tau) d\tau, \qquad (3.20)$$

$$\|\partial_t u^n(t,\cdot)\|_{L^2} \le C_m t^{\frac{\sqrt{\delta}}{2}} \int_1^t M[u]^p(\tau) d\tau.$$
 (3.21)

Combining (3.19)-(3.21) yields

$$M[u^{n}](t) \le C_{m} t^{\frac{\sqrt{\delta}}{2} + 1} \int_{1}^{t} M[u]^{p}(\tau) d\tau.$$
(3.22)

It follows from (3.15) and (3.22) that  $M[\mathcal{N}u](t) \leq C_m t^{\frac{\sqrt{\delta}}{2}+1} (\|(u_0, u_1)\|_D + \int_1^t M[u]^p(\tau) d\tau)$ , which indicates the validness of (3.12), where  $C_{1T} = C_m T^{\frac{\sqrt{\delta}}{2}+1}$  and  $C_{2T} = C_m T^{\frac{\sqrt{\delta}}{2}+1}(T-1)$ .

and

$$\begin{aligned} \||u(\tau,\cdot)|^{p} - |\tilde{u}(\tau,\cdot)|^{p}\|_{L^{1}} &\leq C \||u(\tau,\cdot)|^{p} - |\tilde{u}(\tau,\cdot)|^{p}\|_{L^{2}} \left(\phi(t) - \phi(1) + M\right)^{\frac{n}{2}} \\ &\leq C\tau^{\frac{(m+1)n}{2}} M[u - \tilde{u}](\tau) \left(M[u]^{p-1}(\tau) + M[\tilde{u}]^{p-1}(\tau)\right), \end{aligned}$$
(3.24)

where  $\theta(2p)$  is defined by (3.18). Taking k = 0 in (3.6c), using Duhamel's principle and (3.23)-(3.24), we have

$$\begin{split} \|\mathcal{N}u(t,\cdot) - \mathcal{N}\tilde{u}(t,\cdot)\|_{L^{2}} &\leq C_{m}t^{-(m+1)\frac{n}{2} + \frac{\sqrt{\delta} - \mu + 1}{2}} \int_{1}^{t} \tau^{-\frac{\sqrt{\delta} - \mu - 1}{2}} \\ &\times \left(\||u(\tau,\cdot)|^{p} - |\tilde{u}(\tau,\cdot)|^{p}\|_{L^{1}} + \tau^{(m+1)\frac{n}{2}}\||u(\tau,\cdot)|^{p} - |\tilde{u}(\tau,\cdot)|^{p}\|_{L^{2}}\right)d\tau \\ &\leq C_{m}t^{-(m+1)\frac{n}{2} + \frac{\sqrt{\delta} - \mu + 1}{2}} \\ &\times \int_{1}^{t} \tau^{\frac{(m+1)n}{2} - \frac{\sqrt{\delta} - \mu - 1}{2}} M[u - \tilde{u}](\tau) \left(M[u]^{p-1}(\tau) + M[\tilde{u}]^{p-1}(\tau)\right)d\tau \\ &\leq C_{m}t^{\frac{\sqrt{\delta}}{2} + 1} \int_{1}^{t} M[u - \tilde{u}](\tau) \left(M[u]^{p-1}(\tau) + M[\tilde{u}]^{p-1}(\tau)\right)d\tau. \end{split}$$
(3.25)

By taking k = 1 in (3.6c) and (3.7c), we obtain the estimates of  $\|\nabla (\mathcal{N}u(t, \cdot) - \mathcal{N}\tilde{u}(t, \cdot))\|_{L^2}$ and  $\|\partial_t (\mathcal{N}u(t, \cdot) - \mathcal{N}\tilde{u}(t, \cdot))\|_{L^2}$  as

$$\begin{aligned} \|\nabla \left(\mathcal{N}u(t,\cdot) - \mathcal{N}\tilde{u}(t,\cdot)\right)\|_{L^{2}} &\leq C_{m}t^{\frac{\sqrt{\delta}}{2}+1}\int_{1}^{t}M[u-\tilde{u}](\tau)\left(M[u]^{p-1}(\tau) + M[\tilde{u}]^{p-1}(\tau)\right)d\tau, \\ \|\partial_{t}\left(\mathcal{N}u(t,\cdot) - \mathcal{N}\tilde{u}(t,\cdot)\right)\|_{L^{2}} &\leq C_{m}t^{\frac{\sqrt{\delta}}{2}}\int_{1}^{t}M[u-\tilde{u}](\tau)\left(M[u]^{p-1}(\tau) + M[\tilde{u}]^{p-1}(\tau)\right)d\tau. \end{aligned}$$

$$(3.26)$$

We infer by (3.25) and (3.26) that

$$M[\mathcal{N}u - \mathcal{N}\tilde{u}](t) \le C_m t^{\frac{\sqrt{\delta}}{2} + 1} \int_1^t M[u - \tilde{u}](\tau) \big( M[u]^{p-1}(\tau) + M[\tilde{u}]^{p-1}(\tau) \big) d\tau,$$

which indicates that (3.13) holds, where  $C_{3T} = C_m T^{\frac{\sqrt{\delta}}{2}+1}(T-1)$ .

Applying the same argument, we can deal with the remaining cases, so no further details will be provided.

Case 2: 
$$\frac{\sqrt{\delta}}{2(m+1)} + \frac{1}{2} - \frac{n}{2} = 1$$
, i.e.,  $\delta = (m+1)^2(n+1)^2$ .  
Case 3:  $0 < \frac{\sqrt{\delta}}{2(m+1)} + \frac{1}{2} - \frac{n}{2} < 1$ , i.e.,  $(m+1)^2(n-1)^2 < \delta < (m+1)^2(n+1)^2$ .  
Case 4:  $\frac{\sqrt{\delta}}{2(m+1)} + \frac{1}{2} - \frac{n}{2} = 0$ ,  $n > 1$ , i.e.,  $\delta = (m+1)^2(n-1)^2$ ,  $n > 1$ .  
Case 5:  $\frac{\sqrt{\delta}}{2(m+1)} + \frac{1}{2} - \frac{n}{2} < 0$ ,  $n > 1$ , i.e.,  $0 < \delta < (m+1)^2(n-1)^2$ ,  $n > 1$ .

# 4. The Proof of Theorem 2.1

4.1. Test functions and some asymptotic properties. In this section, we will provide solutions to the conjugate equation of (1.1)

$$\partial_t^2 \Phi(t,x) - t^{2m} \Delta \Phi(t,x) - \partial_t \left(\frac{\mu}{t} \Phi(t,x)\right) + \frac{\nu^2}{t^2} \Phi(t,x) = 0.$$
(4.1)

Inspired by [27], we want to express the solution to (4.1) in terms of Gaussian hypergeometric functions. For  $\beta \in \mathbb{R}$ , define

$$\Phi_{\beta}(t,x) = t^{-\beta+1} \psi_{\beta} \left( \frac{(m+1)^2 |x|^2}{t^{2(m+1)}} \right), \ t \ge 1,$$
(4.2)

where  $\psi_{\beta} \in C^2([0, \frac{1}{m+1}))$ . For brevity, let  $z = \frac{(m+1)^2|x|^2}{t^{2(m+1)}}$ .

**Lemma 4.1.**  $\Phi_{\beta}(t, x)$  is a solution to (4.1) if and only if  $\psi_{\beta}$  is a solution to

$$z(1-z)\psi_{\beta}''(z) + \left[\frac{n}{2} - \left(\frac{\beta-1}{m+1} + \frac{1}{2(m+1)} + \frac{\mu}{2(m+1)} + 1\right)z\right]\psi_{\beta}'(z) - \frac{\beta(\beta+\mu-1)+\nu^2}{4(m+1)^2}\psi_{\beta}(z) = 0.$$
(4.3)

Proof. Substituting

$$\begin{aligned} \partial_t^2 \Phi_\beta(t,x) &= \beta(\beta-1)t^{-\beta-1}\psi_\beta(z) + 4(m+1)^2t^{-\beta-1}z^2\psi_\beta''(z) \\ &+ \left[2(m+1)(\beta-1) + 2(m+1)\beta + 4(m+1)^2\right]t^{-\beta-1}z\psi_\beta'(z), \\ \Delta\Phi_\beta(t,x) &= 4(m+1)^2zt^{-2m}t^{-\beta-1}\psi_\beta''(z) + 2n(m+1)^2t^{-2m}t^{-\beta-1}\psi_\beta'(z), \\ \frac{\partial}{\partial t}\left(\frac{\mu}{t}\Phi_\beta(t,x)\right) &= \left(-\mu + \mu(-\beta+1)\right)t^{-\beta-1}\psi_\beta(z) - 2(m+1)\mu t^{-\beta-1}z\psi_\beta'(z) \end{aligned}$$

into (4.1), we get

$$t^{-\beta-1} \Big\{ \Big[ 4(m+1)^2 z^2 - 4(m+1)^2 z \Big] \psi_{\beta}''(z) \\ + \Big[ \Big( 2(m+1)(2\beta-1) + 4(m+1)^2 \Big) z - 2(m+1)^2 n + 2(m+1)\mu z \Big] \psi_{\beta}'(z) \\ + \Big[ \beta(\beta-1) + \mu + \mu(\beta-1) + \nu^2 \Big] \psi_{\beta}(z) \Big\} = 0,$$

which implies (4.3).

As is well known, when |z| < 1,

$$F(a, b, c; z) = \sum_{k=0}^{\infty} \frac{(a)_k(b)_k}{k!(c)_k} z^k$$
(4.4)

is a solution to the Gauss hypergeometric equation

$$z(1-z)h''(z) + [\gamma - (\alpha + \beta + 1)z]h'(z) - \alpha\beta h(z) = 0,$$
  
where  $(d)_k = \begin{cases} 1, & k = 0, \\ \prod_{j=1}^k (d+j-1), & k \ge 0 \end{cases}$  and  $F(a, b, c; z)$  is the Gauss hypergeometric

function with parameters (a, b, c).

**Lemma 4.2.** Assume that |z| < 1 and  $\psi_{\beta}(z)$  is a solution to (4.3), then

$$\psi_{\beta}(z) = F(a, b, c; z) = \sum_{k=0}^{\infty} \frac{(a)_k(b)_k}{k!(c)_k} z^k,$$
(4.5)

where  $a = \frac{2\beta + \mu - 1 + \sqrt{\delta}}{4(m+1)}$ ,  $b = \frac{2\beta + \mu - 1 - \sqrt{\delta}}{4(m+1)}$  and  $c = \frac{n}{2}$ .

Proof. Consider the quadratic equation

$$s^{2} - \frac{2\beta + \mu - 1}{2(m+1)}s + \frac{\beta(\beta + \mu - 1) + \nu^{2}}{4(m+1)^{2}} = 0,$$

since  $\frac{(2\beta+\mu-1)^2}{4(m+1)^2} - \frac{\beta(\beta+\mu-1)+\nu^2}{(m+1)^2} = \frac{\delta}{4(m+1)^2} > 0$ , then  $a = \frac{2\beta+\mu-1+\sqrt{\delta}}{4(m+1)}$  and  $b = \frac{2\beta+\mu-1-\sqrt{\delta}}{4(m+1)}$  are the two real roots. Moreover,

$$\begin{cases} a+b+1 = \frac{\beta-1}{m+1} + \frac{1}{2(m+1)} + \frac{\mu}{2(m+1)} + 1, \\ ab = \frac{\beta(\beta+\mu-1)+\nu^2}{4(m+1)^2}, \end{cases}$$

then (4.3) is transformed into

$$z(1-z)\psi_{\beta}''(z) + \left[\frac{n}{2} - (a+b+1)z\right]\psi_{\beta}'(z) - ab\psi_{\beta}(z) = 0,$$

which shows (4.5) by (4.4).

 -	-	_	

From Lemmas 4.1 and 4.2, we know that

$$\Phi_{\beta}(t,x) = t^{-\beta+1}\psi_{\beta}(z) = t^{-\beta+1}F(a,b,c;z)$$
(4.6)

is a solution to (4.1), where a, b, c are given by Lemma 4.2.

The following lemma gives the asymptotic behavior of  $\psi_{\beta}(z)$  and  $\psi'_{\beta}(z)$ .

**Lemma 4.3.** (i) If  $\frac{1+\sqrt{\delta}-\mu}{2} < \beta < \frac{(m+1)n-\mu+1}{2}$ , there exists a  $C = C(m, n, \beta, \mu, \nu) > 1$  such that for any  $z \in [0, 1)$ ,

$$1 \le \psi_{\beta}(z) \le C. \tag{4.7}$$

(ii) If  $\beta > \frac{(m+1)(n-2)-\mu+1}{2}$ , there exists a  $C = C(m, n, \beta, \mu, \nu) > 1$  such that for any  $z \in [0, 1)$ ,

$$C^{-1}(1-\sqrt{z})^{\frac{(m+1)(n-2)-\mu+1-2\beta}{2(m+1)}} \le |\psi_{\beta}'(z)| \le C(1-\sqrt{z})^{\frac{(m+1)(n-2)-\mu+1-2\beta}{2(m+1)}}.$$
 (4.8)

*Proof.* (i)  $\frac{1+\sqrt{\delta}-\mu}{2} < \beta < \frac{(m+1)n-\mu+1}{2}$  guarantees that a > 0, b > 0 and c - a - b > 0. By the formula (15.4.20) in [22]:

$$F(a, b, c; 1) = \frac{\Gamma(c)\Gamma(c - a - b)}{\Gamma(c - a)\Gamma(c - b)} \text{ for } Re(c - a - b) > 0,$$

we deduce that (4.7) holds.

(ii) Note the formula (15.5.1) in [22]:

$$\frac{d}{dz}F(a, b, c; z) = \frac{ab}{c}F(a+1, b+1, c+1; z),$$

so

$$\psi'_{\beta}(z) = \frac{ab}{c}F(a+1,b+1,\frac{n}{2}+1;z).$$

Then the formula (15.4.23) in [22]:

$$\lim_{z \to 1^-} \frac{F(a,b,c;z)}{(1-z)^{c-a-b}} = \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} \text{ for } Re(c-a-b) < 0$$

shows that

$$\lim_{z \to 1^{-}} \frac{\psi_{\beta}'(z)}{(1-z)^{\frac{n}{2}+1-(a+1)-(b+1)}} = \lim_{z \to 1^{-}} \frac{2ab}{n} \frac{F(a+1,b+1,\frac{n}{2}+1;z)}{(1-z)^{\frac{n}{2}-a-b-1}}$$
$$= \lim_{z \to 1^{-}} \frac{2ab}{n} \frac{F(a+1,b+1,\frac{n}{2}+1;z)}{(1-z)^{\frac{(m+1)(n-2)-\mu+1-2\beta}{2(m+1)}}} = \frac{\Gamma(\frac{n}{2}+1)\Gamma(a+b+1-\frac{n}{2})}{\Gamma(a+1)\Gamma(b+1)} > 0,$$

where we used that  $\frac{n}{2} + 1 - (a+1) - (b+1) = \frac{(m+1)(n-2)-2\beta-\mu+1}{2(m+1)} < 0$  because  $\beta > \frac{(m+1)(n-2)-\mu+1}{2}$ . Then  $C^{-1}(1-z) \frac{(m+1)(n-2)-\mu+1-2\beta}{2(m+1)} \le |\psi_{\beta}'(z)| \le C(1-z) \frac{(m+1)(n-2)-\mu+1-2\beta}{2(m+1)}$ 

holds for any  $z \in [0, 1)$ , therefore (4.8) holds by noting  $1 + \sqrt{z}$  is bounded from above and below for  $z \in [0, 1)$ .

4.2. **Some preliminary estimates.** Before starting to prove Theorem 2.1, we will provide some preliminary results. Let

$$H_{\beta}(t) = \int_{\mathbb{R}^n} |u(t,x)|^p \Phi_{\beta}(t,x) dx, \qquad (4.9)$$

$$I_{\beta}(t) = \int_{1}^{t} (t-s)sH_{\beta}(s)ds,$$
(4.10)

$$J_{\beta}(t) = \int_{1}^{t} (1+s)^{-3} I_{\beta}(s) ds, \qquad (4.11)$$

where  $\beta \in (\frac{1+\sqrt{\delta}-\mu}{2}, \frac{(m+1)n-\mu+1}{2})$ ,  $\Phi_{\beta}(t, x)$  is a solution of (4.1) as constructed in Section 4.1 and  $t \ge 1$ . We will show that  $J_{\beta}(t)$  blows up before  $e^{c\varepsilon^{-p(p-1)}}$ , hence,  $I_{\beta}(t)$  and  $H_{\beta}(t)$  as well.

**Remark 4.1.** The condition  $\delta < (m+1)^2 n^2$  in Theorem 2.1 guarantees the existence of  $\beta \in (\frac{1+\sqrt{\delta}-\mu}{2}, \frac{(m+1)n-\mu+1}{2})$ , and as mentioned in Section 1, the range of  $\delta$  reflects the competitive relationship between the Fujita indice and the Strauss one.

Now we give some preliminary estimates.

**Lemma 4.4.** For any 
$$\beta \in (\frac{1+\sqrt{\delta}-\mu}{2}, \frac{(m+1)n-\mu+1}{2})$$
 and  $t \ge 1$ ,  
 $t^2 J_{\beta}(t) \le \frac{1}{2} \int_{1}^{t} (t-s)^2 H_{\beta}(s) ds.$ 

*Proof.* Differentiating (4.10), we have

$$I'_{\beta}(t) = \int_{1}^{t} s H_{\beta}(s) ds \quad \text{and} \quad I''_{\beta}(t) = t H_{\beta}(t).$$

Since  $I_{\beta}(1) = I'_{\beta}(1) = 0$ , it follows from integration by parts that

$$\int_{1}^{t} (t-s)^{2} H_{\beta}(s) ds = \int_{1}^{t} (t-s)^{2} s^{-1} I_{\beta}''(s) ds = -\int_{1}^{t} \frac{\partial}{\partial s} \left( (t-s)^{2} s^{-1} \right) I_{\beta}'(s) ds$$
$$= \int_{1}^{t} \frac{\partial^{2}}{\partial s^{2}} \left( (t-s)^{2} s^{-1} \right) I_{\beta}(s) ds = \int_{1}^{t} \frac{2t^{2}}{s^{3}} I_{\beta}(s) ds \ge 2t^{2} J_{\beta}(t).$$

**Lemma 4.5.** Suppose that  $(u_0, u_1) \in D$  be nonnegative, not identically zero and  $supp(u_0, u_1) \subset \{x : |x| \leq M\}$  for some  $M < \frac{1}{m+1}$ , then for any  $\beta \in (\frac{1+\sqrt{\delta}-\mu}{2}, \frac{(m+1)n-\mu+1}{2}) \cap [1-\mu, +\infty), \delta < (m+1)^2n^2$  and  $t \geq 1$ , we have

$$\varepsilon \mathcal{E}_{0,\beta,m}(u_0) + \varepsilon \mathcal{E}_{1,\beta,m}(u_0, u_1)(t-1) + \int_1^t (t-s) H_\beta(s) ds$$

$$= \int_{\mathbb{R}^n} u(t,x) \Phi_\beta(t,x) dx + \int_1^t s^{-\beta} \int_{\mathbb{R}^n} u(s,x) \overline{\psi_\beta} \big(\frac{(m+1)^2 |x|^2}{s^{2(m+1)}}\big) dx ds,$$
(4.12)

where

$$\begin{split} \overline{\psi}_{\beta}(z) &= (2\beta + \mu - 2)\psi_{\beta}(z) + 4(m+1)z\psi_{\beta}'(z), \\ \mathcal{E}_{0,\beta,m}(u_0) &= \int_{\mathbb{R}^n} u_0(x)\psi_{\beta}\big((m+1)^2|x|^2\big)dx \ge 0, \\ \mathcal{E}_{1,\beta,m}(u_0,u_1) &= \int_{\mathbb{R}^n} u_1(x)\psi_{\beta}\big((m+1)^2|x|^2\big)dx \\ &+ \int_{\mathbb{R}^n} u_0(x)\Big[2(m+1)^3|x|^2\psi_{\beta}'\big((m+1)^2|x|^2\big) + (\mu+\beta-1)\psi_{\beta}\big((m+1)^2|x|^2\big)\Big]dx \ge 0. \end{split}$$

**Remark 4.2.** Noting that  $1-\mu \leq \frac{(m+1)n-\mu+1}{2}$ , so  $(\frac{1+\sqrt{\delta}-\mu}{2}, \frac{(m+1)n-\mu+1}{2}) \cap [1-\mu, \infty) \neq \emptyset$ . We claim that  $\mathcal{E}_{0,\beta,m}(u_0)$  and  $\mathcal{E}_{1,\beta,m}(u_0, u_1)$  are nonnegative. Indeed, since  $\beta \in (\frac{1+\sqrt{\delta}-\mu}{2}, \frac{(m+1)n-\mu+1}{2})$  and  $\psi_{\beta}((m+1)^2|x|^2) = F(a, b; \frac{n}{2}; (m+1)^2|x|^2)$ , (4.7) shows that  $\mathcal{E}_{0,\beta,m}(u_0) \geq 0$ ; obviously,  $\psi'_{\beta}((m+1)^2|x|^2) = \frac{2ab}{n}F(a+1,b+1; \frac{n}{2}; (m+1)^2|x|^2) > 0$ ,  $\beta \geq 1-\mu$  and the nonnegativity of  $u_0, u_1$  imply that  $\mathcal{E}_{1,\beta,m}(u_0, u_1) \geq 0$ .

*Proof of Lemma 4.5.* Note (4.1), (4.9) and substitute  $\Phi_{\beta}(t, x)$  into (2.3), we obtain

### Y.- Q. LI AND F. GUO

$$\int_{\mathbb{R}^n} \left( \partial_t u(t,x) \Phi_\beta(t,x) - u(t,x) \partial_t \Phi_\beta(t,x) + \frac{\mu}{t} u(t,x) \Phi_\beta(t,x) \right) dx$$

$$= \int_1^t H_\beta(s) ds + \varepsilon \int_{\mathbb{R}^n} \left( -u_0(x) \partial_t \Phi_\beta(1,x) + \left( \mu u_0(x) + u_1(x) \right) \Phi_\beta(1,x) \right) dx.$$
(4.13)

Since  $\partial_t \Phi_\beta(1,x) = (-\beta + 1)\psi_\beta\left((m+1)^2|x|^2\right) - 2(m+1)^3|x|^2\psi'_\beta\left((m+1)^2|x|^2\right)$  and  $\Phi_\beta(1,x) = \psi_\beta\left((m+1)^2|x|^2\right)$ , then

$$\begin{split} &\int_{\mathbb{R}^n} \left( -u_0(x) \partial_t \varPhi_\beta(1, x) + \left( \mu u_0(x) + u_1(x) \right) \varPhi_\beta(1, x) \right) dx \\ &= \int_{\mathbb{R}^n} u_1(x) \psi_\beta \left( (m+1)^2 |x|^2 \right) dx \\ &+ \int_{\mathbb{R}^n} u_0(x) \left[ 2(m+1)^3 |x|^2 \psi_\beta' \left( (m+1)^2 |x|^2 \right) + (\mu+\beta-1) \psi_\beta \left( (m+1)^2 |x|^2 \right) \right] \\ &= \mathcal{E}_{1,\beta,m}(u_0, u_1) \ge 0. \end{split}$$
(4.14)

Further calculation leads to

$$\int_{\mathbb{R}^n} \left( \partial_t u(t,x) \Phi_\beta(t,x) - u(t,x) \partial_t \Phi_\beta(t,x) + \frac{\mu}{t} u(t,x) \Phi_\beta(t,x) \right) dx$$

$$= \frac{d}{dt} \int_{\mathbb{R}^n} u(t,x) \Phi_\beta(t,x) dx + t^{-\beta} \int_{\mathbb{R}^n} \overline{\psi_\beta}(z) u(t,x) dx.$$
(4.15)

If follows from (4.13)-(4.15) that

$$\frac{d}{dt} \int_{\mathbb{R}^n} u(t,x) \Phi_{\beta}(t,x) dx + t^{-\beta} \int_{\mathbb{R}^n} \overline{\psi_{\beta}}(z) u(t,x) dx = \int_1^t H_{\beta}(s) ds + \varepsilon \mathcal{E}_{1,\beta,m}(u_0,u_1).$$

A further integration by parts over [1, t] provides that

$$\begin{split} \int_{\mathbb{R}^n} u(t,x) \varPhi_{\beta}(t,x) dx &- \varepsilon \mathcal{E}_{0,\beta,m}(u_0) + \int_1^t s^{-\beta} \int_{\mathbb{R}^n} u(s,x) \overline{\psi_{\beta}} \Big( \frac{(m+1)^2 |x|^2}{s^{2(m+1)}} \Big) dx ds \\ &= \varepsilon \mathcal{E}_{1,\beta,m}(u_0,u_1)(t-1) + \int_1^t (t-s) H_{\beta}(s) ds, \end{split}$$

which implies (4.12).

For any q > 1, consider

$$\beta_q = \frac{(m+1)n - \mu + 1}{2} - \frac{m+1}{q} \tag{4.16}$$

and let p' be the conjugate number of p, i.e.,  $\frac{1}{p} + \frac{1}{p'} = 1$ .

**Lemma 4.6.** Under the assumptions of Lemma 4.5, let  $p > \frac{2(m+1)}{(m+1)n-\sqrt{\delta}}$  and when n = 1, we assume  $\mu \ge m$ .

(i) When q > p, there exists a  $C = C(m, n, \beta, \mu, \nu) > 0$  such that

$$\varepsilon \mathcal{E}_{0,\beta_{q},m}(u_{0}) + \varepsilon \mathcal{E}_{1,\beta_{q},m}(u_{0},u_{1})(t-1) + \int_{1}^{t} (t-s)H_{\beta_{q}}(s)ds$$

$$\leq Ct^{-\beta_{q}+1+\frac{(m+1)n}{p'}} \|u(t,\cdot)\|_{L^{p}(\mathbb{R}^{n})} + C\int_{1}^{t} s^{-\beta_{p}+\frac{(m+1)n}{p'}} \|u(s,\cdot)\|_{L^{p}(\mathbb{R}^{n})}ds.$$
(4.17)

(ii) When q = p, there exists a  $C = C(m, n, \beta, \mu, \nu) > 0$  such that

$$\int_{1}^{t} (t-s)H_{\beta_{p}}(s)ds \leq Ct^{-\beta_{p}+1+\frac{(m+1)n}{p'}} \|u(t,\cdot)\|_{L^{p}(\mathbb{R}^{n})} + C\int_{1}^{t} s^{\frac{(m+1)n}{p'}-\beta_{p}} (\log s)^{\frac{1}{p'}} \|u(s,\cdot)\|_{L^{p}(\mathbb{R}^{n})} ds.$$
(4.18)

**Remark 4.3.** The proof of Lemma 4.6 is based on Lemmas 4.3 and 4.5, so we need to guarantee that

$$|z| = \frac{(m+1)^2 |x|^2}{t^{2(m+1)}} < 1,$$
  
$$\beta_p = \frac{(m+1)n - \mu + 1}{2} - \frac{m+1}{p} \in \left(\frac{1 + \sqrt{\delta} - \mu}{2}, \frac{(m+1)n - \mu + 1}{2}\right)$$
(4.19)

and

$$\beta_p \ge 1 - \mu. \tag{4.20}$$

Indeed, the finite propagation speed of waves implies

$$supp \ u(t, \cdot) \subset \{x : |x| \le \phi(t) - \phi(1) + M\},$$
 (4.21)

*note that*  $M < \frac{1}{m+1} = \phi(1)$ *, so*  $|x| \le \frac{t^{m+1}}{m+1} - \frac{1}{m+1} + M < \frac{t^{m+1}}{m+1}$ *, then*  $|z| = \frac{(m+1)^2 |x|^2}{t^{2(m+1)}} < 1$ *. In addition, (4.19) is equal to*  $p > \frac{2(m+1)}{(m+1)n-\sqrt{\delta}}$ *. Hence we need to check (4.20). In fact,* 

$$\beta_p \ge 1 - \mu \iff \frac{2(m+1)}{p} \le (m+1)n + \mu - 1.$$

Since  $p = p_S(n + \frac{\mu}{m+1}, m)$ , by (1.8) we have

$$\frac{2(m+1)}{p} = p[(m+1)n + \mu - 1] - [(m+1)(n-2) + \mu + 3]$$

and

$$p = \frac{(m+1)(n-2)+\mu+3+\sqrt{[(m+1)(n-2)+\mu+3]^2+8(m+1)[(m+1)n+\mu-1]}}{2[(m+1)n+\mu-1]},$$

then

$$\frac{2(m+1)}{p} \le (m+1)n + \mu - 1$$
  

$$\iff p[(m+1)n + \mu - 1] - [(m+1)(n-2) + \mu + 3] \le (m+1)n + \mu - 1$$
  

$$\iff (m+1)^2(n^2 - 2n) + 2(m+1)(n\mu + 1 - \mu) + \mu^2 - 1 \ge 0,$$
(4.22)

this holds obviously for any  $n \ge 2$ . And when n = 1,

$$(4.22) \iff -(m+1)^2 + 2(m+1) + \mu^2 - 1 \ge 0 \iff \mu^2 \ge m^2 \iff \mu \ge m.$$

Now we give the proof of Lemma 4.6.

*Proof*. According to the assumptions on q in (i) and (ii), we know that  $\beta_q \geq 1 - \mu$  and  $\beta_q \in (\frac{1+\sqrt{\delta}-\mu}{2}, \frac{(m+1)n-\mu+1}{2})$ . By Lemma 4.5, we have

$$\begin{split} \varepsilon \mathcal{E}_{0,\beta_q,m}(u_0) &+ \varepsilon \mathcal{E}_{1,\beta_q,m}(u_0,u_1)(t-1) + \int_1^t (t-s)H_{\beta_q}(s)ds \\ &= \int_{\mathbb{R}^n} u(t,x)\Phi_{\beta_q}(t,x)dx + \int_1^t s^{-\beta_q} \int_{\mathbb{R}} u(s,x)\overline{\psi_{\beta_q}}\big(\frac{(m+1)^2|x|^2}{s^{2(m+1)}}\big)dxds \\ &:= \Omega_1(t) + \int_1^t \Omega_2(s)ds, \end{split}$$

where

$$\Omega_1(t) = \int_{\mathbb{R}^n} u(t, x) \Phi_{\beta_q}(t, x) dx,$$
  
$$\Omega_2(t) = t^{-\beta_q} \int_{\mathbb{R}} u(t, x) \overline{\psi}_{\beta_q} \left(\frac{(m+1)^2 |x|^2}{t^{2(m+1)}}\right) dx$$

with  $z = \frac{(m+1)^2 |x|^2}{t^{2(m+1)}}$ . Hölder's inequality implies that

$$\Omega_1(t) \le \left(\int_{\mathbb{R}^n} |u(t,x)|^p dx\right)^{\frac{1}{p}} \left(\int_{\mathbb{R}^n} |\Phi_{\beta_q}(t,x)|^{p'} dx\right)^{\frac{1}{p'}}.$$
(4.23)

By (i) of Lemma 4.3,

$$\Phi_{\beta_q}(t,x) = t^{-\beta_q+1}\psi_{\beta_q}(z) \le Ct^{-\beta_q+1},$$

so

$$\Omega_{1}(t) \leq C \|u(t,\cdot)\|_{L^{p}(\mathbb{R}^{n})} t^{-\beta_{q}+1} \left(\int_{|x| \leq \phi(t)-\phi(1)+M} dx\right)^{\frac{1}{p'}} \\ \leq C t^{-\beta_{q}+1+\frac{(m+1)n}{p'}} \|u(t,\cdot)\|_{L^{p}(\mathbb{R}^{n})}.$$

By Hölder's inequality,

$$\Omega_2(t) \le t^{-\beta_q} \|u(t,\cdot)\|_{L^p(\mathbb{R}^n)} \Big[ \int_{\mathbb{R}^n} (\overline{\psi}_{\beta_q}(z))^{p'} dx \Big]^{\frac{1}{p'}},$$

by Lemma 4.3,  $|(2\beta+\mu-2)\psi_{\beta_q}(z)|\leq C$  and

$$|\psi'_{\beta_q}(z)| \le C(1-\sqrt{z})^{\frac{(m+1)(n-2)-\mu+1-2\beta_q}{2(m+1)}},$$

then

$$\begin{aligned} |\overline{\psi}_{\beta_q}(z)| &= |(2\beta + \mu - 2)\psi_{\beta_q}(z) + 4(m+1)z\psi'_{\beta_q}(z)| \\ &\leq C \Big(1 - \frac{(m+1)|x|}{t^{m+1}}\Big)^{\frac{(m+1)(n-2)-\mu+1-2\beta_q}{2(m+1)}}, \end{aligned}$$

so

$$\Omega_{2}(t) \leq Ct^{-\beta_{q}} \|u(t,\cdot)\|_{L^{p}(\mathbb{R}^{n})} \left[ \int_{|x| \leq \phi(t) - \phi(1) + M} \left( 1 - \frac{(m+1)|x|}{t^{m+1}} \right)^{\frac{(m+1)(n-2)-\mu+1-2\beta_{q}}{2(m+1)}p'} dx \right]^{\frac{1}{p'}} \\
= Ct^{-\beta_{q}} \|u(t,\cdot)\|_{L^{p}(\mathbb{R}^{n})} \left( \int_{0}^{\frac{(m+1)(\phi(t)-\phi(1)+M)}{t^{m+1}}} (1-\rho)^{-\frac{p'}{q'}} \left(\frac{t^{m+1}}{m+1}\rho\right)^{n-1} \frac{t^{m+1}}{m+1} d\rho \right)^{\frac{1}{p'}} \\
\leq Ct^{-\beta_{q} + \frac{(m+1)n}{p'}} \|u(t,\cdot)\|_{L^{p}(\mathbb{R}^{n})} \left( \int_{0}^{\frac{(m+1)(\phi(t)-\phi(1)+M)}{t^{m+1}}} (1-\rho)^{-\frac{p'}{q'}} d\rho \right)^{\frac{1}{p'}},$$
(4.24)

here we calculated that  $\frac{(m+1)(n-2)-\mu+1-2\beta_q}{2(m+1)}p' = -\frac{p'}{q'}$  and  $\rho \leq 1$  because of  $M \leq \frac{1}{m+1} = \phi(1)$ ,

$$\rho = \frac{(m+1)r}{t^{m+1}} = \frac{(m+1)|x|}{t^{m+1}} \le (m+1)\frac{\phi(t) - \phi(1) + M}{t^{m+1}} \le (m+1)\frac{\phi(t)}{t^{m+1}} = 1.$$

When q > p, we have  $-\frac{p'}{q'} < -1$ , so

$$\int_{0}^{\frac{(m+1)(\phi(t)-\phi(1)+M)}{t^{m+1}}} (1-\rho)^{-\frac{p'}{q'}} d\rho \le C \left(1 - \frac{(m+1)(\phi(t)-\phi(1)+M)}{t^{m+1}}\right)^{1-\frac{p'}{q'}} = C \left(\frac{1-M(m+1)}{t^{m+1}}\right)^{1-\frac{p'}{q'}} \le C \left(\frac{1}{t^{m+1}}\right)^{1-\frac{p'}{q'}};$$

while q = p,

$$\int_{0}^{\frac{(m+1)(\phi(t)-\phi(1)+M)}{t^{m+1}}} (1-\rho)^{-\frac{p'}{q'}} d\rho = \int_{0}^{\frac{(m+1)(\phi(t)-\phi(1)+M)}{t^{m+1}}} \frac{1}{1-\rho} d\rho$$
$$= \log \frac{t^{m+1}}{1-(m+1)M} \le C \log t.$$

Then,

$$\Omega_{2}(t) \leq \begin{cases} Ct^{-\beta_{p} + \frac{(m+1)n}{p'}} \|u(t,\cdot)\|_{L^{p}(\mathbb{R}^{n})}, & q > p, \\ Ct^{-\beta_{q} + \frac{(m+1)n}{p'}} (\log t)^{\frac{1}{p'}} \|u(t,\cdot)\|_{L^{p}(\mathbb{R}^{n})}, & q = p. \end{cases}$$
(4.25)

Therefore,

$$\int_{1}^{t} \Omega_{2}(s) ds \leq \begin{cases} C \int_{1}^{t} s^{-\beta_{p} + \frac{(m+1)n}{p'}} \|u(s,\cdot)\|_{L^{p}(\mathbb{R}^{n})} ds, & q > p, \\ C \int_{1}^{t} s^{-\beta_{p} + \frac{(m+1)n}{p'}} (\log s)^{\frac{1}{p'}} \|u(s,\cdot)\|_{L^{p}(\mathbb{R}^{n})} ds, & q = p. \end{cases}$$
(4.26)

Consequently, (4.17) and (4.18) follow from (4.23), (4.26) and the nonnegativity of  $\mathcal{E}_{0,\beta,m}(u_0)$  and  $\mathcal{E}_{1,\beta,m}(u_0, u_1)$ .

4.3. The proof of Theorem 2.1. Now, we will derive a second-order ordinary differential inequality satisfied by  $J_{\beta}(t)$ , and then utilize the following result proved by Zhou [36] to obtain the lifespan estimate.

**Lemma 4.7.** ([36]) Let  $\sigma_0 \ge 0$  and  $U \in C^2([\sigma_0, \infty))$  be nonnegative function. Assume there exists positive numbers c, C and C' such that

$$\begin{cases} U''(\sigma) + 2U'(\sigma) \ge c\sigma^{1-p} (U(\sigma))^p, \\ U(\sigma) \ge C\varepsilon^p \sigma, \\ U'(\sigma) \ge C'\varepsilon^p. \end{cases}$$
(4.27)

Then there exists a  $\varepsilon_0 > 0$  such that for any  $\varepsilon \in (0, \varepsilon_0)$ , U blows up before  $C'' \varepsilon^{-p(p-1)}$  for some C'' > 0.

Now we prove Theorem 2.1.

*Proof*. For  $\sigma > 0$ ,  $\beta_{p+\sigma} = \frac{(m+1)n-\mu+1}{2} - \frac{m+1}{p+\sigma} \in (\frac{1+\sqrt{\delta}-\mu}{2}, \frac{(m+1)n-\mu+1}{2})$  and  $\beta_{p+\sigma} \ge 1-\mu$ . From (4.6) and (4.9), by Lemma 4.3, there exists a C > 0 such that

$$C^{-1}t^{(\beta_p-1)\frac{1}{p}} (H_{\beta_p}(t))^{\frac{1}{p}} \le \|u(t,\cdot)\|_{L^p(\mathbb{R}^n)} \le Ct^{(\beta_p-1)\frac{1}{p}} (H_{\beta_p}(t))^{\frac{1}{p}}.$$

Note

$$-\beta_{p+\sigma} + 1 + \frac{(m+1)n}{p'} + \frac{1}{p}(\beta_p - 1) = \frac{(m+1)n - \beta_p + 1}{p'} - 1$$

and

$$\frac{(m+1)n - \beta_p + 1}{p'} = \frac{1}{p} + 1 + \frac{1}{p^2} \frac{((m+1)n - 1 + \mu)p^2 - ((m+1)(n-2) + \mu + 3)p - 2(m+1)}{2} = \frac{1}{p} + 1,$$

it follows from (i) of Lemma 4.6 that

$$\varepsilon \mathcal{E}_{0,\beta_{p+\sigma},m}(u_{0}) + \varepsilon \mathcal{E}_{1,\beta_{p+\sigma},m}(u_{0},u_{1})(t-1) 
\leq Ct^{-\beta_{p+\sigma}+1+\frac{(m+1)n}{p'}} \|u(t,\cdot)\|_{L^{p}(\mathbb{R}^{n})} + C \int_{1}^{t} s^{-\beta_{p}+\frac{(m+1)n}{p'}} \|u(s,\cdot)\|_{L^{p}(\mathbb{R}^{n})} ds 
\leq Ct^{-\beta_{p+\sigma}+1+\frac{(m+1)n}{p'}+(\beta_{p}-1)\frac{1}{p}} (H_{\beta_{p}}(t))^{\frac{1}{p}} 
+ C \int_{1}^{t} s^{-\beta_{p}+\frac{(m+1)n}{p'}+(\beta_{p}-1)\frac{1}{p}} (H_{\beta_{p}}(t))^{\frac{1}{p}} ds 
= Ct^{1+\frac{1}{p}+\beta_{p}-\beta_{p+\sigma}} (H_{\beta_{p}}(t))^{\frac{1}{p}} + C \int_{1}^{t} s^{\frac{1}{p}} (H_{\beta_{p}}(t))^{\frac{1}{p}} ds.$$
(4.28)

Integrating (4.28) on [1, t] and using Hölder's inequality, we have

$$\varepsilon \mathcal{E}_{0,\beta_{p+\sigma},m}(u_0)(t-1) + \varepsilon \mathcal{E}_{1,\beta_{p+\sigma},m}(u_0,u_1)(\frac{t^2}{2} - t + \frac{1}{2})$$

$$\leq C \int_1^t s^{1+\frac{1}{p}+\beta_p-\beta_{p+\sigma}} (H_{\beta_p}(s))^{\frac{1}{p}} ds + C \int_1^t (t-s)s^{\frac{1}{p}} (H_{\beta_p}(s))^{\frac{1}{p}} ds \qquad (4.29)$$

$$\leq C \left(\int_1^t s H_{\beta_p}(s) ds\right)^{\frac{1}{p}} (t^{p'+1})^{\frac{1}{p'}} = C t^{1+\frac{1}{p'}} (I'_{\beta_p}(t))^{\frac{1}{p}},$$

which implies that

$$I'_{\beta_{p}}(t) \geq Ct^{-p(2-\frac{1}{p})} \varepsilon^{p} \Big( \mathcal{E}_{0,\beta_{p+\sigma},m}(u_{0})(t-1) + \mathcal{E}_{1,\beta_{p+\sigma},m}(u_{0},u_{1}) \frac{t^{2}-2t+1}{2} \Big)^{p}$$
(4.30)  
 
$$\geq C\varepsilon^{p}t, \quad \forall t > 1,$$

then

$$I_{\beta_p}(t) \ge C\varepsilon^p t^2, \quad \forall t > 1.$$
(4.31)

By the definition of  $J_{\beta_p}(t)$ ,  $(1+t)^3 J_{\beta_p}'(t) = I_{\beta_p}(t)$ , so

$$J'_{\beta_p}(t) \ge C(1+t)^{-3}\varepsilon^p t^2 \ge C(1+t)^{-1}\varepsilon^p,$$
(4.32)

then

$$J_{\beta_p}(t) \ge C\varepsilon^p \log(1+t). \tag{4.33}$$

By (ii) of Lemma 4.6,

$$\int_{1}^{t} (t-s)H_{\beta_{p}}(s)ds \leq Ct^{-\beta_{p}+1+\frac{(m+1)n}{p'}} \|u(t,\cdot)\|_{L^{p}(\mathbb{R}^{n})} \\
+ C\int_{1}^{t} s^{\frac{(m+1)n}{p'}-\beta_{p}} (\log s)^{\frac{1}{p'}} \|u(s,\cdot)\|_{L^{p}(\mathbb{R}^{n})}ds \\
\leq Ct^{-\beta_{p}+1+\frac{(m+1)n}{p'}+\frac{\beta_{p}-1}{p}} (H_{\beta_{p}}(t))^{\frac{1}{p}} \\
+ C\int_{1}^{t} s^{\frac{(m+1)n}{p'}-\beta_{p}} (\log s)^{\frac{1}{p'}} s^{\frac{\beta_{p}-1}{p}} (H_{\beta_{p}}(s))^{\frac{1}{p}}ds \\
= Ct^{1+\frac{1}{p}} (H_{\beta_{p}}(t))^{\frac{1}{p}} + C\int_{1}^{t} s^{\frac{1}{p}} (\log s)^{\frac{1}{p'}} (H_{\beta_{p}}(s))^{\frac{1}{p}}ds.$$
(4.34)

Integrating (4.34) on [1, t] and using Hölder's inequality give that

$$\frac{1}{2} \int_{1}^{t} (t-s)^{2} H_{\beta_{p}}(s) ds 
\leq C \int_{1}^{t} s^{1+\frac{1}{p}} \left( H_{\beta_{p}}(s) ds \right)^{\frac{1}{p}} + C \int_{1}^{t} (t-s) s^{\frac{1}{p}} (\log s)^{\frac{1}{p'}} \left( H_{\beta_{p}}(s) \right)^{\frac{1}{p}} ds \qquad (4.35) 
\leq C (I'_{\beta_{p}}(t))^{\frac{1}{p}} (1+t)^{\frac{2p-1}{p}} \left( \log(t+1) \right)^{\frac{1}{p'}},$$

then Lemma 4.4 shows that

$$t^{2}J_{\beta_{p}}(t) \leq \frac{1}{2} \int_{1}^{t} (t-s)^{2}H_{\beta_{p}}(s)ds \leq C(I'_{\beta_{p}}(t))^{\frac{1}{p}}(1+t)^{\frac{2p-1}{p}} \left(\log(t+1)\right)^{\frac{1}{p'}},$$

i.e.,

$$\left(\log(t+1)\right)^{1-p} \left(J_{\beta_p}(t)\right)^p \le C I'_{\beta_p}(t)(1+t)^{2p-1} t^{-2p} \le C I'_{\beta_p}(t)(1+t)^{-1}.$$
(4.36)

By the definition of  $J_{\beta_p}(t)$ ,

$$3(1+t)^2 J'_{\beta_p}(t) + (1+t)^3 J''_{\beta_p}(t) = I'_{\beta_p}(t), \qquad (4.37)$$

combining (4.36), we have

$$\left(\log(t+1)\right)^{1-p} \left(J_{\beta_p}(t)\right)^p \le 3C(1+t)J'_{\beta_p}(t) + C(1+t)^2 J''_{\beta_p}(t).$$
(4.38)

Let  $1 + t = e^{\tau}$  and  $J_0(\tau) = J_{\beta_p}(t) = J_{\beta_p}(e^{\tau} - 1)$ , then

$$\begin{cases} J'_0(\tau) = J'_{\beta_p}(t)(1+t), \\ J''_0(\tau) = J''_{\beta_p}(t)(1+t)^2 + J'_{\beta_p}(t)(1+t). \end{cases}$$
(4.39)

It follows from (4.32), (4.38) and (4.39) that

$$\begin{cases} J_0''(\tau) + 2J_0'(\tau) \ge C \left(J_0(\tau)\right)^p \tau^{1-p}, \\ J_0(\tau) \ge C \varepsilon^p \tau, \\ J_0'(\tau) \ge C \varepsilon^p, \end{cases}$$
(4.40)

by (ii) of Lemma 4.7, we deduced that  $J_0(\tau)$  blows up before  $\tau = C\varepsilon^{-p(p-1)}$ , so  $J_{\beta_p}(t)$  blows up before  $t = e^{C\varepsilon^{-p(p-1)}}$ .

### Y.- Q. LI AND F. GUO

### 5. The Proof of Theorem 2.2

5.1. Simplification of the problem. In the case of  $\delta = 1$ , we claim that when  $n \geq 3$ ,  $p_S(n + \frac{\mu}{m+1}, m) > p_F((m+1)n + \frac{\mu-1-\sqrt{\delta}}{2})$  holds for any  $m \geq 0$ ; and for n = 2,  $p_S(n + \frac{\mu}{m+1}, m) > p_F((m+1)n + \frac{\mu-1-\sqrt{\delta}}{2})$  holds if and only if  $\mu \in [0, a(m))$ , where a(m) is the positive root of the quadratic equation

$$\eta^{2} - [8(m+1)^{3} - 8(m+1)^{2} + 2(m+1) - 1]\eta + 8(m+1) - 8(m+1)^{2} - 2 = 0.$$
(5.1)

Indeed,

$$p_{S}(n + \frac{\mu}{m+1}, m) > p_{F}((m+1)n + \frac{\mu - 1 - \sqrt{\delta}}{2})$$

$$\iff \frac{(m+1)(n-2) + \mu + 3 + \sqrt{[(m+1)(n-2) + \mu + 3]^{2} + 8(m+1)[(m+1)n + \mu - 1]}}{2[(m+1)n + \mu - 1]}$$

$$> 1 + \frac{2}{(m+1)n + \frac{\mu}{2} - 1}$$

$$\iff [(m+1)n - 2(m+1) - 1]\mu^{2}$$

$$+ [2(m+1)^{3}n^{2} + (m+1)^{2}n^{2} - 6(m+1)^{2}n - 2(m+1)n + 6(m+1) - 1]\mu$$

$$+ 2(m+1)^{3}n^{3} - 4(m+1)^{3}n^{2} - 2(m+1)^{2}n^{2} + 8(m+1)^{2}n$$

$$- 2(m+1)n - 4(m+1) + 2 > 0.$$
(5.2)

We find that

$$(m+1)n - 2(m+1) - 1 > 0,$$
  
$$2(m+1)^3n^2 + (m+1)^2n^2 - 6(m+1)^2n - 2(m+1)n + 6(m+1) - 1 > 0$$

and

$$2(m+1)^3n^3 - 4(m+1)^3n^2 - 2(m+1)^2n^2 + 8(m+1)^2n - 2(m+1)n - 4(m+1) + 2 > 0$$

hold for  $n \ge 3$  and any  $m \ge 0$ , so (5.2) is obviously valid since  $\mu \ge 0$ . And when n = 2, it is easy to see that for (5.2) to hold, it is sufficient  $\mu \in [0, a(m))$ , where a(m) is the positive root of (5.1). Therefore, in view of Theorem 2.1, we only need to consider the cases of n = 2 with  $\mu \ge a(m)$  and n = 1 with  $p = p_F(m + \frac{\mu}{2})$ .

We introduce the change of variables

$$u(t,x) = t^{-\frac{\mu}{2}}w(t,x),$$
(5.3)

note  $\delta = 1$ , (1.1) can be transformed into

$$\begin{cases} \partial_t^2 w - t^{2m} \Delta w = t^{-\frac{\mu}{2}(p-1)} |w|^p, & t \ge 1, x \in \mathbb{R}^n, \\ w(1,x) = w_0(x), \ \partial_t w(1,x) = w(1,x), & x \in \mathbb{R}^n, \end{cases}$$
(5.4)

where  $w_0(x) = \varepsilon u_0(x)$ ,  $w(1, x) = \frac{\mu}{2} \varepsilon u_0(x) + \varepsilon u_1(x)$ . Now, we prove that the solution to (5.4) blows up in finite time, and consequently, so does (1.1).

5.2. **The Proof of Theorem 2.2.** The proof is based on the following ordinary differential inequalities, which is called Kato's lemma.

**Lemma 5.1.** ([34]) Let p > 1,  $a \ge 1$ , and (p-1)a = q-2. Supposed  $F \in C^2[0,T)$ satisfies, when  $t \ge T_0 > 0$ , (a)  $F(t) \ge K_0(t+R)^a$ , (b)  $F''(t) \ge K_1(t+R)^{-q}F^p(t)$ ,

with some positive constants  $K_0, K_1, T_0$  and R. Fixing  $K_1$ , there exists a positive constant  $c_0$ , independent of R and  $T_0$  such that if  $K_0 \ge c_0$ , then  $T < \infty$ .

Let

$$F(t) = \int_{\mathbb{R}^n} w(t, x) dx,$$
(5.5)

by Hölder's inequality and the finite propagation speed of waves,

$$|F(t)|^{p} \leq \int_{|x| \leq \phi(t) - \phi(1) + M} |w(t, x)|^{p} dx \left( \int_{|x| \leq \phi(t) - \phi(1) + M} dx \right)^{p-1} \\ \leq C(t+M)^{(m+1)n(p-1)} \int_{\mathbb{R}^{n}} |w(t, x)|^{p} dx,$$
(5.6)

then

$$F''(t) = t^{-\frac{\mu}{2}(p-1)} \int_{\mathbb{R}^n} |w(t,x)|^p dx \ge C(t+M)^{-\frac{\mu}{2}(p-1)-(m+1)n(p-1)} |F(t)|^p.$$
(5.7)

By [4], we know that

$$\lambda(t) = \left(\frac{1}{m+1}\right)^{\frac{1}{2(m+1)}} t^{\frac{1}{2}} K_{\frac{1}{2(m+1)}} \left(\frac{t^{m+1}}{m+1}\right)^{\frac{1}{2(m+1)}} \left(\frac{t^{m+1}}{m+1}\right)^{\frac{1}{2(m+1)}} t^{\frac{1}{2}} K_{\frac{1}{2(m+1)}} \left(\frac{t^{m+1}}{m+1}\right)^{\frac{1}{2(m+1)}} t^{\frac{1}{2}} K_{\frac{1}{2(m+1)}} \left(\frac{t^{m+1}}{m+1}\right)^{\frac{1}{2(m+1)}} t^{\frac{1}{2}} K_{\frac{1}{2(m+1)}} \left(\frac{t^{m+1}}{m+1}\right)^{\frac{1}{2(m+1)}} t^{\frac{1}{2}} K_{\frac{1}{2(m+1)}} \left(\frac{t^{m+1}}{m+1}\right)^{\frac{1}{2(m+1)}} t^{\frac{1}{2(m+1)}} t^{\frac$$

is a solution to

$$\lambda''(t) - t^{2m}\lambda(t) = 0, \ t \ge 1,$$

where

$$K_l(z) = \int_0^\infty e^{-z \cosh y} \cosh(ly) dy, \ l \in \mathbb{R}$$

is the modified Bessel function of order l, which is a solution to

$$z^{2}\frac{d^{2}}{dz^{2}}w(z) + z\frac{d}{dz}w(z) - (z^{2} + l^{2})w(z) = 0.$$

Let

$$G(t) = \int_{\mathbb{R}^n} w(t, x)\psi(t, x)dx,$$
(5.8)

where  $\psi(t, x) = \lambda(t)\varphi(x)$  and

$$\varphi(x) = \begin{cases} \int_{S^{n-1}} e^{x \cdot \omega} d\omega, & n \ge 2, \\ e^x + e^{-x}, & n = 1, \end{cases}$$

so,  $\partial_t^2 \psi - t^{2m} \Delta \psi = 0$ . Furthermore, about  $\lambda(t)$ ,  $\varphi(x)$  and G(t) we have

**Lemma 5.2.** ([4]) There exists C > 0 such that for any  $t \in [1, \infty)$ ,

$$C^{-1}t^{-\frac{m}{2}}e^{-\phi(t)} \le \lambda(t) \le Ct^{-\frac{m}{2}}e^{-\phi(t)}.$$
(5.9)

**Lemma 5.3.** ([34]) Let r > 1, there exists a C = C(m, n, r, M) > 0 such that

$$\int_{|x| \le \phi(t) - \phi(1) + M} \varphi^r(x) dx \le C e^{r\phi(t)} \left(1 + \phi(t)\right)^{\frac{(2-r)(n-1)}{2}}.$$
(5.10)

**Lemma 5.4.** Let w be a solution to (5.4) and the assumptions in Theorem 2.2 hold, then there exists a  $T_0 = T_0(m, n) > 1$  such that

$$G(t) \ge Ct^{-m}, \text{ for all } t \ge T_0.$$
(5.11)

**Remark 5.1.** *The proof of Lemma 5.4 is quite similar to Lemma 3.3 in* [4] *and Lemma 3.2 in* [14], *so it is not repeated here.* 

Utilizing Hölder's inequality and Lemmas 5.2-5.4, for  $t \ge T_0$ , we have

$$\begin{aligned} |G(t)|^{p} &\leq \int_{\mathbb{R}^{n}} |w(t,x)|^{p} dx \Big( \int_{\mathbb{R}^{n}} |\psi(t,x)|^{\frac{p}{p-1}} dx \Big)^{p-1} \\ &\leq C \lambda^{p}(t) e^{p\phi(t)} (1+\phi(t))^{\frac{(p-2)(n-1)}{2}} \int_{\mathbb{R}^{n}} |w(t,x)|^{p} dx \\ &\leq C (t+M)^{-\frac{mp}{2}} (1+\phi(t))^{\frac{(p-2)(n-1)}{2}} \int_{\mathbb{R}^{n}} |w(t,x)|^{p} dx, \end{aligned}$$

therefore,

$$F''(t) = t^{-\frac{\mu}{2}(p-1)} \int_{\mathbb{R}^n} |w(t,x)|^p dx$$
  

$$\geq t^{-\frac{\mu}{2}(p-1)} (t+M)^{\frac{mp}{2}} (1+\phi(t))^{-\frac{(p-2)(n-1)}{2}} |G(t)|^p$$
  

$$\geq C(t+M)^{-\frac{\mu}{2}(p-1)-\frac{mp}{2}-\frac{(m+1)(p-2)(n-1)}{2}},$$
(5.12)

integrating twice we arrive at

$$F(t) \ge C(t+M)^{\max\{-\frac{\mu}{2}(p-1)-\frac{mp}{2}-\frac{(m+1)(n-1)(p-2)}{2}+2,1\}}.$$
(5.13)

In the case of n = 2, since  $p = p_F(2(m+1) + \frac{\mu}{2} - 1)$  and  $\mu \ge a(m)$ , we get

$$-\frac{\mu}{2}(p-1) - \frac{mp}{2} - \frac{(m+1)(n-1)(p-2)}{2} + 2 = \frac{3}{2} - \frac{\mu}{2[2(m+1) + \frac{\mu}{2} - 1]} \le 1,$$

so by (5.13),  $F(t) \ge C(t+M)$ , substituting it into (5.7) gives

$$F''(t) \ge C(t+M)^{-\frac{4(m+1)+\mu}{2(m+1)+\frac{\mu}{2}-1}+1+\frac{2}{2(m+1)+\frac{\mu}{2}-1}} = C(t+M)^{-1},$$

integrating twice and combining (5.7), we get

$$\begin{cases} F''(t) \ge C(t+M)^{-\frac{4(m+1)+\mu}{2(m+1)+\frac{\mu}{2}-1}} |F(t)|^p, \ t \ge T_0, \\ F(t) \ge C(t+M)\log(t+M). \end{cases}$$
(5.14)

Let  $q = \frac{4(m+1)+\mu}{2(m+1)+\frac{\mu}{2}-1}$ , a = 1, then (p-1)a = q-2, so Lemma 5.1 shows F(t) blows up in finite time.

In the case of n = 1, since  $p = p_F(m + \frac{\mu}{2})$ ,

$$-\frac{\mu}{2}(p-1) - \frac{mp}{2} - \frac{(m+1)(n-1)(p-2)}{2} + 2 = 2 - \frac{m}{2} - \frac{2m+2\mu}{2m+\mu} \le 1$$
(5.15)

holds, so  $F(t) \ge C(t+M)$  by (5.13), substituting it into (5.7), we have  $F''(t) \ge C(t+M)^{-2-\frac{4}{2m+\mu}} = C(t+M)^{-1}$ , therefore,

$$\begin{cases} F''(t) \ge C(t+M)^{-2-\frac{4}{2m+\mu}} |F(t)|^p, \ t \ge T_0, \\ F(t) \ge C(t+M) \log(t+M), \end{cases}$$
(5.16)

let  $q = 2 + \frac{4}{2m+\mu}$ , a = 1, then (p-1)a = q-2, so F(t) blows up in finite time.

## Acknowledgments

Y-Q. Li is supported by the Postgraduate Research & Practice Innovation Program of Jiangsu Province (KYCX24\_1786). F. Guo is supported by the Priority Academic Program Development of Jiangsu Higher Education Institutions and the NSF of Jiangsu Province (BK20221320).

### REFERENCES

- [1] J. BARROS-NETO, I.M. GELFAND, Fundamental solutions for the Tricomi operator, *Duke Math. J.*, **98** (1999), 465-483.
- [2] J. BARROS-NETO, I.M. GELFAND, Fundamental solutions for the Tricomi operator. II, Duke Math. J., 111 (2002), 561-584.
- [3] J. BARROS-NETO, I.M. GELFAND, Fundamental solutions for the Tricomi operator. III, *Duke Math. J.*, **128** (2005), 119-140.
- [4] M.F. BEN HASSEN, M. HAMOUDA, M.A. HAMZA, H.K. TEKA, Nonexistence result for the generalized Tricomi equation with the scale-invariant damping, mass term and time derivative nonlinearity, *Asymptot. Anal.*, **128** (2022), 495-515.
- [5] M. D'ABBICCO, Small data solutions for the Euler-Poisson-Darboux equation with a power nonlinearity, J. Differential Equations, 286 (2021), 531-556.
- [6] M. D'ABBICCO, The semilinear Euler-Poisson-Darboux equation: a case of wave with critical dissipation, arXiv:2008.08703, (2020).
- [7] M. D'ABBICCO, S. LUCENTE, NLWE with a special scale-invariant damping in odd space dimension, *Discrete Contin. Dyn. Syst.*, AIMS Proceedings (2015), 312-319.
- [8] M. D'ABBICCO, S. LUCENTE, M. REISSIG, A shift in the Strauss exponent for semilinear wave equations with a not effective damping, J. Differential Equations, 259 (2015), 5040-5073.
- [9] F. FRANKL, On the problems of Chaplygin for mixed sub- and supersonic flows, *Bull. Acad. Sci. URSS. Sér. Math.*, 9 (1945), 121-143.
- [10] A. FRIEDMAN, Partial Differential Equations, Krieger, New York (1976), 1215-1240.
- [11] H. FUJITA, On the blowing up of solutions of the Cauchy problem for  $u_t = \Delta u + u^{1+\alpha}$ , J. Fac. Sci. Univ. Tokyo Sect. I, 13 (1966), 109-124.
- [12] V. GEORGIEV, H. LINDBLAD, C.D.SOGGE, Weighted Strichartz estimates and global existence for semilinear wave equations, *Amer. J. Math.*, **119** (1997), 1291-1319.
- [13] R.T. GLASSEY, Existence in the large for  $\Box u = F(u)$  in two space dimensions, *Math. Z.*, **178** (1981), 233-261.
- [14] M. HAMOUDA, M.A. HAMZA, Blow-up and lifespan estimate for the generalized Tricomi equation with mixed nonlinearities, *Adv. Pure Appl. Math.*, **12** (2021), 54-70.
- [15] K. HAYAKAWA, On nonexistence of global solutions of some semilinear parabolic differential equations, Proc. Jpn. Acad., 49 (1973), 503-505.
- [16] D. HE, I. WITT, H. YIN, On semilinear Tricomi equations with critical exponents or in two space dimensions, J. Differential Equations, 263 (2017), 8102-8137.
- [17] D. HE, I. WITT, H. YIN, On the global solution problem for semilinear generalized Tricomi equations, I, *Calc. Var. Partial Differential Equations*, **56** (2017), 1-24.

#### Y.- Q. LI AND F. GUO

- [18] D. HE, I. WITT, H. YIN, On the global solution problem for semilinear generalized Tricomi equations, II, *Pacific J. Math.*, **314** (2021), 1-24.
- [19] F. JOHN, Blow-up of solutions of nonlinear wave equations in three space dimensions, *Proc. Nat. Acad. Sci. U.S.A.*, 28 (1979), 1559-1560.
- [20] J. LIN, Z. TU, Lifespan of semilinear generalized Tricomi equation with Strauss type exponent, arXiv:1903.11351, (2019).
- [21] W. N. NASCIMENTO, A. PALMIERI, M. REISSIG, Semi-linear wave models with power nonlinearity and scale-invariant time-dependent mass and dissipation, *Math. Nachr.*, 290 (2017), 1779-1805.
- [22] F.W.J. OLVER, D.W. LOZIER, R.F. BOISVERT, C.W. CLARK, NIST Handbook of Mathematical Functions, *U.S. Depart-ment of Commerce*, Washington, 2010.
- [23] A. PALMIERI, Global existence of solutions for semi-linear wave equation with scaleinvariant damping and mass in exponentially weighted spaces, J. Math. Anal. Appl., 461 (2018), 1215-1240.
- [24] A. PALMIERI, On the the critical exponent for the semilinear Euler-Poisson-Darboux-Tricomi equation with power nonlinearity, arXiv:2105.09879, (2021).
- [25] A. PALMIERI, M. REISSIG, A competition between Fujita and Strauss type exponents for blow-up of semi-linear wave equations with scale-invariant damping and mass, J. Differential Equations, 266 (2019), 1176-1220.
- [26] A. PALMIERI, M. REISSIG, Semi-linear wave models with power non-linearity and scaleinvariant time-dependent mass and dissipation, II, *Math. Nachr.*, 291 (2018), 1859-1892.
- [27] A. PALMIERI, Z. TU, Lifespan of semilinear wave equation with scale invariant dissipation and mass and sub-Strauss power nonlinearity, *J. Math. Anal. Appl.*, **470** (2019), 447-469.
- [28] E.L. SHISHKINA, S.M. SITNIK, General form of the Euler-Poisson-Darboux equation and application of transmutation method, *Electron. J. Differential Equations*, (2017), Paper No. 177, 20. pp.
- [29] W.A. STRAUSS, Nonlinear scattering theory at low energy, J. Funct. Anal., 41 (1981), 110-133.
- [30] Y. SUN, Sharp lifespan estimates for subcritical generalized semilinear Tricomi equations, Math. Methods Appl. Sci., 44 (2021), 10239-10251.
- [31] D. TATARU, Strichartz estimates in the hyperbolic space and global existence for the semilinear wave equation, *Trans. Amer. Math. Soc.*, **353** (2001), 795-807.
- [32] J. WIRTH, Solution representations for a wave equation with weak dissipation, *Math. Methods Appl. Sci.*, **27** (2004), 101-124.
- [33] K. YAGDJIAN, A note on the fundamental solution for the Tricomi-type equation in the hyperbolic domain, *J. Differential Equations*, **206** (2004), 227-252.
- [34] B.T. YORDANOV, Q.S. ZHANG, Finite time blow up for critical wave equations in high dimensions, J. Funct. Anal., 231 (2006), 361-374.
- [35] Y. ZHOU, Blow up of solutions to semilinear wave equations with critical exponent in high dimensions, *Chin. Ann. Math.*, **28 B** (2007), 205-212.
- [36] Y. ZHOU, W. HAN, Life-span of solutions to critical semilinear wave equations, *Comm. Partial Differential Equations*, **39** (2014), 439-451.

#### Yuequn Li

SCHOOL OF MATHEMATICAL SCIENCES, NANJING NORMAL UNIVERSITY NANJING 210023, CHINA *Email address*: yqli1214@163.com

#### Fei Guo

SCHOOL OF MATHEMATICAL SCIENCES AND KEY LABORATORY FOR NSLSCS, MINISTRY OF EDUCA-TION, NANJING NORMAL UNIVERSITY, NANJING 210023, CHINA

Email address: guof@njnu.edu.cn