

SEPARATED DETERMINANTAL POINT PROCESSES AND GENERALIZED FOCK SPACES

GIUSEPPE LAMBERTI AND XAVIER MASSANEDA

ABSTRACT. We study conditions so that the determinantal point process Λ_ϕ associated to a generalized Fock space defined by a doubling subharmonic weight ϕ is almost surely a separated sequence in \mathbb{C} . Under a natural assumption on ϕ , we provide a characterization of such processes. Additionally, we emphasize the role of intrinsic repulsion in determinantal processes by comparing Λ_ϕ with the Poisson process of the same first intensity. As an application, we show that the determinantal process Λ_α associated to the canonical weight $\phi_\alpha(z) = |z|^\alpha$, $\alpha > 0$, is almost surely separated if and only if $\alpha < 4/3$. In contrast, the Poisson process Λ_α^P having the same first intensity as Λ_α is almost surely separated if and only if $\alpha < 1$.

1. INTRODUCTION

In this paper we provide conditions so that a particular family of determinantal point processes in the complex plane are almost surely separated. Recall that a sequence $\Lambda = \{\lambda_k\}_{k \geq 1} \subset \mathbb{C}$ is *separated* if

$$\inf_{j \neq k} |\lambda_k - \lambda_j| > 0.$$

Separation plays an important role in many numerical and function theoretic problems, such as the description of interpolating and sampling sequences for various spaces.

Intuitively, a (simple) point process is a random configuration of points, but it is more convenient to think of it as a random measure of the form

$$\tau_\Lambda = \sum_{\lambda \in \Lambda} \delta_\lambda,$$

where Λ is a finite or countable subset of \mathbb{C} . The distribution (or law) of the point process is then determined by the random variables

$$N(B) = \#(\Lambda \cap B) = \int_B d\tau_\Lambda, \quad B \subset \mathbb{C} \text{ compact.}$$

For a background on random measures and point processes we refer the reader to [9].

Date: February 5, 2025.

Key words and phrases. Random point process, separated sequences.

The first author partially supported by the project UBGRS 2.0 (ANR-20-SFRI-0001). Second author partially supported by the Generalitat de Catalunya (grant 2021 SGR 00087) and the Spanish Ministerio de Ciencia e Innovación (project PID2021-123405NB-I00).

We start by recalling the generalized Fock spaces \mathcal{F}_ϕ defined by doubling weights ϕ . Let ϕ be a subharmonic function in \mathbb{C} with doubling Laplacian $\nu = \Delta\phi$, i.e., for which there exists $C > 0$ such that

$$\nu(D(z, 2r)) \leq C \nu(D(z, r)), \quad z \in \mathbb{C}, \quad r > 0.$$

Then, denote by \mathcal{F}_ϕ the generalized Fock space of entire functions f such that

$$\int_{\mathbb{C}} |f(z)|^2 e^{-2\phi(z)} \Delta\phi(z) < +\infty.$$

This is a reproducing kernel Hilbert space whose associated kernel is the Bergman kernel K_ϕ . We refer the reader to the next section for precise definitions.

The focus of this paper is on the determinantal point processes Λ_ϕ associated to the Bergman kernel K_ϕ . We proceed to explain this in further detail. According to a theorem of Macchi and Soshnikov ([20, Theorem 3], see also [12, Lemma 4.5.1]) whenever an Hermitian kernel $K(z, \zeta)$ defines a self-adjoint operator \mathcal{K} on $L^2(\mathbb{C}, \mu)$, which is locally trace class with all eigenvalues in $[0, 1]$, there exists a unique determinantal point process Λ associated with K and μ . This means that for any collections of disjoint sets $B_1, \dots, B_n \subset \mathbb{C}$ one has

$$\mathbb{E} \left[\prod_{k=1}^n \#(B_k \cap \Lambda) \right] = \int_{B_1} \cdots \int_{B_n} \det(K(z_i, z_j))_{1 \leq i, j \leq n} d\mu(z_1) \cdots d\mu(z_n).$$

The integrand is usually called the n^{th} -correlation function of Λ . For the particular case corresponding to $n = 1$, this is

$$\mathbb{E}[N(B)] = \int_B K(z, z) d\mu(z)$$

and the measure $K(z, z) d\mu(z)$ is called the *first intensity* (or average distribution) of the process.

The Bergman kernel K_ϕ defines a projection operator from $L^2(\mathbb{C}, \mu_\phi)$ onto \mathcal{F}_ϕ so it satisfies the hypothesis of the aforementioned theorem by Macchi and Soshnikov (the definition of μ_ϕ is provided in the following section). Consequently, the determinantal point process Λ_ϕ associated to K_ϕ and μ_ϕ is well-defined.

These processes can be seen as generalizations of the process studied in [5], where the authors investigate the Palm measures associated to \mathcal{F}_ϕ , under the assumption that $\Delta\phi$ is uniformly bounded both above and below.

We mention that determinantal processes were first considered by Macchi [14] as a model for fermions, and since then have appeared naturally in a surprising variety of problems (see e.g. [12, Chapter 4]).

Our main result, Theorem 2.1, characterizes the processes Λ_ϕ which are almost surely separated, under a natural restriction on ϕ . As a by-product of its proof, we notice that when Λ_ϕ is a.s. separated, then its upper density is 0. This automatically provides a characterization of the processes Λ_ϕ which are almost surely interpolating for the classical Fock spaces $\mathcal{F}_{\beta|z|^2}$, $\beta > 0$ (see Corollary 2.4). Of course, the analogous result holds as well for the L^p versions of these classical Fock spaces.

A distinctive feature of determinantal processes is their built-in repulsion (i.e. a negative correlation of nearby points), which can be quantified by the decay of the kernel. To assess the role played by this repulsion, we compare Λ_ϕ with the Poisson process Λ_ϕ^P of “totally independent” points having the same first intensity (average distribution) as Λ_ϕ . The existence of such a Poisson process is established, for instance, in [13, Section 3.2]. An adaptation of [11, Theorem 1.2] shows that, by the big fluctuations of the Poisson process, much stricter conditions are required for Λ_ϕ^P to be almost surely separated (Theorem 2.5).

The plan of the paper is as follows. In Section 2 we gather the main properties of the Fock spaces \mathcal{F}_ϕ and their associated processes Λ_ϕ . We also state the main results. The proof of the main result (Theorem 2.1) and its consequences is given in Section 3. The key step in this proof is the estimate of the probability that two or more points fall in a cell of fixed size (Proposition 3.6). Finally, Section 4 is devoted to briefly recall the proof of Theorem 2.5.

A final word about notation: the expression $A \preceq B$ means that there exists a constant $C > 0$, independent of whatever arguments are involved, such that $A \leq CB$. If both $A \preceq B$ and $B \preceq A$ then we write $A \simeq B$.

Acknowledgements: We would like to thank Jordi Marzo and Joaquim Ortega-Cerdà for their help on determinantal processes and for the many valuable discussions.

2. DETERMINANTAL PROCESSES INDUCED BY GENERALIZED FOCK SPACES. RESULTS.

Let ϕ be a subharmonic function with doubling Laplacian and let \mathcal{F}_ϕ be the associated Fock space. In order to express the norm in \mathcal{F}_ϕ it is more convenient to work with a suitable regularization of $\nu = \Delta\phi$.

Following the ideas of M. Christ [7], for each $z \in \mathbb{C}$ denote by $\rho(z)$ (or $\rho_\phi(z)$ if we want to stress the dependency on ϕ) the positive radius such that

$$\nu(D(z, \rho(z))) = 1.$$

It is not difficult to see (e.g. [15, p.869]) that

$$|\rho(z) - \rho(\zeta)| \leq |z - \zeta|, \quad z, \zeta \in \mathbb{C}.$$

In particular, $\rho(z)$ is continuous. The function ρ^{-2} can then be seen as a regularized version of $\Delta\phi$, as described in [7]. Then, define

$$\mathcal{F}_\phi = \left\{ f \in H(\mathbb{C}) : \|f\|_\phi^2 := c_\phi \int_{\mathbb{C}} |f(z)|^2 e^{-2\phi(z)} \frac{dm(z)}{\rho^2(z)} < +\infty \right\}.$$

The constant c_ϕ is chosen so that $\|1\|_\phi = 1$; hence

$$(1) \quad d\mu_\phi(z) := c_\phi e^{-2\phi(z)} \frac{dm(z)}{\rho^2(z)}$$

is a probability measure in \mathbb{C} .

Equipped with the scalar product

$$\langle f, g \rangle_\phi = \int_{\mathbb{C}} f(z) \overline{g(z)} d\mu_\phi(z), \quad f, g \in \mathcal{F}_\phi,$$

the space \mathcal{F}_ϕ is a reproducing kernel Hilbert space whose kernel K_ϕ is also referred to as the Bergman kernel of \mathcal{F}_ϕ . Specifically, this means that K_ϕ is holomorphic in z , anti-holomorphic in ζ , and for any $f \in \mathcal{F}_\phi$,

$$f(z) = \langle f, \overline{K(z, \cdot)} \rangle_\phi.$$

Let now Λ_ϕ be the determinantal point process associated to the Hermitian kernel $K_\phi(z, \zeta)$ and the reference measure μ_ϕ . Its first intensity is, by definition,

$$K_\phi(z, z) d\mu_\phi(z).$$

According to Lemma 21 in [15]; there exists $C > 0$ such that

$$(2) \quad C^{-1} e^{2\phi(z)} \leq K_\phi(z, z) \leq C e^{2\phi(z)} \quad z \in \mathbb{C}.$$

Therefore, by (1), the first intensity of Λ_ϕ is comparable to the regularization of the Laplacian of ϕ :

$$(3) \quad K_\phi(z, z) d\mu_\phi(z) \simeq \frac{dm(z)}{\rho^2(z)}.$$

In particular, given any Borel set $B \subset \mathbb{C}$, the expected value of the counting variable $N_\phi(B) = \#(\Lambda_\phi \cap B)$ is

$$(4) \quad \mathbb{E}[N_\phi(B)] = \int_B K_\phi(z, z) d\mu_\phi(z) \simeq \int_B \frac{dm(z)}{\rho^2(z)}.$$

Since we are looking for conditions so that Λ_ϕ is a.s. separated, we restrict ourselves to the cases where the order of $\mathbb{E}[N_\phi(B)]$ is asymptotically smaller than the area of B as we move away from the origin. This is achieved by imposing that

$$(5) \quad \liminf_{r \rightarrow +\infty} \min_{|z|=r} \rho(z) = +\infty.$$

We are ready to state our main result.

Theorem 2.1. *Let ϕ be a doubling subharmonic function satisfying (5), and let Λ_ϕ be its associated determinantal point process. Then*

$$\mathbb{P}(\Lambda_\phi \text{ is separated}) = \begin{cases} 1 & \text{if } \int_{\mathbb{C}} \frac{dm(z)}{\rho^6(z)} < +\infty, \\ 0 & \text{if } \int_{\mathbb{C}} \frac{dm(z)}{\rho^6(z)} = +\infty. \end{cases}$$

Remark 2.2. There is no restriction in assuming that ϕ is regular: according to [15, Theorem 14], there always exist $\psi \in \mathcal{C}^\infty(\mathbb{C})$ subharmonic and $C > 0$ such that $\Delta\psi$ is a doubling measure, $|\psi - \phi| \leq C$ and

$$\Delta\psi \simeq \frac{1}{\rho_\psi^2} \simeq \frac{1}{\rho_\phi^2},$$

so that $\mathcal{F}_\phi = \mathcal{F}_\psi$ and $\|f\|_\phi \simeq \|f\|_\psi$ for all $f \in \mathcal{F}_\phi$.

When ϕ is regular, the first intensity of Λ_ϕ is thus comparable to $\Delta\phi \, dm$, and the integral above can be rewritten as

$$\int_{\mathbb{C}} (\Delta\phi(z))^3 \, dm(z).$$

We keep the form given in the statement, because that is how it appears in the proof.

Canonical examples of the weights considered in Theorem 2.1 are $\phi_\alpha(z) = |z|^\alpha$, $\alpha \in (0, 2)$. For $\alpha = 2$, the distribution of $\Lambda_\alpha := \Lambda_{\phi_\alpha}$ is invariant by translations (see e.g. [17]). In particular, for any $\varepsilon > 0$ the probabilities $\mathbb{P}[\#(\Lambda_2 \cap D(z, \varepsilon))] > 0$ are strictly positive and independent of $z \in \mathbb{C}$. Then the arguments of Bufetov, Qiu and Shamov for the hyperbolic disk [6, Lemma 3] apply similarly to \mathbb{C} to conclude that Λ_α is a.s. not separated. This explains the restriction $\alpha \in (0, 2)$, which is precisely (5), since $\rho_{\phi_\alpha}(z) \simeq |z|^{1-\frac{\alpha}{2}}$.

Corollary 2.3. *Let Λ_α be the determinantal point process associated to the weight $\phi_\alpha(z) = |z|^\alpha$, $\alpha \in (0, 2)$. Then,*

$$\mathbb{P}(\Lambda_\alpha \text{ is separated}) = \begin{cases} 1 & \text{if } \alpha < 4/3 \\ 0 & \text{if } \alpha \geq 4/3. \end{cases}$$

The proof of Theorem 2.1 shows that, in case Λ_ϕ is separated, its upper density

$$D_+(\Lambda_\phi) := \limsup_{r \rightarrow \infty} \sup_{z \in \mathbb{C}} \frac{\#(\Lambda_\phi \cap D(z, r))}{\pi r^2}$$

is zero. By a well-known theorem by K. Seip and R. Wallstén [18, 19], a sequence Λ in the plane is interpolating for the classical Fock space $\mathcal{F}_{\beta|z|^2}$, $\beta > 0$, if and only if it is separated and $D_+(\Lambda) < \beta$ (see e.g. [22, Chapter 3]).

Corollary 2.4. *Let ϕ be a doubling subharmonic weight for which (5) holds and let Λ_ϕ be its associated determinantal process. Then, for any $\beta > 0$,*

$$\mathbb{P}(\Lambda_\phi \text{ is } \mathcal{F}_{\beta|z|^2}\text{-interpolating}) = \begin{cases} 1 & \text{if } \int_{\mathbb{C}} \frac{dm(z)}{\rho^\beta(z)} < +\infty \\ 0 & \text{if } \int_{\mathbb{C}} \frac{dm(z)}{\rho^\beta(z)} = +\infty. \end{cases}$$

We finish this section by comparing the results above with the analogues for the Poisson process with the same first intensity. Consider

$$d\sigma_\phi(z) = \frac{dm(z)}{\rho^2(z)}$$

and denote by Λ_ϕ^P the Poisson process with underlying measure σ_ϕ . Observe that, by (3), Λ_ϕ^P can be seen as the Poisson point process having the same average distribution as the determinantal process Λ_ϕ .

The Poisson process Λ_ϕ^P is characterized by two properties:

- (i) for any Borel set $A \subset \mathbb{C}$ the counting function $N(A) = \#(A \cap \Lambda_\phi^P)$ has Poisson distribution of parameter $\sigma_\phi(A)$,
- (ii) for two disjoint sets $A, B \subset \mathbb{C}$ the variables $N(A), N(B)$ are independent.

A straightforward adaptation of Theorem 1.2 in [11] provides the following characterization.

Theorem 2.5. *Let ϕ be a doubling subharmonic weight satisfying (5) and let Λ_ϕ^P be the Poisson process with underlying measure σ_ϕ . Then*

$$\mathbb{P}(\Lambda_\phi^P \text{ is separated}) = \begin{cases} 1 & \text{if } \int_{\mathbb{C}} \frac{dm(z)}{\rho^4(z)} < +\infty \\ 0 & \text{if } \int_{\mathbb{C}} \frac{dm(z)}{\rho^4(z)} = +\infty. \end{cases}$$

For the cases $\phi(z) = |z|^\alpha$, $\alpha \in (0, 2)$, this means that Λ_ϕ^P is almost surely separated if and only if $\alpha < 1$.

As in the determinantal case, we have an immediate consequence for (classical) Fock interpolation.

Corollary 2.6. *Let ϕ be a doubling subharmonic weight for which (5) holds, and let Λ_ϕ^P be the Poisson process with underlying measure σ_ϕ . Then, for any $\beta > 0$,*

$$\mathbb{P}(\Lambda_\phi \text{ is } \mathcal{F}_{\beta|z|^2}\text{-interpolating}) = \begin{cases} 1 & \text{if } \int_{\mathbb{C}} \frac{dm(z)}{\rho^4(z)} < +\infty \\ 0 & \text{if } \int_{\mathbb{C}} \frac{dm(z)}{\rho^4(z)} = +\infty. \end{cases}$$

3. DETERMINANTAL PROCESS: PROOF OF THEOREM 2.1 AND COROLLARY 2.4

3.1. Preliminaries. We start this section with some technical, well-known properties of \mathcal{F}_ϕ .

A first elementary property to be used throughout the paper is the following (see [15, Corollary 3]): for any $r > 0$ there exists $c_r > 0$ such that

$$(6) \quad c_r^{-1} \rho(z) \leq \rho(\zeta) \leq c_r \rho(z), \quad \zeta \in D(z, r\rho(z)).$$

There are at least two related distances in \mathbb{C} induced by \mathcal{F}_ϕ . The first is given by the reproducing kernel K_ϕ (see [1, p.128]):

$$(7) \quad d_K(z, \zeta) = \sqrt{1 - \frac{|K_\phi(z, \zeta)|^2}{K_\phi(z, z)K_\phi(\zeta, \zeta)}}, \quad z, \zeta \in \mathbb{C},$$

while the second is the distance d_B associated to the Bergman metric

$$(8) \quad \Delta \log K_\phi(z, z) dz \otimes d\bar{z}.$$

According to Proposition 8 (or 9) in [2], the Bergman distance is the so-called *inner distance* of d_K , that is,

$$d_B(z, \zeta) = \inf \{ \ell_{d_k}(\gamma) : \gamma \text{ curve joining } z \text{ to } \zeta \},$$

where, if $\gamma : [0, 1] \rightarrow \mathbb{C}$,

$$\ell_{d_k}(\gamma) = \sup \left\{ \sum_{i=1}^n d_K(\gamma(t_{i-1}), \gamma(t_i)) : 0 = t_0 < \dots < t_n = 1 \right\}.$$

In particular,

$$(9) \quad d_K(z, \zeta) \leq d_B(z, \zeta), \quad z, \zeta \in \mathbb{C}.$$

Next, we recall some well-known estimates for the reproducing kernel (see [4, Proposition 14] and the references therein).

Proposition 3.1. *There exist positive constants C , ε and η , depending only on ϕ , such that,*

- (a) $|K_\phi(z, \zeta)| \leq C e^{\phi(z)+\phi(\zeta)} e^{-d_B^\varepsilon(z, \zeta)}$ for all $z, \zeta \in \mathbb{C}$,
- (b) $|K_\phi(z, \zeta)| \geq C e^{\phi(z)+\phi(\zeta)}$ for all $\zeta \in D(z, \eta\rho(z))$,
- (c) $C^{-1}/\rho^2(z) \leq \Delta \log K_\phi(z, z) \leq C/\rho^2(z)$.

In [4] the estimate (a) is given in terms of the distance d_ϕ associated to the metric $\rho^{-2}(z)dz \otimes d\bar{z}$ which, by (c), is equivalent to d_B .

Another property to be used in the proof of Theorem 2.1 is the following.

Lemma 3.2. ([4, Lemma 12]; [16, Lemma 2.7]) *For any $\varepsilon > 0$ and $k \geq 0$*

$$\sup_{z \in \mathbb{C}} \int_{\mathbb{C}} \left(\frac{|\zeta - z|}{\rho(z)} \right)^k e^{-d_B^\varepsilon(z, \zeta)} \frac{dm(\zeta)}{\rho^2(\zeta)} < +\infty.$$

It is not difficult to see ([15, Lemma 4]) that for every $r > 0$ there exists $C > 0$ such that

$$(10) \quad C^{-1} \frac{|z - \zeta|}{\rho(z)} \leq d_B(z, \zeta) \leq C \frac{|z - \zeta|}{\rho(z)}, \quad \zeta \in D(z, r\rho(z)).$$

The distance d_K satisfies similar estimates, which are important in the proof of Theorem 2.1.

Lemma 3.3. (a) *For every $r > 0$ there exists $C > 0$ depending only on ϕ such that*

$$d_K(z, \zeta) \leq C \frac{|z - \zeta|}{\rho(z)}, \quad \zeta \in D(z, r\rho(z)).$$

(b) *There exist $\eta > 0$ and a positive constant C such that*

$$d_K(z, \zeta) \geq C \frac{|z - \zeta|}{\rho(z)}, \quad \zeta \in D(z, \eta\rho(z)).$$

Proof. (a) Apply (9) and the upper estimate in (10).

(b) By [2, p. 30], the distance $d_K(z, \zeta)$ is the solution to the extremal problem

$$d_K(z, \zeta) = \frac{\sup\{|F(\zeta)| : F \in \mathcal{F}_\phi, \|F\|_\phi \leq 1, F(z) = 0\}}{\sqrt{K_\phi(\zeta, \zeta)}}.$$

Take, fixed $z \in \mathbb{C}$, the function

$$F_z(\zeta) = \frac{\zeta - z}{\rho(z)} \frac{K_\phi(\zeta, z)}{\sqrt{K_\phi(z, z)}}.$$

By Proposition 3.1(a) and (2),

$$\|F_z\|_\phi^2 \simeq \int_{\mathbb{C}} \frac{|\zeta - z|^2}{\rho^2(z)} \frac{|K_\phi(\zeta, z)|^2}{K_\phi(z, z)} e^{-2\phi(\zeta)} \frac{dm(\zeta)}{\rho^2(\zeta)} \lesssim \int_{\mathbb{C}} \frac{|\zeta - z|^2}{\rho^2(z)} e^{-2d_B^2(z, \zeta)} \frac{dm(\zeta)}{\rho^2(\zeta)},$$

hence, by Lemma 3.2, $\|F_z\|_\phi \lesssim 1$.

Thus, by (2) once more,

$$\begin{aligned} d_K(z, \zeta) &\gtrsim \frac{|F_z(\zeta)|}{\sqrt{K_\phi(\zeta, \zeta)}} = \frac{|\zeta - z|}{\rho(z)} \frac{|K_\phi(\zeta, z)|}{\sqrt{K_\phi(z, z)} \sqrt{K_\phi(\zeta, \zeta)}} \\ &\simeq \frac{|\zeta - z|}{\rho(z)} |K_\phi(\zeta, z)| e^{-\phi(z)} e^{-\phi(\zeta)}. \end{aligned}$$

Then we conclude with Proposition 3.1 (b). \square

3.2. Main Proposition. We begin by considering a standard partition of the plane and studying the probability that two or more points fall in the same cell.

Let

$$T_{n,k} := \left\{ z = re^{i\theta} \in \mathbb{C} : n-1 \leq r < n, \frac{\theta}{2\pi} \in \left[\frac{k-1}{n}, \frac{k}{n} \right) \right\}, \quad n \geq 1, k = 1, \dots, n$$

and let $z_{n,k} = (n-1/2) e^{i2\pi \frac{k-1/2}{n}}$ denote its ‘‘center’’.

Consider the corresponding counting random variables

$$X_{n,k} = \#(\Lambda_\phi \cap T_{n,k}).$$

Observe that, by (4) and (6),

$$(11) \quad \mathbb{E}[X_{n,k}] \simeq \int_{T_{n,k}} \frac{dm(z)}{\rho^2(z)} \simeq \frac{1}{\rho^2(z_{n,k})}.$$

A specific feature of any determinantal process Λ is that the counting random variable $N(B) = \#(\Lambda \cap B)$ can be expressed as a sum of independent Bernoulli variables ξ_j . Moreover, the

parameters λ_j of the variables ξ_j are precisely the eigenvalues of the restriction operator on B (see e.g. [12, Theorem 4.5.3]). Hence, taking $\Lambda = \Lambda_\phi$ and $B = T_{n,k}$, we have

$$X_{n,k} = \sum_{j=1}^{\infty} \xi_j,$$

where $\xi_j \sim \text{Bernoulli}(\lambda_j)$ and λ_j is the j^{th} eigenvalue (arranged in decreasing order) of the restriction operator

$$(12) \quad Tf(z) = \int_{T_{n,k}} f(\zeta) K_\phi(z, \zeta) d\mu_\phi(\zeta), \quad f \in \mathcal{F}_\phi.$$

The key result in the proof of Theorem 2.1 is the estimate of the probability that two or more points fall in the same cell $T_{n,k}$, which, as we shall see soon, is comparable to $\rho^{-6}(z_{n,k})$. We begin with the following general representation of this probability.

Lemma 3.4. *Let $T_{n,k}$ and $X_{n,k}$ as above. Then,*

$$\mathbb{P}(X_{n,k} \geq 2) = \int_{T_{n,k}} \int_{T_{n,k}} [K_\phi(z, z)K_\phi(\zeta, \zeta) - |K_\phi(z, \zeta)|^2] d\mu_\phi(z) d\mu_\phi(\zeta) + O\left((\mathbb{E}[X_{n,k}])^3\right).$$

Remark 3.5. (a) As we shall see in the proof, this identity is valid in great generality. Given a determinantal point process Λ with correlation kernel $K(z, \zeta)$ and reference measure μ , and given $B \subset \mathbb{C}$ and its counting variable $N(B) = \#(\Lambda \cap B)$, the analogue for $\mathbb{P}(N(B) \geq 2)$ also holds. Of course, this is only of interest when $(\mathbb{E}[N(B)])^3$ is much smaller than the double integral.

(b) The identity in the statement can also be rewritten in terms of the distance d_K defined in (7) and the first intensity of the process:

$$\mathbb{P}(X_{n,k} \geq 2) = \int_{T_{n,k}} \int_{T_{n,k}} d_K^2(z, \zeta) K_\phi(z, z) d\mu_\phi(z) K_\phi(\zeta, \zeta) d\mu_\phi(\zeta) + O\left((\mathbb{E}[X_{n,k}])^3\right).$$

(c) The double integral above is exactly $\mathbb{E}[X_{n,k}(X_{n,k} - 1)]$ (see [12, (1.2.4)]). Thus, the result can also be rephrased as

$$\mathbb{P}(X_{n,k} \geq 2) = \mathbb{E}[X_{n,k}^2] - \mathbb{E}[X_{n,k}] + O\left((\mathbb{E}[X_{n,k}])^3\right).$$

Proof. By definition,

$$\begin{aligned} \mathbb{P}(X_{n,k} \geq 2) &= 1 - \mathbb{P}(X_{n,k} = 0) - \mathbb{P}(X_{n,k} = 1) = 1 - \prod_{j=1}^{\infty} (1 - \lambda_j) - \sum_{l=1}^{\infty} \lambda_l \prod_{j \neq l} (1 - \lambda_j) \\ &= 1 - \exp\left(-\sum_{j=1}^{\infty} \log\left(\frac{1}{1 - \lambda_j}\right)\right) - \exp\left(-\sum_{j=1}^{\infty} \log\left(\frac{1}{1 - \lambda_j}\right)\right) \left(\sum_{l=1}^{\infty} \frac{\lambda_l}{1 - \lambda_l}\right). \end{aligned}$$

Observe that all λ_j are very close to 0, because of condition (5), and the fact that, as seen in (11),

$$\mathbb{E}[X_{n,k}] = \text{trace}(T) = \sum_{j=1}^{\infty} \lambda_j \simeq \frac{1}{\rho^2(z_{n,k})}.$$

Plugging the power series

$$e^{-x} = \sum_{n=0}^{\infty} (-1)^{n+1} \frac{x^n}{n!} = 1 - x + \frac{x^2}{2} + \dots$$

$$\log\left(\frac{1}{1-x}\right) = \sum_{n=1}^{\infty} \frac{x^n}{n} = x + \frac{x^2}{2} + \dots$$

$$\frac{x}{1-x} = \sum_{n=1}^{\infty} x^n = x + x^2 + \dots$$

and keeping the terms up to order 2 in the expression above, we get

$$\begin{aligned} \mathbb{P}(X_{n,k} \geq 2) &= \sum_{j=1}^{\infty} \log\left(\frac{1}{1-\lambda_j}\right) - \frac{1}{2} \left(\sum_{j=1}^{\infty} \log\left(\frac{1}{1-\lambda_j}\right) \right)^2 - \\ &\quad - \left(\sum_{l=1}^{\infty} \frac{\lambda_l}{1-\lambda_l} \right) \left(1 - \sum_{j=1}^{\infty} \log\left(\frac{1}{1-\lambda_j}\right) \right) + O\left(\left(\sum_{j=1}^{\infty} \lambda_j \right)^3 \right) \\ &= \sum_{j=1}^{\infty} \left(\lambda_j + \frac{\lambda_j^2}{2} \right) - \frac{1}{2} \left(\sum_{j=1}^{\infty} \left(\lambda_j + \frac{\lambda_j^2}{2} \right) \right)^2 - \left(\sum_{l=1}^{\infty} (\lambda_l + \lambda_l^2) \right) - \\ &\quad - \left(\sum_{l=1}^{\infty} (\lambda_l + \lambda_l^2) \right) \left(1 - \sum_{j=1}^{\infty} \left(\lambda_j + \frac{\lambda_j^2}{2} \right) \right) + O\left(\left(\sum_{j=1}^{\infty} \lambda_j \right)^3 \right) \\ &= \sum_{j,l=1}^{\infty} \lambda_l \lambda_j - \sum_{l=1}^{\infty} \lambda_l^2 + O\left(\left(\sum_{j=1}^{\infty} \lambda_j \right)^3 \right) \\ &= \left(\sum_{l=1}^{\infty} \lambda_l \right)^2 - \sum_{l=1}^{\infty} \lambda_l^2 + O\left(\left(\sum_{j=1}^{\infty} \lambda_j \right)^3 \right). \end{aligned}$$

By the trace formula,

$$\text{trace}(T) = \sum_{j=1}^{\infty} \lambda_j = \int_{T_{n,k}} K_{\phi}(\zeta, \zeta) d\mu_{\phi}(\zeta),$$

and by the spectral theorem for integral Hilbert-Schmidt operators,

$$\sum_{j=1}^{\infty} \lambda_j^2 = \int_{T_{n,k}} \int_{T_{n,k}} |K_{\phi}(z, \zeta)|^2 d\mu_{\phi}(z) d\mu_{\phi}(\zeta).$$

Thus

$$\left(\sum_{j=1}^{\infty} \lambda_j\right)^2 - \sum_{j=1}^{\infty} \lambda_j^2 = \int_{T_{n,k}} \int_{T_{n,k}} [K_{\phi}(z, z)K_{\phi}(\zeta, \zeta) - |K_{\phi}(z, \zeta)|^2] d\mu_{\phi}(z)d\mu_{\phi}(\zeta),$$

as desired. \square

An important consequence of Lemma 3.4 is the following.

Proposition 3.6. *There exists $n_0 = n_0(\phi) \in \mathbb{N}$ such that for all $n \geq n_0$ and $k = 1, \dots, n$*

$$\mathbb{P}(X_{n,k} \geq 2) \simeq \frac{1}{\rho^6(z_{n,k})},$$

where the implicit constants do not depend on n and k .

Proof. We start with the formula of Remark 3.5(b). Observe that, by condition (5), for any $\eta > 0$ there exists n_0 such that $T_{n,k} \subset D(z, \eta\rho(z))$ for all $z \in T_{n,k}$, $n \geq n_0$ and $k = 1, \dots, n$. Then, by Lemma 3.3,

$$d_K(z, \zeta) \simeq \frac{|z - \zeta|}{\rho(z)}, \quad z, \zeta \in T_{n,k}.$$

Therefore

$$\mathbb{P}(X_{n,k} \geq 2) \simeq \int_{T_{n,k}} \int_{T_{n,k}} \frac{|z - \zeta|^2}{\rho^2(z)} K_{\phi}(z, z) d\mu_{\phi}(z) K_{\phi}(\zeta, \zeta) d\mu_{\phi}(\zeta) + O\left((\mathbb{E}[X_{n,k}])^3\right).$$

By (3), (11) and the fact that $\rho(\zeta) \simeq \rho(z_{n,k})$ for any $\zeta \in T_{n,k}$, we get finally

$$\begin{aligned} \mathbb{P}(X_{n,k} \geq 2) &\simeq \int_{T_{n,k}} \int_{T_{n,k}} \frac{|z - \zeta|^2}{\rho^2(z)} \frac{dm(z)}{\rho^2(z)} \frac{dm(\zeta)}{\rho^2(\zeta)} + O\left(\frac{1}{\rho^6(z_{n,k})}\right) \\ &\simeq \frac{1}{\rho^6(z_{n,k})} \left(\int_{T_{n,k}} \int_{T_{n,k}} |z - \zeta|^2 dm(z) dm(\zeta) + 1 \right) \simeq \frac{1}{\rho^6(z_{n,k})}. \end{aligned}$$

\square

Remark 3.7. The lower estimate in Proposition 3.6 can be proved in a more direct way. Since the expected value of $X_{n,k} = \sum_j \xi_j$ is close to zero, the probability that two or more points fall in $T_{n,k}$ is concentrated in the events $\{\xi_1 = 1\}$ and $\{\xi_2 = 1\}$ (since ξ_1, ξ_2 are the two Bernoulli variables with larger parameter). In any case, by the independence of ξ_1 and ξ_2 , we have the estimate

$$\mathbb{P}(X_{n,k} \geq 2) \geq \mathbb{P}((\xi_1 = 1) \cap (\xi_2 = 1)) = \lambda_1 \lambda_2.$$

By definition

$$\lambda_1 = \sup_{\substack{f \in \mathcal{F}_{\phi} \\ \|f\|_{\phi}=1}} \langle Tf, f \rangle_{\phi},$$

and, by (12) and the reproducing property of $K_\phi(z, \zeta)$,

$$\langle Tf, f \rangle_\phi = \int_{T_{n,k}} |f(z)|^2 d\mu_\phi(z).$$

Taking as f the normalized reproducing kernel for the center of $T_{n,k}$,

$$k_{n,k}(z) = \frac{K_\phi(z, z_{n,k})}{\sqrt{K_\phi(z_{n,k}, z_{n,k})}}$$

we see that

$$\lambda_1 \geq \int_{T_{n,k}} \frac{|K_\phi(z, z_{n,k})|^2}{K_\phi(z_{n,k}, z_{n,k})} d\mu_\phi(z).$$

By Proposition 3.1 (b), for $z \in T_{n,k}$, which by n large enough is contained in the disk $D(z_{n,k}, \eta\rho(z_{n,k}))$,

$$(13) \quad \lambda_1 \gtrsim \int_{T_{n,k}} K_\phi(z, z) d\mu_\phi(z) \simeq \int_{T_{n,k}} \frac{dm(z)}{\rho^2(z)} \simeq \frac{1}{\rho^2(z_{n,k})}.$$

In order to estimate λ_2 we use the formula

$$\lambda_2 = \sup_{\substack{\dim(F)=2 \\ F \subset \mathcal{F}_\phi}} \inf_{g \in F} \frac{\langle Tg, g \rangle_\phi}{\|g\|_\phi^2}.$$

Take $F \subset \mathcal{F}_\phi$ to be the subspace generated by

$$g_1(z) = k_{n,k}(z) \quad \text{and} \quad g_2(z) = \frac{z - z_{n,k}}{\rho(z_{n,k})} k_{n,k}(z).$$

Proposition 3.1 (a), together with (2) yield, for some $\varepsilon > 0$,

$$\|g_2\|_\phi^2 = \int_{\mathbb{C}} \frac{|z - z_{n,k}|^2}{\rho^2(z_{n,k})} \frac{|K_\phi(z, z_{n,k})|^2}{K_\phi(z_{n,k}, z_{n,k})} d\mu_\phi(z) \lesssim \int_{\mathbb{C}} \frac{|z - z_{n,k}|^2}{\rho^2(z_{n,k})} e^{-d_B^\varepsilon(z, z_{n,k})} \frac{dm(z)}{\rho^2(z)}.$$

By Lemma 3.2 with $k = 2$, we see that $\|g_2\|_\phi \lesssim 1$.

Also, since g_2 vanishes on $z_{n,k}$, the functions g_1 and g_2 are orthogonal in \mathcal{F}_ϕ . Then, for any function

$$g = ag_1 + bg_2 \in F \quad a, b \in \mathbb{C}$$

one has $\|g\|_\phi^2 = |a|^2 \|g_1\|_\phi^2 + |b|^2 \|g_2\|_\phi^2 \simeq |a|^2 + |b|^2$.

On the other hand, again by the estimates of Proposition 3.1 (a) and (b),

$$\begin{aligned} \langle Tg, g \rangle_\phi &\simeq \int_{T_{n,k}} \left| a + b \frac{\zeta - z_{n,k}}{\rho(z_{n,k})} \right|^2 \frac{|K_\phi(\zeta, z_{n,k})|^2}{K_\phi(z_{n,k}, z_{n,k})} d\mu_\phi(\zeta) \\ &\simeq \int_{T_{n,k}} \left| a + b \frac{\zeta - z_{n,k}}{\rho(z_{n,k})} \right|^2 \frac{dm(\zeta)}{\rho^2(\zeta)} \\ &\simeq \frac{1}{\rho^2(z_{n,k})} \int_{T_{n,k}} \left| a + b \frac{\zeta - z_{n,k}}{\rho(z_{n,k})} \right|^2 dm(\zeta). \end{aligned}$$

Let $D_{n,k} = D(z_{n,k}, 1/4)$ and notice that $D_{n,k} \subset T_{n,k}$. Since 1 and $\zeta - z_{n,k}$ are orthogonal in $L^2(D_{n,k}, dm)$,

$$\begin{aligned} \langle Tg, g \rangle_\phi &\lesssim \frac{1}{\rho^2(z_{n,k})} \int_{D_{n,k}} \left| a + b \frac{\zeta - z_{n,k}}{\rho(z_{n,k})} \right|^2 dm(\zeta) \\ &\simeq \frac{1}{\rho^2(z_{n,k})} \left(|a|^2 + \frac{|b|^2}{\rho^2(z_{n,k})} \int_{D_{n,k}} |\zeta - z_{n,k}|^2 dm(\zeta) \right) \\ &\simeq \frac{1}{\rho^2(z_{n,k})} \left(|a|^2 + \frac{|b|^2}{\rho^2(z_{n,k})} \right). \end{aligned}$$

With this

$$\lambda_2 \geq \inf_{g \in F} \frac{\langle Tg, g \rangle_\phi}{\|g\|_\phi^2} \gtrsim \inf_{a, b \in \mathbb{C}} \frac{\frac{1}{\rho^2(z_{n,k})} \left(|a|^2 + \frac{|b|^2}{\rho^2(z_{n,k})} \right)}{|a|^2 + |b|^2} = \frac{1}{\rho^4(z_{n,k})}.$$

This together with (13) yields finally

$$P(X_{n,k} \geq 2) \geq \lambda_1 \lambda_2 \gtrsim \frac{1}{\rho^2(z_{n,k})} \frac{1}{\rho^4(z_{n,k})} = \frac{1}{\rho^6(z_{n,k})}.$$

3.3. Proof of Theorem 2.1. A direct consequence of Proposition 3.6 is the following.

Proposition 3.8. *Let ϕ and $X_{n,k}$ as above. Then*

$$\mathbb{P}(X_{n,k} \geq 2 \text{ infinitely often}) = \begin{cases} 0 & \text{if } \int_{\mathbb{C}} \frac{dm(z)}{\rho^6(z)} < +\infty \\ 1 & \text{if } \int_{\mathbb{C}} \frac{dm(z)}{\rho^6(z)} = +\infty. \end{cases}$$

Proof. Assume first that $\int_{\mathbb{C}} \frac{dm(z)}{\rho^6(z)} < +\infty$. By Proposition 3.6

$$\sum_{n,k} \mathbb{P}(X_{n,k} \geq 2) \simeq \sum_{n,k} \frac{1}{\rho^6(z_{n,k})} \simeq \sum_{n,k} \int_{T_{n,k}} \frac{dm(z)}{\rho^6(z)} = \int_{\mathbb{C}} \frac{dm(z)}{\rho^6(z)} < \infty,$$

and the result follows from the first Borel-Cantelli lemma [3, Theorem 4.3].

Assume now that $\int_{\mathbb{C}} \frac{dm(z)}{\rho^6(z)} = +\infty$. Observe that if

$$E = \{X_{n,k} \geq 2 \text{ infinitely often}\},$$

then

$$E^c = \{\exists n_0 \in \mathbb{N} : \forall n \geq n_0, \forall k = 1, \dots, n, X_{n,k} \leq 1\} = \bigcup_{n_0=1}^{\infty} \bigcap_{n \geq n_0} \bigcap_{k=1}^n (X_{n,k} \leq 1).$$

Thus, in order to see that $\mathbb{P}(E^c) = 0$ it will be enough to see that for all $n_0 \in \mathbb{N}$

$$(14) \quad \mathbb{P}\left(\bigcap_{n \geq n_0} \bigcap_{k=1}^n (X_{n,k} \leq 1)\right) = 0.$$

According to a general estimate by S. Ghosh [10, Theorem 1.4], for every collection of disjoint Borel sets $\{B_j\}_{j=1}^N$ and integers $\{m_j\}_{j=1}^N$

$$\mathbb{P}\left(\bigcap_{j=1}^N (N_\phi(B_j) \leq m_j)\right) \leq \prod_{j=1}^N \mathbb{P}((N_\phi(B_j) \leq m_j)),$$

where $N_\phi(B) = \#(\Lambda_\phi \cap B)$ indicates the counting function on B .

Taking any $N > n_0$ and applying this estimate to all $T_{n,k}$ with $n_0 \leq n \leq N$, we see that

$$\mathbb{P}\left(\bigcap_{n=n_0}^N \bigcap_{k=1}^n (X_{n,k} \leq 1)\right) \leq \prod_{n=n_0}^N \prod_{k=1}^n \mathbb{P}(X_{n,k} \leq 1).$$

Letting $N \rightarrow \infty$ we get

$$\mathbb{P}\left(\bigcap_{n \geq n_0} \bigcap_{k=1}^n (X_{n,k} \leq 1)\right) \leq \prod_{n=n_0}^{\infty} \prod_{k=1}^n \mathbb{P}(X_{n,k} \leq 1),$$

hence we shall have (14) as soon as we prove that for all $n_0 \in \mathbb{N}$

$$(15) \quad \prod_{n=n_0}^{\infty} \prod_{k=1}^n \mathbb{P}(X_{n,k} \leq 1) = 0.$$

Denoting $\varepsilon_{n,k} = \mathbb{P}(X_{n,k} \geq 2)$ and recalling that, by Proposition 3.6, $\varepsilon_{n,k} \simeq \rho^{-6}(z_{n,k})$, we have

$$\begin{aligned} \prod_{n=n_0}^{\infty} \prod_{k=1}^n \mathbb{P}(X_{n,k} \leq 1) &= \exp\left(-\sum_{n=n_0}^{\infty} \sum_{k=1}^n \log\left(\frac{1}{1-\varepsilon_{n,k}}\right)\right) \\ &\simeq \exp\left(-\sum_{n=n_0}^{\infty} \sum_{k=1}^n \varepsilon_{n,k}\right) \simeq \exp\left(-\sum_{n=n_0}^{\infty} \sum_{k=1}^n \frac{1}{\rho^6(z_{n,k})}\right). \end{aligned}$$

Since the series

$$\sum_{n=1}^{\infty} \sum_{k=1}^n \frac{1}{\rho^6(z_{n,k})} \simeq \int_{\mathbb{C}} \frac{dm(z)}{\rho^6(z)}$$

diverges, we have (15), as desired. \square

Once Proposition 3.8 is settled, we finish the proof of Theorem 2.1 with a standard argument dating back, at least, to the proof of Theorem 2 in [8].

Assume first that $\int_{\mathbb{C}} \frac{dm(z)}{\rho^6(z)} < \infty$. All the arguments above can be applied analogously to the shifted regions ($n \geq 1$, $k = 1, \dots, n$):

$$\begin{aligned} \tilde{T}_0 &= \{z \in \mathbb{C} : |z| < 1/2\} = D(0, 1/2), \\ \tilde{T}_{n,k} &= \left\{z = re^{i\theta} \in \mathbb{C} : n - \frac{1}{2} \leq r < n + \frac{1}{2}, \frac{\theta}{2\pi} \in \left[\frac{k-1/2}{n}, \frac{k+1/2}{n}\right)\right\}, \end{aligned}$$

and the corresponding random variables $\tilde{X}_{n,k} = \#(\Lambda_\phi \cap \tilde{T}_{n,k})$. In particular, Proposition 3.8 holds equally if the random variables $X_{n,k}$ are replaced by the similar random variables $\tilde{X}_{n,k}$.

Consider then the events

$$E = \{X_{n,k} \geq 2 \text{ infinitely often}\}, \quad \tilde{E} = \{\tilde{X}_{n,k} \geq 2 \text{ infinitely often}\},$$

for which $\mathbb{P}(E^c) = \mathbb{P}(\tilde{E}^c) = 1$.

Under the event $E^c \cap \tilde{E}^c$, for all but finitely many n, k , we have at most one point in the regions $T_{n,k}, \tilde{T}_{n,k}$. Thus, the points of Λ_ϕ contained in these regions are separated by a fixed constant ($1/4$ will do). Since the finitely many regions where $X_{n,k}, \tilde{X}_{n,k}$ could be bigger than 1 contain at most a finite number of points of Λ_ϕ , we deduce that Λ_ϕ is almost surely separated.

Notice that we actually prove that Λ_ϕ is the union of a finite sequence and a sequence which is separated by a constant comparable to the size of the cells $T_{n,k}$.

For the case $\int_{\mathbb{C}} \frac{dm(z)}{\rho^6(z)} = \infty$ we can repeat the above arguments to a grid of size $1/l, l \geq 1$. Let

$$T_{n,k}^l := \left\{ z = re^{i\theta} \in \mathbb{C} : \frac{n-1}{l} \leq r < \frac{n}{l}, \frac{\theta}{2\pi} \in \left[\frac{k-1}{ln}, \frac{k}{ln} \right) \right\}, \quad n \geq 1, k = 1, \dots, ln.$$

Then, letting $X_{n,k}^{(l)} = \#(\Lambda_\phi \cap T_{n,k}^l)$, we see, in the same way as in Proposition 3.8, that the events

$$E^l = \{X_{n,k}^{(l)} \geq 2 \text{ infinitely often}\}$$

have all probability 1.

Under the event E^l there are infinitely many couples $\lambda_k, \lambda_j \in \Lambda_\phi$ at a distance smaller than $1/l$. Therefore, under the event $\bigcap_{l=1}^{\infty} E^l$, which still has probability 1, the sequence Λ_ϕ is not separated.

3.4. Proof of Corollary 2.4. Assume that $\int_{\mathbb{C}} \frac{dm(z)}{\rho^6(z)} < +\infty$. As mentioned in Section 2, it will be enough to see that the upper density of Λ_ϕ is zero almost surely.

For any $N \in \mathbb{N}$ consider now the cells of size N ,

$$T_{n,k}^N := \left\{ z = re^{i\theta} \in \mathbb{C} : (n-1) \leq \frac{r}{N} < n, \frac{\theta}{2\pi N} \in [(k-1), k) \right\}, \quad n \geq 1, k = 1, \dots, n$$

and their shifted versions

$$\tilde{T}_0^N = D(0, N/2)$$

$$\tilde{T}_{n,k}^N := \left\{ z = re^{i\theta} \in \mathbb{C} : n - \frac{1}{2} \leq \frac{r}{N} < n + \frac{1}{2}, \frac{\theta}{2\pi N} \in \left[k - \frac{1}{2}, k + \frac{1}{2} \right) \right\}.$$

Consider also the associated counting variables

$$X_{n,k}^{(N)} = \#(\Lambda_\phi \cap T_{n,k}^N), \quad \tilde{X}_{n,k}^{(N)} = \#(\Lambda_\phi \cap \tilde{T}_{n,k}^N).$$

The same arguments above show that the events

$$E = \{X_{n,k}^{(N)} \geq 2 \text{ infinitely often}\}, \quad \tilde{E} = \{\tilde{X}_{n,k}^{(N)} \geq 2 \text{ infinitely often}\}$$

have both probability 0, so that, after removing a finite number of points to Λ_ϕ we obtain a sequence in which the points are separated by a distance $N/4$.

This implies, in particular, that

$$D_+(\Lambda_\phi) = \limsup_{r \rightarrow \infty} \sup_{z \in \mathbb{C}} \frac{\#(\Lambda_\phi \cap D(z, r))}{\pi r^2} \lesssim \frac{1}{N}.$$

Since this is valid for any N , we deduce that, with probability one, Λ_ϕ is separated and $D_+(\Lambda_\phi) = 0$.

4. POISSON PROCESS: PROOF OF THEOREM 2.5

This is a straightforward adaptation of [11, Theorem 1.2], which we briefly recall.

It will be enough to prove the analogue of Proposition 3.8 for the Poisson point process: if

$$X_{n,k} = \#(\Lambda_\phi^P \cap T_{n,k}),$$

then

$$(16) \quad \mathbb{P}(X_{n,k} \geq 2 \text{ infinitely often}) = \begin{cases} 1 & \text{if } \int_{\mathbb{C}} \frac{dm(z)}{\rho^4(z)} < +\infty \\ 0 & \text{if } \int_{\mathbb{C}} \frac{dm(z)}{\rho^4(z)} = +\infty. \end{cases}$$

Once this is proved, the same arguments applied in the determinantal case to shifted and to thinner partitions of the plane yield the result.

By definition of the Poisson process Λ_ϕ^P , the parameter $\mathfrak{p}_{n,k}$ of the Poisson variable $X_{n,k}$ is

$$\mathfrak{p}_{n,k} = \mathbb{E}(X_{n,k}) = \text{Var}(X_{n,k}) = \int_{T_{n,k}} \frac{dm(z)}{\rho^2(z)} \simeq \frac{1}{\rho^2(z_{n,k})}.$$

Thus, we see that

$$\begin{aligned} \mathbb{P}(X_{n,k} \geq 2) &= 1 - \mathbb{P}(X_{n,k} = 0) - \mathbb{P}(X_{n,k} = 1) = 1 - e^{-\mathfrak{p}_{n,k}} - e^{-\mathfrak{p}_{n,k}} \mathfrak{p}_{n,k} \\ &= \frac{\mathfrak{p}_{n,k}^2}{2} (1 + o(1)) \simeq \frac{1}{\rho^4(z_{n,k})}. \end{aligned}$$

Hence

$$\sum_{n=1}^{\infty} \sum_{k=1}^n \mathbb{P}(X_{n,k} \geq 2) \simeq \sum_{n=1}^{\infty} \sum_{k=1}^n \frac{1}{\rho^4(z_{n,k})} \simeq \int_{\mathbb{C}} \frac{dm(z)}{\rho^4(z)},$$

and (16) follows from the second Borel-Cantelli lemma ([3, Theorem 4.4]) since, by definition of Poisson process, the different variables $X_{n,k}$ are independent.

REFERENCES

- [1] J. Agler and J.E. McCarthy, *Pick interpolation and Hilbert function spaces*. Grad. Stud. Math., 44, American Mathematical Society, Providence, RI, 2002.
- [2] N. Arcozzi, R. Rochberg, E. Sawyer, and B.D. Wick, *Distance functions for reproducing kernel Hilbert spaces*. Function spaces in modern analysis, 25–53. Contemp. Math., 547, American Mathematical Society, Providence, RI, 2011.
- [3] P. Billingsley, *Probability and Measure*. 3rd ed., John Wiley & Sons Ltd., Chichester, 1995.
- [4] J. Buckley, X. Massaneda, and J. Ortega-Cerdà, *Inhomogeneous random zero sets*. Indiana Univ. Math. J. 63 (2014), no. 3, 739–781.
- [5] A.I. Bufetov and Y. Qiu, *Determinantal point processes associated with Hilbert spaces of holomorphic functions*. Comm. Math. Phys. 351 (2017), no. 1, 1–44.
- [6] A.I. Bufetov, Y. Qiu, and A. Shamov, *Kernels of conditional determinantal measures and the Lyons-Peres completeness conjecture*. J. Eur. Math. Soc. 23 (2021), 1477–1519.
- [7] M. Christ, *On the $\bar{\partial}$ equation in weighted L^2 norms in \mathbb{C}^1* . J. Geom. Anal. 1 (1991), no. 3, 193–230.
- [8] W.G. Cochran, *Random Blaschke products*. Trans. Amer. Math. Soc. 322 (1990), no. 2, 731–755.
- [9] D.J. Daley and D. Vere-Jones, *An Introduction to the Theory of Point Processes*. Vol. I, 2nd ed., Springer-Verlag, New York, 2003.
- [10] S. Ghosh, *Determinantal processes and completeness of random exponentials: the critical case*. Probab. Theory Related Fields 163 (2015), no. 3–4, 643–665.
- [11] A. Hartmann and X. Massaneda, *Inhomogeneous Poisson processes in the disk and interpolation*. Proc. Edinb. Math. Soc. (2) 67 (2024), no. 3, 749–777.
- [12] J.B. Hough, M. Krishnapur, Y. Peres, and B. Virág, *Zeros of Gaussian analytic functions and determinantal point processes*. University Lecture Series, 51, American Mathematical Society, Providence, RI, 2009.
- [13] G. Last and M. Penrose, *Lectures on the Poisson Process*. Institute of Mathematical Statistics Textbooks, 7, Cambridge University Press, Cambridge, 2018.
- [14] O. Macchi, *The coincidence approach to stochastic point processes*. Adv. Appl. Probab. 7 (1975), 83–122.
- [15] N. Marco, X. Massaneda, and J. Ortega-Cerdà, *Interpolating and sampling sequences for entire functions*. Geom. Funct. Anal. 13 (2003), no. 4, 862–914.
- [16] J. Marzo and J. Ortega-Cerdà, *Pointwise estimates for the Bergman kernel of the weighted Fock space*. J. Geom. Anal. 19 (2009), no. 4, 890–910.
- [17] B. Rider and B. Virág, *Complex determinantal processes and HI noise*. Electron. J. Probab. 12 (2007), no. 45, 1238–1257.
- [18] K. Seip, *Density theorems for sampling and interpolation in the Bargmann-Fock space. I*. J. Reine Angew. Math. 429 (1992), 91–106.
- [19] K. Seip and R. Wallstén, *Density theorems for sampling and interpolation in the Bargmann-Fock space. II*. J. Reine Angew. Math. 429 (1992), 107–113.
- [20] A. Soshnikov, *Determinantal random point fields*. Russian Math. Surveys 55 (2000), no. 5, 923–975.
- [21] P. Walker, *The zeros of the partial sums of the exponential series*. Amer. Math. Monthly 110 (2003), no. 4, 337–339.
- [22] K. Zhu, *Analysis on Fock Spaces*. Graduate Texts in Mathematics, Springer, Boston, MA, 2012.

UNIV. BORDEAUX, CNRS, BORDEAUX INP, IMB, UMR 5251, F-33400, TALENCE, FRANCE

DEPARTAMENT DE MATEMÀTIQUES I INFORMÀTICA, UNIVERSITAT DE BARCELONA, GRAN VIA 585, 08007-BARCELONA, CATALONIA