On a conjecture of Pach-Spencer-Tóth for graph crossing numbers

Kaizhe Chen^{*} Jie Ma[†]

February 5, 2025

Abstract

The crossing number of a graph G denotes the minimum number of crossings in any planar drawing of G. In this short note, we confirm a long-standing conjecture posed by Pach, Spencer, and Tóth over 25 years ago, establishing an optimal lower bound on the crossing number of graphs that satisfy some monotone properties. Furthermore, we address a related open problem introduced by Pach and Tóth in 2000, which explores the interplay between the crossing number of a graph, its degree sequence, and its bisection width.

1 Introduction

Let G be a (simple) graph. An arc is the image of a continuous injective map $[0,1] \to \mathbb{R}^2$. A planar drawing of G is a mapping f that assigns to each vertex v of G a point f(v) in the plane and to each edge xy of G an arc connecting f(x) and f(y), not passing through the image of any other vertex. We assume that no three edges have an interior point in common. The crossing number $\operatorname{cr}(G)$ of G denotes the minimum number of crossings in any planar drawing of G. Throughout this paper, for a graph G, we let n(G) denote the number of vertices in G and let e(G) denote the number of edges in G.

The study of crossing numbers lies at the nexus of discrete geometry and algorithmic design, with applications ranging from geometric embeddings to VLSI layout optimization. Investigating crossing numbers and their variants, such as the rectilinear crossing number, reveals their deep connections to combinatorial geometry, hardness of approximation, and efficient algorithms, making them central to both foundational and applied research in discrete and computational geometry (see [15] for a comprehensive survey). The following famous result, known as *the crossing lemma*, which is proved by Ajtai, Chvátal, Newborn and Szemerédi [2] and independently by Leighton [7], gives a general lower bound for the

^{*}School of the Gifted Young, University of Science and Technology of China, Hefei, Anhui 230026, China. Email: ckz22000259@mail.ustc.edu.cn.

[†]School of Mathematical Sciences, University of Science and Technology of China, Hefei, Anhui 230026, China, and Yau Mathematical Sciences Center, Tsinghua University, Beijing 100084, China. Research supported by National Key Research and Development Program of China 2023YFA1010201 and National Natural Science Foundation of China grant 12125106. Email: jiema@ustc.edu.cn.

crossing number of graphs with given numbers of vertices and edges: Every *n*-vertex graph G with $e \ge 4n$ edges satisfies

$$\operatorname{cr}(G) \ge \frac{1}{64} \frac{e^3}{n^2}.$$
(1)

Improvements on the constants can be found in [12, 1]. Another influential result proved by Garey and Johnson [5] states that the crossing number problem is NP-complete.

Answering a question of Simonovits, Pach, Spencer and Tóth [11] proved that the general lower bound (1) can be improved substantially for graphs satisfying certain monotone properties, as follows. Let G be a graph with n vertices and e edges. Suppose that there are constants $A, \alpha > 0$ such that

any subgraph
$$H$$
 of G satisfies $e(H) \le A \cdot (n(H))^{1+\alpha}$. (2)

Then there exist constants c, c' > 0 depending only on A and α such that

if
$$e \ge cn \log^2 n$$
, then the crossing number of G satisfies $\operatorname{cr}(G) \ge c' \frac{e^{2+1/\alpha}}{n^{1+1/\alpha}}$. (3)

This bound is tight up to a constant factor as shown in [11]. Furthermore, Pach, Spencer and Tóth [11] conjectured that the same statement holds even for all *n*-vertex graphs with $e \ge cn$ edges for a suitable constant c > 0 (see also [14]).

Conjecture 1.1 (Pach-Spencer-Tóth, [11]). Let G be a graph with n vertices and e edges satisfying (2) for some constants $A, \alpha > 0$. Then there exist constants c, c' > 0 depending only on A and α such that if $e \ge cn$, then $\operatorname{cr}(G) \ge c' \frac{e^{2+1/\alpha}}{n^{1+1/\alpha}}$.

The authors of [11] also verified this conjecture for several interesting monotone properties, including the family of $K_{s,t}$ -free graphs as well as the family of C_{2k} -free graphs for $k \in \{2,3\}$. Füredi and Kündgen [4] obtained an improvement on (3) by showing that in the case $\alpha \in (0, 1/2)$, the same bound holds under the weaker condition $e \ge cn \log n$.

In this paper, we resolve Conjecture 1.1 by proving the following theorem. Our argument refines the approach developed in [11, 4].

Theorem 1.2. Let G be a graph with n vertices and e edges. Suppose that there are constants $A, \alpha > 0$ such that any subgraph H of G satisfies $e(H) \leq A (n(H))^{1+\alpha}$. Then there exist constants c, c' > 0 depending only on A and α such that

if
$$e \ge cn$$
, then the crossing number of G satisfies $\operatorname{cr}(G) \ge c' \frac{e^{2+1/\alpha}}{n^{1+1/\alpha}}$.

The following result is a direct consequence of Theorem 1.2, where the cases of $k \in \{2, 3\}$ are first proved in [11, Theorems 3.1 and 3.2].

Corollary 1.3. Let $k \ge 2$ be an integer and G be a graph of n vertices and e edges, which contains no cycle of length 2k. Then there exist two constants c, c' > 0 depending only on k such that if $e \ge cn$, then the crossing number of G satisfies

$$\operatorname{cr}(G) \ge c' \frac{e^{2+k}}{n^{1+k}}.$$

The study of crossing numbers is deeply intertwined with other graph parameters, particularly the *bisection width*, which has proven instrumental in bridging combinatorial and topological properties of graphs. Let G be a graph with n vertices. For any bipartition $V(G) = V_1 \cup V_2$, let $E(V_1, V_2)$ denote the set of edges in G with one endpoint in V_1 and the other endpoint in V_2 . The bisection width, denoted by b(G), of G is defined as

$$b(G) = \min |E(V_1, V_2)|,$$

where the minimum is taken over all bipartitions $V(G) = V_1 \cup V_2$ with $|V_1|, |V_2| \ge n/3$. Leighton [6] was the first to identify a strong correlation between the bisection width and the crossing number of a graph, rooted from the Lipton-Tarjan separator theorem [8] for planar graphs. This was enhanced in the following remarkable theorem, established by Pach, Shahrokhi and Szegedy [10] and independently by Sýkora and Vrt'o [17]: Let G be a graph of n vertices, whose degrees are $d_1, d_2, ..., d_n$. Then it holds that

$$b(G) \le 6.32\sqrt{\operatorname{cr}(G)} + 1.58\sqrt{\sum_{i=1}^{n} d_i^2}.$$
 (4)

We would like to emphasize that the proof of (3) in [11] heavily relies on the use of the above theorem. The example of a *star* shows that the inequality (4) does not remain true if we remove the last term on its right hand side. Pach and Tóth [13, Problem 5] asked whether the dependence of the inequality (4) on the degrees of the vertices can be improved.

Problem 1.4 (Pach-Tóth, [13], Problem 5). Determine the possible values of t such that

$$b(G) = O\left(\sqrt{\operatorname{cr}(G)} + \left(\sum_{i=1}^{n} d_{i}^{t}\right)^{1/t}\right)$$
(5)

holds for every graph G of n vertices with degrees $d_1, d_2, ..., d_n$.

In the next result, we solve this problem by determining all possible values of t for which (5) holds.

Theorem 1.5. Let t be a positive constant. For $0 < t \leq 2$, every graph G of n vertices with degrees $d_1, d_2, ..., d_n$ satisfies

$$b(G) = O\left(\sqrt{\operatorname{cr}(G)} + \left(\sum_{i=1}^{n} d_i^t\right)^{1/t}\right).$$
(6)

For t > 2, there exist infinitely many integers n with a graph G of n vertices having degrees $d_1, d_2, ..., d_n$ such that

$$b(G) \ge \frac{1}{12} n^{1/2 - 1/t} \cdot \left(\sqrt{\operatorname{cr}(G)} + \left(\sum_{i=1}^{n} d_i^t \right)^{1/t} \right).$$
(7)

Therefore, (5) holds if and only if $0 < t \leq 2$.

Finally, we would like to present the following result as a dual theorem of Theorem 1.2, demonstrating the possibility to bound the number of edges in a graph through estimates on the crossing numbers.

Theorem 1.6. Let G be a graph. Suppose that there are constants $N, \alpha > 0$ such that any subgraph H of G with at least N edges satisfies

$$\operatorname{cr}(H) \le \frac{(e(H))^2}{2^{16+3/\alpha}}.$$

Then there exists a constant A which depends only on N and α such that

$$e(G) \le A \cdot (n(G))^{1+\alpha}$$
.

We hope that Theorem 1.6 would facilitate the solution of other unsolved problems in combinatorial and computational geometry, such as the unit distance problem and the k-sets problem (see e.g. [9]). For instance, if one can prove that the crossing number of any unit distance graph G of n vertices and e edges satisfies $cr(G) = O(e^2/\log \log e)$, then Theorem 1.6 implies that $e = n^{1+o(1)}$.

The rest of the paper is organized as follows. In Section 2, we prove Theorem 1.5. In Section 3, we prove Theorem 1.2, from which we derive Corollary 1.3 and Theorem 1.6.

2 Crossing number and bisection width

Proof of Theorem 1.5. First, we consider the case that $0 < t \leq 2$. For any $x \geq 0$, let $f(x) = x^{2/t}$. Then the second derivative of f(x) is

$$f''(x) = \frac{2}{t} \left(\frac{2}{t} - 1\right) x^{2/t - 2} \ge 0.$$

For $1 \leq i \leq n$, let $x_i = d_i^t$. It follows by the Jensen's inequality that

$$\sum_{i=1}^{n} f(x_i) \le (n-1)f(0) + f\left(\sum_{i=1}^{n} x_i\right) = f\left(\sum_{i=1}^{n} x_i\right).$$

That is

$$\left(\sum_{i=1}^n d_i^2\right)^{1/2} \le \left(\sum_{i=1}^n d_i^t\right)^{1/t}$$

Then, the inequality (6) follows from (4).

Now suppose that t > 2. Let G be a graph defined as follows. The vertex set of G is the following subset of integer points in \mathbb{R}^2

$$V(G) = \{(i, j) | i, j \in [n]\}, \text{ where } [n] = \{1, 2, ..., n\}.$$

Two vertices in V(G) are adjacent if and only if the Euclidean distance between them is one (see Figure 1). It is clear that G is a planar graph with n^2 vertices of maximum degree




Figure 1: A drawing of G with n = 5

four. We claim that $b(G) \ge n/3$. Indeed, let $V(G) = V_1 \cup V_2$ be any partition of V(G) such that $|V_1|, |V_2| \ge n^2/3$. It suffices to prove that $|E(V_1, V_2)| \ge n/3$.

Suppose that there is an $i_0 \in [n]$ such that $(i_0, j) \in V_1$ for every $j \in [n]$. Since $|V_1| \le 2n^2/3$, there are at most 2n/3 number of j such that $(i, j) \in V_1$ for every $i \in [n]$. Thus, there are at least n/3 columns containing at least one edge in $E(V_1, V_2)$, and hence $|E(V_1, V_2)| \ge n/3$.

Now, we assume that there does not exist an $i_0 \in [n]$ such that $(i_0, j) \in V_1$ for every $j \in [n]$. That is, each row contains at least one vertex in V_2 . Since $|V_1| \ge n^2/3$, there are at least n/3 rows containing at least one vertex in V_1 . Each such row contains at least one edge in $E(V_1, V_2)$, and hence $|E(V_1, V_2)| \ge n/3$.

Thus, $b(G) \ge n/3$. Since cr(G) = 0, we have

$$\sqrt{\operatorname{cr}(G)} + \left(\sum_{i=1}^{n^2} d_i^t\right)^{1/t} \le \left(\sum_{i=1}^{n^2} 4^t\right)^{1/t} = 4n^{2/t} \le 12n^{2/t-1}b(G),$$

completing the proof.

Remark 2.1. In the proof of Theorem 1.5, for any integer n, we give a counterexample G of n^2 vertices which is planar, i.e., $\operatorname{cr}(G) = 0$. The graph G can be easily turned into counterexamples G' with unbounded $\operatorname{cr}(G')$, by the following operations: choosing any $o(n^2)$ edges of G and replacing each of these edges by a copy of $K_{s,s}$ (for any fixed $s \geq 3$).¹ In this way, we can obtain a non-planar counterexample with $o(n^2)$ number of crossings.

3 Proofs of main results

In this section, we prove Theorem 1.2, Corollary 1.3 and Theorem 1.6.

Proof of Theorem 1.2. Let G be a graph with n vertices and e edges. Suppose that there are constants $A, \alpha > 0$ such that, for any subgraph H of G, we have

$$e(H) \le A(n(H))^{1+\alpha}.$$
(8)

To prove Theorem 1.2, we assume for a contradiction that $e \ge cn$,

$$\operatorname{cr}(G) < c' \frac{e^{2+1/\alpha}}{n^{1+1/\alpha}},$$

¹To be more precise, let S be the set of vertices in these $o(n^2)$ edges. Then G' is obtained from G by replacing each vertex $v \in S$ with an independent set I_v of size s, adding all edges between vertices in I_v and vertices $u \in V(G) \setminus S$ if $uv \in E(G)$, and adding all edges between I_v and I_u if $uv \in E(G)$.

and G is drawn in the plane with exactly cr(G) crossings, where c and c' are suitable constants which will be determined later (see (18)).

First, we split every vertex of G whose degree exceeds $\overline{d} := 2e/n$ into vertices of degree at most \overline{d} , as follows. Let v be a vertex of G such that the degree d of v in G satisfies $d > \overline{d}$. Let $vw_1, vw_2, ..., vw_d$ be the edges incident to v, listed in clockwise order. Replace v by $\lceil d/\overline{d} \rceil$ new vertices, $A_v = \{v_1, v_2, ..., v_{\lceil d/\overline{d} \rceil}\}$, placed in clockwise order on a very small circle C around v. Connect w_j to v_i if and only if $\overline{d}(i-1) < j \leq \overline{d}i$ for all $1 \leq j \leq d$ and $1 \leq i \leq \lceil d/\overline{d} \rceil$. Let the circle C be small enough such that we do not introduce any new crossings. Repeat this procedure until the degree of each vertex is at most \overline{d} , and denote the resulting graph by G'.

By the construction of G', we have e(G') = e(G) = e, and

$$\operatorname{cr}(G') \le \operatorname{cr}(G) < c' \frac{e^{2+1/\alpha}}{n^{1+1/\alpha}}.$$
(9)

Observe that

$$n(G') - n(G) = \sum_{v:d(v) > \overline{d}} (|A_v| - 1) < \sum_{v:d(v) > \overline{d}} \frac{d(v)}{\overline{d}} \le \frac{2e}{\overline{d}} = n$$

We deduce $N := n(G') \in [n, 2n)$.

Now, we break G' into smaller components according to the following algorithm:

Algorithm 1 (Decomposition Algorithm)

- 1: Let $G^0 = G', G_1^0 = G', M_0 = 1$, and $m_0 = 1$.
- 2: For $i \ge 0$, suppose that G^i consists of M_i components $G_1^i, G_2^i, ..., G_{M_i}^i$, each of at most $(2/3)^i N$ vertices. Let $m_i \in [0, M_i]$ be the constant such that

$$\left(\frac{2}{3}\right)^{i+1} N \le n(G_j^i) \le \left(\frac{2}{3}\right)^i N, \text{ for } 1 \le j \le m_i; \tag{10}$$

$$n(G_j^i) < \left(\frac{2}{3}\right)^{i+1} N, \text{ for } m_i < j \le M_i.$$
 (11)

3: Recall the constants A, α from (8). If

$$\left(\frac{2}{3}\right)^i \ge \frac{1}{(2A)^{1/\alpha}} \frac{e^{1/\alpha}}{N^{1+1/\alpha}},$$

then for every $1 \leq j \leq m_i$, delete $b(G_j^i)$ edges from G_j^i to obtain two induced subgraphs, each of at most $(2/3)n(G_j^i)$ vertices. Let G^{i+1} denote the resulting graph on N vertices. Then each component of G^{i+1} has at most $(2/3)^{i+1}N$ vertices. Return to Step 2. 4: Else STOP.

Suppose that the Decomposition Algorithm terminates when i = k. If k = 0, then

$$1 < \frac{1}{(2A)^{1/\alpha}} \frac{e^{1/\alpha}}{N^{1+1/\alpha}}.$$

It follows that

$$e > 2AN^{1+\alpha} \ge 2An^{1+\alpha},$$

which is contradictory to the assumption (8). Thus k > 0, and

$$\left(\frac{2}{3}\right)^k < \frac{1}{(2A)^{1/\alpha}} \frac{e^{1/\alpha}}{N^{1+1/\alpha}} \le \left(\frac{2}{3}\right)^{k-1}.$$
(12)

We will first show that G^k contains less than e/2 edges. The number of vertices of each component of G^k satisfies

$$n(G_j^k) \le \left(\frac{2}{3}\right)^k N < \frac{1}{(2A)^{1/\alpha}} \frac{e^{1/\alpha}}{N^{1+1/\alpha}} N = \left(\frac{e}{2AN}\right)^{1/\alpha}.$$
(13)

Let V_j be the subset of V(G) defined as follows:

$$V_j = \{ v \in V(G) | v \in V(G_j^k) \text{ or } A_v \cap V(G_j^k) \neq \emptyset \},\$$

i.e., V_j is the vertex subset consisting of all 'pre-images' of $V(G_j^k)$. The construction of G' implies that $|V_j| \leq n(G_j^k)$. Let $G[V_j]$ be the subgraph of G induced by V_j . By the assumption (8) and the inequality (13), we have

$$e(G[V_j]) \le A|V_j|^{1+\alpha} \le A\left(n(G_j^k)\right)^{1+\alpha} < An(G_j^k) \cdot \frac{e}{2AN} = \frac{e}{2N}n(G_j^k).$$
(14)

By the construction of G', each edge in G_j^k has a corresponding edge in $G[V_j]$. Thus, we have

$$e(G_j^k) \le e\left(G[V_j]\right)$$

It follows, as desired, that

$$e(G^k) = \sum_{j=1}^{M_k} e(G_j^k) < \frac{e}{2N} \sum_{j=1}^{M_k} n(G_j^k) = \frac{e}{2}.$$

To obtain a contradiction, it now suffices to show that we deleted at most e/2 edges of G' to obtain G^k . For any $0 \le i < k$, the inequality (10) implies that

$$m_i \le \left(\frac{3}{2}\right)^{i+1}.\tag{15}$$

It follows by the Cauchy-Schwarz inequality and (9) that,

$$\sum_{j=1}^{m_i} \sqrt{\operatorname{cr}(G_j^i)} \le \sqrt{m_i \sum_{j=1}^{m_i} \operatorname{cr}(G_j^i)} \le \sqrt{\left(\frac{3}{2}\right)^{i+1} \operatorname{cr}(G')} < \sqrt{\left(\frac{3}{2}\right)^{i+1}} \sqrt{c' \frac{e^{2+1/\alpha}}{n^{1+1/\alpha}}}.$$
 (16)

For any vertex v in G^i , denote by $d(v, G^i)$ the degree of v in G^i . Since G^i is a subgraph of G', we have

$$\max_{v \in V(G^i)} d(v, G^i) \le \max_{v \in V(G')} d(v, G') \le \overline{d},$$

and

$$\sum_{v \in V(G^i)} d(v, G^i) \le \sum_{v \in V(G')} d(v, G') = 2e(G') = 2e.$$

Thus, using the Cauchy-Schwarz inequality again, we deduce that for every $0 \le i < k$

$$\sum_{j=1}^{m_i} \sqrt{\sum_{v \in V(G_j^i)} d^2(v, G_j^i)} \leq \sqrt{m_i \sum_{v \in V(G^i)} d^2(v, G^i)}$$
$$\leq \sqrt{\left(\frac{3}{2}\right)^{i+1}} \sqrt{\max_{v \in V(G^i)} d(v, G^i) \sum_{v \in V(G^i)} d(v, G^i)}$$
$$\leq \sqrt{\left(\frac{3}{2}\right)^{i+1}} \sqrt{\overline{d} \cdot 2e} = \sqrt{\left(\frac{3}{2}\right)^{i+1}} \cdot \frac{2e}{\sqrt{n}}.$$
(17)

Using the theorem (4) and inequalities (16) and (17), the total number σ of edges deleted during the Decomposition Algorithm is

$$\begin{split} \sigma &= \sum_{i=0}^{k-1} \sum_{j=1}^{m_i} b(G_j^i) \le 6.32 \sum_{i=0}^{k-1} \sum_{j=1}^{m_i} \sqrt{\operatorname{cr}(G_j^i)} + 1.58 \sum_{i=0}^{k-1} \sum_{j=1}^{m_i} \sqrt{\sum_{v \in V(G_j^i)} d^2(v, G_j^i)} \\ &< 6.32 \sqrt{c' \frac{e^{2+1/\alpha}}{n^{1+1/\alpha}}} \sum_{i=0}^{k-1} \sqrt{\left(\frac{3}{2}\right)^{i+1}} + \frac{3.16e}{\sqrt{n}} \sum_{i=0}^{k-1} \sqrt{\left(\frac{3}{2}\right)^{i+1}} \\ &= \sqrt{\frac{3}{2}} \frac{\sqrt{(3/2)^k} - 1}{\sqrt{3/2} - 1} \left(6.32 \sqrt{c' \frac{e^{2+1/\alpha}}{n^{1+1/\alpha}}} + \frac{3.16e}{\sqrt{n}} \right) \end{split}$$

By the inequality (12) that $(3/2)^{k-1} \leq \frac{(2A)^{1/\alpha}N^{1+1/\alpha}}{e^{1/\alpha}}$ and the fact that N < 2n, we have

$$\sigma < 6.7 \sqrt{\frac{(2A)^{1/\alpha} N^{1+1/\alpha}}{e^{1/\alpha}}} \left(6.32 \sqrt{c' \frac{e^{2+1/\alpha}}{n^{1+1/\alpha}}} + \frac{3.16e}{\sqrt{n}} \right)$$
$$< 45 \sqrt{2^{1+2/\alpha} A^{1/\alpha} c' e^2} + 22 \sqrt{2^{1+2/\alpha} A^{1/\alpha} n^{1/\alpha} e^{2-1/\alpha}}.$$

Each term on the right hand side of the above inequality is at most e/4, provided that

$$e > cn, \ c' = \frac{1}{(180)^2 2^{1+2/\alpha} A^{1/\alpha}} \quad \text{and} \quad c = (88)^{2\alpha} 2^{\alpha+2} A.$$
 (18)

Hence $\sigma < e/2$, which is a contradiction, completing the proof.

To prove Corollary 1.3, we need the following classic theorem of Bondy-Simonovits [3].

Theorem 3.1 (Bondy-Simonovits, [3]). For a fixed integer $k \ge 2$, every n-vertex graph G that does not contain any cycle of length 2k has at most $100k \cdot n^{1+1/k}$ edges.

Proof of Corollary 1.3. It's a direct consequence of Theorem 1.2 and Theorem 3.1. \Box

To conclude this section, we give a proof of Theorem 1.6.

Proof of Theorem 1.6. Let $N, \alpha > 0$ be constants. We say a graph G is (N, α) -good, if any subgraph H of G with at least N edges satisfies

$$\operatorname{cr}(H) \le \frac{(e(H))^2}{2^{16+3/\alpha}}.$$
 (19)

We point out that if G is (N, α) -good, then any subgraph of G is (N, α) -good as well. Let G be an (N, α) -good graph with n vertices and e edges. We aim to show that $e \leq An^{1+\alpha}$, where $A = \max\{88^22^{1+3/\alpha}, N\}$.

Suppose for a contradiction that $e > An^{1+\alpha}$. Without losing generality, we may further assume that G is the minimum counterexample. Consider any proper subgraph H of G. Since H is also (N, α) -good, we have $e(H) \leq A (n(H))^{1+\alpha}$. This also implies that $e - 1 \leq An^{1+\alpha}$, which further implies that $e \leq An^{1+\alpha} + 1 \leq 2An^{1+\alpha}$. Now we derive that any subgraph H of G satisfies $e(H) \leq 2A (n(H))^{1+\alpha}$. Using Theorem 1.2, if $e \geq cn$, then

$$\operatorname{cr}(G) \ge c' \frac{e^{2+1/\alpha}}{n^{1+1/\alpha}},\tag{20}$$

where

$$c' = \frac{1}{(180)^2 2^{1+2/\alpha} (2A)^{1/\alpha}}$$
 and $c = (88)^{2\alpha} 2^{\alpha+2} (2A)$.

Since $n^2 \ge e > An^{1+\alpha} \ge An$, we have n > A, and hence

$$e > An^{1+\alpha} > A^{1+\alpha}n \ge (88)^{2\alpha}2^{\alpha+3}An = cn.$$

Combining the inequalities (19) and (20), we derive

$$\frac{e^2}{2^{16+3/\alpha}} \ge \operatorname{cr}(G) \ge c' \frac{e^{2+1/\alpha}}{n^{1+1/\alpha}} > c' \frac{e^2 \left(An^{1+\alpha}\right)^{1/\alpha}}{n^{1+1/\alpha}} = \frac{e^2}{(180)^{2}2^{1+3/\alpha}}.$$

Simplifying the above inequality results in the incorrect statement $(180)^2 > 2^{15}$, thereby completing the proof.

References

- [1] E. Ackerman, On topological graphs with at most four crossings per edge, Computational Geometry 85 (2019), 35 pages.
- [2] M. Ajtai, V. Chvátal, M. Newborn and E. Szemerédi, Crossing-free subgraphs, Theory and Practice of Combinatorics, Mathematical Studies, Vol. 60, North-Holland, Amsterdam (1982), 9-12.
- [3] J. A. Bondy and M. Simonovits, Cycles of even length in graphs, J. Combin. Theory Ser. B 16 (1974), 97-105.
- [4] Z. Füredi and A. Kündgen, Moments of graphs in monotone families, J. Graph Theory 57 (2006), 37-48.

- [5] M. Garey and D. Johnson, Crossing number is NP-complete, SIAM Journal on Algebraic and Discrete Methods 4 (1983), 312-316.
- [6] F. T. Leighton, Complexity Issues in VLSI, MIT Press, Cambridge, MA, 1983.
- [7] F. T. Leighton, New lower bound techniques for VLSI, Math. Systems Theory 17 (1984), 47-70.
- [8] R. Lipton and R. Tarjan, A separator theorem for planar graphs, SIAM J. Appl. Math., 36 (1979), 177-189.
- [9] J. Matousek, Lectures on discrete geometry (Vol. 212), Springer Science & Business Media (2013).
- [10] J. Pach, F. Shahrokhi and M. Szegedy, Applications of the crossing number, Algorithmica 16 (1996), 111-117.
- [11] J. Pach, J. Spencer and G. Tóth, New bounds on crossing numbers, ACM Symposium on Computational Geometry (Miami, 1999), Discrete Comput. Geom. 24 (2000), 623-644.
- [12] J. Pach and G. Tóth, Graphs drawn with few crossings per edge, Combinatorica 17 (1997), 427-439.
- [13] J. Pach and G. Tóth, Thirteen problems on crossing numbers, Geombinatorics, 9 (2000), 194-207.
- [14] R. B. Richter and G. Salazar, Crossing numbers, Topics in topological graph theory 128 (2009), 133-150.
- [15] M. Schaefer, The Graph Crossing Number and its Variants: A Survey, the Electronic Journal of Combinatorics (2024), #DS21. (DOI: https://doi.org/10.37236/2713)
- [16] L. A. Székely, Crossing numbers and hard Erdős problems in discrete geometry, Combin. Probab. Comput. 6 (1998), 353-358.
- [17] O. Sýkora and I. Vrt'o, On VLSI layouts of the star graph and related networks, Integration, 17 (1994), 83-93.