# Limitations of deducing measures of limsup sets from measures of finite intersections

Charlie Wilson

February 5, 2025

#### Abstract

Early results by Borel-Cantelli and more recently Erdös-Chung, Frolov and Feng-Li-Shen have provided bounds for the measure of the limsup set in terms of measures of its constituent sets and their intersections. Despite progress being made to give stronger bounds we show there is no hope for finding non-trivial bounds for the limsup measure from the measures of pairwise intersections in general. We go further and construct counterexamples wherein we have two sequences of sets whose individual measures all the way up to k-wise intersection measure  $\mu(A_{i_1} \cap A_{i_2} \cap ... \cap A_{i_k})$  are comparable but one sequence will have limsup measure 1 and the other 0. We later to go on to see that if we control all finite intersections (not just up to k intersections) then we do indeed fix the limsup measure.

#### Keywords

Measure theory, Borel Cantelli, Diophantine approximation, Probability theory

AMS subject classification 28A75 11K60, 37A50.

#### Acknowledgements

Thanks to Mark Holland and Demi Allen for the ongoing support and help with the project. Also thanks to Felipe Ramirez for the advice and suggestions. Thanks also to the backing of DWN, CBR and HBY.

## 1 Introduction

We consider a probability space  $(\Omega, \mathscr{A}, \mu)$ . We are interested in the limsup of sets, wherein we consider a sequence of sets  $(A_i)_{i \in \mathbb{N}}$  and define the limsup to be:

$$\limsup_{n \to \infty} A_n = \bigcap_{n \ge 1} \bigcup_{i \ge n} A_i \; .$$

For ease of notation we may refer to this set as  $\limsup A_i$  or simply the 'limsup' setsurpressing the  $n \to \infty$ . The limsup set is usually viewed as the set of points that lie in infinitely many of the sets  $A_i$ .

In areas such as metric number theory, Diophantine approximation and shrinking target problems in Dynamics we wish to show that a limsup has positive measure or even full measure. There are blackbox tools for doing this such as the Borel Cantelli lemmas or Kochen and Stone (stated and referenced later). Indeed a recent result of Beresnevich and Velani [BV23] states that if  $A_i$  is a sequence of balls having positive measure limsup set, and  $B_i$  is a sequence of balls having identical measures and pairwise measures, then the limsup of  $B_i$  also has psoitive measure.

A question arising therefore, is to ask whether the condition of them being balls is strictly necessary. Consider two sequences of sets  $(A_i)_{i \in \mathbb{N}}$  and  $(B_i)_{i \in \mathbb{N}}$  with the following properties:

$$\mu(A_i) \ge \mu(B_i), \quad \forall i \in \mathbb{N}, \\ \mu(A_i \cap A_j) \le \mu(B_i \cap B_j), \quad \forall i, j \in \mathbb{N}.$$

$$(1)$$

- † Does  $\mu(\limsup B_i) = 1 \implies \mu(\limsup A_i) = 1$ ?
- ‡ If the first fails, does at least:  $\mu(\limsup B_i) = 1 \implies \mu(\limsup A_i) > 0$ ?

Naively we may think the original Kochen Stone Theorem as stated as Theorem 2.1 in Section 2 may indicate truth in these statements. This result is a furthering of the Divergence Borel-Cantelli (also stated in 2) which would suggest that bigger measures of sets and smaller intersection should increase the limsup measure. As mentioned earlier Theorem 2.4, taken from [BV23], shows that if we assume the  $A_i$  and  $B_i$  are actually balls then the second question (‡) is indeed true. It is natural to feel it could be true but indeed we'll see that it is false for more general sets than balls. We will see two sequences of sets satisfying (1) with  $\mu(\limsup B_i) = 1$  and  $\mu(\limsup A_i) = 0$ .

The examples expressed in Section 4 (Proposition 4.1 and Proposition 4.2) shows the answer to the first question (†) is indeed false. As explained later, the ideas generally come from Venn diagrams, which provide intuitive visualizations for the problem. We also give analogous examples in the Lebesgue measure situation.

Similarly the answer to the second question (‡) can be found in Section 5 (Theorem 5.1 and Theorem 5.2). We construct two sequences of sets which satisfy (1) with differing limsup measures thereby answering the second question. It also encourages us to ask about the consequences if we ask for some control of measures of higher intersections. A discussion of this problem can be found at the beginning of Section 6 where we ask which way the inequalities should go for the question to have substance. We ultimately look to the inclusion exclusion principle and note these choices also are supported by the work of Frolov in [Fro12]. We see even in this case the limsup cannot be completely controlled and give examples of this using some elaborate sets with interesting intersection properties, this is done in Theorem 1.1.

**Theorem 1.1.** For a given  $m \in \mathbb{N}$ , there exists  $(A_i)_{i \in \mathbb{N}}$  and  $(B_i)_{i \in \mathbb{N}}$  in the space [0,1]endowed with the Borel sigma algebra and Lebesgue measure satisfying:

$$\mu(A_i) \ge \mu(B_i) \quad \forall i \in \mathbb{N}$$
  

$$\mu(A_i \cap A_j) \le \mu(B_i \cap B_j) \quad \forall i < j \in \mathbb{N}$$
  

$$(-1)^r \mu(A_{i_1} \cap \dots \cap A_{i_r}) \le (-1)^r \mu(B_{i_1} \cap \dots \cap B_{i_r}) \quad \forall 1 \le r \le m, \quad \forall i_1 < \dots < i_r \in \mathbb{N} \quad (2)$$
  

$$\mu(\limsup A_i) = 0$$
  

$$\mu(\limsup B_i) = 1$$

This result clearly furthers the answers to those of  $(\dagger)$  and  $(\ddagger)$  and thus Theorem 5.1 and Theorem 5.2. Readers only interested in this overarching result are advised to skip to Section 6. The sections 4 and 5 are largely included for the easy to follow examples presented and give the intuition to make Section 6 more palatable.

We see then that fixing the measures of up to level k intersections between sets doesn't necessarily fix the limsup measure. A natural question is that if we fix all of the finite intersections, does this indeed imply that the limsups is fixed? We show that in Section 3 this is true, for two sequences of sets which have the measures of the finite level intersections identical then their limsups must have the same measure. Indeed we can even replace the equalities with appropriate inequalities (dependent on the number of sets intersected) and then we can bound the limsup measure of one sequence by the limsup measure of the other. The result is stated below:

**Theorem 1.2.** Let  $(\Omega_1, \mathscr{A}_1, \mu_1)$  and  $(\Omega_2, \mathscr{A}_2, \mu_2)$  be two (potentially distinct) probability spaces. Let  $(A_i)_{i \in \mathbb{N}}$  be a sequence of measurable sets in  $\Omega_1$  and  $(B_i)_{i \in \mathbb{N}}$  be a sequence in  $\Omega_2$ . Suppose  $\mu_1(\limsup A_i) = c$  with  $0 \le c \le 1$ . Furthermore suppose the measure of the finite intersections of both sequences are equal. That is to say: for any number of sets  $k \in \mathbb{N}$ and indices  $i_1, i_2, ..., i_k \in \mathbb{N}$  then  $\mu_1(A_{i_1} \cap A_{i_2} \cap ... \cap A_{i_k}) = \mu_2(B_{i_1} \cap B_{i_2} \cap ... \cap B_{i_k})$ . Then these conditions imply  $\mu_2(\limsup B_i) = c$ .

We see that work by Frolov [Fro12] and references therein (for a full history on these results) give improvements on Borel Cantelli's lemma by looking at higher intersections measures, weaker conditions on the sets and tighter bounds. Section 2 lists some of the state of the art. One of the consequences of this paper is that we can not hope to form a precise formula for the limsup measure dependent entirely on the intersection measure quantities. Specifically, for many cases we can have two sequences of sets with intersection measures up to degree k fixed but one has limsup measure zero and the other limsup measure 1.

So what are the repercussions of this? This really demonstrates that in some 'not so nice' cases there is a need to look at fundamental geometrical and topological properties of the sequences to say things about the limsup measure. Beresnevich and Velani do such things by assuming sets are balls in Theorem 2.3 and Theorem 2.4. It may be interesting and fruitful to look further into deriving inequalities and bounds for the limsup measure dependent on some topological properties.

## 2 Previous results

When asking about the measure of the limsup set the first result in this area would be the Borel Cantelli lemmas which later get refined to the Erdos-Chung inequality [CE52].

One of the original Divergence Borel Cantelli forms says that if the sums of measures of the sets diverges  $(\sum \mu(A_i) = \infty)$  and they are all pairwise independent then the limsup has full measure  $(\mu(\limsup A_i)) = 1$ . Furthermore if the sums of measures converge  $(\sum \mu(A_i) < \infty)$  then the convergence Borel Cantelli says  $(\mu(\limsup A_i)) = 0$ .

We now explore some of the more recent results on this matter. There have been a few approaches made to improve the classical Borel Cantelli results. In this paper we concern ourselves almost entirely with the divergence ones despite work by many others to produce stronger forms of the convergence result. We note that many of the papers giving new bounds for the limsup measure also give examples of sequences of sets where the results stated give better bounds for the measure of the limsup than the original Borel Cantelli.

To this end, work was done by Erdos, Renyi and Chung [CE52] and [ER59] give a specific case of the subsequent result and the general case was done by Kochen and Stone [KS64]:

**Theorem 2.1.** Suppose that  $\sum_{i=1}^{\infty} \mu(A_i) = \infty$  then:

$$\mu(\limsup_{n \to \infty} A_n) \ge \limsup_{n \to \infty} \frac{\left(\sum_{s=1}^n \mu(A_s)\right)^2}{\sum_{s,t=1}^n \mu(A_s \cap A_t)} .$$
(3)

It is clear that this is indeed a strengthening of the result as condition (3) will trivially be satisfied by independent sequences, the presentation on the right hand side is referred to as quasi independence on average. However, there are examples where the  $A_i$  has full limsup measure but the right hand side of the inequality is 0 so sometimes this expression tells us no new information.

Feng, Li and Shen provided a weighted version of this result in [FLS09]. That is for any sequence of weightings  $\{w_n\}_{n\in\mathbb{N}}$  with each  $w_n\in\mathbb{R}$  then provided  $\sum_{n=1}^{\infty}w_n\mu(A_n)=\infty$ :

$$\mu(\limsup_{n \to \infty} \mu(A_n)) \ge \limsup_{n \to \infty} \frac{(\sum_{k=1}^n w_k \mu(A_k))^2}{\sum_{i=1}^n \sum_{j=1}^n w_i w_j \mu(A_i \cap A_j)}$$

There have been some other approaches taken to develop new bounds. Petrov [Pet04] took a variational principle approach to finding a bound. This is omitted from here as doesn't add much additional intuition from the results stated. Another approach is taken by noticing we can express the measure of the limsup in terms of the limits of measures of

unions as follows using continuity of measure:

$$\mu(\limsup_{i \to \infty} A_i) = \mu\left(\bigcap_{n \ge 1} \bigcup_{i \ge n} A_i\right) \tag{4}$$

$$=\lim_{n\to\infty}\mu\left(\bigcup_{i\ge n}A_i\right)\tag{5}$$

$$= \lim_{n \to \infty} \lim_{m \to \infty} \mu\left(\bigcup_{i=n}^{m} A_i\right) .$$
(6)

This then means if we can find suitable bounds for the union of sets we have a natural bound for the limsup. This was the approach taken by Frolov in several papers.

We now examine a paper that uses measures of intersections of more sets. The work of [Fro12] does just this and the paper also gives a fairly comprehensive overview of previous results in this area. This is also talked about later when we investigate controlling higher intersections.

**Theorem 2.2.** Suppose  $\sum_{n=1}^{\infty} \mu(A_n) = \infty$ . Set:

$$s_1(n) = \sum_{i=1}^n \mu(A_i), \qquad s_2(n) = 2 \sum_{1 \le i < j \le n} \mu(A_i \cap A_j), \qquad s_3(n) = 6 \sum_{1 \le i < j < k \le n} \mu(A_i \cap A_j \cap A_k) .$$

Then let:

$$\delta_1(n) = (n-1)s_1(n) - s_2(n), \quad \delta_2(n) = (n-2)s_2(n) - s_3(n).$$

Now, assuming  $\delta_1(n)/n \to \infty$  and  $s_2(n) = o(\delta_1(n) + \delta_2(n))$  as  $n \to \infty$ . Then:

$$\mu(\limsup_{n \to \infty} A_n) \ge \limsup_{n \to \infty} \left( \frac{\delta_1(n)^2}{n(\delta_1(n) + \delta_2(n))} + \frac{s_1(n)}{n} \right) .$$

Further papers by Frolov suggested slightly better bounds (but all of a similar flavour) by improving the initial probability of union bound. These are stated in these papers: [Fro14] and [Fro15a]. A further result was stated in [Fro15b] which under some mild conditions:

$$\mu(\limsup_{n \to \infty} A_n) \ge \limsup_{n \to \infty} \left( \frac{1}{n} \sum_{k=1}^n \mu(A_k) + \frac{\left(\sum_{i=1}^n \mu(A_i^c \cap A_k)\right)}{\sum_{i=1}^n \sum_{j=1}^n \mu(A_i^c \cap A_j \cap A_k)} \right) .$$

Really the key takeaway from all of this is that from a naive approach, we wish for  $\mu(A_i)$  to be large,  $\mu(A_i \cap A_j)$  to be small and  $\mu(A_i \cap A_j \cap A_k)$  to be large in order to have larger limsup measure. Of course this doesn't take into account that the quantities are related and have to obey stipulations. Alas, our later results will indeed demonstrate this naive intuition is false.

More recent work by Beresnevich and Velani give some necessary and sufficient conditions regarding the limsup measure this is really stretching the theory to its limits. The following two theorems are the corresponding Theorems 1 and 3 from [BV23]. We state two of the theorems in full as subtle details in the hypotheses are very important. On the face of it the second result may seem to imply our initial motivating problems († and ‡) in the introduction to be true, it's only upon closer inspection that the theory does not in general apply as it assumes the sets are balls.

**Theorem 2.3.** Let  $(\Omega, \mathscr{A}, \mu, d)$  be a metric measure space equipped with a doubling Borel probability measure  $\mu$ . Let  $\{B_i\}_{i\in\mathbb{N}}$  be a sequence of balls in  $\Omega$  with the radius of  $B_i$  tending to 0 as i goes to infinity and such that  $\exists a, b > 0$  such that  $\mu(aB_i) \leq b\mu(B_i)$  holds for all i sufficiently large. Let  $E_{\infty} := \limsup B_i$ . Then

$$\mu(E_{\infty}) = 1$$

if and only if there exists a constant C > 0 such that for any ball B centred in supp $\mu$  there is a sub-sequence  $\{L_{i,B}\}_{i\in\mathbb{N}}$  of  $\{B_i\}_{i\in\mathbb{N}}$  of balls contained in B (i.e.  $L_{i,B} \subset B$  for all i), such that

$$\sum_{i=1}^{\infty} \mu(L_{i,B}) = \infty, \tag{7}$$

and for infinitely many  $Q \in \mathbb{N}$ 

$$\sum_{s,t=1}^{Q} \mu(L_{s,B} \cap L_{t,B}) \le \frac{C}{\mu(B)} \left( \sum_{s=1}^{Q} \mu(L_{s,B}) \right)^2 .$$
(8)

**Theorem 2.4.** Let  $(\Omega, \mathscr{A}, \mu, d)$  be a metric measure space equipped with a doubling Borel probability measure  $\mu$ . Let  $\{B_i\}$  be a sequence of balls such that  $\exists a, b > 0$  such that  $\mu(aB_i) \leq b\mu(B_i)$  holds for all i sufficiently large holds. Let  $E_{\infty} := \limsup_{i \to \infty} B_i$ . Then

$$\mu(E_{\infty}) > 0,$$

if and only if there exists a sub-sequence  $\{L_i\}_{i\in\mathbb{N}}$  of  $\{B_i\}_{i\in\mathbb{N}}$  and a constant C > 0 such that

$$\sum_{i=1}^{\infty} \mu(L_i) = \infty,$$

and for infinitely many  $Q \in \mathbb{N}$ 

$$\sum_{s,t=1}^{Q} \mu(L_s \cap L_t) \le C \left(\sum_{s=1}^{Q} \mu(L_s)\right)^2$$

We will later see that dropping the condition that the  $B_i$  are balls causes the statements to fail. It perhaps can be extended to convex shapes but our results later show not in general.

The essence of these results is to ask how much we can state about the limsup measure from the measure of the intersections. Our later results show here that unfortunately knowledge of the measures of these intersections is in general not always enough to say statements about limsup measures. There is thus something fundamentally geometric about this quantity.

## 3 The case where all finite intersections are fixed

We consider the case of two sequences with identical measured finite intersections between sets. Here we are allowing arbitrarily many finite intersections to be the same as opposed to just pairwise intersections as discussed in the introduction and in the subsequent chapter.

We now state our first main result again.

**Theorem 1.2.** Let  $(\Omega_1, \mathscr{A}_1, \mu_1)$  and  $(\Omega_2, \mathscr{A}_2, \mu_2)$  be two (potentially distinct) probability spaces. Let  $(A_i)_{i \in \mathbb{N}}$  be a sequence of measurable sets in  $\Omega_1$  and  $(B_i)_{i \in \mathbb{N}}$  be a sequence in  $\Omega_2$ . Suppose  $\mu_1(\limsup A_i) = c$  with  $0 \le c \le 1$ . Furthermore suppose the measure of the finite intersections of both sequences are equal. That is to say: for any number of sets  $k \in \mathbb{N}$ and indices  $i_1, i_2, ..., i_k \in \mathbb{N}$  then  $\mu_1(A_{i_1} \cap A_{i_2} \cap ... \cap A_{i_k}) = \mu_2(B_{i_1} \cap B_{i_2} \cap ... \cap B_{i_k})$ . Then these conditions imply  $\mu_2(\limsup B_i) = c$ .

**Remark.** Perhaps the first surprising thing about this result is that it holds for sequences in possibly different measure spaces and relies solely on the sizes of the sets and intersections not the sigma algebra.

This result provides something of an 'upper limit' for the results seen in Section 2. That is, if we have control of all finite intersections measures then we do control the limsup measure. This is in opposition to later results where we control only up to a fixed number of intersections and this doesn't always give control of the limsup measure.

The proof goes through very cleanly and really is just some simple measure theory footwork.

*Proof.* We first note that by the inclusion-exclusion principle for any finite measure  $\mu$  by fixing the finite intersections measures we also fix the finite unions:

$$\mu\left(\bigcup_{i=1}^{n} A_{i}\right) = \sum_{k=1}^{n} (-1)^{k+1} \sum_{1 \le i_{1} < \dots < i_{k} \le n} \mu(A_{i_{1}} \cap \dots \cap A_{i_{k}}) .$$

Hence we have that for all values of k and n:

$$\mu_1\left(\bigcup_{i=k}^n A_i\right) = \mu_2\left(\bigcup_{i=k}^n B_i\right) \ . \tag{9}$$

We now consider the sequence of sets given by incrementing n. That is for  $r \ge k$ :

$$\tilde{A}_{k,r} = \bigcup_{i=k}^{r} A_i, \quad \tilde{B}_{k,r} = \bigcup_{i=k}^{r} B_i.$$

Hence from (9):

$$\mu_1(A_{k,r}) = \mu_2(B_{k,r})$$
.

Now by the containment property we have that:

 $\tilde{A}_{k,r}\subset \tilde{A}_{k,r+1}\subset \tilde{A}_{k,r+2}\subset \tilde{A}_{k,r+3}\subset \dots \ .$ 

The equivalent for  $\tilde{B}$  also works. The containment property means that we can apply the continuity of measure to deduce:

$$\mu_1\left(\bigcup_{i=k}^{\infty} A_i\right) = \lim_{r \to \infty} \mu_1(\tilde{A}_{k,r}) = \lim_{r \to \infty} \mu_2(\tilde{B}_{k,r}) = \mu_2\left(\bigcup_{i=k}^{\infty} B_i\right)$$

Furthermore, we have again by a similar monotonicity argument (as seen in (6)) that:

$$\mu_1(\limsup_{i \to \infty} A_i) = \lim_{k \to \infty} \lim_{r \to \infty} \mu_1(\tilde{A}_{k,r}) = \lim_{k \to \infty} \lim_{r \to \infty} \mu_2(\tilde{B}_{k,r}) = \mu_2(\limsup_{i \to \infty} B_i) .$$

This can also be viewed as a structural result as we can map the two spaces onto each other via a map which preserves the measures of these sets and their intersections. We can actually adapt this proof further by including inequalities to provide a bounding result. The motivation stems from the inclusion-exclusion principle. That is that the odd terms have coefficient positive and even terms coefficient negative. The inequalities developed reflect this. The use of the inclusion-exclusion principle coefficients will be used in later sections also to motivate our problems.

**Theorem 3.1.** Let  $(\Omega_1, \mathscr{A}_1, \mu_1)$  and  $(\Omega_2, \mathscr{A}_2, \mu_2)$  be two (potentially distinct) probability spaces. Let  $(A_i)_{i \in \mathbb{N}}$  be a sequence of measurable sets in  $\Omega_1$  and  $(B_i)_{i \in \mathbb{N}}$  be a sequence in  $\Omega_2$ . Suppose for any number of sets  $k \in \mathbb{N}$  and indices  $i_1, i_2, ..., i_k \in \mathbb{N}$  then  $\mu_1(A_{i_1} \cap A_{i_2} \cap ... \cap A_{i_k}) \ge \mu_2(B_{i_1} \cap B_{i_2} \cap ... \cap B_{i_k})$  for k odd and  $\mu_1(A_{i_1} \cap A_{i_2} \cap ... \cap A_{i_k}) \le \mu_2(B_{i_1} \cap B_{i_2} \cap ... \cap B_{i_k})$ for k even. Then these conditions imply  $\mu_1(\limsup A_i) \ge \mu_2(\limsup B_i)$  also.

*Proof.* Same as the previous but with equality signs replaced with appropriate inequalities.  $\Box$ 

# 4 Showing a disparity in limsup set measures can be achieved

Here we answer the first question (†) as proposed in the introduction. Throughout this section we list explicit examples showing a disparity in limsup sets.

As discussed in the previous chapter, in order to have any hope of a disparity between sizes of limsups sets of two sequences with equal corresponding measures and pairwise intersections, we must alter the measures of the higher order intersections in our sequence. It is clear why therefore we use Venn diagrams as a visual representation of intersections and measures of these. We also demonstrate an alternate representation of these in the ambient space of [0, 1] endowed with the Lebesgue measure case. **Proposition 4.1.** There exist two sequences of sets  $(A_i)_{i \in \mathbb{N}}$  and  $(B_i)_{i \in \mathbb{N}}$  satisfying:

$$\mu(A_i) \ge \mu(B_i) \quad \forall i \in \mathbb{N},$$
  
$$\mu(A_i \cap A_j) \le \mu(B_i \cap B_j) \quad \forall i \neq j \in \mathbb{N},$$

and also  $\mu(\limsup A_i) < \mu(\limsup B_i)$ .

**Remark.** The idea here is that we construct two Venn diagrams with the properties that the measure of each set involved and pairwise intersections are the same but they have fundamentally different total measures. That is, in one we concentrate lots of the mass into the pairwise intersections and nowhere else and in the other we concentrate the mass on the triple intersection and the individual sets (disjoint from the other sets). Thus, the total measure inside the Venn diagrams differ. We simply construct the sequence by repeating these 3 sets over and over. By fixing the measures of the sets and the pairwise intersections of the 3 sets in the Venn diagram, we ensure these for the sequences formed also.

It's worth noting the inequalities in the conditions can all be made strict with minor tweaks and the numeric values of the limsup measure aren't special, again with tweaks these can take other values in a neighbourhood.

We go about constructing our sequences in two separate lemmas:

**Lemma 4.1.** There exists a sequence of sets  $(A_i)_{i \in \mathbb{N}}$  such that:  $\mu(A_i) = 1/2$ ,

$$\mu(A_i \cap A_j) = \begin{cases} \frac{1}{2} & i \equiv j \mod 3\\ \frac{1}{4} & else \end{cases}$$

and also  $\mu(\limsup(A_i)) = 3/4$ .

*Proof.* We refer to the sets A, B and C from Figure 1 and subsequently construct our sequence of  $(A_i)_{i \in \mathbb{N}}$  as follows:

$$A_i = \begin{cases} A & i \equiv 1 \mod 3\\ B & i \equiv 2 \mod 3\\ C & i \equiv 3 \mod 3 \end{cases}$$

That is that our sequence  $A_1$ ,  $A_2$ ,  $A_3$ ,  $A_4$ ,  $A_5$ ,  $A_6$ ,  $A_7$ ,  $A_8$ ,  $A_9$ ,  $A_{10}$ ,  $A_{11}$ ,  $A_{12}$ ... looks like: A, B, C, A, B, C, A, B, C, A, B, C, ... This ensures that each of A, B, C occurs infinitely often and indeed nothing else occurs infinitely often. Thus  $\mu(\limsup(A_i)) = 3 \times \frac{1}{4} = \frac{3}{4}$ . The intersection properties are very clear from the Figure.

**Lemma 4.2.** There exists a sequence of sets  $(B_i)_{i \in \mathbb{N}}$  such that:  $\mu(B_i) = 1/2$ ,

$$\mu(B_i \cap B_j) = \begin{cases} \frac{1}{2} & i \equiv j \mod 3\\ \frac{1}{4} & else \end{cases}$$

and also  $\mu(\limsup(B_i)) = 1$ .




Figure 1: Venn diagram construction of sets A, B and C

*Proof.* Similar to before refer to sets D, E and F from Figure 2 we construct our sequence of  $(B_i)_{i \in \mathbb{N}}$  as follows:

$$B_i = \begin{cases} D & i \equiv 1 \mod 4\\ E & i \equiv 2 \mod 4\\ F & i \equiv 3 \mod 4 \end{cases}$$

That is that our sequence  $B_1, B_2, B_3, B_4, B_5, B_6, B_7, B_8, B_9, B_{10}, B_{11}, B_{12}...$  looks like: D, E, F, D, E, F, D, E, F, D, E, F, ... This ensures that each of D, E, F occurs infinitely often and indeed nothing else occurs infinitely often. Thus  $\mu(\limsup(B_i)) = 3 \times \frac{1}{4} + \frac{1}{4} = \frac{4}{4} = 1$ . The intersection properties are very clear from the Figure.

*Proof.* (Proposition 4.1) Using the sequences formed in the previous two lemmas for  $(A_i)_{i \in \mathbb{N}}$  and  $(B_i)_{i \in \mathbb{N}}$  satisfy the inequalities and achieve the disparity in measures of the limsup.  $\Box$ 

We further see the explicit examples of these sets in the Lebesgue measure on [0, 1] case to give an alternate perspective and visualisation of these cases.

#### 4.0.1 Lebesgue measure example of the 3 sets

We now show these examples but expressed in the Lebesgue measure setting. We note we only show the first 3 sets of each sequence as we just repeat these in our construction.






Figure 2: Venn diagram construction of sets D, E and F





## 4.1 The case of 4 sets

We now adapt these ideas for the case of 4 sets as opposed to 3 with an effort to increase the disparity in the measure of the limsup sets. Much of the ideas will feel very familiar so we state the results more briefly.

**Proposition 4.2.** There exists two sequences of sets  $(A_i)_{i \in \mathbb{N}}$  and  $(B_i)_{i \in \mathbb{N}}$  satisfying:

$$\mu(A_i) \ge \mu(B_i) \quad \forall i \in \mathbb{N},$$
  
$$\mu(A_i \cap A_j) \le \mu(B_i \cap B_j) \quad \forall i, j \in \mathbb{N},$$

and also  $\mu(\limsup A_i) = \frac{6}{9}$  and  $\mu(\limsup B_i) = 1$ .





Figure 4: Lebesgue construction of  $(B_i)_{i \in \mathbb{N}}$ 

**Lemma 4.3.** There exists a sequence of sets  $(A_i)_{i \in \mathbb{N}}$  such that:  $\mu(A_i) = 1/3$ ,

$$\mu(A_i \cap A_j) = \begin{cases} \frac{1}{3} & i \equiv j \mod 4\\ \frac{1}{9} & else \end{cases}$$

and also  $\mu(\limsup(A_i)) = 2/3$ .

Figure 5: Venn diagram construction of sets A, B, C and D



*Proof.* Referring to the sets in Figure 5, we subsequently construct our sequence of  $(A_i)_{i\in\mathbb{N}}$ 

as follows:


$$A_i = \begin{cases} A & i \equiv 1 \mod 4 \\ B & i \equiv 2 \mod 4 \\ C & i \equiv 3 \mod 4 \\ D & i \equiv 4 \mod 4 \end{cases}$$


That is that our sequence  $A_1$ ,  $A_2$ ,  $A_3$ ,  $A_4$ ,  $A_5$ ,  $A_6$ ,  $A_7$ ,  $A_8$ ,  $A_9$ ,  $A_{10}$ ,  $A_{11}$ ,  $A_{12}$ ... looks like:  $A, B, C, D, A, B, C, D, A, B, C, D, \dots$ . This ensures that each of A, B, C, D occurs infinitely often and indeed nothing else occurs infinitely often. Thus  $\mu(\limsup(A_i)) = 6 \times \frac{1}{9} = \frac{6}{9}$ . It is trivial from the figure that the intersections hold.

**Lemma 4.4.** There exists a sequence of sets  $(B_i)_{i \in \mathbb{N}}$  such that:  $\mu(A_i) = 1/3$ ,

$$\mu(B_i \cap B_j) = \begin{cases} \frac{1}{3} & i \equiv j \mod 4\\ \frac{1}{9} & else \end{cases}$$

and also  $\mu(\limsup(B_i)) = 1$ .





*Proof.* Referring to the sets in Figure 6, we subsequently construct our sequence of  $(B_i)_{i \in \mathbb{N}}$  as follows:

$$B_i = \begin{cases} E & i \equiv 1 \mod 4\\ F & i \equiv 2 \mod 4\\ G & i \equiv 3 \mod 4\\ H & i \equiv 4 \mod 4 \end{cases}$$

That is that our sequence  $B_1$ ,  $B_2$ ,  $B_3$ ,  $B_4$ ,  $B_5$ ,  $B_6$ ,  $B_7$ ,  $B_8$ ,  $B_9$ ,  $B_{10}$ ,  $B_{11}$ ,  $B_{12}$ ... looks like:  $E, F, G, H, E, F, G, H, E, F, G, H, \ldots$ . This ensures that each of E, F, G, H occurs infinitely often and indeed nothing else occurs infinitely often. Thus  $\mu(\limsup(B_i)) = 4 \times \frac{2}{9} + \frac{1}{9} = \frac{9}{9} = 1$ 

*Proof.* (Proposition 4.2) Again the sequences formed in these lemmas do the trick.  $\Box$ 

Similar to before we give the analogous sets in the the Lebesgue measure setting.

#### 4.1.1 Lebesgue measure example of 4 sets

We now want to see how far this idea gets us. We got a greater disparity for the case of 4 sets as opposed to the case of 3 sets, thus we have an expectation that we get a larger difference in the measures of the limsup sets if we have more sets repeated. To this end;







Figure 8: Lebesgue construction of  $(B_i)_{i \in \mathbb{N}}$ 



## 4.2 Arbitrarily many sets repeated

Here we have to leave the pretty world of Venn diagrams behind as the case of 4 Venn diagrams is really the last Venn pictorially feasible as the diagrams become too complicated and elaborate to really gauge whats going on. Alas, I believe the prior two examples give a good enough sense of the situation at hand for us to no longer necessitate a diagram.

**Theorem 4.1.** For any  $\epsilon > 0$  there exists two sequences of sets  $(C_i)_{i \in \mathbb{N}}$  and  $(D_i)_{i \in \mathbb{N}}$  satisfying:

$$\mu(C_i) \ge \mu(D_i) \quad \forall i \in \mathbb{N}$$
$$\mu(C_i \cap C_j) \le \mu(D_i \cap D_j) \quad \forall i, j \in \mathbb{N}$$

and also  $\mu(\limsup C_i) \leq \frac{1}{2} + \epsilon$  and  $\mu(\limsup D_i) = 1$ .

**Remark.** For the proof we start with two setups of a finite collection of sets. One, whose measure is concentrated in the pairwise intersections alone and the other whose measure is concentrated on the intersection of all the sets as well as the areas in only one set (\*i.e.  $A \cap B^c \cap C^c \cap D^c...$ ). Then we construct our sequences by cycling through these (as we did in the previous examples) and find the measure of the limsup accordingly.

*Proof.* To this end we proceed. Again we consider  $C_1, C_2, ..., C_n$  and  $D_1, D_2, ..., D_n$  to be two systems of sets whose measures we will prescribe as follows:

We build each sequence out of a collection of disjoint sets. For the construction of  $C_i$  we will concentrate all of the measure onto the pairwise intersections and nowhere else. Throughout 'a' will be a normalisation factor. We first define  $C_{k,l}$  for  $1 \le k < l \le n$ . These will be disjoint sets each with  $\mu(C_{k,l}) = a$  We then define  $C_i$  to be the collection of  $C_{k,l}$  such that one of the subscripts is equal to i. Explicitly:

$$C_i = \left(\bigcup_{i < l \le n} C_{i,l}\right) \cup \left(\bigcup_{1 \le k < i} C_{k,i}\right) .$$

Hence we attain the  $C_i$  have the following measure properties:

$$\mu(C_i) = (n-1)a \qquad \forall i : 1 \le i \le n,$$
  
$$\mu(C_i \cap C_j) = \mu(C_{i,j}) = a \qquad \forall i, j : 1 \le i < j \le n,$$
  
$$\mu(C_i \cap C_j \cap C_k) = 0 \qquad \forall i, j, k : i < j < k.$$

Trivially as we repeatedly cycle through the sets  $C_i$ , each of the  $C_{k,l}$ , that it is composed of is hit infinitely often. Thus the total limsup measure will be the sum of the measures of these  $C_{k,l}$ . Therefore the limsup measure will be:

$$\mu(\limsup C_i) = \binom{n}{2}a = \frac{n(n-1)}{2}a .$$

We construct the  $D_i$  as follows. Now, we define n + 1 disjoint sets with suitable measures. We let  $\tilde{D}_i$  for  $1 \le i \le n$  have measure (n-2)a each. Also  $\tilde{D}_{n+1}$  with measure a. Then we set:

$$D_i = \tilde{D}_i \cup \tilde{D}_{n+1}$$

The subsequent measure theoretic properties are:

$$\mu(D_i) = (n-1)a \qquad \forall i : 1 \le i \le n,$$
  
$$\mu(D_i \cap D_j) = \mu(\tilde{D}_{n+1}) = a \qquad \forall i, j : 1 \le i \le j \le n,$$
  
$$\mu(D_1 \cap D_2 \cap \dots \cap D_n)) = \mu(\tilde{D}_{n+1}) = a.$$

Here the sets all pairwise intersect on the same region, namely  $\tilde{D}_{n+1}$ , that is there is one region in which they all overlap on and pairwise overlap nowhere else. As before we set the sequences  $(C_i)_{i\in\mathbb{N}}$  and  $(D_i)_{i\in\mathbb{N}}$  to be the cycled versions of these. This means the total limsup will be as follows:

$$\mu(\limsup D_i) = n \times (n-2)a + a = (n^2 - 2n + 1)a = (n-1)^2a.$$

We see that as n tends to  $\infty$  the ratio of these tends to  $\frac{1}{2}$ . Lastly the inequalities stated in the hypothesis are satisfied trivially by their construction.

## 5 Can we improve the disparity in limsup measure?

We've seen cases where the two limsup sets of sequences may differ by roughly a factor of a half, thereby answering the first question (†) in the introduction. Can we do better and solve (‡)?

That is to say can we have two sequences of sets with the usual relation comparing their measures of sets and pairwise intersections such that one has a limsup whose measure is 0 and the other has a measure 1 limsup? This is a genuinely non trivial result as Theorem 2.4 stated earlier implies this to be false assuming the sets are balls.

Well indeed so. Naturally the construction is slightly more involved than the previous. Perhaps something to motivate our next steps is to draw inspiration from the prior examples. That is, we want one of the sequences to be formed from some sets whose intersections concentrate on just one area and then another sequence whose collective intersections are very small. The problem thus is to find two sequences satisfying:

We give examples of these. One in  $\mathbb{R}^2$  and another in  $\mathbb{R}$ .

## 5.1 Example in $\mathbb{R}^2$

Throughout this subsection we are looking at the unit square  $([0, 1]^2)$  endowed with the usual Borel sigma algebra and Lebesgue measure.

**Theorem 5.1.** There exists sequences  $(A_i)_{i \in \mathbb{N}}$  and  $(B_i)_{i \in \mathbb{N}}$  in the measure space described above satisfying the following properties:

$$\mu(A_i) \ge \mu(B_i) \quad \forall i \in \mathbb{N},$$
  

$$\mu(A_i \cap A_j) \le \mu(B_i \cap B_j) \quad \forall i < j \in \mathbb{N},$$
  

$$\mu(\limsup A_i) = 0,$$
  

$$\mu(\limsup B_i) = 1.$$
(10)

**Remark.** We construct each sequence separately in the subsequent two lemmas. We give bounds for the measures at points, as opposed to the explicit values they take as after all, we are only interested in how these measures relate to each other.

Again, with very subtle tweaking the inequalities can be made strict.

**Lemma 5.1.** There exists a sequence  $(A_i)_{i \in \mathbb{N}}$  with the following properties:

For 
$$n \in \mathbb{N}$$
 where  $n = 3(i-1) + k$  with  $k \in \{1, 2, 3\}$ :  $\mu(A_n) = \frac{1}{3i}$ ,  
For  $n, m \in \mathbb{N}$  distinct with  $n = 3(i-1) + k, m = 3(j-1) + l$ :  $\mu(A_n \cap A_m) \leq \frac{1}{9 \max\{i, j\}}, \mu(\limsup A_i) = 0$ .

**Remark.** The idea will be to have really uniform measure intersections between these sets, a natural way to do this is as follows: We repeatedly divide sections by 3 and choose our sets iteratively from these divisions to intersect evenly. In essence at every stage we divide all of our prior intervals into 3 and then pick our subsequent 3 sets to take 1 distinct part of each division.

*Proof.* Let's begin by constructing  $(A_i)$  as follows inductively:

$$A_1 = \left[0, \frac{1}{3}\right) \times [0, 1], \quad A_2 = \left[\frac{1}{3}, \frac{2}{3}\right) \times [0, 1], \quad A_3 = \left[\frac{2}{3}, 1\right] \times [0, 1].$$


As seen in Figure 10, we then divide each of the strips equally into 3 vertical strips. We then shrink these down vertically to a height of  $\frac{1}{2}$ . We select  $A_4$  to be the first strip and every third strip after- this is the dark grey region selected in Figure 10. We select  $A_5$  to be the second strip and then every third strip after. We select  $A_6$  to be the third strip and then every third strip after.

Explicitly written:

$$A_{4} = \begin{bmatrix} 0, \frac{1}{9} \end{bmatrix} \times \begin{bmatrix} 0, \frac{1}{2} \end{bmatrix} \cup \begin{bmatrix} \frac{3}{9}, \frac{4}{9} \end{bmatrix} \times \begin{bmatrix} 0, \frac{1}{2} \end{bmatrix} \cup \begin{bmatrix} \frac{6}{9}, \frac{7}{9} \end{bmatrix} \times \begin{bmatrix} 0, \frac{1}{2} \end{bmatrix},$$
  

$$A_{5} = \begin{bmatrix} \frac{1}{9}, \frac{2}{9} \end{bmatrix} \times \begin{bmatrix} 0, \frac{1}{2} \end{bmatrix} \cup \begin{bmatrix} \frac{4}{9}, \frac{5}{9} \end{bmatrix} \times \begin{bmatrix} 0, \frac{1}{2} \end{bmatrix} \cup \begin{bmatrix} \frac{7}{9}, \frac{8}{9} \end{bmatrix} \times \begin{bmatrix} 0, \frac{1}{2} \end{bmatrix},$$
  


$$A_{6} = \begin{bmatrix} \frac{2}{9}, \frac{3}{9} \end{bmatrix} \times \begin{bmatrix} 0, \frac{1}{2} \end{bmatrix} \cup \begin{bmatrix} \frac{5}{9}, \frac{6}{9} \end{bmatrix} \times \begin{bmatrix} 0, \frac{1}{2} \end{bmatrix} \cup \begin{bmatrix} \frac{8}{9}, 1 \end{bmatrix} \times \begin{bmatrix} 0, \frac{1}{2} \end{bmatrix}.$$




We repeat this idea for subsequent sets, we split each of the strips into three vertical strips. We compress them vertically so they now have a height of  $\frac{1}{3}$ . We pick  $A_7$  to be the first strip followed by every third one thereafter-this is shown by Figure 11. We select  $A_8$  to be the second strip followed by after third thereafter. We select  $A_9$  to be the third strip followed by every third thereafter.

It is clear that inductively we construct the sets in blocks of 3 so that at stage 'i' we take our strips from the 'i - 1'th stage split them into 3 vertical strips. We then shrink them so that they have height  $\frac{1}{i}$ . We select  $A_{3(i-1)+1}$  to be the first strip followed by every third one thereafter. We pick  $A_{3(i-1)+2}$  to be the second strip followed by every third one thereafter. We pick  $A_{3(i-1)+3}$  to be the third and then every third thereafter.

Now we proceed to examine the properties of these. Again, it will be easiest to consider





this in terms of blocks:

$$\underbrace{A_1, A_2, A_3}_{\text{1st block}}, \underbrace{A_4, A_5, A_6}_{\text{2nd block}}, \underbrace{A_7, A_2, A_3}_{\text{3rd block}}, \underbrace{A_{10}, A_{11}, A_{12}}_{\text{4th block}}, \dots$$

Then any set A in say the *i*th block will have measure:  $\mu(A) = \frac{1}{3i}$ .

Now we consider the measures of the intersections. For A,B distinct sets but in the same block its clear by construction that they have null intersection and thus  $\mu(A \cap B) = 0$ . For A in block i and B in block j (with i, j distinct) it is clear from the construction that  $\mu(A \cap B) = \frac{1}{9 \max\{i,j\}}$ .

It is clear by inspection that the  $\limsup(A_i) = 0$  as the maximum height of the sets tends to 0:

$$\mu(\limsup(A_i)) = \mu\left(\bigcap_{t=1}^{\infty}\bigcup_{s=t}^{\infty}A_s\right)$$
$$= \mu\left(\bigcap_{t=1}^{\infty}\bigcup_{s=3t}^{\infty}A_s\right)$$
$$= \mu\left(\bigcap_{t=1}^{\infty}\bigcup_{s=t}^{\infty}[0,1]\times\left[0,\frac{1}{t}\right]\right)$$
$$= \mu\left(\bigcap_{t=1}^{\infty}[0,1]\times\left[0,\frac{1}{t}\right]\right)$$
$$= \mu\left([0,1]\times[0,0]\right)$$
$$= 0.$$

We now turn our attention to constructing the  $(B_i)_{i \in \mathbb{N}}$ . In contrast to the  $(A_i)_{i \in \mathbb{N}}$  we concentrate all the pairwise intersections in one region which will shrink over time.

**Lemma 5.2.** There exists a sequence  $(B_i)_{i \in \mathbb{N}}$  with the following properties:

For 
$$n \in \mathbb{N}$$
 where  $n = 3(i-1) + k$  with  $k \in \{1, 2, 3\}$ :  $\mu(B_n) = \frac{1}{3i}$ ,  
For  $n, m \in \mathbb{N}$  distinct with  $n = 3(i-1) + k, m = 3(j-1) + l$ :  $\mu(B_n \cap B_m) \ge \frac{1}{9 \max\{i, j\}},$   
 $\mu(\limsup B_i) = 1$ .

**Remark.** To reiterate once more, for the construction of the  $B_i$  we simply build them to have measures comparable to that of the  $A_i$ , in order to satisfy the inequalities stated before. This avoids the 'messiness' of using the measure quantities as stated in the statement of the lemma.

*Proof.* Each set  $(B_i)$  is composed of two (or three) rectangles. One 'vertical' rectangle of height 1, aligned on the left hand side with bottom left point at (0,0) and top right at  $(\frac{\mu(A_i)}{3}, 1)$ . The next rectangle(s) are the 'horizontal' ones. They touch the right hand most line of the vertical rectangle and the right hand side of the square. These 'move up' with i and once it reaches the top of the square it starts back at the bottom again. More specifically the first 'horizontal' rectangle is aligned with the x axis and then the next rectangle in the next set lies atop of the previous and so on. Once the rectangles exceed y = 1 we cut them off here and place the rest at the bottom-that is the bottom aligns with y = 0. This is demonstrated in Figure 12.

By construction, the area of the left hand, vertical rectangle is  $\frac{\mu(A_i)}{3}$ .


The total area of the horizontal rectangle(s) is to be  $\frac{2\mu(A_i)}{3}$ . Thus the total measure of  $B_i$  is the same as  $A_i$ . So the height of each horizontal rectangle is  $\frac{2\mu(A_i)}{3-\mu(A_i)}$  with width  $1 - \frac{\mu(A_i)}{3}$ . Therefore the measures of the 'horizontal' rectangle will be  $\frac{2\mu(A_i)}{3-\mu(A_i)} \times (1 - \frac{\mu(A_i)}{3}) = \frac{2\mu(A_i)}{3}$  as desired. As we see below, the sum of the heights diverges so it is clear that this cycles through from bottom to top infinitely often and thus the limsup will have measure 1.


$$\begin{array}{l} \text{Height of } A_i = \frac{2\mu(A_i)}{3 - \mu(A_i)}, \\ \text{sum of these heights} = \sum_{i=1}^{\infty} \frac{2\mu(A_i)}{3 - \mu(A_i)} \leq \sum_{i=1}^{\infty} \frac{2\mu(A_i)}{3} = \frac{2}{3} \sum_{i=1}^{\infty} \mu(A_i) = \infty \ . \end{array}$$

Therefore it diverges and the rectangles cycle through infinitely often.

It is trivial that  $\forall i : \mu(A_i) = \mu(B_i)$  simply by construction. Also for i < j:

$$\mu(B_i \cap B_j) \ge \mu\left(\left[0, \frac{\mu(A_i)}{3}\right] \times [0, 1] \cap \left[0, \frac{\mu(A_j)}{3}\right] \times [0, 1]\right)$$





$$= \mu\left(\left[0, \frac{\mu(A_j)}{3}\right] \times [0, 1]\right) = \frac{\mu(A_j)}{3} \ge \mu(A_i \cap A_j) \ .$$

*Proof.* (Theorem 4.2) We have thus exhibited examples of sequences of sets satisfying (10).  $\Box$ 

We now move onto the general dimension case. We exhibit another examples of sets now in 1D, these generalise to arbitrary dimensions nicely. They will also be key inspiration for sets we construct for the general intersection case.

#### 5.2 Examples in $\mathbb{R}$

Now we construct a new pair of sequences of sets in  $\mathbb{R}$ -specifically [0, 1]. Again we look at the Borel sigma algebra on this and Lebesgue measure. Now we shall have  $(C_i)_{i \in \mathbb{N}}$  in place of  $(A_i)_{i \in \mathbb{N}}$  and  $(D_i)_{i \in \mathbb{N}}$  in place of  $(B_i)_{i \in \mathbb{N}}$  in (10).

**Theorem 5.2.** There exists sequences  $(C_i)_{i \in \mathbb{N}}$  and  $(D_i)_{i \in \mathbb{N}}$  in the measure space described above satisfying:

$$\mu(C_i) \ge \mu(D_i) \quad \forall i \in \mathbb{N},$$
  

$$\mu(C_i \cap C_j) \le \mu(D_i \cap D_j) \quad \forall i < j \in \mathbb{N},$$
  

$$\mu(\limsup C_i) = 0,$$
  

$$\mu(\limsup D_i) = 1.$$
(11)

**Remark.** The standout from the proofs of this theorem and the prior was that in order to gain a disparity in limsup we need in one case to have all of our sets overlapping on a region with mass comparable to the mass of the set and in another case to have all of our measure spread out and well distributed.

It is worth noting here that the inequalities we demand for the measures of  $C_n$  and  $D_n$  can be made strict by making the  $D_n^{(1)}$  slightly larger by say a factor of  $\frac{5}{4}$  and shrinking the widths of the  $D_n^{(2)}$  in the construction.

It is further worth noting that this is not specific to the 1 dimensional Lebesgue measure, there exist models in higher dimensions where this phenomenon also occurs. This can easily be realised by placing ' $\times [0,1]^{n-1}$ ' at the end of the sets and we have ourselves an example in  $\mathbb{R}^n$ .

The constructions will have very similar flavours to the previous in how we construct our examples inductively and concentrate measures

**Lemma 5.3.** There exists a sequence  $(C_i)_{i \in \mathbb{N}}$  with the following properties:

$$\forall n \in \mathbb{N} \quad \mu(C_{2n-1}) = \mu(C_{2n}) = \frac{1}{n+2},$$
  
$$\forall i < j \in \mathbb{N} \quad \mu(C_i \cap C_j) = \begin{cases} 0 & j-i = 1 \text{ and } i \text{ is even} \\ \frac{1}{2}\mu(C_j) & else \end{cases}$$
  
$$\mu(\limsup(C_i)) = 0.$$

*Proof.* We inductively construct the sets in pairs. At each step we will construct the subsequent pair of sets out of the previous pair. Furthermore we will see at each step that  $C_{2n-1}$  and  $C_{2n}$  are disjoint and they lie inside the union of the previous two sets. To be more explicit,  $C_{2n-1}$  and  $C_{2n}$  will be made of  $2^{n-1}$  disjoint sub-intervals of length  $\frac{1}{2^{n-1}(n+2)}$ . They will be equally spread amongst the previous two sets, so that half of the measure of  $C_{2n-1}$  will be contained in  $C_{2n-2}$  and the other half in  $C_{2n-3}$  and the same for  $C_{2n}$ .

We begin the inductive process. At each step we construct a pair of sets from the prior pair such that at step n each set is built from  $2^{n-1}$  disjoint intervals of length  $\frac{1}{2^{n-1}(n+2)}$  hence  $2^n$  endpoints.

First we set  $C_1 = [0, 1/3]$  and  $C_2 = [2/3, 1]$ . Trivially this meets the criterion of being built from  $2^0 = 1$  disjoint intervals. They are also of length  $\frac{1}{3} = \frac{1}{2^0(1+2)}$ .

Continuing the induction process, suppose we have constructed  $C_{2(n-1)-1}$  and  $C_{2(n-1)}$ (with the desired number and lengths of disjoint intervals) then inductively we construct  $C_{2n-1}$  and  $C_{2n}$ . First denote  $\{a_{k,n-1}\}_{1 \le k \le 2^{n-1}}$  to be the left endpoints of all the intervals of  $C_{2(n-1)-1}$  and  $C_{2(n-1)}$ . Similarly, denote  $\{b_{k,n-1}\}_{1 \le k \le 2^{n-1}}$  to be the right endpoints of all the intervals of all the intervals of  $C_{2(n-1)-1}$  and  $C_{2(n-1)-1}$ . We then construct  $C_{2n-1}$  and  $C_{2n}$  as follows:

$$C_{2n-1} = \bigcup_{k=1}^{2^{n-1}} [a_{k,n-1}, a_{k,n-1} + \frac{1}{2^{n-1}(n+2)}],$$
$$C_{2n} = \bigcup_{k=1}^{2^{n-1}} [b_k - \frac{1}{2^{n-1}(n+2)}, b_k].$$


We can see here that the interval lengths are as desired and as  $b_{k,n-1} - a_{k,n-1} = \frac{1}{2^{n-2}(n+1)}$ then  $C_{2n-1}$  and  $C_{2n}$  are indeed disjoint and completely lie within the prior two sets.

A visual aid of this is given in Figure 13. Lets examine the measure properties of these from the fact the intervals are disjoint and additivity of measures:

$$\mu(C_{2n-1}) = \mu(C_{2n}) = 2^{n-1} \times \frac{1}{2^{n-1}(n+2)} = \frac{1}{n+2},$$

For i < j:

$$\mu(C_i \cap C_j) = \begin{cases} 0 & j-i = 1 \text{ and } i \text{ is even} \\ \frac{1}{2}\mu(C_j) & \text{else} \end{cases}$$



Trivially from the nesting of the  $C_i$  and their diminishing measures, the limsup's measure is 0. This is really the the same reasoning as in the previous subsection, using containment properties. It will be important to note furthermore the sum of the measures of  $C_i$  diverges for the next part.

**Lemma 5.4.** There exists a sequence  $(D_i)_{i \in \mathbb{N}}$  with the following properties:

$$\begin{aligned} \forall n \in \mathbb{N} \quad \mu(D_{2n-1}) &= \mu(D_{2n}) = \frac{1}{n+2} \\ \forall i < j \in \mathbb{N} \quad \mu(D_i \cap D_j) \geq \begin{cases} 0 & j-i = 1 \text{ and } i \text{ is even} \\ \frac{1}{2}\mu(D_j) & else \end{cases} \\ \mu(\limsup(D_i)) &= 1 \end{aligned}$$

**Remark.** Again, we allow ourselves to be a bit less precise in this lemma as after all we are only concerned with how the measures compare for those of the  $C_i$ .

*Proof.* We now go about the construction of the  $D_i$ , this will be in a similar vein to previous chapters in how we concentrate much of the measure in one region and have a 'floating' small interval that moves along to achieve full limsup measure. To this end, we go about the construction, again by an inductive construction:

We begin with:

$$D_1 = \left[0, \frac{\mu(C_1)}{2}\right] \cup \left[\frac{\mu(C_1)}{2}, \mu(C_1)\right] = \left[0, \frac{1}{3 \times 2}\right] \cup \left[\frac{1}{3 \times 2}, \frac{1}{3}\right] = \left[0, \frac{1}{6}\right] \cup \left[\frac{1}{6}, \frac{1}{3}\right]$$

Continuing with this, we suppose that we have constructed  $D_{n-1}$  and now go about constructing  $D_n$ . Again we want the interval fixed on the left. This will be:

$$D_n^{(1)} = \left[0, \frac{\mu(C_n)}{2}\right]$$

Now let  $c_{n-1}$  be the rightmost point of  $D_{n-1}$ . We now have two options to add in the next part:

$$D_n^{(2)} = \begin{cases} \left[\frac{\mu(C_n)}{2}, \mu(C_n) - 1 + c_{n-1}\right] \cup \left[c_{n-1}, 1\right] & c_{n-1} + \frac{\mu(C_n)}{2} > 1\\ \left[c_{n-1}, c_{n-1} + \frac{\mu(C_n)}{2}\right] & \text{else} \end{cases}$$

The reason for this is simply so the moving part 'loops back' and keeps re-covering the unit interval. Ultimately we define:

$$D_n = D_n^{(1)} \cup D_n^{(2)}$$

Now examining the measure properties of this: By construction:

$$\mu(D_n) = \mu(C_n)$$

For i < j:

$$\mu(D_i \cap D_j) \ge \mu(D_i^{(1)} \cap D_j^{(1)}) = \mu\left(\left[0, \frac{\mu(C_i)}{2}\right] \cap \left[0, \frac{\mu(C_j)}{2}\right]\right)$$
$$= \mu\left(\left[0, \frac{\mu(C_j)}{2}\right]\right) = \frac{\mu(C_j)}{2} = \mu(C_i \cap C_j)$$

As we will see, the measures of the floating part diverge, ensuring they keep 'looping round' so trivially the limsup has full measure:

$$\sum_{n=1}^{\infty} \mu(D_n^{(2)}) = \sum_{n=1}^{\infty} \frac{\mu(C_n)}{2} = \sum_{n=1}^{\infty} \frac{1}{n+2} = \infty$$

*Proof.* (Theorem 5.2) We have thus exhibited examples of sequences of sets satisfying (11).  $\Box$ 

## 6 Developing the theory to deal with more intersections

We've seen we can have two sequences  $(A_i)_{i \in \mathbb{N}}$  and  $(B_i)_{i \in \mathbb{N}}$  satisfying the following:

$$\mu(A_i) \ge \mu(B_i) \quad \forall i \in \mathbb{N}$$
  
$$\mu(A_i \cap A_j) \le \mu(B_i \cap B_j) \quad \forall i < j \in \mathbb{N}$$
(12)

Such that  $\mu(\limsup A_i) = 0$  and  $\mu(\limsup B_i) = 1$ 

It is therefore only natural to see if we control the triple intersections, or indeed arbitrarily many intersections up to a fixed finite amount can we get the same '0-1' relation of the limsup measures for this. Alternatively we may have a situation where by controlling more it means the structure is more rigid, similar to that of Theorem 2.4 where if we use balls, the 0-1 relation fails. We saw before from Theorem 1.2 that if we control all the finite intersections measures then we do fix the limsup measure. This result suggests that it may be less obvious that a 0-1 relation on the limsups may exists.

The next question is how do we want the inequality to go for higher intersections to make this question substantial. Well lets appeal to existing intuition we have. In previous sections we saw from Theorem 2.1 (by Kochen and Stone) that on the face of it  $\mu(A_i)$  being larger should mean the limsup should have larger measure and similarly the quantities  $\mu(A_i \cap A_j)$ being small should increase the limsup measure. Of course, the examples shown previously saw this is not the case always.

I feel the intuition is less immediately obvious for triple intersections, we appeal to the examples, current state of the art and the inclusion exclusion principle. Unfortunately, the basic form of the Divergence Borel Cantelli and Kochen Stone don't do this. However, we saw from Section 2 that Frolov has done work on this, specifically Theorem 2.2. This result suggested that we would want the measures of the triple intersections to be larger.

Reinforcing this further we recall from (4) seen earlier, again in Section 2 that on the face of it, in order to get bigger limsup measure we would want the finite union bigger as seen before.

Whence, we appeal to the inclusion exclusion principle:

$$\mu\left(\bigcup_{i=1}^{n} A_{i}\right) = \sum_{k=1}^{n} (-1)^{k+1} \sum_{1 \le i_{1} < \dots < i_{k} \le n} \mu(A_{i_{1}} \cap \dots \cap A_{i_{k}})$$

Again, on the naive approach, in order to have the limsup big we want the left hand side to be large, as seen in (4). Thus the signs of the terms on the right hand side are really what we are concerned with here. This theory is consistent with our prior comments from the Kochen Stone as the coefficients of the  $\mu(A_i)$  are +1 (therefore we want them big), and coefficients of the  $\mu(A_i \cap A_j)$  are negative (therefore we want them small). Indeed for any intersection, if the number of distinct sets is even then the coefficient is negative and we want these small for high limsup measure. Similarly if the number of distinct sets in the intersection is odd then the coefficient is positive and we would want these to be small.

This naturally leads us to the following. We construct two sequences of sets  $(A_i)_{i \in \mathbb{N}}$  and  $(B_i)_{i \in \mathbb{N}}$  such that the following intersection properties are satisfied:

$$\mu(A_i) \ge \mu(A_i) \quad \forall i \in \mathbb{N}$$
  

$$\mu(A_i \cap A_j) \le \mu(B_i \cap B_j) \quad \forall i < j \in \mathbb{N}$$
  

$$\mu(A_i \cap A_j \cap A_k) \ge \mu(B_1 \cap B_j \cap B_k) \quad \forall i < j < k \in \mathbb{N}$$
(13)

Indeed we actually go about solving the more general problem:

**Theorem 1.1.** For a given  $m \in \mathbb{N}$ , there exists  $(A_i)_{i \in \mathbb{N}}$  and  $(B_i)_{i \in \mathbb{N}}$  in the space [0,1]endowed with the Borel sigma algebra and Lebesgue measure satisfying:

$$\mu(A_i) \ge \mu(B_i) \quad \forall i \in \mathbb{N}$$
  

$$\mu(A_i \cap A_j) \le \mu(B_i \cap B_j) \quad \forall i < j \in \mathbb{N}$$
  

$$(-1)^r \mu(A_{i_1} \cap \dots \cap A_{i_r}) \le (-1)^r \mu(B_{i_1} \cap \dots \cap B_{i_r}) \quad \forall 1 \le r \le m, \quad \forall i_1 < \dots < i_r \in \mathbb{N} \quad (14)$$
  

$$\mu(\limsup A_i) = 0$$
  

$$\mu(\limsup B_i) = 1$$

**Remark.** Once again, we see that this is not specific to the 1 dimensional Lebesgue measure, analogues models in higher dimensions also occur. This can easily be done by placing  $(\times [0, 1]^{n-1})$  at the end of the sets and we have ourselves an example in  $\mathbb{R}^n$ .

Before we begin this we must define some sequences of sets with some useful intersecting properties. We'll later use these to define the sets in (14).

## 6.1 Construction of $(E_n^{p,q})_{n \in \mathbb{N}}$

Lets go ahead and define the sets:  $(E_n^{p,q})_{n\in\mathbb{N}}$  where  $p,q\in\mathbb{N}$  with  $p\leq q$ . We do this using 'shrinking' similitude's. That is, we re-apply what looks like a series of contractions (though these will alter at each level) and produce some sets with very well behaved intersection properties. Again this is in the ambient space of [0, 1] endowed with the usual topology and Borel measure.

**Proposition 6.1.** There exists a sequence  $(E_n^{p,q})_{n\in\mathbb{N}}$  for  $p,q\in\mathbb{N}$  with  $p\leq q$  with the following properties:

$$\mu(E_i^{p,q}) = \frac{p}{qi} \quad \forall 1 < i \in \mathbb{N}$$

$$\mu(E_i^{p,q} \cap E_j^{p,q}) = \frac{p^2}{q^2 j} \quad \forall 1 < i < j \in \mathbb{N}$$

$$\mu(E_{i_1}^{p,q} \cap E_{i_2}^{p,q} \cap \dots \cap E_{i_n}^{p,q}) = \frac{p^n}{q^n i_n} \quad \forall 1 < i_1 < i_2 < \dots < i_n \in \mathbb{N}$$

$$(15)$$

*Furthermore;* 

$$\limsup_{n \to \infty} (E_n^{p,q}) = 0$$

**Remark.** Perhaps its worth saying here that the first set is a bit annoying and that the properties don't hold here. We can amend this by 'shifting our sets up', that is the new  $E_1^{p,q}$  is the original  $E_2^{p,q}$  and so on. We needn't do that here as in the grand scheme of things this makes practically no difference to the solution later.

The reason for defining the sets in the way we do here is purely to have these nice forms for the intersection measures. Thus when we combine these sets to form our  $A_i$  and  $B_i$  we can achieve the inequalities desired. *Proof.* We begin this by defining two sequences of sets  $(E_n^{p,q})_{n\in\mathbb{N}}$  and  $(F_n^{p,q})_{n\in\mathbb{N}}$ . As is typical we set  $E_1^{p,q} = F_1^{p,q} = [0,1]$ .

For the next set we take the prior  $[0,1] = F_1^{p,q}$  and place q disjoint intervals of equal measure within it such that the total measure is  $\frac{1}{2}$ . Explicitly done this will be:

$$F_2^{p,q} = \left[0, \frac{1}{2q}\right] \cup \left[\frac{1}{q}, \frac{3}{2q}\right] \cup \left[\frac{2}{q}, \frac{5}{2q}\right] \cup \dots \cup \left[\frac{q-1}{q}, \frac{2q-1}{2q}\right]$$

We then construct  $E_2^{p,q}$  from this by simply selecting the first p intervals and discarding the rest. This ensures we have  $\mu(E_2^{p,q}) = \frac{p}{q}\mu(F_2^{p,q}) = \frac{p}{2q}$ .

Next we do  $F_3^{p,q}$ . This is done by taking the intervals of  $F_2^{p,q}$  and placing q distinct and equal length intervals within such that the total measure of these is  $\frac{1}{3}$ . Specifically, each interval in  $F_3^{p,q}$  has length  $\frac{1}{3q^2}$ . Trivially these will all 'fit in' as the lengths of the prior intervals is  $\frac{1}{2q}$  and the total length of the intervals within these is  $\frac{1}{3q}$ .

Now to construct  $E_3^{p,q}$  from these we simply go along and select the first p, then skip the next q-p, then select the next p, then skip the next q-p and so on. Thus the total measure achieved is  $\mu(E_3^{p,q}) = \frac{p}{a}\mu(F_3^{p,q}) = \frac{p}{3q}$ .

We then construct the subsequent intervals inductively: Having constructed  $F_n^{p,q}$  out of  $q^{n-1}$  intervals of equal length with total measure  $\frac{1}{n}$ . We then do a similar procedure to before by placing q smaller intervals into the intervals of  $F_n^{p,q}$ . Each of these new intervals will be of length  $\frac{1}{(n+1)q^n}$ . Thus the total measure is  $q^{n-1} \times q \times \frac{1}{(n+1)q^n} = \frac{1}{n+1}$ . Again we check that there is enough 'room to fit in'. Within one of the intervals from  $F_n^{p,q}$  of length  $\frac{1}{nq^{n-1}}$  we construct  $F_{n+1}^{p,q}$  by placing q intervals of length  $\frac{1}{(n+1)q^n}$ . Thus these combined have total measure  $\frac{1}{(n+1)q^{n-1}}$ . Whence the total measure is strictly less and we can indeed construct  $F_{n+1}^{p,q}$  as desired.

Next for the construction of  $E_{n+1}^{p,q}$  we again go along the intervals of  $F_{n+1}^{p,q}$  and select the first p to be included in  $E_{n+1}^{p,q}$  then reject the next q-p, then include the following p, then reject the next q-p and so on. This implies the measure of  $E_{n+1}^{p,q}$  is  $\frac{p}{q(n+1)}$ . This completes the construction.


Now we examine further the intersection properties too and realise the following:

$$\mu(E_{i}^{p,q}) = \frac{p}{qi} \quad \forall 1 < i \in \mathbb{N}$$

$$\mu(E_{i}^{p,q} \cap E_{j}^{p,q}) = \frac{p^{2}}{q^{2}j} \quad \forall 1 < i < j \in \mathbb{N}$$

$$\mu(E_{i_{1}}^{p,q} \cap E_{i_{2}}^{p,q} \cap ... \cap E_{i_{n}}^{p,q}) = \frac{p^{n}}{q^{n}i_{n}} \quad \forall 1 < i_{1} < i_{2} < ... < i_{n} \in \mathbb{N}$$
(16)

It is finally worth noting that for reasons discussed in prior chapters the limsup of  $E_n^{p,q}$  will be 0. Again by the containment properties and nesting of the  $F_n^{p,q}$ , which have vanishing measure, the limsup of these is null. Furthermore the  $E_n^{p,q}$  are contained in the respective



 $F_n^{p,q}$  and thus the limsup of these is null. Explicitly written:

$$\mu(\limsup_{n \to \infty} (E_n^{p,q})) \le \mu(\limsup_{n \to \infty} (F_n^{p,q}))$$
$$\le \mu(F_n^{p,q}) = \frac{1}{n} \quad \forall n$$

Thus the limsup is 0 measure. Note on the second line we use the containment of the  $F_n^{p,q}$ .

## 6.2 Proof of Theorem 1.1

#### **6.2.1** Construction of $A_n$

The way we go about constructing our sequences to solve this problem is to shrink some of the  $E_n^{p,q}$  sets by a constant and then combine them shifted along so each of the  $E_n^{p,q}$  used doesn't intersect with others. Note that here  $c_i E^{p,q}$  means we multiply each element of our set  $E^{p,q}$  by  $c_i$ . As the c's will always be less than 1 this will mean the sets shrink.

Specifically this is that  $(A_n)_{n \in \mathbb{N}}$  will be:

$$A_n = c_1 E_n^{p_1, q_1} \cup (c_2 E_n^{p_2, q_2} + d_2) \cup (c_3 E_n^{p_2, q_3} + d_3) \cup \dots \cup (c_k E_n^{p_k, q_k} + d_k)$$

The  $p_i$  and  $q_i$  will be chosen later to make inequalities work. Also, k will just be the number of these sets used and will be chosen later. Here the  $c_i$  are the shrinking factors and  $d_i$  are the shifts (both are positive quantities) chosen to ensure there is no overlap between the distinct  $E_n^{p,q}$ . That is to say  $d_i \ge d_{i-1} + c_{i-1}$ . Furthermore we wish for it to lie in the unit interval so  $c_k + d_k \le 1$ .

We now examine the sizes of these sets and the measures of their intersections also. We use what we found in Proposition 6.1, specifically equation (15). To do this we observe that our choices of  $c_i$  and  $d_i$  were such that there was separation between these sets.

**Lemma 6.1.** The sequence  $(A_i)_{i \in \mathbb{N}}$  satisfies:

$$\begin{split} \mu(A_i) &= c_1 \frac{p_1}{iq_1} + c_2 \frac{p_2}{iq_2} + \ldots + c_k \frac{p_k}{iq_k} \quad \forall 1 < i \in \mathbb{N} \\ \mu(A_i \cap A_j) &= c_1 \frac{p_1^2}{jq_1^2} + c_2 \frac{p_2^2}{jq_2^2} + \ldots + c_k \frac{p_k^2}{jq_k^2} \quad \forall 1 < i < j \in \mathbb{N} \\ \mu(A_{i_1} \cap A_{i_2} \cap \ldots \cap A_{i_r}) &= c_1 \frac{p_1^r}{i_r q_1^r} + c_2 \frac{p_2^r}{i_r q_2^r} + \ldots + c_k \frac{p_k^r}{i_r q_k^r} \quad \forall 1 < i_1 < i_2 < \ldots < i_n \in \mathbb{N} \\ \mu(\limsup(A_n)) &= 0 \end{split}$$

*Proof.* We go through each of the rows of the lemma and use (15) to establish the desired equalitites.

$$\begin{split} \mu(A_i) &= \mu(c_1 E_i^{p_1,q_1} \cup (c_2 E_i^{p_2,q_2} + d_2) \cup (c_3 E_i^{p_2,q_3} + d_3) \cup \ldots \cup (c_k E_i^{p_k,q_k} + d_k)) \\ &= \mu(c_1 E_i^{p_1,q_1}) + \mu(c_2 E_i^{p_2,q_2}) + \mu(c_3 E_i^{p_2,q_3}) + \ldots + \mu(c_k E_i^{p_k,q_k})) \\ &= c_1 \mu(E_i^{p_1,q_1}) + c_2 \mu(E_i^{p_2,q_2}) + c_3 \mu(E_i^{p_2,q_3}) + \ldots + c_k \mu(E_i^{p_k,q_k}) \\ &= c_1 \frac{p_1}{iq_1} + c_2 \frac{p_2}{iq_2} + c_3 \frac{p_3}{iq_3} + \ldots + c_k \frac{p_k}{iq_k} \end{split}$$

In a similar vein we find that for i < j:

$$\mu(A_i \cap A_j) = c_1 \mu(E_i^{p_1,q_1} \cap E_j^{p_1,q_1}) + c_2 \mu(E_i^{p_2,q_2} \cap E_j^{p_2,q_2}) + \dots + c_k \mu(E_i^{p_k,q_k} \cap E_j^{p_k,q_k})$$
$$= c_1 \frac{p_1^2}{jq_1^2} + c_2 \frac{p_2^2}{jq_2^2} + \dots + c_k \frac{p_k^2}{jq_k^2}$$

Lastly for  $i_1 < i_2 < \ldots < i_r$ 

$$\mu(A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_r}) = c_1 \frac{p_1^r}{i_r q_1^r} + c_2 \frac{p_2^r}{i_r q_2^r} + \dots + c_k \frac{p_k^r}{i_r q_k^r}$$

It is finally worth noting that for reasons discussed in prior chapters and the previous proof the limsup  $A_n$  has measure 0. As after-all, it is the disjoint combination of  $E_n^{p,q}$  sequences, it thus suffices to look at the limsup measure of each  $E_n^{p,q}$ . This was shown to be zero in Proposition 6.1.

#### **6.2.2** Construction of $B_n$

We do a similar thing with the  $(B_n)_{n\in\mathbb{N}}$  except now we also have a 'floating bit' as seen in previous examples. This will simply be a 'small' interval that moves along to ensure full limsup measure. This will be 'small' in comparison to the other sets as we will see. The float will be called  $H_n^{\delta}$ , where  $H_n^{\delta}$  is the moving interval and  $\delta$  the small scaling factor such that the total width is  $\frac{\delta}{n}$ . We have to add an extra subtlety that  $H_n^{\delta}$  doesn't intersect the  $F_n^{p,q}$  corresponding to the  $E_n^{p,q}$  in  $B_n$ . This ensures that when we intersect sets later, the  $H_n^{\delta}$  don't intersect any of the later  $E^{p,q}$  sets which aren't already intersected. Where we 'miss out' these parts we just continue to place mass after this until the total is  $\delta/n$ . The  $H_n^{\delta}$  ensure full limsup for the  $B_n$ . This is clear as the  $H_n^{\delta}$  keep cycling through the unit interval infinitely often as the total measure sum diverges. Perhaps we should note that the only area not hit by  $H_n^{\delta}$  is the  $F_n$  which shrink to a measure 0 set anyway. Explicitly written:

$$B_n = \tilde{c}_1 E_n^{p_1, q_1} \cup (\tilde{c}_2 E_n^{p_2, q_2} + \tilde{d}_2) \cup (\tilde{c}_3 E_n^{p_2, q_3} + \tilde{d}_3) \cup \dots \cup (\tilde{c}_k E_n^{p_k, q_k} + \tilde{d}_k) \cup H_n^{\delta}$$

Again, the  $p_i$ ,  $q_i$ , k are as before for  $A_i$  and will be chosen later. Similarly the  $\tilde{c}_i$  are to be chosen later and the  $\tilde{d}_i$  are separation constants. Now we observe the following:

**Lemma 6.2.** The sequence  $(B_i)_{i \in \mathbb{N}}$  satisfies:

$$\begin{split} \mu(B_i) &\leq \tilde{c}_1 \frac{p_1}{iq_1} + \tilde{c}_2 \frac{p_2}{iq_2} + \tilde{c}_3 \frac{p_3}{iq_3} + \ldots + \tilde{c}_k \frac{p_k}{iq_k} + \frac{\delta}{i} \quad \forall 1 < i \in \mathbb{N} \\ \mu(B_i \cap B_j) &\geq \tilde{c}_1 \frac{p_1^2}{jq_1^2} + \tilde{c}_2 \frac{p_2^2}{jq_2^2} + \ldots + \tilde{c}_k \frac{p_k^2}{jq_k^2} \quad \forall 1 < i < j \in \mathbb{N} \\ \mu(B_i) &\leq \tilde{c}_1 \frac{p_1}{iq_1} + \tilde{c}_2 \frac{p_2}{iq_2} + \tilde{c}_3 \frac{p_3}{iq_3} + \ldots + \tilde{c}_k \frac{p_k}{iq_k} + \frac{\delta}{i} \\ \mu(B_{i_1} \cap B_{i_2} \cap \ldots \cap B_{i_r}) &\geq \tilde{c}_1 \frac{p_1^r}{i_r q_1^r} + \tilde{c}_2 \frac{p_2^r}{i_r q_2^r} + \ldots + \tilde{c}_k \frac{p_k^r}{i_r q_k^r} \quad \forall 1 < i_1 < i_2 < \ldots < i_n \in \mathbb{N} \\ \mu(B_{i_1} \cap B_{i_2} \cap \ldots \cap B_{i_r}) &\leq \tilde{c}_1 \frac{p_1^r}{i_r q_1^r} + \tilde{c}_2 \frac{p_2^r}{i_r q_2^r} + \ldots + \tilde{c}_k \frac{p_k^r}{i_r q_k^r} + \frac{\delta}{i_r} \quad \forall 1 < i_1 < i_2 < \ldots < i_n \in \mathbb{N} \\ \mu(B_{i_1} \cap B_{i_2} \cap \ldots \cap B_{i_r}) &\leq \tilde{c}_1 \frac{p_1^r}{i_r q_1^r} + \tilde{c}_2 \frac{p_2^r}{i_r q_2^r} + \ldots + \tilde{c}_k \frac{p_k^r}{i_r q_k^r} + \frac{\delta}{i_r} \quad \forall 1 < i_1 < i_2 < \ldots < i_n \in \mathbb{N} \\ \lim \sup(B_n) &= 1 \end{split}$$

**Remark.** Again, we needn't have strict equalities as we are really just concerned with bounding these quantities by the corresponding for  $A_i$ . We omit many details that have already been stated in the prior lemma.

*Proof.* We do a similar process with the B's but now we notice the  $H_n^{\delta}$  term poses a challenge. This means the measure is more difficult to calculate precisely but we may form two obvious bounds listed below. By the way we construct the 'floating bit' we know it won't intersect any of the later E-sets however they could potentially intersect each other as is reflected in the bounds. The lower bound is trivial as we just neglect the  $H_n^{\delta}$  terms.

$$\mu(B_i) \ge \tilde{c}_1 \frac{p_1}{iq_1} + \tilde{c}_2 \frac{p_2}{iq_2} + \tilde{c}_3 \frac{p_3}{iq_3} + \dots + \tilde{c}_k \frac{p_k}{iq_k}$$
$$\mu(B_i) \le \tilde{c}_1 \frac{p_1}{iq_1} + \tilde{c}_2 \frac{p_2}{iq_2} + \tilde{c}_3 \frac{p_3}{iq_3} + \dots + \tilde{c}_k \frac{p_k}{iq_k} + \frac{\delta}{i}$$
$$\mu(B_{i_1} \cap B_{i_2} \cap \dots \cap B_{i_r}) \ge \tilde{c}_1 \frac{p_1^r}{i_r q_1^r} + \tilde{c}_2 \frac{p_2^r}{i_r q_2^r} + \dots + \tilde{c}_k \frac{p_k^r}{i_r q_k^r}$$
$$\mu(B_{i_1} \cap B_{i_2} \cap \dots \cap B_{i_r}) \le \tilde{c}_1 \frac{p_1^r}{i_r q_1^r} + \tilde{c}_2 \frac{p_2^r}{i_r q_2^r} + \dots + \tilde{c}_k \frac{p_k^r}{i_r q_k^r} + \frac{\delta}{i_r}$$

Discussion of the limsup measure was stated earlier on. The manner in which we construct  $H_n^{\delta}$  ensures the sequence is of full measure.

We then examine what this means for our original problem.

### 6.2.3 Using these to rephrase (14)

We see that the inequalities above combined with (14) mean that it suffices to find values that satisfy the following set of inequalities for  $1 \le r \le m$ . (Note earlier on we discussed the fact the limsups attain 0 and 1 respectively). For the purposes of solving (14) where we go up to m intersections, we choose k (the number of  $E^{p,q}$  sets) to also equal m.

For r odd:

$$c_1 \frac{p_1^r}{i_r q_1^r} + c_2 \frac{p_2^r}{i_r q_2^r} + \dots + c_m \frac{p_m^r}{i_r q_m^r} \ge \tilde{c}_1 \frac{p_1^r}{i_r q_1^r} + \tilde{c}_2 \frac{p_2^r}{i_r q_2^r} + \dots + \tilde{c}_k \frac{p_m^r}{i_r q_m^r} + \frac{\delta}{i_r}$$
(17)

For r even:

$$c_1 \frac{p_1^r}{i_r q_1^r} + c_2 \frac{p_2^r}{i_r q_2^r} + \dots + c_m \frac{p_m^r}{i_r q_m^r} \le \tilde{c}_1 \frac{p_1^r}{i_r q_1^r} + \tilde{c}_2 \frac{p_2^r}{i_r q_2^r} + \dots + \tilde{c}_m \frac{p_m^r}{i_r q_m^r}$$
(18)

We go on to define the following values for the  $c_i$ ,  $\tilde{c}_i$ ,  $\delta$  and  $p_i/q_i$  for  $1 \le i \le m$  in order to satisfy these.

We provide one method of finding such suitable constants. Others may be done perhaps by viewing it as a matrix problem. The matrices formed are similar to what are known as the Vandermonde matrices. Perhaps properties of these could be used. Expressed now in terms of matrices for visual convenience we have:

$$\begin{pmatrix} -\frac{p_1}{q_1} & -\frac{p_2}{q_2} & \dots & -\frac{p_m}{q_m} \\ \frac{p_1^2}{q_1^2} & \frac{p_2^2}{q_2^2} & \dots & \frac{p_m^2}{q_m^2} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{(-p_1)^m}{q_1^m} & \frac{(-p_2)^m}{q_2^m} & \dots & \frac{(-p_m)^m}{q_m^m} \end{pmatrix} \underline{\mathbf{c}} \leq \begin{pmatrix} -\frac{p_1}{q_1} & -\frac{p_2}{q_2} & \dots & -\frac{p_m}{q_m} \\ \frac{p_1^2}{q_1^2} & \frac{p_2^2}{q_2^2} & \dots & \frac{p_m^2}{q_m^2} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{(-p_1)^m}{q_1^m} & \frac{(-p_2)^m}{q_2^m} & \dots & \frac{(-p_m)^m}{q_m^m} \end{pmatrix} \underline{\mathbf{c}} \leq \begin{pmatrix} -\frac{p_1}{q_1} & -\frac{p_2}{q_2} & \dots & -\frac{p_m}{q_m} \\ \frac{p_1^2}{q_1^2} & \frac{p_2^2}{q_2^2} & \dots & \frac{p_m^2}{q_m^2} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{(-p_1)^m}{q_1^m} & \frac{(-p_2)^m}{q_2^m} & \dots & \frac{(-p_m)^m}{q_m^m} \end{pmatrix} \underline{\mathbf{c}} \leq \begin{pmatrix} \delta \\ 0 \\ \delta \\ \vdots \\ \delta \end{pmatrix}$$

where:

$$\underline{\mathbf{c}} = \begin{pmatrix} c_1 \\ c_2 \\ c_3 \\ \vdots \\ c_m \end{pmatrix} \quad \underline{\tilde{\mathbf{c}}} = \begin{pmatrix} \tilde{c}_1 \\ \tilde{c}_2 \\ \tilde{c}_3 \\ \vdots \\ \tilde{c}_m \end{pmatrix}$$

#### 6.2.4 Picking adequate constants to satisfy (17) and (18)

We will pick  $q_1 = 10^{b!}$  for some  $b \in \mathbb{N}$  to be chosen later. Then we select  $q_2 = 10^{(b+1)!}$ ,  $q_3 = 10^{(b+2)!}$  and so on up until  $q_m = 10^{(b+m-1)!}$ . For convenience we will take  $p_1 = p_2 = ... = p_m = 1$ . This ensures that when we take powers each element in the row will be of drastically different orders of magnitude so it will be easier to pick 'good'  $c_i$  coefficients (taking values 1 or 2 times a power of 10) which will ensure that the inequalities hold. Furthermore we will take  $\delta$  incredibly small compared to these quantities. The problem then breaks down into picking  $\delta$ , b,  $c_i$  and  $\tilde{c}_i$  for the following:

Lemma 6.3. There exists constants satisfying:

$$\begin{pmatrix} -\frac{1}{10^{b!}} & \cdots & -\frac{1}{10^{(b+m-1)!}} \\ \frac{1}{10^{2\times b!}} & \cdots & \frac{1}{10^{2\times (b+m-1)!}} \\ \vdots & \ddots & \vdots \\ \frac{(-1)^m}{10^{m\times b!}} & \cdots & \frac{(-1)^m}{10^{m\times (b+m-1)!}} \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ c_3 \\ \vdots \\ c_k \end{pmatrix} \leq \begin{pmatrix} -\frac{1}{10^{b!}} & \cdots & -\frac{1}{10^{(b+m-1)!}} \\ \frac{1}{10^{2\times (b+m-1)!}} & \frac{1}{10^{2\times (b+m-1)!}} \\ \vdots & \ddots & \vdots \\ \frac{(-1)^m}{10^{m\times b!}} & \cdots & \frac{(-1)^m}{10^{m\times (b+m-1)!}} \end{pmatrix} \begin{pmatrix} \tilde{c}_1 \\ \tilde{c}_2 \\ \tilde{c}_3 \\ \vdots \\ \tilde{c}_m \end{pmatrix} - \begin{pmatrix} \delta \\ 0 \\ \delta \\ \vdots \\ \delta \end{pmatrix}$$

*Proof.* It is natural to denote this matrix by M

We will first fix the  $c_1$  then  $c_2$  and so on until  $c_m$ . Then pick a b big enough so our choices ensure a proper gap between terms and the desired 'largest term' in a row is indeed largest.

We will want to take the lower left to upper right diagonal elements to be the biggest magnitude terms in their respective rows (when multiplied by the  $c_i$ ).

We go about setting out our goal more precisely. We want the matrix values to be biggest along the reverse diagonal when we multiply out in the usual convention. That is for row t, the largest value is taken by the term in column m - t + 1, specifically  $M_{t,m-t+1}c_{m-t+1}$ . Furthermore from the way we fix the  $q_i$  to be factorial exponents of the power of 10 and the way we'll fix  $c_i$  and b to put 'big gaps' between terms so that the 'biggest term' in the row will be far bigger and will completely dominate the others in the row. Indeed it will dominate the sum of the others by a factor of 10, thereby rendering them negligible in comparison. We will also set  $\tilde{c}_i$  to be of the same magnitude but multiplied by 2 or 1/2 respectively to ensure the inequality holds.

We are able to normalise later but for now go about defining the  $c_i$  as follows:  $c_1 = 2$ .  $c_i = \frac{3+(-1)^{i+1}}{2} \times 10^{\gamma_i}$  for  $2 \le i \le m$ . We further set  $\tilde{c}_1 = 1$  and  $\tilde{c}_i = \frac{3+(-1)^i}{2} \times 10^{\gamma_i}$  for  $2 \le i \le m$ . Then  $\tilde{c}_i = c_i \times \{\frac{1}{2}, 2\}$  dependent on the parity of i.

We want the bottom left  $M_{m,1}c_1$  to be the largest term of this row. Furthermore we want  $M_{m-1,2}c_2$  to be the largest term of its row. Also,  $M_{m-t+1,t}c_t$  should be the largest of the (m-t+1)th row.

Due to the way we pick the elements of M (that is factorial exponents of the power of 10) the columns grow at vastly different rates. This means that say if  $M_{a,b}c_b > M_{a,d}c_d$ , where b < d then as the *d*th column shrinks quicker than the *b*th,  $M_{a+1,b}c_b > M_{a+1,d}c_d$  too-indeed the disparity will be orders of magnitude greater too due to the choice of how they grow.

Also, if  $M_{a,b}c_b < M_{a,d}c_d$  then by similar logic  $M_{a-1,b}c_b < M_{a-1,d}c_d$ . With this in hand it suffices to satisfy the following in order to achieve our goal:

To this end, we go about defining  $\gamma_i$  and  $\alpha_i$  as follows:

$$\gamma_2 = \frac{2m-1}{2}((b+1)! - b!)$$
$$\alpha_2 = \frac{1}{2}((b+1)! - b!)$$

For  $2 \le n \le m$ :

$$\gamma_n = \sum_{i=1}^{n-1} (m - \frac{1}{2} - (i-1))((b+i)! - (b+i-1)!)$$
$$\alpha_n = \frac{1}{2}((b+n-1)! - (b+n-2)!)$$

We now go about actually showing the desired properties of (19) do indeed hold:

$$\begin{split} M_{m-a+1,a}c_a/M_{m-a+1,a+1}c_{a+1} &= 10^{-(m-a+1)\times(b+a-1)!}10^{\gamma_a}/10^{-(m-a+1)\times(b+a)!}10^{\gamma_{a+1}} \\ &= 10^{(m-a+1)((b+a)!-(b+a-1)!)}/10^{(m-\frac{1}{2}-(a-1))((b+a)!-(b+a-1)!)} \\ &= 10^{\frac{1}{2}((b+a)!-(b+a-1)!)} \\ &= 10^{\alpha_{a+1}} \end{split}$$

$$M_{m-a+1,a}c_a/M_{m-a+1,a-1}c_{a-1} = 10^{-(m-a+1)\times(b+a-1)!}10^{\gamma_a}/10^{-(m-a+1)\times(b+a-2)!}10^{\gamma_{a-1}}$$
  
=  $10^{(m-\frac{1}{2}-(a-2))((b+a-1)!-(b+a-2)!)}/10^{(m-a+1)\times((b+a-1)!-(b+a-2)!)}$   
=  $10^{\frac{1}{2}((b+a-1)!-(b+a-2)!)}$   
=  $10^{\alpha_a}$ 

So we have that (19) holds and furthermore by properties discussed before:

$$\sum_{i \in \{1,2,\dots,m\} \setminus a} M_{m-a+1,i} c_i \le 10^{-\alpha_a} \times (m-1) \times M_{m-a+1,a} c_a$$

Thus picking say b = m the  $10^{-\alpha_a}$  term completely dominates the m-1 and thus  $M_{m-a+1,a}c_a$  dominates the sum of the rest of the row by orders of magnitude. Therefore the inequality on this row is controlled by our choice of  $c_i$  and  $\tilde{c}_i$ . This ensures the matrix inequality as stated in the lemma

*Proof.* (of Theorem 1.1) Thus by our choice of these quantities, we have that the original matrix equation and thus the expressions (17) and (18) are satisfied. We can renormalise the  $c_i$  and  $\tilde{c}_i$  if necessary. Furthermore  $\delta$  can then be selected to be smaller than all terms appearing in the expansion. That is to say  $\delta$  will be less than  $M_{a,b}c_b$  so that it won't influence any of the inequalities. This concludes the inequalities and we discussed the limsup properties earlier so have demonstrated the existence of two sequences satisfying (14).

Despite this result holding, it would still be interesting in finding some strengthened forms of the Borel Cantelli lemma and perhaps formulate some new forms dependent on more topological properties similar to that done by [BV23]. This would ultimately go some way in providing new tools to develop theory in areas such as Graph Theory, Diophantine approximation, Dynamical Systems and more.

## Bibliography

- [CE52] K. L. Chung and P. Erdös. "On the application of the Borel-Cantelli lemma". In: *Trans. Amer. Math. Soc.* 72 (1952).
- [ER59] P. Erdös and A Rényi. "On Cantor's series with convergent  $\sum 1/q_n$ ". In: Ann. Univ. Sci. Budap. Rolando Eötvös, Sect. Math 2 (1959).
- [KS64] Simon Kochen and Charles Stone. "A note on the Borel-Cantelli lemma". In: Illinois J. Math. 8 (1964).
- [Pet04] Valentin V. Petrov. "A generalization of the Borel-Cantelli lemma". In: Statist. Probab. Lett. 67.3 (2004).
- [FLS09] Chunrong Feng, Liangpan Li, and Jian Shen. "On the Borel-Cantelli lemma and its generalization". In: C. R. Math. Acad. Sci. Paris 347.21-22 (2009).
- [Fro12] Andrei N. Frolov. "Bounds for probabilities of unions of events and the Borel-Cantelli lemma". In: *Statist. Probab. Lett.* 82 (2012).
- [Fro14] A. N. Frolov. "On inequalities for probabilities of unions of events and the Borel-Cantelli lemma". In: Vestnik St. Petersburg Univ. Math. 47 (2014).
- [Fro15a] A. N. Frolov. "On estimation of probabilities of unions of events with applications to the Borel-Cantelli lemma". In: *Vestnik St. Petersburg Univ. Math.* 48 (2015).
- [Fro15b] Andrei N. Frolov. "On lower and upper bounds for probabilities of unions and the Borel-Cantelli lemma". In: *Studia Sci. Math. Hungar.* 52 (2015).
- [AO21] Rob Arthan and Paulo Oliva. "On the Borel-Cantelli Lemmas, the Erdős-Rényi theorem, and the Kochen-Stone theorem". In: J. Log. Anal. 13 (2021).
- [BV23] Victor Beresnevich and Sanju Velani. "The divergence Borel-Cantelli lemma revisited". In: J. Math. Anal. Appl. 519.1 (2023).