

THE DIMENSION-FREE GEHRING-HAYMAN INEQUALITY FOR QUASIGEODESICS

CHANG-YU GUO, MANZI HUANG* AND XIANTAO WANG

ABSTRACT. A well-known theorem of J. Heinonen and S. Rohde in 1993 states that if $D \subset \mathbb{R}^n$ is quasiconformally equivalent to a uniform domain, then the Gehring-Hayman inequality holds in D : quasihyperbolic geodesics in D minimize the Euclidean length among all curves in D with the same end points, up to a universal dimension-dependent multiplicative constant.

In this paper, we develop a new approach to strengthen the above result in the following three aspects: 1) obtain a dimension-free multiplicative constant in the Gehring-Hayman inequality; 2) relax the class of quasihyperbolic geodesics to more general quasigeodesics; 3) relax the quasiconformal equivalence to more general coarsely quasihyperbolic equivalence.

As a byproduct of our general approach, we are able to prove that the above improved Gehring-Hayman inequality indeed holds in Banach spaces. This answers affirmatively an open problem raised by J. Heinonen and S. Rohde in 1993 and reformulated by J. Väisälä in 2005.

CONTENTS

1. Introduction	2
1.1. Background	2
1.2. Main results	4
1.3. Outline of the proof	6
2. Preliminaries	7
2.1. List of reserved constants	7
2.2. Coarse quasihyperbolicity and quasiconformality	8
2.3. Quasigeodesics, λ -curves and uniform domains	9
2.4. Gromov hyperbolicity and Rips spaces	14
3. Compactness of λ -pairs with \mathfrak{C} -Condition	23
3.1. Main result	23
3.2. Auxiliary lemmas	24
3.3. Proof of Theorem 3.2	33
4. Proofs of the main results	35
5. Proof of Theorem 3.3	36
5.1. Preparation for the proof	37
5.2. ζ -sequence on $\alpha[x_1, y_1]$	40

Date: February 6, 2025.

2020 Mathematics Subject Classification. Primary: 30C65, 30F45; Secondary: 30C20.

Key words and phrases. Gromov hyperbolicity, Gehring-Hayman inequality, Quasiconformal map, Quasihyperbolic geodesic, Uniform domain.

*Corresponding author: Manzi Huang.

C.-Y. Guo is supported by the Young Scientist Program of the Ministry of Science and Technology of China (No. 2021YFA1002200), the NSF of China (No. 12101362 and 12311530037), the Taishan Scholar Project and the NSF of Shandong Province (No. ZR2022YQ01). M. Huang and X. Wang are partly supported by NSF of China (No.12371071 and No.11271182).

5.3. λ -curves associated to ζ -sequence	43
5.4. ξ -sequence on the associated λ -curves	46
5.5. Six-tuples with Property C	50
5.6. A sequence of six-tuples	66
5.7. Iterative construction of new six-tuples	76
5.8. Final proof of Theorem 3.3	79
References	81

1. INTRODUCTION

1.1. **Background.** The unit disk $\mathbb{D} \subset \mathbb{C}$ plays a significant role in planar conformal analysis, as seen from the celebrated Riemannian mapping theorem, it is the distinguished conformal representative of all proper simply connected domains. In particular, many analysis on proper simply connected domains can be done via the analysis of conformal maps on unit disks [31]. Replacing conformal mappings with the less rigid class of quasiconformal mappings, one would naturally expect that the class of quasiconformal images of unit ball inherits similar effects as planar disks in higher dimensions. Partially based on this point of view and on some ideas of F. John [24], O. Martio and J. Sarvas [30] introduced the important class of *uniform domains*, which roughly speaking requires that the domain is quasiconvex and that each pair of points be joinable by a twisted double cone which is not too crooked nor too thin. Since then, uniform domains have received considerably large interests in the past decades and they have formed the “correct” class of nice higher dimensional domains in analysis appearing in surprising many contexts [3, 10, 12, 13, 14, 17, 22, 23, 25, 29].

Another more geometric view of the unit disk $\mathbb{D} \subset \mathbb{C}$ is to regard it as a complete Riemannian 2-manifold of negative curvature -1, when changing the underlying Euclidean metric to standard hyperbolic metric. Due to the Riemann mapping theorem, in terms of hyperbolic geometry, the unit disk becomes the distinguished hyperbolic isometric representative of all proper simply connected domains. In the celebrated work [15], M. Gromov introduced the notion of *Gromov hyperbolic* metric spaces, which generalizes the metric properties of classical hyperbolic geometry and of trees. Recall that a geodesic metric space $X = (X, d)$ is called Gromov hyperbolic if there exists a constant $\delta > 0$ such that each side of a geodesic triangle in X lies in the δ -neighborhood of the other two sides. Gromov hyperbolicity is a large-scale property and turns out to be very useful in geometric group theory and metric geometry [4, 8].

In an important and fundamental work [5], M. Bonk, J. Heinonen and P. Koskela have shown that the above two point of view, relating uniform spaces and Gromov hyperbolic spaces, are indeed conformally equivalent in the sense that there is a one-to-one conformal correspondence between the quasiisometric classes of certain Gromov hyperbolic spaces and the quasisimilarity classes of bounded locally compact uniform spaces. A basic starting point here is to explain uniformity through negative curvature with the aid of quasihyperbolic metric and the quasihyperbolization of a uniform space is Gromov hyperbolic. Based on this, they have developed an interesting uniformization theory, which provides conformal deformations of proper geodesic Gromov hyperbolic spaces into bounded uniform spaces. An fundamental result in [5] implies that Gromov hyperbolic domains in \mathbb{R}^n satisfy the *Gehring-Hayman inequality* and a *ball-separation condition*, quantitatively. In a seminal work [2], Z. Balogh and S. Buckley proved that the converse statement also holds, resulting the following useful geometric characterization:

a domain $X \subset \mathbb{R}^n$ is Gromov hyperbolic with respect to the quasihyperbolic metric k if and only if X satisfies the Gehring-Hayman inequality and a ball-separation condition, quantitatively.

Since our main focus of this paper is related to the Gehring-Hayman inequality, it is useful to recall the definition here.

Definition 1.1. A metric space (X, d) is said to satisfy the Gehring-Hayman inequality if there exists a constant $b > 0$ such that for all $x, y \in X$ and for each quasihyperbolic geodesic $[x, y]$ with end points x and y and each curve γ with the same end points, it holds

$$\ell_d([x, y]) \leq b\ell_d(\gamma). \quad (1.1)$$

The Gehring-Hayman inequality, which was initially proved in [11], states that the hyperbolic geodesic in a simply connected planar domain essentially minimizes the Euclidean length among all curves in the domain with the same end points, up to a universal multiplicative constant. This inequality has later been generalized to quasihyperbolic geodesics in domains $\Omega \subset \mathbb{R}^n$ that are quasiconformally equivalent to uniform domains by J. Heinonen and S. Rohde in [20]. This inequality, together with its conformally deformed variant, has found successful applications in the study of lengths of radii or boundary behavior of conformal and quasiconformal mappings [1, 2, 6, 7, 19, 21, 26]. Important extensions of this inequality to general (locally compact) metric spaces were obtained in [5, 27]. For our later reference, we state the main theorem of J. Heinonen and S. Rohde as follows.

Theorem A ([20, Theorem 1.1]). Let $D \subset \mathbb{R}^n$ be a domain that is K -quasiconformally equivalent to a c -uniform domain. If ϑ is a quasihyperbolic geodesic in D and if γ is any other arc in D with the same end points as ϑ , then

$$\ell(\vartheta) \leq b\ell(\gamma),$$

where $b > 0$ depends only on the dimension n , the uniformity constant c and the quasiconformality coefficient K .

As straightforward applications of Theorem A, J. Heinonen and S. Rohde also obtained the following Gehring-Hayman inequality for image of quasihyperbolic geodesics.

Theorem B ([20, Theorem 1.5]). Suppose that $D \subsetneq \mathbb{R}^n$ is a c -uniform domain, and that $f : D \rightarrow D'$ is M -bi-Lipschitz in the quasihyperbolic metric. If ϑ is a quasihyperbolic geodesic in D , and if γ is any other arc in D with the same end points as ϑ , then

$$\ell(f(\vartheta)) \leq b_1 \ell(f(\gamma)),$$

where b_1 depends only on the dimension n , the uniformity constant c and the quasihyperbolicity coefficient M .

As far as we know, in all the previously mentioned works, the validity of Gehring-Hayman inequality depends (either explicitly or implicitly) on the finite dimensionality of the underlying space. Our aim of this paper is to derive dimension-free estimates in both Theorems A and B. This is partially motivated by extending (part of) the theory of quasiconformal maps established in the seminal work of Heinonen and Koskela [18] to a larger class of non-Loewner metric spaces (such as infinite dimensional Banach spaces), where new methods giving dimension-free estimates are valuable. Notice that the proof of Theorem A (and thus also Theorem B) relies crucially on *the Whitney decomposition of domains* and *the modulus of curve families*. Both these two techniques requires finite dimensionality of the underlying space \mathbb{R}^n (in particular,

the n -dimensional Lebesgue measure \mathcal{L}^n on \mathbb{R}^n) and thus produces an extra dependence on the dimension n . Therefore, in order to obtain dimension-free estimate, a new approach is necessary.

1.2. Main results. To state our main results of this paper, we need a couple of definitions. First of all, we recall the notion of an uniform domain in a general metric space X .

Definition 1.2. A domain D in a metric space $X = (X, d)$ is called c -uniform, $c \geq 1$, if each pair of points z_1, z_2 in D can be joined by a rectifiable curve γ in D satisfying

- (1) $\min_{j=1,2} \{\ell(\gamma[z_j, z])\} \leq c d_D(z)$ for all $z \in \gamma$, and
- (2) $\ell(\gamma) \leq c d(z_1, z_2)$,

where $\ell(\gamma)$ denotes the arc-length of γ with respect to the metric d , $\gamma[z_j, z]$ the part of γ between z_j and z , and $d_D(z) := d(z, \partial D)$. In a c -uniform domain D , any curve $\gamma \subset D$, which satisfies conditions (1) and (2) above, is called a c -uniform curve or a double c -cone curve.

From now on, we assume that E and E' are Banach spaces with dimensions no less than 2 and D is a proper subdomain in E .

Definition 1.3. A homeomorphism $f : D \rightarrow D'$, where $D' \subset E'$, is called C -coarsely M -quasihyperbolic, abbreviated (M, C) -CQH, if for all $x, y \in D$, it holds

$$\frac{1}{M}(k_D(x, y) - C) \leq k_{D'}(f(x), f(y)) \leq M k_D(x, y) + C,$$

where k_D and $k_{D'}$ denote the quasihyperbolic metrics in D and D' , respectively (see Section 2 for the definition). In particular, an $(M, 0)$ -CQH mapping is said to be M -bi-Lipschitz in the quasihyperbolic metric.

Definition 1.4. A curve $\gamma \subset D$ is said to be a c -quasigeodesic, $c \geq 1$, if for all x, y in γ , it holds

$$\ell_k(\gamma[x, y]) \leq c k_D(x, y),$$

where $\ell_k(\gamma[x, y])$ denotes the quasihyperbolic length of $\gamma[x, y]$ (see Section 2 for the definition).

It is clear that, a c -quasigeodesic is a quasihyperbolic geodesic if and only if $c = 1$.

Our main result of this paper reads as follows.

Theorem 1.5. *Suppose that $D \subsetneq \mathbb{R}^n$ is homeomorphic to a c -uniform domain via an (M, C) -CQH mapping. If ϑ is a c_0 -quasigeodesic in D and if γ is any other arc in D with the same end points as ϑ , then*

$$\ell(\vartheta) \leq b\ell(\gamma),$$

where $b = b(c_0, c, M, C)$ depends only on the given parameters c_0, c, M and C .

Theorem 1.5 improves on Theorem A in the following three aspects:

- (1) The constant b is independent of the dimension n .
- (2) ϑ in Theorem 1.5 lies in the larger class of quasigeodesics than quasihyperbolic geodesics.
- (3) The class of CQH mappings is *strictly larger* than the class of quasiconformal mappings. Indeed, by [32, Theorem 6.4] and [33, Theorem 4.14], quasiconformal mappings in Euclidean spaces are quantitatively coarsely quasihyperbolic, but the converse is not true; see Example 2.1 below for a counter-example.

As an application of Theorem 1.5, we obtain the following improved Gehring-Hayman inequality for image of quasigeodesics under bi-Lipschitz quasihyperbolic homeomorphisms.

Theorem 1.6. *Suppose that $D \subsetneq \mathbb{R}^n$ is a c -uniform domain, and that $f : D \rightarrow D'$ is M -bi-Lipschitz in the quasihyperbolic metric. If ϑ is a c_0 -quasigeodesic in D , and if γ is any other arc in D with the same end points as ϑ , then*

$$\ell(f(\vartheta)) \leq b_1 \ell(f(\gamma)),$$

where $b_1 = b_1(c, c_0, M)$ depends only on the given parameters c , c_0 and M .

Theorem 1.6 improves on Theorem B in the following two aspects:

- (1) The constant b_1 is independent of the dimension n .
- (2) ϑ is a quasigeodesic, which is weaker than quasihyperbolic geodesics unless $c_0 = 1$.

Since our new approach is rather general, we are able to prove the dimension-free Gehring-Hayman inequality in the setting of Banach spaces, from which Theorem 1.5 follows as a special case.

Theorem 1.7. *Let E be a Banach space and $D \subsetneq E$ be homeomorphic to a c -uniform domain via an (M, C) -CQH mapping. If ϑ is a c_0 -quasigeodesic in D , and if L is any other arc in D with the same end points as ϑ , then*

$$\ell(\vartheta) \leq b \ell(L),$$

where $b = b(c, c_0, C, M)$ depends only on the given parameters c , c_0 , C and M .

Theorem 1.7 provides an affirmative answer to the following question: *Does Theorem A hold for domains that are freely quasiconformally equivalent to an uniform domain in general Banach spaces?* This question was first asked by J. Heinonen and S. Rohde in [20, Remark 2.25] and then by J. Väisälä in [35, Open problem 5 in Section 6] with a more precise formulation using the class of coarsely quasihyperbolic homeomorphism (CQH). Theorem 1.7 actually answers this question in a slightly stronger form, where quasihyperbolic geodesics are relaxed to the larger class of quasigeodesics.

As an application of Theorem 1.7, we obtain the following improved Gehring-Hayman inequality for image of quasigeodesics in the setting of Banach spaces, from which Theorem 1.6 follows as a special case.

Theorem 1.8. *Let E be a Banach space. Suppose that $D \subsetneq E$ is a c -uniform domain, and that $f : D \rightarrow D'$ is M -bi-Lipschitz in the quasihyperbolic metric. If ϑ is a c_0 -quasigeodesic in D , and if L is any other arc in D with the same end points as ϑ , then*

$$\ell(f(\vartheta)) \leq b_1 \ell(f(L)),$$

where $b_1 = b_1(c, c_0, M)$ depends only on the given parameters c , c_0 and M .

To state our next application, we need to recall the notion of inner uniform domains and John domains.

If the condition (2) in Definition 1.2 is replaced by the weaker inequality

$$\ell(\gamma) \leq c \sigma_D(z_1, z_2),$$

where σ_D is the *inner distance* defined by

$$\sigma_D(z_1, z_2) = \inf\{\ell(\alpha) : \alpha \subset D \text{ is a rectifiable curve joining } z_1 \text{ and } z_2\},$$

then D is said to be *inner uniform* and the corresponding curve γ is called an *inner uniform curve*.

If D only satisfies the condition (1) in Definition 1.2, then it is said to be a c -John domain, and the corresponding curve γ is called a c -John curve or a c -cone curve.

A classical result of Gehring and Osgood [12, Corollary 2] states that each quasihyperbolic geodesic in a uniform domain in \mathbb{R}^n is a uniform curve, quantitatively. An analog of this result for inner uniform domains remain valid: each quasihyperbolic geodesic in a John domain in \mathbb{R}^n that quasiconformally equivalent to a uniform domain is an inner uniform curve, quantitatively; see for example [16, Proposition 3.7]. As an application of Theorem 1.7, we obtain an improved extension of this result to Banach spaces.

Theorem 1.9. *Suppose that $D \subsetneq E$ is an a -John domain which is homeomorphic to a c -uniform domain via an (M, C) -CQH mapping. Then each c_0 -quasigeodesic in D is an inner b -uniform curve with $b = b(a, c, C, M)$. In particular, D is an inner uniform domain.*

The improvement on the corresponding Euclidean result, mentioned before the statement of Theorem 1.9, is again three-fold. Firstly, the inner uniformity coefficient b is dimension-free. Secondly, the class of quasigeodesics is strictly larger than the class of quasihyperbolic geodesics. Thirdly, the class of CQH mappings is strictly larger than the class of quasiconformal mappings.

1.3. Outline of the proof. In this section, we briefly outline our approach and main innovations on the proof of Theorem 1.7. Let ϑ be a quasigeodesic in D with end points w and ϖ and let γ be any other curve in D with the same end points as ϑ . Our aim is to prove

$$\ell(\vartheta) \leq b \ell(L) \quad (*)$$

and we may assume $k_D(w, \varpi) > 1$ as the other case follows from simple computation.

Our strategy for the proof of (*) is to adapt the “*contradiction-compactness argument*” to (locally noncompact) Banach spaces. This type of argument has appeared in many different contexts in geometric analysis or metric geometry, but usually in the finite dimensional setting. To run the contradiction argument, suppose for some large b , $\ell(\vartheta) \geq b \ell(L)$. Then we want to construct some “good subcurves” ϑ_i of ϑ and L_i of L , with the same end points, such that $\frac{\ell(\vartheta_i)}{\ell(L_i)} \gtrsim \mu^i \rightarrow \infty$ as $i \rightarrow \infty$. On the other hand, we can use the quasihyperbolic length of L to give uniform upper bound for $\frac{\ell(\vartheta_i)}{\ell(L_i)} \leq M < \infty$. This contradiction would give the desired inequality.

To realize the above idea, we introduce the new class of λ -curves (see Definition 2.5) and λ -pairs (see Definition 3.1). λ -curves are essentially quasigeodesics with good control on the quasigeodesic coefficient. Comparing with quasihyperbolic geodesics, λ -curves are less sensitive with respect to small variations, which make the coarse analysis much more effective. A λ -pair (γ, L) consists of two curves in D with the same end points, where γ is a λ -curve, and we say that a λ -pair (γ, L) satisfies \mathfrak{C} -condition if $\ell(\gamma) \geq \mathfrak{C}\ell(L)$.

The most important step, to realize the previous strategy, is to prove the following “compactness theorem” (see Theorem 3.2) for λ -pairs: *if a λ -pair (γ, L) satisfies \mathfrak{C} -condition with \mathfrak{C} sufficiently large, then there exists a λ -subpair (γ_1, L_1) of (γ, L) , which satisfies $\mu\mathfrak{C}$ -condition with a large $\mu \gg 1$, quantitatively.* More precisely, it says that if the length of a λ -curve γ is too large compared with a fixed curve L with the same end points, then we may find two points w_1, w_2 on L such that for any λ -curve β connecting these two points, we have $\ell(\beta) \geq \mu\mathfrak{C}\ell(L[w_1, w_2])$. Geometric insight suggests us that in certain small portion L_1 on L , the curve γ has to oscillate a lot and so we may construct a λ -curve γ_1 with end points on L (based on γ and L) such that

$\ell(\gamma_1)/\ell(L_1)$ becomes definitely larger than $\ell(\gamma_1)/\ell(L_1)$. However, to write down a rigorous proof turns out to be not easy at all.

In our approach, the proof of Theorem 3.2 is again based on a contraction argument. One key auxiliary result for the proof is Theorem 3.3, which says that the length of a class of λ -curve is uniformly bounded from above by the inner distance between its end points. This auxiliary result allows us to construct a sequence of points with prescribed metric controls (see Lemmas 3.11 and 3.14), using basic properties of quasihyperbolic metrics, λ -curves, Gromov hyperbolicity and uniformity. A careful analysis on these points would lead to a contradiction.

The proof of Theorem 3.3 is extremely technical and is again by a contradiction argument. The main innovation in the proof is to introduce the notion of *six-tuples* (see Definition 5.13). A six-tuple $\Omega = [w_1, w_2, \gamma, \varpi, w_3, w_4]$ consists of five points $w_1, w_2, \varpi, w_3, w_4$ of special positions with controlled metric properties and a λ -curve γ connecting w_1 and w_2 . We shall analyze the basic properties of λ -curves via studying *six-tuples which satisfies Property C*. The main ingredient is to construct iteratively new six-tuples based on older ones with prescribed metric properties. This step requires very delicate and careful estimates of quasihyperbolic distances using basic properties of λ -curves, Gromov hyperbolicity and CQH-mappings. One feature of our analysis is that we reserve a large amount of constants (see Section 2.1 for the list) in the proof to compensate the lack of compactness in locally noncompact spaces and then derive a contradiction based on a careful selection of these constants.

It would be interesting to know whether Theorems 1.7, 1.8 and 1.9 hold under the weaker assumption that $D \subset E$ is δ -Gromov hyperbolic. We expect that the techniques developed in this paper would be useful in the study of this question.

Structure. The arrangement of this paper is as follows. In Section 2, we introduce the list of reserved constants used in this paper, recall some known results, and develop a series of elementary properties concerning the quasihyperbolic metric, quasigeodesics, λ -curves, solid arcs etc. In Section 3, we prove the compactness theorem for λ -pairs satisfying \mathfrak{C} -Condition, namely Theorem 3.2 mentioned above, assuming the technical Theorem 3.3. Section 4 is devoted to the proofs of main results in this paper. In Section 5, we give a complete proof of Theorem 3.3.

Notations. Throughout this paper, the norm of a vector $z \in E$ is written as $|z|$ and the distance between two points $z_1, z_2 \in E$ is denoted by $|z_1 - z_2|$. We write $[z_1, z_2]$ for the line segment with end points z_1 and z_2 .

Let D and D' be proper domains in E and E' , respectively, and let $f : D \rightarrow D'$ be a homeomorphism. For convenience, in what follows, the symbols x, y, z, \dots represents points in D , while the primes x', y', z', \dots denote their images in D' under f . Similarly, we use $\alpha, \beta, \gamma, \dots$ to denote arcs in D , while the primes $\alpha', \beta', \gamma', \dots$ refer to their images in D' under f . For a set A in D , A' often denotes its image in D' under f . Throughout this paper, we call γ an arc or a curve in D if it is a continuous map from an interval to D and we use $\ell_d(\gamma)$ and $\text{diam}_d(\gamma)$ to denote the d -length and diameter of γ , respectively. When the content is clear, we omit the subscript d .

2. PRELIMINARIES

2.1. List of reserved constants. In the following, we list the constants used in this paper.

- (1) C and M : $\min\{C, M\} \geq 1$;
- (2) c_0 : $c_0 > 1$;

- (3) c_1 and h_1 : $c_1 = \frac{9}{4}M(M+1)$ and $h_1 = C(\frac{9}{4}M+1)$ (see Lemma 2.9);
- (4) c_2 : $c_2 = c_2(c, c_1, h_1) \geq e^{1+c_1+h_1}$ (see Theorem 2.10 and Remark 2.11);
- (5) c_3 : $c_3 = 8c_2^2e^{2c_1+3h_1}$ (see Lemma 2.12 and Remark 2.13);
- (6) c_4 : $c_4 = c_4(c) \geq 8$ (see Theorem 2.14 and Remark 2.15);
- (7) v_0 : $v_0 = v_0(c_0, \lambda, \delta) \geq 16$ (see Lemma 2.17 and Remark 2.18);
- (8) μ_0 and ρ_0 : $\mu_0 = \mu_0(c, C, M, \delta) \geq 16$ and $\rho_0 = \rho_0(c, C, M, \delta) > 0$ (see Lemma 2.22 and Remark 2.23);
- (9) μ_1 and μ_2 : $\mu_1 = 3\mu_0$ (see Lemma 2.24) and $\mu_2 = \frac{1}{4} + \mu_0 + 2v_0$ (see Lemma 2.26);
- (10) μ_3, μ_4 and μ_5 : $\mu_3 = \frac{1}{8}c_0^2c_2c_3c_4^2\mu_2e^{4+C+h_1+M}$, $\mu_4 = 12c_2c_4(1+\mu_3)^2$ and $\mu_5 = 2\mu_4e^{2\mu_4e^{8\mu_4}}$;
- (11) μ_6 and μ_7 : $\mu_6 = \frac{1}{4}e^{\mu_4-2\mu_3}$ and $\mu_7 = e^{\mu_3+3\mu_4}$;
- (12) μ_8 and μ_9 : $\mu_8 = 4(1+c_3)^{3\mu_3+56\mu_1}e^{16\mu_7}$ and $\mu_9 = \mu_8^{\frac{1}{3}}$;
- (13) \mathbf{c} : $\mathbf{c} \geq 64\mu_1$.
- (14) τ_1, τ_2 and τ_3 : $2^{11}\mu_1 \leq 4\tau_1 \leq \tau_2 \leq \frac{1}{4}\tau_3$.

2.2. Coarse quasihyperbolicity and quasiconformality. In the section, we shall construct an example which shows that coarsely quasihyperbolic homeomorphisms are in general not necessarily quasiconformal, even in the finite dimensional case.

Example 2.1. For $s > 0$, let $\mathbb{B}(s) = \{x \in \mathbb{C} : |x| < s\}$, where \mathbb{C} denotes the complex plane, and O is the origin of \mathbb{C} . Set

$$f(x) = \begin{cases} g(|x|)\frac{x}{|x|}, & x \in \mathbb{B}(2) \setminus \{O\}, \\ O, & x = O, \end{cases}$$

where $g : [0, 2) \rightarrow [0, 2)$ is a homeomorphism defined as follows:

$$g(t) = \begin{cases} g_0(t), & t \in [0, \frac{1}{2}), \\ g_n(t), & t \in [1 - \frac{1}{2^n}, 1 - \frac{1}{2^{(n+1)}}) \text{ for } n \in \{1, 2, \dots\}, \\ t, & t \in [1, 2), \end{cases}$$

$g_0(t) = \frac{3}{2}t$, and for each $n \in \{1, 2, \dots\}$,

$$g_n(t) = a_n t + b_n$$

with

$$a_n = \frac{2^{(n+1)!}(2^{(n+2)!-(n+1)!} - 1)}{2^{(n+2)!}(2^{(n+1)!-n!} - 1)} \text{ and } b_n = 1 - \frac{1}{2^{(n+1)!}} - a_n(1 - \frac{1}{2^{n!}}).$$

Then the following statements hold.

- (1) $f : \mathbb{B}(2) \rightarrow \mathbb{B}(2)$ is (1, 4)-CQH.
- (2) $f : \mathbb{B}(1) \rightarrow \mathbb{B}(1)$ is not φ -FQC for any self-homeomorphism $\varphi : [0, +\infty) \rightarrow [0, +\infty)$. In particular, $f : \mathbb{B}(2) \rightarrow \mathbb{B}(2)$ is not quasiconformal by [32, Remark 3.5].

Recall that a homeomorphism $f : \mathbb{B}(1) \rightarrow \mathbb{B}(1)$ is said to be φ -FQC, if for any $x, y \in \mathbb{B}(2)$, it holds

$$\varphi^{-1}(k_{\mathbb{B}(1)}(x, y)) \leq k_{\mathbb{B}(1)}(f(x), f(y)) \leq \varphi(k_{\mathbb{B}(1)}(x, y)). \quad (2.1)$$

Proof. Firstly, we show that f is (1, 4)-CQH. For this, let $x, y \in \mathbb{B}(2)$.

- If $x, y \in \mathbb{B}(1)$, then $f(x), f(y) \in \mathbb{B}(1)$ and [32, Lemma 2.2(2)] give

$$k_{\mathbb{B}(2)}(f(x), f(y)) \leq 2 \leq k_{\mathbb{B}(2)}(x, y) + 2.$$

- If $x, y \in \mathbb{B}(2) \setminus \mathbb{B}(1)$, then $f(x) = x, f(y) = y$, and so,

$$k_{\mathbb{B}(2)}(f(x), f(y)) = k_{\mathbb{B}(2)}(x, y).$$

- If $x \in \mathbb{B}(1), y \in \mathbb{B}(2) \setminus \mathbb{B}(1)$, then $f(x) \in \mathbb{B}(1), f(y) = y$. It follows that

$$k_{\mathbb{B}(2)}(f(x), f(y)) \leq k_{\mathbb{B}(2)}(f(x), O) + k_{\mathbb{B}(2)}(O, f(y)) \leq 4 + k_{\mathbb{B}(2)}(x, y).$$

Therefore, we conclude from the above estimates that for any $x, y \in \mathbb{B}(2)$,

$$k_{\mathbb{B}(2)}(f(x), f(y)) \leq k_{\mathbb{B}(2)}(x, y) + 4.$$

Similarly, we may prove that for any $x, y \in \mathbb{B}(2)$,

$$k_{\mathbb{B}(2)}(x, y) \leq k_{\mathbb{B}(2)}(f(x), f(y)) + 4.$$

This shows that f is $(1, 4)$ -CQH.

Suppose on the contrary that $f: \mathbb{B}(1) \rightarrow \mathbb{B}(1)$ is φ -FQC for a self-homeomorphism $\varphi: [0, +\infty) \rightarrow [0, +\infty)$. To get a contradiction, for each $n \in \mathbb{N}$, select $x_n, y_n \in \mathbb{S}(1 - \frac{1}{2^{n!}})$ with

$$|x_n - y_n| = \frac{1}{2^{n!}}.$$

Then it follows from [32, Lemma 2.2(2)] that

$$k_{\mathbb{B}(1)}(x_n, y_n) \leq 1.$$

On the other hand, the definition of f ensures that both $f(x_n)$ and $f(y_n)$ are on $\mathbb{S}(1 - \frac{1}{2^{(n+1)!}})$, and

$$|f(x_n) - f(y_n)| \geq |x_n - y_n|.$$

This, together with [32, Lemma 2.2(1)] and (2.1), implies that

$$\varphi(1) \geq k_{\mathbb{B}(1)}(f(x_n), f(y_n)) \geq n \log 2,$$

which is impossible if n is sufficiently large. This completes the proof. \square

2.3. Quasigeodesics, λ -curves and uniform domains. We start this section by recalling the definition of quasihyperbolic metric, which was initially introduced by Gehring and Palka [13] for a domain in \mathbb{R}^n and then has been extensively studied in [12]. Since we are concerning it in a Banach space, we shall follow the presentation from [32].

The *quasihyperbolic length* of a rectifiable arc γ in a proper domain $D \subsetneq E$ is defined as

$$\ell_k(\gamma) := \int_{\gamma} \frac{|dz|}{d_D(z)}.$$

For any z_1, z_2 in D , the *quasihyperbolic distance* $k_D(z_1, z_2)$ between z_1 and z_2 is set to be

$$k_D(z_1, z_2) = \inf_{\gamma} \{\ell_k(\gamma)\},$$

where the infimum is taken over all rectifiable arcs γ joining z_1 and z_2 in D .

An arc γ from z_1 to z_2 is called a *quasihyperbolic geodesic* if $\ell_k(\gamma) = k_D(z_1, z_2)$. Clearly, each subarc of a quasihyperbolic geodesic is a quasihyperbolic geodesic. It is a well-known fact that a quasihyperbolic geodesic between any two points in E exists if the dimension of E is finite; see [12, Lemma 1]. This is not true in infinite dimensional spaces; see [33, Example 2.9].

In order to remedy this shortage, J. Väisälä [33] introduced the class of *quasigeodesics* in E ; see Definition 1.4. Moreover, he proved in [33, Theorem 3.3] that for each pair of points z_1 and z_2 in D and each $c > 1$, there exists a c -quasigeodesic in D joining z_1 and z_2 .

For any z_1, z_2 in D , let $\gamma \subset D$ be an arc with end points z_1 and z_2 . Then we have the following elementary estimates (see for instance [32, Section 2])

$$\ell_k(\gamma) \geq \log \left(1 + \frac{\ell(\gamma)}{\min\{d_D(z_1), d_D(z_2)\}} \right) \quad (2.2)$$

and

$$\begin{aligned} k_D(z_1, z_2) &\geq \log \left(1 + \frac{\sigma_D(z_1, z_2)}{\min\{d_D(z_1), d_D(z_2)\}} \right) \\ &\geq \log \left(1 + \frac{|z_1 - z_2|}{\min\{d_D(z_1), d_D(z_2)\}} \right) \geq \left| \log \frac{d_D(z_2)}{d_D(z_1)} \right|. \end{aligned} \quad (2.3)$$

Lemma 2.2. *Suppose that x and y are two points in D with $k_D(x, y) \geq 1$.*

- (1) *Then $\max\{d_D(x), d_D(y)\} \leq 2|x - y|$.*
- (2) *Let γ be an arc in D joining x and y . Then for any $w \in \gamma$, $d_D(w) \leq 2\ell(\gamma)$.*

Proof. To prove the first statement (1), we assume that

$$\max\{d_D(x), d_D(y)\} = d_D(x).$$

Suppose on the contrary that $d_D(x) > 2|x - y|$. Then the segment $[x, y] \subset \mathbb{B}(x, \frac{1}{2}d_D(x))$ and thus for any $z \in [x, y]$,

$$d_D(z) \geq d_D(x) - |x - z| > \frac{1}{2}d_D(x),$$

which implies

$$k_D(x, y) \leq \int_{[x, y]} \frac{|dz|}{d_D(z)} < 1.$$

This contradiction shows that the first statement holds.

To prove the second statement (2), we may assume that γ is rectifiable. Suppose on the contrary that there is some point $w_0 \in \gamma$ such that

$$\ell(\gamma) < \frac{1}{2}d_D(w_0).$$

Then for any $w \in \gamma$, we have

$$d_D(w) \geq d_D(w_0) - \ell(\gamma[w, w_0]) > \frac{1}{2}d_D(w_0).$$

This gives

$$k_D(x, y) \leq \int_{\gamma} \frac{|dw|}{d_D(w)} < 1.$$

This contradiction proves the second statement, and hence, the proof is complete. \square

Next we discuss the coarse length of a curve intrudced by J. Väisälä in [33, Section 4]. Let γ be an (open, closed or half open) arc in D . Write $\bar{x} = (x_0, \dots, x_m)$ with $m \geq 1$, whose coordinates form a finite sequence of m successive points of γ . For $h \geq 0$, we say that \bar{x} is

h -coarse if $k_D(x_{j-1}, x_j) \geq h$ for all $j \in \{1, \dots, m\}$. Let $\Phi_k(\gamma, h)$ be the family of all h -coarse sequences of γ . Set

$$s_k(\bar{x}) := \sum_{j=1}^m k_D(x_{j-1}, x_j) \quad \text{and} \quad \ell_k(\gamma, h) := \sup \{s_k(\bar{x}) : \bar{x} \in \Phi_k(\gamma, h)\}$$

with the convention that $\ell_k(\gamma, h) = 0$ if $\Phi_k(\gamma, h) = \emptyset$. Then the number $\ell_k(\gamma, h)$ is called the h -coarse quasihyperbolic length of γ .

Definition 2.3 (Solid arc). An arc γ in D is called (c, h) -solid with $c \geq 1$ and $h \geq 0$ if

$$\ell_k(\gamma[z_1, z_2], h) \leq c k_D(z_1, z_2)$$

for all pairs of points z_1 and z_2 in γ . We denote the class of all (c, h) -solid arcs in D by $S^{c,h}$. The symbol $S_{xy}^{c,h}$ stands for those arcs in $S^{c,h}$ with end points x, y .

By [34, Section 2.2] or [33, Section 4], we know that for any arc $\gamma \subset D$, $\ell_k(\gamma, 0) = \ell_k(\gamma)$. Notice that γ is a $(c, 0)$ -solid arc if and only if it is a c -quasigeodesic.

Definition 2.4 (Short arc). A curve γ in D is said to be h -short if for all $x, y \in \gamma$,

$$\ell_k(\gamma[x, y]) \leq k_D(x, y) + h.$$

It is clear from Definition 2.4 that every subarc of an h -short arc is h -short, every h -short arc must be h_1 -short for any $h_1 \geq h$, and every quasihyperbolic geodesic is h -short with $h = 0$.

Definition 2.5 (λ -curve). For each pair of distinct points $x, y \in D$ and $\lambda > 0$, we set

$$c_{x,y}^\lambda := \min \left\{ 1 + \frac{\lambda}{2k_D(x, y)}, \frac{9}{8} \right\}.$$

An arc $\gamma \subset D$ with end points x and y is called a λ -curve if it is a $c_{x,y}^\lambda$ -quasigeodesic in D . We use Λ_{xy}^λ to stand for the set of all λ -curves in D with end points x and y and use γ_{xy}^λ to denote a general λ -curve in Λ_{xy}^λ . Throughout this paper, we always assume that $\lambda \in (0, \frac{1}{2}]$.

By [33, Theorem 3.3], we know that for $x, y \in D$ and $\lambda > 0$, if $x \neq y$, then $\Lambda_{xy}^\lambda \neq \emptyset$. For $c_0 \geq 1$, we use Q^{c_0} to denote all c_0 -quasigeodesics in D and $Q_{xy}^{c_0}$ for c_0 -quasigeodesics in D with end points x and y .

The next result shows that all λ -curves possess the following shortness.

Lemma 2.6. *For any two distinct points $x \neq y \in D$, every λ -curve $\gamma \in \Lambda_{xy}^\lambda$ is λ -short.*

Proof. Take any two points $u \neq v \in \gamma$. Then

$$\ell_k(\gamma) \leq \left(1 + \frac{\lambda}{2k_D(x, y)}\right) k_D(u, v) = k_D(u, v) + \frac{\lambda k_D(u, v)}{2k_D(x, y)}.$$

Observe that γ is a $\frac{9}{8}$ -quasigeodesic and thus

$$k_D(u, v) \leq \ell_k(\gamma) \leq \frac{9}{8} k_D(x, y).$$

Combining the above two estimates gives the desired inequality:

$$\ell_k(\gamma[u, v]) \leq k_D(u, v) + \lambda.$$

□

The following lemma shows that every subcurve of a λ -curve is a 2λ -curve.

Lemma 2.7. *Let $x \neq y$ be two distinct points in D and fix a λ -curve $\gamma \in \Lambda_{xy}^\lambda$. Then for any $z_1 \neq z_2 \in \gamma$, the subarc $\gamma[z_1, z_2]$ of γ belongs to $\Lambda_{z_1 z_2}^{2\lambda}$.*

Proof. Let $z_1 \neq z_2 \in \gamma$ and observe that for any $u \neq v \in \gamma$,

$$\ell_k(\gamma[u, v]) \leq \min \left\{ 1 + \frac{\lambda}{2k_D(x, y)}, \frac{9}{8} \right\} k_D(u, v).$$

To prove the lemma, it suffices to show that

$$\min \left\{ 1 + \frac{\lambda}{2k_D(x, y)}, \frac{9}{8} \right\} \leq \min \left\{ 1 + \frac{\lambda}{k_D(z_1, z_2)}, \frac{9}{8} \right\}.$$

As γ is a $\frac{9}{8}$ -quasigeodesic, we have

$$k_D(z_1, z_2) \leq \ell_k(\gamma) \leq \frac{9}{8} k_D(x, y),$$

which clearly implies

$$1 + \frac{\lambda}{2k_D(x, y)} \leq 1 + \frac{\lambda}{k_D(z_1, z_2)}.$$

The proof is thus complete. \square

Lemma 2.8. *Let $x \neq y$ be two distinct points in D . For any curve β connecting x and y and any λ -curve $\gamma \in \Lambda_{xy}^\lambda$, if $\ell(\gamma) > \nu_1 \ell(\beta)$ with $\nu_1 \geq 18e^9$, then for any $w \in \beta$, $d_D(w) \leq 2\ell(\beta)$.*

Proof. By Lemma 2.2, it suffices to show that $k_D(x, y) > 1$. Suppose on the contrary that $k_D(x, y) \leq 1$. Since γ is a $\frac{9}{8}$ -quasigeodesic, it follows from (the proof of) [34, Lemma 2.31] with $c = \frac{9}{8}$ and $C = 1$ that γ is a $18e^9$ -uniform curve. This implies that

$$\ell(\gamma) \leq 18e^9 |x - y| \leq 18e^9 \ell(\beta),$$

which clearly contradicts with the assumption that $\ell(\gamma) > \nu_1 \ell(\beta) \geq 18e^9 \ell(\beta)$. \square

From now on, we make the following convention: For two conditions, we say that Condition Φ quantitatively implies Condition Ψ if Condition Φ implies Condition Ψ and the data of Condition Ψ depends only on that of Condition Φ .

Lemma 2.9. (1) *For any two distinct points $x \neq y \in D$, each element in Λ_{xy}^λ is a $(c_0, 0)$ -solid arc and belongs to $Q_{xy}^{c_0}$ with $c_0 = \frac{9}{8}$.*

(2) *Suppose that $f : D \rightarrow D'$ is (M, C) -CQH. Then the image of any λ -curve under f is a (c_1, h_1) -solid arc, where $c_1 = \frac{9}{4}M(M + 1)$ and $h_1 = C(\frac{9}{4}M + 1)$.*

Proof. By [33, Theorem 4.15], we see that under an (M, C) -CQH mapping, the solidness of an arc implies the solidness of its image, quantitatively. The explicit formulation of c_1 and h_1 in terms of M and C follows from direct computations in [33, Theorems 4.11 and 4.14]. \square

By [33, Theorem 6.22] and Lemma 2.9(1), we have the following result concerning solid arcs and λ -curves in uniform domains.

Theorem 2.10 ([33, Theorem 6.22]). *Suppose that D is a c -uniform domain. For $z_1, z_2 \in D$, let $\gamma \in S_{z_1 z_2}^{c_1, h_1} \cup \Lambda_{z_1 z_2}^\lambda$. Then for some $c_2 = c_2(c, c_1, h_1)$, we have*

(1) for any $z \in \gamma$,

$$\min\{\text{diam}(\gamma[z_1, z]), \text{diam}(\gamma[z, z_2])\} \leq c_2 d_D(z);$$

(2) $\text{diam}(\gamma) \leq c_2 \max\{|z_1 - z_2|, 2(e^{h_1} - 1) \min\{d_D(z_1), d_D(z_2)\}\}$.

Remark 2.11. In the rest of this paper, we assume that

$$c_2 = c_2(c, c_1, h_1) \geq e^{1+c_1+h_1}.$$

Based on Lemma 2.9 and Theorem 2.10, we next prove the following diameter estimate for λ -curves under CQH-mappings in an uniform domain.

Lemma 2.12. *Suppose that $f : D \rightarrow D'$ is (M, C) -CQH, D' is a c -uniform domain, and let $x, y \in D$ be two distinct points. For any λ -curve $\gamma \in \Lambda_{xy}^\lambda$, if there is $x_0 \in \gamma$ such that $|x' - x'_0| = \frac{1}{2}|x' - y'|$, then*

$$\text{diam}(\gamma') \leq c_3 d_{D'}(x'_0),$$

where $c_3 = c_3(c, C, M)$.

Recall that $\gamma' = f(\gamma)$ denotes the image of γ under f .

Proof. Without loss of generality, we assume that $d_{D'}(x') \leq d_{D'}(y')$.

By Lemma 2.9, γ' is a (c_1, h_1) -solid with $c_1 = \frac{9}{4}M(M+1)$ and $h_1 = C(\frac{9}{4}M+1)$. To prove this lemma, we consider two cases: $|x' - y'| < \frac{1}{2}d_{D'}(x')$ and $|x' - y'| \geq \frac{1}{2}d_{D'}(x')$.

In the former case, it follows from Lemma 2.2(1) that $k_{D'}(x', y') < 1$. Since $f^{-1} : D' \rightarrow D$ is (M, C) -CQH as well, we get from Lemma 2.9 that

$$k_D(x, x_0) \leq \ell_k(\gamma) \leq \frac{9}{8}k_D(x, y) \leq \frac{9}{8}(Mk_{D'}(x', y') + C) \leq \frac{9}{8}(M + C).$$

Then, by (2.3), we have

$$\log \frac{d_{D'}(x')}{d_{D'}(x'_0)} \leq k_{D'}(x', x'_0) \leq Mk_D(x, x_0) + C \leq \frac{9}{8}M(M + C) + C,$$

which implies

$$d_{D'}(x') \leq e^{\frac{9}{8}M(M+C)+C} d_{D'}(x'_0).$$

By Theorem 2.10(2), we have

$$\text{diam}(\gamma') \leq c_2 \max\{|x' - y'|, 2e^{h_1} \min\{d_{D'}(x'), d_{D'}(y')\}\}.$$

This, together with our assumptions $|x' - y'| \leq \frac{1}{2}d_{D'}(x')$ and $d_{D'}(x') \leq d_{D'}(y')$, leads to the following estimate

$$\text{diam}(\gamma') \leq 2c_2 e^{h_1} d_{D'}(x') \leq 2c_2 e^{h_1} e^{\frac{9}{8}M(M+C)+C} d_{D'}(x'_0), \quad (2.4)$$

where $c_2 = c_2(c, c_1, h_1)$ is the constant given by Theorem 2.10.

Next, we consider the latter case, that is, $|x' - y'| \geq \frac{1}{2}d_{D'}(x')$. Since $|x' - x'_0| = \frac{1}{2}|x' - y'|$, we have

$$\min\{\text{diam}(\gamma'[x', x'_0]), \text{diam}(\gamma'[y', x'_0])\} \geq \frac{1}{2}|x' - y'|.$$

It follows from this estimate and Theorem 2.10(2) that

$$\text{diam}(\gamma') \leq 4c_2 e^{h_1} |x' - y'| \leq 8c_2 e^{h_1} \min\{\text{diam}(\gamma'[x', x'_0]), \text{diam}(\gamma'[y', x'_0])\}.$$

This, together with Theorem 2.10(1), gives

$$\text{diam}(\gamma') \leq 8c_2^2 e^{h_1} d_{D'}(x'_0). \quad (2.5)$$

Then the desired estimate $\text{diam}(\gamma') \leq c_3 d_{D'}(x'_0)$ follows from (2.4) and (2.5). \square

Remark 2.13. In the rest of the paper, we assume that

$$c_3 = 8c_2^2 e^{2c_1 + 3h_1}.$$

We close this section with the following metric characterization of uniform domains in E , which we shall apply in our later proofs.

Theorem 2.14 ([33, Theorem 6.16]). *For a domain $D \subsetneq E$, the following statements are equivalent:*

- (1) D is a c -uniform domain;
- (2) For any $x, y \in D$, $k_D(x, y) \leq c_4 \log \left(1 + \frac{|x-y|}{\min\{d_D(x), d_D(y)\}} \right)$,

where the constants c and c_4 depend only on each other.

Remark 2.15. In the rest of the paper, we assume that

$$c_4 = c_4(c) \geq 8.$$

2.4. Gromov hyperbolicity and Rips spaces. There are many equivalent definitions of Gromov hyperbolicity and we shall mainly follow the one used in [37].

Definition 2.16. A domain $D \subsetneq E$ is called δ -Gromov hyperbolic, $\delta > 0$, if (D, k_D) is δ -Gromov hyperbolic. This means that for all $x, y, z, p \in D$,

$$(x|z)_p \geq \min\{(x|y)_p, (y|z)_p\} - \delta,$$

where $(x|y)_p$ is the Gromov product defined by

$$2(x|y)_p = k_D(p, x) + k_D(p, y) - k_D(x, y).$$

It is well-known that every inner uniform domain is Gromov hyperbolic; see [36, Remark 2.16] or [5, Theorem 1.11]. Since Gromov hyperbolicity is preserved by CQH mappings [37, Theorem 3.18], it follows that every quasiconformal image of an inner uniform domain in \mathbb{R}^n is Gromov hyperbolic. In particular, this implies by the Riemann mapping theorem that every simply connected proper subdomain of the complex plane is Gromov hyperbolic.

Lemma 2.17. *Suppose that D is δ -Gromov hyperbolic, and $x \neq y \in D$. Then there exists a constant $v_0 = v_0(c_0, \delta) > 0$ such that*

- for any $\gamma \in \Lambda_{xy}^\lambda$, any $\beta \in Q_{xy}^{c_0}$ and any $u_1 \in \gamma$, there is $u_2 \in \beta$ such that $k_D(u_1, u_2) \leq v_0$;
- for any $\gamma \in \Lambda_{xy}^\lambda$, any $\beta \in Q_{xy}^{c_0}$ and any $v_1 \in \beta$, there is $v_2 \in \gamma$ such that $k_D(v_1, v_2) \leq v_0$.

Proof. By [37, Theorem 3.11], we know that in a δ -Gromov hyperbolic domain, if an h -short arc γ and a c -quasigeodesic β have the same end points, then the Hausdorff distance (with respect to the quasihyperbolic metric) between these two arcs is bounded by a constant which depends only on c , h and δ . Then the conclusions follow from the above fact by noticing Lemma 2.6. \square

Remark 2.18. In the rest of the paper, we assume that

$$v_0 = v_0(c_0, \delta) \geq 16.$$

Let x_1, x_2 and x_3 be a triple of points in D . For each $i \in \{1, 2, 3\}$, let α_i denote an arc joining x_i and x_{i+1} , where $x_4 = x_1$. We write $\Delta = (\alpha_1, \alpha_2, \alpha_3)$ to represent a *triangle* in D with sides $\alpha_1, \alpha_2, \alpha_3$ and the points x_1, x_2, x_3 are called the *vertices* of Δ . A triangle Δ in D is called *h-short* if all its three sides are *h-short*.

The following result is a direct consequence of Lemma 2.6.

Lemma 2.19. *Let x_1, x_2 and x_3 be a triple of points in D . For any λ -curves $\gamma_1 \in \Lambda_{x_1x_2}^\lambda$, $\gamma_2 \in \Lambda_{x_2x_3}^\lambda$ and $\gamma_3 \in \Lambda_{x_3x_1}^\lambda$, the triangle $\Delta = (\gamma_1, \gamma_2, \gamma_3)$ is λ -short.*

Next, we recall the definition of Rips space introduced in [37, Section 2.26]. In what follows, we use the notation $k_D(w, \alpha)$ to denote the quasihyperbolic distance between the point $w \in D$ and the curve α , i.e., $k_D(w, \alpha) = \inf\{k_D(w, v) : v \in \alpha\}$.

Definition 2.20 (Rips space). Let $h \geq 0$ be a constant. If there is a constant $C > 0$ such that for every h -short triangle $\Delta = (\alpha_1, \alpha_2, \alpha_3)$ in D and for all $w \in \alpha_i$, $i \in \{1, 2, 3\}$, with $\alpha_4 = \alpha_1$ and $\alpha_5 = \alpha_2$, it holds

$$k_D(w, \alpha_{i+1} \cup \alpha_{i+2}) \leq C,$$

then we call (D, k_D) a (C, h) -Rips space.

The following lemma gives uniform control on distance between “close” points on short triangle in a Rips space.

Lemma 2.21. *Suppose that (D, k_D) is a (C, h) -Rips space. Then for every h -short triangle $\Delta = (\alpha_1, \alpha_2, \alpha_3)$ in D , and for each $i \in \{1, 2, 3\}$, there is $w_i \in \alpha_i$ such that for any $i \neq j \in \{1, 2, 3\}$,*

$$k_D(w_i, w_j) \leq 3C.$$

Proof. Assume that the vertices of $\Delta = (\alpha_1, \alpha_2, \alpha_3)$ are x_1, x_2 and x_3 such that for each $i \in \{1, 2, 3\}$, α_i connects x_i and x_{i+1} , where $x_4 = x_1$. We consider two cases.

Case 1. For any $z \in \alpha_1$, $k_D(z, \alpha_2) \leq C$.

Since Δ is a triangle, there are $w_1 \in \alpha_1 \setminus \{x_1, x_2\}$ and $w_3 \in \alpha_3 \setminus \{x_1, x_3\}$ such that

$$k_D(w_1, w_3) < C.$$

The assumption in this case ensures that there is $w_2 \in \alpha_2 \setminus \{x_2, x_3\}$ such that

$$k_D(w_1, w_2) < 2C.$$

These yield that

$$k_D(w_2, w_3) \leq k_D(w_1, w_2) + k_D(w_1, w_3) \leq 3C.$$

Clearly, w_1, w_2 and w_3 are the desired points as claimed in the lemma.

Case 2. There is $w_0 \in \alpha_1$ such that $k_D(w_0, \alpha_2) > C$.

Since $k_D(x_2, \alpha_2) = 0$, the assumption in this case implies that there are u_1 and $u_2 \in \alpha_1[x_2, w_0]$ such that for any $u \in \alpha_1[x_2, u_1]$, $k_D(u, \alpha_2) \leq C$ and for any $w \in \alpha_1[u_1, u_2] \setminus \{u_1\}$, $k_D(w, \alpha_2) > C$. Then there is $w_1 \in \alpha_1[u_1, u_2] \setminus \{u_1, w_0\}$ such that $k_D(w_1, \alpha_2) \leq \frac{4}{3}C$, which implies that there is $w_2 \in \alpha_2 \setminus \{x_2, x_3\}$ such that

$$k_D(w_1, w_2) \leq \frac{5}{3}C.$$

Since $k_D(w_1, \alpha_2 \cup \alpha_3) \leq C$, we see from the fact $k_D(w_1, \alpha_2) > C$ that $k_D(w_1, \alpha_3) \leq C$, which implies that there is $w_3 \in \alpha_3 \setminus \{x_1, x_3\}$ such that

$$k_D(w_1, w_3) \leq \frac{4}{3}C.$$

By triangle inequality, we have

$$k_D(w_2, w_3) \leq k_D(w_2, w_1) + k_D(w_1, w_3) \leq 3C.$$

These show that the points w_1 , w_2 and w_3 satisfy our requirements and hence the proof is complete. \square

The next result would be useful in the proofs of our later results.

Lemma 2.22. *The following statements hold.*

- (1) *If D is δ -Gromov hyperbolic, then (D, k_D) is a (μ_0, ρ_0) -Rips space with $\mu_0 = \mu_0(\delta)$ and $\rho_0 = \rho_0(\delta)$;*
- (2) *If $f : D \rightarrow D'$ is (M, C) -CQH and D' is a c -uniform domain, then (D, k_D) is a (μ_0, ρ_0) -Rips space with $\mu_0 = \mu_0(c, C, M)$ and $\rho_0 = \rho_0(c, C, M)$. In particular, if D' is c -uniform, then it is a (μ_0, ρ_0) -Rips space with $\mu_0 = \mu_0(c)$ and $\rho_0 = \rho_0(c)$.*

Proof. The first assertion follows from [37, Theorem 2.35].

For the second assertion, note first that by [36, Theorem 2.12], a uniform domain is quantitatively Gromov hyperbolic in the quasihyperbolic metric, and so $(D', k_{D'})$ is δ' -Gromov hyperbolic with $\delta' = \delta'(c)$. Next, by [37, Theorem 3.18], Gromov hyperbolicity is quantitatively preserved by CQH mappings. Consequently, (D, k_D) is δ -Gromov hyperbolic with $\delta = \delta(c, C, M)$. The claim follows then from the first assertion. \square

Remark 2.23. In the rest of the paper, we assume that

$$\mu_0 = \mu_0(c, C, M, \delta) \geq 16 \quad \text{and} \quad \rho_0 = \rho_0(c, C, M, \delta) > 0.$$




FIGURE 1. Illustration of Lemma 2.24

Now, we prove two elementary estimates on distance between certain points in a Rips space.

Lemma 2.24. *Suppose that (D, k_D) is a (μ_0, ρ_0) -Rips space. For each $i \in \{1, 2, 3\}$, let $\{u_i, v_i\}$ denote a pair of points in D , $\gamma_i \in \Lambda_{u_i v_i}^\lambda$ and let $x_i \in \gamma_i \cap \gamma_{i+1}$, where $\gamma_4 = \gamma_1$. Then for each $i \in \{1, 2, 3\}$, there is $w_i \in \gamma_{i+1}[x_i, x_{i+1}]$ such that for any $r \neq s \in \{1, 2, 3\}$,*

$$k_D(w_r, w_s) \leq \mu_1,$$

where $x_4 = x_1$ and $\mu_1 = 3\mu_0$.

Proof. This is a direct consequence of Lemmas 2.6, 2.7, 2.19 and 2.21; see Figure 1. \square

Lemma 2.25. *Suppose that (D, k_D) is a (μ_0, ρ_0) -Rips space. For each $i \in \{1, 2\}$, let $\{u_i, v_i\}$ be a pair of points in D , $\gamma_i \in \Lambda_{u_i v_i}^\lambda$ and let $x \in \gamma_1 \cap \gamma_2$, $y \in \gamma_1[x, v_1]$ and $z \in \gamma_2[x, v_2]$. If $k_D(y, z) \leq \tau$, then for any $w \in \gamma_1[x, y]$, there exists $u \in \gamma_2[x, z]$ such that*

$$k_D(w, u) \leq \mu_0 + 2\lambda + \tau.$$

Proof. Let $w \in \gamma_1[x, y]$ and fix any $\gamma_3 \in \Lambda_{yz}^\lambda$. Since (D, k_D) is a (μ_0, ρ_0) -Rips space, it follows from Lemmas 2.6 and 2.7 that there exists $v \in \gamma_2[x, z] \cup \gamma_3$ satisfying

$$k_D(w, v) \leq \lambda + \mu_0.$$

If $v \in \gamma_2[x, z]$, then we simply take $u = v$. If $v \in \gamma_3$, then we take $u = z$. It follows from Lemma 2.6 and the previous estimate that

$$\begin{aligned} k_D(w, u) &\leq k_D(w, v) + \ell_k(\gamma_3) \leq k_D(w, v) + k_D(y, z) + \lambda \\ &\leq \mu_0 + 2\lambda + \tau. \end{aligned}$$

In either cases, we see that the conclusion of lemma holds. The proof is thus complete. \square

The next result concerns a similar estimate in a Gromov hyperbolic space.

Lemma 2.26. *Suppose that D is δ -Gromov hyperbolic, and $x, y, z \in D$ are distinct points. Then there exists a constant $\mu_2 = \mu_2(\delta, c_0)$ such that for each $\gamma_1 \in Q_{xy}^{c_0}$, $\gamma_2 \in Q_{yz}^{c_0}$, $\gamma_3 \in Q_{xz}^{c_0}$ and each $u \in \gamma_1$, there is $v \in \gamma_2 \cup \gamma_3$ satisfying the inequality*

$$k_D(u, v) \leq \mu_2.$$

Indeed, we may take $\mu_2 = \frac{1}{4} + \mu_0 + 2v_0$ with μ_0 and v_0 given by Lemma 2.22 and Lemma 2.17, respectively.

Proof. Fix $u \in \gamma_1$, $\beta_1 \in \Lambda_{xy}^\lambda$, $\beta_2 \in \Lambda_{yz}^\lambda$ and $\beta_3 \in \Lambda_{xz}^\lambda$ with $\lambda = \frac{1}{4}$. Then Lemma 2.17 shows that there is $u_1 \in \beta_1$ such that

$$k_D(u, u_1) \leq v_0, \tag{2.6}$$

where $v_0 = v_0(c_0, \delta)$ is a uniform constant depending only on c_0 and δ .

By Lemma 2.22(1), (D, k_D) is a (μ_0, ρ_0) -Rips space with $\mu_0 = \mu_0(\delta)$ and $\rho_0 = \rho_0(\delta)$. Then Lemma 2.6 implies that there is $u_2 \in \beta_2 \cup \beta_3$ such that

$$k_D(u_1, u_2) \leq \frac{1}{4} + \mu_0. \tag{2.7}$$

Without loss of generality, we may assume $u_2 \in \beta_3$. Applying Lemma 2.17 once again, we find $v \in \gamma_3$ such that

$$k_D(u_2, v) \leq v_0,$$

which, together with (2.6) and (2.7), yields

$$k_D(u, v) \leq k_D(u, u_1) + k_D(u_1, u_2) + k_D(u_2, v) \leq \frac{1}{4} + \mu_0 + 2v_0.$$

This completes the proof. \square

We now establish the following technical lemma, which compares the length of quasigeodesics with one common end point in Gromov hyperbolic spaces.

Lemma 2.27. *Suppose that D is δ -Gromov hyperbolic, x_1, x_2 and x_3 are points in D , and that $\mu_2 = \mu_2(\delta, c_0)$ is the constant given by Lemma 2.26. Suppose further that there is a constant $\tau \geq 0$ such that $k_D(x_2, x_3) \leq \tau$. If*

$$\min \{k_D(x_1, x_2), k_D(x_1, x_3)\} > 1,$$

then for any $\vartheta \in Q_{x_1 x_2}^{c_0}$ and $\gamma \in Q_{x_1 x_3}^{c_0}$, we have

$$\max \{\ell(\vartheta), \ell(\gamma)\} \leq 4e^{2c_0^2(2\tau+3\mu_2)+\frac{4}{3}\mu_2} \min \{\ell(\vartheta), \ell(\gamma)\}.$$

Proof. Fix $\vartheta \in Q_{x_1 x_2}^{c_0}$ and $\gamma \in Q_{x_1 x_3}^{c_0}$. If $\ell(\vartheta) = \ell(\gamma)$, then there is nothing to prove. Otherwise, we may assume without loss of generality that

$$\ell(\vartheta) > \ell(\gamma).$$

Under this assumption, to prove the lemma, we only need to verify the following inequality:

$$\ell(\vartheta) \leq 4e^{2c_0^2(2\tau+3\mu_2)+\frac{4}{3}\mu_2} \ell(\gamma), \quad (2.8)$$

where $\mu_2 = \mu_2(\delta, c_0)$ is the constant given by Lemma 2.26.

In the following, we divide the discussions into two cases: $k_D(x_1, x_2) \leq 2c_0(2\tau + 3\mu_2)$ and $k_D(x_1, x_2) > 2c_0(2\tau + 3\mu_2)$. Since $k_D(x_1, x_3) > 1$, it follows from Lemma 2.2(2) that

$$d_D(x_1) \leq 2\ell(\gamma). \quad (2.9)$$

In the first case, (2.2) gives

$$2c_0^2(2\tau + 3\mu_2) \geq c_0 k_D(x_1, x_2) \geq \ell_k(\vartheta) \geq \log \left(1 + \frac{\ell(\vartheta)}{\min\{d_D(x_1), d_D(x_2)\}} \right).$$

This, together with (2.9), implies

$$\ell(\vartheta) \leq e^{2c_0^2(2\tau+3\mu_2)} \min\{d_D(x_1), d_D(x_2)\} \leq 2e^{2c_0^2(2\tau+3\mu_2)} \ell(\gamma). \quad (2.10)$$

In the second case, that is, $k_D(x_1, x_2) > 2c_0(2\tau + 3\mu_2)$, we need some preparation. First, we determine a partition of the arc ϑ . For this, we denote x_2 by y_0 . Clearly, there exists a finite sequence of points $\{y_i\}_{i=0}^m$ in ϑ with $m \geq 3$ such that for each $i \in \{1, \dots, m-1\}$, $y_i \in \vartheta[y_{i-1}, x_1]$,

$$k_D(y_{i-1}, y_i) = c_0(2\tau + 3\mu_2) \quad \text{and} \quad 0 < k_D(y_{m-1}, y_m) \leq c_0(2\tau + 3\mu_2), \quad (2.11)$$

where $y_m = x_1$; see Figure 2.

By (2.2), for each $i \in \{1, \dots, m\}$, we have

$$\log \left(1 + \frac{\ell(\vartheta[y_{i-1}, y_i])}{\min\{d_D(y_{i-1}), d_D(y_i)\}} \right) \leq \ell_k(\vartheta[y_{i-1}, y_i]) \leq c_0 k_D(y_{i-1}, y_i) \leq c_0^2(2\tau + 3\mu_2),$$

from which it follows

$$\ell(\vartheta[y_{i-1}, y_i]) \leq e^{c_0^2(2\tau+3\mu_2)} \min\{d_D(y_{i-1}), d_D(y_i)\}. \quad (2.12)$$




FIGURE 2. Illustration for the proof of second case in Lemma 2.27

Next, we are going to find a partition of γ based on $\{y_i\}_{i=1}^{m-1}$. For this, we fix $\beta \in Q_{x_2 x_3}^{c_0}$ and select $p \in \beta$ such that

$$k_D(y_1, \beta) \leq k_D(y_1, p) < k_D(y_1, \beta) + \frac{7}{8}\tau, \quad (2.13)$$

where $k_D(y_1, \beta)$ is the quasihyperbolic distance from y_1 to β .

From the assumption of $k_D(x_2, x_3) < \tau$, we note

$$\ell_k(\beta) \leq c_0 k_D(x_2, x_3) < c_0 \tau,$$

and thus

$$k_D(y_1, p) \geq k_D(y_1, y_0) - k_D(y_0, p) \geq k_D(y_1, y_0) - c_0 \tau.$$

Then we infer from (2.11) and (2.13) that

$$k_D(y_1, \beta) \geq k_D(y_1, p) - \frac{7}{8}\tau \geq c_0(3\mu_2 + \tau) - \frac{7}{8}\tau > \mu_2. \quad (2.14)$$

Since D is δ -Gromov hyperbolic, we know from Lemma 2.26 that there is $q_1 \in \gamma \cup \beta$ such that $k_D(y_1, q_1) \leq \mu_2$. By (2.14), we see that $q_1 \in \gamma$. This implies that

$$k_D(y_1, \gamma) \leq k_D(y_1, q_1) \leq \mu_2. \quad (2.15)$$

Now, we are ready to determine a point in γ based on y_1 . Denote x_3 by w_0 , and let $w_1 \neq w_0$ be a point in γ (see Figure 2) such that

$$k_D(y_1, w_1) < \frac{4}{3}k_D(y_1, \gamma).$$

Then it follows from (2.15) that

$$k_D(y_1, w_1) \leq \frac{4}{3}\mu_2, \quad (2.16)$$

which is the point we need.

To find the next point in γ based on y_2 , we fix $\beta_1 \in Q_{w_1 y_1}^{c_0}$ and derive a lower bound for $k_D(y_2, \beta_1)$ as follows. Since $\ell_k(\beta_1) \leq c_0 k_D(y_1, w_1)$, we infer from (2.11) and (2.16) that

$$k_D(y_2, \beta_1) \geq k_D(y_1, y_2) - \ell_k(\beta_1) \geq c_0\left(\frac{5}{3}\mu_2 + 2\tau\right) > \mu_2. \quad (2.17)$$

Once again, Lemma 2.26 implies that there is $q_2 \in \gamma[x_1, w_1] \cup \beta_1$ such that

$$k_D(y_2, q_2) \leq \mu_2.$$

Then (2.17) ensures that $q_2 \in \gamma[x_1, w_1]$, and thus,

$$k_D(y_2, \gamma[x_1, w_1]) \leq k_D(y_2, q_2) \leq \mu_2. \quad (2.18)$$

Let w_2 be a point in $\gamma[x_1, w_1] \setminus \{w_1\}$ (see Figure 2) such that

$$k_D(y_2, w_2) < \frac{4}{3}k_D(y_2, \gamma[x_1, w_1]).$$

Then it follows from (2.18) that

$$k_D(y_2, w_2) \leq \frac{4}{3}\mu_2, \quad (2.19)$$

which is the point we want.

By repeating this procedure, we may find a finite sequence of points $\{w_i\}_{i=1}^{m-1}$ in γ (see Figure 2) such that for each $i \in \{1, \dots, m-1\}$,

$$w_i \in \gamma[x_1, w_{i-1}] \setminus \{w_{i-1}\} \text{ and } k_D(y_i, w_i) \leq \frac{4}{3}\mu_2. \quad (2.20)$$

In the following, we prove that for each $i \in \{1, \dots, m-1\}$,

$$\ell(\vartheta[y_{i-1}, y_i]) \leq 2e^{c_0^2(2\tau+3\mu_2)+\frac{4}{3}\mu_2}\ell(\gamma[w_{i-1}, w_i]). \quad (2.21)$$

First, (2.20), together with (2.3), ensures that for each $i \in \{1, \dots, m-1\}$, we have

$$\left| \log \frac{d_D(y_i)}{d_D(w_i)} \right| \leq k_D(y_i, w_i) \leq \frac{4}{3}\mu_2,$$

which gives

$$d_D(y_i) \leq e^{\frac{4}{3}\mu_2} d_D(w_i). \quad (2.22)$$

Since

$$k_D(w_{i-1}, w_i) \geq k_D(y_{i-1}, y_i) - k_D(y_{i-1}, w_{i-1}) - k_D(y_i, w_i),$$

we deduce from (2.11), (2.20) and the assumption $k_D(y_0, w_0) = k_D(x_2, x_3) < \tau$ that

$$k_D(w_{i-1}, w_i) \geq \begin{cases} (2c_0 - 1)\tau + \frac{9c_0 - 4}{3}\mu_2, & \text{when } i = 1, \\ 2c_0\tau + \frac{9c_0 - 8}{3}\mu_2, & \text{when } 2 \leq i \leq m-1. \end{cases}$$

Since our choice of c_0 and μ_2 in Section 2.1 implies that

$$k_D(w_{i-1}, w_i) > 1,$$

we know from Lemma 2.2(2) that

$$d_D(w_i) \leq 2\ell(\gamma[w_{i-1}, w_i]),$$

and so, it follows from (2.12) and (2.22) that for each $i \in \{1, \dots, m-1\}$,

$$\ell(\vartheta[y_{i-1}, y_i]) \leq 2e^{c_0^2(2\tau+3\mu_2)+\frac{4}{3}\mu_2}\ell(\gamma[w_{i-1}, w_i]),$$

which proves (2.21). Moreover, by (2.9) and (2.12), we have

$$\ell(\vartheta[y_{m-1}, y_m]) \leq 2e^{c_0^2(2\tau+3\mu_2)}\ell(\gamma). \quad (2.23)$$

Now, we are ready to show (2.8). Since

$$\ell(\vartheta) = \sum_{i=1}^{m-1} \ell(\vartheta[y_{i-1}, y_i]) + \ell(\vartheta[y_{m-1}, y_m]),$$

it follows from (2.21) and (2.23) that

$$\ell(\vartheta) \leq 4e^{c_0^2(2\tau+3\mu_2)+\frac{4}{3}\mu_2} \ell(\gamma). \quad (2.24)$$

In either cases, we conclude from (2.10) and (2.24) that the desired inequality (2.8) holds, and hence, the proof is complete. \square

Based on Lemma 2.27, we now establish the following useful length comparison theorem for quasigeodesics with the same end points in Gromov hyperbolic spaces.

Theorem 2.28. *Suppose that D is δ -Gromov hyperbolic, x_1 and x_2 are points in D . Then for any $\vartheta, \gamma \in Q_{x_1 x_2}^{c_0}$, it holds*

$$\max\{\ell(\vartheta), \ell(\gamma)\} \leq 4e^{8c_0^2\mu_2} \min\{\ell(\vartheta), \ell(\gamma)\}.$$

Proof. Without loss of generality, we may assume that $\min\{d_D(x_1), d_D(x_2)\} = d_D(x_1)$ and $\max\{\ell(\vartheta), \ell(\gamma)\} = \ell(\vartheta)$. Note that if $k_D(x_1, x_2) \leq 1$, then we may get from (2.2) that

$$\log\left(1 + \frac{\ell(\vartheta)}{d_D(x_1)}\right) \leq \ell_k(\vartheta) \leq c_0 k_D(x_1, x_2) \leq c_0,$$

which gives

$$\ell(\vartheta) \leq (e^{c_0} - 1)d_D(x_1). \quad (2.25)$$

On the other hand, we know from (2.3) that

$$\log\left(1 + \frac{|x_1 - x_2|}{d_D(x_1)}\right) \leq k_D(x_1, x_2) \leq 1,$$

which implies that

$$|x_1 - x_2| \leq (e - 1)d_D(x_1).$$

Clearly, for each $x \in [x_1, x_2]$, it holds

$$d_D(x) \geq \min\{d_D(x_1), d_D(x_2)\} - \frac{1}{2}|x_1 - x_2| = d_D(x_1) - \frac{1}{2}|x_1 - x_2| \geq \frac{3-e}{2}d_D(x_1),$$

and so we have

$$k_D(x_1, x_2) \leq \int_{x \in [x_1, x_2]} \frac{|dx|}{d_D(x)} \leq \frac{2|x_1 - x_2|}{(3-e)d_D(x_1)}.$$

Combining this with (2.25) gives

$$\frac{\ell(\vartheta)}{e^{c_0}d_D(x_1)} \leq \ell_k(\vartheta) \leq c_0 k_D(x_1, x_2) \leq \frac{2c_0|x_1 - x_2|}{(3-e)d_D(x_1)},$$

from which it follows

$$\ell(\vartheta) \leq \frac{2c_0e^{c_0}}{3-e}|x_1 - x_2| \leq \frac{2c_0e^{c_0}}{3-e}\ell(\gamma).$$

Thus, we only need to consider the case $k_D(x, y) > 1$. In this case, the result follows directly from Lemmas 2.27 (applied with $x_2 = x_3$). \square

As an application of Theorem 2.28, we prove a technical lemma.

Lemma 2.29. *Suppose that D is δ -Gromov hyperbolic. For $u, v \in D$, let $\beta \subset D$ be a curve connecting u and v , and suppose that for all $w \in \beta$,*

$$d_D(w) \leq 2\ell(\beta). \quad (2.26)$$

Let $u_1 \in \beta$ and $v_1 \in \beta[u_1, v]$. For any λ -curve $\gamma_1 \in \Lambda_{u_1 v_1}^\lambda$, if $\ell(\gamma_1) \geq \mu_9^{\frac{1}{2}}\ell(\beta)$, then there is $w_0 \in \beta[u_1, v_1]$ such that

(1) *for any λ -curve $\gamma_2 \in \Lambda_{u_1 w_0}^\lambda$,*

$$\mu_9^{\frac{1}{4}}\ell(\beta) \leq \ell(\gamma_2) \leq \mu_9^{\frac{1}{2}}\ell(\beta).$$

(2) *for any $z \in \beta[w_0, v_1] \setminus \{w_0\}$, and for any λ -curve $\gamma_3 \in \Lambda_{u_1 z}^\lambda$ satisfying*

$$\ell(\gamma_3) > \mu_9^{\frac{1}{4}}\ell(\beta).$$

Proof. For each $w \in \beta[u_1, v_1]$, let Λ_w be a subclass of $\Lambda_{u_1 w}^\lambda$ consisting of $\gamma_w \in \Lambda_{u_1 w}^\lambda$ with

$$\ell(\gamma_w) \leq 2 \inf_{\gamma \in \Lambda_{u_1 w}^\lambda} \{\ell(\gamma)\}.$$

Then there exist $w_1 \in \beta[u_1, v_1]$ and $\gamma_{w_1} \in \Lambda_{w_1}$ such that

(i) $\ell(\gamma_{w_1}) \leq 2\mu_9^{\frac{1}{4}}\ell(\beta)$;

(ii) for any $w \in \beta[w_1, v_1] \setminus \{w_1\}$, there exists $\gamma_w \in \Lambda_w$ with $\ell(\gamma_w) > 2\mu_9^{\frac{1}{4}}\ell(\beta)$.

Indeed, if not, then by Theorem 2.28, $\ell(\gamma_1) \leq 8e^{11\mu_2}\mu_9^{\frac{1}{4}}\ell(\beta) < \mu_9^{\frac{1}{2}}\ell(\beta)$, contradicting with our assumption.

Now select $w_0 \in \beta[w_1, v_1]$ such that

$$0 < k_D(w_0, w_1) < 1 \quad (2.27)$$

and let $\gamma_{w_0} \in \Lambda_{w_0}$ be given by (ii). Then we deduce from Lemma 2.9(1), (2.26) and (ii) that

$$\begin{aligned} k_D(u_1, w_0) &\geq \frac{8}{9}\ell_k(\gamma_{w_0}) \geq \frac{8}{9} \log \left(1 + \frac{\ell(\gamma_{w_0})}{\min\{d_D(u_1), d_D(w_0)\}} \right) \\ &\geq \frac{8}{9} \log \left(1 + \frac{\ell(\gamma_{w_0})}{2\ell(\beta)} \right) \geq \frac{8}{9} \log \left(1 + \mu_9^{\frac{1}{4}} \right) > 1. \end{aligned}$$

Moreover, by (2.27), we obtain that

$$k_D(u_1, w_1) \geq k_D(u_1, w_0) - k_D(w_1, w_0) \geq \frac{8}{9} \log \left(1 + \mu_9^{\frac{1}{4}} \right) - 1 > 1.$$

The above inequalities yield

$$\min\{k_D(u_1, w_0), k_D(u_1, w_1)\} > 1. \quad (2.28)$$

Since D is δ -Gromov hyperbolic, we infer from (2.27), (2.28), (i) and Lemma 2.27 (replacing x_1, x_2 and x_3 with u_1, w_1 and w_0 , respectively) that

$$\mu_9^{\frac{1}{4}}\ell(\beta) \leq \frac{1}{2}\ell(\gamma_{w_0}) \leq \ell(\gamma_2) \leq e^{9\mu_2}\ell(\gamma_{w_1}) < \mu_9^{\frac{1}{2}}\ell(\beta),$$

which proves (1).

For the second assertion, simply note that by (ii), it holds

$$\mu_9^{\frac{1}{4}}\ell(\beta) < \frac{1}{2}\ell(\gamma_z) \leq \ell(\gamma_3).$$

The proof is thus complete.

□

3. COMPACTNESS OF λ -PAIRS WITH \mathfrak{C} -CONDITION

3.1. **Main result.** Throughout this section, we make the following standing assumptions.

Standing Assumptions. we assume that $f : D \rightarrow D'$ is (M, C) -CQH and D' is a c -uniform domain. Let $\omega \neq \varpi \in D$, and let $\gamma \in \Lambda_{\omega\varpi}^\lambda$ be a fixed element with $\lambda = \frac{1}{4}$.

For the statement of our main result of this section, we need to recall the following definitions.

Definition 3.1 (λ -pair and \mathfrak{C} -Condition). A λ -curve $\gamma \in \Lambda_{uv}^\lambda$ and a curve β in D is called a λ -pair, written as (γ, β) , if the end points of β are u and v . For $u_1 \neq v_1 \in \beta$, the pair $(\gamma[u_1v_1], \beta[u_1, v_1])$ is called a λ -subpair of (γ, β) . A λ -pair (γ, β) is said to satisfy \mathfrak{C} -Condition if there is a constant $\mathfrak{C} > 0$ such that $\ell(\gamma) \geq \mathfrak{C}\ell(\beta)$.

The aim of this section is to prove the following theorem, which plays a key role for the discussions in the next section. In the statement below, the constants μ_5 and μ_6 are defined in Section 2.1.

Theorem 3.2. Suppose that \mathfrak{l} is an arc in D such that (γ, \mathfrak{l}) is a λ -pair and satisfies \mathfrak{C} -Condition (see Figure 3). If $\mathfrak{C} \geq \frac{4}{5}\mu_5$, then there exist points $\omega_1 \neq \varpi_1 \in \mathfrak{l}$ such that for any λ -curve $\gamma[\omega_1\varpi_1] \in \Lambda_{\omega_1\varpi_1}^\lambda$, the λ -subpair $(\gamma[\omega_1\varpi_1], \mathfrak{l}[\omega_1, \varpi_1])$ of (γ, \mathfrak{l}) satisfies $\mu_6\mathfrak{C}$ -Condition, i.e.,

$$\ell(\gamma[\omega_1\varpi_1]) \geq \mu_6\mathfrak{C}\ell(\mathfrak{l}[\omega_1, \varpi_1]).$$




FIGURE 3. Illustration for the proof of Theorem 3.2

The proof of Theorem 3.2 depends on Theorem 3.3 below, which relates the length of a λ -curve and the inner distance between its end points. In the statement below, the constants μ_7 and μ_8 are again defined in Section 2.1.

Theorem 3.3. *Suppose that $f : D \rightarrow D'$ is (M, C) -CQH and D' is a c -uniform domain. For any distinct points $x \neq y \in D$ and any λ -curve $\gamma \in \Lambda_{xy}^\lambda$, if there is $x_0 \in \gamma$ such that*

$$|x' - x'_0| = \frac{1}{2}|x' - y'| \quad \text{and} \quad \ell(\gamma) \leq \mu_7 d_D(x_0), \quad (3.1)$$

then it holds

$$\ell(\gamma) \leq \mu_8 \sigma_D(x, y).$$

As the proof of Theorem 3.3 is rather technical and lengthy, we move it to Section 5. An illustration picture of this result can be found in Figure 4.

3.2. Auxiliary lemmas. In this section, under the **Standing Assumptions** in Section 3.1 and assumptions of Theorem 3.2, we shall establish a series of auxiliary results that are necessary for the proof of Theorem 3.2.

For the convenience of later references, we collect some useful properties in Proposition 3.4 below, where (P_1) comes from Lemma 2.9(1), (P_2) from Lemma 2.9(2), (P_3) from [36, Theorem 2.12] and [37, Theorem 3.18], and finally (P_4) from Lemma 2.22(2).

Proposition 3.4. (P_1) *For any $x \neq y \in D$, every λ -curve $\gamma \in \Lambda_{xy}^\lambda$ is a $(c_0, 0)$ -solid arc and belongs to $Q_{xy}^{c_0}$ with $c_0 = \frac{9}{8}$;*

(P_2) *For any λ -curve $\gamma \in \Lambda_{uv}^\lambda$, its image γ' under f is (c_1, h_1) -solid, where $c_1 = \frac{9}{4}M(M+1)$ and $h_1 = C(\frac{9}{4}M+1)$;*

(P_3) *Both D and D' are δ -Gromov hyperbolic, where $\delta = \delta(c, C, M)$;*

(P_4) *Both (D, k_D) and $(D', k_{D'})$ are (μ_0, ρ_0) -Rips spaces.*

Throughout this section, we let $c_2 = c_2(c, c_1, h_1)$ be the constant given by Theorem 2.10 and $c_4 = c_4(c)$ be a constant given by Theorem 2.14.

The first lemma is an easy consequence of Lemma 2.8.

Lemma 3.5. *For any $w \in \mathfrak{l}$, $d_D(w) \leq 2\ell(\mathfrak{l})$.*

Proof. Note that by our choice of constants in Section 2.1, $\frac{4}{5}\mu_5 > 18e^9$ and thus $\ell(\gamma) > \frac{4}{5}\mu_5\ell(\mathfrak{l}) > 18e^9\ell(\mathfrak{l})$. The lemma follows from Lemma 2.8. \square

Lemma 3.6. $\text{diam}(\gamma') \leq 4c_2e^{h_1}|\omega' - \varpi'|$.

Proof. Lemma 3.5 and the assumption $\ell(\gamma) > \mathfrak{C}\ell(\mathfrak{l})$ in Theorem 3.2 yield

$$\ell(\gamma) \geq \frac{\mathfrak{C}}{2}d_D(\omega).$$

Then we get from Lemma 2.6 and (2.2) that

$$k_D(\omega, \varpi) \geq \ell_k(\gamma) - \frac{1}{4} \geq \log\left(1 + \frac{\ell(\gamma)}{d_D(\omega)}\right) - \frac{1}{4} \geq \log\left(1 + \frac{\mathfrak{C}}{2}\right) - \frac{1}{4}.$$

Since f is (M, C) -CQH and $\mathfrak{C} \geq \frac{4}{5}\mu_5$, it follows

$$k_{D'}(\omega', \varpi') \geq \frac{1}{M}(k_D(\omega, \varpi) - C) \geq \frac{1}{M} \log \left(1 + \frac{\mathfrak{C}}{2} \right) - \frac{1+4C}{4M} > 1.$$

Thus an application of Lemma 2.2(1) gives

$$|\omega' - \varpi'| \geq \frac{1}{2} \max\{d_{D'}(\omega'), d_{D'}(\varpi')\}. \quad (3.2)$$

Since γ' is (c_1, h_1) -solid by Proposition 3.4(P_2) and D' is c -uniform, Theorem 2.10(2) gives

$$\text{diam}(\gamma') \leq c_2 \max\{|\omega' - \varpi'|, 2(e^{h_1} - 1) \min\{d_{D'}(\omega'), d_{D'}(\varpi')\}\},$$

where $c_2 = c_2(c, c_1, h_1)$ is from Theorem 2.10. Then (3.2) guarantees that

$$\text{diam}(\gamma') \leq 4c_2 e^{h_1} |\omega' - \varpi'|.$$

The proof is thus complete. \square

Let z_0 (resp. y_0) be the first point on γ along the direction from ω to ϖ (resp. from ϖ to ω) such that

$$|\omega' - z'_0| = \frac{1}{4} \text{diam}(\gamma') \quad (\text{resp. } |\varpi' - y'_0| = \frac{1}{4} \text{diam}(\gamma')); \quad (3.3)$$

see Figure 3 for an illustration. Then, $z_0 \in \gamma[\omega, y_0]$ and $y_0 \in \gamma[\varpi, z_0]$.

Lemma 3.7. (1) For any $z \in \gamma[\omega, z_0]$, $\text{diam}(\gamma'[\omega', z']) \leq c_2 d_{D'}(z')$, and for any $y \in \gamma[\varpi, y_0]$, $\text{diam}(\gamma'[\varpi', y']) \leq c_2 d_{D'}(y')$.

$$(2) \ell(\gamma[z_0, y_0]) \leq (1 + 4c_2)^{c_4 M} e^{\frac{1}{2} + C} \min\{d_D(z_0), d_D(y_0)\}.$$

Proof. (1). We only show the first claim in statement (1), since the proof of second claim is similar. For this, let $z \in \gamma[\omega, z_0]$. Then the choice of z_0 in (3.3) implies that

$$\text{diam}(\gamma'[\omega', z']) \leq 2|\omega' - z'_0| = \frac{1}{2} \text{diam}(\gamma'),$$

and thus,

$$\text{diam}(\gamma'[\varpi', z']) \geq \frac{1}{2} \text{diam}(\gamma') \geq \text{diam}(\gamma'[\omega', z']).$$

Then we know from Theorem 2.10(1) that

$$\text{diam}(\gamma'[\omega', z']) \leq c_2 d_{D'}(z'),$$

which verifies the desired claim.

(2). Since D' is c -uniform, Theorem 2.14 gives

$$\begin{aligned} k_{D'}(z'_0, y'_0) &\leq c_4 \log \left(1 + \frac{|y'_0 - z'_0|}{\min\{d_{D'}(y'_0), d_{D'}(z'_0)\}} \right) \\ &\leq c_4 \log \left(1 + \frac{\text{diam}(\gamma')}{\min\{d_{D'}(y'_0), d_{D'}(z'_0)\}} \right). \end{aligned}$$

By the choice of z_0 and y_0 in (3.3) and statement (1) of Lemma 3.7, we have

$$k_{D'}(z'_0, y'_0) \leq c_4 \log(1 + 4c_2).$$

Since f is (M, C) -CQH, it follows from Lemmas 2.6 and 2.7 that

$$\ell_k(\gamma[z_0, y_0]) - \frac{1}{2} \leq k_D(z_0, y_0) \leq M k_{D'}(z'_0, y'_0) + C \leq c_4 M \log(1 + 4c_2) + C,$$

and thus, we infer from (2.2) that

$$\log \left(1 + \frac{\ell(\gamma[z_0, y_0])}{\min\{d_D(z_0), d_D(y_0)\}} \right) \leq c_4 M \log(1 + 4c_2) + C + \frac{1}{2}.$$

This proves the second statement, and hence, the proof is complete. \square

Without loss of generality, we assume that

$$\ell(\gamma[\omega, z_0]) \geq \ell(\gamma[\varpi, y_0]). \quad (3.4)$$

Under this assumption, we have the following lemma.

Lemma 3.8. $\ell(\gamma) \leq 2 \left(1 + (1 + 4c_2)^{c_4 M} e^{\frac{1}{2} + C} \right) \ell(\gamma[\omega, z_0]).$

Proof. First, we know from (3.4) and Lemma 3.7(2) that

$$\begin{aligned} \ell(\gamma) &= \ell(\gamma[\omega, z_0]) + \ell(\gamma[\varpi, y_0]) + \ell(\gamma[z_0, y_0]) \\ &\leq 2\ell(\gamma[\omega, z_0]) + (1 + 4c_2)^{c_4 M} e^{\frac{1}{2} + C} \min\{d_D(z_0), d_D(y_0)\}. \end{aligned} \quad (3.5)$$

Next, we show the following estimate:

$$d_D(z_0) \leq 2\ell(\gamma[\omega, z_0]). \quad (3.6)$$

Suppose on the contrary that

$$d_D(z_0) > 2\ell(\gamma[\omega, z_0]). \quad (3.7)$$

Then it holds

$$d_D(\omega) \geq d_D(z_0) - \ell(\gamma[\omega, z_0]) \geq \frac{1}{2}d_D(z_0). \quad (3.8)$$

Consequently, (3.5), (3.7) and (3.8) give

$$\ell(\gamma) \leq \left(1 + (1 + 4c_2)^{c_4 M} e^{\frac{1}{2} + C} \right) d_D(z_0) \leq 2 \left(1 + (1 + 4c_2)^{c_4 M} e^{\frac{1}{2} + C} \right) d_D(\omega).$$

On the other hand, since $\ell(\gamma) \geq \frac{4}{5}\mu_5 \ell(\mathfrak{l})$, Lemma 3.5 implies

$$\ell(\gamma) \geq \frac{2}{5}\mu_5 d_D(\omega).$$

This is impossible as our choice of constant in Section 2.1 implies

$$\frac{2}{5}\mu_5 > 2 \left(1 + (1 + 4c_2)^{c_4 M} e^{\frac{1}{2} + C} \right).$$

Thus (3.6) holds and the desired estimate in Lemma 3.8 follows from (3.5) and (3.6). \square

In the following lemmas, it is helpful to see Figure 3 for an intuition.

Lemma 3.9. *Let $z \in \gamma[\omega, z_0]$ such that $k_D(\omega, z) \geq \mu_3$. Then*

(1) *There exists $q \in \mathfrak{l}$ such that $|\omega' - q'| = \mu_3^{-1} \text{diam}(\gamma'[\omega', z'])$, and for any $u \in \mathfrak{l}[\omega, q] \setminus \{q\}$,*

$$|\omega' - u'| < \mu_3^{-1} \text{diam}(\gamma'[\omega', z']).$$

(2) *For any λ -curve $\gamma_0 \in \Lambda_{\omega q}^\lambda$, there exists $q_0 \in \gamma_0$ such that*

(a) *$|\omega' - q'_0| = \frac{1}{4} \text{diam}(\gamma'_0)$, and for any $v \in \gamma_0[\omega, q_0] \setminus \{q_0\}$,*

$$|\omega' - v'| < \frac{1}{4} \text{diam}(\gamma'_0);$$

(b) $\max\{k_D(z, q_0), k_{D'}(z', q'_0)\} \leq c_0 c_2 c_3 c_4^2 \mu_2 (4 + C + h_1 + M)^3$.

Proof. (1). Since $\mu_3 > 4c_4e^{h_1}$, we see from Lemma 3.6 that

$$|\omega' - \varpi'| > \mu_3^{-1} \text{diam}(\gamma'[\omega', z']).$$

This implies that there exists $q \in \mathfrak{l}$ such that

$$|\omega' - q'| = \mu_3^{-1} \text{diam}(\gamma'[\omega', z']),$$

and for any $u \in \mathfrak{l}[\omega, q] \setminus \{q\}$,

$$|\omega' - u'| < \mu_3^{-1} \text{diam}(\gamma'[\omega', z']).$$

This shows statement (1).

(2). Let $\gamma_0 \in \Lambda_{\omega q}^\lambda$ be a fixed element and let $q_0 \in \gamma_0$ be the first point along the direction from ω to q (see Figure 3) such that

$$|\omega' - q'_0| = \frac{1}{4} \text{diam}(\gamma').$$

Then q_0 satisfies all the requirements in (2a).

For the statement (2b), we shall prove the following estimate.

$$|z' - q'_0| \leq c_0^2 c_2^2 c_3 c_4^2 \mu_2 e^{4+C+CM+h_1+M+M^2} \min\{d_{D'}(q'_0), d_{D'}(z')\}. \quad (3.9)$$

Since statements (1) and (2a) ensure that

$$|z' - \omega'| \leq \text{diam}(\gamma'[\omega', z']) = \mu_3 |\omega' - q'| \leq \mu_3 \text{diam}(\gamma'_0) = 4\mu_3 |\omega' - q'_0|,$$

it follows that

$$\begin{aligned} |z' - q'_0| &\leq |z' - \omega'| + |\omega' - q'_0| \leq (1 + 4\mu_3) |\omega' - q'_0| \\ &\leq (1 + 4\mu_3) \text{diam}(\gamma'_0[\omega', q'_0]). \end{aligned} \quad (3.10)$$

Furthermore, the choice of q_0 in (2a) implies that

$$\text{diam}(\gamma'_0[\omega', q'_0]) \leq \text{diam}(\gamma'_0[q', q'_0]).$$

Since by Proposition 3.4(P_2), γ'_0 is a (c_1, h_1) -solid arc, and since D' is c -uniform, we deduce from (3.10) and Theorem 2.10(1) that

$$\begin{aligned} |z' - q'_0| &\leq (1 + 4\mu_3) \min\{\text{diam}(\gamma'_0[\omega', q'_0]), \text{diam}(\gamma'_0[q', q'_0])\} \\ &\leq c_2(1 + 4\mu_3) d_{D'}(q'_0). \end{aligned}$$

To prove (3.9), it suffices to deduce the following estimate:

$$|z' - q'_0| \leq c_2^2 e^{C+CM+h_1+M+M^2} d_{D'}(z'). \quad (3.11)$$

For this, we consider two cases: $\min\{d_{D'}(\omega'), d_{D'}(q')\} < 2\mu_3 |\omega' - q'|$ and $\min\{d_{D'}(\omega'), d_{D'}(q')\} \geq 2\mu_3 |\omega' - q'|$. In the former case, Proposition 3.4(P_2) and Theorem 2.10(2) give

$$\text{diam}(\gamma'_0) \leq c_2 \max\{|\omega' - q'|, 2(e^{h_1} - 1) \min\{d_{D'}(\omega'), d_{D'}(q')\}\} < 4c_2 e^{h_1} \mu_3 |\omega' - q'|.$$

Thus, we infer from Lemma 3.7(1) and statement (1) that

$$\text{diam}(\gamma'_0) < 4c_2^2 e^{h_1} d_{D'}(z'). \quad (3.12)$$

Moreover, Lemma 3.7(1), (3.12) and the statement (2a) imply

$$|z' - q'_0| \leq |z' - \omega'| + |\omega' - q'_0| \leq c_2 d_{D'}(z') + \frac{1}{4} \text{diam}(\gamma'_0) \leq c_2(1 + c_2 e^{h_1}) d_{D'}(z'). \quad (3.13)$$

In the latter case, that is, $\min\{d_{D'}(\omega'), d_{D'}(q')\} \geq 2\mu_3 |\omega' - q'|$. Lemma 2.2(1) implies

$$k_{D'}(\omega', q') < 1.$$

Since f is (M, C) -CQH, Lemma 2.6 and (2.3) yield

$$\begin{aligned}
\log\left(1 + \frac{|\omega' - q'_0|}{d_{D'}(\omega')}\right) &\leq k_{D'}(\omega', q'_0) \leq Mk_D(\omega, q_0) + C \\
&\leq M\ell_k(\gamma) + C \leq M\left(\frac{1}{4} + k_D(\omega, q)\right) + C \\
&\leq M\left(\frac{1}{4} + C + Mk_{D'}(\omega', q')\right) + C \\
&\leq M\left(\frac{1}{4} + C + M\right) + C.
\end{aligned} \tag{3.14}$$

Furthermore, we infer from statement (1) that

$$d_{D'}(\omega') \geq \min\{d_{D'}(\omega'), d_{D'}(q')\} \geq 2\mu_3|\omega' - q'| = 2\text{diam}(\gamma'[\omega', z']) \geq 2|\omega' - z'|.$$

As a consequence, we have

$$d_{D'}(z') \geq d_{D'}(\omega') - |\omega' - z'| \geq |\omega' - z'|.$$

Then we conclude from (3.14) that

$$|z' - q'_0| \leq |z' - \omega'| + |\omega' - q'_0| \leq 2e^{M(\frac{1}{4} + C + M) + C} d_{D'}(z'). \tag{3.15}$$

Now, (3.11) follows from (3.13) and (3.15), and hence, (3.9) holds.

Note that (3.9) and Theorem 2.14 ensure that

$$\begin{aligned}
k_{D'}(z', q'_0) &\leq c_4 \log\left(1 + \frac{|z' - q'_0|}{\min\{d_{D'}(q'_0), d_{D'}(z')\}}\right) \\
&\leq c_4 \log\left(1 + c_0^2 c_2^2 c_3 c_4^2 \mu_2 e^{4 + C + CM + h_1 + M + M^2}\right) \\
&\leq c_0 c_2 c_3 c_4^2 \mu_2 (4 + C + h_1 + M)^2.
\end{aligned}$$

Since f is (M, C) -CQH, we obtain from the previous estimate that

$$k_D(z, q_0) \leq Mk_{D'}(z', q'_0) + C \leq c_0 c_2 c_3 c_4^2 \mu_2 (4 + C + h_1 + M)^3.$$

This proves (2b), and hence, completes the proof of lemma. \square

Lemma 3.10. *There are an integer m and a finite sequence of points $\{\eta_i\}_{i=1}^{m+1}$ on $\gamma[\omega, z_0]$ such that*

- (1) $m > e^{8\mu_4}$ and $m\mu_4 \leq \ell_k(\gamma[\omega, z_0]) < (m+1)\mu_4$;
- (2) for each $i \in \{2, \dots, m\}$, $\eta_i \in \gamma[\eta_{i-1}, z_0]$,

$$\ell_k(\gamma[\eta_{i-1}, \eta_i]) = \mu_4 \quad \text{and} \quad \mu_4 \leq \ell_k(\gamma[\eta_m, \eta_{m+1}]) < 2\mu_4,$$

where $\eta_1 = \omega$ and $\eta_{m+1} = z_0$;

- (3) for any $i \neq j \in \{1, 2, \dots, m+1\}$,

$$k_{D'}(\eta'_i, \eta'_j) \geq (2M)^{-1}\mu_4;$$

- (4) for each $i \in \{2, \dots, m\}$,

- (a) $\ell(\gamma[\eta_{i-1}, \eta_i]) \leq e^{2\mu_4} d_D(\eta_i)$;
- (b) $\text{diam}(\gamma'[\eta'_{i-1}, \eta'_i]) \leq c_2 d_{D'}(\eta'_i)$;
- (c) $\text{diam}(\gamma'[\eta'_{i-1}, \eta'_i]) \geq e^{(3c_4 M)^{-1}\mu_4} d_{D'}(\eta'_{i-1})$;
- (d) $\text{diam}(\gamma'[\eta'_1, \eta'_i]) \geq c_2^{-1} e^{(3c_4 M)^{-1}\mu_4} \text{diam}(\gamma'[\eta'_1, \eta'_{i-1}])$.

Proof. (1). By Lemmas 3.5, 3.8 and the assumption $\ell(\gamma) > \mathfrak{C}\ell(\mathfrak{l})$, we have

$$\ell(\gamma[\omega, z_0]) \geq \frac{1}{\wp_2} \ell(\gamma) \geq \frac{\mathfrak{C}}{2\wp_2} d_D(w),$$

where $\wp_2 = 2 \left(1 + (1 + 4c_2)^{c_4 M} e^{\frac{1}{2} + C} \right)$. Then it follows from (2.2) that

$$\ell_k(\gamma[\omega, z_0]) \geq \log \left(1 + \frac{\ell(\gamma[\omega, z_0])}{d_D(\omega)} \right) \geq \log \left(1 + \frac{\mathfrak{C}}{2\wp_2} \right) > \mu_4 e^{8\mu_4},$$

where in the last inequality we used the the assumption $\mathfrak{C} > \frac{4}{5}\mu_5 = \frac{8}{5}\mu_4 e^{2\mu_4} e^{8\mu_4}$.

Consequently, there exists an integer m such that (1) holds.

(2). By the last estimate in the proof of (1), there exists a sequence $\{\eta_i\}_{i=1}^{m+1}$ in $\gamma[\omega, z_0]$ satisfying (2).

(3). Let $\{i, j\} \subset \{1, \dots, m+1\}$ with $i < j$. Since f is (M, C) -CQH, we have

$$k_{D'}(\eta'_i, \eta'_j) \geq \frac{1}{M} (k_D(\eta_i, \eta_j) - C).$$

It follows from Lemmas 2.6 and 2.7, together with (2), that

$$k_{D'}(\eta'_i, \eta'_j) \geq \frac{1}{M} (k_D(\eta_i, \eta_j) - C) \geq \frac{1}{M} (\ell_k(\gamma[\eta_i, \eta_j]) - \frac{1}{2} - C) \geq (2M)^{-1} \mu_4,$$

as $\mu_4 > \frac{1}{2} + C$. This proves (3).

(4). For each $i \in \{2, \dots, m+1\}$, (2.2) implies

$$\log \left(1 + \frac{\ell(\gamma[\eta_{i-1}, \eta_i])}{d_D(\eta_i)} \right) \leq \ell_k(\gamma[\eta_{i-1}, \eta_i]).$$

Then (4a) follows from (2).

The statement (4b) follows immediately from Lemma 3.7:

$$\text{diam}(\gamma'[\eta'_i, \eta'_{i-1}]) \leq \text{diam}(\gamma'[\eta'_1, \eta'_{i-1}]) \leq c_2 d_{D'}(\eta'_{i-1}).$$

Since $k_{D'}(\eta'_{i-1}, \eta'_i) > 1$ by (3), it follows from Lemma 2.2(1) and (4b) that

$$d_{D'}(\eta'_{i-1}) \leq 2|\eta'_{i-1} - \eta'_i| \leq 2c_2 d_{D'}(\eta'_i), \quad (3.16)$$

Note that Theorem 2.14 and (3) ensure that

$$(2M)^{-1} \mu_4 \leq k_{D'}(\eta'_{i-1}, \eta'_i) \leq c_4 \log \left(1 + \frac{|\eta'_{i-1} - \eta'_i|}{\min\{d_{D'}(\eta'_{i-1}), d_{D'}(\eta'_i)\}} \right).$$

Thus we see from the previous estimate and (3.16) that

$$\begin{aligned} \text{diam}(\gamma'[\eta'_{i-1}, \eta'_i]) &\geq |\eta'_{i-1} - \eta'_i| \\ &\geq (e^{(2c_4 M)^{-1} \mu_4} - 1) \min\{d_{D'}(\eta'_{i-1}), d_{D'}(\eta'_i)\} \\ &\geq e^{(3c_4 M)^{-1} \mu_4} d_{D'}(\eta'_{i-1}). \end{aligned}$$

This shows (4c).

By Lemma 3.7 and (4c), we have

$$\begin{aligned} \text{diam}(\gamma'[\eta'_1, \eta'_i]) &\geq \text{diam}(\gamma'[\eta'_{i-1}, \eta'_i]) \geq e^{(3c_4 M)^{-1} \mu_4} d_{D'}(\eta'_{i-1}) \\ &\geq c_2^{-1} e^{(3c_4 M)^{-1} \mu_4} \text{diam}(\gamma'[\eta'_1, \eta'_{i-1}]). \end{aligned}$$

This proves (4d) and thus completes the proof of lemma. \square

Lemma 3.11. *Let $u_1 = \omega = \eta_1$. Then the following statements hold.*

- (1) For each $i \in \{2, \dots, m+1\}$, there exists $u_i \in \mathfrak{l}[u_{i-1}, \varpi]$ such that
- (a) $|u'_1 - u'_i| = \mu_3^{-1} \text{diam}(\gamma'[\eta'_1, \eta'_i])$, and for any $u \in \mathfrak{l}[u_1, u_i] \setminus \{u_i\}$,

$$|\eta'_1 - u'| < \mu_3^{-1} \text{diam}(\gamma'[\omega', \eta'_i]);$$
 - (b) $|u'_1 - u'_i| \geq \frac{1}{2\mu_3} d_{D'}(\eta'_1)$.
- (2) For each $i \in \{2, \dots, m+1\}$ and any λ -curve $\gamma_i \in \Lambda_{u_1 u_i}^\lambda$, there exists $u_{0,i} \in \gamma_i$ such that
- (a) $|u'_1 - u'_{0,i}| = \frac{1}{4} \text{diam}(\gamma'_i)$, and for any $u \in \gamma_i[u_1, u_{0,i}]$,

$$|\eta'_1 - u'| < \frac{1}{4} \text{diam}(\gamma'_i).$$
 - (b) $\max\{k_D(\eta_i, u_{0,i}), k_{D'}(\eta'_i, u'_{0,i})\} \leq c_0 c_2 c_3 c_4^2 \mu_2 (4 + C + h_1 + M)^3$;
 - (c) $|u'_1 - u'_{0,i}| \leq c_2 d_{D'}(u'_{0,i})$;
 - (d) $e^{-c_0 c_2 c_3 c_4^2 \mu_2 (4 + C + h_1 + M)^3} d_{D'}(u'_{0,i}) \leq d_{D'}(\eta'_i) \leq e^{c_0 c_2 c_3 c_4^2 \mu_2 (4 + C + h_1 + M)^3} d_{D'}(u'_{0,i})$;
- (3) For each $i \in \{3, \dots, m+1\}$,
- (a) $k_D(u_{i-1}, u_i) \geq (4c_4 M^2)^{-1} \mu_4$;
 - (b) $\max\{d_{D'}(u'_{i-1}), d_{D'}(u'_i)\} \leq 2|u'_{i-1} - u'_i|$.

Proof. (1). Let $i \in \{2, \dots, m+1\}$. Since f is (M, C) -CQH, we know from Lemma 3.10(3) that

$$k_D(\eta_1, \eta_i) \geq \frac{1}{M} (k_{D'}(\eta'_1, \eta'_i) - C) \geq (3M^2)^{-1} \mu_4 > \mu_3.$$

Applying Lemma 3.9(1), with z replaced by η_i , we obtain that there exists $u_i \in \mathfrak{l}$ such that (1a) holds. Furthermore, by Lemma 3.9(2), we also know that for any λ -curve $\gamma_i \in \Lambda_{u_1 u_i}^\lambda$, there exists $u_{0,i} \in \gamma_i$ such that the statements (2a) and (2b) are satisfied.

Before the proof of (1b), we check that for each $i \in \{2, \dots, m+1\}$, $u_i \in \mathfrak{l}[u_{i-1}, \varpi]$. Clearly, we only need to consider the case when $i \in \{3, \dots, m+1\}$. It follows from (1a) and Lemma 3.10(4d) that

$$|u'_1 - u'_i| \geq (c_2 \mu_3)^{-1} e^{(3c_4 M)^{-1} \mu_4} \text{diam}(\gamma'[\eta'_1, \eta'_{i-1}]) = c_2^{-1} e^{(3c_4 M)^{-1} \mu_4} |u'_1 - u'_{i-1}|.$$

Since $c_2^{-1} e^{(3c_4 M)^{-1} \mu_4} > 1$, this implies that $u_i \in \mathfrak{l}[u_{i-1}, \varpi]$.

By Lemma 3.10(4c) and (1a), we know that for any $i \in \{2, \dots, m+1\}$,

$$\mu_3 |u'_1 - u'_i| = \text{diam}(\gamma'[\eta'_1, \eta'_i]) \geq \text{diam}(\gamma'[\eta'_1, \eta'_2]) \geq \frac{1}{2} d_{D'}(\eta'_1), \quad (3.17)$$

from which (1b) follows.

(2). The statements (2a) and (2b) are already verified in the previous proof. For (2c), we know from (2a) that

$$\text{diam}(\gamma'_i[u'_1, u'_{0,i}]) \leq \text{diam}(\gamma'_i[u'_i, u'_{0,i}]).$$

It follows from Proposition 3.4(P₂) and Theorem 2.10(1) that

$$|u'_1 - u'_{0,i}| \leq \text{diam}(\gamma'_i[u'_1, u'_{0,i}]) \leq c_2 d_{D'}(u'_{0,i}),$$

which proves (2c).

The statement (2d) follows immediately from (2b) and the following estimate from (2.3)

$$k_{D'}(\eta'_i, u'_{0,i}) \geq \max \left\{ \log \frac{d_{D'}(u'_{0,i})}{d_{D'}(\eta'_i)}, \log \frac{d_{D'}(\eta'_i)}{d_{D'}(u'_{0,i})} \right\}.$$

(3). Lemma 3.10(4c), Lemma 3.7(1) and (1a) yield

$$\begin{aligned} |u'_i - u'_{i-1}| &\geq |u'_i - u'_1| - |u'_1 - u'_{i-1}| = \mu_3^{-1}(\text{diam}(\gamma'[\eta'_1, \eta'_i]) - \text{diam}(\gamma'[\eta'_1, \eta'_{i-1}])) \\ &\geq \mu_3^{-1}(e^{(3c_4M)^{-1}\mu_4} d_{D'}(\eta'_{i-1}) - \text{diam}(\gamma'[\eta'_1, \eta'_{i-1}])) \\ &\geq \mu_3^{-1}(e^{(3c_4M)^{-1}\mu_4} - c_2) d_{D'}(\eta'_{i-1}). \end{aligned} \quad (3.18)$$

Moreover, for each $i \in \{3, \dots, m+1\}$, by Lemma 3.7(1), (1a) and (3.17), we have

$$\begin{aligned} d_{D'}(u'_{i-1}) &\leq d_{D'}(u'_1) + |u'_1 - u'_{i-1}| = d_{D'}(\eta'_1) + |u'_1 - u'_{i-1}| \\ &\leq (2 + \mu_3^{-1}) \text{diam}(\gamma'[\eta'_1, \eta'_{i-1}]) \\ &\leq c_2(2 + \mu_3^{-1}) d_{D'}(\eta'_{i-1}). \end{aligned}$$

Then, it follows from (2.3), (3.18) and the previous estimate that

$$k_{D'}(u'_{i-1}, u'_i) \geq \log \frac{|u'_{i-1} - u'_i|}{d_{D'}(u'_{i-1})} \geq \log \frac{e^{(3c_4M)^{-1}\mu_4} - c_2}{c_2(1 + 2\mu_3)} > 1.$$

Consequently, we know from Lemma 2.2(1) that

$$\max\{d_{D'}(u'_{i-1}), d_{D'}(u'_i)\} \leq 2|u'_{i-1} - u'_i|.$$

Since f is (M, C) -CQH, we have

$$k_D(u_{i-1}, u_i) \geq \frac{1}{M}(k_{D'}(u'_{i-1}, u'_i) - C) > (4c_4M^2)^{-1}\mu_4,$$

These two estimates give (3a) and (3b), and hence, the proof is complete. \square

Lemma 3.12. *For each $i \in \{2, \dots, m+1\}$, let $\gamma_i \in \Lambda_{u_1 u_i}^\lambda$. Then for any $u \in \gamma_i$, we have*

- (1) $d_{D'}(u') \leq 4c_2(1 + 2\mu_3)d_{D'}(u'_{0,i});$
- (2) $d_{D'}(u') \leq e^{\mu_3} d_{D'}(\eta'_i).$

Proof. (1). Let $u \in \gamma_i$. By statements (1b) and (2a) in Lemma 3.11 and Lemma 3.11(2c), we have

$$\begin{aligned} d_{D'}(u') &\leq d_{D'}(u'_1) + \text{diam}(\gamma'_i) \leq (1 + 2\mu_3) \text{diam}(\gamma'_i) \\ &= 4(1 + 2\mu_3)|u'_1 - u'_{0,i}| \leq 4c_2(1 + 2\mu_3)d_{D'}(u'_{0,i}). \end{aligned}$$

This proves (1).

(2). Note that

$$4c_2(1 + 2\mu_3)e^{c_0 c_2 c_3 c_4^2 \mu_2 (4+C+h_1+M)^3} < e^{\mu_3}.$$

The desired statement (2) follows from (1) and Lemma 3.11(2d). \square

Lemma 3.13. *For each $i \in \{3, \dots, m+1\}$, let $\gamma_{i-1} \in \Lambda_{u_1 u_{i-1}}^\lambda$ be a λ -curve. Then for any $v \in \gamma_{i-1}$, it holds*

$$k_D(u_{0,i}, v) > (4c_4M)^{-1}\mu_4.$$

Proof. Let $i \in \{3, \dots, m+1\}$ and $v \in \gamma_{i-1}$. Since f is (M, C) -CQH, we get from (2.3) that

$$k_D(u_{0,i}, v) \geq M^{-1}(k_{D'}(u'_{0,i}, v') - C) \geq M^{-1}\left(\log \frac{d_{D'}(u'_{0,i})}{d_{D'}(v')} - C\right). \quad (3.19)$$

By Lemma 3.11(2d) and Lemma 3.12(2), we have

$$\frac{d_{D'}(u'_{0,i})}{d_{D'}(v')} \geq e^{-2c_0c_2c_3c_4\mu_2(4+C+h_1+M)^3-\mu_3} \frac{d_{D'}(\eta'_i)}{d_{D'}(\eta'_{i-1})}.$$

Furthermore, Lemmas 3.7 and 3.10(4c) give

$$d_{D'}(\eta'_{i-1}) \leq e^{-(3c_4M)^{-1}\mu_4} \text{diam}(\gamma'[\eta'_{i-1}, \eta'_i]) \leq c_2 e^{-(3c_4M)^{-1}\mu_4} d_{D'}(\eta'_i).$$

Substituting these two estimates in (3.19) leads to

$$k_D(u_{0,i}, v) \geq (4c_4M)^{-1}\mu_4.$$

The proof is complete. \square

Lemma 3.14. *For each $i \in \{3, \dots, m+1\}$ and each $\beta_i \in \Lambda_{u_{i-1}u_i}^\lambda$, there exists $u_{1,i} \in \beta_i$ such that*

- (1) $k_D(u_{0,i}, u_{1,i}) \leq \frac{1}{2} + \mu_0$;
- (2) $\ell(\gamma[\eta_{i-1}, \eta_i]) \leq e^{\mu_3+2\mu_4} d_D(u_{1,i})$;
- (3) $\ell(\beta_i) \geq \frac{1}{2} d_D(u_{1,i})$.

Proof. (1) For each $i \in \{3, \dots, m+1\}$, let $\gamma_{i-1} \in \Lambda_{u_{i-1}u_i}^\lambda$ and $\gamma_i \in \Lambda_{u_1u_i}^\lambda$. Since the triangle $(\gamma_{i-1}, \gamma_i, \beta_i)$ is λ -short by Lemma 2.19, we know from Proposition 3.4(P_4) that there exists $u_{1,i} \in \gamma_{i-1} \cup \beta_i$ such that

$$k_D(u_{0,i}, u_{1,i}) \leq \frac{1}{2} + \mu_0.$$

As $(4c_4M)^{-1}\mu_4 > \frac{1}{2} + \mu_0$, we infer from Lemma 3.13 that $u_{1,i} \in \beta_i$. This proves (1).

(2). By (1) and Lemma 3.11(2b), we know that

$$\begin{aligned} k_D(u_{1,i}, \eta_i) &\leq k_D(u_{1,i}, u_{0,i}) + k_D(u_{0,i}, \eta_i) \\ &\leq \frac{1}{2} + \mu_0 + c_0c_2c_3c_4^2\mu_2(4+C+h_1+M)^3 < \mu_3. \end{aligned}$$

This together with (2.3) implies

$$d_D(\eta_i) \leq e^{k_D(u_{1,i}, \eta_i)} d_D(u_{1,i}) < e^{\mu_3} d_D(u_{1,i}).$$

Then, it follows from the above estimate and Lemma 3.10(4a) that

$$\ell(\gamma[\eta_{i-1}, \eta_i]) \leq e^{2\mu_4} d_D(\eta_i) \leq e^{2\mu_4+\mu_3} d_D(u_{1,i}).$$

This proves (2).

(3). By Lemma 3.11(3a), we have

$$\max\{k_D(u_{i-1}, u_{1,i}), k_D(u_i, u_{1,i})\} \geq \frac{1}{2} k_D(u_{i-1}, u_i) > (8c_4M^2)^{-1}\mu_4 > 1.$$

Then Lemma 2.2(2) ensures that

$$\ell(\beta_i) \geq \frac{1}{2} d_D(u_{1,i}).$$

This proves (3), and hence, completes the proof of lemma. \square

3.3. Proof of Theorem 3.2. In this section, we prove Theorem 3.2 by a contradiction argument and we shall keep all the notations used in Section 3.2.

Suppose that for any $u \neq v$ in \mathbb{I} , there is an element $\beta \in \Lambda_{uv}^\lambda$ such that

$$\ell(\beta) < \mu_6 \mathfrak{C} \ell(\mathbb{I}[u, v]). \quad (3.20)$$

This assumption implies that for each $i \in \{2, \dots, m+1\}$, there is an element in $\beta_i \in \Lambda_{u_{i-1}u_i}^\lambda$ such that

$$\ell(\mathbb{I}[u_{i-1}, u_i]) \geq (\mu_6 \mathfrak{C})^{-1} \ell(\beta_i). \quad (3.21)$$

To reach a contradiction, we make the following claim.

Claim 3.1. For each $i \in \{2, \dots, m+1\}$, $\ell(\mathbb{I}[u_{i-1}, u_i]) \geq e^{\mu_4 - \mu_3} (\mu_6 \mathfrak{C})^{-1} \ell(\gamma[\eta_{i-1}, \eta_i])$.

For the proof, we let $i \in \{2, \dots, m+1\}$ and divide the discussions into two cases.

Case 3.1-1: $\ell(\beta_i) \leq e^{3\mu_4} d_D(u_{1,i})$.

Let $u_{2,i} \in \beta_i$ be such that

$$|u'_{i-1} - u'_{2,i}| = \frac{1}{2} |u'_{i-1} - u'_i|. \quad (3.22)$$

This, together with Proposition 3.4(P_2) and Theorem 2.10(1), implies that

$$\begin{aligned} |u'_{i-1} - u'_{2,i}| &= \min \{ |u'_{i-1} - u'_{2,i}|, |u'_i - u'_{2,i}| \} \\ &\leq \min \{ \text{diam}(\beta_i'[u'_{i-1}, u'_{2,i}]), \text{diam}(\beta_i'[u'_{2,i}, u'_i]) \} \\ &\leq c_2 d_{D'}(u'_{2,i}). \end{aligned} \quad (3.23)$$

To continue the proof, we need a relation between $\ell(\beta_i)$ and $d_D(u_{2,i})$. Firstly, we derive a relation between $\text{diam}(\beta_i')$ and $d_{D'}(u'_{2,i})$. Since Proposition 3.4(P_2), Theorem 2.10(2) and Lemma 3.11(3b) yield

$$\begin{aligned} \text{diam}(\beta_i') &\leq c_2 \max \{ |u'_{i-1} - u'_i|, 2(e^{h_1} - 1) \min \{ d_{D'}(u'_{i-1}), d_{D'}(u'_i) \} \} \\ &\leq 4c_2 e^{h_1} |u'_{i-1} - u'_i|, \end{aligned} \quad (3.24)$$

we get from (3.22) and (3.23) that

$$\text{diam}(\beta_i') \leq 8c_2^2 e^{h_1} d_{D'}(u'_{2,i}). \quad (3.25)$$

Secondly, we establish a relation between $\text{diam}(\beta_i')$ and $d_{D'}(u'_{1,i})$. When $i \in \{3, \dots, m+1\}$, by Lemma 3.11(1a) and (2a), we have

$$\begin{aligned} |u'_{i-1} - u'_i| &\leq |u'_{i-1} - u'_1| + |u'_1 - u'_i| = \mu_3^{-1} \text{diam}(\gamma'[\eta'_1, \eta'_{i-1}]) + |u'_1 - u'_i| \\ &\leq \mu_3^{-1} \text{diam}(\gamma'[\eta'_1, \eta'_i]) + |u'_1 - u'_i| \\ &= 2|u'_1 - u'_i| \leq 2 \text{diam}(\gamma'_i) = 8|u'_1 - u'_{0,i}|. \end{aligned} \quad (3.26)$$

Combining (3.24) with (3.26), we see from Lemma 3.11(2c) that

$$\text{diam}(\beta_i') \leq 32c_2^2 e^{h_1} d_{D'}(u'_{0,i}). \quad (3.27)$$

On the other hand, since f is (M, C) -CQH, it follows from (2.3) and the choice of $u_{1,i}$ (see Lemma 3.14(1)) that

$$\log \frac{d_{D'}(u'_{0,i})}{d_{D'}(u'_{1,i})} \leq k_{D'}(u'_{0,i}, u'_{1,i}) \leq M k_D(u_{0,i}, u_{1,i}) + C \leq M \left(\frac{1}{2} + \mu_0 \right) + C.$$

Then we infer from the above estimate and (3.27) that

$$\text{diam}(\beta_i') \leq 32c_2^2 e^{C+h_1+2\mu_0 M} d_{D'}(u'_{1,i}), \quad (3.28)$$

where in the final estimate we used the fact that $\mu_0 > 16$.

Now, we are ready to establish the necessary relation between $\ell(\beta_i)$ and $d_D(u_{2,i})$. Note that the combination of (3.25) and (3.28) shows that

$$\text{diam}(\beta_i') \leq 32c_2^2 e^{C+h_1+2\mu_0 M} \min\{d_{D'}(u'_{1,i}), d_{D'}(u'_{2,i})\}.$$

Then it follows from (2.3), the assumption that f is (M, C) -CQH and Theorem 2.14 that

$$\begin{aligned} \log \frac{d_D(u_{1,i})}{d_D(u_{2,i})} &\leq k_D(u_{1,i}, u_{2,i}) \leq M k_{D'}(u'_{1,i}, u'_{2,i}) + C \\ &\leq M c_4 \log \left(1 + \frac{\text{diam}(\beta_i')}{\min\{d_{D'}(u'_{1,i}), d_{D'}(u'_{2,i})\}} \right) + C \\ &\leq M c_4 \log \left(1 + 32c_2^2 e^{C+h_1+2\mu_0 M} \right) + C \leq \mu_3. \end{aligned}$$

Combining it with the assumption $\ell(\beta_i) \leq e^{3\mu_4} d_D(u_{1,i})$ yields

$$\ell(\beta_i) \leq \mu_7 d_D(u_{2,i}). \quad (3.29)$$

Notice that (3.22) and (3.29) implies that the assumptions in Theorem 3.3 are satisfied with x , y and x_0 replaced by u_{i-1} , u_i and $u_{2,i}$, respectively. It follows that

$$\ell(\beta_i) \leq \mu_8 \sigma_D(u_{i-1}, u_i).$$

Since $\mu_8 \leq (\frac{4}{5}\mu_5)^{\frac{2}{3}}$ and $\mathfrak{C} \geq \frac{4}{5}\mu_5$, we get from the previous estimate that

$$\ell(\beta_i) \leq \mathfrak{C}^{\frac{2}{3}} \sigma_D(u_{i-1}, u_i). \quad (3.30)$$

Moreover, the statements (2) and (3) in Lemma 3.14 guarantee that

$$\ell(\gamma[\eta_{i-1}, \eta_i]) \leq 2e^{\mu_3+2\mu_4} \ell(\beta_i),$$

which together with (3.30) give

$$\ell(\mathfrak{l}[u_{i-1}, u_i]) \geq \sigma_D(u_{i-1}, u_i) \geq (2\mathfrak{C}^{\frac{2}{3}} e^{\mu_3+2\mu_4})^{-1} \ell(\gamma[\eta_{i-1}, \eta_i]). \quad (3.31)$$

Case 3.1-2: $\ell(\beta_i) > e^{3\mu_4} d_D(u_{1,i})$.

Under this assumption, it follows from (3.21) and Lemma 3.14(2) that

$$\ell(\mathfrak{l}[u_{i-1}, u_i]) \geq e^{3\mu_4} (\mu_6 \mathfrak{C})^{-1} d_D(u_{1,i}) \geq e^{\mu_4-\mu_3} (\mu_6 \mathfrak{C})^{-1} \ell(\gamma[\eta_{i-1}, \eta_i]). \quad (3.32)$$

The claim then follows from (3.31) and (3.32) by noticing that

$$(2\mathfrak{C}^{\frac{2}{3}} e^{\mu_3+2\mu_4})^{-1} \geq e^{\mu_4-\mu_3} (\mu_6 \mathfrak{C})^{-1}.$$

Let us continue the proof based on Claim 3.1. By Claim 3.1 and (3.4), we have

$$\ell(\mathfrak{l}) \geq \sum_{i=2}^{m+1} \ell(\mathfrak{l}[u_{i-1}, u_i]) \geq e^{\mu_4-\mu_3} (\mu_6 \mathfrak{C})^{-1} \ell(\gamma[\omega, z_0]) \geq \frac{1}{2} e^{\mu_4-\mu_3} (\mu_6 \mathfrak{C})^{-1} \ell(\gamma).$$

As $2\mu_6 < e^{\mu_4-\mu_3}$, the above inequality contradicts with our assumption $\ell(\gamma) \geq \mathfrak{C}\ell(\mathfrak{l})$. The proof of Theorem 3.2 is thus complete. \square

4. PROOFS OF THE MAIN RESULTS

The purpose of this section is to prove the main results in this paper, i.e., Theorems 1.7 ~ 1.9, based on Theorem 3.2. For the proof of Theorem 1.7, we need the following useful result bounding $\ell(\gamma)$ in terms of $\ell(\alpha)$. Recall the definition of μ_5 given in Section 2.1.

Proposition 4.1. *Under the **Standing Assumptions** in Section 3.1, let α be an arc in D joining ω and ϖ . Then we have*

$$\ell(\gamma) \leq \frac{4}{5}\mu_5\ell(\alpha).$$

Proof. It suffices to consider the case when α is rectifiable. Suppose on the contrary that

$$\ell(\gamma) > \frac{4}{5}\mu_5\ell(\alpha). \quad (4.1)$$

To reach a contradiction, let $u_1 = \omega$ and $v_1 = \varpi$. Then it follows from Theorem 3.2 and (4.1) that there are points $u_2 \neq v_2$ on α such that $u_2 \in \alpha[u_1, v_2]$, $v_2 \in \alpha[u_2, v_1]$, and there is an element $\gamma_2 \in \Lambda_{u_2v_2}^\lambda$ such that

$$\ell(\gamma_2) \geq \frac{4}{5}\mu_5\mu_6\ell(\alpha[u_2, v_2]). \quad (4.2)$$

Combining Theorem 3.2 with (4.2), we see that there are points $u_3 \neq v_3$ on $\alpha[u_2, v_2]$ such that $u_3 \in \alpha[u_2, v_3]$, $v_3 \in \alpha[u_3, v_2]$, and there is an element $\gamma_3 \in \Lambda_{u_3v_3}^\lambda$ such that

$$\ell(\gamma_3) \geq \frac{4}{5}\mu_5\mu_6^2\ell(\alpha[u_3, v_3]).$$

Repeating the above procedure, we obtain two sequences $\{u_i\}_{i=1}^\infty$ and $\{v_i\}_{i=1}^\infty$ such that for each $i \geq 1$, $u_{i+1} \neq v_{i+1} \in \alpha[u_i, v_i]$, $u_{i+1} \in \alpha[u_i, v_{i+1}]$, $v_{i+1} \in \alpha[u_{i+1}, v_i]$, and there is an element $\gamma_i \in \Lambda_{u_iv_i}^\lambda$ such that

$$\ell(\gamma_i) \geq \frac{4}{5}\mu_5\mu_6^{i-1}\ell(\alpha[u_i, v_i]). \quad (4.3)$$

Select $u_0 \in \alpha$ so that $d_D(u_0) = \min_{u \in \alpha}\{d_D(u)\}$, and set

$$\kappa = \frac{\ell(\alpha)}{d_D(u_0)}.$$

Then $0 < \kappa < \infty$ and

$$\ell_k(\alpha) = \int_\alpha \frac{|dz|}{d_D(z)} \leq \frac{\ell(\alpha)}{d_D(u_0)} = \kappa.$$

Note that by (2.2) and Lemma 2.6, we have

$$\log\left(1 + \frac{\ell(\gamma_i)}{d_D(u_i)}\right) \leq \ell_k(\gamma_i) \leq k_D(u_i, v_i) + \frac{1}{4} \leq \ell_k(\alpha[u_i, v_i]) + \frac{1}{4} \leq \frac{1}{4} + \kappa.$$

On the other hand, Lemma 3.5 guarantees that for each $i \geq 1$,

$$d_D(u_i) \leq 2\ell(\alpha[u_i, v_i]),$$

which gives

$$\ell(\gamma_i) \leq e^{\frac{1}{4} + \kappa} d_D(u_i) \leq 2e^{\frac{1}{4} + \kappa} \ell(\alpha[u_i, v_i]).$$

This, together with (4.3), implies

$$\frac{4}{5}\mu_5\mu_6^{i-1} \leq \frac{\ell(\gamma_i)}{\ell(\alpha[u_i, v_i])} \leq 2e^{\frac{1}{4} + \kappa},$$

which is impossible, as $\lim_{i \rightarrow \infty} \left(\frac{4}{5}\mu_5\mu_6^{i-1}\right) = \infty$. Consequently, the proof is complete. \square

Proof of Theorem 1.7. Let ϑ be a c_0 -quasigeodesic, $c_0 \geq 1$, in D with end points ω and ϖ , and let L be an arc in D with the same end points as ϑ . By Proposition 3.4(P_3), D is δ -Gromov hyperbolic, for some $\delta = \delta(c, C, M)$.

We may assume that L is rectifiable with $\ell(L) < \ell(\vartheta)$. In the following proof, we consider two cases: $k_D(\omega, \varpi) > 1$ and $k_D(\omega, \varpi) \leq 1$.

In the former case, as both ϑ and γ are $(1 + c_0)$ -quasigeodesics, we obtain from Theorem 2.28 that

$$\ell(\vartheta) \leq \lambda_0 \ell(\gamma), \quad (4.4)$$

where $\lambda_0 = 4e^{8(1+c_0)^2\mu_2}$. On the other hand, Proposition 4.1 gives

$$\ell(\gamma) \leq \frac{4}{5}\mu_5 \ell(L).$$

Combining the above two inequalities leads to our desired estimate

$$\ell(\vartheta) \leq \frac{4}{5}\mu_5 \lambda_0 \ell(L). \quad (4.5)$$

In the latter case, that is, $k_D(\omega, \varpi) \leq 1$, then we know from [34, Lemma 2.31] that γ is a $18e^9$ -uniform arc, which implies

$$\ell(\gamma) \leq 18e^9 |\omega - \varpi| \leq 18e^9 \ell(L).$$

Combining it with (4.4) gives

$$\ell(\vartheta) \leq 18\lambda_0 e^9 \ell(L). \quad (4.6)$$

Set $b = \frac{4}{5}\mu_5 \lambda_0$. Then according to our choice of constants in Section 2.1,

$$b \geq 18\lambda_0 e^9$$

and thus we conclude from (4.5) and (4.6) that

$$\ell(\vartheta) \leq b \ell(L).$$

The theorem follows from the arbitrariness of ω and ϖ in D . \square

Proof of Theorem 1.8. By [32, Theorem 4.8] and [33, Theorem 4.11], we know that the image of a c_0 -quasigeodesic under an M -bi-Lipschitz homeomorphism (in the quasihyperbolic metric) is a c'_0 -quasigeodesic with $c'_0 = 2c_0 M^2(1 + M^2)$. As the inverse of an M -bi-Lipschitz homeomorphism is again M -bi-Lipschitz, Theorem 1.8 follows directly from Theorem 1.7. \square

Proof of Theorem 1.9. For ω and ϖ in D , let ϑ be a c_0 -quasigeodesic in D connecting ω and ϖ . First of all, under the assumptions in the theorem, [28, Theorem 1] ensures that ϑ is a b_1 -cone arc, where $b_1 = b_1(a, c, c_0, C, M)$. Secondly, Theorem 1.7 implies that

$$\ell(\vartheta) \leq b_2 \sigma_D(\omega, \varpi),$$

where $b_2 = b_2(c, c_0, C, M)$. Set $b = \max\{b_1, b_2\}$. Then the previous estimates imply that ϑ is a b -inner uniform curve and thus the proof is complete. \square

5. PROOF OF THEOREM 3.3

The purpose of this section is to prove Theorem 3.3, which has played a key role in the proof of Theorem 3.2. As in Section 3, we fix $\lambda = \frac{1}{4}$.

5.1. **Preparation for the proof.** Note that if $k_D(x, y) \leq 1$, then we know from [34, Lemma 2.31] that $\gamma \in \Lambda_{xy}^\lambda$ is a c -uniform arc with $c = 18e^9$, which implies that the theorem holds. Thus, in the following proof, we always assume that

$$k_D(x, y) > 1. \quad (5.1)$$

Let α be a rectifiable curve in D joining x and y ; see Figure 4. Then it follows from Lemma 2.2 and (5.1) that for each $w \in \alpha$,

$$d_D(w) \leq 2\ell(\alpha). \quad (5.2)$$




FIGURE 4. Illustration for the proof of Theorem 3.3 in Section 5.1

We shall prove the theorem by a contradiction. To this end, suppose that

$$\ell(\gamma) \geq \frac{1}{2}\mu_8\ell(\alpha). \quad (5.3)$$

To reach a contradiction, let $x_1 = x$ and y_1 be the first point in α along the direction from x_1 to y (see Figure 4) such that

$$|x'_1 - y'_1| = |x'_1 - y'|. \quad (5.4)$$

It is possible that $y_1 = y$. Let $\gamma_1 \in \Lambda_{x_1 y_1}^\lambda$ be a fixed element and select $y_0 \in \gamma_1$ such that

$$|x'_1 - y'_0| = \frac{1}{2}|x'_1 - y'_1|. \quad (5.5)$$

Then we know from Lemma 2.12 that

$$\text{diam}(\gamma') \leq c_3 d_{D'}(x'_0) \quad \text{and} \quad \text{diam}(\gamma'_1) \leq c_3 d_{D'}(y'_0), \quad (5.6)$$

where c_3 is the constant given by Lemma 2.12.

The following two lemmas concern the point y_0 .

Lemma 5.1. *Let c_4 be the constant given by Theorem 2.14. Then*

- (1) $\ell(\gamma) \leq 8\mu_7(1 + c_3)^{\frac{81}{16}}c_4 M e^{6C+9\mu_2}\ell(\gamma[x_1, y_0])$;
- (2) $d_D(x_0) \leq (1 + c_3)^{c_4 M} e^C d_D(y_0)$.

Proof. (1). We shall apply Lemma 2.27 to prove (1). For this, we need some preparation.

Since (3.1), (5.4), and (5.5) lead to

$$|x'_0 - y'_0| \leq |x'_0 - x'_1| + |x'_1 - y'_0| = |x'_1 - y'_1|,$$

we infer from (5.4) and (5.6) that

$$|x'_0 - y'_0| \leq c_3 \min\{d_{D'}(x'_0), d_{D'}(y'_0)\}.$$

Since D' is c -uniform and f is (M, C) -CQH, we know from Theorem 2.14 and the above estimate that

$$\begin{aligned} k_D(x_0, y_0) &\leq M k_{D'}(x'_0, y'_0) + C \leq c_4 \log \left(1 + \frac{|x'_0 - y'_0|}{\min\{d_{D'}(x'_0), d_{D'}(y'_0)\}} \right) \\ &\leq c_4 M \log(1 + c_3) + C. \end{aligned} \quad (5.7)$$

Next, we estimate $k_D(x_1, x_0)$ and $k_D(x_1, y_0)$. First, we derive a lower bound for $\ell(\gamma[x_1, x_0])$. By the triangle inequality and (3.1), we have

$$\ell(\gamma[x_1, x_0]) \geq d_D(x_0) - d_D(x_1) \geq \mu_7^{-1} \ell(\gamma) - d_D(x_1).$$

This, combining with (5.2) and (5.3), gives

$$\ell(\gamma[x_1, x_0]) \geq \frac{\mu_8 - 4\mu_7}{2\mu_7} \ell(\alpha). \quad (5.8)$$

By Proposition 3.4(P_1), (2.2), (5.2) and (5.8), we have

$$\begin{aligned} k_D(x_1, x_0) &\geq \frac{8}{9} \ell_k(\gamma[x_1, x_0]) \geq \frac{8}{9} \log \left(1 + \frac{\ell(\gamma[x_1, x_0])}{d_D(x_1)} \right) \\ &\geq \frac{8}{9} \log \left(1 + \frac{\mu_8 - 4\mu_7}{4\mu_7} \right) > 1. \end{aligned}$$

Then it follows from (5.7) and the above estimate that

$$\begin{aligned} k_D(x_1, y_0) &\geq k_D(x_1, x_0) - k_D(x_0, y_0) \\ &\geq \frac{8}{9} \log \left(1 + \frac{\mu_8 - 4\mu_7}{4\mu_7} \right) - c_4 M \log(1 + c_3) - C > 1. \end{aligned}$$

Thus we have

$$\min\{k_D(x_1, x_0) > 1, k_D(x_1, y_0)\} > 1. \quad (5.9)$$

Since (D, k_D) is a (μ_0, ρ_0) -Rips space by Proposition 3.4(P_4), and (5.7) and (5.9) hold, we may apply Lemma 2.27 with x_1, x_2 and x_3 replaced by x_1, x_0 and y_0 to deuce

$$\ell(\gamma[x_1, x_0]) \leq 4(1 + c_3)^{\frac{81}{16} c_4 M} e^{\frac{81}{16} C + 9\mu_2} \ell(\gamma[x_1, y_0]). \quad (5.10)$$

Furthermore, by (3.1), (5.9) and Lemma 2.2(2), it holds

$$\ell(\gamma) \leq \mu_7 d_D(x_0) \leq 2\mu_7 \ell(\gamma[x_1, x_0]).$$

This together with (5.10) give

$$\ell(\gamma) \leq 8\mu_7 (1 + c_3)^{\frac{81}{16} c_4 M} e^{6C + 9\mu_2} \ell(\gamma[x_1, y_0]).$$

This proves (1).

(2). By (2.3) and (5.7), we have

$$\log \frac{d_D(x_0)}{d_D(y_0)} \leq k_D(x_0, y_0) \leq c_4 M \log(1 + c_3) + C,$$

from which it follows

$$d_D(x_0) \leq (1 + c_3)^{c_4 M} e^C d_D(y_0).$$

This proves (2) and hence completes the proof of lemma. \square

As an application of Lemma 5.1, we obtain the following result.

Lemma 5.2. *For any λ -curve $\gamma_1 \in \Lambda_{x_1 y_1}^\lambda$, $\ell(\gamma_1) \geq \mu_9^{\frac{1}{2}} \ell(\alpha)$.*

Proof. Fix $\gamma_1 \in \Lambda_{x_1 y_1}^\lambda$. By Lemma 5.1(2) and (3.1), we know that

$$\ell(\gamma_1) \geq \ell(\gamma_1[x_1, y_0]) \geq d_D(y_0) - d_D(x_1) \geq \frac{1}{(1 + c_3)^{c_4 M} e^C \mu_7} \ell(\gamma) - d_D(x_1).$$

Then it follows from (5.2) and (5.3) that

$$\ell(\gamma_1) \geq \frac{\mu_8 - 4(1 + c_3)^{c_4 M} e^C \mu_7}{2(1 + c_3)^{c_4 M} e^C \mu_7} \ell(\alpha) \geq \mu_9^{\frac{1}{2}} \ell(\alpha).$$

The proof is complete. \square

Lemma 5.3. *Let $u_1 \in \alpha[x_1, y_1]$, $\gamma_0 \in \Lambda_{u_1 y_0}^\lambda \in \Lambda_{u_1 y_0}$ and $v_1 \in \gamma_0$. If $k_D(v_1, y_0) \geq 2c_2 c_4 M \log c_3$, then for any $u \in \gamma_0[u_1, v_1]$,*

$$\text{diam}(\gamma'_0[u'_1, u']) \leq c_2 d_{D'}(u'),$$

where c_2 is the constant given by Theorem 2.10.

Proof. Let $u \in \gamma_0[u_1, v_1]$. Since f is (M, C) -CQH and $k_D(v_1, y_0) \geq 2c_2 c_4 M \log c_3$, we deduce from Proposition 3.4(P_1) that

$$\begin{aligned} k_{D'}(u', y'_0) &\geq \frac{1}{M} (k_D(u, y_0) - C) \geq \frac{1}{M} \left(\frac{8}{9} \ell_k(\gamma_0[u, y_0]) - C \right) \\ &\geq \frac{1}{M} \left(\frac{8}{9} \ell_k(\gamma_0[v_1, y_0]) - C \right) \geq \frac{16}{9} c_2 c_4 \log c_3 - \frac{C}{M}. \end{aligned} \quad (5.11)$$

Next, we prove

$$\text{diam}(\gamma'_0[u'_1, u']) \leq \text{diam}(\gamma'_0[u', y'_0]). \quad (5.12)$$

Suppose on the contrary that

$$\text{diam}(\gamma'_0[u'_1, u']) > \text{diam}(\gamma'_0[u', y'_0]). \quad (5.13)$$

Under this contrary assumption, we shall find an upper bound for the quantity $k_{D'}(u', y'_0)$. Since Proposition 3.4(P_2) implies that the image γ'_0 of γ_0 under f is (c_1, h_1) -solid, Theorem 2.10(1) and (5.13) yield

$$|y'_0 - u'| \leq \text{diam}(\gamma'_0[u', y'_0]) \leq c_2 d_{D'}(u'). \quad (5.14)$$

Since $k_D(v_1, y_0) \geq 2c_2 c_4 M \log c_3$, Lemma 2.6 gives

$$k_D(u_1, y_0) \geq \ell_k(\gamma_0) - \frac{1}{4} \geq k_D(v_1, y_0) - \frac{1}{4} \geq 2c_2 c_4 M \log c_3 - \frac{1}{4}.$$

Since f is (M, C) -CQH, the previous estimate implies

$$k_{D'}(u'_1, y'_0) \geq \frac{1}{M} (k_D(u_1, y_0) - C) \geq 2c_2 c_4 \log c_3 - \frac{1 + 4M}{4M} > 1.$$

Thus, by Lemma 2.2(1), we have

$$\max\{d_{D'}(u'_1), d_{D'}(y'_0)\} \leq 2|u'_1 - y'_0|,$$

which, together with Theorem 2.10(2), shows that

$$\begin{aligned} |y'_0 - u'| &\leq \text{diam}(\gamma'_0) \\ &\leq c_2 \max\{|u'_1 - y'_0|, 2(e^{h_1} - 1) \min\{d_{D'}(u'_1), d_{D'}(y'_0)\}\} \\ &\leq 4c_2 e^{h_1} |u'_1 - y'_0|. \end{aligned} \quad (5.15)$$

Moreover, by the choice of y_1 and (5.5), we have

$$|u'_1 - y'_0| \leq |u'_1 - x'_1| + |x'_1 - y'_0| \leq \frac{3}{2}|x'_1 - y'_1|,$$

and thus, we infer from (5.6) and (5.15) that

$$|y'_0 - u'| \leq 6c_2 e^{h_1} |x'_1 - y'_1| \leq 6c_2 c_3 e^{h_1} d_{D'}(y'_0).$$

Then we get from Theorem 2.14, the above estimate and (5.14) that

$$k_{D'}(u', y'_0) \leq c_4 \log \left(1 + \frac{|y'_0 - u'|}{\min\{d_{D'}(u'), d_{D'}(y'_0)\}} \right) \leq c_4 \log(1 + 6c_2 c_3 e^{h_1}). \quad (5.16)$$

Since

$$\frac{16}{9} c_2 c_4 \log c_3 - \frac{C}{M} > c_4 \log(1 + 6c_2 c_3 e^{h_1}),$$

we see that (5.16) contradicts with (5.11). Thus (5.12) holds, and consequently, we infer from Theorem 2.10(1) that

$$\text{diam}(\gamma'_0[u'_1, u']) \leq c_2 d_{D'}(u').$$

The proof is complete. \square

5.2. ζ -sequence on $\alpha[x_1, y_1]$. In this section, we shall construct a finite sequence of points on α , called ζ -sequence, with good controls on the length of λ -curves between successive points.

Proposition 5.4. *There exist an integer $m \geq 2$ and a sequence $\{x_i\}_{i=1}^{m+1}$ of points on α such that the following conclusions hold.*

(a) *For each $i \in \{2, \dots, m\}$ and for any λ -curve $\alpha_{i-1} \in \Lambda_{x_{i-1}x_i}^\lambda$, we have*

$$\mu_9^{\frac{1}{4}} \ell(\alpha) \leq \ell(\alpha_{i-1}) \leq \mu_9^{\frac{1}{2}} \ell(\alpha).$$

(b) *For any $z \in \alpha[x_i, y_1]$, and for each λ -curve $\varsigma_{i-1} \in \Lambda_{x_{i-1}z}^\lambda$,*


$$\ell(\varsigma_{i-1}) > \mu_9^{\frac{1}{4}} \ell(\alpha).$$

(c) *There is an element $\alpha_m \in \Lambda_{x_m x_{m+1}}^\lambda$ such that*

$$0 < \ell(\alpha_m) \leq \mu_9^{\frac{1}{2}} \ell(\alpha).$$

Proof. By Proposition 3.4(P_3), D is δ -Gromov hyperbolic. It follows from (5.2) and Lemma 5.2 that the assumptions in Lemma 2.29 are satisfied for any λ -curve $\varsigma \in \Lambda_{x_1 y_1}^\lambda$ and α . Then it follows that there is x_2 in $\alpha[x_1, y_1]$ such that for any λ -curve $\alpha_1 \in \Lambda_{x_1 x_2}^\lambda$, it holds

$$\mu_9^{\frac{1}{4}} \ell(\alpha) \leq \ell(\alpha_1) \leq \mu_9^{\frac{1}{2}} \ell(\alpha), \quad (5.17)$$


 FIGURE 5. Illustration for the construction of ζ -sequence

and for any $z \in \alpha[x_2, y_1]$, there is a λ -curve $\varsigma_1 \in \Lambda_{x_1 z}^\lambda$ such that

$$\ell(\varsigma_1) \geq \mu_9^{\frac{1}{4}} \ell(\alpha).$$

If there is an element in $\varsigma_{21} \in \Lambda_{x_2 y_1}^\lambda$ such that $\ell(\varsigma_{21}) \leq \mu_9^{\frac{1}{2}} \ell(\alpha)$, then we take $m = 2$, $x_3 = y_1$ and $\alpha_2 = \varsigma_{21}$. Then $\{x_1, x_2, x_3\}$ and α_2 are the desired points and λ -curve satisfying all the requirements of Proposition 5.4.

If for any λ -curve $\varsigma_{21} \in \Lambda_{x_2 y_1}^\lambda$, $\ell(\varsigma_{21}) > \mu_9^{\frac{1}{2}} \ell(\alpha)$, then the similar reasoning as above ensures that there is $x_3 \in \alpha[x_2, y_1]$ such that for any λ -curve $\alpha_2 \in \Lambda_{x_2 x_3}^\lambda$,

$$\mu_9^{\frac{1}{4}} \ell(\alpha) \leq \ell(\alpha_2) \leq \mu_9^{\frac{1}{2}} \ell(\alpha),$$

and for any $z \in \alpha[x_3, y_1]$, and for each λ -curve $\varsigma_2 \in \Lambda_{x_2 z}^\lambda$,

$$\ell(\varsigma_2) \geq \mu_9^{\frac{1}{4}} \ell(\alpha).$$

If there is an element $\varsigma_{31} \in \Lambda_{x_2 y_1}^\lambda$ such that $\ell(\varsigma_{31}) \leq \mu_9^{\frac{1}{2}} \ell(\alpha)$, then we take $m = 3$, $x_4 = y_1$ and $\alpha_3 = \varsigma_{31}$. Then $\{x_1, x_2, x_3, x_4\}$ and α_3 are the desired points and λ -curve satisfying all the requirements of Proposition 5.4.

If for any λ -curve $\varsigma_{31} \in \Lambda_{x_3 y_1}^\lambda$, $\ell(\varsigma_{31}) > \mu_9^{\frac{1}{2}} \ell(\alpha)$, then the similar reasoning as above ensures that there is $x_4 \in \alpha[x_3, y_1]$ such that for any λ -curve $\alpha_3 \in \Lambda_{x_3 x_4}^\lambda$,

$$\mu_9^{\frac{1}{4}} \ell(\alpha) \leq \ell(\alpha_3) \leq \mu_9^{\frac{1}{2}} \ell(\alpha).$$

As α is rectifiable, repeating the above procedure finitely many times, we obtain that there is a sequence of points $\{x_i\}_{i=1}^{m+1}$ on α and a λ -curve $\alpha_m \in \Lambda_{x_m x_{m+1}}^\lambda$ satisfying all the requirements of Proposition 5.4. \square

Lemma 5.5. *There is an element $\varsigma_m \in \Lambda_{x_{m-1} x_{m+1}}^\lambda$ such that*

$$\mu_9^{\frac{1}{4}} \ell(\alpha) < \ell(\varsigma_m) \leq \mu_9^{\frac{3}{4}} \ell(\alpha).$$

Proof. By Proposition 5.4(b), there is an element $\varsigma_m \in \Lambda_{x_{m-1} x_{m+1}}^\lambda$ such that

$$\ell(\varsigma_m) > \mu_9^{\frac{1}{4}} \ell(\alpha).$$

Next, we prove that the λ -curve ς_m simultaneously satisfies

$$\ell(\varsigma_m) \leq \mu_9^{\frac{3}{4}} \ell(\alpha). \quad (5.18)$$

Fix a λ -curves $\alpha_{m-1} \in \Lambda_{x_{m-1} x_m}^\lambda$ and let $\alpha_m \in \gamma_{x_m x_{m+1}}^\lambda$ be given by Proposition 5.4(c). Since (D, k_D) is a (μ_0, ρ_0) -Rips space by Proposition 3.4(P_4), we see from Lemma 2.24 that there exist $\eta_1 \in \alpha_{m-1}$, $\eta_2 \in \alpha_m$ and $\eta_3 \in \varsigma_m$ such that for any $i \neq j \in \{1, 2, 3\}$,

$$k_D(\eta_i, \eta_j) \leq \mu_1, \quad (5.19)$$

where $\mu_1 = 3\mu_0$.

Next, we estimate $\ell(\varsigma_m[x_{m+1}, \eta_3])$ and $\ell(\varsigma_m[x_{m-1}, \eta_3])$. To this end, we divide our discussions into two cases: $\min\{k_D(x_{m+1}, \eta_2), k_D(x_{m+1}, \eta_3)\} > 1$ and $\min\{k_D(x_{m+1}, \eta_2), k_D(x_{m+1}, \eta_3)\} \leq 1$. In the former case, we consider the points η_2, η_3 and x_{m+1} . Since Proposition 3.4(P_3) implies that D is δ -Gromov hyperbolic, we infer from (5.19), Lemma 2.27 and Proposition 5.4(c) that

$$\ell(\varsigma_m[x_{m+1}, \eta_3]) \leq 4e^{25\mu_2} \ell(\alpha_m[x_{m+1}, \eta_2]) \leq 4\mu_9^{\frac{1}{2}} e^{25\mu_2} \ell(\alpha),$$

where in the first inequality, we used the fact $\mu_1 \leq \mu_2$.

In the latter case, that is, $\min\{k_D(x_{m+1}, \eta_2), k_D(x_{m+1}, \eta_3)\} \leq 1$, it follows from (5.19) that

$$k_D(x_{m+1}, \eta_3) \leq \min\{k_D(x_{m+1}, \eta_2), k_D(x_{m+1}, \eta_3)\} + k_D(\eta_2, \eta_3) \leq 1 + \mu_1.$$

Since $\varsigma_m[x_{m+1}, \eta_3]$ is a 2λ -curve by Lemma 2.7, we infer from (2.2) and Proposition 3.4(P_1) that

$$\begin{aligned} \log \left(1 + \frac{\ell(\varsigma_m[x_{m+1}, \eta_3])}{d_D(x_{m+1})} \right) &\leq \ell_k(\varsigma_m[x_{m+1}, \eta_3]) \leq \frac{9}{8} k_D(x_{m+1}, \eta_3) \\ &\leq \frac{9}{8} (1 + \mu_1), \end{aligned}$$

which, together with (5.2), implies that

$$\ell(\varsigma_m[x_{m+1}, \eta_3]) \leq e^{\frac{9}{8}(1+\mu_1)} d_D(x_{m+1}) \leq 2e^{\frac{9}{8}(1+\mu_1)} \ell(\alpha).$$

In either cases, we obtain

$$\ell(\varsigma_m[x_{m+1}, \eta_3]) \leq 4\mu_9^{\frac{1}{2}} e^{25\mu_2} \ell(\alpha). \quad (5.20)$$

Similarly, repeating the above arguments with x_{m+1}, η_2 replaced by x_{m-1}, η_1 , we obtain

$$\ell(\varsigma_m[x_{m-1}, \eta_3]) \leq 4\mu_9^{\frac{1}{2}} e^{25\mu_2} \ell(\alpha). \quad (5.21)$$

Since $8\mu_9^{\frac{1}{2}} e^{25\mu_2} \leq \mu_9^{\frac{3}{4}}$, (5.18) follows from (5.20) and (5.21), and hence, the proof of lemma is complete. \square

For convenience, we make the following notational conventions.

- (1) for each $i \in \{1, 2, \dots, m-1\}$, let $\zeta_i = x_i$ and $\zeta_m = x_{m+1}$.
- (2) For each $i \in \{1, 2, \dots, m-2\}$, we fix an element $\alpha_i \in \Lambda_{\zeta_i \zeta_{i+1}}^\lambda$. Moreover, let $\alpha_{m-1} = \varsigma_m \in \Lambda_{x_{m-1} x_{m+1}}^\lambda$ be given by Lemma 5.5, and $\alpha_m \in \Lambda_{x_m x_{m+1}}^\lambda$ be given by Proposition 5.4(c).

Then we see from Proposition 5.4(a) and Lemma 5.5 that for each $i \in \{1, 2, \dots, m-2\}$,

$$\mu_9^{\frac{1}{4}} \ell(\alpha) < \ell(\alpha_i) \leq \mu_9^{\frac{1}{2}} \ell(\alpha) \quad \text{and} \quad \mu_9^{\frac{1}{4}} \ell(\alpha) < \ell(\alpha_{m-1}) \leq \mu_9^{\frac{3}{4}} \ell(\alpha), \quad (5.22)$$

and thus, we infer from (2.2) and (5.2) that for each $i \in \{1, 2, \dots, m-1\}$,

$$\ell_k(\alpha_i) \geq \log \left(1 + \frac{\ell(\alpha_i)}{d_D(\zeta_i)} \right) \geq \log \left(1 + \frac{\ell(\alpha_i)}{2\ell(\alpha)} \right) > \frac{1}{4} \log \frac{\mu_9}{16}. \quad (5.23)$$

5.3. λ -curves associated to ζ -sequence. Fix any λ -curve $\varsigma \in \Lambda_{xy_1}^\lambda$ and write $\gamma_1 = \varsigma[\zeta_1, y_0]$ and $\gamma_m = \varsigma[y_1, y_0]$. For $i \in \{2, 3, \dots, m-1\}$, let $\gamma_i \in \Lambda_{\zeta_i y_0}^\lambda$ be an element; see Figure 5.

Since, by Proposition 3.4(P_4), (D, k_D) is a (μ_0, ρ_0) -Rips space, it follows from Lemma 2.24 that for each $i \in \{1, 2, \dots, m-1\}$, there exist three points $w_i \in \alpha_i$, $w_{i,1} \in \gamma_i$, $w_{i,2} \in \gamma_{i+1}$ such that

$$k_D(w_i, w_{i,1}) \leq \mu_1, \quad k_D(w_{i,1}, w_{i,2}) \leq \mu_1 \quad \text{and} \quad k_D(w_i, w_{i,2}) \leq \mu_1. \quad (5.24)$$

Lemma 5.6. *For each $i \in \{1, 2, \dots, m-1\}$, we have*

$$\max\{\ell(\gamma_i[\zeta_i, w_{i,1}]), \ell(\gamma_{i+1}[\zeta_{i+1}, w_{i,2}])\} \leq 4\mu_9^{\frac{3}{4}} e^{25\mu_2} \ell(\alpha).$$

Proof. Without loss of generality, we assume $\max\{\ell(\gamma_i[\zeta_i, w_{i,1}]), \ell(\gamma_{i+1}[\zeta_{i+1}, w_{i,2}])\} = \ell(\gamma_i[\zeta_i, w_{i,1}])$, since the proof of other case is similar.

We first assume that $\min\{k_D(\zeta_i, w_i), k_D(\zeta_i, w_{i,1})\} > 1$. It follows from (5.24), Proposition 3.4(P_1), and Lemma 2.27 (applied to ζ_i, w_i and $w_{i,1}$) that

$$\ell(\gamma_i[\zeta_i, w_{i,1}]) \leq 4e^{25\mu_2} \ell(\alpha_i[\zeta_i, w_i]) \stackrel{(5.22)}{\leq} 4\mu_9^{\frac{3}{4}} e^{25\mu_2} \ell(\alpha).$$

In the remaining case, that is, $\min\{k_D(\zeta_i, w_i), k_D(\zeta_i, w_{i,1})\} \leq 1$, it follows from (5.24), (2.2) and Proposition 3.4(P_1) that

$$\begin{aligned} \log \left(1 + \frac{\ell(\gamma_i[\zeta_i, w_{i,1}])}{d_D(\zeta_i)} \right) &\leq \ell_k(\gamma_i[\zeta_i, w_{i,1}]) \leq \frac{9}{8} k_D(\zeta_i, w_{i,1}) \\ &\leq \frac{9}{8} \min\{k_D(\zeta_i, w_i), k_D(\zeta_i, w_{i,1})\} + k_D(w_i, w_{i,1}) \leq \frac{9}{8} (1 + \mu_1), \end{aligned}$$

which, together with (5.2), implies that

$$\ell(\gamma_i[\zeta_i, w_{i,1}]) \leq e^{\frac{9}{8}(1+\mu_1)} d_D(\zeta_i) \leq 2e^{\frac{9}{8}(1+\mu_1)} \ell(\alpha) < 4\mu_9^{\frac{3}{4}} e^{25\mu_2} \ell(\alpha).$$

In either cases, the proof is complete. \square

Next, we are going to find one more point on every γ_i . As Lemma 5.1(2) and (5.2) ensure that for each $i \in \{1, \dots, m\}$,

$$\ell(\gamma_i) \geq d_D(y_0) - d_D(\zeta_i) \geq (1 + c_3)^{-(c_4M+C)} d_D(x_0) - 2\ell(\alpha),$$

we know from (3.1) and (5.3) that

$$\ell(\gamma_i) > \left(\frac{\mu_8}{2\mu_7(1+c_3)^{c_4M+C}} - 2 \right) \ell(\alpha) > 2\mu_9 \ell(\alpha).$$

Thus there exists $w_{0,i} \in \gamma_i$ such that

$$\ell(\gamma_i[\zeta_i, w_{0,i}]) = \mu_9 \ell(\alpha) > 4\mu_9^{\frac{3}{4}} e^{25\mu_2} \ell(\alpha). \quad (5.25)$$

Furthermore, we infer from (5.25) and Lemma 5.6 that $w_{0,1} \in \gamma_1[w_{1,1}, y_0]$, $w_{0,m} \in \gamma_m[w_{m-1,2}, y_0]$, and for each $i \in \{2, \dots, m-1\}$,

$$w_{0,i} \in \gamma_i[w_{i,1}, y_0] \cap \gamma_i[w_{i-1,2}, y_0].$$

Lemma 5.7. *For each $i \in \{1, \dots, m-1\}$,*

$$\min\{k_D(w_{i,1}, w_{0,i}), k_D(w_{i,2}, w_{0,i+1})\} \geq \frac{1}{5} \log \mu_9.$$

Proof. Let $i \in \{1, \dots, m-1\}$. Since Lemma 5.6 and (5.2) lead to

$$d_D(w_{i,1}) \leq \ell(\gamma_i[\zeta_i, w_{i,1}]) + d_D(\zeta_i) < 2(1 + 2\mu_9^{\frac{3}{4}} e^{25\mu_2}) \ell(\alpha),$$

and since Lemma 5.6 and (5.25) give

$$\ell(\gamma_i[w_{i,1}, w_{0,i}]) = \ell(\gamma_i[\zeta_i, w_{0,i}]) - \ell(\gamma_i[\zeta_i, w_{i,1}]) \geq (\mu_9 - 4\mu_9^{\frac{3}{4}} e^{25\mu_2}) \ell(\alpha),$$

we get from Proposition 3.4(P_1), (2.2) and the above two estimates that

$$k_D(w_{i,1}, w_{0,i}) \geq \frac{8}{9} \ell_k(\gamma_i[w_{i,1}, w_{0,i}]) \geq \frac{8}{9} \log \left(1 + \frac{\ell(\gamma_i[w_{i,1}, w_{0,i}])}{d_D(w_{i,1})} \right) \geq \frac{1}{5} \log \mu_9.$$

Arguing similarly as above, replacing γ_i by γ_{i+1} , we obtain

$$k_D(w_{i,2}, w_{0,i+1}) \geq \frac{1}{5} \log \mu_9.$$

This completes the proof of lemma. \square

Lemma 5.8. *For each $i \in \{1, \dots, m-1\}$,*

$$\min\{k_D(y_0, w_{i,1}) - k_D(w_{0,i}, y_0), k_D(y_0, w_{i,2}) - k_D(w_{0,i+1}, y_0)\} \geq \frac{1}{5} \log \mu_9 - \frac{1}{2}.$$

Proof. By Lemmas 2.6, 2.7 and 5.7, we see that for each $i \in \{1, \dots, m-1\}$,

$$\begin{aligned} k_D(y_0, w_{i,1}) &\geq \ell_k(\gamma_i[w_{i,1}, y_0]) - \frac{1}{2} \\ &= \ell_k(\gamma_i[w_{i,1}, w_{0,i}]) + \ell_k(\gamma_i[w_{0,i}, y_0]) - \frac{1}{2} \\ &\geq k_D(w_{0,i}, y_0) + \frac{1}{5} \log \mu_9 - \frac{1}{2}. \end{aligned}$$

Similarly, we can show that for each $i \in \{1, \dots, m-1\}$,

$$k_D(y_0, w_{i,2}) \geq k_D(w_{0,i+1}, y_0) + \frac{1}{5} \log \mu_9 - \frac{1}{2}.$$

This completes the proof. \square

Let $t_1 \in \{1, \dots, m\}$ be such that

$$k_D(y_0, w_{0,t_1}) = \min_{i \in \{1, \dots, m\}} \{k_D(y_0, w_{0,i})\}. \quad (5.26)$$

Then we have the following assertion.

Lemma 5.9. *Suppose that there exist $i \in \{1, \dots, m\}$ and $u_i \in \gamma_i[w_{0,i}, y_0]$ such that $k_D(y_0, u_i) = k_D(y_0, w_{0,t_1})$.*

(1) *If $i \in \{2, \dots, m\}$, then there exists $u_{i-1} \in \gamma_{i-1}$ such that*

$$k_D(y_0, u_{i-1}) = k_D(y_0, w_{0,t_1}) \quad \text{and} \quad k_D(u_i, u_{i-1}) \leq \frac{3}{2} + 2\mu_0 + 2\mu_1.$$

(2) *If $i \in \{1, \dots, m-1\}$, then there exists $u_{i+1} \in \gamma_{i+1}$ such that*

$$k_D(y_0, u_{i+1}) = k_D(y_0, w_{0,t_1}) \quad \text{and} \quad k_D(u_i, u_{i+1}) \leq \frac{3}{2} + 2\mu_0 + 2\mu_1.$$

Proof. We only need to prove (1) since the proof of (2) is similar. Let $i \in \{2, \dots, m\}$. To find the asserted point u_{i-1} , we need some preparation.

Firstly, we determine the position of u_i on γ_i . Since Lemmas 2.6 and 2.7 lead to

$$\ell_k(\gamma_i[y_0, u_i]) \leq k_D(y_0, u_i) + \frac{1}{2},$$

and since Lemma 5.8 and (5.26) give

$$k_D(y_0, w_{i,1}) \geq k_D(w_{0,i}, y_0) + \frac{1}{5} \log \mu_9 - \frac{1}{2} \geq k_D(y_0, u_i) + \frac{1}{5} \log \mu_9 - \frac{1}{2},$$

we obtain from the above two estimates that for $i \in \{2, \dots, m-1\}$, it holds

$$\ell_k(\gamma_i[y_0, w_{i,1}]) \geq k_D(y_0, w_{i,1}) \geq \ell_k(\gamma_i[y_0, u_i]) + \frac{1}{5} \log \mu_9 - 1.$$

Similarly, we know that for $i \in \{2, \dots, m\}$,

$$\ell_k(\gamma_i[y_0, w_{i-1,2}]) \geq \ell_k(\gamma_i[y_0, u_i]) + \frac{1}{5} \log \mu_9 - 1.$$

These ensure that for $i \in \{2, \dots, m-1\}$,

$$u_i \in (\gamma_i[w_{i,1}, y_0] \setminus \{w_{i,1}\}) \cap (\gamma_i[w_{i-1,2}, y_0] \setminus \{w_{i-1,2}\}) \quad (5.27)$$

and

$$u_m \in \gamma_i[w_{m-1,2}, y_0] \setminus \{w_{m-1,2}\}. \quad (5.28)$$

Secondly, we find a point on γ_{i-1} . Since Proposition 3.4(P_4) implies that (D, k_D) is a (μ_0, ρ_0) -Rips space, it follows from (5.24) and Lemma 2.25 (replacing x, y and z with $y_0, w_{i-1,2}$ and $w_{i-1,1}$), together with (5.27) and (5.28), that there exists $v_{i-1} \in \gamma_{i-1}[w_{i-1,1}, y_0]$ such that

$$k_D(u_i, v_{i-1}) \leq \frac{1}{2} + \mu_0 + \mu_1. \quad (5.29)$$

Then we have

$$\begin{aligned} k_D(y_0, u_i) - \frac{1}{2} - \mu_0 - \mu_1 &\leq k_D(y_0, u_i) - k_D(u_i, v_{i-1}) \\ &\leq k_D(y_0, v_{i-1}) \leq k_D(y_0, u_i) + k_D(u_i, v_{i-1}) \\ &\leq k_D(y_0, u_i) + \frac{1}{2} + \mu_0 + \mu_1. \end{aligned} \quad (5.30)$$

Now, we are ready to find the required point. If $k_D(y_0, v_{i-1}) \leq k_D(y_0, u_i)$, then we infer from Lemma 5.8 and (5.26) that

$$k_D(y_0, w_{i-1,1}) \geq k_D(y_0, w_{0,i-1}) + \frac{1}{5} \log \mu_9 - \frac{1}{2} \geq k_D(y_0, u_i) + \frac{1}{5} \log \mu_9 - \frac{1}{2} > k_D(y_0, v_{i-1}),$$

from which we know that there exists $u_{i-1} \in \gamma_{i-1}[w_{i-1,1}, v_{i-1}]$ such that

$$k_D(y_0, u_{i-1}) = k_D(y_0, u_i).$$

Then it follows from Lemmas 2.6 and 2.7, and (5.30) that

$$\begin{aligned} k_D(u_{i-1}, v_{i-1}) &\leq \ell_k(\gamma_{i-1}[y_0, u_{i-1}]) - \ell_k(\gamma_{i-1}[y_0, v_{i-1}]) \\ &\leq k_D(y_0, u_{i-1}) - k_D(y_0, v_{i-1}) + \frac{1}{2} \\ &\leq 1 + \mu_0 + \mu_1, \end{aligned}$$

which, together with (5.29), gives

$$k_D(u_{i-1}, u_i) \leq k_D(u_{i-1}, v_{i-1}) + k_D(v_{i-1}, u_i) \leq \frac{3}{2} + 2\mu_0 + 2\mu_1.$$

If $k_D(y_0, v_{i-1}) > k_D(y_0, u_i)$, then select $u_{i-1} \in \gamma_{i-1}[y_0, v_{i-1}]$ such that

$$k_D(y_0, u_{i-1}) = k_D(y_0, u_i).$$

Then a similar computation as above gives

$$k_D(u_{i-1}, u_i) \leq k_D(u_{i-1}, v_{i-1}) + k_D(v_{i-1}, u_i) \leq \frac{3}{2} + 2\mu_0 + 2\mu_1.$$

The proof of the lemma is thus complete. \square

5.4. ξ -sequence on the associated λ -curves. In this section, we shall find a new sequence of points. By the choice of w_{0,t_1} in (5.26) and Lemma 5.9, we know that for each $i \in \{1, \dots, m\}$, there exists $\xi_i \in \gamma_i[w_{0,i}, y_0]$ (see Figure 5) such that

$$k_D(y_0, \xi_i) = k_D(y_0, w_{0,t_1}), \quad (5.31)$$

and for every $i \in \{1, \dots, m-1\}$,

$$k_D(\xi_i, \xi_{i+1}) \leq \frac{3}{2} + 2\mu_0 + 2\mu_1. \quad (5.32)$$

Lemma 5.10. (i) For each $i \in \{1, \dots, m\}$ and all $z \in \gamma_i[\zeta_i, \xi_i]$, it holds

$$k_D(y_0, z) \geq \log \frac{\mu_8}{2(2 + \mu_9)(1 + c_3)^{c_4 M} e^C \mu_7} - \frac{1}{2}.$$

(ii) Let $i \in \{1, \dots, m-1\}$. For any λ -curve $\beta_i \in \Lambda_{\xi_i, \xi_{i+1}}^\lambda$, it holds

$$\ell_k(\beta_i) \leq \frac{7}{4} + 2\mu_0 + 2\mu_1.$$

(iii) Let $i, j \in \{1, \dots, m-1\}$. For any λ -curve $\beta_i \in \Lambda_{\xi_i, \xi_{i+1}}^\lambda$ and any $u \in \beta_i$, it holds

$$k_D(y_0, \xi_j) - \frac{7}{4} - 2\mu_0 - 2\mu_1 \leq k_D(y_0, u) \leq k_D(y_0, \xi_j) + \frac{7}{4} + 2\mu_0 + 2\mu_1.$$

Proof. (i). First of all, for each $i \in \{1, \dots, m\}$, we have

$$\begin{aligned} d_D(y_0) &\stackrel{\text{Lemma 5.1(2)}}{\geq} (1 + c_3)^{-c_4 M} e^{-C} d_D(x_0) \\ &\stackrel{(3.1)}{\geq} ((1 + c_3)^{c_4 M} e^C \mu_7)^{-1} \ell(\gamma) \\ &\stackrel{(5.3)}{\geq} (2(1 + c_3)^{c_4 M} e^C \mu_7)^{-1} \mu_8 \ell(\alpha). \end{aligned} \tag{5.33}$$

Secondly, we get from (5.2) and (5.25) that

$$d_D(w_0, t_1) \leq d_D(\zeta_{t_1}) + \ell(\gamma_{t_1}[\zeta_{t_1}, w_0, t_1]) \leq (2 + \mu_9) \ell(\alpha). \tag{5.34}$$

Let $z \in \gamma_i[\zeta_i, \xi_i]$. Then we obtain from Lemmas 2.6 and 2.7 that

$$\begin{aligned} k_D(y_0, z) &\geq \ell_k(\gamma_i[y_0, z]) - \frac{1}{2} \geq \ell_k(\gamma_i[y_0, \xi_i]) - \frac{1}{2} \\ &\geq k_D(y_0, \xi_i) - \frac{1}{2} \stackrel{(5.31)}{=} k_D(y_0, w_0, t_1) - \frac{1}{2} \stackrel{(2.3)}{\geq} \log \frac{d_D(y_0)}{d_D(w_0, t_1)} - \frac{1}{2} \\ &\stackrel{(5.33)+(5.34)}{\geq} \log \frac{\mu_8}{2(2 + \mu_9)(1 + c_3)^{c_4 M} e^C \mu_7} - \frac{1}{2}. \end{aligned}$$

This proves (i).

(ii). Let $i \in \{1, \dots, m-1\}$. For any λ -curve $\beta_i \in \Lambda_{\xi_i, \xi_{i+1}}^\lambda$, we have by (5.32) and Lemma 2.6 that

$$\ell_k(\beta_i) \leq k_D(\xi_i, \xi_{i+1}) + \frac{1}{4} \leq \frac{7}{4} + 2\mu_0 + 2\mu_1,$$

which proves (ii).

(iii). Let $i \in \{1, \dots, m-1\}$. For any λ -curve $\beta_i \in \Lambda_{\xi_i, \xi_{i+1}}^\lambda$, let $u \in \beta_i$. By (ii), we know that

$$\begin{aligned} k_D(y_0, \xi_i) - \frac{7}{4} - 2\mu_0 - 2\mu_1 &\leq k_D(y_0, \xi_i) - \ell_k(\beta_i) \\ &\leq k_D(y_0, u) \leq k_D(y_0, \xi_i) + \ell_k(\beta_i) \\ &\leq k_D(y_0, \xi_i) + \frac{7}{4} + 2\mu_0 + 2\mu_1. \end{aligned} \tag{5.35}$$

Since for any $i, j \in \{1, \dots, m-1\}$, $k_D(y_0, \xi_i) = k_D(y_0, \xi_j)$ by (5.31), statement (iii) follows immediately from (5.35). The proof of lemma is thus complete. \square

In the following, for each $i \in \{1, \dots, m\}$, let $\{\xi_i\}$ be the sequence given by (5.31). Then for each $i \in \{1, \dots, m-1\}$, we fix a λ -curve $\beta_i \in \Lambda_{\xi_i, \xi_{i+1}}^\lambda$ and set $\beta_0 = \bigcup_{i=1}^{m-1} \beta_i$; see Figure 5. Fix




FIGURE 6. Illustration for the proofs of Lemmas 5.11 and 5.12

$u, v \in \beta_0$. Then there are $i_1, i_2 \in \{1, \dots, m-1\}$ such that $u \in \beta_{i_1}$ and $v \in \beta_{i_2}$ (see Figure 6). The following result gives a sufficient condition for i_1 and i_2 to be different.

Lemma 5.11. *Let $u \in \beta_{i_1}$ and $v \in \beta_{i_2}$, where $i_1, i_2 \in \{1, \dots, m-1\}$. If $k_D(u, v) \geq 3\mu_1$, then $i_1 \neq i_2$.*

Proof. Suppose on the contrary that $i_1 = i_2$, that is, $u, v \in \beta_{i_1}$. Then we know from Lemma 5.10(ii) that

$$k_D(u, v) \leq \ell_k(\beta_{i_1}) \leq 2 + 2\mu_0 + 2\mu_1 < 3\mu_1.$$

This contradiction implies that $i_1 \neq i_2$. \square

Lemma 5.12. *Suppose that $i_1, i_2 \in \{1, \dots, m-1\}$ with $i_1 < i_2$, $u \in \beta_{i_1}$ and $v \in \beta_{i_2}$. Fix $\varsigma_1 \in \Lambda_{uy_0}^\lambda$, $\varsigma_2 \in \Lambda_{vy_0}^\lambda$ and $\varsigma_3 \in \Lambda_{uv}^\lambda$. Then for each $i \in \{1, 2, 3\}$, there exists a point $\eta_{i_1 i_2, i} \in \varsigma_i$ such that*

(1) for any $r \neq s \in \{1, 2, 3\}$,

$$k_D(\eta_{i_1 i_2, r}, \eta_{i_1 i_2, s}) \leq \mu_1;$$

(2) $k_D(u, \eta_{i_1 i_2, 1}) - \frac{15}{4} - \mu_0 - 6\mu_1 \leq k_D(v, \eta_{i_1 i_2, 2}) \leq k_D(u, \eta_{i_1 i_2, 1}) + \frac{15}{4} + \mu_0 + 6\mu_1;$

(3) $\frac{1}{2}k_D(u, v) - \frac{15}{8} - \frac{1}{2}\mu_0 - 4\mu_1 \leq k_D(u, \eta_{i_1 i_2, 3}) \leq \frac{1}{2}k_D(u, v) + 2 + \frac{1}{2}\mu_0 + 4\mu_1;$

(4) $\frac{1}{2}k_D(u, v) - \frac{15}{8} - \frac{1}{2}\mu_0 - 4\mu_1 \leq k_D(v, \eta_{i_1 i_2, 3}) \leq \frac{1}{2}k_D(u, v) + 2 + \frac{1}{2}\mu_0 + 4\mu_1.$

Proof. (1) Since Proposition 3.4(P_4) implies that (D, k_D) is a (μ_0, ρ_0) -Rips space, we see from Lemma 2.24 that for each $i \in \{1, 2, 3\}$, there exist $\eta_{i_1 i_2, i} \in \varsigma_i$ so that statement (1) holds.

(2) By Lemma 5.10(ii), we have

$$k_D(y_0, \xi_{i_1+1}) - \frac{7}{4} - 2\mu_0 - 2\mu_1 \leq k_D(y_0, \xi_{i_1+1}) - \ell_k(\beta_{i_1})$$

$$\begin{aligned}
&\leq k_D(y_0, u) \leq k_D(y_0, \xi_{i_1+1}) + \ell_k(\beta_{i_1}) \\
&\leq k_D(y_0, \xi_{i_1+1}) + \frac{7}{4} + 2\mu_0 + 2\mu_1,
\end{aligned}$$

and

$$\begin{aligned}
k_D(y_0, \xi_{i_2}) - \frac{7}{4} - 2\mu_0 - 2\mu_1 &\leq k_D(y_0, \xi_{i_2}) - \ell_k(\beta_{i_2}) \\
&\leq k_D(y_0, v) \leq k_D(y_0, \xi_{i_2}) + \ell_k(\beta_{i_2}) \\
&\leq k_D(y_0, \xi_{i_2}) + \frac{7}{4} + 2\mu_0 + 2\mu_1.
\end{aligned}$$

These, together with (5.31), give

$$k_D(y_0, u) - \frac{7}{2} - \mu_0 - 5\mu_1 \leq k_D(y_0, v) \leq k_D(y_0, u) + \frac{7}{2} + \mu_0 + 5\mu_1. \quad (5.36)$$

Moreover, we know from statement (1) of the lemma that

$$k_D(y_0, \eta_{i_1 i_2, 1}) - \mu_1 \leq k_D(y_0, \eta_{i_1 i_2, 2}) \leq k_D(y_0, \eta_{i_1 i_2, 1}) + \mu_1. \quad (5.37)$$

Since $\varsigma_1 \in \Lambda_{uy_0}^\lambda$, we have

$$\begin{aligned}
k_D(u, \eta_{i_1 i_2, 1}) &\leq \ell_k(\varsigma_1[u, \eta_{i_1 i_2, 1}]) = \ell_k(\varsigma_1) - \ell_k(\varsigma_1[y_0, \eta_{i_1 i_2, 1}]) \\
&\stackrel{\text{Lemma 2.6+(5.37)}}{\leq} k_D(y_0, u) - k_D(y_0, \eta_{i_1 i_2, 2}) + \frac{1}{4} + \mu_1 \\
&\stackrel{(5.36)}{\leq} k_D(y_0, v) - k_D(y_0, \eta_{i_1 i_2, 2}) + \frac{15}{4} + \mu_0 + 6\mu_1 \\
&\leq k_D(v, \eta_{i_1 i_2, 2}) + \frac{15}{4} + \mu_0 + 6\mu_1.
\end{aligned}$$

Similarly, we have

$$k_D(v, \eta_{i_1 i_2, 2}) \leq k_D(u, \eta_{i_1 i_2, 1}) + \frac{15}{4} + \mu_0 + 6\mu_1.$$

These prove the second statement (2).

(3) To prove the third statement of the lemma, we note first that

$$\begin{aligned}
k_D(u, \eta_{i_1 i_2, 3}) &\leq k_D(u, \eta_{i_1 i_2, 1}) + \mu_1 && \text{(by (1) of the lemma)} \\
&\leq k_D(v, \eta_{i_1 i_2, 2}) + \frac{15}{4} + \mu_0 + 7\mu_1 && \text{(by (2) of the lemma)} \\
&\leq k_D(v, \eta_{i_1 i_2, 3}) + \frac{15}{4} + \mu_0 + 8\mu_1 && \text{(by (1) of the lemma)}
\end{aligned}$$

Similarly, we may prove

$$k_D(u, \eta_{i_1 i_2, 3}) \geq k_D(v, \eta_{i_1 i_2, 3}) - \frac{15}{4} - \mu_0 - 8\mu_1.$$

These ensure that

$$\begin{aligned}
k_D(u, v) - k_D(u, \eta_{i_1 i_2, 3}) - \frac{15}{4} - \mu_0 - 8\mu_1 &\leq k_D(u, v) - k_D(v, \eta_{i_1 i_2, 3}) \\
&\leq k_D(u, \eta_{i_1 i_2, 3}) \leq \ell_k(\varsigma_3) - \ell_k(\varsigma_3[v, \eta_{i_1 i_2, 3}]) \\
&\leq k_D(u, v) - k_D(v, \eta_{i_1 i_2, 3}) + \frac{1}{4} \\
&\leq k_D(u, v) - k_D(u, \eta_{i_1 i_2, 3}) + 4 + \mu_0 + 8\mu_1.
\end{aligned}$$

Consequently, we get

$$\frac{1}{2}k_D(u, v) - \frac{15}{8} - \frac{1}{2}\mu_0 - 4\mu_1 \leq k_D(u, \eta_{i_1 i_2, 3}) \leq \frac{1}{2}k_D(u, v) + 2 + \frac{1}{2}\mu_0 + 4\mu_1.$$

This proves (3).

(4) The proof of (4) is similar to the one given above and thus it is omitted here.

This completes the proof of lemma. \square

5.5. Six-tuples with Property C. In this section, we use \mathfrak{c} to denote a constant with $\mathfrak{c} \geq 64\mu_1$. Given β_0 , ξ -sequence as in previous sections, we first introduce the following useful definition.

Definition 5.13 (Six-tuple). We call $[w_1, w_2, \gamma_{12}, \varpi, w_3, w_4]$ a *six-tuple* (see Figure 7) if the following conditions are satisfied:

- $w_1 \in D$ and $w_2 \in \beta_0$;
- $\varpi \in \beta_0[w_2, \xi_m]$ and $\gamma_{12} \in \Lambda_{w_1 w_2}^\lambda$;
- $w_3 \in \gamma_{12}$ and $w_4 \in \beta_0[w_2, \varpi]$.

We usually use the symbol $\Omega = [w_1, w_2, \gamma_{12}, \varpi, w_3, w_4]$ to represent a six-tuple.

Property C. Given $\mathfrak{c} \geq 64\mu_1$, a six-tuple $\Omega = [w_1, w_2, \gamma_{12}, \varpi, w_3, w_4]$ is said to satisfy *Property C* with constant \mathfrak{c} if

- (1) $k_D(w_1, w_2) \leq \mathfrak{c}$ and $k_D(w_3, w_4) = \mathfrak{c}$;
- (2) for any $u \in \gamma_{12}$ and any $v \in \beta_0[w_4, \varpi]$, $k_D(u, v) \geq \mathfrak{c} - \frac{1}{4}$.

If a six-tuple Ω satisfies Property C with constant \mathfrak{c} , then we shall write $\Omega^\mathfrak{c} = \Omega$ to indicate its dependence on \mathfrak{c} .

In most situations, we shall simply say that $\Omega^\mathfrak{c} = [w_1, w_2, \gamma_{12}, \varpi, w_3, w_4]$ is a six-tuple with Property C to indicate that the six-tuple $[w_1, w_2, \gamma_{12}, \varpi, w_3, w_4]$ satisfies Property C with constant \mathfrak{c} .




FIGURE 7. Illustration for the definition of a six-tuple $\Omega = [w_1, w_2, \gamma_{12}, \varpi, w_3, w_4]$

The following result provides a sufficient condition for the existence of a six-tuples with Property C.

Lemma 5.14. *Let $w_1 \in D$, $w_2 \in \beta_0$ and $\gamma_{12} \in \Lambda_{w_1 w_2}^\lambda$. Suppose that $0 < k_D(w_1, w_2) \leq \mathbf{c}$ and there is $\varpi \in \beta_0[w_2, \xi_m]$ such that for any $w \in \gamma_{12}$, $k_D(w, \varpi) > \mathbf{c}$. Then there exist two points $w_3 \in \gamma_{12} \setminus \{w_2\}$ and $w_4 \in \beta_0[w_2, \varpi]$ such that $\Omega^c = [w_1, w_2, \gamma_{12}, \varpi, w_3, w_4]$ is a six-tuple with Property C.*

Proof. For the proof, it is helpful to revisit Figure 7 for a couple of times. For convenience, we write $\gamma = \gamma_{12}$ and set

$$\ell_k(\beta_0[w_2, \varpi]) = \nu. \quad (5.38)$$

Then $\nu > \mathbf{c}$.

We start the proof with the following claim.

Claim 5.1. *There are $u_1 \in \gamma \setminus \{w_2\}$ and $v_1 \in \beta_0[w_2, \varpi]$ such that*

- (1) $\ell_k(\beta_0[v_1, \varpi]) \leq \nu - \mathbf{c} + \frac{1}{8}$, and
- (2) *one of the following statements is true:*
 - (a) *The six-tuple $[w_1, w_2, \gamma, \varpi, u_1, v_1]$ satisfies Property C with constant \mathbf{c} .*
 - (b) *There exist $u_2 \in \gamma[u_1, w_1]$ and $q_1 \in \beta_0[v_1, \varpi]$ such that*

$$k_D(q_1, u_2) < \mathbf{c} - \frac{1}{4}.$$

To prove the claim, select $u_1 \in \gamma$ such that

$$\ell_k(\gamma[u_1, w_2]) = \frac{1}{8} \min \{1, k_D(w_1, w_2)\} \leq \frac{\mathbf{c}}{8}. \quad (5.39)$$

Then we see from the assumption $k_D(u_1, \varpi) > \mathbf{c}$ and the choice of u_1 in (5.39) that there exists $v_1 \in \beta_0[w_2, \varpi]$ such that

$$k_D(u_1, v_1) = \mathbf{c}, \quad (5.40)$$

and for any $w \in \beta_0[v_1, \varpi] \setminus \{v_1\}$,

$$k_D(u_1, w) > \mathbf{c}.$$

Since the choice of v_1 ensures that $k_D(u_1, v) \geq \mathbf{c}$, we infer from (5.39) that for any $u \in \gamma[u_1, w_2]$ and any $v \in \beta_0[v_1, \varpi]$, it holds

$$k_D(u, v) \geq k_D(v, u_1) - k_D(u_1, u) \geq k_D(v, u_1) - \ell_k(\gamma[u_1, w_2]) \geq \mathbf{c} - \frac{1}{8}. \quad (5.41)$$

This implies that

$$\ell_k(\beta_0[v_1, w_2]) \geq k_D(v_1, w_2) \geq \mathbf{c} - \frac{1}{8},$$

and so, (5.38) leads to

$$\ell_k(\beta_0[v_1, \varpi]) = \ell_k(\beta_0[w_2, \varpi]) - \ell_k(\beta_0[w_2, v_1]) \leq \nu - \mathbf{c} + \frac{1}{8}.$$

If for any $u \in \gamma[u_1, w_1]$ and any $v \in \beta_0[v_1, \varpi]$, we have

$$k_D(u, v) \geq \mathbf{c} - \frac{1}{4},$$

then it follows from (5.40), (5.41) and the assumption $0 < k_D(w_1, w_2) \leq \mathbf{c}$ that $[w_1, w_2, \gamma, \varpi, u_1, v_1]$ is a six-tuple with Property C. Otherwise, there must exist $u_2 \in \gamma[u_1, w_1]$ and $q_1 \in \beta_0[v_1, \varpi]$ such that

$$k_D(q_1, u_2) < \mathbf{c} - \frac{1}{4}.$$

This proves the claim.

If Claim 5.1(2a) occurs, by setting $w_3 = u_1$ and $w_4 = v_1$, we know that $[w_1, w_2, \gamma, \varpi, w_3, w_4]$ is the required six-tuple with Property C, and the proof of lemma is complete.

If Claim 5.1(2b) occurs, then we make the following claim.

Claim 5.2. *There is $v_2 \in \beta_0[q_1, \varpi]$ such that*

- (1) $\ell_k(\beta_0[v_2, \varpi]) \leq \nu - \mathbf{c} - \frac{1}{8}$, and
- (2) *one of the following statements is true:*
 - (a) *The six-tuple $[w_1, w_2, \gamma, \varpi, u_2, v_2]$ satisfies Property C with constant \mathbf{c} .*
 - (b) *There exist $u_3 \in \gamma[u_1, w_1]$ and $q_2 \in \beta_0[v_2, \varpi]$ such that*

$$k_D(q_2, u_3) < \mathbf{c} - \frac{1}{4}.$$

Since $k_D(u_2, \varpi) > \mathbf{c}$ and $k_D(q_1, u_2) < \mathbf{c} - \frac{1}{4}$, we see from Claim 5.1(2b) that there is $v_2 \in \beta_0[q_1, \varpi]$ such that

$$k_D(u_2, v_2) = \mathbf{c}, \tag{5.42}$$

and for any $w \in \beta_0[v_2, \varpi] \setminus \{v_2\}$,

$$k_D(u_2, w) > \mathbf{c}.$$

Then it follows from Claim 5.1(2b) that

$$\ell_k(\beta_0[v_1, v_2]) \geq k_D(q_1, v_2) \geq k_D(v_2, u_2) - k_D(u_2, q_1) > \frac{1}{4},$$

and thus, we infer from Claim 5.1(1) that

$$\ell_k(\beta_0[v_2, \varpi]) = \ell_k(\beta_0[v_1, \varpi]) - \ell_k(\beta_0[v_1, v_2]) \leq \nu - \mathbf{c} - \frac{1}{8}.$$

If for any $u \in \gamma[u_1, w_1]$ and any $v \in \beta_0[v_2, \varpi]$,

$$k_D(u, v) \geq \mathbf{c} - \frac{1}{4},$$

we see from (5.41), (5.42) and the assumption $0 < k_D(w_1, w_2) \leq \mathbf{c}$ that $[w_1, w_2, \gamma, \varpi, u_2, v_2]$ is a six-tuple with Property C. Otherwise, there are $u_3 \in \gamma[u_1, w_1]$ and $q_2 \in \beta_0[v_2, \varpi]$ such that

$$k_D(q_2, u_3) < \mathbf{c} - \frac{1}{4}.$$

This prove the claim.

If Claim 5.2(2a) occurs, by choosing $w_3 = u_2$ and $w_4 = v_2$, we know that $[w_1, w_2, \gamma, \varpi, w_3, w_4]$ is our required six-tuple satisfying Property C with constant \mathbf{c} , and thus the proof is complete.

If Claim 5.2(2b) occurs, then we make the following claim.

Claim 5.3. *There is $v_3 \in \beta_0[q_2, \varpi]$ such that*

- (1) $\ell_k(\beta_0[v_3, \varpi]) \leq \nu - \mathbf{c} - \frac{3}{8}$, and
- (2) *one of the following statements is true:*
 - (a) *The six-tuple $[w_1, w_2, \gamma, \varpi, u_3, v_3]$ satisfies Property C with constant \mathbf{c} .*
 - (b) *There exist $u_4 \in \gamma[u_1, w_1]$ and $q_3 \in \beta_0[v_3, \varpi]$ such that*

$$k_D(q_3, u_4) < \mathbf{c} - \frac{1}{4}.$$

Continuing this procedure for finitely many times, we arrive at the following conclusion.

Conclusion 5.4. *Let $a = 4\nu - 4\mathfrak{c} + \frac{1}{2}$ and $b = 4\nu - 4\mathfrak{c} + \frac{3}{2}$. Then there are an integer K with $K \in [a, b]$ and $v_K \in \beta_0[q_{K-1}, \varpi]$ such that*

- (1) $\ell_k(\beta_0[v_K, \varpi]) \leq \nu - \mathfrak{c} + \frac{1}{8} - \frac{K-1}{4} \leq \frac{1}{4}$, and
- (2) *one of the following statements is true:*
 - (a) *The six-tuple $[w_1, w_2, \gamma, \varpi, u_K, v_K]$ satisfies Property C with constant \mathfrak{c} .*
 - (b) *There exist $u_{K+1} \in \gamma[u_1, w_1]$ and $q_K \in \beta_0[v_K, \varpi]$ such that*

$$k_D(q_K, u_{K+1}) < \mathfrak{c} - \frac{1}{4}.$$

We assert that the statement (2b) in Conclusion 5.4 cannot occur. Otherwise, since $k_D(u_{K+1}, \varpi) > \mathfrak{c}$, we get that there is $v_{K+1} \in \beta_0[q_K, \varpi]$ such that

$$k_D(u_{K+1}, v_{K+1}) = \mathfrak{c},$$

which leads to

$$\ell_k(\beta_0[v_K, v_{K+1}]) \geq k_D(q_K, v_{K+1}) \geq k_D(v_{K+1}, u_{K+1}) - k_D(u_{K+1}, q_K) > \frac{1}{4}.$$

This contradicts with Conclusion 5.4(1) since $v_{K+1} \in \beta_0[v_K, \varpi]$.

Set $w_3 = u_K$ and $w_4 = v_K$. Then Conclusion 5.4(2a) implies that the six-tuple $[w_1, w_2, \gamma, \varpi, w_3, w_4]$ is our required six-tuple satisfying Property C, and thus, the proof is complete. \square

The following three lemmas concern basic properties of six-tuples with Property C.

Lemma 5.15. *Let γ_1 be given as in Section 5.2 (see Figure 5). Let $w_1 \in \gamma_1$ and $\gamma_{12} = \gamma_1[w_1, w_2]$, $w_2 = \xi_1$, $w_3 \in \gamma_{12}$, $\varpi \in \beta_0[w_2, \xi_m]$ and $w_4 \in \beta_0[w_2, \varpi]$ (see Figure 8). Suppose that the six-tuple $[w_1, w_2, \gamma_{12}, \varpi, w_3, w_4]$ satisfies Property C with constant \mathfrak{c} . Then $k_D(w_1, w_3) \leq 16\mu_1$.*




FIGURE 8. Illustration for the proof of Lemma 5.15

Proof. Fix two λ -curves $\varsigma_3 \in \Lambda_{w_3w_4}^\lambda$ and $\varsigma_4 \in \Lambda_{w_4y_0}^\lambda$. Since (D, k_D) is a (μ_0, ρ_0) -Rips space by Proposition 3.4(P_4), we know from Lemma 2.24 that there exist $u_1 \in \gamma_1[w_3, y_0]$, $u_2 \in \varsigma_3$ and $u_3 \in \varsigma_4$ (see Figure 8) such that for any $r \neq s \in \{1, 2, 3\}$,

$$k_D(u_r, u_s) \leq \mu_1. \quad (5.43)$$

Then we get

$$\begin{aligned} k_D(y_0, u_3) - \mu_1 &\leq k_D(y_0, u_3) - k_D(u_3, u_1) \leq k_D(y_0, u_1) \\ &\leq k_D(u_1, u_3) + k_D(u_3, y_0) \leq k_D(y_0, u_3) + \mu_1. \end{aligned} \quad (5.44)$$

We prove the lemma by contradiction. Suppose that

$$k_D(w_1, w_3) > 16\mu_1. \quad (5.45)$$

To reach a contradiction, we divide the discussions into two cases: $k_D(u_2, w_3) > 5\mu_1$ and $k_D(u_2, w_3) \leq 5\mu_1$. In the former case, there must exist $u_4 \in \varsigma_3[u_2, w_3]$ such that

$$k_D(u_4, w_3) = 3\mu_1. \quad (5.46)$$

Then we see from (5.43) and Lemma 2.25 (with x, y and z replaced by w_3, u_1 and u_2) that there exists $u_5 \in \gamma_1[w_3, u_1]$ such that

$$k_D(u_4, u_5) \leq 1 + \mu_0 + \mu_1. \quad (5.47)$$

We deduce from Lemma 2.7 and (5.46) that

$$\ell_k(\gamma_1[w_3, u_5]) \leq \frac{1}{2} + k_D(w_3, u_5) \leq \frac{1}{2} + k_D(w_3, u_4) + k_D(u_4, u_5) \leq \frac{3}{2} + \mu_0 + 4\mu_1,$$

and thus, (5.45) yields

$$\ell_k(\gamma_1[w_3, u_5]) < \ell_k(\gamma_1[w_3, w_1]).$$

This ensures that

$$u_5 \in \gamma_1[w_1, w_3] \setminus \{w_3\}. \quad (5.48)$$

Then we know from Lemma 2.6, (5.46) and (5.47) that

$$\begin{aligned} k_D(w_4, u_5) &\leq k_D(w_4, u_4) + k_D(u_4, u_5) \\ &\leq \ell_k(\varsigma_3) - k_D(u_4, w_3) + k_D(u_4, u_5) \\ &\leq k_D(w_3, w_4) + \frac{5}{4} + \mu_0 - 2\mu_1. \end{aligned} \quad (5.49)$$

On the other hand, since the six-tuple $[w_1, w_2, \gamma_{12}, \varpi, w_3, w_4]$ satisfies Property C with constant \mathfrak{c} , we obtain from (5.48) that

$$k_D(w_4, u_5) \geq \mathfrak{c} - \frac{1}{4} = k_D(w_3, w_4) - \frac{1}{4}.$$

This contradicts with (5.49), since $\mu_1 = 3\mu_0$ and $\mu_0 \geq 16$.

In the latter case, that is, $k_D(u_2, w_3) \leq 5\mu_1$, it follows from (5.43) that

$$k_D(w_3, u_1) \leq k_D(w_3, u_2) + k_D(u_2, u_1) \leq 6\mu_1. \quad (5.50)$$

As Property C(1) implies

$$k_D(w_1, w_2) \leq k_D(w_3, w_4),$$

we see from Lemma 2.7 and (5.45) that

$$k_D(w_2, w_3) \leq \ell_k(\gamma_1[w_1, w_2]) - k_D(w_1, w_3) \leq k_D(w_1, w_2) + \frac{1}{2} - 16\mu_1$$

$$\leq k_D(w_3, w_4) + \frac{1}{2} - 16\mu_1.$$

Then we get from (5.50) that

$$\begin{aligned} k_D(y_0, w_2) &\leq k_D(y_0, u_1) + k_D(u_1, w_3) + k_D(w_3, w_2) \\ &\leq k_D(w_3, w_4) + k_D(y_0, u_1) + \frac{1}{2} - 10\mu_1. \end{aligned} \quad (5.51)$$

Since $w_4 \in \beta_0$, there must exist $i \in \{1, \dots, m-1\}$ such that $w_4 \in \beta_i \in \Lambda_{\xi_i \xi_{i+1}}^\lambda$ (see Figure 8). Note that $k_D(y_0, \xi_i) = k_D(y_0, \xi_1) = k_D(y_0, w_2)$ by (5.31), and thus, we obtain from Lemma 5.10(ii) that

$$\begin{aligned} k_D(y_0, w_4) &\leq k_D(y_0, \xi_i) + \ell_k(\beta_i) \leq k_D(y_0, \xi_i) + \frac{7}{4} + 2\mu_0 + 2\mu_1 \\ &= k_D(y_0, w_2) + \frac{7}{4} + 2\mu_0 + 2\mu_1. \end{aligned} \quad (5.52)$$

Meanwhile, Lemma 2.6 and (5.43) show that

$$\begin{aligned} k_D(y_0, w_4) &\geq \ell_k(\zeta_4) - \frac{1}{4} \geq k_D(w_4, u_3) + k_D(y_0, u_3) - \frac{1}{4} \\ &\geq k_D(w_4, u_2) - k_D(u_2, u_3) + k_D(y_0, u_3) - \frac{1}{4} \\ &\geq k_D(w_4, w_3) - k_D(w_3, u_2) + k_D(y_0, u_3) - \frac{1}{4} - \mu_1, \end{aligned}$$

which, together with the assumption $k_D(u_2, w_3) \leq 5\mu_1$, yields

$$k_D(y_0, w_4) \geq k_D(w_3, w_4) + k_D(y_0, u_3) - \frac{1}{4} - 6\mu_1.$$

Combining this with (5.52), we get

$$k_D(y_0, w_2) \geq k_D(w_3, w_4) + k_D(y_0, u_3) - 2 - 2\mu_0 - 8\mu_1,$$

and thus, (5.44) gives

$$k_D(y_0, w_2) \geq k_D(w_3, w_4) + k_D(y_0, u_1) - 2 - 2\mu_0 - 9\mu_1,$$

This contradicts with (5.51), and hence the proof is complete. \square

Lemma 5.16. *Suppose that $[w_1, w_2, \gamma_{12}, \varpi, w_3, w_4]$ is a six-tuple satisfying Property C with constant \mathfrak{c} (see Figure 9). For each integer $r \in \{1, 2, 3\}$, let $\gamma_{rs} \in \Lambda_{w_r w_s}^\lambda$ with $s = r + 1$, where $\gamma_{13} = \gamma_{12}[w_1, w_3]$ and $\gamma_{23} = \gamma_{12}[w_2, w_3]$. Then*

- (1) (a) *there exist $x_1 \in \gamma_{23}$, $x_2 \in \gamma_{24}$ and $x_3 \in \gamma_{34}$ such that for any $r \neq s \in \{1, 2, 3\}$,*

$$k_D(x_r, x_s) \leq \mu_1;$$

- (b) $k_D(w_3, x_3) \leq \frac{1}{2} + \mu_1$.

- (2) (a) *there exist $p_1 \in \gamma_{13}$, $p_2 \in \gamma_{14}$ and $p_3 \in \gamma_{34}$ such that for any $r \neq s \in \{1, 2, 3\}$,*

$$k_D(p_r, p_s) \leq \mu_1;$$

- (b) $k_D(w_3, p_3) \leq \frac{1}{2} + \mu_1$.

- (3) $k_D(w_2, w_4) \geq k_D(w_2, w_3) - \frac{5}{4} - 5\mu_1 + \mathfrak{c}$.

- (4) *for any $w \in \gamma_{34}$, it holds*

$$k_D(y_0, w) \leq k_D(y_0, \xi_1) - \min\{k_D(w_2, w_3) + k_D(w_3, w), k_D(w_4, w)\} + \frac{15}{2} + 13\mu_1.$$




FIGURE 9. Illustration for the proof of Lemma 5.16 I

Proof. (1) By Proposition 3.4(P_4), (D, k_D) is a (μ_0, ρ_0) -Rips space, and by Lemma 2.7, γ_{23} is a 2λ -curve. Then Lemma 2.24 guarantees the existence of the points $x_1 \in \gamma_{23}$, $x_2 \in \gamma_{24}$ and $x_3 \in \gamma_{34}$ satisfying statement (1a).

For (1b), we may estimate as follows:

$$\begin{aligned}
k_D(w_3, x_3) + k_D(x_3, w_4) &\leq \ell_k(\gamma_{34}) \leq k_D(w_3, w_4) + \frac{1}{4} && \text{(by Lemma 2.6)} \\
&\leq k_D(w_4, x_1) + \frac{1}{2} && \text{(by Property C)} \\
&\leq k_D(w_4, x_3) + k_D(x_3, x_1) + \frac{1}{2} \\
&\leq k_D(w_4, x_3) + \frac{1}{2} + \mu_1 && \text{(by statement (1a))}
\end{aligned}$$

(2) The proof of (2) is similar to that of (1) and thus is omitted.

(3) Statement (1) ensures that

$$k_D(w_2, x_2) \geq k_D(w_2, w_3) - k_D(w_3, x_3) - k_D(x_3, x_1) - k_D(x_1, x_2) \geq k_D(w_2, w_3) - \frac{1}{2} - 3\mu_1$$

and that

$$k_D(w_4, x_2) \geq k_D(w_4, w_3) - k_D(w_3, x_3) - k_D(x_3, x_2) \geq k_D(w_3, w_4) - \frac{1}{2} - 2\mu_1.$$

These two estimates, together with Lemma 2.6 and Property C(1), imply that

$$\begin{aligned} k_D(w_2, w_4) &\geq \ell_k(\gamma_{24}) - \frac{1}{4} \geq k_D(w_2, x_2) + k_D(x_2, w_4) - \frac{1}{4} \\ &\geq k_D(w_2, w_3) - \frac{5}{4} - 5\mu_1 + \mathbf{c}. \end{aligned}$$

This proves the statement (3).

(4) To prove the statement (4), we shall construct several auxiliary points. By the statement (3), we see that $k_D(w_2, w_4) > 58\mu_1$. It follows from Lemma 5.11 that there exist two integers $r_1 < r_2 \in \{1, \dots, m\}$ such that $w_2 \in \beta_{r_1}$ and $w_4 \in \beta_{r_2}$ (see Figure 9), where β_i are defined in Section 5.4. Let $\varsigma_1 \in \Lambda_{w_2 y_0}^\lambda$ and $\varsigma_2 \in \Lambda_{w_4 y_0}^\lambda$.

Applying Lemma 5.12 (with u and v replaced by w_2 and w_4), we infer that there are $\eta_{r_1 r_2, 1} \in \varsigma_1$, $\eta_{r_1 r_2, 2} \in \varsigma_2$ and $\eta_{r_1 r_2, 3} \in \gamma_{24}$ (see Figure 9) such that for any $r \neq s \in \{1, 2, 3\}$,

$$k_D(\eta_{r_1 r_2, r}, \eta_{r_1 r_2, s}) \leq \mu_1. \quad (5.53)$$

Let $w \in \gamma_{34}$. To establish the desired upper bound for $k_D(y_0, w)$, we consider two cases.

Case 1: $\eta_{r_1 r_2, 3} \in \gamma_{24}[w_2, x_2]$.

Under this assumption, we separate the arguments into two subcases: $w \in \gamma_{34}[w_3, x_3]$ and $w \in \gamma_{34}[x_3, w_4]$. In the former case (see the red w in Figure 9), we know from the statement (1b) and Lemma 2.7 that

$$\max\{k_D(w_3, w), k_D(w, x_3)\} \leq \ell_k(\gamma_{34}[w_3, x_3]) \leq \frac{1}{2} + k_D(w_3, x_3) \leq 1 + \mu_1, \quad (5.54)$$

and thus, we get from the statement (1a) that

$$k_D(w, x_2) \leq k_D(w, x_3) + k_D(x_3, x_2) \leq 1 + 2\mu_1. \quad (5.55)$$

Since (D, k_D) is a (μ_0, ρ_0) -Rips space, it follows from (5.53) and Lemma 2.25 (replacing x , y and z by w_4 , $\eta_{r_1 r_2, 2}$ and $\eta_{r_1 r_2, 3}$) that there exists $x_4 \in \varsigma_2[w_4, \eta_{r_1 r_2, 2}]$ (see Figure 9) such that

$$k_D(x_2, x_4) \leq 1 + \mu_0 + \mu_1.$$

This, together with (5.55), implies

$$k_D(w, x_4) \leq k_D(w, x_2) + k_D(x_2, x_4) \leq 2 + \mu_0 + 3\mu_1.$$

Then

$$k_D(w_4, w) \leq k_D(w_4, x_4) + k_D(x_4, w) \leq k_D(w_4, x_4) + 2 + \mu_0 + 3\mu_1,$$

and thus, we obtain from the above two estimates and Lemma 2.6 that

$$\begin{aligned} k_D(y_0, w) &\leq k_D(y_0, x_4) + k_D(x_4, w) \leq \ell_k(\varsigma_2) - \ell_k(\varsigma_2[w_4, x_4]) + k_D(x_4, w) \\ &\leq k_D(y_0, w_4) - k_D(w_4, x_4) + k_D(x_4, w) + \frac{1}{4} \\ &\leq k_D(y_0, w_4) - k_D(w_4, w) + \frac{17}{4} + 2\mu_0 + 6\mu_1 \\ &\stackrel{\text{Lemma 5.10(iii)}}{\leq} k_D(y_0, \xi_1) - k_D(w_4, w) + 6 + \mu_0 + 9\mu_1. \end{aligned}$$

This proves (4).

In the latter case, that is, $w \in \gamma_{34}[w_4, x_3]$ (see the blue w in Figure 9), since (D, k_D) is a (μ_0, ρ_0) -Rips space, it follows from statement (1a) and Lemma 2.25 (replacing x, y and z by w_4, x_2 and x_3) that there exists $x_5 \in \gamma_{24}[w_4, x_2]$ such that

$$k_D(w, x_5) \leq 1 + \mu_0 + \mu_1. \quad (5.56)$$

Let us consider $w_4, \eta_{r_1 r_2, 2}$ and $\eta_{r_1 r_2, 3}$. As $\eta_{r_1 r_2, 3} \in \gamma_{24}[w_2, x_2]$, it follows from (5.53) and Lemma 2.25 that there exists $x_6 \in \varsigma_2[w_4, \eta_{r_1 r_2, 2}]$ such that

$$k_D(x_5, x_6) \leq 1 + \mu_0 + \mu_1.$$

This, together with the choice of x_5 in (5.56), implies

$$k_D(w, x_6) \leq k_D(w, x_5) + k_D(x_5, x_6) \leq 2(1 + \mu_0 + \mu_1).$$

Combining the above estimate with Lemma 2.6 gives

$$\begin{aligned} k_D(y_0, w) &\leq k_D(y_0, x_6) + k_D(x_6, w) \leq \ell_k(\varsigma_2) - \ell_k(\varsigma_2[w_4, x_6]) + k_D(x_6, w) \\ &\leq k_D(y_0, w_4) - k_D(w_4, x_6) + k_D(x_6, w) + \frac{1}{4} \\ &\leq k_D(y_0, w_4) - k_D(w_4, w) + 2k_D(x_6, w) + \frac{1}{4} \\ &\leq k_D(y_0, w_4) - k_D(w_4, w) + \frac{17}{4} + \mu_0 + 5\mu_1 \\ &\stackrel{\text{Lemma 5.10(iii)}}{\leq} k_D(y_0, \xi_1) - k_D(w_4, w) + 6 + 8\mu_1, \end{aligned}$$

which again proves (4).

Case 2: $\eta_{r_1 r_2, 3} \in \gamma_{24}[w_4, x_2]$.

Under this assumption, we again separate the arguments into two subcases: $w \in \gamma_{34}[w_3, x_3]$ and $w \in \gamma_{34}[w_4, x_3]$. For the former (see the red w in Figure 10), it follows from (5.53) and Lemma 2.25 (replacing x, y and z by $w_2, \eta_{r_1 r_2, 1}$ and $\eta_{r_1 r_2, 3}$) that there exists $x_7 \in \varsigma_1[w_2, \eta_{r_1 r_2, 1}]$ (see Figure 10) such that

$$k_D(x_2, x_7) \leq 1 + \mu_0 + \mu_1.$$

This, together with (5.55), gives

$$k_D(w, x_7) \leq k_D(w, x_2) + k_D(x_2, x_7) \leq 2 + 2\mu_1.$$

By Lemma 2.6, statement (1a) and the above two estimates, we have

$$\begin{aligned} k_D(y_0, w) &\leq k_D(y_0, x_7) + k_D(x_7, w) \leq \ell_k(\varsigma_1) - \ell_k(\varsigma_1[w_2, x_7]) + k_D(x_7, w) \\ &\leq k_D(y_0, w_2) - k_D(w_2, x_7) + k_D(x_7, w) + \frac{1}{4} \\ &\leq k_D(y_0, w_2) - k_D(w_2, x_1) + k_D(x_1, x_2) + k_D(x_2, x_7) + k_D(x_7, w) + \frac{1}{4} \quad (5.57) \\ &\leq k_D(y_0, w_2) - k_D(w_2, x_1) + \frac{13}{4} + \mu_0 + 4\mu_1 \\ &\stackrel{\text{Lemma 5.10(iii)}}{\leq} k_D(y_0, \xi_1) - k_D(w_2, x_1) + 5 + 7\mu_1. \end{aligned}$$

It remains to estimate $k_D(w_2, x_1)$. Note that statements (1a) and (1b) give

$$k_D(x_1, w_3) \leq k_D(x_1, x_3) + k_D(x_3, w_3) \leq \frac{1}{2} + 2\mu_1. \quad (5.58)$$




FIGURE 10. Illustration for the proof of Lemma 5.16 II

It follows from this and (5.54) that

$$k_D(x_1, w_3) + k_D(w_3, w) \leq \frac{3}{2} + 3\mu_1,$$

and thus, we have

$$\begin{aligned} k_D(w_2, x_1) &\geq k_D(w_2, x_1) + k_D(x_1, w_3) + k_D(w_3, w) - \frac{3}{2} - 3\mu_1 \\ &\geq k_D(w_2, w_3) + k_D(w_3, w) - \frac{3}{2} - 3\mu_1. \end{aligned}$$

Substituting this to (5.57), we infer that

$$k_D(y_0, w) \leq k_D(y_0, \xi_1) - (k_D(w_2, w_3) + k_D(w_3, w)) + \frac{13}{2} + 10\mu_1.$$

This proves (4).

In the latter case, that is, $w \in \gamma_{34}[w_4, x_3]$ (see the blue w in Figure 10), it follows from statement (1a) and Lemma 2.25 (replacing x , y and z by w_4 , x_2 and x_3) that there exist $x_8 \in \gamma_{24}[w_4, x_2]$ such that

$$k_D(w, x_8) \leq 1 + \mu_0 + \mu_1. \quad (5.59)$$

If $x_8 \in \gamma_{24}[x_2, \eta_{r_1 r_2, 3}]$ (see the blue x_8 in Figure 10), then (5.53) and Lemma 2.25 imply that there exists $x_9 \in \gamma_{11}[w_2, \eta_{r_1 r_2, 1}]$ such that

$$k_D(x_8, x_9) \leq 1 + \mu_0 + \mu_1,$$

and by (5.59), it holds

$$k_D(w, x_9) \leq k_D(w, x_8) + k_D(x_8, x_9) \leq 2(1 + \mu_0 + \mu_1).$$

Then we obtain from Lemma 2.6 and the above two estimates that

$$\begin{aligned} k_D(y_0, w) &\leq k_D(y_0, x_9) + k_D(x_9, w) \leq \ell_k(\varsigma_1) - \ell_k(\varsigma_1[w_2, x_9]) + k_D(w, x_9) \\ &\leq k_D(y_0, w_2) - k_D(w_2, x_9) + k_D(w, x_9) + \frac{1}{4} \\ &\leq k_D(y_0, w_2) - k_D(w_2, x_8) + k_D(x_8, x_9) + k_D(w, x_9) + \frac{1}{4} \\ &\stackrel{\text{Lemma 5.10(iii)}}{\leq} k_D(y_0, \xi_1) - k_D(w_2, x_8) + 5 + 2\mu_0 + 6\mu_1. \end{aligned} \tag{5.60}$$

Next, we derive a lower bound for $k_D(w_2, x_8)$. Note that by statement (1a) and (5.59), we have

$$k_D(x_2, x_8) \geq k_D(x_3, w) - k_D(x_2, x_3) - k_D(w, x_8) \geq k_D(x_3, w) - (1 + \mu_0 + 2\mu_1),$$

and that by (5.58) and statement (1b), it holds

$$k_D(x_1, w_3) + k_D(w_3, x_3) \leq 1 + 3\mu_1.$$

Combining the above two estimates with Lemma 2.7 gives

$$\begin{aligned} k_D(w_2, x_8) &\geq \ell_k(\gamma_{24}[w_2, x_8]) - \frac{1}{2} \geq k_D(w_2, x_2) + k_D(x_2, x_8) - \frac{1}{2} \\ &\geq k_D(w_2, x_1) - k_D(x_1, x_2) + k_D(x_2, x_8) - \frac{1}{2} \\ &\geq k_D(w_2, x_1) + k_D(x_3, w) - \left(\frac{3}{2} + \mu_0 + 3\mu_1\right) \\ &\geq k_D(w_2, w_3) - k_D(w_3, x_1) + k_D(w, w_3) - k_D(w_3, x_3) - \left(\frac{3}{2} + \mu_0 + 3\mu_1\right) \\ &\geq k_D(w_2, w_3) + k_D(w_3, w) - \left(\frac{5}{2} + \mu_0 + 6\mu_1\right), \end{aligned}$$

Substituting the above estimate to (5.60) yields

$$k_D(y_0, w) \leq k_D(y_0, \xi_1) - (k_D(w_2, w_3) + k_D(w_3, w)) + \frac{15}{2} + 13\mu_1,$$

which proves (4).

If $x_8 \in \gamma_{24}[\eta_{r_1 r_2, 3}, w_4]$ (see the red x_8 in Figure 10), then Lemma 2.25 and (5.53) show that there exists $x_{10} \in \varsigma_2[w_4, \eta_{r_1 r_2, 2}]$ (see Figure 10) such that

$$k_D(x_8, x_{10}) \leq 1 + \mu_0 + \mu_1.$$

Since (5.59) ensures

$$k_D(w, x_{10}) \leq k_D(w, x_8) + k_D(x_8, x_{10}) \leq 2(1 + \mu_0 + \mu_1),$$

we derive from Lemma 2.6 and the above estimate that

$$\begin{aligned}
 k_D(y_0, w) &\leq k_D(y_0, x_{10}) + k_D(x_{10}, w) \leq \ell_k(\varsigma_2) - \ell_k(\varsigma_2[w_4, x_{10}]) + k_D(x_{10}, w) \\
 &\leq k_D(y_0, w_4) - k_D(w_4, x_{10}) + k_D(x_{10}, w) + \frac{1}{4} \\
 &\leq k_D(y_0, w_4) - k_D(w_4, w) + 2k_D(x_{10}, w) + \frac{1}{4} \\
 &\leq k_D(y_0, w_4) - k_D(w_4, w) + \frac{17}{4} + \mu_0 + 5\mu_1 \\
 &\stackrel{\text{Lemma 5.10(iii)}}{\leq} k_D(y_0, \xi_1) - k_D(w_4, w) + 6 + 8\mu_1,
 \end{aligned}$$

which again proves (4). The proof of lemma is thus complete. \square

Let $w_{1,1} \in D$, $\varpi \in \beta_0$, $w_{1,2} \in \beta_0[\xi_1, \varpi]$, $\gamma_{12}^1 \in \Lambda_{w_{1,1}w_{1,2}}^\lambda$, $w_{1,3} \in \gamma_{12}^1$ and $w_{1,4} \in \beta_0[w_{1,2}, \varpi]$. Then $[w_{1,1}, w_{1,2}, \gamma_{12}^1, \varpi, w_{1,3}, w_{1,4}]$ is a six-tuple. Then we let $w_{2,1} = w_{1,3}$, $w_{2,2} = w_{1,4}$, $\gamma_{12}^2 \in \Lambda_{w_{2,1}w_{2,2}}^\lambda$, $w_{2,3} \in \gamma_{12}^2$ and $w_{2,4} \in \beta_0[w_{2,2}, \varpi]$ (see Figure 11). Then $[w_{2,1}, w_{2,2}, \gamma_{12}^2, \varpi, w_{2,3}, w_{2,4}]$ is a six-tuple as well. For $i \in \{1, 2\}$, we set

$$\Omega_i = [w_{i,1}, w_{i,2}, \gamma_{12}^i, \varpi, w_{i,3}, w_{i,4}].$$




FIGURE 11. Illustration for the proof of Lemma 5.17

Lemma 5.17. *If both Ω_1 and Ω_2 satisfy Property C with constant \mathfrak{c} , then*

$$k_D(w_{1,3}, w_{2,3}) = k_D(w_{2,1}, w_{2,3}) \leq \frac{1}{2}(\mathfrak{c} - k_D(w_{1,2}, w_{1,3})) + 28\mu_1.$$

Proof. Suppose on the contrary that

$$k_D(w_{1,3}, w_{2,3}) = k_D(w_{2,1}, w_{2,3}) > \frac{1}{2}(\mathfrak{c} - k_D(w_{1,2}, w_{1,3})) + 28\mu_1. \quad (5.61)$$

Under this contrary assumption, we do some preparation. First of all, since for each $i \in \{1, 2\}$, the six-tuple $\Omega_i = [w_{i,1}, w_{i,2}, \gamma_{12}^i, \varpi, w_{i,3}, w_{i,4}]$ satisfies Property **C** with constant \mathfrak{c} , we know from Property **C(1)** that

$$k_D(w_{i,1}, w_{i,2}) \leq \mathfrak{c} \text{ and } k_D(w_{i,3}, w_{i,4}) = \mathfrak{c}. \quad (5.62)$$

Next, we shall construct several auxiliary points. By (5.61), we see that there is $u_1 \in \gamma_{12}^2[w_{1,3}, w_{2,3}]$ (see Figure 11) such that

$$k_D(w_{1,3}, u_1) = k_D(w_{2,1}, u_1) = \frac{1}{2}(\mathfrak{c} - k_D(w_{1,2}, w_{1,3})) + \frac{1}{2}, \quad (5.63)$$

and so (5.61) gives

$$k_D(w_{2,3}, u_1) \geq k_D(w_{2,3}, w_{1,3}) - k_D(w_{1,3}, u_1) \geq 28\mu_1 - \frac{1}{2}. \quad (5.64)$$

By Lemma 2.6 and (5.62), we have

$$\begin{aligned} k_D(w_{1,2}, w_{1,3}) &\leq \ell_k(\gamma_{12}^1[w_{1,2}, w_{1,3}]) \leq \ell_k(\gamma_{12}^1) - \ell_k(\gamma_{12}^1[w_{1,1}, w_{1,3}]) \\ &\leq k_D(w_{1,1}, w_{1,2}) - k_D(w_{1,1}, w_{1,3}) + \frac{1}{4} \\ &\leq \mathfrak{c} - k_D(w_{1,1}, w_{1,3}) + \frac{1}{4}, \end{aligned}$$

and thus

$$\frac{1}{2}(\mathfrak{c} - k_D(w_{1,2}, w_{1,3})) + \frac{1}{2} \geq \frac{1}{2}k_D(w_{1,1}, w_{1,3}) + \frac{3}{8} \geq \frac{3}{8}. \quad (5.65)$$

By (5.65) and the choice of u_1 in (5.63), we have

$$k_D(w_{2,1}, u_1) \geq \frac{3}{8}.$$

Then it follows from Lemma 2.6, Property **C(1)**, (5.62) and the above estimate that

$$\begin{aligned} k_D(w_{2,3}, u_1) &\leq \ell_k(\gamma_{12}^2[w_{2,3}, u_1]) \leq \ell_k(\gamma_{12}^2) - k_D(w_{2,1}, u_1) \\ &\leq k_D(w_{2,1}, w_{2,2}) - k_D(w_{2,1}, u_1) + \frac{1}{4} \\ &\leq \mathfrak{c} - \frac{1}{8} < k_D(w_{2,3}, w_{2,4}). \end{aligned}$$

Fix $\gamma_{34}^2 \in \Lambda_{w_{2,3}w_{2,4}}^\lambda$. Then the previous inequality ensures that there is $u_2 \in \gamma_{34}^2$ (see Figure 11) such that

$$k_D(w_{2,3}, u_2) = k_D(w_{2,3}, u_1). \quad (5.66)$$

Fix $\varsigma_{12} \in \Lambda_{u_1u_2}^\lambda$. Since (D, k_D) is a (μ_0, ρ_0) -Rips space by Proposition 3.4(P_4), we know from Lemma 2.24 that there exist $v_1 \in \gamma_{12}^2[w_{2,3}, u_1]$, $v_2 \in \gamma_{34}^2[w_{2,3}, u_2]$ and $v_3 \in \varsigma_{12}$ (see Figure 11) such that for any $r \neq s \in \{1, 2, 3\}$,

$$k_D(v_r, v_s) \leq \mu_1. \quad (5.67)$$

The following claim concerns the points v_1 and v_2 .

Claim 5.17-1: $k_D(w_{2,3}, v_1) \leq \frac{1}{2} + 2\mu_1$ and $k_D(w_{2,3}, v_2) \leq \frac{1}{2} + \mu_1$.

Indeed, it follows from Lemma 2.6 and Property C(1) that

$$\begin{aligned} \ell_k(\gamma_{34}^2[w_{2,4}, v_2]) + \ell_k(\gamma_{34}^2[w_{2,3}, v_2]) &\leq \ell_k(\gamma_{34}^2) \leq \mathbf{c} + \frac{1}{4} \\ &\stackrel{\text{Property C(2)}}{\leq} k_D(w_{2,4}, v_1) + \frac{1}{2} \leq k_D(w_{2,4}, v_2) + k_D(v_1, v_2) + \frac{1}{2} \\ &\stackrel{(5.67)}{\leq} \ell_k(\gamma_{34}^2[w_{2,4}, v_2]) + \frac{1}{2} + \mu_1. \end{aligned}$$

By the above estimate and (5.67), we get

$$k_D(w_{2,3}, v_1) \leq k_D(w_{2,3}, v_2) + k_D(v_2, v_1) \leq \ell_k(\gamma_{34}^2[w_{2,3}, v_2]) + k_D(v_2, v_1) \leq \frac{1}{2} + 2\mu_1.$$

This proves the claim.

It follows from Claim 5.17-1 and (5.64) that

$$k_D(u_1, v_1) \geq k_D(u_1, w_{2,3}) - k_D(w_{2,3}, v_1) \geq -1 + 26\mu_1, \quad (5.68)$$

and from Claim 5.17-1, (5.64) and the choice of u_2 in (5.66) that

$$k_D(u_2, v_2) \geq k_D(u_2, w_{2,3}) - k_D(w_{2,3}, v_2) \geq -1 + 27\mu_1. \quad (5.69)$$

To continue the discussions, we need to find more auxiliary points. Fix $\rho_1 \in \Lambda_{u_1 y_0}^\lambda$, $\rho_2 \in \Lambda_{u_2 y_0}^\lambda$ and $\rho_3 \in \Lambda_{u_1 u_2}^\lambda$. By Lemma 2.24, we know that there are $p_1 \in \rho_1$, $p_2 \in \rho_2$ and $p_3 \in \rho_3$ (see Figure 11) such that for any $r \neq s \in \{1, 2, 3\}$,

$$k_D(p_r, p_s) \leq \mu_1. \quad (5.70)$$

To reach a contradiction, we consider two cases, depending on the position of p_3 .

Case 1: $p_3 \in \rho_3[u_2, v_3]$ (see the blue p_3 in Figure 11).

Under this assumption, we know from (5.70) and Lemma 2.25 (replacing x, y and z by u_1, p_1 and p_3) that there exists $q_1 \in \rho_1[u_1, p_1]$ (see Figure 11) such that

$$k_D(v_3, q_1) \leq 1 + \mu_0 + \mu_1. \quad (5.71)$$

Then it follows from (5.67) that

$$k_D(q_1, v_1) \leq k_D(q_1, v_3) + k_D(v_3, v_1) \leq 1 + \mu_0 + 2\mu_1.$$

Next, we derive a lower bound for $k_D(y_0, q_1)$. By the previous inequality, we have

$$k_D(y_0, q_1) \geq k_D(y_0, v_1) - k_D(v_1, q_1) \geq k_D(y_0, v_1) - (1 + \mu_0 + 2\mu_1). \quad (5.72)$$

Moreover, Lemma 2.6 and (5.62) imply

$$\begin{aligned} k_D(w_{1,4}, v_1) &\leq \ell_k(\gamma_{12}^2[w_{1,4}, v_1]) \leq \ell_k(\gamma_{12}^2) - \ell_k(\gamma_{12}^2[w_{1,3}, v_1]) \\ &\leq \mathbf{c} - (k_D(w_{1,3}, u_1) + k_D(u_1, v_1)) + \frac{1}{4} \\ &\stackrel{(5.63)+(5.68)}{\leq} \frac{1}{2}(\mathbf{c} + k_D(w_{1,2}, w_{1,3})) + \frac{3}{4} - 26\mu_1. \end{aligned}$$

Then it follows from (5.72) and the above estimate that

$$\begin{aligned} k_D(y_0, q_1) &\geq k_D(y_0, w_{1,4}) - k_D(w_{1,4}, v_1) - (1 + \mu_0 + 2\mu_1) \\ &\geq k_D(y_0, w_{1,4}) - \frac{1}{2}(\mathbf{c} + k_D(w_{1,2}, w_{1,3})) - \frac{7}{4} - \mu_0 + 24\mu_1, \end{aligned}$$

and thus, we get from Lemma 5.10(iii) that

$$k_D(y_0, q_1) \geq k_D(y_0, \xi_1) - \frac{1}{2}(\mathbf{c} + k_D(w_{1,2}, w_{1,3})) - \frac{7}{2} + 21\mu_1. \quad (5.73)$$

On the other hand, By (5.62) and the choice of u_1 in (5.63), we have

$$\begin{aligned} \frac{1}{2}(\mathbf{c} + k_D(w_{1,2}, w_{1,3})) - \frac{1}{2} &= k_D(w_{1,4}, w_{1,3}) - k_D(w_{1,3}, u_1) \\ &\leq k_D(w_{1,4}, u_1) \leq \ell_k(\gamma_{12}^2) - \ell_k(\gamma_{12}^2[w_{1,3}, u_1]) \\ &\stackrel{\text{Lemma 2.6}}{\leq} k_D(w_{1,3}, w_{1,4}) - k_D(w_{1,3}, u_1) + \frac{1}{4} \\ &= \frac{1}{2}(\mathbf{c} + k_D(w_{1,2}, w_{1,3})) - \frac{1}{4} \\ &= k_D(w_{1,2}, w_{1,3}) + k_D(w_{1,3}, u_1) - \frac{3}{4}. \end{aligned}$$

This implies that

$$\min\{k_D(w_{1,2}, w_{1,3}) + k_D(w_{1,3}, u_1), k_D(w_{1,4}, u_1)\} \geq \frac{1}{2}(\mathbf{c} + k_D(w_{1,2}, w_{1,3})) - \frac{1}{2},$$

and thus, Lemma 5.16(4) gives

$$\begin{aligned} k_D(y_0, u_1) &\leq k_D(y_0, \xi_1) - \min\{k_D(w_{1,2}, w_{1,3}) + k_D(w_{1,3}, u_1), k_D(w_{1,4}, u_1)\} + \frac{15}{2} + 13\mu_1 \\ &\leq k_D(y_0, \xi_1) - \frac{1}{2}(\mathbf{c} + k_D(w_{1,2}, w_{1,3})) + 8 + 13\mu_1. \end{aligned}$$

This, combining with Lemma 2.6, leads to

$$\begin{aligned} k_D(y_0, q_1) &\leq \ell_k(\rho_1) \leq k_D(y_0, u_1) + \frac{1}{4} \\ &\leq k_D(y_0, \xi_1) - \frac{1}{2}(\mathbf{c} + k_D(w_{1,2}, w_{1,3})) + \frac{33}{4} + 13\mu_1. \end{aligned}$$

Since $\mu_1 \geq 48$, this contradicts with (5.73). Thus the proof in this case is complete.

Case 2: $p_3 \in \rho_3[u_1, v_3]$ (see the red p_3 in Figure 11).

Under the assumption, considering the points u_2 , p_2 and p_3 , we know from (5.70) and Lemma 2.25 that there exists $q_2 \in \rho_2[u_2, p_2]$ (see Figure 11) such that

$$k_D(q_2, v_3) \leq 1 + \mu_0 + \mu_1. \quad (5.74)$$

Then it follows from (5.67) that

$$k_D(q_2, v_2) \leq k_D(q_2, v_3) + k_D(v_3, v_2) \leq 1 + \mu_0 + 2\mu_1,$$

and thus, we get

$$k_D(y_0, q_2) \geq k_D(y_0, v_2) - k_D(v_2, q_2) \geq k_D(y_0, v_2) - (1 + \mu_0 + 2\mu_1). \quad (5.75)$$

Next, we derive a lower bound for $k_D(y_0, q_2)$. Since Lemma 2.6, Claim 5.17-1 and (5.62) give

$$\begin{aligned}
k_D(w_{1,4}, v_2) &\leq k_D(w_{1,4}, w_{2,3}) + k_D(w_{2,3}, v_2) \\
&\leq \ell(\gamma_{12}^2) - \ell(\gamma_{12}^2[w_{1,3}, v_1]) + k_D(w_{2,3}, v_2) \\
&\leq k_D(w_{1,3}, w_{1,4}) - (k_D(w_{1,3}, u_1) + k_D(u_1, v_1)) + k_D(w_{2,3}, v_2) + \frac{1}{4} \\
&\leq \mathbf{c} - (k_D(w_{1,3}, u_1) + k_D(u_1, v_1)) + \frac{3}{4} + \mu_1 \\
&\stackrel{(5.63)+(5.68)}{\leq} \frac{1}{2}(\mathbf{c} + k_D(w_{1,2}, w_{1,3})) + \frac{5}{4} - 25\mu_1.
\end{aligned}$$

This, together with (5.75), implies that

$$\begin{aligned}
k_D(y_0, q_2) &\geq k_D(y_0, w_{1,4}) - k_D(w_{1,4}, v_2) - 1 - \mu_0 - 2\mu_1 \\
&\geq k_D(y_0, w_{1,4}) - \frac{1}{2}(\mathbf{c} + k_D(w_{1,2}, w_{1,3})) - \frac{9}{4} - \mu_0 + 23\mu_1,
\end{aligned}$$

and thus, Lemma 5.10(iii) gives

$$k_D(y_0, q_2) \geq k_D(y_0, \xi_1) - \frac{1}{2}(\mathbf{c} + k_D(w_{1,2}, w_{1,3})) - 4 + 20\mu_1. \quad (5.76)$$

On the other hand, since (5.62) yields

$$k_D(w_{2,4}, u_2) \geq k_D(w_{2,4}, w_{2,3}) - k_D(w_{2,3}, u_2) = \mathbf{c} - k_D(w_{2,3}, u_2)$$

and since the choice of u_2 in (5.66) leads to

$$\begin{aligned}
k_D(w_{1,4}, w_{2,3}) + k_D(w_{2,3}, u_2) &\geq k_D(w_{1,4}, w_{1,3}) - (k_D(w_{1,3}, u_1) + k_D(u_1, w_{2,3})) + k_D(w_{2,3}, u_2) \\
&= \mathbf{c} - k_D(w_{1,3}, u_1),
\end{aligned}$$

we have

$$\min\{k_D(u_2, w_{2,4}), k_D(w_{1,4}, w_{2,3}) + k_D(w_{2,3}, u_2)\} \geq \mathbf{c} - \max\{k_D(u_1, w_{1,3}), k_D(u_2, w_{2,3})\}.$$

This, together with Lemma 5.16(4) and the fact $w_{2,2} = w_{1,4}$, implies that

$$\begin{aligned}
k_D(y_0, u_2) &\leq k_D(y_0, \xi_1) - \min\{k_D(u_2, w_{2,4}), k_D(w_{1,4}, w_{2,3}) + k_D(u_2, w_{2,3})\} + \frac{15}{2} + 13\mu_1 \\
&\leq k_D(y_0, \xi_1) + \max\{k_D(u_1, w_{1,3}), k_D(u_2, w_{2,3})\} - \mathbf{c} + \frac{15}{2} + 13\mu_1.
\end{aligned} \quad (5.77)$$

Moreover, we have by (5.67) and (5.74) and Lemma 2.6 that

$$\begin{aligned}
k_D(y_0, q_2) &\leq \ell_k(\rho_2) - \ell_k(\rho_2[u_2, q_2]) \leq k_D(y_0, u_2) - k_D(u_2, q_2) + \frac{1}{4} \\
&\leq k_D(y_0, u_2) - (k_D(u_2, v_2) - k_D(v_2, v_3) - k_D(v_3, q_2)) + \frac{1}{4} \\
&\leq k_D(y_0, u_2) - k_D(u_2, v_2) + \frac{5}{4} + \mu_0 + 2\mu_1.
\end{aligned}$$

It follows from Claim 5.17-1 and (5.77) that

$$\begin{aligned}
k_D(y_0, q_2) &\leq k_D(y_0, u_2) - (k_D(u_2, w_{2,3}) - k_D(w_{2,3}, v_2)) + \frac{5}{4} + \mu_0 + 2\mu_1 \\
&\leq k_D(y_0, u_2) - k_D(w_{2,3}, u_2) + \frac{7}{4} + \mu_0 + 3\mu_1 \\
&\leq k_D(y_0, \xi_1) + \max\{k_D(u_1, w_{1,3}), k_D(u_2, w_{2,3})\} - k_D(w_{2,3}, u_2) \\
&\quad + \frac{37}{4} + \mu_0 + 16\mu_1 - \mathfrak{c}.
\end{aligned}$$

If $\max\{k_D(u_1, w_{1,3}), k_D(u_2, w_{2,3})\} = k_D(u_2, w_{2,3})$, then the above estimate gives

$$\begin{aligned}
k_D(y_0, q_2) &\leq k_D(y_0, \xi_1) + \frac{37}{4} + \mu_0 + 16\mu_1 - \mathfrak{c} \\
&\leq k_D(y_0, \xi_1) - \frac{1}{2}(\mathfrak{c} + k_D(w_{1,2}, w_{1,3})) + \frac{75}{8} + \mu_0 + 16\mu_1,
\end{aligned}$$

where in the last inequality, we used the following estimate

$$\mathfrak{c} + k_D(w_{1,2}, w_{1,3}) - \frac{1}{4} \leq \mathfrak{c} + \ell_k(\gamma_{12}^1) - \frac{1}{4} \stackrel{\text{Lemma 2.6}}{\leq} \mathfrak{c} + k_D(w_{1,1}, w_{1,2}) \stackrel{\text{Property C(1)}}{=} 2\mathfrak{c}.$$

If $\max\{k_D(u_1, w_{1,3}), k_D(u_2, w_{2,3})\} = k_D(u_1, w_{1,3})$, then by (5.63), we have

$$\begin{aligned}
k_D(y_0, q_2) &\leq k_D(y_0, \xi_1) + k_D(w_{1,3}, u_1) - k_D(w_{2,3}, u_2) + \frac{37}{4} + \mu_0 + 16\mu_1 - \mathfrak{c} \\
&\leq k_D(y_0, \xi_1) + k_D(w_{1,3}, u_1) + \frac{37}{4} + \mu_0 + 16\mu_1 - \mathfrak{c} \\
&\leq k_D(y_0, \xi_1) - \frac{1}{2}(\mathfrak{c} + k_D(w_{1,2}, w_{1,3})) + \frac{39}{4} + \mu_0 + 16\mu_1.
\end{aligned}$$

In either case, we obtain

$$k_D(y_0, q_2) \leq k_D(y_0, \xi_1) - \frac{1}{2}(\mathfrak{c} + k_D(w_{1,2}, w_{1,3})) + \frac{39}{4} + \mu_0 + 16\mu_1. \quad (5.78)$$

Combining (5.76) and (5.78), we deduce that $4\mu_1 \leq \mu_0 + 14 \leq \mu_1 + 14$, which is impossible as $\mu_1 \geq 48$. Therefore, the proof is complete. \square

5.6. A sequence of six-tuples. In the rest of this paper, we use τ_1 , τ_2 and τ_3 to denote constants defined in Section 2.1.

let $\Omega_1 = [w_{1,1}, w_{1,2}, \gamma_{12}^1, \varpi, w_{1,3}, w_{1,4}]$ be a six-tuple, where $w_{1,1} \in D$, $w_{1,2} \in \beta_0$, $\gamma_{12}^1 \in \Lambda_{w_{1,1}w_{1,2}}^\lambda$, $\varpi \in \beta_0[w_{1,2}, \xi_m]$, $w_{1,3} \in \gamma_{12}^1$ and $w_{1,4} \in \beta_0[w_{1,2}, \varpi]$ (see Figure 12).

In the rest of this section, we assume that

$$k_D(w_{1,1}, w_{1,2}) = \tau_3 \quad \text{and} \quad k_D(w_{1,1}, w_{1,3}) \leq \tau_1. \quad (5.79)$$

Then

$$k_D(w_{1,2}, w_{1,3}) \geq k_D(w_{1,2}, w_{1,1}) - k_D(w_{1,1}, w_{1,3}) \geq -\tau_1 + \tau_3 \geq \frac{15}{4}\tau_2. \quad (5.80)$$

This implies that there is $w_{2,1} \in \gamma_{12}^1[w_{1,2}, w_{1,3}]$ (see Figure 12) such that

$$k_D(w_{1,3}, w_{2,1}) = \tau_2. \quad (5.81)$$

Based on Ω_1 and $w_{2,1}$, we obtain the second six-tuple $\Omega_2 = [w_{2,1}, w_{2,2}, \gamma_{12}^2, \varpi_2, w_{2,3}, w_{2,4}]$ by setting $w_{2,2} = w_{1,2}$, $\gamma_{12}^2 = \gamma_{12}^1[w_{2,1}, w_{2,2}]$, $\varpi_2 = w_{1,4} \in \beta_0[w_{2,2}, \xi_m]$, $w_{2,3} \in \gamma_{12}^2$ and $w_{2,4} \in \beta_0[w_{2,2}, \varpi_2]$ (see Figure 12).




FIGURE 12. Illustration for the proof of Lemma 5.18

Then we obtain the third six-tuple $\Omega_3 = [w_{3,1}, w_{3,2}, \gamma_{12}^3, \varpi_3, w_{3,3}, w_{3,4}]$ by setting $w_{3,1} = w_{2,3}$, $w_{3,2} = w_{2,4}$, $\gamma_{12}^3 \in \Lambda_{w_{2,3}w_{2,4}}^\lambda$, $\varpi_3 = w_{1,4} \in \beta_0[w_{3,2}, \xi_m]$, $w_{3,3} \in \gamma_{12}^3$ and $w_{3,4} \in \beta_0[w_{3,2}, \varpi_3]$ and the fourth six-tuple $\Omega_4 = [w_{4,1}, w_{4,2}, \gamma_{12}^4, \varpi_4, w_{4,3}, w_{4,4}]$ by setting $w_{4,1} = w_{3,3}$, $w_{4,2} = w_{3,4}$, $\gamma_{12}^4 \in \Lambda_{w_{3,3}w_{3,4}}^\lambda$, $\varpi_4 = w_{1,4} \in \beta_0[w_{4,2}, \xi_m]$, $w_{4,3} \in \gamma_{12}^4$ and $w_{4,4} \in \beta_0[w_{4,2}, \varpi_4]$ (see Figure 12).

By repeating this procedure, we find a finite sequence of six-tuples $\{\Omega_i\}_{i=1}^{m_1}$ with $\Omega_i = [w_{i,1}, w_{i,2}, \gamma_{12}^i, \varpi_i, w_{i,3}, w_{i,4}]$ obtained by setting for each $i \in \{3, \dots, m_1\}$, $w_{i,1} = w_{i-1,3}$, $w_{i,2} = w_{i-1,4}$, $\gamma_{12}^i \in \Lambda_{w_{i-1,3}w_{i-1,4}}^\lambda$, $\varpi_i = w_{1,4} \in \beta_0[w_{i,2}, \xi_m]$, $w_{i,3} \in \gamma_{12}^i$ and $w_{i,4} \in \beta_0[w_{i,2}, \varpi_i]$.

Then we have the following useful estimates.

Lemma 5.18. (1) Suppose that $m_1 \geq 2$ and Ω_1 satisfies Property C with constant τ_3 . Then

$$k_D(w_{1,4}, w_{2,3}) \geq k_D(w_{1,3}, w_{1,4}) + k_D(w_{1,3}, w_{2,3}) - \frac{5}{4} - 4\mu_1.$$

(2) Suppose $m_1 = 2$. If both Ω_1 and Ω_2 satisfy Property C with constant τ_3 , then

(2a) $k_D(w_{2,3}, w_{2,4}) \leq k_D(w_{1,4}, w_{2,3}) + \frac{7}{4} + 4\mu_1 - \tau_2$, and

(2b) for any $\gamma_{34}^2 \in \Lambda_{w_{2,3}w_{2,4}}^\lambda$ and any $w \in \gamma_{34}^2$, it holds $k_D(w_{1,4}, w) > \tau_3$.

(3) Suppose $m_1 \geq 3$ and for each $i \in \{1, \dots, m_1\}$, Ω_i satisfies Property C with constant τ_3 . If

$$k_D(w_{2,3}, w_{m_1,3}) \leq -36\mu_1 - \tau_1 + \tau_2,$$

then

(3a) $k_D(w_{2,3}, w_{m_1,4}) \leq k_D(w_{1,4}, w_{2,3}) + \frac{7}{4} - 32\mu_1 - \tau_1$, and

(3b) for any $\gamma_{34}^{m_1} \in \Lambda_{w_{m_1,3}w_{m_1,4}}^\lambda$ and $w \in \gamma_{34}^{m_1}$, it holds $k_D(w_{1,4}, w) > \tau_3$.

Proof. (1). Fix $\varsigma_{23} \in \Lambda_{w_{1,4}w_{2,3}}^\lambda$ and $\gamma_{34}^1 \in \Lambda_{w_{1,3}w_{1,4}}^\lambda$. Since (D, k_D) is a (μ_0, ρ_0) -Rips space by Proposition 3.4(P_4), we obtain from Lemma 2.24 that there exist $y_1 \in \gamma_{12}^1[w_{1,3}, w_{2,3}]$, $y_2 \in \varsigma_{23}$ and $y_3 \in \gamma_{34}^1$ (see Figure 12) such that for any $r \neq s \in \{1, 2, 3\}$,

$$k_D(y_r, y_s) \leq \mu_1. \quad (5.82)$$

Since Ω_1 satisfies Property C with constant τ_3 , it follows that

$$k_D(w_{1,3}, w_{1,4}) = \tau_3 \leq k_D(y_1, w_{1,4}) + \frac{1}{4},$$

and thus, we get from the above estimate, Lemma 2.6 and (5.82) that

$$\begin{aligned} k_D(w_{1,3}, y_3) &\leq \ell_k(\gamma_{34}^1) - \ell_k(\gamma_{34}^1[w_{1,4}, y_3]) \\ &\leq k_D(w_{1,3}, w_{1,4}) - k_D(w_{1,4}, y_3) + \frac{1}{4} \\ &\leq k_D(y_1, w_{1,4}) - k_D(w_{1,4}, y_3) + \frac{1}{2} \\ &\leq k_D(y_1, y_3) + \frac{1}{2} \leq \frac{1}{2} + \mu_1. \end{aligned}$$

This implies

$$k_D(w_{1,4}, y_3) \geq k_D(w_{1,4}, w_{1,3}) - k_D(w_{1,3}, y_3) \geq k_D(w_{1,3}, w_{1,4}) - \frac{1}{2} - \mu_1 \quad (5.83)$$

and

$$\begin{aligned} k_D(w_{2,3}, y_2) &\geq k_D(w_{2,3}, w_{1,3}) - k_D(w_{1,3}, y_3) - k_D(y_3, y_2) \\ &\geq k_D(w_{2,3}, w_{1,3}) - \frac{1}{2} - 2\mu_1. \end{aligned}$$

Moreover, (5.82) and (5.83) ensure that

$$k_D(w_{1,4}, y_2) \geq k_D(w_{1,4}, y_3) - k_D(y_3, y_2) \geq k_D(w_{1,3}, w_{1,4}) - \frac{1}{2} - 2\mu_1.$$

Therefore, it follows from Lemma 2.6 and the above two estimates that

$$\begin{aligned} k_D(w_{1,4}, w_{2,3}) &\geq \ell_k(\varsigma_{23}) - \frac{1}{4} \geq k_D(w_{2,3}, y_2) + k_D(y_2, w_{1,4}) - \frac{1}{4} \\ &\geq k_D(w_{1,3}, w_{1,4}) + k_D(w_{1,3}, w_{2,3}) - \frac{5}{4} - 4\mu_1. \end{aligned}$$

This proves (1).

(2). The proofs are similar to those used in (3) below and thus will be omitted here.

(3). Since the choice of $w_{2,1}$ in (5.81), the assumption $k_D(w_{2,3}, w_{m_1,3}) \leq -36\mu_1 - \tau_1 + \tau_2$ and Lemma 2.7 guarantee that

$$\begin{aligned} k_D(w_{1,3}, w_{2,3}) &\geq \ell_k(\gamma_{12}^1[w_{1,3}, w_{2,3}]) - \frac{1}{2} \geq \ell_k(\gamma_{12}^1[w_{1,3}, w_{2,1}]) - \frac{1}{2} \\ &\geq \tau_2 - \frac{1}{2} \geq k_D(w_{2,3}, w_{m_1,3}) + 36\mu_1 + \tau_1 - \frac{1}{2}, \end{aligned}$$

and since Property C(1) gives

$$k_D(w_{m_1,3}, w_{m_1,4}) = k_D(w_{1,3}, w_{1,4}) = \tau_3, \quad (5.84)$$

we obtain from the above two estimates that

$$k_D(w_{2,3}, w_{m_1,4}) \leq k_D(w_{m_1,3}, w_{m_1,4}) + k_D(w_{2,3}, w_{m_1,3})$$

$$\begin{aligned}
& \stackrel{(5.84)}{=} k_D(w_{1,3}, w_{1,4}) + k_D(w_{2,3}, w_{m_1,3}) \\
& \leq k_D(w_{1,3}, w_{1,4}) + k_D(w_{1,3}, w_{2,3}) + \frac{1}{2} - 36\mu_1 - \tau_1 \\
& \stackrel{\text{statement (1)}}{\leq} k_D(w_{1,4}, w_{2,3}) + \frac{7}{4} - 32\mu_1 - \tau_1.
\end{aligned}$$

This proves (3a).

Next, we start the proof of (3b) with the following estimate:

$$k_D(w_{1,4}, w_{m_1,4}) \geq -\frac{57}{4} - 25\mu_1 - \tau_1 + 2\tau_3. \quad (5.85)$$

To prove this estimate, we shall construct several auxiliary points, and then, establish some properties concerning these points.

Fix $\varsigma_{2m_1} \in \Lambda_{w_{2,3}w_{m_1,4}}^\lambda$ and $\varsigma_{1m_1} \in \Lambda_{w_{1,4}w_{m_1,4}}^\lambda$. Since (D, k_D) is a (μ_0, ρ_0) -Rips space, we know from Lemma 2.24 that there exist $\omega_1 \in \varsigma_{2m_1}$, $\omega_2 \in \varsigma_{1m_1}$ and $\omega_3 \in \varsigma_{23}$ (see Figure 12) such that for any $r \neq s \in \{1, 2, 3\}$, it holds

$$k_D(\omega_r, \omega_s) \leq \mu_1. \quad (5.86)$$

Then we make the following claim.

Claim 5.18-1: $k_D(w_{m_1,4}, \omega_1) \leq k_D(w_{1,4}, \omega_3) + 2 - 31\mu_1 - \tau_1$.

Indeed, we may estimate using Lemma 2.6 as follows.

$$\begin{aligned}
k_D(w_{1,4}, \omega_3) & \geq k_D(w_{1,4}, w_{2,3}) - k_D(w_{2,3}, \omega_1) - k_D(\omega_1, \omega_3) \\
& \stackrel{\text{statement (3a)}}{\geq} k_D(w_{2,3}, w_{m_1,4}) - k_D(w_{2,3}, \omega_1) - \frac{7}{4} + 31\mu_1 + \tau_1 \\
& \geq \ell_k(\varsigma_{2m_1}) - \ell_k(\varsigma_{2m_1}[w_{2,3}, \omega_1]) - 2 + 31\mu_1 + \tau_1 \\
& \geq k_D(w_{m_1,4}, \omega_1) - 2 + 31\mu_1 + \tau_1.
\end{aligned}$$

This proves the claim.

Next, we would like to find two new auxiliary points. Observe that

$$\begin{aligned}
k_D(w_{1,4}, y_2) & \geq k_D(w_{1,4}, y_3) - k_D(y_3, y_2) \stackrel{(5.82)}{\geq} k_D(w_{1,4}, y_3) - \mu_1 \\
& \stackrel{(5.83)}{\geq} k_D(w_{1,3}, w_{1,4}) - \frac{1}{2} - 2\mu_1 \stackrel{(5.84)}{=} -\frac{1}{2} - 2\mu_1 + \tau_3.
\end{aligned}$$

This implies that there exist $y_4 \in \varsigma_{23}[w_{1,4}, y_2]$ (see Figure 12) such that

$$k_D(w_{1,4}, y_4) = -\frac{1}{2} - 2\mu_1 - \frac{\tau_1}{2} + \tau_3. \quad (5.87)$$

Considering $w_{1,4}$, y_2 and y_3 , it follows from (5.82) and Lemma 2.25 that there exists $y_5 \in \gamma_{34}^1[y_3, w_{1,4}]$ (see Figure 12) such that

$$k_D(y_4, y_5) \leq 1 + \mu_0 + \mu_1. \quad (5.88)$$

Then the choice of y_4 in (5.87) and (5.88) imply

$$k_D(y_5, w_{1,4}) \leq k_D(w_{1,4}, y_4) + k_D(y_4, y_5) \leq \frac{1}{2} + \mu_0 - \mu_1 - \frac{\tau_1}{2} + \tau_3,$$

and thus

$$k_D(w_{1,3}, y_5) \geq k_D(w_{1,3}, w_{1,4}) - k_D(y_5, w_{1,4}) \stackrel{(5.84)}{\geq} -\frac{1}{2} - \mu_0 + \mu_1 + \frac{\tau_1}{2}.$$

These two inequalities, together with (5.80), ensure that

$$\min\{k_D(w_{1,2}, w_{1,3}) + k_D(w_{1,3}, y_5), k_D(y_5, w_{1,4})\} = k_D(y_5, w_{1,4}).$$

Then it follows from Lemma 5.16(4) and the above equation that

$$k_D(y_0, y_5) \leq k_D(y_0, \xi_1) - k_D(y_5, w_{1,4}) + \frac{15}{2} + 13\mu_1. \quad (5.89)$$

Based on Claim 5.16-1, we establish the following claim which concerns the position of ω_3 .

Claim 5.18-2: $\omega_3 \in \varsigma_{23}[w_{2,3}, y_2]$.

Suppose on the contrary that $\omega_3 \in \varsigma_{23}[w_{1,4}, y_2]$ (see the red ω_3 in Figure 12). To reach a contradiction, we first derive a lower bound for $k_D(y_0, \omega_1)$. It follows from (5.82), Lemmas 2.6 and 2.7 that

$$\begin{aligned} k_D(w_{1,4}, \omega_3) &\leq \ell_k(\varsigma_{23}[w_{1,4}, y_2]) \leq k_D(w_{1,4}, y_2) + \frac{1}{2} \\ &\leq k_D(w_{1,4}, y_3) + k_D(y_3, y_2) + \frac{1}{2} \leq \ell_k(\gamma_{34}^1) + k_D(y_3, y_2) + \frac{1}{2} \\ &\leq k_D(w_{1,3}, w_{1,4}) + \frac{3}{4} + \mu_1 \stackrel{(5.84)}{=} \frac{3}{4} + \mu_1 + \tau_3. \end{aligned} \quad (5.90)$$

Since $w_{m_1,4} \in \beta_0$, Lemma 5.10(iii) gives

$$\begin{aligned} k_D(y_0, \omega_1) &\geq k_D(y_0, w_{m_1,4}) - k_D(w_{m_1,4}, \omega_1) \\ &\geq k_D(y_0, \xi_1) - k_D(w_{m_1,4}, \omega_1) - \frac{7}{4} - 2\mu_0 - 2\mu_1 \\ &\stackrel{\text{Claim 5.18-1}}{\geq} k_D(y_0, \xi_1) - k_D(w_{1,4}, \omega_3) - \frac{15}{4} - 2\mu_0 + 29\mu_1 + \tau_1. \end{aligned} \quad (5.91)$$

Next, we find one more point ω_4 and establish lower bounds for $k_D(y_0, \omega_4)$. Considering $w_{1,4}$, y_2 and y_3 , it follows from (5.82) and Lemma 2.25 that there exists $\omega_4 \in \varsigma_{23}[y_3, w_{1,4}]$ (see Figure 12) such that

$$k_D(\omega_3, \omega_4) \leq 1 + \mu_0 + \mu_1, \quad (5.92)$$

which, together with (5.86), shows

$$k_D(\omega_1, \omega_4) \leq k_D(\omega_1, \omega_3) + k_D(\omega_3, \omega_4) \leq 1 + \mu_0 + 2\mu_1. \quad (5.93)$$

On the one hand, we have

$$\begin{aligned} k_D(y_0, \omega_4) &\geq k_D(y_0, \omega_1) - k_D(\omega_1, \omega_4) \\ &\stackrel{(5.91)+(5.93)}{\geq} k_D(y_0, \xi_1) - k_D(w_{1,4}, \omega_3) - \frac{19}{4} + 26\mu_1 + \tau_1, \end{aligned} \quad (5.94)$$

which, together with (5.90), leads to

$$k_D(y_0, \omega_4) \geq k_D(y_0, \xi_1) - \frac{11}{2} + 25\mu_1 + \tau_1 - \tau_3. \quad (5.95)$$

On the other hand, (5.92) gives

$$k_D(w_{1,4}, \omega_4) \geq k_D(w_{1,4}, \omega_3) - k_D(\omega_3, \omega_4) \geq k_D(w_{1,4}, \omega_3) - 1 - \mu_0 - \mu_1,$$

and thus we know from (5.80) and Lemma 5.16(4) that

$$\begin{aligned} k_D(y_0, \omega_4) &\leq k_D(y_0, \xi_1) - \min\{k_D(w_{1,2}, w_{1,3}) + k_D(w_{1,3}, \omega_4), k_D(\omega_4, w_{1,4})\} + \frac{15}{2} + 13\mu_1 \\ &\leq k_D(y_0, \xi_1) - \min\{-\tau_1 + \tau_3, k_D(w_{1,4}, \omega_3) - 1 - \mu_0 - \mu_1\} + \frac{15}{2} + 13\mu_1. \end{aligned}$$

This contradicts with either (5.94) or (5.95), as $\mu_1 = 3\mu_0 \geq 48$. Therefore, Claim 5.18-2 holds.

Based on Claim 5.18-2, we now finish the proof of (5.85). It follows from Claim 5.18-2 that $\omega_3 \in \varsigma_{23}[w_{2,3}, y_2]$ (see the black ω_3 in Figure 12). Considering the points $w_{1,4}$, ω_2 and ω_3 , we know from (5.86) and Lemma 2.25 that there exists $y_6 \in \varsigma_{1m_1}[\omega_2, w_{1,4}]$ such that

$$k_D(y_4, y_6) \leq 1 + \mu_0 + \mu_1. \quad (5.96)$$

Next, we prove a lower bound for $k_D(w_{m_1,4}, y_6)$. It follows from (5.88) and (5.96) that

$$k_D(y_5, y_6) \leq k_D(y_5, y_4) + k_D(y_4, y_6) \leq 2(1 + \mu_0 + \mu_1),$$

and thus,

$$\begin{aligned} k_D(y_0, y_5) &\geq k_D(y_0, w_{m_1,4}) - k_D(w_{m_1,4}, y_6) - k_D(y_6, y_5) \\ &\geq k_D(y_0, w_{m_1,4}) - k_D(w_{m_1,4}, y_6) - 2(1 + \mu_0 + \mu_1) \\ &\stackrel{\text{Lemma 5.10(iii)}}{\geq} k_D(y_0, \xi_1) - k_D(w_{m_1,4}, y_6) - \frac{15}{4} - \mu_0 - 5\mu_1. \end{aligned}$$

This, together with (5.89), yields that

$$\begin{aligned} k_D(w_{m_1,4}, y_6) &\geq k_D(y_5, w_{1,4}) - \frac{45}{4} - \mu_0 - 18\mu_1 \geq k_D(w_{1,4}, y_4) - k_D(y_4, y_5) \\ &\stackrel{(5.87)+(5.88)}{\geq} -\frac{51}{4} - 2\mu_0 - 21\mu_1 - \frac{\tau_1}{2} + \tau_3. \end{aligned} \quad (5.97)$$

Observe that (5.87) and (5.96) give

$$k_D(w_{1,4}, y_6) \geq k_D(w_{1,4}, y_4) - k_D(y_4, y_6) \geq -\frac{3}{2} - \mu_0 - 3\mu_1 - \frac{\tau_1}{2} + \tau_3.$$

This, together with Lemma 2.6 and (5.97), implies

$$\begin{aligned} k_D(w_{m_1,4}, w_{1,4}) &\geq \ell_k(\varsigma_{m_14}) - \frac{1}{4} \geq k_D(w_{m_1,4}, y_6) + k_D(w_{1,4}, y_6) - \frac{1}{4} \\ &\geq -\frac{29}{2} - 25\mu_1 - \tau_1 + 2\tau_3, \end{aligned}$$

which establishes (5.85).

Now, we may finish the proof of (3b) based on (5.85). For this, let $w \in \gamma_{34}^{m_1}$ (see Figure 12). We consider two cases: $k_D(w_{m_1,3}, w) \leq 30\mu_1 + \tau_1$ and $k_D(w_{m_1,3}, w) > 30\mu_1 + \tau_1$.

In the former case, it follows from the assumption $k_D(w_{2,3}, w_{m_1,3}) \leq -36\mu_1 - \tau_1 + \tau_2$ that

$$\begin{aligned} k_D(w_{1,4}, w) &\geq k_D(w_{1,4}, w_{2,3}) - k_D(w_{2,3}, w_{m_1,3}) - k_D(w_{m_1,3}, w) \\ &\geq k_D(w_{1,4}, w_{2,3}) - k_D(w_{2,3}, w_{m_1,3}) - 30\mu_1 - \tau_1 \\ &\geq k_D(w_{1,4}, w_{2,3}) + 6\mu_1 - \tau_2. \end{aligned} \quad (5.98)$$

On the other hand, statement (1) and Lemma 2.7 lead to

$$\begin{aligned} k_D(w_{1,4}, w_{2,3}) &\geq k_D(w_{1,3}, w_{1,4}) + \ell_k(\gamma_{12}^1[w_{1,3}, w_{2,3}]) - \frac{7}{4} - 4\mu_1 \\ &\geq k_D(w_{1,3}, w_{1,4}) + \ell_k(\gamma_{12}^1[w_{1,3}, w_{2,1}]) - \frac{7}{4} - 4\mu_1 \\ &\stackrel{(5.81)+\text{Property C(1)}}{\geq} -\frac{7}{4} - 4\mu_1 + \tau_2 + \tau_3. \end{aligned}$$

Combining this with (5.98) gives

$$k_D(w_{1,4}, w) \geq -\frac{7}{4} + 2\mu_1 + \tau_3 > \tau_3.$$

In the latter case, that is, $k_D(w_{m_1,3}, w) > 30\mu_1 + \tau_1$, it follows from Lemma 2.6 and (5.84) that

$$\begin{aligned} k_D(w, w_{m_1,4}) &\leq \ell_k(\gamma_{34}^{m_1}) - \ell_k(\gamma_{34}^{m_1}[w_{m_1,3}, w]) \\ &\leq k_D(w_{m_1,3}, w_{m_1,4}) - k_D(w_{m_1,3}, w) + \frac{1}{4} \\ &\leq \frac{1}{4} - 30\mu_1 - \tau_1 + \tau_3, \end{aligned}$$

and thus, we infer from (5.85) that

$$k_D(w_{1,4}, w) \geq k_D(w_{1,4}, w_{m_1,4}) - k_D(w_{m_1,4}, w) \geq -\frac{59}{4} + 5\mu_1 + \tau_3 > \tau_3.$$

In either cases, we have verified (3b) and hence the proof is complete. \square

Lemma 5.19. *If the six-tuples Ω_1 and Ω_2 satisfy Property C with constant τ_3 , then*

$$k_D(w_{2,1}, w_{2,3}) \leq 11\mu_1.$$

Proof. The proof consists of several steps. In the first step, we shall establish a lower bound for $k_D(w_{2,3}, y_0)$. Since Ω_1 satisfies Property C with constant τ_3 , we obtain from Property C(1) and (5.79) that

$$k_D(w_{1,1}, w_{1,2}) = k_D(w_{1,3}, w_{1,4}) = \tau_3.$$

Meanwhile, Lemma 2.6 gives

$$k_D(w_{1,2}, w_{1,3}) - \frac{1}{4} \leq \ell_k(\gamma_{12}^1) - \frac{1}{4} \leq k_D(w_{1,1}, w_{1,2}) = k_D(w_{1,3}, w_{1,4}).$$

This implies that

$$\min\{k_D(w_{1,2}, w_{1,3}), k_D(w_{1,3}, w_{1,4})\} \geq k_D(w_{1,2}, w_{1,3}) - \frac{1}{4}.$$

On the other hand, since $w_{1,3} \in \gamma_{34}^1 \in \Lambda_{w_{1,3}w_{1,4}}^\lambda$, we obtain from Lemma 5.16(4), the above estimate and Lemma 2.7 that

$$\begin{aligned} k_D(w_{1,3}, y_0) &\leq k_D(y_0, \xi_1) - \min\{k_D(w_{1,2}, w_{1,3}), k_D(w_{1,3}, w_{1,4})\} + \frac{15}{2} + 13\mu_1 \\ &\leq k_D(y_0, \xi_1) - k_D(w_{1,2}, w_{1,3}) + \frac{31}{4} + 13\mu_1 \\ &\leq k_D(y_0, \xi_1) - \ell_k(\gamma_{12}^1[w_{1,2}, w_{1,3}]) + \frac{33}{4} + 13\mu_1 \\ &\leq k_D(y_0, \xi_1) - k_D(w_{1,2}, w_{2,3}) - k_D(w_{2,3}, w_{1,3}) + \frac{33}{4} + 13\mu_1. \end{aligned}$$

This implies that

$$k_D(y_0, \xi_1) - k_D(w_{1,2}, w_{2,3}) \geq k_D(w_{1,3}, y_0) + k_D(w_{1,3}, w_{2,3}) - \frac{33}{4} - 13\mu_1. \quad (5.99)$$

As $w_{1,2} \in \beta_0$, Lemma 5.10(iii) ensures that

$$\begin{aligned} k_D(w_{2,3}, y_0) &\geq k_D(y_0, w_{1,2}) - k_D(w_{1,2}, w_{2,3}) \\ &\geq k_D(y_0, \xi_1) - k_D(w_{1,2}, w_{2,3}) - \frac{7}{4} - 2\mu_0 - 2\mu_1 \\ &\stackrel{(5.99)}{\geq} k_D(w_{1,3}, y_0) + k_D(w_{1,3}, w_{2,3}) - 10 - 2\mu_0 - 15\mu_1. \end{aligned} \quad (5.100)$$




FIGURE 13. Illustration for the proof of Lemma 5.19

In the second step, we construct several auxiliary points. Fix $\varsigma_{23} \in \Lambda_{w_{2,3}y_0}^\lambda$ and $\varsigma_{13} \in \Lambda_{w_{1,3}y_0}^\lambda$. As (D, k_D) is a (μ_0, ρ_0) -Rips space by Proposition 3.4(P_4), we know from Lemma 2.24 that there exist $p_1 \in \varsigma_{23}$, $p_2 \in \gamma_{12}^1[w_{2,3}, w_{1,3}]$ and $p_3 \in \varsigma_{13}$ such that for any $r \neq s \in \{1, 2, 3\}$,

$$k_D(p_r, p_s) \leq \mu_1. \quad (5.101)$$

Then we make the following claim concerning the position of $w_{2,1}$.

Claim 5.19-I: $w_{2,1} \in \gamma_{12}^1[w_{2,3}, p_2]$ (see Figure 13).

Indeed, observe that

$$\begin{aligned} k_D(w_{2,3}, y_0) &\leq k_D(w_{2,3}, p_2) + k_D(p_2, p_3) + k_D(p_3, y_0) \stackrel{(5.101)}{\leq} k_D(w_{2,3}, p_2) + k_D(p_3, y_0) + \mu_1 \\ &\leq \ell_k(\varsigma_{13}) - \ell_k(\varsigma_{13}[w_{1,3}, p_3]) + k_D(w_{2,3}, p_2) + \mu_1 \\ &\stackrel{\text{Lemma 2.6}}{\leq} k_D(w_{1,3}, y_0) - (k_D(w_{1,3}, p_2) - k_D(p_2, p_3)) + k_D(w_{2,3}, p_2) + \frac{1}{4} + \mu_1 \\ &\leq k_D(w_{1,3}, y_0) - k_D(w_{1,3}, p_2) + k_D(w_{2,3}, p_2) + \frac{1}{4} + 2\mu_1. \end{aligned}$$

Combining this with (5.100), we get

$$k_D(w_{1,3}, p_2) \leq k_D(w_{2,3}, p_2) - k_D(w_{2,3}, w_{1,3}) + \frac{41}{4} + 2\mu_0 + 17\mu_1.$$

On the other hand, by Lemma 2.7, we have

$$k_D(w_{2,3}, p_2) - k_D(w_{2,3}, w_{1,3}) + k_D(w_{1,3}, p_2) \leq \ell_k(\gamma_{12}^1[w_{2,3}, w_{1,3}]) - k_D(w_{2,3}, w_{1,3}) \leq \frac{1}{2}.$$

The above two estimates, together with Lemma 2.7, imply

$$\ell_k(\gamma_{12}^1[w_{1,3}, p_2]) \leq k_D(w_{1,3}, p_2) + \frac{1}{2} \leq \frac{47}{8} + \frac{5}{2}\mu_0 + 9\mu_1 < \tau_2.$$

Then we infer from the choice of $w_{2,1}$ in (5.81) and the previous estimate that

$$\ell_k(\gamma_{12}^1[w_{1,3}, w_{2,1}]) \geq \tau_2 \geq \ell_k(\gamma_{12}^1[w_{1,3}, p_2]).$$

This proves the claim.

Fix $s_{24} \in \Lambda_{w_{2,4}y_0}^\lambda$ and $\gamma_{34}^2 \in \Lambda_{w_{2,3}w_{2,4}}^\lambda$. Then as (D, k_D) is a (μ_0, ρ_0) -Rips space, Lemma 2.24 ensures that there exist $p_4 \in s_{23}$, $p_5 \in \gamma_{34}^2$ and $p_6 \in s_{24}$ such that for any $r \neq s \in \{4, 5, 6\}$,

$$k_D(p_r, p_s) \leq \mu_1. \quad (5.102)$$

Thus we have the following basic estimates.

$$k_D(w_{2,3}, p_4) \leq k_D(w_{2,3}, p_5) + k_D(p_5, p_4) \leq k_D(w_{2,3}, p_5) + \mu_1, \quad (5.103)$$

$$k_D(p_4, y_0) \leq k_D(p_6, y_0) + k_D(p_6, p_4) \leq k_D(p_6, y_0) + \mu_1 \quad (5.104)$$

and

$$k_D(w_{2,4}, p_6) \geq k_D(w_{2,4}, p_5) - k_D(p_5, p_6) \geq k_D(w_{2,4}, p_5) - \mu_1. \quad (5.105)$$

Next, we establish a lower bound for $k_D(w_{2,3}, p_4)$. Note that Lemma 2.6 and (5.105) give

$$\begin{aligned} k_D(w_{2,4}, y_0) &\geq \ell_k(s_{24}) - \frac{1}{4} \geq k_D(w_{2,4}, p_6) + k_D(p_6, y_0) - \frac{1}{4} \\ &\geq k_D(w_{2,4}, p_5) + k_D(p_6, y_0) - \frac{1}{4} - \mu_1. \end{aligned}$$

Since $w_{24}, w_{1,2} \in \beta_0$, applying Lemma 5.10(iii) twice gives

$$k_D(w_{2,4}, y_0) \leq k_D(y_0, \xi_1) + \frac{7}{4} + 2\mu_0 + 2\mu_1 \leq k_D(w_{1,2}, y_0) + \frac{7}{2} + \mu_0 + 5\mu_1.$$

Then we get from the above two estimates that

$$\begin{aligned} k_D(w_{2,4}, p_5) &\leq k_D(w_{1,2}, y_0) - k_D(p_6, y_0) + \frac{15}{4} + \mu_0 + 6\mu_1 \\ &\leq k_D(w_{1,2}, w_{2,3}) + k_D(w_{2,3}, p_4) + k_D(p_4, y_0) - k_D(p_6, y_0) + \frac{15}{4} + \mu_0 + 6\mu_1 \\ &\stackrel{(5.103)+(5.104)}{\leq} k_D(w_{1,2}, w_{2,3}) + k_D(w_{2,3}, p_5) + \frac{15}{4} + \mu_0 + 8\mu_1. \end{aligned}$$

Note that Property C(1) implies

$$k_D(w_{2,4}, p_5) \geq k_D(w_{2,4}, w_{2,3}) - k_D(w_{2,3}, p_5) = \tau_3 - k_D(w_{2,3}, p_5),$$

and we obtain from the above two estimates that

$$k_D(w_{2,3}, p_5) \geq -\frac{1}{2}k_D(w_{1,2}, w_{2,3}) - \frac{15}{8} - \frac{1}{2}\mu_0 - 4\mu_1 + \frac{\tau_3}{2}. \quad (5.106)$$

Furthermore, Lemma 2.6 and (5.79) yield

$$k_D(w_{1,2}, w_{2,3}) \leq \ell_k(\gamma_{12}^1) - \ell_k(\gamma_{12}^1[w_{1,1}, w_{2,3}]) \leq \tau_3 - \ell_k(\gamma_{12}^1[w_{1,3}, w_{2,3}]) + \frac{1}{4}.$$

Substituting the above estimate in (5.106) gives

$$k_D(w_{2,3}, p_5) \geq \frac{1}{2}\ell_k(\gamma_{12}^1[w_{1,3}, w_{2,3}]) - 2 - \frac{1}{2}\mu_0 - 4\mu_1.$$

This, together with (5.102) and Lemma 2.7, implies that

$$\begin{aligned}
k_D(w_{2,3}, p_4) &\geq k_D(w_{2,3}, p_5) - k_D(p_5, p_4) \geq \frac{1}{2}\ell_k(\gamma_{12}^1[w_{1,3}, w_{2,3}]) - 2 - \frac{1}{2}\mu_0 - 5\mu_1 \\
&= \frac{1}{2}\ell_k(\gamma_{12}^1[w_{2,1}, w_{2,3}]) + \frac{1}{2}\ell_k(\gamma_{12}^1[w_{1,3}, w_{2,1}]) - 2 - \frac{1}{2}\mu_0 - 5\mu_1 \\
&\stackrel{(5.81)}{\geq} \frac{1}{2}\ell_k(\gamma_{12}^1[w_{2,1}, w_{2,3}]) - 2 - \frac{1}{2}\mu_0 - 5\mu_1 + \frac{1}{2}\tau_2.
\end{aligned} \tag{5.107}$$

In the third step, we continue to find more auxiliary points. Firstly, we see that there is $p_7 \in \gamma_{12}^1[w_{2,1}, w_{2,3}]$ (see Figure 13) such that

$$k_D(w_{2,3}, p_7) = \frac{1}{2}k_D(w_{2,1}, w_{2,3}). \tag{5.108}$$

Since Ω_2 satisfies Property C with constant τ_3 , $p_7 \in \gamma_{12}^2 = \gamma_{12}^1[w_{2,1}, w_{2,3}]$ and $w_{2,4} \in \beta_0[w_{2,4}, \varpi_2]$, we have

$$k_D(w_{2,3}, w_{2,4}) = \tau_3 \leq k_D(w_{2,4}, p_7) + \frac{1}{4}. \tag{5.109}$$

Secondly, we know from Claim 5.19-I, (5.101) and Lemma 2.25 (replacing x , y and z by $w_{2,3}$, p_2 and p_1) that there is $p_8 \in \varsigma_{23}[w_{2,3}, p_1]$ such that

$$k_D(p_7, p_8) \leq 1 + \mu_0 + \mu_1 \tag{5.110}$$

and so,

$$\begin{aligned}
\ell_k(\varsigma_{23}[w_{2,3}, p_8]) &\leq k_D(w_{2,3}, p_8) + \frac{1}{2} \leq k_D(w_{2,3}, p_7) + k_D(p_7, p_8) + \frac{1}{2} \\
&\stackrel{(5.108)+(5.110)}{\leq} \frac{1}{2}k_D(w_{2,1}, w_{2,3}) + \frac{3}{2} + \mu_0 + \mu_1 \\
&\stackrel{(5.107)}{\leq} k_D(w_{2,3}, p_4) + \frac{7}{2} + \frac{3}{2}\mu_0 + 6\mu_1 - \frac{1}{2}\tau_2 \\
&< \ell_k(\varsigma_{23}[w_{2,3}, p_4]),
\end{aligned}$$

which demonstrates that

$$p_8 \in \varsigma_{23}[w_{2,3}, p_4].$$

Thirdly, considering $w_{2,3}$, p_4 and p_5 , we know from (5.102) and Lemma 2.25 (see Figure 13) that there is $p_9 \in \gamma_{34}^2[w_{2,3}, p_5]$ such that

$$k_D(p_8, p_9) \leq 1 + \mu_0 + \mu_1.$$

Hence (5.110) gives

$$k_D(p_7, p_9) \leq k_D(p_7, p_8) + k_D(p_8, p_9) \leq 2(1 + \mu_0 + \mu_1). \tag{5.111}$$

Now, we are ready to finish the proof. By (5.109), (5.111) and Lemma 2.6, we know that

$$\begin{aligned}
k_D(w_{2,3}, p_9) &\leq \ell_k(\gamma_{34}^2) - k_D(w_{2,4}, p_9) \leq k_D(w_{2,3}, w_{2,4}) - k_D(w_{2,4}, p_9) + \frac{1}{4} \\
&\stackrel{(5.109)}{\leq} k_D(w_{2,4}, p_9) + k_D(p_9, p_7) - k_D(w_{2,4}, p_9) + \frac{1}{2} \stackrel{(5.111)}{\leq} \frac{5}{2} + 2\mu_0 + 2\mu_1.
\end{aligned}$$

This, together with the choice of p_7 in (5.108) and (5.111), gives

$$k_D(w_{2,1}, w_{2,3}) = 2k_D(w_{2,3}, p_7) \leq 2(k_D(w_{2,3}, p_9) + k_D(p_7, p_9)) \leq 9 + 2\mu_0 + 10\mu_1 \leq 11\mu_1.$$

The proof is thus complete. \square

5.7. Iterative construction of new six-tuples. In this section, we shall prove an useful lemma, which gives a way to iteratively construct new six-tuples based on old ones. To this end, for each integer $n \geq 1$, we set

$$\varrho_n = \tau_2 \sum_{i=1}^n \frac{1}{4^i} + 85\mu_1 \sum_{i=1}^n \frac{1}{2^i}. \quad (5.112)$$

Then, it is clear that

$$\frac{85}{2}\mu_1 + \frac{1}{4}\tau_2 < \varrho_n < 85\mu_1 + \frac{1}{3}\tau_2. \quad (5.113)$$

Lemma 5.20. *Let $w_{1,1} \in D$, $w_{1,2} \in \beta_0$, $\varpi \in \beta_0[w_{1,2}, \xi_m]$, $\gamma_{12}^1 \in \Lambda_{w_{1,1}w_{1,2}}^\lambda$, $w_{1,3} \in \gamma_{12}^1$ and $w_{1,4} \in \beta_0[w_{1,2}, \varpi]$ (see Figure 14). Suppose that the six-tuple $\Omega_1 = [w_{1,1}, w_{1,2}, \gamma_{12}^1, \varpi, w_{1,3}, w_{1,4}]$ satisfies Property C with constant τ_3 ,*

$$k_D(w_{1,1}, w_{1,2}) = \tau_3, \quad (5.114)$$

and there is an integer $n \geq 2$ such that

$$k_D(w_{1,1}, w_{1,3}) \leq \varrho_n. \quad (5.115)$$

Then the following statements hold:

- (1) there exist $u_1 \in D$, $u_2 \in \beta_0[w_{1,2}, w_{1,4}]$, $\alpha_{12} \in \Lambda_{u_1 u_2}^\lambda$, $\varpi_1 = w_{1,4} \in \beta_0[u_2, \varpi]$, $u_3 \in \alpha_{12}$, and $u_4 \in \beta_0[u_2, \varpi_1]$ such that the six-tuple $\Omega = [u_1, u_2, \alpha_{12}, \varpi_1, u_3, u_4]$ satisfies Property C with constant τ_3 ;
- (2) $k_D(u_1, u_2) = \tau_3$, $k_D(w_{1,2}, u_2) \geq \tau_3 - \frac{1}{4}$, and
- (3) $k_D(u_1, u_3) \leq \varrho_{n+1}$.

Proof. To find the required points in the lemma, we need to construct several auxiliary six-tuples based on Ω_1 . The construction of the first new six-tuple is included in the following claim.




FIGURE 14. Illustration for the proof of Lemma 5.20

Claim 5.20-I: Let $w_{2,2} = w_{1,2}$. Then there are $w_{2,1} \in \gamma_{12}^1[w_{1,3}, w_{2,2}]$, $\gamma_{12}^2 \in \Lambda_{w_{2,1}w_{2,2}}^\lambda$, $\varpi_2 = w_{1,4}$, $w_{2,3} \in \gamma_{12}^2$ and $w_{2,4} \in \beta_0[w_{2,2}, \varpi_2]$ (see Figure 14) such that

- (1) the six-tuple $\Omega_2 = [w_{2,1}, w_{2,2}, \gamma_{12}^2, \varpi_2, w_{2,3}, w_{2,4}]$ satisfies Property C with constant τ_3 ;
- (2) $k_D(w_{1,3}, w_{2,1}) = \tau_2$, and
- (3) for any $u \in \gamma_{12}^2$ and any $v \in \beta_0[w_{2,4}, \varpi_2]$, $k_D(u, v) \geq \tau_3 - \frac{1}{4}$.

We first find the point $w_{2,1}$. Note that

$$k_D(w_{1,3}, w_{2,2}) \geq k_D(w_{2,2}, w_{1,1}) - k_D(w_{1,1}, w_{1,3}) \stackrel{(5.114)+(5.115)}{\geq} \tau_3 - \varrho_n \stackrel{(5.113)}{>} 3\tau_2 \quad (5.116)$$

and so there exists $w_{2,1} \in \gamma_{12}^1[w_{1,3}, w_{2,2}]$ (see Figure 14) such that

$$k_D(w_{1,3}, w_{2,1}) = \tau_2. \quad (5.117)$$

Then, we know from Lemma 2.6 that

$$\begin{aligned} k_D(w_{2,1}, w_{2,2}) &\leq \ell_k(\gamma_{12}^1[w_{2,1}, w_{2,2}]) = \ell_k(\gamma_{12}^1) - \ell_k(\gamma_{12}^1[w_{1,3}, w_{2,1}]) \\ &\leq k_D(w_{1,1}, w_{1,2}) - k_D(w_{1,3}, w_{2,1}) + \frac{1}{4} \stackrel{(5.114)+(5.117)}{=} \frac{1}{4} - \tau_2 + \tau_3. \end{aligned} \quad (5.118)$$

Set $\gamma_{12}^2 = \gamma_{12}^1[w_{2,1}, w_{2,2}]$ and $\varpi_2 = w_{1,4}$. We shall apply Lemma 5.14 to find the remaining two points $w_{2,3}$ and $w_{2,4}$. For this, we need to show that τ_3 is a lower bound for the quantity $k_D(v, w_{1,4})$ for any $v \in \gamma_{12}^2$.

Since the six-tuple Ω_1 satisfies Property C with constant τ_3 , we see from Lemma 5.16(3) that

$$\begin{aligned} k_D(w_{1,4}, w_{2,2}) &= k_D(w_{1,2}, w_{1,4}) \geq k_D(w_{1,2}, w_{1,3}) - \frac{5}{4} - 5\mu_1 + \tau_3 \\ &\stackrel{(5.116)}{\geq} -\frac{5}{4} - 5\mu_1 - \varrho_n + 2\tau_3. \end{aligned} \quad (5.119)$$

For any $v \in \gamma_{12}^2$, we know from Lemma 2.7 that

$$\begin{aligned} k_D(v, w_{1,4}) &\geq k_D(w_{1,4}, w_{2,2}) - k_D(w_{2,2}, v) \geq k_D(w_{1,4}, w_{2,2}) - \ell_k(\gamma_{12}^1[w_{2,2}, v]) \\ &\geq k_D(w_{1,4}, w_{2,2}) - \ell_k(\gamma_{12}^1[w_{2,2}, w_{2,1}]) \\ &\geq k_D(w_{1,4}, w_{2,2}) - k_D(w_{2,2}, w_{2,1}) - \frac{1}{2} \\ &\stackrel{(5.118)+(5.119)}{\geq} -2 - 5\mu_1 - \varrho_n + \tau_2 + \tau_3 \stackrel{(5.113)}{>} \tau_3 \end{aligned} \quad (5.120)$$

Now, we see from (5.118) and (5.120) that $w_{2,1}$, $w_{2,2}$, γ_{12}^2 and $\varpi_2 = w_{1,4}$ satisfy all the assumptions in Lemma 5.14. It follows that there are $w_{2,3} \in \gamma_{12}^2$ and $w_{2,4} \in \beta_0[w_{2,2}, w_{1,4}]$ such that the six-tuple $\Omega_2 = [w_{2,1}, w_{2,2}, \gamma_{12}^2, w_{1,4}, w_{2,3}, w_{2,4}]$ satisfies Property C with constant τ_3 (see Figure 14). Then Property C(2) guarantees that for any $u \in \gamma_{12}^2$ and any $v \in \beta_0[w_{2,4}, w_{1,4}]$,

$$k_D(u, v) \geq \tau_3 - \frac{1}{4}.$$

This proves the last statement of Claim 5.20-I, and hence, the proof of claim is complete.

The construction of next six-tuple is formulated in the following claim.

Claim 5.20-II: Let $w_{3,1} = w_{2,3}$, $w_{3,2} = w_{2,4}$, $\gamma_{12}^3 \in \Lambda_{w_{3,1}w_{3,2}}^\lambda$ and $\varpi_3 = w_{1,4}$. Then

- (1) $k_D(w_{3,1}, w_{3,2}) = \tau_3$;
- (2) there are $w_{3,3} \in \gamma_{12}^3$ and $w_{3,4} \in \beta_0[w_{3,2}, \varpi_3]$ such that
 - (a) the six-tuple $\Omega_3 = [w_{3,1}, w_{3,2}, \gamma_{12}^3, \varpi_3, w_{3,3}, w_{3,4}]$ satisfies Property C with constant τ_3 ;

- (b) $k_D(w_{1,2}, w_{3,2}) = k_D(w_{2,2}, w_{3,2}) \geq \tau_3 - \frac{1}{4}$, and
- (c) $k_D(w_{2,3}, w_{3,3}) = k_D(w_{3,1}, w_{3,3}) \leq \frac{67}{2}\mu_1 + \frac{1}{2}\varrho_n + \frac{1}{2}\tau_2$.

Since Ω_2 satisfies Property C with constant τ_3 , we infer from Property C(1) that

$$k_D(w_{3,1}, w_{3,2}) = k_D(w_{2,3}, w_{2,4}) = \tau_3.$$

This implies statement (1) of Claim 5.20-II.

Moreover, we know from Lemma 5.18(2b) that for any $w \in \gamma_{34}^2 = \gamma_{12}^3$,

$$k_D(w, w_{1,4}) > \tau_3. \quad (5.121)$$

Thus, we know from statement (1) of Claim 5.20-II, (5.121) and Lemma 5.14 (replacing $w_1, w_2, \varpi, \mathbf{c}$ with $w_{3,1}, w_{3,2}, w_{1,4}, \tau_3$) that there exist $w_{3,3} \in \gamma_{12}^3$ and $w_{3,4} \in \beta_0[w_{3,2}, w_{1,4}]$, such that the six-tuple $\Omega_3 = [w_{3,1}, w_{3,2}, \gamma_{12}^3, w_{1,4}, w_{3,3}, w_{3,4}]$ satisfies Property C with constant τ_3 (see Figure 14). This proves the statement (2a).

Since $w_{1,2} = w_{2,2} \in \gamma_{12}^2$ and $w_{3,2} = w_{2,4} \in \beta_0[w_{2,4}, \varpi_2]$, statement (2b) follows from Claim 5.20-I(3).

Since both Ω_1 and Ω_2 satisfy Property C with constant τ_3 , we know from Lemma 5.19 that

$$k_D(w_{2,1}, w_{2,3}) \leq 11\mu_1, \quad (5.122)$$

and since both Ω_2 and Ω_3 satisfy Property C with constant τ_3 , it follows from Lemma 5.17 that

$$k_D(w_{2,3}, w_{3,3}) = k_D(w_{3,1}, w_{3,3}) \leq \frac{1}{2}(\tau_3 - k_D(w_{2,2}, w_{2,3})) + 28\mu_1. \quad (5.123)$$

Then (5.116), the choice of $w_{2,1}$ in (5.117) and (5.122) show that

$$\begin{aligned} k_D(w_{2,2}, w_{2,3}) &\geq k_D(w_{2,2}, w_{1,3}) - k_D(w_{1,3}, w_{2,1}) - k_D(w_{2,1}, w_{2,3}) \\ &\geq -11\mu_1 - \varrho_n - \tau_2 + \tau_3. \end{aligned}$$

Substituting the above estimate in (5.123) gives

$$k_D(w_{3,1}, w_{3,3}) = k_D(w_{2,3}, w_{3,3}) \leq \frac{67}{2}\mu_1 + \frac{1}{2}\varrho_n + \frac{1}{2}\tau_2.$$

This proves the statement (2c), and hence, the proof of claim is complete.

One more six-tuple is constructed in the following claim.

Claim 5.20-III: *Let $w_{4,1} = w_{3,3}$, $w_{4,2} = w_{3,4}$, $\gamma_{12}^4 \in \Lambda_{w_{4,1}w_{4,2}}^\lambda$ and $\varpi_4 = w_{1,4}$. Then*

- (1) $k_D(w_{4,1}, w_{4,2}) = \tau_3$;
- (2) *there are $w_{4,3} \in \gamma_{14}^2$ and $w_{4,4} \in \beta_0[w_{4,2}, \varpi_4]$ such that*
 - (a) *the six-tuple $\Omega_4 = [w_{4,1}, w_{4,2}, \gamma_{12}^4, \varpi_4, w_{4,3}, w_{4,4}]$ satisfies Property C with constant τ_3 ;*
 - (b) $k_D(w_{1,2}, w_{4,2}) = k_D(w_{2,2}, w_{4,2}) \geq \tau_3 - \frac{1}{4}$, and
 - (c) $k_D(w_{4,1}, w_{4,3}) = k_D(w_{4,1}, w_{4,3}) \leq 45\mu_1 + \frac{1}{4}\varrho_n + \frac{1}{4}\tau_2$.

We have shown that all six-tuples Ω_1, Ω_2 and Ω_3 satisfy Property C. Thus, statement (1) of Claim 5.20-III follows from Property C(1). Moreover, as Claim 5.20-I(2b) shows

$$k_D(w_{3,1}, w_{3,3}) = k_D(w_{2,3}, w_{3,3}) \leq \frac{67}{2}\mu_1 + \frac{1}{2}\varrho_n + \frac{1}{2}\tau_2 \stackrel{(5.113)}{\leq} -36\mu_1 - \tau_1 + \tau_2.$$

we know from Lemma 5.18(3b) that for any $w \in \gamma_{12}^4 = \gamma_{34}^3$,

$$k_D(w, w_{1,4}) > \tau_3. \quad (5.124)$$

Thus we know from Lemma 5.14 (replacing $w_1, w_2, \varpi, \mathbf{c}$ with $w_{4,1}, w_{4,2}, w_{1,4}, \tau_3$), together with the statement (1) of Claim 5.20-III and (5.124), that there exist $w_{4,3} \in \gamma_{12}^4$ and $w_{4,4} \in \beta_0[w_{4,2}, \varpi_4]$, such that the six-tuple $\Omega_4 = [w_{4,1}, w_{4,2}, \gamma_{12}^4, \varpi_4, w_{4,3}, w_{4,4}]$ satisfies Property C with constant τ_3 (see Figure 14). This proves the statement (2a).

Since $w_{2,2} = w_{2,1}$ and $w_{4,2} = w_{3,4} \in \beta_0[w_{2,4}, w_{1,4}]$, we see from Claim 5.20-I(3) that statement (2b) holds.

Since Ω_2 satisfies Property C with constant τ_3 , it follows from Property C(1) that

$$k_D(w_{3,1}, w_{3,2}) = k_D(w_{2,3}, w_{2,4}) = \tau_3,$$

and so, we have

$$k_D(w_{2,3}, w_{3,3}) \geq k_D(w_{3,1}, w_{3,2}) - k_D(w_{3,2}, w_{3,3}) = \tau_3 - k_D(w_{3,2}, w_{3,3}).$$

Since both Ω_3 and Ω_4 satisfy Property C, we obtain from Lemma 5.17 (with $\Omega_1 = \Omega_3$ and $\Omega_2 = \Omega_4$) and the above estimate that

$$\begin{aligned} k_D(w_{3,3}, w_{4,3}) &= k_D(w_{4,1}, w_{4,3}) \leq \frac{1}{2}(\tau_3 - k_D(w_{3,2}, w_{3,3})) + 28\mu_1 \\ &\leq \frac{1}{2}k_D(w_{2,3}, w_{3,3}) + 28\mu_1 \stackrel{\text{Claim 5.20-II(2c)}}{\leq} 45\mu_1 + \frac{1}{4}\varrho_n + \frac{1}{4}\tau_2. \end{aligned}$$

Since $w_{3,3} = w_{4,1}$, the above inequality shows that statement (2c) holds, and hence, the proof of claim is complete.

Let $u_1 = w_{4,1}$, $u_2 = w_{4,2}$, $\alpha_{12} = \gamma_{12}^4$, $\varpi_1 = w_{1,4}$, $u_3 = w_{4,3}$ and $u_4 = w_{4,4}$. Then statements (1) and (2) in Lemma 5.20 follow from Claim 5.20-III (1), (2a) and (2b). Since by (5.112), $45\mu_1 + \frac{1}{4}\varrho_n + \frac{1}{4}\tau_2 \leq \varrho_{n+1}$ when $n \geq 2$, statement (3) in Lemma 5.20 follows from Claim 5.20-III (2c). The proof of lemma is thus complete. \square

5.8. Final proof of Theorem 3.3. In this section, we complete the proof of Theorem 3.3. Towards a contradiction, set

$$\tau_2 = \frac{1}{32} \log \mu_8 \quad \text{and} \quad \tau_3 = \frac{1}{8} \log \mu_8. \quad (5.125)$$

For $i \in \{1, \dots, m\}$, let ξ_i be the points defined in Section 5.4, and $w_{1,2} = \xi_1$ (see Figure 15). Since Lemma 5.10(i) and the choice of ξ_i in (5.31) ensure that

$$k_D(\xi_i, y_0) = k_D(w_{1,2}, y_0) \geq \log \frac{\mu_8}{2(1 + \mu_9)e^C(1 + c_3)^{c_1 M} \mu_7} - \frac{1}{2} \stackrel{(5.125)}{\geq} 2\tau_3 + \frac{1}{2}, \quad (5.126)$$

we infer from the above estimate that there exists $w_{1,1} \in \gamma_1[w_{1,2}, y_0]$ such that

$$k_D(w_{1,1}, w_{1,2}) = \tau_3. \quad (5.127)$$

Let $\gamma_{12}^1 = \gamma_1[w_{1,1}, w_{1,2}]$. Then for any $w \in \gamma_{12}^1$, we know from Lemma 2.7 that

$$k_D(w, \xi_m) \geq \ell_k(\gamma_1[w, \xi_m]) - \frac{1}{2} > \ell_k(\gamma_1[\xi_m, y_0]) - \frac{1}{2} \geq k_D(\xi_m, y_0) - \frac{1}{2} \stackrel{(5.126)}{>} 2\tau_3.$$

Thus it follows from (5.127) and Lemma 5.14 that there exist $w_{1,3} \in \gamma_{12}^1$ and $w_{1,4} \in \beta_0[w_{1,2}, \xi_m]$ such that for $\varpi = w_{1,4}$, the six-tuple $[w_{1,1}, w_{1,2}, \gamma_{12}^1, \varpi, w_{1,3}, w_{1,4}]$ satisfies Property C with constant τ_3 (see Figure 15). Then by Lemma 5.15, it holds

$$k_D(w_{1,1}, w_{1,3}) \leq 16\mu_1 \stackrel{(5.112)}{\leq} \varrho_2,$$




FIGURE 15. Illustration for the proof of Theorem 3.3

and thus, we infer from (5.127) and Lemma 5.20 that there exist $w_{2,1} \in D$, $w_{2,2} \in \beta_0[w_{1,2}, w_{1,4}]$, $\gamma_{12}^2 \in \Lambda_{w_{2,1}, w_{2,2}}^\lambda$, $\varpi_2 = w_{1,4} \in \beta_0[w_{2,2}, w_{1,4}]$, $w_{2,3} \in \gamma_{12}^2$ and $w_{2,4} \in \beta_0[w_{2,2}, \varpi_2]$ such that the six-tuple $[w_{2,1}, w_{2,2}, \gamma_{12}^2, \varpi_2, w_{2,3}, w_{2,4}]$ satisfies Property C with constant τ_3 (see Figure 15). Moreover, it holds

$$k_D(w_{2,1}, w_{2,2}) = \tau_3, \quad k_D(w_{1,2}, w_{2,2}) \geq \tau_3 - \frac{1}{4} \quad \text{and} \quad k_D(w_{2,1}, w_{2,3}) \leq \varrho_3.$$

Now, applying Lemma 5.20 once again, we know that there exist $w_{3,1} \in D$, $w_{3,2} \in \beta_0[w_{2,2}, w_{2,4}]$, $\gamma_{12}^3 \in \Lambda_{w_{3,1}, w_{3,2}}^\lambda$, $\varpi_3 = w_{2,4} \in \beta_0[w_{3,2}, w_{2,4}]$, $w_{3,3} \in \gamma_{12}^3$ and $w_{3,4} \in \beta_0[w_{3,2}, \varpi_3]$ such that the six-tuple $[w_{3,1}, w_{3,2}, \gamma_{12}^3, \varpi_3, w_{3,3}, w_{3,4}]$ satisfies Property C with constant τ_3 (see Figure 15). Moreover, it holds

$$k_D(w_{3,1}, w_{3,2}) = \tau_3, \quad k_D(w_{2,2}, w_{3,2}) \geq \tau_3 - \frac{1}{4} \quad \text{and} \quad k_D(w_{3,1}, w_{3,3}) \leq \varrho_4.$$

Repeating this procedure for m_1 times, where $m_1 = \lceil \frac{\ell_k(\beta_0)}{\tau_3 - \frac{1}{4}} \rceil + 3$, we get six sequences $\{w_{i,1}\}_{i=1}^{m_1}$, $\{w_{i,2}\}_{i=1}^{m_1}$, $\{\gamma_{12}^i\}_{i=1}^{m_1}$, $\{\varphi_i\}_{i=1}^{m_1}$, $\{w_{i,3}\}_{i=1}^{m_1}$ and $\{w_{i,4}\}_{i=1}^{m_1}$ (see Figure 15) such that

- (1) $w_{1,1} \in \gamma_1[y_0, \xi_1]$, $w_{1,2} = \xi_1$, $w_{1,3} \in \gamma_1[w_{1,1}, w_{1,2}]$ and $w_{1,4} \in \beta_0[w_{1,2}, \xi_m]$;
- (2) for each $i \in \{2, \dots, m_1\}$, $w_{i,1} \in D$, $w_{i,2} \in \beta_0[w_{i-1,2}, w_{i-1,4}]$, $\gamma_{12}^i \in \Lambda_{w_{i,1}, w_{i,2}}^\lambda$, $\varpi_i = w_{i-1,4}$, $w_{i,3} \in \gamma_{12}^{i-1}$ and $w_{i,4} \in \beta_0[w_{i,2}, \varpi_i]$;
- (3) for each $i \in \{1, \dots, m_1\}$, the six-tuple $[w_{i,1}, w_{i,2}, \gamma_{12}^i, \varpi_i, w_{i,3}, w_{i,4}]$ satisfies Property C with constant τ_3 , where $w_{0,4} = \xi_m$, and
- (4) for each $i \in \{1, \dots, m_1 - 1\}$, $k_D(w_{i,2}, w_{i+1,2}) \geq \tau_3 - \frac{1}{4}$.

Then summing over all i in (4) gives

$$\sum_{i=1}^{m_1-1} k_D(w_{i,2}, w_{i+1,2}) \geq \left(\tau_3 - \frac{1}{4}\right)(m_1 - 1) \geq \ell_k(\beta_0) + \tau_3 - \frac{1}{4},$$

which is impossible as

$$\sum_{i=1}^{m_1-1} k_D(w_{i,2}, w_{i+1,2}) \leq \ell_k(\beta_0).$$

The proof of Theorem 3.3 is thus complete.

Acknowledgment. We would like to thank Prof. Pekka Koskela for helpful discussions and suggestions that greatly improved our exposition.

REFERENCES

- [1] Z. M. BALOGH AND M. BONK, Lengths of radii under conformal maps of the unit disc, *Proc. Amer. Math. Soc.*, **127** (1999), 801–804.
- [2] Z. M. BALOGH AND S. M. BUCKLEY, Geometric characterizations of Gromov hyperbolicity, *Invent. Math.*, **153** (2003), 261–301.
- [3] R. BANUELOS, On an estimate of Cranston and McConnell for elliptic diffusions in uniform domains, *Probab. Theory Related Fields* **76** (1987), no. 3, 311–323.
- [4] M. BONK AND B. KLEINER, Conformal dimension and Gromov hyperbolic groups with 2-sphere boundary, *Geom. Topol.* **9** (2005), 219–246.
- [5] M. BONK, J. HEINONEN AND P. KOSKELA, Uniformizing Gromov hyperbolic domains, *Asterisque*, **270** (2001), 1–99.
- [6] M. BONK AND P. KOSKELA, Conformal metrics and size of the boundary, *Amer. J. Math.*, **124** (2002), 1247–1287.
- [7] M. BONK, P. KOSKELA AND S. ROHDE, Conformal metrics on the unit ball in Euclidean space, *Proc. London Math. Soc.*, **77** (1998), 635–664.
- [8] M. R. BRIDSON AND A. HAEFLIGER, Metric spaces of non-positive curvature. Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 319. Springer-Verlag, Berlin, 1999.
- [9] S. M. BUCKLEY AND D. A. HERRON, Quasihyperbolic geodesics are hyperbolic quasi-geodesics, *J. Eur. Math. Soc. (JEMS)* **22** (2020), 1917–1970.
- [10] F. W. GEHRING, Uniform domains and the ubiquitous quasidisk, *Jahresber. Deutsch. Math. Verein*, **89** (1987), 88–103.
- [11] F. W. GEHRING AND W. K. HAYMAN, An inequality in the theory of conformal mapping, *J. Math. Pure Appl.*, **41** (1962), 353–361.
- [12] F. W. GEHRING AND B. G. OSGOOD, Uniform domains and the quasi-hyperbolic metric, *J. Analyse Math.*, **36** (1979), 50–74.
- [13] F. W. GEHRING AND B. P. PALKA, Quasiconformally homogeneous domains, *J. Analyse Math.*, **30** (1976), 172–199.
- [14] P. GOLDSTEIN, Z. GROCHULSKA, C.-Y. GUO, P. KOSKELA AND D. NANDI, Characterizations of generalized John domains via homological bounded turning, *Colloq. Math.* **176** (2024), no. 1, 87–105.
- [15] M. GROMOV, Hyperbolic groups, Essays in group theory, Math. Sci. Res. Inst. Publ., 8, Springer, New York, 1987, 75–263.
- [16] C.-Y. GUO, Uniform continuity of quasiconformal mappings onto generalized John domains, *Ann. Acad. Sci. Fenn. Math.* **40** (2015), no. 1, 183–202.
- [17] P. HAJLASZ AND P. KOSKELA, Sobolev met Poincaré, *Mem. Amer. Math. Soc.* **145** (2000), no. 688.
- [18] J. HEINONEN AND P. KOSKELA, Quasiconformal mappings in metric spaces with controlled geometry, *Acta Math.*, **181** (1998), 1–61.
- [19] J. HEINONEN AND R. NÄKKI, Quasiconformal distortion on arcs, *J. Anal. Math.*, **63** (1994), 19–53.
- [20] J. HEINONEN AND S. ROHDE, The Gehring-Hayman inequality for quasihyperbolic geodesics, *Math. Proc. Cambridge Philos. Soc.*, **114** (1993), 393–405.

- [21] D. A. HERRON, Quasiconformal deformations and volume growth, *Proc. London Math. Soc.* (3) **92** (2006), no. 1, 161–199.
- [22] D. A. HERRON, Uniform domains and hyperbolic distance, *J. Anal. Math.* **143** (2021), no. 1, 349–400.
- [23] M. HUANG, Y. LI, S. PONNUSAMY AND X. WANG, The quasiconformal subinvariance property of John domains in R^n and its applications, *Math. Ann.* **363** (2015), no. 1-2, 549–615.
- [24] F. JOHN, Rotation and strain, *Comm. Pure. Appl. Math.* **14** (1961), 391–413.
- [25] P. W. JONES, Quasiconformal mappings and extendability of functions in Sobolev spaces, *Acta Math.*, **147** (1981), 71–88.
- [26] P. KOSKELA AND P. LAMMI, Gehring-Hayman theorem for conformal deformations, *Comment. Math. Helv.*, **88** (2013), 185–203.
- [27] P. KOSKELA, P. LAMMI AND V. MANOJLOVIĆ, Gromov hyperbolicity and quasihyperbolic geodesics, *Ann. Sci. Éc. Norm. Supér.* (4) **47** (2014), no. 5, 975–990.
- [28] Y. LI, Neargeodesic in John domain in Banach spaces, *Internat. J. Math.*, **25** (2014), 1450041.
- [29] G. J. MARTIN, Quasiconformal and bi-Lipschitz homeomorphisms, uniform domains and the quasihyperbolic metric, *Trans. Amer. Math. Soc.* **292** (1985), no. 1, 169–191.
- [30] O. MARTIO AND J. SARVAS, Injectivity theorems in plane and space, *Ann. Acad. Sci. Fenn. Ser. A I Math.* **4** (1978), 383–401.
- [31] CH. POMMERENKE, Boundary behaviour of conformal maps. Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 299. Springer-Verlag, Berlin, 1992.
- [32] J. VÄISÄLÄ, Free quasiconformality in metric spaces. I, *Ann. Acad. Sci. Fenn. Ser. A I Math.*, **15** (1990), 355–379.
- [33] J. VÄISÄLÄ, Free quasiconformality in metric spaces. II, *Ann. Acad. Sci. Fenn. Ser. A I Math.*, **16** (1991), 255–310.
- [34] J. VÄISÄLÄ, Relatively and inner uniform domains, *Conformal Geom. Dyn.*, **2** (1998), 56–88.
- [35] J. VÄISÄLÄ, Quasihyperbolic geodesics in convex domains, *Results in Math.*, **48** (2005), 184–195.
- [36] J. VÄISÄLÄ, Hyperbolic and uniform domains in Banach spaces, *Ann. Acad. Sci. Fenn. Math.* **30** (2005), 261–302.
- [37] J. VÄISÄLÄ, Gromov hyperbolic spaces, *Expo. Math.*, **23** (2005), 187–231.

(C.-Y. Guo) RESEARCH CENTER FOR MATHEMATICS AND INTERDISCIPLINARY SCIENCES, SHANDONG UNIVERSITY, 266237, QINGDAO, P. R. CHINA, AND DEPARTMENT OF PHYSICS AND MATHEMATICS, UNIVERSITY OF EASTERN FINLAND, 80101, JOENSUU, FINLAND

Email address: changyu.guo@sdu.edu.cn

(M. Huang) DMOE-LCSM, SCHOOL OF MATHEMATICS AND STATISTICS, HUNAN NORMAL UNIVERSITY, CHANGSHA, HUNAN 410081, P. R. CHINA

Email address: mzhuang79@yahoo.com.cn

(X. Wang) MOE-LCSM, SCHOOL OF MATHEMATICS AND STATISTICS, HUNAN NORMAL UNIVERSITY, CHANGSHA, HUNAN 410081, P. R. CHINA

Email address: xtwang@hunnu.edu.cn