

ARTICLE TYPE

Three-dimensional simulations of accretion disks in pre-CE systems

Ana L. Juárez-García,^{1,2} Orsola De Marco,^{1,2} Fabio De Colle,³ Diego López-Cámara,^{3,4} Enrique Moreno Méndez,⁵ and Jesús Carrillo-Santamaría⁶¹School of Mathematical and Physical Sciences, Macquarie University, Balaclava Road, North Ryde, Sydney, NSW 2109, Australia²Astrophysics and Space Technologies Research Centre, Macquarie University, Balaclava Road, North Ryde, Sydney, NSW 2109, Australia³Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, A. P. 70-543 04510 D. F. Mexico⁴Investigador por México, CONAHCYT – Universidad Nacional Autónoma de México, Instituto de Astronomía, AP 70-264, CDMX 04510, México⁵Facultad de Ciencias, Universidad Nacional Autónoma de México, A. P. 70-543 04510 D. F. Mexico⁶Instituto de Astronomía, Universidad Nacional Autónoma de México, A. P. 70-264 04510 D. F. México**Author for correspondence:** Ana Juárez-García, Email: analourdes.jurezgarca@hdr.mq.edu.au.

Abstract

Before a binary system enters into a common envelope (CE) phase, accretion from the primary star onto the companion star through Roche Lobe overflow (RLOF) will lead to the formation of an accretion disk, which may generate jets. Accretion before and during the CE may alter the outcome of the interaction. Previous studies have considered different aspects of this physical mechanism. Here we study the properties of an accretion disk formed via 3D hydrodynamic simulations of the RLOF mass transfer between a $7 M_{\odot}$, red supergiant star and a $1.4 M_{\odot}$, neutron star companion. We simulate only the volume around the companion for improved resolution. We use a 1D implicit MESA simulation of the evolution of the system during 30 000 years between the on-set of the RLOF and the CE to guide the binary parameters and the mass-transfer rate, while we simulate only 21 years of the last part of the RLOF in 3D using an ideal gas isothermal equation of state. We expect that a pre-CE disk under these parameters will have a mass of $\sim 5 \times 10^{-3} M_{\odot}$ and a radius of $\sim 40 R_{\odot}$ with a scale height of $\sim 5 R_{\odot}$. The temperature profile of the disk is shallower than that predicted by the formalism of Shakura and Sunyaev, but more reasonable cooling physics would need to be included. We stress test these results with respect to a number of physical and numerical parameters, as well as simulation choices, and we expect them to be reasonable within a factor of a few for the mass and 15% for the radius. We also contextualize our results within those presented in the literature, in particular with respect to the dimensionality of simulations and the adiabatic index. We consider what properties of magnetic fields and jets may be supported by our disk and discuss prospects for future work.

Keywords: accretion, accretion discs – binaries (including multiple): close – hydrodynamics – methods: numerical – stars: evolution

1. Introduction

Massive stars ($M \gtrsim 8 M_{\odot}$) are almost always in binary and multiple systems with $\sim 70\%$ of them involved in various forms of interaction, including tidal interactions and mass transfer, leading eventually to close binaries and mergers (Moe & Di Stefano, 2017). These interacting binary systems can give rise to high-energy phenomena, such as cataclysm variables (e.g., Warner, 1995), type Ia supernovae (e.g., Iben & Tutukov, 1984; Chevalier, 2012), short and long gamma-ray bursts (e.g., Fryer & Woosley, 1998; Brown et al., 2007; Ramirez-Ruiz & Lee, 2009), and gravitational wave emission (e.g., Abbott et al., 2016).

For a certain range of binary and stellar parameters, the massive binary becomes a high-mass X-ray binary, where a red giant or red supergiant (RGS) feeds mass through the inner Lagrangian point, L_1 , to a neutron star (or sometimes a black hole) companion in a phase of wind accretion and, possibly, Roche lobe overflow (RLOF). These systems form an accretion disk around the compact companion, which is X-ray bright and may develop jets. If the mass ratio is large, as is the case if the compact object is a neutron star, mass transfer reduces the orbital separation. In addition, the RGSs expand upon loss of mass, further accelerating the mass transfer. For neutron star – RGS systems, it is likely that unstable mass transfer and a common envelope (CE) phase may result (Ivanova et al., 2013;

Tauris et al., 2017). The outcome of this phase depends on whether or not the CE can be ejected before the neutron star merges with the helium core of the RSG.

Most observed X-ray binaries are undergoing a long-lived phase of stable wind accretion with timescales that depend on the parameters of the system. Some systems may be undergoing RLOF, in which case the evolution timescales are likely much shorter with a possibility of unstable mass transfer and CE (e.g., high mass X-ray binaries may remain in the RLOF state for only 10 000 yr; Savonije 1977, see also Tauris et al. 2017). Although, presumably, X-ray binaries in the stable wind accretion phase are more frequently observed (e.g., Cygnus X-1), it is possible that some may be caught in the faster phase of unstable mass transfer. Dickson (2024) presented a model of X-ray binary M 33 X-7, that they believe to be in an unstable mass transfer phase based on a measurement by Ramachandran et al. (2022) of the donor substantially overfilling its Roche lobe.

One open question in the study of massive CE interactions is the impact of the energetic feedback due to accretion of envelope gas onto the compact companion, before and during the in-spiral in the CE. For neutron stars and black hole companions, this could be critically important for the outcome of the interaction. Shiber et al. (2019) simulated an *ad hoc* jet, emanating from a companion in a low mass CE interaction to

conclude that the jet would aid in unbinding envelope mass with consequences for the binary separation of the post-CE binary. On the other hand López-Cámara *et al.* (2022), using a self-regulated jet, powered by a fraction of the mass accretion rate that reached the inner boundary, showed that the jet would likely be quenched, even if it existed before the companion entered the CE.

A disk that forms during wind accretion and RLOF could survive inside the CE, be destroyed, or be destroyed and reform, as a result of accretion of envelope material on the secondary star (MacLeod & Ramirez-Ruiz, 2015; MacLeod *et al.*, 2017; Moreno Méndez *et al.*, 2017; Chamandy *et al.*, 2018; López-Cámara *et al.*, 2019; Shiber *et al.*, 2019; López-Cámara *et al.*, 2020; Moreno Méndez, 2022). So far previous studies have found that the formation of the accretion disk inside a CE depends on the thermal properties of the envelope (adiabatic index of $\lesssim 1.2$). Murguía-Berthier *et al.* (2017) pointed out that this phase is only a transitory phase, due to the lack of stellar regions (zones of partial ionization where γ is small enough) where the envelope is compressible enough to form a disk.

In this work, we study the formation of the disk around a $1.4 M_{\odot}$ neutron star, caused by RLOF mass transfer from a $7 M_{\odot}$ RGS, undergoing unstable mass transfer. We attempt to gain a quantitative idea of the parameters of an accretion disk to, eventually, determine its fate inside the CE. This work is also intended to contribute to the literature by studying accretion disks in 3D hydrodynamics. The goal is to determine when and how accretion disks form in response to mass accretion through L_1 and as a function of a number of physical and numerical parameters to set this study in the broader context of disk formation (e.g., Makita *et al.*, 2000).

This paper is structured as follows. In Section 2 we outline the overall methodology with Sections 2.1 and 2.2 presenting the governing equations and simulation parameters. In Section 3 we give details of the formation and evolution of the disk and, in Section 3.1 the disk parameters. In Section 4 we discuss the sensitivity of our results to some physical and numerical parameters, while in Section 5 we present our conclusions.

2. Methods

To study the mass transfer phase through the L_1 point, we consider a binary system consisting a $7 M_{\odot}$ red supergiant as the donor star and a $1.4 M_{\odot}$ neutron star as the companion star (see Figure 1 for a cartoon of the setup). We model the evolution of this binary system, between the RLOF phase and the CE phase, using the 1D implicit code MESA (Modules for Experiments in Stellar Astrophysics; version r21.12.1; Paxton *et al.* 2011, 2013, 2015, 2018, 2019).

We then use the 3D hydrodynamic numerical code MEZCAL (De Colle *et al.*, 2012) to simulate the formation of the accretion disk around the companion star using the mass transfer rate given by the MESA model as boundary condition to inject material into the computational domain. The 3D computational domain is represented by a dotted line in Figure 1.

To only simulate the region around the accreting star, the 3D simulation is performed in a co-rotating system of reference, centered on the companion which is represented by a point mass particle encircled by an inflow boundary of radius R_{in} . Mass is injected through a nozzle that represents the L_1 point, located at the center of the left boundary face. Below we motivate the method and the setup and explain the specific assumptions.




Figure 1. Setup cartoon. The donor star is a red super giant of $7 M_{\odot}$ with a radius of $139 R_{\odot}$, orbiting a compact object of $1.41 M_{\odot}$, with a separation of $270 R_{\odot}$. The dotted line (centered on the compact object) represents the computational domain in our 3D simulations.

2.1 The hydrodynamic code and its governing equations

To study the formation and stability of accretion disks during the RLOF phase, we run a series of 3D numerical simulations employing the adaptive mesh refinement code MEZCAL (De Colle *et al.*, 2012). The code integrates the hydrodynamic equations in a rotating frame. Self gravity is not included in these simulations.

We solve the three-dimensional Euler equations for an inviscid gas (see, e.g., Makita *et al.* 2000), that is,

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{v}) = 0, \quad (1)$$

$$\frac{\partial (\rho \mathbf{v})}{\partial t} + \nabla \cdot (\rho \mathbf{v} \mathbf{v} + P \mathbf{I}) = -\rho \mathbf{f}, \quad (2)$$

$$\frac{\partial e}{\partial t} + \nabla \cdot [(e + p) \mathbf{v}] = -\rho \mathbf{v} \cdot \mathbf{f}, \quad (3)$$

representing the evolution of mass density, ρ , gas momentum, $\rho \mathbf{v}$, and gas energy density, e . The variable P represents the pressure, \mathbf{I} is the identity tensor, \mathbf{v} is the velocity vector, and \mathbf{f} is the specific force vector. The specific force vector $\mathbf{f} = (f_x, f_y, f_z)$ represents the gravity forces and fictitious forces associated with the rotating frame.

The binary system consists of a donor star of mass M_1 , and an accreting star of mass M_2 , with a mass ratio defined as $q = M_1/M_2$ during their mass transfer phase. Our simulations are conducted in Cartesian coordinates in three dimensions with different levels of resolution. The binary system has an orbital separation a , with an orbital frequency $\Omega = (G(M_1 + M_2)/a^3)^{1/2}$.

Then, the orbital period is given by $P = 2\pi/\Omega$. The origin of the rotating frame is the accreting star; the donor star is positioned on the left of the accretor (donor position is $(-a, 0, 0)$). We perform the simulation in dimensionless units using a as the length scale and $a\Omega/2\pi$ as the velocity scale. The time unit is the initial period of the binary. This means that quantities can be rescaled based on these units to physical units. In Table 1 we list the scaling factors between code and physical units.

Table 1. Description of units, rescaling factors to physical systems, in terms of the binary separation (a), orbital frequency (Ω), and total mass ($M_{\text{tot}} = M_{\text{don}} + M_{\text{acc}}$, where M_{don} is the mass of the donor star and M_{acc} is the mass of the companion star).

	Code unit	Value in cgs
Length	a	1.85×10^{13} cm
Velocity	$a\Omega/2\pi$	7.75×10^6 cm s $^{-1}$
Mass	M_{tot}	1.67×10^{34} g
Time	$2\pi/\Omega$	0.48 yr
Mass transfer rate	$M_{\text{tot}}\Omega/2\pi$	6.97×10^{27} g s $^{-1}$
Density	M_{tot}/a^3	2.62×10^{-6} g cm $^{-3}$
Pressure	$M_{\text{tot}}\Omega^2/4\pi^2a$	3.89×10^8 dyne cm $^{-2}$

With these assumptions, we can now write out the force vector in dimensionless form at each point in the computational domain, remembering that there is no self-gravity, but that we are operating in the rotating frame:

$$f_x = -2v_y - \left(x + \frac{q}{1+q} \right) + \frac{1/(1+q)}{|\mathbf{r}_1|^3} (x-a) + \frac{q/(1+q)}{|\mathbf{r}_2|^3} x, \quad (4)$$

$$f_y = 2v_x - \gamma + \frac{1/(1+q)}{|\mathbf{r}_1|^3} y + \frac{q/(1+q)}{|\mathbf{r}_2|^3} y, \quad (5)$$

$$f_z = \frac{1/(1+q)}{|\mathbf{r}_1|^3} z + \frac{q/(1+q)}{|\mathbf{r}_2|^3} z, \quad (6)$$

where \mathbf{r}_1 and \mathbf{r}_2 are the distances from the point considered $\mathbf{r} = (x, y, z)$ to the center of each star. In Equations (4) and (5), on the right-hand side, the first term represents the Coriolis force and the second term represents the centrifugal force. The last two terms in Equations (4) and (5), as well as the terms in Equation (6), represent the gravity force of each star.

2.2 Initial conditions

2.2.1 Calculation of the mass transfer rate through L_1 using a 1D implicit code

To determine the mass transfer rate in a binary system between a massive red supergiant donor star and a compact accretor, we use MESA to simulate the evolution of a binary system comprising a $7 M_{\odot}$, solar metallicity main sequence star with a radius of $57 R_{\odot}$ and a $1.4 M_{\odot}$ point mass companion, initially located at an orbital separation of $a = 270 R_{\odot}$. The Roche lobe radius of the primary star (R_{L_1}) is calculated according to the prescription by Eggleton (1983):

$$\frac{R_{L_1}}{a} = \frac{0.49q^{2/3}}{0.6q^{2/3} + \ln(1+q^{1/3})}, \quad (7)$$

where a is the orbital separation, $q = M_1/M_2$ is the mass ratio between the donor star (M_1) and the accreting star (M_2).

We let the MESA simulation run for $\sim 4.0 \times 10^7$ yrs until the primary expands to fill its Roche lobe and starts to transfer mass to the companion. At this time, called time zero in Figure 2, the red supergiant has a helium-burning core surrounded by a shell that experiences hydrogen burning (primarily through the CNO cycle), along with a massive convective hydrogen envelope. The orbital separation is $270.4 R_{\odot}$, the red supergiant has a mass of $6.98 M_{\odot}$, a radius of $134 R_{\odot}$, an effective temperature of 3981 K, and a luminosity of $3645 L_{\odot}$. The companion star has a mass of $1.4 M_{\odot}$, and the mass transfer rate at time zero is $1.4 \times 10^{-8} M_{\odot} \text{ yr}^{-1}$.




Figure 2. Temporal evolution of the mass transfer rate from the primary to the secondary star in our long-term, MESA simulation of the binary system. The initial masses of the stars are $7 M_{\odot}$ and $1.4 M_{\odot}$; they have an initial orbital separation of $270 R_{\odot}$ and an orbital period of 177 days. We indicate with the orange box the mass transfer evolution that we are simulating in 3D.

We then continue the MESA simulation for an additional ~ 30000 yr, during which the mass transfer rate increases to a maximum value of $9.7 \times 10^{-2} M_{\odot} \text{ yr}^{-1}$ (as we can see in Figure 2). After 30000 yrs the mass of the red supergiant is $6.88 M_{\odot}$, its radius is $143 R_{\odot}$ and its effective temperature is ~ 5012 K, while the companion star mass has increased to $1.49 M_{\odot}$, having accreted some mass. The orbital separation is $243 R_{\odot}$. After this point we assume that the mass transfer leads to a CE in a short timescale. See Table 2 for a summary of the initial and final parameters of the simulation.

Table 2. Initial (start of RLOF) and final (the time of CE) binary parameters for 30000 yr of Roche lobe mass transfer, modeled with the MESA code.

Quantity	Initial value	Final value
Donor star	$7.0 M_{\odot}$	$6.88 M_{\odot}$
Accretor star	$1.4 M_{\odot}$	$1.49 M_{\odot}$
Orbital separation	$270 R_{\odot}$	$243 R_{\odot}$
Mass transfer rate	$1.6 \times 10^{-9} M_{\odot} \text{ yr}^{-1}$	$9.7 \times 10^{-2} M_{\odot} \text{ yr}^{-1}$

The 3D simulation described next (Section 2.2.2) instead spans only 21 years. This period of time is taken between 29500 yr and 29521 yr of the 30000 yr stretch of the 1D simulation. Hence, the mass transfer rate in the 3D simulation, is prescribed from the 1D simulation to be between

$2.3 \times 10^{-4} M_{\odot} \text{ yr}^{-1}$ and $9.7 \times 10^{-2} M_{\odot} \text{ yr}^{-1}$. In Section 4.1 we will explore the impact that this mass-transfer rate choice has on the the accretion disk parameters.

2.2.2 Calculation of disk formation using a 3D explicit code

In the 3D simulations, a $1.4 M_{\odot}$ point mass particle represents the neutron star companion, at the centre of the domain. We assume that the donor star, with a mass of $7.0 M_{\odot}$, is located outside the computational domain at a position $(-a, 0, 0)$ from the companion star, where $a = 266 R_{\odot}$ is the binary separation at the start of the hydrodynamic simulation. In this way, the position of L_1 is at $83.1 R_{\odot}$ (5.79×10^{12} cm) from the companion star. The gas is injected through the L_1 point, represented in the simulations by a small rectangular boundary with a variable size given by the cross section of the mass transfer stream, as explained in the following. The setup scheme is shown in Figure 1.

The point-mass companion sits inside a spherical inflow boundary of radius $R_{\text{in}} = 1.3 R_{\odot}$ (9.3×10^{10} cm). The density of gas within this inner boundary is $\rho_{\text{in}} = 1.0 \times 10^{-20} \text{ g cm}^{-3}$ and the pressure is $1.0 \times 10^{-8} \text{ dyne cm}^{-2}$. Grid cells cannot be empty, so we set a low background density $\rho_{\text{bg}} = 2.6 \times 10^{-22} \text{ g cm}^{-3}$, temperature $T_{\text{bg}} = 1.0 \times 10^5 \text{ K}$, and pressure $P_{\text{bg}} = 8.9 \times 10^{-11} \text{ dyne cm}^{-2}$. The sensitivity of our simulations to different values of the background conditions is tested in Section 4.3. The gas adiabatic index is fixed to $\gamma = 1.1$ during the whole simulation, based on Makita *et al.* (2000), MacLeod & Ramirez-Ruiz (2015) and Murguía-Berthier *et al.* (2017). See Section 4.5 for further discussion about the adiabatic index in simulations of accretion disks.

Following the Jackson *et al.* (2017) prescription for optically thin mass transfer, we define an elliptical area centered on L_1 , where most of the material escapes L_1 , referred from now on as the “nozzle”. The nozzle area is defined by $S_{L_1} = \pi \Delta z \Delta y$, where the two dimensions are related to the pressure scale height in the y - and z -direction, and vary as:

$$\Delta y = \frac{\sqrt{2} c_T}{\Omega \sqrt{A-1}}, \quad \Delta z = \frac{\sqrt{2} c_T}{\Omega \sqrt{A}}, \quad (8)$$

where c_T is the isothermal sound speed, $c_T = \sqrt{\gamma * k_B T / \mu}$, Ω is the orbital frequency and A is a dimensionless coefficient that depends on the mass ratio, q (equivalently M_2/M_1 or M_1/M_2 in this equation), defined as

$$A(q) = 4 + \frac{4.16}{-0.96 + q^{1/3} + q^{-1/3}}. \quad (9)$$

We set the size of the nozzle on the x -axis to be two cells thick at the coarsest level of refinement, or $\Delta x = 1.30 R_{\odot}$ (9.05×10^{10} cm). The initial mass injection rate is $\dot{M}_{\text{ini}} = 2.31 \times 10^{-4} M_{\odot} \text{ yr}^{-1}$, has a subsonic velocity only in the x direction with value $v_{L_1} = 77.7 \times 10^3 \text{ cm s}^{-1}$ (Lubow & Shu, 1975; Jackson *et al.*, 2017; Cehula & Pejcha, 2023). The mass injection rate (\dot{M}) is interpolated at each time step (Δt), using the values given by the MESA simulation (see Section 2.2.1). Using the interpolated values, we calculate the nozzle volume

(V_{L_1}), nozzle density as $\rho_{L_1} = \dot{M} \Delta t / V_{L_1}$, and the pressure of the nozzle as $P_{L_1} = \rho_{L_1} c_T^2$, using the effective temperature of the donor star to calculate c_T .

We start the hydrodynamic simulation at $t_{\text{ini}} = 29\,500 \text{ yr}$ after the start of the mass transfer in the MESA simulation and let the hydrodynamic simulation run for 21 yr (or for 44 orbital periods), with a final mass transfer rate of $\dot{M}_{\text{fin}} = 9.7 \times 10^{-2} M_{\odot} \text{ yr}^{-1}$. We also performed eight different simulations, altering different physical and numerical parameters (see Table 3). We will justify the need for these additional simulations in Section 4.

The size of the computational domain depends on the distance between the first Lagrange point and the accreting star, initially at $-d_{L_1} = 83.1 R_{\odot}$ (5.79×10^{12} cm). Hence, the computational box has dimensions $d_{L_1} \leq x \leq 1.25d_{L_1}$, $-1.5d_{L_1} \leq y \leq 1.5d_{L_1}$ and $-0.5d_{L_1} \leq z \leq 0.5d_{L_1}$ with outflow boundary conditions at each of the six faces (except for the location of the nozzle that is technically and inflow boundary). The grid was modified to have better resolution surrounding the companion star and around the nozzle, avoiding numerical problems. We employ (72, 96, 32) cells at the coarsest refinement level, with three levels of refinement (see Section 4.4 for convergence tests), corresponding to a maximum resolution of $\Delta x = \Delta y = \Delta z = 0.65 R_{\odot}$ (4.53×10^{10} cm).

3. Results: Formation and evolution of the disk

In what follows, we give details of the simulation whose input parameters are explained in Section 2.2. We refer to this simulation as the “Reference simulation” (sim-0 in Table 3). Later, we test the results against changes in the input parameters.




Figure 3. Volumetric density rendering showing the accretion disk at four times ($t = 0.3 \text{ yr}$, 1.5 yr , 10.5 yr , and 20.9 yr). We can appreciate the 3D structure of the accretion disk, surrounding the companion star (represented by the white area in the middle of the accretion disk).

In Figure 3 we show a volumetric density rendering that shows the accretion disk forming over 21 yrs. At the start of the simulation, the injected material with velocity in the

Table 3. Simulations' summary: inputs are varied to understand the resilience of the accretion disk parameters to numerical and physical input parameter changes. We vary the initial mass transfer rate (sim-dot-#), injection velocity through L_1 (sim-vel-#), and background temperature and density (sim-bgT-#). We compare each simulation to the reference simulation (sim-0). The majority of the simulations were carried out for 21 yr and start with a donor star of $7 M_\odot$, a companion star of $1.4 M_\odot$, an orbital separation of $266.34 R_\odot$, and Ω of $4.18 \times 10^{-7} \text{ s}^{-1}$.

Model	\dot{M}_{ini} ($M_\odot \text{ yr}^{-1}$)	\dot{M}_{fin} ($M_\odot \text{ yr}^{-1}$)	ρ_{bg} (g cm^{-3})	P_{bg} (dyne cm^{-2})	T_{bg} (K)	v_{L_1} (cm s^{-1})
sim-0	2.3×10^{-4}	9.7×10^{-2}	2.6×10^{-22}	8.9×10^{-11}	1.0×10^5	7.7×10^4
sim-mdot-1	1.4×10^{-8}	1.4×10^{-8}	2.6×10^{-22}	8.9×10^{-11}	1.0×10^5	7.7×10^4
sim-mdot-2	1.1×10^{-5}	6.4×10^{-5}	2.6×10^{-22}	8.9×10^{-11}	1.0×10^5	7.7×10^4
sim-mdot-3	3.7×10^{-3}	9.7×10^{-2}	2.6×10^{-22}	8.9×10^{-11}	1.0×10^5	7.7×10^4
sim-vel-1	2.3×10^{-4}	9.7×10^{-2}	2.6×10^{-22}	8.9×10^{-11}	1.0×10^5	5.7×10^5
sim-vel-2	2.3×10^{-4}	9.7×10^{-2}	2.6×10^{-22}	8.9×10^{-11}	1.0×10^5	6.2×10^5
sim-bgT-1	2.3×10^{-4}	9.7×10^{-2}	2.6×10^{-22}	8.9×10^{-12}	1.0×10^4	7.7×10^4
sim-bgT-2	2.3×10^{-4}	9.7×10^{-2}	2.6×10^{-22}	8.9×10^{-10}	1.0×10^6	7.7×10^4

x -direction (into the computational domain) effectively free falls towards the companion star under its own gravity, but due to its angular momentum, it sweeps around forming a disk. The injected material moves towards the companion, taking ≈ 0.12 yr to reach the internal boundary, and is deflected around the central potential toward the lower density medium. The interaction between the deflected material and the injection stream creates the accretion disk's base structure, seen after 0.3 yr (upper left panel of Figure 3). One year later, the accumulated deflected material forms two high density spiral arms on the orbital plane (see upper right panel).

After 10.5 yr the shock between the spiral arms has created a high density structure with a disk-like shape, surrounding the companion star (see bottom left panel). The material in the disk is orbiting around the companion star, forming bow shocks with the injected material, showing a higher density on the left side of the computational domain than on the right side, consistent with the results of Makita et al. (2000)^a. By the end of the simulation at 20.9 yr (see bottom right panel of Figure 3) the accretion disk has maintained its approximate structure for 10.5 yr, albeit while growing somewhat in radius and scale height, due to the accumulation of mass.

In the next section, we quantify what we have just described qualitatively.

3.1 Accretion disk parameters

In Figure 4, we present orbital and perpendicular density slices of the accretion disk at the same four times shown in Figure 3. Along the orbital plane we see the formation of spiral arms, the injected mass leaves the nozzle and is deflected towards the companion star, due to its gravitational pull, Coriolis force, and centrifugal force. With an increase in the rate of mass transfer, a high density structure can be seen in the last panel of the top row of Figure 4.

The edge-on panels in Figure 4, bottom row, show the formation of the aforementioned bow shocks, due to the interac-

tion between the injection stream and the accretion disk. Also, the edge-on view shows the presence of small, low density outflows along the polar axes, and may suggest the formation of hydrodynamically collimated jets in the future; however, our simulation ends too early to follow their development. These outflows are not visible in the rendered plots as a result of the choice of the color bar limits.

In Figure 5 we plot density profiles along the z -axis of the accretion disk, measured at $26.7 R_\odot$ (1.86×10^{12} cm) from the companion star along the positive and negative side of the x - and y -axes, at $t = 10.5$ yr (blue line), and $t = 21$ yr (red line). The upper panels of Figure 5 show the density profiles measured on the positive and negative sides of the x -axis, panels (a) and (b), respectively. Panels (c) and (d) show the density profiles measured on the positive and negative sides of the y -axis, respectively (these labels are also marked in Figure 4).

Panel (b) shows the material to be more extended in the z -direction because the incoming material from the nozzle is constantly interacting with the disk material. Meanwhile, panel (a) presents a clear description of the disk thickness, with a sharply decreasing density above and below the midplane, just as we can see in the density maps (Figure 4). At 10.5 yr the disk has an average thickness of $15.6 R_\odot$ (1.09×10^{12} cm). By the end of the simulation (21 yr) the disk has reached an average thickness of $20.4 R_\odot$ (1.42×10^{12} cm). We also measure the scale height of the disk at 10.5 yr to be $H = 6.7 R_\odot$ (0.47×10^{12} cm), while at the end of the simulation (21 yr) the disk scale height has slightly decreased to $4.9 R_\odot$ (0.34×10^{12} cm).

Panels (c) and (d) in Figure 5 show that the thickness of the disk perpendicular to the line that joins the two stars is thicker and less defined. Panel (d), includes the interaction of the deflected material that has left the nozzle and encounters the accretion disk, just as we can appreciate in the density maps of the orbital plane. The shape of the vertical density profiles at each of the four locations does not change significantly over 10 years; only the density increases over time due to the constant injection of material.

We measured the mass inside one hundred radial points (Figure 6), measured from the inner boundary (encircling the companion star) to the edge of the computational domain. We

^aSince much of this work is based on the work of Makita et al. (2000) we will continue the comparison throughout this paper and bring it to bear in a discussion in Section 4.5.




Figure 4. Density slices of the accretion disk on the orbital plane (top row) and perpendicular plane (bottom row) of sim-0 at times $t = 0.3$ yr, 1.5 yr, 10.5 yr, and 21.0 yr (from left to right). The letters in the third and fourth upper panels represent the positions used to calculate the density profiles (see text). The axes are in dimensionless code units ($1 \text{ code unit} = 1.85 \times 10^{13} \text{ cm}$).




Figure 5. Density profile of the accretion disk versus height at a radius of $26.7 R_{\odot}$ ($1.86 \times 10^{12} \text{ cm}$). Two times are shown: $t = 10.5$ yr (blue line) and 21 yr (red line). The readings were taken at four symmetric points around the companion star along the x - and y -axis (panel (a): $+x$ -axis, (b): $-x$ -axis, (c): $+y$ -axis, (d): $-y$ -axis).

repeat this calculation for the same times as in Figure 5. At $t = 0.3$ yr (blue line panel) the injected material has just reached the internal boundary, but the disk has not formed yet. Once the disk has formed, at 1.5 yr (green line panel in Figure 6), the enclosed mass increases with radius until it reaches the edge of the disk. At this point, we see a flattening of the slope of the enclosed mass. The nozzle is located at a radius of $74.4 R_{\odot}$ ($5.18 \times 10^{12} \text{ cm}$) from the inner boundary, and it manifests itself in the steepening of the gradient at the largest radius, particularly evident at the last time (bottom right panel in Figure 6). We determine the radius of the disk by locating the first inflection point (black point in all panels). In Table 4 we summarize the disk characteristics discussed so far.

Table 4. Disk properties at different moments in time in sim-0.

Time (yrs)	$R_{\text{disk}} (R_{\odot})$	$M_{\text{disk}} (M_{\odot})$	$H_{\text{disk}} (R_{\odot})$
0.3	5.3	2.33×10^{-6}	—
1.5	41	4.29×10^{-5}	—
10.5	41	5.14×10^{-4}	6.7
21.0	39	5.46×10^{-3}	4.9

Using the inner inflow boundary, we measure the accretion rate onto the companion as a function of time (Figure 7). The accreted mass does not contribute to the mass of the companion. We identify a cubic volume, $2R_{\text{in}}$ on one side, which contains the inflow boundary sphere. The total accretion rate is measured by calculating the mass flux through the cubic boundary, by taking the projection of the velocity of each




Figure 6. Cumulative mass as a function of radius from the companion star, at different times: $t = 0.3$ yr (top left panel, orange line), $t = 1.5$ yr (top right panel, maroon line), $t = 10.5$ yr (bottom left panel, purple line), $t = 20.9$ yr (bottom right panel, blue line). The black symbol in each panel indicates the adopted radius of the disk.

cell liming the boundary onto the normal direction to the boundary (solid line in Figure 7), or

$$\dot{M} = \sum_{i=1}^N \rho_i A_i \vec{v}_i \cdot \hat{n}, \quad (10)$$

where ρ_i is the density of the cell, \vec{v}_i is the velocity vector of the cell, and A_i represents the area of the cell face through which the material will cross in the next time step, and \hat{n} is the vector perpendicular to that face directed into the cubic boundary. By the end of the simulation, at 21 yrs, the mass accretion rate is $4.1 \times 10^{-3} M_{\odot} \text{ yr}^{-1}$. We also carry out the same measurement, but where we project both the velocity vector of each cell and the area of the face that the gas crosses, along the radial direction to the companion – this method results in smaller values for the accretion rate by a factor of 0.6 (dashed line in Figure 7).

We compare the numerical accretion rate obtained above with the mass accretion rate predicted theoretically by the viscosity and density of the disk, defined as:

$$\dot{M} = 3\pi \Sigma \alpha c_s H_{\text{disk}}, \quad (11)$$

where Σ is the surface density of the disk (we assume $\Sigma = M_{\text{disk}}/R_{\text{disk}}^2$), and α is the α -viscosity formalism of Shakura & Sunyaev (1973). For thin disks, we assume $\alpha = 0.1$. The variable c_s is the speed of sound at R_{disk} and H_{disk} is the scale height of the disk. Using the aforementioned results at the end of the simulation, we predict an accretion rate of $3.7 \times 10^{-3} M_{\odot} \text{ yr}^{-1}$, consistent with the accretion rate measured at the end of the simulation.

The material in the nozzle has density (ρ_{L_1}) and pressure (P_{L_1}) defined in terms of the mass injection rate, the volume of the nozzle, and the velocity, as defined in Section 2.2. Assuming an ideal gas, we show, in Figure 8, temperature slices in the orbital and perpendicular planes at the end of the simulation (21 yr). The material inside the nozzle has an average




Figure 7. Mass accretion rate onto the companion (traversing the inner, inflow boundary) as a function of time. The solid line represents the mass accretion rate assuming the inner boundary is a cube with edge length $2R_{\text{in}}$ and the dashed line shows the mass accretion rate calculated projecting the area and velocity of the cell over the spherical inner boundary.

temperature of ~ 5000 K, once the material leaves the nozzle its temperature drops to ~ 3000 K, due to the pressure difference between the nozzle and the material just outside the nozzle. Inside the inflow boundary around the accretor, the high temperature ($\sim 10^7$ K) is due to the imposed low density. We observe a gradient of temperature decreasing radially away from the inner boundary, we can also observe on the perpendicular plane (x - z plane) the interaction between the injected mass and the mass that circles around the companion.




Figure 8. Temperature slices for sim-0 in the orbital (top panel) and perpendicular planes (bottom panel) of the disk at $t = 21$ yr.

In Figure 9, we plot a temperature profile along the positive x -axis at 21 yr, with the internal boundary shown as a black

vertical line and the radius of the disk as a dashed gray line. In the mid-plane along the positive x-direction, the cells closer to the inner boundary have a temperature of 1.1×10^6 K, their location corresponds with the region where the pressure gradients is the highest; the cells located at the edge of the disk have lower temperatures of $\sim 60\,000$ K.




Figure 9. Temperature profile for sim-0 in the mid plane along the positive x-axis at $t = 21$ yr. The disk radius is indicated with a vertical gray dashed line. The inner boundary radius with a vertical black line. The solution for steady disks is indicated by the green line ($T \propto r^{-3/4}$) and the best fit ($\alpha = 1.12 \pm 0.02$) is indicated by the red line and red shaded area (corresponding to an error of 15%).

Following the prescription of [Shakura & Sunyaev \(1973\)](#) for steady disks, the effective temperature profile should follow $T \propto r^{-3/4}$ (green line in Figure 9). This function fits our data with an average percentage error of 38%. The best fit is obtained using $T \propto r^{-1.12 \pm 0.02}$ (red line and shaded area in Figure 9) with an average percentage error of 15%. [Mineshige et al. \(1994\)](#) stated that a value of $\alpha < -3/4$ characterizes X-ray binary systems in the flaring branch on the X-ray hardness-intensity diagram.




Figure 10. Slices of Mach number in the orbital plane (top panels) and perpendicular plane (bottom panels) of the disk at $t = 10.5$ yr (left panels) and $t = 21.0$ yr (right panels) for sim-0.

The material is injected through the nozzle with a subsonic velocity of 7.7×10^4 cm s^{-1} in the x direction (note that the sound speed value near the nozzle ranges between 6 and 8 km s^{-1}). In the upper panels of Figure 10, we show the Mach number at 10.5 yr and at the end of the simulation (21 yr) with slices in the orbital and perpendicular planes. The nozzle is on the left of the domain, seen as a short vertical bar with subsonic velocity. Just outside the nozzle, the velocity becomes highly supersonic ($\mathcal{M} = 27$ at 21 yr). When the injected material approaches the centre of the domain, it circles it and collides with the material that was already in orbit around the companion star, slowing down.

In Figure 11, we plot the gas velocity profile in the mid-plane along the positive and negative x-axes at $t = 10.5$ yr, normalized to the Keplerian velocity ($v_k = \sqrt{GM/r}$). The inner boundary radius, R_{in} , is marked with a black line, while a gray line indicates the radius of the disk, R_{disk} . The cell velocities around the inner boundary are Keplerian on average, though there is quite a bit of scatter in individual cells. Cells at the edge of the disk move with velocities approximately 80% - 90% of the Keplerian value. This is likely due to the pressure support of the gas in the disk.




Figure 11. Velocity profile in the mid-plane along the positive and negative axes at 10.5 yr. The velocity is normalized to the Keplerian velocity (v_k). The inner boundary is marked by the black vertical solid line and the radius of the disk by the vertical gray dashed line.

We further study the disk by examining the locus of the gas cells in the specific energy vs. specific angular momentum plane ([Hayashi et al., 2021](#), Figure 12^b). We calculate the specific orbital energy as

$$e_{orb} = \frac{1}{2}v^2 - \frac{GM}{r}, \quad (12)$$

where v is the magnitude of the velocity in the center of the cell, r is the position of each cell with respect to the companion star, and M is the mass of the companion star. The magnitude of the specific angular momentum for each cell is:

$$j_{orb} = r \cdot v. \quad (13)$$

^bThe time evolution of the specific angular momentum versus energy plot is available at <https://drive.google.com/file/d/10Kh7uXEJaRr8zY5GNmyQr24Vy9wC--G-/view?usp=sharing>




Figure 12. Specific angular momentum versus specific orbital energy for every cell in the computational domain once the disk has formed (sim-0). The integration time corresponds to $t = 10.5$ yrs. The density color table is the same as in Figure 4. The black line is the solution for a circular orbit. The black dashed line and the black dash-dotted line represent orbits with higher eccentricity ($e = 0.9$ and 0.9999 , respectively). The time evolution of this plot is available at <https://drive.google.com/file/d/10Kh7uXEJaRr8zY5GNmyQr24Vy9wC--G-/view?usp=sharing>.

Each cell is color-coded by density (with the same color bar as in Figure 4), where the cells that compose the high density accretion disk are marked in red, and dark blue points represent the low density medium cells. The black line in Figure 12 represents the analytical solution of a particle moving on a circular Keplerian orbit, while the black dashed line and the back dash-dotted line indicate orbits with eccentricity 0.9 and ~ 1 , respectively.

The majority of the high-density cells have negative orbital energy, which means that the material is bound to the companion star. On the other hand, the insert plot, showing a zoom-in near the origin, shows that some of the material has positive orbital energy; hence it is unbound. Some of the gas dropped by the nozzle onto the companion star is not initially bound to the system but slows down upon colliding with gas ahead of it. The cells in the high-density region are dispersed between the black line and the gray dashed line, which means that they move in closed orbits with eccentricities less than 0.9.

We finally attempt to determine whether the mass accretion rate is consistent with a disk viscosity provided by a plausible valued magnetic field. We use the relationship of Wardle (2007), whereby:

$$B \simeq 0.2 \dot{M}_{-7}^{1/2} r_{\text{au}}^{-5/4} \left(\frac{M}{M_{\odot}} \right)^{1/4} G, \quad (14)$$

where \dot{M}_{-7} is the mass accretion rate in units of $10^{-7} M_{\odot} \text{ yr}^{-1}$, r_{au} is the radius in au and M is the mass of the accretor. Using a mass accretion rate of $4 \times 10^{-4} M_{\odot} \text{ yr}^{-1}$, an inner boundary of the simulation of $1.3 R_{\odot}$ and a central mass of $1.4 M_{\odot}$, we obtain a magnetic field at the inner boundary of ~ 8000 G, which at the surface of a 10 km-radius neutron star is 6×10^{13} G (i.e., a magnetar-scale magnetic field) following magnetic flux conservation for a dipole ($B \sim r^{-2}$). Also, an accretion rate of $4 \times 10^{-4} M_{\odot} \text{ yr}^{-1}$ could result in a jet mass loss rate of $4 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$. An escape velocity at the inner boundary of 644 km s^{-1} , would result in a mechanical luminosity of $\sim 3 \times 10^3 L_{\odot}$. On the other hand, if the jet is launched close to the surface of the neutron star (which is not resolved in our numerical simulations), the escape velocity would be $\sim 0.3 c$, and the mechanical luminosity would be $\sim 10^8 L_{\odot}$.

4. Sensitivity of results to the choice of some physical and numerical parameters

4.1 Mass injection rate

The full evolution of the mass transfer rate from the moment of Roche lobe overflow to the moment of CE as seen in the 1D, MESA simulation lasts 30 000 yr (Figure 2) and goes from $1.6 \times 10^{-9} M_{\odot} \text{ yr}^{-1}$ to $9.7 \times 10^{-2} M_{\odot} \text{ yr}^{-1}$ (Table 2). This entire period cannot be modeled in 3D which by necessity can only simulate a much shorter time (21 years for us). The choice was therefore made to model a period of time towards the end of the 30 000 years modeled in 1D between 29 500 and 29 521 years where the mass injection rate goes between 2.3×10^{-4} and $9.7 \times 10^{-2} M_{\odot} \text{ yr}^{-1}$ (see sim-0 in Table 5).

Using sim-0 as reference, we computed three additional simulations with different initial mass transfer rates (called “sim-mdot-#” in Table 3 and Table 5) and hence different start times in the context of the MESA simulation. Therefore, each 3D simulation samples a different part of the \dot{M} vs. time curve in Figure 2. In Table 5 we present a summary of the initial and final parameters of these simulations, noting that, besides the mass transfer rate, all other parameters are the same as the Reference simulation sim-0 – see Section 2.2).

In Figure 13, we present density slices in the orbital plane of these 3D simulations at their final time step (see the fifth column in Table 5) in order to increase the initial mass transfer rate. In the upper left panel, we show the simulation sim-mdot-1, with the smallest initial mass transfer rate of $1.4 \times 10^{-8} M_{\odot} \text{ yr}^{-1}$, running for the first 128 years of the 30 000 yr MESA simulation. As expected, only a small, low mass disk forms at such low mass transfer rate, even if the run time is relatively long.

In sim-mdot-2 with initial mass transfer rate of $1.1 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$ (top right panel), starting towards the end of the 30 000-yr long MESA simulation (only 120 yr before sim-0) and running for 29 yr, a disk forms with a radius of just under $50 R_{\odot}$ and a mass that is not that dissimilar to that of sim-mdot-1, despite the mass injection rate being larger by three orders of magnitude (the run time was four times smaller).

In the lower left panel of Figure 13, we present sim-0, our reference simulation. The initial mass accretion rate is only 20 times higher at the start than the previous one, but it increases far more steeply than for sim-mdot-2. The disk size is of the same order as the one in sim-mdot-2, but the disk mass is ~ 100 times larger.

Finally, sim-mdot-3 is very similar to sim-0 but starts 8 years later with an initial mass transfer rate that is again, ~ 20 times larger. It runs only 13 years to the same end point as sim-0 (see the lower right panel, Figure 13). The disk reaches a similar radius ($\sim 43 R_{\odot}$) and a mass that is 1.6 times larger than the mass of the accretion disk of sim-0, despite the fact that the simulation started 8 years later and therefore ran for only 13 years compared to 21 of the reference simulation and injected slightly less mass. This is due to the fact that disk growth is not only dependent on the mass transfer rate and length of simulation, but also on the specific geometry of the flow, which dictates how much mass accretes through the inner boundary as well as the shape of the disk at the time it is measured, whereby the “edge” of the disk as defined by our criteria (Section 3.1) can vary slightly.

With hindsight, these tests could be performed more systematically so as to gain a better idea of whether the disk parameters as stated in Table 5 are close to what we might expect to be the disk just before CE for a systems such as ours. With these tests as they are we can only state that the disk’s mass and radius are likely reasonable within a factor of $\lesssim 2$ for the mass, and 15% for the radius. Given other sources of uncertainty this is a reasonable and sufficient statement for now.




Figure 13. Densities slices on the orbital plane at the end of the 3D simulations. The models that are shown are: sim-0 (lower left panel), sim-mdot-1 (upper left panel), sim-mdot-2 (upper right panel) and sim-mdot-3 (lower right panel). Note how the density color bar may have different maximum limits.

Table 5. Parameters of a sequence of 3D simulations aiming to assess the resilience of disk parameters (M_{disk} and R_{disk}) to the choice of mass transfer rate and simulation length. The mass-transfer rate, donor mass, accretor mass, and orbital separation are selected at the MESA start time. The disk mass and radius are measured at the end of the simulation.

Model	$\dot{M}_{L1,\text{start}}$ ($M_{\odot} \text{ yr}^{-1}$)	MESA start time (yr)	$\dot{M}_{L1,\text{stop}}$ ($M_{\odot} \text{ yr}^{-1}$)	Length of sim. (yr)	Injected mass (M_{\odot})	Donor mass (M_{\odot})	Accretor mass (M_{\odot})	Orbital separation (R_{\odot})	R_{L1} (R_{\odot})	M_{disk} (M_{\odot})	R_{disk} (R_{\odot})
sim-mdot-1	1.4×10^{-8}	0	1.4×10^{-8}	128	1.8×10^{-6}	6.98	1.40	270	84	6.3×10^{-9}	10
sim-mdot-2	1.1×10^{-5}	29380	6.4×10^{-5}	29	1.1×10^{-3}	6.98	1.40	268	83	5.1×10^{-5}	46
sim-0	2.3×10^{-4}	29500	9.7×10^{-2}	21	8.1×10^{-2}	6.97	1.41	266	83	5.5×10^{-3}	39
sim-mdot-3	3.7×10^{-3}	29508	9.7×10^{-2}	13	5.2×10^{-2}	6.94	1.44	258	81	8.6×10^{-3}	43

4.2 Velocity through the nozzle

The distribution and kinematics of the gas at L_1 (the “nozzle”) is calculated according to the prescription of [Lubow & Shu \(1975\)](#), [Ritter \(1988\)](#), and [Jackson et al. \(2017\)](#). They used Bernoulli’s principle to describe the evolution of the gas moving from the donor’s surface toward L_1 . The gas above the donor photosphere moves with a velocity $v_{L_1} \ll c_T$, where c_T is the isothermal sound speed, while near L_1 , the gas is assumed to reach a velocity comparable to the isothermal sound speed, $v_{L_1} \lesssim c_T$. After the gas passes L_1 , due to the pressure gradient, the gas free falls supersonically into the companion’s Roche Lobe.

In our simulation, we need to set an injection velocity because otherwise gas placed in the nozzle does not enter the computational domain at the prescribed rate. This initial nozzle velocity is therefore arbitrary, and we thus need to ensure that changing its value does not affect the disk parameters.

We test the dependency of the simulation on the velocity of injection (v_{L_1}) at the nozzle, by executing three different simulations using the same parameters as the reference simulation, but setting different nozzle velocities in the x-direction: a subsonic velocity of $7.75 \times 10^4 \text{ cm s}^{-1}$ (sim-0; 0.01 in code units), the isothermal sound speed at the donor’s photosphere, or $5.74 \times 10^5 \text{ cm s}^{-1}$ (sim-vel-1), and a supersonic velocity of $6.16 \times 10^5 \text{ cm s}^{-1}$ (sim-vel-2). See [Table 3](#) for a summary of all the simulations parameters.

The top panel of [Figure 14](#) shows the cumulative mass as a function of radius at 21 years, similar to [Figure 6](#), the nozzle is located at a radius of $74 R_\odot$ ($5.2 \times 10^{12} \text{ cm}$) from the inner boundary. For the reference simulation, sim-0, with the lowest injection velocity, the formation of the disk takes longer since the injected mass needs more time to leave the nozzle and to fall into the companion’s Roche lobe. The accretion disk in this simulation has 11 percent more mass than simulations sim-vel-1 and sim-vel-2. We conclude that relatively small differences in the injection velocity around the isothermal velocity value c_T , do not greatly affect the disk parameters.

The density profile along the z-axis at 21.0 yr ([Figure 14](#), bottom panel) is similar in the three simulations at z-values close to the inner boundary, the disk scale height in the reference simulation at 21 yr ($H_{\text{sim-0}} = 4.9 R_\odot$) is similar to the disk scale height for sim-vel-1 ($H_{\text{sim-vel-1}} = 4.3 R_\odot$) and the disk scale height for sim-vel-2 ($H_{\text{sim-vel-2}} = 5.0 R_\odot$). The density at the edges of the box is an unbound low-density gas interacting with the even lower-density background; this gas does not affect the dynamics/structure of the accretion disk. The arbitrarily assumed velocity injection does not affect the final results of the simulation.

4.3 Sensitivity of the results to background density and temperature

Filling the background with low-density gas is an expedient to ensure that no cell is empty, which would cause the inability to calculate pressure gradients. The value of the background density is $\rho_{\text{bg}} = 2.60 \times 10^{-22} \text{ g cm}^{-3}$ (1×10^{-16} code units), the lowest viable value, below which the gradients are so steep




Figure 14. The cumulative mass as a function of radius (top panel) and the vertical density profile (bottom panel) for models with different injection velocities. Dark green line: $v_{L_1} = 7.75 \times 10^4 \text{ cm s}^{-1}$ (model sim-0), green line: $v_{L_1} = 5.74 \times 10^5 \text{ cm s}^{-1}$ (sim-vel-1), emerald line: $v_{L_1} = 6.16 \times 10^5 \text{ cm s}^{-1}$ (sim-vel-2, which has the necessary velocity to leave the nozzle every time step).

that the code would be forced to take unfeasibly short time steps. As consequence of this background density choice, at the beginning of the simulation a low density rarefaction wave propagates from the discontinuity between the inner boundary and the background moving outward and out of the computational domain. The rarefaction wave leaves the computational box before the injected material reaches the inner boundary. This rarefaction wave does not affect the movement of the injected material, and the evolution of the disk formation shown in Figure 4.




Figure 15. The cumulative mass as a function of radius (top panel) and the vertical density profile (bottom panel) for models with different background temperatures. Purple line: $T_{\text{bg}} = 10^5 \text{ cm s}^{-1}$ (model sim-0), red line: $T_{\text{bg}} = 10^4 \text{ cm s}^{-1}$ (sim-bgT-1), orange line: $T_{\text{bg}} = 10^6 \text{ cm s}^{-1}$ (sim-bgT-2).

We repeat the simulation using the same background density but decreasing and increasing the background temperature to $1 \times 10^4 \text{ K}$ (sim-bgT-1) and $1 \times 10^6 \text{ K}$ (sim-bgT-2), respectively (note that sim-0 has a temperature of $1 \times 10^5 \text{ K}$). The adiabatic index is $\gamma = 1.1$ in all simulations (see Section 4.5 for a discussion about the adiabatic index). In Figure 15, we show the total cumulative mass as a function of radius (top panel) and the density profile of the disk along the z -axis (bottom panel). The vertical density profile of the disk is similar for all three simulations, but the highest temperature (and pressure) simulation results in approximately half of the mass in the disk at 21 years. Increasing the background temperature by an order of magnitude increases the pressure of the background such that it is higher than the pressure inside the nozzle (see fifth column in Table 3), reducing the amount of mass that can flow out from the nozzle. In the case of sim-bgT-2 only half

of the injected material manage to leave the nozzle compared to the reference simulation.

4.4 Convergence tests

We finally test the sensitivity of the simulation to spatial and temporal resolution. Our comparison models have the same parameters as sim-0 (see Table 3) with $(72 \times 96 \times 32)$ cells at the coarsest level; we also preserved the size of the computational box ($-d_{L1} \leq x \leq 1.25d_{L1}$, $-1.5d_{L1} \leq y \leq 1.5d_{L1}$ and $-0.5d_{L1} \leq z \leq 0.5d_{L1}$, where $d_{L1} = 83 R_{\odot}$). We repeated the simulation with increasing levels of refinements: 2 levels (as the low-resolution simulation), 3, 4, and 5 levels (high-resolution simulation). The inner boundary around the companion star has a constant radius of $R_{\text{in}} = 1.3 R_{\odot}$ in all four simulations, and the size of the nozzle is also not resolution dependent. Refinement occurs primarily in the higher density regions close to the accretor, as concentric circles on the orbital plane, and around the nozzle area; this high-resolution refinement expands along the line that connects the nozzle with the inner boundary on the XZ plane.

Table 6. Disk properties for the Reference simulation (sim-0), at 21 yr, with different resolutions.

Resolution (# levels)	R_{disk} (R_{\odot})	M_{disk} (M_{\odot})	H_{disk} (R_{\odot})
2	37	2.68×10^{-6}	7.2
3 (sim-0)	39	5.46×10^{-5}	4.9
4	41	5.81×10^{-4}	4.5
5	42	6.15×10^{-3}	4.6

In Figure 16, we show the convergent behavior of the disk mass and scale height. We measure the properties of the disk at the end of the simulation (21 yr; see Table 6). The results show that the disk properties converge across different resolution levels. For the highest resolution (5 levels) simulation, the disk mass shows a 12% discrepancy compared to the Reference simulation (sim-0; 3 levels of refinement), the disk radius and scale height as measured using the criteria established in Section 3.1 are consistent across the three highest resolution simulations.

4.5 Simulation dimensionality and adiabatic index

The adoption of an isothermal gas was proposed by Lubow & Shu (1975). Armitage & Livio (2000) then showed, using an adiabatic index of $4/3$ in their 2D simulations of a mass transfer into the Roche lobe of a neutron star, that an accretion disk could form. Subsequently, Makita et al. (2000), MacLeod et al. (2017), and Murguía-Berthier et al. (2017) showed, using 3D simulations of mass transfer through a Roche lobe with various system parameters, that the maximum adiabatic index that would allow a disk to form was $\gamma \leq 1.2$, for simulations without radiation or cooling function implemented.

We tested these claims by first performing a 2D comparison simulation of the reference simulation (sim-0, Table 3), and three additional 2D simulations with larger adiabatic indices and then carrying out two 3D simulations with higher




Figure 16. The cumulative mass as a function of radius (top panel) and the vertical density profile (bottom panel) of the accretion disk for different resolutions. sim-0 has 3 levels of refinement.

adiabatic indices than sim-0.

In Figure 17, we present density slices of the four, 2D simulations with $\gamma = 1.1$ (top left panel, a 2D version of sim-0), 1.2 (top right panel), $4/3$ (bottom left panel) and $5/3$ (bottom right panel), after 21 yr. In all simulations, the injected material revolves around the companion and forms a high density disk structure.

In Figure 18 we show density slices of two 3D simulations performed with adiabatic indexes $\gamma = 4/3$ and $\gamma = 5/3$ to investigate the reluctance to form a disk for higher values of γ observed by other authors. This figure should be compared with sim-0 ($\gamma = 1.1$) shown in the last column of Figure 4. For $\gamma = 4/3$, only slightly higher than the value used in sim-0, some of the material gets deflected around the companion, which may prevent the formation of a stable disk. For $\gamma = 5/3$, the pressure gradient dominates over the gravitational force of the companion so that a fraction of the material is deflected away from the center and may inhibit the formation of a disk.

We quantify the difference between 3D simulations with different adiabatic indices (see last row of Table 4). The top panel of Figure 19 shows the cumulative mass as a function of radius, from which we measure the radius and mass of the disk-like structure in each simulation. Even though the simulations have the same mass transfer rates as the reference simulation, the high-density structure formed on the simulation with an adiabatic index $\gamma = 4/3$ is five times smaller ($15.2 R_{\odot}$ or 1.06×10^{12} cm) and two orders of magnitude less




Figure 17. Density in the orbital plane for 2D simulations with different adiabatic indexes. Adiabatic index $\gamma = 1.1$ is on the upper-left panel (a 2D version of the 3D sim-0), $\gamma = 1.2$ in the upper-right panel, $\gamma = 4/3$ in the lower-left panel, and $\gamma = 5/3$ in the lower-right panel. All simulations are plotted at $t=21$ yr.




Figure 18. Density slices in the orbital plane (upper panels) and perpendicular plane (lower panels) for 3D simulations with different adiabatic indexes ($\gamma = 4/3$ left, and $\gamma = 5/3$ right). The simulations are plotted at $t = 21$ yr, and can be compared with last panel of Fig. 4.

massive ($5.37 \times 10^{-5} M_{\odot}$). In the case of the simulation with $\gamma = 5/3$ a radius and a mass value cannot be determined from the cumulative mass plot.

In the bottom panels of Figure 19, we compare the density profile along the z -axis of the simulations with different thermal properties. It is clear that when the adiabatic index is closer to the isothermal value the vertical structure is more defined, as expected from the bottom panels of Figure 18, while for the higher adiabatic indices there is only a very marginal equatorial compression.




Figure 19. The cumulative mass as a function of radius (top panel) and vertical density profile (bottom panel) for 3D models with different adiabatic index. The simulations are plotted at $t = 21$ yr. Density slices of these models are presented in the last column of Figure 4 and in Fig. 18.

These results are consistent with those of Makita et al. (2000), MacLeod & Ramirez-Ruiz (2015), and Murguía-Berthier et al. (2017), who showed that disk formation depends primarily on its thermal properties: a disk forms only with an adiabatic index lower than $\gamma \leq 1.2$ when the gas is cooler and has more compressibility. Although, in the case of $\gamma = 4/3$ a thick disk seems to be formed in our simulations.

5. Discussion and conclusion

The goal of this paper was to study the formation of an accretion disk around a neutron star ($1.4 M_{\odot}$) due to unstable mass transfer from an intermediate mass red supergiant ($7 M_{\odot}$)

through Roche lobe overflow (RLOF). This phase likely immediately precedes a common envelope (CE) in-spiral with concomitant accretion onto the companion and possibly the formation of a jet that may affect the pre-in-spiral, as well as the in-spiral phases. Such feedback may lead to a different outcome than what has been modeled thus far by 3D hydrodynamic simulations without feedback (e.g., Lau et al., 2022).

By necessity, with an explicit 3D simulation, we only model a short time (21 years) of the evolution of the RLOF phase predicted to last 30 000 years by a 1D, implicit model. By carefully choosing to model in 3D the last phases of the mass-transfer before the putative CE in-spiral, we show that the 3D disk mass is likely only a factor of a few smaller than it might be if the entire phase were modeled, and very similar in radius.

We show that the accretion disk in our system grows to a mass of $\sim 5 \times 10^{-3} M_{\odot}$, a radius of $\sim 40 R_{\odot}$, and a scale height of $\sim 5 R_{\odot}$, just before it presumably goes into the CE in-spiral phase. This disk has approximately Keplerian rotation near the inner boundary, while the outer regions have a rotation velocity slightly slower than Keplerian due to pressure support. The temperature profile between the inner boundary and the disk's outer radius can be fit with an exponential law with index -1.1 steeper than -0.75 predicted for accretion disks. An immediate improvement to understanding the disk's temperature (and structure) would be to include explicit cooling. At the one significant figure level, these results are resilient with respect to various physical and numerical choices. The results are well converged with respect to spatial and temporal resolution.

The accretion rate through the inner boundary that surrounds the companion reaches $4 \times 10^{-3} M_{\odot} \text{ yr}^{-1}$, at the end of the simulation, which is consistent with the accretion rate predicted using a Shakura & Sunyaev (1973) formalism with $\alpha = 0.1$. The Eddington limit for mass accretion onto a neutron star of $1.4 M_{\odot}$ is on the order of $10^{-8} M_{\odot} \text{ yr}^{-1}$. X-ray binaries are known to accrete at rates that can be 100 times the Eddington rate. That said the rate modeled for systems accreting in an unstable mass transfer regime can be larger than that (e.g., 1000 times, see Dickson, 2024). Extreme mass accretion rates predicted in the last phases of mass transfer before the CE in-spiral are likely far more complex physical phenomena, with likely extreme X-ray feedback. Moreno Méndez (2022) explained the formation of high mass X-ray binaries, with a black hole as a companion, with hyper accretion rates ($\sim 1 \times 10^8 \dot{M}_{\text{Edd}}$) before the CE phase.

Understanding the potential for jet formation from disks such as this one would be the next critical step because it may impact the early CE, even if the accretion disk and jet do not survive entering the envelope (Murguía-Berthier et al., 2017). Further investigation on the survival of the accretion disk during the CE phase for this system is intended on a follow up study.

Acknowledgments We acknowledge the computing time granted by DGTIC UNAM on the supercomputer Miztli (projects LANCAD-UNAM-DGTIC-281 and LANCAD-

UNAM-DGTIC-321). AJG and FDC acknowledge support from the UNAM-PAPIIT grant IN113424. AJG also acknowledges funding support from Macquarie University through the International Macquarie University Research Excellence Scholarship ('iMQRES') and the Postgraduate Research Fund ('PGRF').

Data Availability Statement The data underlying this article will be shared on reasonable request to the corresponding author.

References

- Abbott, B. P., Abbott, R., Abbott, T., et al. 2016, *Physical review letters*, 116, 241103
- Armitage, P. J., & Livio, M. 2000, *ApJ*, 532, 540
- Brown, G. E., Lee, C.-H., & Moreno Méndez, E. 2007, *ApJ*, 671, L41
- Cehula, J., & Pejcha, O. 2023, *Monthly Notices of the Royal Astronomical Society*, 524, 471
- Chamandy, L., Frank, A., Blackman, E. G., et al. 2018, *MNRAS*, 480, 1898
- Chevalier, R. A. 2012, *The Astrophysical Journal Letters*, 752, L2
- De Colle, F., Granot, J., López-Cámara, D., & Ramirez-Ruiz, E. 2012, *The Astrophysical Journal*, 746, 122
- Dickson, D. 2024, *ApJ*, 975, 130
- Eggleton, P. P. 1983, *ApJ*, 268, 368
- Fryer, C., & Woosley, S. 1998, *The Astrophysical Journal*, 502, L9
- Hayashi, K., Kawaguchi, K., Kiuchi, K., Kyutoku, K., & Shibata, M. 2021, *Phys. Rev. D*, 103, 043007
- Iben, I., J., & Tutukov, A. V. 1984, *ApJS*, 54, 335
- Ivanova, N., Justham, S., Chen, X., et al. 2013, *The Astronomy and Astrophysics Review*, 21, 1
- Jackson, B., Arras, P., Penev, K., Peacock, S., & Marchant, P. 2017, *The Astrophysical Journal*, 835, 145
- Lau, M. Y. M., Hirai, R., González-Bolívar, M., et al. 2022, *MNRAS*, 512, 5462
- López-Cámara, D., De Colle, F., & Moreno Méndez, E. 2019, *MNRAS*, 482, 3646
- López-Cámara, D., De Colle, F., Moreno Méndez, E., Shiber, S., & Iaconi, R. 2022, *MNRAS*, 513, 3634
- López-Cámara, D., Moreno Méndez, E., & De Colle, F. 2020, *Monthly Notices of the Royal Astronomical Society*, 497, 2057
- Lubow, S. H., & Shu, F. H. 1975, *The Astrophysical Journal*, 198, 383
- MacLeod, M., Antoni, A., Murguía-Berthier, A., Macías, P., & Ramirez-Ruiz, E. 2017, *The Astrophysical Journal*, 838, 56
- MacLeod, M., & Ramirez-Ruiz, E. 2015, *The Astrophysical Journal*, 803, 41
- Makita, M., Miyawaki, K., & Matsuda, T. 2000, *Monthly Notices of the Royal Astronomical Society*, 316, 906
- Mineshige, S., Honma, F., Hirano, A., et al. 1994, in *Theory of Accretion Disks — 2*, ed. W. J. Duschl, J. Frank, F. Meyer, E. Meyer-Hofmeister, & W. M. Tscharnutter (Dordrecht: Springer Netherlands), 187–193
- Moe, M., & Di Stefano, R. 2017, *ApJS*, 230, 15
- Moreno Méndez, E. 2022, *arXiv e-prints*, arXiv:2207.14765
- Moreno Méndez, E., López-Cámara, D., & De Colle, F. 2017, *MNRAS*, 470, 2929
- Murguía-Berthier, A., MacLeod, M., Ramirez-Ruiz, E., Antoni, A., & Macías, P. 2017, *The Astrophysical Journal*, 845, 173
- Paxton, B., Bildsten, L., Dotter, A., et al. 2011, *ApJS*, 192, 3
- Paxton, B., Cantiello, M., Arras, P., et al. 2013, *ApJS*, 208, 4
- Paxton, B., Marchant, P., Schwab, J., et al. 2015, *ApJS*, 220, 15
- Paxton, B., Schwab, J., Bauer, E. B., et al. 2018, *ApJS*, 234, 34
- Paxton, B., Smolec, R., Schwab, J., et al. 2019, *ApJS*, 243, 10
- Ramachandran, V., Oskoinova, L. M., Hamann, W. R., et al. 2022, *A&A*, 667, A77
- Ramirez-Ruiz, E., & Lee, W. 2009, *Nature*, 460, 1091
- Ritter, H. 1988, *A&A*, 202, 93
- Savonije, G. 1977, *Astronomy and Astrophysics*, 62, 317
- Shakura, N. I., & Sunyaev, R. A. 1973, *A&A*, 24, 337
- Shakura, N. I., & Sunyaev, R. A. 1973, *Astronomy and Astrophysics*, 24, 337
- Shiber, S., Iaconi, R., De Marco, O., & Soker, N. 2019, *MNRAS*, 488, 5615
- Tauris, T. M., Kramer, M., Freire, P. C. C., et al. 2017, *ApJ*, 846, 170
- Wardle, M. 2007, *Ap&SS*, 311, 35
- Warner, B. 1995, *Cataclysmic Variable Stars*, Cambridge Astrophysics (Cambridge University Press), doi:10.1017/CBO9780511586491

Appendix 1. Numerical considerations

Appendix 1.1 Conservation of mass, energy and angular momentum

As we explain in Sec. 2.2.2, the Reference simulation has constant mass injection through the nozzle, we have external outflow boundaries and the inner inflow boundary. To test for conservation, we run the reference simulation, for 9.5 years after which we switch off the nozzle, change the outflow boundary condition into reflective boundaries, and remove the outflow boundary around the companion star in the middle of the domain. We then run the simulation for an additional 11.5 years, during which we evaluate the conservation of mass, angular momentum, and energy.

For each time step, we calculate the total mass, angular momentum, and energy in the inertial frame of reference. For this we integrate the mass, angular momentum in each cell at every time step. In the case of the total energy, we take into account the kinetic energy of each cell, the potential energy between the fluid and the companion star, and the thermal energy of each cell. The integrated values of mass, angular momentum and energy are shown in Figure 20, where the gray area represents the time when the nozzle and mass injection are on, causing the total mass, energy and angular momentum to increase over time. The dashed light line in each panel represents the first value of mass, energy angular momentum after the nozzle is switched off.

As shown in the top panel of Figure 20, the mass within the reflective boundaries remains constant throughout the simulation, with a maximum variation of 0.02 % at 21 yr. As expected, due to numerical effects such as numerical viscosity, and the way the conservation of the angular momentum equation is discretized, in the middle panel of Figure 20, the total angular momentum grows up to 6% over the 10 years the simulation has reflective boundaries. For the measure of the total energy within the computational box (bottom panel in Figure 20), after turning off the nozzle, there is a decrease in kinetic energy over 0.15 yr. This change is reflected in a step increase in thermal energy of $2.5 \times 10^{18} \text{ erg s}^{-1}$. The energy has a maximum change of 13% after the mass injection finish and remains constant until the end of the simulation. The extent of the non-conservation of the preceding values justifies the approximations made in the simulations.




Figure 20. Total mass (top panel), total angular momentum (middle panel) within the computational domain as a function of time. The bottom panel shows total kinetic energy (Kin energy), potential energy (Pot energy), thermal energy (Therm energy), and total energy (Tot energy) as a function of time. During the gray shadow area, the mass injection is on, and internal inflow boundary is present on the computational box. The dash line in each panel represents the comparative value after the injection is over.