Rewriting modulo in diagrammatic algebras and application to categorification

Léo Schelstraete

Abstract

We develop a rewriting theory suitable for diagrammatic algebras and lay down the foundations of a systematic study of their higher structures. In this paper, we focus on the question of finding bases. As an application, we give the first proof of a basis theorem for graded \mathfrak{gl}_2 -foams, a certain diagrammatic algebra appearing in categorification and quantum topology.

Our approach is algorithmic, combining linear rewriting, higher rewriting and rewriting modulo another set of rules—for diagrammatic algebras, the modulo rules typically capture a categorical property, such as pivotality. In the process, we give novel approaches to the foundations of these theories, including to the notion of confluence. Other important tools include termination rules that depend on contexts, rewriting modulo invertible scalars, and a practical guide to classifying branchings modulo.

This article is written to be accessible to experts on diagrammatic algebras with no prior knowledge on rewriting theory, and vice-versa.

Contents

1	Intr	Introduction 2					
	1.1	State of the art	5				
	1.2	Extended summary	12				
	1.3	Perspectives	18				
2	Linear Gray polygraphs 1						
	2.1	A summary via example	20				
	2.2	<i>n</i> -sesquicategories and their presentations	22				
	2.3	Linear <i>n</i> -sesquicategories and their presentations	28				
	2.4	Gray polygraphs and linear Gray polygraphs	30				
3	Linear Gray rewriting modulo						
	3.1	Coherence modulo from convergence modulo	31				
	3.2	Abstract rewriting modulo	34				
	3.3	Linear rewriting modulo	39				
	3.4	Higher rewriting modulo	48				
	3.5	Higher linear rewriting modulo	53				
	3.6	Summary	59				
4	A ba	asis for graded gl ₂ -foams via rewriting theory	61				
	4.1	Graded \mathfrak{gl}_2 -foams	61				
	4.2	A convergent presentation of graded \mathfrak{gl}_2 -foams	65				
	4.3	Coherence of foam isotopies	67				

4.4	Analysis of monomial local branchings	70
4.5	Addendum: another deformation of \mathfrak{gl}_2 -foams $\ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots$	80

References

1 Introduction

Monoidal categories, and more generally 2-categories, are ubiquitous in mathematics. Their study categorifies classical algebra, exhibiting novel structures and phenomenons [Sel11]; we refer to it as *2-dimensional algebra*, or more loosely speaking, *higher algebra*. While symbols are the language of classical algebra, *string diagrams* (dual to pasting diagrams) are the language of higher algebra. For that reason, linear (strict) 2-categories presented using string diagrams are often called *diagrammatic algebras*.

Diagrammatic algebras are particularly prevalent in representation theory and low-dimensional topology. A classical example is the Temperley–Lieb category: it describes (some of) the representation theory of the quantum group $U_q(\mathfrak{sl}_2)$. It is also at the heart of the definition of the Jones polynomial [Jon85; Kau87], an invariant of knots whose discovery birthed the field of quantum topology. The morphisms of the Temperley–Lieb category are pictured as $\mathbb{C}(q)$ -linear combinations of certain diagrams (see Fig. 1.1). This allows some of its defining relations to be interpreted as certain isotopies of diagrams—a typical feature of diagrammatic algebras.

$$\left[\bigcup_{i=1}^{n} \bigcup_{j=1}^{n} \left[(q+q^{-1}) \right] \right] = \left[(q+q^{-1}) \right]$$

81

Figure 1.1: A relation in the Temperley– Lieb category. Thanks to the diagrammatics, it is easily understood as "evaluate the closed loop to $q + q^{-1}$, and apply a planar isotopy".

Other examples of diagrammatic algebras in representation theory include:

- diagrammatic algebras describing intertwiners of certain representations, such as the Temperley–Lieb category, gl_n-webs [CKM14; Kup96], or the Brauer category;
- diagrammatic categorification, such as the 2-Kac–Moody algebra $\mathcal{U}(\mathfrak{g})$ of Khovanov–Lauda [KL09; KL10] and Rouquier [Rou08] categorifying the quantum group $U_q(\mathfrak{g})$ (where \mathfrak{g} is a simple complex Lie algebra), or the Elias–Williamson category $\mathcal{H}(W)$ [EW14] categorifying the Hecke algebra $H_q(W)$ (where W is a Coxeter group);
- diagrammatic supercategorification, such as the 2-Kac–Moody superalgebra $SU(\mathfrak{g})$ [BE17b] conjecturally categorifying the covering quantum group $U_{q,\pi}(\mathfrak{g})$ [CHW13], or the graded-2-category **GFoam**_d of graded \mathfrak{gl}_2 -foams [SV23] (discussed below).

While the third family of examples is less classical than the other two, it is representative of a general trend: newly discovered diagrammatic algebras (here $SU(\mathfrak{g})$) often arise as variations, or "deformations", of classical diagrammatic algebras (here $U(\mathfrak{g})$), in the loose sense that the presentation of $SU(\mathfrak{g})$ has many similarities with the presentation of $U(\mathfrak{g})$.

Unfortunately, diagrammatic algebras are hard. In particular, finding a hom-basis, i.e. a basis for each hom-space, is hard. Often, a careful study of the presentation leads to a candidate hombasis, shown to generated hom-spaces—the hard part is to show linear independence. A classical solution is to find a concrete faithful representation of the diagrammatic algebra, extrapolating linear independence of the candidate hom-basis from the linear independence of its image. However, in the higher context, finding such a concrete representation is a difficult task—assuming it

categorical property	topological interpretation		
interchange law	planar rectilinear isotopies		
pivotality	planar isotopies		
symmetric structure	rectilinear (non-planar) isotopies		
symmetric pivotality	(non-planar) isotopies		

Table 1: categorical properties of 2-categories and their topological interpretations as string diagrams; see [FY89; JS; JS91; KL80] for proofs (in partial cases) and [Sel11] for a review.

even exists. To give an example, the basis theorem for 2-Kac–Moody algebras $\mathcal{U}(\mathfrak{g})$ was originally proved in the restricted case $\mathfrak{g} = \mathfrak{sl}_n$ by Khovanov–Lauda [KL10] using a representation on the cohomology of flag varieties. The general proof for any type \mathfrak{g} was only sketched a decade later, by work of Webster [Web18]. Outside of the case $\mathfrak{g} = \mathfrak{sl}_2$ [BK22], the analogue statement for 2-Kac–Moody superalgebras remains conjectural.

As one can expect, many subsequent questions about these diagrammatic algebras rely on knowing that hom-spaces have the expected dimension. For instance, if the basis conjecture holds fo 2-Kac–Moody superalgebras, it would follow that they do indeed categorify covering groups—this remains a conjecture outside of special cases [EL16; KK12; KKO13].

The above discussion suggests the following:

QUESTION: Are there generic and intrinsic techniques to study presentations of diagrammatic algebras? In particular, are there generic and intrinsic techniques to find hom-bases in diagrammatic algebras?

Better still, we may hope to implement these techniques on a computer. In classical algebra, these questions and related ones have a long history, which includes Gröbner basis and Buchberger's algorithm for commutative algebras [Buc06], Shirshov's work on PBW-basis in Lie algebras [Shi09], and Bokut's composition lemma [Bok76] and Bergmann's diamond lemma for associative algebras [Ber78]. Each of these approaches can be seen as an instance of (classical) *rewriting theory*: the study of presented (classical) algebraic structures from an algorithmic perspective. By its very nature, the rewriting approach is well-suited to be implemented on a computer. It also gives an abstract tool to find bases, and more generally to find explicit generators of higher relations, known as *syzygys*; see [GHM19] for a modern perspective.

Recent decades has seen the emergence of *higher rewriting theory*, an analogue of classical rewriting theory suitable for 2-categories and *n*-categories; see Subsection 1.1 for a discussion on the literature. Unfortunately, the application of the theory to diagrammatic algebras have had limited success. This is perhaps not surprising, given the many qualitative differences between classical associative algebras and diagrammatic algebras. Rather, we should expect that mimicking classical rewriting theory at the higher level will not be sufficient, and that a richer and more flexible rewriting theory is needed. There have been notable attempts [All18a; DEL21; Dup21; Dup22] in that direction in the later years; we discuss them and relations with our work in Subsection 1.2.8.

In this work, we lay down the foundations of a new rewriting theory:

MAIN RESULT: There exists a rewriting theory sufficiently flexible to be applicable to diagrammatic algebras. In particular, it gives generic and intrinsic techniques to find hom-bases in diagrammatic algebras.

We now discuss some aspects of our theory; a more in-depth introduction can be found in the extended summary (Subsection 1.2). The casual reader may wish to first read the state of the art (Subsection 1.1), which also serves as an introduction to rewriting theory for non-experts.

Inspired by the work of Dupont [Dup22], an important feature of our theory is the ability to rewrite modulo. As pointed out above, it is often the case that certain defining relations of a diagrammatic algebra capture a categorical property, which in turn admits a topological interpretation in terms of string diagrams (see Table 1). For instance, in a pivotal category string diagrams are best understood up to planar isotopies. When rewriting modulo, we partition the set of relations into two sets: *oriented* relations, thought as specific to the given diagrammatic algebra, and *unoriented* relations, thought as intrinsic to the underlying categorical structure. Working modulo has a price: it makes the classification of branchings significantly more involved. We take some time explaining how to deal with that situation, emphasizing the use of the naturality conditions associated with the chosen categorical structure (see Subsection 1.2.6).

Furthermore, we allow the modulo data to contain scalars. For instance, the 2-Kac–Moody superalgebra $SU(\mathfrak{g})$ is a super-2-category [BE17a], in the sense that its interchange law only holds up to sign:

Here $p(\alpha)$ and $p(\beta)$ denotes the extra data of a parity for α and β . In that situation, one would still like to think of string diagrams up to planar rectilinear isotopies, although this equivalence only holds up to sign. Adding scalars to the relations of a presentation is the simplest way of "deforming" that presentation, and appears routinely in newly discovered diagrammatic algebras.

In fact, the "higher part" of our theory is not based on linear (strict) 2-categories, but on *linear* sesquicategories (Section 2): in a nutshell, linear 2-categories without the interchange law. This is both motivated by examples such as SU(g) and by theoretical considerations; see Subsections 1.2.4 and 1.2.7. In analogy with the terminology of [FM18], we call our theory *linear Gray rewriting* modulo.

Contrary to (say) commutative algebras, our approach is not a *one-fit-all* approach, that could be neatly expressed as a single black-box. Instead, our exposition of the theory (Section 3) emphasizes how it is a combination of smaller results. This flexibility allows the use of *context-dependent rewriting rules*, in the sense that local rewriting rules may be conditioned by the global context (see Subsection 3.5.1).

On the theoretical side, we build our theory on the novel notion of \succ -*tamed congruence* (see Subsection 1.2.3 and Lemma 3.16), where \succ is a preorder. \succ -tamed congruence serves as a replacement for confluence in classical rewriting theory. Here is a schematic:



In the second diagram, horizontal positions are used to suggest relative ordering with respect to \succ , reading from left to right. \succ -tamed congruence implies confluence, but not conversely. We explain

in Subsection 1.2.4 why modifying the foundation in this way is necessary and, in hindsight, makes the theory clearer.

Finally, we apply the theory to a diagrammatic algebra arising from categorification and quantum topology, the graded-2-category **GFoam**_d of graded \mathfrak{gl}_2 -foams (Section 4):

Main theorem A (Theorem 4.10). The graded-2-category \mathbf{GFoam}_d has the expected hom-basis.

A graded-2-category [SV23] is analogous to a super-2-category, where the interchange law only holds up to scalar (see Subsection 1.2.1). At the time of writing, our approach is the only known approach to this result; it gives the first application of rewriting theory to quantum topology. The result is of independent interest: indeed, the higher representation theoretic construction of odd Khovanov homology given in [SV23] relies on it.

The rewriting approach allows a classification of the possible "deformations" of \mathbf{GFoam}_d , in a sense to be made precise. This leads to another variant \mathbf{GFoam}'_d of \mathbf{GFoam}_d . Topologically, the diagrammatic algebras \mathbf{GFoam}_d and \mathbf{GFoam}'_d respectively relate to type X and type Y in odd Khovanov homology. We expect that this fact hides a greater correspondence between higher structures; see Remark 4.37. More generally, we expect that rewriting theory will allow an exploration of higher structures in diagrammatic algebras; see Subsection 1.3 for further speculations.

1.0.1 Organisation

The rest of this introduction consists of a state of the art (Subsection 1.1)—which also serves as an introduction to rewriting theory for non-experts—an extended summary (Subsection 1.2) and some perspectives for future directions of research (Subsection 1.3).

Apart for this introduction, the article has three sections. The short Section 2 defines the necessary categorical structures at play; in particular, a suitable notion of presentation for linear sesquicategories given by *linear sesquipolygraphs* (Definition 2.11); see also Subsections 1.1.2 and 2.4.1 in the state of the art and Subsection 1.2.1 in the extended summary. Section 2 begins with a description of the nomenclature using an example: we expect it to be mostly sufficient for the impatient reader.

Section 3 is the heart of the paper: it develops the whole theory, starting from first principles. See also Subsections 1.1.1 and 1.1.3 to 1.1.6 in the state of the art and Subsections 1.2.2 to 1.2.8 in the extended summary.

Finally, Section 4 consists of the proof of the basis theorem for graded \mathfrak{gl}_2 -foams (Main theorem A), exhibiting all the techniques developed in Section 3.

1.0.2 Acknowledgments

The author would like to thank Sigiswald Barbier, Jon Brundan, Ben Elias, Yves Guiraud, Louis-Hadrien Robert, Pedro Vaz and Emmanuel Wagner for their interest, questions and comments. Some string diagrams were done using SaTeX. The author was supported by the Fonds de la Recherche Scientifique–FNRS under the Aspirant Fellowship FC 38559 and by the Max Planck Institute for Mathematics.

1.1 State of the art

We give a restricted state of the art on rewriting theory, suitable for our purpose. This section is also meant as a gentle introduction to rewriting theory for the casual reader.

1.1.1 What is rewriting theory?

The word problem for monoids asks the following question: given a presented monoid G with generators in the set X, is there an algorithm that decides whether two words with letters in X are equal as elements of G? While known to be undecidable in general [Mar47; Pos47], one can hope to solve the word problem in practical cases. *Rewriting theory* suggests the following method, which we illustrate with the symmetric group on three strands (understood as a monoid), defined using its Coxeter presentation:

$$\mathfrak{G}_3 = \langle \sigma, \tau \mid \sigma \sigma = 1, \tau \tau = 1, \sigma \tau \sigma = \tau \sigma \tau \rangle.$$

A rewriting system consists of a choice of orientations on the defining relations. For instance:

$$\mathsf{P}_{\mathfrak{G}_3} \coloneqq (\mathsf{X}, \mathsf{R}), \quad \text{with} \quad \mathsf{X} = \{\sigma, \tau\} \quad \text{and} \quad \mathsf{R} = \{\sigma\sigma \to 1, \tau\tau \to 1, \sigma\tau\sigma \to \tau\sigma\tau\}. \tag{2}$$

Let X^{*} denote the set of words in letters in X. Any rewriting system defines a non-deterministic algorithm, where for each oriented relation $A \rightarrow B$ and words $x, y \in X^*$, the algorithm may perform the reduction $xAy \rightarrow xBy$, but not the reduction $xBy \rightarrow xAy$. In the terminology of rewriting theory, $xAy \rightarrow xBy$ is called a *rewriting step*, and a successive composition of rewriting steps, denoted $a \xrightarrow{*} b$, is called a *rewriting sequence*.

A pair of co-initial rewriting sequences $(f: a \xrightarrow{*} b, g: a \xrightarrow{*} b')$ is called a *branching*, and a pair of a co-terminal rewriting sequences $(f': b \xrightarrow{*} c, g': b' \xrightarrow{*} c)$ is called a *confluence*. A branching that admits a confluence is said to be *confluent*. This is illustrated in the diagrams below, where plain arrows (resp. dotted arrows) denote branchings (resp. confluences):



Figure 1.2 gives examples of confluent branchings in $P_{\mathfrak{G}_3}$.

To solve the word problem, we require two key properties: *termination*, which postulates that any rewriting sequence terminates, and *confluence*, which postulates that every branching is confluent. A rewriting system which is both terminating and confluent is called *convergent*. In that case, the non-deterministic algorithm always produces an output, and this output is always the same. Given a word as input, we call the output its *normal form*. It is not so hard to prove that under convergence, two words are equal in the associated monoid if and only if they have identical normal forms. This provides a solution to the word problem.

We are left with the problem of showing that indeed, the rewriting system $P_{\mathfrak{G}_3}$ is convergent. Termination is not hard: it can be shown using a suitable partial order. On the other hand, confluence must in principle be checked for *every* branching. A branching (f, g) for which f and gare rewriting *steps* (in contrast to rewriting *sequences*) is said to be *local*. For instance, the three branchings in Fig. 1.2, depicted in plain arrows, are local. *Newmann's lemma* (Lemma 3.13) states that assuming termination, confluence follows from confluence of local branchings.



Figure 1.2: Critical branchings in $P_{\mathfrak{G}_3}$.

Still, many local branchings remain, as the following ones (for any two words $x, y \in X^*$):



Each of them admits a "somehow canonical" confluence. The first branching is an independent branching: intuitively, it consists of two rewriting steps that do not interact with each other. The

second branching is a *contextualized branching*: it is of the form x(f, g)y, for (f, g) the first local branching in Fig. 1.2, and the confluence of x(f, g)y is canonically induced from the confluence of (f, g). A critical branching is a local branching which is neither an independent branching nor a contextualized branching. As suggested by our discussion, confluence of local branchings follows from confluence of critical branchings. The latter constitute the minimal amount of computations one needs to perform in a given situation; the rest follows from general considerations. The reader may convince themself that Fig. 1.2 gives a complete list of critical branchings in $P_{\mathfrak{G}_3}$. It follows that the rewriting system $P_{\mathfrak{G}_3}$ is both terminating and locally confluent, and hence convergent by Newmann's lemma.

Polygraphs 1.1.2

What is a presentation of a (strict and small) n-category? As a first example, consider again the rewriting system $P_{\mathfrak{G}_3} = (X, R)$ defined in (2). Viewing a monoid as a category with a single object {*}, the set X consists of generating 1-cells with source and target the object {*}. We encapsulate the latter fact with (trivial) source and target maps $s_0, t_0 : X \to \{*\}$. In this perspective, the set of words X* is the free category generated by X. Similarly, the set R consists of generating 2-cells, and we define maps $s_1, t_1 \colon \mathsf{R} \to \mathsf{X}^*$ setting $s_1(r) = A$ and $t_1(r) = B$ for each oriented relation $r: A \to B$. Reformulated in this way, $\mathsf{P}_{\mathfrak{G}_3}$ defines the data of a 2-polygraph. More generally, an *n*-category can be presented by an (n + 1)-polygraph, with generating (k + 1)-cells P_{k+1} defined on the free *k*-category P_k^* generated by the lower cells:



structure	presentation		structure	presentation
set	1-polygraph		module	linear 1-polygraph
category (monoid)	2-polygraph (with one object)	(linear category (associative algebra)	linear 2-polygraph (with one object)
2-category	3-polygraph		linear 2-category	linear 3-polygraph

Table 2: Low-dimensional *n*-polygraphs

Table 3: Low-dimensional linear *n*-polygraphs

the 2-polygraph
$$\mathsf{P}_{\mathfrak{G}_3}$$
 presenting \mathfrak{G}_3 an $(n+1)$ -polygraph P presenting an *n*-category

See Table 2 for a summary of low-dimensional *n*-polygraphs. The *n*-category presented by a (n+1)-polygraph P is obtained by quotienting P_n^* by the relation $s_n(r) = t_n(r)$ for each $r \in \mathsf{P}_{n+1}$.

Polygraphs were first introduced by Street [Str76], under the name of *computads*; the term *signatures* also appears in the literature. Polygraphs were independently introduced by Burroni [Bur93] to study generalizations of the word problem—the terminology is by now standard in the rewriting community.

1.1.3 Higher rewriting

Polygraphs can be thought as higher-dimensional rewriting systems. Consider once again the 2-polygraph $P_{\mathfrak{G}_3} = (X, R)$ presenting \mathfrak{G}_3 from (2). A word in X is nothing else than a path of generating 1-cells in X:

$$\sigma\tau\sigma\in\mathsf{X}^*\quad\leftrightarrow\quad\ast\overset{\sigma}{\to}_{\mathsf{X}}\ast\overset{\tau}{\to}_{\mathsf{X}}\ast\overset{\sigma}{\to}_{\mathsf{X}}\ast$$

In this way, each element of X is thought as a 1-dimensional rewriting step, and each word in X^* as a rewriting sequence. With this point of view, a generating 2-cell in R is nothing else than a generating 2-dimensional rewriting step between 1-dimensional rewriting sequences:



Similarly, in an *n*-polygraph one thinks of a (k + 1)-cell as a generating (k + 1)-dimensional rewriting step between k-dimensional rewriting sequences.

Higher rewriting theory has witnessed increasing interests in the last two decades, prominently by the French school [GM09; GM13; GM18; Gui06; Laf03; Mim14]. As a major application, let us note the construction of a homology theory for ω -categories, based on polygraphic resolutions [Gue21; LM09; LMW10; Mét03; Mim10]. For further details, we refer the reader to the recent monograph on the subject [Ara+23].

In general, the rewriting theory associated to an (n+1)-polygraph is called (n+1)-dimensional rewriting theory; it presents an n-category. The 1-dimensional case corresponds to rewriting in sets; it is also known as *abstract rewriting*. As we have seen, 2-dimensional rewriting theory corresponds to rewriting in categories, and monoids in particular. In this case, a relation $r: A \to B$, i.e. a generating 2-cell, can always be composed on the left and on right with 1-cells x and y respectively, leading to a new relation

$$x \star_0 r \star_0 y \colon x \star_0 A \star_0 y \to x \star_0 B \star_0 y.$$

This process is called *contextualization* (here \star_0 denotes the 0-composition, i.e. the horizontal composition). We call the data of x and y a *context*, denoted with the letter Γ , and write $\Gamma[r]$ for the relation $x \star_0 r \star_0 y$. In monoids, contextualization amounts to multiplying a relation on the left and on the right with words, as we have seen already in Subsection 1.1.1 with the notion of contextualized branchings.

Three-dimensional rewriting is rewriting in 2-categories. In this case, a relation $r: A \to B$ is a generating 3-cell, and contextualization amounts to first composing horizontally with 1-cells xand y, and then vertically with 2-cells α and β :



We similarly write $\Gamma[r]$ for a contextualization of the relation r. In what follows, we will rarely discuss the cases n > 3, and often say *higher rewriting theory* to refer to the case n = 3. The toolbox of higher rewriting theory resembles the one of rewriting in monoids: Newmann's lemma is (as always) applicable, and one has similar notions of independent and contextualized branchings.

1.1.4 Linear rewriting

Linear rewriting is the algorithmic study of presented modules. Given a commutative ring \Bbbk , a \Bbbk -module M is presented by a set \mathcal{B} , together with a set R of oriented relations on $\langle \mathcal{B} \rangle_{\Bbbk}$, the free \Bbbk -module generated by \mathcal{B} . This is encapsulated by the data of a *linear 1-polygraph*:

$$\langle \mathcal{B} \rangle_{\mathbb{k}} \xleftarrow{s}{t} \mathbf{R}.$$

Similarly, *linear 2-polygraphs* present linear categories, and in particular associative algebras [GHM19]; and linear (n + 1)-polygraphs present linear *n*-categories [All18a]. See Table 2 for a summary of low-dimensional linear *n*-polygraphs. The associated linear *n*-dimensional rewriting theory was studied by Guiraud, Hoffbeck and Malbos in the case n = 2 [GHM19], and by Alleaume in the case n = 3 [All18a], with application to the oriented affine Brauer algebra; see Subsection 1.2.8 for how our approach differs.

The most classical setting for linear rewriting is commutative algebras. For that reason, we call *monomials* the elements of \mathcal{B} . An oriented relation $r: s(r) \to t(r) \in \mathbb{R}$ is assumed to be of the form "rewrite a monomial into a linear combination of monomials", that is:

$$r: b \to_{\mathbb{R}} \lambda_1 b_1 + \ldots + \lambda_n b_n \qquad \text{for } \lambda_1, \ldots, \lambda_n \in \mathbb{K}, b, b_1, \ldots, b_n \in \mathcal{B}$$
(3)

We say that relations in r are *left-monomial*.

A generic rewriting step is of the form

$$\lambda r + v \colon \lambda s(r) + v \to_{\mathbf{R}} \lambda t(r) + v \quad \lambda \in \mathbb{k} \setminus \{0\}, v \in \langle \mathcal{B} \rangle_{\mathbb{k}}$$

and it is said to be *positive* if the monomial $s(r) \in \mathcal{B}$ does not appear in the linear decomposition of v. For example, if $\mathcal{B} = \{a, b, c\}$ and $r: a \to b + c \in \mathbb{R}$ is an oriented relation, then both

$$2a + b = a + (a + b) \longrightarrow_{\mathbb{R}} (b + c) + (a + b)$$
 and $2a + b \longrightarrow_{\mathbb{R}} 2(b + c) + b$

are rewriting steps, but only the latter is positive (note the use of dashed and plain arrows to distinguish the two). To avoid rewriting loops, one must restrict to positive rewriting steps. Otherwise, 0 rewrites into 0:

$$0 = a - a \dashrightarrow_{\mathsf{R}} (b + c) - a \dashrightarrow_{\mathsf{R}} (b + c) - (b + c) = 0.$$

Positivity may look like a minor modification of the theory; in fact, it constitutes the main difficulty of the linear setting. We shall say more about that in Subsection 1.2.4.

We denote \mathbb{R}^{st} the set of rewriting steps and \mathbb{R}^+ the set of positive rewriting steps. Similarly to the abstract setting, we can reduce the study of confluence to the study of *local* confluence. In fact, we can further reduce to local *monomial* confluence, where a *monomial branching* is a branching whose source is a monomial:

Lemma 1.1 (linear Newmann's lemma). Let S = (B; R) be a linear 1-polygraph. If R^+ terminates, then confluence of monomial local R^+ -branchings implies confluence.

1.1.5 Rewriting modulo

Enforcing all relations to be oriented can be too restrictive. Instead, one may wish to rewrite with a set of oriented relations \mathbb{R}^1 , modulo another set of *un*oriented relations \mathbb{E} . More precisely, the working data of *abstract rewriting modulo* is given by two 1-polygraphs

$$X \xleftarrow{s}{t} \mathbb{R}$$
 and $X \xleftarrow{s}{t} \mathbb{E}$,

defined on the same underlying set X. Denote by $\mathbb{E}^{\top} = (\mathbb{E} \cup \mathbb{E}^{-1})^*$ the free groupoid generated by \mathbb{E} . Intuitively, relations in \mathbb{E}^{\top} are *un*oriented rewriting sequences in \mathbb{E} . In this context, a *rewriting step modulo* is a composition $e' \circ r \circ e$ with $r \in \mathbb{R}$ and $e, e' \in \mathbb{E}^{\top}$:

$$\bullet \overset{e}{\leadsto_{\mathbb{E}}} \bullet \overset{r}{\longrightarrow}_{\mathbb{R}} \bullet \overset{e'}{\leadsto_{\mathbb{E}}} \bullet$$

In other words, in between a rewriting step in \mathbb{R} one can apply an arbitrary number of relations in \mathbb{E} , in any direction. The data $\mathbb{S} = (X; \mathbb{R}, \mathbb{E})$ defines an *abstract rewriting system modulo*, and a rewriting step modulo as above is called an \mathbb{S} -rewriting step.

Typically, the modulo data \mathbb{E} will consists of relations thought as being "structural", in the sense of being part of some underlying algebraic structure. For instance, rewriting in commutative algebras is implicitly rewriting in associative algebras modulo commutativity (\mathbb{E} consists of relations $xy \to yx$ for all monomials x and y).

Rewriting modulo allows an inductive approach to the word problem. Indeed, instead of trying to fit the relations $\mathbb{R} \sqcup \mathbb{E}$ into a single convergent rewriting system, one can instead show that \mathbb{R} is convergent "modulo \mathbb{E} " on one hand, and that \mathbb{E} is convergent on the other hand.

Different variants of rewriting modulo have been developed, often with more restrictive modulo rules than the ones described above [Hue77; JK86; JL12; Mar98; PS81; Vir95]. Rewriting modulo is used in [CDM22] to study confluence in Lawvere theories. A higher analogue to rewriting modulo was introduced in [DM22], using the formalism of double categories. In the linear setting, Dupont extended Alleaume's approach [All18a] modulo in order to rewrite modulo pivotality in 2-categories [Dup22]. Based on his theory, he proposed an approach to the basis problem in 2-Kac–Moody algebras in simply-laced cases [Dup21]. An approach to rewriting modulo in super-2-categories was also proposed in [DEL21], motivated by the study of the 2-Kac–Moody superalgebra in the case $g = \mathfrak{sl}_2$; see Subsection 1.2.8 for how our approach differs from [DEL21; Dup21; Dup22].

¹We use blackboard font to refer to abstract rewriting; this is unrelated to the set of real numbers.



Figure 1.3: In a 3-category, a Gray category or a 3-sesquicategory, the interchange law for 2-morphisms respectively holds strictly, holds weakly via interchangers, or does not hold (a priori).

1.1.6 Gray rewriting

Starting with n = 3, the strict and weak notions of an *n*-category start to diverge: while a bicategory is alway equivalent to a 2-category (i.e. a strict bicategory), not every tricategory is equivalent to a 3-category (i.e. a strict tricategory). However, every tricategory is equivalent to a *Gray category* [GPS95]. In that sense, Gray categories provide a simpler notion than tricategories, while retaining their expressivity:



In Gray categories, associativity and unitality hold strictly, as for 3-categories. However, the interchange law for 2-morphisms only holds up to certain coherent 3-morphisms, called *interchangers*; see Fig. 1.3.

In [FM22], Forest and Mimram initiated the study of rewriting theory for *weak n*-categories, starting with Gray categories. It turns out to be easier to consider the more general framework of *n*-sesquicategories [Ara22; FM22; Str96] (called *n*-precategories in [FM22]). As for Gray categories, *n*-sesquicategories are strictly associative and unital structures. However, they do not contain any coherence data for interchange laws; see Fig. 1.3. One can understand an *n*-sesquicategories play a san "unfinished definition" for what a semistrict *n*-category would be. Sesquicategories play a prominent role in the graphical proof-assistant *Globular* and its successor *homotopy.io* [BKV18; BV17; Dor23; RV19].

Recall that an *n*-category is presented by an (n + 1)-polygraph (see Subsection 1.1.2); similarly, an *n*-sesquicategory is presented by an (n + 1)-sesquipolygraph [FM22]. A 0-, 1- and 2-sesquipolygraph is the same as 0-, 1- and 2-polygraph; the distinction only starts to appear with 3-sesquipolygraphs. In particular, a 2-sesquicategory is presented by a 3-sesquipolygraph. A 3-sesquipolygraph which contains generating interchangers is called a *Gray polygraph*; in this case, it presents a 2-category. Note that both Gray polygraphs and 3-polygraphs present 2-categories. However, the former explicitly contains interchangers as generating 3-cells, which alters the associated rewriting theory.

1.2 Extended summary

1.2.1 Linear Gray polygraphs

Graded-2-categories [SV23] are generalizations of 2-categories and super-2-categories, where the interchange law only holds up to scalar:

$$\overset{\alpha}{\bullet}_{\beta} = \mu(\deg \alpha, \deg \beta) \overset{\bullet}{\bullet}_{\alpha} \overset{\bullet}{\bullet}_{\beta}$$
(4)

Here deg α and deg β denotes the extra data of a degree for α , β in some abelian group, and μ associates a scalar to the pair (deg α , deg β). In Section 2, we define *linear n-sesquicategories* and *linear n-sesquipolygraphs* as direct linear analogues of *n*-sesquicategories and *n*-sesquipolygraphs [FM22]. We call graded interchangers the 3-cells capturing the graded interchange law (4). A *linear Gray polygraph* is defined as a linear 3-sesquipolygraph which contains its own graded interchangers. We get a notion of presentation for a graded-2-category, suitable for rewriting theory:

Main definition A (Definition 2.17). *A* presentation *of a graded-2-category is a linear Gray poly-graph.*

As our running example for this extended summary, we define °P, the \mathbb{Z} -linear Gray polygraph of superadjunction. Its underlying 2-polygraph is defined as °P₀ = {*} (a single object), °P₁ = { | } (a single 1-cell) and °P₂ = { \cap , \bigcup } (unit and counit, reading from bottom to top). In addition, the two 2-cells are given a parity

$$p(\bigcap) = p(\bigcup) = 1 \in \mathbb{Z}/2\mathbb{Z},$$

which extends additively to generic 2-cells in ${}^{\circ}\mathsf{P}_{2}^{*}$. Finally, we set ${}^{\circ}\mathsf{P}_{3} = {}^{\circ}\mathsf{R}_{3} \sqcup {}^{\circ}\mathsf{E}_{3}$ where

$$^{\circ}\mathsf{R}_{3} = \left\{ \bigcup \rightarrow \left|, \bigcup \rightarrow -\right| \right\} \text{ and } ^{\circ}\mathsf{E}_{3} = \left\{ \left| \stackrel{\bullet}{\overset{\circ}{\overset{\circ}}}_{\beta} \stackrel{\rightarrow}{\overset{\circ}{\overset{\circ}}}_{\text{for all } \alpha, \beta \in {}^{\circ}\mathsf{P}_{2}^{*}} \stackrel{\bullet}{\overset{\circ}{\overset{\circ}}}_{\beta} \right| \right\}.$$

We denote ${}^{\circ}\mathsf{R}$ and ${}^{\circ}\mathsf{E}$ the linear 3-sesquipolygraphs with ${}^{\circ}\mathsf{P}_{2}^{*}$ as their underlying 2-polygraph and 3-cells ${}^{\circ}\mathsf{R}_{3}$ and ${}^{\circ}\mathsf{E}_{3}$, respectively. Here ${}^{\circ}\mathsf{E}$ is the linear 3-sesquipolygraph of super interchangers on ${}^{\circ}\mathsf{P}_{2}^{*}$ and ${}^{\circ}\mathsf{P} = {}^{\circ}\mathsf{R} \sqcup {}^{\circ}\mathsf{E}$ is a linear Gray polygraph.

Denote $^{\circ}C$ the super-2-category presented by $^{\circ}P$. Superadjunction is the super analogue of the classical notion of adjunction: as such, we expect hom-spaces in $^{\circ}C$ to have the same dimension as its classical analogue. We show this fact using *linear Gray rewriting modulo*; the working data is $^{\circ}S = (^{\circ}R, ^{\circ}E)$, which records the splitting of $^{\circ}P$ into oriented relations $^{\circ}R$ and unoriented relations $^{\circ}E$. (Of course, one can give a much simpler proof, but this is not our point.) In passing, we explain the sign in the second 3-cell of $^{\circ}R_3$ (see the discussion after Lemma 1.3).

Linear Gray polygraphs and related notions are formally defined in Section 2.

1.2.2 Linear rewriting modulo

Our study of rewriting modulo starts with abstract rewriting modulo—rewriting modulo in sets—in Subsections 3.1 and 3.2. We formally introduce the notion of an *abstract rewriting systems modulo* $\mathbb{S} = (X; \mathbb{R}, \mathbb{E})$ (or *abstract RSM* for short); see the notations of Subsection 1.1.5.

We then proceed with linear rewriting modulo—rewriting modulo in k-modules for some commutative ring k—in Subsection 3.3, starting with the notion of a *linear rewriting system modulo* S = (B; R, E) (or *linear RSM* for short). The working data is given by two linear 1-polygraphs R and E defined on the same underlying set of monomials \mathcal{B} , with the notations of Subsection 1.1.4. There are additional conditions, including that R is left-monomial (see (3)), and that relations in E are of the form $b \to \lambda b'$ for some λ an invertible scalar; we say that E is *monomial-invertible*. This later fact means that we allow rewriting *modulo invertible scalars*. For instance, fix two 1-morphisms μ and ν in in ° \mathcal{C} . We get a module ° $\mathcal{C}(\mu, \nu)$ presented by the linear RSM S = (\mathcal{B} ; R, E) where $\mathcal{B} = {}^{\circ}P_{2}^{*}(\mu, \nu)$, R = °R(μ, ν) and E = °E(μ, ν). In other words, °P induces a family of linear RSMs, one for each hom-space, where we rewrite modulo the superinterchange law.

In analogy with the non-modulo setting, we define a set of rewriting steps S^{st} and a set of *positive* rewriting steps S^+ ; the later is the canonical abstract RSM associated to S. Denote NF_S the k-module of normal forms for S^+ , that is, the k-module consisting of elements on which S^+ terminates.

Rewriting fact A (BASIS-FROM-CONVERGENCE THEOREM 3.36). Let M be a k-module presented by a linear rewriting system modulo S = (B; R, E). If S^+ is convergent, then the canonical linear map

 $\mathrm{NF}_{S}/\langle E \rangle_{\Bbbk} \to M$

is an isomorphism. In particular, if B is basis for $NF_S/\langle E \rangle_k$, then it is a basis for M.

In other words, provided we can prove convergence, finding a basis for M reduces to finding a basis for $NF_S/\langle E \rangle_k$. In the example where $M = {}^{\circ}C(\mu, \nu)$, the module of normal forms NF_S is linearly generated by diagrams where each strand has at most one critical point; in the module $NF_S/\langle E \rangle_k$, we view theses diagrams up to the superinterchange law. It is not too difficult to see that any set *B* of representatives for the interchange law constitutes a basis of $NF_S/\langle E \rangle_k$. Provided S^+ is convergent, the set *B* provides a basis for ${}^{\circ}C(\mu, \nu)$ by Rewriting fact A.

The remaining of the theory serves the sole purpose of showing convergence. Termination is often the easy part²: in our running example $^{\circ}S$, the modulo data $^{\circ}E$ preserves the number of generating 2-cells, while rewriting steps in $^{\circ}R$ strictly decrease it. Hence, we focus on showing confluence.

1.2.3 Tamed congruence

Recall the linear Newmann's lemma (Lemma 1.1). An analogous statement can be given in the modulo setting. However, we shall need a more general version, based on the notion of tamed congruence (see (1) in the beginning of the introduction and Subsection 3.2.1 in the text):

Rewriting fact B (TAMED LINEAR NEWMANN'S LEMMA 3.43). Let S = (B; R, E) be a linear RSM and \succ a preorder on B satisfying some conditions. If \succ is terminating and every monomial local S^+ -branching is \succ -tamely S^{st} -congruent, then S^+ is convergent.

In practice, the preorder \succ is the terminating order used to show termination; in our running example °S, the preorder \succ compares the number of generating 2-cells.

Why tamed congruence? This is explained in the next subsection (see also Subsections 1.2.5 and 1.2.7).

²Although not always: see [GM18] for an approach to termination.

1.2.4 Positivity and contextualization

A hom-basis is a collection of bases, one for each hom-space. In principle, linear rewriting modulo could be applied to each hom-space at a time to find a hom-basis. Of course, in practice we would like to leverage the higher structure, that is, the fact that these hom-spaces are related with each other (and themselves) via contexts (see Subsection 1.1.3). More precisely, if (f, g) is a branching and Γ is a context, we would like to say that if (f', g') is a confluence for (f, g), then $(\Gamma[f'], \Gamma[g'])$ is a confluence for $(\Gamma[f], \Gamma[g])$. The fact is essentially trivial for monoids; see Subsection 1.1.1. This is also true for associative algebras and for commutative algebras (i.e. associative algebras modulo commutativity). However, *this is not true in general*. Indeed, positivity is in general *not* preserved by contexts:

WARNING: In general, the contextualization of a positive rewriting step needs not be positive.

Consider our running example °S and denote $\sim_{\circ E}$ the equivalence relation induced by °E, that is, equivalence via the superinterchange law; we call it °E-congruence. Then:

 $\bigcirc | | \not\sim_{\circ \mathsf{E}} | | \bigcirc \text{ while } \bigcirc | \sim_{\circ \mathsf{E}} \bigcirc \bigcirc.$

More formally, we have two 2-cells $v, w \in {}^{\circ}\mathsf{P}_2^l$ and a context Γ such that $v \not\sim_{\circ\mathsf{E}} w$, while $\Gamma[v] \sim_{\circ\mathsf{E}} \Gamma[w]$. In other words, *contextualization does not act freely*. In particular, if $\lambda r + v$ is a rewriting step, it may be that $\Gamma[s(r)]$ is ${}^{\circ}\mathsf{E}$ -congruent (up to sign) to a monomial in the linear decomposition of $\Gamma[v]$, even if s(r) is not ${}^{\circ}\mathsf{E}$ -congruent (up to sign) to a monomial in the linear decomposition of v. That is, it may be that $\lambda\Gamma[r] + \Gamma[v]$ is not positive, even if $\lambda r + v$ is positive.

This gives a theoretical motivation for using 2-sesquicategories as a foundation of our theory, even when rewriting in 2-categories: contextualization *is* free on 2-sesquicategories, and it is the (explicit) addition of a modulo rule that (may) prevents freeness. Note that contextualization is free both for associative algebras and associative algebras modulo commutativity; this perhaps explains why the problem had not been recognised before (to the author's knowledge).

However, while the contextualization of a positive confluence needs not be a positive confluence, it is a tamed congruence:

Rewriting fact C (CONTEXTUALIZATION LEMMA 3.62). Let S = (R, E) be a higher linear rewriting system modulo, that is, a pair of linear 3-sesquipolygraphs satisfying some conditions. Let (f, g) be a monomial local S^+ -branching and Γ a context. If (f, g) is S^+ -confluent, then $\Gamma[f, g]$ is \succ -tamely S^{st} -congruent.

1.2.5 Independent branchings

An *independent branching* (Subsection 3.4.3) is a branching where the sources of the two branchings "do not overlap". This is the higher analogue of an independent branching for monoids; see Subsection 1.1.1. The confluence of independent branchings is trivial for monoids, and one may expect the same for diagrammatic algebras. However, this is not the case! The reason is the same as the one explained in Subsection 1.2.4, and can be solved using tamed congruence:

Rewriting fact D (Lemma 3.65). Let S = (R, E) be a higher linear rewriting system modulo (see Rewriting fact C) and \succ a preorder satisfying some conditions. Every independent S⁺-branching is \succ -tamely Sst-congruent.

1.2.6 How to classify branchings?

The arguments of this subsection are abstract; thus we fix an abstract RSM $S = (X; \mathbb{R}, \mathbb{E})$. Recall that the equivalence relation induced by \mathbb{E} on X is called \mathbb{E} -congruence; subsequently, an unoriented composition of relations in \mathbb{E} is called an \mathbb{E} -congruence.

Working modulo induces a huge number of branchings: indeed, in between the two branches of a branching, one can apply an arbitrary number of modulo relations. To classify branchings, we should not only understand elements of X modulo, but also rewriting sequences modulo. We say that two rewriting sequences f and f' are \mathbb{E} -congruent if there exists \mathbb{E} -congruences e_s and e_t such that:



Two branchings (f, g) and (f', g') are *branchwise* \mathbb{E} -congruent if f (resp. g) is \mathbb{E} -congruent to f' (resp. g'). It is not difficult to see that:

Rewriting fact E (BRANCHWISE CONFLUENCE LEMMA 3.17). Let $\mathbb{S} = (\mathbb{R}, \mathbb{E})$ be an abstract RSM. If (f, g) and (f', g') are branchwise \mathbb{E} -congruent branchings, then (f, g) is confluent if and only if (f', g') is.

Hence, the study of confluence can be done up to branchwise \mathbb{E} -congruence. Consider again our running example °S. What does it mean to understand the rewriting steps in °R up to °E-congruence? The interchange law preserves the set of generating 2-cells; more precisely, any sequence of interchangers between two diagrams *s* and *t* induces a canonical bijection between the generating 2-cells in *s* and the generating 2-cells in *t*. Hence it makes sense to speak about *the* cap and *the* cup associated with a rewriting step in °S:

Lemma 1.2 (characterization of rewriting steps in $^{\circ}$ S). If two rewritings steps in $^{\circ}$ S apply to the same cup and cap, then they are $^{\circ}$ E-congruent.

To show Lemma 1.2, we decompose °E-congruences as compositions of interchangers and proceed inductively. Indeed, in a Gray category, interchangers come with naturality axioms, such as the following ($A: \phi \rightarrow \phi'$ is a 3-cell and β is a 2-cell):



The two arrows labelled "A" are contextualizations of the 3-cell A; intuitively, we apply A "locally on ϕ ". From the rewriting point of view, this naturality axiom gives an \mathbb{E} -congruence between two rewriting steps.

The situation is similar for other modulo data; see Section 4 for an example with pivotality. As a rule of thumb, one should always include naturality axioms in the modulo. For instance, when working modulo pivotality, one should include the pivotal axioms associated to all generators, and not only some of them.

Using the characterization given in Lemma 1.2, topological arguments can be used to deduce the following:

Lemma 1.3 (classification of branchings in °S). Every monomial local °S-branching is branchwise °E-congruent either to an independent branching, or to a contextualization of one the following (trivially positively confluent) branchings:

Combining Lemma 1.3 with Rewriting fact C (CONTEXTUALIZATION LEMMA 3.62), Rewriting fact D (tamed congruence of independent branchings), Rewriting fact E (BRANCHWISE CONFLUENCE LEMMA 3.17) and Rewriting fact B (TAMED LINEAR NEWMANN'S LEMMA 3.43) implies confluence of °S. Hence, °S is convergent.

Because confluence reduces to confluence of the two branchings in Lemma 1.3, we call the later *critical branchings*. Requiring their confluence enforces that the product of the signs in the zigzag relations $^{\circ}R_3$ is a minus sign; the choice given is one of the two possible conventions.

1.2.7 Further rewriting techniques

Our running example °S was very elementary. For instance, it is sometimes not possible to give a satisfactory characterization of rewriting steps solely based on \mathbb{E} -congruence; this is the case of graded \mathfrak{gl}_2 -foams considered in Section 4. Indeed, we may wish to replace "branchwise \mathbb{E} -congruence" in Rewriting fact E by "branchwise \mathbb{S} -confluence" (see Subsection 3.2.3 for a precise definition). Unfortunately, this modified version doesn't hold. However, it holds if we consider tamed congruence instead:

Rewriting fact F (BRANCHWISE TAMED CONGRUENCE LEMMA 3.19). Let $\mathbb{S} = (\mathbb{R}, \mathbb{E})$ be an abstract RSM and \succ a preorder on X satisfying some conditions. If (f, g) and (f', g') are branchwise \succ -tamely congruent branchings, then (f, g) is \succ -tamely congruent if and only if (f', g') is.

This comes down to the fact that tamed congruence is transitive: if f_1 , f_2 and f_3 are three rewriting sequences such that f_1 is tamely congruent to f_2 and f_2 is tamely congruent to f_3 , then f_1 is tamely congruent to f_3 (assuming the hypotheses of Rewriting fact F). This gives an abstract reason for considering tamed congruence over confluence.

We may also wish to restrict rewriting steps depending on the context. For instance, assume we add the following 3-cell to ${}^{\circ}R_3$ in our running example ${}^{\circ}S$:



If we allow all contextualizations $\Gamma[\mathbf{r}]$, our rewriting system modulo does not terminate:

This is a common issue in diagrammatic algebras, including this precise example [All18a] and graded \mathfrak{gl}_2 -foams (Section 4). To solve it, we consider context-dependent rewriting system modulo, described in Definition 3.56.

Yet another tool is the INDEPENDENT REWRITING LEMMA 3.67; we refer to the main text for its description.

1.2.8 Comparison with the literature

Our theory has three main features: it can be (*weakly*, i.e. *Gray*) *higher*, it can be *linear* and it can be *modulo*. In other words, it allows to rewrite modulo in 2-sesquicategories and in linear 2-sesquicategories. We call it (*linear*) *Gray rewriting modulo*.

We now compare it with the existing literature, using the terminology *strict* to emphasize that a rewriting theory does not allow modulo:

- Strict rewriting in 2-sesquicategories, and in particular strict rewriting containing interchangers, recovers Forest and Mimram's Gray rewriting theory [FM22].
- Strict rewriting in 2-categories (see Subsection 1.1.3) is rewriting in 2-sesquicategories modulo interchangers.
- Strict rewriting in linear 2-categories is rewriting in linear 2-sesquicategories modulo interchangers. Alleaume [All18a] also developed a rewriting theory for linear 2-categories.
- Rewriting modulo in linear 2-categories is rewriting modulo in linear 2-sesquicategories with a modulo containing interchangers. Dupont [Dup22] also developed a rewriting modulo theory for linear 2-categories.
- Rewriting modulo in graded-2-categories is rewriting modulo in linear 2-sesquicategories with a modulo containing graded interchangers. Dupont, Ebert and Lauda [DEL21] also developed a rewriting modulo theory for super-2-categories.
- Although we do not describe it explicitly, our work can be used to define rewriting modulo in monoids and associative algebras; in particular, strict rewriting in associative algebras. This latter case recovers the work of Guiraud, Hoffbeck and Malbos [GHM19].

In that sense, our work is an extension of [FM18] to linear rewriting and rewriting modulo; or an extension of [GHM19] to (weakly) higher rewriting and rewriting modulo. We also have important debts toward [All18a; Dup21; Dup22]: the striking connection between rewriting theory and diagrammatic algebras in the first place, and the importance of rewriting modulo. However, our work is distinct in many respects, both theoretically and practically, and even when restricted to their respective setting. Let us highlight two of them:

- Theoretically, our theory is based on tamed congruence. Its importance, even for strict rewriting in linear 2-categories, is highlighted in Subsection 1.2.4.
- Practically, we spend a fair amount of time formalizing how to classify branchings modulo (see Subsection 1.2.6). The ability to confidently classify branchings is a cornerstone of the theory, as if some forgotten critical branchings are not confluent, it prevents confluence altogether.

We stress further differences in the text; see Remarks 3.24 and 3.57 and Footnotes 17, 21 and 23.

Remark 1.4. Our theory works over a commutative ring \Bbbk , and not just a field, although scalars appearing in the modulo must be invertible. However, we stress that the modulo data must be monomial-invertible. In other words, while it incorporates multiplication by an invertible scalar, a relation such as $b \sim_{\texttt{E}} b_1 + b_2$ for distinct monomials b, b_1 and b_2 is not a valid modulo rule (see also Footnote 17 and the related discussion). Extending our work to this more general setting is an important open problem.

1.2.9 Non-degeneracy of graded \mathfrak{gl}_2 -foams

In Section 4 we prove a basis theorem for the graded-2-category \mathbf{GFoam}_d of graded \mathfrak{gl}_2 -foams (Theorem 4.10). This uses the full-strength of linear Gray rewriting modulo, including context-dependent termination and rewriting with a non-coherent modulo. The proof illustrates all the general techniques discussed above.

1.3 Perspectives

A rewriting approach can be heavy: finding a practical convergent rewriting system may require a lot of trial and error, and classifying critical branchings can be laborious. Moreover:

OBSTRUCTION TO REWRITING THEORY: there is no guaranty that a convergent rewriting system exists, and if so, that it is sufficiently reasonable to be used in practice.

However, once established the rewriting perspective provides a rich understanding of the combinatorial structure of the presentation. We give some future directions of research below. As more and more examples are studied, we hope that the theory of linear Gray rewriting modulo will evolve into a standard set of tools, both easy-to-master and powerful, fostering the exploration of still finer higher symmetries.

1.3.1 Examples, examples, examples

Rewriting theory could be applied to many diagrammatic algebras. This would give new (and sometimes only) proofs of their respective hom-basis theorems, in an intrinsic and algorithmic fashion.

1.3.2 Computer implementation

Assuming one has a candidate convergent presentation (see the obstruction above), the rewriting approach is relatively straightforward, at least in principle: enumerate critical branchings and show that each of them is confluent. We expect both of these processes to be implementable on a computer, at least when working modulo (graded) interchangers. This should vastly expend what is meant by a "sufficient reasonable" convergent presentation.

This direction of research should relate with current developments in applied category theory, such as the graphical proof-assistants *Globular* and *homotopy.io* [BKV18; BV17; Dor23; RV19].

1.3.3 A Buchberger's algorithm for diagrammatic algebras?

In some cases, even finding a convergent presentation can be made automatic. For commutative algebras, this is known as the Buchberger's algorithm; in generality, this is known as the Knuth–Bendix completion. In fact, the Buchberger's algorithm itself can be optimized using machine learning [Pei21]. Results along those lines would further help the systematic study of diagrammatic algebras.

1.3.4 Deformation theory

Typically, once a rewriting approach is established for a diagrammatic algebra, it also applies to all of its "deformations", in a loose sense. Consider the rewriting proof given in Subsection 1.2 for our running example °S: the sign in the relations °R was only relevant once we checked confluence of critical branchings (Lemma 1.3). Similarly, the rewriting proof given in Section 4 for graded \mathfrak{gl}_2 -foams tells us how scalars can be chosen in order to get the same basis theorem, leading to precisely two choices, the super-2-category **GFoam**_d and **GFoam**'_d (Subsection 4.5).

Similar ideas have appeared before in the literature [Bar24; Eli22], although not explicitly using rewriting techniques. Making the relationship between rewriting theory and deformation theory, especially in the context of diagrammatic algebras, is an interesting direction of research; see also [Sch24, section 6.6.4] for related comments.

1.3.5 Higher structures

The application of rewriting theory goes beyond finding bases. By considering how rewriting sequences form unoriented cycles, one can extract an understanding of relations *between* relations, or *coherence data*; see Subsection 3.1. Critical branchings of a convergent rewriting system lead to an explicit description of this coherence data. For instance, coherent presentations of Artin monoids can be obtained via rewriting theory, leading to a new proof of Deligne's theorem on categorical actions of Artin monoids [GGM15]. In the linear setting, this coherence data is known as *syzygies*; applications include computation of homological invariants or study of Koszulness [GHM19]. In this work, we only discuss coherence in Subsection 3.1; however, we expect that the coherence results of [FM18; GHM19] can be adapted to our setting. We leave this for future work.

In the recent years, stable homotopy theory and ∞ -categories have become more and more prevalent in link homologies and higher representation theory [Dra+24; HKK16; Liu+24; Liu24; LLS20; LS14a; MR20; SSS20]. At present however, the complexity of the ∞ -setting remains a major obstacle to exploration beyond the simplest cases. Explicit presentations of the coherence data, obtained via rewriting methods, could be an important step forward.

2 Linear Gray polygraphs

This section introduces the necessary categorical structures to present graded-2-categories and define their rewriting theory. This can be understood as a linear analogue to the work of Forest and Mimram [FM22, section 2 and 3]; equivalently, as a generalization of the work of Alleaume on linear n-polygraphs [All18a] to allow weak interchangers. For an introduction to the ideas of this section, see Subsections 1.1.2, 1.1.6 and 1.2.1.

The notion of an *n*-sesquicategory was first defined by Street [Str96] in the case n = 2. The general case was independently introduced by Forest–Mimram [FM22] (following the general theory of Weber [Web13]) under the name of "*n*-precategories" and by Araújo [Ara22, section 1.6] under the name of "*n*-sesquicategories". Although we shall follow Forest and Mimram's presentation, we choose Araújo's terminology to avoid confusion with already existing notions of *n*-precategories in the literature. Enriched category theory provides yet another defining approach to *n*-sesquicategories; see [FM22, section 2.4 and Appendix A].

To motivate the formal definitions, we start with an example in Subsection 2.1. We expect it to be sufficient for the impatient reader. Subsection 2.2 review the notions of n-sesquicategories and n-sesquipolygraphs, following [FM22] (which they respectively call "n-prepolygraphs" and

"*n*-precategories"). We then introduce their linear analogue in Subsection 2.3. Finally, Subsection 2.4 defines Gray polygraphs (still following [FM22]) and linear Gray polygraphs, the latter defining presentations for graded-2-categories.

In this section and as throughout the paper, every categorical structure is assumed to be small.

Notation 2.1. As much as possible, we use calligraphic fonts (e.g. C) for categories, and sans serif fonts (e.g. P) for their presentations. If necessary, blackboard fonts (e.g. P) refer to purely set-theoretic concepts, and typewriter fonts (e.g. P) to purely linear concepts.

2.1 A summary via example

We give a more formal definition of $^{\circ}P$, the \mathbb{Z} -linear 3-sesquipolygraph of superadjunction given in Subsection 1.2.1, introducing further nomenclature along the way. For simplicity, we don't discuss the parity associated to the 2-generators in $^{\circ}P$, and grading in general.

To present higher categories, one needs to provide generators for each categorical level; in this framework, relations are simply the generators at the highest categorical level. In the case of °P, the first three levels are the sets °P₀ = {*}, °P₁ = { | } and °P₂ = { \land, \bigcup }. Each level °P_{*n*+1} comes equipped with a source and target map into the *free n-sesquicategory* (°P_{*n*})* (Subsection 2.2.4) generated by the previous level °P_{*n*}:

$$(^{\circ}\mathsf{P}_{n})^{*} \xleftarrow{s_{n}}{t_{n}} ^{\circ}\mathsf{P}_{n+1}.$$

For objects, the free 0-sesquicategory is the set itself: $({}^{\circ}\mathsf{P}_0)^* = {}^{\circ}\mathsf{P}_0$. In our case, we have ${}^{\circ}\mathsf{P}_0 = \{*\}$ and the maps s_0, t_0 are the trivial maps. The free 1-sesquicategory is given by formal horizontal juxtaposition of 1-generators:

$$(^{\circ}\mathsf{P}_{1})^{*} = \left\{ \underbrace{|\cdots|}_{n} \mid n \in \mathbb{N} \right\}.$$

(If there were more than one object, we should ask the horizontal juxtaposition to be compatible with the 0-source s_0 and the 0-target t_0). The maps s_1, t_1 are the obvious ones, reading s_1 on the bottom and t_1 on the top, respectively. The data ${}^{\circ}\mathsf{P}_{\leq 2} = ({}^{\circ}\mathsf{P}_0, {}^{\circ}\mathsf{P}_1, {}^{\circ}\mathsf{P}_2)$, together with their source and target maps, form the data of a 2-sesquipolygraph (Subsection 2.2.5):



To define the highest level consisting of the defining relations, we must describe the *free* \mathbb{Z} -*linear 2-sesquicategory* $({}^{\circ}\mathsf{P}_2)^l$ generated by ${}^{\circ}\mathsf{P}_{\leq 2}$ (Subsection 2.3.2). First, juxtapose horizontally the 2-generators \bigcap and \bigcup with elements of $({}^{\circ}\mathsf{P}_1)^*$, both on the left and on the right. This process is called *whiskering*. Then, juxtapose vertically an arbitrary number of these whiskered 2-generators, gluing along 1-source s_1 and 1-target t_1 . This defines the 2-morphisms of $({}^{\circ}\mathsf{P}_2)^*$, the free 2-sesquicategory generated by ${}^{\circ}\mathsf{P}_{\leq 2}$. Note that we did not mod out by the interchange relation: indeed, $({}^{\circ}\mathsf{P}_2)^*$ is a 2-sesquicategory, and not a strict 2-category. Finally, we denote $({}^{\circ}\mathsf{P}_2)^l$ the linearization of $({}^{\circ}\mathsf{P}_2)^*$, that is, the linear 2-sesquicategory where each hom-space is the free \mathbb{Z} -module generated by the corresponding hom-set in $({}^{\circ}\mathsf{P}_2)^*$. The set ${}^{\circ}\mathsf{P}_3$ is equipped with source and target maps s_2 and t_2 into $({}^{\circ}\mathsf{P}_2)^l$:



Elements of ${}^{\circ}\mathsf{P}_3$ are pictured as two-frame movies, reading s_2 on the left and t_2 on the right.

In fact, the data of °P can be equivalently described as a *scalar 3-sesquipolygraph* scl(°P) (Subsection 2.2.7). This is because the image of s_2 lies in (°P₂)*, and the image of t_2 lies in $\mathbb{Z}(^{\circ}P_2)^*$ (i.e. elements of (°P₂)* times scalars); we say that °P is a *monomial linear 3-sesquipolygraph* (Subsection 2.3.4). In a general linear *n*-sesquipolygraph, source and target of generating 3-cells can be any linear combinations of 2-cells.

We define $scl(^{\circ}\mathsf{P})_{\leq 2} = {}^{\circ}\mathsf{P}_{\leq 2}$ and

$$\operatorname{scl}(^{\circ}\mathsf{P})_{3} = \left\{ \bigcup \Longrightarrow \left|, \bigcup \rightrightarrows^{-1} \right|, \left| \stackrel{\bullet}{\Rightarrow} \right|_{\beta} \left| \stackrel{(-1)^{p(\alpha)p(\beta)}}{\Rightarrow} \right|_{\alpha} \left| \stackrel{\bullet}{\beta} \right|_{\beta} \right\}_{\text{for all } \alpha, \beta \in (^{\circ}\mathsf{P}_{2})^{*}} \left| \stackrel{\circ}{\beta} \right|_{\beta} \right\}$$

The set $scl(^{\circ}\mathsf{P})_3$ is equipped with source and target maps s_2 and t_2 into $(^{\circ}\mathsf{P}_2)^*$. Moreover, each element is equipped with a scalar, pictured here on top of the arrow; formally, a map $scl: ^{\circ}\mathsf{P}_3 \to \mathbb{Z}$.

$$\operatorname{scl}(^{\circ}\mathsf{P}) \coloneqq \bigcup_{i=0}^{\circ} \mathbb{P}_{0}^{*} \xrightarrow{\circ} \mathbb{P}_{1}^{*} \xrightarrow{\circ} \mathbb{P}_{2}^{*} \xrightarrow{\mathbb{Z}}_{s_{2}}^{*} \operatorname{scl}$$
$$\bigcup_{i=0}^{\circ} \mathbb{P}_{0} \xrightarrow{\circ} \mathbb{P}_{1} \xrightarrow{\circ} \mathbb{P}_{2} \xrightarrow{\circ} \mathbb{P}_{2}^{*} \operatorname{scl}(^{\circ}\mathsf{P}_{3})$$

The last family of 3-generators in ${}^{\circ}\mathsf{P}_3$ and $\mathrm{scl}({}^{\circ}\mathsf{P})_3$ are the interchange generators on ${}^{\circ}\mathsf{P}_{\leq 2}$. They from the 3-sesquipolygraph of interchangers (Subsection 2.4.1), denoted ${}^{\circ}\mathsf{P}_{\leq 2}$ Gray; it can be viewed either as a monomial linear 3-sesquipolygraph or as a scalar 3-sesquipolygraph. Because ${}^{\circ}\mathsf{P}$ contains its own 3-sesquipolygraph of interchangers, it is a *linear Gray polygraph* (Subsection 2.4.2).

The set ${}^{\circ}\mathsf{P}_3$ describes generating relations; to describe *all* relations, we must define the free 3-sesquicategory $({}^{\circ}\mathsf{P}_3)^*$ generated by ${}^{\circ}\mathsf{P}_3$. First, we revisit our terminology: we call horizontal juxtaposition the *0-composition* (as we glue along objects) and vertical juxtaposition the *1-composition* (as we glue along 1-morphisms). We define the *contextualization* of a 3-generator as first 0-whiskering with elements of $({}^{\circ}\mathsf{P}_1)^*$, and then 1-whiskering with elements of $({}^{\circ}\mathsf{P}_2)^*$. This can be pictured as the relevant 0-compositions and 1-compositions on its source and target:

Here $A \in {}^{\circ}\mathsf{P}_3$ is a 3-generator and Γ is a *context*; that is, the data of 1-cells and 2-cells "surrounding A" via 0- and 1-whiskerings. We write $\operatorname{Cont}({}^{\circ}\mathsf{P}_3)$ the set of contextualized 3-generators. Finally, a generic 3-morphism in ${}^{\circ}\mathsf{P}_3^*$ is a 2-composition of contextualized 3-generators, gluing along 2-source and 2-target. This is pictured as a multi-frame movie.

3-morphisms in $({}^{\circ}\mathsf{P}_3)^*$ have the following structure: a 2-composition (pictured as a composition of movies) and actions of respectively $({}^{\circ}\mathsf{P}_1)^*$ and $({}^{\circ}\mathsf{P}_2)^*$ via the 0- and 1-whiskerings (pictured as horizontal and vertical juxtapositions). In general, the (k+1)-cells of an *n*-sesquicategory (Subsection 2.2.2) have a *k*-composition gluing along *k*-cells, and actions of lower cells via whiskerings. *Contextualization* (Subsection 2.2.3) constitutes the combined action of all whiskerings. Given a globular extension P (Subsection 2.2.1), the *n*-cells of the *free n*-sesquicategory P* (Subsection 2.2.4) are given by formal (n - 1)-compositions of contextualized *n*-generators Cont(P).

2.2 *n*-sesquicategories and their presentations

We review the notion of n-sesquicategories and their presentations, following the presentation given in [FM22]. The last subsections introduce graded and scalar variants of n-sesquicategories and their presentations.

2.2.1 *n*-globular sets

Let $n \in \mathbb{N} = \{0, 1, ...\}$. An *n*-globular set C is a diagram of sets and functions as follows:

$$\mathcal{C}_0 \xleftarrow{s_0}{t_0} \mathcal{C}_1 \xleftarrow{s_1}{t_1} \dots \xleftarrow{s_{n-1}}{t_{n-1}} \mathcal{C}_n$$

such that $s_j \circ s_{j+1} = s_j \circ t_{j+1}$ and $t_j \circ s_{j+1} = t_j \circ t_{j+1}$ for each $0 \le j < n$. The maps s_j and t_j are respectively called *source maps* and *target maps*. An element $u \in C_j$ is called a *j-cell*, with $s_{j-1}(u)$ and $t_{j-1}(u)$ respectively its *source* and *target*, which we sometimes simply denote by s(u) and t(u). We refer to j as the *dimension* of u. For $0 \le i < j$, we define the *i-source* of u to be

 $s_i(u) = s_i \circ (\text{any suitable composition of source and target maps})(u),$

where the choice in the bracket does not matter thanks to the properties of source and target maps. Note that the subscript indicates that $s_i(u)$ is an *i*-cell. We define the *i*-target $t_i(u)$ similarly. A morphism of globular sets $f : C \to D$ is a family $f_i : C_i \to D_i$ of functions that commute with the source and target maps. It is an isomorphism if each function f_i is a bijection.

Given an *n*-globular set C, a 0-sphere is an ordered pair of 0-cells in C. For $0 < i \le n$, an *i*-sphere is an ordered pair (f, g) of *i*-cells such that s(f) = s(g) and t(f) = t(g). For $0 \le i < k \le n$ and an *i*-sphere (f, g), we set

$$\mathcal{C}_k(f,g) = \{ u \in \mathcal{C}_k \mid s_i(u) = f, t_i(u) = g \}.$$

The source and target maps restrict to maps $C_k(f,g) \xleftarrow{s_k}{t_k} C_{k+1}(f,g)$.

A globular extension of C is an (n + 1)-globular set P such that $\mathsf{P}_k = \mathcal{C}_k$ for $0 \le k \le n$:

$$\underbrace{\begin{array}{c} \mathcal{C}_0 \xleftarrow{s_0}{t_0} \mathcal{C}_1 \xleftarrow{s_1}{t_1} \dots \xleftarrow{s_{n-1}}{t_{n-1}} \mathcal{C}_n \xleftarrow{s_n}{t_n} \mathsf{P}_{n+1} \\ \mathcal{C} \end{array}}_{\mathcal{C}}$$

For $k \in \mathbb{N}$ such that $0 \le k \le n$, the *k*-restriction of \mathcal{C} is the subglobular set

$$\mathcal{C}_{\leq k} \coloneqq \mathcal{C}_0 \xleftarrow{s_0}{t_0} \mathcal{C}_1 \xleftarrow{s_1}{t_1} \dots \xleftarrow{s_{k-1}}{t_{k-1}} \mathcal{C}_k$$

2.2.2 *n*-sesquicategories

We review the definition of an n-sesquicategory (and n-sesquifunctor) introduced in [FM22, section 2.2] (with minor changes to the presentation).

An *n*-sesquicategory is the data of an *n*-globular set C together with

- identity functions $id^k : C_{k-1} \to C_k$, for $0 < k \le n$,
- composition functions $\star_{k,k} : C_k \times_{k-1} C_k \to C_k$ for $0 < k \le n$,

• left- and right-whisker functions $\star_{i,k} : C_i \times_{i-1} C_k \to C_k$ and $\star_{k,i} : C_k \times_{i-1} C_i \to C_k$ for $0 < i < k \le n$,

satisfying the axioms (i) and (ii) below. Hereabove we abbreviated \times_{C_k} with \times_k . We use similar notations for composition and whiskers, but one cannot confuse one for the other as they have different domain. In fact, this choice of notation emphasizes that whisker functions should be thought of compositions with identities of cells of lower dimension. As such, composition and whiskers have similar properties, and it is sometimes useful to consider $\star_{k,l}$ for all $0 \le k, l \le n$.

Note that for $0 \le k, l \le n$, the function $\star_{k,l}$ is defined on $(u, v) \in C_k \times C_l$ if and only if $s_i(u) = t_i(v)$, where $i = \min(k, l) - 1$. In that case, we say that u and v are *i-composable*, and we write $u \star_i v$, or even $u \star v$, for $u \star_{k,l} v$. While the notation $u \star_{k,l} v$ emphasizes the dimension of the respective cells, the notation $u \star_i v$ emphasizes the dimension of the compatibility condition. Also, for u an (i - 1)-cell we write id_u instead of $\mathrm{id}^i(u)$. The axioms of an n-sesquicategory are as follows:

(i) for $0 < k \le n$, with $f \in \mathcal{C}_{k-1}$ and $\alpha, \beta, \gamma \in \mathcal{C}_k$ suitably k-composable:

$$t_{k-1}(\mathrm{id}_f) = f = s_{k-1}(\mathrm{id}_f)$$
$$s_{k-1}(\alpha \star_{k,k} \beta) = s_{k-1}(\beta) \qquad t_{k-1}(\alpha \star_{k,k} \beta) = t_{k-1}(\alpha)$$
$$\mathrm{id}_{t(\alpha)} \star_{k,k} \alpha = \alpha = \alpha \star_{k,k} \mathrm{id}_{s(\alpha)}$$
$$\alpha \star_{k,k} (\beta \star_{k,k} \gamma) = (\alpha \star_{k,k} \beta) \star_{k,k} \gamma$$

(ii) for $0 < i < k, k' \leq n$, with $x \in C_{i-1}$, $f, g \in C_i, \phi \in C_{k-1}$ and $A \in C_k$, $B \in C_{k'}$ suitably composable:

$$f \star_{i,k} (g \star_{i,k} A) = (f \star_{i,i} g) \star_{i,k} A \qquad (A \star_{k,i} f) \star_{k,i} g = A \star_{k,i} (f \star_{i,i} g)$$

$$id_{x} \star_{i,i} A = A \qquad A \star_{i,i} id_{x} = A$$

$$(f \star_{i,k} A) \star_{k,i} g = f \star_{i,k} (A \star_{k,i} g)$$

$$s_{k-1}(f \star_{i,k} A) = f \star_{i,k-1} s_{k-1}(A) \qquad s_{k-1}(A \star_{k,i} f) = s_{k-1}(A) \star_{k-1,i} f$$

$$t_{k-1}(f \star_{i,k} A) = f \star_{i,k-1} t_{k-1}(A) \qquad t_{k-1}(A \star_{k,i} f) = t_{k-1}(A) \star_{k-1,i} f$$

$$f \star_{i,k} id_{\phi} = id_{f \star_{i,k-1} \phi} \qquad id_{\phi} \star_{k,i} f = id_{\phi \star_{k-1,i} f}$$

$$f \star_{i,\max(k,k')} (A \star_{k,k'} B) = (f \star_{i,k} A) \star_{k,k'} (f \star_{i,k'} B)$$

$$(A \star_{k,k'} B) \star_{\max(k,k'),i} f = (A \star_{k,i} f) \star_{k,k'} (B \star_{k',i} f)$$

An *n*-sesquifunctor between two *n*-sesquicategories is a morphism between the underlying globular sets, preserving identities and compositions as expected. It is an isomorphism if the underlying morphism of globular sets is an isomorphism. This ends the definition of an *n*-sesquicategory and of an *n*-sesquifunctor. \diamond

Remark 2.2 (low-dimensional cases). A 0-sesquicategory is simply a set, and a 1-sesquicategory a category. The distinction with strict *n*-categories only appears when $n \ge 2$. Contrary to strict *n*-categories, 2-cells of an *n*-sesquicategories cannot be horizontally composed. Instead, they can be whiskered with 1-cells, understood as acting as identity 2-cells. In particular, in a 3-sesquicategory there is a priori no relationship between the two sides of the (2-dimensional) interchange law;

see Fig. 1.3. We use string diagrammatics to picture 2-cells of an *n*-sesquicategory. String diagrams are equipped with a Morse function on the generators, as for graded-2-categories. 3-cells are then pictured as *movies*, i.e. paths of 2-cells, and 4-cells as *movie moves*, i.e. paths of paths of 2-cells.

Remark 2.3. With the single-index notation, the relations above become:

(i) for $0 < k \le n$, with $f \in \mathcal{C}_{k-1}$ and $\alpha, \beta, \gamma \in \mathcal{C}_k$ suitably k-composable:

$$t_{k-1}(\mathrm{id}_f) = f = s_{k-1}(\mathrm{id}_f)$$
$$s_{k-1}(\alpha \star_{k-1} \beta) = s_{k-1}(\beta) \qquad t_{k-1}(\alpha \star_{k-1} \beta) = t_{k-1}(\alpha)$$
$$\mathrm{id}_{t(\alpha)} \star_{k-1} \alpha = \alpha = \alpha \star_{k-1} \mathrm{id}_{s(\alpha)}$$
$$\alpha \star_{k-1} (\beta \star_{k-1} \gamma) = (\alpha \star_{k-1} \beta) \star_{k-1} \gamma$$

(ii) for $0 < i < k, k' \leq n$, with $x \in C_{i-1}$, $f, g \in C_i$, $\phi \in C_{k-1}$ and $A \in C_k$, $B \in C_{k'}$ suitably composable:

$$f \star_{i-1} (g \star_{i-1} A) = (f \star_{i-1} g) \star_{i-1} A \quad (A \star_{i-1} f) \star_{i-1} g = A \star_{i-1} (f \star_{i-1} g)$$

$$id_x \star_{i-1} A = A \qquad A \star_{i-1} id_x = A$$

$$(f \star_{i-1} A) \star_{i-1} g = f \star_{i-1} (A \star_{i-1} g)$$

$$s_{k-1}(f \star_{i-1} A) = f \star_{i-1} s_{k-1}(A) \qquad s_{k-1}(A \star_{i-1} f) = s_{k-1}(A) \star_{i-1} f$$

$$t_{k-1}(f \star_{i-1} A) = f \star_{i-1} t_{k-1}(A) \qquad t_{k-1}(A \star_{i-1} f) = t_{k-1}(A) \star_{i-1} f$$

$$f \star_{i-1} id_{\phi} = id_{f \star_{i-1} \phi} \qquad id_{\phi} \star_{i-1} f = id_{\phi \star_{i-1} f}$$

$$f \star_{i-1} (A \star_{\min(k,k')-1} B) = (f \star_{i-1} A) \star_{\min(k,k')-1} (f \star_{i-1} B)$$

$$(A \star_{\min(k,k')-1} B) \star_{i-1} f = (A \star_{i-1} f) \star_{\min(k,k')-1} (B \star_{i-1} f)$$

Remark 2.4. Note that if we let $i \in \mathbb{N}$ be such that $0 \le i < n$ and (f, g) be an (i - 1)-sphere³ in C, the identity and composition functions restrict as follows:

$$\operatorname{id}^{i+1} \colon \mathcal{C}_i(f,g) \to \mathcal{C}_{i+1}(f,g)$$
$$\star_{i+1,i+1} \colon \mathcal{C}_{i+1}(f,g) \times_i \mathcal{C}_{i+1}(f,g) \to \mathcal{C}_{i+1}(f,g)$$

It follows from the axioms of an *n*-sesquicate gory that, equipped with the maps id^{i+1} and $\star_{i+1,i+1}$, the 1-globular set

$$\mathcal{C}_i(f,g) \xleftarrow{s_i \\ t_i} \mathcal{C}_{i+1}(f,g)$$

defines a 1-category.

2.2.3 Contexts

Let C be an *n*-sesquicategory and let $\Box = (f, g)$ an (i - 1)-sphere for $0 < i \le n$. A *context in* C with boundary \Box is a formal composition

$$\Gamma \coloneqq v_i \star_{i-1} (\dots \star_1 (v_1 \star_0 \Box \star_0 w_1) \star_1 \dots) \star_{i-1} w_i \tag{6}$$

³For i = 0, we abuse notation and assume there exists a single (-1)-sphere \Box , and denote $C_i(\Box) \coloneqq C_i$.

where v_j, w_j are *j*-cells suitably composable. We set $s_{i-1}(\Gamma) \coloneqq s_{i-1}(w_i)$ and $t_{i-1}(\Gamma) \coloneqq t_{i-1}(v_i)$. For instance, if $\Box = (f, g)$ is a 1-sphere we picture Γ as

We write $Cont(\Box)$ the set of contexts in C with boundary \Box .

For $k \in \mathbb{N}$ such that $i \leq k \leq n$ and for each k-cell A in C such that $(s_{i-1}(A), t_{i-1}(A)) = \Box$, we write

$$\Gamma[A] \coloneqq v_i \star_{i-1} (\dots \star_1 (v_1 \star_0 A \star_0 w_1) \star_1 \dots) \star_{i-1} w_i.$$
(8)

We call $\Gamma[A]$ a *contextualization* of A. Any context Γ defines a function

$$\Gamma: \mathcal{C}_k(\Box) \to \mathcal{C}_k(s_{i-1}(\Gamma), t_{i-1}(\Gamma)).$$

We call this function *contextualization* with Γ .

Recall from Remark 2.4 how (i - 1)-spheres in C defines categories of *i*-cells and (i + 1)-cells. It follows from the axioms of an *n*-sesquicategory that contextualization suitably commutes with source, target, identity and composition:

$$\begin{array}{c} \mathcal{C}_{i}(\Box) \xleftarrow{s_{i}} \mathcal{C}_{i+1}(\Box) \\ \Gamma \downarrow & \downarrow \Gamma \\ \mathcal{C}_{i}(s_{i-1}(\Gamma), t_{i-1}(\Gamma)) \xleftarrow{s_{i}} \mathcal{C}_{i+1}(s_{i-1}(\Gamma), t_{i-1}(\Gamma)) \end{array}$$

In other words, contextualization with Γ defines a functor of categories.

A globular extension P of an n-sesquicate gory C is a globular extension for the underlying globular set of C. Given $A \in \mathsf{P}_{n+1}$ and

$$\Gamma \in \operatorname{Cont}\left(\left(s_{n-1}(A), t_{n-1}(A)\right)\right),\,$$

we define $\Gamma[A]$ as in Eq. (8). We write Cont(P) the set of such (n + 1)-cells:

$$\operatorname{Cont}(\mathsf{P}) \coloneqq \big\{ \Gamma[A] \mid A \in \mathsf{P}_{n+1} \text{ and } \Gamma \in \operatorname{Cont}\left((s_{n-1}(A), t_{n-1}(A)) \right) \big\}.$$

This defines a globular extension of C, also denoted $\operatorname{Cont}(\mathsf{P})$ by abuse of notation, which canonically extends P , in the sense that there is a canonical inclusion $\mathsf{P} \subset \operatorname{Cont}(\mathsf{P})$ that commutes with the source and target maps.

Remark 2.5 (low-dimensional cases). If n = 0 and P is a globular extension of a set C_0 , then $\operatorname{Cont}(\mathsf{P}) = \mathsf{P}$. If n = 1 and P is a globular extension of a category $C_0 \rightleftharpoons_{t_0}^{s_0} C_1$, then

$$\operatorname{Cont}(\mathsf{P}) =$$

$$\{v \star_{1,2} A \star_{2,1} w \mid A \in \mathsf{P}, v, w \in \mathcal{C}_1, s_0(v) = t_0(A) \text{ and } s_0(A) = t_0(v)\}.$$

Diagrammatically:

$$v \star_{1,2} A \star_{2,1} w = \begin{vmatrix} \cdots \\ v \\ \cdots \end{vmatrix} \begin{vmatrix} \cdots \\ A \\ \cdots \\ \cdots \end{vmatrix} \begin{vmatrix} \cdots \\ w \\ \cdots \\ \cdots \end{vmatrix}$$

If n = 2, then elements of Cont(P) are as in Eq. (5), with $A \in P$.

2.2.4 Free *n*-sesquicategories

Let C be an *n*-sesquicategory and P a globular extension of C. Denote by P_{n+1}^* the set consisting of formal identities id_v for each *n*-cell $v \in C_n$ and formal compositions

$$u_1 \star_{n+1,n+1} u_2 \star_{n+1,n+1} \ldots \star_{n+1,n+1} u_d$$

with $u_i \in \text{Cont}(\mathsf{P})$ and $s(u_i) = t(u_{i-1})$ for $1 \le i \le d$. Formal compositions of length zero (i.e. d = 0) are identities, and we regard the above up to the usual identity axioms. This defines an (n + 1)-globular extension of \mathcal{C} , denoted P^* , with source and target maps defined as $s(\mathrm{id}_v) = t(\mathrm{id}_v) = v$, and

$$s(u_1 \star_{n+1,n+1} u_2 \star_{n+1,n+1} \dots \star_{n+1,n+1} u_d) = s(u_d)$$

and
$$t(u_1 \star_{n+1,n+1} u_2 \star_{n+1,n+1} \dots \star_{n+1,n+1} u_d) = t(u_1).$$

We let $\star_{n+1,n+1}$: $\mathsf{P}^*_{n+1} \times_{\mathcal{C}_n} \mathsf{P}^*_{n+1} \to \mathsf{P}^*_{n+1}$ be the formal juxtaposition of suitably *n*-composable elements of P^*_{n+1} , and $\star_{i,n+1}$: $\mathcal{C}_i \times_{i-1} \mathsf{P}^*_{n+1} \to \mathsf{P}^*_{n+1}$ for 0 < i < n+1 be defined as $f \star_{i,n+1} \operatorname{id}_a = \operatorname{id}_{f \star_{i,n} a}$ and

$$f \star_{i,n+1} (u_1 \star_{n+1,n+1} u_2 \star_{n+1,n+1} \dots \star_{n+1,n+1} u_k) = (f \star_{i,n+1} u_1) \star_{n+1,n+1} (f \star_{i,n+1} u_2) \star_{n+1,n+1} \dots \star_{n+1,n+1} (f \star_{i,n+1} u_k).$$

We similarly define $\star_{n+1,i}$. This makes P^{*} into an (n + 1)-sesquicategory, the free (n + 1)-sesquicategory generated by P. We sometimes abuse notation and write P^{*} to denote the set P^{*}_{n+1}.

Remark 2.6 (low-dimensional case). Recall the setting of Remark 2.5. If n = 0, then P^{*} is the free category whose morphisms are formal compositions of elements in P. If n = 1, P^{*} is the free 2-sesquicategory whose 2-morphisms are formal vertical compositions, or 1-composition, of elements in Cont(P). If n = 2, then P^{*} is the free 3-sesquicategory whose 3-morphisms are formal 2-compositions of elements in Cont(P), which we picture as sequences of movies.

2.2.5 *n*-sesquipolygraphs

An *n*-sesquipolygraph P [FM22, section 2.5] consists of the following data:



defined inductively as follows:

- P_0 is a set, and $\mathsf{P}_0^* = \mathsf{P}_0$,
- P_{i+1} is a globular extension for the *i*-globular set $\mathsf{P}_0^* \succeq \ldots \Leftarrow \mathsf{P}_i^*$, and P_{i+1}^* is the free (i+1)-sesquicategory generated by P_{i+1} .

An n-sesquipolygraph provide a notion of presentation for n-sesquicategories, which we now describe.

Given an *n*-sesquicategory C, an equivalence relation \sim on C_n is said to be *higher* [FM22, section 2.6]⁴ if whenever $u \sim v$, we have

- s(u) = s(v) and t(u) = t(v),
- $\Gamma[u] \sim \Gamma[v]$ for each context with boundary (s(u), t(u)) = (s(v), t(v)).

Any (n+1)-sesquicategory C defines a higher equivalence relation on the underlying n-sesquicategory $C_{\leq n}$, setting $\sim_{\mathcal{C}}$ to be the smallest higher equivalence relation such that $s(u) \sim_{\mathcal{C}} t(u)$ for all $u \in C_{n+1}$. We write $[\mathcal{C}]_{\sim}$ for the n-sesquicategory obtained by quotienting $\mathcal{C}_{\leq n}$ with $\sim_{\mathcal{C}}$. if P is an (n+1)-sesquipolygraph, we similarly define \sim_{P} and $[\mathsf{P}]_{\sim}$ starting with the (n+1)-sesquicategory P^* . Then:

Definition 2.7 ([FM22, section 2.6]). A presentation of an *n*-sesquicategory C is the data of an (n + 1)-prepolygraph P such that $[P]_{\sim}$ is isomorphic to C.

Remark 2.8 (low-dimensional cases). A 0-sesquipolygraph is a set. A 1-sesquipolygraph is the same as 1-polygraph, which is the same as a 1-globular set. A 2-sesquipolygraph is the same as a 2-polygraph (see Subsection 1.1.2). For n > 2, n-sesquipolygraphs and n-polygraphs are distinct notions.

2.2.6 Graded *n*-sesquicategories and their presentations

Let G be an abelian group. We extend all the above to the graded case; setting $G = \{*\}$ recovers the previously introduced notions. A set \mathbb{P} is said to be *graded* if it is equipped with a degree function deg: $\mathbb{P} \to G$. A function between graded sets is *homogeneous* if it preserves the degree functions.

A graded *n*-globular set is an *n*-globular set C such that C_n is a graded set. A graded *n*-precategory is an *n*-precategory whose underlying *n*-globular set is graded, such that $\star_{n,n}$ is additive with respect to the grading, and such that the action of a *k*-cell (k < n) on *n*-cells preserves the grading. If C is an *n*-sesquicategory and P is a graded extension of C, then P^{*} is a graded *n*-sesquicategory, where P^{*}_n inherits a grading by setting

$$\deg\left(\Gamma_1[x_1]\star_{n+1,n+1}\ldots\star_{n+1,n+1}\Gamma_m[x_m]\right) \coloneqq \deg(x_1)+\ldots+\deg(x_m)$$

Here $x_k \in \mathsf{P}$ and each Γ_k is a context in \mathcal{C} . A graded *n*-sesquipolygraph P is the data of a (n-1)-prepolygraph $\mathsf{P}_{\leq n-1}$ and a graded extension P_n .

Given a graded *n*-globular set C, a homogeneous extension is a graded globular extension P such that the source and target maps $s_n, t_n \colon P \to C_n$ are homogeneous functions. A graded homogeneous (n + 1)-sesquipolygraph P is the data of a graded *n*-sesquipolygraph $P_{\leq n}$ and a homogeneous extension P_n . Schematically, a graded homogeneous (n + 1)-sesquipolygraph is the following data:



Given a graded set \mathbb{P} , an equivalence relation \sim on \mathbb{P} is *homogeneous* if

⁴In [FM22, section 2.6], a higher equivalence relation is called a congruence.

 $x \sim y$ implies $\deg(x) = \deg(y)$.

If a higher equivalence relation \sim on an *n*-precategory C is homogeneous, the quotient *n*-sesquicategory $[C]_{\sim}$ is graded. Note that the higher equivalence relation induced by a graded homogeneous (n + 1)-sesquipolygraph on the underlying graded *n*-sesquicategory is homogeneous.

Definition 2.9. A presentation of a graded *n*-sesquicategory is the data of a graded homogeneous (n + 1)-sesquipolygraph P such that $[P]_{\sim}$ is isomorphic to C.

2.2.7 Scalar *n*-sesquipolygraphs

A set is said to be *scalar* if it is (\mathbb{k}, \cdot) -graded; that is, a scalar set is a set \mathbb{P} equipped with a function scl: $\mathbb{P} \to \mathbb{k}$. Given a scalar set \mathbb{P} , we write \sim_{scl} , or \equiv_{scl} depending on context, for the smallest homogeneous equivalence relation on \mathbb{P} ; that is, we have $p \sim_{\text{scl}} q$ if and only if scl(p) = scl(q). If the image of scl consists of invertible scalars $\mathbb{k}^{\times} \subset \mathbb{k}$, we say that \mathbb{P} is *scalar-invertible*.

The following are restatements of graded definitions introduced in the previous section.

An *n*-globular set (resp. an *n*-sesquipolygraph) P is *scalar* if P_n is a scalar set. An *n*-sesquicategory \mathcal{C} is *scalar* if \mathcal{C}_n is a scalar set and for every *n*-cells α, β and context Γ , we have

 $\operatorname{scl}(\Gamma[\alpha]) = \operatorname{scl}(\alpha)$ and $\operatorname{scl}(\alpha \star_{n-1} \beta) = \operatorname{scl}(\alpha)\operatorname{scl}(\beta).$

Given a scalar *n*-sesquipolygraph P, the free *n*-sesquicategory P^* generated by P is canonically scalar. Finally, we say that P is a *scalar-invertible n-sesquipolygraph* if P_n is scalar-invertible.

The linear n-sesquicategory presented by a scalar n-sesquipolygraph is defined in Definition 2.13.

2.3 Linear *n*-sesquicategories and their presentations

In this section, we extend the notion of *n*-sesquicategories to the linear case. In fact, we work in the generality of graded linear structures; setting $G = \{*\}$ provides the analogous non-graded linear notions.

Notation 2.10. We fix throughout the section an abelian group G, a commutative ring \Bbbk and a \mathbb{Z} -bilinear map $\mu: G \times G \to \Bbbk^{\times}$. The word "graded" always refers to G-graded, and "linear" to \Bbbk -linear. Given a homogeneous element v, we write $\deg(v)$ is grading.

2.3.1 Linear *n*-sesquicategories

A graded linear *n*-sesquicategory is an *n*-sesquicategory C where, for each (n - 1)-sphere (f, g), the set $C_n(f, g)$ has the structure of a graded k-module, such that the *n*-composition is bilinear and whiskering *n*-cells with a *j*-cell for j < n is linear. In other words:

$$\begin{aligned} (\lambda' u' + v') \star_{n,n} (\lambda u + v) \\ &= \lambda' \lambda(u' \star_{n,n} u) + \lambda'(u' \star_{n,n} v) + \lambda(v' \star_{n,n} u) + v' \star_{n,n} v, \\ &\qquad x \star_{j,n} (\lambda u + v) = \lambda(x \star_{j,n} u) + x \star_{j,n} v, \\ &\qquad (\lambda u + v) \star_{j,n} x = \lambda(u \star_{n,j} x) + v \star_{n,j} x, \end{aligned}$$

with scalars λ, λ' in \mathbb{k} , (n-1)-cells f, g, h in \mathcal{C}_{n-1} , *n*-cells u, v (resp. u', v') in $\mathcal{C}_n(f, g)$ (resp. $\mathcal{C}_n(g, h)$), and a *j*-cell *x* suitably *j*-composable.

2.3.2 Free linear *n*-sesquicategories

Let C be an *n*-sesquicategory and P a graded globular extension of C. The *free graded linear n-pre*category generated by P is the linear *n*-sesquicategory P^l such that $P_{\leq n}^l \coloneqq C_{\leq n}$ and for each *n*-sphere (f,g) in C, $P_{n+1}^l(f,g)$ is the free k-module generated by $P_{n+1}^*(f,g)$.

The k-module $\mathsf{P}_{n+1}^{l}(f,g)$ inherits a grading (of k-module) from the grading (of set) of the set $\mathsf{P}_{n+1}^{*}(f,g)$. Extending (bi)linearly the operations $\star_{n+1,n+1}$, $\star_{k,n+1}$ and $\star_{n+1,k}$ ($k \leq n$) on P_{n+1}^{*} defines a structure of graded linear (n+1)-sesquicategory on P^{l} .

2.3.3 Linear *n*-sesquipolygraphs

A graded linear (n+1)-sesquipolygraph⁵ P is the data of a graded *n*-sesquipolygraph $P_{\leq n}$, together with a homogeneous extension P_{n+1} of the graded linear *n*-sesquicategory $P_{\leq n}^l$:



Given a graded k-module P, an equivalence relation \sim on P is *linear homogeneous* if whenever $v \sim w$ for $v, w \in P$, the following two conditions hold:

- if v and w are homogeneous, deg(v) = deg(w),
- $\lambda v + u \sim \lambda w + u$ for all scalar $\lambda \in \mathbb{k}$ and $u \in P$.

If a higher equivalence relation \sim on a graded linear *n*-sesquicategory C is linear homogeneous, the quotient *n*-sesquicategory $[C]_{\sim}$ is a graded linear *n*-sesquicategory. Note that the higher equivalence relation induced by a graded linear (n + 1)-sesquipolygraph P on the underlying graded linear *n*-sesquicategory $P_{< n-1}^l$ is linear homogeneous.

Definition 2.11. A presentation of a graded linear *n*-sesquicategory is the data of a graded linear (n + 1)-sesquipolygraph P such that $[P]_{\sim}$ is isomorphic to C.

Remark 2.12 (low-dimensional cases). A linear 1-sesquipolygraph is the same as a linear 1-polygraph, and a linear 2-sesquipolygraph is the same as a linear 2-polygraph. For n > 2, linear n-sesquipolygraphs and linear n-polygraphs [All18a] are distinct notions.

2.3.4 Monomial linear *n*-sesquipolygraphs

A graded linear (n + 1)-sesquipolygraph P is monomial if:

$$s_n(r) \in \mathsf{P}_n^*$$
 and $t_n(r) \in \Bbbk \mathsf{P}_n^*$ for all $r \in \mathsf{P}_{n+1}$,

where $\mathbb{k}\mathsf{P}_n^*$ is the subset of P_n^l consisting of vectors of the form λb for $\lambda \in \mathbb{k}$ and $b \in \mathsf{P}_n^*$. We further say it is *monomial-invertible* if $t_n(r) \in \mathbb{k}^{\times}\mathsf{P}_n^*$. We have the following canonical bijection:

$$\left\{ \begin{array}{c} \text{scalar(-invertible)} \\ \text{graded homogeneous} \\ (n+1)\text{-sesquipolygraph} \end{array} \right\} \xrightarrow[\text{scl}]{\text{lin}} \left\{ \begin{array}{c} \text{monomial(-invertible)} \\ \text{graded linear} \\ (n+1)\text{-sesquipolygraph} \end{array} \right\}$$

⁵One can more generally define graded linear (n, p)-sesquipolygraphs, adapting the approach of [All18a].

Given a scalar graded homogeneous (n + 1)-sesquipolygraph P, its *linearization* is the monomial graded linear (n + 1)-sesquipolygraph lin(P) defined as $lin(P)_{\leq n} = P_{\leq n}$ and $lin(P)_{n+1} = P_{n+1}$ as sets, but with the following source and target maps:

 $lin(s_n)(r) = s_n(r)$ and $lin(t_n)(r) = scl(r)t_n(r)$.

The inverse $scl := lin^{-1}$ is defined analogously.

Definition 2.13. Given a scalar graded homogeneous (n+1)-sesquipolygraph P, the linear *n*-sesquicategory presented by P is the linear *n*-sesquicategory presented by lin(P).

2.4 Gray polygraphs and linear Gray polygraphs

This section reviews the notion of a *Gray polygraph*, as can be extracted from [FM22], and then introduces its linear analogue called a *linear Gray polygraph*.

2.4.1 Gray polygraphs

Let Q be a 2-sesquipolygraph. The 3-sesquipolygraph of interchangers is the 3-sesquipolygraph QGray such that $QGray_{\leq 2} = Q$ and $QGray_3$ consists of interchange generators, defined for each 0-composable α , f, β with α : $f \Rightarrow f', g \in P_1^*, \beta$: $h \Rightarrow h' \in P_2$, as the 3-cell

$$X_{\alpha,g,\beta} \colon (\alpha \star_0 g \star_0 h') \star_1 (f \star_0 g \star_0 \beta) \Rrightarrow (f' \star_0 g \star_0 \beta) \star_1 (\alpha \star_0 g \star_0 h),$$

pictured as:

Definition 2.14. A Gray polygraph⁶ is a 3-sesquipolygraph P such that

$$P_{\leq 2}$$
Gray $\subset P$.

In other words, P contains its own 3-sesquipolygraph of interchangers.

One checks that if P is a Gray polygraph, then $[\mathsf{P}]_{\sim}$ is a 2-category. This leads to the following definition:

Definition 2.15. A presentation of a 2-category C is the data of a Gray polygraph P such that $[P]_{\sim}$ is isomorphic to C.

2.4.2 Linear Gray polygraphs

Recall Notation 2.10. Let Q be a graded 2-sesquipolygraph. The 3-sesquipolygraph of (G, μ) -graded interchangers is the scalar-invertible 3-sesquipolygraph QGray, equipped with the function

$$\mathsf{QGray} \to \mathbb{k}^{\times},$$
$$X_{\alpha,g,\beta} \mapsto \mu(\deg \alpha, \deg \beta).$$

We abuse notation and similarly denote QGray the associated monomial-invertible linear *n*-sesquipolygraph.

⁶What we call a Gray polygraph is the underlying 3-sesquipolygraph (or 3-prepolygraph in their terminology) of what is called a Gray presentation in [FM22].

Definition 2.16. A (G, μ) -linear Gray polygraph is a graded linear 3-sesquipolygraph P such that P contains its own monomial graded 3-sesquipolygraph of (G, μ) -graded interchangers.

Similarly, a (G, μ) -scalar Gray polygraph is a scalar graded 3-sesquipolygraph P such that P contains its own scalar graded 3-sesquipolygraph of (G, μ) -graded interchangers. In other words, a (G, μ) -scalar Gray polygraph is precisely the same data as a monomial (G, μ) -linear Gray polygraph.

One checks that if P is a (G, μ) -linear Gray polygraph, then $[P]_{\sim}$ is a (G, μ) -graded-2-category⁷. This leads to the following definition:

Definition 2.17. A presentation of a (G, μ) -graded-2-category C is the data of a (G, μ) -linear Gray polygraph P such that $[P]_{\sim}$ is isomorphic to C.

Remark 2.18. If $G = \{*\}$ is trivial, a $(\{*\}, id)$ -graded-2-category is just a linear 2-category. Hence, a $(\{*\}, id)$ -linear Gray polygraph, and in particular a $(\{*\}, id)$ -scalar Gray polygraph, defines a presentation of a linear 2-category.

3 Linear Gray rewriting modulo

This section introduces *linear Gray rewriting modulo*, the rewriting theory of graded-2-categories. It has three main features: it allows *modulo*, it is *linear*, and it is *higher*. Each of these features requires its own treatment, in addition with a description of how they combine.

We refer to the extended summary (Subsection 1.2) for an introduction to this section and comparison with the literature. We sometimes use graded \mathfrak{gl}_2 -foams (Section 4) as a source of (counter)examples; the reader is then referred to Subsection 4.1 for the definitions.

We remind the reader that every categorical structure is assumed to be small.

Notation 3.1. We fix the same notations G, \Bbbk and μ as in Notation 2.10 throughout the section. As much as possible, we follow the font conventions introduced in Notation 2.1; in addition, we use blackboard fonts (e.g. \mathbb{P}) for abstract rewriting systems and typewriter fonts (e.g. \mathbb{P}) for linear rewriting systems. Equivalence relations are denoted either with the symbol \sim or the symbol \equiv ; the former case is typically used for congruence of a rewriting system, while the latter is typically used for an equivalence relation on rewriting sequences (see Subsection 3.2 for both notions). In particular, \equiv is typically one "categorical dimension" higher than \sim .

3.1 Coherence modulo from convergence modulo

Given a groupoid \mathcal{P} , one may be interested in understanding its set of connected components $\pi_0(\mathcal{P})$,⁸ or its $\pi_1(\mathcal{P})$, that is, its *coherence*. In this section, we describe how one can use convergence to answer both questions.

Let S be a category. The notion of a higher equivalence relation on *n*-sesquicategories defined in Subsection 2.2.5 specializes to categories: an equivalence relation \equiv on the set of morphisms, such that if $f \equiv f'$, then (i) s(f) = s(f') and t(f) = t(f'), and (ii) if g and h are suitably composable morphisms, then $g \circ f \circ h \equiv g \circ f' \circ h$. In this context, we call such an equivalence relation an *abstract equivalence*, and say that two morphisms f and g are \equiv -equivalent if $f \equiv g$. We choose this terminology to avoid confusion with the notion of \equiv -congruence, used extensively

⁷To save space, we will not recall the definition of a (G, μ) -graded-2-category [SV23] (although see Subsection 1.2.1 for an informal definition); one can take Definition 2.17 as a definition.

⁸That is, its Grothendieck group $K_0(\mathcal{P})$.

in the next section. In particular, if $\equiv \equiv \equiv_{dis.}$ is the discrete abstract equivalence (see Definition 3.6), \equiv -equivalence reduces to equality, while \equiv -congruence reduces to congruence (see the next sections for the definitions).

We denote S^{\top} the localisation of the category S. Given a groupoid P, we fix the following data for the section:

a category
$$S$$
 such that $S^{\top} = P$, a wide⁹ subgroupoid $\mathcal{E} \subset S$,
and an abstract equivalence \equiv on S . (*)

We use the same notation \equiv for the smallest abstract equivalence on S^{\top} containing \equiv .¹⁰ Morphisms in S are depicted with plain arrow $x \to y$ (or $x \longrightarrow y$). Morphisms in \mathcal{E} are depicted with unoriented wiggly lines $x \sim y$ (or $x \rightsquigarrow y$). Unspecified morphisms $x \to_S y$ (resp. $x \sim_{\mathcal{E}} y$) indicate the mere existence of a morphism in S (resp. in \mathcal{E}) between x and y. Note the use of subscripts to specify the category, used extensively in what follows.

Recall from Subsection 2.2.7 the notion of scalar category and of its associated abstract equivalence \equiv_{scl} .

Definition 3.2. The category \mathcal{P} is said to be \equiv -coherent modulo \mathcal{E} if every endomorphism is \equiv -equivalent to a morphism in \mathcal{E} , up to conjugation. That is, for every endomorphism $f: x \to_{\mathcal{P}} x$, there exist morphisms $g: x \to_{\mathcal{P}} y$ and $e: y \sim_{\mathcal{E}} y$, such that $f \equiv g^{-1} \circ e \circ g$:

$$f \underbrace{\frown}_{x} \equiv x \underbrace{g}_{g^{-1}} y \underbrace{\frown}_{y} e$$

If \mathcal{P} is scalar, we say that it is scalar-coherent modulo \mathcal{E} whenever it is \equiv_{scl} -coherent modulo \mathcal{E} .

Remark 3.3. If \mathcal{E} is discrete (it does not contain any morphism apart from identities; we write $\mathcal{E} = \emptyset$), then \mathcal{P} is \equiv -coherent modulo \mathcal{E} if and only if parallel morphisms are \equiv -equivalent, which recovers the usual notion of coherence for a category. In particular, if \mathcal{P} is scalar with scalars scl: $\mathcal{P}_1 \to \mathbb{K}$, then \mathcal{P} is scalar-coherent if for every endomorphism $f: x \to_{\mathcal{P}} x$ we have scl(f) = 1.

Lemma 3.4 (transitivity of coherence modulo). Let C be a groupoid, \equiv an abstract equivalence on C and $\mathcal{E} \subset \mathcal{D}$ wide subgroupoids of C. If C is \equiv -coherent modulo \mathcal{D} and \mathcal{D} is \equiv -coherent modulo \mathcal{E} , then C is \equiv -coherent modulo \mathcal{E} .

An *S*-branching is a pair (f, f') with $f : x \to_S y$ and $f' : x \to_S y'$. An *S*-confluence is a pair (g, g') with $g : y \to_S z$ and $g' : y' \to_S z$. A branching (f, f') is said to be \equiv -confluent if there exists a confluence (g, g') such that $f' \circ f \equiv g' \circ g$:



We say that S is \equiv -confluent if every branching is \equiv -confluent.

We say that S is *terminating* if every infinite sequence $(f_n)_{n \in \mathbb{N}}$ of morphisms in S with $t(f_n) = s(f_{n+1})$ eventually terminates in morphisms in \mathcal{E} . We say that S is \equiv -convergent if it

⁹Containing all objects; equivalently, containing all identities.

¹⁰In particular, if $f \equiv g$ in S, then we both have $f \equiv g$ and $f^{-1} \equiv g^{-1}$ in S^{\top} .

is both \equiv -confluent and terminating. An object $y \in S$ for which $y \to_S z$ implies $y \sim_{\mathcal{E}} z$ is called an *S*-normal form. We denote NF_S the set of *S*-normal forms. If $x \to_S y$ and y is an *S*-normal form, we say that y is an *S*-normal form for x.

Note that the notions of termination, convergence and normal form all depend on the data of \mathcal{E} . To stress this dependency, we sometimes add *modulo* \mathcal{E} , as in *terminating modulo* \mathcal{E} or *convergent modulo* \mathcal{E} .

The *Church–Rosser property* is a classical property of ≡-confluent categories (see for instance [Ara+23, p. 1.3.14]):

Lemma 3.5 (Church–Rosser property). Let $S \equiv and \mathcal{E}$ as in (*). If S is \equiv -confluent, then every morphism $f \in \mathcal{P} = S^{\top}$ can be decomposed as $f \equiv h^{-1} \circ g$ for some $g, h \in S$.

Proof. By definition, f decomposes as $f = h_1 \circ g_1^{-1} \circ h_2 \circ g_2^{-1} \circ \ldots \circ h_k \circ g_k^{-1}$ for some $k \in \mathbb{N}$ and $g_i, h_i \in S$. Inductively applying \equiv -confluence leads to the desired decomposition.

Definition 3.6. The discrete abstract equivalence on S is the abstract equivalence $\equiv_{\text{dis.}}$ such that every pair of parallel morphisms is equivalent with respect to $\equiv_{\text{dis.}}$.

We simply denote $\equiv_{dis.}$ -confluence as *confluence*, and $\equiv_{dis.}$ -convergence as *convergence*. Note that \equiv -confluence (resp. \equiv -convergence) for any abstract equivalence \equiv implies confluence (resp. convergence).

Finally, the following two propositions explain how convergence leads to a description of $\pi_0(\mathcal{P})$ and $\pi_1(\mathcal{P})$, respectively.

Proposition 3.7. Let \mathcal{P} , \mathcal{S} and \mathcal{E} as in (*). If \mathcal{S} is convergent modulo \mathcal{E} , then the canonical mapping $NF_{\mathcal{S}} \to \pi_0(\mathcal{P})$ sending an \mathcal{S} -normal form to its connected component induces a bijection

$$\operatorname{NF}_{\mathcal{S}_{\pi_0}(\mathcal{E})} \xrightarrow{\sim} \pi_0(\mathcal{P}),$$

where the quotient identifies objects belonging to the same connected component of \mathcal{E} .

Proof. If we have $f: x \to_S y$, then x and y belong to the same connected component of \mathcal{P} . In particular, since S is terminating the mapping $NF_S \to \pi_0(\mathcal{P})$ is surjective. Assume then that x and y are S-normal forms belonging to the same connected component of \mathcal{P} . By the Church–Rosser property (Lemma 3.5), there exists $f: x \to_S z$ and $g: y \to_S z$. Since x and y are S-normal forms, f and g must be in \mathcal{E} , so that x and y belong to the same connected component of \mathcal{E} . \Box

Proposition 3.8. Let S, \equiv and \mathcal{E} as in (*). If S is \equiv -convergent modulo \mathcal{E} , then $\mathcal{P} = S^{\top}$ is \equiv -coherent modulo \mathcal{E} .

Proof. Thanks to the Church–Rosser property (Lemma 3.5), for any endomorphism $f: x \to_{\mathcal{P}} x$ of \mathcal{P} there exist $f_1, f_2: x \to_{\mathcal{S}} z$ morphisms in \mathcal{S} such that $f \equiv f_1^{-1} \circ f_2$. Since \mathcal{S} is terminating, there exists an \mathcal{S} -normal form y and a morphism $h: z \to_{\mathcal{S}} y$. Because \mathcal{S} is \equiv -confluent, the branching $(h \circ f_1, h \circ f_2)$ admits a \equiv -confluence. Since y is an \mathcal{S} -normal form, this confluence is in \mathcal{E} , and there exists $e \in \mathcal{E}$ such that $e \circ h \circ f_1 \equiv h \circ f_2$. Setting $g \coloneqq h \circ f_1$ gives $e \circ g \equiv g \circ f$, which concludes.

3.2 Abstract rewriting modulo

Recall that 1-polygraph, 1-sesquipolygraph and 1-globular set are identical notions (Remark 2.8). Recall also the notion of the free category \mathbb{P}^* generated by a 1-polygraph \mathbb{P} (Subsection 2.2.4). We denote \mathbb{P}^{\top} the localisation of \mathbb{P}^* .

Definition 3.9. An abstract rewriting system (abstract RS) is the data $(\mathbb{P}; \equiv)$ of a 1-polygraph \mathbb{P} together with an abstract equivalence \equiv on \mathbb{P}^* .

Unpacking the definition, an abstract RS is the data of a set X of 0-cells, called *elements*, and a set \mathbb{P} of 1-cells, called *rewriting steps*, equipped with *source* and *target maps*:

$$X \xleftarrow{s}{t} \mathbb{P},$$

together with an abstract equivalence \equiv on the free category \mathbb{P}^* generated by \mathbb{P} . Note that we abuse notation and denote \mathbb{P} both the 1-polygraph and the set of relations. A morphism in \mathbb{P}^* (resp. in \mathbb{P}^\top) is called a *rewriting sequence* (resp. a *congruence*). If a rewriting sequence (resp. a congruence) decomposes in $n \mathbb{P}$ -rewriting steps, we call the number n its *length*. We write $x \to_{\mathbb{P}} y$ to refer to an unspecified rewriting step with source x and target y (note that there could be more than one rewriting step between given source and target), or to indicate the existence of such a rewriting step. Similarly, we write $x \xrightarrow{*}_{\mathbb{P}} y$ (resp. $x \sim_{\mathbb{P}} y$) to denote a rewriting sequence (resp. a congruence), and say that x rewrites into (resp. is congruent to) y.

$$x \to_{\mathbb{P}} y$$
 $x \xrightarrow{*}_{\mathbb{P}} y$ $x \sim_{\mathbb{P}} y$
rewriting step rewriting sequence congruence

Note that following these notations, $x \xrightarrow{*}_{\mathbb{P}} y$ coincides with $x \to_{\mathbb{P}^*} y$, and $x \sim_{\mathbb{P}} y$ coincides with $x \to_{\mathbb{P}^+} y$.

Let $\mathbb{R} = (X, \mathbb{R})$ and $\mathbb{E} = (X, \mathbb{E})$ be two 1-polygraphs with the same underlying set of elements. We define the following 1-polygraph:

$${}_{\mathbb{E}}\mathbb{R}_{\mathbb{E}} \coloneqq \mathbb{E}^{\top} imes_{X} \mathbb{R} imes_{X} \mathbb{E}^{\top}.$$

In other words, a rewriting step in $_{\mathbb{E}}\mathbb{R}_{\mathbb{E}}$ is a triple $(e, r, e') \in \mathbb{E}^{\top} \times \mathbb{R} \times \mathbb{E}^{\top}$ with t(e) = s(r) and t(r) = s(e'). The source and target maps are defined as s(e, r, e') = s(e) and t(e, r, e') = t(e').

Definition 3.10. An abstract rewriting system modulo (abstract RSM) is the data $\mathbb{S} = (\mathbb{R}, \mathbb{E}; \equiv)$ of two 1-polygraphs $\mathbb{R} := (X, \mathbb{R})$ and $\mathbb{E} = (X, \mathbb{E})$, together with an abstract equivalence \equiv on $(\mathbb{E}\mathbb{R}_{\mathbb{E}})^* \cup \mathbb{E}^{\top}$. In that case, we say that \mathbb{S} is an abstract RSM modulo \mathbb{E} on the set X.¹¹

An S-rewriting sequence is either an \mathbb{E} -congruence (in which case it has *length zero*) or an $\mathbb{E}\mathbb{R}_{\mathbb{E}}$ -rewriting sequence (in which case it has the same length as the $\mathbb{E}\mathbb{R}_{\mathbb{E}}$ -rewriting sequence).¹² We denote S* the set of S-rewriting sequences. Note that $(\mathbb{E}\mathbb{R}_{\mathbb{E}})^* \cup \mathbb{E}^{\top} = \mathbb{S}^*$. We similarly denote $x \to_{\mathbb{S}} y$ (resp. $x \stackrel{*}{\to}_{\mathbb{S}} y$, resp. $x \sim_{\mathbb{S}} y$) an S-rewriting step (resp. an S-rewriting sequence, resp. an S-congruence).

¹¹ In [DM22; Dup22], the authors allow for a more general definition, where S is any abstract RS such that $\mathbb{R} \subset \mathbb{S} \subset \mathbb{E}\mathbb{R}_{\mathbb{E}}$. We do not work in this generality. Moreover, we impose \mathbb{R} and \mathbb{E} to be defined on the same set of elements; see Footnote 21 for further comments on this.

¹² Our terminology differs from [Dup21; Dup22], where S-rewriting sequences coincide with $_{\mathbb{E}}\mathbb{R}_{\mathbb{E}}$ -rewriting sequences. In particular, our notions of branching and confluence do not explicitly depend on the modulo data, while the notion of termination does; this is the converse of [Dup21; Dup22].

An abstract RSM inherits all the notions and results introduced in the previous section, setting (\ast) as

$$\mathcal{S} \coloneqq \mathbb{S}^*$$
 and $\mathcal{E} \coloneqq \mathbb{E}^\top$.

For instance, we say that S is *terminating* (or *terminating modulo* E) if S is terminating modulo \mathcal{E} ; equivalently, S is terminating if there is no infinite sequence $\{f_n\}_{n\in\mathbb{N}}$ such that f_n is an S-rewriting step and $t(f_n) = s(f_{n+1})$. (Note that with our conventions, one *cannot* replace "step" by "sequence" in the above characterization.)

In contrast with the previous section, an abstract RSM provides a notion of *locality*. A *local* \mathbb{S} -branching is a pair (f, g) with $f: x \to_{\mathbb{S}} y$ and $f: x \to_{\mathbb{S}} y'$ two rewriting steps in \mathbb{S} :



We say that S is *locally* \equiv -*confluent* if all local branchings are \equiv -confluent. An S-*local triple* is a triple [f, e, g] with \mathbb{R} -rewriting steps f and g and \mathbb{E} -congruence e, such that s(f) = s(e) and t(e) = s(g). Note that every S-local triple defines a local S-branching $(f, g \circ e)$, and that if every S-local triple is \equiv -confluent, then S is \equiv -confluent. Local S-branchings and S-local triples should be thought as essentially identical notions, one being more suited for general statements while the other being better suited for explicit computations.

Given an abstract RSM $\mathbb{S} = (\mathbb{R}, \mathbb{E}, \equiv)$, an abstract RSM $\mathbb{T} = (\mathbb{R}_1, \mathbb{E}, \equiv_1)$ is a *sub-abstract RSM* of \mathbb{S} if $\mathbb{R}_1 \subset \mathbb{R}$ and for all $f, g \in \mathbb{T}^*$, $f \equiv_1 g$ implies that $f \equiv g$. If an \mathbb{S} -branching (f, g) admits a \mathbb{T} -congruence (f', g') such that $f' \circ f \equiv g' \circ g$, we say that (f, g) is (\mathbb{T}, \equiv) -confluent. We say that an \mathbb{S} -congruence h is (\mathbb{T}, \equiv) -confluent (resp. (\mathbb{T}, \equiv) -congruent) if there exists a \mathbb{T} -confluence (f, g) with s(f) = s(h), s(g) = t(h), such that $f \equiv g \circ h$ (resp. a \mathbb{T} -congruence h' such that $h \equiv h'$). We use similar notations to indicate that a given notion related to \mathbb{S} restricts to a notion related to \mathbb{T} .

Remark 3.11 (scalar abstract RSM). Recall from Subsection 2.2.7 the notion of scalar 1-polygraph, scalar relation and free scalar 1-category. An abstract RSM $S = (\mathbb{R}, \mathbb{E}, \equiv)$ is said to be *scalar* if \mathbb{R} is scalar, \mathbb{E} is scalar-invertible and $\equiv \equiv \equiv_{scl}$ (that is, $r \equiv s$ if and only if scl(r) = scl(s)).

Remark 3.12 (quotient). Given an abstract RSM $\mathbb{S} = (X; \mathbb{R}, \mathbb{E}; \equiv)$, we can define its *quotient* abstract RS $[\mathbb{S}]^{\mathbb{E}} := ([X]^{\mathbb{E}}; [\mathbb{R}]^{\mathbb{E}}; [\equiv]^{\mathbb{E}})$, where $[X]^{\mathbb{E}}$ is the set of \mathbb{E} -congruence classes and $[\mathbb{R}]^{\mathbb{E}}$ and $[\equiv]^{\mathbb{E}}$ are defined analogously. In that case, \mathbb{S} is terminating modulo \mathbb{E} if and only if $[\mathbb{S}]^{\mathbb{E}}$ is terminating, and if \mathbb{E} is \equiv -coherent, then \mathbb{S} is \equiv -confluent if and only if $[\mathbb{S}]^{\mathbb{E}}$ is $[\equiv]^{\mathbb{E}}$ -confluent. In other words, when \mathbb{E} is coherent, the theory of rewriting modulo reduces to rewriting (without modulo) on its quotient.

However, explicitly working with the modulo data highlights some of the difficulties of higher linear rewriting (see Subsection 1.2.4 and Subsection 3.5). Moreover, one does not always have the luxury of a coherent modulo data; indeed, the modulo data for graded \mathfrak{gl}_2 -foams (Section 4) is not coherent (see Remark 3.26).

The next subsections describe how confluence can be achieved from a local analysis.

3.2.1 Tamed Newmann's lemma

The classical Newmann's lemma states that under termination, confluence follows from local confluence. It readily extends to modulos:

Lemma 3.13 (Newmann's lemma). Let $\mathbb{S} = (\mathbb{R}, \mathbb{E}; \equiv)$ be an abstract RSM. If \mathbb{S} is a terminating and locally \equiv -confluent, then it is \equiv -confluent.

However, local confluence turns out to be too restrictive for some purposes, especially in the linear context. In this subsection, we introduce the weaker notion of \succ -*tamed* \equiv -*congruence*, for which an analogue of Newmann's lemma still holds.

Recall that a binary transitive relation on X is called a *preorder*. Let $\mathbb{E} = (X, \mathbb{E})$ be an abstract RS. A preorder \succ is said to be \mathbb{E} -invariant if $(x' \sim_{\mathbb{E}} x \text{ and } x \succ y \text{ and } y \sim_{\mathbb{E}} y')$ implies $(x' \succ y')$. We shall always assume that preorders on X are \mathbb{E} -invariant. If $M \subset X$ is a set of elements in X, we write $x \succ M$ if $x \succ y$ for all $y \in M$. If $f = f_n \circ \ldots \circ f_1$ is a sequence of composable arrows on X, we write $x \succ f$ to mean $x \succ \{s(f_1), t(f_1), \ldots, t(f_n)\}$.

Definition 3.14. Let $\mathbb{S} = (\mathbb{R}, \mathbb{E}; \equiv)$ be an abstract RSM on the set X and \succ a preorder on X. An \mathbb{S} -branching (f, g) of source \bullet is said to be \succ -tamely \equiv -congruent (resp. \succ -tamely \equiv -confluent) if there exists a \equiv -congruence h (resp. \equiv -confluence (f', g')) such that $\bullet \succ h$ (resp. $\bullet \succ f'^{-1} \circ g'$).

In particular, \succ -tameness implies $\bullet \succ t(f)$ and $\bullet \succ t(g)$. Here is a schematic for a \succ -tamed congruence, where horizontal positions are used to suggest relative orderings with respect to \succ :



Our notion of \succ -tameness is reminiscent of the notion of *confluence by decreasingness* as introduced by van Oostrom [vOos94], and as appearing in [All18a; Dup21; Dup22] in the context of higher linear rewriting. Indeed, S-rewriting steps inherit a preorder from \succ by stating that $h_1 \succ h_2$ if and only if $s(h_1) \succ s(h_2)$ and $s(h_1) \succ t(h_2)$. With this choice, a \succ -tame \equiv -confluence is decreasing in the sense of [vOos94]. However, moving the order from rewriting steps to elements allows a meaningful weakening of the notion to congruence. As far as we are aware, this has not appeared in the literature.

Definition 3.15. Let S be an abstract RSM. A preorder \succ on X is said to be compatible with S if $x \to_S y$ implies $x \succ y$.

We denote $\succ_{\mathbb{S}}$ the minimal \mathbb{E} -invariant preorder compatible with \mathbb{S} . Note that if an \mathbb{E} -invariant preorder \succ is compatible with \mathbb{S} , then minimal elements for \succ are \mathbb{S} -normal forms. In particular, if \succ is well-founded (we shall say that \succ is *terminating*) then \mathbb{S} is terminating. The converse holds if $\succ = \succ_{\mathbb{S}}$.

Note also that if an \mathbb{E} -invariant preorder \succ is compatible with \mathbb{S} , \equiv -confluence implies \succ -tame \equiv -confluence, and that irrespective of whether \succ is compatible, \succ -tame \equiv -confluence always implies \succ -tame \equiv -congruence.
Lemma 3.16 (tamed Newmann's lemma). Let $\mathbb{S} = (\mathbb{R}, \mathbb{E}; \equiv)$ be an abstract RSM and \succ an \mathbb{E} -invariant preorder compatible with \mathbb{S} . If \succ is terminating and every local \mathbb{S} -branching is \succ -tamely \equiv -congruent, then \mathbb{S} is \equiv -confluent.

Proof. Since \succ is terminating, we can proceed by induction on \succ (see e.g. [Ara+23, section 1.3.9]) to show that the following property holds for every $x \in X$:

P(x): every S-branching with source y such that $x \succ y$ is S-confluent.

If x is minimal for \succ , then in particular x is an S-normal form, and so P(x) automatically holds. Consider then x generic and assume that P(y) holds whenever $x \succ y$.

If h is an S-congruence such that $x \succ h$, one can use the induction hypothesis to show that h is S-confluent. (Recall that h being S-confluent means that there exists a confluence (f, g) with s(f) = s(h), s(g) = t(h), such that $f \equiv g \circ h$.) In particular, every local S-branching with source x is S is confluent.

Consider a (not necessarily local) S-branching (f, g). Decomposing f and g into S-rewriting steps $f_m \circ \ldots \circ f_1$ and $g_n \circ \ldots \circ g_1$ respectively gives a local S-branching (f_1, g_1) , S-confluent by the previous paragraph. The rest of the confluence can be completed using induction on \succ , as shown in the following diagram:



This concludes.

3.2.2 Branchwise E-congruence and Newmann's lemma

Following the (tamed) Newmann's lemma, we wish to study confluence (or tamed congruence) of local branchings. In principle, working modulo makes it a difficult task. Indeed, given that the length of the \mathbb{E} -congruence in a local triple is not limited, the number of local branchings is in general infinite. To circumvent this problem, rewriting steps need to be understood *up to* \mathbb{E} -congruence, similarly to how elements in X are understood up to \mathbb{E} -congruence. In practice, one has canonical ways to do so, coming from naturality axioms with regard to the modulo: naturality of interchangers when working modulo interchange (see Subsection 3.4.6), naturality of the pivotal structure when working up to isotopies (see Lemma 4.25), and so on. This subsection formalizes this situation.

Let $\mathbb{S} = (\mathbb{R}, \mathbb{E}; \equiv)$ be an abstract RSM. Two S-rewriting sequences f and g are said to be (\mathbb{E}, \equiv) -congruent if there exist \mathbb{E} -congruences $e_s : s(f) \sim_{\mathbb{E}} s(g)$ and $e_t : t(f) \sim_{\mathbb{E}} t(g)$ such that

 $e_t \circ f \equiv g \circ e_s$:



An S-branching (f, g) is said to be (\mathbb{E}, \equiv) -congruent if there exists an \mathbb{E} -congruent $e: t(f) \sim_{\mathbb{E}} t(g)$ such that $e \circ f \equiv g$. Two S-branchings (f, f') and (g, g') are said to be *branchwise* (\mathbb{E}, \equiv) -congruent if they have the same source and (f, g) (resp. (f', g')) is (\mathbb{E}, \equiv) -congruent.

The following lemma states that "confluence is preserved under branchwise $\mathbb E\text{-congruence}$ ":

Lemma 3.17. Let $\mathbb{S} = (\mathbb{R}, \mathbb{E}; \equiv)$ be an abstract RSM. If (f, g) and (f', g') are branchwise (\mathbb{E}, \equiv) -congruent \mathbb{S} -branchings, then (f, g) is \equiv -confluent if and only if (f', g') is. \Box

The proof of the above lemma fits into one picture:



In practice, one works with S-local triples. Two S-local triples [f, e, g] and [f', e', g'] are said to be (\mathbb{E}, \equiv) -congruent if f (resp. g) is (\mathbb{E}, \equiv) -congruent to f' (resp. g') such that the relevant square of \mathbb{E} -congruences is \equiv -equivalent:



In other words, [f, e, g] and [f', e', g'] represent branchwise (\mathbb{E}, \equiv) -congruent S-branchings. For local triples, Lemma 3.17 says that [f, e, g] is (\mathbb{S}, \equiv) -confluent if and only if [f', e', g'] is.

3.2.3 Branchwise confluence and tamed Newmann's lemma

Replacing \mathbb{E} -congruence with \equiv -confluence (resp. \succ -tamed \equiv -congruence) defines the following analogous branchwise notions:

Definition 3.18. Let $\mathbb{S} = (\mathbb{R}, \mathbb{E})$ be an abstract RSM and \succ an \mathbb{E} -invariant preorder on X. Two \mathbb{S} -branchings (f, g) and (f', g') are branchwise \equiv -confluent (resp. branchwise \succ -tamely \equiv -congruent) if they have the same source and the branchings (f, f') and (g, g') are respectively \equiv -confluent (resp. \succ -tamely \equiv -congruent).

Contrary to branchwise \mathbb{E} -congruence, working up to branchwise confluence does not preserve confluence. However, the following lemma states that "tamed congruence is preserved under branchwise tamed congruence". Having applications in mind, we state it with respect to a sub-abstract RSM.

Lemma 3.19 (BRANCHWISE TAMED CONGRUENCE LEMMA). Let $\mathbb{S} = (\mathbb{R}, \mathbb{E})$ be an abstract RSM and \succ an \mathbb{E} -invariant preorder on X compatible with \mathbb{S} . Let also $\mathbb{T} \subset \mathbb{S}$ be a sub-abstract RSM. If (f, g) and (f', g') are branchwise \succ -tamely (\mathbb{T}, \equiv) -congruent \mathbb{S} -branchings, then (f, g) is (\mathbb{T}, \equiv) -tamely (\mathbb{T}, \equiv) -congruent if and only if (f', g') is.

The proof of the above lemma fits into one picture:



The BRANCHWISE TAMED CONGRUENCE LEMMA 3.19 is an important practical tool, as it can greatly simplify the study of tamed congruence. We will use it heavily in Section 4.

3.2.4 Branchwise rewriting

As we can rewrite elements, we can similarly rewrite a rewriting sequence into another rewriting sequence, or a branching (f, g) into another branching (f', g'). The latter appears as a special case of branchwise \equiv -confluence, with (f, g) branchwise \equiv -confluent to a trivial branching.

Definition 3.20. Let S be an abstract RSM. We say that an S-branching (f,g) rewrites into a branching (f',g') if there exist S-rewriting sequences connecting t(f) with t(f'), s(f) = s(g) with s(f') = s(g') and t(g) with t(g'), such that the relevant squares are \equiv -equivalent:



Lemma 3.21. Let \mathbb{S} be an abstract RSM and \succ an \mathbb{E} -invariant preorder compatible with \mathbb{S} . Let also (f,g) be an \mathbb{S} -branching that rewrites into another \mathbb{S} -branching (f',g'). If (f',g') is \succ -tamed \equiv -congruent, then so is (f,g).

Branchwise rewriting is another practical tool used in Section 4. We shall come back to it in Subsection 3.4.5 and Subsection 3.5.5.

3.3 Linear rewriting modulo

This section generalizes linear rewriting theory to the modulo setting. A lot of constructions and results parallel the non-modulo setting, following [GHM19]; see Subsection 1.2.8 for further details on the literature.

As in Notation 2.10, we fix \Bbbk a commutative ring and denote \Bbbk^{\times} its invertible elements. Given a set \mathcal{B} , we write $\langle \mathcal{B} \rangle_{\Bbbk}$ the free \Bbbk -module generated by \mathcal{B} .

3.3.1 Linear rewriting systems modulo

Recall that linear 1-polygraph and linear 1-sesquipolygraph are identical notions. Unpacking the definition, a linear 1-polygraph is the data of a set \mathcal{B} of 0-cells, called *monomials*, and a set P of 1-cells, called *relations*, equipped with *source* and *target maps s* and *t*:

$$\langle \mathcal{B} \rangle_{\Bbbk} \xleftarrow{s}{t} \mathsf{P}.$$

Elements in $V \coloneqq \langle \mathcal{B} \rangle_{\Bbbk}$ are called *vectors*. We fix a choice of monomials \mathcal{B} (and hence vectors) for the reminder of the section. The \Bbbk -module presented by $(\mathcal{B}, \mathbb{P})$ is the module $\langle \mathcal{B} \rangle_{\Bbbk} / \langle \mathbb{P} \rangle_{\Bbbk}$. Note that $\langle \mathbb{P} \rangle_{\Bbbk}$ can be viewed as a globular extension of $\langle \mathcal{B} \rangle_{\Bbbk}$, extending *s* and *t* linearly. If $\mathbb{P}^{=}$ denotes the reflexive closure of \mathbb{P} , we write $\mathbb{P}^{l} := \langle \mathbb{P}^{=} \rangle_{\Bbbk}$, viewed as a globular extension:

$$\langle \mathcal{B} \rangle_{\mathbb{k}} \stackrel{s}{\underset{t}{\longleftarrow}} \mathbb{P}^{l}.$$

Explicitly, relations in \mathbb{P}^l are of the form $\sum_i \lambda_i r_i + v$ for $\lambda_i \in \mathbb{K}$, $r_i \in \mathbb{P}$ and $v \in V$, with source $\sum_i \lambda_i s(r_i) + v$ and target $\sum_i \lambda_i t(r_i) + v$. As is explained in Subsection 3.3.2, \mathbb{P}^l should be thought of the set of congruences associated to P, analogous to \mathbb{P}^{\top} in the abstract case.

We say that R is *left-monomial* if for all $r \in \mathbb{R}$, we have $s(r) \in \mathcal{B}$; in other words, each r is of the form $b \xrightarrow{r}_{\mathbb{R}} \sum_{i} \lambda_{i} b_{i}$, with $\lambda_{i} \in \mathbb{k}$ and $b, b_{i} \in \mathcal{B}$. We say that R is *adapted* if it is left-monomial and we have $s(r) \notin \operatorname{supp}(t(r))$ for every $r \in \mathbb{R}$. Then:

Definition 3.22. A linear rewriting system (linear RS) *is the data* (\mathcal{B} ; P) *of an adapted linear 1-polygraph* P *on a set* \mathcal{B} .

We sometimes leave \mathcal{B} implicit, and call P a linear RS. The adaptedness condition is not an important restriction. Indeed, if $\mu b + \sum_{i} \mu_{i} b_{i} = 0$ is some relation in a k-module presentation, we can rewrite it as $b = -\mu^{-1} \sum_{i} \mu_{i} b_{i}$ (provided that μ is invertible). Doing so with every relation gives an adapted linear 1-polygraph presenting the given k-module (provided we can always find such an invertible scalar μ).

We now extend to modulos the notion of linear RS. Let E be another set of linear relations on \mathcal{B} :

$$\langle \mathcal{B} \rangle_{\Bbbk} \stackrel{s}{\xleftarrow{t}} \mathsf{E}.$$

Denote $\Bbbk^{\times}\mathcal{B}$ the subset of $\langle \mathcal{B} \rangle_{\Bbbk}$ consisting of vectors of the form λb for $\lambda \in \Bbbk^{\times}$ and $b \in \mathcal{B}$. We say that E is *monomial-invertible* if it is left-monomial and for all $e \in E$, we have $t(e) \in \Bbbk^{\times}\mathcal{B}$; in other words, each e is of the form $b \xrightarrow{e}_{\mathsf{E}} \lambda b'$, with $\lambda \in \Bbbk^{\times}$ and $b, b' \in \mathcal{B}$. This coincides with the notion of monomial-invertible linear 1-polygraph defined in Subsection 2.3.4. If we drop the condition that scalars are invertible, we simply say that E is *monomial*. We will always assume that the modulo data is monomial-invertible.

Similarly to abstract RSM, we write $u \sim_{\mathbf{E}} v$ if there exists $e \in \mathbf{E}^l$ such that s(e) = u and t(e) = v, and in that case we say that u and v are E-congruent. In the linear context, we furthermore have a notion of *projective* E-congruence, an equivalence relation on the set of monomials \mathcal{B} , defined as $b \sim_{\mathbf{E}} b'$ if and only if there exists $\lambda \in \mathbb{k}^{\times}$ such that $b \sim_{\mathbf{E}} \lambda b'$. For v a vector in V, we set

$$\widetilde{\operatorname{supp}}_{\mathsf{E}}(v) \coloneqq \left\{ b \in \mathcal{B} \mid b \stackrel{\cdot}{\sim}_{\mathsf{E}} b' \text{ for some } b' \in \operatorname{supp}(v) \right\}$$

and call $\widetilde{\operatorname{supp}}_{\mathsf{E}}(v)$ the E-projective support of v.

A set R of linear relations on \mathcal{B} is said to be E-*adapted* if it is left-monomial and we have $s(r) \notin \widetilde{\operatorname{supp}}_{\mathsf{E}}(t(r))$ for every $r \in \mathsf{R}$.

Definition 3.23. A linear rewriting system modulo (linear RSM) is the data S = (B; R, E) of two linear 1-polygraphs R := (B; R) and E := (B; E), such that E is monomial-invertible and R is E-adapted.¹³

We sometimes leave \mathcal{B} implicit, and call (R, E) a linear RSM. Note that if $E = \emptyset$ (i.e. the set of relations in E is empty), we recover the notion of linear rewriting system. The *module presented* by S is the module $\langle \mathcal{B} \rangle_{\Bbbk} / \langle R \sqcup E \rangle_{\Bbbk}$. We write:

$$[-]_{\mathtt{S}} \colon \langle \mathcal{B} \rangle_{\Bbbk} \to \langle \mathcal{B} \rangle_{\Bbbk} / \langle \mathtt{R} \sqcup \mathtt{E} \rangle_{\Bbbk}$$

the associated quotient map. Finally, we write $S^l := (R \sqcup E)^l$.

We conclude this subsection with a few remarks; some appeal to concepts only defined in the next subsections.

Remark 3.24. The (E-)adaptedness condition prevents "obvious" obstructions to termination. Indeed, if $s(r) \in \hat{supp}_{E}(t(r))$ then there exists an infinite sequence of positive S-rewriting steps, all of type r. Moreover, without this assumption the Church–Rosser property for positive rewritings steps (Lemma 3.31) does not hold; see Remark 3.32. Given how fundamental this result is, we choose to enforce the adaptedness condition in the definition, in contrast with the abstract case. This is only a choice of presentation; indeed, in practice one eventually wishes to work with terminating rewriting systems, which implies adaptedness. In [GHM19], "left-monomial" encompasses both our "left-monomial" and "adapted" notions. However, the adaptedness condition is dropped in [All18a] (in the context of strict higher linear rewriting) and in [Dup21; Dup22] (in the context of strict higher linear rewriting modulo).¹⁴ This motivates our change of terminology, hoping to avoid further confusion. Note that while the adaptedness condition is easy to check in the non-modulo setting, it can be more involved in the modulo setting, as one needs to scan through projective E-congruence classes.

Remark 3.25. Recall the notion of scalar abstract RSM from Remark 3.11. A linear RSM (\mathcal{B} ; R, E) is *monomial* if R is monomial. The bijection between scalar 1-polygraphs and monomial linear 1-polygraphs (see Subsection 2.3.4) extends to rewriting systems:

sending $\mathbb{S} = (X; \mathbb{R}, \mathbb{E}, \equiv)$ to $S = (\mathcal{B}; \mathbb{R}, \mathbb{E})$, with $\mathcal{B} = \langle X \rangle_{\mathbb{k}}$, $\mathbb{R} = \lim(\mathbb{R})$ and $\mathbb{E} = \lim(\mathbb{E})$. (Caveat: we did not impose the adaptedness condition in the abstract case; see also Remark 3.24.) It is canonical in the sense that a property holds for \mathbb{S} if and only the suitable analogue holds for S; for instance, \mathbb{S} is scalar-confluent if and only if S^+ is confluent.

Remark 3.26. Recall the notion of quotient of an abstract abstract RSM from Remark 3.12. If E is scalar-coherent in the sense that $scl(E)^{\top}$ is scalar-coherent, we can similarly define a quotient for a linear RSM $S = (\mathcal{B}; R, E)$, working on the $[\mathcal{B}]^{E}$ of *projective* E-congruence classes. This applies for instance when E is *scalar-free*, in the sense that $s(e), t(e) \in \mathcal{B}$ for all $e \in E$.

Coherence of E is necessary to define the quotient, as otherwise an element $b \in \mathcal{B}$ may be a zero divisor in the module $\langle \mathcal{B} \rangle_{\Bbbk} / \langle E \rangle_{\Bbbk}$. The case of graded \mathfrak{gl}_2 -foams, described Section 4, provides an example where E is not fully coherent: interchanging two identical dots gives a scalar $XY \in \Bbbk$ (for \Bbbk as defined in Definition 4.4).

¹³Our definition differs from that in [Dup21; Dup22], as already commented in Footnotes 11 and 12 for the abstract case; see Remark 3.24 for a further difference, specific to the linear case.

¹⁴In particular, Lemma 4.2.9 in [All18a] and Lemma 1.1.5 in [Dup22] are not correct as stated. However, in both cases they eventually impose the condition of "exponentiation freedom", which implies adaptedness. (This condition is missing in [All18a], but this is corrected in [All18b].)

Remark 3.27. We have chosen not to equip a linear RSM with the extra data of an abstract equivalence \equiv , as we did in the abstract case. In other words, a linear RSM is implicitly equipped with the discrete abstract equivalence $\equiv_{\text{disc.}}$, identifying any two parallel relations. While this is unnecessary for our purpose, there is no obstruction in generalizing linear RSMs to arbitrary equivalences on relations.

3.3.2 Positive rewriting steps

Compared to the abstract case, the linear case requires a specific notion of rewriting step:

Definition 3.28. Let P be a linear RS. A P-rewriting step is an element $\alpha \in \langle \mathsf{P} \rangle_{\Bbbk}$ of the form

$$\alpha = \lambda r + v, \quad \lambda \in \mathbb{k} \setminus \{0\}, r \in \mathsf{P}, v \in \langle \mathcal{B} \rangle_{\mathbb{k}}.$$

In that case, we say that α is of type r. The P-rewriting step α is said to be positive¹⁵ if

$$s(r) \notin \operatorname{supp}(v).$$

We write a P-rewriting step as $\alpha : s(\alpha) \to_{P} t(\alpha)$, and a positive P-rewriting step as $\alpha : s(\alpha) \to_{P} t(\alpha)$. The set of (resp. positive) P-rewriting steps is denoted \mathbb{P}^{st} (resp. \mathbb{P}^{+}).

If v = 0, we say that α is monomial; in that case, α is necessarily positive.

Note that P^{st} is necessarily symmetric; indeed, if α is a P-rewriting step then so is its inverse

$$\alpha^{-1} = (-\lambda)r + \lambda(s(r) + t(r)) + v,$$

as defined in the proof above. Hence, P^{st} does not terminate and cannot provide a suitable reduction algorithm. However, if α is positive then α^{-1} cannot be positive; indeed, the assumptions $s(r) \notin \operatorname{supp}(t(r))$ (adaptedness of P) and $s(r) \notin \operatorname{supp}(v)$ (positiveness of α) imply that

$$s(r) \in \operatorname{supp}(\lambda(s(r) + t(r)) + v).$$

This makes positive P-rewriting steps suitable candidates to define a reduction algorithm.

These notions are readily extended to modulos:

Definition 3.29. Let S = (R, E) be a linear RSM. An S-rewriting step is a composition

$$u \sim_{\mathbf{E}} \lambda s(r) + v \longrightarrow_{\mathbf{R}} \lambda t(r) + v \sim_{\mathbf{E}} w,$$

where the middle arrow is a R-rewriting step. This S-rewriting step is said to be positive if

$$s(r) \notin \widetilde{\operatorname{supp}}_{\mathsf{E}}(v).$$

We write an S*-rewriting step as* \rightarrow *s, and a positive* P*-rewriting step as* \rightarrow *s:*

$$\begin{array}{ll} s(\alpha) \dashrightarrow_{\mathbf{S}} t(\alpha) & s(\alpha) \rightarrow_{\mathbf{S}} t(\alpha) \\ \text{(not necessarily positive)} & positive \\ rewriting step & rewriting step \end{array}$$

The set of (resp. positive) S-rewriting steps is denoted S^{st} (resp. S^+).

If v = 0, we say that α is monomial; in that case, α is necessarily positive.

¹⁵This terminology appears e.g. in [CDM22] other references [All18a; Dup22; GHM19] use the terminology *elementary relation* for rewriting step, and rewriting step for positive rewriting step.

As in the non-modulo context, one can argue that S^{st} is symmetric, with positive S-rewriting steps providing an answer to this formal obstruction to termination.

Note that the positivity condition is $s(r) \notin \widetilde{\text{supp}}_{\mathsf{E}}(v)$, and not $s(r) \notin \text{supp}(v)$. In other words, a positive S-rewriting step is *not* a composition as above such that the middle arrow is a positive R-rewriting step. The positivity condition is stronger, and depends on E. Otherwise, positive S-rewriting steps do not provide a suitable solution to termination.¹⁶ This contrasts with the abstract setting.

Note that for the positivity condition to make sense, E must be monomial. Extending the theory to non-monomial modulo rules remains a non-trivial question.¹⁷

Both S^{st} and S^+ provide an abstract RSM associated to S, namely respectively $(\langle \mathcal{B} \rangle_{\Bbbk}; S^{st}, E^{st})$ and $(\langle \mathcal{B} \rangle_{\Bbbk}; S^+, E^{st})^{18}$. As such, an S-rewriting step (resp. a positive S-rewriting step) in the sense of Definition 3.29 is the same as an S^{st} -rewriting step (resp. an S^+ -rewriting step), and we shall use the two terminologies interchangeably. Similarly, a *positive* S-*rewriting sequence* denotes an S^+ -rewriting sequence.

The following lemma shows that both S-rewriting steps and positive S-rewriting steps are suited to study S-congruence:

Lemma 3.30. Let $S = (\mathcal{B}; R, E)$ be a linear RSM, $u, v \in \langle \mathcal{B} \rangle_{\Bbbk}$ and recall the notation $[u]_{S}, [v]_{S}$ for their respective image in $\langle \mathcal{B} \rangle_{\Bbbk} / \langle R \sqcup E \rangle_{\Bbbk}$. The following statements are equivalent:

- (i) $[u]_{\mathbf{S}} = [v]_{\mathbf{S}}$ in $\langle \mathcal{B} \rangle_{\mathbb{k}} / \langle \mathbf{R} \sqcup \mathbf{E} \rangle_{\mathbb{k}}$;
- (*ii*) u and v are S^{st} -congruent;
- (iii) u and v are S^+ -congruent.

In particular, $S^l = (S^+)^\top = (S^{st})^\top$.

The lemma implies that there is no distinction between the properties of S^+ -congruence, an S^{st} -congruence or an S^l -congruence; note however that a *given* S^{st} -congruence may not be positive. In order to prove the lemma, we need the following Church–Rosser property for positive rewriting steps, which generalizes modulo Lemma 3.1.2 in [GHM19]. The last statement is explained and used in Subsection 3.3.4, and can be ignored for the purpose of this subsection.

Lemma 3.31. Let S = (B; R, E) be a linear RSM. If f is an S-rewriting step, then there exist positive S-rewriting steps g, h of length at most one such that $f = h^{-1} \circ g$:



Moreover, we have $f \succeq^{\text{rel}} w$ for any E-invariant linear preorder \succ on \mathcal{B} (see Definition 3.42).

¹⁶For instance, consider the linear RSM $\mathbf{S} = (\mathcal{B}; \mathbf{R}, \mathbf{E})$ with $\mathcal{B} = \{a, a', b\}, \mathbf{R} = \{a \to b\}$ and $\mathbf{E} = \{a \sim a'\}$, and the S-rewriting sequence $0 = a - a \sim_{\mathbf{E}} a - a' \rightarrow_{\mathbf{R}} b - a' \sim_{\mathbf{E}} b - a \rightarrow_{\mathbf{R}} b - b = 0$.

¹⁷ In [Dup21; Dup22], a positive S-rewriting step *is* defined as a composition as above such that the middle arrow is a positive R-rewriting step, and E is not constrained to be monomial. In [DEL21], Dupont's work is applied with a non-monomial E.

¹⁸We may also associate the abstract RSM ($\langle \mathcal{B} \rangle_{\Bbbk}$; \mathbb{R}^{st} , \mathbb{E}^{st}); this would essentially lead to the same abstract rewriting theory as ($\langle \mathcal{B} \rangle_{\Bbbk}$; \mathbb{S}^{st} , \mathbb{E}^{st}). However, ($\langle \mathcal{B} \rangle_{\Bbbk}$; \mathbb{R}^{+} , \mathbb{E}^{st}) and ($\langle \mathcal{B} \rangle_{\Bbbk}$; \mathbb{S}^{+} , \mathbb{E}^{st}) are in general very different abstract RSMs!

Proof. Let $\lambda r + v$ be the R-rewriting step associated to f, with $r \in \mathbb{R}$, $\lambda \in \mathbb{k} \setminus \{0\}$ and $v \in \langle \mathcal{B} \rangle_{\mathbb{k}}$. Extracting s(r) from the decomposition of v, we let $\mu \in \mathbb{k}$ and $v' \in \langle \mathcal{B} \rangle_{\mathbb{k}}$ such that $v \sim_{\mathbb{E}} \mu s(r) + v'$ and $s(r) \notin \widetilde{supp}_{\mathsf{E}}(v')$. Set

$$w \coloneqq (\lambda + \mu)t(r) + v'.$$

Then $s(f) \sim_{\mathbb{E}} (\lambda + \mu)s(r) + v'$ (resp. $t(f) \sim_{\mathbb{E}} \mu s(r) + (\lambda t(r) + v')$) is either equal to w if $\mu = -\lambda$ (resp. if $\mu = 0$), or there exists a positive rewriting step $(\lambda + \mu)s(r) + v' \rightarrow_{\mathbb{R}} w$ (resp. $\mu s(r) + (\lambda t(r) + v') \rightarrow_{\mathbb{R}} w$):

$$\lambda s(r) + \mu s(r) + v' \xrightarrow{}_{\mathbf{E}} s(f) \xrightarrow{f}_{--\stackrel{f}{\xrightarrow{}}_{\mathbf{R}}} t(f) \xrightarrow{}_{\mathbf{E}} \lambda t(r) + \mu s(r) + v'$$

Here we use the fact that $s(r) \notin \widetilde{\operatorname{supp}}_{\mathsf{E}}(t(r))$ (E-adaptedness) to ensure that $s(r) \notin \widetilde{\operatorname{supp}}_{\mathsf{E}}(\lambda t(r) + v')$. Finally, it follows from

$$\dot{\widetilde{\operatorname{supp}}}_{\mathsf{E}}\left(\lambda s(r) + \mu s(r) + v'\right) \cup \dot{\widetilde{\operatorname{supp}}}_{\mathsf{E}}\left(\lambda t(r) + \mu s(r) + v'\right) \supset \dot{\widetilde{\operatorname{supp}}}_{\mathsf{E}}\left((\lambda + \mu)t(r) + v'\right)$$

that $f \succcurlyeq^{\text{rel}} w$ for any E-invariant linear preorder \succ on \mathcal{B} .

Proof of Lemma 3.30. (iii) \Leftrightarrow (ii) is given by Lemma 3.31. If $u \to_{\mathbf{S}} v$, then [u] = [v], so (ii) \Rightarrow (i). To show (i) \Rightarrow (ii), assume that [u] = [v]. In that case, $u \sim_{\mathbf{E}} v + \sum_{i=0}^{n} \lambda_i(s(r_i) - t(r_i))$ for some scalars $\lambda_i \in \mathbb{K}$ and relations $r_i \in \mathbf{R}$. Write $u_j = \sum_{i \leq j} \lambda_i(t(r_i) - s(r_i))$ and $\alpha_j : u_j \to u_{j-1}$ the obvious S-rewriting step. Successively applying the α_j 's defines an S-rewriting sequence

$$u \sim v + u_n \xrightarrow{\alpha_n} v + u_{n-1} \xrightarrow{\alpha_n} \dots \xrightarrow{\alpha_n} v.$$

This concludes.

Remark 3.32. Without the adaptedness condition, Lemma 3.31 does not hold: for instance, one can consider the linear RS $P = \{a \rightarrow 2a\}$ and the non-positive rewriting step $a + a \rightarrow 2a + a$ as a counterexample.

3.3.3 Basis from convergence

Fix S = (B; R, E) a linear RSM. In this subsection, we explain how convergence of S^+ can provide a basis for the underlying module. Unsurprisingly, finding a basis is closely related to understanding congruence. As in the abstract setting, normal forms are prime candidates for congruence representatives. To get a candidate basis, we look at monomial normal forms:

Definition 3.33. Let S = (B; R, E) be a linear RSM. A monomial S^+ -normal form is a monomial $b \in B$ which is a normal form for S^+ ; that is, we have $b \not\sim_E s(r)$ for all $r \in R$. We denote BNF_S the set of monomial S^+ -normal forms.

Remark 3.34. A linear combination of S⁺-normal forms is an S⁺-normal form, and monomials in the support of an S⁺-normal form are S⁺-normal forms. The zero vector 0 is always an S⁺-normal form. In other words, NF_S is a k-module and NF_S = $\langle BNF_S \rangle_k$.

Proposition 3.35. Let S = (B; R, E) be a linear RSM. If S^+ is convergent modulo, the canonical linear map

$$\mathrm{NF}_{S}/\langle \mathrm{E}
angle_{\Bbbk} \to \langle \mathcal{B}
angle_{\Bbbk}/\langle \mathrm{R} \sqcup \mathrm{E}
angle_{\Bbbk}$$

is an isomorphism. In particular, if $B \subset \mathcal{B}NF_S$ is such that $[B]_E$ is a basis for the module $NF_S/\langle E \rangle_k$, then $[B]_S$ is a basis for $\langle \mathcal{B} \rangle_k / \langle \mathbb{R} \sqcup E \rangle_k$, the module presented by S.

Proof. By definition, $\pi_0(S^+)$ denotes the k-module of S^+ -congruence classes. Lemma 3.30 can be reformulated as stating that the canonical linear map $\pi_0(S^+) \to \langle \mathcal{B} \rangle_k / \langle R \sqcup E \rangle_k$ is an isomorphism. Proposition 3.7 concludes.

In the situation of the above proposition, finding a basis reduces to finding a basis of the module $NF_S/\langle E \rangle_{\Bbbk}$. Since $(\mathcal{B}NF_S, E)$ is a monomial linear RS, we may instead consider the scalar RS $(\mathcal{B}NF_S, \operatorname{scl}(E)^{\top})$ (see Remark 3.25). If E^l is coherent, that is, if $\operatorname{scl}(E)^{\top}$ is scalar-coherent, then any choice of projective E-congruence representatives on $\mathcal{B}NF_S$ defines a basis of $NF_S/\langle E \rangle_{\Bbbk}$.

This leads to the BASIS-FROM-CONVERGENCE THEOREM:

Theorem 3.36 (BASIS-FROM-CONVERGENCE THEOREM). Let $S = (\mathcal{B}; R, E)$ be a linear RSM. If S^+ is convergent modulo and if $scl(E)^{\top}$ is scalar-coherent on the set $\mathcal{B}NF_S$ of monomial S^+ -normal forms, then the module $\langle \mathcal{B} \rangle_{\Bbbk} / \langle R \sqcup E \rangle_{\Bbbk}$ presented by S is free, and any choice of projective E-congruence representatives on $\mathcal{B}NF_S$ defines a basis.

Setting $E = \emptyset$, the Basis-From-Convergence Theorem 3.36 becomes:

Corollary 3.37. Let $(\mathcal{B}; P)$ be a linear RS. If P^+ is convergent, then $\mathcal{B}NF_P$ is a basis for the module $\langle \mathcal{B} \rangle_{\Bbbk} / \langle P \rangle_{\Bbbk}$ presented by P.

3.3.4 Termination order

This section extends to the linear case the notion of abstract compatible preorder (Subsection 3.2.1).

Given a linear RSM S = (B; R, E), we say that a relation \succ on B is E-*invariant* if it is invariant with respect to projective E-congruence (recall Subsection 3.3.1) in the sense of Subsection 3.2.1; that is, if

$$(a' \sim_{\mathsf{E}} a \text{ and } a \succ b \text{ and } b \sim_{\mathsf{E}} b') \text{ implies } (a' \succ b').$$

In the presence of a linear RSM, We shall always assume that relations are E-invariant.

Definition 3.38. Let S = (B; R, E) be a linear RSM and \succ an E-invariant preorder on B. We say that \succ is compatible with S if

 $s(r) \succ b$ for all $r \in \mathbb{R}$ and $b \in \operatorname{supp}(t(r))$.

We denote \succ_{S} the smallest E-invariant preorder on \mathcal{B} compatible with S.

Recall from Subsection 2.3.3 the notion of a linear relation. An E-invariant preorder \succ on \mathcal{B} induces an \mathbb{E}^l -invariant linear relation \succ^+ on $\langle \mathcal{B} \rangle_{\Bbbk}$, setting $u \succ^+ v$ whenever the following two conditions hold:

- (a) $\operatorname{supp}_{\mathbf{E}}(u) \neq \operatorname{supp}_{\mathbf{E}}(v);$
- (b) for every a in $\widetilde{\operatorname{supp}}_{\mathsf{E}}(v) \setminus \widetilde{\operatorname{supp}}_{\mathsf{E}}(u)$, there exists $b \in \widetilde{\operatorname{supp}}_{\mathsf{E}}(u) \setminus \widetilde{\operatorname{supp}}_{\mathsf{E}}(v)$ such that $b \succ a$.

This definition corresponds to setting $u \succ^+ v$ if and only if $\widetilde{\operatorname{supp}}_{\mathsf{E}}(u) \succ^{\operatorname{set}} \widetilde{\operatorname{supp}}_{\mathsf{E}}(v)$ where $\succ^{\operatorname{set}}$ is the multi-set relation induced by \succ ,¹⁹ defined on the power set $\mathcal{P}(\mathcal{B})$ as:

 $M \succ^{\text{set}} N \quad \Leftrightarrow \quad M \neq N \text{ and } \forall y \in N \setminus M, \exists x \in M \setminus N \text{ such that } x \succ y.$

Equivalently, $M \succ^{\text{set}} N$ if and only if one can go from M to N by a sequence of moves consisting in removing an element b and adding elements a_i with $b \succ a_i$. This last interpretation motivates the definition of \succ^+ , designed precisely such that the following holds:

Lemma 3.39. Let S = (B; R, E) be a linear RSM and \succ a relation on B. If \succ is a strict²⁰ E-invariant preorder on B compatible with S in the linear sense of Definition 3.38, then \succ^+ is a strict E^l -invariant preorder on $\langle B \rangle_k$ compatible with S^+ in the abstract sense of Definition 3.15.

Proof. It is shown in [BN98, lemma 2.5.4] that if \succ is a strict preorder, so is \succ^{set} ; hence if \succ is a strict preorder, so is \succ^+ . Note that without the strictness condition, \succ^+ may not even be transitive.

Consider then $r: s(r) \rightarrow_{\mathbb{R}} t(r)$. Strictness (or E-adaptedness) implies that

$$\dot{\widetilde{\operatorname{supp}}}_{\mathsf{E}}(s(r)) \setminus \dot{\widetilde{\operatorname{supp}}}_{\mathsf{E}}(t(r)) = \dot{\widetilde{\operatorname{supp}}}_{\mathsf{E}}(s(r))$$

so that $s(r) \succ^+ t(r)$. The general case follows from the following lemma.

Lemma 3.40. Let S = (B; R, E) be a linear RSM and \succ a relation on B. For vectors $u, v \in \langle B \rangle_{\Bbbk}$, we have:

$$u \succ^+ v \quad \Rightarrow \quad \lambda u + w \succ^+ \lambda v + w$$

for all $\lambda \in \mathbb{k} \setminus \{0\}$ and $w \in \langle \mathcal{B} \rangle_{\mathbb{k}}$ such that $\dot{\operatorname{supp}}_{\mathsf{E}}(u) \cap \dot{\operatorname{supp}}_{\mathsf{E}}(w) = \emptyset$.

Proof. Since $\dot{supp}_{\mathsf{E}}(\lambda u + w) = \dot{supp}_{\mathsf{E}}(u) \sqcup \dot{supp}_{\mathsf{E}}(w)$, we have:

$$\begin{split} \widetilde{\operatorname{supp}}_{\mathsf{E}}(\lambda u + w) \setminus \widetilde{\operatorname{supp}}_{\mathsf{E}}(\lambda v + w) \supset \widetilde{\operatorname{supp}}_{\mathsf{E}}(u) \setminus \widetilde{\operatorname{supp}}_{\mathsf{E}}(v), \\ \dot{\widetilde{\operatorname{supp}}}_{\mathsf{E}}(\lambda v + w) \setminus \dot{\widetilde{\operatorname{supp}}}_{\mathsf{E}}(\lambda u + w) \subset \dot{\widetilde{\operatorname{supp}}}_{\mathsf{E}}(v) \setminus \dot{\widetilde{\operatorname{supp}}}_{\mathsf{E}}(u). \end{split}$$

We now relate to termination:

Lemma 3.41. Let S = (B; R, E) be a linear RSM and \succ a preorder on B. If \succ is E-invariant and compatible with S, we have the following implications:

 \succ is terminating on $\mathcal{B} \iff \succ^+$ is terminating on $\langle \mathcal{B} \rangle_{\Bbbk} \Rightarrow S^+$ terminates.

Moreover, if $\succ = \succ_{s}$ (see Definition 3.38) then the converse of the last implication holds.

Proof. Note that if a preorder terminates, it is necessarily strict. It is shown in [BN98, theorem 2.5.5] that \succ is terminating if and only if \succ^{set} is terminating. In that case, Lemma 3.39 implies that S⁺ terminates.

It remains to show that if S^+ terminates, then \succ_S is terminating on \mathcal{B} . We proceed by contraposition and assume that there exists an infinite sequence

$$b_0 \succ_{\mathsf{S}} b_1 \succ_{\mathsf{S}} b_2 \succ_{\mathsf{S}} \dots b_n \succ_{\mathsf{S}} b_{n+1} \succ_{\mathsf{S}} \dots$$

in \mathcal{B} . Let $r_n \in \mathbb{R}$ such that $s(r_n) \sim_{\mathbb{E}} b_n$ and $b_{n+1} \in \widetilde{\operatorname{supp}}_{\mathbb{E}}(t(r_n))$. We construct a sequence of \mathbb{S}^+ -rewriting steps starting with $v_0 = b_0$ and defining v_{n+1} recursively by applying r_m on v_n (possibly after an E-congruence), for m the biggest index possible. The assumption on m ensures that we always have $b_{m+1} \in \operatorname{supp}(v_{n+1})$, so that this process does not end. Hence, \mathbb{S}^+ does not terminate.

¹⁹See e.g. [BN98, Definition 2.5.3]; we only use the special case of sets, and therefore denote it \succ^{set} .

²⁰A preorder \succ is *strict* if $b \neq b$ for all $b \in \mathcal{B}$.

We conclude the subsection with the following notion, which appears in the statement of Lemma 3.31:

Definition 3.42. Let \succ be a relation on a set \mathcal{B} . the relative relation \succeq^{rel} induced by \succ is the following relation on the power set $\mathcal{P}(\mathcal{B})$:

$$M \succcurlyeq^{\mathrm{rel}} N \quad \Leftrightarrow \quad \forall b \in \mathcal{B}, (b \succ M) \Rightarrow (b \succ N).$$

If $S = (\mathcal{B}; R, E)$ is a linear RSM and \succ an E-invariant preorder on \mathcal{B} , we set $u \succeq^{\text{rel}} v$ if and only if $\widetilde{\text{supp}}_{E}(u) \succeq^{\text{rel}} \widetilde{\text{supp}}_{E}(v)$.

Note that equivalently, $M \succeq^{\text{rel}} N$ if and only if $\forall L \in \mathcal{P}(\mathcal{B}), (L \succ M)$ implies $(L \succ N)$.

3.3.5 Tamed linear Newmann's lemma

The linear structure of a linear RSM S = (B; R, E) induces canonical types of S⁺-local triples. Let [f, e, g] be a S⁺-local triple such that f and g is of type r_1 and r_2 , respectively. Then:

- if $s(r_1) \dot{\not{\sim}}_{\mathsf{E}} s(r_2)$, one says that [f, e, g] is additive,
- if $s(r_1) \sim_{\mathbf{E}} s(r_2)$, one says that [f, e, g] is intersecting.

Recall the notion of branchwise E-congruence for branchings (Subsection 3.2.2). A S⁺-local triple [f, e, g] is additive if, up to branchwise E-congruence, it has of the following form:

$$[f, e, g] = [\lambda_1 r_1 + \lambda_2 s(r_2) + v, \text{id}, \lambda_1 s(r_1) + \lambda_2 r_2 + v],$$

An additive branching has a canonical Sst-confluence:



On the other hand, [f, e, g] is intersecting if, up to branchwise E-congruence, it has the following form:

$$[f, e, g] = [\lambda_1 r_1 + w, e_r + w, \lambda_2 r_2 + w],$$

where $e_r \colon \lambda_1 s(r_1) \sim_{\mathsf{E}} \lambda_2 s(r_2)$. We say that [f, e, g] is *monomial* if it is intersecting and if, up to branchwise E-congruence, it has the form $\lambda_1 = 1$ and w = 0 with the notations above. That is, [f, e, g] is monomial if, up to branchwise E-congruence, it has the following form:

$$[f, e, g] = [r_1, e_r, \lambda r_2].$$

where $e_r : s(r_1) \sim_{\mathsf{E}} \lambda s(r_2)$.

Note that both branches of a monomial branching are monomial rewriting steps. By definition, every intersecting branching is of the form $\lambda_1[f, e, g] + w$ for [f, e, g] a monomial branching. Still by definition, additive, intersecting and monomial branchings are positive branchings.

We end the subsection with a linear analogue of the tamed Newmann's lemma (Lemma 3.13). Given a linear RSM S = (B; R, E) and a preorder \succ on B, we say " \succ -tameness" to refer to \succ ⁺-tameness.

Theorem 3.43 (TAMED LINEAR NEWMANN'S LEMMA). Let $S = (\mathcal{B}; R, E)$ be a linear RSM and \succ an E-invariant preorder on \mathcal{B} compatible with S. If \succ is terminating and every monomial local S^+ -branching is \succ -tamely S^{st} -congruent, then S^+ is convergent.

The TAMED LINEAR NEWMANN'S LEMMA 3.43 reduces the study of convergence to tamed congruence of monomial local branchings. The TAMED LINEAR NEWMANN'S LEMMA 3.43 and the BASIS-FROM-CONVERGENCE THEOREM 3.36 are the two main results of linear rewriting modulo theory, allowing one to deduce bases from a local analysis. The hypothesis that \succ is terminating is necessary; see [GHM19, Remark 4.2.4] for a counterexample.

Our proof can be understood as a generalization of [GHM19, Theorem 4.2.1, part (ii)], working with tamed Sst-congruence instead of S⁺-confluence.

Proof. It follows from hypothesis and Lemma 3.40 that every intersecting S^+ -branching is \succ -tamely S^{st} -congruent. Consider then an additive S^+ -local triple as pictured above. We wish to show that its canonical S^{st} -confluence is tamed, that is:

$$\lambda_1 s(r_1) + \lambda_2 s(r_2) + w \succ^+ \lambda_1 t(r_1) + \lambda_2 t(s_2) + w.$$

This happens precisely if $s(r_2) \notin \widetilde{\operatorname{supp}}_{\mathsf{E}}(t(r_1))$ or if $s(r_1) \notin \widetilde{\operatorname{supp}}_{\mathsf{E}}(t(r_2))$. (These two cases correspond respectively to the upper branch or lower branch being positive.) Assuming otherwise, the compatibility of \succ implies that $s(r_2) \succ s(r_1)$ and $s(r_1) \succ s(r_2)$. This contradicts the assumption that \succ terminates.We conclude that every local S⁺-branching is \succ -tamely Sst-congruent.

Note then that:

Lemma 3.44. A branching is \succ -tamely Sst-congruent if and only if it is \succ -tamely S⁺-congruent.

Proof. This follows from Lemma 3.31 and the fact that for $M, N_1, N_2, L \in \mathcal{P}(\mathcal{B})$, if $M \geq^{\text{rel}} N_1 \cup N_2$ and $N_1 \geq^{\text{rel}} L$, then $M \geq^{\text{rel}} N_1 \cup N_2$.

It follows that every local S⁺-branching is \succ -tamely S⁺-congruent. We can now apply the tamed (abstract) Newmann's lemma (Lemma 3.16) and conclude.

3.4 Higher rewriting modulo

This section introduces (weak) higher rewriting modulo, extending the work of Forest and Mimram [FM22], who studied the non-modulo and context-agnostic (see below) setting. Although we restrict to 3-dimensional rewriting, many notions are not specific to the 3-dimensional case; in particular, this section could be adapted to the 2-dimensional case (rewriting modulo in categories, including monoids).

To some extent, it serves as a blueprint for the next section, which deals with higher *linear* rewriting modulo.

3.4.1 Higher rewriting system modulo

Recall the notion of 3-sesquicategory (Subsection 2.2.2), 3-sesquipolygraph and higher equivalence (Subsection 2.2.5). For C a 3-sesquicategory and \equiv a higher equivalence on C, we say that \equiv satisfies the independence axiom if the following holds:

independence axiom: for every pair of 1-composable 3-cells $f: \phi \to \phi'$ and $g: \psi \to \psi'$ in P*, the 3-cells $(\phi' \star_1 g) \star_2 (f \star_1 \psi)$ and $(f \star_1 \psi') \star_2 (\phi \star_1 g)$ are \equiv -equivalence:



The independence axiom captures the interchange of 3-cells in P*.

Definition 3.45. A higher rewriting system (higher RS) ($P; \equiv$) is the data of a 3-sesquipolygraph P, together with a higher equivalence \equiv on P^{*} (see Subsection 2.2.5) satisfying the independence axiom. A higher rewriting system modulo (higher RSM) S = (R, E; \equiv) is the data of two 3-sesquipolygraphs R and E with the same underlying 2-sesquipolygraph R₂ = E₂,²¹ together with a higher equivalence \equiv on R^{*} \cup E^T satisfying the independence axiom.

Recall the notion of contexts (Subsection 2.2.3). Given a higher RSM $S = (R, E; \equiv)$ every choice of 1-sphere \Box in R_1^* defines an abstract RSM $S(\Box) = (\mathbb{R}(\Box), \mathbb{E}(\Box); \equiv)$ on the underlying set $X(\Box)$, where

$$X(\Box) = \mathsf{R}_2^*(\Box) = \mathsf{E}_2^*(\Box),$$
$$\mathbb{R}(\Box) = \operatorname{Cont}(\mathsf{R}_3)(\Box) \quad \text{and} \quad \mathbb{E}(\Box) = \operatorname{Cont}(\mathsf{E}_3)(\Box).$$

We abuse notation and write \equiv for its restriction on $(\mathsf{R}^* \cup \mathsf{E}^\top)(\Box)$. Moreover, every context Γ on \Box defines a morphism of abstract RSM

$$\Gamma \colon \mathbb{S}(\Box) \to \mathbb{S}(s_1(\Gamma), t_1(\Gamma)).$$
(9)

(A morphism of abstract RSM is a pair of morphisms of 1-globular sets, the latter denoting a pair of functions commuting with the source and target maps; see Subsection 2.2.1.) We say that a branching (f,g) is a *contextualization* of another branching (f',g') whenever $\Gamma[f',g'] = (f,g)$.

We think of a higher RSM as a category of abstract RSMs, with morphisms given by contexts. In principle, one could deal with each abstract RSM $S(\Box)$ independently, using the tools of abstract rewriting theory modulo (Subsection 3.2). In practice, one would like to leverage the fact that these abstract RSMs gather together in a category, and relate to one another (and themself) via contextualization. In that sense, higher rewriting theory (modulo) is nothing else than the study of morphisms of abstract rewriting systems (modulo).

²¹ In [Dup21; Dup22], Dupont allows the more general definition $E_2 \subset R_2$ in the context of linear 3-polygraph. We prefer to avoid this generality, as it leads to ambiguity for the statement "E is convergent". For instance, for the linear 3-polygraphs E and R in [Dup21; Dup22], we have that E is convergent when viewed on E_2 , but not when viewed on R_2 . Similar issues appear in [DEL21].

In many practical situations, termination can only be obtained with context-dependent termination rule, where r being a rewriting rule does not imply that $\Gamma[r]$ is. For that reason, we introduce the following notion:

Definition 3.46. Let $S = (R, E; \equiv)$ be a higher RSM. An abstract sub-system \mathbb{T} is the data of a family of subsets $\mathbb{T}(\Box) \subset R_3^*(\Box)$ for each 1-sphere \Box in R_1^* .

We think of \mathbb{T} as a family of sub-abstract RSMs $\mathbb{T}(\Box) \subset \mathbb{S}(\Box)$ modulo $\mathsf{E}(\Box)$ and on the set $X(\Box)$, and write $\mathbb{T} \subset \mathsf{S}$ to emphasize this point. We say that \mathbb{T} is \equiv -confluent (resp. terminating, or *terminating modulo* E) if every sub-abstract RSM $\mathbb{T}(\Box)$ is \equiv -confluent (resp. terminating modulo $\mathsf{E}(\Box)$), and similarly for other notions of abstract rewriting theory.

One can always consider S as its own abstract sub-system, so that the notion of abstract subsystem generalizes the notion of higher RSM. We refer to the case $\mathbb{T} = S$ as *context-dependent*, and to the case $\mathbb{T} \subsetneq S$ as *context-agnostic*.

3.4.2 Compatibility and contextualization

The notion of compatible abstract preorder generalizes verbatim to the higher setting:

Definition 3.47. Let $S = (R, E; \equiv)$ be a higher RSM and $\mathbb{T} \subset S$ an abstract sub-system. A preorder on \mathbb{R}_2^* is said to be compatible with \mathbb{T} if it is compatible with every sub-abstract RSM $\mathbb{T}(\Box)$, in the sense of Definition 3.15. We denote $\succ_{\mathbb{T}}$ the smallest preorder compatible with \mathbb{T} .

E-invariance of \succ is defined similarly.

We now describe contextualization:

Lemma 3.48. Let $S = (R, E; \equiv)$ be a higher RSM and $\mathbb{T} \subset S$ an abstract sub-system. Let (f, g) be a local \mathbb{T} -branching and Γ a context. Assume (f, g) admits a (\mathbb{T}, \equiv) -confluence (f', g') such that $\Gamma[f', g']$ belongs to \mathbb{T} . Then $\Gamma[f, g]$ is \mathbb{T} -confluent.

Proof. Since \equiv is a higher equivalence, $f' \circ f \equiv g' \circ g$ implies that $\Gamma[f' \circ f] \equiv \Gamma[g' \circ g]$.

In particular, in the context-agnostic case:

Lemma 3.49. Let $S = (R, E; \equiv)$ be a higher RSM. Let (f, g) be a local S-branching and Γ a context. If (f, g) is (S, \equiv) -confluent, then $\Gamma[f, g]$ is (S, \equiv) -confluent.

3.4.3 Independent branchings

The higher structure of a higher RS (P; \equiv) induces canonical types of local P-branchings. A local P-branching is *independent* if it is of the form $(f,g) = (f \star_1 \psi, \phi \star_1 g)$, for P-rewriting steps $f: \phi \to \phi'$ and $g: \psi \to \psi'$:



When working modulo, that is with a higher RSM $S = (R, E; \equiv)$, independent branchings refer to independent R-branchings. Note that the property of being an independent branching is preserved by context: if (f, g) is an independent branching, so is $\Gamma[f, g]$. A local S-branching (or an S-local triple) is said to be *overlapping* if it is not branchwise E-congruent to an independent R-branching.

An independent branching always have a canonical S-confluence, given by

$$(g',f') \coloneqq (\phi' \star_1 \mathsf{g},\mathsf{f} \star_1 \psi')$$

The following is tautological, thanks to the independence axiom:

Lemma 3.50. Let $S = (R, E; \equiv)$ be a higher RSM and $\mathbb{T} \subset S$ an abstract sub-system. Let (f, g) is an independent \mathbb{T} -branching. If its canonical \mathbb{T} -confluence (g', f') is in \mathbb{T} , then it defines a (\mathbb{T}, \equiv) -confluence for (f, g).

In particular, in the context-agnostic case:

Lemma 3.51. Let $S = (R, E; \equiv)$ be a higher RSM. Every independent S-branching is (S, \equiv) -confluent.

3.4.4 Higher Newmann's lemma

In the context-agnostic case, the above results can be gathered in a single black-box:

Lemma 3.52 (higher Newmann's lemma). Let $S = (R, E; \equiv)$ be a higher RSM. If S is terminating and if every overlapping local S-branching is, up to branchwise E-congruence, a contextualization of $a(S, \equiv)$ -confluent branching, then S is convergent.

Proof. This follows from the (abstract) Newmann's lemma (Lemma 3.13), in combination with the branchwise confluence lemma (Lemma 3.17), confluence of contextualizations (Lemma 3.49) and confluence of independent branchings (Lemma 3.51). \Box

3.4.5 Independent rewriting

Recall the abstract notion of rewriting branchings as introduced in Subsection 3.2.4. In the higher setting, one often wants to rewrite part of a diagram away from a given branching. This gives the notion of *independent rewriting*. Given rewriting steps $f: \phi \to \phi_1$, $g: \phi \to \phi_2$ and $h: \psi \to \psi'$, setting

$$f = \mathsf{f} \star_1 \psi, \quad g = \mathsf{g} \star_1 \psi \quad \text{and} \quad h = \phi \star_1 \mathsf{h}$$

defines three pairs of branchings (f, g), (f, h) and (h, g), the latter two being independent branchings. Define also

$$f' = \mathsf{f} \star_1 \psi', \quad g' = \mathsf{g} \star_1 \psi', \quad h_1 = \phi_1 \star_1 \mathsf{h} \quad \text{and} \quad h_2 = \phi_2 \star_1 \mathsf{h}.$$

We say that the branching (f, g) rewrites into the branching (f', g') via the triple (h_1, h, h_2) (and similarly if the vertical positions of f, g and h are swapped), as pictured below:



We have the following lemma:

Lemma 3.53. Let $S = (R, E; \equiv)$ be a higher RSM, $\mathbb{T} \subset S$ an abstract sub-system and \succ an E-invariant preorder on \mathbb{R}_2^* compatible with \mathbb{T} . In the situation above, assume (f, g) is a \mathbb{T} -branching. If h, h_1 and h_2 are in \mathbb{T} and if (f', g') is \succ -tamed (\mathbb{T}, \equiv) -congruent, then so is (f, g).

In particular, in the context-agnostic case:

Lemma 3.54. Let $S = (R, E; \equiv)$ be a higher RSM and \succ an E-invariant preorder on R_2^* compatible with S. In the situation above, if (f', g') is \succ -tamed (S, \equiv) -congruent, then so is (f, g).

3.4.6 Gray rewriting system modulo

A *Gray rewriting system* ($P; \equiv$) is the data of a higher RS (P, \equiv) such that P is a Gray polygraph and \equiv verifies the interchange naturality axiom:

interchange naturality axioms: for all 0-composable A, g, β with $A: \phi \Rightarrow \phi': f \Rightarrow f' \in \mathsf{P}_3, g \in \mathsf{P}_1^*$ and $\beta: h \Rightarrow h' \in \mathsf{P}_2$, and for all 0-composable α, g, B with $\alpha: f \Rightarrow f' \in \mathsf{P}_2, g \in \mathsf{P}_1^*$ and $B: \psi \Rightarrow \psi': h \Rightarrow h' \in \mathsf{P}_3$, we have

$$\begin{aligned} X_{\phi',g,\beta} \star_2 \left(\left(A \star_0 g \star_0 h' \right) \star_1 \left(f \star_0 g \star_0 \beta \right) \right) \\ &\equiv \left(\left(f' \star_0 g \star_0 \beta \right) \star_1 \left(A \star_0 g \star_0 h \right) \right) \star_2 X_{\phi,g,\beta} \end{aligned}$$

and
$$\begin{aligned} X_{\alpha,g,\psi'} \star_2 \left(\left(\alpha \star_0 g \star_0 h' \right) \star_1 \left(f \star_0 g \star_0 B \right) \right) \\ &\equiv \left(\left(f' \star_0 g \star_0 B \right) \star_1 \left(\alpha \star_0 g \star_0 h \right) \right) \star_2 X_{\alpha,g,\psi'} \end{aligned}$$

where $X_{\phi',g,\beta}$ denotes a composition of interchange generators, interchanging 2-cells in ψ with the 2-cell β , and similarly for the other Xs. These interchange naturality axioms can be pictured as:



A *Gray rewriting system modulo* (Gray RSM) $S = (R, E; \equiv)$ is the data of a higher RSM such that E is a Gray polygraph.

3.5 Higher linear rewriting modulo

This section finally introduces (weak) higher linear rewriting modulo, building on all the previous sections. In particular, linear Gray rewriting modulo is defined in Subsection 3.5.6. Similarly to Subsection 3.4, while we restrict to 3-dimensional rewriting, many notions are not specific to the 3-dimensional case; in particular, this could be adapted to the 2-dimensional case (rewriting modulo in linear categories, including algebras).

3.5.1 Higher linear rewriting system modulo

Recall the notions introduced in Subsection 2.3.

Definition 3.55. A higher linear rewriting system (*higher linear RS*) is the same data as a linear 3-sesquipolygraph P. A higher linear rewriting system modulo (*higher linear RSM*) S = (R, E) is the data of a left-monomial 3-sesquipolygraph R and a monomial-invertible 3-sesquipolygraph E, such that $R_{\leq 2} = E_{\leq 2}$.

As in the abstract higher case, a higher linear RSM S = (R, E) defines a category of linear 1-polygraphs $S(\Box) = (\mathcal{B}(\Box), R(\Box), E(\Box))$, setting

$$\mathcal{B}(\Box) \coloneqq \mathsf{R}_2^*(\Box) = \mathsf{E}_2^*(\Box),$$
$$\mathsf{R}(\Box) \coloneqq \operatorname{Cont}(\mathsf{R})(\Box) \quad \text{and} \quad \mathsf{E}(\Box) \coloneqq \operatorname{Cont}(\mathsf{E})(\Box),$$

for each 1-sphere \Box in $\mathsf{R}_1^* = \mathsf{E}_1^*$. Each context Γ on \Box defines a morphism of linear 1-polygraphs:

$$\Gamma \colon \mathbf{S}(\Box) \to \mathbf{S}(s_1(\Gamma), t_1(\Gamma)).$$

Contrary to the abstract higher case however, $S(\Box)$ *needs not be a linear RSM*, as we did not impose the adaptedness condition appearing in Definition 3.23.

As for the abstract higher case, we define a notion of sub-systems, imposing now the adaptedness condition:

Definition 3.56. Let S = (R, E) be a higher linear RSM. A linear sub-system T is the data of a family of sub-sets $T(\Box) \subset R_3^*(\Box)$ for each 1-sphere \Box in R_1^* . Moreover, we assume the adaptedness condition:

 $s(r) \notin \widetilde{\operatorname{supp}}_{\mathsf{E}}(t(r)) \quad \forall r \in \mathsf{T}.$

We say that S is adapted if S is a linear sub-system of itself.

As in the higher abstract case, we think of T as defining a family of sub-linear RSM $T(\Box) \subset S(\Box)$ modulo $E(\Box)$, and write $T \subset S$ to emphasize that point. We say that T is \equiv -confluent (resp. terminating, or terminating modulo E) if every sub-linear RSM $T(\Box)$ is confluent (resp. terminating modulo $E(\Box)$), and similarly for other notions of linear rewriting theory.

Note that adaptedness *must* be defined via a linear sub-system, since in general, it depends on context: if r is adapted, $\Gamma[r]$ needs not be.

A T-rewriting step has the following general form (compare with Definition 3.29):

$$\alpha \colon u \sim_{\mathbf{E}} \lambda \Gamma[s(r)] + v \longrightarrow_{\mathbf{T}} \lambda \Gamma[t(r)] + v \sim_{\mathbf{E}} w.$$

We say that α is of type r^{22} As before, α is positive if $\Gamma[s(r)] \notin \widetilde{\operatorname{supp}}_{\mathsf{E}}(v)$. Note that contextualization needs not preserve positivity: we may have

 $\Gamma'[\Gamma[s(r)]] \in \dot{\widetilde{\operatorname{supp}}}_{\mathsf{E}}(\Gamma'[v]), \quad \text{ even if } \quad \Gamma[s(r)] \notin \dot{\widetilde{\operatorname{supp}}}_{\mathsf{E}}(v).$

However, it is true that if α is monomial (that is, if v = 0) then $\Gamma[\alpha]$ is monomial.

Remark 3.57. In [All18a; All18b], [Dup21; Dup22] and [DEL21], the authors used the notion of quasi-termination to deal with context-dependent termination issues, and suggested a basis theorem based on quasi-terminating normal forms. Our approach gives a different way to deal with the same issue, avoiding the use of quasi-terminating normal forms altogether. It is also more general, as it applies to settings that are not quasi-terminating, such as the setting of graded \mathfrak{gl}_2 -foams studied in Section 4.

Quasi-terminating normal forms are typically badly behaved. For instance, a monomial in the support of a quasi-normal form needs not be a quasi-normal form, and a linear combination of quasi-normal forms needs not be a quasi-normal form; compare with Remark 3.34. Extracting a basis from a quasi-terminating system is hazardous. For instance, one can consider the quasi-terminating linear rewriting system (non-modulo) ($\mathcal{B} = \{a, b, c\}, \mathbb{R} = \{a \rightarrow b + c, b \rightarrow a - c\}$): the set of monomial quasi-normal forms is $\mathcal{B}NF_{\mathbb{R}} = \mathcal{B} = \{a, b, c\}$, which is not a basis of the module presented by \mathbb{R} .

3.5.2 Strong compatibility

Recall that for a linear RSM S = (B; R, E), a preorder \succ is compatible (Definition 3.38) if for all $r \in R$, $s(r) \succ b$ for all $b \in supp_{\mathsf{E}}(t(r))$. With the hypothesis of Definition 3.58, we say that T is *compatible* is each linear RSM $T(\Box)$ is compatible, that is, if

$$s(r) \succ b$$
, for all $r \in T$ and $b \in \widetilde{\operatorname{supp}}_{\mathsf{F}}(t(r))$.

As we shall see, compatibility is *not* sufficient for higher rewriting, and a stronger condition is required:

Definition 3.58. Let S = (R, E) be a higher linear RSM and T a linear-sub system. An E-invariant preorder \succ on R_2^* is said to be strongly compatible with T if:

$$\Gamma[s(r)] \succ b$$
, for all $r \in \mathsf{R}$, context Γ such that $\Gamma[r] \in \mathsf{T}$, and $b \in \Gamma[\widetilde{\operatorname{supp}}_{\mathsf{E}}(t(r))]$.

Strong compatibility is indeed stronger than compatibility:

Lemma 3.59. Let S = (R, E) be a higher linear RSM, T a linear-sub system and \succ an E-invariant preorder on \mathbb{R}_2^* . If \succ is strongly compatible with T, then \succ is compatible with T.

Proof. It suffices to choose the trivial context in the strong compatibility condition.

However, the converse of Lemma 3.59 does not hold! At best, one can say that since $\Gamma[r] \in T$, compatibility implies that $\Gamma[s(r)] = s(\Gamma[r]) \succ b$ for all $b \in \widetilde{\operatorname{supp}}_{\mathsf{E}}(t(\Gamma[r])) = \widetilde{\operatorname{supp}}_{\mathsf{E}}(\Gamma[t(r)])$. This is *not* sufficient to conclude, as the inclusion

$$\widetilde{\operatorname{supp}}_{\mathsf{E}}(\Gamma[t(r)]) \subset \Gamma[\widetilde{\operatorname{supp}}_{\mathsf{E}}(t(r))]$$

²²This is a slight abuse of notation: as belonging to a linear RSM, α is of type $\Gamma[r]$ according to Definition 3.28.

needs not be an equality! Consider more generically a vector w and the inclusion

$$\widetilde{\operatorname{supp}}_{\mathsf{E}}(\Gamma[w]) \subset \Gamma[\widetilde{\operatorname{supp}}_{\mathsf{E}}(w)]$$

Recall that contextualization needs not not act freely (see Subsection 1.2.4): there may exist monomials b_1, b_2 such that $b_1 \not\sim_{\mathsf{E}} b_2$ but $\Gamma[b_1] \sim_{\mathsf{E}} \Gamma[b_2]$. In particular, if $w = \lambda_1 b_1 + \lambda_2 b_2$ and λ_1, λ_2 have well-chosen scalars, we have $\Gamma[w] = 0$ and thus $\Gamma[b_1] \notin \widetilde{\operatorname{supp}}_{\mathsf{E}}(\Gamma[w])$.

Remark 3.60. Super \mathfrak{gl}_2 -foams give an explicit example where compatibility does not imply strong compatibility. Consider \Bbbk as in Definition 4.4 with X = Z = 1 and Y = -1; this is the super case. We let S = (R, E) and T be as in Subsection 4.2. Let $\widetilde{T} \subset T$ be the linear sub-system of T consisting only in rewriting steps of type nc; that is, \widetilde{T} consists in rewriting steps of type nc for which the two strands are distinct. Let $\widetilde{\succ}$ be the preorder on S_2^* which compares the number of *open shadings*—shadings that are not enclosed by a strand. The preorder is $\widetilde{\succ}$ compatible with \widetilde{T} . Compatibility is clear for $\Gamma[nc] \in \widetilde{T}$ such that both strands are not closed, as the number of open shadings strictly decreases. On the other hand, if at least one of the strand is closed, then the target of $\Gamma[nc]$ is zero:



Hence $\Gamma[\mathsf{nc}]$ is trivially compatible with \succ . However, the same situation prevents the preorder \succeq from being strongly compatible with $\widetilde{\mathsf{T}}$, as the number of open shadings is the same for $\Gamma[s(\mathsf{nc})]$ and b, for all $b \in \Gamma[supp_{\mathsf{E}}(t(\mathsf{nc}))]$.

3.5.3 Contextualization

We now study how confluence, tamed congruence and contextualization relates.

Lemma 3.61. Let S = (R, E) be a higher linear RSM, $T \subset S$ a linear sub-system and \succ an E-invariant preorder on R_2^* strongly compatible with T. Consider a composition $g \star_2 f$, where f is a monomial S-rewriting step and g is a positive S-rewriting sequence. If Γ is a context such that both $\Gamma[f]$ and $\Gamma[g]$ are in T, then $\Gamma[g]$ is \succ -tamed by $\Gamma[s(f)]$.

Proof. Decomposing g as a composition of positive S-rewriting steps, the situation is:

$$s(f) \xrightarrow{f} v_0 \xrightarrow{g_0} v_1 \xrightarrow{g_1} \cdots \xrightarrow{g_{n-1}} v_n$$

$$\Gamma[s(f)] \xrightarrow{\Gamma[f]} \Gamma[v_0] \xrightarrow{\Gamma[g_0]} \Gamma[v_1] \xrightarrow{\Gamma[g_1]} \cdots \xrightarrow{\Gamma[g_{n-1}]} \Gamma[v_n]$$

We wish to show that $\Gamma[s(f)] \succ \operatorname{supp}_{\mathsf{E}}(\Gamma[v_i])$ for all $0 \le i \le n$. Rather, given the inclusion $\dot{\operatorname{supp}}_{\mathsf{E}}(\Gamma[w]) \subset \Gamma\left[\operatorname{supp}_{\mathsf{E}}(w)\right]$ which holds for generic vector w, it suffices to show

$$\Gamma[s(f)] \succ \Gamma\left[\dot{\operatorname{supp}}_{\mathsf{E}}(v_i)\right]$$
(10)

for all $0 \le i \le n$.

When i = 0, we have $v_0 = t(f)$. By hypothesis $\Gamma[f] \in T$, so that (10) follows from strong compatibility. It remains to show that

$$\Gamma\left[\overset{\cdot}{\operatorname{supp}}_{\mathsf{E}}(v_i)\right] \succeq^{\operatorname{rel}} \Gamma\left[\overset{\cdot}{\operatorname{supp}}_{\mathsf{E}}(v_{i+1})\right].$$

(Recall the definition of relative relation from Definition 3.42.) Write $g_i = \lambda r + v$, its canonical decomposition as a positive S-rewriting step. On the one hand:

$$\Gamma\left[\overset{\cdot}{\operatorname{supp}}_{\mathsf{E}}(v_{i+1})\right] = \Gamma\left[\overset{\cdot}{\operatorname{supp}}_{\mathsf{E}}(t(r)+v)\right] \subset \Gamma\left[\overset{\cdot}{\operatorname{supp}}_{\mathsf{E}}(t(r))\right] \cup \Gamma\left[\overset{\cdot}{\operatorname{supp}}_{\mathsf{E}}(v)\right].$$

On the other hand, thanks to the positivity of g_i , we have $\widetilde{\operatorname{supp}}_{\mathsf{E}}(s(r) + v) = \{s(r)\} \sqcup \overset{\cdot}{\operatorname{supp}}_{\mathsf{E}}(v)$, and hence

$$\Gamma\left[\overset{\cdot}{\operatorname{supp}}_{\mathsf{E}}(v_i)\right] = \Gamma\left[\overset{\cdot}{\operatorname{supp}}_{\mathsf{E}}(s(r)+v)\right] = \{\Gamma[s(r)]\} \cup \Gamma\left[\overset{\cdot}{\operatorname{supp}}_{\mathsf{E}}(v)\right]$$

Because $\Gamma[g_i] \in T$, strong compatibility implies that $\Gamma[s(r)] \succ \Gamma\left[\overset{\cdot}{\operatorname{supp}}_{\mathsf{E}}(t(r))\right]$. This concludes.

Corollary 3.62 (CONTEXTUALIZATION LEMMA). Let S = (R, E) be a higher linear RSM, $T \subset S$ a linear sub-system and \succ an E-invariant preorder on R_2^* strongly compatible with T. Let (f, g) be a monomial local S-branching which admits a positive S-confluence (f', g'). If Γ is a context such that both $\Gamma[f, g]$ and $\Gamma[f', g']$ belong to T, then $\Gamma[f', g']$ is a \succ -tamed T^{st} -congruence for $\Gamma[f, g]$.

In general, $\Gamma[f', g']$ needs not be positive,²³ as the example (11) below illustrates (it is taken from Lemma 4.32). Recall that plain arrows denote positive rewriting steps, while dashed arrows denote not-necessarily-positive rewriting steps. In the example below, the graded interchange law is part of the modulo. On the left-hand side, the branches of the confluence (labelled "dd") evaluate to zero, so that only the other diagram remains. The two diagrams on the top (resp. on the bottom) are *not* projectively congruent modulo, so that both branches of the confluence are positive. The same confluent branching is pictured on the right-hand side, contextualized with a cap and cup. Because the graded interchange law is part of the modulo, dots can now freely move between the top and the bottom. In particular, the two diagrams on the top (resp. on the bottom) are now projectively congruent modulo, and the branches of the confluence are not positive anymore.



In the context-agnostic setting, the CONTEXTUALIZATION LEMMA 3.62 reduces to the following:

Corollary 3.63. Let S = (R, E) be an adapted higher linear RSM and \succ an E-invariant preorder on R_2^* strongly compatible with S. Let (f, g) be a monomial local S-branching and Γ a context. If (f, g) is positively S-confluent, then $\Gamma[f, g]$ is \succ -tamely S-congruent.

²³ In particular, there are related gaps in the proofs of [All18a, lemma 4.2.12] and [Dup22, theorem 2.2.9].

3.5.4 Independent branchings

As we have seen, for higher RSM the statement that independent branchings are \equiv -confluent is tautological (Lemma 3.51). As we now explain, the linear case is more subtle.

Fix P a higher linear RS and (f, g) an independent local P⁺-branching. Without loss of generality, this means that $f = f \star_1 s(g)$ and $g = s(f) \star_1 g$, for some monomial P⁺-rewriting steps f and g. The P-confluence $(g', f') = (t(f) \star_1 g, f \star_1 t(g))$ defines a canonical P-confluence for (f, g). Write:

$$f: s(f) \to \lambda_1 x_1 + \ldots + \lambda_m x_m$$
 and $g: s(g) \to \mu_1 y_1 + \ldots + \mu_n y_n$,

where the x_i 's (resp. y_j 's) are monomials with pairwise distinct E-projective classes. Then the P-confluence (g', f') can be explicitly decomposed as a Pst-confluence, setting $f' = f'_n \star_2 \ldots \star_2 f'_1$ and $g' = g'_m \star_2 \ldots \star_2 g'_1$, where:

$$f'_j = \sum_{k < j} \mu_k(t(\mathbf{f}) \star_1 y_k) + \mu_j(\mathbf{f} \star_1 y_j) + \sum_{j < k} \mu_k(s(\mathbf{f}) \star_1 y_k)$$

and
$$g'_i = \sum_{k < i} \lambda_k(x_k \star_1 t(\mathbf{g})) + \lambda_i(x_i \star_1 \mathbf{g}) + \sum_{i < k} \lambda_k(x_k \star_1 s(\mathbf{g})).$$

Lemma 3.64 (independent branchings, context-dependent case). Let S = (R, E) be an adapted higher linear RSM, $T \subset S$ a linear sub-system and \succ an E-invariant preorder strongly compatible with T. Let (f, g) be an independent local T^+ -branchings. If for each $1 \le j \le n$ (resp. $1 \le i \le m$) and with the notations above, the S^+ -rewriting step $f \star_1 y_j$ (resp. $x_i \star_1 g$) is in T, then the canonical T^{st} -confluence (g', f') is \succ -tamed by s(f) = s(g).

Proof. For each $1 \leq j \leq n$, the E-projective supports of $s(f'_j)$ and $t(f'_j)$ are contained in the following set:

$$M \coloneqq \bigcup_{1 \le j \le n} \dot{\widetilde{\operatorname{supp}}}_{\mathsf{E}} \big(t(\mathsf{f}) \star_1 y_j \big) \cup \bigcup_{1 \le j \le n} \dot{\widetilde{\operatorname{supp}}} \big(s(\mathsf{f}) \star_1 y_j \big).$$

It follows from strong compatibility that $s(f) \star_1 s(g) \succ s(f) \star_1 y_j$ (since $g = s(f) \star_1 g$ belongs to T) and that $s(f) \star_1 y_j \succ t(f) \star_1 y_j$ (since $f \star_1 y_i$ belongs to T). By transitivity of the preorder \succ , we have $s(f) \star_1 s(g) \succ t(f) \star_1 y_j$. Hence:

$$s(\mathsf{f}) \star_1 s(\mathsf{g}) \succ M$$

In other words, f' is \succ -tamed by $s(f) \star_1 s(g)$.

An analogous argument shows that g' is \succ -tamed by $s(f) \star_1 s(g)$. This concludes.

Note the necessity of strong compatibility in the proof of Lemma 3.64: $s(f) \star_1 s(g) \succ s(f) \star_1 y_j$ does *not* follow from compatibility alone, as while y_j is in the E-projective support of t(g), it may be that $s(f) \star_1 y_j$ is *not* in the support of $s(f) \star_1 t(g)$.

The proof of Lemma 3.64 resembles the proof of Theorem 4.2.1, part (iii) in [GHM19], with tameness appearing implicitly. In our terminology, their setting (associative algebra) restrict to context-agnostic systems, and as contextualization acts freely, there is no distinction between compatibility and strong compatibility.

The following is a direct corollary of Lemma 3.64:

Lemma 3.65 (independent branchings, context-agnostic case). Let S = (R, E) be an adapted higher linear RSM and \succ an E-invariant preorder on R_2^* strongly compatible with S. Every independent S^+ -branching is \succ -tamely S^{st} -congruent.

In the context-dependent case, one cannot rely on such a general statement. However:

Lemma 3.66. Let S = (R, E) be a higher linear RSM, $T \subset S$ a linear sub-system and \succ an E-invariant preorder on R_2^* strongly compatible with T. Assume given another linear sub-system $B \subset T$, such that B^+ is convergent. If every $(S \setminus T)$ -rewriting step is B-congruent, then every independent T-branching is \succ -tamely T^{st} -congruent.

This will be the situation of graded \mathfrak{gl}_2 -foams.

Proof. We adapt the proof of Lemma 3.64, and borrow its notations. Assume, up to reordering the y_j 's, that $f \star_1 y_j$ is in T for $1 \leq j \leq k$ and that $f \star_1 y_j$ is *not* in T for $k < j \leq m$. By the same argument as in the proof of Lemma 3.64, $f'_k \star_2 \ldots \star_2 f'_1$ is a Tst-rewriting sequence \succ -tamed by $s(f) \star_1 s(g)$. On the other hand, by the hypothesis of the lemma, $f'_m \star_2 \ldots \star_2 f'_{k+1}$ is B-congruent. Arguing similarly for g', we get a T-congruence for (f, g) given as a B-congruence "sandwiched" between two \succ -tamed Tst-congruence. Since B⁺ is convergent, we can replace the B-congruence by a B⁺-confluence. Compatibility of \succ with B and transitivity of \succ^+ conclude.

3.5.5 Independent rewriting

We consider the linear analogue of abstract independent rewriting defined in Subsection 3.4.5. Let us use the same notations with f, g and h monomial rewriting steps, which decompose as:

$$f: \phi \to_{\mathsf{R}} \lambda_1 x_1 + \ldots + \lambda_l x_l,$$

g: $\phi \to_{\mathsf{R}} \mu_1 y_1 + \ldots + \mu_m y_m,$
and h: $\psi \to_{\mathsf{R}} \nu_1 z_1 + \ldots + \nu_n z_n,$

where x_i 's (resp. y_j 's and $z'_k s$) are monomials belonging to pairwise distinct E-projective classes. In these decompositions, the independent rewriting of (f, g) into (f', g') via the triple (h, h_1, h_2) is pictured as follows:

$$\begin{array}{c} & \underset{f=\mathsf{f}\star_{1}\psi}{\overset{h_{1}=\sum_{i}\lambda_{i}(x_{i}\star_{1}\mathsf{h})}{f'=\sum_{k}\nu_{k}(\mathsf{f}\star_{z_{k}})}} \sum_{i,k}\lambda_{i}\nu_{k}(x_{i}\star_{1}z_{k}) \\ \phi \star_{1}\psi \xrightarrow{h=\phi\star_{1}\mathsf{h}}{\overset{h=\phi\star_{1}\mathsf{h}}{\longrightarrow}} \sum_{k}\nu_{k}(\phi\star_{1}z_{k}) \\ g=\mathsf{g}\star_{1}\psi \xrightarrow{j}{\overset{j}{\longrightarrow}} \mu_{j}(y_{j}\star_{1}\psi) \xrightarrow{g'=\sum_{k}\nu_{k}(\mathsf{g}\star_{z_{k}})}{\overset{h=\phi\star_{1}\mathsf{h}}{\longrightarrow}} \sum_{j,k}\mu_{j}\nu_{k}(y_{j}\star_{1}z_{k}) \end{array}$$

Lemma 3.67 (INDEPENDENT REWRITING LEMMA). Let S = (R, E) be a higher linear RSM, $T \subset S$ a linear sub-system and \succ an E-invariant preorder on R_2^* strongly compatible with T. Let f, g and h monomial R^+ -rewriting steps as above, such that:

- (i) $f = f \star_1 \psi$, $g = g \star_1 \psi$ and $h = \phi \star_1 h$ are in T, and for each $1 \le i \le l$ (resp. $1 \le j \le m$, resp. $1 \le k \le n$), the rewriting step $x_i \star_1 h$ (resp. $y_j \star_1 h$) is in T,
- (ii) for each $1 \le k \le n$, the branching $(f \star_1 z_k, g \star_1 z_k)$ is \succ -tamely T^{st} -congruent,

then (f, g) is \succ -tamely T^{st} -congruent.

Proof. Since $h = \phi \star_1 h$ is in T, strong compatibility implies that $\phi \star_1 \psi \succ \phi \star_1 z_k$ (for all suitable k). Let h'_k be a Tst-congruence for $(f \star_1 z_k, g \star_1 z_k) \succ$ -tamed by its source $\phi \star_1 z_k$, and hence by $\phi \star_1 \psi$ by transitivity of \succ . It follows that the Tst-congruence $\sum_k \nu_k h'_k$ is \succ -tamed by $\phi \star_1 \psi$.

Moreover, strong compatibility implies that $\phi \star_1 \psi \succ x_i \star_1 \psi$ (since $f = f \star_1 \psi$ belongs to T) and $x_i \star_1 \phi \succ x_i \star_1 z_k$ (since $x_i \star_h$ belongs to T), for all suitable *i* and *k*-hence $\phi \star_1 \psi \succ x_i \star_1 z_k$ for all suitable *i* and *k*, by transitivity of \succ . It follows that $h_1 = \sum_i \lambda_i (x_i \star_1 h)$ admits a decomposition as a Tst-congruence \succ -tamed by $\phi \star_1 \psi$. Similarly, $h_2 = \sum_j \lambda_j (y_j \star_1 h)$ admits a Tst-congruence \succ -tamed by $\phi \star_1 \psi$, using the hypotheses that $g = g \star_1 \psi$ and $y_i \star_1 h$ (for all suitable *i*) belong to T.

We conclude that the triple $(h_1, \sum_k \nu_k h'_k, h_2)$ defines a Tst-congruence for (f, g), \succ -tamed by its source $\phi \star_1 \psi$.

Lemma 3.68 (independent rewriting lemma, context-agnostic case). Let S = (R, E) be a higher linear RSM and \succ an E-invariant preorder on R_2^* strongly compatible with S. If for each $1 \le k \le n$, the branching $(f \star_1 z_k, g \star_1 z_k)$ is \succ -tamely Tst-congruent, then (f, g) is \succ -tamely Tst-congruent.

3.5.6 Linear Gray rewriting system modulo

Linear Gray rewriting modulo is simply higher linear rewriting modulo where the modulo contains graded interchangers. Below are the formal definitions.

Recall the notations G and μ from Notation 2.10:

Definition 3.69. A (G, μ) -graded linear Gray rewriting system modulo is a higher linear RSM S = (R, E) such that E is a (G, μ) -graded linear Gray polygraph.

In particular, we can specialize to the scalar case:

Definition 3.70. $A(G, \mu)$ -scalar Gray rewriting system modulo S := (R, E; scl) *is the data of two scalar G-graded 3-sesquipolygraphs* (R; scl) *and* (E; scl) *with* $R_2^* = E_2^*$, *such that* E *is a* (G, μ) -*scalar Gray polygraph.*

Remark 3.71. Gray rewriting modulo corresponds to rewriting modulo in strict 2-categories, and linear Gray rewriting corresponds to rewriting in graded-2-categories. In particular, linear Gray rewriting where μ is trivial corresponds to rewriting in linear strict 2-categories.

3.6 Summary

Given a presented linear 2-sesquicategory, how can one use rewriting theory to find a basis? This section summarizes the main tools developed throughout this section. While we focus on the linear case (which is more involved), the same ideas apply for higher (non-linear) rewriting modulo.

3.6.1 The setup

Let C be a linear 2-sesquicategory presented by a linear 3-sesquipolygraph P in the sense of Definition 2.11. Assume the following choices of data has been made:

- (a) a splitting of P as $P = R \sqcup E$, defining a higher linear RSM S = (R, E) presenting C (see Definition 3.55),
- (b) a linear sub-system $T \subset S$ (see Definition 3.56),
- (c) a preorder \succ strongly compatible with T (see Definition 3.58).

Typically, C would be a graded-2-category and E would contain the 3-sesquipolygraph of graded-interchangers, making the data S = (R, E) a linear Gray rewriting system. To find a basis using the BASIS-FROM-CONVERGENCE THEOREM 3.36, the following needs to be checked:

- (α) S and T present the same underlying (family of) module(s): it suffices to show that every $(S \setminus T)^+$ -rewriting step is T-congruent, so that two vectors are S-congruent if and only if they are T-congruent (see Lemma 3.30).
- (β) scl(E)^{\top} is scalar-coherent on β NF_T: recall that β NF_T denotes the set of monomial T⁺normal forms (Definition 3.33), and that scl(E)^{\top} is scalar-coherent on β NF_T (Definition 3.2 and Remark 3.3) if for all $b \in \beta$ NF_T, the existence of an E-congruence $b \sim_{\mathsf{E}} \lambda b$ for some scalar $\lambda \in \mathbb{k}$ implies that $\lambda = 1$.

Showing scalar-coherence of $\operatorname{scl}(\mathsf{E})^{\top}$ on $\mathcal{B}\operatorname{NF}_T$ can be done either using ad-hoc arguments, or by apply higher rewriting modulo theory to $\operatorname{scl}(\mathsf{E})^{\top}$, together with Proposition 3.8 (coherence from convergence).

(γ) T⁺ *is convergent*: we use the TAMED LINEAR NEWMANN'S LEMMA 3.43, showing on one hand the order \succ is terminating,²⁴ and on the other hand that every monomial local T⁺-branching is \succ -tamely Tst-congruent.

The analysis of monomial local T⁺-branching is the hardest task, which we now discuss. Of course, in the context-agnostic case T = S, (α) is automatic.

Remark 3.72. In the example of superadjunction °P given in the extended summary (Subsection 1.2), we have:

- (α) we are in a context-agnostic setting, so (α) is automatic;
- (β) using the coherence theorem for interchangers [Sch24, Theorem A.3.1], one can check that $scl({}^{\circ}E)^{\top}$ is scalar-coherent on ${}^{\circ}P_{2}^{*}$, and a fortiori on $\mathcal{B}NF_{{}^{\circ}S}$;
- (γ) as said in Subsection 1.2.3, the order \succ compares the number of generating 2-cells.

3.6.2 How to classify monomial local branchings?

Working modulo makes classifying monomial local branchings difficult, as it considerably increases the number of rewriting steps. However, in the context of diagrammatic algebra, the modulo data typically has a topological interpretation, which can be leveraged: e.g. rectilinear isotopies when working modulo interchangers, or planar isotopies when working modulo a pivotal structure. We describe here a general strategy to classify local branchings, to be adapted depending on the example at hand:

- (i) Understand coherence of the modulo. In other words, provide a topological or combinatorial description of when two monomial 2-cells are projectively E-congruent. this can be done either using ad-hoc arguments, or by applying higher rewriting modulo theory to scl(E). In practice, (β) in the previous section comes as a byproduct.
- (ii) Describe naturalities of the modulo. The modulo typically captures some underlying categorical structure, which should come with natural compatibilities with the rewriting steps of the rewriting system. For instance, a Gray RSM comes with interchange naturalities (see Subsection 3.4.6). We call these E-naturalities.

²⁴While this will not be the case in this paper, defining a terminating order may be hard in general. For this problem, the method of derivation of Guiraud and Malbos may be useful [GM18].

- (iii) Characterize rewriting steps modulo. In other words, provide a topological or combinatorial description of when two rewriting steps are E-congruent. (One needs not show that this characterization captures *all* E-congruences.) To do so, use the coherence statement to express isotopies in terms of E-naturalities. In fact, given the BRANCHWISE TAMED CONGRUENCE LEMMA 3.19, one can instead characterize when two rewriting steps are \succ -tamed T⁺-congruent.
- (iv) Classify monomial local branchings. Using the characterization above, provide a list of monomial local T⁺-branchings, called *critical branchings*, such that every monomial local T⁺-branching rewrites into (a linear combination of) either an independent branching or a branchwise ≻-tamely Tst-congruent to a contextualization of a critical branching.

In the last point (iv), three main tools are at play: the INDEPENDENT REWRITING LEMMA 3.67, the BRANCHWISE TAMED CONGRUENCE LEMMA 3.19 and the CONTEXTUALIZATION LEMMA 3.62. The results of Subsection 3.5.4 also help to deal with independent branchings. Hopefully, this leaves only a few critical branchings for which an explicit computation is needed, and one can conclude that every monomial local T^+ -branching is \succ -tamely T^{st} -congruent.

Remark 3.73. If E is scalar-free (Remark 3.26), the strategy described above greatly simplifies. For instance, this is the case of linear Gray rewriting modulo scalar-free (or non-graded) interchangers, which coincides with rewriting in linear strict 2-categories [All18a]. However, even in that case it is useful to explicitly consider the interchange law as part of the modulo, given the caveats related to freeness of contextualization, described in Subsection 1.2.4 (see also Footnote 23).

Remark 3.74. In the example of superadjunction $^{\circ}\mathsf{P}$ given in the extended summary (Subsection 1.2), this process was described in Subsection 1.2.6. We have that (i) as said above, $\mathrm{scl}(^{\circ}\mathsf{E})^{\top}$ is scalar-coherent, (ii) naturalities are interchange naturalities, (iii) the characterization was given in Lemma 1.2, and (iv) the classification was given in Lemma 1.3.

4 A basis for graded \mathfrak{gl}_2 -foams via rewriting theory

In this section, we apply linear Gray rewriting modulo as developed in Section 3 to a certain graded-2-category **GFoam**_d, the graded-2-category of \mathfrak{gl}_2 -foams, and show that it has the appropriate basis. The strategy follows the blueprint given in Subsection 3.6.2. Subsection 4.1 reviews the graded-2-category **GFoam**_d (Definition 4.8) and states the basis theorem (Theorem 4.10). Subsection 4.2 then defines the working data for the rewriting theory. The core of the proof is given in Subsection 4.3 and Subsection 4.4, respectively dealing with coherence of the modulo data and confluence of monomial local branchings. We conclude with an addendum (Subsection 4.5), showing that **GFoam**_d admits a variant **GFoam**'_d satisfying the same basis theorem.

4.1 Graded \mathfrak{gl}_2 -foams

Fix a positive integer $d \in \mathbb{N}$. We review the graded-2-category **GFoam**_d of \mathfrak{gl}_2 -foams. To fit our purpose, we describe it with a linear Gray polygraph GFoam_d. First, we describe the underlying 2-polygraph (GFoam_d)_{≤ 2}.

The objects of $\mathsf{GFoam}_d \operatorname{are}^{25}$

$$(\mathsf{GFoam}_d)_0 \coloneqq \bigsqcup_{k \in \mathbb{N}} \{\lambda \in \{1, 2\}^k \mid \lambda_1 + \ldots + \lambda_k = d\}.$$

²⁵In [SV23], (GFoam_d)₀ is denoted $\underline{\Lambda}_d$.

For each $\lambda \in (\mathsf{GFoam}_d)_0$ with k coordinates, we define a label on its coordinates

$$l_{\lambda} \colon \{1, \ldots, k\} \to \{1, \ldots, d\}$$

by setting $l_{\lambda}(i) = \sum_{j < i} \lambda_j + 1$. For instance, $l_{(1,1,2,1)} = (1, 2, 3, 5)$. In other words, the label $l_{\lambda}(i)$ is a sort of "weighted coordinate", where coordinate with value 2 counts double.

The 1-cells of GFoam_d are

$$(\mathsf{GFoam}_d)_1 \coloneqq \left\{ (\dots, \underset{i}{1}, \underset{i+1}{1}, \dots) \ \bigwedge_i \ (\dots, \underset{i}{2}, \dots) \ , \ (\dots, \underset{i}{2}, \dots) \ \downarrow \ (\dots, \underset{i}{1}, \underset{i+1}{1}, \dots) \right\}$$

Here the value l_{λ} is given below the corresponding coordinate. We read 1-morphism from right to left: for instance if d = 2, then the first 1-cell is a 1-morphism from (2) to (1, 1). The label *i*, called the *colour* of the strand, has value $1 \le i \le d - 1$.

The 2-cells of GFoam_d are



Each 2-cell in $(\mathsf{GFoam})_2$ is equipped with a \mathbb{Z}^2 -degree, denoted below its diagram.

We must explain "legal label". If a diagram $D \in (\mathsf{GFoam})_2$ has a region labelled by an element $\lambda \in (\mathsf{GFoam})_0$, then λ induces a label on each of the regions of D, following the rule given in $(\mathsf{GFoam})_1$.

Definition 4.1. Let $D \in (\mathsf{GFoam})_2$ and $\lambda \in (\mathsf{GFoam})_0$, where λ labels one of the regions of D. We say that λ is legal if:

- (i) the labels induced by λ on the other regions of D are labels in (GFoam)₀;
- (ii) if D is a dot with colour i, then $\lambda_i = 1$, where λ_i is the coordinate i of λ .

In the remaining of the section, we omit writing objects in diagrams.

Remark 4.2. The diagrams above is a string diagrammatics for certain singular surfaces called \mathfrak{gl}_2 -*foams*, or simply *foam*. A foam is made of *1*-*facets* and *2*-*facets*; two 1-facets and one 2-facet can join at a singular line called a *seam*. The string diagrammatics encodes \mathfrak{gl}_2 -foams via their

seams. Another diagrammatics, the *shading diagrammatics*, encodes \mathfrak{gl}_2 -foams via their 2-facets:



The legal condition for dots (Definition 4.1) translates as saying that dots cannot sit on 2-facets, or in the shading diagrammatics, that dots cannot sit on shaded regions.

Definition 4.3 (Terminology). A strand labelled by a colour *i* is called an *i*-strand. We similarly define *i*-dots. A region shaded with a colour *i* is called a *i*-shading. Two colours *i* and *j* are said to be distant if |i - j| > 1. Given a colour *i* and a diagram ψ , we say that *i* is distant from ψ if for each *j*-strand (resp. *j*-dot) in ψ , we have |i - j| > 1 (resp. $j \neq i, i + 1$).

The following gives the structural data that makes \mathbf{GFoam}_d a (G, μ) -graded-2-category:

Definition 4.4. Let \Bbbk be a commutative ring together with three invertible elements X, Y and Z in \Bbbk^{\times} such that $X^2 = Y^2 = 1$. Given this data, let μ be the following bilinear form for the abelian group $G := \mathbb{Z}^2$:

$$\mu \colon \mathbb{Z}^2 \times \mathbb{Z}^2 \to \mathbb{k}^{\times},$$
$$((a,b), (c,d)) \mapsto X^{ac} Y^{bd} Z^{ad-bc}.$$

We can now define $(GFoam)_3$, the set of generating relations. Foreseeing the rewriting, we decompose it as

$$(\mathsf{GFoam})_3 = \mathsf{E}_3 \sqcup \mathsf{R}_3,$$

and write E and R the associated linear 3-sesquipolygraphs with $(E)_{\leq 2} = (R)_{\leq 2} = (GFoam_d)_{\leq 2}$.

4.1.1 Foams isotopies

Recall the convention that we omit objects: all the 3-cells below can be labelled by any object, as long as the label is legal (Definition 4.1).

We define E as a scalar Gray polygraph, decomposing its 3-cells as

$$\mathsf{E}_3 = \mathsf{FGray}_3 \sqcup \mathsf{X}_3 \sqcup \mathsf{Z}_3.$$

Here FGray is the scalar 3-sesquipoly graph of (\mathbb{Z}^2, μ) -graded interchangers on $(\mathsf{GFoam}_d)_{\leq 2}$ (see Subsection 2.4.2). In other words:

$$\mathsf{FGray}_3 \coloneqq \left\{ \left| \begin{array}{c} \bullet_{\alpha} \\ \bullet_{\beta} \\ \bullet_{\beta} \\ \text{for all } \alpha, \beta \in (\mathsf{GFoam})_2^* \end{array} \right| \left| \begin{array}{c} \bullet_{\beta} \\ \bullet_{\beta} \\ \bullet_{\beta} \\ \text{for all } \alpha, \beta \in (\mathsf{GFoam})_2^* \end{array} \right| \right\}$$

Here we use the convention for scalar 3-sesquipolygraph already used in Subsection 2.1, writing the associated scalar on top of the arrow. The corresponding linear relation is the graded-interchange law (4).

The scalar 3-sesquipolygraph X has $(X)_{\leq 2} = (GFoam_d)_{\leq 2}$ and 3-cells



where we recall the terminology of Definition 4.3. The scalar associated to each 3-cell in X_3 is 1; hence we don't write anything above the corresponding arrow.

The scalar 3-sesquipolygraph Z has $(Z)_{\leq 2} = (\mathsf{GFoam}_d)_{\leq 2}$ and 3-cells

$$\mathsf{Z}_{3} \coloneqq \left\{ \bigcap_{i}^{i} \stackrel{i}{\Rightarrow} \stackrel{i}{\downarrow} \bigcap_{i}^{X} \stackrel{i}{\Rightarrow} \stackrel{i}{\uparrow} \bigcap_{i}^{Z^{2}} \stackrel{i}{\Rightarrow} \stackrel{i}{\uparrow} \stackrel{i}{\bigcap} \stackrel{YZ^{2}}{\stackrel{i}{\Rightarrow}} \stackrel{i}{\downarrow} \right\}$$

zigzag relations

Note that the graded-interchange law, the braid-like relations, the pitchfork relations and the zigzag relations encode a certain "braid-like pivotal" structure in \mathbf{GFoam}_d . This will be our modulo data (Subsection 4.3).

Topologically, these relations encode certain isotopies of singular surfaces called foam isotopies.

Definition 4.5 (Terminology). A 3-cell in E^{\top} is called a foam isotopy, or simply an isotopy; accordingly, we shall say that two diagrams are isotopic if they are the source and target of a 3-cell in E^{\top} .

The data of dots and strands in a given diagram is preserved under foam isotopies; that is, if $e: D_0 \rightarrow D_1$ is an isotopy, then there is a canonical bijection between the set of dots and strands of D_0 , with the set of dots and strands of D_1 . The following statement is shown in Subsection 4.3:

Proposition 4.6 (coherence of foam isotopies). If two parallel morphisms in E^{\top} define the same bijection on dots and strands, then they have the same associated scalar.

Definition 4.7 (Terminology). Following on Definition 4.5, we say that two dots in a diagram are isotopic if there exists an isotopic diagram in which the two dots are next to each other. Similarly, a dot and a strand, or two strands, can be isotopic.

4.1.2 The graded-2-category of \mathfrak{gl}_2 -foams

Finally, we define the linear 3-sesquipolygraph R as $(R)_{\leq 2} = (GFoam_d)_{\leq 2}$ and 3-cells given in Fig. 4.1.

Definition 4.8. The (G, μ) -graded-2-category \mathbf{GFoam}_d is the (G, μ) -graded-2-category presented by the linear Gray polygraph $\mathsf{GFoam}_d = \mathsf{E} \sqcup \mathsf{R}$.



Figure 4.1: 3-cells in R₃. Recall the convention that we omit objects: all the 3-cells can be labelled by any object, as long as the label is legal (Definition 4.1). In the last two cases, the wiggly lines are only visual aids, and are not part of the data. Each 3-cell has its notation, depicted above the arrow, so that $R_3 = \{dd, dm, bb_{\circlearrowright}, bb_{\circlearrowright}, nc, sq\}$.

4.1.3 The basis theorem

We review the notion of a reduced family ([SV23]). As it will not be essential for us, we only give a quick definition, and refer to [SV23, Section 2.5] for details.

Recall Remark 4.2. Given a foam $F: W \to W'$, write $\mathfrak{sl}(F)$ the surface obtained by removing its 2-facets. Similarly, given two parallel 1-morphisms W and W' in \mathbf{GFoam}_d , one can associate a certain union of circles S^1 denoted $\mathfrak{sl}(W \sqcup_{\partial} W')$. Write $\pi_0(\mathfrak{sl}(W \sqcup_{\partial} W'))$ the set of connected components in $\mathfrak{sl}(W \sqcup_{\partial} W')$.

A foam $F: W \to W'$ is *reduced* if $\mathfrak{sl}(F)$ is a union of disks, with at most one dot on each disk. Up to foam isotopies, a reduced foam is characterised by a subset $\delta \subset \pi_0(\partial \mathfrak{sl}(W \sqcup_{\partial} W'))$: a dot lies on a disk D if and only if ∂D belongs to δ . In that case, we say that F is δ -dotted.

Definition 4.9. Let W and W' be two parallel 1-morphisms in \mathbf{GFoam}_d . A reduced family is a family $(F_{\delta})_{\delta \subset \pi_0(\mathfrak{sl}(W \sqcup_{\partial} W'))}$, where $F_{\delta} \colon W \to W'$ is a δ -dotted reduced foam.

Theorem 4.10 (Basis theorem for graded \mathfrak{gl}_2 -foams). Let W and W' be two parallel 1-morphisms in **GFoam**_d. Any reduced family defines a basis of the \Bbbk -module $\operatorname{Hom}_{\mathbf{GFoam}_d}(W, W')$.

4.2 A convergent presentation of graded \mathfrak{gl}_2 -foams

This section sets up the "working data" for the rewriting theory of graded \mathfrak{gl}_2 -foams. This is the data (a), (b) and (c) as described in Subsection 3.6.1.

4.2.1 A linear Gray RSM for graded \mathfrak{gl}_2 -foams

Our rewriting approach to Theorem 4.10 is based on the following linear Gray RSM:

$$\mathsf{S} \coloneqq (\mathsf{R}, \mathsf{E}).$$

The following is a restatement of Definition 4.8:

Lemma 4.11. The linear Gray RSM S := (R, E) presents the graded-2-category \mathbf{GFoam}_d .

We further define B, a linear sub-3-sesquipolygraph of R, with $B_{<2} \coloneqq R_{<2}$ and

$$\mathsf{B}_3 \coloneqq \{\mathsf{dd}, \mathsf{dm}, \mathsf{bb}_{\circlearrowright}, \mathsf{bb}_{\circlearrowright}\}.$$

We write R := Cont(R), E := Cont(E) and B := Cont(B) the associated (family of) linear RSs, and S := (R, E) the associated (family of) linear RSM.

Note that:

Lemma 4.12. The following are S-rewriting steps:



4.2.2 A context-dependent linear sub-system

The linear Gray RSM S is not terminating, and hence not suited for a reduction algorithm. For instance, we have the following infinite sequence:

$$\int \prod_{i=1}^{nc} (i) = (i) + (i) + (i) = (i) + (i) = (i) + (i) + (i) = (i) + (i) = (i) + (i) = (i) + (i) = (i) + (i) + (i) = (i) + (i) = (i) + (i) + (i) = (i) + (i) + (i) = (i) + (i) + (i) + (i) + (i) = (i) + ($$

To avoid this obstruction to termination, we define a linear sub-system of S that prevents one from applying a neck-cutting relation (nc) on two pieces of the same strand. Taking into account the analogue obstruction for the squeezing relation (sq) leads to the following definition:

Definition 4.13. The (family of) linear RSM(s) T is the linear sub-system $T \subset S$ (Definition 3.56) such that for each $r \in R_3$, $\Gamma[r] \in T$ if and only if either $r \in B$, or r = nc (resp. sq) of colour i (resp. (i, i + 1)), and the two pieces of *i*-strands belong to distinct strands in $\Gamma[r]$.

It is not hard to check that T is adapted, and hence is indeed a linear sub-system in the sense of Definition 3.56; in fact, S is already adapted. Note that T is *not* of form Cont(T) for some sub-Gray RSM $T \subset S$.

The following lemma justifies why T is a suitable candidate to prove Theorem 4.10:

Lemma 4.14. A foam is a T^+ -normal form is and only if it is reduced. In particular, any choice of foam isotopy representatives for BNF_{T^+} is a reduced family.

The proof of Lemma 4.14 is given in [Sch24, Proposition 1.6.5]; in order to keep the focus on the rewriting theory, we omit it here.

Given a foam F, we associate to F the following data (recall Definition 4.3):

 $#sh_i(F) :=$ number of *i*-shadings in the shading diagrammatics of *F*, $#cl_i(F) :=$ number of closed *i*-strands in the string diagrammatics of *F*, $#d_i(F) :=$ number of *i*-dots in *F*.

Each of these data defines a partial order on foams. Let \succ be the lexicographic order induced by these partial orders:

 $\succ \coloneqq (\#\mathrm{sh}_1, \ldots, \#\mathrm{sh}_{d-1}, \#\mathrm{cl}_1, \ldots, \#\mathrm{cl}_{d-1}, \#\mathrm{d}_1, \ldots, \#\mathrm{d}_d).$

By definition, \succ is terminating.

Proposition 4.15. The preorder \succ is strongly compatible (Definition 3.58) with T.

4.2.3 A basis for graded \mathfrak{gl}_2 -foams

The following is a corollary of the coherence of foam isotopies (Proposition 4.6):

Corollary 4.16. E^{\top} is scalar-coherent on reduced foams.

Proof. One can check that a reduced foam, expressed in string diagrammatics, does not have any closed strands (see [Sch24, Corollary 1.6.8]), nor isotopic dots with the same colour. Hence, foam isotopies on reduced foams all induce the same bijection on dots and strands. The statement then follows from Proposition 4.6.

We already know that \succ is terminating and strongly compatible with T⁺ (Proposition 4.15) and that E^T is scalar-coherent on monomial T-normal forms (Lemma 4.14 and Corollary 4.16). In order to show Theorem 4.10 using the BASIS-FROM-CONVERGENCE THEOREM 3.36, it remains to show the coherence of foam isotopies (Proposition 4.6)—this is done in Subsection 4.1.1—and to check that, on one hand:

Lemma 4.17. The (family of) linear RSM(s) T and S present the same underlying (family of) module(s).

and on the other hand, thanks to the TAMED LINEAR NEWMANN'S LEMMA 3.43:

Proposition 4.18. *Every monomial local* T^+ *-branching is* \succ *-tamely* T^{st} *-congruent.*

Both statements are shown in Subsection 4.4.

4.3 Coherence of foam isotopies

Recall the terminology introduced in Definitions 4.3, 4.5 and 4.7 and the scalar 3-sesquipolygraphs defined in Subsection 4.1.1. Let

 $\overline{\mathsf{E}} = \mathsf{FGray} \sqcup \mathsf{X}.$

In this subsection, we prove the coherence of foam isotopies (Proposition 4.6) using the higher Newmann's lemma (Lemma 3.52) for the Gray RSM (Z, \overline{E}) . In that regard, the strategy of proof is close to the one described for superadjunction in Subsection 1.2.

We start by providing the analogous coherence result for the modulo data:

Lemma 4.19 (coherence of foam isotopies, except zigzag relations). If two parallel morphisms in \overline{E}^{\top} define the same bijection on dots and strands, then they have the same associated scalar.

Proof. Since μ is symmetric, the scalar of a relation in FGray^{\top} only depends on how generating 2-cells vertically permute (see [SV23, Section 2.1], and also the coherence of interchangers in Gray categories [Sch24, Appendix A]). In fact, it only depends on how generating 2-cells with a non-trivial grading vertically permute. 3-cells in X^{\top} have trivial associated scalar. While they do not preserve the data of the set of generating 2-cells, they do preserve the data of the set of non-trivially graded generating 2-cells. That is, if $e: D_0 \rightarrow D_1$ is a 3-cell in X, there is a canonical way to identify the non-trivially graded generating 2-cells of D_0 with those of D_1 . In other words, the permutation of non-trivially graded generating 2-cells is a well-defined data associated to any 3-cell in $\overline{\mathsf{E}}^{\top}$, and this data determines the associated scalar. Finally, we check that this permutation data is itself determined by the bijection on dots and strands.

The modulo \overline{E}^{\top} can be understood as encompassing a braided-like structure. Indeed, \overline{E}^{\top} is equivalently generated by the braided-like relations and the following 3-morphisms:



which capture the naturality of the braiding with respect to the cap, the cup and the dot. We say "braided-like" to emphasize the restriction on the labels.

Recall that any Gray RSM has canonical interchange naturalities (Subsection 3.4.6). For (Z, \overline{E}) , we further have the following braided-like naturalities:

Lemma 4.20 (braided-like \overline{E} -naturalities). For any relation $f: \psi_0 \to \psi_1$ in Z and i any colour distant from ψ_0 (see Definition 4.3), we have the following \overline{E} -congruences:



Our next step is to upgrade our knowledge of \overline{E} -naturalities to a general characterization of \overline{E} -congruence classes, leveraging our understanding of coherence in \overline{E} :

Lemma 4.21 (characterization of \overline{E} -congruence classes for Z). If (f, g) is a local Z-branching such that f and g are of the same type with identical cup and cap, then (f, g) is \overline{E} -scalar-congruent.

Proof. Let [f, e, g] be a Z-local triple with f and g as in the statement of the lemma. Thanks to coherence of the modulo (Lemma 4.19), we can choose the isotopy $e: s(f) \to s(g)$ such that each step either does not overlap s(f), or consists of a braided-like \overline{E} -naturality (Lemma 4.20). This concludes.

With this characterization at hand, we can classify local Z-branchings and reduce scalarconfluence to four explicit branchings, the critical branchings:

Lemma 4.22. Every local Z-branching is, up to branchwise \overline{E} -congruence, either an independent branching or a contextualization of one the following scalar-confluent branchings:



Proof. Let $[f, e, g] = [\Gamma_f[r_f], e, \Gamma_f[r_g]]$ be a Z-local triple, with $r_f, r_g \in Z_3$. We wish to simplify [f, e, g] by replacing it with a branchwise \overline{E} -congruent branching using the characterization given in Lemma 4.21. Given a cup \cup and a cap \cap belonging to the same strand S, we refer to a "strand between the \cup and \cap " to mean a strand crossing S on the piece of strand connecting \cup to \cap .

Consider the cup and cap of $s(r_g)$, respectively denoted \cup_g and \cap_g , as sitting in s(f). They necessarily belong to the same strand S, possibly with strands between \cup_g and \cap_g . Using isotopies in $\overline{\mathsf{E}}$, we can slide these strands away, so that no strand lies between \cup_g and \cap_g . Moreover, we can do so without adding strands between \cup_f and \cap_f , the cup and cap of $s(r_f)$ (viewed as sitting in s(f)). Thanks to the characterization given in Lemma 4.21, this does not change the branchwise $\overline{\mathsf{E}}$ -congruence class of [f, e, g].

If the cups and caps of $s(r_f)$ and $s(r_g)$ are disjoint, we can further use interchanges to move the cup and cap of $s(r_g)$ one below the other, again without affecting $s(r_f)$. This shows that [f, e, g] is branchwise $\overline{\mathsf{E}}$ -congruent to an independent Z-branching. If $s(r_f)$ and $s(r_g)$ have precisely one cap or one cup in common, we find the four critical branchings given in the lemma.

Moreover:

Lemma 4.23. Z terminates modulo \overline{E} .

Proof. Z strictly diminishes the number of caps and cups, which is kept constant by \overline{E} .

We now have all the ingredients to show Proposition 4.6:

Proof of Proposition 4.6. Using the higher Newmann's lemma (Lemma 3.52), Lemma 4.22 and termination (Lemma 4.23), we conclude that Z is convergent \overline{E} . Hence, Z^{\top} is scalar-coherent modulo \overline{E}^{\top} (coherence modulo from convergence modulo; Proposition 3.8).

To show the proposition, it suffices to show that if a loop e in E^{\top} defines the identity bijection of dots and strands, then its associated scalar is one. Since Z^{\top} is scalar-coherent modulo \overline{E}^{\top} , we can write e as $e = g \circ h \circ g^{-1}$ with h in \overline{E}^{\top} . We conclude using coherence of \overline{E}^{\top} (Lemma 4.19). \Box

4.4 Analysis of monomial local branchings

We denote $A := T \setminus B$; that is, A consists of the rewriting step of type nc and sq which are in T.

We study of monomial local T⁺-branchings. It is roughly divided in three parts: both branches are of type B (Proposition 4.26), one branch is of type B and the other is of type A (Lemma 4.35), and both branches are of type A (Lemma 4.36). The general strategy (with some variations) is the one described in Subsection 3.6.2: give naturalities of the modulo, characterize rewriting steps modulo, and finally enumerate monomial local branchings.

Throughout we use the terminology of Definitions 4.3, 4.5 and 4.7.

4.4.1 Foam isotopy naturalities

Lemma 4.24 (braided-like E-naturalities). The relations in $R_3 \sqcup \{\overline{dm}, \overline{nc}, \overline{sq}\}$ satisfy the same braided-like naturalities as pictured in Lemma 4.20.

Proof. This follows from the fact that if i is distant from ψ_0 , then i is distant from ψ_1 (using the same notations as in Lemma 4.20). This is straightforward in most cases; we only detail type dm. Let j be the colour of the migrating dot in ψ_0 . By assumption, the associated j-strand crosses an i-strand, so that we have |i - j| > 1. In particular $j + 1 \neq i, i + 1$ and the (j + 1)-dot can still slide through the i-strand once it has migrated.

For $f: \psi_0 \to \psi_1$ a 3-cell in $\mathsf{R}_3 \sqcup \{\overline{\mathsf{dm}}, \overline{\mathsf{nc}}, \overline{\mathsf{sq}}\}\)$, we write $f: 0 \not \to {}^{\mathsf{I}} \not \oplus$ the 3-cell obtained by rotating each diagram by a half-turn. Most 3-cells rotate to themselves, except for dm (resp. $\overline{\mathsf{dm}}\)$, which rotate to $\overline{\mathsf{dm}}\)$ (resp. $\overline{\mathsf{dm}}\)$).

Lemma 4.25 (pivotal E-naturalities). Let $f: \psi_0 \to \psi_1$ be a 3-cell in $\mathsf{R}_3 \sqcup \{\overline{\mathsf{dm}}, \overline{\mathsf{nc}}, \overline{\mathsf{sq}}\}$. Then the following squares commute:



Proof. The statement is trivial for types dd, bb_{\odot} and bb_{\odot} , and follows from graded interchange for types dm. For types nc and sq, it comes down to the following computations:



This also implies the lemma for the overlined types dm, \overline{nc} and \overline{sq} , using coherence of E.

4.4.2 Confluence of monomial local B⁺-branchings

We begin our study of confluence with positive branchings in the context-agnostic linear subsystem $B \subset T$, which derives from a linear Gray RSM: B = Cont(B). In fact, we show that in that case, confluences can also be taken in B:

Proposition 4.26. *Every monomial local* B^+ *-branching is* \succ *-tamely* B^{st} *-congruent.*

Note that since \succ is strongly compatible with T (Proposition 4.15), it is also strongly compatible with B. As a preliminary step, we give a topological characterization of E-congruence classes:

Lemma 4.27 (characterization of E-congruence classes for B). Let (f, g) be a monomial local B⁺branching, with f and g of the same type. The following holds:

- (a) type dd: (f, g) is E-congruent;
- (b) type dm: If f and g have isotopic i-dot and i-strand, then (f, g) is E-congruent;
- (c) type bb_{\bigcirc} and bb_{\bigcirc} : If f and g have isotopic *i*-strand, then (f, g) is E-congruent.

We call the data associated to each type its combinatorial data, and say that this combinatorial data characterizes the E-congruence class of the type.

Proof. Denote ψ the local picture of the rewriting step f; that is, $\psi = s(r)$ for $r \in B$ and $f = \Gamma[r]$ for some context Γ . In each type, we use coherence of E (Proposition 4.6) to present the isotopy e as a composition of E-naturalities, as described in Lemma 4.24 and Lemma 4.25:

- (a) Trivial, since both f and g rewrites to zero.
- (b) Through the isotopy e, the i-dot starts and ends next to the i-strand. Hence, we can choose e such that the i-dot always remains next to the i-strands. In that case, the only isotopies that overlap ψ are braided-like naturalities.
- (c) Given that the bubble ψ starts and ends without any strand crossing it, we can choose the isotopy e such D always crosses strands "at once". In that case, the only isotopies that overlap ψ are braided-like isotopies.

As B is context-agnostic, the case of independent B⁺-branchings comes for free (Lemma 3.65):

Lemma 4.28. *Each independent* B^+ *-branchings is* \succ *-tamely* B^{st} *-congruent.*

We can now prove the proposition:

Proof of Proposition 4.26. We use the BRANCHWISE TAMED CONGRUENCE LEMMA 3.19 and the fact that independent B^+ -branchings are \succ -tamely B^{st} -congruent (Lemma 4.28) without further mention.

Let [f, e, g] be a monomial B-local triple. If f and g are of type dd, bb_{\circlearrowright} or bb_{\circlearrowright} (not necessarily both of the same type), then either they have the same combinatorial data and hence are E-congruent, or their combinatorial data are disjoint. We can isotope them away from each other, and [f, e, g] is branchwise E-congruent to an independent branching. A similar argument applies if f is of type dm and g is of type bb_{\circlearrowright} .

Assume f is of type dm and g is of type dd. If their combinatorial data are disjoint, [f, e, g] is branchwise E-congruent to an independent branching. Otherwise, they share the data of an *i*-dot.

In that case, [f, e, g] is E-congruent to a contextualization of the following local B⁺-branching, shown to be B⁺-confluent:



It follows from the CONTEXTUALIZATION LEMMA 3.62 that [f, e, g] is \succ -tamely B-confluent.

Assume finally that f is of type dm and g is of type bb_{\circlearrowright} . A similar reasoning reduces the statement to the following B⁺-confluence:



This concludes.

Finally, it follows from the TAMED LINEAR NEWMANN'S LEMMA 3.43 that:

Corollary 4.29. B^+ is convergent.

A *bubble* is an endomorphism of an identity 1-morphism, possibly viewed inside a bigger diagram. One can check (see [Sch24, Lemma 1.6.7]) that any bubble can be "evaluated" using B^+ -rewriting steps, in the sense that it rewrites into a sum of diagrams, each consisting only of dots. Proposition 4.26 shows that this evaluation is uniquely defined up to E-congruence, so that we can speak of *the* B^+ -rewriting sequence evaluating a bubble:

Definition 4.30. For each bubble ϕ , its bubble evaluation is the B⁺-rewriting sequence, well-defined up to E-congruence, which rewrites ϕ into a sum of dots:

$$bb^*: \quad \phi \xrightarrow{*}_{\mathsf{B}} \sum dots.$$

A B-rewriting sequence defined as a contextualized bb* is said to be of type bb*.

4.4.3 Characterizing neck-cutting and squeezing relation up to B-confluence

In order to study rewriting steps of type nc and sq, we would like to give a topological characterization of their E-congruence classes akin to the one given for B in Lemma 4.27. In fact, it will be easier to characterize their branchwise B^+ -confluence classes.

Proposition 4.31 (characterization of branchwise B-confluence classes for nc and sq). Let (f, g) be a monomial local S⁺-branching with f and g of type nc and label i (resp. of type sq and label (i, i + 1)). If f and g apply to the same i-strand(s), then (f, g) is B⁺-confluent.

We call the data of the *i*-strand(s) the *combinatorial data* of type nc (resp. sq), and say that it *characterizes* their B^+ -confluence class. Before proving Proposition 4.31, we show the following elementary B^+ -confluences:
Lemma 4.32 (spatial-like B⁺-confluence). The following branchings are B⁺-confluent, where ϕ is an arbitrary bubble and the dotted wiggly line denotes either a neck-cutting or a squeezing relation:



Proof. We shall see that in each case, it suffices to evaluate the bubble ϕ (see Definition 4.30) and apply some additional dot migrations to achieve B⁺-confluence. Denote

$$\mathsf{bb}^* \colon \quad \textcircled{\phi} \xrightarrow{*}_\mathsf{B} \sum_{\delta} \ \textcircled{\delta}$$

the bubble evaluation of ϕ . Denote by δ a generic union of dots appearing in this bubble evaluation. Consider the first branching. We compare



If δ only consists of *j*-dots with $j \neq i, i+1$, then δ can slide from top to bottom, without additional scalar. If δ contains at least two *j*-dots with j = i, i+1, then both sides rewrite to zero, possibly migrating an *i*-dot into a (i+1)-dot first. Finally, If δ contains exactly one *j*-dot with j = i, i+1, then both sides rewrites to a single diagram consisting of a (i+1)-dot on top and a (i+1)-dot on the bottom (again using dot migrations).

The other branchings are treated similarly. For the second and fourth branchings, one can use dot migrations to rewrite *i*- and (i+1)-dots into (i+2)-dots, allowing them to cross the *i*-strands. For the third branching, the fact that a dot cannot sit on a shaded region (for certain labels, see Remark 4.2) prevents (i + 1)- and (i + 2)-dots, and *i*-dots can be treated as before, sliding then first across the (i + 1)-strand.

We can now prove the proposition:

Proof of Proposition 4.31. Let [f, e, g] a local triple as in the proposition. Since B⁺ is convergent (Corollary 4.29), it suffices to show that [f, e, g] is B-congruent. As in Lemma 4.27, denote ψ the local picture of the rewriting step f. That is, $\psi = s(nc)$ is two vertical pieces of *i*-strands (resp. $\psi = s(sq)$ is four vertical pieces of strands), with $f = \Gamma[nc]$ (resp. $f = \Gamma[sq]$) for some context Γ . The main idea is to treat ψ as an extra formal generator, performing only isotopies independent of ψ , or E-naturalities and B-confluences as described in Lemma 4.24 (braided-like E-naturalities), Lemma 4.25 (pivotal E-naturalities), and Lemma 4.32 (spatial-like B⁺-confluences). This does not change whether [f, e, g] is B-congruent, thanks to the BRANCHWISE TAMED CONGRUENCE LEMMA 3.19. We describe the procedure in more details for type nc; the type sq is analogous. An example is given below, picturing only the two *i*-strands and some bubble ϕ :



First, we move the two *i*-strands in s(f) closer to one another, so that they remain parallel throughout, except possibly close to their endpoints. This procedure may require evaluating bubbles (Definition 4.30). Doing so does not affect what we need to demonstrate, thanks to the INDEPENDENT REWRITING LEMMA 3.67.

Then, ψ can be slid along the two parallel *i*-strands using pivotal E-naturalities (Lemma 4.25). Doing so, it may cross distant strands or bubbles: both go through thanks to braided-like E-naturalities (Lemma 4.24) and spatial-like B⁺-confluences (Lemma 4.32). Applying the same procedure to s(g) eventually leads to the same diagram, up to isotopies independent of ψ .

Lemma 4.33. Every monomial $(S \setminus T)^+$ -rewriting step is E-congruent to a B-congruence.

Proof. By Proposition 4.31, a monomial $(S \setminus T)^+$ -rewriting step f of type nc is branchwise B⁺-confluent to a rewriting step which, up to contextualization, has the following form:



which is readily seen to be B⁺-confluent. The CONTEXTUALIZATION LEMMA 3.62 together with the BRANCHWISE TAMED CONGRUENCE LEMMA 3.19 implies that f is \succ -tamely Bst-congruent. A similar argument holds for $(S \setminus T)^+$ -rewriting step of type sq, reducing the argument to the following rewriting step, readily seen to be B⁺-confluent (see e.g. [SV23, Lemma 2.24]):



This concludes.

As a direct corollary, we have that T and S present the same underlying module (Lemma 4.17). Moreover:

 \square

Lemma 4.34. Every independent local T^+ -branching is \succ -tamely T^{st} -congruent.

Proof. This follows from Lemma 3.66.

4.4.4 Confluence of monomial local (A⁺, B⁺)-branchings

Recall the notation $A = T \setminus B$. We use the terminology (A^+, B^+) -branching to refer to a branching which has one branch in A^+ and the other in B^+ .

Lemma 4.35. Every monomial local (A^+, B^+) -branching is \succ -tamely T^{st} -congruent.

Proof. Throughout we use that independent local T^+ -branchings are \succ -tamely T^{st} -congruent (Lemma 4.34) without further mention. We also stop explicitly mentioning the BRANCHWISE TAMED CONGRUENCE LEMMA 3.19.

Let [f, e, g] be a monomial (A^+, B^+) -local triple. As a result of the characterization of B (Lemma 4.27) and the characterization of $\{nc, sq\}$ (Proposition 4.31), we can freely choose the combinatorial representatives of f and g. For instance, if f and g have distinct combinatorial data then we can choose the combinatorial representatives of f and g so that [f, e, g] is an independent branching and hence \succ -tamely Tst-congruent. In particular, if g is of type dd then [f, e, g] is automatically \succ -tamely Tst-congruent. A similar reasoning applies if g is of type dm, choosing to apply the dot migration away from where the neck-cutting or the squeezing relation happens.

Assume that f is of type nc and g is of type bb₀, such that the strand in its combinatorial data (call it s_1) is one of the two *i*-strands in the combinatorial data of f (call them s_1 and s_2). The fact that s_1 is closed forces $s := s_1 = s_2$, so that f is in fact not a T⁺-rewriting step to start with. If instead g is of type bb₀ sharing part of its combinatorial data with f, then [f, e, g] is branchwise \succ -tamely Tst-congruent to a contextualization of the T⁺-confluent branching pictured in Fig. 4.2. Since the given T⁺-confluence is in B, the CONTEXTUALIZATION LEMMA 3.62 applies.



Figure 4.2: Critical branching between types B and type nc.

Similar arguments can be given if f is of type sq and g is of type bb₍₎ (resp. bb₍₎), showing that [f, e, g] is branchwise \succ -tamely Tst-congruent to a contextualization of the T⁺-confluent branching pictured in Fig. 4.3. In that case however, the T⁺-confluences are not in B, and more care is needed to apply the CONTEXTUALIZATION LEMMA 3.62; in fact, it *does not* apply to the first of the two critical branchings in Fig. 4.3. One can give yet another critical branching to deal with this case. Instead, we describe another argument that does not require further computation and works for all critical branchings. This argument is essentially the same argument as used in Lemma 3.66.

Note that each confluence in Fig. 4.2 and Fig. 4.3 contains at most one A^+ -rewriting step; called it r. Assume that for some context Γ , one of these confluences fails to verify the hypothesis of the CONTEXTUALIZATION LEMMA 3.62. That means that $\Gamma[r]$ is not in A, and hence by Lemma 4.33, [f, e, g] is B-congruent. We conclude that [f, e, g] is B⁺-confluent, thanks to the convergence of B (Corollary 4.29).

4.4.5 Confluence of monomial local A⁺-branchings

Lemma 4.36. Every monomial local A^+ -branching is \succ -tamely T^{st} -congruent.

Proof. In what follows, we use that independent local T^+ -branchings are \succ -tamely T^{st} -congruent (Lemma 4.34), the BRANCHWISE TAMED CONGRUENCE LEMMA 3.19 and the CONTEXTUALIZATION



Figure 4.3: Critical branchings between types B and type sq.

LEMMA 3.62 without explicit mentions.

Let [f, e, g] be a monomial A⁺-local triple. As in the proof of Lemma 4.35, the characterization of $\{nc, sq\}$ Proposition 4.31 implies that we can freely choose the combinatorial representatives of f and g.

Contrary to the proof of Lemma 4.35 however, even if f and g have distinct combinatorial data, there may not exist combinatorial representatives for which f and g are independent. On the other hand, we shall see that in most cases, we can choose the combinatorial representatives of f and g such that [f, e, g] rewrites into an independent branching, in the sense of Subsection 3.5.5. Only two cases will not follow this scheme: they are the critical branchings given in Fig. 4.4.

Consider first the case where both f and g are of type nc. If their respective colour are i and j, we can assume that either j = i or j = i + 1 (otherwise, we can choose combinatorial representatives such that [f, e, g] is an independent branching). In these cases, we can choose combinatorial representatives such that $[f, e, g] = \Gamma[f', e', g']$ for some context Γ and [f', e', g'] is encoded in the diagram on the left-hand side below (the two wiggly lines encode f and g):



If j = i, then $\Gamma[f', e', g']$ rewrites via two neck-cuttings into a linear combination of branchings $\Gamma[f'', e'', g'']$ with f'' and g'' associated to the same combinatorial data, as pictured in the schematic on the right-hand side above (we only picture the *i*-strands, leaving the dots implicit). If Γ does not connect any of the *i*-strands involved, then all monomial rewriting steps involved are in T, and we can apply the INDEPENDENT REWRITING LEMMA 3.67. Otherwise, it means that only one, or none, neck-cutting is necessary to get to the same situation, and the INDEPENDENT REWRITING LEMMA 3.67 is still applicable. In any case, [f, e, g] is \succ -tamely Tst-congruent.

If j = i + 1, the T⁺-confluence of [f', e', g'] is the first critical branching pictured in Fig. 4.4. As this confluence only uses one rewriting step of type A, we conclude as in Lemma 4.35 that $[f, e, g] = \Gamma[f', e', g']$ is \succ -tamely Tst-congruent.

Consider now the case where f is of type sq and g is of type nc. Denote (i, i + 1) and j their respective colour. As before, we can assume that j = i - 1, i, i + 1 or i + 2. Then, up to choice of combinatorial representatives and vertical symmetry, [f, e, g] is equal to $\Gamma[f', e', g']$ for some context Γ and [f', e', g'] as pictured below:



If j = i + 2, we can further isotope the branching to get an independent branching. If j = i or if j = i + 1, we can rewrite it using neck-cuttings and conclude similarly as above.

The last case j = i - 1 leads to the second critical branching pictured in Fig. 4.4. Note that flipping everything vertically leads to the same critical branching because j and i + 1 are distant colours. The confluence uses two rewriting steps of type A, one of type sq and the other of type nc; we refer to them simply as "sq" and "nc". If Γ ["sq"] does not belong to T, then f does not belong to

T either; hence if $[f, e, g] = \Gamma[f', e', g']$ is a T⁺-branching, $\Gamma[$ "sq"] necessarily belong to T. On the other hand, if $\Gamma[$ "nc"] does not belong to T, we can replace it with a B⁺-confluence. We conclude that [f, e, g] is \succ -tamely Tst-congruent.

Finally, consider the case where both f and g are of type sq, with respective colours (i, i + 1) and (j, j + 1). Without loss of generality we can assume that $j \ge i$, and furthermore that either j = i, j = i+1 or j = i+2. Each case will allow different choices of combinatorial representatives. To help the exposition, we fix the positions of the i and (i + 1)-strands associated to f, simply called the i- and (i + 1)-strands below, and discuss how the j- and (j + 1)-strands associated to g, simply called the j- and (j + 1)-strands below, can be isotoped with regard to the i- and (i + 1)-strands.

If j = i + 2, then j is adjacent to i + 1. In particular, the two j-strands cannot be isotoped through the (i + 1)-strands. As there exists an isotopy joining the j and (j + 1)-strands, the four strands must lie on one side of the (i + 1)-strands (at least partially for the (j + 1)-strands), as pictured in the following schematic (replacing wiggly lines with straight lines to avoid clutter):



On the side where the *j*-strands are, there is one *i*-strand. Given that *i* is distant from *j* and *j*+1, it can be isotoped through the *j* and (j+1)-strands. In that way, [f, e, g] is branchwise E-congruent to an independent branching.

If j = i + 1, then the (i + 1)-strands and (j + 1)-strands cannot intersect. This leads to three possible schematics, up to symmetries:



In the first schematic, one of the (i + 1)-strands coincides with one of the *j*-strands. The first two schematic are independent branchings on the nose. The last one rewrites into an independent branching, as pictured.

Finally, if j = i, either have the (i + 1)-strands and (j + 1)-strands do not intersect, or they coincide. Below we only picture the two schematics for which rewriting the branching is necessary:



This concludes.





Figure 4.4: Critical branchings in types {nc, sq}

4.5 Addendum: another deformation of \mathfrak{gl}_2 -foams

L

In this addendum, we use our rewriting approach to classify "deformations" of \mathfrak{gl}_2 -foams, leading to a variant **GFoam**_d of **GFoam**_d satisfying the same basis theorem (Theorem 4.10).

The structure of graded-2-category on \mathbf{GFoam}_d is already somehow universal. Indeed, the abelian group \mathbb{Z}^2 is isomorphic to the abelian group presented by generators $D, \cup^L, \cup^R, \cap^L, \cap^R$ and relations

$$\cap^{R} + \cup^{R} = 0$$
$$\cap^{L} + \cup^{L} = 0$$
$$\cup^{R} + \cap^{L} + D = 0$$
$$\cup^{L} + \cap^{R} = D$$
$$\cup^{L} + \cup^{R} + \cap^{L} + \cap^{R} = 0$$

obtained from taking the "abelianization" of the defining local relations of \mathbf{GFoam}_d . Hence, if we assume that the grading is independent on the colours of the generators, then the \mathbb{Z}^2 -grading on \mathbf{GFoam}_d is the most general one. Moreover, a symmetric bilinear map $\mu \colon \mathbb{Z}^2 \times \mathbb{Z}^2 \to \mathbb{k}^{\times}$ is determined by its values on the generators of $\mathbb{Z}^2 \times \mathbb{Z}^2$, with relations:

$$\mu((1,0),(1,0))^2 = \mu((0,1),(0,1))^2 = 1$$
 and $\mu((1,0),(0,1))\mu((0,1),(1,0)) = 1$.

This gives the parameters X, Y and Z.

Let us now look how the defining relations could be deformed. For simplicity, we assume that the braid-like relations, pitchfork relations and dot slide (i.e. relations captured by X) remain scalar-free. Up to normalization, we can further assume that the scalars of the zigzag relations, the dot migration (dm) and the evaluation of counter-clockwise bubble (bb_{\odot}) keep the same scalars.

Going over all the critical branchings, one finds exactly one extra possibility where nc remains as it is, and we have:

This defines an a priori distinct (\mathbb{Z}^2, μ) -graded-2-category **GFoam**'_d. Renormalizing the right-ward cap, we can also define **GFoam**'_d by only modifying the zigzag relations as follows:

$$\int \bigcup_{i} = \bigcup_{i} = X \bigwedge_{i} = X YZ^{2} \bigwedge_{i} = XZ^{2} \bigvee_{i} = XZ^{$$

This other graded deformation ([Sch24, Definition 6.6.9]) verifies the same basis theorem (Theorem 4.10), by exactly the same rewriting proof. In particular, it verifies the same categorification theorem [SV23, Theorem 2.29].

Remark 4.37. As noted in [SV23, Remark 3.24], GFoam_d defines "type X" odd Khovanov homology, while GFoam'_d defines "type Y" odd Khovanov homology, although these two variants are isomorphic. The existence of these variants is explained by so-called "ladybug squares", certain squares in the hypercube that compose to zero. Interestingly, the same ladybug squares are at the heart of *Khovanov homotopy type* [LS14a], a stable homotopy refinement of Khovanov homology. Most of the refinement is canonical, except on ladybug squares, for which a choice has to be made. This failure of canonicity is what allows Khovanov homotopy type to be a strictly stronger invariant than Khovanov homology [LS14b; See12].

References

- [All18a] C. Alleaume, Rewriting in Higher Dimensional Linear Categories and Application to the Affine Oriented Brauer Category, *J. Pure Appl. Algebra*, vol. 222, no. 3 (2018), pp. 636–673 (cit. on pp. 3, 9, 10, 17, 19, 29, 36, 41, 42, 54, 56, 61).
- [All18b] C. Alleaume, Higher-dimensional linear rewriting and coherence in categorification and representation theory, Ph.D. thesis, Université de Lyon, 2018 (cit. on pp. 41, 54).
- [Ara+23] D. Ara, A. Burroni, Y. Guiraud, P. Malbos, F. Métayer, and S. Mimram, Polygraphs: From Rewriting to Higher Categories, preprint 2023, arXiv: 2312.00429 [cs, math] (cit. on pp. 8, 33, 37).
- [Ara22] M. Araújo, Simple String Diagrams and N-Sesquicategories, Theory Appl. Categ., vol. 38 (2022), Paper No. 34, 1284–1325 (cit. on pp. 11, 19).
- [Bar24] S. Barbier, Diagram Categories of Brauer Type, preprint 2024, arXiv: 2406.18436 [math] (cit. on p. 19).
- [BE17a] J. Brundan and A. P. Ellis, Monoidal Supercategories, *Comm. Math. Phys.*, vol. 351, no. 3 (2017), pp. 1045–1089 (cit. on p. 4).
- [BE17b] J. Brundan and A. P. Ellis, Super Kac–Moody 2-categories, *Proc. London Math. Soc.*, vol. 115, no. 5 (2017), pp. 925–973 (cit. on p. 2).
- [Ber78] G. M. Bergman, The Diamond Lemma for Ring Theory, Adv. in Math., vol. 29, no. 2 (1978), pp. 178–218 (cit. on p. 3).
- [BK22] J. Brundan and A. Kleshchev, Odd Grassmannian Bimodules and Derived Equivalences for Spin Symmetric Groups, preprint 2022, arXiv: 2203.14149 [math] (cit. on p. 3).
- [BKV18] K. Bar, A. Kissinger, and J. Vicary, Globular: An Online Proof Assistant for Higher-Dimensional Rewriting, Log. Methods Comput. Sci., vol. 14, no. 1 (2018), Paper No. 8, 16 (cit. on pp. 11, 18).
- [BN98] F. Baader and T. Nipkow, Term Rewriting and All That, Cambridge University Press, Cambridge, 1998, xii+301 (cit. on p. 46).
- [Bok76] L. A. Bokut', Imbeddings into Simple Associative Algebras, Algebra i Logika, vol. 15, no. 2 (1976), pp. 117–142, 245 (cit. on p. 3).
- [Buc06] B. Buchberger, An Algorithm for Finding the Basis Elements of the Residue Class Ring of a Zero Dimensional Polynomial Ideal, J. Symbolic Comput., vol. 41, no. 3-4 (2006), pp. 475–511 (cit. on p. 3).
- [Bur93] A. Burroni, Higher-Dimensional Word Problems with Applications to Equational Logic, in: *Theoretical Computer Science*, vol. 115, 1, 1993, pp. 43–62 (cit. on p. 8).
- [BV17] K. Bar and J. Vicary, Data Structures for Quasistrict Higher Categories, in: 2017 32nd Annual ACM/IEEE Symposium on Logic in Computer Science (LICS), IEEE, [Piscataway], NJ, 2017, p. 12 (cit. on pp. 11, 18).
- [CDM22] C. Chenavier, B. Dupont, and P. Malbos, Confluence of Algebraic Rewriting Systems, Math. Structures Comput. Sci., vol. 32, no. 7 (2022), pp. 870–897 (cit. on pp. 10, 42).
- [CHW13] S. Clark, D. Hill, and W. Wang, Quantum Supergroups I. Foundations, Transformation Groups, vol. 18, no. 4 (2013), pp. 1019–1053 (cit. on p. 2).

[CKM14] S. Cautis, J. Kamnitzer, and S. Morrison, Webs and Quantum Skew Howe Duality, Math. Ann., vol. 360, no. 1-2 (2014), pp. 351-390 (cit. on p. 2). [DEL21] B. Dupont, M. Ebert, and A. D. Lauda, Super Rewriting Theory and Nondegeneracy of Odd Categorified Sl(2), 2021, arXiv: 2102.00276 [math] (cit. on pp. 3, 10, 17, 43, 49, 54). [DM22] B. Dupont and P. Malbos, Coherent Confluence modulo Relations and Double Groupoids, J. Pure Appl. Algebra, vol. 226, no. 10 (2022), Paper No. 107037, 57 (cit. on pp. 10, 34). [Dor23] C. Dorn, Associative n-Categories, preprint 2023, arXiv: 1812.10586 [math] (cit. on pp. 11, 18). [Dra+24] A. Dranowski, M. Guo, A. Lauda, and A. Manion, Spectral 2-Actions, Foams, and Frames in the Spectrification of Khovanov Arc Algebras, preprint 2024, arXiv: 2402.11368 [math] (cit. on p. 19). [Dup21] B. Dupont, Rewriting modulo Isotopies in Khovanov-Lauda-Rouquier's Categorification of Quantum Groups, Adv. Math., vol. 378 (2021), Paper No. 107524, 75 (cit. on pp. 3, 10, 17, 34, 36, 41, 43, 49, 54). B. Dupont, Rewriting modulo Isotopies in Pivotal Linear (2,2)-Categories, J. Algebra, [Dup22] vol. 601 (2022), pp. 1-53 (cit. on pp. 3, 4, 10, 17, 34, 36, 41-43, 49, 54, 56). [EL16] A. P. Ellis and A. D. Lauda, An Odd Categorification of $U_q(\mathfrak{sl}_2)$, Quantum Topol., vol. 7, no. 2 (2016), pp. 329-433 (cit. on p. 3). [Eli22] B. Elias, A Diamond Lemma for Hecke-type Algebras, Trans. Amer. Math. Soc., vol. 375, no. 3 (2022), pp. 1883-1915 (cit. on p. 19). [EW14] B. Elias and G. Williamson, The Hodge Theory of Soergel Bimodules, Ann. of Math. (2), vol. 180, no. 3 (2014), pp. 1089–1136 (cit. on p. 2). [FM18] S. Forest and S. Mimram, Coherence of Gray Categories via Rewriting, in: 3rd International Conference on Formal Structures for Computation and Deduction, vol. 108, LIPIcs. Leibniz Int. Proc. Inform. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2018, Art. No. 15, 16 (cit. on pp. 4, 17, 19). [FM22] S. Forest and S. Mimram, Rewriting in Gray Categories with Applications to Coherence, Math. Structures Comput. Sci., vol. 32, no. 5 (2022), pp. 574-647 (cit. on pp. 11, 12, 17, 19, 20, 22, 26, 27, 30, 48). [FY89] P. J. Freyd and D. N. Yetter, Braided Compact Closed Categories with Applications to Low-Dimensional Topology, Adv. Math., vol. 77, no. 2 (1989), pp. 156-182 (cit. on p. 3). [GGM15] S. Gaussent, Y. Guiraud, and P. Malbos, Coherent Presentations of Artin Monoids, Compos. Math., vol. 151, no. 5 (2015), pp. 957-998 (cit. on p. 19). Y. Guiraud, E. Hoffbeck, and P. Malbos, Convergent Presentations and Polygraphic [GHM19] Resolutions of Associative Algebras, Math. Z., vol. 293, no. 1-2 (2019), pp. 113-179 (cit. on pp. 3, 9, 17, 19, 39, 41-43, 48, 57). [GM09] Y. Guiraud and P. Malbos, Higher-Dimensional Categories with Finite Derivation Type, Theory Appl. Categ., vol. 22 (2009), No. 18, 420-478 (cit. on p. 8).

[GM13]	Y. Guiraud and P. Malbos, Identities among Relations for Higher-Dimensional Rewriting Systems, in: <i>OPERADS 2009</i> , vol. 26, Sémin. Congr. Soc. Math. France, Paris, 2013, pp. 145–161 (cit. on p. 8).
[GM18]	Y. Guiraud and P. Malbos, Polygraphs of Finite Derivation Type, <i>Math. Structures Comput. Sci.</i> , vol. 28, no. 2 (2018), pp. 155–201 (cit. on pp. 8, 13, 60).
[GPS95]	R. Gordon, A. J. Power, and R. Street, Coherence for Tricategories, <i>Mem. Amer. Math. Soc.</i> , vol. 117, no. 558 (1995), pp. vi+81 (cit. on p. 11).
[Gue21]	L. Guetta, Homology of Categories via Polygraphic Resolutions, <i>J. Pure Appl. Algebra</i> , vol. 225, no. 10 (2021), Paper No. 106688, 33 (cit. on p. 8).
[Gui06]	Y. Guiraud, Termination Orders for 3-Polygraphs, C. R. Math. Acad. Sci. Paris, vol. 342, no. 4 (2006), pp. 219–222 (cit. on p. 8).
[HKK16]	P. Hu, D. Kriz, and I. Kriz, Field Theories, Stable Homotopy Theory, and Khovanov Homology, <i>Topology Proc.</i> , vol. 48 (2016), pp. 327–360 (cit. on p. 19).
[Hue77]	G. Huet, Confluent Reductions: Abstract Properties and Applications to Term Rewriting Systems, in: <i>18th Annual Symposium on Foundations of Computer Science</i> <i>(Providence, R.I., 1977)</i> , IEEE, Long Beach, CA, 1977, pp. 30–45 (cit. on p. 10).
[JK86]	JP. Jouannaud and H. Kirchner, Completion of a Set of Rules modulo a Set of Equations, <i>SIAM J. Comput.</i> , vol. 15, no. 4 (1986), pp. 1155–1194 (cit. on p. 10).
[JL12]	JP. Jouannaud and J. Li, Church-Rosser Properties of Normal Rewriting, in: <i>Computer Science Logic 2012</i> , vol. 16, LIPIcs. Leibniz Int. Proc. Inform. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2012, pp. 350–365 (cit. on p. 10).
[Jon85]	V. F. R. Jones, A Polynomial Invariant for Knots via von Neumann Algebras, <i>Bull. Amer. Math. Soc. (N.S.)</i> , vol. 12, no. 1 (1985), pp. 103–111 (cit. on p. 2).
[JS]	A. Joyal and R. Street, The Geometry of Tensor Calculus II, available on Ross Street's website (cit. on p. 3).
[JS91]	A. Joyal and R. Street, The Geometry of Tensor Calculus. I, <i>Adv. Math.</i> , vol. 88, no. 1 (1991), pp. 55–112 (cit. on p. 3).
[Kau87]	L. H. Kauffman, State Models and the Jones Polynomial, <i>Topology</i> , vol. 26, no. 3 (1987), pp. 395–407 (cit. on p. 2).
[KK12]	SJ. Kang and M. Kashiwara, Categorification of Highest Weight Modules via Khovanov-Lauda-Rouquier Algebras, <i>Invent. Math.</i> , vol. 190, no. 3 (2012), pp. 699–742 (cit. on p. 3).
[KKO13]	SJ. Kang, M. Kashiwara, and Sj. Oh, Supercategorification of Quantum Kac-Moody Algebras, <i>Adv. Math.</i> , vol. 242 (2013), pp. 116–162 (cit. on p. 3).
[KL09]	M. Khovanov and A. Lauda, A Diagrammatic Approach to Categorification of Quantum Groups I, <i>Represent. Theory</i> , vol. 13, no. 14 (2009), pp. 309–347 (cit. on p. 2).
[KL10]	M. Khovanov and A. Lauda, A Categorification of Quantum $sl(n)$, Quantum Topol. (2010), pp. 1–92 (cit. on pp. 2, 3).
[KL80]	G. M. Kelly and M. L. Laplaza, Coherence for Compact Closed Categories, <i>J. Pure Appl. Algebra</i> , vol. 19 (1980), pp. 193–213 (cit. on p. 3).
[Kup96]	G. Kuperberg, Spiders for Rank 2 Lie Algebras, <i>Comm. Math. Phys.</i> , vol. 180, no. 1 (1996), pp. 109–151 (cit. on p. 2).

[Laf03]	Y. Lafont, Towards an Algebraic Theory of Boolean Circuits, <i>J. Pure Appl. Algebra</i> , vol. 184, no. 2-3 (2003), pp. 257–310 (cit. on p. 8).
[Liu+24]	Y. L. Liu, A. Mazel-Gee, D. Reutter, C. Stroppel, and P. Wedrich, A Braided Monoidal $(\infty, 2)$ -Category of Soergel Bimodules, preprint 2024, arXiv: 2401.02956 [math] (cit. on p. 19).
[Liu24]	Y. L. Liu, Braiding on Complex Oriented Soergel Bimodules, preprint 2024, arXiv: 2407.04891 [math] (cit. on p. 19).
[LLS20]	T. Lawson, R. Lipshitz, and S. Sarkar, Khovanov Homotopy Type, Burnside Category and Products, <i>Geom. Topol.</i> , vol. 24, no. 2 (2020), pp. 623–745 (cit. on p. 19).
[LM09]	Y. Lafont and F. Métayer, Polygraphic Resolutions and Homology of Monoids, <i>J. Pure Appl. Algebra</i> , vol. 213, no. 6 (2009), pp. 947–968 (cit. on p. 8).
[LMW10]	Y. Lafont, F. Métayer, and K. Worytkiewicz, A Folk Model Structure on Omega-Cat, <i>Adv. Math.</i> , vol. 224, no. 3 (2010), pp. 1183–1231 (cit. on p. 8).
[LS14a]	R. Lipshitz and S. Sarkar, A Khovanov Stable Homotopy Type, J. Amer. Math. Soc., vol. 27, no. 4 (2014), pp. 983–1042 (cit. on pp. 19, 80).
[LS14b]	R. Lipshitz and S. Sarkar, A Steenrod Square on Khovanov Homology, <i>J. Topol.</i> , vol. 7, no. 3 (2014), pp. 817–848 (cit. on p. 80).
[Mar47]	A. Markoff, On the Impossibility of Certain Algorithms in the Theory of Associative Systems, <i>C. R. (Doklady) Acad. Sci. URSS (N.S.)</i> , vol. 55 (1947), pp. 583–586 (cit. on p. 6).
[Mar98]	C. Marché, Normalized Rewriting: A Unified View of Knuth-Bendix Completion and Gröbner Bases Computation, in: <i>Symbolic Rewriting Techniques (Ascona, 1995)</i> , vol. 15, Progr. Comput. Sci. Appl. Logic, Birkhäuser, Basel, 1998, pp. 193–208 (cit. on p. 10).
[Mét03]	F. Métayer, Resolutions by Polygraphs, <i>Theory Appl. Categ.</i> , vol. 11 (2003), No. 7, 148–184 (cit. on p. 8).
[Mim10]	S. Mimram, "Computing Critical Pairs in 2-Dimensional Rewriting Systems", in: <i>RTA 2010: Proceedings of the 21st International Conference on Rewriting Techniques and Applications</i> , vol. 6, LIPIcs. Leibniz Int. Proc. Inform. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2010, pp. 227–241 (cit. on p. 8).
[Mim14]	S. Mimram, Towards 3-Dimensional Rewriting Theory, <i>Log. Methods Comput. Sci.</i> , vol. 10, no. 2 (2014), 2:1, 47 (cit. on p. 8).
[MR20]	A. Manion and R. Rouquier, Higher Representations and Cornered Heegaard Floer Homology, preprint 2020, arXiv: 2009.09627 [math] (cit. on p. 19).
[Pei21]	D. J. Peifer, Reinforcement Learning in Buchberger's Algorithm, ProQuest LLC, Ann Arbor, MI, 2021, 170 pp. (cit. on p. 18).
[Pos47]	E. L. Post, Recursive Unsolvability of a Problem of Thue, <i>J. Symbolic Logic</i> , vol. 12 (1947), pp. 1–11 (cit. on p. 6).
[PS81]	G. E. Peterson and M. E. Stickel, Complete Sets of Reductions for Some Equational Theories, <i>J. Assoc. Comput. Mach.</i> , vol. 28, no. 2 (1981), pp. 233–264 (cit. on p. 10).
[Rou08]	R. Rouquier, 2-Kac–Moody Algebras, preprint 2008, arXiv: 0812.5023 (cit. on p. 2).

[RV19]	D. Reutter and J. Vicary, High-Level Methods for Homotopy Construction in Associative n-Categories, in: 2019 34th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS), IEEE, [Piscataway], NJ, 2019, [13 pp.] (Cit. on pp. 11, 18).
[Sch24]	L. Schelstraete, Odd Khovanov Homology, Higher Representation Theory and Higher Rewriting Theory, Ph.D. thesis, UCLouvain, 2024, 264 pp., available at arXiv:2410.11405 (cit. on pp. 19, 60, 66–68, 72, 80).
[See12]	C. Seed, Computations of the Lipshitz-Sarkar Steenrod Square on Khovanov Homology, preprint 2012, arXiv: 1210.1882 [math] (cit. on p. 80).
[Sel11]	P. Selinger, A Survey of Graphical Languages for Monoidal Categories, in: <i>New Structures for Physics</i> , vol. 813, Lecture Notes in Phys. Springer, Heidelberg, 2011, pp. 289–355 (cit. on pp. 2, 3).
[Shi09]	A. I. Shirshov, Some Algorithmic Problems for Lie Algebras, in: <i>Selected Works of A.I. Shirshov</i> , ed. by L. Bokut, I. Shestakov, V. Latyshev, and E. Zelmanov, Basel: Birkhäuser, 2009, pp. 125–130 (cit. on p. 3).
[SSS20]	S. Sarkar, C. Scaduto, and M. Stoffregen, An Odd Khovanov Homotopy Type, <i>Adv. Math.</i> , vol. 367 (2020), p. 107112 (cit. on p. 19).
[Str76]	R. Street, Limits Indexed by Category-Valued 2-Functors, Journal of Pure and Applied Algebra, vol. 8, no. 2 (1976), pp. 149–181 (cit. on p. 8).
[Str96]	R. Street, Categorical Structures, in: <i>Handbook of Algebra, Vol. 1</i> , vol. 1, Handb. Algebr. Elsevier/North-Holland, Amsterdam, 1996, pp. 529–577 (cit. on pp. 11, 19).
[SV23]	L. Schelstraete and P. Vaz, Odd Khovanov Homology and Higher Representation Theory, preprint 2023, arXiv: 2311.14394 [math] (cit. on pp. 2, 5, 12, 31, 61, 65, 68, 74, 80).
[Vir95]	P. Viry, Rewriting modulo a Rewrite System, Pisa, IT: Università di Pisa, 1995 (cit. on p. 10).
[vOos94]	V. van Oostrom, Confluence by Decreasing Diagrams, <i>Theoret. Comput. Sci.</i> , vol. 126, no. 2 (1994), pp. 259–280 (cit. on p. 36).
[Web13]	M. Weber, Free Products of Higher Operad Algebras, <i>Theory Appl. Categ.</i> , vol. 28 (2013), No. 2, 24–65 (cit. on p. 19).
[Web18]	B. Webster, Unfurling Khovanov-Lauda-Rouquier Algebras, preprint 2018, arXiv: 1603.06311 [math] (cit. on p. 3).