

# An overview of dualities in non-commutative harmonic analysis

Yulia Kuznetsova

## 1 Introduction

The aim of this mini-course is to give an overview of tools of the non-commutative harmonic analysis. By volume limitations, it cannot be complete, but tries to be as wide as possible and accessible to starting PhD students.

Most of the general theory presented below can be found in the books of Jacques Dixmier [4] and Gerald Folland [6]. On spherical theory, the recommended reference is Sigurdur Helgason, *Groups and Geometric Analysis* [8]. And on quantum groups, Thomas Timmermann, *An Invitation to Quantum Groups and Duality* [19]. Other sources are research articles.

### *Duality and Fourier transform*

Duality is one of the main tools in classical harmonic analysis. The Fourier transform  $\mathcal{F}$  takes convolution to pointwise multiplication, thus allowing to simplify differential equations and to estimate their solutions. Keep in mind that solutions of differential equations are convolutions with Green functions, so we are particularly interested in convolution operators  $T : f \mapsto f * k$ ; the kernel  $k$  can be a function, measure, or a distribution.

This theory is fully available on an Abelian locally compact group  $G$ : the group of characters of  $G$

$$\widehat{G} = \{\phi : G \rightarrow \mathbb{T} \text{ continuous homomorphism}\}$$

is also an Abelian LCG, and  $\widehat{\widehat{G}} \simeq G$ . The Fourier transform  $\mathcal{F}$  satisfies  $\mathcal{F}(f * g) = \mathcal{F}(f)\mathcal{F}(g)$  and is isometric on  $L^2$ , thus it is easy to say when  $T$  is bounded on  $L^2$ : exactly when  $\mathcal{F}(k)$  is bounded.

---

University of Franche-Comté, Besançon, France

If  $G$  is non-abelian, such a duality is not possible. The first natural analogue of the group of characters is  $\widehat{G}$ : the set of all unitary irreducible representations of  $G$ . But  $\widehat{G}$  is no more a group. Three solutions exist:

- Work with  $\widehat{G}$  as a topological or a measured space  
→ Plancherel measure, leads to an isometry on  $L^2$  in many cases
- Consider only finite-dimensional representations  
→ tensor categories, especially useful for compact groups
- Study group algebras  
→ Duality of quantum groups

And a special domain is the study of subclasses of operators which admit Fourier-like methods (analysis on symmetric spaces).

I will speak of all of them except for the second one which could not enter in the volume available. A good reference for this approach is [15].

## 2 Unitary dual

Abelian groups are well described by their characters having scalar values, and the Fourier transform of a function is again a scalar-valued function. In the non-commutative case, representations act on Hilbert spaces, possibly of infinite dimension, and the Fourier transform will be an operator-valued function on  $\widehat{G}$ .

Recall that every unitary representation  $\pi$  of  $G$  on a Hilbert space  $H_\pi$  generates a representation of  $L^1(G)$  by

$$\pi(f) = \int_G f(g)\pi(g)dg; \quad (1)$$

the integral is understood as an operator such that for every  $x, y \in H_\pi$

$$\langle \pi(f)x, y \rangle = \int_G f(g)\langle \pi(g)x, y \rangle dg.$$

This representation extends by continuity to  $C^*(G)$ . Conversely, every \*-representation of  $C^*(G)$  defines a unitary representation of  $G$  (provided that  $\pi(C^*(G))H$  is dense in  $H$ ). A representation  $\pi$  is called irreducible if it has no nontrivial closed invariant subspaces. If  $G$  is abelian, every irreducible  $\pi$  is one-dimensional.

Representations  $\pi, \rho$  of  $G$  on Hilbert spaces  $H_\pi, H_\rho$  are called equivalent if there exists a unitary  $T : H_\pi \rightarrow H_\rho$  such that  $T\pi(g) = \rho(g)T$  for every  $g \in G$ .

**Definition 1.**  $\widehat{G}$  is the set of equivalence classes of *unitary irreducible* representations.

**Nilpotent case**

It is usually a complicated task to describe  $\widehat{G}$ . For a nilpotent simply connected Lie group  $G$ , a powerful orbit method is available. Examples of such groups include strictly upper-triangular matrices and Heisenberg groups  $\mathbb{R} \ltimes \mathbb{R}^n$ .

Let  $\mathfrak{g}$  denote the Lie algebra of  $G$ . At  $X \in \mathfrak{g}$ , the *adjoint action* of  $g \in G$  is

$$Ad(g) : \exp(tX) \mapsto g \exp(tX) g^{-1}.$$

On  $\mathfrak{g}^*$ , this generates the *coadjoint action*:

$$Ad^*(g) : \lambda \mapsto \lambda \circ Ad(g).$$

We say that  $\lambda \sim \mu$  if  $\lambda = Ad^*(g)\mu$  for some  $g$ . This is an equivalence relation. Without going into further detail, let us state

**Theorem 1 (Kirillov [12])**  $\widehat{G}$  is in bijection with  $\mathfrak{g}^* / \sim$ .

**Heisenberg group**

This example will help to see the identification above. We set  $\mathbb{H} = \mathbb{R} \ltimes \mathbb{R}^2$ , or the group of matrices  $\begin{pmatrix} 1 & a & c \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix}$ .

Its Lie algebra is  $\mathfrak{g} = \langle X, Y, Z \rangle$  with

$$X = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad Y = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \quad Z = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

so that  $[X, Y] = Z$  and  $[X, Z] = [Y, Z] = 0$ .

The space  $\mathfrak{g}^*$  is 3-dimensional and the coadjoint action (we do not calculate it here) is

$$Ad^*(a, b, c) : (\mu, \nu, \lambda) \mapsto (\mu + b\lambda, \nu - a\lambda, \lambda).$$

We see that the orbit of  $(\mu, \nu, 0)$  is only this point, and if  $\lambda \neq 0$ , the orbit is the plane  $\Omega_\lambda = \{(\mu, \nu, \lambda) : (\mu, \nu) \in \mathbb{R}^2\}$ . Denote  $\Omega = \{\Omega_\lambda : \lambda \in \mathbb{R}\}$ .

For  $\mu, \nu \in \mathbb{R}$ , the one-dimensional representation corresponding to  $(\mu, \nu, 0)$  is given by

$$\pi_{\mu, \nu}(a, b, c) = e^{2\pi i(a\mu + b\nu)};$$

for  $\lambda \in \mathbb{R}$ , infinite-dimensional irreducibles act on  $f \in L^2(\mathbb{R})$  as

$$\pi_\lambda(a, b, c)(f)(t) = e^{2\pi i\lambda(c+bt)} f(t+a).$$

By direct calculations, one can verify that for  $f \in L^1(\mathbb{H}) \cap L^2(\mathbb{H})$ ,

$$\|f\|_2^2 = \int_{\mathbb{R}} Tr[\pi_\lambda(f)^* \pi_\lambda(f)] |\lambda| d\lambda.$$

The measure

$$\mu(\{\pi_{\mu,\nu}\}) = 0, \quad \mu|_{\Omega} = |\lambda|d\lambda \quad (2)$$

is called the Plancherel measure on  $\widehat{\mathbb{H}}$ , and we will discuss it below in the general case.

**Topology on the unitary dual**

This dual is more often considered as a measure space, but one starts with a topology. For  $\pi \in \widehat{G}$ , its kernel  $\ker \pi$  is an ideal in  $C^*(G)$ . We define the *closure* of any  $E \subset \widehat{G}$ :

$$\overline{E} = \{\pi \in \widehat{G} : \pi \equiv 0 \text{ on } I_E = \bigcap_{\rho \in E} \ker \rho\}.$$

There exists (by Kuratowski theorem) a topology on  $\widehat{G}$  in which this is the closure map. One calls it the Fell, or hull-kernel topology.  $\widehat{G}$  is not always Hausdorff.

If  $G$  is *abelian*, then  $C^*(G) \simeq C_0(\widehat{G})$ . For  $\rho \in \widehat{G}$ , we have  $\ker \rho = \{f \in C_0(\widehat{G}) : f(\rho) = 0\}$ , so that  $I_E = \{f \in C_0(\widehat{G}) : f|_E \equiv 0\}$ . Then

$$\overline{E} = \{\pi \in \widehat{G} : f(\pi) = 0 \text{ for every } f \in C_0(\widehat{G}) \text{ vanishing on } E\}.$$

Clearly this is the usual closure of  $E$ .

## 2.1 Plancherel measure

For  $f \in L^1(G)$  and  $\pi \in \widehat{G}$ , set  $\widehat{f}(\pi) = \pi(f)$ , as given by the formula (1): this defines the noncommutative Fourier transform. Sometimes one sets instead  $\pi(f) = \int_G f(g)\pi(g^{-1})dg$ .

For groups of so called ‘‘Type I’’, there is a direct analogy with the classical case:

**Theorem 2** *Let  $G$  be a unimodular Type I group. There exists a unique positive measure  $\mu$  on  $\widehat{G}$  such that for every  $f \in L^1(G) \cap L^2(G)$*

$$\|f\|_2^2 = \int_{\widehat{G}} \text{Tr}(\widehat{f}(\pi)^* \widehat{f}(\pi)) d\mu(\pi). \quad (3)$$

There exists a non-unimodular version [6, Theorem 7.50] but the group should be ‘‘almost’’ Type I.

Let us now define the term ‘‘Type I’’. Recall that a subalgebra  $A$  in  $B(H)$  (the space of bounded operators on a Hilbert space  $H$ ) is called a von Neumann algebra if it is closed in strong or, equivalently, weak operator topology. This happens if and only if  $A$  is equal to its second commutant  $A''$ ; and for a set  $S \subset B(H)$ , its commutant is defined as  $S' = \{y \in B(H) : yx = xy \ \forall x \in S\}$ . Next, a von Neumann algebra  $A$  is a factor if  $A \cap A' = \mathbb{C}1$ .

Now, being Type I means: if  $\pi(G)''$  is a factor, then it is of type I, that is, isomorphic to  $B(H)$  (of finite or infinite dimension). This is also equivalent to the following: if  $\pi$  is irreducible, then  $\pi(C^*(G))$  contains all compact operators. If  $G$  is not of Type I, then it has necessarily both Type II and Type III representations [7]. An example of a non-Type I group is the free group of two generators.

Other terms for this class are GCR and postliminal groups. These notions have appeared independently and have different definitions, but have turned to be the same classes of groups as Type I.

In relation to the dual, the following facts are known. For any  $G$ , if  $\pi_1 = \pi_2 \in \widehat{G}$ , then clearly  $\ker \pi_1 = \ker \pi_2$ . For Type I groups, the converse is true. Moreover, a second countable group  $G$  is Type I if and only if the map  $\pi \mapsto \ker \pi$  is injective, and if and only if the topology of  $\widehat{G}$  is  $T_0$ .

Finally, if  $G$  is Type I and  $\pi$  irreducible, then the von Neumann algebra  $\pi(G)''$  admits a trace.

- Every connected semisimple and nilpotent Lie group is Type I.
- A discrete group is Type I if and only if it is almost abelian (that is, has an abelian subgroup of finite index).

### **Compact groups**

In this case,  $\widehat{G}$  is discrete: if  $\chi_\pi : g \mapsto \text{Tr} \pi(g)$  is the character of  $\pi$ , then  $\rho(\chi_\pi) = 0$  if  $\pi \neq \rho$  and  $\pi(\chi_\pi) \neq 0$ , so that  $\widehat{G} \setminus \{\pi\}$  is closed, thus  $\{\pi\}$  open.

The Plancherel theorem takes the following form:

$$\|f\|_2^2 = \sum_{\pi \in \widehat{G}} d_\pi \text{Tr}(\pi(f)\pi(f)^*).$$

### **Inverse Fourier transform**

The Plancherel identity implies that for  $f, g \in L^1 \cap L^2(G)$

$$\begin{aligned} \langle f, g \rangle &= \int_{\widehat{G}} \text{Tr}(\pi(g)^* \pi(f)) d\mu(\pi) \\ &= \int_{\widehat{G}} \int_G \text{Tr}(\overline{g(x)} \pi(x)^* \pi(f)) dx d\mu(\pi), \end{aligned}$$

so that

$$f(x) = \int_{\widehat{G}} \text{Tr}(\pi(x)^* \pi(f)) d\mu(\pi).$$

This allows in particular to define pseudo-differential operators on such groups [17].

### **Plancherel measure: the support**

The support of  $\mu$  (intersection of all closed sets of full measure) is the set of  $\rho \in \widehat{G}$  weakly contained in  $\lambda$ , that is, such that for every  $\xi \in H_\rho$ ,  $\varepsilon > 0$ , and compact  $K \subset G$  there exist  $f_k \in L^2(G)$  with

$$\sup_{g \in K} \left| \langle \rho(g)\xi, \xi \rangle - \sum_{k=1}^n \tilde{f}_k * f_k(g) \right| < \varepsilon.$$

Here  $\tilde{f}(x) = \bar{f}(x^{-1})$ .

$G$  is amenable iff  $\text{supp } \mu = \widehat{G}$ . This is not the case, for example, for  $SL_2(\mathbb{R})$ .

**Example: Heisenberg group**

Recall that  $\widehat{\mathbb{H}} = \{\pi_{\mu, \nu}\} \cup \{\pi_\lambda\}$ , and only  $\pi_\lambda$  enter in the Plancherel formula (2). For  $k \in L^1(\mathbb{H}) \cap L^2(\mathbb{H})$ ,

$$\begin{aligned} \pi_\lambda(k)f(t) &= \int_H k(a, b, c) e^{2\pi i \lambda(c+bt)} f(t+a) da db dc \\ &= \int_{\mathbb{R}} \mathcal{F}_2 \mathcal{F}_3(k)(x-t, -\lambda, -\lambda t) f(x) dx \end{aligned}$$

so that  $\widehat{k}(\pi_\lambda)$  is an integral operator with the kernel as above.

**Plancherel measure: non-unimodular case**

The Plancherel measure still exists, with Formula (3) and (4) valid, but the group should be ‘‘almost Type I’’ [6, Theorem 7.50] and we have to adjust the definition of the Fourier transform setting

$$\widehat{f}(\pi) = \pi(f)D_\pi,$$

with  $\pi(f)$  as in (1). Here  $D_\pi$  are self-adjoint operators on  $H_\pi$  such that  $D_\pi \pi(g) = \Delta(g)^{1/2} \pi(g) D_\pi$  for every  $g \in G$ . Their existence is not obvious, but becomes clearer knowing that every  $\pi$  can be realized on the space  $H_\pi = L^2(G, V_\pi)$  where  $V_\pi$  is also a Hilbert space; one sets then  $D_\pi f(g) = \Delta(g)^{-1/2} f(g)$  for  $f \in H_\pi$  and  $g \in G$ .

The Plancherel identity for  $f, g \in L^1 \cap L^2(G)$  becomes

$$\langle f, g \rangle = \int_{\widehat{G}} \int_G \text{Tr}(D_\pi \overline{g(x)} \pi(x)^* \pi(f) D_\pi) dx d\mu(\pi),$$

and the inverse Fourier transform

$$f(x) = \int_{\widehat{G}} \text{Tr}(\pi(x)^* \pi(f) D_\pi^2) d\mu(\pi).$$

**‘ $ax+b$ ’ group**

$G$  is the group of matrices

$$\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}$$

with  $a > 0$ ,  $b \in \mathbb{R}$ , which correspond to the affine maps on the real line:  $g_{a,b}x = ax + b$ .

We could consider also  $G_n = \{(a, b) : a \in \mathbb{R}, b \in \mathbb{R}^n\}$ , acting the same way.  $G$  is non-unimodular: the left Haar measure is  $\frac{1}{a^2}dad b$ , the right one  $\frac{1}{a}dad b$ , the modular function is  $a^{-1}$ .

One can describe  $\widehat{G}$  by the orbit method, arriving at:  $\widehat{G} = \{\pi_\lambda : \lambda \in \mathbb{R}\} \cup \{\pi^+, \pi^-\}$ , acting as

$$\pi_\lambda(a, b) = a^{i\lambda};$$

$\pi^+, \pi^-$  act on  $H^+ = L^2(0, +\infty)$  and  $H^- = L^2(-\infty, 0)$  respectively by the same formula:

$$\pi^\pm(a, b)\phi(t) = \sqrt{a} e^{2\pi i b t} \phi(at).$$

The Plancherel measure  $\mu$  on  $\widehat{G}$  is such that  $\mu(\{\pi_\lambda\}) = 0$  for every  $\lambda$ , and

$$\|f\|_2^2 = Tr[\widehat{f}(\pi^+) * \widehat{f}(\pi^+)] + Tr[\widehat{f}(\pi^-) * \widehat{f}(\pi^-)]$$

for  $f \in L^2(G)$ . The adjustment operators are  $D^\pm \phi(s) = \sqrt{|s|} \phi(s)$ , and  $\widehat{f}(\pi^\pm) = \pi^\pm(f) D^\pm$ . One verifies directly that  $\widehat{f}(\pi^+)$  is an integral operator with the kernel

$$k_f^+(s, t) = \frac{\sqrt{s}}{t} \mathcal{F}_2 f(s^{-1}t, -s),$$

the Fourier transform being applied in the second variable only. A similar formula holds for  $\widehat{f}(\pi^-)$ .

## 2.2 Disintegration

The Plancherel measure has much more properties than just the isometry of the Fourier transform on  $L^2(G)$ . The next property to discuss is the decomposition of representations into direct integrals of irreducibles [4].

Let  $(\Omega, \mu)$  be a measured space and  $H_\omega$  a Hilbert space for every  $\omega \in \Omega$ . A *measurable field* of Hilbert spaces is a linear subspace  $\Gamma \subset \prod_{\omega \in \Omega} H_\omega$  such that

- $\omega \mapsto \|x(\omega)\|$  is measurable for every  $x \in \Gamma$ ;
- there exist  $(x_n) \subset \Gamma$  such that  $\{x_n(\omega)\}_{n \in \mathbb{N}}$  is total in  $H_\omega$  for every  $\omega$ ;
- if  $\omega \mapsto \langle x(\omega), y(\omega) \rangle$  is measurable for every  $x \in \Gamma$ , then  $x \in \Gamma$ .

We denote

$$\Gamma := \int^\oplus H_\omega d\mu(\omega).$$

If  $A(\omega) \in B(H_\omega)$  for every  $\omega$  and  $(A(\omega)x(\omega)) \in \Gamma$  for every  $x \in \Gamma$ , then  $A = \int^\oplus A(\omega) d\mu(\omega)$  is said to be a measurable field of operators on  $\Gamma$ . It is bounded if and only if

$$\|A\| = \operatorname{esssup}_{\omega \in \Omega} \|A(\omega)\| < \infty.$$

***Disintegration of representations***

If  $\pi_\omega$  is a unitary representation of  $G$  on  $H_\omega$  for every  $\omega \in \Omega$  and  $(\pi_\omega(g))$  is a measurable field of operators for every  $g \in G$ , then  $\pi = \int^\oplus \pi_\omega d\mu(\omega)$ , the direct integral of  $\pi_\omega$ , is a representation of  $G$  on  $\int^\oplus H_\omega$ . Conversely, if  $G$  is a second countable locally compact group, then every unitary representation  $\pi$  of  $G$  on a separable  $H$  is unitarily equivalent to a direct integral  $\int^\oplus \pi_\omega d\mu(\omega)$  where  $\pi_\omega$  are irreducible for almost all  $\omega$  [4, 8.5.2], [6, 7.37].

This disintegration is not unique, and this can be seen on the example of the most important representation: of the regular one. The left regular representation  $\lambda$  of  $G$  acts at  $f \in L^2(G)$  as

$$[\lambda(g)f](h) = f(g^{-1}h).$$

***A non-uniqueness example*** [21]. Let  $\mathbb{F}_2$  be the free group of two generators  $a$  and  $b$ . Then  $\lambda$  has two decompositions into irreducibles

$$\lambda = \int_{\mathbb{T}} U_\theta d\theta = \int_{\mathbb{T}} V_\theta d\theta$$

such that none of  $U_\theta, V_{\theta'}$  are equivalent.

We introduce briefly these representations. Let  $G_a, G_b$  denote all words in  $\mathbb{F}_2$  which end by  $a, a^{-1}$  or  $b, b^{-1}$  respectively, and set  $G'_a = G_a \cup \{e\}$ ,  $G'_b = G_b \cup \{e\}$ . For  $\theta \in \mathbb{T}$ , the space  $H_\theta$  is the closed linear span of the indicator functions  $\{\delta_g : g \in G'_b\}$  which are its orthonormal basis;  $U_\theta$  acts as

$$\begin{aligned} U_\theta(a)\delta_e &= \theta\delta_e, & U_\theta(a)\delta_g &= \delta_{ag}, & g \in G_b; \\ U_\theta(b)\delta_g &= \delta_{bg}. \end{aligned}$$

For  $V_\theta$ , the construction is the same with  $a$  and  $b$  interchanged. One can show that  $U_\theta, V_{\theta'}$  are irreducible and pairwise inequivalent.

***Disintegration over the dual***

To achieve uniqueness, or otherwise stated, to integrate over the dual, one has to return to Type I groups. Multiplicities occur in this process:

**Theorem 3** [6, 7.40] *Let  $G$  be a second countable Type I group,  $\sigma$  a unitary representation of  $G$  on a separable  $H$ . Then there exist finite pairwise orthogonal measures  $\mu_n$ ,  $n \in \mathbb{N} \cup \{\infty\}$ , on  $\widehat{G}$ , such that*

$$\sigma \simeq \int^\oplus \pi d\mu_1(\pi) \oplus 2 \int^\oplus \pi d\mu_2(\pi) \oplus \cdots \oplus |\mathbb{N}| \int^\oplus \pi d\mu_\infty(\pi).$$

*These measures are unique up to equivalence.*



In particular, for the regular representation

$$\lambda = \int_{\pi \in \widehat{G}} \pi \otimes I_{\overline{H}_\pi} d\mu(\pi) \quad (4)$$

with the Plancherel measure  $\mu$ ; the group should be second countable and Type I [6, 7.44, 7.50]. If  $G$  is compact, we recover the familiar decomposition

$$\lambda = \bigoplus_{\pi \in \widehat{G}} d_\pi \pi.$$

### ***Translation-invariant operators***

Not only  $\lambda(g)$  but every translation invariant operator admits such a disintegration. Recall that convolution operators are translation invariant; often, converse is true [20], [5, Corollary 3.2.1]. If  $T$  is bounded on  $L^2$ , then:

$$T \text{ is left invariant} \Leftrightarrow T \in \lambda(G)'' ,$$

where  $S''$  is the bicommutant in  $B(L^2(G))$ , or in other terms, the von Neumann algebra generated by  $S$ .

In the same assumptions on  $G$  as above, for every  $T \in \lambda(G)''$

$$T = \int_{\widehat{G}}^{\oplus} T_\pi \otimes I_{\overline{H}_\pi} d\mu(\pi). \quad (5)$$

Only the support of  $\lambda$  enters in this decomposition. If  $Tf = k * f$ , then

$$T_\pi = \widehat{k}(\pi) = \int_G k(x)\pi(x)dx.$$

### ***Heisenberg group: sub-Laplacian***

One can verify that, on  $L^2(\mathbb{R})$ , for every  $\lambda \in \mathbb{R}$

$$\pi_\lambda(X) = \sqrt{|\lambda|} \frac{\partial}{\partial t}, \quad \pi_\lambda(Y) = i \operatorname{sgn} \lambda \sqrt{|\lambda|} t, \quad \pi_\lambda(Z) = 2\pi i \lambda.$$

If  $\mathcal{L} = X^2 + Y^2$  is the sub-Laplacian, then, even if it is unbounded, the disintegration (5) holds for it [5], with

$$\mathcal{L}_{\pi_\lambda} f(t) = |\lambda| (f''(t) - t^2 f(t)). \quad (6)$$

## **3 Symmetric spaces**

Let  $G$  be a semisimple Lie group,  $K$  its maximal compact subgroup. By Iwasawa decomposition,  $G = KAN$  with  $A \simeq \mathbb{R}^l$  abelian,  $N$  nilpotent. Let  $\mathfrak{a} \simeq \mathbb{R}^l$  and  $\mathfrak{g}$  denote the Lie algebras of  $A$  and  $G$ .

The principal series representations  $\pi_{\sigma,\lambda}$  of  $G$  are parametrized by irreducibles  $\sigma \in \widehat{K}$  and  $\lambda \in \mathfrak{a}^*$ ; if  $\sigma = 1$ , they have  $K$ -invariant vectors and are called spherical. One can realize them on subspaces of  $L^2(K)$ .

For every  $\lambda \in \mathfrak{a}^*$ , the function  $\phi_\lambda$  on  $G$  is the following coefficient of  $\pi_{1,\lambda}$ :

$$\phi_\lambda(x) = \langle \pi_\lambda(x)1, 1 \rangle_{L^2(K)}.$$

For every  $x \in G$  one can write uniquely  $x = k_x a_x n_x$ ; set  $H(x) = \log a_x$ , then

$$\phi_\lambda(x) = \int_K e^{-(i\lambda+\rho)(H(x^{-1}k))} dk,$$

where  $\rho \in \mathfrak{a}^*$  is defined by the algebraic structure of  $\mathfrak{g}$ .

**Spherical transform**

For  $f$  on  $G$ , we define the spherical transform

$$\mathcal{H}(f)(\lambda) = \int_G f(x)\phi_\lambda(x^{-1})dx,$$

so that  $\mathcal{H}(f)$  is a function on  $\mathfrak{a}^*$ . It has many similarities to the classical Fourier transform: For  $f, g$  in the space  $C^\infty(K \backslash G/K)$  of  $K$ -biinvariant smooth functions on  $G$ , we have

- $\mathcal{H}(f * g) = \mathcal{H}(f)\mathcal{H}(g)$ ;
- $\|\mathcal{H}(f)\|_2 = \|f\|_2$ ;
- inversion formula:  $f(x) = \int_{\mathfrak{a}^*} \mathcal{H}(f)(\lambda)\phi_\lambda(x)|\mathbf{c}(\lambda)|^{-2}d\lambda$ ,

where  $\mathbf{c}$  is the important Harish-Chandra  $\mathbf{c}$ -function on  $G$ .

**Spherical functions and differential operators**

Every  $\phi_\lambda$  is an eigenfunction of all differential operators on  $C^\infty(G)$  invariant with respect to  $G$  on the left and  $K$  on the right. In particular, the structure of a Riemannian manifold on  $G/K$  gives rise to its Laplace-Beltrami operator  $\mathcal{L}$  which can be considered as an operator on  $C^\infty(G)$  in the above class. It is convenient to denote  $\mathcal{L}_\rho = \mathcal{L} + |\rho|^2$ , then

$$\mathcal{L}_\rho\phi_\lambda = |\lambda|^2\phi_\lambda.$$

For every function of  $\mathcal{L}_\rho$ , we have also  $F(\mathcal{L}_\rho)\phi_\lambda = F(|\lambda|^2)\phi_\lambda$ . Being left invariant, it is a convolution operator  $f \mapsto f * k_F$ , and the inversion formula for the spherical transform implies that

$$k_F(x) = \int_{\mathfrak{a}^*} F(|\lambda|^2)\phi_\lambda(x)|\mathbf{c}(\lambda)|^{-2}d\lambda.$$

Fourier analysis methods then apply, since this is just an integral over  $\mathfrak{a}^* \simeq \mathbb{R}^l$ . The complicated part is the spherical function, for which one seeks simpler approximations; it is known for example, that for “good”  $x = e^H \in A$  and  $\|H\| \rightarrow \infty$ ,

$$\phi_\lambda(e^H) \sim \sum_{k=1}^n \mathbf{c}(s_k \lambda) e^{(is_k \lambda - \rho)(H)},$$

where  $\{s_k\}$  form a certain group of reflections on  $\mathfrak{a}$ , called the Weyl group.

**Examples**

If  $G = SL_n(\mathbb{R})$ , then  $K = SO_n(\mathbb{R})$ ,  
 $A = \{ \text{diagonal matrices in } G \text{ with positive diagonal} \}$ ,  
 $N = \{ \text{upper-triangular matrices with diagonal } 1 \}$ , and  
 $G/K \simeq AN = \{ \text{upper-triangular matrices in } G \text{ with positive diagonal} \}$ .

The ‘ax+b’ group can be identified with  $G/K$  where  $G = SO(2, 1)_o$  is the connected component of the identity of  $SO(2, 1)$ , and  $K = SO(2)$ .

## 4 Traces and weights

A *weight* on a  $C^*$  or von Neumann algebra  $A$  is a map  $\tau : A^+ \rightarrow [0, +\infty]$  such that

- $\tau$  is additive
- $\tau(tx) = t\tau(x)$  for  $t \in \mathbb{R}_+$ ,  $x \in A^+$ , with  $0 \cdot (+\infty) = 0$ .  
 If in addition  $\tau(xx^*) = \tau(x^*x)$  for all  $x \in A$ , then  $\tau$  is a *trace*.

A trace  $\tau$  is *faithful* if  $\tau(x) = 0$  only for  $x = 0$ , and *semifinite* if for any  $x \in A^+$  there exists  $y \in A^+$  such that  $y \leq x$  and  $\tau(y) < \infty$ .

★ On  $A = B(H)$ : the usual trace  $Tr$

★ On  $A = C_0(G)$ :  $\tau(f) = \int_G f$ , so a weight can be viewed as a non-commutative integral

**Plancherel weight / trace:** On  $A = C_r^*(G)$  or  $VN(G)$ , we can set  $\tau(L_k) = k(e)$  if  $k = f^* * f$ , or equivalently

$$\tau(L_f^* L_f) = \tau(L_{f^* * f}) = \|f\|_2^2.$$

If  $G$  is unimodular, this is a trace. In general, just a weight.

We can see that the equality (3) is in fact a disintegration of  $\tau(L_f^* L_f)$  into  $\tau_\pi(\widehat{f}(\pi)^* \widehat{f}(\pi))$  where every  $\tau_\pi$  is the usual trace.

**Example:** Consider again the Heisenberg group  $\mathbb{H}$  and the sub-Laplacian  $\mathcal{L} = X^2 + Y^2$  on it. For any  $u \in \mathbb{R}$ , by (3)

$$\tau(I_{(0,u)}(\mathcal{L})) = \int_{\mathbb{R}} \tau_\lambda(I_{(0,u)}(\mathcal{L}_{\pi_\lambda})) |\lambda| d\lambda;$$

the operator  $\mathcal{L}_{\pi_\lambda}$  is given by (6) and its eigenvalues are known, so one can show [1] that

$$\tau(I_{(0,u)}(\mathcal{L})) = \int_{\mathbb{R}} \#\{k : s_{k,\lambda} < u\} |\lambda| d\lambda \lesssim u^{3/2}. \quad (7)$$

### **Non-commutative $L_p$ spaces**

Let  $M$  be a von Neumann algebra with a faithful semifinite normal trace  $\tau$  (normal means that  $\tau(\sup x_i) = \sup \tau(x_i)$  for every bounded increasing net in  $A^+$ ). For  $x \in M$ , define

$$\|x\|_p = \tau(|x|^p)^{1/p}$$

and  $L_p(M)$  as the completion of  $\{x \in M : \|x\|_p < \infty\}$ .

For  $p = \infty$ ,  $\|\cdot\|_\infty$  is the usual norm of  $M = L^\infty(M)$ .

★ If  $M = L^\infty(X)$  with  $\tau(f) = \int_X f d\tau$ , then  $L^p(M) = L^p(X)$ .

★ If  $M = \lambda(G)'' = VN(G)$ , then with the Plancherel weight, for  $f \in L^2(G)$

$$\|L_f\|_{L^2(M)} = \tau(L_f^* L_f)^{1/2} = \|f\|_2.$$

### **Weak $L_p$ spaces**

One can define *generalized singular numbers*  $\mu_t(x)$  for  $x \in M$ ,  $t \in \mathbb{R}_+$ :

$$\mu_t(x) = \inf\{\lambda > 0 : \tau(I_{(|\lambda|, +\infty)}(x)) < t\}$$

with the help of spectral projections of  $x$ , and set

$$\|x\|_{p,\infty} = \sup_{t>0} t^{1/p} \mu_t(x).$$

The obtained completed space  $L^{p,\infty}(M)$  is called the non-commutative weak  $L_p$  space, or Lorentz space.

For a decreasing function  $f \geq 0$  on  $\mathbb{R}_+$ , we get  $\mu_t(f) = f(t)$ . If  $x$  is a compact operator on a Hilbert space, then  $\mu_n(x) = \lambda_n(|x|)$  are the eigenvalues of  $|x| = (x^*x)^{1/2}$ .

### **Non-unimodular case**

If  $G$  is not unimodular, the Plancherel weight is not a trace. Non-commutative  $L^p$  spaces can be defined still, with complex interpolation available for them; but the Lorentz spaces not, it is an open problem.

There are three definitions of non-tracial  $L^p$  spaces, each of them with its advantages and disadvantages: by Haagerup (exposed by Terp [18]), Izumi [10] and Hilsum [9]. We will not go further here.

### **Quantum groups**

This notion generalizes both  $C_0(G)$  and  $C_r^*(G)$ , and provides a duality between them. An equivalent theory is built upon von Neumann algebras rather than  $C^*$ -algebras, and includes  $L^\infty(G)$  and  $VN(G)$ . Many new examples, not coming from groups, are included in this theory.

**Definition 1 (Kustermans, Vaes [13])** A *locally compact quantum group* (LCQG) is a von Neumann algebra  $M$  with a map  $\Delta : M \rightarrow M \otimes M$ , called comultiplication, and two weights  $\phi, \psi$  on  $M$  such that

- $\Delta$  is a unital normal  $*$ -homomorphism;
- $\Delta$  is coassociative:  $(1 \otimes \Delta)\Delta = (\Delta \otimes 1)\Delta$ ;
- $\phi, \psi$  are normal and semifinite;
- $\phi$  is left invariant and  $\psi$  is right invariant: for  $\omega \in M_*^+$  and  $x \in M^+$ , if  $\phi(x)$  or  $\psi(x)$  respectively is finite, then

$$\begin{aligned}\phi(\omega \otimes 1)\Delta(x) &= \phi(x)\omega(1), \\ \psi(1 \otimes \omega)\Delta(x) &= \psi(x)\omega(1).\end{aligned}$$

On  $M = L^\infty(G)$ , the weights are integrals with respect to the left and right Haar measures. On  $M = L^\infty(G)$ , they are Plancherel weights.

**Duality:** There exists a dual LCQG  $(\widehat{M}, \widehat{\Delta}, \widehat{\phi}, \widehat{\psi})$  such that  $\widehat{\widehat{M}} = M$ . The algebras  $L^\infty(G)$  and  $VN(G)$  are duals of each other. There are also analogues of the Fourier transform:  $*$ -homomorphisms with dense ranges

$$\lambda : M_* \rightarrow \widehat{M}, \quad \widehat{\lambda} : \widehat{M}_* \rightarrow M.$$

For  $M = L^\infty(G)$ , we have  $M_* = L^1(G)$  and  $\lambda : f \mapsto L_f \in VN(G)$ . The map  $\lambda$  is thus the left regular representation of  $L^1(G)$ . If  $G$  is abelian, then  $VN(G)$  is identified with  $L^\infty(\widehat{G})$  via the usual Fourier transform, with  $\widehat{L_f g} = \widehat{f * g} = \widehat{f}\widehat{g}$ , so modulo this identification  $L_f$  is the operator of multiplication by  $\widehat{f}$  and thus deserves its name as the Fourier transform of  $f$ .

For  $1 \leq p \leq 2$ , there are  $L_p$ -Fourier transforms (Cooney [3], Caspers [2]):  $\mathcal{F}_p$  maps  $L^p(M)$  to  $L^{p'}(\widehat{M})$ , with  $1/p + 1/p' = 1$ . In the case of  $M = L^\infty(G)$ , we have  $L^p(M) = L^p(G)$ , and  $\mathcal{F}_p : f \mapsto L_f \Delta^{1/p'} \in L^{p'}(VN(G))$ , where we need to multiply first by a power of the modular function.

$M$  is said of Kac type if  $\phi, \psi$  are traces. The example above shows that in the non-Kac case, in particular for a non-unimodular group,  $\mathcal{F}_p$  are different operators for different  $p$ , as opposed to the classical case where only their domains are different.

**Hausdorff-Young inequality**

If  $G$  is an abelian locally compact group, then for  $1 \leq p \leq 2$   $\|\widehat{f}\|_{p'} \leq C_p \|f\|_p$ . On quantum groups this is also true [2]:

$$\|\mathcal{F}_p(f)\|_{L^{p'}(\widehat{M})} \leq \|f\|_{L^p(M)},$$

with an equality for  $p = 2$ . In the Kac case, a stronger inequality holds [22], involving the weak norm  $\|f\|_{L^{p,p'}(M)}$ .

## 5 Multipliers

In the classical case, a multiplier with a symbol  $m$  is the operator  $T_m = \mathcal{F}^{-1}(m\widehat{f})$ . On a quantum group of Kac type, one can adopt the same approach: for  $m \in \widehat{M}$ , we set

$$T_m(y) = \mathcal{F}^{-1}(m\mathcal{F}(y)),$$

$y \in M$ . We assume from now that both  $M$  and  $\widehat{M}$  are of Kac type. Up to my knowledge, there is no theory of non-Kac multipliers, as those would require a coordination of  $\mathcal{F}_p$  transforms for different  $p$ . An exception is given by spherical multipliers on symmetric spaces: these groups are non-unimodular, but the multipliers in question are described by the spherical transform which reduces to a commutative convolution subalgebra of functions on  $G/K$ .

Say that  $T_m$  is an  $L^p - L^q$  Fourier multiplier if it extends to a bounded map from  $L^p(M)$  to  $L^q(M)$ .

**Theorem 4 (H. Zhang [22])** *For  $1 < p \leq 2 \leq q < \infty$  and  $r$  such that  $1/r = 1/p - 1/q$ , every  $m \in L^{r,\infty}(\widehat{M})$  is an  $L^p - L^q$  Fourier multiplier with*

$$\|T_m\|_{L^p(M) \rightarrow L^q(M)} \leq C_{p,q} \|m\|_{L^{r,\infty}(\widehat{M})}.$$

**Example:** The case of  $M = L^\infty(G)$  with a unimodular group  $G$  was considered before by Akylzhanov and Ruzhansky [1]. They give also the following example: on the Heisenberg group  $\mathbb{H}$  with the sub-Laplacian  $\mathcal{L} = X^2 + Y^2$ ,

$$\|(1 - \mathcal{L})^{-s}\|_{L^{r,\infty}(VN(\mathbb{H}))} \lesssim \sup_{u>0} u^{3/2r} (1 + u)^{-s},$$

which is finite as soon as  $s \geq \frac{3}{2r}$ , so that in these assumptions  $(1 - \mathcal{L})^{-s}$  is bounded from  $L^p(\mathbb{H})$  to  $L^q(\mathbb{H})$ . The estimate of the weak  $L^r$  norm is based on (7).

In the dual case of  $M = VN(G)$  on a unimodular group  $G$ , the symbol of a multiplier is a bounded function  $m : G \rightarrow \mathbb{C}$ . The map

$$T_m : L_f \mapsto L_{mf} \tag{8}$$

has the above form: here  $\mathcal{F} : L_f \mapsto f$  is the Fourier transform of  $M$ , which is the inverse to that of  $\widehat{M} = L^\infty(G)$ , so that  $L_{mf} = \mathcal{F}^{-1}(m\mathcal{F}(f))$ .

**Free group of  $n$  generators**  $G = \mathbb{F}_n$

Let  $|g|$  be the length of the reduced word  $g \in G$ , and

$$m_t(g) = e^{-t|g|}.$$

For positivity reasons,  $T_{m_t}$  is bounded on  $L^p(VN(G))$  for every  $p \in [1, +\infty]$ . We can calculate the weak  $L^r$  norm of  $m_t$ .

We have  $|m_t(g)| \geq \alpha$  iff  $|g| \leq -\frac{1}{t} \log \alpha = \frac{1}{t} |\log \alpha|$  since  $m_t < 1$ . The next formula is valid for  $0 < \alpha < 1$ , otherwise  $\lambda_m(\alpha) = 0$ :

$$\lambda_m(\alpha) = |\{g \in G : |m(g)| > \alpha\}| \leq (2n)^{t^{-1} |\log \alpha|}.$$

$$\|m_t\|_{L^{r,\infty}(L^\infty(G))} = \sup_{\alpha > 0} \alpha \lambda_m^{1/r}(\alpha) \leq \sup_{0 < \alpha < 1} \alpha^{1-(tr)^{-1} \log(2n)}$$

and this is finite iff  $1 - (tr)^{-1} \log(2n) \geq 0$ , that is, iff  $t \geq \frac{\log(2n)}{r}$ . Under this condition and  $p, q$  as in Theorem 4,  $T_m$  is an  $L^p - L^q$  multiplier.

Even more is true, on any discrete group (H. Zhang, [22]): if  $|\xi| \leq m_t$  and  $1 < p < \infty$ ,  $\frac{1}{p^*} = |\frac{1}{2} - \frac{1}{p}|$ , then  $\|T_\xi\|_{L^p(VN(G))} \lesssim \|\xi\|_{L^{p^*,\infty}(G)} \leq \|m_t\|_{L^{p^*,\infty}(G)}$ .

By different methods, another estimate was obtained by Junge, Palazuelos, Parcet, and Perrin [11]: For  $1 < p \leq q < \infty$ ,  $\|T_{m_t}\|_{L^p(VN(G)) \rightarrow L^q(VN(G))} \leq 1$  as soon as  $t > C \log \frac{q-1}{p-1}$ .

For  $L^p$  *multipliers*, a major result on  $\mathbb{R}^d$  is

**Theorem 5 (Hörmander–Mikhlin, 1956 – 1960)** *If  $m$  is  $\frac{d}{2} + 1$  times differentiable and with some  $C_m > 0$ , with usual notations for  $\alpha \in \mathbb{N}^d$ ,*

$$\max_{0 \leq |\alpha| \leq \lfloor \frac{d}{2} \rfloor + 1} \sup_{\xi \in \mathbb{R}^d} |\xi^{|\alpha|} \partial^\alpha m(\xi)| \leq C_m, \quad (9)$$

then  $m$  is an  $L^p(\mathbb{R}^d)$  multiplier for every  $1 < p < \infty$ .

Briefly, the estimate depends on a finite number of derivatives of  $m$  which should decrease polynomially. It is applicable in particular to  $e^{-|\xi|}$  or to  $(1 + |\xi|^2)^{-r}$  with  $r$  large enough. It turns out that similar assumptions have sense for multipliers on group algebras.

**Free group  $\mathbb{F}_\infty$  of a countable set of generators**

Let  $\{g_k\}_{k \in \mathbb{N}}$  be the generators of  $G$ . Let  $m$  be a function on  $\mathbb{R}^d$ , satisfying (9). For  $g = g_{i_1}^{k_1} \dots g_{i_n}^{k_n}$ , set

$$\tilde{m}(g) = \begin{cases} m(k_1, \dots, k_d) & \text{if } n \leq d; \\ m(k_1, \dots, k_n, 0, \dots, 0) & \text{if } n < d. \end{cases}$$

Then  $T_{\tilde{m}}$  is bounded on  $L^p(VN(\mathbb{F}_\infty))$  (Mei, Ricard, Xu [14]).

**Multipliers on the group algebra of  $SL_n(\mathbb{R})$**

For  $g \in SL_n(\mathbb{R})$ , set  $|g| := \max(\|g - e\|, \|g^{-1} - e\|)$ . Let  $X_1, \dots, X_{n^2-1}$  be the standard basis of

$$\mathfrak{sl}_n(\mathbb{R}) = \{A \in M_n(\mathbb{R}) : Tr A = 0\}.$$

For  $\gamma = (j_1, \dots, j_{|\gamma|})$  and a function  $m$  denote

$$d_g^\gamma m(g) = X_{j_1} \dots X_{j_{|\gamma|}} m.$$

**Theorem 6 (Parcet, Ricard, De la Salle)** [16] *If for all  $|\gamma| \leq \lfloor \frac{n^2}{2} \rfloor + 1$  and  $g \in G$*

$$|g|^{|\gamma|} |d_g^\gamma m(g)| \leq C_m,$$

*then  $T_m$  is completely bounded on  $L^p(VN(G))$  for  $1 < p < \infty$ .*

These examples close our brief survey.

## References

1. R. Akyzhanov, M. Ruzhansky.  $L_p - L_q$  multipliers on locally compact groups, *Journal of Functional Analysis* 278 (3), 108324, 2020.
2. M. Caspers. The  $L_p$ -Fourier transform on locally compact quantum groups. *J. Operator Theory*, 69(1):161–193, 2013.
3. T. Cooney. A Hausdorff-Young inequality for locally compact quantum groups. *Internat. J. Math.*, 21(12):1619–1632, 2010.
4. J. Dixmier,  $C^*$ -algebras, North-Holland, 1969.
5. V. Fischer, M. Ruzhansky, *Quantization on Nilpotent Lie Groups*, Birkhäuser, 2016.
6. G. Folland, *A course in Abstract Harmonic Analysis*, CRC Press, 1995.
7. J. Glimm, Type I  $C^*$ -algebras, *Annals of Math.* 73 no. 3 (1961), 572–612.
8. S. Helgason, *Groups and Geometric Analysis*, AMS, 1984.
9. M. Hilsaum, Les espaces  $L^p$  d'une algèbre de von Neumann définies par la dérivée spatiale, *J. Funct. Anal.* 40 (1981), 151–169.
10. H. Izumi, Constructions of non-commutative  $L^p$ -spaces with a complex parameter arising from modular actions, *Internat. J. Math.* 8 (1997), 1029–1066.
11. M. Junge, C. Palazuelos, J. Parcet, M. Perrin, Hypercontractivity in group von Neumann algebras, *American Mathematical Society* 249 (1183), 2014.
12. A. A. Kirillov, *Lectures on the Orbit Method*, AMS, 2004.
13. J. Kustermans, S. Vaes, Locally compact quantum groups, *Ann. Sci. Ecole Norm. Sup.* (4), 33(6):837–934, 2000.
14. T. Mei, E. Ricard, Q. Xu, A Mihklin multiplier theory for free groups and amalgamated free products of von Neumann algebras. *Adv. Math.*, 403 (2022), Paper No. 108394, 32 pp.
15. S. Neshveyev, L. Tuset. *Compact quantum groups and Their Representation Categories*, SMF, *Cours Spécialisés*, 2013.
16. J. Parcet, E. Ricard, M. de la Salle, Fourier multipliers in  $SL_n(\mathbb{R})$ , *Duke Math. J.* 171 (2022), 1235–1297.
17. M. Ruzhansky, V. Turunen, *Pseudo-Differential Operators and Symmetries*, Birkhäuser, 2010.
18. M. Terp,  $L^p$  spaces associated with von Neumann algebras. Notes, Report No. 3a + 3b, *Kobenhavns Universitets Matematiske Institut*, Juni 1981.
19. Th. Timmermann, *An Invitation to Quantum Groups and Duality*, EMS, 2008.
20. J. G. Wendel, Left centralizers and isomorphisms of group algebras, *Pacif. J. Math.* 2 (1952), 251–261.
21. H. Yoshizawa, Some remarks on unitary representations of the free group, *Osaka Math. J.* 3 no. 1 (1951), 55–63.
22. H. Zhang,  $L_p - L_q$  Fourier Multipliers on Locally Compact Quantum Groups, *J. Fourier Anal. Appl.* 29, 46 (2023).