

MINIMAL TETRAHEDRA AND AN ISOPERIMETRIC GAP THEOREM IN NON-POSITIVE CURVATURE

CORNELIA DRUȚU, URS LANG, PANOS PAPASOGLU, AND STEPHAN STADLER

ABSTRACT. We investigate isoperimetric inequalities for Lipschitz 2-spheres in $\text{CAT}(0)$ spaces, proving bounds on the volume of efficient null-homotopies. In one dimension lower, it is known that a quadratic inequality with a constant smaller than $\mathbf{c}_2 = 1/(4\pi)$ – the optimal constant for the Euclidean plane – implies that the underlying space is Gromov hyperbolic, and a linear inequality holds. We establish the first analogous gap theorem in higher dimensions: if a proper $\text{CAT}(0)$ space satisfies a Euclidean inequality for 2-spheres with a constant below the sharp threshold $\mathbf{c}_3 = 1/(6\sqrt{\pi})$, then the space also admits an inequality with an exponent arbitrarily close to 1. As a corollary we obtain a similar result for Lipschitz surfaces of higher genus. Towards our main theorem we prove a (non-sharp) Euclidean isoperimetric inequality for null-homotopies of 2-spheres, apparently missing in the literature. A novelty in our approach is the introduction of minimal tetrahedra, which we demonstrate satisfy a linear inequality.

1. INTRODUCTION

1.1. Background and main results. Isoperimetric filling inequalities in dimension $n + 1$ control the volume needed in order to fill an n -cycle S by an $(n + 1)$ -chain V . Their specific form is intimately related to the geometry of the underlying metric space. A central role is played by isoperimetric inequalities of the Euclidean type

$$\text{vol}_{n+1}(V) \leq \text{const} \cdot \text{vol}_n(S)^{1+1/n}.$$

By a fundamental result of Wenger [Wen05], which builds on earlier work of Gromov [Gro83], such inequalities hold in particular in all Banach spaces and $\text{CAT}(0)$ spaces. (See also [Sch20] for the optimal inequality for 2-cycles in Hadamard manifolds.) It is natural to expect that for $\text{CAT}(0)$ spaces, Euclidean isoperimetric inequalities continue to hold when the classes of admissible cycles and fillings are restricted to Lipschitz spheres and balls, respectively. However, to our knowledge, for $n > 1$ this is not recorded in the literature, and we will provide the result for $n = 2$ (with a non-optimal constant) in Theorem C below. In the case $n = 1$ – when circles are filled by discs – much more is known. It has recently been shown that non-positive curvature is equivalent to a Euclidean isoperimetric inequality with the sharp constant $\mathbf{c}_2 := 1/(4\pi)$ [LW18b, SW24]. Moreover,

Date: February 6, 2025.

a length space that admits a quadratic isoperimetric inequality for curves with a constant strictly smaller than \mathbf{c}_2 is necessarily Gromov hyperbolic [Wen08a, LWY20, SW24]. These sharp results were predated by the general observation, originally due to Gromov [Gro87], that length spaces with a subquadratic isoperimetric inequality must in fact satisfy a linear isoperimetric inequality [Ol'91, Pap95a, Pap95b, Bow95, Dru01].

The main result of this paper is a first result of this kind in higher dimension: a sharp isoperimetric gap theorem for fillings of 2-spheres by 3-balls in CAT(0) spaces.

Theorem A. *For a proper CAT(0) space X , the following are equivalent:*

- (1) *There exists a constant $c < \mathbf{c}_3 := 1/(6\sqrt{\pi})$ such that every Lipschitz 2-sphere $\hat{S} \subset X$ of large area admits a filling by a Lipschitz 3-ball $\hat{B} \subset X$ with volume*

$$\text{vol}(\hat{B}) \leq c \cdot \text{area}(\hat{S})^{3/2}.$$

- (2) *For every $\delta > 0$ there exists a constant $C = C(\delta)$ such that every Lipschitz 2-sphere $S \subset X$ extends to a Lipschitz 3-ball $B \subset X$ with volume*

$$\text{vol}(B) \leq C \cdot \text{area}(S)^{1+\delta}.$$

- (3) *The asymptotic rank of X is at most 2.*

The notion of *asymptotic rank* appearing in the last item stems from Gromov's discussion of a variety of related invariants in [Gro93, §6.B₂], and was further investigated in [Kle99, Wen06a, Wen11, Des16, KL20, GL23, LS24]. For proper CAT(0) spaces with cocompact isometry group, the asymptotic rank equals the maximal n such that X contains an n -flat, that is, an isometric copy of \mathbb{R}^n . In particular, the isoperimetric inequality in the second item of Theorem A holds for the universal cover X of any compact manifold of non-positive curvature provided that X contains no 3-flat. For a general CAT(0) space X , the asymptotic rank is at most n if and only if no asymptotic cone of X contains an $(n+1)$ -flat, and it is at most 1 if and only if X is Gromov hyperbolic. We refer to the beginning of Section 7 for the actual definition in the general case. Notice that the asymptotic rank is a quasi-isometry invariant.

Gromov conjectured that proper cocompact CAT(0) spaces of asymptotic rank at most n admit linear isoperimetric inequalities

$$\text{vol}_{n+1}(V) \leq \text{const} \cdot \text{vol}_n(S)$$

for fillings of n -cycles by $(n+1)$ -chains (compare [Gro93, §6.B₂, p. 128, (b)]). Apart from the (hyperbolic) rank one case (see [Lan00] for $n > 1$), this is presently known for symmetric spaces or homogeneous Hadamard manifolds (see [Gro93, §5.D, (5)(b'₁)] and [Leu14, Isl23]). For general CAT(0) spaces of asymptotic rank at most n , where $n \geq 2$, the best known result in this direction is Wenger's sub-Euclidean inequality [Wen11], stating that every n -cycle of mass s admits a filling with mass at most $o(s^{1+1/n})$ as $s \rightarrow \infty$.

In particular, since the isoperimetric inequalities in dimensions up to the asymptotic rank are at best Euclidean, the isoperimetric profiles detect the asymptotic rank. Furthermore, a family of linear isoperimetric inequalities for n -cycles with bounded density quotients was deduced in [GL23]. The result is restated in Theorem 2.9 below and provides in addition a bound on the filling radius, which will enter the proof of Theorem A. These inequalities, as well as those in [Wen05, Wen11], actually refer to the chain complex of metric integral currents in the sense of Ambrosio–Kirchheim, which comprises the Lipschitz singular chains and has favorable compactness properties (see Section 2.6 for further information). By contrast, Theorem A concerns Lipschitz mappings and null-homotopies of spheres, the use of integral currents will thus be limited to the proofs of two intermediate results.

In the terminology of [Gro99a, Section 6.D], the second assertion in Theorem A is referred to as an isoperimetric inequality of *infinite rank*. Inequalities of this type are also discussed in [Pap09, Wen11]. A priori, this assertion is weaker than the conjectured linear bound (possibly $C = C(\delta) \rightarrow \infty$ as $\delta \rightarrow 0$), and proving the latter, if correct, seems to require a strategy different from ours. However, a linear inequality will be obtained for so-called minimal tetrahedra in Theorem D below, which constitutes a crucial ingredient of Theorem A. Currently we do not know how to adapt Theorem A to homological fillings of general 2-cycles, as the decomposition argument used in the proof requires good topological control. Yet, we obtain the following corollary via Gromov’s systolic inequality for closed surfaces.

Corollary B. *Let X be a proper CAT(0) space of asymptotic rank at most 2. For every $\delta > 0$ and every integer $g \geq 0$ there exists a constant $C_g = C_g(\delta)$ such that every closed Lipschitz surface $\Sigma \subset X$ of genus g extends to a Lipschitz handlebody $H \subset X$ with volume*

$$\text{vol}(H) \leq C_g \cdot \text{area}(\Sigma)^{1+\delta}.$$

We would like to point out the formal similarity between our results and another family of filling problems in quantitative topology, originally raised by Gromov [Gro99b]. Specifically, given a null-cobordant closed Riemannian manifold M of bounded geometry (say, with sectional curvature $|K| \leq 1$ and injectivity radius at least 1), one wants to minimize the volume of compact Riemannian manifolds W of bounded geometry with $\partial W = M$. An upper bound of order $O(\text{vol}(M)^{1+\delta})$ for every $\delta > 0$ was established in [CDMW18], and an analogous result in the PL category has recently been shown in [MW23]. Notably, as in our case, Gromov originally suggested a linear inequality.

1.2. Strategy of proof. We now outline the proof of Theorem A. It clearly suffices to prove the implications (1) \Rightarrow (3) and (3) \Rightarrow (2). We show that the first holds in all dimensions, with the sharp constant \mathbf{c}_{n+1} , for Lipschitz n -spheres and asymptotic rank at most n (see Proposition 7.1). The most novel and challenging part is the implication (3) \Rightarrow (2). Let us refer to the

inequality in (2) as a δ -isoperimetric inequality. Thus, if the asymptotic rank of a proper CAT(0) space is at most 2, then we want to show a δ -isoperimetric inequality for all $\delta > 0$.

The first ingredient is the case $\delta = 1/2$ which corresponds to the Euclidean isoperimetric inequality for 2-spheres. The result holds for general CAT(0) spaces, regardless of the asymptotic rank (see Theorem 3.2):

Theorem C. *Let X be a CAT(0) space. Then there exists a constant $C > 0$ such that every Lipschitz 2-sphere $S \subset X$ bounds a Lipschitz 3-ball $B \subset X$ with volume*

$$\text{vol}(B) \leq C \cdot \text{area}(S)^{3/2}.$$

The argument uses Wenger's approach from [Wen08b] for general integral n -cycles and homological fillings, and the restriction to 2-spheres allows us to control the topology of the filling. The proof works more generally for spaces that admit certain *Lipschitz coning inequalities*¹. The latter hold, for instance, if X is a 2-connected Riemannian manifold with a geometric action of a combable group; see [ECH⁺92, Theorem 10.3.6]. However, it remains unclear how to generalize Theorem C to higher dimensions.

Returning to the proof of Theorem A, observe that the Euclidean isoperimetric inequality provides the desired implication for all 2-spheres up to a certain area, depending on δ . Furthermore, this threshold can be made arbitrarily large by increasing the constant in the δ -isoperimetric inequality. The proof is indirect. Assuming that the conclusion fails for some $\delta \in (0, 1/2)$, we would like to select a Lipschitz 2-sphere with minimal area among all spheres that violate the δ -isoperimetric inequality for a suitable large constant. Due to the lack of compactness, we need to work with a sphere $f: S^2 \rightarrow X$ of nearly minimal area instead.

These assumptions allow us to prove that f satisfies an *intrinsic quadratic isoperimetric inequality* for subdiscs of sufficiently large area. We then combine this with an elaborate decomposition process to subdivide the sphere, after a small deformation, into a controlled number of relatively large pieces with effective bounds on their circumference and area. This is a crucial point in our argument. As Gromov remarks in [Gro93, §5.D, (5)(a)], the main difficulty in extending the iterative scheme for the filling of circles (used, for example, in [Gro93, §5.A''₃] or [Bow95]) to higher dimensions is the lack of nice canonical subdivisions of n -dimensional manifolds for $n \geq 2$.

The next step in the proof is to deform f into a Lipschitz sphere \bar{f} that is *piecewise minimal*, whereby we use the δ -isoperimetric inequality for spheres of small area to bound the volume of the homotopy. Intuitively, one may think of \bar{f} as being piecewise geodesic on the 1-skeleton of a triangulation of S^2 and sending 2-simplices to minimal discs. After coning off the 0-skeleton and inserting further minimal discs into the geodesic triangles thus obtained,

¹There is $C > 0$ such that for $n = 1, 2$, every Lipschitz n -sphere S in an r -ball bounds a Lipschitz $(n + 1)$ -ball B with $\text{vol}_{n+1}(B) \leq Cr \text{vol}_n(S)$.

it remains to fill a controlled number of *minimal tetrahedra*. This is accomplished by the following linear isoperimetric inequality (see Theorem 4.3). Altogether, we obtain a filling contradicting the assumption on f , which finishes the proof of Theorem A.

Theorem D. *Let X be a proper CAT(0) space of asymptotic rank at most 2. There exists a constant $\mu > 0$ such that each minimal tetrahedron $\tau \subset X$ satisfies a linear isoperimetric inequality for fillings by balls,*

$$\text{Fillvol}(\tau) \leq \mu \cdot \text{area}(\tau).$$

This result represents the key novel idea of this paper. It crucially exploits properties of minimal discs established in [Sta21], the aforementioned bound on the filling radius from [GL23], and an idea of White [Whi84]. As a by-product of the above strategy, the converse holds as well: a linear isoperimetric for minimal tetrahedra implies that the asymptotic rank is at most 2.

1.3. Organization. In Section 2 we introduce our notation and recall some basic facts from metric geometry. We then collect relevant background from the theory of minimal surfaces in metric spaces, the theory of metric integral currents, and quantitative dimension theory. Section 3 contains the proof of Theorem C. In Section 4, we introduce *minimal tetrahedra* – Lipschitz 2-spheres composed of four minimal discs filling the four geodesic triangles spanned by a quadruple of points – and give the proof of Theorem D. Section 5 discusses some auxiliary quantitative triangulation results for (pseudo-)metric discs. Section 6 is devoted to the construction of certain homotopies of spheres that improve their intrinsic structure. In the final Section 7, we complete the proof of Theorem A and derive Corollary B.

1.4. Acknowledgements. The first author is grateful to the Max Planck Institute for Mathematics in Bonn for its hospitality during several research visits, while part of the work at this paper was carried out. The second author was partially supported by Swiss National Science Foundation Grant 197090. The last author was partially supported by DFG grant SPP 2026.

2. PRELIMINARIES

2.1. Basics. In the Euclidean space \mathbb{R}^n , we denote the open unit ball by U^n , the closed unit ball by B^n , and its boundary, the unit sphere, by S^{n-1} . The Lebesgue measure of B^n is denoted by ω_n . A *disc* D is a topological space homeomorphic to B^2 .

We denote the distance in a metric space X by d . The closed ball in X of radius r and center $x \in X$ is denoted by

$$B(x, r) = \{y \in X : d(x, y) \leq r\}.$$

If closed balls in X are compact, then X is called *proper*. For compact subsets $A, A' \subset X$ we denote by $|A, A'|_H$ their Hausdorff distance.

For $n \geq 0$, the n -dimensional Hausdorff measure on X is denoted by \mathcal{H}^n . The normalizing constant is chosen so that on the Euclidean space \mathbb{R}^n the Hausdorff measure \mathcal{H}^n equals the Lebesgue measure.

A continuous curve in X is called *rectifiable* if it has finite length. An *arc* is a homeomorphism from a compact interval to a subset of X . A *geodesic* is an isometric map from an interval into X . We refer to the images of the corresponding maps by the same names. A *Jordan curve* Γ in a metric space X is a closed subset homeomorphic to S^1 .

The space X is called a *geodesic space* if any pair of points in X is connected by a geodesic. A $\text{CAT}(0)$ *space* X is a complete geodesic space such that distances on a geodesic triangle in X are bounded from above by the corresponding distances on its comparison triangle in \mathbb{R}^2 . We assume some familiarity with $\text{CAT}(0)$ geometry and refer the reader to [BGS85, Bal95, BH99] for more information.

2.2. Lipschitz maps, area and volume. Let $U \subset \mathbb{R}^n$ be an open set, and let X be a metric space. Every Lipschitz map $f: U \rightarrow X$ has a *metric differential* at almost every point $x \in U$ in the following sense. There exists a unique semi-norm on \mathbb{R}^n , denoted $|df_x|$, such that

$$\lim_{x' \rightarrow x} \frac{d(f(x'), f(x)) - |df_x|(x' - x)}{\|x' - x\|} = 0.$$

The map $x \mapsto |df_x|$ into the space of semi-norms has a Borel measurable representative. The *Jacobian* $\text{Jac}(s)$ of a semi-norm s on \mathbb{R}^n is the Hausdorff n -measure in (\mathbb{R}^n, s) of the Euclidean unit cube if s is a norm, and $\text{Jac}(s) = 0$ otherwise. The *volume* of a Lipschitz map $f: U \rightarrow X$ is given by the *area formula*

$$\text{vol}_n(f) := \int_U \text{Jac}(|df_x|) dx = \int_X \#(f^{-1}\{y\}) d\mathcal{H}^n(y).$$

See [Kir94, KS93, LW17]. If X is a $\text{CAT}(0)$ space, then for almost every $x \in U$ where the metric differential is non-degenerate, the norm $|df_x|$ is induced by an inner product on \mathbb{R}^n (see [Wen06b, Lemma 4.3]). If $n = 1$ or $n = 2$, we will speak of *length* or *area* instead of volume and write $\text{length}(f)$ or $\text{area}(f)$, respectively. The volume of a Lipschitz map from a relatively open subset U of S^n is defined similarly.

A Lipschitz map from an n -sphere or $(n + 1)$ -ball to X will be called a *Lipschitz n -sphere* or *Lipschitz $(n + 1)$ -ball*, respectively. For convenience, we will sometimes also refer to their images in X as Lipschitz n -spheres or Lipschitz $(n + 1)$ -balls, often denoted by $S \subset X$ and $B \subset X$, respectively. The volumes $\text{vol}_n(S)$ and $\text{vol}_{n+1}(B)$ are then understood to be the volumes of the corresponding maps or, equivalently, as the Hausdorff measures counted with multiplicities. If the dimension is clear from the context, we omit the subscript and simply write $\text{vol}(S)$ and $\text{vol}(B)$.

We say that a Lipschitz ball $\bar{f}: B^{n+1} \rightarrow X$ *fills* the Lipschitz sphere $f := \bar{f}|_{S^n}$, or that S *bounds* B where $S = f(S^n)$ and $B = \bar{f}(B^{n+1})$. For a

given Lipschitz sphere $f: S^n \rightarrow X$, we define its *filling volume* by

$$\text{Fillvol}(f) := \inf\{\text{vol}(\bar{f}) \mid \bar{f}: B^{n+1} \rightarrow X \text{ a Lipschitz ball filling } f\}.$$

If no such \bar{f} exists, then $\text{Fillvol}(f) = \infty$.

In order to build Lipschitz balls of small volume that fill a given Lipschitz sphere, it is useful to decompose the sphere. To do this, we need some control on the fibers of Lipschitz functions. The following result, which will be used repeatedly, is a slight variation of [ABC13, Theorem 2.5].

Proposition 2.1. *Let $f: B^2 \rightarrow \mathbb{R}$ be a non-constant Lipschitz function. We denote its fibers $f^{-1}(t)$ by Π_t . Then for almost every value $t \in \mathbb{R}$ the following holds.*

- (1) $\mathcal{H}^1(\Pi_t)$ is finite;
- (2) The differential of f is non-trivial at \mathcal{H}^1 -almost every point in Π_t ;
- (3) Π_t decomposes as a disjoint union of rectifiable Jordan curves, rectifiable arcs with boundary points on S^1 , and an \mathcal{H}^1 -negligible set.

Proof. Extend f to a Lipschitz function $F: \mathbb{R}^2 \rightarrow \mathbb{R}$ with compact support. Now [ABC13, Theorem 2.5] shows that, for almost every t , the fiber $F^{-1}(t)$ has finite \mathcal{H}^1 measure, every connected component Γ of $F^{-1}(t)$ with $\mathcal{H}^1(\Gamma) > 0$ is a Jordan curve, and the union of the remaining connected components has \mathcal{H}^1 measure zero. Furthermore, every such Jordan curve Γ admits a unit speed parametrization $s \mapsto \gamma(s)$ such that $(\nabla F(\gamma(s)), \dot{\gamma}(s))$ is a positively oriented basis of \mathbb{R}^2 for almost every s . By the coarea formula, t can be chosen in addition such that $F^{-1}(t) \cap S^1$ is a finite set. Then every Jordan curve $\Gamma \subset F^{-1}(t)$ is either contained in B^2 or intersects B^2 in finitely many arcs or points. \square

The above statement holds true in similar form for maps defined on S^2 . In this case, the fibers generically decompose into a disjoint union of Jordan curves and an \mathcal{H}^1 -negligible set, no arcs appear.

2.3. Plateau problem and minimal discs. Recall that U^2 denotes the open unit disc in \mathbb{R}^2 . Let X be a CAT(0) space and $\Gamma \subset X$ a closed rectifiable curve. The Plateau problem for Γ asks to find a disc of minimal area spanning Γ . For Jordan curves it was solved in every complete metric space by Lytchak-Wenger [LW17] and the result was extended to general curves by Creutz [Cre22]. For a Lipschitz parametrization $\gamma: S^1 \rightarrow \Gamma$, consider the family of competitors

$$\Lambda(\gamma, X) := \{u \in W^{1,2}(U^2, X) \mid u \text{ spans } \gamma\},$$

where $W^{1,2}(U^2, X)$ denotes the Sobolev space of finite energy maps, and an element u is said to *span* γ if its trace $\text{tr}(u) \in L^2(S^1, \Gamma)$ has a continuous representative homotopic to γ . We will not recall the relevant definitions from Sobolev theory with values in metric spaces here, but refer the reader to [LW17] for details. For our purposes, it is enough to know that every finite energy map has a well-defined area, and every Lipschitz map $v: B^2 \rightarrow X$

restricts to an element in $W^{1,2}(U^2, X)$; its trace is simply $v|_{S^1}$ and its area is $\text{area}(v)$ as defined above. Plateau's problem then asks to find an area minimizer $v \in \Lambda(\gamma, X)$, meaning

$$\text{area}(v) = \inf\{\text{area}(u) \mid u \in \Lambda(\gamma, X)\}.$$

The following solution is well-known to experts and can be assembled from existing results in the literature. The key point is that it yields a Lipschitz map on the closed disc, whereas the (quasi-conformal) solution to the Plateau problem from [LW17, Cre22] is only locally Lipschitz in the interior.

Recall that a *disc retract* is a Peano continuum homeomorphic to a non-separating subset of the plane. A CAT(0) *disc retract* is a CAT(0) space homeomorphic to a disc retract.

Theorem 2.2. *Let X be a proper CAT(0) space and $\Gamma \subset X$ a closed rectifiable curve. Then there exists a Lipschitz disc $v: B^2 \rightarrow X$ that solves the Plateau problem for Γ . Moreover, v satisfies the following properties.*

- (monotonicity of density ratios)

$$\pi \leq \frac{\text{area}(v(B^2) \cap B(x, r))}{r^2} \leq \frac{\text{area}(v(B^2) \cap B(x, s))}{s^2}$$

for $x \in v(B^2)$ and $0 < r \leq s < d(x, v(S^1))$.

- v admits a factorization as $v = v' \circ v''$ with $v'': B^2 \rightarrow Z$ and $v': Z \rightarrow X$, where the space Z is a CAT(0) disc retract and v'' is a surjective Lipschitz map with connected fibers. Furthermore, v' is area minimizing among Lipschitz maps from Z and preserves the length of every rectifiable curve.

Recall that the intrinsic metric space Z_u associated to a map $u: B^2 \rightarrow X$ is the quotient metric space obtained from the pseudo-metric d_u on B^2 defined by

$$d_u(z_1, z_2) = \inf\{\text{length}(u \circ \gamma) \mid \gamma \text{ a path in } B^2 \text{ joining } z_1 \text{ and } z_2\}.$$

Proof. By [Cre22, Theorem 4.3] there exists a solution to the Plateau problem $u \in \Lambda(\gamma, X)$ that is locally $\frac{1}{3}$ -Hölder continuous and extends to a $\frac{1}{27}$ -Hölder continuous map on B^2 which we will still denote by u . Since u is harmonic, it is locally Lipschitz in U^2 by [KS93]. Moreover, the associated intrinsic metric space Z_u is CAT(0), the natural projection $\pi_u: B^2 \rightarrow Z_u$ has connected fibers, and Z_u is a disc retract [LW24, PS19]. By [LW18a, Corollary 3.2], we have $\pi_u \in W^{1,2}(U^2, Z_u)$ and the approximate metric differentials of u and π_u coincide almost everywhere. In particular, $\text{area}_u(W) = \text{area}_{\pi_u}(W)$ for every Borel set $W \subset U^2$. Let $c: S^1 \rightarrow \partial Z_u$ be a Lipschitz parametrization representing $\text{tr}(\pi_u)$. The proof of [Sta21, Lemma 30] provides a surjective Lipschitz map $\varphi: B^2 \rightarrow Z_u$ with connected fibers that extends c . Then

$$\text{area}(\varphi|_W) = \mathcal{H}^2(\varphi(W))$$

for every Borel set $W \subset B^2$ by the aforementioned area formula. Set $v = \bar{u} \circ \varphi$, where $\bar{u}: Z_u \rightarrow X$ is the unique 1-Lipschitz map such that $u = \bar{u} \circ \pi_u$. Then, for any ball $B(x, r) \subset X$,

$$\begin{aligned} \text{area}(u(B^2) \cap B(x, r)) &= \text{area}(\bar{u}(Z_u) \cap B(x, r)) = \mathcal{H}^2(\bar{u}^{-1}(B(x, r))) \\ &= \text{area}_\varphi(v^{-1}(B(x, r))) = \text{area}_v(v^{-1}(B(x, r))) = \text{area}(v(B^2) \cap B(x, r)). \end{aligned}$$

In particular, v satisfies the required monotonicity formula since u does [Sta21, Corollary 63]. \square

We call a Lipschitz disc as above a *minimal disc*.

2.4. Disc decompositions. Let $\Sigma \subset \mathbb{R}^2$ be a compact connected planar surface with $k+1 \geq 1$ boundary components. Note that $\mathbb{R}^2 \setminus \Sigma$ has precisely one unbounded component and Σ has one *outer boundary circle* σ_0 and k *inner boundary circles* $\sigma_1, \dots, \sigma_k$. If k is positive, then we call such a surface a *disc with k holes*. We call it a *polygonal disc with k holes* if in addition every boundary circle is a piecewise geodesic Jordan curve.

The following provides a weak surrogate for a triangulation. Given a disc with holes Σ , we define a *disc decomposition* of Σ by means of a connected finite Lipschitz graph $\Gamma \subset \Sigma$ with $\partial\Sigma \subset \Gamma$ such that each vertex has valence at least three, adjacent vertices are connected by a unique edge, and the closure of each component of $\Sigma \setminus \Gamma$ is a disc. In particular, we have a covering by discs D_i :

$$\Sigma = D_1 \cup \dots \cup D_n.$$

We call a disc decomposition *polygonal* if the edges of Γ are piecewise geodesic. Similarly, we define (polygonal) disc decompositions of 2-spheres.

Recall that a map between topological spaces is called *monotone* if it has connected fibers.

Lemma 2.3. *Let $D \subset \mathbb{R}^2$ be a closed disc with rectifiable boundary. Then every Lipschitz parametrization $\gamma: S^1 \rightarrow \partial D$ extends to a surjective monotone Lipschitz map $\varphi: B^2 \rightarrow D$.*

Proof. Denote by Y the interior of D equipped with the induced path metric. Since ∂D is rectifiable, any pair of points in Y can be joined by a polygon of length at most half the length of ∂D . Thus, the completion \bar{Y} is compact and therefore a CAT(0) disc [LW18b, Proposition 12.1, Lemma 12.2]. Denote by $f: \bar{Y} \rightarrow D$ the natural surjective 1-Lipschitz map. Since $\bar{Y} \setminus \partial\bar{Y}$ is a length space [LW18b, Theorem 1.3], and f is a local isometry in the interior, f preserves the length of every rectifiable curve in \bar{Y} . Since f restricts to a homeomorphism between the interiors, f must be a homeomorphism. Thus, $f^{-1} \circ \gamma$ is a Lipschitz parametrization of $\partial\bar{Y}$. By [Sta21, Lemma 30], $f^{-1} \circ \gamma$ extends to a monotone surjective Lipschitz map $\psi: B^2 \rightarrow \bar{Y}$. Thus, the composition $\varphi = f \circ \psi$ has the required properties. \square

The following proposition will often be used to reparametrize Lipschitz spheres.

Proposition 2.4. *Let $\Gamma \subset S^2$ be a finite Lipschitz graph. Then, for every $\epsilon > 0$, there exists a graph $P \subset S^2$ and a monotone Lipschitz map $F: S^2 \rightarrow S^2$ with the following properties.*

- (1) *The edges of P are piecewise geodesics;*
- (2) *P is ϵ -Hausdorff close to Γ ;*
- (3) *F restricts to a homeomorphism $P \rightarrow \Gamma$;*
- (4) *F is uniformly ϵ -close to the identity.*

Proof. After possibly inserting finitely many Lipschitz arcs, we may assume that Γ is connected and the closure of every component of $S^2 \setminus \Gamma$ is a closed disc of diameter at most $\frac{\epsilon}{3}$. To define P , start with Γ and replace every edge e by a δ -close piecewise geodesic e' , for some $\delta \ll \epsilon$, such that the new edges only intersect in the original vertices. Then the boundary of every component V of $S^2 \setminus P$ is a Jordan curve consisting of finitely many geodesic segments. By [Tuk80], the closure \bar{V} is a bilipschitz disc. Its diameter is at most $\frac{\epsilon}{3} + 2\delta$.

To define F , we first send every vertex of Γ to itself. On the edges of P , we choose for every edge e' a bilipschitz map to the closed unit interval and compose it with a constant speed parametrization of the corresponding edge e of Γ . Now we use Lemma 2.3 to find monotone Lipschitz extensions of F to the components of $S^2 \setminus P$. By construction, F is monotone Lipschitz and sends P homeomorphically to Γ . Moreover, it maps every component of $S^2 \setminus P$ to the corresponding component of $S^2 \setminus \Gamma$. Therefore $d(F(x), x) < \epsilon$ for all points $x \in S^2$. \square

For $\epsilon \in (0, 1)$ define $\varrho_\epsilon: B^2 \rightarrow B^2$ by

$$\varrho_\epsilon(x) = \begin{cases} \frac{1}{1-\epsilon} \cdot x & \text{if } \|x\| \leq 1 - \epsilon, \\ \frac{x}{\|x\|} & \text{else.} \end{cases}$$

Let D be a bilipschitz disc. Then we call a map $\psi_\epsilon: D \rightarrow D$ an ϵ -squeeze if $\psi_\epsilon = \beta \circ \varrho_\epsilon \circ \beta^{-1}$ for some bilipschitz map $\beta: B^2 \rightarrow D$. Note that in this case, if $f: D \rightarrow X$ a Lipschitz map to a metric space X , then the pseudo-metric spaces (D, d_f) and $(D, d_{f \circ \psi_\epsilon})$ are naturally isometric. Moreover, if γ is a path in D with $\partial\gamma \subset \partial D$, then there exists a path $\tilde{\gamma}$ in D with $\partial\tilde{\gamma} \subset \partial D$ and such that ψ_ϵ maps $\tilde{\gamma}$ monotonically onto γ .

Lemma 2.5. *Let $\Sigma \subset S^2$ be a disc with $k \geq 1$ holes and rectifiable boundary. Let $f: \Sigma \rightarrow X$ be a Lipschitz map. Let $s > 0$ be such that $d_f(x, \sigma_0) < s$ for every $x \in \sigma_1 \cup \dots \cup \sigma_k$. Then, for every $\epsilon > 0$ there exist a disc with k holes $\Sigma' \subset S^2$ and piecewise geodesic boundary components $\sigma'_0, \dots, \sigma'_k$, a surjective monotone Lipschitz map $\psi: S^2 \rightarrow S^2$ that sends Σ' onto Σ , and k pairs of piecewise geodesic paths $\beta_j^\pm \subset \Sigma'$ with the following properties.*

- (1) $|\Sigma', \Sigma|_H < \epsilon$ and ψ is ϵ -close to the identity;
- (2) $\text{length}_{f \circ \psi}(\sigma'_i) = \text{length}_f(\sigma_i)$, $i = 0, \dots, k$;
- (3) $d_{f \circ \psi}(x', \sigma'_0) < s$ for every $x' \in \sigma'_1 \cup \dots \cup \sigma'_k$;
- (4) $\text{length}_{f \circ \psi}(\beta_j^\pm) < s$, $j = 1, \dots, k$;

- (5) $|\beta_j^-, \beta_j^+|_H < \epsilon$ and $\beta_j^- \cap \beta_j^+ = \emptyset$, $j = 1, \dots, k$;
 (6) β_j^\pm joins σ_j' to $\sigma_0' \cup \Lambda_{j-1}$ and $\beta_j^\pm \cap (\sigma_0' \cup \Lambda_{j-1} \cup \sigma_j') = \partial\beta_j^\pm$ where $\Lambda_l = \bigcup_{i=1}^l (\beta_i^- \cup \beta_i^+)$ for $1 \leq l \leq k$.

In particular, the 1-rectifiable set $\Gamma' = \partial\Sigma' \cup \Lambda_k$ carries a natural graph structure and induces a polygonal disc decomposition of Σ' .




FIGURE 1. A polygonal resolution of a disc with two holes.

Proof. We prove the statement by induction on k . For $k = 1$, choose an arc $\gamma \subset \Sigma$ joining σ_1 to σ_0 with $\gamma \cap \partial\Sigma = \partial\gamma$ and $\text{length}_f(\gamma) < s$. For $\epsilon > 0$, apply Proposition 2.4 to the finite Lipschitz graph $\Gamma_1 = \partial\Sigma \cup \gamma$ to obtain an ϵ -close piecewise geodesic graph P_1 and a map $F: S^2 \rightarrow S^2$ ϵ -close to the identity. We define β^- to be the edge of P_1 corresponding to γ , and Σ' to be the annulus with boundary $P_1 \setminus \beta^-$. We then add an auxiliary piecewise geodesic $\alpha \subset \Sigma'$ to P_1 such that the closures of the components of $\Sigma \setminus (P_1 \cup \alpha)$ are bilipschitz discs. Next, we choose an ϵ -squeeze on each of those discs to define a monotone Lipschitz map $\psi': \Sigma' \rightarrow \Sigma'$. We then find a piecewise geodesic β^+ , disjoint, but ϵ -Hausdorff close to β^- , that is mapped monotonically onto an arc in $N_\epsilon(\beta^-) \cap P_1$ by ψ' . We obtain the induction base by setting $\psi = F \circ \psi'$. The induction step is similar. We ignore the boundary circle σ_k and apply the induction hypothesis to the disc with $(k - 1)$ holes. So we may assume that Σ has piecewise geodesic boundary and we have already found $k - 1$ pairs of piecewise geodesic paths $\beta_j^\pm \subset \Sigma$ with the right properties. To proceed, we choose an arc γ with $\text{length}_f(\gamma) < s$ joining σ_k to $\sigma_0 \cup \Lambda_{k-1}$ where $\Lambda_l = \bigcup_{i=1}^l (\beta_i^- \cup \beta_i^+)$ for $1 \leq l \leq k$. As above, we apply Proposition 2.4 to replace γ by a piecewise geodesic β_k^- and then use ϵ -squeezes to produce β_k^+ . \square

2.5. Nagata dimension. We recall the definition of the Nagata dimension (see [LS05]). A family $\mathcal{B} = \{B_i\}_{i \in I}$ of subsets of a metric space X is called

D -bounded if every B_i has diameter at most D . We say that \mathcal{B} has s -multiplicity at most m if every set of diameter $\leq s$ in X meets no more than m members of the family. The Nagata dimension of X is at most n if and only if there exists a constant c such that for every $s > 0$, X admits a cs -bounded covering with s -multiplicity at most $n + 1$.

If $\{B_i\}_{i \in I}$ is a D -bounded covering of a subset $S \subset X$ with s -multiplicity at most $n + 1$, and if $\epsilon \in (0, s/2)$, then the corresponding family of the open ϵ -neighborhoods $U_\epsilon(B_i)$ forms a $(D + 2\epsilon)$ -bounded cover of $U_\epsilon(S)$ with $(s - 2\epsilon)$ -multiplicity at most $n + 1$. In particular, with a slight adjustment of the constant c , the coverings in the above definition can be taken open.

The Nagata dimension will be used through the following lemma.

Lemma 2.6. *Suppose that X is a proper CAT(0) space, and $K \subset X$ is a compact subset with a cs -bounded covering with s -multiplicity at most $n + 1$, for some integer n and positive constants c and s . Then there exist*

- (1) *a finite simplicial complex Σ of dimension at most n , metrized as a subcomplex of some simplex of edge length s in a Euclidean space;*
- (2) *a constant $L > 0$ depending only on n and c , and an L -Lipschitz map $\psi: K \rightarrow \Sigma$;*
- (3) *a Lipschitz map $\phi: \Sigma \rightarrow X$ with $\phi(\Sigma^{(0)}) \subset K$ that is L -Lipschitz on every simplex of Σ , such that $\phi \circ \psi$ is L -Lipschitz and satisfies $d(x, \phi \circ \psi(x)) \leq Ls$ for all $x \in K$.*

Proof. By the remark preceding the lemma, we may assume that the given covering is finite, as K is compact. Now one can proceed as in [BWY23, Proposition 6.1], using a subordinate Lipschitz partition of unity. The fact that ϕ is Lipschitz, with a constant that is possibly not bounded in terms of n and c , is not stated there; here, it holds since ϕ is locally Lipschitz and Σ is compact. \square

We will also use the following related concept. The Assouad dimension of a metric space X is defined as the infimum of all (real) numbers $\beta > 0$ with the property that there exists some $C > 1$ such that, for every $\epsilon \in (0, 1)$, every set of diameter D in X can be covered by no more than $C\epsilon^{-\beta}$ sets of diameter at most ϵD (see [Hei01, Definition 10.15]). Unlike the Nagata dimension, this is in general not an integer. However, by [LDR15, Theorem 1.1], the Nagata dimension (as well as the Hausdorff dimension) is at most the Assouad dimension. The result is quantitative, in that the constant c in the definition of the Nagata dimension depends only on the Assouad dimension and the respective constant C .

2.6. Metric integral currents. For a proper metric space X , we let $\mathbf{I}_{*,c}(X)$ denote the chain complex of metric integral currents with compact support in X , see [AK00, Lan11]. As the use of currents will be limited to the proofs of Theorem 4.3 and Proposition 7.1, we give only a brief outline and refer to [GL23] for a more detailed introduction.

Formally, a current $T \in \mathbf{I}_{n,c}(X)$ is an $(n+1)$ -linear functional on the space of $(n+1)$ -tuples of real-valued Lipschitz functions on X , subject to some further conditions. As a basic example, every compact set $B \subset \mathbb{R}^n$ with finite perimeter, in particular every closed ball or polytope, induces a current $T = \llbracket B \rrbracket \in \mathbf{I}_{n,c}(\mathbb{R}^n)$ defined by

$$\llbracket B \rrbracket(f_0, f_1, \dots, f_n) := \int_B f_0 \det[\partial_j f_i]_{i,j=1}^n dx,$$

where the partial derivatives exist almost everywhere by Rademacher's theorem. Notice that for smooth functions f_i , this corresponds to the classical (de Rham) current given by the integration of the differential n -form $f_0 df_1 \wedge \dots \wedge df_n$ on B .

For a current $T \in \mathbf{I}_{n,c}(X)$, the boundary $\partial T \in \mathbf{I}_{n-1,c}(X)$ (if $n \geq 1$) and the push-forward $\varphi_{\#}T \in \mathbf{I}_{n,c}(Y)$ under a Lipschitz map $\varphi: X \rightarrow Y$ into another proper metric space Y are defined by the relations

$$\begin{aligned} \partial T(f_0, \dots, f_{n-1}) &:= T(1, f_0, \dots, f_{n-1}), \\ \varphi_{\#}T(g_0, \dots, g_n) &:= T(g_0 \circ \varphi, \dots, g_n \circ \varphi). \end{aligned}$$

Note that $\partial(\varphi_{\#}T) = \varphi_{\#}(\partial T)$. Associated with $T \in \mathbf{I}_{n,c}(X)$ is a compactly supported Borel measure $\|T\|$ on X with finite mass $\mathbf{M}(T) := \|T\|(X)$, and the support $\text{spt}(T)$ of T is defined as the support of $\|T\|$. If $\varphi: X \rightarrow Y$ is a λ -Lipschitz map, then $\mathbf{M}(\varphi_{\#}T) \leq \lambda^n \mathbf{M}(T)$. In the case of the above example $T = \llbracket B \rrbracket$, the measure $\|T\|$ is just the restriction of the Lebesgue measure to B . If X is a CAT(0) space and $u: B^2 \rightarrow X$ is a Lipschitz disc with connected fibers, then $\|u_{\#}\llbracket B^2 \rrbracket\|(B) = \text{area}(u(B^2) \cap B)$ for every Borel set $B \subset X$.

Every singular Lipschitz n -chain $\sum_{i=1}^N a_i \sigma_i$ in X , where $a_i \in \mathbb{Z}$ and the maps $\sigma_i: \Delta^n \rightarrow X$ are Lipschitz on an n -simplex $\Delta^n \subset \mathbb{R}^n$, induces a corresponding current $\sum_{i=1}^N a_i \sigma_{i\#}\llbracket \Delta^n \rrbracket \in \mathbf{I}_{n,c}(X)$. If X is locally Lipschitz contractible in a suitable sense, then the resulting chain map from singular Lipschitz chains to compactly supported integral currents induces an isomorphism of the respective homology theories (see [RS09] and [Mit19]). Moreover, with some additional assumptions on X , the subspace induced by Lipschitz n -chains is dense in $\mathbf{I}_{n,c}(X)$ with respect to the metric $(T, T') \mapsto \mathbf{N}(T - T')$, where $\mathbf{N}(T) := \mathbf{M}(T) + \mathbf{M}(\partial T)$ (see [Gol22, Theorem 1.3] and [BWY23, Corollary 1.5]). One of the main benefits from passing to $\mathbf{I}_{n,c}(X)$ is the following compactness theorem (see [AK00, Theorem 5.2 and Theorem 8.5]).

Theorem 2.7. *Given a sequence of integral currents $T_k \in \mathbf{I}_{n,c}(X)$ with supports in a fixed compact set $K \subset X$ and $\sup_k \mathbf{N}(T_k) < \infty$, there exists a subsequence converging weakly, that is, pointwise as functionals, to an integral current $T \in \mathbf{I}_{n,c}(X)$ with support in K .*

The mass is lower semi-continuous with respect to weak convergence: $T_k \rightarrow T$ implies that $\mathbf{M}(T) \leq \liminf_{k \rightarrow \infty} \mathbf{M}(T_k)$. For a cycle $S \in \mathbf{I}_{n,c}(X)$ of

dimension $n \geq 1$, a current $V \in \mathbf{I}_{n+1,c}(X)$ is a *filling* of S if $\partial V = S$. If X is a CAT(0) space, then a geodesic cone construction shows that such a V exists (see [Wen06b, Theorem 4.1]). A filling is *minimizing* if it has minimal mass among all fillings of its boundary cycle. The first part of the following analogue of Theorem 2.2 now follows directly from Theorem 2.7 together with the fact that the nearest point retraction $X \rightarrow K$ is 1-Lipschitz. For the assertion about the density, see [Wen06b, Corollary 4.4 and (4.28)].

Theorem 2.8. *Let X be a proper CAT(0) space, and let $S \in \mathbf{I}_{n,c}(X)$ be a cycle with support in a compact ball K . Then there exists a minimizing filling $V \in \mathbf{I}_{n,c}(X)$ of S with support in K . Furthermore, every minimizing filling V satisfies*

$$\omega_{n+1} \leq \frac{\|V\|(B(x, r))}{r^{n+1}} \leq \frac{\|V\|(B(x, s))}{s^{n+1}}$$

whenever $x \in \text{spt}(V)$ and $0 < r \leq s < d(x, \text{spt}(S))$.

See also [Wen05, Theorem 1.6], where X is not assumed to be proper.

Towards a linear isoperimetric inequality in dimensions above the asymptotic rank, the following result for a restricted class of cycles was proved (in a more general form) in [GL23, Theorem 1.1 and Theorem 7.3]. We say that $S \in \mathbf{I}_{n,c}(X)$ has *c-controlled density*, for some constant $c > 0$, if $\|S\|(B(x, r))/r^n \leq c$ for all $x \in X$ and $r > 0$.

Theorem 2.9. *Let X be a proper CAT(0) space of asymptotic rank at most $n \geq 1$. Then for all $c > 0$ there exists $\varrho = \varrho(X, n, c) > 0$ such that every cycle $S \in \mathbf{I}_{n,c}(X)$ with c-controlled density has a filling $V \in \mathbf{I}_{n+1,c}(X)$ such that $\mathbf{M}(V) \leq \varrho \mathbf{M}(S)$ and the support of V is within distance at most ϱ from $\text{spt}(S)$.*

The bound on the filling radius will be used in the proof of Theorem 4.3.

3. THE EUCLIDEAN ISOPERIMETRIC INEQUALITY FOR SPHERES

We first recall Reshetnyak's majorization theorem [Reš68] which implies the sharp Euclidean isoperimetric inequality for curves.

Proposition 3.1. *Let X be a CAT(0) space. Then for every closed unit speed curve $\alpha: [0, l] \rightarrow X$ there exist a compact convex set $C \subset \mathbb{R}^2$, a surjective closed unit speed curve $\beta: [0, l] \rightarrow \partial C$ that is simple except in the degenerate case when C is a segment of length $l/2$, and a 1-Lipschitz map $f: C \rightarrow X$ such that $f(\beta(t)) = \alpha(t)$ for all $t \in [0, l]$. In particular,*

$$\text{area}(f) \leq \text{area}(C) \leq \frac{1}{4\pi} \cdot l^2.$$

We now turn to the Euclidean isoperimetric inequality for 2-spheres, Theorem C, which we restate as follows.

Theorem 3.2. *Let X be a CAT(0) space. Then there exists a constant $C > 0$ such that every Lipschitz 2-sphere $f: S^2 \rightarrow X$ extends to a Lipschitz ball $\bar{f}: B^3 \rightarrow X$ with volume*

$$\text{vol}(\bar{f}) \leq C \cdot \text{area}(f)^{3/2}.$$

The following argument is similar to Wenger's proof of Gromov's filling inequality [Wen08b].

Proof. Let $f: S^2 \rightarrow X$ be a Lipschitz 2-sphere with image $S \subset X$. We will decompose f into finitely many Lipschitz 2-spheres $S = \bigcup_{a \in \mathcal{A}} S_a$ such that each S_a can be efficiently filled by coning. Our decomposition process works as follows. If $\Gamma \subset S^2$ is a bilipschitz Jordan curve, then we fill $f(\Gamma)$ by a Lipschitz disc $D \subset X$ using the isoperimetric inequality, thereby obtaining two Lipschitz 2-spheres $S^\pm \subset X$ that intersect in D . If we then fill both of them by Lipschitz 3-balls, we obtain a corresponding ball-filling for S . In the following, length, area and vol of the image Z of a Lipschitz map will denote the appropriate Hausdorff measure of Z , counted with multiplicities. For instance, if $f: S^2 \rightarrow X$ is a Lipschitz 2-sphere with image S and the map f is clear from the context, then we simply write

$$\text{area}(S) = \int_X \#f^{-1}(x) d\mathcal{H}^2(x).$$

Note that, by the area formula, this coincides with $\text{area}(f)$.

Sublemma 3.3. *There exist $q \in (0, 1)$ and $C > 0$, such that we can fill finitely many disjoint Jordan curves $\Gamma_j \subset S$ by Lipschitz discs D_j to produce a decomposition*

$$S \cup \bigcup_j D_j = \bigcup_{a \in \mathcal{A}} S_a \cup \bigcup_{b \in \mathcal{B}} S_b$$

into Lipschitz 2-spheres with the following properties.

- (1) $\sum_{a \in \mathcal{A}} \text{Fillvol}(S_a) \leq C \cdot \text{area}(S)^{3/2}$;
- (2) $\sum_{b \in \mathcal{B}} \text{area}(S_b) \leq q \cdot \text{area}(S)$.

Proof. The following three ingredients are needed.

- By the coarea formula, if $x \in X$, then

$$\text{length}(S \cap \partial B(x, r)) \leq \frac{d}{dr} \text{area}(S \cap B(x, r))$$

for almost every $r > 0$.

- For almost every point $p \in S^2$ we have

$$\liminf_{r \rightarrow 0} \frac{\text{area}(S \cap B(f(p), r))}{r^2} \geq \pi.$$

- For any Lipschitz 2-sphere $\varphi: S^2 \rightarrow B(x, r)$, coning off at the point x provides a Lipschitz 3-ball $\bar{\varphi}: B^3 \rightarrow B(x, r)$ with

$$\text{vol}(\bar{\varphi}) \leq \frac{r}{3} \cdot \text{area}(\varphi).$$

The proof of [Wen08b, Proposition] provides for every $\epsilon \in (0, 1)$ finitely many disjoint balls $B_i := B(x_i, r_i)$, $i = 1, \dots, k$, such that

- (1) $\text{area}(S \cap B_i) \geq \frac{\pi}{4} \epsilon \cdot r_i^2$;
- (2) $\text{length}(S \cap \partial B_i) \leq 2(\pi\epsilon)^{1/2} \cdot \text{area}(S \cap B_i)^{1/2}$;
- (3) $\sum_{i=1}^k \text{area}(S \cap B_i) \geq \frac{1}{25} \cdot \text{area}(S)$.

By Proposition 2.1, we may additionally assume that $f^{-1}(\partial B_i)$ is a disjoint union of Jordan curves up to a \mathcal{H}^1 -negligible set and that the gradient of the function $x \mapsto d(x_i, f(x))$ is non-zero for \mathcal{H}^1 -almost every point in $f^{-1}(\partial B_i)$. We choose an auxiliary $\varrho \in (0, \frac{1}{2})$ small enough, such that the inflated balls $(1 + \varrho)B_i := B(x_i, (1 + \varrho)r_i)$ are still disjoint. Let $\hat{F} := \{\hat{\Gamma}_j\}$ be the family of Jordan curves contained in $\bigcup_{i=1}^k f^{-1}(\partial B_i)$. Let $\hat{\Gamma}_j$ be such a Jordan curve with $f(\hat{\Gamma}_j) \subset \partial B_i$. For one of the components of $S^2 \setminus \hat{\Gamma}_j$, denoted Ω_j , the gradient of $x \mapsto d(x_i, f(x))$ is an inner normal vector of Ω_j at \mathcal{H}^1 -almost all points of Γ_j . Since f is Lipschitz continuous, only finitely many $\hat{\Gamma}_j$ have such an outer domain Ω_j with $f(\Omega_j) \not\subset (1 + \varrho)B_i$. We remove all other Jordan curves from \hat{F} and denote the remaining finite family by $F = \{\Gamma_j\}$. By Proposition 2.4, we may assume that F consists of bilipschitz Jordan curves. Next, we fill every Lipschitz circle $f|_{\Gamma_j}$ by a Lipschitz disc D_j such that $\text{area}(D_j) \leq \frac{1}{4\pi} \cdot \text{length}(\partial D_j)^2$. We denote by $\{D_j\}_{j \in J_i}$ the family of discs with boundary in ∂B_i and assume without loss of generality that $D_j \subset B_i$. By property (2), their total area is controlled by

$$\sum_{j \in J_i} \text{area}(D_j) \leq \frac{1}{4\pi} \cdot \text{length}(S \cap \partial B_i)^2 \leq \epsilon \cdot \text{area}(S \cap B_i).$$

By inserting the discs D_j , we have decomposed f into finitely many Lipschitz 2-spheres

$$S \cup \bigcup_j D_j = \bigcup_{a \in \mathcal{A}} S_a \cup \bigcup_{b \in \mathcal{B}} S_b$$

such that each S_a lies in $(1 + \varrho)B_i$ for some $i = i(a) \in \{1, \dots, k\}$ and each S_b intersects the balls B_i only in some of the inserted discs D_j . Denote by $\mathcal{A}_i \subset \mathcal{A}$ the subset of spheres in $(1 + \varrho)B_i$. Using the coning inequality and the properties of our decomposition, we estimate the filling volumes as

follows:

$$\begin{aligned}
 \sum_{a \in \mathcal{A}} \text{Fillvol}(S_a) &\leq \sum_i \sum_{a \in \mathcal{A}_i} \frac{(1+\varrho)r_i}{3} \cdot \text{area}(S_a) \\
 &\leq \sum_i \frac{(1+\varrho)r_i}{3} \cdot \left(\text{area}(S \cap (1+\varrho)B_i) + \sum_{j \in J_i} \text{area}(D_j) \right) \\
 &\leq \sum_i \frac{(1+\varrho)r_i}{3} \cdot (\text{area}(S \cap (1+\varrho)B_i) + \epsilon \cdot \text{area}(S \cap B_i)) \\
 &\leq \sum_i r_i \cdot \text{area}(S \cap (1+\varrho)B_i) \leq \sum_i \frac{2}{\sqrt{\pi\epsilon}} \cdot \text{area}(S \cap (1+\varrho)B_i)^{3/2} \\
 &\leq \frac{2}{\sqrt{\pi\epsilon}} \cdot \text{area}(S)^{3/2}.
 \end{aligned}$$

Next, we estimate the total area of the remaining spheres:

$$\begin{aligned}
 \sum_{b \in \mathcal{B}} \text{area}(S_b) &\leq \frac{24}{25} \cdot \text{area}(S) + \sum_j \text{area}(D_j) \\
 &\leq \frac{24}{25} \cdot \text{area}(S) + \epsilon \cdot \sum_i \text{area}(S \cap B_i) \leq \frac{49}{50} \cdot \text{area}(S)
 \end{aligned}$$

if $\epsilon \leq \frac{1}{50}$. This completes the proof of the Sublemma. □

To prove the theorem we will apply the Sublemma iteratively. Suppose that $S \subset B(o, R)$ for some $R > 0$ and $o \in X$. Set $\lambda = q^{3/2} \in (0, 1)$. Choose $l \in \mathbb{N}$ such that $\frac{R}{3} \cdot \lambda^l \leq \text{area}(S)^{1/2}$. Applying the Sublemma l times in $B(o, R)$, we can estimate the filling volume of S as follows:

$$\begin{aligned}
 \text{Fillvol}(S) &\leq \sum_{a \in \mathcal{A}_1} \text{Fillvol}(S_a) + \dots + \sum_{a \in \mathcal{A}_l} \text{Fillvol}(S_a) + \sum_{b \in \mathcal{B}_l} \text{Fillvol}(S_b) \\
 &\leq C \cdot \left(\text{area}(S)^{3/2} + \sum_{b \in \mathcal{B}_1} \text{area}(S_b)^{3/2} + \dots + \sum_{b \in \mathcal{B}_{l-1}} \text{area}(S_b)^{3/2} + \frac{R}{3} \cdot \sum_{b \in \mathcal{B}_l} \text{area}(S_b) \right) \\
 &\leq C \cdot \left(\text{area}(S)^{3/2} + \lambda \cdot \text{area}(S)^{3/2} + \dots + \lambda^{l-1} \cdot \text{area}(S)^{3/2} + \frac{R}{3} \cdot \lambda^l \cdot \text{area}(S) \right) \\
 &\leq C \cdot (1 + \lambda + \dots + \lambda^{l-1} + 1) \cdot \text{area}(S)^{3/2} \\
 &\leq C' \cdot \text{area}(S)^{3/2}
 \end{aligned}$$

with $C' = C \cdot \frac{2-\lambda}{1-\lambda}$, independent of l . □

4. MINIMAL TRIANGLES AND TETRAHEDRA

We assume that X is a proper CAT(0) space. Recall the notion of a minimal disc provided by Theorem 2.2. For us, minimal discs filling geodesic triangles will play a special role, and we call such a disc a *minimal triangle*. The purpose of this section is to prove the linear isoperimetric inequality

for minimal tetrahedra stated in Theorem D. A *minimal tetrahedron* is a Lipschitz 2-sphere composed of four minimal triangles as faces, six geodesics as edges, and four points as vertices.

A key observation for the proof is that minimal triangles and tetrahedra in X have Nagata dimension at most 2. This is a consequence of the following a priori density bound for minimal triangles.

Lemma 4.1. *Let $u: B^2 \rightarrow X$ be a minimal triangle. For all $x \in X$ and $r > 0$,*

$$\text{area}(u(B^2) \cap B(x, r)) \leq \frac{3\pi}{2} \cdot r^2.$$

Proof. By definition, u is a solution to the Plateau problem for a geodesic triangle $\Delta \subset X$. If Δ is a Jordan curve, then the statement is a consequence of [Sta21, Proposition 80] because geodesic triangles have total curvature at most 3π . However, the argument from [Sta21] also covers the case of a general triangle Δ . We recall the necessary steps, but refer the reader to [Sta21, Lemmata 78 and 79] for more details. By Theorem 2.2, we can decompose u as $u = u' \circ u''$ with $u'': B^2 \rightarrow Z$ and $u': Z \rightarrow X$. The space Z is a CAT(0) disc retract and u'' is a surjective Lipschitz map with connected fibers. Furthermore, u' is area minimizing among Lipschitz maps from Z , and preserves the length of every rectifiable curve. In particular, u' is 1-Lipschitz and therefore ∂Z is a geodesic triangle whose sides are mapped isometrically by u' . We now extend u' to a minimal plane in a larger CAT(0) space \hat{X} . We obtain \hat{X} in two steps. We first glue a half-line to every vertex of Δ to obtain a CAT(0) space X' . Note that in X' every side s of Δ is contained in a complete geodesic c_s that intersects X precisely in s . We then glue a flat half-plane along the boundary to every geodesic c_s to obtain \hat{X} . By Reshetnyak's gluing theorem, \hat{X} is CAT(0). In a similar way, we extend Z to a CAT(0) plane \hat{Z} . Note that the map u' extends to a map $f: \hat{Z} \rightarrow \hat{X}$ that is locally isometric on $\hat{Z} \setminus Z$. The map f is a *proper intrinsic minimal plane* in the sense of [Sta21, Definition 41]. In particular, f satisfies the monotonicity of area densities for all radii $r > 0$ [Sta21, Proposition 62]. By construction, the area growth of f is equal to the area growth of a flat cone over a circle of length 3π :

$$\lim_{r \rightarrow \infty} \frac{\text{area}(f(\hat{Z}) \cap B(x, r))}{\pi r^2} = \frac{3}{2}.$$

Hence, we obtain the required inequality. \square

In what follows, we use the dimension theory definitions as provided in Section 2.5.

Lemma 4.2. *Let $u: B^2 \rightarrow X$ be a minimal triangle. Then $u(B^2)$ has Assouad dimension at most 2 and hence also Nagata dimension at most 2, both with absolute constants.*

Proof. Let $\epsilon \in (0, 1)$, and let $Y \subset u(B^2)$ be a subset of diameter $s > 0$. Put $r := \frac{\epsilon}{6}s$, and choose a maximal set $Z \subset Y$ of distinct points at mutual distance $> 3r$. The balls $B(z, 3r)$ with $z \in Z$ cover Y , and their diameter is at most ϵs . Let $Z' \subset Z$ denote the set of points at distance at most r from the geodesic triangle $u(S^1)$. Choosing a closest point projection $\pi: Z' \rightarrow u(S^1)$, we get a set $\pi(Z')$ of distinct points at mutual distance $> r$. Furthermore, the intersection of $\pi(Z')$ with each side of $u(S^1)$ has diameter at most s . It follows that

$$\#Z' \leq 3\left(\frac{s}{r} + 1\right) \leq \frac{21}{\epsilon} \leq \frac{21}{\epsilon^2}.$$

On the other hand, if $z \in Z \setminus Z'$, then $\text{area}(u(B^2) \cap B(z, r)) \geq \pi r^2$ by Theorem 2.2), and the r -neighborhood of $Z \setminus Z'$ in $u(B^2)$ has diameter at most $s + 2r \leq \frac{4}{3}s$ and area at most $\frac{3\pi}{2} \cdot (\frac{4}{3}s)^2 = \frac{8\pi}{3}s^2$ by Lemma 4.1. Hence,

$$\#(Z \setminus Z') \leq \frac{8\pi s^2}{3\pi r^2} < \frac{96}{\epsilon^2}.$$

This shows that $u(B^2)$ can be covered by $117\epsilon^{-2}$ sets of diameter ϵs . Thus, $u(B^2)$ has Assouad dimension at most 2, hence, by [LDR15, Theorem 1.1], also Nagata dimension at most 2, with an absolute implicit constant. \square

We are now ready for the proof of Theorem D, which we restate for convenience. The argument uses an idea of White [Whi83, Whi84].

Theorem 4.3. *Let X be a proper CAT(0) space of asymptotic rank at most 2. There exists a constant $\mu > 0$ such that every minimal tetrahedron $f: S^2 \rightarrow X$ satisfies a linear isoperimetric inequality for fillings by balls,*

$$\text{Fillvol}(f) \leq \mu \cdot \text{area}(f).$$

Proof. Let $f: S^2 \rightarrow X$ be a minimal tetrahedron with image τ . By definition, S^2 is parametrized by four positively oriented bilipschitz discs $\varphi_1, \dots, \varphi_4: B^2 \rightarrow S^2$ such that each $u_i := f \circ \varphi_i: B^2 \rightarrow X$ is a minimal triangle. Accordingly, the integral cycle $S := f_{\#}(\partial[B^3]) \in \mathbf{I}_{2,c}(X)$ induced by f can be written as the sum $\sum_i u_{i\#}[B^2]$. By Lemma 4.1, we have

$$\|u_{\#}[B^2]\|(B(x, r)) \leq \frac{3\pi}{2} \cdot r^2$$

for every minimal triangle u , every $x \in X$ and $r > 0$. Thus, S has uniformly controlled density. Hence, by Theorem 2.9, S admits a filling by an integral 3-current $V \in \mathbf{I}_{3,c}(X)$ whose support lies within uniformly bounded distance $s > 0$ from τ . By Lemma 4.2, τ has Nagata dimension at most 2. Since τ is compact, and by the remark preceding Lemma 2.6, there exists an absolute constant c and a finite cs -bounded cover of the closed s -neighborhood K of τ with s -multiplicity at most 3. Applying Lemma 2.6 to K , we find a finite simplicial complex Σ of dimension at most 2, metrized as a subcomplex of some simplex of edge length s in a Euclidean space, and Lipschitz maps $\psi: K \rightarrow \Sigma$ and $\phi: \Sigma \rightarrow X$. The push-forwards by ψ of the filling V , and

hence also of the cycle ∂V , are zero as currents. Since Σ is piecewise Euclidean, it is locally conical. In particular, by [RS09, Corollary 1.4] or [Mit19, Theorem 1.3], the homology theories induced by compactly supported integral currents and by Lipschitz chains, respectively, are isomorphic. Thus, the Lipschitz cycle $\psi \circ f$ is homologically trivial. Now we cone off the 1-skeleton of Σ to get a simply connected complex $\hat{\Sigma}$. We can arrange that $\hat{\Sigma}$ is again metrized as a subcomplex of some simplex of edge length s in a Euclidean space. We extend ϕ to a Lipschitz map $\hat{\phi}: \hat{\Sigma} \rightarrow X$, by first choosing an arbitrary image point for the new vertex of $\hat{\Sigma}$, and then extending to all of $\hat{\Sigma}$, via coning. By Hurewicz, $\psi \circ f$ bounds a Lipschitz 3-ball Ψ in $\hat{\Sigma}$, which has volume zero. Then, $\hat{\phi} \circ \Psi$ is a Lipschitz 3-ball of volume zero that fills $v \circ f$, where $v = \phi \circ \psi$. We combine $\hat{\phi} \circ \Psi$ with the geodesic homotopy h from f to $v \circ f$ to obtain a Lipschitz 3-ball filling f . By Lemma 2.6, the map v is L -Lipschitz where $L > 0$ depends only on c , and has controlled displacement, $d(x, v(x)) \leq L \cdot s$ for all $x \in K$. In particular, $\text{vol}(h) \leq L \cdot s \cdot \text{area}(f)$. Thus, the result holds for $\mu := L \cdot s$. \square

5. QUANTITATIVE TRIANGULATION OF DISCS

In this section, the goal is to prove the necessary quantitative triangulation result for discs. The arguments are inspired by [Pap96, Pap09]. Other quantitative triangulation results of surfaces have recently appeared in [CR22, NR23]. Our setting differs in that the underlying topology of the metric spaces in question may be complicated. To preserve the good topological properties of surfaces, we will instead work with pseudo-metrics.

Let X be a complete metric space, Z a complete length space homeomorphic to a closed disc and $\varphi: Z \rightarrow X$ a Lipschitz map. We denote by d_φ the associated *intrinsic pseudo-metric* on Z ,

$$d_\varphi(x, y) := \inf_{\gamma} \text{length}(\varphi \circ \gamma)$$

where the infimum runs over all paths γ in Z joining x and y . We define the *intrinsic radius* of Z by

$$\text{rad}_\varphi(Z) := \max_{x \in Z} d_\varphi(x, \partial Z).$$

Moreover, for a curve $\gamma \subset Z$ we will use $\text{length}_\varphi(\gamma) := \text{length}(\varphi \circ \gamma)$ to denote its φ -length.

By a *separating arc* α of Z we mean a simple arc $\alpha \subset Z$ that intersects ∂Z precisely in the two (distinct) endpoints.

The following proposition is analogous to the result for van Kampen diagrams on p. 796 in [Pap96].

Proposition 5.1. *Suppose that $\text{rad}_\varphi(Z) \leq r \leq \text{length}_\varphi(\partial Z)/17$. Then there exists a separating arc α of Z such that the two discs D_1, D_2 with $D_1 \cup D_2 = Z$ and $D_1 \cap D_2 = \alpha$ satisfy*

$$\text{length}_\varphi(\partial D_1) \leq 17r, \quad \text{length}_\varphi(\partial D_2) \leq \text{length}_\varphi(\partial Z) - r.$$

Proof. We denote by d'_φ the intrinsic pseudo-metric on ∂Z induced by $\varphi|_{\partial Z}$, thus $d'_\varphi(x, y) \geq d_\varphi(x, y)$ is the shortest φ -length of a subarc of ∂Z connecting $x, y \in \partial Z$. Suppose first that there exists a point $p \in \partial Z$ such that

$$q \in \partial Z, d_\varphi(p, q) \leq 4r \Rightarrow d'_\varphi(p, q) \leq 6r.$$

Let $\bar{p} \in \partial Z$ be a point with $d'_\varphi(p, \bar{p}) = \frac{1}{2} \text{length}_\varphi(\partial Z)$, and note that $d_\varphi(p, \bar{p}) > 4r$ because $d'_\varphi(p, \bar{p}) > 6r$. Thus, the set $S_{3r}^\varphi(p) = \{d_\varphi(p, \cdot) = 3r\}$ separates p from \bar{p} . Hence, there exists a simple arc β in $S_{3r}^\varphi(p)$ with endpoints $p_1, p_2 \in \partial Z$ that still separates p from \bar{p} in Z . Since $\text{rad}_\varphi(Z) \leq r$, there exists a point $q \in \beta$ at d_φ -distance at most r from both connected components of $\partial Z \setminus \{p, \bar{p}\}$. For $i = 1, 2$, choose a point q_i closest to q in the component containing p_i . Since $d_\varphi(p, q) = 3r$, we have $2r \leq d_\varphi(p, q_i) \leq 4r$, hence $2r \leq d'_\varphi(p, q_i) \leq 6r$ and $4r \leq d'_\varphi(q_1, q_2) \leq 12r$. There exists a separating arc from q_1 to q_2 of φ -length less than or equal to $2r$, and the resulting discs D_1 and D_2 containing p and \bar{p} , respectively, satisfy

$$\text{length}_\varphi(\partial D_1) \leq 14r, \quad \text{length}_\varphi(\partial D_2) \leq \text{length}_\varphi(\partial Z) - 4r + 2r.$$

In the remaining case, we have that for every $p \in \partial Z$ there exists a $q \in \partial D$ with $d_\varphi(p, q) \leq 4r$ and $d'_\varphi(p, q) \geq 6r$. We fix p, q such that $d'_\varphi(p, q)$ is minimal among all such pairs. If $d'_\varphi(p, q) \leq 9r$, we choose a separating arc from p to q of φ -length less than $5r$ to get a decomposition with

$$\text{length}_\varphi(\partial D_1) \leq 14r, \quad \text{length}_\varphi(\partial D_2) \leq \text{length}_\varphi(\partial Z) - 6r + 5r.$$

If $d'_\varphi(p, q) > 9r$, let $\beta \subset \partial Z$ be a subarc from p to q of φ -length $d'_\varphi(p, q)$, and let $p' \in \beta$ be the point with $d'_\varphi(p, p') = 9r$. There exists a $q' \in \partial Z$ with $d_\varphi(p', q') \leq 4r$ and $d'_\varphi(p', q') \geq d'_\varphi(p, q)$ by the choice of p and q . In particular, $q' \notin \beta$, thus any path in Z from p to q separates p' and q' . It follows that

$$d_\varphi(p, p') < d_\varphi(p, q) + d_\varphi(p', q') \leq 8r,$$

thus there exists a separating arc from p to p' of φ -length less than $8r$. This gives a decomposition satisfying the assertion of the lemma. \square

Recall the notion of a disc decomposition from Section 2.4.

Corollary 5.2. *Let $r \geq \text{rad}_\varphi(Z)$. Then there exists a disc decomposition $Z = D_1 \cup \dots \cup D_n$ induced by an embedded graph $\Gamma \subset Z$ such that*

- (1) $n \leq \text{length}_\varphi(\partial Z)/r$, provided that $\text{length}_\varphi(\partial Z) \geq r$;
- (2) $\text{length}_\varphi(\partial D_i) \leq 17r$ for $i = 1, \dots, n$.

Furthermore, $D_i \cap D_j$ is either empty, a point, or a simple arc for $i \neq j$.

Proof. If $\text{length}_\varphi(\partial Z) \leq 17r$, then the result holds for $n = 1$. If $\text{length}_\varphi(\partial Z) > 17r$, we use Proposition 5.1 and induction. \square




FIGURE 2. Example of a disc decomposition.

6. IMPROVING 2-SPHERES

This is a technical section that provides a tool needed in the proof of our main result. The setting we have in mind here is that we already know how to find fillings of small volume for all Lipschitz 2-spheres of area at most $A > 0$. Our aim is then to understand how to improve the intrinsic structure of a Lipschitz 2-sphere S of area only a little bit larger than A .

We begin with a useful calculus fact.

Lemma 6.1. *Given $\delta > 0$, there exists $\lambda = \lambda(\delta) \in (0, 1)$ such that*

$$(x + y)^{1+\delta} > (x + a)^{1+\delta} + (y + a)^{1+\delta}$$

whenever $0 < x \leq y$ and $a \leq \lambda x$.

Proof. Dividing the desired inequality by $(y + a)^{1+\delta}$, and setting $a := \lambda x$ and $q := x/(y + a)$, we see that it suffices to find λ so that

$$((1 - \lambda)q + 1)^{1+\delta} > ((1 + \lambda)q)^{1+\delta} + 1.$$

Since $(1 + a)^{1+\delta} \geq 1 + (1 + \delta)a$ for $a \geq 0$, and since $(1 + \lambda)q \leq 1$, this can be further reduced to the inequality

$$(1 + \delta)(1 - \lambda) > 1 + \lambda,$$

which holds for $\lambda < \delta/(2 + \delta)$. \square

Let again X be a CAT(0) space. For $M, \delta > 0$ we consider all Lipschitz spheres $f: S^2 \rightarrow X$ such that

$$\text{Fillvol}(f) \geq M \cdot \text{area}(f)^{1+\delta}.$$

For $\epsilon \geq 0$ we call such a sphere (M, δ, ϵ) -minimal if every Lipschitz sphere $f': S^2 \rightarrow X$ with $\text{area}(f') \leq \text{area}(f) - \epsilon$ satisfies the reverse inequality $\text{Fillvol}(f') \leq M \cdot \text{area}(f')^{1+\delta}$.

This notion is related to the intrinsic isoperimetric profile of a sphere. We say that a Lipschitz sphere $f: S^2 \rightarrow X$ is *intrinsically* (c, θ) -isoperimetric, for some $c > 0$ and $\theta \geq 0$, if every disc $D \subset S^2$ with $\text{area}_f(D) \leq \frac{1}{2} \text{area}(f)$ and $\text{area}_f(D) \geq \theta$ satisfies $\text{area}_f(D) \leq c \cdot \text{length}_f(\partial D)^2$.

Lemma 6.2. *For every $\delta > 0$ and $\theta \geq 0$ there exist $c = c(\delta) > 0$ and $\epsilon(\delta, \theta) \in [0, \theta]$ such that the following holds for every $M > 0$. If $f: S^2 \rightarrow X$ is an (M, δ, ϵ) -minimal Lipschitz sphere, then f is intrinsically (c, θ) -isoperimetric.*

Proof. Given $\delta > 0$ and $\theta \geq 0$, let $\lambda = \lambda(\delta) \in (0, 1)$ be the constant from Lemma 6.1 and put

$$c := \frac{1}{4\pi\lambda}, \quad \epsilon := (1 - \lambda)\theta.$$

Let $f: S^2 \rightarrow X$ be an (M, δ, ϵ) -minimal Lipschitz sphere and suppose, contrary to the assertion, that there exists a decomposition $S^2 = D_1 \cup D_2$ into two discs along a Jordan curve $\alpha \subset S^2$ such that

$$\theta \leq x := \text{area}_f(D_1) \leq y := \text{area}_f(D_2)$$

and $\text{area}_f(D_1) > c \cdot \text{length}_f(\alpha)^2$. After reparametrizing f using Proposition 2.4, we can assume that α is piecewise geodesic. Then, by Proposition 3.1 and the choice of c , we can fill the closed curve $f|_\alpha$ by a Lipschitz disc of area

$$a \leq \frac{1}{4\pi} \cdot \text{length}_f(\alpha)^2 \leq \lambda \cdot \text{area}_f(D_1) = \lambda x$$

in X . Since α is piecewise geodesic, this creates two Lipschitz 2-spheres f_1, f_2 with

$$\text{area}(f) = x + y, \quad \text{area}(f_1) = x + a, \quad \text{area}(f_2) = y + a.$$

Note that $x - a \geq (1 - \lambda)x \geq (1 - \lambda)\theta = \epsilon$ and hence

$$\text{area}(f_i) \leq \text{area}(f) - x + a \leq \text{area}(f) - \epsilon.$$

Thus, by (M, δ, ϵ) -minimality of f and Lemma 6.1, we have

$$\begin{aligned} \text{Fillvol}(f) &\leq \text{Fillvol}(f_1) + \text{Fillvol}(f_2) \\ &\leq M \cdot (x + a)^{1+\delta} + M \cdot (y + a)^{1+\delta} \\ &< M \cdot (x + y)^{1+\delta} = M \cdot \text{area}(f)^{1+\delta}, \end{aligned}$$

contradicting the assumption on f . □

We will now work towards the main result of this section, Proposition 6.5. We begin with a modified version of the classical Besicovitch inequality [Gro83]. In the following we denote by $Q = [0, 1]^2$ the unit square.

Lemma 6.3 (Besicovitch). *Let X be a complete metric space and $\varphi: Q \rightarrow X$ a Lipschitz disc. Suppose that the opposite sides of the square have intrinsic distance at least a and b , respectively. Then $\text{area}(\varphi) \geq a \cdot b$.*

Proof. Let us first assume that φ is bilipschitz. Denote by $Z = (Q, d_\varphi)$ the associated intrinsic disc and by $\pi_\varphi: Q \rightarrow Z$ the induced canonical projection which is also bilipschitz. By [LW18a, Corollary 3.2], the metric derivatives of φ and π_φ coincide almost everywhere. Thus, $\text{area}(\varphi) = \text{area}(\pi_\varphi)$. Since φ is bilipschitz, the area formula implies $\text{area}(\pi_\varphi) = \mathcal{H}^2(Z)$. Hence, we obtain the claim from the classical Besicovitch inequality which shows $\mathcal{H}^2(Z) \geq a \cdot b$. If φ is only Lipschitz, we consider for $\epsilon > 0$ the auxiliary bilipschitz map $\varphi_\epsilon: Q \rightarrow X \times \mathbb{R}^3$ defined by $\varphi_\epsilon(x) = (\varphi(x), \epsilon x)$. Note that $d_{\varphi_\epsilon} \geq d_\varphi$ and $\text{area}(\varphi_\epsilon) \rightarrow \text{area}(\varphi)$ for $\epsilon \rightarrow 0$. This concludes the proof. \square

The proof of Proposition 6.5 also employs the following decomposition lemma for spheres analogous to [Pap09, Proposition 2.3]. Since our setting is slightly different, we provide a complete argument.

Similarly as for spheres, we say that a Lipschitz disc $f: D \rightarrow X$ is *intrinsically (c, θ) -isoperimetric* if every subdisc $D' \subset D$ with $\text{area}_f(D') \geq \theta$ satisfies $\text{area}_f(D') \leq c \cdot \text{length}_f(\partial D')^2$.

Lemma 6.4. *Let $f: S^2 \rightarrow X$ be a Lipschitz 2-sphere. Then there exists a simple closed curve $\alpha \subset S^2$ such that the two discs H_1, H_2 with $H_1 \cup H_2 = S^2$ and $H_1 \cap H_2 = \alpha$ satisfy*

$$\text{area}_f(H_i) \geq \frac{1}{4} \text{area}(f) \quad (i = 1, 2), \quad \text{length}_f(\alpha) \leq 4\sqrt{\text{area}(f)}.$$

Moreover, if f is intrinsically (c, θ) -isoperimetric for some $c > 0$ and $\theta \in [0, \frac{1}{4} \text{area}(f)]$, then both $f|_{H_1}$ and $f|_{H_2}$ are intrinsically $(3c, \theta)$ -isoperimetric.

Proof. Since f is Lipschitz, there exists a Jordan curve α in S^2 such that the resulting discs H_1 and H_2 both have f -area at least $\frac{1}{4} \text{area}(f)$. Denote by \mathcal{F} the family of all such Jordan curves in S^2 and let \bar{L} denote the infimal f -length of curves in \mathcal{F} . Let $\alpha \in \mathcal{F}$ be ϵ -minimal, i.e. $\text{length}_f(\alpha) < \bar{L} + \epsilon$. Denote by H_1 and H_2 the discs associated to α and assume that $\text{area}_f(H_1) \geq \text{area}_f(H_2)$. Note that if c is an arc in H_1 joining two points on α , and \bar{c} is the shortest subarc of α joining the same points, then $\text{length}_f(c) \geq \text{length}_f(\bar{c}) - \epsilon$. This shows that if we decompose α into four arcs of equal length, then opposite arcs have distance in H_1 at least $\frac{1}{4} \text{length}_f(\alpha) - \epsilon$. By Lemma 6.3, we conclude that

$$\text{area}_f(H_1) \geq \left(\frac{1}{4} \text{length}_f(\alpha) - \epsilon \right)^2.$$

Choosing $\epsilon < (1 - \frac{\sqrt{3}}{2})\sqrt{\text{area}(f)}$, we obtain the desired inequalities.

Suppose now that f is intrinsically (c, θ) -isoperimetric with $c > 0$ and $\theta \in [0, \frac{1}{4} \text{area}(f)]$, and $D \subset H_i$ is a subdisc. We only need to consider the

case that $\text{area}_f(D) > \frac{1}{2} \text{area}(f)$. Then

$$\text{area}_f(D) \leq \frac{3}{4} \text{area}(f) \leq 3 \text{area}_f(S^2 \setminus D) \leq 3c \cdot \text{length}_f(\partial D)^2$$

as claimed. \square

Now we come to the main goal of this section. The result is inspired by [Pap96] (see p. 803 therein).

Proposition 6.5. *Let X be a proper CAT(0) space. For $\delta, \epsilon \in (0, 1]$ and $M > 0$, let $f: S^2 \rightarrow X$ be an (M, δ, ϵ) -minimal Lipschitz 2-sphere with $\text{area}(f) \geq 4$. Suppose further that f is intrinsically (c, θ) -isoperimetric for some $\theta \in [0, 1]$ and $c \geq 1$. Then we find for every $s > 0$ with $\frac{\theta}{3c} \leq s^2 \leq \frac{1}{8c} \text{area}(f)$ a Lipschitz 2-sphere $f': S^2 \rightarrow X$ with $\text{area}(f') \leq \text{area}(f)$ and a Lipschitz homotopy h from f to f' with*

$$\text{vol}(h) \leq 12M \cdot c \cdot s^{2\delta} \cdot \text{area}(f);$$

moreover, f' admits a disc decomposition $S^2 = D_1 \cup \dots \cup D_n$, induced by a finite Lipschitz graph $\Gamma \subset S^2$, such that

- (1) $n \leq 14 \cdot \text{area}(f)/s^2$,
- (2) $\text{length}_{f'}(\partial D_i) \leq 34s$,
- (3) $\text{area}_{f'}(D_i) \leq 3c \cdot (34s)^2$

for $i = 1, \dots, n$. In particular, $\text{length}_{f'}(\Gamma) \leq 238 \cdot \text{area}(f)/s$.

Proof. Let α and H_1, H_2 be given as in Lemma 6.4. Then

$$\text{length}_f(\partial H_i) \leq 4\sqrt{\text{area}(f)} \leq 8\sqrt{\text{area}_f(H_i)}, \quad i = 1, 2.$$

We will provide the necessary structure by treating the discs $f|_{H_i}$, $i = 1, 2$, individually. Let $\varphi: D \rightarrow X$ be either one of these discs.

Consider the Lipschitz functions $g: D \rightarrow \mathbb{R}$ and $g_s: D \rightarrow \mathbb{R}/s\mathbb{Z}$ defined by $g(x) := d_\varphi(x, \partial D)$ and $g_s(x) := g(x) \bmod s$. By Proposition 2.1, we find $s_1 \in (\frac{1}{2}s, s)$ such that the s_1 -fiber satisfies $g_s^{-1}(s_1) = \hat{\Pi} \cup N$ where $\hat{\Pi}$ is a disjoint union of Jordan curves $\hat{\Pi} = \bigcup_{j \in J} \alpha_j$, $J \subset \mathbb{N}$, and N is \mathcal{H}^1 -negligible. Moreover, by the coarea inequality, we can choose s such that

$$\text{length}_\varphi(\hat{\Pi}) \leq \frac{2 \text{area}(\varphi)}{s}.$$

We let $\Omega_j \subset D$ denote the Jordan domain of α_j , and we define J_s as the set of all $j \in J$ such that α_j does not lie in the closure of a domain Ω_k with $\text{length}_\varphi(\alpha_k) < s$. We assume $J_s = \{1, \dots, m\}$ and set

$$\Pi = \bigcup_{j \in J_s} \alpha_j.$$

In particular, for each $j \in J_s$ the curve α_j itself has $\text{length}_\varphi(\alpha_j) \geq s$. Hence,

$$m \leq \frac{2 \text{area}(\varphi)}{s^2}.$$

Define $s_0 = 0$ and $s_i = s_1 + (i - 1)s$ for $i \geq 2$. We have $g_s^{-1}(s_1) = \bigcup_{i=1}^{\infty} g^{-1}(s_i)$. The regions $g^{-1}[s_{i-1}, s_i]$ have φ -radius at most s .

By Proposition 2.4, we can deform Π to be piecewise geodesic without changing its φ -length, at the cost of precomposing f with a monotone Lipschitz map that is uniformly close to the identity map on S^2 . This deformation requires a Lipschitz homotopy of zero volume. In particular, the resulting Lipschitz sphere is still intrinsically (c, θ) -isoperimetric and (M, δ, ϵ) -minimal. Hence, we can assume that Π itself consists of piecewise geodesic Jordan curves.

We now define $\varphi' : D \rightarrow X$ such that $\varphi' = \varphi$ on $D \setminus \bigcup_{j \in J \setminus J_s} \Omega_j$ and $\varphi'|_{\Omega_j}$ is a minimal disc if $j \notin J_s$ and Ω_j is not contained in a larger Jordan domain Ω_k with $k \notin J_s$. In particular, for such a minimal disc $\varphi'|_{\Omega_j}$ we have $\text{area}_{\varphi'}(\Omega_j) \leq \frac{1}{4\pi}s^2$ and hence $\text{rad}_{\varphi'}(\Omega_j) \leq \frac{1}{2\pi}s$ by Proposition 3.1 and [Sta21, Lemma 51]. We conclude that the φ' -radii of the components of $D \setminus \Pi$ are bounded by $\frac{2\pi+1}{2\pi}s \leq 2s$.

The closure of each component of $D \setminus \Pi$ is a disc with holes. By Lemma 2.5, after a small deformation, we may assume that there are m pairs of disjoint piecewise geodesic arcs β_j^{\pm} in D of φ' -length at most $2s$ such that the graph $\tilde{\Gamma} = \Pi \cup \bigcup_{j=1}^m (\beta_j^- \cup \beta_j^+)$ induces a disc decomposition of D into discs D_l of φ' -radius at most $2s$.




FIGURE 3. Illustration of the decomposition process.

Now we apply Corollary 5.2 with $r = 2s$ to each of those discs. Thus, either $\text{length}_{\varphi}(\partial D_l) < 2s$ and we set $O_l := D_l$, or $\text{length}_{\varphi}(\partial D_l) \geq 2s$ and we obtain a decomposition into discs

$$D_l = O_{1l} \cup \dots \cup O_{pl}$$

induced by a graph $\Gamma_l \subset D_l$ with $p \leq \frac{1}{2s} \text{length}_{\varphi'}(\partial D_l)$ and $\text{length}_{\varphi'}(\partial O_{il}) \leq 34s$. Renumbering, we obtain a disc decomposition

$$D = O_1 \cup \dots \cup O_{\tilde{n}}$$

induced by a graph $\hat{\Gamma} = \tilde{\Gamma} \cup \bigcup_l \Gamma_l$. Since there are $2m+1$ discs D_l , and since $\text{area}(\varphi) \geq \text{area}(f)/4 \geq 2s^2$, we have

$$\begin{aligned} \tilde{n} &\leq \frac{1}{2s} \cdot \sum_l \text{length}_{\varphi}(\hat{\partial} D_l) + (2m+1) \\ &\leq \frac{1}{s} \left(\text{length}_{\varphi}(\Pi) + m \cdot 2s + \frac{1}{2} \cdot \text{length}_{\varphi}(\partial D) \right) + (2m+1) \\ &\leq \frac{1}{s} \left(\frac{2 \text{area}(\varphi)}{s} + \frac{4 \text{area}(\varphi)}{s} + 4\sqrt{\text{area}(\varphi)} \right) + \frac{5 \text{area}(\varphi)}{s^2} \\ &\leq \frac{14 \text{area}(\varphi)}{s^2}. \end{aligned}$$

Hence, by applying the above construction to both discs $f|_{H_i}$, $i = 1, 2$, we obtain a disc decomposition

$$S^2 = D_1 \cup \dots \cup D_n$$

induced by $\Gamma = \hat{\Gamma}_1 \cup \alpha \cup \hat{\Gamma}_2$, where $\hat{\Gamma}_i$ denotes the graph constructed for $f|_{H_i}$. (If it happens that $\Gamma = \alpha$, then we introduce an arbitrary vertex on α , so that Γ is a graph.) We have already established properties (1) and (2). The remaining property (3) follows because $\theta \leq \text{area}(f)/4$, and therefore both $f|_{H_1}$ and $f|_{H_2}$ are intrinsically $(3c, \theta)$ -isoperimetric by Lemma 6.4.

It remains to construct the homotopy h . Again, we consider $f|_{H_1}$ and $f|_{H_2}$ separately and continue with the above setup. For each of the minimal discs $\varphi'|\Omega_j$ in the definition of φ' , the union with $\varphi|\Omega_j$ defines a Lipschitz 2-sphere σ_j with

$$\text{area}(\sigma_j) \leq 2 \text{area}_{\varphi}(\Omega_j) \leq 6cs^2 \leq \frac{3}{4} \cdot \text{area}(f) \leq \text{area}(f) - \epsilon$$

by the properties of f and the choice of s . Thus, by assumption, we have

$$\text{Fillvol}(\sigma_j) \leq M \cdot \text{area}(\sigma_j)^{1+\delta}.$$

Filling all spheres σ_j , we obtain h and the bound

$$\begin{aligned} \text{vol}(h) &\leq M \cdot \sum_j \text{area}(\sigma_j)^{1+\delta} \leq M \cdot \sum_j (2 \cdot \text{area}_{\varphi}(\Omega_j))^{1+\delta} \\ &\leq 4M \cdot \sum_j \text{area}_{\varphi}(\Omega_j) \cdot (3cs^2)^{\delta} \leq 12M \cdot c \cdot s^{2\delta} \cdot \text{area}(\varphi). \end{aligned}$$

Note that h keeps the boundary of φ fixed. Applying the construction to both $f|_{H_1}$ and $f|_{H_2}$, we get the desired homotopy of f . This completes the proof. \square

7. PROOF OF MAIN THEOREM

In this section we will prove Theorem A and Corollary B.

By definition, the asymptotic rank of a CAT(0) space X equals the supremum of all n for which there exist sequences of positive numbers $\lambda_k \rightarrow \infty$, $\epsilon_k \rightarrow 0$, and maps ϕ_k from the Euclidean unit ball $B^n \subset \mathbb{R}^n$ into X with

$$|\lambda_k^{-1}d(\phi_k(y), \phi_k(y')) - \|y - y'\|| \leq \epsilon_k$$

for all $y, y' \in B^n$. This means that the sets $\phi_k(B^n)$, equipped with (the restrictions of) the rescaled metrics $\lambda_k^{-1}d$, converge in the Gromov–Hausdorff topology to B^n .

Recall further that ω_n denotes the Lebesgue measure of B^n . We let

$$\mathbf{c}_n := \frac{\text{vol}_n(B^n)}{\text{vol}_{n-1}(S^{n-1})^{n/(n-1)}} = \frac{\omega_n}{(n\omega_n)^{n/(n-1)}} = \frac{1}{(n^n\omega_n)^{1/(n-1)}}$$

denote the constant in the optimal isoperimetric inequality in \mathbb{R}^n . In particular $\mathbf{c}_2 = 1/(4\pi)$ and $\mathbf{c}_3 = 1/(6\sqrt{\pi})$.

We start with the proof of the implication (1) \Rightarrow (3) in Theorem A, which holds more generally in all dimensions. This uses similar arguments as in [Wen05, Sect. 4].

Proposition 7.1. *Let X be a proper CAT(0) space. Suppose that there exist constants $c < \mathbf{c}_{n+1}$ and $v > 0$ such that every Lipschitz n -sphere $S \subset X$ with $\text{vol}_n(S) > v$ bounds a Lipschitz $(n+1)$ -ball $B \subset X$ with $\text{vol}_{n+1}(B) \leq c \cdot \text{vol}_n(S)^{(n+1)/n}$. Then the asymptotic rank of X is at most n .*

Proof. Suppose for contradiction that the asymptotic rank of X is at least $n+1$. Then there exists a sequence of $(1 + \epsilon_k)$ -bi-Lipschitz embeddings

$$\phi_k : B^{n+1} \cap (\epsilon_k \mathbb{Z})^{n+1} \rightarrow X_k := (X, \lambda_k^{-1}d),$$

where $\lambda_k \rightarrow \infty$ and $\epsilon_k \rightarrow 0$. By the generalized Kirszbraun theorem [LS97], every ϕ_k can be extended to a $(1 + \epsilon_k)$ -Lipschitz map $\varphi_k : B^{n+1} \rightarrow X_k$. Define $S_k := (\varphi_k)_\# \partial[B^{n+1}] \in \mathbf{I}_{n,c}(X_k)$, and note that $\text{spt}(S_k) \subset \varphi_k(B^{n+1}) \subset B(x_k, 2)$ for $x_k := \varphi_k(0)$. By Theorem 2.8 there exists a minimizing filling $V_k \in \mathbf{I}_{n+1,c}(X_k)$ of S_k with support in $B(x_k, 2)$. It follows from the assumption that for sufficiently large k ,

$$\mathbf{M}(V_k) \leq c \cdot \mathbf{M}(S_k)^{(n+1)/n} \leq c(1 + \epsilon_k)^{n+1} \cdot \text{vol}_n(S^n)^{(n+1)/n}.$$

Since the φ_k are uniformly Lipschitz, and by the lower density bound for V_k , the sets $K_k := \varphi_k(B^{n+1}) \cup \text{spt}(V_k)$ are equi-compact. Hence, by Gromov's theorem [Gro81], they admit isometric embeddings $\psi_k : K_k \rightarrow Z$ into a single compact metric space Z . After passing to subsequences, we may assume that the $\psi_k(K_k)$ converge to a compact set $K \subset Z$ with respect to the Hausdorff distance in Z , the $(\psi_k)_\# V_k$ converge weakly to a current $V \in \mathbf{I}_{n+1,c}(Z)$ with support in K (recall Theorem 2.7), and the $\psi_k \circ \varphi_k$ converge uniformly to an isometric embedding $\iota : B^{n+1} \rightarrow Z$ with image in K . This implies that $(\psi_k \circ \varphi_k)_\# [B^{n+1}] \rightarrow T := \iota_\# [B^{n+1}] \in \mathbf{I}_{n+1,c}(Z)$ weakly. Note that

$\partial T = \partial V$, as $\partial(\psi_k \circ \varphi_k)_\# \llbracket B^{n+1} \rrbracket = (\psi_k)_\# S_k = \partial(\psi_k)_\# V_k$. By the lower semi-continuity of mass under weak convergence, and since $c < \mathbf{c}_{n+1}$, we have

$$\mathbf{M}(V) \leq \liminf_{k \rightarrow \infty} \mathbf{M}(V_k) \leq c \cdot \text{vol}_n(S^n)^{(n+1)/n} < \omega_{n+1}.$$

Note that K is the Gromov–Hausdorff limit of the sets $K_k \subset X_k$ and thus embeds isometrically into any ultralimit (X_ω, x_ω) of the (X_k, x_k) . Since (X_ω, x_ω) is a CAT(0) space, and the image of $\iota(B^{n+1})$ is a closed convex subset, we infer that there exists a 1-Lipschitz retraction $\pi: K \rightarrow \iota(B^{n+1})$. The cycle $\pi_\# V - T$ vanishes, because any $(n+2)$ -dimensional filling in $\iota(B^{n+1})$ is zero. Hence, $\mathbf{M}(T) = \mathbf{M}(\pi_\# V) \leq \mathbf{M}(V) < \omega_{n+1}$. This contradicts the fact that $\mathbf{M}(T) = \mathbf{M}(\iota_\# \llbracket B^{n+1} \rrbracket) = \omega_{n+1}$. \square

It is clear that (2) implies (1) in Theorem A. The implication (3) \Rightarrow (2) is more elaborate and we need some preparation. The following lemma allows us to deform a general Lipschitz disc to a *piecewise minimal* Lipschitz disc, while keeping control on boundary length and area.

Lemma 7.2. *Let X be a CAT(0) space and $\varphi: D \rightarrow X$ a Lipschitz disc with $\text{area}(\varphi) \leq a \cdot t^2$ and $\text{length}_\varphi(\partial D) \leq t$ for some constants $a, t > 0$. Let $V = \{v_1, \dots, v_n\} \subset \partial D$ for $n \geq 1$ be a cyclically ordered set and denote by $e_i \subset \partial D$ the arc between v_i and v_{i+1} , where $v_{n+1} := v_1$. Moreover, let $D_i \subset D$ denote the closed disc bounded by e_i and the geodesic $v_i v_{i+1}$. Let $\psi: D \rightarrow X$ be a Lipschitz disc such that*

- ψ agrees with φ on V ;
- ψ maps the whole disc D_i to the geodesic in X from $\varphi(v_i)$ to $\varphi(v_{i+1})$;
- ψ maps every geodesic $v_1 v_i$ to the geodesic in X from $\varphi(v_1)$ to $\varphi(v_i)$;
- ψ restricts to a minimal triangle T_i on every triangle $v_1 v_i v_{i+1} \subset D$.

Then there exists a Lipschitz homotopy $h: S^1 \times [0, 1] \rightarrow X$ from $\varphi|_{S^1}$ to $\psi|_{S^1}$ with $\text{area}(h) \leq \frac{1}{\pi} \cdot t^2$, and the discs φ and ψ together with the homotopy h form a Lipschitz 2-sphere $\sigma: S^2 \rightarrow X$ with $\text{area}(\sigma) \leq (a+1) \cdot t^2$.

Proof. For $i = 1, \dots, n$, let α_i denote the loop composed of $\varphi|_{e_i}$ and the geodesic in X from $\varphi(v_{i+1})$ to $\varphi(v_i)$. To get h , we use Proposition 3.1 to fill each α_i by a Lipschitz disc of area


$$a_i \leq \frac{1}{4\pi} \cdot \text{length}_\varphi(\alpha_i)^2 \leq \frac{1}{\pi} \cdot \text{length}_\varphi(e_i)^2.$$

The union of these discs defines h and we have

$$\text{area}(h) \leq \sum_{i=1}^n a_i \leq \frac{1}{\pi} \cdot \text{length}_\varphi(\partial D)^2 \leq \frac{t^2}{\pi}.$$

Furthermore, the Lipschitz disc ψ satisfies

$$\text{area}(\psi) \leq \sum_{i=1}^n \text{area}(T_i) \leq \sum_{i=1}^n \frac{t}{2} \cdot \text{length}_\varphi(e_i) \leq \frac{t^2}{2}.$$

FIGURE 4. Decomposition of D .

Hence, the Lipschitz 2-sphere σ formed by ϕ and ψ together with h has

$$\text{area}(\sigma) \leq \left(a + \frac{1}{2} + \frac{1}{\pi}\right) \cdot t^2 < (a + 1) \cdot t^2$$

as required. \square

Now we are ready for the proof of the implication (3) \Rightarrow (2) in Theorem A.

Theorem 7.3. *Let X be a proper CAT(0) space of asymptotic rank at most 2. For every $\delta > 0$, there exists a constant C such that every Lipschitz 2-sphere $S \subset X$ extends to a Lipschitz 3-ball $B \subset X$ with volume*

$$\text{vol}(B) \leq C \cdot \text{area}(S)^{1+\delta}.$$

Proof. Suppose for contradiction that there exists $\delta \in (0, 1/2)$ such that for every $M > 0$ we find a Lipschitz 2-sphere $f: S^2 \rightarrow X$ with

$$\text{Fillvol}(f) \geq M \cdot \text{area}(f)^{1+\delta}.$$

Let $c := c(\delta) \geq 1$ and $\epsilon := \epsilon(\delta, 1) \in (0, 1]$ be the constants from Lemma 6.2 for $\theta = 1$. We choose f to be (M, δ, ϵ) -minimal, for some (large) M , to be determined later. By the lemma, f is intrinsically $(c, 1)$ -isoperimetric. Since $\text{Fillvol}(f) \leq C \cdot \text{area}(f)^{1+1/2}$ by Theorem 3.2, we have

$$\text{area}(f) \geq \left(\frac{M}{C}\right)^{2/(1-2\delta)}.$$

To arrive at a contradiction we will construct a filling of f by a Lipschitz ball of small volume. To do this, we will homotop f to a Lipschitz sphere consisting of a controlled number of minimal triangles, which we then fill using the linear isoperimetric inequality for minimal tetrahedra.

By Proposition 6.5, for $\text{area}(f) \geq 4$ and $s > 0$ with $\frac{1}{3c} \leq s^2 \leq \frac{1}{8c} \text{area}(f)$, we find a Lipschitz 2-sphere $f': S^2 \rightarrow X$ with $\text{area}(f') \leq \text{area}(f)$ and a

Lipschitz homotopy h from f to f' with

$$\text{vol}(h) \leq 12M \cdot c \cdot s^{2\delta} \cdot \text{area}(f);$$

moreover, there exists a disc decomposition $S^2 = D_1 \cup \dots \cup D_n$, induced by a graph $\Gamma \subset S^2$, such that

$$n \leq \frac{14 \text{area}(f)}{s^2}, \quad \text{length}_{f'}(\partial D_i) \leq 34s, \quad \text{area}_{f'}(D_i) \leq 3c \cdot (34s)^2$$

for $i = 1, \dots, n$. The value of s will be specified later. Applying Lemma 7.2 to each of the Lipschitz discs $f'|_{D_i}$ and the corresponding set of vertices on ∂D_i , we obtain Lipschitz 2-spheres σ_i with

$$\text{area}(\sigma_i) \leq 4c \cdot (34s)^2 \leq \frac{3}{4} \cdot \text{area}(f) \leq \text{area}(f) - \epsilon,$$

assuming that $6200c \cdot s^2 \leq \text{area}(f)$. Thus, since f is (M, δ, ϵ) -minimal,

$$\text{Fillvol}(\sigma_i) \leq M \cdot \text{area}(\sigma_i)^{1+\delta}.$$

We can arrange that the Lipschitz homotopies $\partial D_i \times [0, 1] \rightarrow X$ provided by Lemma 7.2 agree on every common edge of two adjacent discs. Filling each sphere σ_i by a Lipschitz 3-ball, we then obtain a Lipschitz homotopy $h': S^2 \times [0, 1] \rightarrow X$ from f' to a Lipschitz 2-sphere \bar{f} such that

$$\text{vol}(h') \leq \frac{14 \text{area}(f)}{s^2} \cdot M \cdot (4 \cdot 34^2)^{3/2} \cdot c^2 \cdot s^{2+2\delta}.$$

Combining h and h' , we get a Lipschitz homotopy \bar{h} from f to \bar{f} with volume

$$\text{vol}(\bar{h}) \leq \text{vol}(h) + \text{vol}(h') \leq 5 \cdot 10^6 c^2 \cdot M \cdot s^{2\delta} \cdot \text{area}(f).$$

By construction, the Lipschitz sphere \bar{f} is piecewise minimal, that is, composed of finitely many minimal triangles spanned by geodesic triangles T_1, \dots, T_m in X , each with vertices in $f'(\partial D_i)$ for some i and, hence, circumference at most $34s$. Moreover, every triangle T_l has one side corresponding to an edge in Γ , and so the number of triangles m is bounded by twice the number E of edges in Γ . By Euler's formula, we have $E \leq 3n$ and therefore $m \leq 6n$.

Choose a point $o \in f'(\Gamma)$. It follows from Proposition 6.5 that

$$\text{diam}(f'(\Gamma)) \leq \frac{1}{2} \cdot \text{length}_{f'}(\Gamma) \leq \frac{119 \cdot \text{area}(f)}{s}.$$

Hence, by taking the geodesics from o to the vertices of T_l and inserting minimal triangles, we obtain a minimal tetrahedron τ_l with

$$\text{area}(\tau_l) \leq 17 \cdot 119 \cdot \text{area}(f) + (17s)^2 \leq 2024 \cdot \text{area}(f).$$

(recall that we have added the assumption $6200c \cdot s^2 \leq \text{area}(f)$). By Theorem 4.3, we have

$$\text{Fillvol}(\tau_l) \leq 2024\mu \cdot \text{area}(f).$$

Again, since the number m of tetrahedra is at most $6n$, we obtain a Lipschitz 3-ball $\Phi: B^3 \rightarrow X$ filling \bar{f} with

$$\text{vol}(\Phi) \leq 2 \cdot 10^5 \mu \cdot \frac{\text{area}(f)^2}{s^2}.$$

Taking the homotopy \bar{h} into account, we get a Lipschitz 3-ball $\Psi: B^3 \rightarrow X$ filling f with

$$\begin{aligned} \text{vol}(\Psi) &\leq \text{vol}(\bar{h}) + \text{vol}(\Phi) \\ &\leq 5 \cdot 10^6 c^2 \cdot M \cdot s^{2\delta} \cdot \text{area}(f) + 2 \cdot 10^5 \mu \cdot \frac{\text{area}(f)^2}{s^2}. \end{aligned}$$

Now we choose $s = \text{area}(f)^{(1-\delta)/2}$ and arrive at

$$\text{Fillvol}(f) \leq \left(\frac{5 \cdot 10^6 c^2}{\text{area}(f)^{\delta^2}} + \frac{2 \cdot 10^5 \mu}{M} \right) \cdot M \cdot \text{area}(f)^{1+\delta}.$$

Recall that $\text{area}(f) \geq (M/C)^{2/(1-2\delta)}$. Thus, if

$$M > \max \left\{ C \cdot (10^7 c^2)^{(1-2\delta)/(2\delta^2)}, 4 \cdot 10^5 \mu \right\},$$

then $\text{Fillvol}(f) < M \cdot \text{area}(f)^{1+\delta}$, in contradiction to the assumption. \square

We conclude this section with

Proof of Corollary B. We fix $\delta > 0$ and prove the claim by induction on g where the case $g = 0$ is the content of Theorem A. Now suppose we know the result for Lipschitz surfaces of genus at most $g - 1$. We equip every closed surface Σ with a Riemannian metric induced by an embedding $\iota: \Sigma \hookrightarrow \mathbb{R}^3$. Let $f: \Sigma_g \rightarrow X$ be a closed Lipschitz surface of genus g . For $\epsilon > 0$ consider the related bilipschitz map $f_\epsilon: \Sigma_g \rightarrow X \times \mathbb{R}^3$ with $f_\epsilon(x) = (f(x), \epsilon \iota(x))$. Choose ϵ small enough such that $\text{area}(f_\epsilon) \leq 2 \text{area}(f)$. We will apply the systolic inequality for closed surfaces [Gro83, Gro96]. Since Σ_g equipped with the induced path metric d_{f_ϵ} is bilipschitz to the Riemannian manifold Σ_g , the systolic inequality holds for $(\Sigma_g, d_{f_\epsilon})$. Thus, we find a non-contractible non-separating simple closed curve $\alpha \subset \Sigma_g$ such that

$$\text{length}_{f_\epsilon}(\alpha) \leq \sigma_g \cdot \sqrt{\text{area}(f_\epsilon)}$$

where σ_g denotes the systolic ratio. By the isoperimetric inequality, we can fill $f_\epsilon \circ \alpha$ by a disc $D \subset X \times \mathbb{R}^3$ with $\text{area}(D) \leq \frac{1}{4\pi} \cdot \text{length}(f_\epsilon \circ \alpha)^2$. Inserting the disc D cuts Σ_g into a surface Σ'_ϵ of genus $g - 1$ with

$$\text{area}_{f_\epsilon}(\Sigma'_\epsilon) = \text{area}_{f_\epsilon}(\Sigma) + \frac{\sigma_g^2}{2\pi} \cdot \text{area}_{f_\epsilon}(\Sigma).$$

We denote by Σ' the projection of Σ'_ϵ to X . By induction, we obtain a Lipschitz handlebody $\bar{f}: H_{g-1} \rightarrow X$ filling Σ' such that

$$\text{vol}(H_{g-1}) \leq C_{g-1} \cdot \text{area}(\Sigma')^{1+\delta}.$$

Now H_{g-1} can be Lipschitz parametrized by a genus g handlebody H_g filling Σ_g . We obtain

$$\begin{aligned} \operatorname{vol}(H_g) &= \operatorname{vol}(H_{g-1}) \leq C_{g-1} \cdot \operatorname{area}(\Sigma')^{1+\delta} \\ &\leq C_{g-1} \cdot \operatorname{area}(\Sigma'_\epsilon)^{1+\delta} \\ &\leq C_{g-1} \left(2 + \frac{\sigma_g^2}{\pi}\right)^{1+\delta} \cdot \operatorname{area}(\Sigma)^{1+\delta}. \end{aligned}$$

This proves the claim with $C_g := C_{g-1} \left(2 + \frac{\sigma_g^2}{\pi}\right)^{1+\delta}$. \square

REFERENCES

- [ABC13] G. Alberti, S. Bianchini, and G. Crippa. Structure of level sets and Sard-type properties of Lipschitz maps. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 12(4):863–902, 2013.
- [AK00] L. Ambrosio and B. Kirchheim. Currents in metric spaces. *Acta Math.*, 185(1):1–80, 2000.
- [Bal95] W. Ballmann. *Lectures on spaces of nonpositive curvature*, volume 25 of *DMV Seminar*. Birkhäuser Verlag, Basel, 1995. With an appendix by Misha Brin.
- [BGS85] W. Ballmann, M. Gromov, and V. Schroeder. *Manifolds of non-positive curvature*, volume 61 of *Progress in Math.* Birkhauser, 1985.
- [BH99] M. Bridson and A. Haefliger. *Metric spaces of non-positive curvature*, volume 319 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin, 1999.
- [Bow95] B. H. Bowditch. A short proof that a subquadratic isoperimetric inequality implies a linear one. *Michigan Math. J.*, 42(1):103–107, 1995.
- [BWY23] G. Basso, S. Wenger, and R. Young. Undistorted fillings in subsets of metric spaces. *Adv. Math.*, 423:Paper No. 109024, 54, 2023.
- [CDMW18] G. R. Chambers, D. Dotterer, F. Manin, and S. Weinberger. Quantitative null-cobordism. *J. Amer. Math. Soc.*, 31(4):1165–1203, 2018. With an appendix by Manin and Weinberger.
- [CR22] P. Creutz and M. Romney. Triangulating metric surfaces. *Proc. Lond. Math. Soc. (3)*, 125(6):1426–1451, 2022.
- [Cre22] P. Creutz. Plateau’s problem for singular curves. *Comm. Anal. Geom.*, 30(8):1779–1792, 2022.
- [Des16] D. Descombes. Asymptotic rank of spaces with bicomings. *Math. Z.*, 284(3-4):947–960, 2016.
- [Dru01] C. Druțu. Cônes asymptotiques et invariants de quasi-isométrie pour des espaces métriques hyperboliques. *Ann. Inst. Fourier (Grenoble)*, 51(1):81–97, 2001.
- [ECH⁺92] David B. A. Epstein, James W. Cannon, Derek F. Holt, Silvio V. F. Levy, Michael S. Paterson, and William P. Thurston. *Word processing in groups*. Jones and Bartlett Publishers, Boston, MA, 1992.
- [GL23] T. Goldhirsch and U. Lang. Characterizations of higher rank hyperbolicity. *Math. Z.*, 305(1):Paper No. 13, 26, 2023.
- [Gol22] T. Goldhirsch. Lipschitz chain approximation of metric integral currents. *Anal. Geom. Metr. Spaces*, 10(1):40–49, 2022.
- [Gro81] M. Gromov. Groups of polynomial growth and expanding maps. *Inst. Hautes Études Sci. Publ. Math.*, 53:53–73, 1981.
- [Gro83] M. Gromov. Filling Riemannian manifolds. *J. Differential Geom.*, 18(1):1–147, 1983.

- [Gro87] M. Gromov. Hyperbolic groups. In *Essays in group theory*, volume 8 of *Math. Sci. Res. Inst. Publ.*, pages 75–263. Springer, New York, 1987.
- [Gro93] M. Gromov. Asymptotic invariants of infinite groups. In *Geometric group theory, Vol. 2 (Sussex, 1991)*, volume 182 of *London Math. Soc. Lecture Note Ser.*, pages 1–295. Cambridge Univ. Press, Cambridge, 1993.
- [Gro96] M. Gromov. Systoles and intersystolic inequalities. In *Actes de la Table Ronde de Géométrie Différentielle (Luminy, 1992)*, volume 1 of *Sémin. Congr.*, pages 291–362. Soc. Math. France, Paris, 1996.
- [Gro99a] M. Gromov. *Metric structures for Riemannian and non-Riemannian spaces*, volume 152 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 1999. Based on the 1981 French original [MR0682063 (85e:53051)], With appendices by M. Katz, P. Pansu and S. Semmes, Translated from the French by Sean Michael Bates.
- [Gro99b] M. Gromov. Quantitative homotopy theory. In *Prospects in mathematics (Princeton, NJ, 1996)*, pages 45–49. Amer. Math. Soc., Providence, RI, 1999.
- [Hei01] J. Heinonen. *Lectures on analysis on metric spaces*. Universitext. Springer-Verlag, New York, 2001.
- [Isl23] H. Isleifsson. Linear isoperimetric inequality for homogeneous hadamard manifolds. *J. Topol. Anal.*, 2023.
- [Kir94] B. Kirchheim. Rectifiable metric spaces: local structure and regularity of the Hausdorff measure. *Proc. Amer. Math. Soc.*, 121(1):113–123, 1994.
- [KL20] B. Kleiner and U. Lang. Higher rank hyperbolicity. *Invent. Math.*, 221(2):597–664, 2020.
- [Kle99] B. Kleiner. The local structure of spaces with curvature bounded above. *Math. Z.*, 231:409–456, 1999.
- [KS93] N. Korevaar and R. Schoen. Sobolev spaces and harmonic maps for metric space targets. *Comm. Anal. Geom.*, 1(3-4):561–659, 1993.
- [Lan00] U. Lang. Higher-dimensional linear isoperimetric inequalities in hyperbolic groups. *Internat. Math. Res. Notices*, (13):709–717, 2000.
- [Lan11] U. Lang. Local currents in metric spaces. *J. Geom. Anal.*, 21(3):683–742, 2011.
- [LDR15] E. Le Donne and T. Rajala. Assouad dimension, Nagata dimension, and uniformly close metric tangents. *Indiana Univ. Math. J.*, 64(1):21–54, 2015.
- [Leu14] E. Leuzinger. Optimal higher-dimensional Dehn functions for some CAT(0) lattices. *Groups Geom. Dyn.*, 8(2):441–466, 2014.
- [LS97] U. Lang and V. Schroeder. Kirszbraun’s theorem and metric spaces of bounded curvature. *Geom. Funct. Anal.*, 7(3):535–560, 1997.
- [LS05] U. Lang and T. Schlichenmaier. Nagata dimension, quasisymmetric embeddings, and Lipschitz extensions. *Int. Math. Res. Not.*, (58):3625–3655, 2005.
- [LS24] A. Lytchak and S. Stadler. Curvature bounds of subsets in dimension two. *J. Differential Geom.*, 127(3):1245–1265, 2024.
- [LW17] A. Lytchak and S. Wenger. Area minimizing discs in metric spaces. *Arch. Ration. Mech. Anal.*, 223(3):1123–1182, 2017.
- [LW18a] A. Lytchak and S. Wenger. Intrinsic structure of minimal discs in metric spaces. *Geom. Topol.*, 22(1):591–644, 2018.
- [LW18b] A. Lytchak and S. Wenger. Isoperimetric characterization of upper curvature bounds. *Acta Math.*, 221(1):159–202, 2018.
- [LW24] A. Lytchak and S. Wagner. Curvature bounds on length-minimizing discs. *Geom. Dedicata*, 218(2):Paper No. 49, 21, 2024.
- [LWY20] A. Lytchak, S. Wenger, and R. Young. Dehn functions and Hölder extensions in asymptotic cones. *J. Reine Angew. Math.*, 763:79–109, 2020.
- [Mit19] A. Mitsuishi. The coincidence of the homologies of integral currents and of integral singular chains, via cosheaves. *Math. Z.*, 292(3-4):1069–1103, 2019.

- [MW23] F. Manin and S. Weinberger. Quantitative pl bordism. *arXiv:2311.16389*, 2023.
- [NR23] D. Ntalampekos and M. Romney. Polyhedral approximation of metric surfaces and applications to uniformization. *Duke Math. J.*, 172(9):1673–1734, 2023.
- [OI'91] A. Yu. Ol'shanskii. Hyperbolicity of groups with subquadratic isoperimetric inequality. *Internat. J. Algebra Comput.*, 1(3):281–289, 1991.
- [Pap95a] P. Papasoglu. On the sub-quadratic isoperimetric inequality. In *Geometric group theory (Columbus, OH, 1992)*, volume 3 of *Ohio State Univ. Math. Res. Inst. Publ.*, pages 149–157. de Gruyter, Berlin, 1995.
- [Pap95b] P. Papasoglu. Strongly geodesically automatic groups are hyperbolic. *Invent. Math.*, 121(2):323–334, 1995.
- [Pap96] P. Papasoglu. On the asymptotic cone of groups satisfying a quadratic isoperimetric inequality. *J. Differential Geom.*, 44(4):789–806, 1996.
- [Pap09] P. Papasoglu. Cheeger constants of surfaces and isoperimetric inequalities. *Trans. Amer. Math. Soc.*, 361(10):5139–5162, 2009.
- [PS19] A. Petrunin and S. Stadler. Metric-minimizing surfaces revisited. *Geom. Topol.*, 23(6):3111–3139, 2019.
- [Reš68] Ju. G. Rešetnjak. Non-expansive maps in a space of curvature no greater than K . *Sibirsk. Mat. Ž.*, 9:918–927, 1968.
- [RS09] C. Riedweg and D. Schäppi. Singular (Lipschitz) homology and homology of integral currents. *arXiv:0902.3831*, 2009.
- [Sch20] F. Schulze. Optimal isoperimetric inequalities for surfaces in any codimension in Cartan-Hadamard manifolds. *Geom. Funct. Anal.*, 30(1):255–288, 2020.
- [Sta21] S. Stadler. The structure of minimal surfaces in CAT(0) spaces. *J. Eur. Math. Soc. (JEMS)*, 23(11):3521–3554, 2021.
- [SW24] S. Stadler and S. Wenger. Isoperimetric inequalities vs. upper curvature bounds. *arXiv:2309.11329*, to appear in *Geom. Topol.*, 2024.
- [Tuk80] P. Tukia. The planar Schönflies theorem for Lipschitz maps. *Ann. Acad. Sci. Fenn. Ser. A I Math.*, 5(1):49–72, 1980.
- [Wen05] S. Wenger. Isoperimetric inequalities of Euclidean type in metric spaces. *Geom. Funct. Anal.*, 15(2):534–554, 2005.
- [Wen06a] S. Wenger. Filling invariants at infinity and the Euclidean rank of Hadamard spaces. *Int. Math. Res. Not.*, pages Art. ID 83090, 33, 2006.
- [Wen06b] S. Wenger. Filling invariants at infinity and the Euclidean rank of Hadamard spaces. *Int. Math. Res. Not.*, pages Art. ID 83090, 33, 2006.
- [Wen08a] S. Wenger. Gromov hyperbolic spaces and the sharp isoperimetric constant. *Invent. Math.*, 171(1):227–255, 2008.
- [Wen08b] S. Wenger. A short proof of Gromov's filling inequality. *Proc. Amer. Math. Soc.*, 136(8):2937–2941, 2008.
- [Wen11] S. Wenger. The asymptotic rank of metric spaces. *Comment. Math. Helv.*, 86(2):247–275, 2011.
- [Whi83] B. White. Existence of least-area mappings of N -dimensional domains. *Ann. of Math. (2)*, 118(1):179–185, 1983.
- [Whi84] B. White. Mappings that minimize area in their homotopy classes. *J. Differential Geom.*, 20(2):433–446, 1984.

MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, ANDREW WILES BUILDING,
WOODSTOCK ROAD, OXFORD OX2 6GG, UK

Email address: `drutu@maths.ox.ac.uk`

DEPARTMENT OF MATHEMATICS, ETH ZÜRICH, RÄMISTRASSE 101, 8092 ZÜRICH,
SWITZERLAND

Email address: `lang@math.ethz.ch`

MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, ANDREW WILES BUILDING,
WOODSTOCK ROAD, OXFORD OX2 6GG, UK

Email address: `papazoglou@maths.ox.ac.uk`

MAX PLANCK INSTITUTE FOR MATHEMATICS, VIVATSGASSE 7, 53111 BONN, GER-
MANY

Email address: `stadler@mpim-bonn.mpg.de`