

THE LAGRANGIAN APPROACH TO THE COMPRESSIBLE PRIMITIVE EQUATIONS

MATTHIAS HIEBER, YOSHIKI IIDA, ARNAB ROY, AND TAREK ZÖCHLING

ABSTRACT. This article develops the hydrostatic Lagrangian approach to the compressible primitive equations. A fundamental aspect in the analysis is the investigation of the compressible hydrostatic Lamé and Stokes operators. Local strong well-posedness for large data and global strong well-posedness for small data are established under various assumptions on the pressure law, both in the presence and absence of gravity.

1. INTRODUCTION

The primitive equations of geophysical fluid dynamics are regarded as a standard model for oceanic and atmospheric dynamics and are derived from the Navier-Stokes equations by assuming a hydrostatic balance for the pressure term. They have been introduced in a series of papers by Lions, Temam and Wang [23, 24, 25].

Very roughly speaking, the model developed by Lions, Temam and Wang couples the compressible primitive equations describing the dynamics of the atmosphere and the incompressible primitive equations for the dynamics of the ocean through a nonlinear interface condition. In the analysis of the compressible equation for the atmosphere they introduced a new coordinate system, the pressure coordinate system, in which, assuming the pressure of the atmosphere at the interface to be constant, one essentially obtains an incompressible primitive equation. This assumption is, however, not physical and hence not true in realistic situations. In their pioneering work Lions, Temam and Wang proved the existence of a weak solution to the primitive equations and also to the coupled system; its uniqueness remains an open problem for both systems until today.

Since then the compressible and incompressible primitive equations have been the subject of intensive mathematical investigations. Indeed, a celebrated result of Cao and Titi [4] asserts that the incompressible primitive equations subject to Neumann boundary conditions are globally strongly well-posed in the three-dimensional setting for arbitrarily large data belonging to H^1 . For related results concerning different approaches, different boundary conditions and also critical spaces, we refer to [17, 19, 22].

The situation for the compressible primitive equations is much less understood. Given a cylindrical domain $\Omega = G \times (0, 1)$, where $G = (0, 1)^2$ the compressible primitive equations are given by the set of equations

$$(1.1) \quad \left\{ \begin{array}{ll} \partial_t \varrho + \operatorname{div}(\varrho u) = 0, & \text{on } \Omega \times (0, T), \\ \varrho(\partial_t v + u \cdot \nabla v) - \mu \Delta v - \mu' \nabla_{\mathbb{H}} \operatorname{div}_{\mathbb{H}} v + \nabla_{\mathbb{H}} p = 0, & \text{on } \Omega \times (0, T), \\ \partial_z p = -g \varrho, & \text{on } \Omega \times (0, T), \\ p = \varrho^\gamma, & \text{on } \Omega \times (0, T), \\ v(0) = v_0, \quad \varrho(0) = \varrho_0 & \end{array} \right.$$

for some $\gamma \geq 1$. Here $u = (v, w) : \Omega \rightarrow \mathbb{R}^3$ denotes the velocity, $p : \Omega \rightarrow \mathbb{R}$ the atmospheric pressure and $\varrho : \Omega \rightarrow \mathbb{R}$ the atmospheric density. Here we use $\nabla_{\mathbb{H}}$, $\operatorname{div}_{\mathbb{H}}$ for the horizontal gradient and the horizontal divergence, that is $\nabla_{\mathbb{H}} := (\partial_x, \partial_y)^T$ and $\operatorname{div}_{\mathbb{H}} := \nabla_{\mathbb{H}} \cdot$. The Lamé coefficients μ and μ' are assumed to satisfy the standard conditions $\mu > 0$ and $\mu + \mu' > 0$.

The above system is supplemented with the boundary conditions

$$(1.2) \quad v|_{G \times \{1\}} = 0, \quad (\partial_z v)|_{G \times \{0\}} = 0, \quad w|_{G \times \{0\} \cup G \times \{1\}} = 0 \quad \text{and} \quad u, p, \varrho \text{ are periodic on } \partial G \times [0, 1].$$

The case $\gamma = 1$ is of particular interest. Indeed, the law of an ideal gas tells us that $p = R\rho T$, where T denotes the temperature of the atmosphere and R the gas constant. This gives us in the isothermal situation the relation $p = c\rho$ for some constant c . The articles of Lions, Temam and Wang [23], [24] also treat this situation. For both classical and recent results on the compressible Navier-Stokes equations, we refer to the works of Lions [26] and Feireisl [15] for weak solutions, and those of Matsumura and Nishida [30] and Enomoto and Shibata [12] for strong solutions. In particular, for the case of the exponent $\gamma = 1$, we refer to the work of Hoff [20] and Danchin and Mucha [10].

The existence of global weak solutions for the compressible system with degenerate viscosities without gravity goes back to the work of Liu and Titi [27]. Ersoy, Ngom and Sy also constructed a global weak solution to some two dimensional version of compressible primitive equations in [14] and [13]. Local strong well-posedness in the three dimensional setting with $\gamma > 1$ was established by Liu and Titi [29] in two cases: with gravity but without vacuum and with vacuum but without gravity. Furthermore, they showed in [28] a global, strong well-posedness result for the compressible primitive equations for data close to an incompressible flow in the situation without gravity. Note that all these results are based on energy methods.

Inspired by the Lagrangian approach to the compressible Navier-Stokes equations, see e.g., [8, 9], we introduce here for the first time the *hydrostatic Lagrangian approach* to the compressible primitive equations. It differs from the usual Lagrangian approach since we define here the flow along the vertical average \bar{v} of the horizontal velocity v , i. e., the flow X is defined as the solution of the equation

$$(1.3) \quad \begin{cases} \partial_t X(t, y_H) &= \bar{v}(t, X(t, y_H)), \\ X(0, y_H) &= y_H, \end{cases}$$

where y_H denotes the horizontal coordinate.

This hydrostatic Lagrangian approach allows us to prove first local, strong well-posedness of (1.1) subject to (1.2) including gravity and general values of γ for initial data $\varrho_0 \in H^{1,q}(\Omega)$ and $v_0 \in B_{qp}^{2(1-1/p)}(\Omega; \mathbb{R}^2)$ subject to suitable compatibility conditions and $p, q > 2$. Moreover, for the case $\gamma = 1$ we obtain a global strong well-posedness result for small data even in the situation of gravity.

Some words about our approach are in order. The hydrostatic Lagrangian transformation leads us to a new set of equations, where the linearized part is given by the *compressible, hydrostatic Stokes operator* A_{CHS} , see Section 6. We show that $-A_{\text{CHS}}$ admits a bounded \mathcal{H}^∞ -calculus on $H_{\text{per}}^{1,q}(G; \mathbb{R}) \times L^q(\Omega; \mathbb{R}^2)$ of an \mathcal{H}^∞ -angle strictly less than $\pi/2$. This allows us to perform a maximal regularity approach to equation (1.1) implying our local and global existence results even in the case of gravity. Here we use the fact that the compressible hydrostatic Stokes operator is invertible when restricted to the space of mean value free functions and that the maximal regularity estimates can be shown in this case to be valid on the time interval $[0, \infty)$.

The boundedness of the \mathcal{H}^∞ -calculus of $-A_{\text{CHS}}$ on $H_{\text{per}}^{1,q}(G; \mathbb{R}) \times L^q(\Omega; \mathbb{R}^2)$ is based on perturbation theory of the *hydrostatic Lamé operator* $-A_{\text{HL}}$ defined and investigated in Section 4. Indeed, we prove in Section 4 that $-A_{\text{HL}}$ is parameter elliptic on the cylindrical domain Ω . Combining the \mathcal{H}^∞ -result for parameter elliptic, cylindrical boundary value problems given in [31] with the perturbation result given in [18], we conclude that $-A_{\text{HL}}$ admits a bounded \mathcal{H}^∞ -calculus on $L^q(\Omega; \mathbb{R}^2)$ of angle strictly less than $\pi/2$.

This article is structured as follows. In Section 2 we present after some preliminaries the three main results of this article. The hydrostatic Lagrangian transform is introduced and investigated then in Section 3. In Section 4 we introduce and study the hydrostatic Lamé operator which leads in Section 5 to our local, strong well-posedness results. The proof of our global strong well-posedness results for small data to the compressible primitive equations is given in Section 6 in various settings.

2. PRELIMINARIES AND MAIN RESULTS

Consider a cylindrical domain $\Omega = G \times (0, 1)$, where $G = (0, 1)^2$. For an integrable function $f: \Omega \rightarrow \mathbb{R}^n$, with $n \in \{1, 2\}$, we define the vertical average of f by $\bar{f} := \int_0^1 f(\cdot, \cdot, \xi) \, d\xi$.

1. *The case $\gamma = 1$ with gravity*

We start with the case $\gamma = 1$ meaning that $p = c\rho$ for some $c > 0$. The pressure law is then given by the law of ideal gas, which reads as $p = RT\rho$, where T denotes temperature and $R > 0$ the gas constant. In the isothermal situation we obtain the relation $p = c\rho$. We consider the set of equations

$$(2.1) \quad \left\{ \begin{array}{ll} \partial_t \rho + \operatorname{div}(\rho u) = 0, & \text{on } \Omega \times (0, T), \\ \rho(\partial_t v + u \cdot \nabla v) - \mu \Delta v - \mu' \nabla_{\mathbb{H}} \operatorname{div}_{\mathbb{H}} v + \nabla_{\mathbb{H}} p = 0, & \text{on } \Omega \times (0, T), \\ \partial_z p = -g\rho, & \text{on } \Omega \times (0, T), \\ p = c\rho, & \text{on } \Omega \times (0, T), \\ v(0) = v_0, \quad \rho(0) = \rho_0 & \end{array} \right.$$

for some $c > 0$. The system is supplemented by the boundary conditions given in (1.2).

For simplicity of the notation we set $c = g = 1$ and deduce then from (2.1)₃ and (2.1)₄ that

$$(2.2) \quad p(t, x, y, z) = \rho(t, x, y, z) = \xi(t, x, y) e^{-z},$$

where ξ denotes the pressure evaluated at the surface $G \times \{0\}$. In the following, we use the explicit vertical dependence of the density to obtain a two dimensional version of the continuity equation by averaging in vertical direction. To this end, we consider the transformation

$$(2.3) \quad z' = \frac{1 - e^{-z}}{\delta} \iff z = \log\left(\frac{1}{1 - \delta z'}\right) \quad \text{for } \delta := 1 - e^{-1} > 0,$$

and introduce the new variables

$$\widehat{v}(t, x, y, z') := v(t, x, y, z(z')) \quad \text{as well as} \quad \widehat{w}(t, x, y, z') := w(t, x, y, z(z')).$$

Note that the surface pressure ξ remains unchanged and (2.2) implies that $p = \rho = \xi \cdot (1 - \delta z')$. Hence, naming the new variables as before and averaging the continuity equation in vertical direction, we find the transformed system

$$(2.4) \quad \left\{ \begin{array}{ll} \partial_t \xi + \bar{v} \cdot \nabla_{\mathbb{H}} \xi + \xi \operatorname{div}_{\mathbb{H}} \bar{v} = 0, & \text{on } \Omega \times (0, T), \\ \xi(\partial_t v + \bar{v} \cdot \nabla_{\mathbb{H}} v) - \frac{\mu \Delta_{\mathbb{H}} v}{1 - \delta z} - \mu \partial_z \left(\frac{1 - \delta z}{\delta^2} \partial_z v \right) - \frac{\mu' \nabla_{\mathbb{H}} \operatorname{div}_{\mathbb{H}} v}{1 - \delta z} = F(\xi, v), & \text{on } \Omega \times (0, T), \\ \partial_z \xi = 0, & \text{on } \Omega \times (0, T), \end{array} \right.$$

where F is given by

$$F(\xi, v) = -\xi((v - \bar{v}) \cdot \nabla_{\mathbb{H}} v + \frac{(1 - \delta z)}{\delta} w \partial_z v) - \nabla_{\mathbb{H}} \xi.$$

Note that the transform (2.3) leaves the boundary conditions (1.2) unchanged.

As for the incompressible primitive equations, where the vertical velocity is determined by the divergence free condition, see e. g. [19], we find that w is determined by the horizontal velocity v and the density at the surface ξ . More precisely, comparing (1.1)₁ and (2.4)₁ yields

$$(2.5) \quad \xi w = - \int_0^z \operatorname{div}_{\mathbb{H}}(\xi \tilde{v}) dz',$$

where $\tilde{v} = v - \bar{v}$. Finally, we prescribe the initial data as

$$(\xi(0), v(0)) = (\xi_0, v_0).$$

2. *The case $\gamma = 2$ with gravity*

We consider the system (1.1) for the case $p = \rho^2$ and set $g = 1$. In that case the relevant quantity for the density is still given by its value at the surface ξ . However, the representation of ρ by ξ differs from

the one given in (2.2). For $\gamma = 2$ we obtain $\varrho = \xi + \frac{1}{2}z$ and after averaging the system reads as

$$(2.6) \quad \begin{cases} \partial_t \xi + \bar{v} \cdot \nabla_{\mathbb{H}} \xi + \xi \operatorname{div}_{\mathbb{H}} \bar{v} + \frac{1}{2} \overline{z \operatorname{div}_{\mathbb{H}} v} = 0, & \text{on } G \times (0, T), \\ (\xi + \frac{1}{2}z)(\partial_t v + u \cdot \nabla v) - \mu \Delta v - \mu' \nabla_{\mathbb{H}} \operatorname{div}_{\mathbb{H}} v + (2\xi + z) \nabla_{\mathbb{H}} \xi = 0, & \text{on } \Omega \times (0, T), \\ \partial_z \xi = 0, & \text{on } \Omega \times (0, T). \end{cases}$$

The system is again supplemented by the boundary conditions given in (1.2). Here we can also recover the vertical velocity w by comparing (1.1)₁ and (2.6)₁ which leads to

$$\xi w = - \int_0^z \operatorname{div}_{\mathbb{H}}(\xi \bar{v}) + \frac{1}{2} z \widetilde{\operatorname{div}_{\mathbb{H}} v} dz'.$$

Finally, we prescribe the initial data as

$$(\xi(0), v(0)) = (\xi_0, v_0).$$

3. The case of general pressure law without gravity

We finally investigate the system (1.1) without gravity but with a generalized pressure law of the form

$$(2.7) \quad p = P(\varrho), \quad \text{where } P \in C^\infty \quad \text{with } c_1 \leq P'(s) \leq c_2 \text{ for all } s \in \operatorname{Im} \varrho,$$

where c_1 and c_2 are positive constants. The corresponding system reads then as

$$(2.8) \quad \begin{cases} \partial_t \varrho + \operatorname{div}(\varrho u) = 0, & \text{on } \Omega \times (0, T), \\ \varrho(\partial_t v + u \cdot \nabla v) - \mu \Delta v - \mu' \nabla_{\mathbb{H}} \operatorname{div}_{\mathbb{H}} v + \nabla_{\mathbb{H}} p = 0, & \text{on } \Omega \times (0, T), \\ \partial_z p = 0, & \text{on } \Omega \times (0, T), \\ p = P(\varrho), & \text{on } \Omega \times (0, T), \end{cases}$$

supplemented by the boundary conditions (1.2).

We deduce from (2.8)₃ as well as (2.8)₄ that $P'(\varrho) \cdot \partial_z \varrho = 0$. The general pressure law (2.7) then yields $\partial_z \varrho = 0$. In particular, the density is independent of the vertical variable and therefore it is sufficient to consider its value at the surface $\xi(t, x, y) := \varrho(t, x, y, 0)$. After averaging the continuity equation of (2.8), taking into account the boundary conditions for w , the above system reads as

$$(2.9) \quad \begin{cases} \partial_t \xi + \operatorname{div}_{\mathbb{H}}(\xi \bar{v}) = 0, & \text{on } G \times (0, T), \\ \xi(\partial_t v + u \cdot \nabla v) - \mu \Delta v - \mu' \nabla_{\mathbb{H}} \operatorname{div}_{\mathbb{H}} v + P'(\xi) \nabla_{\mathbb{H}} \xi = 0, & \text{on } \Omega \times (0, T), \\ \partial_z \xi = 0, & \text{on } \Omega \times (0, T). \end{cases}$$

Here, the vertical velocity can be deduced as in (2.5). The system (2.9) is supplemented by the boundary conditions (1.2) and by the initial conditions

$$(\xi(0), v(0)) = (\xi_0, v_0).$$

Remark 2.1. Let us note that the transformed systems (2.4) as well as (2.6) are equivalent to the original system (1.1), where the pressure and density are related by the formula $p = \varrho^\gamma$ for $\gamma \in \{1, 2\}$. In the case of $\gamma = 1$, the transform (2.3) is invertible for all times and the density can be recovered from (2.2). In the situation of $\gamma = 2$, we can recover the density from $\varrho = \xi + \frac{1}{2}z$ and in the case of no gravity it even holds true that $\varrho = \xi$. In all cases regularity and positivity properties transfer directly from ξ to ϱ as $z \in (0, 1)$.

At this point, some words about our approach for solving (2.4), (2.6) and (2.8) are in order. Let $s \in \mathbb{R}$ and $p, q \in (1, \infty)$. In the sequel, we denote by $W^{s,q}(\Omega)$ the fractional Sobolev spaces, by $H^{s,q}(\Omega)$ the Bessel potential spaces and by $B_{qp}^s(\Omega)$ the Besov spaces, where $\Omega \subset \mathbb{R}^3$ is an open. As usual, we set $H^{0,q}(\Omega) := L^q(\Omega)$ and note that $H^{s,q}(\Omega)$ coincides with the classical Sobolev space $W^{m,q}(\Omega)$ provided $s = m \in \mathbb{N}$. For more information on function spaces we refer e.g. to [3] and [33].

In the following, we need terminology to describe periodic boundary conditions on $\Gamma_l = \partial G \times [0, 1]$ as well as on ∂G . Given $s \in [0, \infty)$ and $p, q \in (1, \infty)$ we define the spaces

$$\mathbf{H}_{\text{per}}^{s,q}(\Omega) := \overline{\mathbf{C}_{\text{per}}^\infty(\overline{\Omega})}^{\|\cdot\|_{\mathbf{H}^{s,q}(\Omega)}} \quad \text{and} \quad \mathbf{H}_{\text{per}}^{s,q}(G) := \overline{\mathbf{C}_{\text{per}}^\infty(\overline{G})}^{\|\cdot\|_{\mathbf{H}^{s,q}(G)}},$$

where horizontal periodicity is modeled by the function spaces $\mathbf{C}_{\text{per}}^\infty(\overline{\Omega})$ and $\mathbf{C}_{\text{per}}^\infty(\overline{G})$ defined in [19]. Of course, we interpret $\mathbf{H}_{\text{per}}^{0,q}$ as L^q . The Besov spaces $\mathbf{B}_{qp,\text{per}}^s(\Omega)$ and $\mathbf{B}_{qp,\text{per}}^s(G)$ are defined in a similar manner. For more information regarding spaces equipped with periodic boundary conditions, see also [16, 17, 19]. Concerning the solution and data spaces we first define the spaces

$$\mathbf{X}_0 := \mathbf{H}_{\text{per}}^{1,q}(G; \mathbb{R}) \times L^q(\Omega; \mathbb{R}^2) \quad \text{and}$$

$$\mathbf{X}_1 := \mathbf{H}_{\text{per}}^{1,q}(G; \mathbb{R}) \times \mathbf{Y}, \quad \text{where } \mathbf{Y} := \{v \in \mathbf{H}_{\text{per}}^{2,q}(\Omega; \mathbb{R}^2) : v = 0 \text{ on } \Gamma_u \text{ and } \partial_z v = 0 \text{ on } \Gamma_b\},$$

where we $\Gamma_u = G \times \{1\}$ and $\Gamma_b = G \times \{0\}$. Then given $0 < T \leq \infty$, we define the data space $\mathbb{E}_{0,T}$ and the solution space $\mathbb{E}_{1,T}$ by

$$\mathbb{E}_{0,T} := L^p(0, T; \mathbf{X}_0) \quad \text{and} \quad \mathbb{E}_{1,T} := \mathbf{H}^{1,p}(0, T; \mathbf{X}_0) \cap L^p(0, T; \mathbf{X}_1).$$

Moreover, we introduce the solution and data spaces concerning only the velocity by

$$\mathbb{E}_{0,T}^v := L^p(0, T; L^q(\Omega; \mathbb{R}^2)), \quad \mathbb{E}_{1,T}^v = \mathbf{H}^{1,p}(0, T; L^q(\Omega; \mathbb{R}^2)) \cap L^p(0, T; \mathbf{Y}).$$

We now in the position to formulate our main theorems concerning the local well-posedness of (2.4) and (2.6) as well as the global well-posedness for small data of (2.4), even in the presence of gravity. Furthermore, in the absence of gravity, i. e., $g = 0$, we also obtain local and global well-posedness of (2.8), where the pressure and density are related by a monotone increasing function.

Our requirements on the set of initial data are collected in the following assumption.

Assumption (A): Let $p, q \in (2, \infty)$ such that $1 \neq \frac{1}{p} + \frac{1}{2q}$ and $\frac{1}{2} \neq \frac{1}{p} + \frac{1}{2q}$ and assume that

- (i) $\xi_0 \in \mathbf{H}_{\text{per}}^{1,q}(G; \mathbb{R})$ satisfies $M_1 \leq \xi_0 \leq M_2$ for some constants $M_1, M_2 > 0$ and all $(x, y) \in G$.
- (ii) $v_0 \in \mathbf{B}_{qp,\text{per}}^{2(1-1/p)}(\Omega; \mathbb{R}^2)$ satisfies $v_0|_{\Gamma_u} = 0$ if $\frac{1}{2q} + \frac{1}{p} < 1$ and $(\partial_z v_0)|_{\Gamma_b} = 0$ if $\frac{1}{2} + \frac{1}{2q} + \frac{1}{p} < 1$.

Theorem 2.2 (local strong well-posedness).

Let $T > 0$ and assume that (ξ_0, v_0) satisfy assumption (A). Then there exists $0 < a = a(v_0) \leq T$ such that

- (a) the system (2.4) admits a unique, strong solution $(\xi, v) \in \mathbb{E}_{1,a}$ satisfying

$$\xi \in \mathbf{H}^{1,p}(0, a; \mathbf{H}_{\text{per}}^{1,q}(G; \mathbb{R})), \quad v \in \mathbf{H}^{1,p}(0, a; L^q(\Omega, \mathbb{R}^2)) \cap L^p(0, a; \mathbf{Y})$$

and there exist constants $M_1^*, M_2^* > 0$ such that $M_1^* \leq \xi \leq M_2^*$ for all $t \in (0, a)$.

- (b) The assertion (a) holds true also for the system (2.6).

Next, we state our result concerning global well-posedness for small initial data.

Theorem 2.3 (global strong well-posedness for small data).

Assume that (ξ_0, v_0) satisfy assumption (A) and in addition that

$$\frac{1}{|G|} \int_G \xi_0 \, d(x, y) = \bar{\xi} > 0 \quad \text{and} \quad \|(\xi_0 - \bar{\xi}, v_0)\|_{\mathbf{H}^{1,q}(G) \times \mathbf{B}_{qp}^{2(1-1/p)}(\Omega)} \leq \delta$$

for some $\delta > 0$. Then there exists $\eta_0 > 0$ such that for all $\eta \in (0, \eta_0)$ there exists $\delta_0 > 0$ and $C > 0$ such that for all $\delta \in (0, \delta_0)$ the system (2.4) admits a unique, strong solution $(\xi, v) \in \mathbb{E}_{1,\infty}$ satisfying

$$\begin{aligned} & \|\xi - \bar{\xi}\|_{L^\infty(\mathbb{R}_+; \mathbf{H}^{1,q}(G))} + \|e^{\eta(\cdot)} \nabla_{\mathbf{H}} \xi\|_{\mathbf{H}^{1,p}(\mathbb{R}_+; L^q(G))} + \|e^{\eta(\cdot)} (\partial_t \xi)\|_{L^p(\mathbb{R}_+; L^q(G))} \\ & + \|e^{\eta(\cdot)} v\|_{L^p(\mathbb{R}_+; \mathbf{H}^{2,q}(\Omega))} + \|e^{\eta(\cdot)} (\partial_t v)\|_{L^p(\mathbb{R}_+; L^q(\Omega))} + \|e^{\eta(\cdot)} v\|_{L^\infty(\mathbb{R}_+; \mathbf{B}_{qp}^{2(1-1/p)}(\Omega))} \leq C\delta. \end{aligned}$$

Moreover, $\xi(t, x, y) \geq \bar{\xi}/2$ for all $t \in \mathbb{R}_+$ and all $(x, y) \in G$.

Finally, we obtain the following result dealing with the case of a general pressure law but without gravity.

Theorem 2.4 (local strong and global strong well-posedness for small data).

Assume that (ξ_0, v_0) satisfy assumption (A).

- (a) Given $T > 0$ there exists $0 < a = a(v_0) \leq T$ such that the system (2.9) admits a unique, strong solution $(\xi, v) \in \mathbb{E}_{1,a}$ satisfying $M_1^* \leq \xi \leq M_2^*$ for some constants $M_1^*, M_2^* > 0$ and all $t \in (0, a)$.
- (b) There exists $\eta_0 > 0$ such that for all $\eta \in (0, \eta_0)$ there exists $\delta_0 > 0$ and $C > 0$ such that for all $\delta \in (0, \delta_0)$ the system (2.9) admits a unique, strong solution $(\xi, v) \in \mathbb{E}_{1,\infty}$ satisfying

$$\begin{aligned} & \|\xi - \bar{\xi}\|_{L^\infty(\mathbb{R}_+; H^{1,q}(G))} + \|e^{\eta(\cdot)} \nabla_{\mathbb{H}} \xi\|_{H^{1,p}(\mathbb{R}_+; L^q(G))} + \|e^{\eta(\cdot)} (\partial_t \xi)\|_{L^p(\mathbb{R}_+; L^q(G))} \\ & + \|e^{\eta(\cdot)} v\|_{L^p(\mathbb{R}_+; H^{2,q}(\Omega))} + \|e^{\eta(\cdot)} (\partial_t v)\|_{L^p(\mathbb{R}_+; L^q(\Omega))} + \|e^{\eta(\cdot)} v\|_{L^\infty(\mathbb{R}_+; B_{qp}^{2(1-1/p)}(\Omega))} \leq C\delta. \end{aligned}$$

Moreover, $\xi(t, x, y) \geq \bar{\xi}/2$ for all $t \in \mathbb{R}_+$ and all $(x, y) \in G$.

Remark 2.5. Note that the results given in the three theorems above are formulated for the reformulated systems (2.4), (2.6) and (2.9). In fact, the density evaluated at the surface ξ is the relevant quantity for the describing the density ϱ . The above Remark 2.1 shows that this is equivalent to solving the original systems (1.1) and (2.8). Hence, the three well-posedness results stated above carry over to the original system (1.1) and (2.8)

Remark 2.6. By an approach similar to the one described for the case $\gamma = 2$, we verify that the assertions of Theorem 2.2(a) hold true also for the general power law $p = \varrho^\gamma$, where $\gamma \geq 1$. However, for simplicity of the presentation we prove here in detail only the cases $\gamma \in \{1, 2\}$.

Remark 2.7. (a) The total mass of the system (1.1) subject to $\gamma \geq 1$ or (2.8) is conserved, i. e., we have $\int_\Omega \rho = \int_\Omega \rho_0$.

(b) Consider the case with gravity and $\gamma = 1$. Define the energy $E(t)$ by

$$E(t) := \int_\Omega \left(\frac{1}{2} \xi |v|^2 + e(\xi) \right) \quad \text{with} \quad e(\xi) = \xi \log \xi + 1 - \xi.$$

Taking inner products of (2.4)₂ with v yields the energy balance

$$E(t) + \int_0^t \left(\mu \|(1 - \delta z)^{-\frac{1}{2}} \nabla_{\mathbb{H}} v\|_2^2 + \frac{\mu}{\delta^2} \|(1 - \delta z)^{\frac{1}{2}} \partial_z v\|_2^2 + \mu' \|(1 - \delta z)^{-\frac{1}{2}} \operatorname{div}_{\mathbb{H}} v\|_2^2 \right) = E(0),$$

for all $t \in (0, a]$, where $a > 0$ denotes the local existence time of the solution given in Theorem 2.2.

(c) In the case without gravity and the general pressure law (2.7) we define the energy $\tilde{E}(t)$ by

$$\tilde{E}(t) := \int_\Omega \left(\frac{1}{2} \xi |v|^2 + \tilde{e}(\xi) \right) \quad \text{with} \quad \tilde{e}(\xi) = \xi \int_1^\xi \frac{P(\xi)}{\xi^2} d\xi - P(1)(\xi - 1).$$

Then taking inner products of (2.9)₂ with v yields the energy balance

$$\tilde{E}(t) + \int_0^t \left(\mu \|\nabla v\|_2^2 + \mu' \|\operatorname{div}_{\mathbb{H}} v\|_2^2 \right) = \tilde{E}(0),$$

for all $t \in (0, a]$, where $a > 0$ denotes the local existence time given in Theorem 2.4

Remark 2.8. We may identify the domain $G = (0, 1)^2$ subject to periodic boundary conditions with the two dimensional torus \mathbb{T}^2 . Then the results described in the three theorem above hold true also for $\mathbb{T}^2 \times (0, 1)$. Furthermore, Theorem 2.2 as well as Remark 2.7 also hold true for $G = \mathbb{R}^2$. In that case the domain is given by a layer $\Omega_L := \mathbb{R}^2 \times (0, 1)$.

3. HYDROSTATIC LAGRANGIAN COORDINATES

The purpose of this transformation is to mitigate the hyperbolic effects in the continuity equation. We term it the ‘‘hydrostatic Lagrangian’’ as it defines the flow along \bar{v} , i.e., the solution X of the following differential equation

$$(3.1) \quad \begin{cases} \partial_t X(t, y_{\mathbb{H}}) = \bar{v}(t, X(t, y_{\mathbb{H}})), \\ X(0, y_{\mathbb{H}}) = y_{\mathbb{H}}, \end{cases}$$

where $y_H = (y_1, y_2) \in G$. The flow $X(t, \cdot) : G \rightarrow G$ is a well-defined C^1 -diffeomorphisms provided that \bar{v} is regular enough. Note that X is only a two dimensional flow. Solving (3.1) leads to

$$(3.2) \quad X(t, y_H) = y_H + \int_0^t \bar{v}(s, X(s, y_H)) ds,$$

for $y_H \in G$. Denoting by $Y(t, \cdot)$, the inverse of $X(t, \cdot)$, we find that

$$\nabla_H Y(t, X(t, y_H)) = [\nabla_H X]^{-1}(t, y_H).$$

We observe that the diffeomorphisms X and Y depend on \bar{v} and more precisely on v .

For X as in (3.2) and p' denoting the Hölder conjugate of p , we see due to $\|\bar{v}\|_{\mathbb{E}_{1,T}^v} \leq C\|v\|_{\mathbb{E}_{1,T}^v}$ that

$$(3.3) \quad \sup_{t \in [0, T]} \|\nabla_H X - \mathbb{I}_2\|_{H^{1,q}(G)} \leq C \int_0^T \|\nabla_H \bar{v}(t, \cdot)\|_{H^{1,q}(G)} \leq CT^{1/p'} \|v\|_{\mathbb{E}_{1,T}^v}$$

The embedding $W^{1,q}(G) \hookrightarrow L^\infty(G)$ yields $\sup_{t \in [0, T]} \|\nabla_H X - \mathbb{I}_2\|_{L^\infty(G)} \leq CT^{1/p'} \|v\|_{\mathbb{E}_{1,T}^v}$. In particular, there exists T' sufficiently small such that

$$(3.4) \quad \sup_{t \in [0, T']} \|\nabla_H X - \mathbb{I}_2\|_{L^\infty(G)} \leq \frac{1}{2}.$$

A Neumann series argument thus guarantees the invertibility of $\nabla_H X(t, \cdot)$ for all $t \in [0, T']$. Thus $\nabla_H Y(t, \cdot)$ exists on $[0, T']$. At the same time, we conclude by (3.2) that $\partial_t \nabla_H X(t, y_H) = \nabla_H \bar{v}(t, y_H)$. Therefore,

$$(3.5) \quad \|\partial_t \nabla_H X(t, \cdot)\|_{L^p(0, T; H^{1,q}(G))} \leq C\|v\|_{\mathbb{E}_{1,T}^v}.$$

In summary, there exists a constant $C > 0$ such that

$$(3.6) \quad \|\nabla_H X\|_{H^{1,p}(0, T; H^{1,q}(G))} + \|\nabla_H X\|_{L^\infty(0, T; H^{1,q}(G))} \leq C.$$

Since $p, q > 2$ the spaces $H^{1,p}(0, T; H^{1,q}(G))$ and $L^\infty(0, T; H^{1,q}(G))$ are Banach algebras. The estimate (3.6) yields the existence of a constant $C > 0$ such that

$$\begin{aligned} \|\det \nabla_H X\|_{W^{1,p}(0, T; H^{1,q}(G))} + \|\det \nabla_H X\|_{L^\infty(0, T; H^{1,q}(G))} &\leq C \text{ and} \\ \|\text{Cof} \nabla_H X\|_{W^{1,p}(0, T; H^{1,q}(G))} + \|\text{Cof} \nabla_H X\|_{L^\infty(0, T; H^{1,q}(G))} &\leq C. \end{aligned}$$

Thanks to (3.4) we find that $\det \nabla_H X \geq C > 0$ on $(0, T) \times G$ for some constant $C > 0$ provided $T \leq T'$. The representation

$$(3.7) \quad Z = [\nabla_H X]^{-1} = \frac{1}{\det \nabla_H X} (\text{Cof} \nabla_H X)^\top$$

then results in

$$\|Z\|_{H^{1,p}(0, T; H^{1,q}(G))} + \|Z\|_{L^\infty(0, T; H^{1,q}(G))} \leq C \text{ and } \|Z - \mathbb{I}_2\|_{L^\infty(0, T; H^{1,q}(G))} \leq CT^{1/p'}.$$

We now consider the change of variables defined by

$$V(t, y_H, z) := v(t, X(t, y_H), z), \quad W(t, y_H, z) := w(t, X(t, y_H), z), \quad \zeta(t, y_H) := \xi(t, X(t, y_H)).$$

Note that by this change of variables the initial values remain unchanged. The transformed equations in the case of $\gamma = 1$ then read as

$$(3.8) \quad \left\{ \begin{array}{ll} \partial_t \zeta + \xi_0 \operatorname{div}_H \bar{V} = F_1(\zeta, V), & \text{on } G \times (0, T), \\ \partial_t V - \frac{\mu \Delta_H V}{(1 - \delta z) \xi_0} - \frac{\mu}{\xi_0} \partial_z \left(\frac{1 - \delta z}{\delta^2} \partial_z V \right) - \frac{\mu' \nabla_H \operatorname{div}_H V}{(1 - \delta z) \xi_0} = F_2(\zeta, V), & \text{on } \Omega \times (0, T), \\ \partial_z \zeta = 0, & \text{on } \Omega \times (0, T), \\ V = 0, & \text{on } \Gamma_u \times (0, T), \\ \partial_z V = 0, & \text{on } \Gamma_b \times (0, T), \\ (\zeta(0), V(0)) = (\xi_0, v_0), & \end{array} \right.$$

where the right-hand sides F_1 and F_2 for $i = 1, 2$ are given by

$$(3.9) \quad \begin{aligned} F_1(\zeta, V) &= -(\zeta - \xi_0) \operatorname{div}_H \bar{V} - \zeta \nabla_H \bar{V} : [\mathbb{Z}^\top - \mathbb{I}_2], \\ (F_2(\zeta, V))_i &= \frac{\mu}{(1 - \delta z)\xi_0} \left(\sum_{l,j,k} \frac{\partial^2 V_i}{\partial y_l \partial y_k} (Z_{k,j} - \delta_{k,j}) Z_{l,j} + \sum_{l,k} \frac{\partial^2 V_i}{\partial y_l \partial y_k} (Z_{l,k} - \delta_{k,l}) + \sum_{l,j,k} Z_{l,k} \frac{\partial V_i}{\partial y_k} \frac{\partial Z_{k,j}}{\partial y_l} \right) \\ &\quad + \frac{\mu'}{(1 - \delta z)\xi_0} \left(\sum_{l,j,k} \frac{\partial^2 V_j}{\partial y_l \partial y_k} (Z_{k,j} - \delta_{k,j}) Z_{l,i} + \sum_{l,j} \frac{\partial^2 V_j}{\partial y_l \partial y_j} (Z_{l,i} - \delta_{k,i}) + \sum_{l,j,k} Z_{l,i} \frac{\partial V_j}{\partial y_k} \frac{\partial Z_{k,j}}{\partial y_l} \right) \\ &\quad + (1 - \frac{\zeta}{\xi_0}) (\partial_t V)_i - \frac{\zeta}{\xi_0} \left(\sum_{l,j} \tilde{V}_l \frac{\partial V_i}{\partial y_k} Z_{l,j} - \frac{(1 - \delta z)}{\delta} (W \partial_z V)_i \right) - \frac{(Z^\top \nabla_H \zeta)_i}{\xi_0}. \end{aligned}$$

Here we set $\tilde{V} := V - \bar{V}$. The differences occurring in the above calculations for the case $\gamma = 2$ are being briefly discussed in the proof of [Theorem 2.2](#).

4. LINEAR THEORY: THE HYDROSTATIC LAMÉ OPERATOR

We start by analyzing the linear equation

$$(4.1) \quad \begin{cases} \partial_t \zeta + \xi_0 \operatorname{div}_H \bar{V} = G_1, & \text{on } G \times (0, T), \\ \zeta(0, \cdot) = \xi_0, & \text{on } G, \end{cases}$$

for the density $\zeta : G \rightarrow \mathbb{R}$ as well as the hydrostatic Lamé equation

$$(4.2) \quad \begin{cases} \partial_t V - \frac{\mu \Delta_H V}{(1 - \delta z)\xi_0} - \frac{\mu}{\xi_0} \partial_z \left(\frac{1 - \delta z}{\delta^2} \partial_z V \right) - \frac{\mu' \nabla_H \operatorname{div}_H V}{(1 - \delta z)\xi_0} + \omega V = G_2, & \text{on } \Omega \times (0, T), \\ V = 0, & \text{on } \Gamma_u \times (0, T), \\ \partial_z V = 0, & \text{on } \Gamma_b \times (0, T), \\ V(0, \cdot) = v_0, & \text{on } \Omega, \end{cases}$$

for the velocity $V : \Omega \rightarrow \mathbb{R}^2$. Here $0 < T \leq \infty$ is given and (ζ, V) are subject to periodic boundary conditions on the lateral boundary.

We associate with the left hand side of (4.2) the hydrostatic Lamé operator as follows: Let $\xi_0 \in H_{\text{per}}^{1,q}(G; \mathbb{R})$ and assume that $\xi_0 \geq c$ for some $c > 0$. Define functions $a : \Omega \rightarrow \mathbb{R}$ and $b : \Omega \rightarrow \mathbb{R}$ by

$$(4.3) \quad a(x, y, z) := \frac{1}{(1 - \delta z)\xi_0} \quad \text{and} \quad b(x, y, z) := \frac{(1 - \delta z)}{\delta^2 \xi_0}$$

and define the differential operator $\mathcal{A}(x, y, z, D)$ by

$$\mathcal{A}(x, y, z, D)v := \mu a(x, y, z) \Delta_H v + \partial_z (\mu b(x, y, z) \partial_z v) + \mu' a(x, y, z) \nabla_H \operatorname{div}_H v,$$

subject to boundary conditions

$$(4.4) \quad v = 0 \text{ on } \Gamma_u \quad \text{and} \quad \partial_z v = 0 \text{ on } \Gamma_b.$$

For $q > 1$ consider the L^q -realization of $\mathcal{A}(x, y, z, D)$ in $L^q(\Omega; \mathbb{R}^2)$ subject to (4.4) defined by

$$\begin{aligned} A_{\text{HL}} v &:= \mathcal{A}(x, y, z, D)v, \\ D(A_{\text{HL}}) &:= \{v \in H_{\text{per}}^{2,q}(\Omega; \mathbb{R}^2) : v = 0 \text{ on } \Gamma_u \quad \text{and} \quad \partial_z v = 0 \text{ on } \Gamma_b\} \end{aligned}$$

We call A_{HL} the *hydrostatic Lamé operator*.

Our aim is to show that $-A_{\text{HL}} + \omega$ admits a bounded \mathcal{H}^∞ -calculus on $L^q(\Omega; \mathbb{R}^2)$ of \mathcal{H}^∞ -angle strictly less than $\pi/2$ provided ω is large enough. In order to do so, we define the operator $\mathcal{A}_1(z, D)$ and its L^q -realization given by

$$\mathcal{A}_1(z, D) := (1 - \delta z)\xi_0 \cdot \mathcal{A}(x, y, z, D) \quad \text{and} \quad A_1 v := \mathcal{A}_1(z, D)v, \quad \text{with} \quad D(A_1) = D(A_{\text{HL}}).$$

We then treat the hydrostatic Lamé operator $-A_{\text{HL}}$ as multiplicative perturbation of A_1 using the results in [\[18\]](#). To this end, we show that the principal part $\mathcal{A}_{1,\#}(z, \xi)$ of the symbol $\mathcal{A}_1(z, \xi)$ is parameter elliptic

in the cylindrical domain Ω . The key property in the context of cylindrical parameter ellipticity is that the operator $\mathcal{A}_1(z, D)$ is *cylindrical*, meaning that it completely resolves into parts of which one acts only on $G = (0, 1)^2$ and the other acts only on $(0, 1)$. The operator $\mathcal{A}_1(z, D)$ is represented as

$$\mathcal{A}_1(z, D)v = \mu\Delta_{\mathbb{H}}v + \mu'\nabla_{\mathbb{H}}\operatorname{div}_{\mathbb{H}}v + \mu\frac{(1-\delta z)^2}{\delta^2}\partial_{zz}v - \mu\frac{(1-\delta z)}{\delta}\partial_zv = \mathcal{A}_2(D_{\mathbb{H}})v + \mathcal{A}_3(z, D_z)v,$$

where we define $\mathcal{A}_2(D_{\mathbb{H}})$ as well as $\mathcal{A}_3(z, D_z)$ by

$$\mathcal{A}_2(D_{\mathbb{H}})v := \mu\Delta_{\mathbb{H}}v + \mu'\nabla_{\mathbb{H}}\operatorname{div}_{\mathbb{H}}v \quad \text{and} \quad \mathcal{A}_3(z, D_z)v := \mu\frac{(1-\delta z)^2}{\delta^2}\partial_{zz}v - \mu\frac{(1-\delta z)}{\delta}\partial_zv.$$

The $L^q(G; \mathbb{R}^2)$ -realization of $\mathcal{A}_2(D_{\mathbb{H}})$ is defined by

$$\begin{aligned} A_2v &:= \mathcal{A}_2(D_{\mathbb{H}})v, \\ \mathcal{D}(A_2) &:= \{H^{2,q}(G; \mathbb{R}^2) : v \text{ periodic on } \partial G\} \end{aligned}$$

and coincides with the classical Lamé operator on $(0, 1)^2$ subject to periodic boundary conditions. The $L^q(0, 1; \mathbb{R}^2)$ -realization of $\mathcal{A}_3(z, D_z)$ is defined as

$$\begin{aligned} A_3v &:= \mathcal{A}_3(z, D_z)v, \\ \mathcal{D}(A_3) &:= \{v \in H^{2,q}(0, 1; \mathbb{R}^2) : v|_{z=1} = 0 \text{ and } \partial_zv|_{z=0} = 0\}. \end{aligned}$$

We now say that the principal part $-\mathcal{A}_{1,\#}(z, \xi)$ is parameter elliptic of angle $\Phi_{-\mathcal{A}_1} = \max\{\Phi_{-\mathcal{A}_2}, \Phi_{-\mathcal{A}_3}\}$ in the cylindrical domain Ω if and only if the principal part of the two dimensional Lamé operator $-\mathcal{A}_{2,\#}(\xi_{\mathbb{H}})$ is parameter elliptic of angle $\Phi_{-\mathcal{A}_2}$ in the sense of [31, Definition 7.4] and the boundary value problem induced by A_3 is parameter elliptic of angle $\Phi_{-\mathcal{A}_3}$ in the sense of [31, Definition 8.2], meaning that the principal part $-\mathcal{A}_{3,\#}(z, \xi_3)$ is parameter elliptic of angle $\Phi_{-\mathcal{A}_3}$ for all $z \in (0, 1)$ and the boundary value problem induced by A_3 satisfies the Lopatinski-Shapiro condition for all $\Phi > \Phi_{-\mathcal{A}_3}$.

For more information on bounded \mathcal{H}^∞ -calculus and parameter elliptic operators in the classical and in the cylindrical sense, we refer to [2, 5, 6, 11, 16, 17, 21, 31, 32].

Lemma 4.1. *Let $q \in (1, \infty)$. The principal part $-\mathcal{A}_{1,\#}(z, \xi)$ is parameter elliptic of angle $\Phi_{-\mathcal{A}_1} = 0$ in the cylindrical domain $\Omega = G \times (0, 1)$. Moreover, there exists $\omega_0 \in \mathbb{R}$ such that for all $\omega > \omega_0$ the operator $-\mathcal{A}_1 + \omega$ admits a bounded \mathcal{H}^∞ -calculus on $L^q(\Omega; \mathbb{R}^2)$ with \mathcal{H}^∞ -angle $\Phi_{-\mathcal{A}_1+\omega}^\infty < \pi/2$.*

Proof. Consider the two dimensional Lamé operator $-\mathcal{A}_2$ on $G = (0, 1)^2$ subject to periodic boundary conditions. Its discrete symbol is given by

$$-\mathcal{A}_{2,\#}(k_{\mathbb{H}}) = \begin{pmatrix} \mu|k_{\mathbb{H}}|^2 + \mu'k_1^2 & \mu'k_1k_2 \\ \mu'k_1k_2 & \mu|k_{\mathbb{H}}|^2 + \mu'k_2^2 \end{pmatrix}, \quad k_{\mathbb{H}} \in \mathbb{Z}^2.$$

Its eigenvalues $\lambda_{1/2}(k_{\mathbb{H}})$ of $-\mathcal{A}_{2,\#}(k_{\mathbb{H}})$ are computed to be

$$\lambda_{1/2}(k_{\mathbb{H}}) = \frac{2\mu + \mu'}{2}|k_{\mathbb{H}}|^2 \pm \frac{\mu'}{2}|k_{\mathbb{H}}|^2, \quad k_{\mathbb{H}} \in \mathbb{Z}^2.$$

We see that $\lambda_{1/2}(k_{\mathbb{H}}) > 0$ for all $k_{\mathbb{H}} \in \mathbb{Z}^2 \setminus \{0\}$ and it thus follows that $-\mathcal{A}_{2,\#}(k_{\mathbb{H}})$ is parameter elliptic of angle $\Phi_{-\mathcal{A}_2} = 0$ in the sense of [31, Definition 7.4]. Next, consider the function $b_1 : \Omega \rightarrow \mathbb{R}$ defined by

$$b_1(x, y, z) := b_1(z) := \frac{(1-\delta z)^2}{\delta^2}.$$

Since b_1 is continuous, it satisfies the smoothness assumptions needed for applying [6]. The principal part $-\mathcal{A}_{3,\#}(z, \xi_3)$ is given by $-\mathcal{A}_{3,\#}(z, \xi_3) = -b_1(z)\xi_3^2$ for all $z \in (0, 1)$ and all $\xi_3 \in \mathbb{R}$. Hence, $-\mathcal{A}_{3,\#}(z, \xi_3)$ is parameter elliptic of angle $\Phi_{-\mathcal{A}_3} = 0$ and even normally strongly elliptic for all $z \in (0, 1)$. In view of the standard boundary conditions involved, it follows that the boundary value problem associated to $-\mathcal{A}_3$ is normally elliptic and satisfies the conditions in [6]. It thus follows that $-\mathcal{A}_{1,\#}(z, \xi)$ is parameter elliptic of angle $\Phi_{-\mathcal{A}_1} = 0$.

To establish the boundedness of the \mathcal{H}^∞ -calculus of $-\mathcal{A}_1$ up to a shift, we note that the lower order coefficient of A_1 only depends on the vertical variable and in particular belongs to $L^\infty(\Omega)$. Moreover, the

coefficients of the corresponding boundary operator are even constant. The assertion thus follows from the Hölder continuity of b_1 and the results in [31, Section 8] and [5]. \square

We are now in position to apply [18, Theorem 3.1] to establish the boundedness of the \mathcal{H}^∞ -calculus of hydrostatic Lamé operator $-A_{\text{HL}}$ up to a shift.

Proposition 4.2. *Let $q \in (2, \infty)$ and assume that $\xi_0 \in H_{\text{per}}^{1,q}(G; \mathbb{R})$ satisfies $\xi_0 \geq c > 0$ in G for some $c > 0$. Then there exists $\omega_0 \in \mathbb{R}$ such that for all $\omega > \omega_0$ the operator $-A_{\text{HL}} + \omega$ admits a bounded \mathcal{H}^∞ -calculus on $L^q(\Omega; \mathbb{R}^2)$ with \mathcal{H}^∞ -angle $\Phi_{-A_{\text{HL}} + \omega}^\infty < \pi/2$.*

Proof. A Sobolev embedding yields that $((1 - \delta z) \cdot \xi_0)^{-1} \in \text{BUC}^\rho(\bar{\Omega})$ for some $\rho \in (0, 1)$. Define the multiplication operator M on $L^q(\Omega; \mathbb{R}^2)$ by

$$(Mv)(x, y, z) := ((1 - \delta z) \cdot \xi_0)^{-1} v(x, y, z), \quad \text{with } D(M) := L^q(\Omega; \mathbb{R}^2).$$

Then M is boundedly invertible and admits a \mathcal{H}^∞ -calculus of angle $\Phi_M^\infty = 0$ on $L^q(\Omega; \mathbb{R}^2)$. We then write

$$\omega - A_{\text{HL}} = \omega + M \cdot (-A_1 + \omega_1) - M \cdot \omega_1,$$

where $\omega_1 \geq 0$ denotes the shift obtained from Lemma 4.1 and the product is defined as in [18]. The commutator estimate

$$\| [M, (-A_1 + \omega_1 + \mu_1)^{-1}] (\mu_2 + M)^{-1} \| \leq \frac{C}{(1 + |\mu_2|)^{1-\alpha} |\mu_1|^{1+\beta}}, \quad \text{for all } (\mu_1, \mu_2) \in \Sigma_{\Phi_{-A_1 + \omega_1}^\infty} \times \Sigma_\pi,$$

holds true whenever $C, \alpha \geq 0, \beta > 0$ and $\alpha + \beta < 1$. Hence, there exists $\omega_0 \in \mathbb{R}$ such that the operator $\omega + M \cdot (-A_1 + \omega_1)$ admits a bounded \mathcal{H}^∞ -calculus of angle strictly less than $\pi/2$ on $L^q(\Omega; \mathbb{R}^2)$ for all $\omega > \omega_0$. The assertion then follows from [18, Theorem 3.1]. \square

The above Proposition 4.2 implies maximal $L^p - L^q$ -regularity estimates for the solution of (4.2). We verify that the trace space $X_\gamma^v = (X_0^v, X_1^v)_{1-1/p}$ for $X_0^v = L^q(\Omega; \mathbb{R}^2)$ and $X_1^v = Y$ is characterized by

$$X_\gamma^v := \begin{cases} \{v_0 \in B_{qp, \text{per}}^{2(1-1/p)}(\Omega; \mathbb{R}^2) : (\partial_z v_0)|_{\Gamma_b} = 0, v_0|_{\Gamma_u} = 0\}, & \frac{1}{2} + \frac{1}{2q} + \frac{1}{p} < 1, \\ \{v_0 \in B_{qp, \text{per}}^{2(1-1/p)}(\Omega; \mathbb{R}^2) : v_0|_{\Gamma_u} = 0\}, & \frac{1}{2q} + \frac{1}{p} < 1, \\ B_{qp, \text{per}}^{2(1-1/p)}(\Omega; \mathbb{R}^2), & 1 < \frac{1}{2q} + \frac{1}{p}. \end{cases}$$

We then obtain the following maximal regularity estimate for the equation (4.2).

Corollary 4.3. *Let $p, q \in (2, \infty)$, $G_2 \in \mathbb{E}_{0, \infty}^v$ and v_0 as in assumption (A). Then there exists $\omega \geq 0$ such that the equation (4.2) admits a unique, strong solution*

$$V \in \mathbb{E}_{1, \infty}^v := H^{1,p}(\mathbb{R}_+; L^q(\Omega; \mathbb{R}^2)) \cap L^p(\mathbb{R}_+; Y),$$

satisfying

$$\|V\|_{\mathbb{E}_{1, \infty}^v} \leq C(\|G_2\|_{\mathbb{E}_{0, \infty}^v} + \|v_0\|_{B_{qp}^{2(1-1/p)}(\Omega)})$$

for some $C > 0$. In particular, the solution operator

$$(4.5) \quad S_V : \mathbb{E}_{0, \infty}^v \times X_\gamma^v \rightarrow \mathbb{E}_{1, \infty}^v, \quad (G_2, v_0) \mapsto V$$

is an isomorphism.

Furthermore, we obtain the following well-posedness result for equation (4.1).

Corollary 4.4. *Let $p, q \in (2, \infty)$ and $V \in \mathbb{E}_{1, \infty}^v$ be the unique solution of (4.2). Assume that $G_1 \in L^p(\mathbb{R}_+; H_{\text{per}}^{1,q}(G; \mathbb{R}))$ and $\xi_0 \in H_{\text{per}}^{1,q}(G; \mathbb{R})$. Then (4.1) admits a unique, strong solution*

$$\zeta \in H^{1,p}(\mathbb{R}_+; H_{\text{per}}^{1,q}(G; \mathbb{R}))$$

satisfying

$$\|\zeta\|_{H^{1,p}(\mathbb{R}_+; H^{1,q}(G, \mathbb{R}))} \leq C(\|G_1\|_{L^p(\mathbb{R}_+; H^{1,q}(G, \mathbb{R}))} + \|V\|_{\mathbb{E}_{1, \infty}^v} + \|\xi_0\|_{H^{1,q}(G)})$$

for some $C > 0$.

5. LOCAL-WELLPOSEDNESS

The proof of [Theorem 2.2](#) is based on the estimates given in [Proposition 4.2](#) and on the estimates on the nonlinearities F_1 and F_2 given in the sequel.

Lemma 5.1. *Let $\tau < 1$ and $p, q \in (2, \infty)$. Then there exist $T \in (0, \tau]$, a constant $\delta > 0$, depending only on p and q , as well as a constant $C > 0$, depending only on p, q and τ , such that*

$$\|F_1(\zeta, V)\|_{L^p(0,T;H^{1,q}(G))} \leq T^\delta \|V\|_{\mathbb{E}_{1,T}^v} \quad \text{and} \quad \|F_2(\zeta, V)\|_{\mathbb{E}_{0,T}^v} \leq C(T^\delta + T^\delta \|V\|_{\mathbb{E}_{1,T}^v} + \|V\|_{\mathbb{E}_{1,T}^v}^2),$$

whenever $(\zeta, V) \in \mathbb{E}_{1,T}$.

Proof. We start by estimating $\zeta \in H^{1,p}(0, T; H_{\text{per}}^{1,q}(G; \mathbb{R}))$. By Sobolev embedding and Hölder's inequality we obtain

$$(5.1) \quad \|\zeta\|_{L^p(0,T;H^{1,q}(G))} \leq CT^{1/p}, \quad \|\zeta - \xi_0\|_{L^\infty(0,T;H^{1,q}(G))} \leq CT^{1/p'} \quad \text{and} \quad \|\zeta\|_{L^\infty(0,T;H^{1,q}(G))} \leq C.$$

In view of $H^{1,q}(G)$ being a Banach algebra for $q > 2$ and estimate (5.1) we see that

$$\begin{aligned} \|F_1\|_{L^p(0,T;H^{1,q}(G))} &\leq \|\zeta - \xi_0\|_{L^\infty(0,T;H^{1,q}(G))} \|\operatorname{div}_H \bar{V}\|_{L^p(0,T;H^{1,q}(G))} \\ &\quad + \|\zeta\|_{L^\infty(0,T;H^{1,q}(G))} \|Z - \mathbb{I}_2\|_{L^\infty(0,T;H^{1,q}(G))} \|\nabla_H \bar{V}\|_{L^p(0,T;H^{1,q}(G))} \\ &\leq CT^{1/p'} \|V\|_{\mathbb{E}_{1,T}^v}. \end{aligned}$$

In order to estimate F_2 observe that by [17, Section 6] we have $\|V \cdot \nabla_H V\|_{L^q(\Omega)} \leq C \|V\|_{H^{1+1/q,q}(\Omega)}^2$. Hence,

$$\|\zeta \sum_{l,j} \tilde{V}_l \frac{\partial V_i}{\partial y_k} Z_{l,j}\|_{L^p(0,T;L^q)} \leq C \|\zeta\|_{L^\infty(0,T;L^\infty(G))} \|Z\|_{L^\infty(0,T;L^\infty(G))} \|V\|_{L^{2p}(0,T;H^{1+1/q,q}(\Omega))}^2 \leq C \|V\|_{\mathbb{E}_{1,T}^v}^2$$

by the embeddings

$$(5.2) \quad \mathbb{E}_{1,T}^v \hookrightarrow L^{2p}(0, T; H^{2-1/p,q}(\Omega)) \hookrightarrow L^{2p}(0, T; H^{1+1/q,q}(\Omega)),$$

for $1/p + 1/q \leq 1$. Next, we estimate the nonlinear term including the vertical velocity, i. e., $W \cdot \partial_z V$, anisotropically and obtain

$$\|\frac{\zeta}{\xi_0} W \cdot \partial_z V\|_{L^p(0,T;L^q(\Omega))} \leq C \|\zeta W\|_{L_t^\infty L_H^q L_z^\infty} \|V\|_{L_t^p L_H^\infty H_z^{1,q}} \leq C \|\zeta W\|_{L_t^\infty L_H^q L_z^\infty} \|V\|_{\mathbb{E}_{1,T}^v}.$$

Combining the embeddings $H_z^{1,q} \hookrightarrow L_z^\infty$, $\mathbb{E}_{1,T}^v \hookrightarrow L^\infty(0, T; L_H^\infty(G; L_z^q(0, 1)))$ for $1/p + 1/q < 1$ with the representation (2.5), we obtain

$$\begin{aligned} \|\zeta W\|_{L_t^\infty L_H^q L_z^\infty} &\leq C \|\tilde{V} Z^\top \nabla_H \zeta + \zeta Z^\top : \nabla_H \tilde{V}\|_{L^\infty(0,T;L^q(\Omega))} \\ &\leq C \|\tilde{V}\|_{L^\infty(0,T;L_H^\infty L_z^q)} \|Z\|_{L^\infty(0,T;L^\infty(G))} \|\nabla_H \zeta\|_{L^\infty(0,T;L^q(G))} \\ &\quad + C \|\zeta\|_{L^\infty(0,T;L^\infty(G))} \|Z\|_{L^\infty(0,T;L^\infty(G))} \|\nabla_H \tilde{V}\|_{L^\infty(0,T;L^q(\Omega))} \\ &\leq C \|V\|_{\mathbb{E}_{1,T}^v}. \end{aligned}$$

Next, observe that

$$\left\| \left(1 - \frac{\zeta}{\xi_0}\right) (\partial_t V) \right\|_{L^p(0,T;L^q(\Omega))} \leq C \|\zeta - \xi_0\|_{L^\infty(0,T;H^{1,q}(G))} \|V\|_{H^{1,p}(0,T;L^q(\Omega))} \leq CT^{1/p'} \|V\|_{\mathbb{E}_{1,T}^v}.$$

We estimate the remaining terms by properties of the Lagrangian transformation given in [Section 3](#) and obtain

$$\begin{aligned} \left\| \sum_{l,j,k} \frac{\partial^2 V_i}{\partial y_l \partial y_k} (Z_{k,j} - \delta_{k,j}) Z_{l,j} \right\|_{L^p(0,T;L^q(\Omega))} &\leq C \|V\|_{L^p(0,T;H^{2,q}(\Omega))} \|Z - \mathbb{I}_2\|_{L^\infty(0,T;H^{1,q}(G))} \|Z\|_{L^\infty(0,T;H^{1,q}(G))} \\ &\leq CT^{1/p'} \|V\|_{\mathbb{E}_{1,T}^v}, \end{aligned}$$

and similarly

$$\left\| \sum_{l,k} \frac{\partial^2 V_i}{\partial y_l \partial y_k} (Z_{l,k} - \delta_{k,l}) \right\|_{L^p(0,T;L^q(\Omega))} \leq C \|V\|_{L^p(0,T;H^{2,q}(\Omega))} \|Z - \mathbb{I}_2\|_{L^\infty(0,T;H^{1,q}(G))} \leq CT^{1/p'} \|V\|_{\mathbb{E}_{1,T}^v}.$$

Furthermore, observe that

$$\begin{aligned} \left\| \sum_{l,j,k} Z_{l,k} \frac{\partial V_i}{\partial y_k} \frac{\partial Z_{k,j}}{\partial y_l} \right\|_{L^p(0,T;L^q(\Omega))} &\leq C \|Z\|_{L^\infty(0,T;L^\infty(G))} \|\nabla_{\text{H}} V\|_{L^p(0,T;H^{1,q}(\Omega))} \sum_{l,j,k} \left\| \frac{\partial Z_{k,j}}{\partial y_l} \right\|_{L^\infty(0,T;L^q(G))} \\ &\leq CT^{1/p'} \|V\|_{\mathbb{E}_{1,T}^v}, \end{aligned}$$

where we used that for $1 \leq j, k, l \leq 2$ and all $y_{\text{H}} \in G$ one has $\frac{\partial Z_{k,j}}{\partial y_l}(0, y_{\text{H}}) = 0$. Hence,

$$\left\| \frac{\partial Z_{k,j}}{\partial y_l} \right\|_{L^\infty(0,T;L^q(G))} \leq T^{1/p'} \left\| \frac{\partial Z_{k,j}}{\partial y_l} \right\|_{H^{1,p}(0,T;L^q(G))} \leq T^{1/p'} \|Z\|_{H^{1,p}(0,T;H^{1,q}(G))} \leq CT^{1/p'}.$$

Finally, (5.1) allows us to estimate the pressure term as

$$\left\| \frac{\zeta}{\xi_0} Z^\top \nabla_{\text{H}} \zeta \right\|_{L^p(0,T;L^q(\Omega))} \leq C \|\zeta\|_{L^\infty(0,T;L^\infty(G))} \|Z\|_{L^\infty(0,T;L^\infty(G))} \|\zeta\|_{L^p(0,T;H^{1,q}(G))} \leq CT^{1/p'}.$$

□

We are now in position to prove our first local well-posedness result.

Proof of Theorem 2.2. Let us remark first that in the sequel we implicitly consider a shifted linearized equations in order to exploit Corollary 4.3. As the resulting linear term added on the right-hand side can be estimated easily, we do not explicitly mention this shift in the remainder of the proof.

Observe next that Corollary 4.3 guarantees the existence of the reference solution V_* to (3.8) with homogeneous right hand side and with initial value v_0 . In the following, the prescript $_0$ indicates homogeneous initial data. We then set

$$\mathbb{B}_r := \{(\zeta, V) \in \mathbb{E}_{1,T}: V - V_* \in {}_0\mathbb{E}_{1,T}^v, \text{ and } \|V - V_*\|_{\mathbb{E}_{1,T}^v} \leq r\}.$$

For $(\zeta_1, V_1) \in \mathbb{B}_r$, we denote by

$$(\zeta, V) := \Psi(\zeta_1, V_1) \in \mathbb{E}_{1,T}$$

the solution to the linearized equation (3.8) with right-hand sides $F_1(\zeta_1, V_1)$ and $F_2(\zeta_1, V_1)$ and guaranteed by Corollary 4.3, Corollary 4.4 as well as by Lemma 5.1. We show next that Ψ is a self-map and a contraction.

Notice that $V - V_* \in {}_0\mathbb{E}_{1,T}^v$ and we hence only need to show that $\|V - V_*\|_{\mathbb{E}_{1,T}^v} \leq r$. Corollary 4.3 as well as Lemma 5.1 yield

$$\begin{aligned} \|V - V_*\|_{\mathbb{E}_{1,T}^v} &\leq C \|F_2(\zeta_1, V_1)\|_{\mathbb{E}_{0,T}^v} \leq C(T^\delta + T^\delta \|V_1\|_{\mathbb{E}_{1,T}^v} + \|V_1\|_{\mathbb{E}_{1,T}^v}^2) \\ &\leq C(T^\delta + T^\delta \|V_*\|_{\mathbb{E}_{1,T}^v} + T^\delta \|V_1 - V_*\|_{\mathbb{E}_{1,T}^v} + \|V_*\|_{\mathbb{E}_{1,T}^v}^2 + \|V_1 - V_*\|_{\mathbb{E}_{1,T}^v}^2) \\ &\leq CT^\delta + CT^\delta (\|V_*\|_{\mathbb{E}_{1,T}^v} + r) + C(\|V_*\|_{\mathbb{E}_{1,T}^v}^2 + r^2), \end{aligned}$$

where the constant $C > 0$ is independent of T . Observing that $\|V_*\|_{\mathbb{E}_{0,T}^v} \rightarrow 0$ as $T \rightarrow 0$, the corresponding terms are getting arbitrary small by choosing $T > 0$ appropriately small. Hence, for appropriate T and r , the solution map Ψ is a self-map. The contraction property follows the same way by using the a priori estimate on ζ given in Corollary 4.4 and the estimate on F_1 given in Lemma 5.1.

Finally, we choose T sufficiently small such that $X(t, \cdot)$ is a C^1 -diffeomorphisms from G into G and set $Y(t, \cdot) = X^{-1}(t, \cdot)$, see Section 3. Then, defining

$$\xi(t, x, y) := \zeta(t, Y(t, x, y)) \quad \text{and} \quad v(t, x, y, z) := V(t, Y(t, x, y), z)$$

we verify that $(\xi, v) \in \mathbb{E}_{1,T}$ satisfies the system (2.4). Since the vertical transform (2.3) is invertible for all times, we find a solution to (1.1). The uniqueness property follows from the uniqueness of solutions to the system (3.8). The proof of assertion a) of Theorem 2.2 is complete.

We finally provide a proof of assertion b). After introducing Lagrangian coordinates for the system (2.6), we see that the hydrostatic Lamé operator in this case reads as

$$B := \mu c(x, y, z) \Delta + \mu' c(x, y, z) \nabla_{\mathbb{H}} \operatorname{div}_{\mathbb{H}} \\ D(B) := \{v \in H_{\text{per}}^{2,q}(\Omega; \mathbb{R}^2) : v = 0 \text{ on } \Gamma_u \text{ and } \partial_z v = 0 \text{ on } \Gamma_b\}$$

Here $c : \Omega \rightarrow \mathbb{R}$ is given by $c(x, y, z) := \frac{1}{\xi_0 + (1/2)z}$. Arguing as in Section 4, we see that $-B + \omega$ admits a bounded \mathcal{H}^∞ -calculus on $L^q(\Omega; \mathbb{R}^2)$ of \mathcal{H}^∞ -angle strictly less than $\pi/2$. Moreover, a result similar to Corollary 4.4 is true for the transformed equation (2.6)₁, which reads as

$$\partial_t \zeta + \zeta_0 \operatorname{div}_{\mathbb{H}} \bar{V} + \frac{1}{2} \int_0^1 z \operatorname{div}_{\mathbb{H}} V \, dz = -(\zeta - \zeta_0) \operatorname{div}_{\mathbb{H}} \bar{V} - \zeta \nabla_{\mathbb{H}} \bar{V} : [Z^\top - \mathbb{I}_2] - \frac{1}{2} \int_0^1 z \nabla_{\mathbb{H}} V : [Z^\top - \mathbb{I}_2] \, dz.$$

We then may conclude local well-posedness of equation (2.6). In fact, the estimates of the nonlinear terms are deduced similarly to the ones given in Lemma 5.1. In particular, the transformed pressure term $(2\xi + z)(Z^\top \nabla_{\mathbb{H}} \xi)$ may be estimated following the arguments given in the proof of Lemma 5.1. \square

6. GLOBAL IN TIME EXISTENCE

In order to prove the assertions of Theorem 2.3 we transform the system (1.1) by a slightly different Lagrange transform compared to the one introduced in Section 3. More precisely, we define the transformed density and velocity by

$$\zeta(t, y_{\mathbb{H}}) := \xi(t, X(t, y_{\mathbb{H}})) - \bar{\xi} \quad \text{and} \quad V(t, y_{\mathbb{H}}, z) := v(t, X(t, y_{\mathbb{H}}), z),$$

where $\bar{\xi} = \frac{1}{|\bar{G}|} \int_{\bar{G}} \xi_0 > 0$. The reason for this modification is our aim to linearize the system (2.4) around the constant steady state $(\bar{\xi}, 0)$. The respective system is then given by

$$(6.1) \quad \left\{ \begin{array}{ll} \partial_t \zeta + \bar{\xi} \operatorname{div}_{\mathbb{H}} \bar{V} = F_1(\zeta, V), & \text{on } G \times (0, \infty), \\ \partial_t V - \frac{\mu \Delta_{\mathbb{H}} V}{(1 - \delta z) \bar{\xi}} - \frac{\mu}{\bar{\xi}} \partial_z \left(\frac{1 - \delta z}{\delta^2} \partial_z V \right) - \frac{\mu' \nabla_{\mathbb{H}} \operatorname{div}_{\mathbb{H}} V}{(1 - \delta z) \bar{\xi}} + \frac{\nabla_{\mathbb{H}} \zeta}{\bar{\xi}} = F_2(\zeta, V), & \text{on } \Omega \times (0, \infty), \\ \partial_z \zeta = 0, & \text{on } \Omega \times (0, \infty), \\ V = 0, & \text{on } \Gamma_u \times (0, \infty), \\ \partial_z V = 0, & \text{on } \Gamma_b \times (0, \infty), \\ (\zeta(0), V(0)) = (\xi_0 - \bar{\xi}, v_0), & \end{array} \right.$$

where the right-hand sides F_1 and F_2 for $i = 1, 2$ are given by

$$(6.2) \quad F_1(\zeta, V) = -\zeta \operatorname{div}_{\mathbb{H}} \bar{V} - (\zeta + \bar{\xi}) \nabla_{\mathbb{H}} \bar{V} : [Z^\top - \mathbb{I}_2], \\ (F_2(\zeta, V))_i = \frac{\mu}{(1 - \delta z) \bar{\xi}} \left(\sum_{l,j,k} \frac{\partial^2 V_i}{\partial y_l \partial y_k} (Z_{k,j} - \delta_{k,j}) Z_{l,j} + \sum_{l,k} \frac{\partial^2 V_i}{\partial y_l \partial y_k} (Z_{l,k} - \delta_{k,l}) + \sum_{l,j,k} Z_{l,k} \frac{\partial V_i}{\partial y_k} \frac{\partial Z_{k,j}}{\partial y_l} \right) \\ + \frac{\mu'}{(1 - \delta z) \bar{\xi}} \left(\sum_{l,j,k} \frac{\partial^2 V_j}{\partial y_l \partial y_k} ((Z_{k,j} - \delta_{k,j}) Z_{l,i} + \sum_{l,j} \frac{\partial^2 V_j}{\partial y_l \partial y_j} (Z_{l,i} - \delta_{k,i}) + \sum_{l,j,k} Z_{l,i} \frac{\partial V_j}{\partial y_k} \frac{\partial Z_{k,j}}{\partial y_l} \right) \\ - \frac{\zeta (\partial_t V)_i}{\bar{\xi}} - \frac{\zeta + \bar{\xi}}{\bar{\xi}} \left(\sum_{l,j} \tilde{V}_l \frac{\partial V_i}{\partial y_k} Z_{l,j} - (W \partial_z V)_i \right) - \frac{1}{\bar{\xi}} \sum_j \frac{\partial \zeta}{\partial y_j} (Z_{j,i} - \delta_{j,i}).$$

6.1. The compressible hydrostatic Stokes operator.

We introduce now the compressible hydrostatic Stokes operator and discuss its properties. Recall that $X_0 := H_{\text{per}}^{1,q}(G, \mathbb{R}) \times L^q(\Omega, \mathbb{R}^2)$ and $X_1 := H_{\text{per}}^{1,q}(G, \mathbb{R}) \times Y$. For $\bar{\xi}$ given as above, define functions $a_{\bar{\xi}} : \Omega \rightarrow \mathbb{R}$ and $b_{\bar{\xi}} : \Omega \rightarrow \mathbb{R}$ by

$$(6.3) \quad a_{\bar{\xi}}(x, y, z) := a_{\bar{\xi}}(z) := \frac{1}{(1 - \delta z) \bar{\xi}} \quad \text{and} \quad b_{\bar{\xi}}(x, y, z) := b_{\bar{\xi}}(z) := \frac{(1 - \delta z)}{\delta^2 \bar{\xi}}.$$

Furthermore, define the differential operator $\mathcal{A}_{\bar{\xi}}$ by

$$\mathcal{A}_{\bar{\xi}}v := \mu a_{\bar{\xi}}(z)v + \partial_z(\mu b_{\bar{\xi}}(z)\partial_z v) + \mu' a_{\bar{\xi}}(z)\nabla_{\mathbb{H}} \operatorname{div}_{\mathbb{H}} v,$$

subject to boundary conditions

$$(6.4) \quad v = 0 \text{ on } \Gamma_u \text{ and } \partial_z v = 0 \text{ on } \Gamma_b.$$

For $q > 1$ consider the L^q -realization of $\mathcal{A}_{\bar{\xi}}$ in $L^q(\Omega; \mathbb{R}^2)$ subject to (6.4) defined by

$$(6.5) \quad \begin{aligned} A_{\mathbb{H}\mathbb{L}, \bar{\xi}}v &:= \mathcal{A}_{\bar{\xi}}v \\ \mathbb{D}(A_{\mathbb{H}\mathbb{L}, \bar{\xi}}) &:= \{v \in \mathbb{H}_{\text{per}}^{2,q}(\Omega; \mathbb{R}^2) : v = 0 \text{ on } \Gamma_u \text{ and } \partial_z v = 0 \text{ on } \Gamma_b\} \end{aligned}$$

We call $A_{\mathbb{H}\mathbb{L}, \bar{\xi}}$ the *hydrostatic Lamé operator* subject to $\bar{\xi}$. Moreover, we define the *compressible, hydrostatic Stokes operator* A_{CHS} in X_0 by

$$(6.6) \quad \begin{aligned} A_{\text{CHS}} &:= \begin{pmatrix} 0 & -\bar{\xi} \operatorname{div}_{\mathbb{H}} \operatorname{avg} \\ -\nabla_{\mathbb{H}} & A_{\mathbb{H}\mathbb{L}, \bar{\xi}} \end{pmatrix}, \\ \mathbb{D}(A_{\text{CHS}}) &:= X_1 = \mathbb{H}_{\text{per}}^{1,q}(G; \mathbb{R}) \times \mathbb{D}(A_{\mathbb{H}\mathbb{L}, \bar{\xi}}), \end{aligned}$$

where $\operatorname{avg}(v) = \int_0^1 v(\cdot, z) dz$.

For the investigation of the linear system (6.1) we show first that the compressible hydrostatic Stokes operator $-A_{\text{CHS}}$ admits a bounded \mathcal{H}^∞ -calculus up to a shift.

Proposition 6.1. *Let $q \in (1, \infty)$. Then there exists $\omega > 0$ such that the operator $-A_{\text{CHS}} + \omega$ admits a bounded \mathcal{H}^∞ -calculus on X_0 with \mathcal{H}^∞ -angle $\Phi_{-A_{\text{CHS}} + \omega}^\infty < \pi/2$.*

Proof. Arguing similarly as in the proof of Proposition 4.2, we conclude that there exists a $\omega_0 \in \mathbb{R}$ such that for all $\omega > \omega_0$ the operator $-A_{\mathbb{H}\mathbb{L}, \bar{\xi}} + \omega$ admits a bounded \mathcal{H}^∞ -calculus on $L^q(\Omega; \mathbb{R}^2)$ of angle $\Phi_{-A_{\mathbb{H}\mathbb{L}, \bar{\xi}} + \omega}^\infty < \pi/2$. We next decompose A_{CHS} into

$$(6.7) \quad A_{\text{CHS}} = \begin{pmatrix} 0 & -\bar{\xi} \operatorname{div}_{\mathbb{H}} \operatorname{avg} \\ 0 & A_{\mathbb{H}\mathbb{L}, \bar{\xi}} \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ -\nabla_{\mathbb{H}} & 0 \end{pmatrix} =: A_0 + A_1.$$

In order to show the bounded \mathcal{H}^∞ -calculus of $-A_{\text{CHS}}$, note that $\bar{\xi} \operatorname{div}_{\mathbb{H}} \operatorname{avg}$ is a bounded operator from Y to $\mathbb{H}_{\text{per}}^{1,q}(G; \mathbb{R})$. Hence, A_0 admits a bounded \mathcal{H}^∞ -calculus on $\mathbb{H}_{\text{per}}^{1,q}(G; \mathbb{R}) \times L^q(\Omega; \mathbb{R}^2)$. Furthermore, since $\nabla_{\mathbb{H}}$ maps $\mathbb{H}_{\text{per}}^{1,q}(G; \mathbb{R}^2)$ boundedly to $L^q(\Omega; \mathbb{R}^2)$, it follows that A_{CHS} is a bounded perturbation of A_0 and that thus A_{CHS} admits a bounded \mathcal{H}^∞ -calculus on X_0 . \square

We prove next that the compressible hydrostatic Stokes operator \mathcal{A} is invertible when defined in a suitable subspace X_m of X_0 . To this end, we study the following resolvent problem within the L^q -setting: given $\lambda \in \mathbb{C}$ satisfying $\operatorname{Re} \lambda \geq 0$, consider the equation

$$(6.8) \quad \left\{ \begin{aligned} \lambda \zeta + \operatorname{div}_{\mathbb{H}} \bar{V} &= f_1, & \text{in } G, \\ \lambda V - \frac{\mu \Delta_{\mathbb{H}} V}{(1 - \delta z)} - \mu \partial_z \left(\frac{1 - \delta z}{\delta^2} \partial_z V \right) - \frac{\mu' \nabla_{\mathbb{H}} \operatorname{div}_{\mathbb{H}} V}{(1 - \delta z)} + \nabla_{\mathbb{H}} \zeta &= f_2, & \text{on } \Omega, \\ \partial_z \zeta &= 0, & \text{on } \Omega, \\ V &= 0, & \text{on } \Gamma_u, \\ \partial_z V &= 0, & \text{on } \Gamma_b, \end{aligned} \right.$$

where we have set $\bar{\xi} = 1$ for simplicity of the representation.

Integrating the first equation of (6.8) and using the periodic boundary conditions we see that for the case $\lambda = 0$ we need to impose the compatibility condition $\int_G f_1 d(x, y) = 0$. Therefore, it is natural to consider the space

$$L_0^q(G; \mathbb{R}) := \{f \in L^q(G; \mathbb{R}) : \int_G f d(x, y) = 0\}$$

and to restrict the operator A_{CHS} to the subspace $X_m := \mathbb{H}_{\text{per}}^{1,q}(G; \mathbb{R}) \cap L_0^q(G; \mathbb{R}) \times L^q(\Omega; \mathbb{R}^2)$ of X_0 and equipped with the domain $\mathbb{D}(A_{\text{CHS}}) \cap X_m$.

Lemma 6.2. *Let $q \in [2, \infty)$ and $(f_1, f_2) \in X_m = H_{\text{per}}^{1,q}(G; \mathbb{R}) \cap L_0^q(G; \mathbb{R}) \times L^q(\Omega, \mathbb{R}^2)$. Then there exists a unique solution $(\zeta, V) \in D(A_{\text{CHS}}) \cap X_m = H_{\text{per}}^{1,q}(G; \mathbb{R}) \cap L_0^q(G; \mathbb{R}) \times Y$ of (6.8) with $\lambda = 0$ and a constant $C > 0$ such that*

$$(6.9) \quad \|\zeta\|_{H^{1,q}(G)} + \|V\|_{H^{2,q}(\Omega)} \leq C (\|f_1\|_{H^{1,q}(G)} + \|f_2\|_{L^q(\Omega)}).$$

Proof. The proof is inspired by [19, Chapter 3]. Note, however, that we need to consider here the hydrostatic Lamé type operator $A_{\text{HL}, \bar{\xi}}$ instead of the Laplacian as in [19, Chapter-3] and that we have to deal with the non-homogeneous divergence equation (6.8)₁. We subdivide the proof into two steps.

Step 1: the case $q = 2$

The existence of a unique solution (ζ, V) to (6.8) combined with the estimate (6.9) for the case $q = 2$ follows from the Babuška-Brezzi theory. In fact, consider the weak formulation of the equation (6.8). For this we introduce the function spaces

$$\mathcal{V} = \{\varphi \in H_{\text{per}}^{1,2}(\Omega; \mathbb{R}^2) : \varphi = 0 \text{ on } \Gamma_u\} \quad \text{and} \quad \mathcal{W} = L_0^2(G; \mathbb{R}),$$

and consider the functions $a_{\bar{\xi}}$ and $b_{\bar{\xi}}$ defined as in (6.3) for $\bar{\xi} = 1$. Suppose now that (ζ, v) is a strong solution of (6.8) for $\lambda = 0$. Multiplying equation (6.8)₁ and (6.8)₂ with $(\psi, \varphi) \in \mathcal{W} \times \mathcal{V}$ and integrating by parts we verify that the weak formulation of (6.8) reads as

$$(6.10) \quad \begin{cases} (\text{div}_{\text{H}} \bar{v}, \psi)_{L^2(G)} = (f_1, \psi)_{L^2(G)}, \\ \mu (a_1 \nabla_{\text{H}} v, \nabla_{\text{H}} \varphi)_2 + \mu (b_1 \partial_z v, \partial_z \varphi)_2 + \mu' (a_1 \text{div}_{\text{H}} v, \text{div}_{\text{H}} \varphi)_2 - (\zeta, \text{div}_{\text{H}} \bar{\varphi})_{L^2(G)} = (f_2, \varphi)_2. \end{cases}$$

We then define the sesquilinear forms $\mathcal{C} : \mathcal{V} \times \mathcal{V} \rightarrow \mathbb{C}$ and $\mathcal{D} : \mathcal{V} \times \mathcal{W} \rightarrow \mathbb{C}$ by

$$\begin{aligned} \mathcal{C}(v, \varphi) &:= \mu (a_1 \nabla_{\text{H}} v, \nabla_{\text{H}} \varphi)_{L^2(\Omega)} + \mu (b_1 \partial_z v, \partial_z \varphi)_{L^2(\Omega)} + \mu' (a_1 \text{div}_{\text{H}} v, \text{div}_{\text{H}} \varphi)_{L^2(\Omega)}, \\ \mathcal{D}(v, \psi) &:= -(\psi, \text{div}_{\text{H}} \bar{v})_{L^2(G)}, \end{aligned}$$

and set

$$\langle \chi_1, \psi \rangle := (-f_1, \psi)_{L^2(G)}, \quad \langle \chi_2, \psi \rangle := (f_2, \varphi)_{L^2(\Omega)}.$$

Then \mathcal{C} and \mathcal{D} are bounded sesquilinear forms, since a_1 and b_1 are uniformly bounded and non-degenerate. Observe that $\chi_1 \in \mathcal{W}^*$ and $\chi_2 \in \mathcal{V}^*$. This allows us to rephrase equation (6.10) as

$$(6.11) \quad \begin{cases} \mathcal{D}(v, \psi) = \langle \chi_1, \psi \rangle, & \psi \in \mathcal{W}, \\ \mathcal{C}(v, \varphi) + \mathcal{D}(\varphi, \zeta) = \langle \chi_2, \psi \rangle, & \varphi \in \mathcal{V}. \end{cases}$$

We use the identity: $\text{curl}_{\text{H}}^{\top}(\text{curl}_{\text{H}} v) = \nabla_{\text{H}} \text{div}_{\text{H}} v - \Delta_{\text{H}} v$ to obtain

$$\mathcal{C}(v, v) = \mu a_1 \|\text{curl}_{\text{H}} v\|_{L^2(\Omega)}^2 + \mu b_1 C \|\partial_z v\|_{L^2(\Omega)}^2 + (\mu + \mu') \|\text{div}_{\text{H}} v\|_{L^2(\Omega)}^2.$$

The coerciveness of \mathcal{C} will follow from the fact that $\mu > 0$, $\mu + \mu' > 0$ and the coefficients a_1, b_1 are uniformly bounded from below. The inf-sup condition for \mathcal{D} follows from [19, Lemma 3.2]. We apply the Babuška-Brezzi theorem to conclude that there exists a unique solution $(\zeta, v) \in \mathcal{W} \times \mathcal{V}$ to (6.11). Employing the method of difference quotients we obtain higher regularity estimates and thus $(\zeta, v) \in H_{\text{per}}^{1,2}(G, \mathbb{R}) \cap L_0^2(G; \mathbb{R}) \times Y$.

Step 2: the case $q > 2$

We reduce the problem to a two-dimensional Stokes equation and an elliptic boundary value problem. Multiplying (6.8)₂ with $(1 - \delta z)$ and taking vertical averages yields

$$(6.12) \quad \begin{cases} -\mu \Delta_{\text{H}} \bar{v} + \nabla_{\text{H}} \tilde{\zeta} = \overline{(1 - \delta z) f_2} + \mu \frac{(1 - \delta)^2}{\delta^2} (\partial_z v)|_{\Gamma_u} + \mu \frac{1}{\delta} v|_{\Gamma_b} + \mu \bar{v} + \mu' \nabla_{\text{H}} f_1, & \text{in } G, \\ \text{div}_{\text{H}} \bar{v} = f_1, & \text{in } G, \end{cases}$$

where $\tilde{\zeta} = (1 - \frac{\delta}{2}) \zeta$. If ζ is known, the velocity v may then be determined by the elliptic problem

$$(6.13) \quad \begin{cases} -a_1 \mu \Delta_{\text{H}} v - \mu \partial_z (b_1 \partial_z v) - a_1 \mu' \nabla_{\text{H}} \text{div}_{\text{H}} v = f_2 - \nabla_{\text{H}} \zeta, & \text{on } \Omega, \\ v = 0, & \text{on } \Gamma_u \\ \partial_z v = 0, & \text{on } \Gamma_b. \end{cases}$$

Assume now that $2 < q \leq 4$. Then $f_1 \in H_{\text{per}}^{1,2}(G; \mathbb{R}) \cap L_0^2(G; \mathbb{R})$, $f_2 \in L^2(\Omega; \mathbb{R}^2)$ and there exists a unique solution $(\zeta, v) \in H_{\text{per}}^{1,2}(G; \mathbb{R}) \cap L_0^2(G; \mathbb{R}) \times Y$ to (6.8). Note that (6.12) combined with (6.13) are equivalent to (6.8). Hence, equations (6.12) and (6.13) have a unique, strong solution as well. By Sobolev embeddings and the trace theorem, each term on the right-hand side of (6.12)₁ belongs to $L^q(G; \mathbb{R}^2)$ and we also see that the right-hand side f_1 of (6.12)₂ lies in $H_{\text{per}}^{1,q}(G; \mathbb{R})$. Applying [19, Lemma 3.5], we deduce that $\tilde{\zeta} \in H_{\text{per}}^{1,q}(G; \mathbb{R}) \cap L_0^q(G; \mathbb{R})$, which implies the same regularity for ζ . Thus each term on the right-hand side of (6.13) lies in $L^q(\Omega; \mathbb{R}^2)$. Hence there exists a unique, strong solution $v \in Y$ to (6.13) by the results in [6] and thus $(\zeta, v) \in D(A_{\text{CHS}}) \cap X_m$. We now may repeat this argument for $4 < q < \infty$ to conclude that the solution constructed in the L^2 -framework admits $H^{1,q} - H^{2,q}$ -regularity. \square

The following lemma concerns the solvability of the resolvent problem (6.8) for $\lambda \in \mathbb{C} \setminus \{0\}$ satisfying $\text{Re } \lambda \geq 0$.

Lemma 6.3. *Let $\lambda \in \mathbb{C} \setminus \{0\}$ with $\text{Re } \lambda \geq 0$, $q \in [2, \infty)$, $(f_1, f_2) \in X_m$. Then there exists a unique solution $(\zeta, V) \in D(A_{\text{CHS}}) \cap X_m$ of (6.8) satisfying*

$$(6.14) \quad \|\zeta\|_{H^{1,q}(G)} + |\lambda| \|V\|_{L^q(\Omega)} + \|V\|_{H^{2,q}(\Omega)} \leq C (\|f_1\|_{H^{1,q}(G)} + \|f_2\|_{L^q(\Omega)}).$$

for some $C > 0$.

Proof. As before we first the case $q = 2$ and use then a bootstrap argument to obtain the case $q > 2$.

Step 1: the case $q = 2$

Let us fix $\lambda \in \mathbb{C} \setminus \{0\}$ with $\text{Re } \lambda \geq 0$. Using (6.8)₁ we find $\zeta = \frac{1}{\lambda}(f_1 - \text{div}_H \bar{v})$ and hence the equation (6.8) can be rewritten as

$$(6.15) \quad \lambda v - a_1 \mu \Delta v - \mu \partial_z (b_1 \partial_z v) - a_1 \mu' \nabla_H \text{div}_H v - \frac{1}{\lambda} \nabla_H \text{div}_H \bar{v} = f_1 - \frac{1}{\lambda} \nabla_H f_2, \text{ in } \Omega,$$

where a_1 and b_1 are given as in the proof of Lemma 6.2. Consider the weak formulation of the above problem given by

$$(6.16) \quad \begin{aligned} \lambda(v, \varphi)_2 + \mu (a_1 \nabla_H v, \nabla_H \varphi)_2 + \mu (b_1 \partial_z v, \partial_z \varphi)_2 + \mu' (a_1 \text{div}_H v, \text{div}_H \varphi)_2 \\ + \frac{1}{\lambda} (\text{div}_H \bar{v}, \text{div}_H \bar{\varphi})_{L^2(G)} = (f_2 - \frac{1}{\lambda} \nabla_H f_1, \varphi)_2 \end{aligned}$$

for $\varphi \in \mathcal{V}$. We rephrase the equation (6.16) as

$$\mathcal{C}_\lambda(v, \varphi) = (f_\lambda, \varphi)_{L^2(\Omega)}, \quad \text{for all } \varphi \in \mathcal{V},$$

and note that by Hölder's inequality it is clear that $\mathcal{C}_\lambda : \mathcal{V} \times \mathcal{V} \rightarrow \mathbb{C}$ is a bounded sesquilinear form. Since a_1 and b_1 are uniformly bounded from below, we conclude that \mathcal{C}_λ is coercive as in the proof of Lemma 6.2. Now we apply Lax-Milgram theorem to conclude that there exists a unique solution $v \in \mathcal{V}$. We employ the method of difference quotients to obtain higher regularity estimates and to obtain $v \in H_{\text{per}}^2(\Omega; \mathbb{R}^2)$ as well as $\zeta \in H_{\text{per}}^1(G; \mathbb{R}) \cap L_0^2(G; \mathbb{R})$ from (6.8)₁.

Step 2: the case $q > 2$

Averaging (6.15) we obtain the following inhomogeneous two-dimensional Lamé type equation in G

$$(6.17) \quad -\mu \Delta_H \bar{v} - (\mu' + \lambda') \nabla_H \text{div}_H \bar{v} = \overline{(1 - \delta z) f_2} - \lambda' \nabla_H f_2 + \mu \frac{(1 - \delta)^2}{\delta^2} (\partial_z v)|_{\Gamma_u} + \mu \frac{1}{\delta} v|_{\Gamma_b} + \mu \bar{v} - \overline{\lambda(1 - \delta z)v},$$

where $\lambda' = \frac{1}{\lambda} (1 - \frac{\delta}{2}) > 0$. Note that if \bar{v} is known we may find V by solving the equation

$$(6.18) \quad \begin{cases} \lambda v - a_1 \mu \Delta_H v - \mu \partial_z (b_1 \partial_z v) - a_1 \mu' \nabla_H \text{div}_H v = f_2 + \frac{1}{\lambda} \nabla_H \text{div}_H \bar{v}, & \text{on } \Omega, \\ v = 0, & \text{on } \Gamma_u, \\ \partial_z v = 0, & \text{on } \Gamma_b. \end{cases}$$

Note that solving (6.8) is equivalent to solving the two-dimensional problem (6.17) as well as the elliptic boundary value problem (6.18). Now a bootstrap argument as in the proof of Lemma 6.2 can be used to deduce the existence of a unique solution (ζ, v) to (6.8) within the L^q -framework. \square

We are finally in position to prove that the part of A_{CHS} on $X_m := H_{\text{per}}^{1,q}(G; \mathbb{R}) \cap L_0^q(G; \mathbb{R}) \times L^q(\Omega; \mathbb{R}^2)$ generates an exponentially stable semigroup.

Proposition 6.4. *Let $q \in [2, \infty)$. The compressible hydrostatic Stokes operator A_{CHS} defined on X_m generates an exponentially stable semigroup $(e^{tA_{\text{CHS}}})_{t \geq 0}$ on X_m .*

Proof. Combining Lemma 6.2 and Lemma 6.3 we see that $\{\lambda \in \mathbb{C} : \text{Re } \lambda \geq 0\} \subset \rho(A_{\text{CHS}})$. Moreover, Proposition 4.2 implies that there exist constant $C > 0$, $\omega \geq 0$ and $\varphi \in [0, \pi/2)$ such that

$$\|(\lambda - A_{\text{CHS}})^{-1}\|_{\mathcal{L}(X_m)} \leq C,$$

for all $\lambda \in \omega + \sum_{\pi-\varphi}$. Moreover, the set $\{\lambda \in \mathbb{C} : \text{Re } \lambda \geq 0\} \setminus \{\omega + \sum_{\pi-\varphi}\}$ is compact and we deduce the existence of a constant $C > 0$ such that

$$\|(\lambda - A_{\text{CHS}})^{-1}\|_{\mathcal{L}(X_m)} \leq C \quad \text{for all } \lambda \in \mathbb{C} \text{ with } \text{Re } \lambda \geq 0.$$

We thus conclude that the spectral bound $s(A_{\text{CHS}})$ of A_{CHS} satisfies $s(A_{\text{CHS}}) = -\eta_0$ for some $\eta_0 > 0$. Since the spectral bound of generators of analytic semigroups coincide with the growth bound of the associated semigroups, see e. g., [1, Theorem 5.1.12], the proof is complete. \square

The fact that the spectral bound of A_{CHS} is strictly negative allows us to sharpen the assertion of Proposition 6.1 and to deduce by [6, Proposition 2.7] that $-A_{\text{CHS}} + \omega$ admits a bounded \mathcal{H}^∞ -calculus of \mathcal{H}^∞ -angle $\Phi_{A_{\text{CHS}} + \omega}^\infty < \pi/2$ for all $\omega > -\eta_0$.

Corollary 6.5. *Let $q \in [2, \infty)$ and η_0 as above. Then $-A_{\text{CHS}} + \omega$ admits a bounded \mathcal{H}^∞ -calculus on X_m of \mathcal{H}^∞ -angle $\Phi_{A_{\text{CHS}} + \omega}^\infty < \pi/2$ for all $\omega > -\eta_0$*

Corollary 6.5 allows now to deduce maximal L^p - L^q -regularity estimates for the linearized compressible system

$$(6.19) \quad \left\{ \begin{array}{ll} \partial_t \zeta + \bar{\xi} \text{div}_{\text{H}} \bar{V} = f_1, & \text{on } G \times (0, \infty), \\ \partial_t V - \frac{\mu \Delta_{\text{H}} V}{(1-\delta z)\bar{\xi}} - \frac{\mu}{\bar{\xi}} \partial_z \left(\frac{1-\delta z}{\delta^2} \partial_z V \right) - \frac{\mu' \nabla_{\text{H}} \text{div}_{\text{H}} V}{(1-\delta z)\bar{\xi}} + \frac{\nabla_{\text{H}} \zeta}{\bar{\xi}} = f_2, & \text{on } \Omega \times (0, \infty), \\ \partial_z \zeta = 0, & \text{on } \Omega \times (0, \infty), \\ V = 0, & \text{on } \Gamma_u \times (0, \infty), \\ \partial_z V = 0, & \text{on } \Gamma_b \times (0, \infty), \\ (\zeta(0), V(0)) = (\xi_0 - \bar{\xi}, v_0). \end{array} \right.$$

Note that the trace space $X_\gamma = (X_0, X_1)_{1-1/p}$ for the initial data (ξ_0, v_0) is given by

$$X_\gamma := H_{\text{per}}^{1,q}(G; \mathbb{R}) \cap L_0^q(G; \mathbb{R}) \times X_\gamma^v.$$

It is now natural to decompose a given integrable function $f: G \rightarrow \mathbb{R}$ into

$$f = f_m + f_{\text{avg}}, \quad \text{with} \quad \int_G f_m = 0 \quad \text{and} \quad f_{\text{avg}} = \frac{1}{|G|} \int_G f.$$

Integrating the first equation of (6.19), we deduce that

$$\partial_t \zeta_{\text{avg}} = f_{1,\text{avg}} \quad \text{for } t > 0 \quad \text{and} \quad \zeta_{\text{avg}}(0) = 0.$$

Thus $\zeta_{\text{avg}} \in L^\infty(\mathbb{R}_+)$ provided $f_{1,\text{avg}} \in L^1(\mathbb{R}_+)$. In this case, the set of equations (6.19) is transformed into the set of equations

$$(6.20) \quad \left\{ \begin{array}{ll} \partial_t \zeta_m + \bar{\xi} \operatorname{div}_H \bar{V} = f_1 - f_{1,\text{avg}}, & \text{on } G \times (0, \infty), \\ \partial_t V - \frac{\mu \Delta_H V}{(1-\delta z)\bar{\xi}} - \frac{\mu}{\bar{\xi}} \partial_z \left(\frac{1-\delta z}{\delta^2} \partial_z V \right) - \frac{\mu' \nabla_H \operatorname{div}_H V}{(1-\delta z)\bar{\xi}} + \frac{\nabla_H \zeta_m}{\bar{\xi}} = f_2, & \text{on } \Omega \times (0, \infty), \\ \partial_z \zeta = 0, & \text{on } \Omega \times (0, \infty), \\ V = 0, & \text{on } \Gamma_u \times (0, \infty), \\ \partial_z V = 0, & \text{on } \Gamma_b \times (0, \infty), \\ (\zeta_m(0), V(0)) = (\xi_0 - \bar{\xi}, v_0). \end{array} \right.$$

Proposition 6.6. *Let $\bar{\xi} > 0$, $\eta_0 > 0$ as above and $p, q \in [2, \infty)$ such that $1 \neq \frac{1}{p} + \frac{1}{2q}$ as well as $\frac{1}{2} \neq \frac{1}{p} + \frac{1}{2q}$. Moreover, assume that*

$$e^{\eta(\cdot)}(f_1, f_2) \in \mathbb{E}_{0,\infty}, \quad f_{1,\text{avg}} \in L^1(\mathbb{R}_+) \quad \text{as well as} \quad (\xi_0, v_0) \in X_\gamma.$$

Then the system (6.20) admits a unique, strong solution (ζ_m, v) for all $\eta \in (0, \eta_0)$ satisfying

$$e^{\eta(\cdot)}(\zeta_m, V) \in \mathbb{E}_{1,\infty} \quad \text{and} \quad \zeta_{\text{avg}} \in L^\infty(\mathbb{R}_+).$$

Furthermore, there exists a constant $C > 0$, depending only on p, q and η , such that

$$(6.21) \quad \begin{aligned} & \|e^{\eta(\cdot)}(\zeta_m, V)\|_{\mathbb{E}_{1,\infty}} + \|\zeta_{\text{avg}}\|_{L^\infty(\mathbb{R}_+)} + \|e^{\eta(\cdot)} \partial_t \zeta_{\text{avg}}\|_{L^\infty(\mathbb{R}_+)} \\ & \leq C (\|(\xi_0 - \bar{\xi}, v_0)\|_{X_\gamma} + \|e^{\eta(\cdot)}(f_1, f_2)\|_{\mathbb{E}_{0,\infty}} + \|f_{1,\text{avg}}\|_{L^1(\mathbb{R}_+)}). \end{aligned}$$

Proof. The proof is a direct consequence of Corollary 6.5. It yields that the equation (6.20) admits a unique solution (ζ_m, V) satisfying (6.21). \square

6.2. Estimates for the nonlinear terms.

In the following, we estimate the nonlinear terms given in (6.2). Given X as in Section 3, we note that there is a constant $C > 0$ such that for all $\varepsilon \in (0, \frac{1}{2C})$ we have

$$\begin{aligned} \|\nabla_H X - \mathbb{I}_2\|_{L^\infty(\mathbb{R}_+ \times G)} & \leq C \|\nabla_H X - \mathbb{I}_2\|_{L^\infty(\mathbb{R}_+; H^{1,q}(G))} \leq C \int_0^\infty e^{-\eta t} e^{\eta t} \|\nabla_H \bar{V}(t, \cdot)\|_{H^{1,q}(G)} dt \\ & \leq C \left(\int_0^\infty e^{-p' \eta t} dt \right)^{1/p'} \|e^{\eta(\cdot)} V\|_{\mathbb{E}_{1,\infty}} \leq C \varepsilon \leq \frac{1}{2}. \end{aligned}$$

Next, we conclude that $\det \nabla_H X \geq C > 0$ in $\mathbb{R}_+ \times G$ and obtain from (3.7) that $\|Z\|_{L^\infty(\mathbb{R}_+; H^{1,q}(G))} \leq C$. Therefore,

$$\|Z - \mathbb{I}_2\|_{L^\infty(\mathbb{R}_+; H^{1,q}(G))} \leq \|Z\|_{L^\infty(\mathbb{R}_+; H^{1,q}(G))} \|\nabla_H X - \mathbb{I}_2\|_{L^\infty(\mathbb{R}_+; H^{1,q}(G))} \leq C \varepsilon.$$

Notice that $\partial_t \nabla_H X = \nabla_H \bar{V}$ and hence

$$\|\partial_t \nabla_H X\|_{L^p(\mathbb{R}_+; H^{1,q}(G))} \leq \|e^{\eta(\cdot)} \nabla_H \bar{V}\|_{L^p(\mathbb{R}_+; H^{1,q}(G))} \leq C \varepsilon.$$

We also have

$$\partial_t (Z - \mathbb{I}_2) = \partial_t Z = -(\nabla_H X)^{-1} (\partial_t \nabla_H X) (\nabla_H X)^{-1} = -Z (\partial_t \nabla_H X),$$

and thus

$$\|\partial_t (Z - \mathbb{I}_2)\|_{L^p(\mathbb{R}_+; H^{1,q}(G))} = \|\partial_t Z\|_{L^p(\mathbb{R}_+; H^{1,q}(G))} \leq C \varepsilon.$$

We fix now $\eta \in (0, \eta_0)$ and choose ε so small that the transform X is indeed a C^1 -diffeomorphism. Moreover, for $\varepsilon > 0$ define

$$(6.22) \quad \mathbb{B}_\varepsilon := \{(\zeta, V) \in \mathbb{E}_{1,\infty} \cap X_m : \|e^{\eta(\cdot)}(\zeta_m, V)\|_{\mathbb{E}_{1,\infty}} + \|\zeta_{\text{avg}}\|_{L^\infty(\mathbb{R}_+)} + \|e^{\eta(\cdot)} \partial_t \zeta_{\text{avg}}\|_{L^\infty(\mathbb{R}_+)} \leq \varepsilon\}.$$

We then obtain the following estimates for the nonlinear terms.

Proposition 6.7. *Let $p, q \in (2, \infty)$. There exists a constant $C > 0$, depending only on p, q and η , such that for all ε sufficiently small and all $(\zeta, V) \in \mathbb{B}_\varepsilon$, the functions F_1, F_2 and $F_{1,\text{avg}}$ satisfy*

$$\|e^{\eta(\cdot)}(F_1, F_2)\|_{\mathbb{E}_{0,\infty}} + \|F_{1,\text{avg}}\|_{L^1(\mathbb{R}_+)} \leq C\varepsilon^2.$$

Proof. We note that for every $(\zeta, V) \in \mathbb{B}_\varepsilon$ we have

$$(6.23) \quad \|\zeta\|_{L^\infty(\mathbb{R}_+; H^{1,q}(G))} \leq \|e^{\eta(\cdot)}\zeta_m\|_{L^\infty(\mathbb{R}_+; H^{1,q}(G))} + \|\zeta_{\text{avg}}\|_{L^\infty(\mathbb{R}_+)} \leq C\varepsilon,$$

as well as

$$(6.24) \quad \|V\|_{L^\infty(\mathbb{R}_+; H^{1,q}(\Omega))} + \|\nabla V\|_{L^p(\mathbb{R}_+; L^\infty(\Omega))} \leq C\varepsilon,$$

due to the embedding $\mathbb{E}_1^q \hookrightarrow L^\infty(\mathbb{R}_+; H^{1,q}(\Omega))$. Recalling that

$$e^{\eta t} F_1 = -e^{\eta t} \zeta \operatorname{div}_H \bar{V} - e^{\eta t} (\zeta + \bar{\xi}) \nabla_H \bar{V} : [Z^\top - \mathbb{I}_2],$$

and hence

$$F_{1,\text{avg}} = -\frac{1}{|G|} \int_G \zeta \operatorname{div}_H \bar{V} \, d(x, y) - \frac{1}{|G|} \int_G (\zeta + \bar{\xi}) \nabla_H \bar{V} : [Z^\top - \mathbb{I}_2] \, d(x, y),$$

we obtain

$$\begin{aligned} \|e^{\eta t} F_1\|_{L^p(\mathbb{R}_+; H^{1,q}(G))} &\leq \|(\zeta + \bar{\xi})\|_{L^\infty(\mathbb{R}_+; H^{1,q}(G))} \|Z^\top - \mathbb{I}_2\|_{L^\infty(\mathbb{R}_+; H^{1,q}(G))} \|e^{\eta(\cdot)} \nabla_H \bar{V}\|_{L^p(\mathbb{R}_+; H^{1,q}(G))} \\ &\quad + \|\zeta\|_{L^\infty(\mathbb{R}_+; H^{1,q}(G))} \|e^{\eta(\cdot)} \operatorname{div}_H \bar{V}\|_{L^p(\mathbb{R}_+; H^{1,q}(G))} \\ &\leq C\varepsilon^2. \end{aligned}$$

Next, we estimate the second term of $F_{1,\text{avg}}$ as

$$\begin{aligned} &\left\| \int_G (\zeta + \bar{\xi}) \nabla_H \bar{V} : [Z^\top - \mathbb{I}_2] \right\|_{L^1(\mathbb{R}_+)} \\ &\leq \|(\zeta + \bar{\xi})\|_{L^\infty(\mathbb{R}_+ \times G)} \|Z^\top - \mathbb{I}_2\|_{L^\infty(\mathbb{R}_+ \times G)} \int_0^\infty \int_G \nabla_H \bar{V} \, d(x, y) \, dt \\ &\leq C \|(\zeta + \bar{\xi})\|_{L^\infty(\mathbb{R}_+; H^{1,q}(G))} \|Z^\top - \mathbb{I}_2\|_{L^\infty(\mathbb{R}_+; H^{1,q}(G))} \int_0^\infty e^{-\eta t} e^{\eta t} \|\nabla_H \bar{V}\|_{L^q(G)} \, dt \\ &\leq C\varepsilon \left(\int_0^\infty e^{-p'\eta t} \, dt \right)^{1/p'} \|e^{\eta(\cdot)} \nabla_H \bar{V}\|_{L^p(\mathbb{R}_+; L^q(G))} \\ &\leq C\varepsilon^2. \end{aligned}$$

The first term of $F_{1,\text{avg}}$ can be estimated similarly. The same is true for F_2 since all nonlinear terms are at least quadratic in terms of V, ζ or $Z - \mathbb{I}_2$. We only show this in detail for the third term of the first line of F_2 and obtain

$$\begin{aligned} \left\| e^{\eta(\cdot)} \left(\sum_{l,j,k} Z_{l,k} \frac{\partial V_i}{\partial y_k} \frac{\partial Z_{k,j}}{\partial y_l} \right) \right\|_{L^p(\mathbb{R}_+; L^q(\Omega))} &= \left\| e^{\eta(\cdot)} \left(\sum_{l,j,k} Z_{l,k} \frac{\partial V_i}{\partial y_k} \frac{\partial (Z_{k,j} - \delta_{k,j})}{\partial y_l} \right) \right\|_{L^p(\mathbb{R}_+; L^q(\Omega))} \\ &\leq C \|Z\|_{L^\infty(\mathbb{R}_+; L^\infty(G))} \|e^{\eta(\cdot)} V\|_{\mathbb{E}_{1,\infty}^q} \|Z - \mathbb{I}_2\|_{L^\infty(\mathbb{R}_+; H^{1,q}(G))} \\ &\leq C\varepsilon^2. \end{aligned}$$

Hence,

$$\|e^{\eta(\cdot)} F_2\|_{L^p(\mathbb{R}_+; L^q(\Omega))} \leq C\varepsilon^2. \quad \square$$

We are finally in position to give a proof of [Theorem 2.3](#).

Proof of Theorem 2.3. Let \mathbb{B}_ε be given as in (6.22). For $(\zeta_1, V_1) \in \mathbb{B}_\varepsilon$ denote by $(\zeta, V) := \Psi(\zeta_1, V_1) \in \mathbb{E}_{1,\infty}$ the unique solution of the linearized equation (6.1) with right-hand sides $F_1(\zeta_1, V_1)$ and $F_2(\zeta_1, V_1)$ guaranteed by [Proposition 6.7](#). We will show that Ψ is a selfmap and a contraction. Choosing the initial data small enough, i. e.,

$$\|(\zeta_0 - \bar{\xi}, v_0)\|_{X_\gamma} \leq \frac{\varepsilon}{2C},$$

where $C > 0$ denotes the constant given in [Proposition 6.6](#). The estimates (6.21) and [Proposition 6.7](#) yield

$$\begin{aligned} & \|e^{\eta(\cdot)}(\zeta_m, V)\|_{\mathbb{E}_{1,\infty}} + \|\zeta_{\text{avg}}\|_{L^\infty(\mathbb{R}_+)} + \|e^{\eta(\cdot)}\partial_t(\zeta_{\text{avg}})\|_{L^\infty(\mathbb{R}_+)} \\ & \leq C(\|(\xi_0 - \bar{\xi}, v_0)\|_{X_\gamma} + \|e^{\eta(\cdot)}(F_1, F_2)\|_{\mathbb{E}_{0,\infty}} + \|F_{1,\text{avg}}\|_{L^\infty(\mathbb{R}_+)}) \leq \frac{\varepsilon}{2} + C\varepsilon^2. \end{aligned}$$

We conclude that for $\varepsilon > 0$ small enough, Ψ is a self-map, i. e., $\Psi: \mathbb{B}_\varepsilon \rightarrow \mathbb{B}_\varepsilon$. The contraction property follows in the same way.

Finally, since ε may be chosen to be sufficiently small, the transform $X(t, \cdot)$ is a C^1 -diffeomorphism from G into G and we set $Y(t, \cdot) = X^{-1}(t, \cdot)$. Defining

$$\xi(t, x, y) := \zeta(t, Y(t, x, y)) \quad \text{and} \quad v(t, x, y, z) := V(t, Y(t, x, y), z)$$

we verify that (ξ, v) is a global, strong solution to system (2.4) satisfying $(\xi, v) \in \mathbb{E}_{1,\infty}$. \square

Proof of [Theorem 2.4](#). We will not provide a detailed proof of [Theorem 2.4](#) as its strategy is similar to the one described in the proofs of [Theorem 2.2](#) and [Theorem 2.3](#). A short discussion of the main differences is nevertheless in order. The hydrostatic Lagrange transformation yields the operator $A = \frac{\mu}{\xi_0}\Delta + \frac{\mu'}{\xi_0}\nabla_{\text{H}} \operatorname{div}_{\text{H}}$ on $L^q(\Omega; \mathbb{R}^2)$ and we show as in [Section 4](#) that $-A + \omega$ admits a bounded \mathcal{H}^∞ -calculus on $L^q(\Omega; \mathbb{R}^2)$ of \mathcal{H}^∞ -angle strictly less than $\pi/2$. The estimates on the nonlinear terms differ only by the pressure term. For the local well-posedness we estimate the transformed pressure $P'(\zeta)(Z^\top \nabla_{\text{H}} \zeta)$ as in the proof of [Lemma 5.1](#).

To deduce the global well-posedness result for small data we rewrite the transformed pressure term as

$$P'(\zeta + \bar{\xi})(Z^\top \nabla_{\text{H}} \zeta) = \nabla_{\text{H}}(P(\zeta)) + P'(\zeta + \bar{\xi}) \sum_j \frac{\partial \zeta}{\partial y_j} (Z_{j,i} - \delta_{j,i}).$$

Setting $\tilde{\zeta} = P(\zeta)$, we solve for $\tilde{\zeta}$ similarly as in [Theorem 2.3](#). The pressure law (2.7) ensures then the invertibility of P and hence for proving the assertion it is sufficient to solve the resulting system for $\tilde{\zeta}$. \square

Acknowledgements This research is supported by the Basque Government through the BERC 2022-2025 program and by the Spanish State Research Agency through BCAM Severo Ochoa excellence accreditation CEX2021-01142-S funded by MICIU/AEI/10.13039/501100011033 and through Grant PID2023-146764NB-I00 funded by MICIU/AEI/10.13039/501100011033 and cofunded by the European Union. Matthias Hieber acknowledges the support by DFG project FOR 5528. Arnab Roy would like to thank the Alexander von Humboldt-Foundation, Grant RYC2022-036183-I funded by MICIU/AEI/10.13039/501100011033 and by ESF+. Yoshiki Iida was supported by JSPS KAKENHI Grant Number JP24KJ2080, and partially supported by Top Global University Project of Waseda University. He is also grateful to Fachbereich Mathematik, TU Darmstadt for its hospitality while visiting Professor Matthias Hieber.

REFERENCES

- [1] W. Arendt, Ch. Batty, M. Hieber, F. Neubrander, *Vector-Valued Laplace Transforms and Cauchy Problems*. Monographs in Mathematics, vol. 96, 2nd-edition, Birkhäuser, Basel, 2011.
- [2] H. Amann, *Linear and Quasilinear Parabolic Problems*. Monographs in Mathematics, vol. 89, Birkhäuser, 1995.
- [3] H. Amann, *Linear and Quasilinear Parabolic Problems. Vol. II. Function Spaces*. Monographs in Mathematics, vol. 106, Birkhäuser/Springer, Cham, 2019.
- [4] C. Cao, E.S. Titi, Global well-posedness of the three-dimensional viscous primitive equations of large scale ocean and atmosphere dynamics. *Ann. of Math.* **166** (2007), 245–267.
- [5] R. Denk, G. Dore, M. Hieber, J. Prüss, A. Venni, New thoughts on old results of R. T. Seeley. *Math. Ann.* **328** (2004), 545–583.
- [6] R. Denk, M. Hieber, J. Prüss, *\mathcal{R} -Boundedness, Fourier Multipliers and Problems of Elliptic and Parabolic Type*. Mem. Amer. Math. Soc. **788** 2003.
- [7] R. Denk, M. Hieber, J. Prüss, Optimal L^p - L^q -estimates for parabolic boundary value problems with inhomogeneous data. *Math. Z.* **257** (2007), 193–224.
- [8] R. Danchin, A Lagrangian approach for the compressible Navier-Stokes equations. *Ann. de l'Institut Fourier* **64** (2014), 753–791.
- [9] R. Danchin, Fourier analysis methods for the compressible Navier-Stokes equations. *Handbook of Math. Anal. in Mechanics of Viscous Fluids*, Y. Giga, A. Novotny (eds.), Springer, 2018, 1843–1903.

- [10] R. Danchin, P. Mucha, Compressible Navier-Stokes equations with ripped density. *Comm. Pure Appl. Math.* **76** (2023), 3437–3492.
- [11] R. Denk, T. Nau, Discrete Fourier multipliers and cylindrical boundary-value problems. *Proc. Roy. Soc. Edinburgh* **143** (2013), 1163–1183.
- [12] Y. Enomoto, Y. Shibata, Global Existence of Classical Solutions and Optimal Decay Rate for Compressible Flows via the Theory of Semigroups. *Handbook of Math. Anal. in Mechanics of Viscous Fluids*, Y. Giga, A. Novotny (eds.), Springer, 2018, 2085–2181.
- [13] M. Ersoy, T. Ngom, Existence of a global weak solution to compressible primitive equations. *C.R. Acad. Sci.* **350**, (2012), 379–382.
- [14] M. Ersoy, T. Ngom, M. Sy, Compressible primitive equations: formal derivation and stability of weak solutions. *Nonlinearity* **24**, (2011), 79–96.
- [15] E. Feireisl, A. Novotný and H. Petzeltová, On the existence of globally defined weak solutions to the Navier-Stokes equations, *J. Math. Fluid Mech.* **3** (2001), no. 4, 358–392.
- [16] Y. Giga, M. Gries, M. Hieber, A. Hussein, T. Kashiwabara, Bounded \mathcal{H}^∞ -calculus for the hydrostatic Stokes operator on L^p -spaces and applications. *Proc. Amer. Math. Soc.* **145** (2017), 3865–3876.
- [17] Y. Giga, M. Gries, M. Hieber, A. Hussein, T. Kashiwabara, Analyticity of solutions to the primitive equations. *Math. Nachr.* **293** (2020), 284–304.
- [18] R. Haller-Dintelmann, M. Hieber, \mathcal{H}^∞ -calculus for products of non-commuting operators. *Math. Z.* **251** (2005), 85–100.
- [19] M. Hieber, T. Kashiwabara, Global strong well-posedness of the three dimensional primitive equations in L^p -spaces. *Arch. Rational Mech. Anal.* **221** (2016), 1077–1115.
- [20] D. Hoff, Dynamics of singularity surfaces for compressible, viscous flows in two space dimensions. *Comm. Pure Appl. Math.* **55** (2002), 1365–1407.
- [21] T. Hytönen, J. van Neerven, M. Veraar, L. Weis, *Analysis in Banach spaces, vol. II*, Springer, Cham, 2016.
- [22] I. Kukavica, M. Ziane, On the regularity of the primitive equations of the ocean. *Nonlinearity* **20** (2007), 2739–2753.
- [23] J.-L. Lions, R. Temam, S.H. Wang, New formulations of the primitive equations of atmosphere and applications. *Nonlinearity* **5** (1992), 237–288.
- [24] J.-L. Lions, R. Temam, S.H. Wang, On the equations of the large-scale ocean. *Nonlinearity* **5** (1992), 1007–1053.
- [25] J.-L. Lions, R. Temam, S.H. Wang, Mathematical theory for the coupled atmosphere-ocean models. (CAO III). *J. Math. Pures Appl. (9)* **74** (1995), 105–163.
- [26] P.-L. Lions, Mathematical topics in fluid mechanics. Vol. 2, *Oxford Lecture Series in Mathematics and its Applications*, Oxford Science Publications, 10, Oxford Univ. Press, New York, 1998.
- [27] X. Liu, E. Titi, Global existence of weak solutions to the compressible primitive equations of atmospheric dynamics with degenerate viscosities. *SIAM J. Math. Anal.* **51** (2019), 1913–1964.
- [28] X. Liu, E. Titi, Zero Mach number limit of the compressible primitive equations part I: Well-prepared initial data. *Arch. Rational Mech. Anal.* **238** (2020), 705–747.
- [29] X. Liu, E. Titi, Local well-posedness of strong solutions to the three-dimensional compressible primitive equations, *Arch. Rational Mech. Anal.* **241** (2021), 729–764.
- [30] A. Matsumura, T. Nishida, Initial boundary value problems for the equations of motion of compressible viscous and heat-conductive fluids, *Comm. Math. Phys.* **89** (1983), 445–464.
- [31] T. Nau, *L^p -Theory of Cylindrical Boundary Value Problems*. Springer Spektrum, Wiesbaden, 2012, xvi+188 pp.
- [32] J. Prüss, G. Simonett, *Moving Interfaces and Quasilinear Parabolic Evolution Equations*. Monographs in Mathematics, vol. 105, Birkhäuser, 2016.
- [33] H. Triebel, *Interpolation Theory, Function Spaces, Differential Operators*. North-Holland, 1978.

TECHNISCHE UNIVERSITÄT DARMSTADT, FACHBEREICH MATHEMATIK, SCHLOSSGARTENSTR. 7, 64289 DARMSTADT, GERMANY

Email address: hieber@mathematik.tu-darmstadt.de

DEPARTMENT OF PURE AND APPLIED MATHEMATICS, GRADUATE SCHOOL OF FUNDAMENTAL SCIENCE AND ENGINEERING, WASEDA UNIVERSITY, 169-8555 TOKYO, JAPAN

Email address: yoshiki-i737@asagi.waseda.jp

BASQUE CENTER FOR APPLIED MATHEMATICS (BCAM), ALAMEDA DE MAZARREDO 14, 48009 BILBAO, SPAIN.

IKERBASQUE, BASQUE FOUNDATION FOR SCIENCE, PLAZA EUSKADI 5, 48009 BILBAO, BIZKAIA, SPAIN.

Email address: aroy@bcamath.org

TECHNISCHE UNIVERSITÄT DARMSTADT, FACHBEREICH MATHEMATIK, SCHLOSSGARTENSTR. 7, 64289 DARMSTADT, GERMANY

Email address: zoechling@mathematik.tu-darmstadt.de