THE DIRICHLET PROBLEM FOR ELLIPTIC EQUATIONS WITH SINGULAR DRIFT TERMS

S. HOFMANN AND A. PATHAK

CONTENTS

1. Introduction, history, and statement of main result	1
2. Notation, terminology, and some known geometric results	3
2.1. Some geometric results	7
3. Preliminaries	11
3.1. A priori estimates	12
3.2. Existence of $Y^{1,2}$ solutions with data in $\operatorname{Lip}_c(\partial \Omega)$, in the small constant case	16
3.3. Weak maximum principle and continuity of solutions with $\operatorname{Lip}_{c}(\partial \Omega)$ data	17
3.4. Existence of elliptic measure in the small constant case, and a local ampleness property	
3.5. Existence of elliptic measure in the large constant case, and an approximation result 19	
3.6. The Green function: existence, estimates, and consequences	21
4. Perturbing L_0 in a small constant case	25
5. Two key lemmas	29
6. Proof of Theorem 1.7: the extrapolation argument	35
References	44

18

ABSTRACT. We establish L^p solvability of the Dirichlet problem, for some finite p, in a 1sided chord-arc domain Ω (i.e., a uniform domain with Ahlfors-David regular boundary), for elliptic equations of the form

 $Lu = -\operatorname{div}(A\nabla u) + \mathbf{B} \cdot \nabla u =: L_0 u + \mathbf{B} \cdot \nabla u = 0,$

given that the analogous result holds (typically with a different value of *p*) for the homogeneous second order operator L_0 . Essentially, we assume that $|\mathbf{B}(X)| \leq \operatorname{dist}(X, \partial \Omega)^{-1}$, and that $|\mathbf{B}(X)|^2 \operatorname{dist}(X, \partial \Omega) dX$ is a Carleson measure in Ω .

1. INTRODUCTION, HISTORY, AND STATEMENT OF MAIN RESULT

In this work, we treat the Dirichlet problem with data in L^p (denoted $(D)_p$), for the equation Lu = 0, where L is an elliptic operator with a drift term, i.e.,

(1.1)
$$Lu := -\operatorname{div}(A\nabla u) + \mathbf{B} \cdot \nabla u =: L_0 u + \mathbf{B} \cdot \nabla u,$$

Date: February 7, 2025.

²⁰²⁰ Mathematics Subject Classification. 35J15, 35J25, 42B37.

Key words and phrases. Dirichlet problem, elliptic equations, elliptic measure, drift terms, Carleson measures.

The first author was partially supported by NSF grant DMS-2349846.

S. HOFMANN AND A. PATHAK

in the setting of a "1-sided chord arc domain", i.e., a uniform domain $\Omega \subset \mathbb{R}^{n+1}$ with (*n*-dimensional) Ahlfors-David regular boundary (all notation and terminology will be defined in the sequel). Our results are perturbative in nature: we assume that the Dirichlet problem is solvable for the purely second order operator L_0 , with data in L^p for some finite p, and show that the analogous solvability result holds (for a possibly different p) for the operator L. Writing $\delta(X) := \operatorname{dist}(X, \partial \Omega)$, we assume in particular that for some $M_0 < \infty$, **B** satisfies

(1.2)
$$|\mathbf{B}(X)| \leq \sqrt{M_0} \,\delta(X)^{-1} \,, \qquad \text{a.e. } X \in \Omega \,,$$

and that $d\mu(X) := |\mathbf{B}(X)|^2 \delta(X) dX$ is a Carleson measure in Ω , i.e.,

(1.3)
$$\|\mathbf{B}\|_{T^{2,\infty}(\Omega)} := \sup_{B} r_B^{-n} \iint_{\Omega \cap B} |\mathbf{B}(X)|^2 \,\delta(X) \, dX \leq M_0 < \infty \,,$$

where the supremum runs over all balls $B \subset \mathbb{R}^{n+1}$ centered on $\partial\Omega$, and r_B is the radius of B. More precisely, we consider a slightly stronger version of (1.3), in which **B** is replaced by its essential supremum on a Whitney ball, i.e., for $X \in \Omega$, set

(1.4)
$$\mathbf{B}_*(X) := \operatorname{ess\,sup}_{Y: |X-Y| < \delta(X)/4} |\mathbf{B}(Y)|,$$

and we suppose that

(1.5)
$$M_1 := \|\mathbf{B}_*\|_{T^{2,\infty}(\Omega)} := \sup_B r_B^{-n} \iint_{\Omega \cap B} \left(\mathbf{B}_*(X)\right)^2 \delta(X) \, dX < \infty,$$

We observe that (1.5) implies each of (1.2) and (1.3) (with $M_0 \leq M_1$).

The principal part of the operator is assumed to be uniformly elliptic, i.e., *A* is an $(n + 1) \times (n + 1)$ matrix of (real) bounded measurable coefficients, not necessarily symmetric, defined on $\Omega \subset \mathbb{R}^{n+1}$, satisfying the uniform ellipticity condition

(1.6)
$$\lambda |\xi|^2 \le \langle A(X)\xi,\xi\rangle := \sum_{i,j=1}^{n+1} A_{ij}(X)\xi_j\xi_i, \quad ||A||_{L^{\infty}(\Omega)} \le \Lambda,$$

for some $\lambda > 0$, $\Lambda < \infty$, and for all $\xi \in \mathbb{R}^{n+1}$, and a.e. $X \in \Omega$.

Our main result is as follows.

Theorem 1.7. Let $\Omega \subset \mathbb{R}^{n+1}$, $n \ge 2$, be a 1-sided chord arc domain (CAD), and let L_0 and L be defined as in (1.1), where the coefficients of the drift term, and of the principal part, satisfy (respectively) (1.5) and (1.6). Suppose that for some $p_0 < \infty$, the Dirichlet problem $(D)_{p_0}$ is solvable for the equation $L_0u = 0$ in Ω . Then there is a finite p (possibly larger than p_0), such that $(D)_p$ is solvable for Lu = 0 in Ω .

Of course, this result is quantitative, i.e., the exponent p, as well as the constants that will appear in our estimates for solutions of $(D)_p$, will depend only on the "allowable parameters": dimension, the constants in (1.5) and (1.6), the 1-sided chord arc constants for Ω and its boundary, the exponent p_0 , and the constants in the quantitative estimates for solutions of $(D)_{p_0}$ for L_0 .

A previous result of this type, valid in Lipschitz domains, had been proved by J. L. Lewis and the first named author of the present paper [HL] (see also [KP], for related results). In the Lipschitz setting, one can exploit the use of a pullback mechanism to reduce matters to the case that Ω is the half-space \mathbb{R}^{n+1}_+ . Such an approach is clearly unavailable in the much more general setting of a 1-sided chord arc domain, the boundary of which need not be given locally as a graph. In [HL], the authors proved a parabolic version of Theorem 1.7, valid in "regular Lip(1,1/2) domains" (see, e.g., [BHMN] for the definition) again using a pullback argument. Indeed, the parabolic case was the main focus of the work [HL]. The elliptic analogue (again, in a Lipschitz domain), can be gleaned from the corresponding parabolic arguments, but in [HL], the authors presented a much simpler putative proof in the elliptic case, based on a claimed proof of doubling of the elliptic measure for *L*. In fact, the claimed proof of doubling was erroneous¹, so that the purely elliptic argument in [HL] is not valid. On the other hand, the parabolic arguments of [HL] are quite delicate and complicated (and lengthy), so a fringe benefit of our work here is that we give a direct (and simpler) proof in the elliptic setting. Of course, our main contribution, as noted above, is that we are able to obtain new results in a much more general class of domains than those treated in [HL].

Outline of the paper. The paper is organized as follows. In Section 2, we introduce our notation and terminology, and we record some previously known geometric facts. In Section 3, we present a number of preliminary results², well-known for the homogeneous second order $L_0 = -\operatorname{div} A \nabla$, which we extend to the operator $L = L_0 + \mathbf{B} \cdot \nabla$ under certain conditions (e.g., sometimes, but not always, smallness in the pointwise size condition (1.2)). We also give an approximation result, which will allow us to work with operators for which the drift coefficient \mathbf{B} has been truncated near the boundary, and is thus (qualitatively) bounded. Of course, all of our quantitative estimates will be independent of such truncations. There is a considerable amount of material to review in Section 3, but on the other hand, most of it is fairly routine and standard, so the expert reader could probably skim this section rapidly, at least on a first reading. Sections 4, 5, and 6 are the heart of the matter. In Section 4, we present a small constant perturbation result, in the spirit of [FKP, Theorem 2.5], in which we show that the elliptic measures for L and L_0 , denoted by ω and ω_0 , are mutually absolutely continuous in the sense of A_{∞} , given that **B** is small in the sense of a certain ω_0 -adapted Carleson measure condition. In Section 5, we prove two lemmas that will play a key role in removing the smallness assumption in Section 4. Finally, we carry out the latter main step in Section 6, removing the small constant restriction via the method of "extrapolation of Carleson measures", a technique introduced in [LM] and and used also in [HL].

2. NOTATION, TERMINOLOGY, AND SOME KNOWN GEOMETRIC RESULTS

In the sequel, we shall assume that $\Omega \subset \mathbb{R}^{n+1}$ is an open set (typically with additional properties to be specified below), we let B = B(X, r) denote the Euclidean ball in \mathbb{R}^{n+1} of radius *r*, centered at *X*, and for $X \in \partial \Omega$, we let

$$\Delta = B \cap \partial \Omega$$

denote the corresponding surface ball on the boundary. We list further notation as follows.

• We use the letters c, C to denote harmless positive constants, not necessarily the same at each occurrence, which depend only on the "allowable parameters": dimension, and the constants appearing in the hypotheses of the main theorem. We shall also sometimes write $a \leq b$ and $a \approx b$ to mean, respectively, that $a \leq Cb$ and $0 < c \leq a/b \leq C$, where the constants *c* and *C* are as above, unless explicitly noted to the contrary. At times, we shall designate by *M* a particular constant whose value will remain unchanged throughout the

¹Indeed, the argument depended in turn on a claimed proof of a pointwise upper bound for the Green function, which is valid (as we shall show in the sequel) when M_0 is sufficiently small in (1.2), but false in general: the second named author has presented a counter-example [P].

²e.g., existence of solutions and elliptic measure, continuity of solutions up to the boundary, Green function estimates, etc.

proof of a given lemma or proposition, but which may have a different value during the proof of a different lemma or proposition.

- Given a domain Ω ⊂ ℝⁿ⁺¹, we shall typically use lower case letters x, y, z, etc., to denote points on ∂Ω, and capital letters X, Y, Z, etc., to denote generic points in ℝⁿ⁺¹ (especially those in ℝⁿ⁺¹ \ ∂Ω).
- Given a Euclidean ball B or surface ball Δ , its radius will be denoted r_B or r_{Δ} , respectively.
- Given a Euclidean or surface ball B = B(X, r) or $\Delta = \Delta(x, r)$, its concentric dilate by a factor of $\kappa > 0$ will be denoted by $\kappa B := B(X, \kappa r)$ or $\kappa \Delta := \Delta(x, \kappa r)$.
- For $X \in \Omega$, we set $\delta(X) := \operatorname{dist}(X, \partial \Omega)$.
- We let \mathcal{H}^n denote *n*-dimensional Hausdorff measure, and let $\sigma := \mathcal{H}^n|_{\partial\Omega}$ denote the surface measure on $\partial\Omega$.
- For a Borel set E ⊂ ℝⁿ⁺¹, we let 1_E denote the usual indicator function of E, i.e. 1_E(x) = 1 if x ∈ E, and 1_E(x) = 0 if x ∉ E.
- For a Borel set E ⊂ ℝⁿ⁺¹, we let int(E) denote the interior of E. If E ⊂ ∂Ω, then int(E) will denote the relative interior, i.e., the largest relatively open set in ∂Ω contained in E. Thus, ∂E := E \ int(E) will denote the boundary of a set E ⊂ ∂Ω.
- For a Borel set *E*, we denote by C(E) (resp. Lip(*E*)) the space of continuous (resp. Lipschitz continuous) functions on *E*, and by $C_c(E)$ (resp. Lip_c(*E*)) the subspace of C(E) (resp. Lip(*E*)) with compact support in *E*.
- For a Borel subset $E \subset \partial \Omega$, we set $\oint_E f d\sigma := \sigma(E)^{-1} \int_E f d\sigma$.
- We shall use the letter *I* (and sometimes *J*) to denote a closed (n + 1)-dimensional Euclidean cube with sides parallel to the co-ordinate axes, and we let $\ell(I)$ denote the side length of *I*. We use *Q* to denote a dyadic "cube" on $\partial\Omega$. The latter exist, given that $\partial\Omega$ is ADR (cf. [DS1], [Chr]), and enjoy certain properties which we enumerate in Lemma 2.4 below.

Definition 2.1. (Ahlfors-David regular, or ADR). We say that a closed set $E \subset \mathbb{R}^{n+1}$ is *n*-dimensional ADR (or simply ADR) ("Ahlfors-David regular") if there is some uniform constant *C* such that

(2.2)
$$\frac{1}{C}r^n \leq \mathcal{H}^n(E \cap B(x,r)) \leq Cr^n, \quad \forall r \in (0,R_0), x \in E,$$

where R_0 is the diameter of E (which may be infinite). When $E = \partial \Omega$, the boundary of a domain Ω , we shall sometimes for convenience simply say that " Ω has the ADR property" to mean that $\partial \Omega$ is ADR.

Definition 2.3. (Doubling) We say that a measure m defined on $\partial\Omega$ is *doubling* (with doubling constant N_{db}), if for every surface ball $\Delta \subset \partial\Omega$, we have

$$\mathfrak{m}(2\Delta) \leq N_{db} \mathfrak{m}(\Delta)$$

Lemma 2.4. (Existence and properties of the "dyadic grid") [Chr, HK]. Suppose that $E \subset \mathbb{R}^{n+1}$ is an ADR set. Then there exist constants $a_0 > 0$, s > 0 and $C_1 < \infty$, depending only on n and the ADR constant, such that for each $k \in \mathbb{Z}$, there is a collection of Borel sets ("cubes")

$$\mathbb{D}_k := \{ Q_j^k \subset E : j \in \mathfrak{I}_k \},\$$

where \Im_k denotes some (possibly finite) index set depending on k, satisfying

4

- (i) $E = \bigcup_{i} Q_{i}^{k}$ for each $k \in \mathbb{Z}$.
- (ii) If $m \ge k$ then either $Q_i^m \subset Q_i^k$ or $Q_i^m \cap Q_i^k = \emptyset$.
- (iii) For each (j,k) and each m < k, there is a unique i such that $Q_j^k \subset Q_i^m$.
- (*iv*) diam $(Q_i^k) \le C_1 2^{-k}$.
- (v) Each Q_i^k contains some "surface ball" $\Delta(x_i^k, a_0 2^{-k}) := B(x_i^k, a_0 2^{-k}) \cap E$.
- (vi) $\mathcal{H}^n(\{x \in Q_j^k : \operatorname{dist}(x, E \setminus Q_j^k) \le \vartheta 2^{-k}\}) \le C_1 \vartheta^s \mathcal{H}^n(Q_j^k)$, for all k, j and for all $\vartheta \in (0, a_0)$.

A few remarks are in order concerning this lemma.

- In the setting of a general space of homogeneous type, this lemma has been proved by Christ [Chr] (see also [HK]), with the dyadic parameter 1/2 replaced by some constant δ ∈ (0, 1). In fact, one may always take δ = 1/2 (see [HMMM, Proof of Proposition 2.12]). In the presence of the Ahlfors-David property, the result already appears in [DS1, DS2]. Some predecessors of this construction have appeared in [D1] and [D2].
- For our purposes, we may ignore those k ∈ Z such that 2^{-k} ≥ diam(E), in the case that the latter is finite.
- We shall denote by $\mathbb{D} = \mathbb{D}(E)$ the collection of all relevant Q_i^k , i.e.,

$$\mathbb{D}:=\cup_k\mathbb{D}_k,$$

where, if diam(*E*) is finite, the union runs over those *k* such that $2^{-k} \leq \text{diam}(E)$.

• Properties (*iv*) and (*v*) imply that there are uniform constants *c*, *C* such that for each cube $Q \in \mathbb{D}_k$, there is a point $x_Q \in Q$, a Euclidean ball $B(x_Q, r_k)$ and a surface ball $\Delta(x_Q, r_k) := B(x_Q, r_k) \cap E$ with $r_k = c2^{-k} \approx \operatorname{diam}(Q)$ and

(2.5)
$$\Delta(x_Q, r_k) \subset Q \subset \Delta(x_Q, Cr_k).$$

We shall refer to the point x_0 as the "center" of Q.

• For a dyadic cube $Q \in \mathbb{D}_k$, we shall set $\ell(Q) = 2^{-k}$, and we shall refer to this quantity as the "length" of Q. Evidently, $\ell(Q) \approx \operatorname{diam}(Q)$.

For future reference, with $E = \partial \Omega$, we set

$$(2.6) \qquad \qquad \mathbb{D}_{Q} := \left\{ Q' \in \mathbb{D}(\partial\Omega) : Q' \subset Q \right\}$$

Definition 2.7. (Tree (or "stopping time tree")). A tree $S \subset \mathbb{D}$ is a collection of cubes such that

- (1) S has a unique maximal (or "top") cube Q(S), i.e., $Q \subset Q(S)$ for all $Q \in S$.
- (2) **S** is *semi-coherent*, i.e., if $Q \in \mathbf{S}$, and $Q \subset Q' \subset Q(\mathbf{S})$, then $Q' \in \mathbf{S}$.

Definition 2.8. (Corkscrew condition). Following [JK], we say that a domain $\Omega \subset \mathbb{R}^{n+1}$ satisfies the "Corkscrew condition" if for some uniform constant c > 0 and for every ball B = B(x, r), with $x \in \partial \Omega$ and $0 < r < \operatorname{diam}(\partial \Omega)$, there is a ball $B(X_B, cr) \subset B(x, r) \cap \Omega$. The point $X_B \subset \Omega$ is called a "Corkscrew point" relative to B.

Remark 2.9. We note that, on the other hand, every $X \in \Omega$, with $\delta(X) < \text{diam}(\partial\Omega)$, may be viewed as a Corkscrew point, relative to some ball *B* centered on $\partial\Omega$. Indeed, set $r = 2\delta(X)$, fix $\hat{x} \in \partial\Omega$ such that $|X - \hat{x}| = \delta(X)$, and let $B := B(\hat{x}, r)$.

Definition 2.10. (Harnack Chain condition). Again following [JK], we say that Ω satisfies the Harnack Chain condition if there are uniform positive constants c, C such that for every $\rho > 0$, $K \ge 1$, and pair of points $X, X' \in \Omega$ with $\delta(X)$, $\delta(X') \ge \rho$ and $|X - X'| < K\rho$, there is a chain of open balls $B_1, \ldots, B_N \subset \Omega$, $N \le C(K)$, with $X \in B_1, X' \in B_N, B_k \cap B_{k+1} \ne \emptyset$ and $c \operatorname{diam}(B_k) \le \operatorname{dist}(B_k, \partial \Omega) \le C \operatorname{diam}(B_k)$. The chain of balls is called a "Harnack Chain".

Definition 2.11. (1-sided NTA, aka "uniform" domain). We say that a domain $\Omega \subset \mathbb{R}^{n+1}$ is 1-sided NTA (*Non-tangentially accessible*) (aka "uniform") if it satisfies the Corkscrew and Harnack Chain conditions.

Definition 2.12. (1-sided CAD). We say that a connected open set $\Omega \subset \mathbb{R}^{n+1}$ is a 1-sided CAD (*1-sided Chord-arc domain*), if it is a uniform (1-sided NTA) domain, and if $\partial \Omega$ is ADR.

Definition 2.13. $(Y^{1,2}, W^{1,2} \text{ and } Y^{1,2}_0, W^{1,2}_0)$. As usual, $W^{1,2}(\Omega) := \{f \in L^2(\Omega) : \nabla f \in L^2(\Omega)\}$, endowed with the norm $||f||_{W^{1,2}(\Omega)} := ||f||_{L^2(\Omega)} + ||\nabla f||_{L^2(\Omega)}$, and $W^{1,2}_0(\Omega)$ is the completion of $C_0^{\infty}(\Omega)$ in the $W^{1,2}(\Omega)$ norm.

Also as usual, in \mathbb{R}^d set $2^* := 2d/(d-2)$, thus for us, d = n + 1 and $2^* := 2(n+1)/(n-1)$. Then $Y^{1,2}(\Omega) := \{f \in L^{2^*}(\Omega) : \nabla f \in L^2(\Omega)\}$, with norm

$$||f||_{Y^{1,2}(\Omega)} := ||f||_{L^{2^*}(\Omega)} + ||\nabla f||_{L^2(\Omega)}.$$

We let $Y_0^{1,2}(\Omega)$ denote the completion of $C_0^{\infty}(\Omega)$ in the $Y^{1,2}(\Omega)$ norm.

Definition 2.14. (Non-tangential approach regions and maximal functions). Given $z \in \partial \Omega$, we define a non-tangential approach region ("cone") with vertex at z by

(2.15)
$$\Gamma(z) := \{Y \in \Omega : |z - Y| < 10\delta(Y)\},\$$

and for a measurable function H defined in Ω , let

(2.16)
$$N_*H(z) := \sup_{Y \in \Gamma(z)} |H(Y)|,$$

denote the non-tangential maximal function of H.

Definition 2.17. (Dirichlet problem with data in L^p).

$$(D)_p \begin{cases} Lu = 0 \text{ in } \Omega\\ u|_{\partial\Omega} = f \in L^p(\partial\Omega)\\ N_*u \in L^p(\partial\Omega), \end{cases}$$

where $u|_{\partial\Omega} = f$ is understood in the sense of nontangential convergence.

Definition 2.18. Continuous Dirichlet problem.

$$(D) \begin{cases} Lu = 0 \text{ in } \Omega\\ u|_{\partial\Omega} = f \in C_c(\partial\Omega)\\ u \in C(\overline{\Omega}). \end{cases}$$

Remark 2.19. As is well known, solvability of the continuous Dirichlet problem, in conjunction with the weak maximum principle, implies the existence of elliptic measure via the Riesz representation theorem.

Definition 2.20. $(A_{\infty} \text{ and weak-}A_{\infty} \text{ for elliptic measure})$. Let m be a doubling measure on $\partial \Omega$. We say that elliptic measure ω is (locally) in weak- A_{∞} on $\partial \Omega$, with respect to m, and we write $\omega \in \text{weak-}A_{\infty}(m)$, if there are uniform positive constants C and s such that for every

ball B = B(x, r) centered on $\partial \Omega$, with radius $r < \text{diam}(\partial \Omega)/4$, and associated surface ball $\Delta = B \cap \partial \Omega$,

(2.21)
$$\omega^X(E) \le C\left(\frac{\mathfrak{m}(E)}{\mathfrak{m}(\Delta)}\right)^s \omega^X(2\Delta), \quad \forall X \in \Omega \setminus 4B, \forall \text{Borel } E \subset \Delta;$$

equivalently, if for every ball *B* and surface ball $\Delta = B \cap \partial \Omega$ as above, and for each point $X \in \Omega \setminus 4B$, $\omega^X \in \text{weak}-A_{\infty}(\mathfrak{m}, \Delta)$ with uniformly controlled weak- A_{∞} constants, i.e., (2.21) holds uniformly, with Δ replaced by $\Delta' = B' \cap \partial \Omega$, and with $E \subset \Delta'$, for all $B' \subset B$ and all $X \in \Omega \setminus 4B$.

If, in addition, ω^X satisfies the doubling property (Definition 2.3), for $X \in \Omega \setminus 4B$, locally for all $\Delta' = B' \cap \partial\Omega$, with $B' \subset B$, then we say that elliptic measure ω is (locally) in A_{∞} on $\partial\Omega$ with respect to m, and we write $\omega \in A_{\infty}(m)$.

In the special case that $m = \sigma$, we will sometimes simply write $\omega \in \text{weak}-A_{\infty}$, or $\omega \in A_{\infty}$.

Remark 2.22. It is well known that solvability of $(D)_p$ for some finite p is equivalent to the property that ω^X belongs (locally) to weak- A_∞ (or to A_∞ , in the case that ω^X is doubling) with respect to σ , in the sense of Definition 2.20, and in fact the Poisson kernel $k^X := d\omega^X/d\sigma$ satisfies a reverse Hölder condition with exponent q = p/(p-1). See, e.g., [H, HLe].

2.1. **Some geometric results.** We record some known geometric facts that will be useful in the sequel.

Let us first make a standard Whitney decomposition of Ω . Let $\mathcal{W} = \mathcal{W}(\Omega)$ denote a collection of (closed) dyadic Whitney cubes of Ω , so that the cubes in \mathcal{W} form a pairwise non-overlapping covering of Ω , which satisfy

(2.23)
$$4 \operatorname{diam}(I) \le \operatorname{dist}(4I, \partial \Omega) \le \operatorname{dist}(I, \partial \Omega) \le 40 \operatorname{diam}(I), \quad \forall I \in \mathcal{W}$$

(just dyadically divide the standard Whitney cubes, as constructed in [Ste, Chapter VI], into cubes with side length 1/8 as large) and also

$$(1/4) \operatorname{diam}(I_1) \leq \operatorname{diam}(I_2) \leq 4 \operatorname{diam}(I_1),$$

whenever I_1 and I_2 touch.

We fix a small parameter $\tau_0 > 0$, so that for any $I \in W$, and any $\tau \in (0, \tau_0]$, the concentric dilate

(2.24)
$$I^*(\tau) := (1+\tau)I$$

still satisfies the Whitney property

(2.25)
$$\operatorname{diam} I \approx \operatorname{diam} I^*(\tau) \approx \operatorname{dist} \left(I^*(\tau), E\right) \approx \operatorname{dist}(I, E), \quad 0 < \tau \le \tau_0.$$

We recall that in [HM2, Section 3, and Appendix A], there is a construction of Whitney regions, "Carleson boxes" (and Carleson tents), and "sawtooth" subdomains corresponding to a tree S, with the following properties.

Proposition 2.26. [HM2, Section 3, and Appendix A]. Let Ω be a 1-sided chord arc domain. Then given a tree $\mathbf{S} \subset \mathbb{D}(\partial \Omega)$, there is a "sawtooth" subdomain $\Omega_{\mathbf{S}} \subset \Omega$, satisfying the following properties:

- (1) $\Omega_{\mathbf{S}}$ is a 1-sided chord arc domain.
- (2) $\Omega_{\mathbf{S}} = \operatorname{int} (\bigcup_{Q \in \mathbf{S}} U_Q)$, where U_Q is a "Whitney region" relative to Q.

(3) The Whitney regions are of the form $U_Q = \bigcup_{I \in W(Q)} I^*$, where I^* is the concentrically fattened version of I defined above, and $W(Q) \subset W$, with

 $\ell(Q) \approx \ell(I) \approx \operatorname{dist}(Q, I) \approx \operatorname{dist}(Q, I^*) \approx \operatorname{dist}(\partial \Omega, I^*), \quad \forall I \in \mathcal{W}(Q).$

(4) For the Whitney regions, we have

(2.27) $\operatorname{dist}(U_Q, Q) \approx \operatorname{dist}(U_Q, \partial \Omega) \approx \operatorname{diam}(Q) \approx \operatorname{diam}(U_Q) \approx |U_Q|^{1/(n+1)}.$

(5) The Whitney regions enjoy the bounded overlap property

$$\sum_{Q \in \mathbb{D}(\partial \Omega)} \mathbf{1}_{U_Q}(X) \lesssim 1, \qquad \forall X \in \Omega.$$

- (6) In the special case that $\mathbf{S} = \mathbb{D}_{Q_0}$ for some $Q_0 \in \mathbb{D}$, we use the notation $R_{Q_0} :=$ int $(\cup_{Q \subset Q_0} U_Q)$, and we refer to R_{Q_0} as the "Carleson box" relative to Q_0 . For each $Q \in \mathbb{D}$, the Carleson box R_Q has the property that there is a ball $B_Q = B(x_Q, r_Q)$, with $r_Q \approx \ell(Q)$, such that $R_Q \supset 2B_Q \cap \Omega$, where $2B_Q = B(x_Q, 2r_Q)$ is the concentric double of B_Q .
- (7) By (2) (or (6)), and (4), it follows that

$$\operatorname{diam}(\Omega_{\mathbf{S}}) \leq \operatorname{diam}(Q(\mathbf{S}))$$
.

In particular, diam(R_Q) \leq diam(Q).

(8) Given a surface ball $\Delta = B \cap \partial \Omega$, where B = B(x, r) is centered on $\partial \Omega$, there is a "Carleson Tent" T_{Δ} associated to Δ , such that

$$B \cap \Omega \subset (5/4)B \cap \Omega \subset T_{\Delta} \subset \kappa B \cap \Omega,$$

where (5/4)B = B(x, 5r/4), and $\kappa B = B(x, \kappa r)$ for some sufficiently large, uniform constant κ . Moreover, T_{Δ} is a 1-sided chord arc domain. Finally, if $B' \subset B$, and $\Delta' = B' \cap \partial\Omega$, then $T_{\Delta'} \subset T_{\Delta}$.

Furthermore, all of the implicit and explicit constants in properties (1), (3), (4), (5), (6), (7) and (8) depend only on dimension and on the 1-sided chord arc constants for Ω .

Remark 2.28. We caution the reader that our notation differs from that of [HM2]. In the latter reference, the subdomains $\Omega_{\mathbf{S}}$ are denoted $\Omega_{\mathcal{F},Q}$, where, translating to the notation of the present paper, we have $Q = Q(\mathbf{S})$, the maximal cube in \mathbf{S} , and $\mathcal{F} = \{Q_j\}_j$ denotes the collection of sub-cubes of $Q = Q(\mathbf{S})$ that are maximal with respect to the property that $Q_j \notin \mathbf{S}$.

We recall some further results from [HM2] that will be useful in the sequel.

Proposition 2.29. [HM2, Proposition 6.1]. Suppose that Ω is a 1-sided chord-arc domain. Fix $Q_0 \in \mathbb{D}$, let $\mathbf{S} \subset \mathbb{D}_{Q_0}$ be a tree, and let $\mathcal{F} = \{Q_j\}_j$ denote the collection of sub-cubes of Q_0 that are maximal with respect to the property that $Q_j \notin \mathbf{S}$. Then

 $(2.30) Q_0 \setminus (\cup_{\mathcal{F}} Q_j) \subset Q_0 \cap \partial \Omega_{\mathbf{S}} \subset \partial \Omega \cap \partial \Omega_{\mathbf{S}} \subset \overline{Q_0} \setminus (\cup_{\mathcal{F}} \operatorname{int}(Q_j)) .$

Remark 2.31. Note that (2.30) implies in particular that

(2.32) $Q_0 \cap \partial \Omega_{\mathbf{S}} = [Q_0 \setminus (\cup_{\mathcal{F}} \operatorname{int}(Q_j))] \cap \partial \Omega_{\mathbf{S}}$

 $= \left[\mathcal{Q}_0 \setminus \left(\cup_{\mathcal{F}} \mathcal{Q}_j \right) \right] \bigcup \left(\partial \Omega_{\mathbf{S}} \cap \left(\cup_{\mathcal{F}} \left[\mathcal{Q}_j \setminus \operatorname{int}(\mathcal{Q}_j) \right] \right) \right) \, .$

Proposition 2.33. [HM2, Proposition 6.3]. Suppose that $\partial\Omega$ is ADR, and that m is a doubling measure on $\partial\Omega$ (Definition 2.3). Then $\partial Q := \overline{Q} \setminus \operatorname{int}(Q)$ has m-measure 0, for every $Q \in \mathbb{D}$.

Proposition 2.34. [HM2, Proposition 6.7]. Suppose that Ω is a 1-sided chord-arc domain. Fix $Q_0 \in \mathbb{D}$, let $\mathbf{S} \subset \mathbb{D}_{Q_0}$ be a tree, and let $\mathcal{F} = \{Q_j\}_j$ denote the collection of sub-cubes of Q_0 that are maximal with respect to the property that $Q_j \notin \mathbf{S}$. Then for each $Q_j \in \mathcal{F}$, there is a surface ball $P_j \subset \partial \Omega_{\mathbf{S}}$, satisfying

(2.35)
$$\operatorname{diam}(P_{i}) \approx \operatorname{dist}(P_{i}, Q_{i}) \approx \operatorname{dist}(P_{i}, \partial\Omega) \approx \ell(Q_{i}),$$

where the uniform implicit constants depend only on the 1-sided chord-arc constants for Ω (in particular, on the parameters fixed in the construction of the Whitney regions U_Q , and the sawtooth regions Ω_S).

Let $x_j^* \in \partial \Omega_S$ denote the center of the surface ball P_j in Proposition 2.34, and let $t_j \approx \ell(Q_j)$ be its radius.

Proposition 2.36. [HM2, Proposition 6.12]. Let Ω , Q_0 , \mathbf{S} , \mathcal{F} and P_j be as in Proposition 2.34. For $Q_j \in \mathcal{F}$, let $MP_j := B(x_j^*, Mt_j) \cap \partial \Omega_{\mathbf{S}}$ be the M-fold concentric dilate of the surface ball P_j on $\partial \Omega_{\mathbf{S}}$. Then for M large enough, there is a surface ball $\Delta_* := B(x_*, t_*) \cap \partial \Omega_{\mathbf{S}}$, with $t_* \approx \ell(Q_0)$, and $x_* \in \partial \Omega_{\mathbf{S}}$, satisfying

(2.37)
$$\Delta_{\star} \subset \left(Q_0 \cap \partial \Omega_{\mathbf{S}}\right) \bigcup \left(\cup_{Q_j \in \mathcal{F}} MP_j\right),$$

where *M* and the various implicit constants depend only on the 1-sided chord-arc constants for Ω (in particular, on the parameters fixed in the construction of the Whitney regions U_Q , and the sawtooth regions Ω_S).

We now observe that, in light of Remark 2.31, we may deduce the following subtle (but eventually useful) self-improvement of Proposition 2.36.

Corollary 2.38. Let Ω , Q_0 , S, \mathcal{F} , Δ_* , P_j and MP_j be as in Proposition 2.36. Then for a possibly larger choice of M (but still uniform, and depending only on allowable parameters), we have the following improvement of (2.37):

(2.39)
$$\Delta_{\star} \subset \left(Q_0 \setminus \left(\cup_{\mathcal{F}} Q_j\right)\right) \bigcup \left(\cup_{Q_j \in \mathcal{F}} MP_j\right).$$

Proof. Using (2.32), we need only observe that for each *j*,

$$\partial \Omega_{\mathbf{S}} \cap |Q_i \setminus \operatorname{int}(Q_i)| \subset \partial \Omega_{\mathbf{S}} \cap Q_i \subset MP_i,$$

by (2.35) and the definition of MP_i , provided that we fix M large enough.

Lemma 2.40 (Local Hardy Inequality). Let $\Omega \subset \mathbb{R}^{n+1}$ be a 1-sided CAD, and consider any ball B = B(x, r), with $x \in \partial \Omega$. Set $\Omega_r := \Omega \cap B(x, r)$, and $\Omega_{Mr} := \Omega \cap B(x, Mr)$, for M > 0. Then there is a constant $\kappa_0 \in (1, \infty)$, depending only on the 1-sided CAD constants, such that for any $u \in W^{1,2}(\Omega_{2\kappa_0 r})$ with vanishing trace on $2\kappa_0 \Delta := B(x, 2\kappa_0 r) \cap \partial \Omega$,

(2.41)
$$\iint_{\Omega_r} \left(\frac{u(Y)}{\delta(Y)}\right)^2 dY \lesssim \iint_{\Omega_{\kappa_0 r}} |\nabla u(Y)|^2 dY,$$

where the implicit constant depends only on dimension and the 1-sided CAD constants.

Remark 2.42. A global version of this result, in which Ω_r and $\Omega_{\kappa_0 r}$ are replaced by Ω , and $u \in W_0^{1,2}(\Omega)$, appears in [A], and for that result, the corkscrew and Harnack chain conditions are not required, rather only "uniform thickness" of the complement of Ω (in particular, Ahlfors-David regularity of the boundary suffices). A more general result appears in [L]. Lemma 2.40 will be a routine consequence of these global results.

Proof of Lemma 2.40. Let φ be a smooth bump function adapted to the ball B = B(x, r), i.e., $\varphi \in C_0^{\infty}(2B), \varphi \equiv 1$ on $B, 0 \le \varphi \le 1$, and $|\nabla \varphi| \le r^{-1}$, and similarly, let η be be a smooth bump function adapted to the ball $B(x, \kappa_0 r)$, with $\eta \equiv 1$ on $B(x, \kappa_0 r)$, and $\supp(\eta) \subset B(x, 2\kappa_0 r)$. Then $u = u\eta$ on $\Omega_{\kappa_0 r}$, so replacing u by $u\eta$, we may assume that $u \in W_0^{1,2}(\Omega)$, and hence by density, we may suppose that $u \in C_0^{\infty}(\Omega)$.

Observe further that in particular, $u\varphi \in W_0^{1,2}(\Omega)$. Thus, by the global result of [A] (or [L]), we see that

$$\begin{split} \iint_{\Omega_r} \left(\frac{u(Y)}{\delta(Y)}\right)^2 dY &\leq \iint_{\Omega} \left(\frac{u(Y)\varphi(Y)}{\delta(Y)}\right)^2 dY \\ &\lesssim \iint_{\Omega} |\nabla u|^2 \varphi^2 dY + \iint_{\Omega} |\nabla \varphi|^2 u^2 dY =: I + II \,. \end{split}$$

Term I clearly satisfies the desired bound in (2.41).

To handle term *II*, first note that by the properties of φ ,

$$II \lesssim r^{-2} \iint_{\Omega_{2r}} u^2 dY.$$

We choose a collection $C \subset \mathbb{D}(\partial \Omega)$, of uniformly bounded cardinality depending only on the 1-sided CAD constants, with $\ell(Q) \approx r$ for all $Q \in C$, and such that

$$\Omega_{2r} \subset \bigcup_{Q \in C} R_Q,$$

where R_Q is the "Carleson box" associated to Q, as in Proposition 2.26 (6). Our goal is now to prove that

(2.43)
$$r^{-2} \iint_{R_Q} u^2 dY \lesssim \iint_{R_Q^*} |\nabla u|^2 dY,$$

uniformly for each $Q \in C$, where $R_Q^* = \operatorname{int} \left(\bigcup_{Q' \subset Q} U_{Q'}^* \right)$, and $U_{Q'}^*$ is a fattened version of the Whitney region $U_{Q'}$, which still retains the same Whitney properties: in particular, $\operatorname{diam}(U_{Q'}^*) \approx \operatorname{dist}(U_{Q'}^*, \partial \Omega) \approx \ell(Q')$. Indeed, observe that by construction, there is a uniform constant κ_0 such that $R_Q^* \subset \Omega_{\kappa_0 r}$ for each $Q \in C$, thus, (2.43) implies the desired bound in (2.41).

We now turn to the proof of (2.43). For any $Q \in \mathbb{D}$, and any $f \in L^1(U_Q^*)$, let $[f]_Q$ and $[f]_Q^*$ denote, respectively, the mean value of f on U_Q , and on the fattened Whitney region U_Q^* , i.e.,

$$[f]_{Q} := |U_{Q}|^{-1} \iint_{U_{Q}} f \, dY, \qquad [f]_{Q}^{*} := |U_{Q}^{*}|^{-1} \iint_{U_{Q}^{*}} f \, dY$$

Fix an arbitrary $Q_0 \in C$, and observe that by definition of the Carleson regions,

$$r^{-2} \iint_{\mathcal{R}_{Q_0}} u^2 dY \leq r^{-2} \sum_{Q \subset Q_0} \iint_{U_Q} u^2 dY \leq III + IV,$$

where

$$III = r^{-2} \sum_{Q \subseteq Q_0} \iint_{U_Q} \left| u - [u]_Q \right|^2 dY,$$

and

$$IV = r^{-2} \sum_{Q \subset Q_0} \iint_{U_Q} \left| \int_Q \sum_{Q' \subset Q} 1_{Q'}(x) \left([u]_{Q'} - [u]_{Q'_{child}} \right) d\sigma(x) \right|^2 dY,$$

where for $Q' \subset Q$ with $Q \ni x$, we let Q'_{child} denote the child of Q' that contains x, and where we have replaced $[u]_Q = [u]_Q - u(x)$ by a telescoping sum, using the fact that we have reduced to the case that $u \in C_0^{\infty}(\Omega)$.

By Poincaré's inequality (see, e.g., [HM2, Section 4])

$$\iint_{U_Q} \left| u - [u]_Q \right|^2 dY \lesssim \ell(Q)^2 \iint_{U_Q^*} |\nabla u|^2 dY,$$

where as above U_Q^* is a fattened version of U_Q . Since the fattened Whitney regions retain the bounded overlap property, we easily obtain (2.43) for term *III*.

Similarly (again see [HM2, Section 4]),

$$\left| [u]_{Q'} - [u]_{Q'_{child}} \right| \leq \ell(Q') [|\nabla u|]_{Q'}^* ,$$

and therefore

$$\begin{split} IV \lesssim r^{-2} \sum_{\mathcal{Q} \subset \mathcal{Q}_0} |U_{\mathcal{Q}}| \, \left(\sum_{\mathcal{Q}' \subset \mathcal{Q}} \frac{\sigma(\mathcal{Q}')}{\sigma(\mathcal{Q})} \, \ell(\mathcal{Q}') \, |U_{\mathcal{Q}'}^*|^{-1} \iint_{U_{\mathcal{Q}'}^*} |\nabla u| \, dY \right)^2 \\ \lesssim r^{-2} \sum_{\mathcal{Q} \subset \mathcal{Q}_0} \ell(\mathcal{Q})^{1-n} \, \left(\iint_{R_{\mathcal{Q}}^*} |\nabla u| \, dY \right)^2 \,, \end{split}$$

where we have used that $|U_Q| \approx |U_Q^*| \approx \ell(Q)^{n+1}$, and $\sigma(Q) \approx \ell(Q)^n$, for each $Q \in \mathbb{D}(\partial\Omega)$. Moreover $|R_Q^*| \approx \ell(Q)^{n+1}$, so by Cauchy-Schwarz, we see in turn that

$$IV \lesssim r^{-2} \sum_{Q \subseteq Q_0} \ell(Q)^2 \iint_{R_Q^*} |\nabla u|^2 dY \approx \sum_{k=0}^{\infty} 2^{-2k} \sum_{Q \in \mathbb{D}_k(Q_0)} \iint_{R_Q^*} |\nabla u|^2 dY,$$

where $\mathbb{D}_k(Q_0)$ is the collection of subcubes of Q_0 with $\ell(Q) = 2^{-k}\ell(Q_0) \approx 2^{-k}r$. Note that by definition, R_Q^* is contained in a union of fat Whitney regions $U_{Q'}^*$ with $Q' \subset Q$, thus, for distinct $Q_1, Q_2 \in \mathbb{D}_k(Q_0)$, any pair of corresponding sub-cubes $Q'_1 \subset Q_1$ and $Q'_2 \subset Q_2$ are disjoint. Since the regions $U_{Q'}^*$ have bounded overlaps, we find that

$$\sum_{Q \in \mathbb{D}_k(Q_0)} \iint_{R_Q^*} |\nabla u|^2 dY \lesssim \iint_{R_{Q_0}^*} |\nabla u|^2 dY$$

uniformly for each k, so we may sum the geometric series to obtain the desired bound for IV.

3. Preliminaries

In this section, for operators of the form $L = -\operatorname{div} A\nabla + \mathbf{B} \cdot \nabla = L_0 + \mathbf{B} \cdot \nabla$, we verify that certain standard estimates for solutions of the homogeneous second order equation $L_0 u = 0$, continue to hold for solutions of Lu = 0, assuming that **B** satisfies (1.2) (but not necessarily the Carleson measure condition (1.3)). For some (but not all) of these results, we shall require the "small constant" version of (1.2), namely

(3.1)
$$|\mathbf{B}(X)| \leq \frac{\varepsilon_0}{\delta(X)}, \quad \text{a.e. } x \in \Omega,$$

where ε_0 will be sufficiently small, depending only on dimension, the ellipticity parameters λ and Λ , and the 1-sided chord-arc constants for Ω .

In addition, we obtain solvability of the Dirichlet problem with continuous data, and consequent existence of elliptic measure, under certain conditions.

Some of the results in this section hold for both $L = -\operatorname{div} A \nabla + \mathbf{B} \cdot \nabla$ and it's adjoint L^* , and for those results, we shall consider more generally the operator

(3.2)
$$\mathcal{L}u := -\operatorname{div}(A\nabla u + \mathbf{B}_1 u) + \mathbf{B}_2 \cdot \nabla u.$$

Taking $\mathbf{B}_1 = 0$, $\mathbf{B}_2 = \mathbf{B}$, we recover the operator *L*, and taking $\mathbf{B}_2 = 0$, $\mathbf{B}_1 = \mathbf{B}$ (and replacing *A* by its transpose A^{T}), we recover L^* .

We begin with some standard *a priori* estimates.

3.1. A priori estimates.

Lemma 3.3 (De Giorgi-Nash-Moser estimates and Harnack's inequality). Let $\Omega \subset \mathbb{R}^{n+1}$ be an open set, let \mathcal{L} be defined as in (3.2), and suppose that (1.2) holds for each of \mathbf{B}_1 and \mathbf{B}_2 . Consider a ball B = B(X, r) such that $3B \subset \Omega$. Let $u \in W^{1,2}(2B)$ be a weak solution of $\mathcal{L}u = 0$ in 2B. Then

(3.4)
$$\sup_{B} |u|^2 \lesssim |2B|^{-1} \iint_{2B} |u|^2 dY,$$

and

$$(3.5) |u(x) - u(y)| \leq |x - y|^{\alpha} \sup_{2B} |u|, \forall x, y \in B.$$

If in addition, u is non-negative in 2B, then

$$(3.6) \qquad \qquad \sup_{B} u \lesssim \inf_{B} u.$$

Moreover, the implicit constants and the Hölder exponent α depend only on dimension, ellipticity of A, and the constant in (1.2).

Proof. Since $3B \subset \Omega$, we have $r \leq \delta(X)$ in 2*B*, and therefore $|\mathbf{B}_1(X)| + |\mathbf{B}_2(X)| \leq r^{-1}$, for a.e. $X \in 2B$. By scale invariance, we may assume r = 1, and thus we have reduced to the case that $||\mathbf{B}||_{L^{\infty}(2B)} \leq 1$. In this form, the results stated may all be found in [GT, Chapter 8].

Remark 3.7. Note that the smallness condition (3.1) is not needed in Lemma 3.3, nor will it be needed in the next result, Lemma 3.8.

Lemma 3.8 (interior Caccioppoli's inequality). Let $\Omega \subset \mathbb{R}^{n+1}$ be an open set, let \mathcal{L} be defined as in (3.2), and suppose that (1.2) holds for each of \mathbf{B}_1 and \mathbf{B}_2 . Consider a ball B = B(X, r) such that $3B \subset \Omega$. Let u be a $W^{1,2}$ solution of $\mathcal{L}u = 0$ in 2B. Then

$$\iint_{B} |\nabla u|^2 dY \lesssim r^{-2} \iint_{2B} u^2 dY$$

where the implicit constant depends only on ellipticity of A and the constant in (1.2).

Proof. The proof is standard. Let φ be a smooth bump function adapted to the ball B = B(x, r), i.e., $\varphi \in C_0^{\infty}(2B)$, $\varphi \equiv 1$ on $B, 0 \le \varphi \le 1$, and $|\nabla \varphi| \le r^{-1}$, so that

$$\begin{split} \iint_{B} |\nabla u|^{2} dY &\leq \iint_{\Omega} |\nabla u|^{2} \varphi^{2} dY \lesssim \iint_{\Omega} A \nabla u \cdot \nabla u \varphi^{2} dY \\ &= \iint_{\Omega} A \nabla u \cdot \nabla (u \varphi^{2}) dY - 2 \iint_{\Omega} A \nabla u \cdot \nabla \varphi \, u \varphi \, dY =: I + II \,. \end{split}$$

12

As usual, for a small number γ at our disposal,

$$|II| \leq \gamma \iint_{\Omega} |\nabla u|^2 \varphi^2 dY + \frac{1}{\gamma} \iint_{\Omega} |\nabla \varphi|^2 u^2 dY$$

Choosing $\gamma > 0$ small enough, we may hide the small term. With γ now fixed, the second term satisfies the desired bound, since $|\nabla \varphi|^2 \leq r^{-2}$.

Since $3B \subset \Omega$, we have $r \leq \delta(X)$ for all $X \in 2B$, hence by (1.2),

$$|\mathbf{B}_1(X)| + |\mathbf{B}_2(X)| \leq r^{-1}$$
, a.e. $X \in 2B$.

Thus, since $\mathcal{L}u = 0$ in 2B,

$$\begin{split} |I| &= \Big| \iint_{\Omega} \mathbf{B}_2 \cdot \nabla u \, u \varphi^2 dY \, + \, \iint_{\Omega} u \, \mathbf{B}_1 \cdot \nabla (u \varphi^2) dY \, \Big| \\ &\lesssim \frac{1}{r} \iint_{\Omega} |\nabla u| \, u \varphi^2 dY \, \lesssim \gamma \iint_{\Omega} |\nabla u|^2 \varphi^2 dY \, + \, \frac{1}{\gamma} \frac{1}{r^2} \iint_{2B} u^2 dY \, , \end{split}$$

where again $\gamma > 0$ is at our disposal. Again we may hide the small term with γ fixed small enough.

Lemma 3.9 (Caccioppoli's inequality at the boundary). Let $\Omega \subset \mathbb{R}^{n+1}$ be a 1-sided CAD, and consider any ball B = B(x, r), with $x \in \partial \Omega$. Let \mathcal{L} be defined as in (3.2), and suppose that (3.1) holds for each of \mathbf{B}_1 and \mathbf{B}_2 , with ε_0 sufficiently small, depending only on the ellipticity of A. Set $\Omega_r := \Omega \cap B(x, r)$, and $\Omega_{2r} := \Omega \cap B(x, 2r)$, and let $u \in W^{1,2}(\Omega_{2r})$ be a solution of $\mathcal{L}u = 0$ in Ω_{2r} , with vanishing trace on $2\Delta := B(x, 2r) \cap \partial \Omega$. Then

(3.10)
$$\iint_{\Omega_r} |\nabla u|^2 dY \lesssim r^{-2} \iint_{\Omega_{2r}} u^2 dY$$

where the implicit constants depend only on the allowable parameters.

Proof. The proof is again standard. As above, choose $\varphi \in C_0^{\infty}(2B)$, with $\varphi \equiv 1$ on B, $0 \le \varphi \le 1$, and $|\nabla \varphi| \le r^{-1}$. We then have

$$\iint_{\Omega_r} |\nabla u|^2 dY \leq \iint_{\Omega} |\nabla u|^2 \varphi^2 dY \lesssim \iint_{\Omega} A \nabla u \cdot \nabla u \varphi^2 dY$$
$$= \iint_{\Omega} A \nabla u \cdot \nabla (u \varphi^2) \, dY - 2 \iint_{\Omega} A \nabla u \cdot \nabla \varphi \, u \varphi \, dY =: I + II.$$

As usual, for a small number γ at our disposal,

$$|II| \lesssim \gamma \iint_{\Omega} |\nabla u|^2 \varphi^2 dY + \frac{1}{\gamma} \iint_{\Omega} |\nabla \varphi|^2 u^2 dY$$

Choosing $\gamma > 0$ small enough, we may hide the small term. With γ now fixed, the second term satisfies the desired bound in (3.10).

Since $\mathcal{L}u = 0$ in Ω_{2r} , and $u\varphi^2 \in W_0^{1,2}(\Omega_{2r})$, using (3.1), we also have

$$\begin{split} |I| &= \left| \iint_{\Omega} \mathbf{B}_{2} \cdot \nabla u \, u \varphi^{2} dY + \iint_{\Omega} u \, \mathbf{B}_{1} \cdot \nabla (u \varphi^{2}) dY \right| \leq \varepsilon_{0} \iint_{\Omega} (|\nabla u|\varphi + |\nabla \varphi|u) \, \frac{u\varphi}{\delta} \, dY \\ &\lesssim \varepsilon_{0} \left(\iint_{\Omega} |\nabla u|^{2} \varphi^{2} dY + \iint_{\Omega} |\nabla \varphi|^{2} u^{2} dY + \iint_{\Omega} \left(\frac{u\varphi}{\delta} \right)^{2} dY \right) \\ &=: \varepsilon_{0} \left(I_{1} + I_{2} + I_{3} \right). \end{split}$$

It is enough to deal with I_1 and I_2 , since, by the global Hardy inequality in [A] and [L], we have

$$I_3 \lesssim \int_{\Omega} \left(|\nabla u| \varphi + |\nabla \varphi| u \right)^2 dY \approx I_1 + I_2.$$

In turn, as was the case for term *II* treated above, $\varepsilon_0 I_1$ may be hidden, provided ε_0 is small enough, and term I_2 yields the desired estimate in (3.10).

Lemma 3.11 (Hölder continuity at the boundary and Carleson's estimate). Let $\Omega \subset \mathbb{R}^{n+1}$ be a 1-sided CAD, and consider any ball $B = B(x_0, r)$, with $x_0 \in \partial \Omega$. Set $\Omega_r := \Omega \cap B(x_0, r)$, and $\Omega_{2r} := \Omega \cap B(x_0, 2r)$. Let \mathcal{L} be defined as in (3.2), and suppose that (3.1) holds for each of \mathbf{B}_1 and \mathbf{B}_2 , with ε_0 sufficiently small. If $u \in W^{1,2}(\Omega_{2r})$ is a non-negative solution to $\mathcal{L}u = 0$ in Ω_{2r} , with vanishing trace on $2\Delta := B(x, 2r) \cap \partial \Omega$, then for some $\alpha > 0$,

(3.12)
$$u(X) \lesssim \left(\frac{\delta(X)}{r}\right)^{\alpha} \sup_{\Omega_{2r}} u, \qquad \forall X \in \Omega_r,$$

and if X_B is a corkscrew point relative to B = B(x, r), then

$$(3.13) \qquad \qquad \sup_{Y \in \Omega_r} u \leq u(X_B)$$

where the implicit constants and the exponent α depend only on ellipticity, dimension, and the 1-sided chord arc constants.

Proof. It suffices to prove (3.12). Indeed, the fact that (3.12) holds for every ball *B* centered on $\partial\Omega$ yields (3.13), by the proof in [CFMS, Theorem 1.1]³, which carries over to the present setting. The only ingredients required are Harnack's inequality, the Corkscrew and Harnack Chain conditions, and (3.12), all of which are available to us. We omit the details.

Turning to (3.12), we shall follow the argument in [HL, pp. 12-14]. Let κ and κ_0 be the constants in in Proposition 2.26 (8), and in Lemma 2.40, respectively. Fix $M \gg \kappa \kappa_0 =: \kappa_1$, and let $X \in \Omega_r$, with $\delta(X) \leq r/M$ (if $\delta(X) > r/M$, then there is nothing to prove). Choose $\hat{x} \in \partial\Omega$ such that $|X - \hat{x}| = \delta(X)$, and set $\rho = \delta(X)$. Then by (3.4) and Lemma 2.40, we have

(3.14)
$$u^2(X) \leq \rho^{-n-1} \iint_{B(X,\rho/4)} u^2 dY \leq \rho^{-n-1} \iint_{\Omega_{\rho}(\hat{x})} u^2 dY \leq \rho^{1-n} \iint_{\Omega_{\kappa_0\rho}(\hat{x})} |\nabla u|^2 dY,$$

where $\Omega_{\rho}(\hat{x}) := B(\hat{x}, \rho) \cap \Omega$.

For $0 < t \le r/M$, set $\Delta_t := B(\hat{x}, t) \cap \partial \Omega$, and let $T(t) := T_{\Delta_t}$ be the Carleson tent associated to Δ_t as in Proposition 2.26 (8), so that (with *M* chosen large as above)

(3.15)
$$\Omega_t(\hat{x}) \subset \Omega_{5t/4}(\hat{x}) \subset T(t) \subset \Omega_{\kappa t}(\hat{x}) \subset \Omega_{2\kappa t}(\hat{x}) \subset \Omega_{3r/2}.$$

By Proposition 2.26 (8), T(t) is a 1-sided CAD⁴ for each *t*. For *t* momentarily fixed, we may solve the Dirichlet problem in $T(t/\kappa_1)$ for $L_0u = 0$, with data $u_0 = u$ (in the trace sense) on $\partial T(t/\kappa_1)$. Set $w := u - u_0$, so that $w \in W_0^{1,2}(T(t/\kappa_1))$. Using that $L_0u = 0$ and $\mathcal{L}u = 0$, we see that

$$\iint_{T(t/\kappa_1)} |\nabla w|^2 dY \approx \iint_{T(t/\kappa_1)} A\nabla w \cdot \nabla w \, dY = \iint_{T(t/\kappa_1)} A\nabla u \cdot \nabla w \, dY$$
$$= -\iint_{T(t/\kappa_1)} \mathbf{B}_2 \cdot \nabla u \, w \, dY - \iint_{T(t/\kappa_1)} u \, \mathbf{B}_1 \cdot \nabla w \, dY$$

³see also [JK, Lemma 4.4], where the same argument is used

⁴Indeed, it is for this reason that we have gone to the trouble of replacing $\Omega_t(\hat{x})$ by T(t).

$$\begin{split} \lesssim \, \varepsilon_0 \, \iint_{T(t/\kappa_1)} \left(|\nabla u| \, \frac{w}{\delta'} + |\nabla w| \, \frac{u}{\delta} \right) dY \\ \lesssim \, \varepsilon_0 \left(\iint_{T(t)} |\nabla u|^2 dY + \iint_{T(t/\kappa_1)} |\nabla w|^2 dY \right) \,, \end{split}$$

where $\delta'(Y) := \operatorname{dist}(Y, \partial T(t)) \leq \delta(Y)$, and we have used (3.1) and either the global Hardy inequality of [A] (or [L]) in the domain $T(t/\kappa_1)$ (to handle the term w/δ'), or else (3.15), the local Hardy inequality Lemma 2.40, and then (3.15) again, to handle the term u/δ . To guide the reader through the latter step, we observe that (3.15) yields that $T(t/\kappa_1) \subset \Omega_{t/\kappa_0}(\hat{x})$ (since $\kappa_1 := \kappa \kappa_0$), whence application of Lemma 2.40 yields an integral over $\Omega_t(\hat{x}) \subset T(t)$.

Hiding the small term involving w, we obtain

(3.16)
$$\iint_{T(t/\kappa_1)} |\nabla w|^2 dY \lesssim \varepsilon_0 \iint_{T(t)} |\nabla u|^2 dY.$$

Given s > 0 and $f \in W^{1,2}(\Omega_{2r})$, set

$$\Phi(f,s) := s^{1-n} \iint_{T(s)} |\nabla f|^2 dY,$$

and observe that for $s < t/(4\kappa_1^2)$, letting X_t denote a corkscrew point relative to the ball $B(\hat{x}, t/(\kappa_0\kappa_1))$, and using (3.15) and the case $\mathbf{B}_1 = 0 = \mathbf{B}_2$ of Lemma 3.9, we have

$$\begin{split} \Phi(u_0,s) &\leq s^{1-n} \iint_{\Omega_{\kappa s}(\hat{x})} |\nabla u_0|^2 \, dY \leq s^{-n-1} \iint_{\Omega_{2\kappa s}(\hat{x})} u_0^2 \, dY \\ &\leq \left(\frac{s}{t}\right)^{2\alpha_0} \sup_{\Omega_{t/(\kappa_0\kappa_1)}(\hat{x})} u_0^2 \leq \left(\frac{s}{t}\right)^{2\alpha_0} u_0^2(X_t) \leq \left(\frac{s}{t}\right)^{2\alpha_0} t^{-n-1} \iint_{\Omega_{t/(\kappa_0\kappa_1)}(\hat{x})} u_0^2 \, dY \\ &\leq \left(\frac{s}{t}\right)^{2\alpha_0} t^{1-n} \iint_{\Omega_{t/\kappa_1}(\hat{x})} |\nabla u_0|^2 \, dY \leq \left(\frac{s}{t}\right)^{2\alpha_0} \Phi(u_0, t/\kappa_1) \,, \end{split}$$

where in the second line of the display, we have used that Lemma 3.11 (with some Hölder exponent $\alpha_0 > 0$) and (3.4) hold for u_0 , and in the last line we have used Lemma 2.40, applied in the 1-sided CAD $T(t/\kappa_1)$.

Combining the last estimate with (3.16), we deduce that for $s < t/(4\kappa\kappa_0)$,

$$\begin{split} \Phi(u,s) \, &\lesssim \, \Phi(u_0,s) \, + \, \Phi(w,s) \, \lesssim \, \left(\frac{s}{t}\right)^{2\alpha_0} \Phi(u_0,t/\kappa_1) \, + \, \left(\frac{t}{s}\right)^{n-1} \Phi(w,t/\kappa_1) \\ &\lesssim \, \left[\left(\frac{s}{t}\right)^{2\alpha_0} \, + \, \varepsilon_0 \left(\frac{t}{s}\right)^{n-1} \right] \Phi(u,t) \, . \end{split}$$

Set $\theta = s/t$, so that first choosing $\theta < 1/(4\kappa_1^2)$ sufficiently small, and then choosing ε_0 depending on θ , we see that for all t < r/M

$$\Phi(u,\theta t) \le \frac{1}{2}\Phi(u,t)\,.$$

Iterating the latter estimate starting with t = r/M, and using (3.14) and (3.15), we find that for some $\alpha > 0$,

$$u^2(X) \lesssim \Phi(u, \kappa_0 \rho) \lesssim \left(\frac{\rho}{r}\right)^{2\alpha} \Phi(u, r/M).$$

Moreover, by (3.15) and Lemma 3.9,

$$\Phi(u, r/M) \lesssim r^{-n-1} \iint_{\Omega_{2\kappa r/M}(\hat{x})} u^2 dY \lesssim \sup_{\Omega_{2r}} u^2.$$

Since we had set $\rho = \delta(X)$, the lemma now follows.

Our next task is to discuss existence of solutions.

3.2. Existence of $Y^{1,2}$ solutions with data in $\text{Lip}_c(\partial \Omega)$, in the small constant case. In this subsection, we construct $Y^{1,2}$ solutions to the Dirichlet problem with data in $\text{Lip}_c(\partial \Omega)$, in the small constant case. Specifically, we consider the operator

$$(3.17) L = -\operatorname{div} A \nabla + \mathbf{B} \cdot \nabla,$$

in the case that the constant ε_0 in (3.1) is small enough. The first step is to construction solutions to the Poisson-Dirichlet problem, with vanishing trace:

We state this construction as a lemma, with estimates, for future reference in the sequel.

Lemma 3.19. Let *L* be as in (3.17), with $|\mathbf{B}(X)| \leq \varepsilon_0 / \delta(X)$. If ε_0 is small enough, depending only on the allowable parameters, then given $F \in Y^{-1,2}(\Omega)$, there is a unique solution $w \in Y_0^{1,2}(\Omega)$ to the problem (PD). Moreover,

$$(3.20) ||w||_{L^{2^*}(\Omega)} \le ||w||_{Y_0^{1,2}(\Omega)} \le ||F||_{Y^{-1,2}(\Omega)} \le ||F||_{L^{2*}(\Omega)},$$

where the implicit constants depend only on allowable parameters.

Proof. The technique is standard. Indeed, if ε_0 is small enough, depending only on dimension, ellipticity of *A*, and the 1-sided CAD constants, then the bilinear form

$$\mathfrak{B}(w,v) := \iint_{\Omega} \left(A \nabla w \cdot \nabla v + \mathbf{B} \cdot \nabla w \, v \right) dY$$

is bounded and coercive on $Y_0^{1,2}(\Omega)$ (here, we use the global Hardy inequality of [A] or [L] to deal with a term involving v/δ). We omit the routine details. Thus by Lax-Milgram, given $F \in Y^{-1,2}(\Omega)$, we see that there is a unique $w \in Y_0^{1,2}(\Omega)$ such that

(3.21)
$$\mathfrak{B}(w,v) = \langle F,v \rangle, \qquad \forall v \in Y_0^{1,2}(\Omega),$$

where $\langle \cdot, \cdot \rangle$ denotes the duality pairing of $Y_0^{1,2}(\Omega)$ and its dual $Y^{-1,2}(\Omega)$. In particular, if we set v = w, then

$$\|\nabla w\|_{L^2(\Omega)}^2 \approx \iint_{\Omega} A \nabla w \cdot \nabla w \, dY = \mathfrak{B}(w, w) + O\left(\varepsilon_0 \|\nabla w\|_{L^2(\Omega)}^2\right) \,,$$

where in the last step we have used (3.1) and the global Hardy inequality. For ε_0 chosen small enough, we may hide the small term to obtain

$$\|w\|_{Y_0^{1,2}(\Omega)}^2 \lesssim \|\nabla w\|_{L^2(\Omega)}^2 \lesssim \mathfrak{B}(w,w) \lesssim \|F\|_{Y^{-1,2}(\Omega)} \|w\|_{Y_0^{1,2}(\Omega)},$$

by (3.21) and Sobolev embedding. Consequently, again using Sobolev embedding, we obtain (3.20).

16

We now give the usual construction (see, e.g., [K, page 5]) of the solution to the Dirichlet problem with compactly supported Lipschitz data. Let $f \in \text{Lip}_c(\mathbb{R}^{n+1})$. Then $Lf \in Y^{-1,2}(\Omega)$ in the weak sense (again, this observation uses the Hardy inequality of [A]), so there is a unique $w \in Y_0^{1,2}(\Omega)$ such that (3.21) holds with F = Lf, i.e., Lw = Lf in the weak sense. Setting u = f - w, we see that $u \in Y^{1,2}(\Omega)$, Lu = 0 in Ω , and that $u|_{\partial\Omega} = f|_{\partial\Omega}$ in the trace sense, and moreover, given any $f_0 \in \text{Lip}_c(\partial\Omega)$, the solution u does not depend on the particular Lip_c extension of f_0 to \mathbb{R}^{n+1} .

3.3. Weak maximum principle and continuity of solutions with $\operatorname{Lip}_c(\partial\Omega)$ data. Let us now observe that the solutions *u* constructed as above satisfy the weak maximum principle. We follow the argument in [GT, pp 179-180]: given $f \in \operatorname{Lip}_c(\mathbb{R}^{n+1})$, let *u* be the solution constructed above, and let $f_0 \in \operatorname{Lip}_c(\partial\Omega)$ denote the restriction of *f* to $\partial\Omega$. Set $M := \sup_{\partial\Omega} f_0$, and observe that we seek to show that $(u - M)^+ = 0$ in Ω . Note that $(u - M)^+ \in Y_0^{1,2}(\Omega)$, thus, since $\nabla u = \nabla (u - M)^+$ in the set $\{X \in \Omega : (u(X) - M)^+ \neq 0\}$, and since Lu = 0 in the weak sense, we have

$$\begin{split} \iint_{\Omega} |\nabla (u - M)^{+}|^{2} dY &\approx \iint_{\Omega} A \nabla u \cdot \nabla (u - M)^{+} dY \\ &= -\iint_{\Omega} \mathbf{B} \cdot \nabla u (u - M)^{+} dY = \iint_{\Omega} \mathbf{B} \cdot \nabla (u - M)^{+} (u - M)^{+} dY \\ &\lesssim \varepsilon_{0} \iint_{\Omega} |\nabla (u - M)^{+}|^{2} dY \,, \end{split}$$

where in the last step we have used the Hardy inequality. For ε_0 small enough, we see that $\nabla (u - M)^+ = 0$ in Ω , and since the boundary trace of $(u - M)^+$ is zero, we must have $u \le M$ in Ω .

Next, we discuss continuity up to the boundary for the solutions to the Dirichlet problem constructed in Subsection 3.2. The following lemma will be useful in several circumstances.

Lemma 3.22. Let *L* be as in (3.17), where **B** satisfies (1.2). Suppose further that the conclusion of Lemma 3.11 holds for *L*, and that for each $f \in \text{Lip}_c(\partial\Omega)$, the equation Lu = 0 has a solution $u \in Y^{1,2}(\Omega)$ (or in $W^{1,2}(\Omega)$), with data $u_{\lfloor \partial \Omega \rfloor} = f$ (in the trace sense), for which the weak maximum principle holds. Then for every compact set $K \subset \overline{\Omega}$,

$$|u(Y) - f(x)| \le C_{K,f} |Y - x|^{\alpha/2}, \qquad Y \in K, x \in \partial\Omega,$$

where α is the Hölder exponent in (3.12), and where the constant $C_{K,f}$ may depend upon K and f, but otherwise depends only on the constants in Lemma 3.11.

Remark 3.24. In particular, Lemma 3.22 applies in the small constant case, by the construction above and Lemma 3.11. On the other hand, note that we do *not* impose smallness of the constant in (1.2), and in the sequel, we shall also use Lemma 3.22 in the absence of smallness. The same remark applies to Corollary 3.25 below.

Proof of Lemma 3.22. Let $f \in \text{Lip}_c(\partial\Omega)$, and let u_f be the corresponding solution. Let $x_0 \in \partial\Omega$, and $Y \in K$, and set $r = |Y - x_0|$. Since we allow the constant in (3.23) to depend on K and on f, we may suppose that r < 1. Let $\varphi \in C_0^{\infty}(B(x_0, 2))$ be a smooth bump function with $\varphi \equiv 1$ on $B(x_0, 1)$, and let $\Phi_r \in C^{\infty}(\mathbb{R}^{n+1})$, with $0 \le \Phi_r \le 1$, $\Phi_r \equiv 0$ on $B(x_0, 2\sqrt{r})$, and $\Phi_r \equiv 1$ on $\mathbb{R}^{n+1} \setminus B(x_0, 4\sqrt{r})$.

Set $\tilde{f} := f - f(x_0)\varphi$, define $f_r := \tilde{f}\Phi_r$, and let $u_{\tilde{f}}$ and u_{f_r} denote the corresponding solutions. Note that $\sup_{\partial\Omega} |\tilde{f} - f_r| \leq_f \sqrt{r}$, therefore by the maximum principle,

$$\sup_{\Omega} |u_{\tilde{f}} - u_{f_r}| \lesssim_f \sqrt{r},$$

Also by the maximum principle, we have $|u_{f_r}| \leq u_{|f_r|} \leq ||f||_{\infty}$, and since f_r vanishes on $\Delta(x_0, 2\sqrt{r}) = B(x_0, 2\sqrt{r}) \cap \partial\Omega$, as does the trace of the solution $v := 1 - u_{\varphi}$, we therefore have

$$|u_{\varphi}(Y) - 1| + |u_{f_r}(Y)| \leq_f \left(\frac{\delta(Y)}{\sqrt{r}}\right)^{\alpha}, \quad Y \in B(x_0, \sqrt{r}) \cap \Omega,$$

by Lemma 3.11. Since $\delta(Y) \le |Y - x_0| = r$, we obtain

$$|u_f(Y) - f(x_0)| \le |u_{\tilde{f}}(Y)| + |f(x_0)(u_{\varphi}(Y) - 1)| \le_f r^{\alpha/2} + r^{1/2} \le_f r^{\alpha/2}.$$

Corollary 3.25. Under the same hypotheses as in Lemma 3.22, we have

(3.26)
$$|u(Y) - u(X)| \le C_{K,f} |Y - X|^{\alpha/4}, \qquad X, Y \in K,$$

where α is the smaller of the two Hölder exponent in (3.5) and (3.12).

Proof. Since the constants in (3.26) are allowed to depend on *K* and *f*, we may assume that r := |X - Y| < 1/2, and that $\delta(X), \delta(Y) < 1$.

Suppose first that $r \leq \frac{1}{2} \max(\delta^2(X), \delta^2(Y))$. In this case, $\delta(X) \approx \delta(Y)$, so by (3.5), we have

$$|u(Y) - u(X)| \lesssim_f \left(\frac{|X - Y|}{\delta(X)}\right)^{\alpha} \lesssim_f r^{\alpha/2}$$

On the other hand, suppose that $r \ge \frac{1}{2} \max(\delta^2(X), \delta^2(Y))$, and fix $\hat{x}, \hat{y} \in \partial \Omega$ such that $\delta(X) = |X - \hat{x}|$ and $\delta(Y) = |Y - \hat{y}|$. Then

$$|u(Y) - u(X)| \le |u(Y) - f(\hat{y})| + |f(\hat{y}) - f(\hat{x})| + |u(X) - f(\hat{x})| =: I + II + III.$$

By Lemma 3.22, since $\delta(Y) + \delta(X) \leq r^{1/2}$ in the present case,

 $I + III \leq_{K,f} r^{\alpha/4}$.

By the triangle inequality, $|\hat{y} - \hat{x}| \le \delta(Y) + \delta(X) + |X - Y| \le r^{1/2}$, so that $II \le_f r^{1/2}$.

3.4. Existence of elliptic measure in the small constant case, and a local ampleness property. We establish solvability of the Dirichlet problem with continuous, compactly supported data on $\partial\Omega$, and consequently, we obtain existence of elliptic measure, for the operator *L* defined in (3.17), under the smallness assumption (3.1). To be precise, we have the following.

Lemma 3.27. Let *L* be as in (3.17), and suppose that (3.1) holds for **B**. If ε_0 is small enough, depending only on allowable parameters, then the continuous Dirichlet problem is solvable for the equation Lu = 0 in Ω , and therefore elliptic measure exists for *L* in Ω .

Proof. The proof is standard. Given a nonnegative function $f \in C_c(\partial\Omega)$, let $\{f_m\}_m$ be a sequence of nonnegative functions in $\operatorname{Lip}_c(\partial\Omega)$, such that $f_m \to f$ uniformly on $\partial\Omega$. Set $u_m := u_{f_m}$, the solution of the Dirichlet problem with data f_m , as constructed in Subsection 3.2. Since these solutions satisfy the weak maximum principle (Subsection 3.3), we have

$$||u_m - u_j||_{L^{\infty}(\Omega)} = ||u_{f_m - f_j}||_{L^{\infty}(\Omega)} \le ||f_m - f_j||_{L^{\infty}(\partial\Omega)},$$

18

thus, u_m converges uniformly in Ω , and therefore by Caccioppoli's inequality, u_m converges also in $W_{loc}^{1,2}(\Omega)$. Let *u* denote the limit. Then Lu = 0 in the weak sense. Indeed, for $\Phi \in C_0^{\infty}(\Omega)$,

$$0 = \iint_{\Omega} \left[A \nabla u_m \cdot \nabla \Phi + \mathbf{B} \cdot \nabla u_k \Phi \right] dY \to \iint_{\Omega} \left[A \nabla u \cdot \nabla \Phi + \mathbf{B} \cdot \nabla u \Phi \right] dY$$

Now let $x \in \partial \Omega$, $Y \in \Omega$, and observe that

$$|u(Y) - f(x)| \le |u(Y) - u_m(Y)| + |u_m(Y) - f_m(x)| + |f_m(x) - f(x)| =: I + II + III.$$

Let $\epsilon > 0$. Since $u_m \to u$ uniformly in Ω , and $f_m \to f$ uniformly on $\partial\Omega$, we can fix $m = m(\epsilon)$ such that $I + III \leq \epsilon$, uniformly for all $Y \in \Omega$ and $x \in \partial\Omega$. With this choice of *m* now fixed, we find that $II \leq \epsilon$, either by application of Lemma 3.22 in the case that *Y*, *x* lie in a suitable neighborhood of supp(f_m), or else by application of (3.12), in the case that *x* is away from supp(f_m) and *Y* is near enough to $\partial\Omega$.

Thus, we have shown that $u \in C(\overline{\Omega})$ with $u \equiv f$ on $\partial\Omega$. Moreover, u satisfies the weak maximum principle $||u||_{L^{\infty}(\Omega)} \leq ||f||_{L^{\infty}(\partial\Omega)}$, as may be seen by taking limits of the analogous inequality for each u_m, f_m . As usual, it then follows that the mapping $f \to u(X)$ defines a bounded linear functional on $C_c(\partial\Omega)$, for each $X \in \Omega$, so that

$$u(X) = \int_{\partial\Omega} f \, d\omega^X \,,$$

for some Radon measure ω^X (the elliptic measure for L), by Riesz representation.

In the next lemma, we suppose that ε_0 is small enough that elliptic measure exists, by Lemma 3.27.

Lemma 3.28. Let Ω be a 1-sided CAD, and set $L = -\operatorname{div} A\nabla + \mathbf{B} \cdot \nabla$, where **B** satisfies (3.1). Let ω denote elliptic measure for L in Ω . If ε_0 is small enough, depending only on dimension, ellipticity of A, and the 1-sided CAD constants, then there is a uniform constant $c \in (0, 1)$ with the same dependence, such that for every $x \in \partial \Omega$, and every $r \in (0, \operatorname{diam}(\partial \Omega))$, if $Y \in \Omega_{r/2} := \Omega \cap B(x, r/2)$, then

(3.29)
$$\omega^{r}(\Delta(x,r)) \ge c > 0.$$

Proof. Let $Y \in \Omega_{r/2}$. Set $u(Y) := 1 - \omega^Y(\Delta(x, r))$. Suppose first that $\delta(Y) \le \eta r$, where $\eta > 0$ is a sufficiently small number to be chosen. Then by Lemma 3.11, $u(Y) \le C\eta^{\alpha} \le 1/2$, provided that η is small enough. With this choice of η now fixed, we see that $\omega^Y(\Delta(x, r)) \ge 1/2$, for all $Y \in \Omega_{r/2} \cap \{\delta(Y) \le \eta r\}$. In particular, the latter bound holds if $Y = Y_0$ is a Corkscrew point relative to the ball $B(x, \eta r)$. Since $\delta(Y_0) \approx_{\eta} r$, we see that (3.29) holds for all $Y \in \Omega_{r/2} \cap \{\delta(Y) > \eta r\}$, with $c = c(\eta)$, by Harnack's inequality.

3.5. Existence of elliptic measure in the large constant case, and an approximation result. In this subsection, we show that there is an elliptic measure for an operator L as in (3.17), with **B** satisfying (1.2), provided that L can be suitably approximated by operators

$$L_k := -\operatorname{div} A \nabla + \mathbf{B}_k \cdot \nabla = L_0 + \mathbf{B}_k \cdot \nabla,$$

for which the hypotheses of Lemma 3.22 (and thus also of Corollary 3.25) hold for each L_k with constants that are uniform in k. Specifically, we shall set

$$\mathbf{B}_k := \mathbf{B} \mathbf{1}_{\Omega_k},$$

where $\Omega_k := \{X \in \Omega : \delta(X) > 1/k\}$. Then $||\mathbf{B}_k||_{L^{\infty}(\Omega)} \le k \sqrt{M_0}$, by (1.2), and $L_k = L_0$ in the border strip $\{X \in \Omega : \delta(X) \le 1/k\}$.

In the next pair of Lemmata, we let ω_k^X and ω^X denote the respective elliptic measures, at the point X, for the operators L_k and L in Ω .

Lemma 3.31. Let L be as in (3.17), with **B** satisfying (1.2). Define L_k as above. Then for each positive integer k, the continuous Dirichlet problem is solvable for the equation $L_k u = 0$ in Ω , the solution satisfies the weak maximum principle, and consequently elliptic measure ω_k exists for L_k .

Lemma 3.32. Let L be as in (3.17), with **B** satisfying (1.2). Define L_k as above, and suppose that the hypotheses of Lemma 3.22 hold for each L_k , and that Lemma 3.11 holds for each L_k , with constants that are uniform in k. Then elliptic measure ω exists for L, and a subsequence $\omega_{k_k}^X$ converges weakly to ω^X , for each $X \in \Omega$.

Proof of Lemma 3.31. We follow a familiar strategy. First consider $f \in \operatorname{Lip}_c(\mathbb{R}^{n+1})$. Since \mathbf{B}_k is (qualitatively) bounded, by [KS, Lemma 4.2] in the case that Ω is bounded (respectively, [M, Theorem 5.3] in the case that Ω is unbounded), there is a solution w of the equation $L_k w = L_k f$, belonging to $W_0^{1,2}(\Omega)$ (resp., $Y_0^{1,2}(\Omega)$), with constants that may depend upon k. Set u := f - w, so that $u \in W^{1,2}(\Omega)$ (resp., $Y^{1,2}(\Omega)$) is a weak solution of $L_k u = 0$, with $u|_{\partial\Omega} = f$ in the trace sense. Then by [GT, Lemma 8.1] (resp., [M, Theorem 5.1]), the weak maximum principle holds:

(3.33)
$$\sup_{\Omega} |u| \le \sup_{\partial \Omega} |f|, \quad f \in \operatorname{Lip}_{c}(\mathbb{R}^{n+1}).$$

We now proceed as in the proof of Lemma 3.27: given a nonnegative function $f \in C_c(\partial\Omega)$, we let $\{f_m\}_m$ be a sequence in $\operatorname{Lip}_c(\partial\Omega)$, such that $f_m \to f$ uniformly on $\partial\Omega$. Set $u_m := u_{f_m}$, the solution of the Dirichlet problem with data f_m , as constructed above. Since these solutions satisfy the weak maximum principle (3.33), we have

$$\|u_m - u_j\|_{L^{\infty}(\Omega)} = \|u_{f_m - f_j}\|_{L^{\infty}(\Omega)} \le \|f_m - f_j\|_{L^{\infty}(\partial\Omega)},$$

thus, u_m converges uniformly in Ω , and therefore by Caccioppoli's inequality, u_m converges also in $W_{loc}^{1,2}(\Omega)$. The limit u is therefore a solution of $L_k u = 0$ in Ω .

Recall that by construction, $L_k = L_0$ when $\delta(X) \le 1/k$, and therefore Lemma 3.11 and Lemma 3.22 apply (with constants that may depend upon k) to the pair u_m and f_m . The rest of the proof now follows that of Lemma 3.27, essentially verbatim, so we leave the remaining details to the reader.

Proof of Lemma 3.32. Let $f \in \operatorname{Lip}_c(\partial\Omega)$, and set u_k be the $Y^{1,2}$ (or $W^{1,2}$) solution of $L_k u_k = 0$, with data f. By assumption, the hypotheses of Lemma 3.11 hold uniformly in k for each L_k , thus by Lemma 3.22 and Corollary 3.25, the family $\{u_k\}$ is equicontinuous on each compact set $K \subset \overline{\Omega}$, and moreover, the solutions u_k are uniformly bounded on Ω , by the weak maximum principle. Consequently, for each such K, there is a subsequence of u_{k_j} converging uniformly on K, and in particular, on each bounded subdomain Ω' compactly contained in Ω . Observe that $L_k = L$ in any such Ω' , for all k large enough (depending on Ω'). Hence, by Caccioppoli's inequality (Lemma 3.8), the convergence holds also in $W^{1,2}(\Omega')$. By choosing a countable collection of compact sets that exhaust $\overline{\Omega}$ (obviously, this step is unnecessary when Ω is bounded), we find a subsequential limit u defined on all of Ω . Note that

$$||u||_{L^{\infty}(\Omega)} \leq \sup_{k} ||u_{k}||_{L^{\infty}(\Omega)} \leq ||f||_{L^{\infty}(\Omega)},$$

since the solutions u_k satisfy the weak maximum principle.

Let $\Phi \in C_0^{\infty}(\Omega)$. Then $\mathbf{B}_k = \mathbf{B}$, hence $L_k = L$, in the support of Φ , provided k is large enough. Thus $Lu_k = 0$ in supp (Φ) , and by the $W^{1,2}$ convergence on compactly contained subdomains, we conclude that u is a weak solution of Lu = 0 in Ω .

We now claim that $u \in C(\overline{\Omega})$, and that u = f on $\partial\Omega$. Continuity in Ω is immediate from (3.5), since Lu = 0. Let us now verify continuity up to the boundary. Let $x \in \partial\Omega$, and let $Y \in \Omega \cap B(x, 1) =: \Omega_1(x)$. By re-numbering, we may assume that $u_k \to u$ uniformly on a compact set K_1 containing $\Omega_1(x)$. We then have

$$|u(Y) - f(x)| \le |u(Y) - u_k(Y)| + |u_k(Y) - f(x)| \le |u(Y) - u_k(Y)| + C_{K_1, f}|Y - x|^{\alpha/2}$$

since Lemma 3.11, and hence the quantitative conclusion of Lemma 3.22, apply to u_k with constants independent of k. Letting $k \to \infty$, we find that (3.23) holds for u, thus in particular $u \in C(\overline{\Omega})$, with $u|_{\partial\Omega} = f$, as claimed.

As usual, existence of the elliptic measure ω now follows from the Riesz representation theorem. Finally, the claimed weak convergence is simply the statement that for $f \in C_c(\partial\Omega)$, and for each $X \in \Omega$, if u_k and u denote, respectively, the solutions to $L_k u_k = 0$ and Lu = 0with data f, then for some subsequence,

$$\int_{\partial\Omega} f \, d\omega_{k_j}^X = u_{k_j}(X) \to u(X) = \int_{\partial\Omega} f \, d\omega^X \,.$$

3.6. The Green function: existence, estimates, and consequences.

Lemma 3.34 (Green function existence, properties and estimates). Let $\Omega \subset \mathbb{R}^{n+1}$ be a bounded 1-sided CAD, and let $L = -\operatorname{div} A\nabla + \mathbf{B} \cdot \nabla$. Suppose that **B** satisfies (3.1), and is qualitatively bounded. Then there exists a unique Green function G(X, Y) defined on $\Omega \times \Omega \setminus \{X = Y\}$, satisfying

(3.35)
$$G(Z, \cdot), G(\cdot, Z) \in W^{1,2}(\Omega \setminus B(Z, r)) \quad \forall Z \in \Omega, r > 0;$$

(3.36)
$$G(Z, \cdot)|_{\partial\Omega} \equiv 0 \equiv G(\cdot, Z)|_{\partial\Omega}$$
 in the trace sense, $\forall Z \in \Omega$;

$$(3.37) G(X,Y) \ge 0, \forall X, Y \in \Omega, X \neq Y$$

Moreover, setting $\mathfrak{G}F(X) := \iint_{\Omega} G(X, Y)F(Y)dy$ and $\mathfrak{G}^*F(Y) := \iint_{\Omega} G(X, Y)F(X)dX$, we have

(3.38)
$$\mathfrak{G}F, \,\mathfrak{G}^*F \in W^{1,2}_0(\Omega), \quad \forall F \in L^\infty(\Omega);$$

(3.39)
$$L \mathfrak{G} F = F$$
 and $L^* \mathfrak{G}^* F = F$, $\forall F \in L^{\infty}(\Omega)$,

in the weak sense. In addition, for every $\Phi \in \operatorname{Lip}_{c}(\mathbb{R}^{n+1})$, and $X \in \Omega$, we have

(3.40)
$$u(X) - \Phi(X) = -\iint_{\Omega} \left[A(Z)\nabla_Z G(X,Z) + \mathbf{B}(Z)G(X,Z) \right] \cdot \nabla \Phi(Z) \, dZ$$

where *u* is the $W^{1,2}$ solution of Lu = 0 with $u|_{\partial\Omega} = \Phi|_{\partial\Omega}$ in the trace sense.

Finally, if ε_0 is sufficiently small, depending only on n, ellipticity of A, and the 1-sided CAD constants, then

 $(3.41) |X - Y|^{1-n} \leq G(X, Y) \leq |X - Y|^{1-n}, |X - Y| \leq \frac{1}{2}\delta(Y),$

where the implicit constants also depend only on the allowable parameters. In particular, ε_0 and the implicit constants in (3.41) do **not** depend upon the qualitative L^{∞} bound for **B**.

Proof of Lemma 3.34. By the qualitative assumption, the existence of a unique Green function, satisfying properties (3.35) - (3.40), follows from the results in [KS] (or the results of [S]).⁵ For these results, the 1-sided CAD assumption is not needed.

On the other hand, property (3.41) is the only quantitative conclusion in the Lemma, and to prove it, we cannot use the pointwise bounds of [KS] off the shelf, since those estimates depend quantitatively on impermissible quantities; however, we can and do use the aforementioned existence and qualitative properties of the Green function proved in [KS].

Let us now give the proof of (3.41). We first establish the right hand inequality. We recall the standard upper and lower Sobolev exponents in \mathbb{R}^d , namely, with d = n + 1,

$$2^* = 2d/(d-2) = 2(n+1)/(n-1), \quad 2_* = 2d/(d+2) = 2(n+1)/(n+3).$$

First note that by Lemma 3.19, if ε_0 is small enough, then for any given $F \in Y^{-1,2}$, there is a unique $w \in Y_0^{1,2}(\Omega)$ solving Lw = F in the weak sense. Moreover, w satisfies the bound (3.20), so in particular

$$\|w\|_{L^{2^*}(\Omega)} \lesssim \|F\|_{L^{2^*}(\Omega)},$$

with implicit constants depending only on allowable parameters.

On the other hand, by [KS, Lemma 4.2, and Propositions 5.3 and 6.13], given $F \in W^{-1,2}(\Omega)$, there is a unique $w_1 \in W^{1,2}_0\Omega$ such that $Lw_1 = F$ in the weak sense. By Sobolev embedding, $w_1 \in Y^{1,2}_0(\Omega)$, hence by uniqueness, for $F \in L^{\infty}(\Omega)$, we have $w_1 = w$, the solution constructed in Lemma 3.19, which therefore satisfies (3.42). Moreover, for $F \in L^{\infty}(\Omega)$, we have the formula

$$w(X) = \iint_{\Omega} G(X, Y) F(Y) dY.$$

Suppose now that $r := |X - Y| \le \delta(Y)/4$. Let $B_1 := B(X, r/4)$, $B_2 = B(Y, r/4)$, and set $F := 1_{B_2}$. Then by Harnack's inequality (3.6), and (3.42),

$$|B_1|^{1/2^*}|B_2|G(X,Y) \lesssim \left(\iint_{B_1} \left(\iint_{B_2} G(Z,W) \, dW\right)^{2^*} dZ\right)^{1/2^*} \lesssim |B_2|^{1/2_*},$$

i.e, in ambient dimension n + 1,

$$G(X,Y) \leq |B_1|^{-1/2^*} |B_2|^{-1/2^*} \approx r^{1-n} = |X-Y|^{1-n}.$$

Using Harnack again, we may extend the estimate from the regime $|X - Y| \le \delta(Y)/4$ to $|X - Y| \le \delta(Y)/2$.

We now turn to the left hand inequality in (3.41). We follow the argument in [HL]. Note that by Lemma 3.27, the elliptic measure for *L* exists. Consequently, for $\Phi \in C_0^{\infty}(\mathbb{R}^{n+1})$, with $\Phi(X) = 0$, (3.40) becomes

(3.43)
$$\int_{\partial\Omega} \Phi \, d\omega^X = -\iint_{\Omega} \left[A(Y) \nabla_Y G(Y, X) + \mathbf{B}(Y) G(X, Y) \right] \cdot \nabla \Phi(Y) \, dY \,,$$

⁵In the case of an unbounded domain, one could substitute the results of [M] for those of [KS, S]. We have not done this here, since we shall use the Green function only in bounded subdomains of our original domain.

Now set $s = |X - Y| \le \delta(Y)/2$, and choose $\Phi \in C_0^{\infty}(\mathbb{R}^{n+1})$ with $0 \le \Phi \le 1$, $\Phi \equiv 0$ in B(X, s/4), and $\Phi \equiv 1$ on $\overline{\Omega} \setminus B(X, s/2)$, with $|\nabla \Phi| \le 1/s$. Since the left hand side of (3.43) equals 1, and since $\delta(Z) \ge s$ in the support of $\nabla \Phi$, we have

$$1 \lesssim \frac{1}{s} \iint_{s/4 \le |X-Z| \le s/2} \left(|\nabla_Z G(X,Z)| + s^{-1} G(X,Z) \right) dZ \lesssim G(X,Y) s^{n-1}$$

by Harnack's inequality, where we have used Cauchy-Schwarz and Caccioppoli's inequality to handle the gradient term. The left hand inequality in (3.41) follows.

In the next lemma, and in the sequel, we use the notation

$$(3.44) \qquad \Delta_Y := B(Y, 10\delta(Y)) \cap \partial\Omega, \qquad Y \in \Omega.$$

Note that Δ_Y is essentially a surface ball. Indeed, if $\hat{y} \in \partial \Omega$ is chosen so that $|Y - \hat{y}| = \delta(Y)$, then $\Delta(\hat{y}, 9\delta(Y)) \subset \Delta_Y \subset \Delta(\hat{y}, 11\delta(Y))$.

Lemma 3.45 ("CFMS" estimates). Suppose that Ω is a bounded 1-sided CAD, and let $L = -\operatorname{div} A \nabla + \mathbf{B} \cdot \nabla$, where **B** satisfies (3.1), and is qualitatively bounded. If ε_0 is sufficiently small, depending only on *n*, ellipticity of *A*, and the 1-sided CAD constants, then for all $X, Y \in \Omega$ such that $|X - Y| \ge \delta(Y)/4$ we have

(3.46)
$$\frac{G(X,Y)}{\delta(Y)} \lesssim \frac{\omega^X(\Delta_Y)}{\sigma(\Delta_Y)},$$

and for all $X, Y \in \Omega$ such that $X \in \Omega \setminus B(Y, 50\kappa_0\delta(Y))$

(3.47)
$$\frac{\omega^X(\Delta_Y)}{\sigma(\Delta_Y)} \lesssim \frac{G(X,Y)}{\delta(Y)},$$

where the implicit constants depend only on allowable parameters, and κ_0 is the constant fixed in Lemma 2.40.

Proof. We follow the argument in [K, p. 10]. Given $Y \in \Omega$, and using the right hand inequality in (3.41), we see that for $|X - Y| = \delta(Y)/4$,

$$\frac{G(X,Y)}{\delta(Y)} \lesssim \delta(Y)^{-n} \lesssim \frac{\omega^X(\Delta_Y)}{\sigma(\Delta_Y)}$$

by Lemma 3.28 (and Harnack's inequality), and ADR. Estimate (3.46) now follows by the maximum principle.

To prove (3.47), we set $R := \delta(Y)$, and fix $Y \in \Omega$ and $X \in \Omega \setminus B(Y, 20R)$. Choose a smooth bump function $\Phi \in C_0^{\infty}(B(Y, 11R))$, with $0 \le \Phi \le 1$, $\Phi \equiv 1$ on B(Y, 10R), and $|\nabla \Phi| \le R^{-1}$. Choose $\hat{y} \in \partial \Omega$ such that $|Y - \hat{y}| = \delta(Y) = R$, and for any r > 0 set $\Omega_r := \Omega \cap B(\hat{y}, r)$. Note that $\Phi(X) = 0$, so that by (3.40) and Lemma 3.27,

$$\begin{split} \omega^{X}(\Delta_{Y}) &\leq \int_{\partial\Omega} \Phi \, d\omega^{X} = -\iint_{\Omega} \left[A(Z) \nabla_{Z} G(X,Z) \, + \, \mathbf{B}(Z) G(X,Z) \right] \cdot \nabla \Phi(Z) \, dZ \\ &\lesssim \frac{1}{R} \iint_{\Omega_{12R}} \left(|\nabla_{Z} G(X,Z)| \, + \, \frac{G(X,Z)}{\delta(Z)} \right) \, dZ \\ &\lesssim R^{(n-1)/2} \left(\iint_{\Omega_{12\kappa_{0}R}} |\nabla_{Z} G(X,Z)|^{2} dZ \right)^{1/2} \end{split}$$

$$\lesssim R^{(n-1)/2} R^{-1} \left(\iint_{\Omega_{24\kappa_0 R}} G^2(X,Z) dZ \right)^{1/2} \lesssim R^{n-1} G(X,Y) \,,$$

where we have used Cauchy-Schwarz and the local Hardy inequality (Lemma 2.40), then Caccioppoli at the boundary (Lemma 3.9) and finally the Carleson estimate (Lemma 3.11, specifically (3.13)). Since $R = \delta(Y)$, the ADR property yields (3.47).

Lemma 3.48. Under the same hypotheses as in Lemma 3.45, for any ball B = B(x, r), centered on $\partial\Omega$, setting $\Delta = B \cap \partial\Omega$, we have

$$\omega^X(2\Delta) \, \lesssim \, \omega^X(\Delta) \,, \qquad X \in \Omega \setminus 4B$$

where the implicit constant depends only on the allowable parameters.

Proof. The lemma will be an easy corollary of Lemma 3.45, and the proof is the standard one. Let κ_0 be the fixed constant in Lemma 2.40 and Lemma 3.45, and let M be a sufficiently large number to be fixed momentarily. Given a ball B = B(x, r), let $X \in \Omega \setminus 4B$, let Y be a corkscrew point relative to the ball B(x, r/M), and cover 2Δ by a collection of surface balls $C = {\Delta_k}_k$, such that $\Delta_k = B_k \cap \partial\Omega$, and each B_k is centered on $\partial\Omega$, and has radius $r_{B_k} = r/(M^2\kappa_0)$. By the ADR property of $\partial\Omega$, we may do this in such a way that C has cardinality $\#(C) \approx (M^2\kappa_0)^n$. For each k, let Y_k be a corkscrew point relative to MB_k . Choosing M large enough, depending only on the constants in the Corkscrew condition, we have $\Delta_Y \subset \Delta$, and also $\Delta_k \subset \Delta_{Y_k}$, and furthermore, $X \in \Omega \setminus B(Y_k, 50\kappa_0\delta(Y_k))$. Note that, since κ_0 and M have been fixed, we have

$$\delta(Y) \approx \delta(Y_k) \approx r, \quad \forall k,$$

and thus also, by ADR,

$$\sigma(2\Delta) \approx \sigma(\Delta) \approx \sigma(\Delta_{Y_k}) \approx \sigma(\Delta_Y) \approx r^n$$

By Lemma 3.45, and then Harnack's inequality, we then have

$$\frac{\omega^X(\Delta_{Y_k})}{r^n} \lesssim \frac{G(X,Y_k)}{r} \lesssim \frac{G(X,Y)}{r}$$

Consequently, since 2Δ is covered by the collection *C*, we see that

$$\frac{\omega^X(2\Delta)}{r^n} \leq \sum_k \frac{\omega^X(\Delta_{Y_k})}{r^n} \lesssim \sum_k \frac{G(X,Y)}{r} \lesssim \frac{\omega^X(\Delta_Y)}{r^n} \leq \frac{\omega^X(\Delta)}{r^n},$$

where in the last two steps we have used (3.46) and the fact that the cardinality of *C* is bounded by a universal constant, and that $\Delta_Y \subset \Delta$.

Finally, we record two results that hold for the homogeneous second order operator $L_0 = -\operatorname{div} A\nabla$. These results are well known. We refer the reader to, e.g., [K, pp. 10-11]: the proofs given there carry over routinely to our setting, and we omit the details.

Lemma 3.49. (Comparison Principle/Boundary Harnack Principle) Let Ω be a 1-sided CAD, let $\Delta = B \cap \partial \Omega$ be a surface ball centered on $\partial \Omega$, and let T_{Δ} be the associated Carleson tent, as in Proposition 2.26 (8). Suppose that u, v are two non-negative $W^{1,2}$ solutions of the equation $L_0 u = L_0 v = 0$ in the tent $T_{2\Delta}$, vanishing in the trace sense on $\partial T_{2\Delta} \cap \partial \Omega$ (and hence continuously on $\partial T_{\Delta} \cap \partial \Omega$, by Lemma 3.11 and properties of the tents). Let X_B be a corkscrew point relative to the ball B. Then

$$\sup_{X\in T_{\Delta}} \frac{u(X)}{v(X)} \lesssim \frac{u(X_B)}{v(X_B)},$$

where the implicit constant depends only on ellipticity, dimension, and the 1-side CAD constants.

Lemma 3.50. (Pole Change formula) Let Ω be a 1-sided CAD, let B be a ball centered on $\partial \Omega$, and let X_B be a corkscrew point relative to B. Let ω_0 denote elliptic measure for L_0 . Then for every Borel set $E \subset \Delta := B \cap \partial \Omega$, we have

$$\frac{\omega_0^X(E)}{\omega_0^X(\Delta)} \approx \omega_0^{X_B}(E), \qquad \forall X \in \Omega \setminus 2B.$$

Lemma 3.50 follows from Lemma 3.45, along with Lemma 3.49 applied to the Green function $G_{L_0}(X, Y)$ in the adjoint variable Y. We remark that we do not know whether Lemma 3.50 holds for L (as opposed to L_0), even in the small constant case. The obstacle is that we do not know whether Lemma 3.49 holds for non-negative solutions of the adjoint equation $L^*u = 0$ (in particular, for the Green function for L, in the adjoint variable).

4. Perturbing L_0 in a small constant case

In this section, we prove an auxiliary result in a suitable "small constant" case, analogous to [FKP, Theorem 2.5], in which we show that elliptic measure for L belongs to A_{∞} with respect to the elliptic measure for L_0 . Before stating the precise result, let us record some notation and a couple of definitions.

In this section, we assume that Ω is a *bounded* 1-sided CAD, with diam(Ω) =: r_0 . We let $X_0 \in \Omega$ be a fixed corkscrew point with respect to the entire boundary, which we can view as $\Delta_{2r_0} = \partial \Omega$, where Δ_{2r_0} is any surface ball on $\partial \Omega$ of radius $2r_0$, and arbitrary center on $\partial \Omega$. Thus

$$\delta(X_0) \approx r_0$$

Set $L_0 := -\operatorname{div} A\nabla$, $L := L_0 + \mathbf{B} \cdot \nabla$, let G_0 and G denote, respectively, the Green functions for L_0 and L in Ω , and let ω_0^X and ω^X denote the corresponding elliptic measures, at any point $X \in \Omega$. In the special case that $X = X_0$, we shall often simply write $\omega_0 := \omega_0^{X_0}$, and $\omega := \omega^{X_0}$. We ensure the existence of G and ω , by first working with truncations of \mathbf{B} , as in Subsection 3.5. After establishing appropriate quantitative estimates for elliptic measure that hold uniformly for all the truncations, we may then pass to the limit. We shall return to the latter point shortly, in more detail (see the beginning of the proof of Theorem 4.8).

Definition 4.1. (ε -smallness) We say that **B** is " ε -small with respect to ω_0 " if

(4.2)
$$|\mathbf{B}(Y)| \le \frac{\varepsilon}{\delta(Y)}, \quad \text{a.e. } Y \in \Omega,$$

and if for every $\Delta = B \cap \partial \Omega$, with $B = B(x, r), x \in \partial \Omega, 0 < r < r_0$,

(4.3)
$$\iint_{B\cap\Omega} |\mathbf{B}(Y)|^2 G_0(X_0, Y) \, dY \leq \varepsilon^2 \omega_0(\Delta)$$

Recall that, associated to any point $X \in \Omega$, we use the notation

$$\Delta_X := B(X, 10\delta(X)) \cap \partial\Omega$$

which essentially defines a surface ball on $\partial \Omega$.

Definition 4.4. ((m, c_0 , θ)-ampleness) Let m be a doubling Borel measure on $\partial\Omega$, and let c_0 , $\theta \in (0, 1)$. We say that an elliptic measure ω has the (m, c_0 , θ)-ampleness property if for all $X \in \Omega$, and for any Borel set $F \subset \Delta_X$,

(4.5)
$$\mathfrak{m}(F) \ge (1-\theta)\mathfrak{m}(\Delta_X) \implies \omega^X(F) \ge c_0.$$

We shall require the following lemma, which is a routine extension of [BL, Lemma 2.2]. For the purposes of Lemma 4.6, we suppose that L is an elliptic operator (possibly with a drift term) defined on Ω , for which the continuous Dirichlet problem (Definition 2.18) is solvable, the weak maximum principle holds, and for which positive solutions satisfy Harnack's inequality.

Lemma 4.6. Suppose that Ω is an open set with an ADR boundary. Let L be an elliptic operator as in the preceeding paragraph, and let ω denote the elliptic measure associated to L. Let m be a doubling Borel measure on $\partial\Omega$, with doubling constant N_{db} , and suppose that ω has the $(\mathfrak{m}, \mathfrak{c}_0, \theta)$ -ampleness property for some constants $\mathfrak{c}_0, \theta \in (0, 1)$. Then, given $s \in (0, 1)$, there is a constant $\eta = \eta(\theta, s, N_{db})$ such that for any surface ball $\Delta = B \cap \partial\Omega$ with radius less than $\frac{1}{2} \operatorname{diam}(\partial\Omega)$, and for any Borel set $F \subset \Delta$, we have the implication

(4.7)
$$\mathfrak{m}(F) \ge (1 - \eta)\mathfrak{m}(\Delta) \implies \omega^{Z}(\frac{1}{2}\Delta) \le C_{s}\omega^{Z}(F) + s\omega^{Z}(\Delta), \quad \forall Z \in \Omega \setminus 2B.$$

We omit the proof of Lemma 4.6, which originally appeared as [BL, Lemma 2.2], in the special case that L is the Laplacian, and $m = \sigma$, the surface measure on $\partial \Omega$. The proof carries over to the present setting essentially unchanged: in fact, the argument in [BL] requires only that L and m satisfy the properties stated above.

Our main result in the section is the following analogue of [FKP, Theorem 2.5]. We use the notation and terminology discussed above, in the preamble to this section.

Theorem 4.8. Let Ω be a bounded 1-sided CAD, with diam $(\Omega) = r_0 < \infty$. Then there is a constant $\varepsilon_0 > 0$, depending on dimension, ellipticity of the coefficient matrix A, and the 1-sided CAD constants, such that if **B** is ε -small with respect to $\omega_0 := \omega_0^{X_0}$, with $\varepsilon \le \varepsilon_0$, then $\omega \in A_{\infty}(\omega_0)$ in the sense of Definition 2.20 with $\mathfrak{m} = \omega_0$, and all constants implicit in the A_{∞} condition depend only on the same allowable parameters. In particular, $\omega^{X_0} \in A_{\infty}(\omega_0, \partial\Omega)$.

Proof. We shall work with the operator L_k , with truncated drift coefficient \mathbf{B}_k , as in Subsection 3.5. Note that the smallness conditions of Definition 4.1 hold uniformly in k. Then elliptic measure exists, by Lemma 3.27. Also, \mathbf{B}_k is (qualitatively) bounded (depending on k), so the conclusions of Lemmas 3.34, 3.45, and 3.48 are all valid, and all the quantitative bounds (explicit and implicit) in those Lemmas hold uniformly in k. Moreover, the hypotheses of Lemma 3.32 also hold. Note further that since $L_0 = L_k$ near the boundary, ω_0 and ω_k are (qualitatively) mutually absolutely continuous, hence $h_k^X := d\omega_k^X/d\omega_0$ exists, for each $X \in \Omega$. Recall that $\omega_0 := \omega_0^{X_0}$. We shall show that $h_k^X \in L^2(\omega_0)$, uniformly in k, with an L^2 bound that in turn implies the claimed A_{∞} property. We may then invoke Lemma 3.32 to deduce that $h^X := d\omega_k^X/d\omega_0$ exists, and belongs to $L^2(\omega_0)$, with the same bound that held uniformly for all h_k^X , whence the conclusion of Theorem 4.8 follows.

For the rest of the proof, we therefore assume that $\mathbf{B} = \mathbf{B}_k$ for some $k \ge 1$, but we suppress the subscript *k* to simplify the notation. We may then invoke Lemmas 3.34, 3.45, and 3.48 as needed. Of course, all of the constants appearing in our estimates will be independent of the truncation, which allows passage to the limit.

By Lemma 3.48, for $\Delta = B \cap \partial \Omega$, with B centered on $\partial \Omega$, we have the doubling estimate

(4.9)
$$\omega^{Z}(\Delta) \leq N_{1} \, \omega^{Z} \left(\frac{1}{2}\Delta\right), \qquad \forall Z \in \Omega \setminus 2B,$$

uniformly for all $\varepsilon \leq \varepsilon_0$, for a sufficiently small $\varepsilon_0 > 0$.

27

Claim. We claim that for all $\varepsilon \leq \varepsilon_0$, with ε_0 sufficiently small,

(4.10)
$$\|h^X\|_{L^2(\partial\Omega,\omega_0)} \lesssim \omega_0(\Delta_X)^{-1/2} + \varepsilon$$

for some uniform implicit constant depending only on n, the 1-sided CAD constants, and ellipticity of A.

Let us take the claim for granted momentarily. Fix $X \in \Omega$, and let $E \subset \Delta_X$ be a Borel set satisfying

(4.11)
$$\omega_0(E) \le \theta \omega_0(\Delta_X),$$

where $\theta \in (0, 1)$ will be chosen momentarily. Then for all $\varepsilon \leq \varepsilon_0$,

$$\begin{split} \omega^{X}(E) &= \int_{E} h^{x} d\omega_{0} \leq \omega_{0}(E)^{1/2} \, \|h^{X}\|_{L^{2}(\partial\Omega,\omega_{0})} \leq \left(\frac{\omega_{0}(E)}{\omega_{0}(\Delta_{X})}\right)^{1/2} + \, \omega_{0}(E)^{1/2} \varepsilon \\ &\lesssim \, \theta^{1/2} + \varepsilon \, \leq \, \left(\theta^{1/2} + \varepsilon_{0}\right) \omega^{X}(\Delta_{X}) \,, \end{split}$$

by (4.10), (4.11), and then Lemma 3.28. Choosing θ and ε_0 sufficiently small, we obtain that $\omega^X(E) \le \frac{1}{2}\omega^X(\Delta_X)$, whence it follows that for some uniform $c_0 > 0$,

$$\omega_0(F) \ge (1-\theta)\,\omega_0(\Delta_X) \implies \omega^X(F) \ge \frac{1}{2}\omega^X(\Delta_X) \ge c_0$$

again by Lemma 3.28, provided that $\varepsilon \leq \varepsilon_0$; i.e., ω is (ω_0, c_0, θ) -ample. Fixing this choice of θ , and recalling that N_1 denotes the doubling constant for ω , we may therefore apply Lemma 4.6, with $\mathfrak{m} = \omega_0$, and with $s = 1/(2N_1)$, to deduce that there is a uniform choice of $\eta \in (0, 1)$ such that for any surface ball $\Delta = B \cap \partial \Omega$ with radius less than $\frac{1}{2} \operatorname{diam}(\partial \Omega)$, for any Borel set $F \subset \Delta$, and for every $Z \in \Omega \setminus 2B$,

$$\begin{split} \omega_0(F) &\geq (1 - \eta) \,\omega_0(\Delta) \implies \frac{1}{N_1} \omega^Z(\Delta) \leq \omega^Z \left(\frac{1}{2} \Delta\right) \leq C_s \omega^Z(F) + s \omega^Z(\Delta) \\ &= C_s \omega^Z(F) + \frac{1}{2N_1} \omega^Z(\Delta) \,. \end{split}$$

Thus, hiding the last term, we have

(4.12)
$$\omega_0(F) \ge (1 - \eta)\,\omega_0(\Delta) \implies \omega^Z(F) \gtrsim \omega^Z(\Delta), \quad \forall Z \in \Omega \setminus 2B,$$

where the implicit constant is uniform. Since (4.12) holds also with Δ replaced by $\Delta' = B' \cap \partial \Omega$, whenever $B' \subset B$, we see that $\omega^Z \in A_{\infty}(\Delta, \omega_0)$ for each $Z \in \Omega \setminus 2B$. In particular, taking $Z = X_0$, we obtain the conclusion of Theorem 4.8, modulo (4.10).

It therefore remains only to verify the claimed estimate (4.10). To this end, let $f \in L^2(\partial\Omega, \omega_0)$ be non-negative, with $||f||_{L^2(\partial\Omega, \omega_0)} \leq 1$, and set

$$u_0(X) = u_{0,f}(X) := \int_{\partial\Omega} f d\omega_0^X, \quad u(X) = u_f(X) := \int_{\partial\Omega} f d\omega^X = \int_{\partial\Omega} f h^X d\omega_0.$$

By Lemma 3.50,

$$\frac{d\omega_0^X}{d\omega_0} \approx \omega_0 (\Delta_X)^{-1} \, .$$

and therefore, since ω_0^X is a probability measure,

(4.13)
$$u_0(X) \le \left(\int_{\partial \Omega} f^2 d\omega_0^X \right)^{1/2} \le \|f\|_{L^2(\omega_0)} \, \omega_0(\Delta_X)^{-1/2} \le \omega_0(\Delta_X)^{-1/2} \, .$$

In particular, if $X = X_0$, we simply have

(4.14)
$$u_0(X_0) \le \left(\int_{\partial\Omega} f^2 d\omega_0\right)^{1/2} \le 1$$

Observe also that

 $||h^X||_{L^2(\omega_0)} = \sup u_f(X),$

where the supremum runs over all non-negative $f \in L^2(\omega_0)$ with $||f||_{L^2(\omega_0)} \leq 1$. Consequently, to prove (4.10), it is enough to show that

$$(4.15) |u(X) - u_0(X)| \leq u_0(X) + \varepsilon ||h^X||_{L^2(\omega_0)} + \varepsilon$$

Since $u_0 - u = 0$ on $\partial \Omega$, we may follow the idea of [FKP] to write, using (3.40),

$$\begin{split} F(X) &:= u_0(X) - u(X) = \iint_{\Omega} G(X,Y) \mathbf{B}(Y) \cdot \nabla u_0(Y) \, dY \\ &= \iint_{|X-Y| < \delta(Y)/4} \dots \, dY \, + \, \iint_{\Omega \cap \{|X-Y| \ge \delta(Y)/4\}} \dots \, dY \, =: \, I + II \,, \end{split}$$

With X fixed, set $r = \delta(X)$. Observe that in term I, $|X - Y| < \delta(Y)/4$ implies that $\delta(Y)/4 < \delta(X)/3$. Thus, by (4.2) and the pointwise estimate for the Green function (in ambient dimension \mathbb{R}^{n+1}),

$$\begin{split} |I| \, &\lesssim \, \varepsilon \, r^{-1} \, \iint_{|X-Y| < r/3} |X-Y|^{1-n} \, |\nabla u_0(Y)| \, dY \\ &\lesssim \, \varepsilon \sum_{k=1}^{\infty} 2^{-k} \, \iint_{2^{-k-1}r \leq |X-Y| < 2^{-k}r} (2^{-k}r)^{-n} \, |\nabla u_0(Y)| \, dY \, =: \, \varepsilon \sum_{k=1}^{\infty} \, 2^{-k} I_k \, . \end{split}$$

By the inequalities of Cauchy-Schwarz, Caccioppoli, and Harnack, $I_k \leq u_0(X)$, and summing the geometric series, we then find that

$$|I| \leq \varepsilon u_0(X)$$

We consider now term *II*. For notational convenience, let Ω^X denote the domain of integration in term *II*, i.e.,

$$\Omega^X := \{Y \in \Omega : |X - Y| \ge \delta(Y)/4\},\$$

and set

$$\Omega_0 := \{ Y \in \Omega : |X_0 - Y| \ge \delta(X_0)/4 \}, \quad \Omega^{X, X_0} := \Omega^X \cap \Omega_0.$$

We then make the further splitting

$$II = \iint_{\Omega^{X,X_0}} \dots \, dY + \iint_{\Omega^X \cap \{|X_0 - Y| < \delta(X_0)/4\}} \dots \, dY =: II_1 + II_2 \, .$$

Observe that $\delta(Y) \approx \delta(X_0)$ in term II_2 , and thus also $|X - Y| \gtrsim \delta(X_0)$. Hence, by Lemma 3.34 and (4.2),

$$\begin{split} |H_2| \, \lesssim \, \varepsilon \delta(X_0)^{-n} \iint_{|X_0 - Y| < \delta(X_0)/4} |\nabla u_0(Y)| \, dY \\ \\ \lesssim \varepsilon \left(\delta(X_0)^{-n-1} \iint_{|X_0 - Y| < \delta(X_0)/2} |u_0(Y)|^2 \, dY \right)^{1/2} \, \lesssim \, \varepsilon u_0(X_0) \lesssim \varepsilon \,, \end{split}$$

where in the last three steps we have used the inequalities of Cauchy-Schwarz and Caccioppoli, then Harnack, and finally (4.14).

29

By Lemma 3.45, setting $\mathcal{G}_0(Y) := G_0(X_0, Y)$, we see that

$$|II_{1}| \lesssim \iint_{\Omega^{X,X_{0}}} \frac{\omega^{X}(\Delta_{Y})}{\omega_{0}(\Delta_{Y})} |\mathbf{B}(Y)| |\nabla u_{0}(Y)| \mathcal{G}_{0}(Y) dY$$

$$\lesssim \left(\iint_{\Omega} H^{X}(Y)^{2} |\mathbf{B}(Y)|^{2} \mathcal{G}_{0}(Y) dY\right)^{1/2} \left(\iint_{\Omega_{0}} |\nabla u_{0}(Y)|^{2} \mathcal{G}_{0}(Y) dY\right)^{1/2}$$

$$=: \mathbf{A} \cdot \mathbf{B}.$$

where $H^X(Y) := \frac{\omega^X(\Delta_Y)}{\omega_0(\Delta_Y)}$. Recalling Definition 2.14, we observe that for $z \in \partial \Omega$,

 $N_*H^X(z) \approx \mathcal{M}_{\omega_0}h^X(z)$.

Then by (4.3) and Carleson's embedding lemma (with respect to the doubling measure ω_0),

$$\mathbf{A} \lesssim \varepsilon \|\mathcal{M}_{\omega_0} h^X\|_{L^2(\omega_0)} \lesssim \varepsilon \|h^X\|_{L^2(\omega_0)}.$$

Moreover, since $u_0 = f$ on $\partial \Omega$, we have as in [DJK] that

$$\mathbf{B} \lesssim \|f\|_{L^2(\omega_0)} \le 1$$

Thus, $|II_1| \leq \varepsilon ||h^X||_{L^2(\omega_0)}$, so adding together our estimates for *I*, *II*₁ and *II*₂, we see that (4.15) holds. In turn, as noted above, this proves the claim (4.10), and therefore also Theorem 4.8.

5. Two key lemmas

In this section, we prove two lemmas that are central to our proof of Theorem 1.7. The first is a stopping-time lemma that plays a key role in the proof of Theorem 1.7 in Section 6, specifically, in our use of the method of "extrapolation of Carleson measures", a bootstrapping technique to lift the Carleson measure constant, developed by J. L. Lewis [LM], and based on the corona construction of Carleson [Car] and Carleson and Garnett [CG]. This technique was also used in [HL] (see in addition [AHLT], [AHMTT], [HM1], [HM2]).

We shall require some additional notation. Throughout this section, M is a fixed constant, eventually to be chosen depending only on allowable parameters. Given $M \ge 2$, and a cube $Q \in \mathbb{D}(\partial\Omega)$, we set

$$\mathbb{D}^{M}(Q) := \left\{ Q' : \frac{1}{M} \le \frac{\ell(Q')}{\ell(Q)} \le M \text{ and } \operatorname{dist}(Q, Q') \le M \operatorname{diam}(Q) \right\} ,$$

and define the *M*-fattened Whitney region

(5.1)
$$U_Q^M = \bigcup_{Q' \in \mathbb{D}^M(Q)} U_{Q'} \,.$$

Note that the fattened Whitney regions enjoy the bounded overlap property

$$\sum_{Q} 1_{U_{Q}^{M}}(Y) \lesssim_{M} 1, \qquad Y \in \Omega$$

Recall that $\mathbb{D}_Q := \{Q' \in \mathbb{D}(\partial \Omega) : Q' \subset Q\}$. For $x \in \partial \Omega$, define the "dyadic cone with aperture *M*, truncated at height $\ell(Q)$ " by

(5.2)
$$\Upsilon^M_Q(x) := \bigcup_{Q' \in \mathbb{D}_Q: x \in Q'} U^M_{Q'}.$$

We further define a "truncated dyadic-conical square function" by

(5.3)
$$\mathcal{H}_{Q}^{M}(\mathbf{F})^{2}(x) := \sum_{Q' \in \mathbb{D}_{Q}} \mathbb{1}_{Q'}(x) \iint_{U_{Q'}^{M}} |\mathbf{F}(Y)|^{2} \delta(Y)^{1-n} dY,$$

and its "shortened" version

(5.4)
$$\mathcal{A}_{Q}^{M,short}(\mathbf{F})^{2}(x) := \sum_{\mathcal{Q}' \in \mathbb{D}_{Q} \setminus \{\mathcal{Q}\}} \mathbb{1}_{\mathcal{Q}'}(x) \iint_{\mathcal{Q}'} |\mathbf{F}(Y)|^{2} \delta(Y)^{1-n} \, dY \,,$$

Set

(5.5)
$$\mathfrak{A}_{Q}^{M}(\mathbf{F})^{2}(x) := \iint_{\Upsilon_{Q}^{M}(x)} |\mathbf{F}(Y)|^{2} \delta(Y)^{1-n} dY,$$

and observe that by the bounded overlap property,

(5.6)
$$\mathcal{R}_Q^M(\mathbf{F})(x) \approx_M \mathfrak{A}_Q^M(\mathbf{F})(x), \qquad x \in \partial \Omega,$$

(of course, the direction $\mathfrak{A}_Q^M(\mathbf{F})(x) \le \mathcal{A}_Q^M(\mathbf{F})(x)$ is trivial) where

Given a tree S with maximal element Q(S) = Q, we also define the dyadic square function restricted to the tree by

(5.7)
$$\mathcal{A}^{M}_{\mathbf{S}}(\mathbf{F})^{2}(x) := \sum_{\mathcal{Q}' \in \mathbf{S}} \mathbb{1}_{\mathcal{Q}'}(x) \iint_{U^{M}_{\mathcal{Q}'}} |\mathbf{F}(Y)|^{2} \delta(Y)^{1-n} dY,$$

Let x_Q denote the "center" of Q as in (2.5), and set $B_Q^M := B(x_Q, CM \operatorname{diam}(Q))$, with C large enough that $\bigcup_{Q' \in \mathbb{D}_Q} U_{Q'}^M \subset B_Q^M$. Then in the special case that $\mathbf{F} = \mathbf{B}_*$, we note for future reference that by definition,

$$(5.8) \quad \int_{Q} \mathcal{A}_{Q}^{M}(\mathbf{B}_{*})^{2} d\sigma = \int_{Q} \sum_{Q' \in \mathbb{D}_{Q}} \mathbb{1}_{Q'}(x) \iint_{U_{Q'}^{M}} |\mathbf{B}_{*}(Y)|^{2} \delta(Y)^{1-n} dY$$
$$\approx_{M} \frac{1}{\sigma(Q)} \sum_{Q' \in \mathbb{D}_{Q}} \iint_{U_{Q'}^{M}} |\mathbf{B}_{*}(Y)|^{2} \delta(Y) dY$$
$$\lesssim_{M} \frac{1}{\sigma(Q)} \iint_{B_{Q}^{M} \cap \Omega} |\mathbf{B}_{*}(Y)|^{2} \delta(Y) dY \leq CM_{1} =: M_{2},$$

uniformly for every $Q \in \mathbb{D}(\partial \Omega)$, where M_1 is the constant in (1.5), and C depends on M and the various underlying constants.

We turn now to the stopping time lemma.

Lemma 5.9. Fix
$$M \ge 2$$
. Let $Q \in \mathbb{D}(\partial \Omega)$, and consider constants $a \ge 0$, $b > 0$. Suppose that

(5.10)
$$\int_{Q} \mathcal{R}_{Q}^{M}(\mathbf{B}_{*})^{2} d\sigma \leq (a+b),$$

then either

(5.11)
$$\int_{Q} \mathcal{R}_{Q}^{M,short}(\mathbf{B}_{*})^{2} d\sigma \leq a,$$

or, there is a tree **S** with maximal cube $Q(\mathbf{S}) = Q$ such that

(5.12)
$$\mathcal{A}^M_{\mathbf{S}}(\mathbf{B}_*)^2(x) \le 2b, \qquad \sigma \text{ - a.e. } x \in Q,$$

(5.13)
$$\sum_{\mathcal{F}_{bad}} \sigma(Q_j) \le \frac{a+b}{a+2b} \sigma(Q),$$

where $\mathcal{F}_{bad} := \{Q_j \in \mathcal{F} : \int_{Q_j} \mathcal{A}_{Q_j}^{M,short}(\mathbf{B}_*)^2 d\sigma > a\}$, and $\mathcal{F} = \{Q_j\}_j$ denotes the collection of sub-cubes of $Q = Q(\mathbf{S})$ that are maximal with respect to the property that $Q_j \notin \mathbf{S}$.

Proof. For convenience of notation, we set

$$\mathcal{H}_Q^M := \mathcal{R}_Q^M(\mathbf{B}_*)^2, \qquad \mathcal{H}_Q^{M,short} := \mathcal{R}_Q^{M,short}(\mathbf{B}_*)^2,$$

and similarly with Q replaced by any other cube $Q' \in \mathbb{D}_Q$. We also set

$$\xi_{\mathcal{Q}'} := \iint_{U_{\mathcal{Q}'}^M} |\mathbf{B}_*(Y)|^2 \delta(Y)^{1-n} \, dY \,, \quad \mathcal{Q}' \in \mathbb{D}_{\mathcal{Q}}$$

so that, e.g., $\mathcal{H}_Q^M(x) = \sum_{Q' \in \mathbb{D}_Q} 1_{Q'}(x) \xi_{Q'}$. Note in particular that

(5.14)
$$\mathcal{H}_Q^M(x) = \mathcal{H}_Q^{M,short}(x) + \xi_Q, \qquad \forall x \in Q.$$

For $Q'' \subset Q$, we also set

$$\mathcal{H}^M_{\mathcal{Q}'',\mathcal{Q}}(x):=\sum_{\mathcal{Q}':\mathcal{Q}''\subset\mathcal{Q}'\subset\mathcal{Q}}\mathbf{1}_{\mathcal{Q}'}(x)\,\xi_{\mathcal{Q}'}\,,$$

so that, in particular, $\mathcal{H}_{Q,Q}^{M}(x) = \xi_Q$ for $x \in Q$. We consider two cases. **Case 1**: $\xi_Q \ge b$. In this case, (5.10) and (5.14) imply that

$$\int_{Q} H_{Q}^{M,short} d\sigma \leq a \,,$$

i.e., (5.11) holds.

Case 2: $\xi_Q < b$. In this case, we subdivide dyadically, to select a family $\mathcal{F} := \{Q_j\}_j \subset \mathbb{D}_Q$ of cubes that are maximal with respect to the property that

(5.15)
$$\sum_{Q':Q_j \subset Q' \subset Q} \xi_{Q'} = \int_{Q_j} \mathcal{H}^M_{Q_j,Q} \, d\sigma > 2b$$

Set $\mathbf{S} := \{Q' \in \mathbb{D}_Q : Q' \text{ is not contained in any } Q_j \in \mathcal{F}\}$. If $x \in Q \setminus \bigcup_{\mathcal{F}} Q_j$, then

(5.16)
$$\int_{Q''} \mathcal{H}_{Q'',Q}^M d\sigma \le 2b, \quad \forall Q'' \ni x.$$

so letting $Q'' \downarrow x$, we see that (5.12) holds for a.e. such x.

On the other hand, suppose now that $x \in Q_j$ for some $Q_j \in \mathcal{F}$. In this case, by definition (5.7),

(5.17)
$$\mathcal{A}_{\mathbf{S}}^{M}(\mathbf{B}_{*})^{2}(x) = \mathcal{H}_{Q_{j}^{*},Q}^{M}(x) = \sum_{\mathcal{Q}': \mathcal{Q}_{j}^{*} \subset \mathcal{Q}' \subset \mathcal{Q}} \xi_{\mathcal{Q}'}, \quad \forall x \in Q_{j},$$

where Q_j^* denotes the dyadic parent of Q_j . Moreover, $\mathcal{H}_{Q_j^*,Q}^M(x)$ is constant on Q_j^* (indeed, $\mathcal{H}_{Q_j^*,Q}^M(x) = \sum_{Q':Q_j^* \subset Q' \subset Q} \xi_{Q'}$ on Q_j^*), and therefore

$$\mathcal{H}^{M}_{\mathcal{Q}^{*}_{j},\mathcal{Q}}(x) = \int_{\mathcal{Q}^{*}_{j}} \mathcal{H}^{M}_{\mathcal{Q}^{*}_{j},\mathcal{Q}} \, d\sigma \leq 2b \,,$$

by maximality of Q_j , for all $x \in Q_j^*$, in particular, for all $x \in Q_j$. Combining the latter estimate with the identity in (5.17), we obtain (5.12).

31

It remains to verify (5.13). To this end, note that by the definition of \mathcal{F}_{bad} , along with the stopping time criterion (5.15), we see that for each $Q_j \in \mathcal{F}_{bad}$,

$$\begin{aligned} (a+2b)\sigma(Q_j) &< \int_{Q_j} \left(\mathcal{A}_{Q_j}^{M,short}(\mathbf{B}_*)^2 + \mathcal{H}_{Q_j,Q}^M \right) d\sigma \\ &= \int_{Q_j} \left(\sum_{Q' \in \mathbb{D}_{Q_j} \setminus \{Q_j\}} \mathbf{1}_{Q'}(x) \xi_{Q'} + \sum_{Q': Q_j \subset Q' \subset Q} \mathbf{1}_{Q'}(x) \xi_{Q'} \right) d\sigma \\ &= \int_{Q_j} \mathcal{H}_Q^M d\sigma = \int_{Q_j} \mathcal{H}_Q^M (\mathbf{B}_*)^2 d\sigma \,. \end{aligned}$$

Summing over $Q_j \in \mathcal{F}_{bad}$, we obtain

$$(a+2b)\sum_{\mathcal{F}_{bad}}\sigma(\mathcal{Q}_j) \leq \sum_{\mathcal{F}}\int_{\mathcal{Q}_j}\mathcal{R}_Q^M(\mathbf{B}_*)^2\,d\sigma \leq \int_{\mathcal{Q}}\mathcal{R}_Q^M(\mathbf{B}_*)^2\,d\sigma \leq (a+b)\sigma(\mathcal{Q})\,,$$

by hypothesis, and (5.13) follows.

The next lemma provides control of a square function in a sawtooth region, in terms of a dyadic square function in the original domain. If Ω_S denotes the sawtooth domain, set

(5.18)
$$\Gamma_{\star}(z) := \{Y \in \Omega_{\mathbf{S}} : |z - Y| < 10\delta_{\star}(Y)\}, \quad z \in \partial \Omega_{\mathbf{S}}$$

where $\delta_{\star}(Y) := \text{dist}(Y, \partial \Omega_{S})$, and define the corresponding conical square function

(5.19)
$$\mathcal{A}_{\star}(\mathbf{B})^{2}(z) := \iint_{\Gamma_{\star}(z)} |\mathbf{B}(Y)|^{2} \,\delta_{\star}^{1-n}(Y) \,dY$$

Let M denote the parameter in the definition of the dyadic conical square function, as above.

Lemma 5.20. There is a choice of M large enough, depending only on the allowable parameters, such that for every $Q_0 \in \mathbb{D}(\partial\Omega)$, and for any tree **S** with top cube $Q(\mathbf{S}) = Q_0$, we have

(5.21)
$$\operatorname{ess\,sup}_{z\in\partial\Omega_{\mathbf{S}}}\mathcal{A}_{\star}(\mathbf{B})(z) \leq \|\mathcal{A}_{\mathbf{S}}^{M}(\mathbf{B})\|_{L^{\infty}(Q_{0})} + \operatorname{ess\,sup}_{Y\in\Omega_{\mathbf{S}}}\delta(Y)|\mathbf{B}(Y)|,$$

where the implicit constant also depends only on the allowable parameters, and where $\Omega_{\mathbf{S}} =$ int $(\bigcup_{Q \in \mathbf{S}} U_Q)$ is the usual "sawtooth domain" associated to \mathbf{S} .

Proof. We note that since $\Omega_{\mathbf{S}} \subset \Omega$, we have the trivial inequality

$$\delta_{\star}(Y) \le \delta(Y) \,,$$

which we shall use repeatedly without further explanation. We further note that by Proposition 2.26 (7),

$$\delta(Y) \le N\ell(Q_0), \qquad \forall Y \in R_{Q_0},$$

for some uniform constant N, hence in particular for all $Y \in \Omega_{\mathbf{S}}$.

Suppose first that $z \in \partial\Omega \cap \partial\Omega_{\mathbf{S}}$. Let $\mathcal{F} := \{Q_j\}_j$ denote the collection of sub-cubes of Q_0 that are maximal with respect to the property that $Q_j \notin \mathbf{S}$. Observe that $Q_0 \setminus (\cup_{\mathcal{F}} Q_j) \subset \partial\Omega \cap \partial\Omega_{\mathbf{S}}$, and $\partial\Omega \cap \partial\Omega_{\mathbf{S}} \setminus (Q_0 \setminus (\cup_{\mathcal{F}} Q_j))$ has \mathcal{H}^n measure zero, by Propositions 2.29 and 2.33. Thus, we may assume that $z \in Q_0 \setminus (\cup_{\mathcal{F}} Q_j)$. Let $Y \in \Gamma_{\star}(z)$ so that

(5.22)
$$\delta(Y) \le |z - Y| < 10\delta_{\star}(Y) \le 10\delta(Y)$$

(where in the first inequality we have used that $z \in \partial\Omega$). Choose $Q \subset Q_0$, with $z \in Q$, and $\delta(Y) \leq N\ell(Q) < 2\delta(Y)$. Then $Y \in U_Q^M$ for M chosen large enough (depending on N). Since there is such a cube Q for each $Y \in \Gamma_{\star}(z)$, we see that $\Gamma_{\star}(z) \subset \Upsilon_{Q_0}^M(z)$. Moreover, $\delta_{\star}(Y) \approx \delta(Y)$, by (5.22). Consequently,

$$\mathcal{A}_{\star}(\mathbf{B})(z) \leq \mathfrak{A}_{Q_0}^M(\mathbf{B})(z) \leq \mathcal{A}_{\mathbf{S}}^M(\mathbf{B})(z),$$

as desired, since $z \in Q_0 \setminus (\cup_{\mathcal{F}} Q_j)$ implies that $Q \in \mathbf{S}$ for every Q with $z \in Q \subset Q_0$.

Next, suppose that $z \in \partial \Omega_{\mathbf{S}} \setminus \partial \Omega$. Then $z \in \partial \Omega_{\mathbf{S}} \cap \Omega$, so $\delta(z) > 0$. Fix $Z_1 \in \Omega_{\mathbf{S}}$ such that $|Z_1 - z| < \delta(z)/2000$. Then by definition of $\Omega_{\mathbf{S}}, Z_1 \in U_{Q_1}$ for some $Q_1 \in \mathbf{S}$, hence

(5.23)
$$\ell(Q_1) \approx \delta(Z_1) \approx \delta(z) \approx \operatorname{dist}(z, Q_1).$$

Let $Y \in \Gamma_{\star}(z)$. We consider three cases. Let us first summarize some geometric information in each case, which we will then use to deduce (5.21).

Case 1: $|z - Y| < \delta(z)/2$. Then $\delta(Y) \approx \delta(z) \approx \ell(Q_1)$, by (5.23).

Case 2: $|z - Y| \ge \delta(z)/2$, and $\delta(Y) \le 4\delta(z)$. By definition of $\Gamma_{\star}(z)$, this means that

$$\frac{1}{8}\delta(Y) \le \frac{1}{2}\delta(z) \le |z - Y| < 10\delta_{\star}(Y) \le 10\delta(Y).$$

Thus, $\delta(Y) \approx \delta_{\star}(Y)$, and $|z - Y| \approx \delta(z) \approx \delta(Y)$, hence $Y \in U_{Q_1}^M$ for *M* large enough.

Case 3: $|z - Y| \ge \delta(z)/2$, and $\delta(Y) > 4\delta(z)$. In this case, $\delta(Y) < \frac{4}{3}|z - Y|$, and therefore

$$\delta(Y) \lesssim |z - Y| < 10\delta_{\star}(Y) \le 10\delta(Y) \,,$$

since $Y \in \Gamma_{\star}(z)$. In particular, $\delta(Y) \approx \delta_{\star}(Y)$. Choose Q such that $Q_1 \subset Q \subset Q_0$, with $\delta(Y) \leq N\ell(Q) < 2\delta(Y)$. Then using (5.23), we see that

 $\delta(Y) \leq \operatorname{dist}(Y, Q) \leq \operatorname{dist}(Y, Q_1) \leq |Y - z| + \operatorname{dist}(z, Q_1) + \operatorname{diam}(Q_1) \leq_N \delta(Y).$

Thus, $Y \in U_O^M$ for *M* chosen large enough.

We now write

$$\Gamma_{\star}(z) = \Gamma_1(z) \cup \Gamma_2(z) \cup \Gamma_3(z),$$

where Γ_1, Γ_2 and Γ_3 are the subsets of Γ_{\star} corresponding to Cases 1, 2 and 3, respectively. This induces a corresponding splitting

$$\mathcal{A}_{\star}(\mathbf{B})^{2}(z) = \sum_{i=1}^{3} \mathcal{A}_{i}(\mathbf{B})^{2}(z) := \sum_{i=1}^{3} \iint_{\Gamma_{i}(z)} |\mathbf{B}(Y)|^{2} \,\delta_{\star}^{1-n}(Y) \,dY \,.$$

Observe that for $Y \in \Gamma_2(z) \cup \Gamma_3(z)$ (thus, as in Case 2 or Case 3), we have shown that $\delta_{\star}(Y) \approx \delta(Y)$ and that $Y \in U_Q^M$ for some Q with $Q_1 \subset Q \subset Q_0$, where $Q_1 \in \mathbf{S}$. Consequently, for any $x \in Q_1$,

$$\mathcal{A}_{2}(\mathbf{B})^{2}(z) + \mathcal{A}_{3}(\mathbf{B})^{2}(z)$$

$$\lesssim \sum_{Q:Q_{1} \subset Q \subset Q_{0}} \iint_{U_{Q}^{M}} |\mathbf{B}(Y)|^{2} \,\delta^{1-n}(Y) \,dY \leq \mathcal{A}_{\mathbf{S}}^{M}(\mathbf{B})^{2}(x) \leq ||\mathcal{A}_{\mathbf{S}}^{M}(\mathbf{B})||_{L^{\infty}(Q_{0})}^{2},$$

as desired.

It remains to consider $\mathcal{A}_1(\mathbf{B})(z)$. Set $\Xi := \operatorname{ess\,sup}_{Y \in \Omega_S} \delta(Y)|\mathbf{B}(Y)|$. In Case 1, we have $|z - Y| < \delta(z)/2$, and $\delta(Y) \approx \delta(z) \approx \ell(Q_1)$. Moreover, for $Y \in \Gamma_{\star}(z)$, we have $|z - Y| \approx \delta_{\star}(Y)$. Therefore,

$$\begin{aligned} \mathcal{A}_{1}(\mathbf{B})^{2}(z) &= \iint_{\Gamma_{\star}(z) \cap \{|z-Y| < \delta(z)/2\}} |\mathbf{B}(Y)|^{2} \, \delta_{\star}^{1-n}(Y) \, dY \\ &\lesssim \frac{\Xi^{2}}{\ell(Q_{1})^{2}} \iint_{|z-Y| \leq \ell(Q_{1})} |z-Y|^{1-n} \, dY \, \lesssim \, \Xi^{2} \,, \end{aligned}$$

since our ambient dimension is n + 1. We conclude by summing our estimates for $\mathcal{R}_i(\mathbf{B})(z)$, i = 1, 2, 3.

In the following remark and corollary, we shall continue to use the same notation as in Lemma 5.20.

Remark 5.24. Fix a cube $Q_0 \in \mathbb{D}(\partial \Omega)$, let **S** be a tree with top cube $Q(\mathbf{S}) = Q_0$, and consider the associated sawtooth domain $\Omega_{\mathbf{S}}$ and Carleson region R_{Q_0} as in Proposition 2.26. By the properties of the Whitney regions U_Q , and of the sawtooth and Carleson regions, we may (and do) fix a point $X_0 \in U_{Q_0} \subset \Omega_{\mathbf{S}} \subset R_{Q_0}$, such that

$$\operatorname{dist}(X_0, \partial U_{Q_0}) \approx \delta_{\star}(X_0) \approx \operatorname{dist}(X_0, \partial R_{Q_0}) \approx \delta(X_0) \approx \ell(Q_0) \approx \operatorname{diam}(\Omega_{\mathbf{S}}) \approx \operatorname{diam}(R_{Q_0})$$

Thus, X_0 is a corkscrew point for Ω_S at the scale $r_0 := \operatorname{diam}(\Omega_S) \approx \ell(Q_0)$, with respect to any center $z \in \partial \Omega_S$.

With the point X_0 fixed, we let $\omega_{0,\star} := \omega_{L_0,\Omega_S}^{X_0}$ and $\mathcal{G}_{0,\star} := \mathcal{G}_{L_0,\Omega_S}(X_0,\cdot)$ denote the elliptic measure and Green function for the operator $L_0 = -\operatorname{div} A\nabla$, in the domain Ω_S , with pole at X_0 .

With *M* now fixed as in Lemma 5.20, we shall simply write $\mathcal{A}_{\mathbf{S}}(\mathbf{B}) := \mathcal{A}_{\mathbf{S}}^{M}(\mathbf{B})$.

Corollary 5.25. Let Q_0 , \mathbf{S} , and $\Omega_{\mathbf{S}}$ be as in Lemma 5.20, and suppose that \mathbf{B} is truncated in Ω , as in Subsection 3.5. Let X_0 be the corkscrew point fixed in Remark 5.24, and let $\omega_{0,\star}$ and $\mathcal{G}_{0,\star}$ be the elliptic measure and Green function as described above. Suppose that

(5.26)
$$\|\mathcal{A}_{\mathbf{S}}(\mathbf{B})\|_{L^{\infty}(Q_0)} + \operatorname{ess\,sup}_{Y \in \Omega_{\mathbf{S}}} \delta(Y)|\mathbf{B}(Y)| \le \beta$$

Then

(5.27)
$$|\mathbf{B}(Y)| \le \frac{\beta}{\delta_{\star}(Y)}, \quad \text{a.e. } Y \in \Omega_{\mathbf{S}},$$

and for every $\Delta_{\star} = B \cap \partial \Omega_{\mathbf{S}}$, with B = B(z, r), $z \in \partial \Omega_{\mathbf{S}}$, $0 < r < r_0$,

(5.28)
$$\iint_{B\cap\Omega_{\mathbf{S}}} |\mathbf{B}(Y)|^2 \mathcal{G}_{0,\star}(Y) \, dY \leq C\beta^2 \omega_{0,\star}(\Delta_{\star})$$

where $C \ge 1$ depends only on the allowable parameters; i.e., **B** is " $\sqrt{C\beta}$ -small with respect to $\omega_{0,\star}$ " in the sense of Definition 4.1, in the domain $\Omega_{\mathbf{S}}$.

Proof. Note that since **B** has been truncated in Ω as in Subsection 3.5, and hence is qualitatively bounded, then by construction, $\mathcal{A}_{\star}(\mathbf{B})$ is continuous on $\partial \Omega_{\mathbf{S}}$ (see (5.19)). Consequently, (5.21) self-improves to give

(5.29)
$$\sup_{z \in \partial \Omega_{\mathbf{S}}} \mathcal{A}_{\star}(\mathbf{B})(z) \leq \|\mathcal{A}_{\mathbf{S}}(\mathbf{B})\|_{L^{\infty}(Q_0)} + \operatorname{ess\,sup}_{Y \in \Omega_{\mathbf{S}}} \delta(Y)|\mathbf{B}(Y)| \leq \beta,$$

i.e., the essential supremum on the left hand side has now been replaced by a supremum; the upper bound on the right hand side is simply the hypothesis (5.26).

The bound (5.27) is trivial, since $\delta_{\star}(Y) \leq \delta(Y)$ for all $Y \in \Omega_{\mathbf{S}}$.

34

~ ~

To establish (5.28), we write

$$\iint_{B\cap\Omega_{\mathbf{S}}} |\mathbf{B}(Y)|^{2} \mathcal{G}_{0,\star}(Y) dY$$
$$= \iint_{B\cap\Omega_{\mathbf{S}}\cap\{|X_{0}-Y|>\delta(Y)/4\}} \dots dY + \iint_{B\cap\Omega_{\mathbf{S}}\cap\{|X_{0}-Y|\leq\delta(Y)/4\}} \dots dY =: I + II.$$

(It may be that *II* is vacuous, unless *B* has radius $r \approx r_0$). By standard Green function estimates in \mathbb{R}^{n+1} , and (5.26),

$$II \lesssim \frac{\beta^2}{r_0^2} \iint_{|X_0 - Y| \le r_0} |X_0 - Y|^{1 - n} dY \lesssim \beta^2,$$

since $r_0 := \operatorname{diam}(\Omega_{\mathbf{S}}) \approx \delta(X_0) \approx \delta(Y)$ in the regime $|X_0 - Y| \leq \delta(Y)/4$. Furthermore, by Lemma 3.45 applied in $\Omega_{\mathbf{S}}$, setting $\Delta_{\star,Y} := B(Y, 10\delta_{\star}(Y)) \cap \partial\Omega_{\mathbf{S}}$, and using (5.18) and (5.19), we see that

$$\begin{split} I &\lesssim \iint_{B \cap \Omega_{\mathbf{S}}} |\mathbf{B}(Y)|^{2} \,\omega_{0,\star}(\Delta_{\star,Y}) \,\delta_{*}^{1-n}(Y) \,dY \\ &= \int_{\partial \Omega_{\mathbf{S}}} \iint_{\Omega_{\mathbf{S}} \cap \{|z-Y| < 10\delta_{\star}(Y)\}} |\mathbf{B}(Y)|^{2} \,\delta_{*}^{1-n}(Y) \,dY \,d\omega_{0,\star}(z) \\ &= \int_{\partial \Omega_{\mathbf{S}}} \mathcal{A}_{\star}(\mathbf{B})^{2}(z) \,d\omega_{0,\star}(z) \,\lesssim \,\beta^{2} \,, \end{split}$$

by (5.29), since $\omega_{0,\star}$ is a probability measure.

6. Proof of Theorem 1.7: The extrapolation argument

We now arrive at the heart of the matter. In this section, we give the proof of Theorem 1.7. As mentioned above in Section 5, our approach will be based on the method of "extrapolation of Carleson measures".

We first require some preliminaries. For $X \in \Omega$, let ω^X denote the elliptic measure at X for the operator $L = -\operatorname{div} A \nabla + \mathbf{B} \cdot \nabla$ in Ω .

An important tool for us will be the following result of [BL], which provides a (necessary and sufficient) criterion for elliptic measure to belong to weak- A_{∞} . We recall the notation of (3.44): for $X \in \Omega$, we set $\Delta_X := B(X, 10\delta(X)) \cap \partial\Omega$.

Lemma 6.1 ([BL]). Let $L := -\operatorname{div} A\nabla + \mathbf{B} \cdot \nabla$, where **B** satisfies (1.2), and suppose that the continuous Dirichlet problem (Definition 2.18) is solvable for L, and that the solutions satisfy the weak maximum principle (thus, elliptic measure exists). For $X \in \Omega$, define Δ_X as in (3.44). Assume that $\partial\Omega$ is ADR, and that there are uniform positive constants η and γ such that for each $X \in \Omega$ with $\delta(X) < 2 \operatorname{diam}(\partial\Omega)$, the following implication holds for the Borel subsets of Δ_X :

(6.2)
$$E \subset \Delta_X, \quad \sigma(E) \ge (1 - \eta)\sigma(\Delta_X) \implies \omega^X(E) \ge \gamma.$$

Then elliptic measure for L is locally in weak- A_{∞} in the sense of Definition 2.20, with uniform constants depending only on η , γ , n, λ , Λ , ADR, and the constant in (1.2).

Remark 6.3. Clearly, the assumption $\delta(X) < 2 \operatorname{diam}(\partial \Omega)$ imposes a meaningful restriction only in the case that Ω is unbounded and $\partial \Omega$ is bounded.

Although not stated explicitly in this form, Lemma 6.1 is proved in [BL]: it follows from the combination of [BL, Lemma 2.2] and its proof, and [BL, Lemma 3.1]. The results in [BL] are stated only for the case that L is the Laplacian, but in fact those results may be applied much more generally: the proof in [BL] requires only Harnack's inequality and the weak maximum principle. In our setting, we recall that Harnack's inequality is available to us, by Lemma 3.3, and that the weak maximum principle, solvability of the continuous Dirichlet problem, and existence of elliptic measure are also available, by Lemma 3.31, if we truncate **B** near the boundary as in Subsection 3.5.

Thus, we consider truncations \mathbf{B}_k and the corresponding operators L_k as in Subsection 3.5. We then show that (6.2) holds, with the constants η and γ uniform in k (that is, independent of the truncation), depending only on allowable parameters. In particular, taking $E = \Delta_X$ in (6.2), we see that $\omega_k^X(\Delta_X) \ge \gamma$, uniformly in k, and therefore for any $x \in \partial\Omega$, $r < \operatorname{diam} \partial\Omega$, and for all $Y \in \Omega \cap B(x, r/2)$, we also have $\omega^Y(\Delta(x, r)) \ge c\gamma$. Thus, Lemma 3.11 holds for L_k , uniformly in k, by a standard iteration argument. We may therefore apply Lemma 3.32 to obtain that ω^X , the elliptic measure for L exists, and also satisfies (6.2). We then deduce from Lemma 6.1 that the weak- A_∞ property holds for the elliptic measure ω , as desired.

In the sequel, we henceforth implicitly assume that **B** has been truncated near the boundary, but we shall suppress the subscript k, for simplicity of notation. All of our quantitative bounds will depend only on allowable parameters, and then we may eventually pass to the limit as described above.

By Lemma 6.1, it suffices to show that there are uniform positive constants η and γ such that for each $X \in \Omega$ with $\delta(X) < 2 \operatorname{diam}(\partial \Omega)$ (see Remark 6.3), and for Δ_X as in (3.44), we have the implication (6.2) for any Borel subset $E \subset \Delta_X$.

We begin by reducing matters to a dyadic version of (6.2). To this end, let $X \in \Omega$, and let $\hat{x} \in \partial \Omega$ be a touching point for X, i.e., $|X - \hat{x}| = \delta(X)$. Choose a cube $Q \in \mathbb{D}(\partial \Omega)$, with $\hat{x} \in Q$, and such that $\ell(Q) \approx \delta(X)$ and $Q \subset \Delta_X$. By ADR, $\sigma(\Delta_X) \approx \sigma(Q)$, so matters are reduced to verifying that there are uniform positive constants η and γ such that for Borel subsets of any cube $Q \in \mathbb{D}(\partial \Omega)$,

(6.4)
$$E \subset Q, \quad \sigma(E) \ge (1 - \eta)\sigma(Q) \implies \omega^{X}(E) \ge \gamma.$$

Indeed, (6.4) for some η and γ , with $Q \subset \Delta_X$ as above, implies (6.2) for a somewhat smaller η (on the order of $c\eta$, with c = c(n, ADR)), and for the same γ .

Remark 6.5. In the construction in [HM2] of the Whitney regions U_Q (see Proposition 2.26), one may choose the associated parameters in such a way that for any $X \in \Omega$, and for $Q \subset \Delta_X$ with $\ell(Q) \approx \delta(X)$, as in the preceding discussion, we have $X \in U_Q$.

Set $\mathcal{A}_Q(\cdot) := \mathcal{A}_Q^M(\cdot)$, where the latter is defined in (5.3), and *M* has been fixed as in Lemma 5.20. Set $\Theta(Q) := \int_Q \mathcal{A}_Q(\mathbf{B}_*)^2 d\sigma$, and recall that by (5.8),

(6.6)
$$\sup_{Q\in\mathbb{D}}\Theta(Q) := \sup_{Q\in\mathbb{D}} \oint_{Q} \mathcal{A}_{Q}(\mathbf{B}_{*})^{2} d\sigma \leq M_{2},$$

where $M_2 = CM_1$, with M_1 as in (1.5), and with C depending only on allowable parameters.

The proof will proceed by induction. The inductive parameter will be a number $a \in [0, M_2)$.

 $\begin{array}{c|c} \hline H[a] \end{array} \begin{array}{l} There \ exist \ positive \ numbers \ \gamma_a \ and \ \eta_a \ such \ that \ for \ any \ given \ Q \in \\ \mathbb{D}(\partial\Omega), \ if \end{array} \\ \hline (6.7) \qquad \Theta(Q) \ \leq \ a\sigma(Q), \\ and \ if \ E \subset Q \ is \ a \ Borel \ set, \ then \\ (6.8) \qquad \sigma(E) \geq (1 - \eta_a)\sigma(Q) \implies \inf_{X \in U_Q} \omega^X(E) \geq \gamma_a \,, \end{array}$

Our proof strategy is as follows. We first verify H[0] (this will be easy), and then show that there is a uniform number b > 0 such that $H[a] \Longrightarrow H[a + b]$. After iterating this procedure on the order of $M_1/b \approx M_2/b$ times, we then find that $H[M_2]$ holds, hence by (6.6), we see that the conclusion (6.4) holds for all $Q \in \mathbb{D}(\partial\Omega)$ with $\eta = \eta_{M_2}$, and with $\gamma = \gamma_{M_2}$, for all $X \in U_Q$. Consequently, as observed above, using Remark 6.5, we find that (6.2) holds for all $X \in \Omega$ (for a somewhat smaller, but still fixed and uniform value of η). In turn, Lemma 6.1 then yields the desired weak- A_{∞} property.

Thus, it remains to verify H[0], and that $H[a] \Longrightarrow H[a + b]$, for each $a \in [0, M_2)$, and for some uniform b > 0.

Proof of H[0]. Fix Q_0 such that $\Theta(Q_0) = 0$. Then by definition, $\mathbf{B}_* \equiv 0$ in the "Carleson box" R_{Q_0} Consequently, $L := L_0$ in R_{Q_0} . Recall (see Proposition 2.26 (6)), that there is a ball $B_{Q_0} = (x_{Q_0}, r_{Q_0})$, with $r_{Q_0} \approx \ell(Q_0)$, such that $2B_{Q_0} \cap \Omega \subset R_{Q_0}$, where as usual $2B_{Q_0}$ denotes the double of B_{Q_0} . Let X_0 be the corkscrew point fixed in Remark 5.24, and let $\omega_{L_0}^{X_0}$ and $\tilde{\omega}_{L_0}^{X_0}$ denote elliptic measure at the point X_0 for L_0 , in the domains Ω and R_{Q_0} respectively. Similarly, let G_{L_0} denote the Green function for L_0 in Ω , and let \tilde{G}_{L_0} denote the Green function for L_0 in the domain R_{Q_0} . By the usual comparison principle, aka "boundary Harnack principle" (i.e., Lemma 3.49), along with Harnack's inequality and the Harnack Chain property,

$$G_{L_0}(X_0, Y) \approx \widetilde{G}_{L_0}(X_0, Y), \quad \forall Y \in \frac{3}{2}B_{Q_0} \cap \Omega.$$

Hence, for any Borel set $F \subset \Delta_{Q_0} := B_{Q_0} \cap \partial \Omega$, by the CFMS estimates for L_0 (Lemma 3.45) we have

(6.9)
$$\omega_{L_0}^{X_0}(F) \approx \widetilde{\omega}_{L_0}^{X_0}(F) \,.$$

Let $E \subset Q_0$ with $\sigma(E) > (1 - \eta_0)\sigma(Q_0)$. If we choose η_0 small enough, then by the A_{∞} property for ω_{L_0} and Lemma 3.28 (and Harnack's inequality and the Harnack Chain condition), using (6.9) we see that

$$\widetilde{\omega}_{L_0}^{X_0}(E \cap \Delta_{Q_0}) \approx \omega_{L_0}^{X_0}(E \cap \Delta_{Q_0}) \gtrsim \omega_{L_0}^{X_0}(Q_0) \gtrsim 1.$$

Let ω_L denote elliptic measure for L in Ω . Since $L = L_0$ in R_{Q_0} , by Harnack's inequality, the Harnack Chain condition, and the maximum principle we have

$$\inf_{X \in U_{\mathcal{Q}_0}} \omega_L^X(E) \gtrsim \omega_L^{X_0}(E) \ge \widetilde{\omega}_{L_0}^{X_0}(E \cap \Delta_{\mathcal{Q}_0}) \gtrsim 1.$$

Thus, H[0] holds.

The next (and main) step is to show that H[a] implies H[a + b], for a sufficiently small (but uniform) choice of b > 0.

 $H[a] \Longrightarrow H[a + b]$. Let $a \ge 0$, let b be a sufficiently small positive number to be chosen, and consider a cube Q_0 such that

(6.10)
$$\Theta(Q_0) \le (a+b)\,\sigma(Q_0)\,,$$

with $\Theta(Q)$ defined as in (6.6). We apply Lemma 5.9 to Q_0 . Then either (5.11) holds, or both (5.12) and (5.13) hold. The former case is easy to handle.

Indeed, if (5.11) holds, then by definition of \mathcal{A}_Q^{short} (recall that we have suppressed the parameter M, which has now been fixed), there is at least one dyadic child of Q, call it Q', such that $\Theta(Q') \leq a$. Consequently, we may apply the induction hypothesis H[a] to Q'. Suppose that $E \subset Q$, with $\sigma(E) \geq (1 - \eta)\sigma(Q)$, where $\eta > 0$ is chosen small enough, depending only on dimension and ADR, so that $\sigma(E \cap Q') \geq (1 - \eta_a)\sigma(Q')$. Applying H[a] in Q', and using Harnack's inequality, we see that

$$\inf_{X\in U_Q}\omega^X(E)\gtrsim \inf_{X\in U_{Q'}}\omega^X(E\cap Q')\geq \gamma_a\,,$$

thus, we obtain H[a + b] in the present case with $\eta_{a+b} = \eta$, and $\gamma_{a+b} = c\gamma_a$, for some c > 0 depending only on allowable parameters.

In order to treat the main case that both (5.12) and (5.13) hold, we shall need to discuss some preliminary matters.

Recall (see Lemma 5.9) that $\mathcal{F} = \{Q_j\}_j$ is the family of sub-cubes of Q_0 that are maximal with respect to the property that $Q_j \notin S$. As in [HM2], we define a projection operator $\mathcal{P}_{\mathcal{F}}$ with respect to the family \mathcal{F} as follows: given a Borel measure m defined on $\partial\Omega$, and a Borel subset $F \subset \partial\Omega$, we set

(6.11)
$$\mathcal{P}_{\mathcal{F}}\mathfrak{m}(F) := \mathfrak{m}\left(F \setminus (\cup_{\mathcal{F}} Q_j)\right) + \sum_{\mathcal{F}} \frac{\sigma(F \cap Q_j)}{\sigma(Q_j)} \mathfrak{m}(Q_j).$$

We note that in particular, for any dyadic cube Q that is *not* contained in any $Q_j \in \mathcal{F}$, we have

(6.12)
$$\mathcal{P}_{\mathcal{F}}\mathfrak{m}(Q) = \mathfrak{m}(Q), \quad Q \not\subset Q_i, \text{ for any } Q_i \in \mathcal{F}.$$

Lemma 6.13. (Essentially, [HM2, Lemma 6.15]) (Dyadic sawtooth lemma for projections). Suppose that Ω is a 1-sided chord-arc domain, and let $L_0 := -\operatorname{div} A\nabla$ be uniformly elliptic in Ω . Fix $Q_0 \in \mathbb{D}(\partial\Omega)$, let $\mathcal{F} = \{Q_j\} \subset \mathbb{D}_{Q_0}$ be a family of pairwise disjoint dyadic cubes that are maximal with respect to the property that they are do not belong to a stopping time tree **S** with top cube $Q(\mathbf{S}) = Q_0$. Let $\mathcal{P}_{\mathcal{F}}$ be the corresponding projection operator, and fix $X_0 \in U_{Q_0}$ as in Remark 5.24. Let $\omega_0 = \omega_0^{X_0}$ and $\omega_{0,\star} = \omega_{0,\star}^{X_0}$ denote the respective elliptic measures (for the operator L_0) for the domains Ω and $\Omega_{\mathbf{S}}$, with fixed pole at X_0 . Let $\nu = \nu^{X_0}$ be the measure on Q_0 defined by

(6.14)
$$\nu(F) = \omega_{0,\star} \left(F \setminus (\cup_{\mathcal{F}} Q_j) \right) + \sum_{Q_j \in \mathcal{F}} \frac{\omega_0(F \cap Q_j)}{\omega_0(Q_j)} \,\omega_{0,\star}(P_j), \qquad F \subset Q_0$$

where $P_j \subset \partial \Omega_s$ is the surface ball satisfying (2.35), whose existence is guaranteed by Proposition 2.34 (that is, by [HM2, Lemma 6.7]). Then

(6.15)
$$\mathcal{P}_{\mathcal{F}}\nu(F) = \omega_{0,\star} \left(F \setminus (\cup_{\mathcal{F}} Q_j) \right) + \sum_{Q_j \in \mathcal{F}} \frac{\sigma(F \cap Q_j)}{\sigma(Q_j)} \,\omega_{0,\star}(P_j), \qquad F \subset Q_0.$$

Moreover, there exists $\rho > 0$ *such that for all* $Q \in \mathbb{D}_{O_0}$ *and* $F \subset Q$ *, we have*

(6.16)
$$\left(\frac{\mathcal{P}_{\mathcal{F}}\omega_0(F)}{\mathcal{P}_{\mathcal{F}}\omega_0(Q)}\right)^{\rho} \lesssim \frac{\mathcal{P}_{\mathcal{F}}\nu(F)}{\mathcal{P}_{\mathcal{F}}\nu(Q)} \lesssim \frac{\mathcal{P}_{\mathcal{F}}\omega_0(F)}{\mathcal{P}_{\mathcal{F}}\omega_0(Q)}.$$

We remark that in (6.14) and in (6.15), we are using that $Q_0 \setminus (\bigcup_{\mathcal{F}} Q_j) \subset \partial \Omega \cap \partial \Omega_S$ (see Proposition 2.29), so that $\omega_{0,\star}(F \setminus (\bigcup_{\mathcal{F}} Q_j))$ makes sense.

The preceeding lemma is a dyadic version of the "Main Lemma" of [DJK], from which its proof is adapted. In the present form, it has essentially been proved: see [HM2, Lemma 6.15]. The latter result was stated only for the Laplacian, but the proof carries through essentially verbatim to the case stated above: one requires only that solutions to the operator $L_0u = 0$ satisfy the standard estimates in the Lemmata 3.11, 3.28, 3.34, 3.45, 3.48, 3.49, and 3.50. We mention that there is an extraneous qualitative hypothesis in the statement of [HM2, Lemma 6.15], namely that Ω has exterior corkscrew points in a *qualitative* sense (i.e., only at small scales, with no quantitative limitation on how small the scales might be): in [HM2], the lemma was applied in approximating domains which had the qualitative property, but the quantitative estimates were uniform for all the approximating domains, and did not depend on this qualitative assumption; in any case this hypothesis is not needed in the proof, as the interested reader may readily verify by consulting [HM2].

We now turn to the main case, in which both (5.12) and (5.13) hold. For S as in Lemma 5.9, by (5.12) and the definition (5.7) of $\mathcal{A}_{S}(\cdot) = \mathcal{A}_{S}^{M}(\cdot)$, we have that whenever $x \in Q \in S$,

$$\iint_{U_Q^M} |\mathbf{B}_*(Y)|^2 \delta(Y)^{1-n} \, dY \leq \mathcal{A}_{\mathbf{S}}(\mathbf{B}_*)^2(x) \leq 2b \,,$$

and thus also, by the trivial containment $U_Q \subset U_Q^M$,

(6.17)
$$\iint_{U_Q} |\mathbf{B}_*(Y)|^2 \delta(Y)^{1-n} \, dY \le 2b \,, \qquad \forall \, Q \in \mathbf{S} \,.$$

Since U_Q is a union of fattened Whitney cubes, we may construct a covering of U_Q by a collection *C* of (n + 1)-dimensional Euclidean cubes, each of side length $\ell(I) = \ell(Q)/N$, so that $U_Q = \bigcup_{I \in C} I$, and where we fix *N* large enough that $|Z - Y| < \delta(Y)/4$ for any pair of points *Y*, $Z \in I$; here, we have used that $\delta(Y) \approx \ell(Q)$, for all $Y \in U_Q$.

With this choice of N now fixed, using (6.17), we see that for each such I,

$$\frac{1}{|I|} \iint_{I} |\delta(Y)\mathbf{B}_{*}(Y)|^{2} dY \leq b$$

In particular, there is some fixed $Y_I \in I$ such that $\delta(Y_I)|\mathbf{B}_*(Y_I)| \leq \sqrt{b}$. By definition (1.4) of \mathbf{B}_* , we see that

$$\delta_{\star}(Y)|\mathbf{B}(Y)| \le \delta(Y)|\mathbf{B}(Y)| \le \sqrt{b}$$
, a.e. $Y \in I$,

hence for a.e. $Y \in U_Q$, and hence for a.e. $Y \in \Omega_S$, where as above, $\delta_{\star}(Y) := \operatorname{dist}(Y, \partial \Omega_S)$. Moreover, by (5.12) and Corollary 5.25, choosing $b = c\varepsilon^2$ for some small enough but fixed constant c > 0, we see that **B** is ε -small with respect to $\omega_{0,\star}$ in Ω_S , in the sense of Definition 4.1, where as above $\omega_{0,\star} := \omega_{L_0,\Omega_S}^{X_0}$ denotes the elliptic measure for the operator $L_0 = -\operatorname{div} A\nabla$, in the domain Ω_S , with pole at the fixed point $X_0 \in U_{Q_0}$ in Remark 5.24. Thus, by Theorem 4.8, applied in the domain Ω_S , with $\varepsilon \leq \varepsilon_0$, we find that

(6.18)
$$\omega_{L,\Omega_{\mathbf{S}}}^{X_0} =: \omega_{\star} \in A_{\infty}(\partial \Omega_{\mathbf{S}}, \omega_{0,\star}),$$

where $\omega_{L,\Omega_{\mathbf{S}}}^{X}$ is elliptic measure for $L = L_0 + \mathbf{B} \cdot \nabla$ in the domain $\Omega_{\mathbf{S}}$.

Remark 6.19. In agreement with our previous usage (as in (6.18) for example), we formalize some notation to distinguish elliptic measures in different domains, and for different operators. We let $\omega^X := \omega_{L,\Omega}^X$, and $\omega_{\star}^X := \omega_{L,\Omega_S}^X$ denote elliptic measure for *L* at the point *X*, in the domains Ω and Ω_S respectively. Similarly $\omega_0^X := \omega_{L_0,\Omega}^X$, and $\omega_{0,\star}^X := \omega_{L_0,\Omega_S}^X$ will denote elliptic measure for L_0 at the point *X*, in the domains Ω and Ω_S . In the special case that $X = X_0$, we simply write $\omega = \omega_{\star}^{X_0}$, $\omega_{\star} = \omega_{\star}^{X_0}$, $\omega_0 = \omega_0^{X_0}$ and $\omega_{0,\star} = \omega_{0,\star}^{X_0}$, where $X_0 \in U_{Q_0}$ has been fixed as in Remark 5.24.

Proceeding with the proof of Theorem 1.7, we recall that we are assuming that H[a] holds, and we seek to establish the conclusion of H[a + b], for the cube Q_0 satisfying (6.10), where as above, we have now fixed $b = c\varepsilon^2$ with $\varepsilon \le \varepsilon_0$.

Suppose then that $E \subset Q_0$, such that

(6.20)
$$\sigma(E) \ge (1 - \eta)\sigma(Q_0).$$

As above, let $X_0 \in U_{Q_0}$ be as in Remark 5.24. Our goal is to show that there is a uniform number $\gamma > 0$ such that

(6.21)
$$\omega(E) := \omega^{X_0}(E) \ge \gamma,$$

provided that $\eta = \eta_{a+b} > 0$ is small enough. By the Harnack chain condition and Harnack's inequality, (6.21) immediately implies the analogous estimate with X_0 replaced by an arbitrary $X \in U_{Q_0}$, and with γ replaced by $c\gamma =: \gamma_{a+b}$, for some uniform constant *c*, and H[a+b] will then follow.

We now turn to the proof of (6.21). Recall that we have applied Lemma 5.9 to Q_0 , so that, by (5.13), since $a < M_2$ (see (6.6)),

$$\sigma\left(\bigcup_{\mathcal{F}_{bad}} Q_j\right) \leq \frac{a+b}{a+2b} \,\sigma(Q_0) \leq \frac{M_2+b}{M_2+2b} \,\sigma(Q_0) =: (1-\theta) \,\sigma(Q_0) \,.$$

Consequently, either

Case 1:

(6.22)
$$\sigma\left(\cup_{\mathcal{F}_{good}} Q_j\right) \geq \frac{\theta}{2} \sigma(Q_0),$$

where $\mathcal{F}_{good} := \mathcal{F} \setminus \mathcal{F}_{bad}$, and $\theta = \theta(M_2, b)$ is a uniform positive number, or

Case 2:

(6.23)
$$\sigma\left(Q_0\setminus\left(\cup_{\mathcal{F}}Q_j\right)\right)\geq \frac{\theta}{2}\sigma(Q_0).$$

Proof of $H[a] \implies H[a + b]$ *in Case 1*. Suppose that (6.20) and (6.22) hold. Note that by definition,

$$\mathcal{F}_{good} := \{ Q_j \in \mathcal{F} : \ \int_{Q_j} \mathcal{A}_{Q_j}^{short} (\mathbf{B}_*)^2 d\sigma \le a \}.$$

Then for $\eta > 0$ small enough, by pigeon-holing and the definition of $\mathcal{A}_{Q_j}^{short}$ (see (5.4)), there is an "extra good" subset $\mathcal{F}_{eg} \subset \mathcal{F}_{good}$ such that there is a child Q'_j of each $Q_j \in \mathcal{F}_{eg}$, satisfying the following three estimates:

(6.24)
$$\sigma\left(\cup_{Q_j\in\mathcal{F}_{eg}}Q'_j\right)\gtrsim_{a,\theta,\eta}\sigma(Q_0).$$

(6.25)
$$\int_{\mathcal{Q}'_j} \mathcal{A}_{\mathcal{Q}'_j} (\mathbf{B}_*)^2 d\sigma \leq a,$$

and

(6.26)
$$\sigma(E \cap Q'_i) \ge (1 - \eta_a)\sigma(Q'_i).$$

Note that we may therefore apply H[a] to such Q'_i , to deduce that

(6.27)
$$\inf_{X \in U_{Q'_j}} \omega^X(E \cap Q'_j) \ge \gamma_a \,.$$

By a covering lemma argument, we may extract a further "extra good separated" subcollection $\mathcal{F}_{egs} \subset \mathcal{F}_{eg}$, such that for some sufficiently large constant $N_0 \gg 1$ to be chosen momentarily (depending as usual only on allowable parameters),

(6.28)
$$\operatorname{dist}(Q_{j_1}, Q_{j_2}) \ge N_0 \max(\operatorname{diam} Q_{j_1}, \operatorname{diam} Q_{j_2})$$

for distinct $Q_{j_1}, Q_{j_2} \in \mathcal{F}_{egs}$, and

$$(6.29) \quad \sum_{Q_j \in \mathcal{F}_{egs}} \sigma(Q_j) = \sigma\left(\cup_{Q_j \in \mathcal{F}_{egs}} Q_j\right) \ge \sum_{Q_j \in \mathcal{F}_{egs}} \sigma(Q'_j) = \sigma\left(\cup_{Q_j \in \mathcal{F}_{egs}} Q'_j\right) \gtrsim_{a,\theta,\eta,N_0} \sigma(Q_0),$$

where as above, Q'_{j} is a child of Q_{j} satisfying (6.25) and (6.26).

We recall the surface balls $P_j \subset \partial \Omega_s$ given in Proposition 2.34. Note that we may (and do) choose N_0 large enough, depending on the implicit constants in (2.35), such that for any distinct $Q_{j_1}, Q_{j_2} \in \mathcal{F}_{egs}$, we have

(6.30)
$$\operatorname{dist}(P_{j_1}, P_{j_2}) \ge \max(\operatorname{diam} Q_{j_1}, \operatorname{diam} Q_{j_2}).$$

We recall that $E \subset Q_0$ is a Borel set satisfying (6.20), and we remind the reader of the notation in Remark 6.19. Since $\Omega_S \subset \Omega$,

$$(6.31) \quad \omega(E) \geq \sum_{Q_j \in \mathcal{F}_{egs}} \omega(E \cap Q'_j)$$
$$= \sum_{Q_j \in \mathcal{F}_{egs}} \int_{\partial \Omega_{\mathbf{S}}} \omega^{Y}(E \cap Q'_j) \, d\omega_{\star}(Y) \geq \sum_{Q_j \in \mathcal{F}_{egs}} \int_{P_j} \omega^{Y}(E \cap Q'_j) \, d\omega_{\star}(Y, , \mathbb{C})$$

where in the last step we have used (6.30). Note that by (6.27), (2.35) and Harnack's inequality,

$$\omega^Y(E\cap Q'_j)\gtrsim \gamma_a\,.$$

Hence,

(6.32)
$$\omega(E) \gtrsim \sum_{Q_j \in \mathcal{F}_{egs}} \omega_{\star}(P_j) = \omega_{\star} \left(\bigcup_{Q_j \in \mathcal{F}_{egs}} P_j \right) =: \omega_{\star}(E_{\mathbf{S}}),$$

where $E_{\mathbf{S}} := \bigcup_{Q_j \in \mathcal{F}_{egs}} P_j$.

We have therefore reduced matters, in Case 1, to proving

$$(6.33) \qquad \qquad \omega_{\star}(E_{\mathbf{S}}) \gtrsim 1.$$

A remark is in order concerning the proof of (6.33). Even though **B** satisfies a small constant version of (1.5) in Ω_S , we cannot simply prove a "small constant" analogue of Theorem 1.7 in Ω_S (which would yield (6.33) directly) because we do not know that L_0 is "good" in Ω_S , i.e., it is not clear that $\omega_{0,\star}$ enjoys the A_{∞} property with respect to surface measure on $\partial \Omega_S$. By hypothesis, such a property holds in Ω , but it seems unlikely that it can be transferred, in general, to the subdomain Ω_S .

Instead, our strategy is to use Lemma 6.13, that is, the dyadic version of the sawtooth lemma (the "Main Lemma") of [DJK], but here we encounter another obstacle, namely that it is not clear that Lemma 6.13 may be applied directly to *L*. Thus, we do something a bit more indirect, inspired by ideas in [FKP]: we shall use (6.18) (which we had deduced from Theorem 4.8, our analogue of [FKP, Theorem 2.5]), to compare ω_{\star} to $\omega_{0,\star}$, and then we apply Lemma 6.13 to the operator L_0 , to compare ω_0 and $\omega_{0,\star}$.

Let us now proceed to establish (6.33). The first step is now easy. We view $\partial \Omega_S$ as a surface ball of radius $r = 2 \operatorname{diam}(\Omega_S) \approx \ell(Q_0)$, so that by (6.18), there is a uniform exponent $\alpha > 0$ such that

(6.34)
$$\omega_{\star}(E_{\mathbf{S}}) \gtrsim \left(\frac{\omega_{0,\star}(E_{\mathbf{S}})}{\omega_{0,\star}(\partial\Omega_{\mathbf{S}})}\right)^{\alpha} \omega_{\star}(\partial\Omega_{\mathbf{S}}).$$

Since ω_{\star} and $\omega_{0,\star}$ are probability measures on $\partial\Omega_{\rm S}$, it now suffices to prove that

$$(6.35) \qquad \qquad \omega_{0,\star}(E_{\mathbf{S}}) \gtrsim 1\,,$$

which we shall do, in turn, using Lemma 6.13.

To this end, define $\mathcal{P}_{\mathcal{F}}$ as in (6.11), let v be as in (6.14), so that $\mathcal{P}_{\mathcal{F}}v$ is given by (6.15). Set

(6.36)
$$E_1 := E \cap \left(\cup_{Q_i \in \mathcal{F}_{egs}} Q'_i \right) \,.$$

We then have that

$$\mathcal{P}_{\mathcal{F}} v(E_1) = \sum_{Q_j \in \mathcal{F}_{egs}} \frac{\sigma(E \cap Q'_j)}{\sigma(Q_j)} \, \omega_{0,\star}(P_j) \approx \sum_{Q_j \in \mathcal{F}_{egs}} \omega_{0,\star}(P_j) = \omega_{0,\star}(E_{\mathbf{S}}) \,,$$

where in the next-to-last step, we have used that, in particular, (6.26) holds for each $Q_j \in \mathcal{F}_{egs}$, while the last step is just the definition of E_s , along with the separation property (6.30). Thus, to prove (6.35), it is equivalent to show that

$$\mathcal{P}_{\mathcal{F}} v(E_1) \gtrsim 1.$$

To this end, we first note that by (6.16), applied with $Q = Q_0$,

$$\mathcal{P}_{\mathcal{F}}\nu(E_1) \gtrsim \left(\frac{\mathcal{P}_{\mathcal{F}}\omega_0(E_1)}{\mathcal{P}_{\mathcal{F}}\omega_0(Q_0)}\right)^{\rho} \mathcal{P}_{\mathcal{F}}\nu(Q_0).$$

It therefore suffices to verify the following pair of claims.

Claim 1:

(6.37)
$$\frac{\mathcal{P}_{\mathcal{F}}\omega_0(E_1)}{\mathcal{P}_{\mathcal{F}}\omega_0(Q_0)} \gtrsim 1,$$

and

Claim 2:

$$(6.38) \qquad \qquad \mathcal{P}_{\mathcal{F}}\nu(Q_0) \gtrsim 1\,,$$

where in each case the implicit constants depend only on allowable parameters.

Proof of Claim 1. By hypothesis, $\omega_0 \in A_{\infty}(\sigma, Q_0)$. By definition of E_1 (6.36), and of $\mathcal{P}_{\mathcal{F}}$ (6.11),

$$\mathcal{P}_{\mathcal{F}}\omega_0(E_1) = \sum_{Q_j \in \mathcal{F}_{egs}} \frac{\sigma(E \cap Q'_j)}{\sigma(Q_j)} \,\omega_0(Q_j)$$

42

$$\approx \sum_{Q_j \in \mathcal{F}_{egs}} \omega_0(Q_j) = \omega_0(\cup_{\mathcal{F}_{egs}} Q_j) \gtrsim \omega_0(Q_0) = \mathcal{P}_{\mathcal{F}} \omega_0(Q_0),$$

where in the second line of the display, we have used (6.26), then (6.29) and the A_{∞} property of ω_0 , and finally (6.12). Thus, Claim 1 (6.37) holds.

Proof of Claim 2. By the formula for $\mathcal{P}_{\mathcal{F}}v$ in (6.15),

(6.39)
$$\mathcal{P}_{\mathcal{F}} \mathcal{V}(Q_0) = \omega_{0,\star} \left(Q_0 \setminus (\cup_{\mathcal{F}} Q_j) \right) + \sum_{\mathcal{F}} \omega_{0,\star}(P_j)$$
$$\gtrsim \omega_{0,\star} \left(Q_0 \setminus (\cup_{\mathcal{F}} Q_j) \right) + \sum_{\mathcal{F}} \omega_{0,\star}(MP_j),$$

where in the last line we have used the doubling property of $\omega_{0,\star}$ to replace the surface ball P_j by the concentric dilate MP_j defined in Proposition 2.36, where M is the dilation factor in Corollary 2.38. Let Δ_{\star} be the surface ball on $\partial\Omega_S$ given by Proposition 2.36. Then by (6.39) and Corollary 2.38,

(6.40)
$$\mathcal{P}_{\mathcal{F}}\nu(Q_0) \gtrsim \omega_{0,\star}(\Delta_{\star}) \gtrsim 1,$$

where in the second inequality we have used Lemma 3.28, Harnack's inequality and the Harnack Chain property, all applied in Ω_S . Of course, (6.40) yields Claim 2 (6.38).

We have now completed the proof of the implication $H[a] \implies H[a+b]$ in the scenario of Case 1, i.e., (6.22).

Proof of H[*a*] \implies *H*[*a* + *b*] *in Case 2*. We now turn to Case 2 (6.23). Recall that *E* satisfies (6.20). Hence, for $\eta = \eta(\theta)$ small enough, (6.23) yields

(6.41)
$$\sigma(E') := \sigma\left(E \setminus \left(\cup_{\mathcal{F}} Q_j\right)\right) \ge \frac{\theta}{4} \sigma(Q_0),$$

where $E' := E \setminus (\cup_{\mathcal{F}} Q_j)$. Since $E' \subset \partial \Omega \cap \partial \Omega_S$, by the maximum principle and then (6.18), we find that for some uniform exponent $\alpha > 0$,

(6.42)
$$\omega(E) \ge \omega_{\star}(E') \gtrsim \left(\frac{\omega_{0,\star}(E')}{\omega_{0,\star}(\partial\Omega_{\mathbf{S}})}\right)^{\alpha} \omega_{\star}(\partial\Omega_{\mathbf{S}}),$$

Since ω_{\star} and $\omega_{0,\star}$ are probability measures on $\partial \Omega_{\rm S}$, it now suffices to prove that

(6.43)
$$\omega_{0,\star}(E') \gtrsim 1.$$

To this end, note that by definition of $\mathcal{P}_{\mathcal{F}} v$ (6.15), and then (6.16) applied with $Q = Q_0$,

$$\omega_{0,\star}(E') = \mathcal{P}_{\mathcal{F}}\nu(E') \gtrsim \left(\frac{\mathcal{P}_{\mathcal{F}}\omega_0(E')}{\mathcal{P}_{\mathcal{F}}\omega_0(Q_0)}\right)^{\rho} \mathcal{P}_{\mathcal{F}}\nu(Q_0).$$

By definition of $\mathcal{P}_{\mathcal{F}}$ (6.11), and (6.12), we have

$$\frac{\mathcal{P}_{\mathcal{F}}\omega_0(E')}{\mathcal{P}_{\mathcal{F}}\omega_0(Q_0)} = \frac{\omega_0(E')}{\omega_0(Q_0)} \approx 1$$

where in the last step we have used (6.41) and that by hypothesis, $\omega_0 \in A_{\infty}(Q_0, \sigma)$. Moreover, exactly as in (6.40), $\mathcal{P}_{\mathcal{F}}\nu(Q_0) \gtrsim 1$, so that (6.43) follows. This concludes the proof that $H[a] \Longrightarrow H[a+b]$ in Case 2.

Thus, $H[a] \Longrightarrow H[a + b]$ in both cases, and the proof of Theorem 1.7 is complete.

43

S. HOFMANN AND A. PATHAK

References

- [A] A. Ancona, On strong barriers and an inequality of Hardy for domains in \mathbb{R}^n , J. London Math. Soc. (2) **34** (1986), 274-290. 9, 10, 14, 15, 16, 17
- [AHLT] P. Auscher, S. Hofmann, J.L. Lewis and P. Tchamitchian, Extrapolation of Carleson measures and the analyticity of Kato's square-root operators, *Acta Math.* 187 (2001), no. 2, 161–190. 29
- [AHMTT] P. Auscher, S. Hofmann, C. Muscalu, T. Tao and C. Thiele, Carleson measures, trees, extrapolation, and T(b) theorems, Publ. Mat. 46 (2002), no. 2, 257–325. 29
- [BL] B. Bennewitz and J.L. Lewis, On weak reverse Hölder inequalities for nondoubling harmonic measures, *Complex Var. Theory Appl.* 49 (2004), no. 7–9, 571-582. 26, 35, 36
- [BHMN] S. Bortz, S. Hofmann, J. M. Martell and K. Nyström, Solvability of the L^p Dirichlet problem for the heat equation is equivalent to parabolic uniform rectifiability in the case of a parabolic Lipschitz graph, preprint, arXiv:2306.17291 3
- [CFMS] L. Caffarelli, E. Fabes. S. Mortola, AND S. Salsa, Boundary behavior of nonnegative solutions of elliptic operators in divergence form, *Indiana Univ. Math. J.* 30 (1981), 621-640. 14
- [Car] L. Carleson, Interpolation by bounded analytic functions and the corona problem, *Ann. of Math. (2)* **76** (1962), 547–559. **29**
- [CG] L. Carleson and J. Garnett, Interpolating sequences and separation properties, J. Analyse Math. 28 (1975), 273–299. 29
- [Chr] M. Christ, A *T*(*b*) theorem with remarks on analytic capacity and the Cauchy integral, *Colloq. Math.*, LX/LXI (1990), 601–628. 4, 5
- [DJK] B.E. Dahlberg, D.S. Jerison and C.E. Kenig, Area integral estimates for elliptic differential operators with nonsmooth coefficients, *Ark. Mat.* 22 (1984), no. 1, 97–108. 29, 39, 42
- [D1] G. David, Morceaux de graphes lipschitziens et intégrales singulières sur une surface. (French) [Pieces of Lipschitz graphs and singular integrals on a surface] *Rev. Mat. Iberoamericana* 4 (1988), no. 1, 73-114. 5
- [D2] G. David, Wavelets and singular integrals on curves and surfaces. Lecture Notes in Mathematics, 1465. Springer-Verlag, Berlin, 1991. 5
- [DS1] G. David and S. Semmes, Singular integrals and rectifiable sets in \mathbb{R}^n : Beyond Lipschitz graphs, *Asterisque* **193** (1991). **4**, **5**
- [DS2] G. David and S. Semmes, Analysis of and on Uniformly Rectifiable Sets, Mathematical Monographs and Surveys 38, AMS 1993. 5
- [FKP] R. A. Fefferman, C. E. Kenig and J. Pipher, The theory of weights and the Dirichlet problem for elliptic equations, *Ann. of Math.* (2) **134** (1991), no. 1, 65–124. 3, 25, 26, 28, 42
- [GT] D. Gilbarg and N. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, 1983. 12, 17, 20
- [HK] T. Hytönen and A. Kairema. Systems of dyadic cubes in a doubling metric space, *Colloquium Mathematicum* 126 (2012), no. 1, 1–33. 4, 5
- [H] S. Hofmann. Quantitative absolute continuity of harmonic measure and the Dirichlet problem: a survey of recent progress, *Acta Mathematica Sinica, English Series* 35 (2019) (Special Volume in honor of the 65th birthday of Carlos Kenig), 1011-1026. 7
- [HLe] S. Hofmann and P. Le. BMO solvability and absolute continuity of harmonic measure J. Geom. Anal. 28 (2018), no. 4, 3278–3299. 7
- [HL] S. Hofmann and J.L. Lewis, The Dirichlet problem for parabolic operators with singular drift terms, *Memoirs of the Amer. Math. Soc.* Vol. 151, No. 719, May 2001. 2, 3, 14, 22, 29
- [HM1] S. Hofmann and J.M. Martell, A_∞ estimates via extrapolation of Carleson measures and applications to divergence form elliptic operators, *Trans. Amer. Math. Soc.* 364 (2012), no. 1, 65–101 29
- [HM2] S. Hofmann and J.M. Martell, Uniform rectifiability and harmonic measure I: Uniform rectifiability implies Poisson kernels in L^p, Ann. Sci. École Norm. Sup. 47 (2014), no. 3, 577–654. 7, 8, 9, 11, 29, 36, 38, 39
- [HMMM] S. Hofmann, D. Mitrea, M. Mitrea, A. Morris, L^p-Square Function Estimates on Spaces of Homogeneous Type and on Uniformly Rectifiable Sets, *Mem. Amer. Math. Soc.* 245, Number 1159, January 2017, 108 pages. 5
- [JK] D. Jerison and C. Kenig. Boundary behavior of harmonic functions in nontangentially accessible domains, Adv. in Math. 46 (1982), 80-147. 5, 6, 14

- [K] C. Kenig, Harmonic analysis techniques for second order elliptic boundary value problems, CBMS Regional Conference Series in Mathematics, 83. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1994. 17, 23, 24
- [KP] C. Kenig and J. Pipher, The Dirichlet problem for elliptic equations with drift terms, Publ. Matemàtiques 45 (2001), 199-217. 2
- [KS] S. Kim and G. Sakellaris, Green's function for second order elliptic equations with singular lower order coefficients, *Communications in Partial Differential Equations* 44 (2019), 228-270. 20, 22
- [L] J. Lewis, Uniformly fat sets, Trans. Amer. Math. Soc. 308 (1988), 177-196. 9, 10, 14, 15, 16
- [LM] J. Lewis and M. Murray, The method of layer potentials for the heat equation in time-varying domains, *Mem. Amer. Math. Soc.* **114** (1995), no. 545. 3, 29
- [M] M. Mourgoglou, Regularity theory and Green's function for elliptic equations with lower order terms in unbounded domains, *Calc. Var.* **62**, 266 (2023). 20, 22
- [P] A. Pathak, A counterexample for pointwise upper bounds on Green's function with a singular drift at boundary, preprint *arXiv:2405.13313.* **3**
- [S] G. Sakellaris, On scale-invariant bounds for the Green's function for second-order elliptic equations with lower-order coefficients and applications, *Analysis and PDE* **14** (2021), 251–299. **22**
- [Ste] E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, Princeton, NJ, 1970. 7

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211, USA *Email address*: hofmanns@missouri.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211, USA *Email address*: ap7mx@mail.missouri.edu