

EXISTENCE AND STABILITY OF VARIOUS MODELS DESCRIBING BEHAVIOUR OF THE THERMOVISCOELASTIC-RATE TYPE FLUIDS

MIROSLAV BULÍČEK AND JAKUB WOŹNICKI

ABSTRACT. Viscoelastic rate-type fluid models are used to describe the behaviour of many complex materials from engineering up to application in biomaterials and medicine. A classical model that belongs to the category of viscoelastic rate-type fluid models is the Giesekus model. Furthermore, in all these applications, the heat conduction frequently takes place and all material coefficients depend heavily on the temperature, and therefore, we introduce here a thermodynamically compatible model for viscoelastic rate-type and heat-conducting fluid for which we show the existence of global weak solution in two-dimensional setting whenever the initial energy and entropy are controlled in natural norms.

1. INTRODUCTION

Our main goal in this paper is to study the equations describing the motion of incompressible viscoelastic and heat conducting fluids and in particular to establish the long-time and the large-data theory. We focus here only on the planar case, i.e., in what follows, the fluid occupies the Lipschitz domain $\Omega \subset \mathbb{R}^2$ and $T > 0$ always denotes the length of time interval. The motion of incompressible fluid is in general described by the system of partial differential equations of the form

$$(1.1) \quad \begin{aligned} \partial_t u + \operatorname{div}_x(u \otimes u) - \operatorname{div}_x \mathbb{T} &= 0, \\ \operatorname{div}_x u &= 0, \end{aligned}$$

that is supposed to be satisfied in $\Omega_T := (0, T) \times \Omega$. Here, $u : \Omega_T \rightarrow \mathbb{R}^2$ is the unknown velocity field and $\mathbb{T} : \Omega_T \rightarrow \mathbb{R}^{2 \times 2}$ denotes the Cauchy stress tensor. The above system must be completed by initial and boundary condition, but most importantly, the constitutive law for \mathbb{T} must be given.

2020 *Mathematics Subject Classification.* 35A01, 35Q35, 76A10, 76D03.

Key words and phrases. non-Newtonian fluids, Giesekus model, thermo-visco-elasticity, global-in-time and large-data existence theory.

Miroslav Bulíček was supported by the project No. 20-11027X financed by GAČR.

Jakub Woźnicki was supported by National Science Center, Poland through the project no. 2023/49/N/ST1/02737.

We shall consider the classical form

$$(1.2) \quad \mathbb{T} := -p\mathbb{I} + 2\nu Du + 2g(\mathbb{B} - \mathbb{I}),$$

where p is the unknown pressure, ν denotes the kinematic viscosity, $Du := \frac{1}{2}((\nabla u) + (\nabla u)^T)$ is the symmetric velocity gradient and $\mathbb{B} : \Omega_T \rightarrow \mathbb{R}^{2 \times 2}$ is the so-called extra stress. The first two terms on the right hand side of (1.2) describe the classical Newtonian fluid, but the last term, the extra stress, represents the elastic effects. Hence, we need to add an equation for \mathbb{B} and we shall assume the so-called rate type model, i.e., we write the constitutive equations for $\overset{\nabla}{\mathbb{B}}$, which denotes the Oldroyd upper convective derivative given as

$$(1.3) \quad \overset{\nabla}{\mathbb{B}} := \partial_t \mathbb{B} + (u \cdot \nabla_x) \mathbb{B} - \nabla_x u \mathbb{B} - \mathbb{B} (\nabla_x u)^T,$$

which is one of the possible objective derivatives applied to \mathbb{B} . The system (1.1) will be completed by the following equation

$$(1.4) \quad \overset{\nabla}{\mathbb{B}} + \delta(\mathbb{B}^2 - \mathbb{B}) + \gamma(\mathbb{B} - \mathbb{I}) = 0.$$

The system (1.1)–(1.4) with $\delta := 0$ and $\gamma > 0$ is usually named Oldroyd-B model, see [24], and belongs among one of the classical models of viscoelasticity. Unfortunately, any mathematical theory which would lead to the global well-posed theory is missing. The other model with $\gamma := 0$ and $\delta > 0$ proposed by Giesekus, see [18], is also widely used and contrary to the Oldroyd-B model, it seems to have much better mathematical properties. Note also that beside the upper convective derivative (1.3), one can also consider a much more general class of derivatives, see [21], but it usually does not affect the qualitative analysis. Similarly, one may also consider models of second order, see [10], but again, once we are able to deal with the classical Giesekus model, it seems that further generalisations are rather straightforward. Therefore, we do not consider them here. Hence, the system, we are interested in, reads as

$$(1.5) \quad \begin{cases} \partial_t u + \operatorname{div}_x(u \otimes u) - \operatorname{div}_x \mathbb{T} = 0, \\ \operatorname{div}_x u = 0, \\ \overset{\nabla}{\mathbb{B}} + \delta(\theta)(\mathbb{B}^2 - \mathbb{B}) = 0, \\ -p\mathbb{I} + 2\nu(\theta)Du + 2g(\theta)(\mathbb{B} - \mathbb{I}) = \mathbb{T}, \end{cases}$$

and we assume that all material coefficients may depend on the temperature $\theta : \Omega_T \rightarrow \mathbb{R}$. Once, we want to add the thermal effect into the system (the dependence of coefficients on the temperature),

we must also impose the validity of the first law of thermodynamics, i.e., the conservation of energy. One of the possible equivalent form (valid for sufficiently smooth solution) is the equation for the internal energy $e : \Omega_T \rightarrow \mathbb{R}$, where we shall assume the Fourier law, i.e.,

$$(1.6) \quad \partial_t e + \operatorname{div}_x(ue) - \operatorname{div}_x(\kappa(\theta)\nabla_x\theta) = \mathbb{T} : Du,$$

where κ denotes the heat conductivity. Note here that the symbol $\mathbb{A} : \mathbb{B}$ denotes the matrix scalar product, i.e., $\mathbb{A} : \mathbb{B} = \sum_{i,j=1}^2 \mathbb{A}_{ij}\mathbb{B}_{ij}$.

It remains to specify the storage and dissipative mechanisms of the fluid, or to say it differently, it remains to specify the relation between the internal energy e , the temperature θ and the tensor \mathbb{B} . Further, this relation must be such that one can identify an entropy in the system, which fulfils the second law of thermodynamics. Here, we closely follow the approach developed in [25] and in [19], where the Helmholtz free energy plays the crucial role. However, we may also refer to the work [14] for the first systematic approach or to [20], where the entropy estimates were derived for the first time.

Hence, we assume here that the Helmholtz free energy of the fluid is the form

$$(1.7) \quad \psi = -c_v\theta(\ln(\theta) - 1) + g(\theta)(\operatorname{tr}(\mathbb{B}) - 2 - \ln \det \mathbb{B}).$$

Here, the first term, corresponds to the classical law when $e = c_v\theta$ with c_v being the heat capacity, and the second term connects the elastic and the temperature effects. Note that if g is constant, this choice is very much classical in the theory of viscoelastic rate type fluids, however, one may also consider much more general setting, see [1].

Having introduced the free energy, we can then define the entropy as

$$\eta := -\partial_\theta\psi = c_v \ln \theta - g'(\theta)(\operatorname{tr}(\mathbb{B}) - 2 - \ln \det \mathbb{B}),$$

and the internal energy is then related by the formula

$$(1.8) \quad e := \psi + \theta\eta = \theta + (g(\theta) - \theta g'(\theta))(\operatorname{tr}(\mathbb{B}) - 2 - \ln \det \mathbb{B}).$$

Using these definitions, we see that

$$\begin{aligned} \partial_t e &= \partial_t\psi + \partial_t\theta\eta + \theta\partial_t\eta = \partial_t\theta(\partial_\theta\psi + \eta) + \partial_\mathbb{B}\psi\partial_t\mathbb{B} + \theta\partial_t\eta = \partial_\mathbb{B}\psi\partial_t\mathbb{B} + \theta\partial_t\eta \\ \implies \partial_t\eta &= \frac{\partial_t e - \partial_\mathbb{B}\psi\partial_t\mathbb{B}}{\theta} = \frac{\partial_t e - \partial_t\mathbb{B} : g(\theta)(\mathbb{I} - \mathbb{B}^{-1})}{\theta}. \end{aligned}$$

Thus, dividing (1.6) by θ , taking the scalar product of the third equation in (1.5) by $\frac{g(\theta)(\mathbb{I}-\mathbb{B}^{-1})}{\theta}$ and subtracting the result, we obtain the following identity for the entropy

$$(1.9) \quad \partial_t \eta + \operatorname{div}_x(\eta u) - \operatorname{div}_x(\kappa(\theta) \nabla_x(\ln \theta)) = \frac{\kappa(\theta) |\nabla_x \theta|^2}{\theta^2} + \frac{2\nu(\theta) |Du|^2}{\theta} + \frac{\delta(\theta) |\mathbb{B} - \mathbb{I}|^2}{\theta}.$$

Here, it is essential that the right-hand side is nonnegative and the term $\kappa(\theta) \nabla_x(\ln \theta)$ represents the entropy flux. It also directly follows from the procedure above that the equation for the internal energy (1.6) and the equation for the entropy (1.9) are interchangeable in (1.5) for sufficiently regular solutions, which we will frequently use.

Finally, we shall always assume that \mathbb{B} is of the form

$$\mathbb{B} = \mathbb{F} \mathbb{F}^T, \quad \mathbb{F} \in \mathbb{R}^{2 \times 2}, \quad \det(\mathbb{F}) > 0.$$

The above identification can be for example guaranteed, by imposing the following equation for \mathbb{F}

$$(1.10) \quad \partial_t \mathbb{F} + \operatorname{div}_x(\mathbb{F} \otimes u) - \nabla_x u \mathbb{F} + \frac{1}{2} \delta(\theta) (\mathbb{F} \mathbb{F}^T \mathbb{F} - \mathbb{F}) = 0.$$

In fact, multiplying this identity by \mathbb{F}^T on the right and multiplying the transposed version of (1.10) by \mathbb{F} on the left, adding the result together, we see that $\mathbb{B} := \mathbb{F} \mathbb{F}^T$ then satisfies (1.4) with $\gamma = 0$.

Hence, we have already introduced the complete set of equations and we close the problem by imposing the following boundary and initial conditions

$$(1.11) \quad u|_{\partial\Omega} = 0, \quad u(0, x) = u_0(x), \quad \mathbb{B}(0, x) = \mathbb{B}_0 = \mathbb{F}_0 \mathbb{F}_0^T, \quad \nabla_x \theta \cdot \mathbf{n} = 0, \quad \theta(0, x) = \theta_0.$$

Furthermore, the initial conditions are assumed to be such that they have bounded kinetic and internal initial energy and that the initial entropy is also under control, i.e.,

$$\|u_0\|_2, \|\mathbb{F}_0\|_2, \|\operatorname{tr}(\mathbb{B}_0) - 2 - \ln \det \mathbb{B}_0\|_1, \|\ln \det \mathbb{B}_0\|_\infty, \|\theta_0\|_1, \|\ln(\theta_0)\|_1 < +\infty \quad \det \mathbb{F}_0 > 0 \text{ a.e. in } \Omega.$$

Finally, we introduce three specific choices of the form of the Helmholtz free energy and in particular the function g :

(P1) If g is a constant function then the entropy depends only on temperature and

$$\eta = c_v \ln(\theta).$$

(P2) If $g(\theta) = c\theta$, then the internal energy depends only on temperature and

$$e = c_v\theta.$$

(P3) If g is a general function, then the equations connecting temperature and the elastic tensor are coupled in a non-trivial way.

Our aim is to develop the theory related to the three cases above, in particular to the most difficult case (P3). To simplify the presentation, we prove in the paper in the general case (P3) only the weak sequential stability of solutions, see Section 3. Although it is not the full proof of existence, it is clear that the full proof can be established. This is in fact very much supported by our second result of the paper, that is connected to the case (P1) and is presented in Section 4, where we show the existence of weak solution. Section 2 is devoted to the basic notation, and in Appendix A we state some classical, auxiliary propositions.

To end this introductory part, we recall available results and emphasise the key novelty of the paper. We start by recalling the results without thermal effects. As we have already mentioned, the result for the full Oldroyd-B model is an open problem. The only available result (for global solutions) is due to Lions and Masmoudi [22], where the existence is shown even in three dimensional setting but only for the so-called co-rotational case. On the other hand, for the Giesekus model (or for even more general class), the first serious attempt for global theory was done in [23]. Although the key idea there is correct, there were still some gaps in the proof, that had to be corrected. The first rigorous result for such a class of fluids was established in [1] with one proviso, an additional stress diffusion term $\Delta\mathbb{B}$ was added to the problem, which simplified the analysis a little bit. The problem without stress diffusion was finally solved in [6] for the two-dimensional setting and later in [7] for the three-dimensional setting and a much more complex class of models. For compressible setting, we refer also to a particular result in [4]. For problems with temperature, the situation is more delicate. Even for problems without viscoelastic effects, the theory is relatively new, and here we refer to [3,9,11] for relevant results. The problems with elastic effects and temperature were first rigorously treated in [8], where, however, only spherical stresses were assumed, i.e. $\mathbb{B} = b\mathbb{I}$. Much more general results was the established in [2], where the authors again assumed a modification of the Giesekus model by adding the stress diffusion into the system and by focusing only on the linear case (P2). Here, in this paper, we, however, do not need any stress diffusion term in the equation nor the simple form of the free energy. Furthermore, it is clear from the proof that we are also

able to cover the three-dimensional case and also a more complex model - to such a setting will be devoted our forth coming paper.

To end this part, it is essential to mention that the paper heavily relies not only on the results mentioned above, but also on the following two very classical techniques. First, we should mention the theory of renormalisation for the transport equation developed in [13], which is used when dealing with the equation for \mathbb{B} . Second, the compensated compactness method applied to heat conductive fluids, which was introduced by Feireisl, see [15] and further generalised and extended to other cases, see [16, 17] and [5]. These methods are used to deduce the compactness of temperature θ and tensor \mathbb{B} .

2. PRELIMINARIES

In what follows, $x \in \Omega$ always denotes the spatial variable, while $t \in (0, T)$ is reserved for the time variable. For vector manipulations, we write $a \cdot b$ for a scalar product whenever $a, b \in \mathbb{R}^d$. Similarly, by $A : B$, we denote the scalar product between two matrices $A, B \in \mathbb{R}^{d \times d}$, and their matrix product by AB . At last, by $A \cdot \cdot B$, we will mean the scalar product between $A, B \in \mathbb{R}^{d \times d \times d}$. In addition, the symbol \otimes is reserved for the tensorial product, i.e., for $a, b \in \mathbb{R}^d$ we denote $a \otimes b \in \mathbb{R}_{\text{sym}}^{d \times d}$ as $(a \otimes b)_{ij} := a_i b_j$ for $i, j = 1, \dots, d$. For a matrix $A = (A_{ij})_{i,j=1}^d$ and a vector $b = (b_k)_{k=1}^d$ we define the third order tensorial product as

$$(A \otimes b)_{ijk} = A_{ij} b_k,$$

and its divergence as

$$\operatorname{div}_x(A \otimes b) = \sum_{j=1}^d \partial_{x_j} (b_j A).$$

We use the standard notation for Sobolev and Lebesgue function space and frequently do not distinguish between scalar-, vector- or matrix-valued functions. In addition, to shorten the notation, we frequently use the following simplifications. When $f \in L^p(\Omega)$, we simplify it to $f \in L_x^p$. Similarly, if $f \in L^p(0, T; L^q(\Omega))$, $f \in L^p(0, T; W^{1,q}(\Omega))$ or $f \in L^p(0, T; W_0^{1,q}(\Omega))$, then we write $f \in L_t^p L_x^q$, $f \in L_t^p W_x^{1,q}$ or $f \in L_t^p W_{0,x}^{1,q}$ respectively (here, $W^{1,q}(\Omega)$ and $W_0^{1,q}(\Omega)$ are the usual Sobolev spaces). Moreover, for $p \in [1, +\infty]$ we denote by p' its Hölder conjugate. To further shorten the notation we introduce the function

$$f(\mathbb{B}) = \operatorname{tr}(\mathbb{B}) - 2 - \ln \det \mathbb{B},$$

which is the one frequently appearing throughout the paper, as well as the one present in the definition of the Helmholtz free energy (1.7). To end, we introduce the assumptions concerning the function ν, κ, δ, g appearing in the equations (1.5) - (1.9). For the case (P1) we assume, that there exists $C > 0$ such that

$$(2.1) \quad C^{-1} \leq \kappa(w) \leq C,$$

$$(2.2) \quad C^{-1} \leq \nu(w) \leq C,$$

$$(2.3) \quad C^{-1} \leq \delta(w) \leq C$$

for any $w \in \mathbb{R}$. And for the case (P3), that there exist $c_\nu, C_\nu, c_\kappa, C_\kappa > 0$ such that

$$(2.4) \quad c_\kappa \leq \kappa(w) \leq C_\kappa,$$

$$(2.5) \quad c_\nu \leq \nu(w) \leq C_\nu.$$

for any $w \in \mathbb{R}$,

$$(2.6) \quad \delta(w) \sim w$$

for any $w \in \mathbb{R}$. Moreover, for a function g we assume that

$$(G1) \quad g \in C^2(\mathbb{R}),$$

$$(G2) \quad g \text{ is concave,}$$

$$(G3) \quad g \text{ is increasing,}$$

$$(G4) \quad \max_{w \geq 0} g(w) < +\infty, \min_{w \geq 0} g(w) = g(0) > 0,$$

$$(G5) \quad g'(w) \text{ is a bounded function,}$$

$$(G6) \quad w g'(w) \text{ and } h_\lambda(w) \text{ are bounded functions, where } h_\lambda(w) := \int_0^w z^\lambda g''(z) \, dz.$$

3. WEAK SEQUENTIAL STABILITY OF STRONG SOLUTIONS FOR THE CASE (P3)

We start by simplifying the equations (1.5) - (1.6) to

$$(3.1) \quad \begin{cases} \partial_t u + \operatorname{div}_x(u \otimes u) - \operatorname{div}_x \mathbb{T} = 0, \\ \operatorname{div}_x u = 0, \\ \partial_t \mathbb{F} + \operatorname{div}_x(\mathbb{F} \otimes u) - \nabla_x u \mathbb{F} + \frac{1}{2} \delta(\theta)(\mathbb{F} \mathbb{F}^T \mathbb{F} - \mathbb{F}) = 0, \\ -p \mathbb{I} + 2\nu(\theta) \mathbb{D}u + 2g(\theta)(\mathbb{F} \mathbb{F}^T) = \mathbb{T}, \\ \partial_t e + u \cdot \nabla_x e - \operatorname{div}_x(\kappa(\theta) \nabla_x \theta) = (2\nu(\theta) Du + 2g(\theta) \mathbb{F} \mathbb{F}^T) : Du. \end{cases}$$

with the initial and boundary conditions

$$(3.2) \quad u|_{\partial\Omega} = 0, \quad u(0, x) = u_0(x), \quad \mathbb{F}(0, x) = \mathbb{F}_0, \quad \nabla_x \theta \cdot \mathbf{n} = 0, \quad \theta(0, x) = \theta_0.$$

One can observe that (1.5)₃ follows formally from (3.1)₃ by multiplying it by \mathbb{F}^T , its transposition by \mathbb{F} , and adding both equations together. Let us state the main theorem that we are going to prove.

Theorem 3.1. *Assume $(u_n, \mathbb{F}_n, \theta_n) \in C^1((0, T) \times \Omega; \mathbb{R}^2) \times C^1((0, T) \times \Omega; \mathbb{R}^{2 \times 2}) \times C^{1,2}((0, T) \times \Omega; \mathbb{R}_+)$ with $\det \mathbb{F}_n > 0$ is a sequence of strong solutions to the system (3.1). Then, there exists a subsequence (which we do not relabel) such that*

$$(3.3) \quad u_n \overset{*}{\rightharpoonup} u \quad \text{weakly}^* \text{ in } L_t^\infty L_x^2,$$

$$(3.4) \quad u_n \rightharpoonup u \quad \text{weakly in } L_t^2 W_{0,x}^{1,2},$$

$$(3.5) \quad u_n \rightharpoonup u \quad \text{weakly in } L_{t,x}^4,$$

$$(3.6) \quad u_n \rightarrow u \quad \text{strongly in } C([0, T]; L^p(\Omega)), 1 \leq p < 4,$$

$$(3.7) \quad \mathbb{F}_n \overset{*}{\rightharpoonup} \mathbb{F} \quad \text{weakly}^* \text{ in } L_t^\infty L_x^2,$$

$$(3.8) \quad \mathbb{F}_n \rightharpoonup \mathbb{F} \quad \text{weakly in } L_{t,x}^4,$$

$$(3.9) \quad \mathbb{F}_n \rightarrow \mathbb{F} \quad \text{strongly in } L_{t,x}^p, 1 \leq p < 4,$$

$$(3.10) \quad \theta_n \rightarrow \theta \quad \text{strongly in } L_{t,x}^{2-\varepsilon}, \varepsilon \in (0, 1),$$

$$(3.11) \quad \nabla_x \theta_n \rightharpoonup \nabla_x \theta \quad \text{weakly in } L_{t,x}^{\frac{4}{3}-\varepsilon}, \varepsilon \in \left(0, \frac{1}{3}\right),$$

$$(3.12) \quad \ln(\theta_n) \rightharpoonup \ln(\theta) \quad \text{weakly in } L_t^2 W_x^{1,2},$$

$$(3.13) \quad \ln(\theta_n) \rightarrow \ln(\theta) \quad \text{strongly in } L_{t,x}^{3-\varepsilon}, \varepsilon \in (0, 1),$$

$$(3.14) \quad \text{tr}(\mathbb{F}_n \mathbb{F}_n^T) \rightharpoonup \text{tr}(\mathbb{F} \mathbb{F}^T) \quad \text{weakly in } L_{t,x}^2,$$

$$(3.15) \quad \ln \det(\mathbb{F}_n \mathbb{F}_n^T) \overset{*}{\rightharpoonup} \ln \det(\mathbb{F} \mathbb{F}^T) \quad \text{weakly}^* \text{ in } L_t^\infty L_x^{\frac{4}{3}-\varepsilon}, \varepsilon \in \left(0, \frac{1}{3}\right).$$

where (u, \mathbb{F}, θ) is a weak solution to the system (3.1), that is

$$(3.16) \quad \int_0^T \int_\Omega -u \cdot \partial_t \phi - u \otimes u : \nabla_x \phi + \nu(\theta) Du : \nabla_x \phi + g(\theta) \mathbb{F} \mathbb{F}^T : \nabla_x \phi \, dx \, dt = \int_\Omega u_0(x) \cdot \phi(0, x) \, dx,$$

for any $\phi \in C_c^1([0, T] \times \Omega; \mathbb{R}^2)$ with $\operatorname{div}_x \phi = 0$,

(3.17)

$$\int_0^T \int_{\Omega} -\mathbb{F} : \partial_t \phi - \mathbb{F} \otimes u : \nabla_x \phi - \nabla_x u \mathbb{F} : \phi + \frac{1}{2} \delta(\theta) (\mathbb{F} \mathbb{F}^T \mathbb{F} - \mathbb{F}) : \phi \, dx \, dt = \int_{\Omega} \mathbb{F}_0(x) : \phi(0, x) \, dx,$$

for any $\phi \in C_c^1([0, T] \times \Omega; \mathbb{R}^{2 \times 2})$,

(3.18)

$$\int_0^T \int_{\Omega} -(\ln(\theta) - g'(\theta) f(\mathbb{F} \mathbb{F}^T)) \partial_t \phi - u (\ln(\theta) - g'(\theta) f(\mathbb{F} \mathbb{F}^T)) \nabla_x \phi + \kappa(\theta) \nabla_x \ln \theta \nabla_x \phi \, dx \, dt \geq 0$$

for any $\phi \in C_c^1([0, T] \times \Omega)$, $\phi \geq 0$. Moreover $\theta > 0$ and $\det \mathbb{F} > 0$.

Proof. We begin by recovering the equation for $\mathbb{B}_n := \mathbb{F}_n \mathbb{F}_n^T$. Namely, multiply the equation (3.1)₃ by \mathbb{F}_n^T from the right and the transpose of the mentioned equation by \mathbb{F}_n from the left. Adding the results together we obtain

$$(3.19) \quad \partial_t \mathbb{B}_n + u_n \nabla_x \mathbb{B}_n + \delta(\theta_n) (\mathbb{B}_n^2 - \mathbb{B}_n) = \nabla_x u_n \mathbb{B}_n + \mathbb{B}_n (\nabla_x u_n)^T.$$

With this, we may apply the material derivative to the equation (1.8) to get

$$\partial_t e_n + u_n \nabla_x e_n = \partial_{\mathbb{B}} \psi(\theta_n, \mathbb{B}_n) (\partial_t \mathbb{B}_n + u_n \nabla_x \mathbb{B}_n) + \theta_n (\partial_t \eta_n + u_n \nabla_x \eta_n),$$

which together with the equations (3.1)₅, (3.19) and the fact that $f'(\mathbb{B}_n) = \mathbb{I} - \mathbb{B}_n^{-1}$ implies

$$(3.20) \quad \partial_t \eta_n + u_n \cdot \nabla_x \eta_n - \operatorname{div}_x (\kappa(\theta_n) \nabla_x (\ln \theta_n)) = \kappa(\theta_n) \frac{|\nabla_x \theta_n|^2}{\theta_n^2} + \nu(\theta_n) \frac{|Du_n|^2}{\theta_n} + \delta(\theta_n) \frac{|\mathbb{B}_n - \mathbb{I}|^2}{\theta_n}.$$

Before moving forward, let us prove a structural lemma connected to equation (3.19) and (3.20).

Lemma 3.2. *Suppose $\eta := \ln(\theta) - g'(\theta) f(\mathbb{B})$, \mathbb{B} are the strong solutions to the equation (3.20) and (3.19) respectively. Then, for any $\lambda > 0$*

$$(3.21) \quad \begin{aligned} & \frac{1}{\lambda} \partial_t \theta^\lambda - \partial_t [h_\lambda(\theta) f(\mathbb{B})] + \frac{1}{\lambda} u \cdot \nabla_x \theta^\lambda - u \cdot \nabla_x [h_\lambda(\theta) f(\mathbb{B})] + \delta(\theta) |\mathbb{B} - \mathbb{I}|^2 (g'(\theta) \theta^\lambda - h_\lambda(\theta)) \\ & - \operatorname{div}_x (\kappa(\theta) \nabla_x (\ln \theta)) \theta^\lambda - 2(\mathbb{B} - \mathbb{I}) : Du (g'(\theta) \theta^\lambda - h_\lambda(\theta)) \\ & = \left(\kappa(\theta) \frac{|\nabla_x \theta|^2}{\theta^2} + \nu(\theta) \frac{|Du|^2}{\theta} + \delta(\theta) \frac{|\mathbb{B} - \mathbb{I}|^2}{\theta} \right) \theta^\lambda, \end{aligned}$$

where

$$h_\lambda(\theta) = \int_0^\theta z^\lambda g''(z) \, dz.$$

Proof. Multiplying the equation (3.19) by $\mathbb{I} - \mathbb{B}^{-1}$ and using the fact that $f'(\mathbb{B}) = \mathbb{I} - \mathbb{B}^{-1}$ gets us

$$(3.22) \quad \partial_t f(\mathbb{B}) + u \cdot \nabla_x f(\mathbb{B}) + \delta(\theta) |\mathbb{B} - \mathbb{I}|^2 = 2(B - \mathbb{I}) : Du.$$

Thus, by plugging (3.22) multiplied by $g'(\theta)$ into (3.20), we obtain

$$\begin{aligned} & \partial_t \theta \left(\frac{1}{\theta} - g''(\theta) f(\mathbb{B}) \right) + u \cdot \nabla_x \theta \left(\frac{1}{\theta} - g''(\theta) f(\mathbb{B}) \right) \\ & - \operatorname{div}_x (\kappa(\theta) \nabla_x (\ln \theta)) + g'(\theta) \delta(\theta) |\mathbb{B} - \mathbb{I}|^2 - 2g'(\theta) (\mathbb{B} - \mathbb{I}) : Du \\ & = \kappa(\theta) \frac{|\nabla_x \theta|^2}{\theta^2} + \nu(\theta) \frac{|Du|^2}{\theta} + \delta(\theta) \frac{|\mathbb{B} - \mathbb{I}|^2}{\theta}. \end{aligned}$$

Hence, after we multiply the equation above by θ^λ we get

$$(3.23) \quad \begin{aligned} & \frac{1}{\lambda} \partial_t \theta^\lambda - \partial_t [h_\lambda(\theta)] f(\mathbb{B}) + \frac{1}{\lambda} u \cdot \nabla_x \theta^\lambda - u \cdot \nabla_x [h_\lambda(\theta)] f(\mathbb{B}) \\ & - \operatorname{div}_x (\kappa(\theta) \nabla_x (\ln \theta)) \theta^\lambda + g'(\theta) \theta^\lambda \delta(\theta) |\mathbb{B} - \mathbb{I}|^2 - 2g'(\theta) \theta^\lambda (\mathbb{B} - \mathbb{I}) : Du \\ & = \left(\kappa(\theta) \frac{|\nabla_x \theta|^2}{\theta^2} + \nu(\theta) \frac{|Du|^2}{\theta} + \delta(\theta) \frac{|\mathbb{B} - \mathbb{I}|^2}{\theta} \right) \theta^\lambda. \end{aligned}$$

To end the proof we simply subtract (3.22) multiplied by $h_\lambda(\theta)$ from (3.23). \square

Now we may proceed to prove some basic bounds for our sequence. Notice that multiplying (3.1)₁ by u_n adding it to (3.1)₅ and integrating the result yields (here we note that $e_n > 0$ as $\theta_n, f(\mathbb{B}_n) > 0$ and the concavity of g implies $g(\theta_n) - \theta_n g'(\theta_n) \geq g(0) > 0$)

$$(3.24) \quad \|u_n\|_{L_t^\infty L_x^2}^2 + \|\theta_n\|_{L_t^\infty L_x^1} + g(0) \|f(\mathbb{B}_n)\|_{L_t^\infty L_x^1} \leq \|u_n\|_{L_t^\infty L_x^2}^2 + \|e_n\|_{L_t^\infty L_x^1} = \|u_0\|_{L_x^2}^2 + \|e_0\|_{L_x^1}.$$

Moving forward, we deduce from (3.20), and the definition of δ (2.6) that

$$\begin{aligned} & \int_0^T \int_\Omega \|\mathbb{B}_n - \mathbb{I}\|^2 dx d\tau \leq \int_0^T \int_\Omega \partial_t \eta_n + u_n \cdot \nabla_x \eta_n - \operatorname{div}_x (\kappa(\theta_n) \nabla_x (\ln \theta_n)) dx d\tau \\ & = \left(\int_\Omega \ln(\theta_n) - g'(\theta_n) f(\mathbb{B}_n) dx \right) \Big|_0^T, \end{aligned}$$

which together with the inequalities $g'(w), f(\mathbb{B}) > 0$, $\ln x \leq x - 1$, the bound (3.24), and the Assumptions (G4), (G6) implies

$$(3.25) \quad \int_0^T \int_\Omega \|\mathbb{B}_n - \mathbb{I}\|^2 dx d\tau \leq C(\|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}).$$

In particular, we obtain

$$(3.26) \quad \|\mathbb{F}_n \mathbb{F}_n^T\|_{L^2_{t,x}} \leq C(\|u_0\|_{L^2_x}, \|\theta_0\|_{L^1_x}, \|\ln(\theta_0)\|_{L^1_x}, \|f(\mathbb{B}_0)\|_{L^1_x}),$$

$$(3.27) \quad \|\mathbb{F}_n\|_{L^4_{t,x}} \leq C(\|u_0\|_{L^2_x}, \|\theta_0\|_{L^1_x}, \|\ln(\theta_0)\|_{L^1_x}, \|f(\mathbb{B}_0)\|_{L^1_x}).$$

Now, we may again multiply (3.1)₁ by u_n and integrate to get

$$\frac{1}{2} \int_{\Omega} |u_n(t, x)|^2 dx + \int_0^t \int_{\Omega} 2\nu(\theta_n) |Du_n|^2 dx d\tau = - \int_0^t \int_{\Omega} g(\theta_n) \mathbb{F}_n \mathbb{F}_n^T : \nabla_x u_n dx d\tau + \frac{1}{2} \int_{\Omega} |u_0|^2 dx.$$

Hence, we may use (3.26) and Young's inequality to deduce

$$\begin{aligned} \int_0^t \int_{\Omega} 2\nu(\theta_n) |Du_n|^2 dx dt &\leq \frac{\|g\|_{\infty}}{8c_{\nu}} \int_0^t \int_{\Omega} |\mathbb{F}_n \mathbb{F}_n^T|^2 dx d\tau + 2c_{\nu} \int_0^t \int_{\Omega} |Du_n|^2 dx d\tau + \frac{1}{2} \|u_0\|_2^2 \\ &\leq 2c_{\nu} \int_0^t \int_{\Omega} |Du_n|^2 dx d\tau + C(\|u_0\|_{L^2_x}, \|\theta_0\|_{L^1_x}, \|\ln(\theta_0)\|_{L^1_x}, \|f(\mathbb{B}_0)\|_{L^1_x}). \end{aligned}$$

Thus

$$(3.28) \quad \|Du_n\|_{L^2_{t,x}} \leq C(\|u_0\|_{L^2_x}, \|\theta_0\|_{L^1_x}, \|\ln(\theta_0)\|_{L^1_x}, \|f(\mathbb{B}_0)\|_{L^1_x}),$$

and in particular by bound (2.5), Korn's inequality, the interpolation, and the Sobolev embeddings

$$(3.29) \quad \|\sqrt{\nu(\theta_n)} Du_n\|_{L^2_{t,x}} \leq C(\|u_0\|_{L^2_x}, \|\theta_0\|_{L^1_x}, \|\ln(\theta_0)\|_{L^1_x}, \|f(\mathbb{B}_0)\|_{L^1_x}),$$

$$(3.30) \quad \|\nabla_x u_n\|_{L^2_{t,x}} \leq C(\|u_0\|_{L^2_x}, \|\theta_0\|_{L^1_x}, \|\ln(\theta_0)\|_{L^1_x}, \|f(\mathbb{B}_0)\|_{L^1_x}),$$

$$(3.31) \quad \|u_n\|_{L^4_{t,x}} \leq C(\|u_0\|_{L^2_x}, \|\theta_0\|_{L^1_x}, \|\ln(\theta_0)\|_{L^1_x}, \|f(\mathbb{B}_0)\|_{L^1_x}),$$

$$(3.32) \quad \|u_n\|_{L^2_t L^b_x} \leq C(\|u_0\|_{L^2_x}, \|\theta_0\|_{L^1_x}, \|\ln(\theta_0)\|_{L^1_x}, \|f(\mathbb{B}_0)\|_{L^1_x}), \quad b < +\infty.$$

For later use, let us also note that the bounds (3.29), and (3.27), as well as the equation (3.1)₁ we may deduce that

$$(3.33) \quad \|\partial_t u_n\|_{L^2((0,T);(W_0^{1,2}(\Omega))^*)} \leq C(\|u_0\|_{L^2_x}, \|\theta_0\|_{L^1_x}, \|\ln(\theta_0)\|_{L^1_x}, \|f(\mathbb{B}_0)\|_{L^1_x}).$$

Moving forward, we multiply (3.1)₃ by \mathbb{F}_n and integrate to obtain

$$(3.34) \quad \|\mathbb{F}_n(t)\|_2^2 + \int_0^t \int_{\Omega} \delta(\theta_n) |\mathbb{F}_n \mathbb{F}_n^T|^2 dx d\tau = \int_0^t \int_{\Omega} 2\mathbb{F}_n \mathbb{F}_n^T : \nabla_x u_n dx d\tau + \int_0^t \int_{\Omega} \delta(\theta_n) |\mathbb{F}_n|^2 dx d\tau + \|\mathbb{F}_0\|_2^2,$$

which by using Hölder inequality, Young inequality and a matrix inequality $|\mathbb{F}|^4 \leq 2|\mathbb{F}\mathbb{F}^T|^2$ implies

$$\begin{aligned} \|\mathbb{F}_n(t)\|_2^2 + \frac{1}{2} \int_0^t \int_\Omega \delta(\theta_n) |\mathbb{F}_n|^4 dx d\tau \\ \leq 2\|\mathbb{F}_n \mathbb{F}_n^T\|_2 \|\nabla_x u_n\|_2 + \int_0^t \int_\Omega \delta(\theta_n) dx d\tau + \frac{1}{4} \int_0^t \int_\Omega \delta(\theta_n) |\mathbb{F}_n|^4 dx d\tau + \|\mathbb{F}_0\|_2^2. \end{aligned}$$

Hence, by (3.26), (3.30), the definition of δ (2.6) and the bounds (3.24)

$$(3.35) \quad \|\mathbb{F}_n\|_{L_t^\infty L_x^2}^2 + \int_0^t \int_\Omega \delta(\theta_n) |\mathbb{F}_n|^4 dx d\tau \leq C(\|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}).$$

In particular, coming back with the obtained bound to the inequality (3.34) gives us

$$(3.36) \quad \int_0^t \int_\Omega \delta(\theta_n) |\mathbb{F}_n \mathbb{F}_n^T|^2 dx d\tau \leq C(\|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}).$$

To derive some compactness for θ_n we use Lemma 3.2 to deduce for any $1 > \lambda > 0$.

$$\begin{aligned} (1-\lambda) \int_0^t \int_\Omega \kappa(\theta_n) \frac{|\nabla_x \theta_n|^2}{\theta_n^{2-\lambda}} dx d\tau \\ \leq \frac{1}{\lambda} \int_\Omega \theta_n^\lambda(t) dx d\tau - \int_\Omega h_\lambda(\theta_n(t)) f(\mathbb{B}_n(t)) dx \\ + \int_0^t \int_\Omega \delta(\theta_n) |\mathbb{B}_n - \mathbb{I}|^2 (g'(\theta_n) \theta_n^\lambda - h_\lambda(\theta_n)) dx d\tau \\ - \int_0^t \int_\Omega 2(\mathbb{B}_n - \mathbb{I}) : Du_n (g'(\theta_n) \theta_n^\lambda - h_\lambda(\theta_n)) dx d\tau, \end{aligned}$$

which together with Hölder inequality, (3.24), (3.36), (3.28), (3.25) and the assumption (G6) implies

$$(3.37)$$

$$\int_0^t \int_\Omega \frac{|\nabla_x \theta_n|^2}{\theta_n^\lambda} dx dt \leq C(\lambda, \|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}), \quad \text{for any } \lambda \in (1, 2).$$

With this, we may infer some uniform bounds on θ_n and $\nabla_x \theta_n$. Let us recall an interpolation inequality

$$(3.38) \quad \|v\|_{L^a(\Omega)}^a \leq C \left(\|v\|_{L^{\frac{2}{2-\lambda}}(\Omega)}^a + \|v\|_{L^{\frac{2}{2-\lambda}}(\Omega)}^{s-2} \|\nabla_x v\|_{L^2(\Omega)}^2 \right),$$

where $\lambda > 1$ is sufficiently close to 1 and a is defined as

$$a := 2 + \frac{2}{2-\lambda}.$$

Hence, with the use of (3.38) and the bounds (3.24), (3.37)

$$\begin{aligned}
(3.39) \quad & \int_0^T \int_{\Omega} \theta_n^{3-\lambda} dx dt = \int_0^T \int_{\Omega} (\theta_n^{\frac{2-\lambda}{2}})^s dx dt \lesssim \int_0^T \|\theta_n^{\frac{2-\lambda}{2}}\|_{\frac{2}{2-\lambda}}^s + \|\theta_n^{\frac{2-\lambda}{2}}\|_{\frac{2}{2-\lambda}}^{s-2} \|\nabla_x (\theta_n^{\frac{2-\lambda}{2}})\|_2^2 dt \\
& = \int_0^T \left(\int_{\Omega} \theta_n dx \right)^{3-\lambda} dt + \int_0^T \left(\int_{\Omega} \theta_n dx \right) \left(\int_{\Omega} \frac{|\nabla_x \theta_n|^2}{\theta_n^\lambda} dx \right) dt \\
& \leq C(\lambda, \|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}).
\end{aligned}$$

for any $\lambda \in (1, 2)$. Then, with the use of Young's inequality, (3.39) and (3.37) we may infer

$$\begin{aligned}
(3.40) \quad & \int_0^T \int_{\Omega} |\nabla_x \theta_n|^{2-\frac{2}{3}\lambda} dx dt = \int_0^T \int_{\Omega} \left(\frac{|\nabla_x \theta_n|^2}{\theta_n^\lambda} \right)^{\frac{2-\frac{2}{3}\lambda}{2}} \theta_n^{\frac{(2-\frac{2}{3}\lambda)\lambda}{2}} dx dt \\
& \leq \int_0^T \int_{\Omega} \frac{|\nabla_x \theta_n|^2}{\theta_n^\lambda} + \theta_n^{\frac{(2-\frac{2}{3}\lambda)\lambda}{2-(2-\frac{2}{3}\lambda)}} dx dt \\
& = \int_0^T \int_{\Omega} \frac{|\nabla_x \theta_n|^2}{\theta_n^\lambda} + \theta_n^{3-\lambda} dx dt \\
& \leq C(\lambda, \|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}),
\end{aligned}$$

for any $\lambda \in (1, 2)$. In short, (3.39) and (3.40) give us

$$(3.41) \quad \|\theta_n\|_{L_{t,x}^{2-\lambda}} \leq C(\lambda, \|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}),$$

$$(3.42) \quad \|\nabla_x \theta_n\|_{L_{t,x}^{\frac{4}{3}-\lambda}} \leq C(\lambda, \|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}),$$

for any $\lambda \in (0, 1)$. Now, our goal will be to establish the regularity of the sequence $f(\mathbb{B}_n)$. First, due to the general inequality for positive, define matrices $|\text{tr}(\mathbb{F})| \leq \sqrt{2}|\mathbb{F}|$ we have

$$(3.43) \quad \|\text{tr}(\mathbb{B}_n)\|_{L_{t,x}^2} \leq \sqrt{2}\|\mathbb{B}_n\|_{L_{t,x}^2} \leq C(\|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}),$$

$$(3.44) \quad \|\text{tr}(\mathbb{B}_n)\|_{L_t^\infty L_x^1} \leq \sqrt{2}\|\mathbb{B}_n\|_{L_t^\infty L_x^1} \leq C(\|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}).$$

To estimate the second term, we multiply the equation (3.19) by \mathbb{B}_n^{-1} and get

$$(3.45) \quad \partial_t \ln \det \mathbb{B}_n + u_n \cdot \nabla_x \ln \det \mathbb{B}_n + \delta(\theta_n) \text{tr}(\mathbb{B}_n - \mathbb{I}) = 0.$$

From (3.41) and the definition of δ (2.6) we can see that

$$(3.46) \quad \|\delta(\theta_n)\|_{L_{t,x}^{2-\varepsilon}} \leq C(\varepsilon, \|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}),$$

for some small $\varepsilon > 0$. Thus, by Hölder inequality, the bounds (3.46), (3.36), and the matrix inequality $|\operatorname{tr}(\mathbb{F})| \leq \sqrt{2}|\mathbb{F}|$ we obtain

$$(3.47) \quad \int_0^T \int_{\Omega} |\delta(\theta_n) \operatorname{tr}(\mathbb{B}_n - \mathbb{I})|^{\frac{4-2\varepsilon}{3-\varepsilon}} dx dt \leq \|\delta(\theta_n)\|_{L^{2-\varepsilon}} \left(\int_0^T \int_{\Omega} \delta(\theta_n) |\operatorname{tr}(\mathbb{B}_n - \mathbb{I})|^2 dx dt \right)^{\frac{2-\varepsilon}{3-\varepsilon}} \\ \leq C(\varepsilon, \|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}).$$

Hence, we may multiply (3.45) by $|\ln \det \mathbb{B}_n|^{\frac{5-3\varepsilon}{4-2\varepsilon}-2} \ln \det \mathbb{B}_n$, and integrate to get

$$\frac{4-2\varepsilon}{5-3\varepsilon} \int_{\Omega} |\ln \det \mathbb{B}_n|^{\frac{5-3\varepsilon}{4-2\varepsilon}}(t) dx \leq \frac{4-2\varepsilon}{5-3\varepsilon} \int_{\Omega} |\ln \det \mathbb{B}_0|^{\frac{5-3\varepsilon}{4-2\varepsilon}} dx + \int_0^t \int_{\Omega} |\ln \det \mathbb{B}_n|^{\frac{1-\varepsilon}{4-2\varepsilon}} \delta(\theta_n) |\operatorname{tr}(\mathbb{B}_n - \mathbb{I})| dx d\tau.$$

Due to the Hölder inequality, (3.47) and the bounds (3.24), (3.43) the above implies

$$(3.48) \quad \|\ln \det \mathbb{B}_n\|_{L_t^\infty L_x^{\frac{5-3\varepsilon}{4-2\varepsilon}}} \lesssim \|\ln \det \mathbb{B}_0\|_\infty + \|\ln \det \mathbb{B}_n\|_1 \left(\int_0^T \int_{\Omega} |\delta(\theta_n) \operatorname{tr}(\mathbb{B}_n - \mathbb{I})|^{\frac{4-2\varepsilon}{3-\varepsilon}} dx d\tau \right)^{\frac{3-\varepsilon}{4-2\varepsilon}} \\ \leq \|\ln \det \mathbb{B}_0\|_\infty + (\|f(\mathbb{B}_n)\|_1 + \|\operatorname{tr}(\mathbb{B}_n) - 2\|_1) \left(\int_0^T \int_{\Omega} |\delta(\theta_n) \operatorname{tr}(\mathbb{B}_n - \mathbb{I})|^{\frac{4-2\varepsilon}{3-\varepsilon}} dx d\tau \right)^{\frac{3-\varepsilon}{4-2\varepsilon}} \\ \leq C(\varepsilon, \|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}, \|\ln \det \mathbb{B}_0\|_{L_x^\infty}).$$

In fact, with the obtained bound we may apply the same reasoning and multiply the equation (3.45) by $|\ln \det \mathbb{B}_n|^{\frac{(5-3\varepsilon)(1-\varepsilon)}{(4-2\varepsilon)^2}-1} \ln \det \mathbb{B}_n$. In the end, by a recursive argument

$$\|\ln \det \mathbb{B}_n\|_{L_t^\infty L_x^\lambda} \leq C(\lambda, \|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}, \|\ln \det \mathbb{B}_0\|_{L_x^\infty}),$$

for any $\lambda < \frac{4-2\varepsilon}{3-\varepsilon}$, or easier

$$(3.49) \quad \|\ln \det \mathbb{B}_n\|_{L_t^\infty L_x^{\frac{4}{3}-\varepsilon}} \leq C(\varepsilon, \|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}, \|\ln \det \mathbb{B}_0\|_{L_x^\infty}),$$

for any small $\varepsilon > 0$. Thus, with the use of (3.43), (3.49)

$$(3.50) \quad \|f(\mathbb{B}_n)\|_{L_{t,x}^{\frac{4}{3}-\varepsilon}} \leq C(\varepsilon, \|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}, \|\ln \det \mathbb{B}_0\|_{L_x^\infty}),$$

for any small $\varepsilon > 0$. We switch our focus towards the compactness of the temperature. Notice that integrating (3.20) gives us

$$(3.51) \quad \int_0^T \int_{\Omega} |\nabla_x \ln(\theta_n)|^2 dx dt = \int_0^T \int_{\Omega} \frac{|\nabla_x \theta_n|^2}{\theta_n^2} dx dt \leq C(\|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}).$$

and

$$\int_{\Omega} \ln(\theta_n) \, dx \geq \int_{\Omega} \ln(\theta_0) \, dx - C(\|g'(\theta_0)f(\mathbb{B}_0)\|_1) \geq -C(\|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}).$$

By (3.24) we may estimate

$$\int_{\{x:\theta_n(t,x)>1\}} \ln(\theta_n(t)) \, dx \leq \int_{\{x:\theta_n(t)>1\}} \theta_n(t) - 1 \, dx \leq C(\|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}).$$

Hence,

$$(3.52) \quad \|\ln(\theta_n)\|_{L_t^\infty L_x^1} \leq C(\|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}),$$

and we may utilize Poincaré–Wirtinger inequality as well as (3.51) and (3.52) to deduce

$$(3.53) \quad \|\ln(\theta_n)\|_{L_t^2 W_x^{1,2}} \leq C(\|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}).$$

Hence, by the above, (3.50), and the Assumption (G5),

$$\|\eta_n\|_{L_{t,x}^{\frac{5}{4}}} \leq C(\|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}, \|\ln \det \mathbb{B}_0\|_{L_x^\infty}).$$

Now denote

$$q_n = \eta_n u_n - \kappa(\theta_n) \nabla_x (\ln(\theta_n)).$$

Then, by (3.53), (3.31), and Hölder's inequality

$$\|\ln(\theta_n) u_n\|_{L_{t,x}^{\frac{4}{3}}} \leq C(\|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}).$$

Similarly, by (3.43), (3.31), and Hölder's inequality

$$\|\operatorname{tr}(\mathbb{B}_n) u_n\|_{L_{t,x}^{\frac{4}{3}}} \leq C(\|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}),$$

and by (3.49), (3.32), Hölder's inequality

$$\|\ln \det \mathbb{B}_n u_n\|_{L_t^2 L_x^{\frac{5}{4}}} \leq C(\|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}, \|\ln \det \mathbb{B}_0\|_{L_x^\infty}).$$

Thus

$$(3.54) \quad \|\eta_n u_n\|_{L_{t,x}^{\frac{5}{4}}} \leq C(\|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}, \|\ln \det \mathbb{B}_0\|_{L_x^\infty}).$$

In particular, the bound above, together with (3.51) implies

$$\|q_n\|_{L_{t,x}^{\frac{5}{4}}} \leq C(\|u_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}, \|\ln(\theta_0)\|_{L_x^1}, \|f(\mathbb{B}_0)\|_{L_x^1}, \|\mathbb{F}_0\|_{L_x^2}, \|\ln \det \mathbb{B}_0\|_{L_x^\infty}).$$

On the other hand, we may utilize the equation (3.20) to deduce

$$\begin{aligned} \|\operatorname{div}_{t,x}(\eta_n, q_n)\|_{L^1_{t,x}} &= \|\partial_t \eta_n + \operatorname{div}_x q_n\|_{L^1_{t,x}} = \left\| \kappa(\theta_n) \frac{|\nabla_x \theta_n|^2}{\theta_n^2} + \nu(\theta_n) \frac{|Du_n|^2}{\theta_n} + \delta(\theta_n) \frac{|\mathbb{B}_n - \mathbb{I}|^2}{\theta_n} \right\|_{L^1_{t,x}} \\ &\leq C(\|u_0\|_{L^2_x}, \|\theta_0\|_{L^1_x}, \|\ln(\theta_0)\|_{L^1_x}, \|f(\mathbb{B}_0)\|_{L^1_x}). \end{aligned}$$

Thus, by Banach–Alaoglu theorem and the Sobolev embeddings, there exist some $\bar{\eta}, \bar{q} \in L^{\frac{5}{4}}_{t,x}$ such that

$$\begin{aligned} \eta_n &\rightharpoonup \bar{\eta} \quad \text{weakly in } L^{\frac{5}{4}}_{t,x} \\ q_n &\rightharpoonup \bar{q} \quad \text{weakly in } L^{\frac{5}{4}}_{t,x} \end{aligned}$$

$$\{\operatorname{div}_{t,x}(\eta_n, q_n)\}_{n \in \mathbb{N}} \text{ compact in } (W_0^{1,3}((0, T) \times \Omega))^*.$$

In a similar way, with the use of equation (3.22) and the bounds (3.43), (3.49) and the Sobolev embeddings, there exist $\overline{f(B)}, \overline{f(B)u} \in L^{\frac{5}{4}}_{t,x}$ such that

$$\begin{aligned} f(\mathbb{B}_n) &\rightharpoonup \overline{f(\mathbb{B})} \quad \text{weakly in } L^{\frac{5}{4}}_{t,x}, \\ f(\mathbb{B}_n)u_n &\rightharpoonup \overline{f(\mathbb{B})u} \quad \text{weakly in } L^{\frac{5}{4}}_{t,x}, \end{aligned}$$

$$\{\operatorname{div}_{t,x}(f(\mathbb{B}_n), f(\mathbb{B}_n)u_n)\}_{n \in \mathbb{N}} \text{ compact in } (W_0^{1,3}((0, T) \times \Omega))^*.$$

And due to the boundedness of $g'(w)$, there exists $\overline{g'(\theta)f(\mathbb{B})} \in L^{\frac{5}{4}}_{t,x}$, such that

$$g'(\theta_n)f(\mathbb{B}_n) \rightharpoonup \overline{g'(\theta)f(\mathbb{B})} \quad \text{weakly in } L^{\frac{5}{4}}_{t,x}.$$

Now consider the sequence $\{\theta_n^{\frac{1}{3}}\}_{n \in \mathbb{N}}$. By (3.41)

$$\|\theta_n^{\frac{1}{3}}\|_{L^{5+\varepsilon}_{t,x}} \leq C(\varepsilon, \|u_0\|_{L^2_x}, \|\theta_0\|_{L^1_x}, \|\ln(\theta_0)\|_{L^1_x}, \|f(\mathbb{B}_0)\|_{L^1_x}, \|\mathbb{F}_0\|_{L^2_x}), \quad \varepsilon \in (0, 1),$$

and by (3.37)

$$\|\nabla_x(\theta_n^{\frac{1}{3}})\|_{L^2_{t,x}} = \frac{1}{3}\|\theta_n^{-\frac{2}{3}}\nabla_x \theta_n\|_{L^2_{t,x}} \leq C(\|u_0\|_{L^2_x}, \|\theta_0\|_{L^1_x}, \|\ln(\theta_0)\|_{L^1_x}, \|f(\mathbb{B}_0)\|_{L^1_x}, \|\mathbb{F}_0\|_{L^2_x}).$$

Thus

$$\|\operatorname{curl}_{t,x}(\theta_n^{\frac{1}{3}}, 0, 0)\|_{L^2_{t,x}} = \|\nabla_x^\perp(\theta_n^{\frac{1}{3}})\|_{L^2_{t,x}} \leq C(\|u_0\|_{L^2_x}, \|\theta_0\|_{L^1_x}, \|\ln(\theta_0)\|_{L^1_x}, \|f(\mathbb{B}_0)\|_{L^1_x}, \|\mathbb{F}_0\|_{L^2_x}).$$

In particular, by the Banach–Alaoglu theorem and the Sobolev embeddings, there exists some $\overline{\theta^{\frac{1}{3}}} \in L_{t,x}^{5+\varepsilon}$ such that

$$\begin{aligned} \theta_n^{\frac{1}{3}} &\rightharpoonup \overline{\theta^{\frac{1}{3}}} \quad \text{weakly in } L_{t,x}^{5+\varepsilon}, \\ \{\text{curl}_{t,x}(\theta_n^{\frac{1}{3}}, 0, 0)\}_{n \in \mathbb{N}} &\text{ compact in } (W_0^{1,2}((0, T) \times \Omega))^*. \end{aligned}$$

Hence, by the div-curl lemma A.2

$$(3.55) \quad \eta_n \theta_n^{\frac{1}{3}} \rightharpoonup \overline{\eta_n \theta^{\frac{1}{3}}} \quad \text{weakly in } L_{t,x}^1,$$

$$(3.56) \quad f(\mathbb{B}_n) \theta_n^{\frac{1}{3}} \rightharpoonup \overline{f(\mathbb{B}) \theta^{\frac{1}{3}}} \quad \text{weakly in } L_{t,x}^1.$$

Here, let us note that $w \mapsto -g'(w)$ and $w \mapsto w^{\frac{1}{3}}$ are increasing functions. Thus, for any $w \in \mathbb{R}$

$$0 \leq f(\mathbb{B}_n)(-g'(\theta_n) + g'(w))(\theta_n^{\frac{1}{3}} - w^{\frac{1}{3}})$$

In particular, for $w := \left(\overline{\theta^{\frac{1}{3}}}\right)^3$ we obtain

$$0 \leq f(\mathbb{B}_n) \left(-g'(\theta_n) + g' \left(\left(\overline{\theta^{\frac{1}{3}}} \right)^3 \right) \right) \left(\theta_n^{\frac{1}{3}} - \overline{\theta^{\frac{1}{3}}} \right).$$

Thus,

$$-f(\mathbb{B}_n)g'(\theta_n)\overline{\theta^{\frac{1}{3}}} \leq -f(\mathbb{B}_n)g'(\theta_n)\theta_n^{\frac{1}{3}} + f(\mathbb{B}_n)g' \left(\left(\overline{\theta^{\frac{1}{3}}} \right)^3 \right) \left(\theta_n^{\frac{1}{3}} - \overline{\theta^{\frac{1}{3}}} \right),$$

and due to (3.56) we may converge in integral formulation for the inequality above, to get

$$(3.57) \quad \int_0^T \int_{\Omega} -\overline{g'(\theta)f(\mathbb{B})\theta^{\frac{1}{3}}} dx dt \leq \int_0^T \int_{\Omega} -\overline{g'(\theta)f(\mathbb{B})\theta^{\frac{1}{3}}} dx dt.$$

Having this, we may apply (3.55), and (3.57) to get

$$\begin{aligned} (3.58) \quad \int_0^T \int_{\Omega} \overline{\ln(\theta)\theta^{\frac{1}{3}}} dx dt &= \int_0^T \int_{\Omega} \overline{\eta\theta^{\frac{1}{3}}} dx dt + \int_0^T \int_{\Omega} \overline{g'(\theta)f(\mathbb{B})\theta^{\frac{1}{3}}} dx dt \\ &= \lim_{n \rightarrow +\infty} \int_0^T \int_{\Omega} \eta_n \theta_n^{\frac{1}{3}} dx dt + \int_0^T \int_{\Omega} \overline{g'(\theta)f(\mathbb{B})\theta^{\frac{1}{3}}} dx dt \\ &= \int_0^T \int_{\Omega} \overline{\ln(\theta)\theta^{\frac{1}{3}}} dx dt - \int_0^T \int_{\Omega} \overline{g'(\theta)f(\mathbb{B})\theta^{\frac{1}{3}}} dx dt + \int_0^T \int_{\Omega} \overline{g'(\theta)f(\mathbb{B})\theta^{\frac{1}{3}}} dx dt \\ &\geq \int_0^T \int_{\Omega} \overline{\ln(\theta)\theta^{\frac{1}{3}}} dx dt. \end{aligned}$$

Now, notice that $w \mapsto \ln(w)$ and $w \mapsto w^{\frac{1}{3}}$ are increasing functions. Thus, for any $w \in L^1((0, T) \times \Omega; \mathbb{R}_+)$, such that $\ln(w) \in L^2_{t,x}$

$$\int_0^T \int_{\Omega} (\ln(\theta_n) - \ln(w))(\theta_n^{\frac{1}{3}} - w^{\frac{1}{3}}) dx dt \geq 0.$$

Due to (3.58) we may converge in the above, and deduce

$$(3.59) \quad \int_0^T \int_{\Omega} (\overline{\ln(\theta)} - \ln(w))(\overline{\theta^{\frac{1}{3}}} - w^{\frac{1}{3}}) dx dt \geq 0.$$

Consider

$$w := e^{\overline{\ln \theta} - \lambda h}, \quad \lambda > 0, h \in L^{\infty}_{t,x} \text{ - arbitrary.}$$

Indeed, since $h \in L^{\infty}_{t,x}$, then $e^{-\lambda h} \in L^{\infty}_{t,x}$, and by the weak lower-semicontinuity of a convex function

$$\phi \mapsto \int_0^T \int_{\Omega} e^{\phi} dx dt$$

we can deduce

$$\|\exp(\overline{\ln \theta})\|_{L^1_{t,x}} \leq \liminf_{n \rightarrow \infty} \|\exp(\ln \theta_n)\|_{L^1_{t,x}} \leq C(\|u_0\|_{L^2_x}, \|\theta_0\|_{L^1_x}, \|f(\mathbb{B}_0)\|_{L^1_x}).$$

Thus, w is an admissible function and we obtain

$$\int_0^T \int_{\Omega} (\overline{\theta^{\frac{1}{3}}} - e^{\frac{1}{3}\overline{\ln \theta} - \lambda h})h dx dt \geq 0.$$

After converging with $\lambda \rightarrow 0^+$ and using the arbitrariness of h we get

$$\overline{e^{\frac{1}{3}\ln \theta}} = \overline{\theta^{\frac{1}{3}}} = e^{\frac{1}{3}\overline{\ln \theta}} \text{ a.e. in } (0, T) \times \Omega.$$

Now let us put $w := \theta_n$ in (3.59). Then, again by (3.58),

$$\lim_{n \rightarrow +\infty} \int_0^T \int_{\Omega} |(\overline{\ln(\theta)} - \ln(\theta_n))(\overline{\theta^{\frac{1}{3}}} - \theta_n^{\frac{1}{3}})| dx dt = \lim_{n \rightarrow +\infty} \int_0^T \int_{\Omega} (\overline{\ln(\theta)} - \ln(\theta_n))(\overline{\theta^{\frac{1}{3}}} - \theta_n^{\frac{1}{3}}) dx dt = 0.$$

Hence, up to the subsequence,

$$(\overline{\ln(\theta)} - \ln(\theta_n))(e^{\frac{1}{3}\overline{\ln \theta}} - e^{\frac{1}{3}\ln \theta_n}) = (\overline{\ln(\theta)} - \ln(\theta_n))(\overline{\theta^{\frac{1}{3}}} - \theta_n^{\frac{1}{3}}) \rightarrow 0 \text{ a.e. in } (0, T) \times \Omega.$$

Since $x \mapsto e^{\frac{1}{3}x}$ is a strictly increasing function, the above is only possible if

$$\ln(\theta_n) \rightarrow \overline{\ln \theta} \text{ a.e. in } (0, T) \times \Omega,$$

which in turn implies that

$$(3.60) \quad \theta_n \rightarrow \theta := e^{\overline{\ln \theta}} \text{ a.e. in } (0, T) \times \Omega.$$

Due to Banach–Alaoglu’s theorem, Lebesgue’s dominated convergence theorem, Aubin–Lions’ lemma A.1 and (3.24), (3.33), (3.29) – (3.32), (3.35), (3.27), (3.36), (3.41), (3.42), (3.60)

$$\begin{aligned}
(3.61) \quad & u_n \overset{*}{\rightharpoonup} u && \text{weakly}^* \text{ in } L_t^\infty L_x^2, \\
(3.62) \quad & u_n \rightharpoonup u && \text{weakly in } L_t^2 W_{0,x}^{1,2}, \\
(3.63) \quad & u_n \rightharpoonup u && \text{weakly in } L_{t,x}^4, \\
(3.64) \quad & u_n \rightarrow u && \text{strongly in } C([0, T]; L^p(\Omega)), 1 \leq p < 4, \\
(3.65) \quad & \mathbb{F}_n \overset{*}{\rightharpoonup} \mathbb{F} && \text{weakly}^* \text{ in } L_t^\infty L_x^2, \\
(3.66) \quad & \mathbb{F}_n \rightharpoonup \mathbb{F} && \text{weakly in } L_{t,x}^4, \\
(3.67) \quad & \nabla_x u_n \mathbb{F}_n \rightharpoonup \overline{\nabla_x u \mathbb{F}} && \text{weakly in } L_{t,x}^{\frac{4}{3}}, \\
(3.68) \quad & \mathbb{F}_n \mathbb{F}_n^T \rightharpoonup \overline{\mathbb{F} \mathbb{F}^T} && \text{weakly in } L_{t,x}^2, \\
(3.69) \quad & |\mathbb{F}_n|^2 \rightharpoonup \overline{|\mathbb{F}|^2} && \text{weakly in } L_{t,x}^2, \\
(3.70) \quad & \mathbb{F}_n \mathbb{F}_n^T \mathbb{F}_n \rightharpoonup \overline{\mathbb{F} \mathbb{F}^T \mathbb{F}} && \text{weakly in } L_{t,x}^{\frac{4}{3}}, \\
(3.71) \quad & |\mathbb{F}_n \mathbb{F}_n^T|^2 \overset{*}{\rightharpoonup} \overline{|\mathbb{F} \mathbb{F}^T|^2} && \text{weakly}^* \text{ in } \mathcal{M}([0, T] \times \overline{\Omega}), \\
(3.72) \quad & \nabla_x u_n \mathbb{F}_n \mathbb{F}_n^T \overset{*}{\rightharpoonup} \overline{\nabla_x u \mathbb{F} \mathbb{F}^T} && \text{weakly}^* \text{ in } \mathcal{M}([0, T] \times \overline{\Omega}), \\
(3.73) \quad & |Du_n|^2 \overset{*}{\rightharpoonup} \overline{|Du|^2} && \text{weakly}^* \text{ in } \mathcal{M}([0, T] \times \overline{\Omega}), \\
(3.74) \quad & g(\theta_n) \mathbb{F}_n \mathbb{F}_n^T \rightharpoonup \overline{g(\theta) \mathbb{F} \mathbb{F}^T} && \text{weakly in } L_{t,x}^2, \\
(3.75) \quad & \delta(\theta_n) |\mathbb{F}_n|^2 \rightharpoonup \overline{\delta(\theta) |\mathbb{F}|^2} && \text{weakly in } L_{t,x}^{\frac{4}{3}-\varepsilon}, \\
(3.76) \quad & \delta(\theta_n) \mathbb{F}_n \mathbb{F}_n^T \mathbb{F}_n \rightharpoonup \overline{\delta(\theta) \mathbb{F} \mathbb{F}^T \mathbb{F}} && \text{weakly in } L_{t,x}^{\frac{14}{11}-\varepsilon}, \\
(3.77) \quad & \delta(\theta_n) |\mathbb{F}_n \mathbb{F}_n^T|^2 \overset{*}{\rightharpoonup} \overline{\delta(\theta) |\mathbb{F} \mathbb{F}^T|^2} && \text{weakly}^* \text{ in } \mathcal{M}([0, T] \times \overline{\Omega}), \\
(3.78) \quad & \nu(\theta_n) |Du_n|^2 \overset{*}{\rightharpoonup} \overline{\nu(\theta) |Du|^2} && \text{weakly}^* \text{ in } \mathcal{M}([0, T] \times \overline{\Omega}), \\
(3.79) \quad & \theta_n \rightarrow \theta && \text{strongly in } L_{t,x}^{2-\varepsilon}, \\
(3.80) \quad & \delta(\theta_n) \rightarrow \delta(\theta) && \text{strongly in } L_{t,x}^{2-\varepsilon}, \\
(3.81) \quad & g(\theta_n) \overset{*}{\rightharpoonup} g(\theta) && \text{weakly}^* \text{ in } L_{t,x}^\infty, \\
(3.82) \quad & \nu(\theta_n) \overset{*}{\rightharpoonup} \nu(\theta) && \text{weakly}^* \text{ in } L_{t,x}^\infty,
\end{aligned}$$

where we have used the notation that $\overline{h(u)}$ is the limit of the sequence $h(u_n)$, for any function h . Actually, due to Egorov's theorem and (3.60) we may identify

$$\begin{aligned}\overline{g(\theta)\mathbb{F}\mathbb{F}^T} &= g(\theta)\overline{\mathbb{F}\mathbb{F}^T}, \\ \overline{\delta(\theta)|\mathbb{F}|^2} &= \delta(\theta)\overline{|\mathbb{F}|^2}, \\ \overline{\delta(\theta)\mathbb{F}\mathbb{F}^T\mathbb{F}} &= \delta(\theta)\overline{\mathbb{F}\mathbb{F}^T\mathbb{F}},\end{aligned}$$

almost everywhere in $(0, T) \times \Omega$, and

$$\begin{aligned}\overline{\delta(\theta)|\mathbb{F}\mathbb{F}^T|^2} &= \delta(\theta)\overline{|\mathbb{F}\mathbb{F}^T|^2}, \\ \overline{\nu(\theta)|Du|^2} &= \nu(\theta)\overline{|Du|^2},\end{aligned}$$

as measures. Hence, we may integrate (3.1)₁, (3.1)₃ and converge with $n \rightarrow \infty$, to obtain

(3.83)

$$\int_0^T \int_{\Omega} -u \cdot \partial_t \phi - u \otimes u : \nabla_x \phi + \nu(\theta) Du : \nabla_x \phi + g(\theta) \overline{\mathbb{F}\mathbb{F}^T} : \nabla_x \phi \, dx \, dt = \int_{\Omega} u_0(x) \cdot \phi(0, x) \, dx,$$

for any $\phi \in C_c^1([0, T] \times \Omega; \mathbb{R}^2)$ with $\operatorname{div}_x \phi = 0$,

(3.84)

$$\int_0^T \int_{\Omega} -\mathbb{F} : \partial_t \phi - \mathbb{F} \otimes u : \nabla_x \phi - \overline{\nabla_x u \mathbb{F}} : u + \frac{1}{2} \delta(\theta) (\overline{\mathbb{F}\mathbb{F}^T\mathbb{F}} - \mathbb{F}) : \phi \, dx \, dt = \int_{\Omega} \mathbb{F}_0 : \phi(0, x) \, dx,$$

for any $\phi \in C_c^1([0, T] \times \Omega; \mathbb{R}^{2 \times 2})$. Now, we will closely follow [6, Subsection 6.4], to show

$$(3.85) \quad \mathbb{F}_n \rightarrow \mathbb{F} \text{ strongly in } L^2((0, T) \times \Omega).$$

As the proof is extremely similar ([6] works with the equations (3.83) - (3.84) for g, ν, δ constant), we will skip the details and briefly explain the differences in the approach. [6] states that it is enough to show that

$$(3.86) \quad \int_0^T \int_{\Omega} -(\overline{|\mathbb{F}|^2} - |\mathbb{F}|^2) \partial_t \phi - u(\overline{|\mathbb{F}|^2} - |\mathbb{F}|^2) \nabla_x \phi \, dx \, dt \leq \int_0^T \int_{\Omega} L(\overline{|\mathbb{F}|^2} - |\mathbb{F}|^2) \phi \, dx \, dt,$$

for any $\phi \in C_c^\infty((-\infty, T) \times \Omega)$, $\phi \geq 0$, and some $L \in L^2((0, T) \times \Omega)$. In fact, (3.86) concludes the argument via renormalization and the inequality dependent on $\|L\|_{L^1}$ (cf. [6, (6.79)] and [6, (6.86)]), thus it is enough to assume $L \in L^1((0, T) \times \Omega)$ and $L(\overline{|\mathbb{F}|^2} - |\mathbb{F}|^2) \in L^1((0, T) \times \Omega)$, which is what we will be able to obtain. To show (3.86) itself, one needs to be able to test (3.1)₃ by $\mathbb{F}_n \phi$ and (3.84) by $\mathbb{F} \phi$ and then converge with $n \rightarrow +\infty$ (cf. [6, (6.57)], [6, (6.58)]). The first case is straightforward,

as we work with a strong formulation. For (3.84) one needs to be more careful than in [6], since $\delta(\theta)\overline{\mathbb{F}\mathbb{F}^T\mathbb{F}}$ does not belong to $L^{\frac{4}{3}}$, and therefore we cannot test by functions which are only in L^4 . To remedy this, we may choose a slightly more sophisticated function in the form of the properly mollified

$$\frac{\mathbb{F}}{1 + \varepsilon|\mathbb{F}|}$$

which is in L^∞ . Then, due to (3.35) we can see that $\delta(\theta)\overline{\mathbb{F}\mathbb{F}^T\mathbb{F}} : \mathbb{F} \in L^1(\Omega)$ and converge with $\varepsilon \rightarrow 0^+$ using Lebesgue's dominated convergence theorem to obtain what we need (cf. [6, (6.58)]). In the end, following line by line the proof in [6] we are able to obtain (3.84) with

$$L := 1 + 2\nabla_x u + 2|\mathbb{F}|^2 + \delta(\theta),$$

which, as mentioned above, is enough to deduce (3.85). The convergence (3.85) is enough to conclude (3.16), (3.17) from (3.83) and (3.84). To show (3.18) notice that (3.85) implies that up to the subsequence

$$(3.87) \quad \mathbb{F}_n \rightarrow \mathbb{F}, \quad \text{a.e. in } (0, T) \times \Omega.$$

Thus, (3.43), (3.49), (3.87), (3.53), (3.51), (3.60), used with Banach–Alaoglu theorem, Vitali's convergence theorem and interpolation, imply that

$$\begin{aligned} \ln(\theta_n) &\rightharpoonup \ln(\theta) && \text{weakly in } L_t^2 W_x^{1,2}, \\ \ln(\theta_n) &\rightarrow \ln(\theta) && \text{strongly in } L_{t,x}^{3-\varepsilon}, \varepsilon \in (0, 1), \\ \text{tr}(\mathbb{F}_n \mathbb{F}_n^T) &\rightharpoonup \text{tr}(\mathbb{F} \mathbb{F}^T) && \text{weakly in } L_{t,x}^2, \\ \ln \det(\mathbb{F}_n \mathbb{F}_n^T) &\overset{*}{\rightharpoonup} \ln \det(\mathbb{F} \mathbb{F}^T) && \text{weakly}^* \text{ in } L_t^\infty L_x^{\frac{4}{3}-\varepsilon}, \varepsilon \in \left(0, \frac{1}{3}\right), \end{aligned}$$

and, since $g'(w)$ is a bounded function,

$$\begin{aligned} g'(\theta_n)\text{tr}(\mathbb{F}_n \mathbb{F}_n^T) &\rightharpoonup g'(\theta)\text{tr}(\mathbb{F} \mathbb{F}^T) && \text{weakly in } L_{t,x}^2, \\ g'(\theta_n)\ln \det(\mathbb{F}_n \mathbb{F}_n^T) &\overset{*}{\rightharpoonup} g'(\theta)\ln \det(\mathbb{F} \mathbb{F}^T) && \text{weakly}^* \text{ in } L_t^\infty L_x^{\frac{4}{3}-\varepsilon}, \varepsilon \in \left(0, \frac{1}{3}\right). \end{aligned}$$

At last, to conclude (3.18) we notice that the strong convergence of u_n (3.64), together with boundedness of $g'(w)$, (3.60), (3.87) and (3.55) implies

$$u_n(\ln(\theta_n) - g'(\theta_n)f(\mathbb{F}_n \mathbb{F}_n^T)) \rightharpoonup u(\ln(\theta) - g'(\theta)f(\mathbb{F} \mathbb{F}^T)) \text{ weakly in } L_{t,x}^{\frac{5}{4}},$$

which is enough to converge in the integrated form of (3.20). The last thing we need to show is that indeed $\theta > 0$ and $\det \mathbb{F} > 0$. Notice that by (3.60) and (3.87)

$$(3.88) \quad \theta \geq 0, \quad \det \mathbb{F} \geq 0, \quad \text{a.e. in } (0, T) \times \Omega.$$

On the other hand, Fatou's lemma together with (3.52), (3.49) implies

$$\|\ln \theta\|_{L_t^\infty L_x^1} + \|\ln \det(\mathbb{F} \mathbb{F}^T)\|_{L_t^\infty L_x^1} \leq C,$$

which, together with (3.88), is enough to deduce

$$\theta > 0, \quad \det \mathbb{F} > 0.$$

□

4. EXISTENCE OF THE WEAK SOLUTIONS FOR THE CASE (P1)

Let us state the main result of this section.

Theorem 4.1. *Suppose that we are in the situation (P1) and κ, ν, δ satisfy the Assumptions (2.1) - (2.3). Then, there exists a weak solution to the system (1.5), (1.9). That is, there exists a triple (u, \mathbb{F}, θ) , such that $u \in L_t^\infty L_x^2 \cap L_t^2 W_{0,x}^{1,2}$, $\mathbb{F} \in L_t^\infty L_x^2 \cap L_{t,x}^4$, $\theta \in L_{t,x}^{2-\varepsilon} \cap L^{\frac{4}{3}-3\varepsilon} W_x^{1, \frac{4}{3}-3\varepsilon}$ ($\varepsilon \in (0, 1)$), and they satisfy the integral equalities*

$$(4.1) \quad \int_0^T \int_\Omega -u \cdot \partial_t \phi - u \otimes u : \nabla_x \phi + \nu(\theta) Du : \nabla_x \phi + \mathbb{F} \mathbb{F}^T : \nabla_x \phi \, dx \, dt = \int_\Omega u_0(x) \cdot \phi(0, x) \, dx,$$

for any $\phi \in C_c^1([0, T) \times \Omega; \mathbb{R}^2)$ with $\operatorname{div}_x \phi = 0$,

$$(4.2) \quad \int_0^T \int_\Omega -\mathbb{F} : \partial_t \phi - \mathbb{F} \otimes u : \nabla_x \phi - \nabla_x u \mathbb{F} : \phi + \frac{1}{2} \delta(\theta) (\mathbb{F} \mathbb{F}^T \mathbb{F} - \mathbb{F}) : \phi \, dx \, dt = \int_\Omega \mathbb{F}_0(x) : \phi(0, x) \, dx,$$

for any $\phi \in C_c^1([0, T) \times \Omega; \mathbb{R}^{2 \times 2})$,

$$(4.3) \quad - \int_0^T \int_\Omega \theta \partial_t \phi \, dx \, dt + \int_0^T \int_\Omega \theta u \cdot \nabla_x \phi \, dx \, dt + \int_0^T \int_\Omega \kappa(\theta) \nabla_x \theta \cdot \nabla_x \phi \, dx \, dt \\ - \int_0^T \int_\Omega \nu(\theta) |Du|^2 \phi \, dx \, dt - \int_0^T \int_\Omega |\mathbb{F} \mathbb{F}^T - \mathbb{I}|^2 \phi \, dx \, dt = \int_\Omega \theta_0(x) \phi(0, x) \, dx$$

for any $\phi \in C_c^1([0, T) \times \overline{\Omega})$. Moreover $\theta > 0$ and $\det \mathbb{F} > 0$.

Since the proof of the Theorem 4.1 is very similar to the one conducted in [1, 6], and does not include many new techniques, when compared to the proof of the Theorem 3.1 we will skip most of the technical detail and simply focus on the differences in the approach.

4.1. Approximating scheme. To prove our main theorem we will use a four-step approximation scheme. Our first goal will be to prove the existence of a solution to the system

$$\begin{cases} \partial_t u + \operatorname{div}_x(u \otimes u) - \operatorname{div}_x \mathbb{T} = 0, \\ \operatorname{div}_x u = 0, \\ \partial_t \mathbb{F} + \operatorname{div}_x(\mathbb{F} \otimes u) - \nabla_x u \mathbb{F} + \frac{1}{2} \delta(\theta)(\mathbb{F} \mathbb{F}^T - \mathbb{F}) = \varepsilon \Delta \mathbb{F}, \\ -p \mathbb{I} + \nu(\theta) \mathbb{D}u + (\mathbb{F} \mathbb{F}^T - \mathbb{I}) = \mathbb{T}, \\ \partial_t \theta + u \cdot \nabla_x \theta - \operatorname{div}_x(\kappa(\theta) \nabla_x \theta) = \nu(\theta) |Du|^2 + |\mathbb{F} \mathbb{F}^T - \mathbb{I}|^2. \end{cases}$$

for a fixed $\varepsilon \in (0, 1)$. We also set the initial and boundary conditions as

$$(4.4) \quad u|_{\partial\Omega} = 0, \quad u(0, x) = u_0(x), \quad \mathbb{F}(0, x) = \mathbb{F}_0, \quad \nabla_x \mathbb{F} \cdot \mathbf{n} = 0, \quad \nabla_x \theta \cdot \mathbf{n} = 0, \quad \theta(0, x) = \theta_0^r,$$

for $r > 0$ and

$$\theta_0^r(x) = \begin{cases} \theta_0(x), & \text{whenever } r \leq \theta_0(x) \leq r^{-1}, \\ 1, & \text{otherwise.} \end{cases}$$

To do so we will employ the Galerkin approximation, but most importantly we shall split the convergences of the linear approximation of θ and u , \mathbb{F} , as Galerkin systems allow only to test by the linear functions of the variables, and as can be seen by our considerations in Lemma 3.2, the expected bounds on the temperature are obtained via testing by a non-linear function.

4.2. Galerkin system. Fix s big enough, so that

$$(4.5) \quad W^{s-1,2}(\Omega) \hookrightarrow L^\infty(\Omega)$$

and let $\{\omega_n\}_{n \in \mathbb{N}}$, $\{A_n\}_{n \in \mathbb{N}}$, $\{\phi_m\}_{m \in \mathbb{N}}$ be the orthogonal bases of $W_{0,\operatorname{div}}^{s,2}(\Omega; \mathbb{R}^2)$, $W^{s,2}(\Omega; \mathbb{R}^{2 \times 2})$, $W^{s,2}(\Omega; \mathbb{R})$ respectively, which are also orthonormal in L^2 , and whose projections are continuous.

Set

$$(4.6) \quad u_{nm}(t, x) = \sum_{i=1}^n \alpha_i(t) \omega_i(x), \quad \mathbb{F}_{nm}(t, x) = \sum_{i=1}^n \beta_i(t) A_i(x), \quad \theta_{nm}(t, x) = \sum_{i=1}^m \gamma_i(t) \phi_i(x),$$

which satisfy the equations for any $j = 1, \dots, n$ and $i = 1, \dots, m$

$$(4.7) \quad \frac{d}{dt} \int_{\Omega} u_{nm} \cdot \omega_j \, dx - \int_{\Omega} u_{nm} \otimes u_{nm} : \nabla_x \omega_j \, dx \\ + \int_{\Omega} \nu(\theta_{nm}) Du_{nm} : \nabla_x \omega_j \, dx + \int_{\Omega} \mathbb{F}_{nm} \mathbb{F}_{nm}^T : \nabla_x \omega_j \, dx = 0,$$

$$(4.8) \quad \frac{d}{dt} \int_{\Omega} \mathbb{F}_{nm} : A_j \, dx - \int_{\Omega} \mathbb{F}_{nm} \otimes u_{nm} : \nabla_x A_j \, dx - \int_{\Omega} (\nabla_x u_{nm} \mathbb{F}_{nm}) : A_j \, dx \\ + \frac{1}{2} \int_{\Omega} \delta(\theta_{nm}) \mathbb{F}_{nm} \mathbb{F}_{nm}^T \mathbb{F}_{nm} : A_j \, dx - \frac{1}{2} \int_{\Omega} \delta(\theta_{nm}) \mathbb{F}_{nm} : A_j \, dx + \varepsilon \int_{\Omega} \nabla_x \mathbb{F}_{nm} : \nabla_x A_j \, dx = 0,$$

$$(4.9) \quad \frac{d}{dt} \int_{\Omega} \theta_{nm} \phi_i \, dx + \int_{\Omega} u_{nm} \cdot \nabla_x \theta_{nm} \phi_i \, dx + \int_{\Omega} \kappa(\theta_{nm}) \nabla_x \theta_{nm} \cdot \nabla_x \phi_i \, dx \\ - \int_{\Omega} \nu(\theta_{nm}) |Du_{nm}|^2 \phi_i \, dx - \int_{\Omega} |\mathbb{F}_{nm} \mathbb{F}_{nm}^T - \mathbb{I}|^2 \phi_i \, dx = 0.$$

4.3. Convergence with $m \rightarrow +\infty$. Following the argumentation in [6, Appendix B] (estimate (B.15)), by multiplying (4.7) and (4.8) by α_j and β_j respectively, after summing over j we get

(4.10)

$$\|u_{nm}\|_{L_t^\infty L_x^2} + \|\mathbb{F}_{nm}\|_{L_t^\infty L_x^2} + \|Du_{nm}\|_{L_{t,x}^2} + \|\mathbb{F}_{nm}\|_{L_{t,x}^4} + \varepsilon \|\nabla_x \mathbb{F}_{nm}\|_{L_{t,x}^2} \leq C(\|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2}),$$

which, by Korn's inequality, implies

(4.11)

$$\|u_{nm}\|_{L_t^\infty L_x^2} + \|\mathbb{F}_{nm}\|_{L_t^\infty L_x^2} + \|\nabla_x u_{nm}\|_{L_{t,x}^2} + \|\mathbb{F}_{nm}\|_{L_{t,x}^4} + \varepsilon \|\nabla_x \mathbb{F}_{nm}\|_{L_{t,x}^2} \leq C(\|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2}).$$

Hence, by the orthogonality of the bases $\{\omega_n\}_{n \in \mathbb{N}}$ and $\{A_n\}_{n \in \mathbb{N}}$, the embedding (4.5), and (4.11) we get

$$(4.12) \quad \|u_{nm}\|_{L_t^\infty W_x^{1,\infty}} \leq \|u_{nm}\|_{L_t^\infty W_x^{s,2}} \leq C(n) \|u_{nm}\|_{L_t^\infty L_x^2} \leq C(n, \|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2})$$

$$(4.13) \quad \|\mathbb{F}_{nm}\|_{L_t^\infty W_x^{1,\infty}} \leq \|\mathbb{F}_{nm}\|_{L_t^\infty W_x^{s,2}} \leq C(n) \|\mathbb{F}_{nm}\|_{L_t^\infty L_x^2} \leq C(n, \|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2})$$

Now, let us multiply (4.9) by $\gamma_j(t)$ and sum all of the equalities over j . Then, we obtain

$$\frac{1}{2} \frac{d}{dt} \|\theta_{nm}\|_{L_x^2} + \|\sqrt{\kappa(\theta_{nm})} \nabla_x \theta_{nm}\|_{L_x^2} = \int_{\Omega} \nu(\theta_{nm}) |Du_{nm}|^2 \theta_{nm} + \int_{\Omega} |\mathbb{F}_{nm} \mathbb{F}_{nm}^T - \mathbb{I}|^2 \theta_{nm}.$$

Thus, with the use of Hölder's inequality, (4.12) as well as the bounds (2.1), we arrive at

$$(4.14) \quad \frac{1}{2} \frac{d}{dt} \|\theta_{nm}\|_{L_x^2} + \sqrt{C^{-1}} \|\nabla_x \theta_{nm}\|_{L_x^2} \leq (\|u_{nm}\|_{L_t^\infty W_x^{1,\infty}} + \|\mathbb{F}_{nm}\|_{L_t^\infty L_x^\infty}) \|\theta_{nm}\|_{L_x^2},$$

which after applying Grönwall's inequality leads to

$$(4.15) \quad \|\theta_{nm}\|_{L_x^2}(t) \leq e^{C(n)t} \|\theta_0\|_{L_x^2}.$$

We would to deduce strong convergences as $m \rightarrow +\infty$, therefore we need the bounds on the time derivatives. We start with the velocity. Using the equation (4.7) and applying the bounds (4.12) together with Hölder's inequality, we get

$$\begin{aligned} \|(\alpha_j)'\|_{L_t^2} &= \|(\partial_t u_{nm}, \omega_j)\|_{L_t^2} \\ &= \|(u_{nm} \otimes u_{nm}, \nabla_x \omega_j) - (\nu(\theta_{nm}) D u_{nm}, \nabla_x \omega_j) - (\mathbb{F}_{nm} \mathbb{F}_{nm}^T, \nabla_x \omega_j)\|_{L_t^2} \\ &\leq C(n, \|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2}) \end{aligned}$$

Hence

$$(4.16) \quad \|\partial_t u_{nm}\|_{L_t^2 W_x^{1,\infty}} \leq C(n, \|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2}).$$

At the same time, with the help of the Fundamental Theorem of Calculus and Hölder's inequality, we may deduce

$$(4.17) \quad |\alpha_j(t) - \alpha_j(s)| \leq \int_s^t |(\alpha_j)'(\tau)| d\tau \lesssim |t - s|^{\frac{1}{2}}, \quad \text{for any } t, s, \in \mathbb{R}.$$

A similar argument can be made for \mathbb{F}_{nm} , and we arrive at

$$(4.18) \quad \|\partial_t \mathbb{F}_{nm}\|_{L_t^2 W_x^{1,\infty}} \leq C(n, \|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2})$$

and

$$(4.19) \quad |\beta_j(t) - \beta_j(s)| \lesssim |t - s|^{\frac{1}{2}}, \quad \text{for any } t, s, \in \mathbb{R}.$$

Thus, by the Banach–Alaoglu theorem and (4.16), as well as the Arzelà–Ascoli theorem and (4.17), we get that there exists $u_n \in C([0, T]; W^{1,\infty}(\Omega)) \cap L^2((0, T); W_{0,\text{div}}^{1,2}(\Omega))$, such that (up to the subsequence)

$$(4.20) \quad \partial_t u_{nm} \xrightarrow{*} \partial_t u_n \quad \text{weakly* in } L_t^2 W_x^{1,\infty},$$

$$(4.21) \quad u_{nm} \rightarrow u_n \quad \text{strongly in } C([0, T]; W^{1,\infty}(\Omega)).$$

By an analogous argument, we obtain the existence of $\mathbb{F}_n \in C([0, T]; W^{1, \infty}(\Omega))$, such that (up to the subsequence)

$$(4.22) \quad \partial_t \mathbb{F}_{nm} \xrightarrow{*} \partial_t \mathbb{F}_n \quad \text{weakly* in } L_t^2 W_x^{1, \infty},$$

$$(4.23) \quad \mathbb{F}_{nm} \rightarrow \mathbb{F}_n \quad \text{strongly in } C([0, T]; W^{1, \infty}(\Omega)).$$

As for the bounds on the time derivative of the temperature. We can use the orthogonality of $\{\phi_i\}$, the continuity of projection, and Hölder's inequality to get for any $f \in W^{1, 2}(\Omega)$

$$\begin{aligned} & \left| \int_{\Omega} \partial_t \theta_{nm} f \, dx \right| \leq \\ & \left| \int_{\Omega} u_{nm} \cdot \nabla_x \theta_{nm} f \, dx + \int_{\Omega} \kappa(\theta_{nm}) \nabla_x \theta_{nm} \cdot \nabla_x f \, dx \right| \\ & + \left| \int_{\Omega} \nu(\theta_{nm}) |Du_{nm}|^2 f \, dx + \int_{\Omega} |\mathbb{F}_{nm} \mathbb{F}_{nm}^T - \mathbb{I}|^2 f \, dx \right| \lesssim \\ & \|u_{nm}\|_{L_x^\infty} \|\theta_{nm}\|_{W_x^{1, 2}} \|f\|_{L_x^2} + \|\theta_{nm}\|_{W_x^{1, 2}} \|f\|_{W_x^{1, 2}} + \|u_{nm}\|_{W_x^{1, \infty}} \|f\|_{L_x^2} + \|\mathbb{F}\|_{L_x^\infty} \|f\|_{L_x^2}. \end{aligned}$$

Thus, integrating over time and using Hölder's, Minkowski's inequalities, and utilizing (4.12), (4.15), and (4.14) we obtain

$$(4.24) \quad \begin{aligned} \|\partial_t \theta_{nm}\|_{L^2((0, T); (W^{1, 2}(\Omega))^*)} & \lesssim \int_0^T \|u_{nm}\|_{L_x^\infty}^2 \|\theta_{nm}\|_{W_x^{1, 2}}^2 + \|\theta_{nm}\|_{W_x^{1, 2}}^2 + \|u_{nm}\|_{W_x^{1, \infty}}^2 + \|\mathbb{F}\|_{L_x^\infty}^2 \, dt \\ & \leq C(n, \|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2}, \|\theta_0\|_{L_x^2}) \end{aligned}$$

Hence, using the Banach–Alaoglu theorem and (4.14), (4.15), (4.24), we deduce

$$(4.25) \quad \partial_t \theta_{nm} \rightharpoonup \partial_t \theta_n \quad \text{weakly in } L^2((0, T); (W^{1, 2}(\Omega))^*),$$

$$(4.26) \quad \theta_{nm} \xrightarrow{*} \theta_n \quad \text{weakly* in } L_t^\infty L_x^2,$$

$$(4.27) \quad \theta_{nm} \rightharpoonup \theta_n \quad \text{weakly in } L_t^2 W_x^{1, 2}.$$

And by the Aubin–Lions lemma (A.1)

$$(4.28) \quad \theta_{nm} \rightarrow \theta_n \quad \text{strongly in } L_t^2 L_x^2.$$

Since in every product, at least one of the factors converges strongly, we can easily infer from the convergences (4.20) - (4.23) and (4.25) - (4.28) the equalities

$$(4.29) \quad \frac{d}{dt} \int_{\Omega} u_n \cdot \omega_j \, dx - \int_{\Omega} u_n \otimes u_n : \nabla_x \omega_j \, dx \\ + \int_{\Omega} \nu(\theta_n) Du_n : \nabla_x \omega_j \, dx + \int_{\Omega} \mathbb{F}_n \mathbb{F}_n^T : \nabla_x \omega_j \, dx = 0,$$

$$(4.30) \quad \frac{d}{dt} \int_{\Omega} \mathbb{F}_n : A_j \, dx - \int_{\Omega} \mathbb{F}_n \otimes u_n : \nabla_x A_j \, dx - \int_{\Omega} (\nabla_x u_n \mathbb{F}_n) : A_j \, dx \\ + \frac{1}{2} \int_{\Omega} \delta(\theta_n) \mathbb{F}_n \mathbb{F}_n^T \mathbb{F}_n : A_j \, dx - \frac{1}{2} \int_{\Omega} \delta(\theta_n) \mathbb{F}_n : A_j \, dx + \varepsilon \int_{\Omega} \nabla_x \mathbb{F}_n : \nabla_x A_j \, dx = 0,$$

for any $j = 1, \dots, n$, and for any $\phi \in W^{1,2}(\Omega; \mathbb{R})$

$$(4.31) \quad \langle \partial_t \theta_n, \phi \rangle + \int_{\Omega} u_n \cdot \nabla_x \theta_n \phi \, dx + \int_{\Omega} \kappa(\theta_n) \nabla_x \theta_n \cdot \nabla_x \phi \, dx \\ - \int_{\Omega} \nu(\theta_n) |Du_n|^2 \phi \, dx - \int_{\Omega} |\mathbb{F}_n \mathbb{F}_n^T - \mathbb{I}|^2 \phi \, dx = 0.$$

4.4. Convergence with $n \rightarrow +\infty$. For the variables u_n and \mathbb{F}_n we have already obtained the general bounds (4.11). For their time derivatives, bounds (4.18), (4.16) depend on n , therefore we must proceed differently. Here, again, similarly as in [6, Appendix B] (estimates (B.19) and (B.21)) we may obtain

$$(4.32) \quad \|\partial_t u_n\|_{L^2((0,T);(W_{0,\text{div}}^{1,2}(\Omega))^*)} + \|\partial_t \mathbb{F}_n\|_{L^{\frac{4}{3}}((0,T);(W^{1,2}(\Omega;\mathbb{R}^{2 \times 2}))^*)} \leq C(\|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2}).$$

As for the temperature, we would like to test (4.31) by $\theta_n^{\lambda-1}$ for some arbitrary $0 < \lambda < 1$ as was explained in Lemma 3.2. To do so properly, one needs to show that in fact $\theta_n \geq r$. It can be done in a similar manner as in [2], i.e. by testing (4.31) by the negative part of $\theta_n - r$. We get

$$(4.33) \quad \theta_n(x, t) \geq r, \quad \text{for a.e. } x \in \Omega \text{ and all } t \in (0, T).$$

We proceed by proving a simple $L_t^\infty L_x^1$ estimate. We may test (4.31) by a function $\phi \equiv 1$. Then, after integrating the equation in $(0, t)$, by Lemma A.3, the condition $\text{div}_x u_n = 0$, and the positiveness of θ_n (4.33), we get

$$\|\theta_n(t)\|_{L_x^1} = \int_0^t \int_{\Omega} |Du_n|^2 + |\mathbb{F}_n \mathbb{F}_n^T - \mathbb{I}|^2 \, dx \, d\tau + \|\theta_n(0)\|_{L_x^1}.$$

Thus, after applying the inequality (4.10), we arrive at the bound

$$(4.34) \quad \|\theta_n\|_{L_t^\infty L_x^1} \leq C(\|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}).$$

Now, we can finally test (4.31) by $\theta_n^{\lambda-1}$. Performing similar operations as in Lemma 3.2, we obtain

$$(4.35) \quad \int_0^t \int_\Omega \frac{|\nabla_x \theta_n|^2}{\theta_n^\lambda} dx dt \leq C(\lambda, \|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}), \quad \text{for any } \lambda \in (1, 2).$$

With this (similarly as under (3.37)), we may infer uniform bounds on θ_n and $\nabla_x \theta_n$

$$(4.36) \quad \|\theta_n\|_{L_{t,x}^{2-\lambda}} \leq C(\lambda, \|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}),$$

$$(4.37) \quad \|\nabla_x \theta_n\|_{L_{t,x}^{\frac{4}{3}-\lambda}} \leq C(\lambda, \|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}),$$

for any $\lambda \in (0, 1)$. Now we can estimate a time derivative of θ_n , to later use the Aubin-Lions lemma A.1. This is done similarly to (4.24), but by testing (4.31) with a function $\phi \in W^{s,2}(\Omega)$ (see (4.5)),

One simply gets

$$(4.38) \quad \|\partial_t \theta_n\|_{L^1((0,T);(W^{s,2}(\Omega))^*)} \leq C(\lambda, \|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}).$$

In consequence, we may apply the Banach–Alaoglu theorem, the Aubin–Lions lemma A.1 to (4.36), (4.37), (4.38) and obtain, that (up to the subsequence)

$$(4.39) \quad \theta_n \rightharpoonup \theta \quad \text{weakly in } L_{t,x}^{2-\lambda},$$

$$(4.40) \quad \theta_n \rightharpoonup \theta \quad \text{weakly in } L_t^{\frac{4}{3}-\lambda} W_x^{1, \frac{4}{3}-\lambda},$$

$$(4.41) \quad \theta_n \rightarrow \theta \quad \text{strongly in } L_t^{\frac{4}{3}-\lambda} L_x^{\frac{8-6\lambda}{2+3\lambda}}.$$

for all $\lambda > 0$, sufficiently close to 0. Similarly, due to (4.32) and (4.11)

$$(4.42) \quad u_n \rightharpoonup u \quad \text{weakly* in } L_t^\infty L_x^2,$$

$$(4.43) \quad u_n \rightharpoonup u \quad \text{weakly in } L_t^2 W_{0,x}^{1,2},$$

$$(4.44) \quad \partial_t u_n \rightharpoonup \partial_t u \quad \text{weakly in } L^2((0, T); (W_{0,\text{div}}^{1,2}(\Omega))^*),$$

$$(4.45) \quad u_n \rightarrow u \quad \text{strongly in } L_{t,x}^q, \text{ for } q \in [1, 4),$$

$$(4.46) \quad \mathbb{F}_n \rightharpoonup \mathbb{F} \quad \text{weakly* in } L_t^\infty L_x^2,$$

$$(4.47) \quad \mathbb{F}_n \rightharpoonup \mathbb{F} \quad \text{weakly in } L_t^2 W_x^{1,2},$$

$$(4.48) \quad \partial_t \mathbb{F}_n \rightharpoonup \partial_t \mathbb{F} \quad \text{weakly in } L^{\frac{4}{3}}((0, T); (W^{1,2}(\Omega; \mathbb{R}^{2 \times 2}))^*),$$

$$(4.49) \quad \mathbb{F}_n \rightarrow \mathbb{F} \quad \text{strongly in } L_{t,x}^q, \text{ for } q \in [1, 4).$$

The convergences (4.39) - (4.49) allow us to easily converge in all of the terms (4.29), (4.30), (4.31) except

$$\int_{\Omega} \nu(\theta_n) |Du_n|^2 \phi \, dx, \quad \int_{\Omega} |\mathbb{F}_n \mathbb{F}_n^T - \mathbb{I}|^2 \phi \, dx.$$

In particular, for the u and \mathbb{F} we obtain the equations

$$(4.50) \quad \langle \partial_t u, \omega \rangle - \int_{\Omega} u \otimes u : \nabla_x \omega \, dx + \int_{\Omega} \nu(\theta) Du : \nabla_x \omega \, dx + \int_{\Omega} \mathbb{F} \mathbb{F}^T : \nabla_x \omega \, dx = 0,$$

for any $\omega \in W_0^{1,2}(\Omega)$.

$$(4.51) \quad \langle \partial_t \mathbb{F}, A \rangle - \int_{\Omega} \mathbb{F} \otimes u : \nabla_x A \, dx - \int_{\Omega} (\nabla_x u \mathbb{F}) : A \, dx \\ + \frac{1}{2} \int_{\Omega} \delta(\theta) \mathbb{F} \mathbb{F}^T \mathbb{F} : A \, dx - \frac{1}{2} \int_{\Omega} \delta(\theta) \mathbb{F} : A \, dx + \varepsilon \int_{\Omega} \nabla_x \mathbb{F} : \nabla_x A \, dx = 0,$$

for any $A \in W^{1,2}(\Omega; \mathbb{R}^{2 \times 2})$. To deal with the problematic terms, we test (4.50) by u and integrate the equation in $(0, t)$, to obtain

$$(4.52) \quad \int_{\Omega} |u|^2(t) \, dx + \int_0^t \int_{\Omega} \nu(\theta) |Du|^2 \, dx \, d\tau + \int_0^t \int_{\Omega} \mathbb{F} \mathbb{F}^T : \nabla_x u \, dx \, d\tau = \int_{\Omega} |u_0|^2 \, dx.$$

In comparison, we test (4.29) by u_n (i.e. multiply the equation by α_j and sum over j), and integrate over $(0, t)$ to get

$$\int_{\Omega} |u_n(t)|^2 \, dx + \int_{\Omega} \nu(\theta_n) |Du_n|^2 \, dx + \int_{\Omega} \mathbb{F}_n \mathbb{F}_n^T : \nabla_x u_n \, dx = \int_{\Omega} |u_n(0)|^2 \, dx.$$

We may perform an analogous argument, testing (4.30) and (4.51) by \mathbb{F}_n and \mathbb{F} respectively. After adding the corresponding equations together, we may use (4.49) and Lemma A.4, to deduce

$$\begin{aligned} & \limsup_{n \rightarrow +\infty} \int_0^t \int_{\Omega} \nu(\theta_n) |Du_n|^2 dx d\tau + \int_0^t \int_{\Omega} |\mathbb{F}_n \mathbb{F}_n^T|^2 dx d\tau \\ & \leq \int_0^t \int_{\Omega} \nu(\theta) |Du|^2 dx d\tau + \int_0^t \int_{\Omega} |\mathbb{F} \mathbb{F}^T|^2 dx d\tau, \end{aligned}$$

which by weak lower semi-continuity implies

$$\begin{aligned} \lim_{n \rightarrow +\infty} \int_0^t \int_{\Omega} \nu(\theta_n) |Du_n|^2 dx d\tau + \int_0^t \int_{\Omega} |\mathbb{F}_n \mathbb{F}_n^T|^2 dx d\tau \\ = \int_0^t \int_{\Omega} \nu(\theta) |Du|^2 dx d\tau + \int_0^t \int_{\Omega} |\mathbb{F} \mathbb{F}^T|^2 dx d\tau. \end{aligned}$$

This is enough to conclude

$$\int_0^T \int_{\Omega} |\sqrt{\nu(\theta_n)} Du_n - \sqrt{\nu(\theta)} Du|^2 dx dt + \int_0^T \int_{\Omega} |\mathbb{F}_n \mathbb{F}_n^T - \mathbb{F} \mathbb{F}^T|^2 dx dt = 0.$$

Thus, we may converge from (4.31) properly and obtain the equality

$$(4.53) \quad \begin{aligned} & - \int_0^T \int_{\Omega} \theta \partial_t \phi dx dt + \int_0^T \int_{\Omega} \theta u \cdot \nabla_x \phi dx dt + \int_0^T \int_{\Omega} \kappa(\theta) \nabla_x \theta \cdot \nabla_x \phi dx dt \\ & - \int_0^T \int_{\Omega} \nu(\theta) |Du|^2 \phi dx dt - \int_0^T \int_{\Omega} |\mathbb{F} \mathbb{F}^T - \mathbb{I}|^2 \phi dx dt = \int_{\Omega} \theta_0(x) \phi(0, x) dx, \end{aligned}$$

for any $\phi \in C_c^1([0, T] \times \overline{\Omega})$.

4.5. Convergence with $r \rightarrow 0^+$. Let us denote by $(u_{r\varepsilon}, \mathbb{F}_{r\varepsilon}, \theta_{r\varepsilon})$ the solution of the approximating scheme, proven to exist in the previous subsections. Since all of the bounds (4.32), (4.11), (4.36), (4.37), (4.38) are uniform in r performing the same calculations as above we may obtain that (up to the subsequence)

$$(4.54) \quad \theta_{r\varepsilon} \rightharpoonup \theta_{\varepsilon} \quad \text{weakly in } L_{t,x}^{2-\lambda},$$

$$(4.55) \quad \theta_{r\varepsilon} \rightharpoonup \theta_{\varepsilon} \quad \text{weakly in } L_t^{\frac{4}{3}-\lambda} W_x^{1, \frac{4}{3}-\lambda},$$

$$(4.56) \quad \theta_{r\varepsilon} \rightarrow \theta_{\varepsilon} \quad \text{strongly in } L_t^{\frac{4}{3}-\lambda} L_x^{\frac{8-6\lambda}{2+3\lambda}}.$$

for all $\lambda > 0$, sufficiently close to 0, and

$$\begin{aligned}
u_{r\varepsilon} &\rightharpoonup u_\varepsilon && \text{weakly* in } L_t^\infty L_x^2, \\
u_{r\varepsilon} &\rightharpoonup u_\varepsilon && \text{weakly in } L_t^2 W_{0,x}^{1,2}, \\
\partial_t u_{r\varepsilon} &\rightharpoonup \partial_t u_\varepsilon && \text{weakly in } L^2((0, T); (W_{0,\text{div}}^{1,2}(\Omega))^*) \\
u_{r\varepsilon} &\rightarrow u_\varepsilon && \text{strongly in } L_{t,x}^q, \text{ for } q \in [1, 4), \\
\mathbb{F}_{r\varepsilon} &\rightharpoonup \mathbb{F}_\varepsilon && \text{weakly* in } L_t^\infty L_x^2, \\
\mathbb{F}_{r\varepsilon} &\rightharpoonup \mathbb{F}_\varepsilon && \text{weakly in } L_t^2 W_x^{1,2}, \\
\partial_t \mathbb{F}_{r\varepsilon} &\rightharpoonup \partial_t \mathbb{F}_\varepsilon && \text{weakly in } L^{\frac{4}{3}}((0, T); (W^{1,2}(\Omega; \mathbb{R}^{2 \times 2}))^*) \\
\mathbb{F}_{r\varepsilon} &\rightarrow \mathbb{F}_\varepsilon && \text{strongly in } L_{t,x}^q, \text{ for } q \in [1, 4).
\end{aligned}$$

As well as the equations

$$\langle \partial_t u_\varepsilon, \omega \rangle - \int_\Omega u_\varepsilon \otimes u_\varepsilon : \nabla_x \omega \, dx + \int_\Omega \nu(\theta_\varepsilon) Du_\varepsilon : \nabla_x \omega \, dx + \int_\Omega \mathbb{F}_\varepsilon \mathbb{F}_\varepsilon^T : \nabla_x \omega \, dx = 0,$$

for any $\omega \in W_0^{1,2}(\Omega)$,

$$\begin{aligned}
\langle \partial_t \mathbb{F}_\varepsilon, A \rangle - \int_\Omega \mathbb{F}_\varepsilon \otimes u_\varepsilon : \nabla_x A \, dx - \int_\Omega (\nabla_x u_\varepsilon \mathbb{F}_\varepsilon) : A \, dx \\
+ \frac{1}{2} \int_\Omega \delta(\theta_\varepsilon) \mathbb{F}_\varepsilon \mathbb{F}_\varepsilon^T \mathbb{F}_\varepsilon : A \, dx - \frac{1}{2} \int_\Omega \delta(\theta_\varepsilon) \mathbb{F}_\varepsilon : A \, dx + \varepsilon \int_\Omega \nabla_x \mathbb{F}_\varepsilon : \nabla_x A \, dx = 0,
\end{aligned}$$

for any $A \in W^{1,2}(\Omega; \mathbb{R}^{2 \times 2})$,

$$\begin{aligned}
- \int_0^T \int_\Omega \theta_\varepsilon \partial_t \phi \, dx \, dt + \int_0^T \int_\Omega \theta_\varepsilon u_\varepsilon \cdot \nabla_x \phi \, dx \, dt + \int_0^T \int_\Omega \kappa(\theta_\varepsilon) \nabla_x \theta_\varepsilon \cdot \nabla_x \phi \, dx \, dt \\
- \int_0^T \int_\Omega \nu(\theta_\varepsilon) |Du_\varepsilon|^2 \phi \, dx \, dt - \int_0^T \int_\Omega |\mathbb{F}_\varepsilon \mathbb{F}_\varepsilon^T - \mathbb{I}|^2 \phi \, dx \, dt = \int_\Omega \theta_0(x) \phi(0, x) \, dx,
\end{aligned}$$

for any $\phi \in C_c^1([0, T] \times \overline{\Omega})$. Therefore, our main goal in this subsection will be to show that after the convergence

$$\theta_\varepsilon > 0, \quad \text{a.e. in } (0, T) \times \Omega.$$

To this end, let us notice that due to (4.33) and (4.56)

$$(4.57) \quad \theta_\varepsilon \geq 0, \quad \text{a.e. in } (0, T) \times \Omega.$$

To get the strict inequality we first formally test (4.53) by $\theta_{r\varepsilon}^{-1}$ (we skip the details as it is a standard procedure involving mollification in time and a function approximating a time interval). We get

$$\int_{\Omega} \ln(\theta_{r\varepsilon}(t)) \, dx \geq \int_{\Omega} \ln(\theta_0^r) \, dx,$$

which since

$$\left| \int_{\Omega} \ln(\theta_0^r) \, dx \right| \leq \int_{\Omega} |\ln(\theta_0^r)| \, dx \leq \int_{\Omega} |\ln(\theta_0)| \, dx \leq C,$$

implies

$$(4.58) \quad \int_{\Omega} \ln(\theta_{r\varepsilon}(t)) \, dx > -C.$$

At the same time, we may test (4.53) by an approximation of the interval $(0, t)$, and, after applying the Lebesgue's differentiation theorem, similarly as in (4.34) obtain

$$\|\theta_{r\varepsilon}\|_{L_t^\infty L_x^1} \leq C(\|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}).$$

Thus, applying a simple inequality

$$\ln(x) \leq x - 1, \quad \text{for } x > 0.$$

we get

$$(4.59) \quad \int_{\{\theta_{r\varepsilon} > 1\}} \ln(\theta_{r\varepsilon}(t)) \, dx \leq \int_{\{\theta_{r\varepsilon} > 1\}} \theta_{r\varepsilon} - 1 \, dx \leq C(\|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}).$$

Combining both (4.58) and (4.59) grants us

$$\|\ln(\theta_{r\varepsilon})\|_{L_t^\infty L_x^1} \leq C(\|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}),$$

which by Fatou's lemma implies

$$(4.60) \quad \|\ln(\theta_\varepsilon)\|_{L_t^\infty L_x^1} \leq C(\|u_0\|_{L_x^2}, \|\mathbb{F}_0\|_{L_x^2}, \|\theta_0\|_{L_x^1}).$$

Combining both (4.60) and (4.57) we arrive at

$$(4.61) \quad \theta_\varepsilon > 0, \quad \text{a.e. in } (0, T) \times \Omega.$$

4.6. Convergence with $\varepsilon \rightarrow 0^+$. This is arguably the easiest part of the proof. Due to the fact that bounds (4.32), (4.11), (4.36), (4.37), (4.38) are uniform in ε performing the same calculations

as before we may obtain that (up to the subsequence)

$$\begin{aligned} \theta_\varepsilon &\rightharpoonup \theta && \text{weakly in } L_{t,x}^{2-\lambda}, \\ \theta_\varepsilon &\rightharpoonup \theta && \text{weakly in } L_t^{\frac{4}{3}-\lambda} W_x^{1, \frac{4}{3}-\lambda}, \\ \theta_\varepsilon &\rightarrow \theta && \text{strongly in } L_t^{\frac{4}{3}-\lambda} L_x^{\frac{8-6\lambda}{2+3\lambda}}. \end{aligned}$$

for all $\lambda > 0$, sufficiently close to 0, and

$$\begin{aligned} u_\varepsilon &\rightharpoonup u && \text{weakly}^* \text{ in } L_t^\infty L_x^2, \\ u_\varepsilon &\rightharpoonup u && \text{weakly in } L_t^2 W_{0,x}^{1,2}, \\ \partial_t u_\varepsilon &\rightharpoonup \partial_t u && \text{weakly in } L^2((0, T); (W_{0,\text{div}}^{1,2}(\Omega))^*), \\ u_\varepsilon &\rightarrow u && \text{strongly in } L_{t,x}^q, \text{ for } q \in [1, 4), \\ \mathbb{F}_\varepsilon &\rightharpoonup \mathbb{F} && \text{weakly}^* \text{ in } L_t^\infty L_x^2, \\ \mathbb{F}_\varepsilon &\rightharpoonup \mathbb{F} && \text{weakly in } L_{t,x}^4, \\ \partial_t \mathbb{F}_\varepsilon &\rightharpoonup \partial_t \mathbb{F} && \text{weakly in } L^{\frac{4}{3}}((0, T); (W^{1,2}(\Omega; \mathbb{R}^{2 \times 2}))^*), \\ \varepsilon \nabla_x \mathbb{F}_\varepsilon &\rightarrow 0 && \text{strongly in } L_{t,x}^2. \end{aligned}$$

The only problem is the loss of the Sobolev regularity of \mathbb{F} and the appropriate bound. Therefore, we cannot apply Aubin–Lions lemma A.1 to deduce strong convergence of \mathbb{F}_ε sequence. To remedy this situation we simply follow [6] to deduce

$$\mathbb{F}_\varepsilon \rightarrow \mathbb{F} \quad \text{strongly in } L_{t,x}^2.$$

We carried out the discussion about the procedure under (3.85), therefore we skip the details here.

In fact, this is even easier as $\delta(\theta_\varepsilon) \mathbb{F}_\varepsilon \mathbb{F}_\varepsilon^T \mathbb{F}_\varepsilon \in L_{t,x}^{\frac{4}{3}}$. Combining all of the convergences we arrive at

$$\langle \partial_t u, \omega \rangle - \int_\Omega u \otimes u : \nabla_x \omega \, dx + \int_\Omega \nu(\theta) Du : \nabla_x \omega \, dx + \int_\Omega \mathbb{F} \mathbb{F}^T : \nabla_x \omega \, dx = 0,$$

for any $\omega \in W_0^{1,2}(\Omega)$,

$$\begin{aligned} \langle \partial_t \mathbb{F}, A \rangle - \int_\Omega \mathbb{F} \otimes u : \nabla_x A \, dx - \int_\Omega (\nabla_x u \mathbb{F}) : A \, dx \\ + \frac{1}{2} \int_\Omega \delta(\theta) \mathbb{F} \mathbb{F}^T \mathbb{F} : A \, dx - \frac{1}{2} \int_\Omega \delta(\theta) \mathbb{F} : A \, dx = 0, \end{aligned}$$

for any $A \in W^{1,2}(\Omega; \mathbb{R}^{2 \times 2})$, and

$$\begin{aligned} & - \int_0^T \int_{\Omega} \theta \partial_t \phi \, dx \, dt + \int_0^T \int_{\Omega} \theta u \cdot \nabla_x \phi \, dx \, dt + \int_0^T \int_{\Omega} \kappa(\theta) \nabla_x \theta \cdot \nabla_x \phi \, dx \, dt \\ & - \int_0^T \int_{\Omega} \nu(\theta) |Du|^2 \phi \, dx \, dt - \int_0^T \int_{\Omega} |\mathbb{F} \mathbb{F}^T - \mathbb{I}|^2 \phi \, dx \, dt = \int_{\Omega} \theta_0(x) \phi(0, x) \, dx, \end{aligned}$$

for any $\phi \in C_c^1([0, T] \times \overline{\Omega})$. Which ends the proof.

APPENDIX A. AUXILIARY PROPOSITIONS

Lemma A.1. (Generalized Aubin–Lions lemma, [26, Lemma 7.7]) *Denote by*

$$W^{1,p,q}(I; X_1, X_2) := \left\{ u \in L^p(I; X_1); \frac{du}{dt} \in L^q(I; X_2) \right\}.$$

Then if X_1 is a separable, reflexive Banach space, X_2 is a Banach space and X_3 is a metrizable locally convex Hausdorff space, X_1 embeds compactly into X_2 , X_2 embeds continuously into X_3 , $1 < p < \infty$ and $1 \leq q \leq \infty$, we have

$$W^{1,p,q}(I; X_1, X_3) \text{ embeds compactly into } L^p(I; X_2).$$

In particular any bounded sequence in $W^{1,p,q}(I; X_1, X_3)$ has a convergent subsequence in $L^p(I; X_2)$.

Lemma A.2. (The div-curl lemma, [12]) *Let $\Omega \subset \mathbb{R}^n$ be an open and bounded domain with a Lipschitz boundary and let $p, q \in (1, \infty)$ with $\frac{1}{p} + \frac{1}{q} = 1$. Suppose $u_k \in L^p(\Omega; \mathbb{R}^n)$, $v_k \in L^q(\Omega; \mathbb{R}^n)$ are sequences such that*

$$u_k \rightharpoonup u \text{ weakly in } L^p(\Omega; \mathbb{R}^n), \quad v_k \rightharpoonup v \text{ weakly in } L^q(\Omega; \mathbb{R}^n),$$

and

$$u_k \cdot v_k \text{ is equiintegrable.}$$

Finally, assume that

$$\operatorname{div} u_k \rightarrow \operatorname{div} u \text{ strongly in } (W_0^{1,\infty}(\Omega))^*, \quad \operatorname{curl} v_k \rightarrow \operatorname{curl} v \text{ strongly in } (W_0^{1,\infty}(\Omega; M^{n \times n}))^*.$$

Then,

$$u_k \cdot v_k \rightharpoonup u \cdot v \text{ weakly in } L^1(\Omega).$$

Lemma A.3. *Let $1 < p, q < +\infty$. Suppose $\psi : \mathbb{R} \rightarrow \mathbb{R}$ is a Lipschitz function. For $r \in \mathbb{R}$ we define*

$$\Psi(x) = \int_r^x \psi(s) \, ds, \quad x \in \mathbb{R}.$$

Then, for any $u \in W^{1,p,p}(I; W^{1,q}(\Omega)) := \{u \in L^p(I; W^{1,q}(\Omega)); \frac{du}{dt} \in (L^p(I; W^{1,q}(\Omega)))^\}$ it holds*

$$\int_{t_1}^{t_2} \langle \partial_t u, \psi(u) \rangle \, dt = \int_{\Omega} \Psi(u(t_2)) \, dx - \int_{\Omega} \Psi(u(t_1)) \, dx, \quad t_1, t_2 \in I.$$

Lemma A.4. *Let $1 \leq p < \infty$ and $\{u_n\}_{n \in \mathbb{N}}$ be a sequence such that $u_n \rightarrow u$ in $L_t^p L_x^p$. Then, there exists a subsequence $\{u_{n_k}\}_{k \in \mathbb{N}}$, such that*

$$\text{for a.e. } t \in (0, T) \quad u_{n_k}(t, x) \rightarrow u(t, x) \text{ in } L_x^p$$

Moreover, if $\{u_{n_k}\}_{k \in \mathbb{N}}$ is bounded in $L_t^\infty L_x^2$ we have for a.e. $t \in (0, T)$

$$\int_{\Omega} |u(t, x)|^2 \, dx \leq \liminf_{k \rightarrow \infty} \int_{\Omega} |u_{n_k}(t, x)|^2 \, dx$$

REFERENCES

- [1] M. Bathory, M. Bulíček, and J. Málek. Large data existence theory for three-dimensional unsteady flows of rate-type viscoelastic fluids with stress diffusion. *Adv. Nonlinear Anal.*, 10(1):501–521, 2021.
- [2] M. Bathory, M. Bulíček, and J. Málek. Coupling the Navier-Stokes-Fourier equations with the Johnson-Segalman stress-diffusive viscoelastic model: global-in-time and large-data analysis. *Math. Models Methods Appl. Sci.*, 34(3):417–476, 2024.
- [3] M. Bulíček, E. Feireisl, and J. Málek. A Navier-Stokes-Fourier system for incompressible fluids with temperature dependent material coefficients. *Nonlinear Anal. Real World Appl.*, 10(2):992–1015, 2009.
- [4] M. Bulíček, E. Feireisl, and J. Málek. On a class of compressible viscoelastic rate-type fluids with stress-diffusion. *Nonlinearity*, 32(12):4665–4681, 2019.
- [5] M. Bulíček, A. Jüngel, M. Pokorný, and N. Zamponi. Existence analysis of a stationary compressible fluid model for heat-conducting and chemically reacting mixtures. *J. Math. Phys.*, 63(5):Paper No. 051501, 48, 2022.
- [6] M. Bulíček, T. Los, Y. Lu, and J. Málek. On planar flows of viscoelastic fluids of Giesekus type. *Nonlinearity*, 35(12):6557–6604, 2022.
- [7] M. Bulíček, T. Los, and J. Málek. On three-dimensional flows of viscoelastic fluids of Giesekus type. *Nonlinearity*, 38(1):015004 (42pp), 2025.
- [8] M. Bulíček, J. Málek, V. Průša, and E. Süli. On incompressible heat-conducting viscoelastic rate-type fluids with stress-diffusion and purely spherical elastic response. *SIAM J. Math. Anal.*, 53(4):3985–4030, 2021.
- [9] M. Bulíček, J. Málek, and K. R. Rajagopal. Mathematical analysis of unsteady flows of fluids with pressure, shear-rate, and temperature dependent material moduli that slip at solid boundaries. *SIAM J. Math. Anal.*, 41(2):665–707, 2009.

- [10] J. Burgers. Mechanical considerations—model systems—phenomenological theories of relaxations and viscosity. *First Report on Viscosity and Plasticity (New York: Nordemann Publishing Company)*, pages 5–67, 1939.
- [11] L. Consiglieri. Weak solutions for a class of non-Newtonian fluids with energy transfer. *J. Math. Fluid Mech.*, 2(3):267–293, 2000.
- [12] S. Conti, G. Dolzmann, and S. Müller. The div-curl lemma for sequences whose divergence and curl are compact in $W^{-1,1}$. *C. R. Math. Acad. Sci. Paris*, 349(3-4):175–178, 2011.
- [13] R. J. DiPerna and P.-L. Lions. Ordinary differential equations, transport theory and Sobolev spaces. *Invent. Math.*, 98(3):511–547, 1989.
- [14] M. Dressler, B. J. Edwards, and H. C. Öttinger. Macroscopic thermodynamics of flowing polymeric liquids. *Rheologica Acta*, 38(2):117–136, 1999.
- [15] E. Feireisl. *Dynamics of viscous compressible fluids*, volume 26 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, Oxford, 2004.
- [16] E. Feireisl and A. Novotný. *Singular limits in thermodynamics of viscous fluids*. Advances in Mathematical Fluid Mechanics. Birkhäuser/Springer, Cham, second edition, 2017.
- [17] E. Feireisl and A. Novotný. *Mathematics of open fluid systems*. Nečas Center Series. Birkhäuser/Springer, Cham, [2022] ©2022.
- [18] H. Giesekus. A simple constitutive equation for polymer fluids based on the concept of deformation-dependent tensorial mobility. *Journal of Non-Newtonian Fluid Mechanics*, 11(1):69–109, 1982.
- [19] J. Hron, V. Miloš, V. Průša, O. Souček, and K. Tůma. On thermodynamics of incompressible viscoelastic rate type fluids with temperature dependent material coefficients. *International Journal of Non-Linear Mechanics*, 95:193–208, 2017.
- [20] D. Hu and T. Lelièvre. New entropy estimates for Oldroyd-B and related models. *Commun. Math. Sci.*, 5(4):909–916, 2007.
- [21] M. Johnson and D. Segalman. A model for viscoelastic fluid behavior which allows non-affine deformation. *Journal of Non-Newtonian Fluid Mechanics*, 2(3):255–270, 1977.
- [22] P. L. Lions and N. Masmoudi. Global solutions for some Oldroyd models of non-Newtonian flows. *Chinese Ann. Math. Ser. B*, 21(2):131–146, 2000.
- [23] N. Masmoudi. Global existence of weak solutions to macroscopic models of polymeric flows. *Journal de Mathématiques Pures et Appliquées*, 96(5):502–520, 2011.
- [24] J. G. Oldroyd. On the formulation of rheological equations of state. *Proc. Roy. Soc. London Ser. A*, 200:523–541, 1950.
- [25] K. Rajagopal and A. Srinivasa. A thermodynamic frame work for rate type fluid models. *Journal of Non-Newtonian Fluid Mechanics*, 88(3):207–227, 2000.
- [26] T. Roubíček. *Nonlinear partial differential equations with applications*, volume 153 of *International Series of Numerical Mathematics*. Birkhäuser/Springer Basel AG, Basel, second edition, 2013.

MATHEMATICAL INSTITUTE, FACULTY OF MATHEMATICS AND PHYSICS, CHARLES UNIVERSITY, SOKOLOVSKÁ 83,
186 75, PRAGUE, CZECH REPUBLIC

Email address: `mbul8060@karlin.mff.cuni.cz`

FACULTY OF MATHEMATICS, INFORMATICS AND MECHANICS, UNIVERSITY OF WARSAW, STEFANA BANACHA 2,
02-097 WARSAW, POLAND; INSTITUTE OF MATHEMATICS OF POLISH ACADEMY OF SCIENCES, JANA I JĘDRZEJA
ŚNIADECKICH 8, 00-656 WARSAW, POLAND

Email address: `jw.woznicki@student.uw.edu.pl`