Quantum integrable model for the quantum cohomology/quantum K-theory of flag varieties and the double β -Grothendieck polynomials

Jirui Guo

School of Mathematical Sciences,
Institute for Advanced Study,
Key Laboratory of Intelligent Computing and Applications,
Tongji University, Shanghai 200092, China

jrkwok@tongji.edu.cn

Abstract

A GL(n) quantum integrable system generalizing the asymmetric five vertex spin chain is shown to encode the ring relations of the equivariant quantum cohomology and quantum K-theory ring of flag varieties. We also show that the Bethe ansatz states of this system generate the double β -Grothendieck polynomials.

Contents

1	Introduction	2
2	The $\operatorname{GL}(n)$ asymmetric five vertex model	4
3	Bethe ansatz equations and the equivariant quantum cohomology/K-theory ring of flag varieties	6
4	Relationship with the double β -Grothendieck polynomials	12
A	Quantum cohomology/K-theory of flag varieties and the double $\beta\text{-Grothendieck}$ polynomials	25
Re	eferences	27

1 Introduction

The mathematical structure and exact solutions of quantum integrable systems have proved to be useful tools in providing information about physical phenomena, as well as revealing certain algebraic results. For example, in [1], it was shown that the underlying Frobenius algebra of the asymmetric six vertex model, which depends on a parameter β , is exactly the equivariant quantum cohomology of the Grassmannians when $\beta = 0$, and the equivariant quantum K-theory ring of the Grassmannians when $\beta = -1$. The Bethe/Gauge correspondence of this model was later studied in [2].

This work aims at generalizing the idea of [1], establishing a relationship between a GL(n) integrable model and the equivariant quantum cohomology/quantum K-theory ring of general flag varieties. We will also show that the Bethe ansatz states of this model generate the double β -Grothendieck polynomials. When $\beta = 0$ these polynomials reduce to the double Schubert polynomials representing Schubert classes in the equivariant cohomology ring, and when $\beta = -1$ they reduce to the double Grothendieck polynomials representing Schubert classes in the equivariant K-theory ring.

The relationships between the quantum cohomology/K-theory of flag varieties and Toda lattices were discussed in [3] and [4]. Moreover, it was shown that quantum K-theory ring of the cotangent bundle of flag varieties can be realized by the XXZ model, and in certain limit it reduces to the quantum K-theory ring of flag varieties [5]. Double β -Grothendieck polynomials can also be realized by the partition functions of certain 2d lattice models [6–8].

Our main results are as follows:

• The Bethe ansatz equations of the GL(n) quantum integrable system defined by the R-matrix (2.3)

$$R^{(n)}(x,y) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & I_{n-1} & 0 \\ 0 & (1+\beta(x\ominus y))I_{n-1} & (x\ominus y)I_{n-1} & 0 \\ 0 & 0 & 0 & R^{(n-1)}(x,y) \end{pmatrix},$$

where

$$R^{(2)}(x,y) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 + \beta(x \ominus y) & x \ominus y & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

are derived (Theorem 3.1).

• It is shown that the Bathe ansatz equations give rise to the Whitney type ring relations of the equivariant quantum cohomology ring of partial flag varieties when $\beta = 0$ in terms of the Chern classes (Eq.(3.22)):

$$c(S_i) \cdot c(S_{i+1}/S_i) = c(S_{i+1}) + (-1)^{k_{n-i-1}-k_{n-i}} q_{n-i} c(S_{i-1}),$$

and give rise to the ring relations of the equivariant quantum K-theory ring of partial flag varieties when $\beta = -1$ in terms of the λ_y classes (Eq.(3.27)):

$$\lambda_y(\mathcal{S}_i) * \lambda_y(\mathcal{S}_{i+1}/\mathcal{S}_i) = \lambda_y(\mathcal{S}_{i+1}) + \frac{q_{n-i}}{1 - q_{n-i}} y^{k_{n-i-1} - k_{n-i}} \left(\lambda_y(\mathcal{S}_{i-1}) - \lambda_y(\mathcal{S}_i)\right) * \det(\mathcal{S}_{i+1}/\mathcal{S}_i),$$

where S_i is the *i*th tautological bundle of the flag variety.

• Given the matrix elements $B_i(u)$ of the monodromy matrix as in Eq.(2.22), and the pseudo vacuum state $|\Omega^{(0)}\rangle$ as in (3.2), the expansion coefficients in the natural basis of the quantum state $B_{N-1}(\sigma_1)\cdots B_1(\sigma_{N-1})|\Omega^{(0)}\rangle$ are exactly the double β -Grothendieck polynomials $\mathcal{G}_{\pi}^{(\beta)}$ (Theorem 4.6):

$$B_{N-1}(\sigma_1)\cdots B_1(\sigma_{N-1})|\Omega^{(0)}\rangle = \sum_{\pi\in S_N} \mathcal{G}_{\pi}^{(\beta)}(\sigma_1,\cdots,\sigma_{N-1};\ominus t_1,\cdots,\ominus t_{N-1})|\omega_N\pi^{-1}\rangle,$$

where ω_N is the permutation with maximal length in the permutation group S_N .

• When all the instances of $e_l(\sigma_1^{(i)}, \dots, \sigma_{N-i}^{(i)})$, $l=1,\dots,N-i$, are identified with $e_l(x_1,\dots,x_{N-i})$ for $i=1,\dots,N-1$, the on-shell Bethe ansatz state (3.4) (in the full flag case) has the expansion (Theorem 4.11):

$$\begin{split} |\psi^{(0)}\rangle &= \sum_{i_1, \dots, i_{k_1}} \psi^{(1)}_{i_1 \dots i_{k_1}} B_{i_1}(\sigma^{(1)}_1) \dots B_{i_{k_1}}(\sigma^{(1)}_{k_1}) |\Omega^{(0)}\rangle \\ &= \sum_{w \in S_N} \mathcal{G}^{(\beta)}_w(x_1, \dots, x_{N-1}; \ominus t_1, \dots, \ominus t_{N-1}) |\omega_N w^{-1}\rangle \,, \end{split}$$

where x_i can be interpreted as the first Chern class of the line bundle S_i/S_{i-1} when $\beta = 0$, and $1 - x_i$ can be interpreted as the K-theory class of S_i/S_{i-1} when $\beta = -1$ (This means $\sigma_a^{(N-i)}$ ($\beta = 0$) or $1 - \sigma_a^{(N-i)}$ ($\beta = -1$), $a = 1, \dots, i$, should be interpreted as the Chern roots of S_i).

This paper is organized as follows. In Sec. 2, the basic ingredients of the GL(n) quantum integrable system generalizing the asymmetric five vertex spin chain are introduced. In Sec. 3, the Bethe ansatz equations are derived, which are shown to give rise to the ring relations of the equivariant quantum cohomology/K-theory ring of the flag varieties. Sec. 4 consists of a series of propositions leading to our main results on the relationship between the quantum integrable model under study

and the double β -Grothendieck polynomials. In the appendix, we present some background on the quantum cohomology/K-theory of flag varieties and the double β -Grothendieck polynomials.

2 The GL(n) asymmetric five vertex model

In this section, we present the basic ingredients of the GL(n) asymmetric five vertex model. Let $V = \mathbb{C}^n$, we define the R-matrix $R^{(n)}(x,y) \in End(V \otimes V)$ as follows:

For n = 2, the R-matrix is defined as

$$R^{(2)}(x,y) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 + \beta(x \ominus y) & x \ominus y & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \tag{2.1}$$

where β is a formal variable, and

$$x \ominus y \equiv \frac{x - y}{1 + \beta y}. (2.2)$$

For n > 2, under the decomposition $V \cong \mathbb{C} \oplus \mathbb{C}^{n-1}$, $R^{(n-1)}(x,y)$ is defined in the following block diagonal form,

$$R^{(n)}(x,y) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & I_{n-1} & 0 \\ 0 & (1+\beta(x\ominus y))I_{n-1} & (x\ominus y)I_{n-1} & 0 \\ 0 & 0 & 0 & R^{(n-1)}(x,y) \end{pmatrix},$$
(2.3)

where I_r stands for the identity matrix of size r. Alternatively, the entries of the R-matrix can be written as

$$R_{ij,kl}^{(n)}(x,y) = (1 + \delta_{k < l}\beta(x \ominus y))\delta_{il}\delta_{jk} + \delta_{k > l}(x \ominus y)\delta_{ik}\delta_{jl}, \tag{2.4}$$

where i, j are row indices, k, l are colomn indices, $\{i, k\}$ and $\{j, l\}$ are indices for the first and second factor of $V \otimes V$ respectively. We also define

$$\delta_{i < j} = \begin{cases} 1, & \text{if } i < j, \\ 0, & \text{if } i \ge j, \end{cases} \quad \delta_{i > j} = \begin{cases} 1, & \text{if } i > j, \\ 0, & \text{if } i \le j. \end{cases}$$

The R-matrix (2.3) appeared in [8] as the $q \to 0$ limit of a Drinfeld twist of the R-matrix for an evaluation module of $U_q(\hat{\mathbf{sl}}_{n+1})$, but here we show that $R^{(n)}(x,y)$ satisfies the Yang-Baxter equation by a direct computation:

Theorem 2.1. The R-matrix defined by Eq.(2.3) satisfies the Yang-Baxter equation

$$R_{12}^{(n)}(x,y)R_{13}^{(n)}(x,z)R_{23}^{(n)}(y,z) = R_{23}^{(n)}(y,z)R_{13}^{(n)}(x,z)R_{12}^{(n)}(x,y)$$
(2.5)

for all n > 1, where $R_{ab}^{(n)}$ acts on the ath and bth factor of $V \otimes V \otimes V$.

Proof. A direct computation shows that Eq.(2.5) holds for n = 2. Now suppose $R^{(n-1)}$ satisfies the Yang-Baxter equation for some n > 2. Let i_{ab} denote the identity map from the ath factor to the bth factor of $V \otimes V \otimes V$. Plugging (2.3) into Eq.(2.5), we find that Eq.(2.5) decomposes into

the following identities:

$$i_{21}i_{32} = i_{31}, \quad i_{13} = i_{23}i_{12}, \tag{2.6}$$

$$i_{12}i_{31}i_{23} = i_{32}i_{13}i_{21}, \quad i_{21}i_{13}i_{32} = i_{23}i_{31}i_{12},$$
 (2.7)

$$(x-y)i_{31}i_{23} + (y-z)i_{21} = i_{21}(x-z), \quad (x-z)i_{12} = (x-y)i_{32}i_{13} + (y-z)i_{12},$$
 (2.8)

$$R_{12}^{(n-1)}(x,y)i_{31}R_{23}^{(n-1)}(y,z) = i_{32}R_{13}^{(n-1)}(x,z)i_{21},$$
(2.9)

$$i_{12}R_{13}^{(n-1)}(x,z)i_{23} = R_{23}^{(n-1)}(y,z)i_{13}R_{12}^{(n-1)}(x,y), (2.10)$$

$$(x-z)(1+\beta y)R_{12}^{(n-1)}(x,y)i_{32} = (y-z)(1+\beta x)i_{31}i_{12} + (x-y)(1+\beta z)i_{32}R_{13}^{(n-1)}(x,z), \quad (2.11)$$

$$(y-z)(1+\beta x)i_{21}i_{13} + (x-y)(1+\beta z)R_{13}^{(n-1)}(x,z)i_{23} = (x-z)(1+\beta y)i_{23}R_{12}^{(n-1)}(x,y), \quad (2.12)$$

$$R_{12}^{(n-1)}(x,y)R_{13}^{(n-1)}(x,z)R_{23}^{(n-1)}(y,z) = R_{23}^{(n-1)}(y,z)R_{13}^{(n-1)}(x,z)R_{12}^{(n-1)}(x,y).$$
(2.13)

Eq.(2.6)-(2.8) are direct consequences of the definition of i_{ab} . Eq.(2.9) and (2.10) can be easily derived from

$$\sum_{k,l} R_{ij,lk}^{(m)}(x,y) R_{kl,st}^{(m)}(y,z) = R_{ij,st}^{(m)}(x,z)$$
(2.14)

and

$$\sum_{k,l} R_{ij,lk}^{(m)}(y,z) R_{kl,st}^{(m)}(x,y) = R_{ij,st}^{(m)}(x,z)$$
(2.15)

respectively, while Eq.(2.11) and (2.12) can be easily derived from

$$(y-z)(1+\beta x)R^{(m)}(x,x) + (x-y)(1+\beta z)R^{(m)}(x,z) = (x-z)(1+\beta y)R^{(m)}(x,y),$$
(2.16)

where $m \geq 2$. Eq.(2.14)-(2.16) can all be proved by simple induction on m. Finally, Eq.(2.13) is the induction hypothesis. Therefore, by induction, the Yang-Baxter equation holds for all $n \geq 2$. \square

To construct an integrable system, we take N copies of $V \cong \mathbb{C}^n$ and label them by $V_i, i = 1, \dots, N$. Let $V_a \cong V$ be an auxiliary space. The monodromy matrix is defined by

$$T_a^{(n)}(u) = R_{aN}(u, t_N) R_{a, N-1}(u, t_{N-1}) \cdots R_{a2}(u, t_2) R_{a1}(u, t_1), \tag{2.17}$$

where R_{ai} acts on $V_a \otimes V_i$. As in the usual QISM method for quantum integrable systems, the Yang-Baxter equation (2.5) leads to the following RTT relation:

$$R_{ab}^{(n)}(x,y)T_a^{(n)}(x)T_b^{(n)}(y) = T_b^{(n)}(y)T_a^{(n)}(x)R_{ab}^{(n)}(x,y).$$
(2.18)

We define the transfer matrix to be

$$t^{(n)}(u) = \text{Tr}_{V_a}(M_n T_a^{(n)}(u)), \tag{2.19}$$

where $M_n = \text{diag}(b_0, b_1, \dots, b_{n-1})$ acts on V_a , imposing a twisted periodic boundary condition. In the following, we also need

$$M_l := \operatorname{diag}(b_{n-l}, \dots, b_{n-1}), \quad l = 1, \dots, n.$$
 (2.20)

Eq.(2.18) tells us that¹

$$[t(u), t(v)] = 0.$$
 (2.21)

Under the decomposition $V_a = \mathbb{C} \oplus \mathbb{C}^{n-1}$, the monodromy matrix can be written in the following form:

$$T_a^{(n)}(u) = \begin{pmatrix} A(u) & B_1(u) & \cdots & B_{n-1}(u) \\ C_1(u) & D_{11}(u) & \cdots & D_{1,n-1}(u) \\ \vdots & \vdots & \ddots & \vdots \\ C_{n-1}(u) & D_{n-1,1}(u) & \cdots & D_{n-1,n-1}(u) \end{pmatrix}$$
(2.22)

with $A(u), B_i(u), C_i(u), D_{ij}(u)$ being operators acting on the Hilbert space $\mathcal{H} = V_1 \otimes V_2 \otimes \cdots \otimes V_N$. The RTT relation (2.18) also yields the following commutation relations:

$$A(x)B_{i}(y) = \frac{1}{y \ominus x} \left[B_{i}(y)A(x) - B_{i}(x)A(y) \right], \tag{2.23}$$

$$D_{ij}(x)B_k(y) = \frac{1}{x-y} \left[\sum_{s,l} \left((1+\beta y)B_s(y)D_{il}(x)R_{ls,jk}^{(n-1)}(x,y) \right) - (1+\beta x)B_j(x)D_{ik}(y) \right], \quad (2.24)$$

$$B_i(x)B_j(y) = \sum_{r,s} B_r(y)B_s(x)R_{sr,ij}^{(n-1)}(x,y).$$
(2.25)

3 Bethe ansatz equations and the equivariant quantum cohomology/K-theory ring of flag varieties

In this section, we derive the Bethe ansatz equations of the GL(n) five vertex model introduced in the last section following the method proposed in [9], and discuss their relationship with the equivariant quantum cohomology and quantum K-theory ring of flag varieties.

Let $V = \mathbb{C}^n$ be spanned by v_0, v_1, \dots, v_{n-1} , where $v_i := |i\rangle$ is the (i+1)th unit vector of \mathbb{C}^n . Define

$$|n_1, n_2, \cdots, n_N\rangle := v_{n_1} \otimes v_{n_2} \otimes \cdots \otimes v_{n_N},$$

then a basis of $V_1 \otimes V_2 \otimes \cdots V_N$ can be chosen as

$$\{|n_1, n_2, \cdots, n_N\rangle \mid 0 \le n_i \le n-1 \}.$$
 (3.1)

We call the basis (3.1) the natural basis of the Hilbert space $\mathcal{H} = V_1 \otimes V_2 \otimes \cdots \otimes V_N$. The pseudo vacuum is taken to be

$$|\Omega^{(0)}\rangle = |0, 0, \cdots, 0\rangle \in V_1 \otimes V_2 \otimes \cdots \otimes V_N. \tag{3.2}$$

It is easy to compute

$$A(u) |\Omega^{(0)}\rangle = |\Omega^{(0)}\rangle, \quad D_{ij}(u) |\Omega^{(0)}\rangle = \prod_{k=1}^{N} (u \ominus t_k) \delta_{ij} |\Omega^{(0)}\rangle.$$
 (3.3)

¹Notice that for any diagonal matrix M acting on the auxiliary space, we have $R_{ab}^{(n)}(x,y)M_aM_b=M_bM_aR_{ab}^{(n)}(x,y)$.

The Bethe ansatz state is taken to be of the form

$$|\psi^{(0)}\rangle = \sum_{i_1,\dots,i_{k_1}} \psi_{i_1\dots i_{k_1}}^{(1)} B_{i_1}(\sigma_1^{(1)}) \cdots B_{i_{k_1}}(\sigma_{k_1}^{(1)}) |\Omega^{(0)}\rangle, \qquad (3.4)$$

where $\sigma_{\alpha_1}^{(1)}$, $\alpha_1 = 1, \dots, k_1$, are parameters to be determined, and the expansion coefficients $\psi_{i_1 \dots i_{k_1}}^{(1)}$ depend on n-2 sets of parameters

$$\sigma_{\alpha_2}^{(2)}, \cdots, \sigma_{\alpha_{n-1}}^{(n-1)}$$

with $\alpha_s = 1, 2, \dots, k_s$, where $\sigma_a^{(s)} \neq \sigma_b^{(s)}$ for $a \neq b$ and $1 \leq s \leq n-1$. Now we can prove the following

Theorem 3.1. If the parameters $\sigma_{\alpha_1}^{(1)}, \dots, \sigma_{\alpha_{n-1}}^{(n-1)}$ satisfy the following Bethe ansatz equations:

$$\prod_{a=1}^{k_1} (1 + \beta(\sigma_a^{(1)} \ominus \sigma_{\alpha_1}^{(1)})) \prod_{i=1}^{N} (\sigma_{\alpha_1}^{(1)} \ominus t_i) = (-1)^{k_1 - 1} q_1 \prod_{b=1}^{k_2} (\sigma_b^{(2)} \ominus \sigma_{\alpha_1}^{(1)}), \tag{3.5}$$

$$\prod_{a=1}^{k_m} (1 + \beta(\sigma_a^{(m)} \ominus \sigma_{\alpha_m}^{(m)})) \prod_{i=1}^{k_{m-1}} (\sigma_{\alpha_m}^{(m)} \ominus \sigma_i^{(m-1)}) = (-1)^{k_m - 1} q_m \prod_{b=1}^{k_{m+1}} (\sigma_b^{(m+1)} \ominus \sigma_{\alpha_m}^{(m)}), \quad (3.6)$$

$$\prod_{a=1}^{k_{n-1}} \left(1 + \beta(\sigma_a^{(n-1)} \ominus \sigma_{\alpha_{n-1}}^{(n-1)})\right) \prod_{i=1}^{k_{n-2}} \left(\sigma_{\alpha_{n-1}}^{(n-1)} \ominus \sigma_i^{(n-2)}\right) = (-1)^{k_{n-1}-1} q_{n-1}, \tag{3.7}$$

where $m = 2, \dots, n-2$, $\alpha_s = 1, \dots, k_s$, $q_s := b_{s-1}/b_s$, $s = 1, \dots, n-1$, then the state (3.4) is a common eigenstate of the transfer matrices t(u) defined by (2.19).

Proof. From the commutation relations (2.23) and (2.24), one can compute

$$A(u) |\psi^{(0)}\rangle = \prod_{a=1}^{k_1} (\sigma_a^{(1)} \ominus u)^{-1} |\psi^{(0)}\rangle + |\phi_A\rangle,$$

$$D_{ij}(u) |\psi^{(0)}\rangle = \frac{\prod_{l=1}^{N} (u \ominus t_l)}{\prod_{a=1}^{k_1} (u \ominus \sigma_a^{(1)})}$$

$$\times \sum_{i_1, \dots, i_{k_1}} \sum_{j_1, \dots, j_{k_1}} \left\{ [T_{ij}^{(n-1)}(u; \sigma^{(1)})]_{i_1 \dots i_{k_1}, j_1 \dots j_{k_1}} \psi_{j_1 \dots j_{k_1}}^{(1)} B_{i_1}(\sigma_1^{(1)}) \dots B_{i_{k_1}}(\sigma_{k_1}^{(1)}) \right\} |\Omega^{(0)}\rangle + |\phi_D\rangle,$$

$$(3.8)$$

where $|\phi_A\rangle$ and $|\phi_D\rangle$ are terms that are not in the subspace spanned by states of the form $B_{i_1}(\sigma_1^{(1)})\cdots B_{i_{k_1}}(\sigma_{k_1}^{(1)})|\Omega^{(0)}\rangle$ (unwanted terms), and

$$[T_{ij}^{(n-1)}(u;\sigma^{(1)})]_{i_1\cdots i_{k_1},j_1\cdots j_{k_1}} = \sum_{\mu_1\cdots\mu_{k_1-1}} R_{ii_{k_1},\mu_{k_1-1}j_{k_1}}^{(n-1)}(u,\sigma_{k_1}^{(1)})\cdots R_{\mu_2i_2,\mu_1j_2}^{(n-1)}(u,\sigma_2^{(1)})R_{\mu_1i_1,jj_1}^{(n-1)}(u,\sigma_1^{(1)}).$$

$$(3.9)$$

Eq.(3.9) suggests that $T^{(n-1)}$ can be viewed as a monodromy matrix of a GL(n-1) integrable system with R-matrix given by $R^{(n-1)}$. Correspondingly, the transfer matrix can be defined as

$$t^{(n-1)}(u;\sigma^{(1)}) = \text{Tr}(M_{n-1}T^{(n-1)}(u;\sigma^{(1)})),$$

where M_l is defined by (2.20). Let

$$\psi_{i_1\cdots i_{k_1}}^{(1)} = \langle i_1 i_2 \cdots i_{k_1} | \psi^{(1)} \rangle, \quad i_s = 1, \cdots, n-1.$$

If $|\psi^{(1)}\rangle$ is an eigenstate of $t^{(n-1)}(u;\sigma^{(1)})$ with eigenvalue $\Lambda^{(1)}(u)$, i.e.

$$t^{(n-1)}(u;\sigma^{(1)})|\psi^{(1)}\rangle = \Lambda^{(1)}(u)|\psi^{(1)}\rangle, \qquad (3.10)$$

then, as a consequence of Eq.(3.8)

$$\operatorname{Tr}(M_{n-1}D(u))|\psi^{(0)}\rangle = \frac{\prod_{l=1}^{N}(u\ominus t_{l})}{\prod_{a=1}^{k_{1}}(u\ominus\sigma_{a}^{(1)})}\Lambda^{(1)}(u)|\psi^{(0)}\rangle + |\tilde{\phi}_{D}\rangle,$$

where $|\tilde{\phi}_D\rangle$ is the sum of the unwanted terms. Then

$$t^{(n)}(u) |\psi^{(0)}\rangle = [b_0 A(u) + \text{Tr}(M_{n-1} D(u)] |\psi^{(0)}\rangle) = \Lambda^{(0)}(u) |\psi^{(0)}\rangle + |\phi^{(n)}\rangle,$$

where

$$\Lambda^{(0)}(u) = b_0 \prod_{a=1}^{k_1} (\sigma_a^{(1)} \ominus u)^{-1} + \frac{\prod_{l=1}^{N} (u \ominus t_l)}{\prod_{a=1}^{k_1} (u \ominus \sigma_a^{(1)})} \Lambda^{(1)}(u), \tag{3.11}$$

and $|\phi^{(n)}\rangle$ is the sum of the unwanted terms. As in the usual Bethe ansatz method, vanishing of the unwanted terms $|\phi^{(n)}\rangle$ amounts to $\Lambda^{(0)}(u)$ being holomorphic, i.e.

$$\operatorname{Res}(\Lambda^{(0)}(u), \sigma_a^{(1)}) = 0$$
 (3.12)

for all $a = 1, \dots, k_1$. Eq.(3.12) gives us the equations for $\sigma_a^{(1)}$:

$$\prod_{\alpha=1}^{k_1} (1 + \beta \sigma_{\alpha}^{(1)}) \prod_{i=1}^{N} (\sigma_a^{(1)} \ominus t_i) \Lambda^{(1)}(\sigma_a^{(1)}) = (-1)^{k_1 - 1} b_0 (1 + \beta \sigma_a^{(1)})^{k_1}, \quad a = 1, \dots, k_1.$$
 (3.13)

Thus we have reduced the eigenvalue problem for the GL(n) system to the eigenvalue problem (3.10) for a GL(n-1) system. Clearly, we can continue to reduce the rank in exactly the same way. At each step, a set of parameters $\sigma_{\alpha_m}^{(m)}$, $\alpha_m = 1, \dots, k_m$, has to be introduced to construct the Bethe ansatz state

$$|\psi^{(m-1)}\rangle = \sum_{i_1,\dots,i_{k_m}} \psi_{i_1,\dots,i_{k_m}}^{(m)} B_{i_1}^{(m-1)}(\sigma_1^{(m)}) \dots B_{i_{k_m}}^{(m-1)}(\sigma_{k_m}^{(m)}) |\Omega^{(m-1)}\rangle, \qquad (3.14)$$

where

$$\psi_{i_1,\dots,i_{k_m}}^{(m)} = \langle i_1,\dots,i_{k_m} | \psi^{(m)} \rangle,$$
 (3.15)

and $B_i^{(m-1)}$'s are the B_i operators defined as matrix elements of the monodromy matrix at the corresponding step as in (2.22). As before, for (3.14) to be an eigenstate, the following equations have to be satisfied:

$$\prod_{b=1}^{k_m} (1 + \beta \sigma_b^{(m)}) \prod_{a=1}^{k_{m-1}} (\sigma_{\alpha_m}^{(m)} \ominus \sigma_a^{(m-1)}) \Lambda^{(m)} (\sigma_{\alpha_m}^{(m)}) = (-1)^{k_m - 1} b_{m-1} (1 + \beta \sigma_{\alpha_m}^{(m)})^{k_m}, \quad \alpha_m = 1, \dots, k_m,$$
(3.16)

accordingly, the eigenvalues satisfy

$$\Lambda^{(m-1)}(u) = b_{m-1} \prod_{\alpha=1}^{k_m} (\sigma_{\alpha}^{(m)} \ominus u)^{-1} + \frac{\prod_{a=1}^{k_{m-1}} (u \ominus \sigma_a^{(m-1)})}{\prod_{b=1}^{k_m} (u \ominus \sigma_b^{(m)})} \Lambda^{(m)}(u)$$
(3.17)

for $m = 2, \dots, n - 2$.

This procedure continues until we get a GL(2) system with R-matrix given by Eq.(2.1). This GL(2) system can be solved exactly by the algebraic Bethe ansatz method as for spin chains, from which we get Eq.(3.7) and

$$\Lambda^{(n-2)}(u) = b_{n-2} \prod_{\alpha=1}^{k_{n-1}} (\sigma_{\alpha}^{(n-1)} \ominus u)^{-1} + b_{n-1} \frac{\prod_{a=1}^{k_{n-2}} (u \ominus \sigma_a^{(n-2)})}{\prod_{b=1}^{k_{n-1}} (u \ominus \sigma_b^{(n-1)})}.$$
 (3.18)

Eq.(3.18) then allows us to solve all the eigenvalues $\Lambda^{(m)}$, $m=0,1,\cdots,n-3$, from Eq.(3.17) and (3.11). Plugging these eigenvalues in Eq.(3.16) and (3.13), we get the equations Eq.(3.6) and (3.5).

Now let us restrict to the cases in which $k_{n-1} < \cdots < k_2 < k_1 < N$ and compare Eq.(3.5)-(3.7) with the ring relations of the quantum cohomology and quantum K-theory ring of flag varieties. The reader may refer to the appendix and references therein for a review of the quantum cohomology and quantum K-theory ring of flag varieties.

When $\beta = 0$, we can identify $\sigma_a^{(n-i)}$ with the Chern roots of \mathcal{S}_i , the *i*th tautological bundle of the flag variety $\operatorname{Fl}(k_{n-1}, \dots, k_2, k_1; N)$, i.e. the *j*th elementary symmetric polynomial $e_j(\sigma_1^{(n-i)}, \dots, \sigma_{k_{n-i}}^{(n-i)})$ is identified with the *j*th Chern class of \mathcal{S}_i :

$$e_j(\sigma_1^{(n-i)}, \dots, \sigma_{k_{n-i}}^{(n-i)}) = c_j(S_i), \quad j = 1, \dots, k_{n-i}.$$

Under this identification, it was checked in [10, 11] that Eq.(3.5)-(3.7) specialized to $\beta = 0$ are exactly the ring relations of the equivariant quantum cohomology of $Fl(k_{n-1}, \dots, k_2, k_1; N)$ with the equivariant parameters given by (t_1, \dots, t_N) .

When $\beta = -1$, we can identify $1 - \sigma_a^{(n-i)}$ with the Chern roots of S_i , i.e. the jth elementary symmetric polynomial $e_j(1 - \sigma_1^{(i)}, \dots, 1 - \sigma_{k_{n-i}}^{(i)})$ is identified with $\wedge^j S_i$ in the Grothendieck group of the flag variety $Fl(k_{n-1}, \dots, k_2, k_1; N)$:

$$e_j(1-\sigma_1^{(n-i)},\cdots,1-\sigma_{k_{n-i}}^{(n-i)})=\wedge^j S_i, \quad j=1,\cdots,k_{n-i}.$$

It was argued in [11] that Eq.(3.5)-(3.7) specialized to $\beta = -1$ are the ring relations of the (predicted) equivariant quantum K-theory ring of $Fl(k_{n-1}, \dots, k_2, k_1; N)$ with the equivariant parameters given by $(1 - t_1, \dots, 1 - t_N)$.

Indeed, as in [11], one can use Vieta's formula² to convert Eq.(3.5)-(3.7) into the ring relations of the equivariant quantum cohomology ring ($\beta = 0$) or the equivariant K-theory ring ($\beta = -1$) of the flag variety Fl($k_{n-1}, \dots, k_2, k_1; N$).

²Vieta's formula: If r_1, r_2, \dots, r_n are roots of the polynomial $a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$, then $e_l(r_1, \dots, r_n) = (-1)^l a_{n-l}/a_n$.

When $\beta = 0$, Eq.(3.5)-(3.7) reduces to

$$\prod_{\alpha=1}^{k_{m-1}} (\sigma_a^{(m)} - \sigma_\alpha^{(m-1)}) = (-1)^{k_m - 1} q_m \prod_{\gamma=1}^{k_{m+1}} (\sigma_\gamma^{(m+1)} - \sigma_a^{(m)})$$
(3.19)

for $a=1,\dots,k_m, m=1,2,\dots,n$, where we have defined $\sigma_i^{(0)}=t_i, k_0=N, k_n=0$. Eq.(3.19) tells us that $\sigma_a^{(m)}, a=1,\dots,k_m$, are roots of the polynomial

$$\sum_{i=0}^{k_{m-1}} (-1)^{i} x^{k_{m-1}-i} e_{i}(\sigma^{(m-1)}) + (-1)^{k_{m}-k_{m+1}} q_{m} \sum_{j=0}^{k_{m+1}} (-1)^{j} x^{k_{m+1}-j} e_{j}(\sigma^{(m+1)}).$$
 (3.20)

Notice that the coefficient of $x^{k_{m-1}-l}$ in (3.20) is³

$$(-1)^{l}e_{l}(\sigma^{(m-1)}) + (-1)^{k_{m-1}-k_{m}+l}q_{m}e_{k_{m+1}-k_{m}+l}(\sigma^{(m+1)}).$$

Assume the other $k_{m-1} - k_m$ roots of (3.20) are $\eta_b^{(m)}$, $b = 1, \dots, k_{m-1} - k_m$, then Vieta's formula yields

$$\sum_{i=0}^{l} e_i(\sigma^{(m)}) e_{l-i}(\eta^{(m)}) = e_l(\sigma^{(m-1)}) + (-1)^{k_{m-1}-k_m} q_m e_{k_{m+1}-k_{m-1}+l}(\sigma^{(m+1)}).$$
 (3.21)

If we interpret $\sigma_a^{(m)}$, $a=1,\dots,k_m$, as Chern roots of \mathcal{S}_{n-m} and $\eta_b^{(m)}$, $b=1,\dots,k_{m-1}-k_m$, as Chern roots of $\mathcal{S}_{n-m+1}/\mathcal{S}_{n-m}$, i.e. $e_i(\sigma^{(m)})=c_i(\mathcal{S}_{n-m}), e_i(\eta^{(m)})=c_i(\mathcal{S}_{n-m+1}/\mathcal{S}_{n-m})$, then Eq.(3.21) becomes⁴

$$c(S_i) \cdot c(S_{i+1}/S_i) = c(S_{i+1}) + (-1)^{k_{n-i-1}-k_{n-i}} q_{n-i} c(S_{i-1}), \tag{3.22}$$

for $i=1,2,\cdots,n-1$, which is the quantum Whitney relation of the equivariant quantum cohomology ring $QH_T^*(\mathrm{Fl}(k_{n-1},\cdots,k_2,k_1;N))$.

When $\beta \neq 0$, let $X_a^{(m)} = 1 + \beta \sigma_a^{(m)}$, then Eq.(3.5)-(3.7) can be written as

$$\prod_{\alpha=1}^{k_{m-1}} \left(X_a^{(m)} - X_{\alpha}^{(m-1)} \right) \prod_{l=1}^{k_m} \left(X_l^{(m)} \right) \\
= (-1)^{k_m - 1} q_m \beta^{k_{m-1} - k_{m+1}} \prod_{s=1}^{k_{m-1}} \left(X_s^{(m-1)} \right) \prod_{\gamma=1}^{k_{m+1}} \left(X_{\gamma}^{(m+1)} - X_a^{(m)} \right) \left(X_a^{(m)} \right)^{k_m - k_{m+1}} \tag{3.23}$$

for $a=1, \dots, k_m, m=1, 2, \dots, n$, where $X_i^{(0)}=1+\beta t_i, k_0=N, k_n=0$. Eq.(3.23) implies that $X_a^{(m)}, a=1, \dots, k_m$, are roots of the polynomial $\sum_{l=0}^{k_{m-1}} a_l x^l$, where

$$a_{k_{m-1}-l} = (-1)^{l} e_{l}(X^{(m-1)}) e_{k_{m}}(X^{(m)})$$

$$+ (-1)^{k_{m-1}-k_{m+1}+l} q_{m} \beta^{k_{m-1}-k_{m+1}} e_{k_{m-1}}(X^{(m-1)}) e_{l-k_{m-1}+k_{m}}(X^{(m+1)}).$$

Assume the roots of $\sum_{l=0}^{k_{m-1}} a_l x^l$ are $X_a^{(m)}, a = 1, \dots, k_m$, and $Y_b^{(m)}, b = 1, \dots, k_{m-1} - k_m$, then

³We set $e_l(x_1, \dots, x_s) = 0$ for l < 0 or l > k.

 $^{^4}S_0$ is the trivial vector bundle of rank zero, so $c(S_0)=1$. S_n is the trivial vector bundle of rank N, but since it carries the T-action, we have $c(S_n)=\sum_{i=0}^N e_i(t_1,\cdots,t_N)$.

Vieta's formula yields

$$\begin{aligned} &e_{k_m}(X^{(m)}) \sum_{i+j=l} e_i(X^{(m)}) e_j(Y^{(m)}) \\ = &e_{k_m}(X^{(m)}) e_l(X^{(m-1)}) + q_m(-\beta)^{k_{m-1}-k_{m+1}} e_{k_{m-1}}(X^{(m-1)}) e_{l-k_{m-1}+k_m}(X^{(m+1)}). \end{aligned} \tag{3.24}$$

Taking $l = k_{m-1}$ in Eq.(3.24), we get

$$e_{k_{m-1}}(X^{(m-1)}) = e_{k_m}(X^{(m)})e_{k_{m-1}-k_m}(Y^{(m)}).$$
(3.25)

Eq.(3.24) and (3.25) together give us the relation

$$\sum_{i+j=l} e_i(X^{(m)})e_j(Y^{(m)}) = e_l(X^{(m-1)}) + q_m(-\beta)^{k_{m-1}-k_{m+1}}e_{l-k_{m-1}+k_m}(X^{(m+1)})e_{k_{m-1}-k_m}(Y^{(m)}).$$
(3.26)

In the case $\beta = -1$, (3.26) was interpreted as the quantum Whitney relation of the equivariant quantum K-theory ring of the flag variety in [11]. The idea is to identify $e_i(X^{(m)})$ as $\wedge^i \mathcal{S}_{n-m}$ and make the following identification

$$e_l(Y^{(m)}) = \begin{cases} \wedge^l (\mathcal{S}_{n-m+1}/\mathcal{S}_{n-m}), & l < k_{m-1} - k_m, \\ \frac{1}{1 - q_m} \det(\mathcal{S}_{n-m+1}/\mathcal{S}_{n-m}), & l = k_{m-1} - k_m. \end{cases}$$

Then Eq.(3.26) for $\beta = -1$ can be written in terms of λ_y class as⁵

$$\lambda_{y}(S_{i}) * \lambda_{y}(S_{i+1}/S_{i}) = \lambda_{y}(S_{i+1}) + \frac{q_{n-i}}{1 - q_{n-i}} y^{k_{n-i-1}-k_{n-i}} (\lambda_{y}(S_{i-1}) - \lambda_{y}(S_{i})) * \det(S_{i+1}/S_{i}), \quad (3.27)$$

which is the quantum Whitney relation of the equivariant quantum K-theory ring $QK^*(Fl(k_{n-1},\dots,k_2,k_1;N))$ proposed in [11].

Remark 1. Notice that in [11], the analogue of the Bethe ansatz equations were derived from the $3d \mathcal{N} = 2$ gauged linear sigma model (GLSM) for flag varieties compactified on a circle. These GLSMs are quiver gauge theories, whose quiver diagram is as follows in our convention [12]

$$(k_{n-1}) \leftarrow (k_{n-2}) \leftarrow (k_1) \leftarrow N$$

where circles represent $U(k_i)$ gauge groups, square represents U(N) global symmetry and arrows represent chiral fields in bifundamental representations. The Higgs branch of the GLSM is a nonlinear sigma model with target space being the flag variety, the equations for vacuum states on the Coulomb branch are exactly the Bethe ansatz equations we obtained above in the case of $\beta = -1$. Though ring relations of quantum K-theory ring of general partial flag varieties have not been rigorously established, it was checked in [11] that the ring relations obtained from the GLSMs match those of the special cases, such as incidence varieties and full flag varieties, in which exact results have been proved.

Remark 2. Theorem 3.1 shows that the Bethe ansatz equations of the quantum integrable model defined by the R-matrix (2.3) and monodromy matrix (2.17) reduce to the vacuum equations on

⁵For a vector bundle \mathcal{E} , the λ_y class is defined as $\lambda_y(\mathcal{E}) = \sum_{i=0}^{\operatorname{rank}(\mathcal{E})} y^i \wedge^i \mathcal{E}$ in the Grothendieck group.




Figure 1: The left action of $B_k(u)$ on $|n_1, n_2, \dots, n_N\rangle$ for N=12 and a specific choice of (m_1, \dots, m_{N-1}) in the expansion (4.2).

the Coulomb branch of the 2d or 3d GLSM for flag varieties when $\beta = 0$ or $\beta = -1$ respectively. Therefore, we have established a Bethe/Gauge correspondence in the sense of [13, 14].

4 Relationship with the double β -Grothendieck polynomials

In this section, we consider the case of full flag variety, i.e. n = N and $k_{N-i} = i$ for $i = 1, \dots, N-1$. We will show that the Bethe ansatz states generate the double β -Grothendieck polynomials (Theorem 4.6 and 4.11). The double β -Grothendieck polynomials $\mathcal{G}_w^{(\beta)}(\boldsymbol{x};\boldsymbol{y})$ for $w \in S_N$ and $\boldsymbol{x} = (x_1, \dots, x_N), \boldsymbol{y} = (y_1, \dots, y_N)$ are defined in the appendix, see Eq.(A.1)-(A.3).

In addition to Eq.(2.2), let us also define

$$\ominus t := \frac{-t}{1 + \beta t} \tag{4.1}$$

and $\ominus \mathbf{t} := (\ominus t_1, \dots, \ominus t_N)$ for $\mathbf{t} = (t_1, \dots, t_N)$. We use ω_N to denote the permutation of N elements with maximal length.

From Eq.(2.4), (2.17) and the fact that $B_k(u) = [T_a^{(n)}(u)]_{0,k}$, it is easy to compute

$$B_{k}(u) | n_{1}, n_{2}, \dots, n_{N} \rangle =$$

$$= \sum_{m_{1}=0}^{N-1} \dots \sum_{m_{N-1}=0}^{N-1} R_{aN}(u, t_{N})_{0, m_{N-1}} \dots R_{a2}(u, t_{2})_{m_{2}, m_{1}} R_{a1}(u, t_{1})_{m_{1}, k} | n_{1}, n_{2}, \dots, n_{N} \rangle$$

$$= \sum_{m_{1}=0}^{N-1} \dots \sum_{m_{N-1}=0}^{N-1} \bigotimes_{i=1}^{N} \left[\delta_{m_{i}, n_{i}} (1 + \beta(u \ominus t_{i}) \delta_{m_{i} > m_{i-1}}) | m_{i-1} \rangle + \delta_{n_{i} < m_{i}} (u \ominus t_{i}) \delta_{m_{i}, m_{i-1}} | n_{i} \rangle \right],$$

$$(4.2)$$

where $m_0 = k, m_N = 0$. The state (4.2) vanishes unless $m_i \ge n_i$ for all $1 \le i \le N$, therefore it vanishes when $n_N \ne 0$. Also, $m_i = m_{i-1}$ if $m_i > n_i$.

The left action of $B_k(u)$ on $|n_1, n_2, \dots, n_N\rangle$ can then be depicted as Fig.1 (for a specific (m_1, \dots, m_{N-1})). We see for any (m_1, \dots, m_{N-1}) in the expansion (4.2), the effect of the operator $B_k(u)$ is to annihilate a $|0\rangle$ state, create a $|k\rangle$ state, and permute n_i among the sites. If we write




Figure 2: The right action of $B_k(u)$ on $\langle n_1, n_2, \dots, n_N |$ for N = 12 and a specific choice of (m_2, \dots, m_N) in the expansion (4.4).

the expansion (4.2) as $B_k(u) | \mathbf{n} \rangle = \sum_{\mathbf{m}} C_{\mathbf{m}} | \psi_{\mathbf{m}} \rangle$, then for any $n_i < m_i$, there is a factor $(u \ominus t_i)$ in $C_{\mathbf{m}}$, and for any $m_i > m_{i-1}$, there is a factor $(1 + \beta(u \ominus t_i))$ in $C_{\mathbf{m}}$.

For $w \in S_N$, let us define

$$|w\rangle := |w(012\cdots N - 1)\rangle, \tag{4.3}$$

then, from (4.2) and the discussion above, we have the following

Lemma 4.1. For any $\pi \in S_N$, we have the expansion

$$B_{\pi(1)}(u_1)B_{\pi(2)}(u_2)\cdots B_{\pi(N-1)}(u_{N-1})|\Omega^{(0)}\rangle = \sum_{w\in S_N} C_w^{\pi}(u_1,\cdots,u_{N-1})|w\rangle,$$

where $C_w^{\pi} \in \mathbb{C}[u_1, \cdots, u_{N-1}, t_1, \cdots, t_N]$ and $|\Omega^{(0)}\rangle$ is defined by Eq.(3.2).

Similarly, the right action of $B_k(u)$ on the dual state $\langle n_1, n_2, \cdots, n_N |$ is

$$\langle n_{1}, n_{2}, \cdots, n_{N} | B_{k}(u)$$

$$= \sum_{m_{2}=0}^{N-1} \cdots \sum_{m_{N}=0}^{N-1} \langle n_{1}, n_{2}, \cdots, n_{N} | R_{aN}(u, t_{N})_{0, m_{N}} \cdots R_{a2}(u, t_{2})_{m_{3}, m_{2}} R_{a1}(u, t_{1})_{m_{2}, k}$$

$$= \sum_{m_{2}=0}^{N-1} \cdots \sum_{m_{N}=0}^{N-1} \bigotimes_{i=1}^{N} \left[\delta_{m_{i}, n_{i}} (1 + \beta(u \ominus t_{i}) \delta_{m_{i} < m_{i+1}}) \langle m_{i+1} | + \delta_{n_{i} < m_{i}} (u \ominus t_{i}) \delta_{m_{i}, m_{i+1}} \langle n_{i} | \right],$$

$$(4.4)$$

where $m_1 = k$, $m_{N+1} = 0$. The state (4.4) vanishes unless $m_i \ge n_i$ for all $1 \le i \le N$ and there is some n_a such that $n_a = k = m_a$ and $m_j > n_j$ for all j < a. Moreover, $m_i = m_{i+1}$ if $m_i > n_i$.

The right action of $B_k(u)$ on $\langle n_1, n_2, \dots, n_N |$ can therefore be depicted as Fig.2 (for a specific choice of (m_2, \dots, m_N)). We see for any (m_2, \dots, m_N) in the expansion (4.4), the effect of the operator $B_k(u)$ is to annihilate a $\langle k |$ state, create a $\langle 0 |$ state on the Nth site, and permute n_i among the other sites. If we write the expansion (4.4) as $\langle n | B_k(u) = \sum_{m} \langle \psi_m | C_m$, then for any $n_i < m_i$, there is a factor $(u \ominus t_i)$ in C_m , and for any $m_i < m_{i+1}$, there is a factor $(1 + \beta(u \ominus t_i))$ in C_m .

For $w \in S_N$, define

$$\langle w| := \langle w(012\cdots N - 1)|, \tag{4.5}$$

then, from the expansion Eq.(4.4), one can prove

Lemma 4.2. If $(n_1, n_2, \dots, n_{N-1})$ satisfies: (i) $0 \le n_i < k$; (ii) $n_{l+1} \ge n_{l+2} \ge \dots \ge n_{N-1}$ for some $l \ge 0$, then

$$\langle \pi(n_1, \dots, n_l), k, n_{l+1}, \dots, n_{N-1} | B_k(u) = \prod_{i=1}^l (u \ominus t_i) \langle \pi(n_1, \dots, n_l), n_{l+1}, \dots, n_{N-1}, 0 |$$

for any $\pi \in S_l$.

Proof. Given the constraints on n_i , the only possible (m_2, m_3, \dots, m_N) in the expansion (4.4) is

$$m_i = \begin{cases} k, & 2 \le i \le l+1, \\ n_i, & l+2 \le i \le N. \end{cases}$$

Since $n_{\pi(i)} < m_i$ for $1 \le i \le l$, we have the factor $\prod_{i=1}^{l} (u \ominus t_i)$ from (4.4).

Let us define

$$\rho_k := s_1 s_2 \cdots s_k \in S_N, \tag{4.6}$$

where s_i is the simple transposition of the *i*th and (i+1)th elements, and ρ_0 is defined to be the identity permutation. Let ∂_i^{β} be the β -divided difference operator defined by Eq.(A.1):

$$\partial_i^{\beta} f(x_1, \dots, x_N) := \frac{(1 + \beta x_{i+1}) f(x_1, \dots, x_N) - (1 + \beta x_i) f(x_1, \dots, x_{i+1}, x_i, \dots, x_N)}{x_i - x_{i+1}}.$$

Generally, for $w = s_{i_1} s_{i_2} \cdots s_{i_{l(w)}}$, we have the definition⁶

$$\partial_w^{\beta} := \partial_{i_1}^{\beta} \partial_{i_2}^{\beta} \cdots \partial_{i_{l(w)}}^{\beta}. \tag{4.7}$$

Now we can use Lemma 4.2 to prove the following

Lemma 4.3.

$$\langle 1, 2, \cdots, k, 0, k+1, \cdots, N-1 | B_{N-1}(\sigma_1) \cdots B_1(\sigma_{N-1}) = \mathcal{G}_{\rho_{b_1}^{-1}\omega_N}^{(\beta)}(\boldsymbol{\sigma}; \ominus \boldsymbol{t}) \langle \Omega^{(0)} | .$$

Proof. By repeated use of Lemma 4.2, one can compute

$$\langle 1, 2, \cdots, k, 0, k+1, \cdots, N-1 | B_{N-1}(\sigma_1) \cdots B_1(\sigma_{N-1})$$

$$= \prod_{a=1}^{N-k-1} \prod_{i=1}^{N-a} (\sigma_a \ominus t_i) \prod_{b=N-k}^{N-2} \prod_{j=1}^{N-b-1} (\sigma_b \ominus t_j) \langle \Omega^{(0)} | .$$
(4.8)

Notice that $\sigma\ominus t=\frac{\sigma-t}{1+\beta t}=\sigma+\ominus t+\beta\sigma(\ominus t),$ Therefore, from Eq.(A.2)

$$\prod_{i+j\leq N} (\sigma_i \ominus t_j) = \prod_{i+j\leq N} (\sigma_i + \ominus t_j + \beta \sigma_i (\ominus t_j)) = \mathcal{G}_{\omega_N}^{(\beta)}(\boldsymbol{\sigma}; \ominus \boldsymbol{t}),$$

⁶In the following, if a polynomial $f(\boldsymbol{x}; \boldsymbol{y})$ depends on two sets of variables \boldsymbol{x} and \boldsymbol{y} as in the case of double β -Grothendieck polynomials, ∂_w^{β} in $\partial_w^{\beta} f(\boldsymbol{x}; \boldsymbol{y})$ is understood to act on the first set of variables, see Eq.(A.1). The same convention applies to the operator $\bar{\partial}_w^{\beta}$ defined by Eq.(4.27).

then, from (4.8) and the definition of the β -Grothendieck polynomials (A.1)-(A.3), we get

$$\langle 1, 2, \cdots, k, 0, k+1, \cdots, N-1 | B_{N-1}(\sigma_1) \cdots B_1(\sigma_{N-1})$$

$$= \frac{\mathcal{G}_{\omega_N}^{(\beta)}(\boldsymbol{\sigma}; \ominus \boldsymbol{t})}{(\sigma_{N-1} \ominus t_1)(\sigma_{N-2} \ominus t_2) \cdots (\sigma_{N-k} \ominus t_k)} \langle \Omega^{(0)} |$$

$$= \partial_{N-1}^{\beta} \partial_{N-2}^{\beta} \cdots \partial_{N-k}^{\beta} \mathcal{G}_{\omega_N}^{(\beta)}(\boldsymbol{\sigma}; \ominus \boldsymbol{t}) \langle \Omega^{(0)} |$$

$$= \mathcal{G}_{\omega_N s_{N-k} \cdots s_{N-2} s_{N-1}}^{(\beta)}(\boldsymbol{\sigma}; \ominus \boldsymbol{t}) \langle \Omega^{(0)} |$$

$$= \mathcal{G}_{s_k \cdots s_2 s_1 \omega_N}^{(\beta)}(\boldsymbol{\sigma}; \ominus \boldsymbol{t}) \langle \Omega^{(0)} | = \mathcal{G}_{\rho_k^{-1} \omega_N}^{(\beta)}(\boldsymbol{\sigma}; \ominus \boldsymbol{t}) \langle \Omega^{(0)} |,$$

where in the second to last step, we used $\omega_N s_{N-i} = s_i \omega_N$.

Lemma 4.4. If $\pi \in S_N$ and $l(s_i\pi) > l(\pi)$, then

$$(\langle s_i \pi | + \beta \langle \pi |) B_{N-1}(\sigma_1) B_{N-2}(\sigma_2) \cdots B_{i+1}(\sigma_{N-i-1}) B_i(\sigma_{N-i}) B_{i-1}(\sigma_{N-i+1})$$

= $\langle \pi | B_{N-1}(\sigma_1) B_{N-2}(\sigma_2) \cdots B_{i+1}(\sigma_{N-i-1}) B_{i-1}(\sigma_{N-i}) B_i(\sigma_{N-i+1}).$

Proof. First we show, for

$$\langle \psi | = \langle n_1, \cdots, n_{a-1}, i-1, n_{a+1}, \cdots, n_{b-1}, i, n_{b+1}, \cdots, n_N |$$
 (4.9)

and

$$\langle s_i \psi | = \langle n_1, \dots, n_{a-1}, i, n_{a+1}, \dots, n_{b-1}, i-1, n_{b+1}, \dots, n_N |$$
 (4.10)

with $n_l < i - 1, l \neq a, b$, we have

$$(\langle s_i \psi | + \beta \langle \psi |) B_i(x) B_{i-1}(y) = \langle \psi | B_{i-1}(x) B_i(y). \tag{4.11}$$

Given the following expansions in the natural basis according to (4.4)

$$\langle \psi | B_{i-1}(x) = \sum_{\boldsymbol{m} \in M_{i-1}} C_{\boldsymbol{m}}^{(i-1)} \langle \psi_{\boldsymbol{m}}^{(i-1)} |,$$

$$\langle \psi | B_{i}(x) = \sum_{\boldsymbol{m} \in M_{i}} C_{\boldsymbol{m}}^{(i)} \langle \psi_{\boldsymbol{m}}^{(i)} |,$$

$$\langle s_{i}\psi | B_{i}(x) = \sum_{\boldsymbol{m} \in \bar{M}_{i}} \bar{C}_{\boldsymbol{m}}^{(i)} \langle \bar{\psi}_{\boldsymbol{m}}^{(i)} |,$$

where M_{i-1}, M_i and \bar{M}_i are the sets in which $\mathbf{m} = (m_1, \dots, m_N)$ takes value such that the corresponding coefficients $C_{\mathbf{m}}^{(i-1)}, C_{\mathbf{m}}^{(i)}$ and $\bar{C}_{\mathbf{m}}^{(i)}$ in (4.4) are nonzero. $\langle \psi_{\mathbf{m}}^{(i-1)} |, \langle \psi_{\mathbf{m}}^{(i)} |$ and $\langle \bar{\psi}_{\mathbf{m}}^{(i)} |$ are members of the natural basis (3.1). There is a map $\phi: M_{i-1} \to \bar{M}_i$ defined by

$$\phi(\mathbf{m})_j = \begin{cases} i, & \text{if } m_j = i - 1, \\ i - 1, & \text{if } m_j = i, \\ m_j, & \text{if } m_j \neq i \text{ and } m_j \neq i - 1. \end{cases}$$

 ϕ is a bijection since $n_l < i-1$ for $l \neq a,b$. Moreover, since we have $n_j < m_j$ for $1 \leq j < b$ and

 $^{^{7}}l(w)$ denotes the length of the permutation w.

 $m \in M_i$, we have an injective map $\tau: M_i \to \bar{M}_i$ defined by

$$\tau(\boldsymbol{m})_j = \begin{cases} i, & 1 \le j \le a, \\ i-1, & a+1 \le j \le b, \\ m_j, & j > b. \end{cases}$$

Eq.(4.4) implies, for $\mathbf{m} \in \text{im}\tau$,

$$C_{\phi^{-1}(\mathbf{m})}^{(i-1)} = (1 + \beta(x \ominus t_a))\bar{C}_{\mathbf{m}}^{(i)}, \quad C_{\tau^{-1}(\mathbf{m})}^{(i)} = (x \ominus t_a)\bar{C}_{\mathbf{m}}^{(i)}. \tag{4.12}$$

On the other hand, if $m \notin \text{im}\tau$, then

$$C_{\phi^{-1}(\mathbf{m})}^{(i-1)} = \bar{C}_{\mathbf{m}}^{(i)}. \tag{4.13}$$

Therefore,

$$\langle \psi | B_{i-1}(x) = \sum_{\phi^{-1}(m) \in M_{i-1}} C_{\phi^{-1}(m)}^{(i-1)} \langle \psi_{\phi^{-1}(m)}^{(i-1)} |$$

$$= \sum_{m \in \text{im}\tau} (1 + \beta(x \ominus t_a)) \bar{C}_{m}^{(i)} \langle \psi_{\phi^{-1}(m)}^{(i-1)} | + \sum_{m \notin \text{im}\tau} \bar{C}_{m}^{(i)} \langle \psi_{\phi^{-1}(m)}^{(i-1)} |,$$

$$\langle \psi | B_{i}(x) = \sum_{\bar{m} \in M_{i}} C_{\bar{m}}^{(i)} \langle \psi_{\bar{m}}^{(i)} | = \sum_{\bar{m} \in M_{i}} (x \ominus t_a) \bar{C}_{\tau(\bar{m})}^{(i)} \langle \psi_{\bar{m}}^{(i)} | = \sum_{m \in \text{im}\tau} (x \ominus t_a) \bar{C}_{m}^{(i)} \langle \psi_{\tau^{-1}(m)}^{(i)} |,$$

$$\langle s_{i}\psi | B_{i}(x) = \sum_{m \in \bar{M}_{i}} \bar{C}_{m}^{(i)} \langle \bar{\psi}_{m}^{(i)} | = \sum_{m \in \text{im}\tau} \bar{C}_{m}^{(i)} \langle \bar{\psi}_{m}^{(i)} | + \sum_{m \notin \text{im}\tau} \bar{C}_{m}^{(i)} \langle \bar{\psi}_{m}^{(i)} |,$$

$$(4.14)$$

where, according to (4.4),

$$\langle \psi_{\phi^{-1}(\boldsymbol{m})}^{(i-1)} | = \langle n_1, \cdots, n_{a-1}, i, n_{a+1}, \cdots, n_{b-1}, n_{s_1}, \cdots, n_{s_{N-b+1}} |,$$

$$\langle \bar{\psi}_{\tau^{-1}(\boldsymbol{m})}^{(i)} | = \langle n_1, \cdots, n_{a-1}, i-1, n_{a+1}, \cdots, n_{b-1}, n_{s_1}, \cdots, n_{s_{N-b+1}} |,$$

$$\langle \bar{\psi}_{\boldsymbol{m}}^{(i)} | = \langle n_1, \cdots, n_{a-1}, i-1, n_{a+1}, \cdots, n_{b-1}, n_{s_1}, \cdots, n_{s_{N-b+1}} |,$$

with some $n_{s_1}, \dots, n_{s_{N-b+1}}$ determined by \boldsymbol{m} . From the equations above, we see $\langle \bar{\psi}_{\boldsymbol{\tau}^{-1}(\boldsymbol{m})}^{(i)} | = \langle \bar{\psi}_{\boldsymbol{m}}^{(i)} |$ and $\langle \psi_{\phi^{-1}(\boldsymbol{m})}^{(i-1)} | B_i(y) = \langle \bar{\psi}_{\boldsymbol{m}}^{(i)} | B_{i-1}(y)$, from which, together with Eq.(4.14), we compute

$$\begin{aligned} & (\langle s_{i}\psi| + \beta \langle \psi|)B_{i}(x)B_{i-1}(y)) \\ &= \sum_{\boldsymbol{m} \notin \operatorname{im}\tau} \bar{C}_{\boldsymbol{m}}^{(i)} \langle \bar{\psi}_{\boldsymbol{m}}^{(i)}| \, B_{i-1}(y) + \sum_{\boldsymbol{m} \in \operatorname{im}\tau} \bar{C}_{\boldsymbol{m}}^{(i)} \langle \bar{\psi}_{\boldsymbol{m}}^{(i)}| \, B_{i-1}(y) + \beta \sum_{\boldsymbol{m} \in \operatorname{im}\tau} (x \ominus t_{a}) \bar{C}_{\boldsymbol{m}}^{(i)} \langle \bar{\psi}_{\boldsymbol{m}}^{(i)}| \, B_{i-1}(y) \\ &= \sum_{\boldsymbol{m} \notin \operatorname{im}\tau} \bar{C}_{\boldsymbol{m}}^{(i)} \langle \psi_{\phi^{-1}(\boldsymbol{m})}^{(i-1)}| \, B_{i}(y) + \sum_{\boldsymbol{m} \in \operatorname{im}\tau} (1 + \beta(x \ominus t_{a})) \bar{C}_{\boldsymbol{m}}^{(i)} \langle \psi_{\phi^{-1}(\boldsymbol{m})}^{(i-1)}| \, B_{i}(y) \\ &= \sum_{\phi^{-1}(\boldsymbol{m}) \in M_{i-1}} C_{\phi^{-1}(\boldsymbol{m})}^{(i-1)} \langle \psi_{\phi^{-1}(\boldsymbol{m})}^{(i-1)}| \, B_{i}(y) \\ &= \langle \psi| \, B_{i-1}(x)B_{i}(y), \end{aligned}$$

where (4.12) and (4.13) were used in the third equality. Thus Eq.(4.11) is proved.

Now assume, for
$$k \geq i+1$$
 and $\langle \varphi | = \langle n_1, \cdots, n_N | \text{ with } \{n_1, \cdots, n_N\} = \{1, 2, \cdots, k-1\}$

 $1, 0, \cdots, 0$, we have

$$(\langle s_i \varphi | + \beta \langle \varphi |) B_{k-1}(\sigma_1) \cdots B_i(\sigma_{k-i}) B_{i-1}(\sigma_{k-i+1}) = \langle \varphi | B_{k-1}(\sigma_1) \cdots B_{i-1}(\sigma_{k-i}) B_i(\sigma_{k-i+1}),$$

$$(4.15)$$

where $\langle \varphi |$ and $\langle s_i \varphi |$ are defined in the same way as Eq.(4.9) and (4.10). Now let $\langle \psi |$ be a state of the form

$$\langle \psi | = \langle n_1, \cdots, n_{a-1}, i-1, n_{a+1}, \cdots, n_{b-1}, i, n_{b+1}, \cdots, n_N |$$

with $n_l \leq k$ and $n_i \neq n_j$ for $i \neq j$ unless $n_i = n_j = 0$. We expand the following states in the natural basis according to (4.4):

$$\langle \psi | B_k(u) = \sum_{\boldsymbol{m} \in M_1} C_{\boldsymbol{m}} \langle \psi_{\boldsymbol{m}} | + \sum_{\boldsymbol{m} \in M_2} C'_{\boldsymbol{m}} \langle \psi'_{\boldsymbol{m}} |,$$
 (4.16)

$$\langle s_i \psi | B_k(u) = \sum_{\boldsymbol{m} \in \bar{M}_1} \bar{C}_{\boldsymbol{m}} \langle s_i \psi_{\boldsymbol{m}} |, \qquad (4.17)$$

where M_2 contains \boldsymbol{m} 's satisfying $i-1=n_a < m_a=m_{a+1}=\cdots=m_b=i$, and M_1 consists of all the other \boldsymbol{m} 's such that the corresponding coefficients are nonzero. There is a one-to-one correspondence between M_1 and \bar{M}_1 given by $\tilde{\phi}: M_1 \to \bar{M}_1$ with $\tilde{\phi}$ defined by

$$\tilde{\phi}(\boldsymbol{m})_{j} = \begin{cases} i, & \text{if } m_{j} = i - 1, \\ i - 1, & \text{if } m_{j} = i, \\ m_{j}, & \text{if } m_{j} \neq i \text{ and } m_{j} \neq i - 1, \end{cases}$$

and from Eq.(4.4)

$$C_{\boldsymbol{m}} = \begin{cases} (1 + \beta(u \ominus t_a)) \bar{C}_{\tilde{\phi}(\boldsymbol{m})}, & \text{if } m_a = i - 1, m_b = i, n_s < m_s = m_b \text{ for } a < s < b, \\ \bar{C}_{\tilde{\phi}(\boldsymbol{m})}, & \text{otherwise,} \end{cases}$$

and there is an injective map $\tilde{\tau}: M_2 \to M_1$ given by

$$\tilde{\tau}(\boldsymbol{m})_j = \begin{cases} i-1, & \text{if } m_j = i \text{ and } j \leq a, \\ m_j, & \text{otherwise,} \end{cases}$$

Notice that Eq.(4.4) also implies

$$C_{\boldsymbol{m}} = \begin{cases} (1 + \beta(u \ominus t_a)) \bar{C}_{\tilde{\phi}(\boldsymbol{m})}, & \text{if } \boldsymbol{m} \in \text{im}\tilde{\tau}, \\ \bar{C}_{\tilde{\phi}(\boldsymbol{m})}, & \text{otherwise.} \end{cases}$$
(4.18)

and, if $m \in \text{im}\tilde{\tau}$, then

$$\langle s_i \psi_{\tilde{\phi}(\boldsymbol{m})} | = \langle \psi'_{\tilde{\tau}^{-1}(\boldsymbol{m})} |, \quad C'_{\tilde{\tau}^{-1}(\boldsymbol{m})} = (u \ominus t_a) \bar{C}_{\tilde{\phi}(\boldsymbol{m})}.$$

Therefore

$$\sum_{\boldsymbol{m}\in M_{1}} \bar{C}_{\tilde{\phi}(\boldsymbol{m})} \langle s_{i}\psi_{\tilde{\phi}(\boldsymbol{m})}| + \beta \sum_{\boldsymbol{m}\in M_{2}} C'_{\boldsymbol{m}} \langle \psi'_{\boldsymbol{m}}|
= \sum_{\boldsymbol{m}\in M_{1}\backslash \text{im}\tilde{\tau}} \bar{C}_{\tilde{\phi}(\boldsymbol{m})} \langle s_{i}\psi_{\tilde{\phi}(\boldsymbol{m})}| + \sum_{\boldsymbol{m}\in \text{im}\tilde{\tau}} \bar{C}_{\tilde{\phi}(\boldsymbol{m})} \langle s_{i}\psi_{\tilde{\phi}(\boldsymbol{m})}| + \beta \sum_{\boldsymbol{m}\in M_{2}} C'_{\boldsymbol{m}} \langle \psi'_{\boldsymbol{m}}|
= \sum_{\boldsymbol{m}\in M_{1}\backslash \text{im}\tilde{\tau}} \bar{C}_{\tilde{\phi}(\boldsymbol{m})} \langle s_{i}\psi_{\boldsymbol{m}}| + \sum_{\boldsymbol{m}\in \text{im}\tilde{\tau}} \bar{C}_{\tilde{\phi}(\boldsymbol{m})} \langle s_{i}\psi_{\tilde{\phi}(\boldsymbol{m})}| + \beta(u\ominus t_{a}) \sum_{\boldsymbol{m}\in \text{im}\tilde{\tau}} \bar{C}_{\tilde{\phi}(\boldsymbol{m})} \langle s_{i}\psi_{\tilde{\phi}(\boldsymbol{m})}|
= \sum_{\boldsymbol{m}\in M_{1}} C_{\boldsymbol{m}} \langle s_{i}\psi_{\boldsymbol{m}}|,$$

$$(4.19)$$

where in the last equality, we used Eq.(4.18). Consequently,

$$(\langle s_{i}\psi| + \beta \langle \psi|)B_{k}(u))$$

$$= \sum_{\boldsymbol{m}\in M_{1}} \bar{C}_{\tilde{\phi}(\boldsymbol{m})} \langle s_{i}\psi_{\tilde{\phi}(\boldsymbol{m})}| + \beta \sum_{\boldsymbol{m}\in M_{2}} C'_{\boldsymbol{m}} \langle \psi'_{\boldsymbol{m}}| + \beta \sum_{\boldsymbol{m}\in M_{1}} C_{\boldsymbol{m}} \langle \psi_{\boldsymbol{m}}|$$

$$= \sum_{\boldsymbol{m}\in M_{1}} C_{\boldsymbol{m}} (\langle s_{i}\psi_{\boldsymbol{m}}| + \beta \langle \psi_{\boldsymbol{m}}|),$$

$$(4.20)$$

where the second equality is due to (4.19). From the induction hypothesis Eq.(4.15), we have

$$(\langle s_i \psi_{\boldsymbol{m}} | + \beta \langle \psi_{\boldsymbol{m}} |) B_{k-1}(\sigma_1) \cdots B_i(\sigma_{k-i}) B_{i-1}(\sigma_{k-i+1}) = \langle \psi_{\boldsymbol{m}} | B_{k-1}(\sigma_1) \cdots B_{i-1}(\sigma_{k-i}) B_i(\sigma_{k-i+1}),$$

$$(4.21)$$

which implies

$$(\langle s_{i}\psi| + \beta \langle \psi|)B_{k}(u)B_{k-1}(\sigma_{1})\cdots B_{i}(\sigma_{k-i})B_{i-1}(\sigma_{k-i+1})$$

$$= \sum_{\boldsymbol{m}\in M_{1}} C_{\boldsymbol{m}}(\langle s_{i}\psi_{\boldsymbol{m}}| + \beta \langle \psi_{\boldsymbol{m}}|)B_{k-1}(\sigma_{1})\cdots B_{i}(\sigma_{k-i})B_{i-1}(\sigma_{k-i+1})$$

$$= \sum_{\boldsymbol{m}\in M_{1}} C_{\boldsymbol{m}} \langle \psi_{\boldsymbol{m}}| B_{k-1}(\sigma_{1})\cdots B_{i-1}(\sigma_{k-i})B_{i}(\sigma_{k-i+1})$$

$$= \langle \psi| B_{k}(u)B_{k-1}(\sigma_{1})\cdots B_{i-1}(\sigma_{k-i})B_{i}(\sigma_{k-i+1}),$$

$$(4.22)$$

where the first equality is due to (4.20), the second equality is due to (4.21), and in the last step we used Eq.(4.16) and the fact that, for $m \in M_2$, i is to the left of i-1 in $\langle \psi'_{m}|$, so (notice that B_l with l > i cannot move i-1 to the left of i)

$$\langle \psi'_{m} | B_{k-1}(\sigma_1) \cdots B_{i-1}(\sigma_{k-i}) B_i(\sigma_{k-i+1}) = 0, \quad m \in M_2.$$

Combining Eq.(4.11), (4.15) and (4.22), the lemma is proved by induction.

In order to relate the quantum states of the integrable system to the double β -Grothendieck polynomials, we need the commutation relations among the different B_i operators. From Eq.(2.4) and (2.25), we have the commutation relations

$$B_i(x)B_i(y) = B_i(y)B_i(x), \tag{4.23}$$

$$B_{i}(x)B_{j}(y) = \frac{B_{j}(y)B_{i}(x) - B_{j}(x)B_{i}(y)}{y \ominus x} = (1 + \beta x)\frac{B_{j}(x)B_{i}(y) - B_{j}(y)B_{i}(x)}{x - y}.$$
 (4.24)

Let us define

$$\bar{\partial}_i^{\beta} = (1 + \beta x_i)\partial_i, \tag{4.25}$$

where ∂_i is the ordinary divided difference operator defined by $\partial_i^{\beta}|_{\beta=0}$. It is easy to check that

$$\bar{\partial}_i^{\beta} = \partial_i^{\beta} + \beta. \tag{4.26}$$

Similar to Eq.(4.7), for $w = s_{i_1} s_{i_2} \cdots s_{i_{l(w)}}$, we have

$$\bar{\partial}_w^{\beta} := \bar{\partial}_{i_1}^{\beta} \bar{\partial}_{i_2}^{\beta} \cdots \bar{\partial}_{i_{l(w)}}^{\beta}. \tag{4.27}$$

From Eq.(4.24), (4.25) and Lemma 4.4, we can prove

Lemma 4.5. For $\pi \in S_N$, if $l(s_i\pi) > l(\pi)$, then

$$\langle s_i \pi | B_{N-1}(\sigma_1) \cdots B_i(\sigma_{N-i}) B_{i-1}(\sigma_{N-i+1}) = \partial_{N-i}^{\beta} \langle \pi | B_{N-1}(\sigma_1) \cdots B_i(\sigma_{N-i}) B_{i-1}(\sigma_{N-i+1}).$$

Proof.

$$(\langle s_i \pi | + \beta \langle \pi |) B_{N-1}(\sigma_1) \cdots B_i(\sigma_{N-i}) B_{i-1}(\sigma_{N-i+1})$$

$$= \langle \pi | B_{N-1}(\sigma_1) \cdots B_{i-1}(\sigma_{N-i}) B_i(\sigma_{N-i+1})$$

$$= (\partial_{N-i}^{\beta} + \beta) \langle \pi | B_{N-1}(\sigma_1) \cdots B_i(\sigma_{N-i}) B_{i-1}(\sigma_{N-i+1}),$$

where the first equality is due to Lemma 4.4, and the second equality is due to Eq.(4.24), (4.25) and (4.26).

Theorem 4.6. For $\pi \in S_N$,

$$B_{N-1}(\sigma_1)B_{N-2}(\sigma_2)\cdots B_1(\sigma_{N-1})\left|\Omega^{(0)}\right\rangle = \sum_{w\in S_N} \mathcal{G}_w^{(\beta)}(\boldsymbol{\sigma};\ominus\boldsymbol{t})\left|\omega_N w^{-1}\right\rangle,$$

where ω_N is the permutation in S_N with maximal length.

Proof. Lemma 4.1 suggests that we only need to show

$$\langle \omega_N w^{-1} | B_{N-1}(\sigma_1) \cdots B_1(\sigma_{N-1}) | \Omega^{(0)} \rangle = \mathcal{G}_w^{(\beta)}(\boldsymbol{\sigma}; \ominus \boldsymbol{t}). \tag{4.28}$$

Lemma 4.3 shows Eq.(4.28) is valid for $\omega_N w^{-1} = \rho_k, k = 0, 1, \dots, N-1$. Because any permutation $w \in S_N$ can be written as $w = \pi \rho_k$ for some $\pi \in S_{N-1}$ acting on the last N-1 elements and some $k = 0, \dots, N-1$, it remains to show the validity of Eq.(4.28) for $\omega_N w^{-1} = \pi \rho_k$ with $\pi \in S_{N-1}$ acting on $\{1, 2, \dots, N-1\}$. We show this by induction on the length of π . Let us assume

$$\langle \tilde{\pi} \rho_k | B_{N-1}(\sigma_1) \cdots B_2(\sigma_{N-2}) B_1(\sigma_{N-1}) | \Omega^{(0)} \rangle = \mathcal{G}_{(\tilde{\pi} \rho_k)^{-1} \omega_N}^{(\beta)}(\boldsymbol{\sigma}; \boldsymbol{\ominus t})$$

for $\tilde{\pi} \in S_{N-1}$ with $l(\tilde{\pi}) \leq m-1$ for some $m \geq 1$. Now, for $\pi \in S_{N-1}$ with $l(\pi) = m$, there exists

 $\tilde{\pi} \in S_{N-1}$ with $l(\tilde{\pi}) = m-1$, such that $\pi = s_i \tilde{\pi}$. Then

$$\langle \pi \rho_{k} | B_{N-1}(\sigma_{1}) \cdots B_{1}(\sigma_{N-1}) | \Omega^{(0)} \rangle$$

$$= \langle s_{i} \tilde{\pi} \rho_{k} | B_{N-1}(\sigma_{1}) \cdots B_{1}(\sigma_{N-1}) | \Omega^{(0)} \rangle$$

$$= \partial_{N-i}^{\beta} \langle \tilde{\pi} \rho_{k} | B_{N-1}(\sigma_{1}) \cdots B_{1}(\sigma_{N-1}) | \Omega^{(0)} \rangle$$

$$= \partial_{N-i}^{\beta} \mathcal{G}_{\rho_{k}^{-1} \tilde{\pi}^{-1} \omega_{N}}^{(\beta)} (\boldsymbol{\sigma}; \ominus \boldsymbol{t})$$

$$= \mathcal{G}_{\rho_{k}^{-1} \tilde{\pi}^{-1} \omega_{N} s_{N-i}}^{(\beta)} (\boldsymbol{\sigma}; \ominus \boldsymbol{t}) = \mathcal{G}_{\rho_{k}^{-1} \tilde{\pi}^{-1} s_{i} \omega_{N}}^{(\beta)} (\boldsymbol{\sigma}; \ominus \boldsymbol{t})$$

$$= \mathcal{G}_{(\pi \rho_{k})^{-1} \omega_{N}}^{(\beta)} (\boldsymbol{\sigma}; \ominus \boldsymbol{t}),$$

where the second equality is due to Lemma 4.5.

Lemma 4.7. For any $w \in S_{N-1}$,

$$B_{w(1)}(\sigma_1)\cdots B_{w(N-1)}(\sigma_{N-1}) = \bar{\partial}_{w^{-1}\omega_{N-1}}^{\beta} B_{N-1}(\sigma_1)\cdots B_1(\sigma_{N-1}).$$

Proof. Let $w = \pi \omega_{N-1}$. When $\pi = s_k$, from Eq.(4.24),

$$B_{w(1)}(\sigma_1) \cdots B_{w(N-1)}(\sigma_{N-1})$$

$$= B_{N-1}(\sigma_1) \cdots B_k(\sigma_{N-k-1}) B_{k+1}(\sigma_{N-k}) \cdots B_1(\sigma_{N-1})$$

$$= \bar{\partial}_{N-k-1}^{\beta} B_{N-1}(\sigma_1) \cdots B_1(\sigma_{N-1})$$

$$= \bar{\partial}_{\omega_{N-1} s_k \omega_{N-1}}^{\beta} B_{N-1}(\sigma_1) \cdots B_1(\sigma_{N-1}).$$

The lemma can then be proved by induction on the length of π .

Remark 3. Lemma 4.7 allows us to deduce the matrix elements of $B_{w(1)}(\sigma_1) \cdots B_{w(N-1)}(\sigma_{N-1})$ from Theorem 4.6.

Example 1. For N=3, we have, by Eq.(4.2) or Theorem 4.6,

$$\begin{split} &B_{2}(\sigma_{1})B_{1}(\sigma_{2}) |000\rangle \\ &= |210\rangle + (\sigma_{1} \ominus t_{1}) |120\rangle + \left[(\sigma_{1} \ominus t_{1}) + (\sigma_{2} \ominus t_{2}) + \beta(\sigma_{1} \ominus t_{1})(\sigma_{2} \ominus t_{2}) \right] |201\rangle \\ &+ (\sigma_{1} \ominus t_{1})(\sigma_{1} \ominus t_{2}) |102\rangle + (\sigma_{1} \ominus t_{1})(\sigma_{2} \ominus t_{1}) |021\rangle + (\sigma_{1} \ominus t_{1})(\sigma_{1} \ominus t_{2})(\sigma_{2} \ominus t_{1}) |012\rangle \\ &= \mathcal{G}_{123}^{(\beta)}(\sigma_{1}, \sigma_{2}; \ominus t_{1}, \ominus t_{2}) |210\rangle + \mathcal{G}_{213}^{(\beta)}(\sigma_{1}, \sigma_{2}; \ominus t_{1}, \ominus t_{2}) |120\rangle + \mathcal{G}_{132}^{(\beta)}(\sigma_{1}, \sigma_{2}; \ominus t_{1}, \ominus t_{2}) |201\rangle \\ &+ \mathcal{G}_{312}^{(\beta)}(\sigma_{1}, \sigma_{2}; \ominus t_{1}, \ominus t_{2}) |102\rangle + \mathcal{G}_{231}^{(\beta)}(\sigma_{1}, \sigma_{2}; \ominus t_{1}, \ominus t_{2}) |201\rangle + \mathcal{G}_{321}^{(\beta)}(\sigma_{1}, \sigma_{2}; \ominus t_{1}, \ominus t_{2}) |102\rangle + \mathcal{G}_{321}^{(\beta)}(\sigma_{1}, \sigma_{2}; \ominus t_{1}, \ominus t_{2}) |201\rangle + \mathcal{G}_{321}^{(\beta)}(\sigma_{1}, \sigma_{2}; \ominus t_{1}, \ominus t_{2}) |1012\rangle \,, \end{split}$$

while

$$B_{1}(\sigma_{1})B_{2}(\sigma_{2})|000\rangle$$

$$=(1+\beta(\sigma_{1}\ominus t_{1}))|120\rangle+[(\sigma_{2}\ominus t_{1})(1+\beta(\sigma_{1}\ominus t_{2}))+(\sigma_{1}\ominus t_{2})(1+\beta(\sigma_{1}\ominus t_{1}))]|102\rangle (4.29)$$

$$+(\sigma_{1}\ominus t_{1})(\sigma_{2}\ominus t_{1})(1+\beta(\sigma_{1}\ominus t_{2}))|012\rangle.$$

It is easy to verify that the coefficients in Eq.(4.29) match the conclusion of Lemma 4.7 since we

have

$$\begin{split} \bar{\partial}_{1}^{\beta}\mathcal{G}_{213}^{(\beta)}(\sigma_{1},\sigma_{2};\ominus t_{1},\ominus t_{2}) &= 1 + \beta(\sigma_{1}\ominus t_{1}), \\ \bar{\partial}_{1}^{\beta}\mathcal{G}_{321}^{(\beta)}(\sigma_{1},\sigma_{2};\ominus t_{1},\ominus t_{2}) &= (\sigma_{1}\ominus t_{1})(\sigma_{2}\ominus t_{1})(1+\beta(\sigma_{1}\ominus t_{2})), \\ \bar{\partial}_{1}^{\beta}\mathcal{G}_{312}^{(\beta)}(\sigma_{1},\sigma_{2};\ominus t_{1},\ominus t_{2}) &= (\sigma_{2}\ominus t_{1})(1+\beta(\sigma_{1}\ominus t_{2})) + (\sigma_{1}\ominus t_{2})(1+\beta(\sigma_{1}\ominus t_{1})), \\ \bar{\partial}_{1}^{\beta}\mathcal{G}_{231}^{(\beta)}(\sigma_{1},\sigma_{2};\ominus t_{1},\ominus t_{2}) &= \bar{\partial}_{1}^{\beta}\mathcal{G}_{132}^{(\beta)}(\sigma_{1},\sigma_{2};\ominus t_{1},\ominus t_{2}) &= \bar{\partial}_{1}^{\beta}\mathcal{G}_{132}^{(\beta)}(\sigma_{1},\sigma_{2};\ominus t_{1},\ominus t_{2}) &= 0. \end{split}$$

Now we would like to apply Theorem 4.6 and Lemma 4.7 to the Bethe ansatz state (3.4). First, we show the following

Theorem 4.8. For any $w \in S_n$, we have the following form of the Cauchy identity

$$\mathcal{G}_w^{(eta)}(oldsymbol{x};\ominusoldsymbol{t}) = \sum_{v \in S_n} \mathcal{G}_v^{(eta)}(oldsymbol{x};\ominusoldsymbol{z})ar{\partial}_v^{eta} \mathcal{G}_w^{(eta)}(oldsymbol{z};\ominusoldsymbol{t}).$$

Proof. The generalized Cauchy identity Eq.(A.7) can be rewritten as (notice that $\ominus \ominus z = z$)

$$\mathcal{G}_w^{(\beta)}(\ominus t; \boldsymbol{x}) = \sum_{v \in S_n} \mathcal{G}_v^{(\beta)}(\boldsymbol{x}; \ominus \boldsymbol{z}) \mathcal{G}_{v^{-1}, w}^{(\beta)}(\ominus t; \boldsymbol{z}). \tag{4.30}$$

From Eq.(A.4), (A.5) and (A.6), we get

$${\mathcal G}_{v^{-1},w}^{(eta)}(\ominus oldsymbol{t};oldsymbol{z}) = ar{\partial}_{v,oldsymbol{z}}^{eta}{\mathcal G}_{1,w}^{(eta)}(\ominus oldsymbol{t};oldsymbol{z}) = ar{\partial}_{v}^{eta}{\mathcal G}_{w^{-1}}^{(eta)}(oldsymbol{z};\ominus oldsymbol{t}).$$

Plugging the identity above in (4.30), we arrive at

$${\mathcal G}_{w^{-1}}^{(eta)}({m x};\ominus{m t}) = \sum_{v\in S_n} {\mathcal G}_v^{(eta)}({m x};\ominus{m z}) ar\partial_v^eta {\mathcal G}_{w^{-1}}^{(eta)}({m z};\ominus{m t}).$$

Then the theorem is proved by replacing w^{-1} with w.

Let $L_n(t,\beta)$ be the submodule of $\mathbb{C}(t,\beta)[x_1,\cdots,x_n]$ defined by

$$L_n(t,\beta) = \operatorname{span}_{\mathbb{C}(t,\beta)} \{ x_1^{i_1} x_2^{i_2} \cdots x_n^{i_n} \mid 0 \le i_j \le n - j \}.$$
 (4.31)

Corollary 4.9. For any $F(x; t, \beta) \in L_n(t, \beta)$, we have

$$F(\boldsymbol{x};\boldsymbol{t},\beta) = \sum_{v \in S_n} \mathcal{G}_v^{(\beta)}(\boldsymbol{x};\ominus \boldsymbol{z}) \bar{\partial}_v^{\beta} F(\boldsymbol{z};\boldsymbol{t},\beta). \tag{4.32}$$

Proof. Since $\{\mathcal{G}_w^{(\beta)}(\boldsymbol{x};\ominus \boldsymbol{t})\mid w\in S_n\}$ is a $\mathbb{C}(\boldsymbol{t},\beta)$ -basis of $L_n(\boldsymbol{t},\beta)$ (as are the Schubert and Grothendieck polynomials [17, Proposition 2.7]), Eq.(4.32) follows from Theorem 4.8 and linearity of $\bar{\partial}_v^{\beta}$.

In section 3, we have seen that $\sigma_a^{(N-i)}$ $(\beta=0)$ or $1-\sigma_a^{(N-i)}$ $(\beta=-1)$, $a=1,\cdots,i$, can be identified with the Chern roots of \mathcal{S}_i when $\sigma_a^{(N-i)}$'s satisfy the Bethe ansatz equations. We will show that the Bethe ansatz state (3.4) generates the double β -Grothendieck polynomials when this identification is implemented. For this purpose, we need the following

Lemma 4.10. For $\pi \in S_n$, if $e_l(\sigma_1, \dots, \sigma_{n-1})$ are identified with $e_l(x_1, \dots, x_{n-1})$, $l = 0, 1, \dots, n-1$, then

$$\sum_{w \in S_{n-1}} \mathcal{G}_w^{(\beta)}(x_1, \cdots, x_{n-2}; \ominus \sigma_1, \cdots, \ominus \sigma_{n-2}) \cdot \bar{\partial}_w^{\beta} \mathcal{G}_{\pi}^{(\beta)}(\sigma_1, \cdots, \sigma_{n-1}; \ominus t_1, \cdots, \ominus t_{n-1})$$
$$= \mathcal{G}_{\pi}^{(\beta)}(x_1, \cdots, x_{n-1}; \ominus t_1, \cdots, \ominus t_{n-1}).$$

Proof. Let $L_n(\boldsymbol{t},\beta)$ be the submodule defined by (4.31). Then, since $\{\mathcal{G}_w^{(\beta)}(\boldsymbol{x};\ominus \boldsymbol{t}) \mid w \in S_n\}$ and $\{e_{i_1}(x_1)e_{i_2}(x_1,x_2)\cdots e_{i_{n-1}}(x_1,\cdots,x_{n-1})\mid 0\leq i_j\leq j\}$ are both $\mathbb{C}(\boldsymbol{t},\beta)$ -bases of $L_n(\boldsymbol{t},\beta)$, we have the expansion

$$\mathcal{G}_{\pi}^{(\beta)}(\sigma_1, \cdots, \sigma_{n-1}; \ominus t_1, \cdots, \ominus t_{n-1})$$

$$= \sum_{i_1, i_2, \cdots, i_{n-1}} C_{i_1 \cdots i_{n-1}}^{\pi}(t, \beta) e_{i_1}(\sigma_1) e_{i_2}(\sigma_1, \sigma_2) \cdots e_{i_{n-1}}(\sigma_1, \cdots, \sigma_{n-1}),$$

where $C_{i_1\cdots i_{n-1}}(\boldsymbol{t},\beta)\in\mathbb{C}(\boldsymbol{t},\beta)$.

Therefore

$$\begin{split} &\sum_{w \in S_{n-1}} \mathcal{G}_{w}^{(\beta)}(x_{1}, \cdots, x_{n-2}; \ominus \sigma_{1}, \cdots, \ominus \sigma_{n-2}) \cdot \bar{\partial}_{w}^{\beta} \mathcal{G}_{\pi}^{(\beta)}(\sigma_{1}, \cdots, \sigma_{n-1}; \ominus t_{1}, \cdots, \ominus t_{n-1}) \\ &= \sum_{w \in S_{n-1}} \mathcal{G}_{w}^{(\beta)}(x_{1}, \cdots, x_{n-2}; \ominus \sigma_{1}, \cdots, \ominus \sigma_{n-2}) \\ & \cdot \bar{\partial}_{w}^{\beta} \left[\sum_{i_{1}, i_{2}, \cdots, i_{n-1}} C_{i_{1} \cdots i_{n-1}}^{\pi}(t, \beta) e_{i_{1}}(\sigma_{1}) e_{i_{2}}(\sigma_{1}, \sigma_{2}) \cdots e_{i_{n-1}}(\sigma_{1}, \cdots, \sigma_{n-1}) \right] \\ &= \sum_{i_{1}, i_{2}, \cdots, i_{n-1}} C_{i_{1} \cdots i_{n-1}}^{\pi}(t, \beta) e_{i_{n-1}}(\sigma_{1}, \cdots, \sigma_{n-1}) \\ & \cdot \left[\sum_{w \in S_{n-1}} \mathcal{G}_{w}^{(\beta)}(x_{1}, \cdots, x_{n-2}; \ominus \sigma_{1}, \cdots, \ominus \sigma_{n-2}) \cdot \bar{\partial}_{w}^{\beta} \left(e_{i_{1}}(\sigma_{1}) e_{i_{2}}(\sigma_{1}, \sigma_{2}) \cdots e_{i_{n-2}}(\sigma_{1}, \cdots, \sigma_{n-2}) \right) \right] \\ &= \sum_{i_{1}, i_{2}, \cdots, i_{n-1}} C_{i_{1} \cdots i_{n-1}}^{\pi}(t, \beta) e_{i_{1}}(x_{1}) e_{i_{2}}(x_{1}, x_{2}) \cdots e_{i_{n-2}}(x_{1}, \cdots, x_{n-2}) e_{i_{n-1}}(\sigma_{1}, \cdots, \sigma_{n-1}) \\ &= \sum_{i_{1}, i_{2}, \cdots, i_{n-1}} C_{i_{1} \cdots i_{n-1}}^{\pi}(t, \beta) e_{i_{1}}(x_{1}) e_{i_{2}}(x_{1}, x_{2}) \cdots e_{i_{n-2}}(x_{1}, \cdots, x_{n-2}) e_{i_{n-1}}(x_{1}, \cdots, x_{n-1}) \\ &= \mathcal{G}_{x}^{(\beta)}(x_{1}, \cdots, x_{n-1}; \ominus t_{1}, \cdots, \ominus t_{n-1}), \end{split}$$

where the second equality is due to $\bar{\partial}_i^{\beta}(f \cdot g) = f(\bar{\partial}_i^{\beta}g)$ if f is symmetric in $(\sigma_1, \dots, \sigma_{n-1})$, the third equality is due to Corollary 4.9, and $e_{i_{n-1}}(\sigma_1, \dots, \sigma_{n-1})$ is identified with $e_{i_{n-1}}(x_1, \dots, x_{n-1})$ in the fourth equality.

Example 2. Notice that $\mathcal{G}_{12}^{(\beta)}(x_1; \ominus \sigma_1) = 1$, $\mathcal{G}_{21}^{(\beta)}(x_1; \ominus \sigma_1) = x_1 \ominus \sigma_1$. In the case n = 3, following Example 1, one can check

$$\mathcal{G}_{12}^{(\beta)}(x_1;\ominus\sigma_1)\mathcal{G}_{321}^{(\beta)}(\sigma_1,\sigma_2;\ominus t_1,\ominus t_2) + \mathcal{G}_{21}^{(\beta)}(x_1;\ominus\sigma_1)\bar{\partial}_1^{\beta}\mathcal{G}_{321}^{(\beta)}(\sigma_1,\sigma_2;\ominus t_1,\ominus t_2)$$

$$= (\sigma_1\ominus t_1)(\sigma_2\ominus t_1)(x_1\ominus t_2) = (x_1\ominus t_1)(x_2\ominus t_1)(x_1\ominus t_2)$$

$$= \mathcal{G}_{321}^{(\beta)}(x_1,x_2;\ominus t_1,\ominus t_2),$$

$$\mathcal{G}_{12}^{(\beta)}(x_{1};\ominus\sigma_{1})\mathcal{G}_{231}^{(\beta)}(\sigma_{1},\sigma_{2};\ominus t_{1},\ominus t_{2}) = (\sigma_{1}\ominus t_{1})(\sigma_{2}\ominus t_{1}) = (x_{1}\ominus t_{1})(x_{2}\ominus t_{1})$$

$$=\mathcal{G}_{231}^{(\beta)}(x_{1},x_{2};\ominus t_{1},\ominus t_{2}),$$

$$\mathcal{G}_{12}^{(\beta)}(x_{1};\ominus\sigma_{1})\mathcal{G}_{312}^{(\beta)}(\sigma_{1},\sigma_{2};\ominus t_{1},\ominus t_{2}) + \mathcal{G}_{21}^{(\beta)}(x_{1};\ominus\sigma_{1})\bar{\partial}_{1}^{\beta}\mathcal{G}_{312}^{(\beta)}(\sigma_{1},\sigma_{2};\ominus t_{1},\ominus t_{2})$$

$$=\frac{x_{1}(\sigma_{1}+\sigma_{2}-t_{1}-t_{2})+(t_{1}t_{2}-\sigma_{1}\sigma_{2})}{(1+\beta t_{1})(1+\beta t_{2})} = (x_{1}\ominus t_{1})(x_{2}\ominus t_{2})$$

$$=\mathcal{G}_{312}^{(\beta)}(x_{1},x_{2};\ominus t_{1},\ominus t_{2}),$$

$$\mathcal{G}_{12}^{(\beta)}(x_{1};\ominus\sigma_{1})\mathcal{G}_{132}^{(\beta)}(\sigma_{1},\sigma_{2};\ominus t_{1},\ominus t_{2}) = \frac{x_{1}+x_{2}-t_{1}-t_{2}+\beta(x_{1}x_{2}-t_{1}t_{2})}{(1+\beta t_{1})(1+\beta t_{2})}$$

$$=\mathcal{G}_{132}^{(\beta)}(x_{1},x_{2};\ominus t_{1},\ominus t_{2}),$$

$$\mathcal{G}_{12}^{(\beta)}(x_{1};\ominus\sigma_{1})\mathcal{G}_{213}^{(\beta)}(\sigma_{1},\sigma_{2};\ominus t_{1},\ominus t_{2}) + \mathcal{G}_{21}^{(\beta)}(x_{1};\ominus\sigma_{1})\bar{\partial}_{1}^{\beta}\mathcal{G}_{213}^{(\beta)}(\sigma_{1},\sigma_{2};\ominus t_{1},\ominus t_{2})$$

$$=(\sigma_{1}\ominus t_{1})+(x_{1}\ominus\sigma_{1})(1+\beta(\sigma_{1}\ominus t_{1})) = x_{1}\ominus t_{1}$$

$$=\mathcal{G}_{213}^{(\beta)}(x_{1},x_{2};\ominus t_{1},\ominus t_{2}),$$

where we have made the following identification

$$e_1(\sigma_1, \sigma_2) = e_1(x_1, x_2), \quad e_2(\sigma_1, \sigma_2) = e_2(x_1, x_2).$$

Finally, we show that an on-shell Bethe state satisfies the following

Theorem 4.11. When all the instances of $e_l(\sigma_1^{(i)}, \dots, \sigma_{N-i}^{(i)})$, $l = 1, \dots, N-i$, are identified with $e_l(x_1, \dots, x_{N-i})$ for $i = 1, \dots, N-1$, the Bethe ansatz state (3.4) (in the case $k_i = N-i$) has the expansion

$$|\psi^{(0)}\rangle = \sum_{w \in S_N} \mathcal{G}_w^{(\beta)}(x_1, \cdots, x_{N-1}; \ominus t_1, \cdots, \ominus t_{N-1}) |\omega_N w^{-1}\rangle.$$
 (4.33)

Proof. We prove by induction on the number of sites. For a two-site system, it is easy to check that, when $\sigma_1^{(1)} = x_1$,

$$\langle 01 | B_1(\sigma_1^{(1)}) | 00 \rangle = \sigma_1^{(1)} \ominus t_1 = x_1 \ominus t_1 = \mathcal{G}_{21}^{(\beta)}(x_1; \ominus t_1),$$

$$\langle 10 | B_1(\sigma_1^{(1)}) | 00 \rangle = 1 = \mathcal{G}_{12}^{(\beta)}(x_1; \ominus t_1),$$

so Eq.(4.33) holds for N=2. Now assume Eq.(4.33) holds for $N=2,3,\cdots,m$ with $m\geq 2$, then we have, from Eq.(3.14) and (3.15), for $w_l\in S_l$ (notice that $|\psi^{(N-s)}\rangle$ is the Bethe state of a $\mathrm{GL}(s)$ system with s sites),

$$\langle \omega_{2}w_{2}^{-1}|\psi^{(N-2)}\rangle = \mathcal{G}_{w_{2}}^{(\beta)}(x_{1};\ominus\sigma_{1}^{(N-2)}),$$

$$\langle \omega_{3}w_{3}^{-1}|\psi^{(N-3)}\rangle = \mathcal{G}_{w_{3}}^{(\beta)}(x_{1},x_{2};\ominus\sigma_{1}^{(N-3)},\ominus\sigma_{2}^{(N-3)}),$$

$$\vdots$$

$$\langle \omega_{m}w_{m}^{-1}|\psi^{(N-m)}\rangle = \mathcal{G}_{w_{m}}^{(\beta)}(x_{1},\cdots,x_{m-1};\ominus\sigma_{1}^{(N-m)},\cdots,\ominus\sigma_{m-1}^{(N-m)})$$

Eq.(3.14) and (3.15) yields

$$|\psi^{(N-m-1)}\rangle = \sum_{v_m \in S_m} \langle v_m | \psi^{(N-m)} \rangle B_{v_m(1)}^{(N-m-1)}(\sigma_1^{(N-m)}) \cdots B_{v_m(m)}^{(N-m-1)}(\sigma_m^{(N-m)}) | \Omega^{(N-m-1)} \rangle.$$
(4.34)

It is easy to verify that

$$\begin{split} &\langle \omega_{m+1}w_{m+1}^{-1}|\,B_{v_m(1)}^{(N-m-1)}(\sigma_1^{(N-m)})\cdots B_{v_m(m)}^{(N-m-1)}(\sigma_m^{(N-m)})\,|\Omega^{(N-m-1)}\rangle\\ =&\bar{\partial}_{v_m^{-1}\omega_m}^{\beta}\,\langle \omega_{m+1}w_{m+1}^{-1}|\,B_m^{(N-m-1)}(\sigma_1^{(N-m)})\cdots B_1^{(N-m-1)}(\sigma_m^{(N-m)})\,|\Omega^{(N-m-1)}\rangle\\ =&\bar{\partial}_{v_m^{-1}\omega_m}^{\beta}\,\mathcal{G}_{w_{m+1}}^{(\beta)}(\sigma_1^{(N-m)},\cdots,\sigma_m^{(N-m)};\ominus\sigma_1^{(N-m-1)},\cdots,\ominus\sigma_m^{(N-m-1)}), \end{split}$$

where the first equality is due to Lemma 4.7 and the second equality is due to Theorem 4.6. As $\langle v_m|\psi^{(N-m)}\rangle=\mathcal{G}_{v_m^{-1}\omega_m}^{(\beta)}(x_1,\cdots,x_{m-1};\ominus\sigma_1^{(N-m)},\cdots,\ominus\sigma_{m-1}^{(N-m)})$ from the induction hypothesis, Lemma 4.10 and Eq.(4.34) then yield

$$\begin{split} &\langle \omega_{m+1}w_{m+1}^{-1}|\psi^{(N-m-1)}\rangle\\ &=\sum_{v_m\in S_m} \langle v_m|\psi^{(N-m)}\rangle \, \langle \omega_{m+1}w_{m+1}^{-1}|\, B_{v_m(1)}^{(N-m-1)}(\sigma_1^{(N-m)})\cdots B_{v_m(m)}^{(N-m-1)}(\sigma_m^{(N-m)})\, |\Omega^{(N-m-1)}\rangle\\ &=\sum_{v_m\in S_m} \mathcal{G}_{v_m^{-1}\omega_m}^{(\beta)}(x_1,\cdots,x_{m-1};\ominus\sigma_1^{(N-m)},\cdots,\ominus\sigma_{m-1}^{(N-m)})\\ &\qquad \cdot \bar{\partial}_{v_m^{-1}\omega_m}^{\beta} \mathcal{G}_{w_{m+1}}^{(\beta)}(\sigma_1^{(N-m)},\cdots,\sigma_m^{(N-m)};\ominus\sigma_1^{(N-m-1)},\cdots,\ominus\sigma_m^{(N-m-1)})\\ &=\mathcal{G}_{w_{m+1}}^{(\beta)}(x_1,\cdots,x_m;\ominus\sigma_1^{(N-m-1)},\cdots,\ominus\sigma_m^{(N-m-1)}). \end{split}$$

Notice that $\sigma_i^{(0)} = t_i$, The proof is thus complete by induction.

Remark 4. As discussed in Section 3, when the Bethe ansatz equations Eq.(3.5)-(3.7) are taken into account, we should interpret $\sigma_a^{(N-i)}$ ($\beta=0$) or $1-\sigma_a^{(N-i)}$ ($\beta=-1$), $a=1,\dots,i$, as the Chern roots of S_i . Therefore, x_i should be interpreted as the first Chern class of S_i/S_{i-1} when $\beta=0$ in the quantum cohomology ring, and $1-x_i$ should be interpreted as the K-theory class of S_i/S_{i-1} when $\beta=-1$ in the K-theory ring. In this sense, we regard the Bethe ansatz state (4.33) as on-shell.

Remark 5. More generally, for $\beta \neq 0$, we expect $1 + \beta x_i$ to represent S_i/S_{i-1} in the connective K-theory ring of the flag variety [18].

Remark 6. In the case of full flag variety, $i_a \neq i_b$ for $a \neq b$ in Eq.(3.14). In the case of partial flag variety $Fl(k_{n-1}, k_{n-2}, \dots, k_1; N)$, it is possible that $i_a = i_b$ for some $a \neq b$. Because of $[B_i(x), B_i(y)] = 0$ (Eq.(4.23)), we expect the expansion coefficients of the Bethe ansatz state to be given by the β -Grothendieck polynomials indexed by S_N/W , where W is generated by the simple transpositions s_i for $i \notin \{k_{n-1}, k_{n-2}, \dots, k_1\}$, i.e. β -Grothendieck polynomials symmetric in $(x_{k_s+1}, \dots, x_{k_{s-1}})$ for all $s = 1, \dots, n$.

Acknowledgement

The author would like to thank Xiang-Mao Ding, Leonardo Mihalcea and Eric Sharpe for useful comments. This work is partially supported by the National Natural Science Foundation of China (Grant No. 12475005), the Natural Science Foundation of Shanghai (Grant No. 24ZR1468600), and the Fundamental Research Funds for the Central Universities.

A Quantum cohomology/K-theory of flag varieties and the double β -Grothendieck polynomials

In this appendix, we give a brief review of the quantum cohomology/K-theory ring of flag varieties, and the double β -Grothendieck polynomials. The reader may refer to e.g. [15, 17, 19–21] and references therein for more information.

Given $E = \mathbb{C}^N$ and $a = (a_1, a_2, \dots, a_l)$ such that $0 < a_1 < a_2 < \dots < a_l < N$, the flag variety $\mathrm{Fl}(a_1, a_2, \dots, a_l; N)$ is defined to be the variety of partial flags

$$V_1 \subset V_2 \subset \cdots \subset V_l \subset E$$

such that $\dim(V_i) = a_i$ for all i.

There is a sequence of vector bundles on $Fl(a_1, a_2, \dots, a_l; N)$ (the tautological sequence):

$$S_1 \subset S_2 \subset \cdots \subset S_l \subset E$$

with rank(S_i) = a_i . Let S_N be the group of permutations of N elements, and let l(w) be the length⁸ of a permutation $w \in S_N$. The Schubert varieties are subvarieties of $Fl(a_1, a_2, \dots, a_l; N)$ indexed by the set $S_N(a) = S_N/W_a$, where $W_a \subset S_N$ is the subgroup generated by the simple transpositions $s_i = (i, i+1)$ for $i \notin \{a_1, \dots, a_l\}$. The codimension of the Schubert variety indexed by the representative $w \in S_N(a)$ with shortest length is equal to l(w).

Let $\Omega_w^{(a)}$ denote the cohomology class Poincaré dual to the Schubert variety indexed by w, then the Schubert classes $\Omega_w^{(a)}$ form a basis for the cohomology ring $H^*(\mathrm{Fl}(a_1, a_2, \dots, a_l; N), \mathbb{Z})$.

Let $\mathrm{Fl}(N)=\mathrm{Fl}(1,2,\cdots,N-1;N)$ denote the full flag variety. The cohomology of $\mathrm{Fl}(N)$ has the following presentation

$$H^*(\mathrm{Fl}(N)) = \mathbb{Z}[x_1, \cdots, x_N]/(e_1^N, \cdots, e_N^N),$$

where $e_i^N = e_i(x_1, \dots, x_N)$ is the *i*th elementary symmetric polynomial in N variables. In this presentation, x_i is identified with the Chern class $c_1(S_i/S_{i-1})$ and the Schubert class Ω_w is represented by the Schubert polynomial $X_w(x_1, \dots, x_N)$, which means that the multiplication rules of the Schubert classes are the same as those of the Schubert polynomials modulo e_i^N , $i = 1, \dots, N$.

The quantum elementary polynomials E_i^N are defined via the Givental-Kim determinant. Let

$$\Gamma_N = \begin{pmatrix} x_1 & q_1 & 0 & \cdots & 0 \\ -1 & x_2 & q_2 & \cdots & 0 \\ 0 & -1 & x_3 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & x_N \end{pmatrix},$$

then E_i^N is defined as the coefficient of λ^i in the characteristic polynomial $\det(1 + \lambda \Gamma_N)$. The quantum cohomology ring of the full flag variety can be presented as

$$QH^*(\operatorname{Fl}(N)) = \mathbb{Z}[x_1, \cdots, x_N]/(E_1^N, \cdots, E_N^N).$$

⁸The length of a permutation $w \in S_N$ is the number of pairs (i,j) such that i < j and w(i) > w(j).

Ring relations of partial flag varieties can be presented similarly, see [19].

The K-theory ring of the flag variety $K(\operatorname{Fl}(a_1, a_2, \dots, a_l; N))$ is the Grothendieck group of the coherent sheaves of $\operatorname{Fl}(a_1, a_2, \dots, a_l; N)$. A basis of $K(\operatorname{Fl}(a_1, a_2, \dots, a_l; N))$ can be chosen as the structure sheaves of the Schubert varieties $\{\mathcal{O}_w | w \in S_n(a)\}$, which can be represented by the Grothendieck polynomials $G_w(x_1, \dots, x_N)$. Ring relations can be found in e.g. [20]. One way to present the ring relations of the (quantum) K-theory ring of $\operatorname{Fl}(a_1, a_2, \dots, a_l; N)$ is to encapsulate them in the (quantum) Whitney relations [11].

For the $T=(\mathbb{C}^*)^N$ action on $E=\mathbb{C}^N$ with characters (equivariant parameters) (t_1,\cdots,t_N) , we can define the equivariant quantum cohomology ring $QH_T^*(\mathrm{Fl}(a;E))$ and equivariant quantum K-theory ring $QK_T(\mathrm{Fl}(a;E))$. The representatives of the Schubert classes are given by the double Schubert polynomials $X_w(x_1,\cdots,x_N;t_1,\cdots,t_N)$ and the double Grothendieck polynomials $G_w(x_1,\cdots,x_N;t_1,\cdots,t_N)$ respectively.

The double Schubert and double Grothendieck polynomials are special cases of a one-parameter family of polynomials called the double β -Grothedieck polynomials $\mathcal{G}_w^{(\beta)}, w \in S_N$ [15]. Let \mathbb{K} be a field of zero characteristic and β be a formal variable, $\boldsymbol{x} = (x_1, \dots, x_N), \boldsymbol{y} = (y_1, \dots, y_N)$, define the β -divided-difference operator ∂_i^{β} acting on $\mathbb{K}(\beta)[x_1, \dots, x_N, y_1, \dots, y_N]$ by

$$\partial_{i}^{\beta} f(x_{1}, \dots, x_{N}; \boldsymbol{y}) := \frac{(1 + \beta x_{i+1}) f(x_{1}, \dots, x_{N}; \boldsymbol{y}) - (1 + \beta x_{i}) f(x_{1}, \dots, x_{i+1}, x_{i}, \dots, x_{N}; \boldsymbol{y})}{x_{i} - x_{i+1}}.$$
(A.1)

If $w = \omega_N$ is the permutation in S_N with maximal length, then we define

$$\mathcal{G}_{\omega_N}^{(\beta)}(\boldsymbol{x};\boldsymbol{y}) = \prod_{i+j \le N} (x_i + y_j + \beta x_i y_j), \tag{A.2}$$

and for other $w \in S_N$, $\mathcal{G}_w^{(\beta)}$ can be defined recursively by

$$\mathcal{G}_{ws_i}^{(\beta)}(\boldsymbol{x};\boldsymbol{y}) = \partial_i^{\beta} \mathcal{G}_w^{(\beta)}(\boldsymbol{x};\boldsymbol{y})$$
(A.3)

if $l(ws_i) = l(w) - 1$. The double β -Grothendieck polynomials reduce to the double Schubert polynomials when $\beta = 0$, and reduce to the double Grothendieck polynomials when $\beta = -1$. The β -Grothendieck polynomials satisfy (see Lemma 5.8 of [18])

$$\mathcal{G}_w^{(\beta)}(\boldsymbol{x};\boldsymbol{y}) = \mathcal{G}_{w^{-1}}^{(\beta)}(\boldsymbol{y};\boldsymbol{x}). \tag{A.4}$$

Moreover, the β -Grothendieck polynomials can be generated by the generalized Hecke algebra with generators u_1, \dots, u_{n-1} satisfying [15]

$$u_i u_j = u_j u_i, |i - j| > 1;$$
 $u_i u_{i+1} u_i = u_{i+1} u_i u_{i+1};$ $u_i^2 = \beta u_i.$

The dual double β -Grothendieck polynomials $\mathcal{H}_w^{(\beta)}(\boldsymbol{x};\boldsymbol{y})$ can be defined recursively by setting $\mathcal{H}_{w_N}^{(\beta)}(\boldsymbol{x};\boldsymbol{y}) = \mathcal{G}_{w_N}^{(\beta)}(\boldsymbol{x};\boldsymbol{y})$ and

$$\mathcal{H}_{ws}^{(eta)}(oldsymbol{x};oldsymbol{y}) = ar{\partial}_i^eta \mathcal{H}_w^{(eta)}(oldsymbol{x};oldsymbol{y})$$

if $l(ws_i) = l(w) - 1$ [16], where $\bar{\partial}_i^{\beta} = \partial_i^{\beta} + \beta$.

More generally, for $v, w \in S_N$, one can define the biaxial double β -Grothendieck polynomials

 $\mathcal{G}_{v,w}^{(\beta)}(\boldsymbol{x};\boldsymbol{y})$ as follows (see Definition 2.6 of [8]):

$$\mathcal{G}_{1,w}^{(\beta)}(\boldsymbol{x};\boldsymbol{y}) = \mathcal{G}_{w}^{(\beta)}(\boldsymbol{x};\boldsymbol{y}). \tag{A.5}$$

If $l(vs_i) = l(v) + 1$, set

$$\mathcal{G}_{vs_{i},w}^{(\beta)}(\boldsymbol{x};\boldsymbol{y}) = \bar{\partial}_{i,\boldsymbol{y}}^{\beta} \mathcal{G}_{v,w}^{(\beta)}(\boldsymbol{x};\boldsymbol{y}), \tag{A.6}$$

where $\bar{\partial}_{i,\boldsymbol{y}}^{\beta}$ is the $\bar{\partial}_{i}^{\beta}$ operator acting on the set of variables $\boldsymbol{y}=(y_{1},y_{2},\cdots,y_{N})$. We have the following generalized Cauchy identity:

Theorem A.1 (Theorem 7.1 of [8]). For any $w \in S_N$,

$$\mathcal{G}_w^{(\beta)}(\boldsymbol{x};\boldsymbol{y}) = \sum_{v \in S_N} \mathcal{G}_v^{(\beta)}(\boldsymbol{y};\boldsymbol{z}) \mathcal{G}_{v^{-1},w}^{(\beta)}(\boldsymbol{x};\ominus \boldsymbol{z}), \tag{A.7}$$

where $\ominus z_i = -z_i/(1+\beta z_i)$.

References

- [1] V. Gorbounov and C. Korff, "Quantum integrability and generalised quantum Schubert calculus," Advances in Mathematics **313**, 282-356 (2014), [arXiv:1408.4718 [math.RT]].
- [2] K. Ueda and Y. Yoshida, "3d N=2 Chern-Simons-matter theory, Bethe ansatz, and quantum K-theory of Grassmannians," JHEP **08** (2020) 157, [arXiv:1912.03792 [hep-th]].
- [3] A. Givental and B. Kim, "Quantum cohomology of flag manifolds and Toda lattices," Commun. Math. Phys. 168, 609-642 (1995), [arXiv:hep-th/9312096].
- [4] A. Givental and Y. P. Lee, "Quantum K-theory on flag manifolds, finite difference Toda lattices and quantum groups," Invent. Math. 151, 193-219 (2003), [arXiv:math/0108105].
- [5] P. Koroteev, P. P. Pushkar, A. V. Smirnov and A. M. Zeitlin, "Quantum K-theory of quiver varieties and many-body systems," Selecta Math. New Ser. 27 (2021) 87, [arXiv:1705.10419 [math.AG]].
- [6] V. Buciumas, T. Scrimshaw and K. Weber, "Colored five-vertex models and Lascoux polynomials and atoms," Journal of the London Mathematical Society, 102 (2019), [arXiv:1908.07364 [math.CO]].
- [7] V. Buciumas and T. Scrimshaw, "Double Grothendieck Polynomials and Colored Lattice Models," International Mathematics Research Notices (2020), [arXiv:2007.04533 [math.CO]].
- [8] B. Brubaker, C. Frechette, A. Hardt, E. Tibor, K. Weber, "Frozen pipes: lattice models for Grothendieck polynomials," Algebraic Combinatorics, 6 (2023) no. 3, pp. 789-833, [arXiv:2007.04310 [math.CO]].
- [9] P. P. Kulish and N. Y. Reshetikhin, "Diagonalisation of GL(N) invariant transfer matrices and quantum N-wave system (Lee model)," J. Phys. A **16** (1983) L591-L596.
- [10] J. Guo, "Quantum Sheaf Cohomology and Duality of Flag Manifolds," Commun. Math. Phys. 374, no.2, 661-688 (2019), [arXiv:1808.00716 [hep-th]].

- [11] W. Gu, L. Mihalcea, E. Sharpe, W. Xu, H. Zhang and H. Zou, "Quantum K theory rings of partial flag manifolds," J. Geom. Phys. 198, 105127 (2024), [arXiv:2306.11094 [hep-th]].
- [12] R. Donagi and E. Sharpe, "GLSM's for partial flag manifolds," J. Geom. Phys. 58 (2008) 1662-1692, [arXiv:0704.1761 [hep-th]].
- [13] N. Nekrasov and S. Shatashvili, "Supersymmetric Vacua and Bethe Ansatz," Nuclear Physics B Proceedings Supplements 192 (2009): 91-112.
- [14] N. Nekrasov and S. Shatashvili, "Quantum Integrability and Supersymmetric Vacua," Progress of Theoretical Physics Supplement 177 (2009): 105-119.
- [15] S. Fomin and A. N. Kirillov, "Yang-Baxter equation, symmetric functions and Grothendieck polynomials," [arXiv:hep-th/9306005].
- [16] A. Lascoux and M. P. Schützenberger, "Symmetry and flag manifolds," in Invariant Theory (Montecatini, 1982), vol. 996 of Lecture Notes in Math., Springer, Berlin, 1983, pp. 118-144.
- [17] C. Lenart and M. Toshiaki, "Quantum Grothendieck Polynomials," [arXiv:math/0608232].
- [18] T. Hudson, "A Thom-Porteous formula for connective K -theory using algebraic cobordism," Journal of K-theory 14 (2013): 343-369, [arXiv:1310.0895 [math.AG]].
- [19] A. S. Buch, "Quantum Cohomology of Partial Flag Manifolds," [arXiv:math/0303245 [math.AG]].
- [20] T. Maeno, S. Naito and D. Sagaki, "A presentation of the torus-equivariant quantum K-theory ring of flag manifolds of type A, Part I: the defining ideal," [arXiv:2302.09485[math.QA]].
- [21] T. Maeno, S. Naito and D. Sagaki, "A presentation of the torus-equivariant quantum K-theory ring of flag manifolds of type A, Part II: quantum double Grothendieck polynomials," [arXiv:2305.17685 [math.QA]].