Distribution learning via neural differential equations: minimal energy regularization and approximation theory

Youssef Marzouk^{*} Zhi Ren^{*}

Jakob Zech[†]

February 7, 2025

Abstract

Neural ordinary differential equations (ODEs) provide expressive representations of invertible transport maps that can be used to approximate complex probability distributions, e.g., for generative modeling, density estimation, and Bayesian inference. We show that for a large class of transport maps T, there exists a time-dependent ODE velocity field realizing a straight-line interpolation (1 - t)x + tT(x), $t \in [0, 1]$, of the displacement induced by the map. Moreover, we show that such velocity fields are minimizers of a training objective containing a specific minimum-energy regularization. We then derive explicit upper bounds for the C^k norm of the velocity field that are polynomial in the C^k norm of the corresponding transport map T; in the case of triangular (Knothe–Rosenblatt) maps, we also show that these bounds are polynomial in the C^k norms of the associated source and target densities. Combining these results with stability arguments for distribution approximation via ODEs, we show that Wasserstein or Kullback–Leibler approximation of the target distribution to any desired accuracy $\epsilon > 0$ can be achieved by a deep neural network representation of the velocity field whose size is bounded in terms of ϵ , the dimension, and the smoothness of the source and target densities. The same neural network ansatz yields guarantees on the value of the regularized training objective.

Contents

1	Intro	oduction	2
	1.1	Contributions	3
	1.2	Related work	4
2	Prel	iminaries	7
	2.1	Notation and definitions	7
	2.2	Problem setup	8
3	Exis	tence and structure of minimizers	9
	3.1	Existence	10
	3.2	Properties of $\Omega_{[0,1]}$	11
	3.3	Regularized solutions	13

*Massachusetts Institute of Technology, Cambridge, MA 02139, USA (ymarz@mit.edu, zren@mit.edu)

[†]Heidelberg University, 69120 Heidelberg, Germany (jakob.zech@uni-heidelberg.de)

4	Regularity of the velocity field <i>f</i>	14
	4.1 General transports	15
	4.2 Triangular transports	18
5	Stability in the velocity field	25
	5.1 Wasserstein distance	25
	5.2 KL-divergence	26
6	Neural network approximation	27
	6.1 Wasserstein distance	28
	6.2 KL-divergence	30
7	Discussion and future work	31
A	Comments on training	32
	A.1 Training algorithm	32
	A.2 Exact computation of the Jacobian	33
B	Knothe–Rosenblatt construction of triangular transport maps	34
C Auxiliary results		

1 Introduction

Sampling from an arbitrary probability distribution is a central problem in computational statistics and machine learning. Transportation of measure (Villani, 2008) offers a useful approach to this problem: the idea is to construct a measurable map that *pushes forward* a relatively simple source distribution (usually chosen to be uniform or standard Gaussian) to the target probability distribution. One can then simulate from the target distribution by drawing samples from the source distribution and evaluating the transport map. This construction is useful for both generative modeling (Grathwohl et al., 2019; Kingma and Dhariwal, 2018) and variational inference (Moselhy and Marzouk, 2012; Rezende and Mohamed, 2015). When the map is invertible, one can also estimate the *density* of the target measure by evaluating the density of the pushforward of the source distribution under the transport map.

Many parameterizations of such transports, ranging from monotone polynomial-based approximations (Marzouk et al., 2016; Zech and Marzouk, 2022a,b; Baptista et al., 2023) to input-convex neural networks (Huang et al., 2021; Wang et al., 2023) have been proposed. In the machine learning literature, normalizing flows (Tabak and Vanden-Eijnden, 2010; Kobyzev et al., 2020) represent transport maps by composing a sequence of relatively simple invertible functions. It is typically required that the Jacobian determinant of the resulting map be easily computable, as it appears in expressions for the pushforward or pullback densities via the change-of-variables formula. Common constructions for normalizing flows including planar and radial flows (Rezende and Mohamed, 2015), affine coupling flows (Dinh et al., 2015), autoregressive flows (Kingma et al., 2016), and neural autoregressive flows (Huang et al., 2018).

More recent work in deep learning has elucidated connections between deep neural networks and differential equations (Sonoda and Murata, 2019; Lu et al., 2018; Ruthotto and Haber, 2020). In particular, neural ordinary differential equations (neural ODEs) (Chen et al., 2018) are ODEs whose velocity fields are represented by neural networks. A neural ODE can be understood intuitively as the "continuous-time limit" of a normalizing flow, insofar as the flow map of the ODE is given by the composition of infinitely many incremental transformations. Neural ODEs can be used to represent distributions (Grathwohl et al., 2019)— again for the purposes of generative modeling, inference, or density estimation—as follows. Let π denote the target distribution from which we wish to sample and let ρ denote the source distribution. Solving the initial value problem

$$\begin{cases} \frac{d}{dt}X(x,t) &= f\left(X(x,t),t\right)\\ X(x,0) &= x \end{cases}$$
(1.1)

up to time t = 1, for some initial condition $x \in \Omega_0$, yields a flow map $x \mapsto X^f(x, 1)$. The goal is to learn a Lipschitz continuous velocity field f, parameterized as a neural network, such that $x \sim \rho$ implies $X^f(x, 1) \sim \pi$; in other words, the flow map pushes forward the source distribution to the target distribution.

Such dynamical representations of transport enjoy several desirable properties. Invertibility of $x \mapsto X^f(x, 1)$ is guaranteed for any Lipschitz f satisfying suitable boundary conditions, as one can solve (1.1) backward in time. Therefore, and in contrast to other methods that directly parameterize the displacement (such as invertible neural networks (Behrmann et al., 2019), normalizing flows (Kobyzev et al., 2020), or other transport maps (Baptista et al., 2023)), no further restrictions need to be imposed on the vector field f that is to be learned. The density η of the ODE state at time t, i.e., the density of $X(\cdot, t)_{\sharp}\rho$, can also easily be computed as it obeys the dynamics

$$\frac{d\log\eta(x,t)}{dt} = -\mathrm{tr}\left(\nabla_X f(X(x,t),t)\right),\,$$

which is known as the *instantaneous change of variables formula*; see Chen et al. (2018).

To understand the usefulness of neural ODEs in learning distributions, there are at least three natural questions to ask. First, it is known that when the source and target measures are well-behaved (for example, both absolutely continuous with respect to the Lebesgue measure), there are in general infinitely many transport maps that push forward one measure onto the other. Moreover, even if we require the time-one flow map $x \mapsto X^f(x, 1)$ to be a particular transport map T, there are in general still infinitely many velocity fields f that realize T. It has thus been observed (Onken et al., 2021; Finlay et al., 2020) that without any form of regularization, learned ODE trajectories connecting x to T(x) may be very irregular. It is therefore natural to ask how we can regularize the training objective to improve the training process. Second, given a regularized training objective, we would like to characterize the structure of its minimizers, and in particular to quantify how well a neural network of a given size (e.g., width, depth, sparsity) can approximate the velocity field corresponding to an optimal solution. Third, we would like to know how to bound the distance between the target measure and the pushforward of the source measure under the ODE-induced flow map, when a neural network is used to approximate the velocity field f.

Our work is the first attempt to address these questions in a unified way. One of our key goals is to obtain explicit rates for distribution approximation using regularized neural ODEs, linking properties of the source and target measures to bounds on the size of a deep neural network representation of the velocity field that achieves a given distributional approximation error. Furthermore, we aim to connect this distribution approximation error to the value of the regularized training objective.

1.1 Contributions

We summarize the main contributions of this paper as follows:

• Realizing straight-line trajectories. For a large class of transport maps T, we show that there exists a corresponding time-dependent velocity field f(x,t) achieving the interpolation (1 - t)x + tT(x)

and hence producing straight-line trajectories. In a Lagrangian frame, these trajectories have constant velocity and hence zero acceleration.

- *Regularity of the space-time domain.* We show that, under certain conditions, the space-time domain covered by such ODE trajectories is a Lipschitz domain, and thus admits suitable extensions that enable the application of existing neural network approximation results.
- *Minimum energy regularization.* We propose a new regularization scheme for ODE velocity fields, based on penalizing the average kinetic energy of trajectories. We characterize the minimizers of the resulting optimization problem (whose objective is the sum of a divergence and this penalty), and show that these time-dependent velocity fields take the straight-line interpolation form above.
- Regularity of the ODE velocity field. We derive explicit upper bounds for the C^k norm of a straightline velocity field that are polynomial in the C^k norm of the corresponding transport map T. This development involves applying a multivariate Faà di Bruno formula in Banach spaces, a procedure that may be of independent interest.
- Explicit links to C^k -smooth densities. For specific transports T, in particular triangular (Knothe-Rosenblatt) transport maps, we then construct upper bounds on the C^k norm of the corresponding straight-line velocity field that have polynomial dependence on the C^k norm of the source and target densities. Along the way, we obtain an explicit upper bound for the C^k norm of the Knothe-Rosenblatt map that depends polynomially on the C^k norms of the source and target densities.
- *Distributional stability.* We relate the approximation error in the velocity field to the error in the distribution induced by the time-one flow map of the resulting ODE, in both Wasserstein distance and Kullback–Leibler (KL) divergence.
- Neural network approximation and optimization. Combining our analysis of the regularity of the velocity field with the preceding stability results, we show that approximation of the target distribution to any desired accuracy $\varepsilon > 0$, in Wasserstein distance or KL divergence, can be achieved by a deep neural network representation of the velocity field whose depth, width, and sparsity are bounded explicitly in terms of ε and the smoothness and dimension of the source and target densities. We then provide guarantees for minimizers of the regularized optimization problem over such neural network classes, showing that there exist velocity fields that render the objective—and hence both the distribution approximation error and the departure from minimum kinetic energy—arbitrarily small.

1.2 Related work

Several papers have studied the approximation power of discrete normalizing flows. Kong and Chaudhuri (2020) investigate basic flow structures (e.g., planar flows, radial flows, Sylvester flows, Householder flows) for L^1 approximation of a target density on \mathbb{R}^d . The authors establish a universal approximation result for d = 1, but show partially negative results for d > 1: there exist distributions that cannot be exactly coupled by such flows, and there are other distributions for which accurate approximation requires composing together a prohibitive number of layers. Teshima et al. (2020a), on the other hand, show that normalizing flows based on the so-called "affine coupling" construction are universal approximators of diffeomorphisms, and consequently that their pushforward distributions converge weakly to any desired target as the complexity of the flow increases in suitable way. Additional universal approximation results have been developed for more specific flow architectures (Huang et al., 2018). However, none of these works characterizes the *rate*

of convergence of the approximation—i.e., how the distribution approximation error that can be achieved scales with the size of the model.

For triangular (Knothe–Rosenblatt) transport maps on $[0, 1]^d$, Zech and Marzouk (2022a) develop a complete approximation theory under the assumption of analytic source and target densities, for neural network or sparse polynomial representations of the map. These results encompass approximation of the maps themselves, but also distribution approximation via the pushforward of a given source distribution by the approximate map. Zech and Marzouk (2022b) extend this analysis to triangular transport maps in infinite dimensions, i.e., on $[0,1]^{\infty}$. Baptista et al. (2024) provide a general framework for analyzing the error of variational approximations of distributions realized using transport maps, and shows how this framework can be applied in specific situations to derive approximation rates.

The preceding works considered approximation of the transport map itself, i.e., direct representations of the 'displacement' of points from the support of the source to the support of the target. In contrast, other recent efforts have addressed approximation issues in *dynamic* representations of transport. Most of these have focused on neural ODEs. Teshima et al. (2020b) show that neural ODEs are universal approximators of smooth diffeomorphisms on \mathbb{R}^d in appropriate Sobolev norms. Li et al. (2019) adapt ideas from dynamical systems to show that neural ODEs are universal approximators of continuous functions from \mathbb{R}^d to \mathbb{R}^m (hence, not only diffeomorphisms) in a L^2 sense, for $d \ge 2$. Yet these universal approximation results do not characterize approximation *rates*, e.g., relate bounds on function approximation error to the size of the network representing the velocity field. Moreover, we note that universal function approximation results are not necessarily relevant to distribution learning: that is, universal approximation of distributions does not require universal function approximation.

Ruiz-Balet and Zuazua (2023) study distribution approximation with neural ODEs, and prove universal approximation for certain target distributions in Wasserstein-1 distance, using an approach that is discrete and constructive. Specifically, they analyze neural ODE-type models from a controllability perspective, explicitly constructing piecewie constant approximations of the target density using a neural network velocity field with ReLU activations. Their analysis does not consider higher-order smoothness, however, and their velocity construction is different from that considered here. In subsequent work, Álvarez-López et al. (2024) show that ReLU velocity fields chosen to be piecewise constant in time can approximate a target distribution arbitrarily closely in relative entropy.

As mentioned earlier, several other works identify neural ODE velocity fields via a *regularized* training objective, i.e., by minimizing a linear combination of a statistical divergence (or negative log-likelihood) and some regularization term. Finlay et al. (2020) argue that a good way of measuring the regularity of the velocity field is through the "acceleration" experienced by a "particle" $X^f(x,t)$ starting at some point x. This acceleration is the total derivative of f in time:

$$\frac{Df(X,t)}{Dt} = \nabla_X f(X,t) \cdot \frac{\partial X}{\partial t} + \frac{\partial f(X,t)}{\partial t} = \nabla_X f(X,t) \cdot f(X,t) + \frac{\partial f(X,t)}{\partial t}.$$
 (1.2)

When this term is zero, particle trajectories will be straight lines. Since the Jacobian matrix $\nabla_X f(X,t)$ is in general not easily accessible, Finlay et al. (2020) choose to implicitly penalize this term by penalizing $|f(X,t)|^2$ and the Frobenius norm $\|\nabla_X f(X,t)\|_F^2$ instead. Similar to Grathwohl et al. (2019), stochastic methods are used to estimate $\|\nabla_X f\|_F^2$ and $\operatorname{tr}(\nabla_X f(X,t))$ in training, where the latter is used in the change of variables formula for the log-likelihood. In more recent work, Onken et al. (2021) propose a discretize-then-optimize approach to training neural ODEs, where a ResNet structure is used to implement the underlying neural network in the ODE. This approach enables exact computation of the Jacobian matrix as well as its trace from the recursive structure of the ResNet. Then, automatic differentiation is used to update the parameters in the neural network, instead of solving the adjoint equation as in Grathwohl et al. (2019) and Finlay et al. (2020). By adopting this discretize-then-optimize approach, we propose to penalize (1.2) directly. We will show that the velocity field f making this term zero is the unique velocity field that yields the minimal kinetic energy among all velocity fields f that produce the same transport map T at time one; hence it is termed the *minimal energy regularization*.

While the neural ODE framework learns the velocity field by minimizing the KL divergence from the target distribution to the pushforward distribution of the source under ODE flow maps, another recent line of research aims to specify the velocity field *a priori* using conditional expectations and to learn the velocity field directly via least-squares regression (Albergo et al., 2023; Albergo and Vanden-Eijnden, 2023; Lipman et al., 2023; Liu et al., 2023). However, the same velocity field approximation questions and distributional stability questions are present in that setting as well. We note that the approximation results we develop in this work are independent of training scheme, and in particular, the straight-line velocity fields we analyze here are central to the rectified flows proposed in Liu et al. (2023).

Indeed, there are a variety of related *distributional stability* results in recent literature (cf. Section 5), addressing the question of how error in the pushforward distribution under an ODE flow map (in some distance or divergence) is controlled by error in the velocity field (in some norm). Benton et al. (2024) show that W_2 error in distribution is controlled by L^2 approximation error of the true velocity field and the time-averaged spatial Lipschitz constant of the approximate flow. In a study of the convergence of continuous normalizing flows, Gao et al. (2024) obtain a stability result almost exactly the same as that in Benton et al. (2024), where W_2 error in distribution is controlled by L^2 error in velocity field times the exponential of spatial Lipschitz constant of the velocity field. Li et al. (2024) analyze a discrete-time version of the probability flow ODE, where TV error in distribution is bounded by terms involving the L^2 error in the score function and in its Jacobian. Again in the setting of probability flow ODEs, Huang et al. (2024) start in continuous time and then considers discretization using a Runge-Kutta scheme. At the continuous level, it is shown that TV error in distribution is controlled by L^2 error in the approximation of the score function and the first and second derivatives of the estimated score; at the discrete level, it is shown that the *p*-th order integrator also converges under an additional assumption that the estimated score function's first p+1 derivatives are bounded. In our work, we show that W^p error in the distribution, for $p \in [1,\infty]$, is controlled by the space-time L^{∞} (i.e., C^{0}) error and spatial Lipschitz constant of f, on compact domains; these are further related to properties of the densities. In addition, we obtain that distribution approximation error in KL is controlled by the C^1 norm of f, again on compact domains.

There are also some results linking properties of the velocity field (e.g., Lipschitz constant in space or time) to properties of the underlying densities. In Benton et al. (2024), the time-averaged spatial Lipschitz constant is related to assumptions on the regularity of all the intermediate distributions between t = 0 and t = 1, along with some Gaussian smoothing; an upper bound is obtained that depends on properties of the chosen interpolant. We note that their regularity assumption is rather different than the C^k smoothness we assume here, as they do not consider higher-order smoothness of the velocity field. Gao et al. (2024) focus on flow matching with linear interpolation. It is shown that the Lipschitz constant of the target velocity field in both the space and time variables is bounded under assumptions of log-concavity/convexity of distributions and boundedness of the domain. In addition, they show that the velocity field itself grows at most linearly with respect to the space variable. No Gaussian smoothing is used in their setting, but they require an early stopping before reaching time t = 1. Both Benton et al. (2024) and Gao et al. (2024) are specific to certain stochastic interpolants, which are different than the straight-line ansatz we analyze here. Also, they do not consider higher-order smoothness of the velocity field or derive upper bounds that are explicit in the densities.

With regard to neural network approximation results, Gao et al. (2024) construct neural network classes

that capture the Lipschitz properties of the velocity field, and derive rates of approximation in the L^{∞} sense. Our companion paper (Marzouk et al., 2024) derives explicit neural network approximation rates for general C^k velocity fields, but its main focus is on statistical finite sample guarantees for neural ODEs trained through likelihood maximization, different from our focus here.

There are also stochastic differential equation (SDE) and specifically neural SDE methods for distribution learning (Tzen and Raginsky, 2019; Song et al., 2021). However, they are rather different than the deterministic ODE approach, and again are not the focus of this work.

2 **Preliminaries**

2.1 Notation and definitions

ODEs and flow maps. We write X(x,t) or $X_f(x,t)$ for the solution of (1.1) with initial condition x at time t = 0, i.e.,

$$X_f(x,t) = x + \int_0^t f(X_f(x,s),s)ds.$$
 (2.1)

Given an initial distribution π_0 , we write π_t or $\pi_{f,t}$ for the pushforward measure $X_f(\cdot, t)_{\sharp}\pi_0$ and $\pi(x, t)$ or $\pi_f(x, t)$ for the corresponding density.

Vectors and multiindices. For $x = (x_1, \ldots, x_d)^{\top} \in \mathbb{R}^d$, |x| is the Euclidean norm. With $\mathbb{N} = \{0, 1, 2, \ldots\}$, we denote multiindices by bold letters such as $v = (v_1, v_2, \ldots, v_d) \in \mathbb{N}^d$, and we use the standard multiindex notations $|v| = \sum_{i=1}^d v_i$ and $v! = \prod_{i=1}^d (v_i!)$. Additionally, $x^v = \prod_{i=1}^d x_i^{v_i}$ and $x_{[k]} \coloneqq (x_1, \ldots, x_k) \in \mathbb{R}^k$ for all $k \leq d$. For two multiindices $v, w \in \mathbb{N}^d, w \prec v$ if and only if one of the following holds: (i) |w| < |v|, (ii) |w| = |v| and there exists a k < d such that $w_1 = v_1, \ldots, w_k = v_k$, but $w_{k+1} < v_{k+1}$.

Derivatives. For $f \in C^1(\mathbb{R}^d, \mathbb{R}^m)$, $\nabla f : \mathbb{R}^d \to \mathbb{R}^{d \times m}$ is the gradient. In case f depends on multiple variables, we write, for example, $\nabla_x f(x,t)$. For a multiindex $\boldsymbol{v} = (v_1, v_2, \dots, v_d) \in \mathbb{N}^d$, where $\mathbb{N} = \{0, 1, 2, \dots\}$, we write $D^{\boldsymbol{v}} f(x) = \frac{\partial^{|\boldsymbol{v}|}}{\partial x_1^{v_1} \dots \partial x_d^{v_d}} f(x)$ for the partial derivative and similarly to the notation above, $D_x^{\boldsymbol{v}} f(x,t)$.

Function spaces. For two Banach spaces X, Y and $n \in \mathbb{N}$ we denote by $\mathcal{L}^n(X; Y)$ the space of all *n*-linear maps from $X^n \to Y$, and by $\mathcal{L}^n_{sym}(X; Y)$ the space of all symmetric *n*-linear maps from $X^n \to Y$ (i.e., $A \in \mathcal{L}^n_{sym}$ iff $A(x_{\sigma(1)}, \ldots, x_{\sigma(n)})$ is independent of the permutation σ of $\{1, \ldots, n\}$). The norms on these spaces are defined as

$$\|A\|_{\mathcal{L}^{n}(X;Y)} \coloneqq \sup_{\|x_{i}\|_{X} \le 1} \|A(x_{1},\dots,x_{n})\|_{Y}, \qquad \|A\|_{\mathcal{L}^{n}_{sym}(X;Y)} \coloneqq \sup_{\|x\|_{X} \le 1} \|A(x,\dots,x)\|_{Y}.$$

We recall that if $A \in \mathcal{L}^n_{sym}(X;Y)$ and $B \in \mathcal{L}^n(X;Y)$ such that $A(x^n) = B(x^n)$ for all $x \in X$, then, see e.g., (Chae, 1985, 14.13),

$$\|A\|_{\mathcal{L}^{n}_{sym}(X;Y)} \le \|B\|_{\mathcal{L}^{n}(X;Y)} \le \exp(n)\|A\|_{\mathcal{L}^{n}_{sym}(X;Y)}.$$
(2.2)

Recall that for $f \in C^k(X, Y)$, the k-th Fréchet derivative $D^k f(x)$ of f at $x \in X$ belongs to $\mathcal{L}^k_{sym}(X; Y)$. For $S \subseteq X$ open and $f \in C^k(S, Y)$ we write

$$\|f\|_{C^k(S)} \coloneqq \sup_{n \le k} \sup_{x \in S} \|D^n f(x)\|_{\mathcal{L}^n(X;Y)}.$$
(2.3)

For $X = \mathbb{R}^d$ and $S \subseteq X$, we use the usual notation $W^{k,p}(S)$, $k \in \mathbb{N}$, $p \in [1, \infty]$, to denote functions with weak derivative up to order k belonging to $L^p(S)$. As a norm, we will use

$$\|f\|_{W^{k,p}(S)} = \begin{cases} \left(\sum_{|\boldsymbol{\alpha}| \le k} \|D^{\boldsymbol{\alpha}}f\|_{L^{p}(S)}^{p}\right)^{\frac{1}{p}} & \text{if } 1 \le p < \infty\\ \max_{|\boldsymbol{\alpha}| \le k} \|D^{\boldsymbol{\alpha}}f\|_{L^{\infty}(S)} & \text{if } p = \infty. \end{cases}$$

Divergences between distributions. Let $(\Omega, \mathcal{F}, \mu)$ be a probability space. For two probability measures ρ and π such that $\rho \ll \mu, \pi \ll \mu$, the information divergences we consider are the following:

- KL (Kullback–Leibler) divergence: Assuming also $\rho \ll \pi$, we define $KL(\rho, \pi) = \int_{\Omega} \log \frac{d\rho}{d\pi}(x)\rho(dx)$.
- If (Ω, m) is also a metric space, then the Wasserstein distance of order p is defined as:

$$W_p(\rho,\pi) = \left(\inf_{\gamma \in \Gamma(\rho,\pi)} \int_{\Omega \times \Omega} m(x,y)^p \gamma(dxdy)\right)^{\frac{1}{p}},$$

where $\Gamma(\rho, \pi)$ is the set of all measures on $\Omega \times \Omega$ with marginals ρ and π . In \mathbb{R}^d , this is simply $W_p(\rho, \pi) = \left(\inf_{\gamma \in \Gamma(\rho, \pi)} \int_{\mathbb{R}^d \times \mathbb{R}^d} |x - y|^p \gamma(dxdy)\right)^{\frac{1}{p}}$.

2.2 Problem setup

In the following we denote by π a *target measure* and by ρ a *source measure* on \mathbb{R}^d . Our general goal is to sample from the target. The source measure is an auxiliary measure that is easy to sample from, and may be chosen at will. Throughout we work under the following assumptions:

Assumption 2.1 (compact support). With $\Omega_0 \coloneqq \operatorname{supp}(\pi)$ and $\Omega_1 \coloneqq \operatorname{supp}(\rho)$, it holds that $\Omega_0, \Omega_1 \subseteq \mathbb{R}^d$ are compact and convex sets. Both ρ and π are absolutely continuous with respect to the Lebesgue measure.

By abuse of notation, we denote the (Lebesgue-) densities of ρ and π by $\rho(x)$ and $\pi(x)$, respectively.

Assumption 2.2 (regularity). There exist constants $L_1 > 1$, $L_2 > 0$ and $k \in \mathbb{N}, k \geq 2$ such that $\|\pi\|_{C^k(\Omega_0)} \leq L_1$, $\|\rho\|_{C^k(\Omega_1)} \leq L_1$ and $\inf_{x \in \Omega_0} \pi(x) \geq L_2$, $\inf_{x \in \Omega_1} \rho(x) \geq L_2$.

We consider two types of problems:

- P1 The target measure π is known through a collection of iid samples. This is the problem considered in, e.g., Grathwohl et al. (2019); Finlay et al. (2020); Onken et al. (2021). The goal is to learn a velocity field f in (1.1) such that with initial distribution chosen to be the target, π₀ = π, the time-one distribution satisfies π_{f,1}(·) = X_f(·, 1)[#]π ≈ ρ. Since the flow map x → X_f(x, 1) is by construction invertible (and its inverse can be evaluated by solving (1.1) backwards in time), one can draw new samples from the source measure and apply the inverse of the flow map to generate (approximate) samples from π. The learned flow map can also be used, without inversion, to estimate the density of π.
- P2 The target measure is known up to a normalizing constant; that is, we can evaluate the unnormalized target density $\tilde{\pi}$. This setting is ubiquitous in Bayesian statistics, since the posterior normalizing constant of a Bayesian model is usually unavailable. This problem is in the reverse direction of the previous problem (Marzouk et al., 2016; Rezende and Mohamed, 2015; Moselhy and Marzouk, 2012); that is, we choose the initial distribution to be the source distribution, $\pi_0 = \rho$, and learn a velocity field f such that $\pi_{f,1}(\cdot) = X_f(\cdot, 1)_{\sharp}\rho \approx \pi$.

From the approximation and algorithmic perspectives, there is no essential difference between problems P1 and P2 above. In both cases, algorithms for learning the velocity field f require: (*i*) a sample from the chosen initial distribution and (*ii*) the ability to evaluate the *desired* time-one density up to a normalizing constant. For the rest of the paper, we will thus adopt the setting of P1 (with initial distribution for the ODE system chosen to be the target π). Our results can be translated to P2 simply by exchanging π and ρ .

The objective functional considered in this work takes the following form:

$$J(f) = \mathcal{D}(X_f(\cdot, 1)_{\sharp}\pi, \rho) + R(f).$$
(2.4)

The first part of the objective is an information divergence between two probability distributions (for example, KL, Wasserstein, etc.). The second part is a regularization term that follows from the discussion in Section 1: we would like the trajectory of the ODE, starting from any initial condition, to be a straight line with constant velocity. In other words, we would like the acceleration in a Lagrangian frame, df(X(x,t),t)/dt, to be zero for all $x \in \Omega_0$ and $t \in [0, 1]$. To this end, we integrate the squared acceleration from (1.2) along the trajectory of a particle $x \in \Omega_0$:

$$R(x,t) = \int_0^t \left| \left(\nabla_X f(X(x,s),s) \right) f(X(x,s),s) + \partial_s f(X,s) \right|^2 ds,$$
(2.5)

and

$$R(f) = \int_{\Omega_0} \int_0^1 \left| \left(\nabla_X f(X(x,s),s) \right) f(X(x,s),s) + \partial_s f(X,s) \right|^2 ds dx$$
(2.6)

We comment here that while our theoretical analysis works for general divergence \mathcal{D} , KL-divergence is the most common objective used in practice. For this purpose, we derived the training algorithm for it in Appendix A.

By the Picard--Lindelöf theorem (Arnold and Silverman, 1978), existence and uniqueness of solutions to the ODE (1.1) requires that the velocity field f(x,t) be continuous in t and Lipschitz continuous in x. Therefore, searching over the space of functions

$$\mathcal{V} = \left\{ f : \mathbb{R}^d \times [0,1] \to \mathbb{R}^d : f \text{ is Lipschitz continuous in } x, \text{ continuous in } t \right\},$$
(2.7)

we have the following optimization problem:

$$\begin{array}{ll} \underset{f \in \mathcal{V}}{\text{minimize}} & J(f) \\ \text{with} & R(f) \text{ defined in (2.6)} \end{array}$$
(OP)

Remark 2.3. In practice, the conditions of the Picard–Lindelöf theorem will always be satisfied for a neural network of finite size with Lipschitz activation functions. In particular, these conditions hold true for ReLU networks, which is what we consider in our theoretical analysis.

3 Existence and structure of minimizers

The objective J(f) is nonnegative, since it is the sum of an information divergence and a nonnegative regularizer. Moreover, as we will see, the optimal solution will make both terms in the objective J(f) zero under our assumptions. First, we state necessary and sufficient conditions on a transport map T such that there exists a velocity field f whose time-one flow map yields straight-line trajectories from x to T(x).

3.1 Existence

Lemma 3.1. Let $\Omega_0 \subseteq \mathbb{R}^d$ be convex and $T \in C^1(\Omega_0, \mathbb{R}^d)$ such that $\det \nabla_x T(x) \neq 0$ for all $x \in \Omega_0$. Then T is injective.

Proof of Lemma 3.1. Assume not. Then there exist $x, y \in \Omega_0$ such that $x \neq y$ and T(x) = T(y). For $s \in [0,1]$ set $f(s) \coloneqq T((1-s)x + sy)$. Since f(0) = f(1), by the mean value theorem there exists $s \in (0,1)$ such that f'(s) = 0. Then $f'(s) = \nabla_x T((1-s)x + sy)(y-x) = 0$. Since $v = y - x \neq 0$, we have $\det(\nabla_x T((1-s)x + sy)) = 0$, which is a contradiction.

Denote in the following, for $x \in \Omega_0 \subset \mathbb{R}^d$, $t \in [0, 1]$, and a map $T : \Omega_0 \to \mathbb{R}^d$,

$$T_t(x) \coloneqq (1-t)x + tT(x), \tag{3.1}$$

i.e., $[0,1] \ni t \mapsto (T_t(x),t)$ parameterizes the straight line of constant velocity between the points (x,0) and (T(x),1) in $\mathbb{R}^d \times [0,1]$. We refer to $t \mapsto T_t(x)$ as the *displacement interpolation* of T. We now investigate under which conditions these lines do not cross for different x, which is necessary for $T_t(x)$ to be expressible as a flow X(x,t) solving (1.1) for a certain f. In other words, we check under what conditions the map $\Omega_0 \ni x \mapsto T_t(x)$ is injective for all $t \in [0,1]$. To state the following lemma, for $A \in \mathbb{R}^{d \times d}$ we let $\sigma(A) := \{\lambda \in \mathbb{R} : \det(A - \lambda I) = 0\}$ denote its spectrum.

Assumption 3.2. It holds that $T \in C^1(\Omega_0, \mathbb{R}^d)$ and

$$(-\infty, 0] \cap \sigma(\nabla_x T(x)) = \emptyset \qquad \forall x \in \Omega_0.$$
(3.2)

Lemma 3.3. Let $\Omega_0 \subseteq \mathbb{R}^d$ be convex and $T \in C^1(\Omega_0, \mathbb{R}^d)$. Then $det(\nabla_x T_t(x)) > 0$ for all $x \in \Omega_0$ and all $t \in [0, 1]$, iff (3.2) holds.

Proof of Lemma 3.3. Since $\nabla_x T_t(x) = (1-t)I + t\nabla_x T(x)$, the map $t \mapsto \det(\nabla_x T_t(x)) \in \mathbb{R}$ is continuous. Because of $\nabla_x T_0(x) = I$, to prove the lemma it is sufficient to show that for every $x \in \Omega_0$, $\sigma(\nabla_x T(x)) \cap (-\infty, 0] = \emptyset$ iff $\det(\nabla_x T_t(x)) \neq 0$ for all $t \in [0, 1]$.

Fix $x \in \Omega_0$. Assume for contradiction that for some $t \in [0, 1]$, we have $\det(\nabla_x T_t(x)) = 0$. Then there exists $v \neq 0$ such that $\nabla_x T_t(x)v = 0 \in \mathbb{R}^d$. Thus $\nabla_x T(x)v = -\frac{1-t}{t}v$ and hence $-\frac{1-t}{t} \in (-\infty, 0]$ is an eigenvalue of $\nabla_x T(x)$. The reverse implication follows similarly. Assume that $s \in \sigma(\nabla_x T(x)) \cap (-\infty, 0]$. Then there exists $v \neq 0$ such that $\nabla_x T(x)v = sv$. Since $t \mapsto -\frac{1-t}{t} : (0,1] \to (-\infty,0]$ is bijective, we can find $t \in (0,1]$ such that $\nabla_x T(x)v = -\frac{1-t}{t}v$, implying $v \in \ker(\nabla_x T_t(x))$ and thus $\det(\nabla_x T_t(x)) = 0$. \Box

Combining the previous two statements establishes the existence of a velocity field such that the timeone flow map of the ODE (1.1) realizes the map $x \mapsto T(x)$, and the ODE dynamics produce straight-line trajectories of constant speed.

Theorem 3.4. Let $k \in \mathbb{N}$ and let $\Omega_0 \subseteq \mathbb{R}^d$ be convex and compact. Assume that $T \in C^k(\Omega_0, \mathbb{R}^d)$ for some $k \geq 2$ satisfies (3.2). With T_t in (3.1) set

$$\Omega_{[0,1]} \coloneqq \left\{ (T_t(x), t) : x \in \Omega_0, \ t \in [0,1] \right\} \subset \mathbb{R}^{d+1}.$$
(3.3)

Then there exists a unique $f : \Omega_{[0,1]} \to \mathbb{R}^d$ such that the solution $X : \Omega_0 \times [0,1] \to \mathbb{R}^d$ of the ODE (1.1) satisfies $X(x,t) = T_t(x)$ for all $x \in \Omega_0$, $t \in [0,1]$. Moreover $\Omega_{[0,1]}$ is simply connected and $f \in C^k(\Omega_{[0,1]}^\circ)$.

A key element of the proof of Theorem 3.4 is that the velocity field f corresponding to the displacement interpolation T_t can be defined implicitly in terms of T: f((1-t)x + tT(x), t) = T(x) - x.

Proof of Theorem 3.4. By (1.1) and because $X(x,t) = T_t(x) = (1-t)x + tT(x)$, we have for $x \in \Omega_0$ and $t \in [0,1]$

$$T(x) - x = \frac{d}{dt}X(x,t) = f(X(x,t),t) = f(T_t(x),t).$$
(3.4)

By Lemma 3.1 and Lemma 3.3, the map

$$(x,t) \mapsto G(x,t) \coloneqq (T_t(x),t) \in \Omega$$

is injective on $\Omega_0 \times [0, 1]$. Thus (3.4) uniquely defines f at each point $G(x, t) \in \Omega$. By construction, this f yields a flow map X as in (1.1) satisfying $X(x, t) = T_t(x)$.

The map $G : \Omega_0 \times [0,1] \to \Omega$ is a continuous bijection, and since $\Omega_0 \times [0,1] \subseteq \mathbb{R}^{d+1}$ is compact, $G^{-1} : \Omega \to \Omega_0 \times [0,1]$ is also a continuous bijection. Since $\Omega_0 \times [0,1]$ is a convex set, it is simply connected. Hence, the homotopy equivalent set $\Omega_{[0,1]}$ must also be simply connected. Moreover, the interior Ω° of Ω is the image of $\Omega_0^\circ \times (0,1)$ under G.

It remains to show $f \in C^k(\Omega_{[0,1]}^\circ)$. Fix $x \in \Omega_0^\circ$ and $t \in (0,1)$. Then

$$\nabla_{(x,t)}G(x,t) = \begin{pmatrix} \nabla_x T_t(x) & T(x) - x \\ 0 & 1 \end{pmatrix} \in \mathbb{R}^{(d+1) \times (d+1)},$$

and this matrix is regular by Lemma 3.3. Since $G \in C^k(\Omega_0 \times [0,1])$, the inverse function theorem (see, e.g., Spivak (1965)[Theorem 2.11]) implies that G^{-1} locally belongs to C^k in a neighbourhood of G(x,t). Since $x \in \Omega_0^{\circ}$ and $t \in (0,1)$ were arbitrary, we have $G^{-1} \in C^k(\Omega_{[0,1]}^{\circ}, \mathbb{R}^{d+1})$. Denote $G^{-1} = (F, E)$ such that $F : \Omega_{[0,1]} \to \mathbb{R}^d$ corresponds to the first d components of G^{-1} . By (3.4), for all $(y,s) \in \Omega_{[0,1]}^{\circ}$ it holds that f(y,s) = T(F(y,s)) - F(y,s), so that f belongs to $C^k(\Omega_{[0,1]}^{\circ})$ as a composition of two C^k functions. \Box

Remark 3.5. Note that $f \in C^k(\Omega_{[0,1]}^\circ)$ means only that f is C^k on the interior of $\Omega_{[0,1]}$. To show that the derivatives are well-defined on the boundary of $\Omega_{[0,1]}$ and that f can be extended to a C^k function outside of $\Omega_{[0,1]}$, certain regularity conditions of the domain are required, which will be discussed in later parts of this section.

3.2 Properties of $\Omega_{[0,1]}$

The set $\Omega_{[0,1]}$ is simply connected, but unlike $\Omega_0 \times [0,1]$, it need not be convex:

Example 3.6 (Rotation). Let $\Omega_0 = \{x \in \mathbb{R}^2 : |x| \leq 1\}$ be the unit disc and let $T : \mathbb{R}^2 \to \mathbb{R}^2$ be the rotation by $\alpha \in [0, 2\pi)$ around $0 \in \mathbb{R}^2$. Then

$$\nabla_x T(x) = \begin{pmatrix} \cos(\alpha) & -\sin(\alpha) \\ \sin(\alpha) & \cos(\alpha) \end{pmatrix}.$$
(3.5)

The spectrum of this matrix consists of the two values $\exp(\pm i\alpha)$, where i denotes the imaginary root of -1. Thus (3.2) holds iff $\alpha \neq \pi$. If $\alpha = \pi$, then T is the negative identity, and thus $T_{1/2}(x) = \frac{1}{2}x - \frac{1}{2}x = 0$ for all $x \in \Omega_0$, so that the all straight lines connecting x and T(x) for $x \in \Omega_0$, meet at $t = \frac{1}{2}$ in the midpoint 0 of the disc. For all $\alpha \in [0, 2\pi) \setminus \{\pi\}$, by Theorem 3.4 and with $\Omega_{[0,1]}$ as in (3.3), there exists a vector field $f \in C^{\infty}(\Omega_{[0,1]})$ such that $T_t(x) = X(x,t)$ for X as in (1.1). One can check that

$$\Omega_{[0,1]} = \left\{ (x,t) \in \mathbb{R}^2 \times [0,1] : |x| \le \sqrt{\sin\left(\frac{\pi}{2} - \frac{\alpha}{2}\right)^2 + \left[t\cos\left(\frac{\pi}{2} - \frac{\alpha}{2}\right) + (1-t)\cos\left(\frac{\pi}{2} + \frac{\alpha}{2}\right)\right]^2} \right\},$$

which is convex if $\alpha = 0$ and nonconvex for all $\alpha \in (0, 2\pi)$.

To approximate the velocity field f from Theorem 3.4 with a neural network, we also need to understand the regularity of the domain $\Omega_{[0,1]}$ on which f is defined. As we will see, $\Omega_{[0,1]}$ is a Lipschitz domain.

Definition 3.7. A bounded domain Ω is called a Lipschitz domain if there exist numbers $\delta > 0$, M > 0, $J \in \mathbb{N}$, and a finite cover of open sets $\{U_j\}_{j=1}^J$ of $\partial\Omega$ such that:

- For every pair of points $x_1, x_2 \in \Omega$ such that $|x_1 x_2| < \delta$ and $dist(x_i, \partial \Omega) < \delta$, i = 1, 2, there exists an index j such that $x_i \in U_j$, i = 1, 2, and $dist(x_i, \partial U_j) > \delta$, i = 1, 2.
- For each *j* there exists some coordinate system $\{\zeta_{j,1}, \ldots, \zeta_{j,d}\}$ in U_j such that the set $\Omega \cap U_j$ consists of all points satisfying $\zeta_{j,d} \leq f_j(\zeta_{j,2}, \ldots, \zeta_{j,d-1})$, where $f_j : \mathbb{R}^{d-1} \to \mathbb{R}$ is a Lipschitz function with Lipschitz constant M.

To show that $\Omega_{[0,1]}$ is a Lipschitz domain, we first need an auxiliary result, Theorem C.2 in Appendix C, establishing that the image of a Lipschitz domain under a sufficiently regular map remains a Lipschitz domain. We can then show the following:

Theorem 3.8. Consider the setting of Theorem 3.4. Then $\Omega_{[0,1]} \subset \mathbb{R}^{d+1}$ in (3.3) is a Lipschitz domain.

Proof of Theorem 3.8. To show that $\Omega_{[0,1]}$ is a Lipschitz domain, first we observe that $\Omega_{[0,1]}$ is the image of $\Omega_0 \times [0,1]$ under the map $(x,t) \to G(x,t) := (tT(x)+(1-t)x,t)$ for $x \in \Omega_0, t \in [0,1]$. Since $\Omega_0 \times [0,1]$ is a product of two convex sets, which is convex, Lemma C.1 shows that the cylinder $\Omega_0 \times [0,1]$ is a Lipschitz domain. To apply Theorem C.2, we need to find a C^1 -diffeomorphism from an open neighborhood \mathcal{O} of $\Omega_0 \times [0,1]$ onto its image. In the context of Theorem 3.4, we have $\det(\nabla_x T_t(x)) > 0$ for all $(x,t) \in \Omega_0 \times [0,1]$. Since $\Omega_0 \times [0,1]$ is a compact set, the infimum of the continuous function $(x,t) \to \det(\nabla_x T_t(x))$ is achieved at some point in the set and thus we can conclude that $\inf_{(x,t)\in\Omega_0 \times [0,1]} \det(\nabla_x T_t(x)) > 0$.

On the other hand, since $T \in C^k(\Omega_0, \mathbb{R}^d)$ for some $k \ge 2$, it follows that $T \in W^{k,\infty}(\Omega_0, \mathbb{R}^d)$. By the extension theorem 3.9, T can be extended to a function $\tilde{T} \in W^{k,\infty}(\mathbb{R}^d, \mathbb{R}^d)$. Since $k \ge 2$, Sobolev embedding shows that $\tilde{T} \in C^1(\mathbb{R}^d, \mathbb{R}^d)$. Now consider the map $\tilde{T}_t(x) = t\tilde{T}(x) + (1-t)x$ for $(x,t) \in \mathbb{R}^{d+1}$. It is clear that $\tilde{T}_t(x)$ is C^1 in (x,t) and also $\tilde{T}_t(x)|_{\Omega_0 \times [0,1]} = T_t(x)$. By the continuity of determinant operator and $\inf_{(x,t)\in\Omega_0\times [0,1]} \det(\nabla_x T_t(x)) > 0$, it follows that there exists an open neighborhood $\mathcal{O} \subset \mathbb{R}^{d+1}$ of $\Omega_0 \times [0,1]$ such that $\det(\nabla_x \tilde{T}_t(x)) > 0$ for all $x \in \mathcal{O}$. Without loss of generality, we can assume \mathcal{O} is convex. This is because we can choose the neighborhood $\Omega_0 \times [0,1] \cup \{B_\epsilon((x,t))|(x,t) \in \partial(\Omega_0 \times [0,1])\}$, which is an open and convex set that can be made arbitrarily close to $\Omega_0 \times [0,1]$ when $\epsilon \to 0$.

Now, consider the extension of G, G(x,t) = (tT(x) + (1-t)x, t) for $(x,t) \in O$. We have

$$\nabla_{(x,t)}\tilde{G}(x,t) = \begin{pmatrix} \nabla_x \tilde{T}_t(x) & \tilde{T}(x) - x \\ 0 & 1 \end{pmatrix} \in \mathbb{R}^{(d+1) \times (d+1)}$$

is a regular matrix for fixed $(x,t) \in O$. Then, the same arguments as in the proof of Theorem 3.4 show that $\tilde{G}(x,t)$ has a global inverse and \tilde{G}^{-1} is C^1 . Therefore, we have a C^1 -diffeomorphism from O onto its image, and Theorem C.2 shows that $\Omega_{[0,1]} = \{(T_t(x),t)\}$ for $x \in \Omega_0, t \in [0,1]$ is a Lipschitz domain.

For Sobolev functions on Lipschitz domains, we have the following extension theorem:

Theorem 3.9 ((Stein, 1970, Chap. 3)). Let $D \subset \mathbb{R}^d$ be a Lipschitz domain.¹ Then there exists a linear operator \mathcal{E} mapping functions on D to functions on \mathbb{R}^d with the following properties:

¹The result in Stein (1970) is stated in terms of so-called "minimally smooth domains," which is a generalization of the notion of Lipschitz domains.

- $\mathcal{E}(f)|_D = f$, that is, \mathcal{E} is an extension operator.
- \mathcal{E} maps $W^{k,p}(D)$ continuously into $W^{k,p}(\mathbb{R}^d)$ for all $1 \leq p \leq \infty$ and all nonnegative integer k.

Combining Theorem 3.4 and 3.8, the extension result in Theorem 3.9 shows that the velocity field f of Theorem 3.4 can be extended to a function in $W^{k,\infty}$ on all of \mathbb{R}^{d+1} .

3.3 Regularized solutions

In the theorem below, we show that the straight-line connections between x and T(x), for a transport map T that pushes forward π to ρ , yield the minimal average kinetic energy, which is why we name the construction *minimal energy regularization*. Here, the "average kinetic energy" is the squared magnitude of the ODE velocity averaged along trajectories, given a distribution π on the initial condition. For any ODE velocity field g(x, t), this quantity can be written in either Lagrangian or Eulerian frames as follows:

$$\int_{\Omega_0} \int_0^1 \pi(x) |g(X_g(x,t),t)|^2 dt \, dx = \int_{\mathbb{R}^d} \int_0^1 \pi_{g,t}(x) |g(x,t)|^2 dt \, dx.$$

Theorem 3.10. Let $\Omega_0, \Omega_1 \subseteq \mathbb{R}^d$ with Ω_0 convex, and let $T \in C^1(\Omega_0, \Omega_1)$ satisfy (3.2). Assume that π and ρ are probability measures on Ω_0, Ω_1 , respectively, and that $T_{\sharp}\pi = \rho$. Then with

$$\mathcal{H} \coloneqq \left\{ g \in \mathcal{V} : X_g(\cdot, 1) |_{\Omega_0} = T \right\}$$

and f from Theorem 3.4, it holds that

$$f = \underset{g \in \mathcal{H}}{\operatorname{arg\,min}} \int_{\mathbb{R}^d} \int_0^1 \pi_{g,t}(x) |g(x,t)|^2 \, \mathrm{d}t \, \mathrm{d}x.$$

Proof of Theorem 3.10. By Theorem 3.4, we know the existence of velocity fields that realize these constructions. We then bound the average kinetic energy from below, using Lagrangian coordinates, as follows:

$$\begin{split} &\int_{\mathbb{R}^d} \int_0^1 \pi_{g,t}(x) |g(x,t)|^2 dt dx = \int_{\Omega_0} \int_0^1 \pi_{g,0}(x) |g(X(x,t),t)|^2 dt dx \\ &= \int_{\Omega_0} \int_0^1 \pi(x) |\partial_t X(x,t)|^2 dt dx \ge \int_{\Omega_0} \pi(x) \left(\int_0^1 |\partial_t X(x,t)| dt \right)^2 dx \quad \ge \int_{\Omega_0} \pi(x) |X(x,1) - X(x,0)|^2 dx \\ &= \int_{\Omega_0} \pi(x) |X(x,1) - x|^2 dx = \int_{\Omega_0} \pi(x) |T(x) - x|^2 dx, \end{split}$$

where the second inequality is due to Jensen's inequality, and equality holds iff $\partial_t X(x,t) = X(x,1) - X(x,0) = T(x) - x$. Then the optimal choice of X is given by X(x,t) = x + t(T(x) - x), which is exactly the displacement interpolation. As a result, the optimal choice for f is given by f(X(x,t),t) = T(x) - x, which is the straight line construction from Theorem 3.4.

Remark 3.11. This construction has important connections to the fluid dynamics formulation of optimal transport (Benamou and Brenier, 2000). Theorem 3.10 shows that for a fixed transport map T, the straight-line construction gives the minimal average kinetic energy. The optimal transport map T is then just the transport map T that minimizes $\int_{\mathbb{R}^d} \pi(x) |T(x) - x|^2 dx$, which is the L²-Wasserstein distance.

With this machinery developed, we are now ready to prove that under our assumptions on the measures π and ρ , (OP) admits optimal solutions that realize the displacement interpolation of transport maps T that push forward π to ρ .

Theorem 3.12. Let π and ρ be measures supported on Ω_0 and Ω_1 , respectively, and let Assumptions 2.1 and 2.2 be satisfied. Then there exists at least one velocity field $f \in \mathcal{V}$ that achieves the global minimum of zero in the optimization problem (OP). Moreover, all global minimizers of (OP) take the form f(y,t) = T(x) - x, with $y = T_t(x)$, for some transport map T such that $T_{\sharp}\pi = \rho$, where $(y,t) \in \Omega_{[0,1]} \coloneqq \{(T_t(x),t) : x \in \Omega_0, t \in [0,1]\} \subset \mathbb{R}^{d+1}$ and T satisfies (3.2).

To show the existence of a velocity field that achieves J(f) = 0, in the proof of Theorem 3.12 we consider a velocity field that realizes the optimal transport map.

Proof of Theorem 3.12. Clearly, the objective function is bounded from below by zero. We first show that the velocity field corresponding to the optimal transport map achieves this minimum value. The existence of optimal transport map (associated with quadratic cost) under our assumptions and the fact that the map can be written as the gradient of a convex potential ϕ are well-established results in the theory of optimal transport maps (see e.g., Villani (2008) and Brenier (1991)). By Assumptions 2.1,2.2 and Theorem C.3, the optimal transport map is given by $T(x) = \nabla \phi(x)$ for some $\phi \in C^{k+2}(\Omega_0)$ that is strictly convex. Therefore, $\nabla T(x)$ has real and nonnegative eigenvalues.

Since $\phi \in C^{k+2}(\Omega_0)$, the following Monge-Ampere equation is satisfied in the classical sense (Brenier (1991)), :

$$\det(\nabla^2 \phi(x)) = \frac{\pi(x)}{\rho(\nabla \phi(x))}, \forall x \in \Omega_0.$$

Since the densities are both bounded away from zero, we can conclude from the Monge-Ampere equation that det $\nabla T(x) = \det \nabla^2 \phi(x) > 0$, $\forall x \in \Omega_0$. In particular, Assumption 3.2 is satisfied and Theorem 3.4 establishes the existence of a velocity field such that the time-one flow map of the ODE realizes this T(x) and the ODE dynamics yield straight line trajectories.

Now, suppose that there is a continuous velocity field f that achieves zero loss in (A.1). Since the densities are continuous and bounded from below by a constant, the expectation $\mathbb{E}_{\pi}[R(x,1)] = 0$ implies that $R(x,1) = 0 \forall x \in \Omega_0$. That is, along each trajectory X(x,t) starting from $x \in \Omega_0$, we have $\frac{df(X(x,t),t)}{dt} = 0$, i.e., f is constant in time along each trajectory. In other words, f(X(x,t),t) = g(x) for some function g. Now consider the ODE $\frac{dX}{dt} = f(X,t) = g(x)$; the solution is X(t) = g(x)t + C, where C is constant in t. To make the KL-divergence zero, we must have X(1) = T(x) for some transport map T such that $T_{\sharp}\pi = \rho$, and we also have X(0) = x as the initial condition. Solving the equations gives g(x) = T(x) - x. That is, the velocity field must take the form T(x) - x for some transport map T.

4 Regularity of the velocity field *f*

As we have seen in Theorem 3.4, for a transport $T : \Omega_0 \to \Omega_1$ as in Assumption 3.2, there exists a unique velocity field $f : \Omega_{[0,1]} \to \mathbb{R}^d$ such that $T(x) = X_f(x,1)$ for all $x \in \Omega_0$. This f is the unique minimizer of the objective (A.1). Furthermore we have an explicit formula for f: With $G : \Omega_0 \times [0,1] \to \Omega_{[0,1]}$, $G(x,t) \coloneqq (tT(x) + (1-t)x, t)$ define $F : \Omega_{[0,1]} \to \Omega_0$ as the first d components of G^{-1} , then

$$f(y,s) = T(F(y,s)) - F(y,s) \qquad \forall (y,s) \in \Omega_{[0,1]}.$$
(4.1)

Based on (4.1), in this section we investigate the regularity of the velocity field f.

As we will see, f inherits the regularity of T, in the sense that $T \in C^k(\Omega_0)$ implies $f \in C^k(\Omega_{[0,1]})$. Essentially, this follows by the inverse function theorem, which yields $G^{-1} \in C^k(\Omega_{[0,1]})$ so that f in (4.1) is a composition of C^k functions. Since the approximability of f by neural networks crucially depends on $\|f\|_{C^k(\Omega_{[0,1]})}$ (see Section 6 ahead), we will carefully derive a precise bound on this norm. We proceed as follows: In Section 4.1, we discuss regularity of f for arbitrary transport maps T. Subsequently, in Section 4.2 we deepen the discussion in the special case of triangular transport maps (which yield triangular velocity fields f).

4.1 General transports

To bound the norm of f in $C^k(\Omega_{[0,1]})$, our proof strategy is to first upper bound $||F||_{C^k(\Omega_{[0,1]})}$, and then use a version of Faá di Bruno's formula to estimate the norm of the composition of F with T.

Since F is obtained as the first d components of the inverse $G : \Omega_{[0,1]} \to \Omega_0 \times [0,1]$, we first provide some abstract results about how to bound the kth derivative of an inverse function. We start by recalling Faá di Bruno's formula in Banach spaces. For completeness we have added the proof in Appendix C, but emphasize that the argument is the same as for real valued functions. In the following, for a multilinear map $A \in L^n(X, Y)$ we write $A(x^n)$ to denote $A(x, \ldots, x)$.

Theorem 4.1 (Faá di Bruno). Let $k \in \mathbb{N}$. Let X, Y and Z be three Banach spaces, and let $F \in C^k(X,Y)$ and $G \in C^k(Y,Z)$.

Then for all $0 \le n \le k$ and with $T_n := \{ \boldsymbol{\alpha} \in \mathbb{N}^n : \sum_{j=1}^n j\alpha_j = n \}$, for all $x, h \in X$ the nth derivative $[D^n(G \circ F)](x)(h^n) \in Z$ of $G \circ F$ at x evaluated at $h^n \in X^n$ equals

$$\sum_{\boldsymbol{\alpha}\in T_n} \frac{n!}{\boldsymbol{\alpha}!} [D^{|\boldsymbol{\alpha}|}G](F(x)) \left(\underbrace{\frac{[DF(x)](h)}{1!}, \dots, \frac{[DF(x)](h)}{1!}}_{\alpha_1 \text{ times}}, \dots, \underbrace{\frac{[D^nF(x)](h^n)}{n!}, \dots, \frac{[D^nF(x)](h^n)}{n!}}_{\alpha_n \text{ times}}\right).$$

Additionally we need the inverse function theorem, the proof of which can also be found in Appendix C.

Theorem 4.2 (Inverse function theorem). Let $k \ge 1$, let X, Y be two Banach spaces, and let $F \in C^k(X, Y)$. At every $x \in X$ for which $DF(x) \in L^1(X, Y)$ is an isomorphism, there exists an open neighbourhood $O \subseteq Y$ of F(x) and a function $G \in C^k(O, X)$ such that F(G(y)) = y for all $y \in O$.

Moreover, for every $n \leq k$ there exists a continuous function $C_n : \mathbb{R}^{n+1}_+ \to \mathbb{R}_+$ (independent of F, G, O) such that for y = F(x) with x as above

$$\|D^{n}G(y)\|_{\mathcal{L}^{n}_{\rm sym}(Y;X)} \leq C_{n}(\|[DF(x)]^{-1}\|_{\mathcal{L}^{1}_{\rm sym}(Y;X)}, \|DF(x)\|_{\mathcal{L}^{1}_{\rm sym}(X;Y)}, \dots, \|D^{n}F(x)\|_{\mathcal{L}^{n}_{\rm sym}(X;Y)}).$$
(4.2)

We start by giving a bound on the derivatives of the composition of functions.

Corollary 4.3. Let $S_1 \subseteq X$, $S_2 \subseteq Y$, $S_3 \subseteq Z$ be three open subsets of the Banach spaces X, Y and Z. Suppose $k \in \mathbb{N}$ and $F \in C^k(S_1, S_2)$ and $G \in C^k(S_2, S_3)$. Then $\|G \circ F\|_{C^k(S_1)} \leq k^k \exp(k) \|G\|_{C^k(S_2)} \max\{1, \|F\|_{C^k(S_1)}\}^k$. *Proof.* By Theorem 4.1, for all $x \in S_1$

$$\|D^{n}(G \circ F)](x)\|_{\mathcal{L}^{n}_{\mathrm{sym}(X;Y)}} \leq \|G\|_{C^{n}(S_{2})} \sum_{\boldsymbol{\alpha} \in T_{n}} \frac{n!}{\boldsymbol{\alpha}!} \prod_{j=1}^{n} \frac{\|F\|_{C^{j}(S_{1})}^{\alpha_{j}}}{(j!)^{\alpha_{j}}}$$
$$\leq \|G\|_{C^{n}(S_{2})} \max\{1, \|F\|_{C^{j}(S_{1})}\}^{n} \sum_{\boldsymbol{\alpha} \in T_{n}} \frac{n!}{\boldsymbol{\alpha}!} \prod_{j=1}^{n} \frac{1}{(j!)^{\alpha_{j}}}, \qquad (4.3)$$

where we used $\sum_{j=1}^{n} \alpha_j \leq n$ for all $\alpha \in T_n$, and that $||F||_{C^j(S_1)} \leq ||F||_{C^n(S_1)}$ for all $j \leq n$ by definition of the norm. By Lemma C.6, the last sum in (4.3) is bounded by n^n . Finally (2.2) and the definition of the C^n -norm in (2.3) imply claim.

We next use Faa di Bruno's formula, to bound the derivatives of the inverse G^{-1} of a function G.

Proposition 4.4. Consider the setting of Theorem 4.2. Let $S \subseteq X$ be open such that $DG(x) \in L(X,Y)$ is an isomorphism for each $x \in S$ and let there be a continuous function $F : G(S) \to X$ such that F(G(y)) = y for all $y \in G(S)$. Additionally assume that for some $\gamma > 0$

$$\sup_{x \in S} \| [DG(x)]^{-1} \|_{\mathcal{L}^1(Y;X)} \le \gamma \quad and \quad \| G \|_{C^k(S)} \le \gamma.$$
(4.4)

Then

$$\|D^n F(y)\|_{\mathcal{L}^n(Y,X)} \le e^k k^{k^2} \gamma^{3k-2} \qquad \forall y \in G(S).$$

$$(4.5)$$

Proof of Proposition 4.4. We proceed by induction over n, and will show that for all $y \in G(S)$ and all $1 \le n \le k$

$$\|D^{n}F(y)\|_{\mathcal{L}_{\text{sym}^{n}(X;Y)}} \le n^{n^{2}}\gamma^{3n-2}.$$
(4.6)

Then (2.2) implies the claim.

For n = 1, $\mathrm{Id} = D(G \circ F)(y)$ for all $y \in G(S)$. By the chain rule, $(D(G \circ F)(y) \circ (DF(y)) = \mathrm{Id}$, so that $DF(y) = [DG(F(y))]^{-1}$. Thus $\|DF(y)\|_{\mathcal{L}^1(X;Y)} \leq \gamma$ by (4.4). This implies (4.6) for n = 1.

Now let $n \ge 1$. Then for any $y \in G(S)$ by (C.3)

$$\begin{split} \|D^{n}F(y)\|_{\mathcal{L}^{n}_{\mathrm{sym}(Y;X)}} \leq & \|([DG](F(y)))^{-1}\|_{\mathcal{L}^{1}(Y;X)} \\ & \cdot \left(\sum_{\pmb{\alpha}\in\bar{T}_{n}}\frac{n!}{\alpha!}\|D^{|\pmb{\alpha}|}G(F(y))\|_{\mathcal{L}^{|\pmb{\alpha}|}(X;Y)}\prod_{m=1}^{n-1}\left(\frac{\|D^{m}F(y)\|_{\mathcal{L}_{\mathrm{sym}}m(Y;X)}}{m!}\right)^{\alpha_{m}}\right). \end{split}$$

Using the induction Assumption (4.6) to bound $||D^m F(y)||_{\mathcal{L}_{sym}^m(Y;X)} \le m^{m^2} \gamma^{3m-2}$ for $1 \le m < n$, we find for $y \in G(S)$

$$\begin{split} \|D^n F(y)\|_{\mathcal{L}_{\operatorname{sym}^n(Y;X)}} &\leq \gamma^2 \left(\sum_{\boldsymbol{\alpha}\in\bar{T}_n} \frac{n!}{\boldsymbol{\alpha}!} \prod_{m=1}^{n-1} \left(\frac{m^m \gamma^{3m-2}}{m!} \right)^{\alpha_m} \right) \\ &\leq \gamma^2 m^{\sum_{m=1}^{n-1} m^2 \alpha_m} \gamma^{3n-4} \left(\sum_{\boldsymbol{\alpha}\in\bar{T}_n} \frac{n!}{\boldsymbol{\alpha}!} \prod_{m=1}^{n-1} \frac{1}{(m!)^{\alpha_m}} \right) \\ &\leq \gamma^{3n-2} n^{(n-1)n} \left(\sum_{\boldsymbol{\alpha}\in\bar{T}_n} \frac{n!}{\boldsymbol{\alpha}!} \prod_{m=1}^{n-1} \frac{1}{(m!)^{\alpha_m}} \right), \end{split}$$

where we have used $\sum_{m=1}^{n-1} m\alpha_m = n$ and $\sum_{m=1}^{n-1} \alpha_m \ge 2$ for all $\alpha \in \overline{T}_n$. The term in the sum is bounded by n^n according to Lemma C.6. Thus for $2 \le n \le k$,

$$||D^n F(y)||_{\mathcal{L}^n_{\text{sym}}(Y;X)} \le n^{n^2} \gamma^{3n-2}$$

which shows (4.6) and concludes the proof.

We now present our first bound on $||f||_{C^k}$. The estimate depends on $||T||_{C^k}$ as well as

$$c_T := \sup_{t \in [0,1]} \sup_{x \in \Omega_0} \| (\nabla T_t(x))^{-1} \|_{\mathbb{R}^{d \times d}}.$$
(4.7)

We subsequently discuss situations in which we can give precise bounds on this constant.

Theorem 4.5. Let Assumption 2.1 be satisfied. Let $k \in \mathbb{N}$ and let $T \in C^k(\Omega_0, \Omega_1)$ satisfy Assumption 3.2. Then for the velocity field $f : \Omega_{[0,1]} \to \mathbb{R}^d$ in (4.1) it holds with

$$\gamma := \max\{2, 1 + c_T\}(1 + \|T\|_{C^k(\Omega_0)} + \sup_{x \in \Omega_0} \|x\|),$$

that

$$||f||_{C^k(\Omega_{[0,1]})} \le 2k^{k^3+k} e^{k^2+k} \gamma^{3k^2-2k+1}$$

Proof of Theorem 4.5. Due to $f = T \circ F - F$ (cp. (4.1)),

$$\|f\|_{C^{k}(\Omega_{[0,1]})} \leq \|T \circ F\|_{C^{k}(\Omega_{[0,1]})} + \|F\|_{C^{k}(\Omega_{[0,1]})}.$$
(4.8)

Moreover, since F is given as the first d components of $G^{-1}: \Omega_{[0,1]} \to \Omega_0 \times [0,1]$, it holds $||F||_{C^k(\Omega,\mathbb{R}^d)} \le ||G^{-1}||_{C^k(\Omega,\mathbb{R}^{d+1})}$. We start by bounding the latter norm.

By definition of $G(x,t) = (T_t(x), t) = (tT(x) + (1-t)x, t),$

$$DG(x,t) = \begin{pmatrix} \nabla_x T_t(x) & T(x) - x \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} t\nabla T(x) + (1-t)I & T(x) - x \\ 0 & 1 \end{pmatrix}.$$

An application of Lemma C.7 and the assumption that $c_T = \sup_{x \in \Omega_0} \|(\nabla_x T_t(x))^{-1}\| < \infty$ gives for all $(x,t) \in \Omega_0 \times [0,1]$

$$\|[DG(x,t)]^{-1}\|_{2} \le 1 + (1 + \|T - \mathrm{Id}\|_{L^{\infty}(\Omega_{0})})\|(\nabla_{x}T_{t}(x))^{-1}\|_{2} \le 1 + (1 + \|T - \mathrm{Id}\|_{L^{\infty}(\Omega_{0})})c_{T}$$

Next we bound the derivatives of G. For n = 1,

$$\begin{aligned} \|DG(x,t)\|_{\mathcal{L}^{1}(\mathbb{R}^{d+1};\mathbb{R}^{d+1})} &\leq \|\nabla_{x}T_{t}(x)\|_{2} + \|T(x) - x\|_{2} + 1 \leq \|T\|_{C^{1}(\Omega_{0})} + 1 + \|T - \mathrm{Id}\|_{C^{0}(\Omega_{0})} + 1 \\ &\leq 2 + 2\|T\|_{C^{1}(\Omega_{0})} + \max_{x \in \Omega_{0}} \|x\|_{2}. \end{aligned}$$

For $n \ge 2$, we first write $G(x,t) = (G_1(x,t), G_2(x,t))$ where $G_1(x,t) = T_t(x)$ and $G_2(x,t) = t$. Then for $2 \le n \le k$ and $(x,t) \in \Omega_0 \times [0,1]$

$$\|D^{n}G(x,t)\|_{\mathcal{L}^{n}(\mathbb{R}^{d+1};\mathbb{R}^{d+1})} \leq \|D^{n}G_{1}(x,t)\|_{\mathcal{L}^{n}(\mathbb{R}^{d+1};\mathbb{R}^{d})} + \|D^{n}G_{2}(x,t)\|_{\mathcal{L}^{n}(\mathbb{R}^{d+1};\mathbb{R})}$$

The second term is bounded by 1 since $t \in [0, 1]$. For the first term, due to $D_t^2 G_1(x, t) \equiv 0$,

$$\begin{split} \|D^{n}G_{1}(x,t)\|_{\mathcal{L}^{n}(\mathbb{R}^{d+1};\mathbb{R}^{d+1})} &\leq \|D_{x}^{n}G_{1}(x,t)\|_{\mathcal{L}^{n}(\mathbb{R}^{d};\mathbb{R}^{d}))} + \|D_{x}^{n-1}(T(x)-x)\|_{\mathcal{L}^{n-1}(\mathbb{R}^{d};\mathbb{R}^{d})} \\ &\leq \|D_{x}^{n}(x)\|_{\mathcal{L}^{n}(\mathbb{R}^{d};\mathbb{R}^{d})} + \|D_{x}^{n}T(x)\|_{\mathcal{L}^{n}(\mathbb{R}^{d};\mathbb{R}^{d})} + \|D_{x}^{n-1}T(x)\|_{\mathcal{L}^{n-1}(\mathbb{R}^{d};\mathbb{R}^{d})} + \|D_{x}^{n-1}(x)\|_{\mathcal{L}^{n-1}(\mathbb{R}^{d};\mathbb{R}^{d})} \\ &\leq \|D_{x}^{n}T(x)\|_{\mathcal{L}^{n}(\mathbb{R}^{d};\mathbb{R}^{d})} + \|D_{x}^{n-1}T(x)\|_{\mathcal{L}^{n-1}(\mathbb{R}^{d};\mathbb{R}^{d})} + 1. \end{split}$$

We conclude with $M := \max_{x \in \Omega_0} \|x\|_2$ that for all $0 \le n \le k$ and all $(x, t) \in \Omega_0 \times [0, 1]$

$$\|D^{n}G(x,t)\|_{\mathcal{L}^{n}_{\text{sym}\,\mathbb{R}^{d+1};\mathbb{R}^{d+1}}} \leq 2(\|T\|_{C^{n}(\Omega_{0})} + 1 + M) \leq 2(\|T\|_{C^{k}(\Omega_{0})} + 1 + M).$$

With

$$\gamma := \max\{2, 1 + c_T\}(\|T\|_{C^k(\Omega_0)} + 1 + M)\}$$

Proposition 4.4 then implies

$$||F||_{C^k(\Omega_{[0,1]})} \le ||G^{-1}||_{C^k(\Omega_{[0,1]})} \le e^k k^{k^2} \gamma^{3k-2}.$$

Furthermore, Corollary 4.3 and (4.8) yield

$$\|f\|_{C^{k}(\Omega_{[0,1]})} \le k^{k} e^{k} \|T\|_{C^{k}(\Omega_{0})} (e^{k} k^{k^{2}} \gamma^{3k-2})^{k} + e^{k} k^{k^{2}} \gamma^{3k-2} \le 2k^{k^{3}+k} e^{k^{2}+k} \gamma^{3k^{2}-2k+1}.$$

Our main observation is, that $||f||_{C^k(\Omega_0)}$ behaves at worst polynomial in $||T||_{C^k(\Omega_0)}$ and c_T in (4.7). One important instance where we can give a precise bound on c_T , is if ∇T is close to identity matrix $I_d \in \mathbb{R}^{d \times d}$ in the sense $||\nabla T(x) - I_d||_2 < 1$. Since T is a transport pushing forward the source π to the target ρ , this condition can be interpreted as the source and the target not being too different.

Lemma 4.6. Suppose that $\sup_{x \in \Omega_0} \|\nabla T(x) - I_d\|_2 = \delta < 1$, where I_d is d-by-d identity matrix. Then the constant in (4.7) satisfies $c_T \leq \frac{1}{1-\delta}$.

Proof. The assumption implies that for all $t \in [0, 1]$

$$\|\nabla_x T_t(x) - I_d\|_2 = \|t\nabla T(x) + (1-t)I_d - (tI_d + (1-t)I_d)\|_2 = t\|\nabla T(x) - I_d\|_2 \le \delta.$$

Since for any $A \in \mathbb{R}^{d \times d}$ with $||A - I_d||_2 = \delta < 1$ we have $A^{-1} = \sum_{j \in \mathbb{N}} (I - A)^j$ and thus $||A^{-1}||_2 \le \frac{1}{1 - \delta}$, the claim follows.

Another instance where c_T can be bounded is if T is a triangular transport. We next discuss this case in more detail.

4.2 Triangular transports

A special type of transport map commonly used in practice is the so-called Knothe-Rosenblatt (KR) map. To avoid further techincalities, throughout this section we restrict ourselves to measures on *d*-dimensional cubes, i.e.

$$\Omega_0 = \Omega_1 = [0, 1]^d. \tag{4.9}$$

The KR map, is the unique transport satisfying *triangularity* and *monotonicity*. To formally introduce these notions, recall that we use the notation convention $x_{[i]} = (x_i)_{i=1}^j$.

Definition 4.7. We say that a map $T = (T_j)_{j=1}^d : [0,1]^d \to [0,1]^d$ is **triangular** iff T_j depends only on $x_{[j]}$ (but not on x_{j+1}, \ldots, x_d) for each $j = 1, \ldots, d$. A triangular map T is called **monotone** iff for each $j = 1, \ldots, d$. A triangular map $T_j(x_{[j]}) > 0$ for all $x_{[j]} \in [0,1]^j$.

Under rather mild assumptions on the reference and target one can show existence and uniqueness of the KR map (Santambrogio (2015)). Moreover, it allows an explicit construction, which we recall in Appendix B. The goal of the present section is to derive bounds on the norms of the velocity field associated with the KR-map.

We start our analysis by pointing out that triangularity of the transport T implies a similar structure for the corresponding velocity field f:

Lemma 4.8. Consider the setting of Theorem 3.4 and let Ω_0 , Ω_1 be as in (4.9). If $T : \Omega_0 \to \Omega_1$ is triangular, then $f : \Omega_{[0,1]} \to \mathbb{R}^d$ is triangular in the sense $f(x,t) = (f_i(x_{[i]},t))_{i=1}^d$.

Proof. Let the solution $X : \Omega_0 \times [0,1] \to \mathbb{R}^d$ of (1.1) satisfy X(x,t) = tT(x) + (1-t)x. Then for the velocity field $f : \Omega_{[0,1]} \to \mathbb{R}^d$ in (1.1) it holds f(X(x,t),t) = T(x) - x, i.e. for each $i = 1, \ldots, d$

$$f_i(X(x,t),t) = T_i(x_{[i]}) - x_i \qquad \forall (x,t) \in \Omega_0 \times [0,1].$$

To prove the lemma we show that for all $i \in \{1, ..., d\}$ it holds

$$\frac{\partial}{\partial X_j} f_i(X(x,t),t) = 0 \qquad \forall j > i, \quad \forall (x,t) \in \Omega_0 \times [0,1].$$
(4.10)

Fix $i \in \{1, \ldots, d\}$. To prove (4.10), we proceed by induction over $j = i + 1, \ldots, d$ starting with j = d. Note that the triangularity of T implies that also $X(x,t) = (X_l(x_{[l]},t))_{l=1}^d$ inherits this triangular structure. Hence $\frac{\partial}{\partial x_d}X_l(x_{[l]}) = 0$ for all l < d. Consequently

$$\frac{\mathrm{d}}{\mathrm{d}x_d}f_i(X(x,t),t) = \frac{\partial}{\partial X_d}f_i(X(x,t),t)\frac{\partial}{\partial x_d}X_d(x,t) = 0 \qquad \forall (x,t) \in \Omega_0 \times [0,1].$$

By the monotonicity assumption on T it holds $\frac{\partial T_d(x)}{\partial x_d} > 0$, and therefore

$$\frac{\partial}{\partial x_d} X_d(x,t) = t \frac{\partial}{\partial x_d} T_d(x) + (1-t) > 0 \qquad \forall (x,t) \in \Omega_0 \times [0,1].$$

Hence $\frac{\partial f_i(X(x,t),t)}{\partial X_d} = 0$ for all $(x,t) \in \Omega_0 \times [0,1]$.

Now suppose (4.10) is true for all j = k + 1, ..., d and some $k \ge i$. Then, using $\frac{\partial X_j(x,t)}{\partial x_k} = 0$ whenever k > j and as well as (4.10) whenever j > k

$$\frac{\mathrm{d}}{\mathrm{d}x_k}f_i(X(x,t),t) = \sum_{j=1}^d \frac{\partial}{\partial X_j}f_i(X(x,t),t)\frac{\partial}{\partial x_k}X_j(x,t) = \frac{\partial}{\partial X_k}f_i(X(x,t),t)\frac{\partial}{\partial x_k}X_k(x,t) = 0,$$

for $(x,t) \in \Omega_0 \times [0,1]$. Since as before $\frac{\partial X_k(x_{[k]},t)}{\partial x_k} > 0$ we find $\frac{\partial f_i(X(x,t),t)}{\partial X_j} = 0$.

In Theorem 3.8 we showed that the domain $\Omega_{[0,1]}$ (cp. (3.3)) of the velocity field f is a Lipschitz domain. For triangular maps and if (4.9) it even holds $\Omega_{[0,1]} = [0,1]^d \times [0,1]$, i.e. the trajectories of the solutions to (1.1) cover the whole d + 1 dimensional cube: **Proposition 4.9.** Let $T : [0,1]^d \to [0,1]^d$ be a monotone, triangular and bijective map. Then

$$\Omega_{[0,1]} = [0,1]^d \times [0,1].$$

Proof. It is easy to see, that for a monotone triangular map, $T : [0,1]^d \to [0,1]^d$ being bijective is equivalent to $x_j \mapsto T_j(x_{[j]})$ being bijective from $[0,1] \to [0,1]$ for each $j = 1, \ldots, d$, see e.g., (Zech and Marzouk, 2022b, Lemma 3.1).

Denote $T_t(x) = (1-t)T(x) + tx$ and additionally $T_{t,j}(x) = (1-t)T_j(x_{[j]}) + tx_j$ $(x,t) \in \Omega_0 \times [0,1]$. For every $t \in [0,1]$, $T_t : [0,1]^d \to [0,1]^d$ is a convex combination of two monotone, triangular bijective maps. Hence $T_t : [0,1]^d \to [0,1]^d$ is also triangular. Moreover, for each $j \in \{1,\ldots,d\}$ and $t \in [0,1]$, $x_j \mapsto T_{t,j}(x_{[j]})$ is a convex combination of two monotonically increasing maps that bijectively map [0,1]onto itself, and thus this function has the same property. In all this shows that $T_t : [0,1]^d \to [0,1]^d$ is monotone, triangular and bijective for each $t \in [0,1]$, which implies the claim.

We wish to apply Theorem 4.5 to bound $||f||_{C^k(\Omega)}$. To do so, it remains to bound $||T||_{C^k([0,1]^d,[0,1]^d)}$ and c_T as below.

Lemma 4.10. Let π , ρ be densities supported on $[0,1]^d$ and satisfy regularity Assumption 2.2. Let $T \in C^1([0,1]^d, [0,1]^d)$ be the Knothe-Rosenblatt map such that $T_{\sharp}\pi = \rho$. Then the constant c_T from (4.7) satisfies

$$c_T := \sup_{x \in [0,1]^d} \| (\nabla T_t(x))^{-1} \|_{R^{d \times d}} \le (\frac{L_1}{L_2})^{2d} \max\{1, \|T\|_{C^1([0,1]^d)}\}^{d-1}$$

Proof. In this proof we use the notation and construction of the transport map provided in appendix **B**. In particular π_i is the marginal density of π in (x_1, \ldots, x_i) , and with $F_{\rho,i}$, $F_{\pi,i}$ as in (B.2), we let

$$G_{\rho,i}(x_{[i-1]}, \cdot) = F_{\rho,i}(x_{[i-1]}, \cdot)^{-1}$$

By construction, the Jacobian ∇T is a triangular matrix. We shall compute the diagonal entries of the Jacobian matrix, which are the eigenvalues. By (B.3)

$$T_i(x) = G_{\rho,i}(T_1(x_1), ..., T_{i-1}(x_{[i-1]}), F_{\pi,i}(x)).$$

Taking derivatives in x_i , we have

$$\partial_{x_i} T_i(x) = \partial_{x_i} G_{\rho,i}(T_1(x_1), \dots, T_{i-1}(x_{[i-1]}), \pi_i(x)) \partial_{x_i} F_{\pi,i}(x).$$

Recall that $F_{\pi,i}(x)$ is the CDF of x_i when viewing $x_{[i-1]}$ as fixed, thus we have $\partial_{x_i} F_{\pi,i}(x) = \pi_i(x)$. Note that $G_{\rho,i}(x_{[i-1]}, F_{\rho,i}(x)) = x_i$. Taking derivative in x_i , we have

$$(\partial_{x_i} G_{\rho,i})(x_{[i-1]}, F_{\rho,i}(x))(\partial_{x_i} F_{\rho,i}(x)) = 1.$$

Note that $F_{\rho,i}(x_{[i-1]}, \cdot) : [0,1] \to [0,1]$ is a bijection. We make the substitution $y_i = F_{\rho,i}(x)$ and we have for all $(x_{[i-1]}, y_i) \in [0,1]^{i-1} \times [0,1]$,

$$(\partial_{x_i} G_{\rho,i})(x_{[i-1]}, y_i) = \frac{1}{\partial_{x_i} F_{\rho,i}(x_{[i-1]}, G_{\rho,i}(x_{[i-1]}, y_i))}.$$

Hence

$$\partial_{x_i} T_i(x_{[i]}) = \frac{\pi_i(x_{[i]})}{\rho_i(T_1(x_1), \dots, T_{i-1}(x_{[i-1]}), G_{\rho,i}(T_1(x_1), \dots, T_{i-1}(x_{[i-1]}), F_{\pi,i}(x)))}.$$

By Assumption 2.2, ρ and π are bounded from above and below by L_1 and L_2 . Thus $\pi_i(x_{[i]})$ and $\rho_i(x_{[i]})$ are bounded from above and below by $\frac{L_1}{L_2}$ and $\frac{L_2}{L_1}$, and $\partial_{x_i}T_i(x_{[i]})$ is bounded from below by $\frac{L_2^2}{L_1^2}$. Note that the diagonal entries of $\nabla_x T_t(x)^{-1} = [(1-t)I_{d\times d} + t\nabla_F T(F(y,t))]^{-1}$ are exactly $\{\frac{1}{1-t+t\sigma_i}\}_{i=1}^d$, which are lower bounded by $\frac{L_2^2}{(1-t)L_2^2+tL_1^2} \ge \frac{L_2^2}{L_1^2}$.

Therefore, we have $\inf_{x \in \Omega_0} \det(\nabla_x T(x)) \ge \left(\frac{L_2}{L_1}\right)^{2d}$. Applying Lemma C.9 then gives the result. \Box

When T is the KR triangular map, $||T||_{C^K}([0,1]^d, [0.1]^d)$ can be computed explicitly in terms of densities as we see below.

Theorem 4.11. Let $\Omega_0 = \Omega_1 = [0,1]^d$ and let π , ρ satisfy Assumption 2.2 with the constants $k \geq 2$, $0 < L_1 \leq L_2 < \infty$.

Then there exist constants $C_{k,d}$ (depending on k and d but independent of ρ , π) and $\beta_d > 0$ (depending on d but independent of k, ρ , π) such that the KR map T pushing forward π to ρ satisfies

$$||T||_{C^{k}([0,1]^{d},[0,1]^{d})} \le C_{k,d} \left(\frac{L_{1}}{L_{2}}\right)^{\beta_{d}k^{d+1}}.$$
(4.11)

Proof. Throughout this proof we use the notation and explicit construction of the KR map introduced in Appendix B: The i-th component of the KR map can then be expressed as

$$T_i(x_{[i]}) = G_{\rho,i}(T_1(x_1), \dots, T_{i-1}(x_{[i-1]}), F_{\pi,i}(x_{[i]})).$$
(4.12)

Here $F_{\rho,i}(x_{[i]}) = \int_0^{x_i} \rho_i(x_{[i-1]}, t_i) dt_i$ with $\rho_i = \frac{\hat{\rho}_i}{\hat{\rho}_{i-1}}$, where

$$\hat{\rho}_i(x_1, \dots, x_i) = \int_{[0,1]^{d-i}} \rho(x_1, \dots, x_d) dx_{i+1} \dots dx_d.$$
(4.13)

The function $F_{\pi,i}(x_{[i]})$ is defined analogous with ρ replaced by π . Finally, $x_i \mapsto G_{\rho,i}(x_{[i]})$ is defined as the inverse of $x_i \mapsto F_{\rho,i}(x_{[i]})$ on [0, 1], i.e.

$$G_{\rho,i}(x_{[i-1]}, F_{\rho,i}(x_{[i]})) = x_i.$$
(4.14)

The proof proceeds as follows: In step 1 we bound $||F_{\pi,i}||_{C^k([0,1]^i)}$, $||F_{\rho,i}||_{C^k([0,1]^i)}$, and in step 2 we bound $||G_{\rho,i}||_{C^k([0,1]^d)}$. In Step 3 we use induction over *i* to bound the norm of T_i in (4.12).

Step 1. Fix $i \in \{1, \dots, d\}$. In this step we show

$$\max\{\|F_{\pi,i}\|_{C^{k}([0,1]^{i})}, \|F_{\rho,i}\|_{C^{k}([0,1]^{i})}\} \le C\left(\frac{L_{1}}{L_{2}}\right)^{k+1}$$
(4.15)

for some constant C depending on d and k but independent of π and ρ . Since our assumptions on ρ and π are identical, it suffices to prove (4.15) for ρ .

Fix a multiindex $v \in \mathbb{N}_{0}^{d}$. If $v_{i} = 0$, then $D^{v}F_{\rho,i} = \int_{0}^{x_{i}} D^{v}\rho_{i}(x_{[i-1]}, t_{i})dt_{i}$. Otherwise with $v' = v - e_{i}$ (where $e_{i} = (\delta_{i,j})_{j=1}^{d}$) it holds $D^{v}F_{\rho,i} = D^{v'}\rho_{i}(x_{[i-1]}, x_{i})$. In either case $\|D^{v}F_{\rho,i}\|_{L^{\infty}([0,1]^{i})} \leq \|\rho_{i}\|_{C^{k}([0,1]^{i})}$. Thus $\|F_{\rho,i}\|_{C^{k}([0,1]^{i})} \leq \|\rho_{i}\|_{C^{k}([0,1]^{i})}$.

Similarly, Assumption 2.2 implies

$$\hat{\rho}_i(x_1, \dots, x_i) = \int_{[0,1]^{d-i}} \rho(x_1, \dots, x_d) dx_{i+1} \dots dx_d > \int_{[0,1]^{d-i}} L_2 dx_{i+1} \dots dx_d > L_2$$

and

$$D^{\boldsymbol{v}}\hat{\rho}_i(x_1,...,x_i) = \int_{[0,1]^{d-i}} D^{\boldsymbol{v}}\rho(x_1,...x_d) dx_{i+1}...dx_d \le \int_{[0,1]^{d-i}} L_1 dx_{i+1}...dx_d = L_1 dx_{i+1}...dx_d$$

for all $(x_1, ..., x_i) \in [0, 1]^i$ and multi-index $\boldsymbol{v} \in \mathbb{N}_0^d$ with $|\boldsymbol{v}| \leq k$. Thus we can conclude that $\|\hat{\rho}_i\|_{C^k([0,1]^i)} \leq L_1$ and Lemma C.10 gives $\|D^{n-j}(\frac{1}{\hat{\rho}_{i-1}})\|_{L^\infty([0,1]^{i-1})} \leq C_{n-j} \frac{L_1^{n-j}}{L_2^{n-j+1}}$ for some constant C_{n-j} that is independent of L_1, L_2 . By the Leibniz rule

$$D^n \rho_i = \sum_{j=0}^n \binom{n}{j} D^j \hat{\rho}_i D^{n-j} \left(\frac{1}{\hat{\rho}_{i-1}}\right)$$

and thus

$$\|D^n \rho_i\|_{L^{\infty}([0,1]^i)} \le CL_1 \frac{L_1^n}{L_2^{n+1}} = C \frac{L_1^{n+1}}{L_2^{n+1}} \qquad \forall n \in \{0, \dots, k\}$$

for some constant C that depends on k but is independent of L_1 , L_2 . In all this shows (4.15) for ρ .

Step 2. Fix $i \in \{1, \ldots, d\}$. In this step we show

$$\max\{\|G_{\pi,i}\|_{C^{k}([0,1]^{i})}, \|G_{\rho,i}\|_{C^{k}([0,1]^{i})}\} \le C\left(\frac{L_{1}}{L_{2}}\right)^{(k+1)(3k-2)}$$
(4.16)

for some constant C depending on d and k but independent of π and ρ . As before, by symmetry it suffices to provide the bound for π .

For $x_{[i]} \in [0,1]^i$ define

$$\tilde{F}_{\rho,i}(x_{[i]}) := (x_{[i-1]}, F_{\rho,i}(x_{[i]})) \in [0,1]^i.$$
(4.17)

Since $x_i \mapsto \tilde{F}_{\rho,i}(x_{[i]})$ is bijective from $[0,1] \to [0,1]$ for every fixed $x_{[i-1]} \in [0,1]^{i-1}$, the map $\tilde{F}_{\rho,i}$: $[0,1]^i \to [0,1]^i$ is a bijection. So is its inverse which we denote by $\tilde{G}_{\rho,i}: [0,1]^i \to [0,1]^i$. It holds for all $x_{[i]} \in [0,1]^i$ that

$$\tilde{G}_{\rho,i}(\tilde{F}_{\rho,i}(x_{[i]})) = x_{[i]}$$

Due to (4.14) and the definition of $\tilde{F}_{\rho,i}$ in (4.17), the *i*th component of $\tilde{G}_{\rho,i}$ is given by $G_{\rho,i}$ and thus

$$\|G_{\rho,i}\|_{C^k([0,1]^i,[0,1])} \le \|\tilde{G}_{\rho,i}\|_{C^k([0,1]^i,[0,1]^i)}.$$

In the following we wish to apply Prop. 4.4 to bound the right-hand side, which will yield a bound on the left-hand side. To this end we first derive a bound on the norm of $\tilde{F}_{\rho,i}$. By (4.17) and (4.15)

$$\|\tilde{F}_{\rho,i}\|_{C^{k}([0,1]^{i},[0,1]^{i})} \leq \|F_{\rho,i}\|_{C^{k}([0,1]^{i},[0,1])} + (i-1) \leq C \left(\frac{L_{1}}{L_{2}}\right)^{k+1}.$$
(4.18)

For the last inequality we used $\frac{L_1}{L_2} \ge 1$, so that $i-1 \le k-1$ can be absorbed into the k-dependent constant C. Next we bound $\|[D\tilde{F}_{\rho,i}]^{-1}\|_2$. It holds

$$\begin{pmatrix} 1 & \dots & 0 & 0 \\ \vdots & \ddots & & \\ 0 & & 1 & 0 \\ \partial_{x_1} F_{\rho,i} & \partial_{x_2} F_{\rho,i} & \dots & \partial_{x_i} F_{\rho,i} \end{pmatrix}$$

and

$$\begin{pmatrix} 1 & \dots & 0 & 0 \\ \vdots & \ddots & & \\ 0 & 1 & 0 \\ -\frac{\partial_{x_1}F_{\rho,i}}{\partial_{x_i}F_{\rho,i}} & -\frac{\partial_{x_2}F_{\rho,i}}{\partial_{x_i}F_{\rho,i}} & \dots & -\frac{1}{\partial_{x_i}F_{\rho,i}} \end{pmatrix}.$$

Since $\partial_{x_i} F_{\rho,i}(x_{[i]}) = \rho_i(x_{[i]}) \ge L_2$ and $\|\partial_{x_j} F_{\rho,i}(x_{[i]})\| \le L_1$ for all $x_{[i]} \in [0,1]^i$ and $j \in \{1,\ldots,i\}$, we have

$$\left\|\frac{\partial_{x_j}F_{\rho,i}}{\partial_{x_i}F_{\rho,i}}\right\|_{L^{\infty}([0,1]^i)} \le \frac{L_1}{L_2} \qquad \forall j \in \{1,\dots,i\}$$

Thus for all $x_{[i]} \in [0,1]^i$

$$\|[D\tilde{F}_{\rho,i}(x_{[i]})]^{-1}\|_{2} \le \|[D\tilde{F}_{\rho,i}(x_{[i]})]^{-1}\|_{F} \le \sqrt{(i-1) + i(\frac{L_{1}}{L_{2}})^{2}} \le \sqrt{2i\frac{L_{1}}{L_{2}}}$$

Since $\frac{L_1}{L_2} \ge 1$, we conclude that there exists a constant C depending on k, but independent of L_1 , L_2 such that

$$\max\{\|[D\tilde{F}_{\rho,i}]^{-1}\|_{2}, \|\tilde{F}_{\rho,i}\|_{C^{k}([0,1]^{i},[0,1]^{i})}\} \le C\frac{L_{1}^{k+1}}{L_{2}^{k+1}}.$$

Thus by Proposition 4.4

$$\|G_{\rho,i}\|_{C^{k}([0,1]^{i},[0,1])} \leq \|\tilde{G}_{\rho,i}\|_{C^{k}([0,1]^{i},[0,1]^{i})} \leq e^{k}k^{k^{2}} \left(C\frac{L_{1}^{k+1}}{L_{2}^{k+1}}\right)^{3k-2} \leq C\left(\frac{L_{1}}{L_{2}}\right)^{(k+1)(3k-2)}$$

This shows (4.16) for ρ .

Step 3. We finish the proof by showing that for all $i \in \{1, \ldots, d\}$

$$\|T_i\|_{C^k([0,1]^i,[0,1])} \le C\left(\frac{L_1}{L_2}\right)^{k^i(k+1) + (k+1)(3k-2)(\sum_{j=0}^{i-1} k^j)}.$$
(4.19)

For $x_{[i]} \in [0,1]^i$ define

$$\tilde{T}_i(x_{[i]}) := (T_1(x_1), \dots, T_{i-1}(x_{[i-1]}), F_{\pi,i}(x_{[i]})) \in [0,1]^i.$$
(4.20)

By (4.12) it holds $T_i = G_{\rho,i} \circ \tilde{T}_i$, and thus Corollary 4.3 implies

$$||T_i||_{C^k([0,1]^i)} \le C ||G_{\rho,i}||_{C^k([0,1]^i)} \max\{1, ||\tilde{T}_i||_{C^k([0,1]^i)}\}^k$$
(4.21)

for a k-dependent constant C.

We proceed by induction over i, and start with i = 1. In this case (4.15), (4.16) and (4.21) yield

$$\|T_1\|_{C^k([0,1])} = \|G_{\rho,1} \circ F_{\pi,1}\|_{C^k([0,1])} \le C\left(\frac{L_1}{L_2}\right)^{(k+1)(3k-2)+k(k+1)},$$

and thus (4.19) is satisfied. For the induction step assume the statement is true for $i - 1 \ge 1$. By (4.15), (4.20) and the induction hypothesis

$$\begin{split} \|\tilde{T}_{i}\|_{C^{k}([0,1]^{i})} &\leq C \left(\sum_{j=1}^{i-1} \|T_{j}\|_{C^{k}([0,1]^{j})} + \|F_{\pi,i}\|_{C^{k}([0,1]^{i})} \right) \\ &\leq C \left(\frac{L_{1}}{L_{2}} \right)^{k^{i-1}(k+1) + (k+1)(3k-2)(\sum_{j=0}^{i-2} k^{j})}, \end{split}$$

where again C may depend on k (or $i \le k$) but not on L_1, L_2 . Then (4.16) and (4.21) imply

$$\begin{split} \|T_i\|_{C^k([0,1]^i)} &\leq C \|G_{\rho,i}\|_{C^k([0,1]^i)} \max\{1, \|\tilde{T}_i\|_{C^k([0,1]^i)}\}^k \\ &\leq C \Big(\frac{L_1}{L_2}\Big)^{(k+1)(3k-2)} \Big(\frac{L_1}{L_2}\Big)^{k(k^{i-1}(k+1)+(k+1)(3k-2)(\sum_{j=0}^{i-2}k^j))} \\ &= C \left(\frac{L_1}{L_2}\right)^{k^i(k+1)+(k+1)(3k-2)(\sum_{j=0}^{i-1}k^j)}. \end{split}$$

Finally, by putting all the estimates together, we obtain the following upper bound for the C^k norm of velocity field.

Theorem 4.12. Let $\Omega_0 = \Omega_1 = [0,1]^d$ and π, ρ satisfy Assumption 2.2 with constant $k \ge 2, 0 < L_1 \le L_2 < \infty$. Let $T : [0,1]^d \to [0,1]^d$ be the KR map pushing forward π to ρ and $f : [0,1]^d \times [0,1] \to [0,1]^d$ be the velocity field in (2.7) that corresponds to the displacement interpolation between x and T(x). Then, there exists constants $C_{k,d}$ that only depends on k, d and β_d that only depends on d, such that the following holds:

$$\|f\|_{C^k([0,1]^d \times [0,1])} \le C_{k,d} \left(\frac{L_1}{L_2}\right)^{\beta_d k^{d+3}}$$

Proof. The proof of the theorem requires a combination of Theorem 4.5, Lemma.4.10 and Theorem 4.11.

First, by Lemma.4.10 and Theorem 4.11, there exists constant C_d and β'_d such that $||T||_{C^1([0,1]^d,[0,1])} \leq C_d(\frac{L_1}{L_2})^{\beta'_d}$ and

$$c_T \leq (\frac{L_1}{L_2})^{2d} \max\{1, (C_d(\frac{L_1}{L_2})^{\beta'_d})^{d-1}\} \leq \max\{(\frac{L_1}{L_2})^{2d}, C_d^{d-1}(\frac{L_1}{L_2})^{\beta'_d(d-1)}\}.$$

By renaming $C_d = \max\{1, C_d^{d-1}\}$ and $\beta'_d = \max\{2d, \beta'_d(d-1)\}$, we can simplify the above expression as $c_T \leq C_d(\frac{L_1}{L_2})^{\beta'_d}$. Note it holds true that $c_T \geq 1$.

By lemma.4.10 and Theorem 4.11, there exists constants $C_{k,d}$ and β_d and

$$\gamma = \max\{2, 1 + c_T\}(1 + \|T\|_{C^k} + \sup_{x \in \Omega_0} \|x\|) \le (1 + c_T)(1 + C_{k,d}(\frac{L_1}{L_2})^{\beta_d k^{d+1}} + \sqrt{d})$$

$$\le 2c_T(1 + C_{k,d}(\frac{L_1}{L_2})^{\beta_d k^{d+1}} + \sqrt{d}) \le 2c_T C_{k,d}(\frac{L_1}{L_2})^{\beta_d k^{d+1}}$$

$$\le C_d(\frac{L_1}{L_2})^{\beta'_d} C_{k,d}(\frac{L_1}{L_2})^{\beta_d k^{d+1}} \le C_{k,d}(\frac{L_1}{L_2})^{\beta_d k^{d+1}}.$$

where we absorb constants whenever possible. In particular, we used $\frac{L_1}{L_2} > 1$ to absorb into $C_{k,d}$ in the third inequality and we used k > 1 to absorb everything in the exponent into a d-dependent β_d in the last inequality.

Finally, applying Theorem 4.5, we obtain

$$\|f\|_{C^{k}([0,1]^{d}\times[0,1])} \leq 2k^{k^{3}+k}e^{k^{2}+k}\gamma^{3k^{2}-2k+1} \leq C_{k,d}(\frac{L_{1}}{L_{2}})^{\beta_{d}k^{d+1}3k^{2}} \leq C_{k,d}(\frac{L_{1}}{L_{2}})^{\beta_{d}k^{d+3}}.$$

5 Stability in the velocity field

In the previous sections we showed existence of velocity fields f that yield flow maps realizing a (triangular) transport that pushes forward π to ρ . In practice, a suitable velocity field g is obtained by minimizing the objective (2.4) over some parametrized function class such as the set of Neural Networks of a certain architecture. In general, such g will only approximate f, and it is therefore of interest to understand how errors in the approximation of f propagate to errors in the distributions realized by the corresponding flow map. This is the purpose of the present section.

5.1 Wasserstein distance

First, we present results when the divergence between probability distributions is measured by Wasserstein distance. That is, we take $\mathcal{D} = W_p$ in (2.4), where W_p is the *p*-Wasserstein distance.

Theorem 5.1. Let $f, g \in \mathcal{V}$ (cp. (2.7)) and $||f - g||_{C^0(\mathbb{R}^d \times [0,1])} < \infty$. Assume that L > 0 is such that $x \mapsto f(x,t)$ has Lipschitz constant L for all $t \in [0,1]$. Let $X_f, X_g : \mathbb{R}^d \times [0,1] \to \mathbb{R}^d$ be as in (2.1). Then

$$\|X_f(\cdot, 1) - X_g(\cdot, 1)\|_{C^0(\mathbb{R}^d)} \le \|f - g\|_{C^0(\mathbb{R}^d \times [0, 1])} e^L.$$
(5.1)

The idea of the proof of Theorem 5.1 is to apply Grönwall's inequality to the evolution of the error $|X_f(x,t) - X_g(x,t)|$ over time. We point out that this stability result merely requires g to approximate f uniformly, however f is additionally assumed to be Lipschitz continuous in space.

Proof of Theorem 5.1. Fix $x \in \mathbb{R}^d$. Then for all $s \in [0, 1]$

$$\begin{aligned} |f(X_f(x,s),s) - g(X_g(x,s),s)| &\leq |f(X_f(x,s),s) - f(X_g(x,s),s)| + |f(X_g(x,s),s) - g(X_g(x,s),s)| \\ &\leq L|X_f(x,s) - X_g(x,s)| + ||f - g||_{C^0(\mathbb{R}^d \times \mathbb{R})}, \end{aligned}$$

where we used the spatial Lipschitz continuity of f. Hence for $t \in [0, 1]$

$$|X_f(x,t) - X_g(x,t)| = \left| \int_0^t f(X_f(x,s),s) - g(X_g(x,s),s)ds \right|$$

$$\leq \int_0^t L|X_f(x,s) - X_g(x,s)|ds + t||f - g||_{C^0(\mathbb{R}^d \times \mathbb{R})}$$

Using Grönwall's inequality, we get $|X_f(x,1) - X_g(x,1)| \le ||f - g||_{C^0(\mathbb{R}^d \times \mathbb{R})} e^L$ as claimed.

Next we show how an approximation of the velocity field affects the difference in distributions in terms of the Wasserstein distance W_p . In the following corollary, we denote by $|\text{supp}(\pi)|$ the Lebesgue measure of the support of π .

Corollary 5.2. Let $f, g \in \mathcal{V}$ and X_f, X_g be as in Theorem 5.1. Let π be a probability distribution on \mathbb{R}^d . Then for any $p \in [1, \infty)$

$$W_p(X_f(\cdot,1)_{\sharp}\pi, X_g(\cdot,1)_{\sharp}\pi) \le \|f-g\|_{C^0(\mathbb{R}^d \times \mathbb{R})} e^L |\operatorname{supp}(\pi)|^{1/p}.$$

Moreover, for $p = \infty$ holds $W_{\infty}(X_f(\cdot, 1)_{\sharp}\pi, X_g(\cdot, 1)_{\sharp}\pi) \le ||f - g||_{C^0(\mathbb{R}^d \times \mathbb{R})} e^L$.

Proof. Let $F : \mathbb{R}^d \to \mathbb{R}^d \times \mathbb{R}^d$ via $F(x) := (X_f(x, 1), X_g(x, 1))$. Observe that $\gamma := F_{\sharp}(\pi \otimes \pi)$ is then a probability distribution on $\mathbb{R}^d \times \mathbb{R}^d$ with marginals $X_f(\cdot, 1)_{\sharp}\pi$ and $X_g(\cdot, 1)_{\sharp}\pi$, i.e. it is a coupling of these measures. If $p < \infty$ then by definition of the Wasserstein distance

$$W_p(X_f(\cdot,1)_{\sharp}\pi, X_g(\cdot,1)_{\sharp}\pi)^p \leq \int_{\mathbb{R}^d \times \mathbb{R}^d} \|x-y\|^p \, \mathrm{d}\gamma(x,y)$$
$$= \int_{\mathbb{R}^d} \|X_f(x,1) - X_g(x,1)\|^p \, \mathrm{d}\pi(x)$$
$$\leq |\mathrm{supp}(\pi)|(\|f-g\|_{C^0(\mathbb{R}^d \times \mathbb{R})} e^L)^p.$$

The case $p = \infty$ is obtained with the usual adjustment of arguments.

5.2 KL-divergence

In this subsection, we measure the distance in the KL-divergence, i.e. $\mathcal{D} = \mathcal{D}_{KL}$ in (2.4). Unlike for the Wasserstein distance, for $\mathcal{D}_{KL}(X(\cdot, 1)_{\sharp}\pi, \rho)$ to be finite, we need in particular $X(\cdot, 1)_{\sharp}\pi \ll \rho$. We restrict ourselves to distributions on cubes, and consider $\Omega_0 = \Omega_1 = [0, 1]^d$.

Most of the work regarding the approximation distributions in KL-divergence using ODE flow maps has already been studied in our companion paper Marzouk et al. (2024); we include some of the relevant results here for the sake of completeness. In Marzouk et al. (2024), an ansatz space

$$C_{\text{ansatz}}^{k}(r) = \{ f = (f_1, ..., f_d)^T : f_j = \tilde{f}_j x_j (1 - x_j), \tilde{f}_j \in C^k([0, 1]^d \times [0, 1], [0, 1]^d) \}$$

$$\cap \{ f \in C^2([0, 1]^d \times [0, 1], [0, 1]^d) : \| f \|_{C^2([0, 1]^d \times [0, 1], [0, 1]^d)} \leq r \}$$

was proposed. Its definition ensures that all push-forward distributions $X_f(\cdot, 1)_{\sharp}\pi$ are supported on $[0, 1]^d$ for any $f \in C^k_{\text{ansatz}}(r)$. In Theorem 4.12, we showed the velocity field corresponding to the straightline interpolation of Knothe-Rosenblatt maps f^{Δ} lies in $C^k([0, 1]^d \times [0, 1], [0, 1]^d)$ and in Theorem 9 of Marzouk et al. (2024), it is shown that

$$\frac{f_j^{\Delta}(x_1,\cdots,x_j)}{x_j(1-x_j)} \in C^k([0,1]^d \times [0,1], [0,1]).$$

Therefore, by choosing r to be large enough, for example, by taking $r = C_{k,d} (\frac{L_1}{L_2})^{\beta_d k^{d+3}}$ corresponding to the upper bound in Theorem 4.12, it suffices to consider an approximating element in $C_{\text{ansatz}}^k(r)$.

We emphasize that bounding discrepancies in Wasserstein distance only requires C^0 control of the velocity fields; however, controlling discrepancies in the KL-divergence, requires C^1 control of the velocity fields, as stated in the next theorem.

Theorem 5.3. Let π , ρ satisfy Assumptions 2.1 and 2.2 with $\Omega_0 = \Omega_1 = [0, 1]^d$. Let f^{Δ} as in Theorem 4.12 and $g \in C^2_{ansatz}(r)$. Then

$$\mathcal{D}_{KL}(X_g(\cdot, 1)_{\sharp}\pi, X_{f^{\Delta}}(\cdot, 1)_{\sharp}\pi) \le C \|f^{\Delta} - g\|_{C^1([0,1]^d)}^2$$

for some constant C that depends on L_1, L_2, d .

Proof. By Lemma 6, Theorem 7 and Theorem 8 of Marzouk et al. (2024) there exists a constant $C_{L_1,L_2,d}$ that depends on L_1, L_2, d such that

$$\|X_g(\cdot,1)_{\sharp}\pi - X_{f^{\Delta}}(\cdot,1)_{\sharp}\pi\|_{C^0([0,1]^d)} = \|X_g(\cdot,1)_{\sharp}\pi - \rho\|_{C^0([0,1]^d)} \le C_{L_1,L_2,d}\|f^{\Delta} - g\|_{C^1([0,1]^d)}$$

To get an upper bound for $\mathcal{D}_{KL}(X_q(\cdot, 1)_{\sharp}\pi, X_{f^{\Delta}}(\cdot, 1)_{\sharp}\pi)$, we bound

$$\begin{aligned} \mathcal{D}_{KL}(X_g(\cdot,1)_{\sharp}\pi, X_{f^{\Delta}}(\cdot,1)_{\sharp}\pi) &= \mathbb{E}_{\pi_g(\cdot,1)} \Big[\log \frac{\pi_g(\cdot,1)}{\rho(x)} \Big] \le \log \mathbb{E}_{\pi_g(x,1)} [\frac{\pi_g(x,1)}{\rho(x)}] = \log \int_{[0,1]^d} \frac{\pi_g(x,1)^2}{\rho(x)} dx \\ &= \log \left(\int_{[0,1]^d} \frac{(\rho(x) - \pi_g(x,1))^2}{\rho(x)} dx + 1 \right) \le \log \left(\frac{C_{L_1,L_2,d}^2}{L_2} \| f^{\Delta} - g \|_{C^1([0,1]^d)}^2 + 1 \right). \end{aligned}$$
The fact that $\log(1+x) \le x$ for all $x \ge 0$ gives the result.

The fact that $\log(1+x) \le x$ for all $x \ge 0$ gives the result.

6 Neural network approximation

In Section 4.2 we studied the regularity of the velocity field corresponding to straight-line trajectories realizing the Knothe-Rosenblatt map at time t = 1. Building on earlier works on neural network approximation theory such as Yarotsky (2017); Yarotsky and Zhevnerchuk (2020); Marzouk et al. (2024) in the present section we conclude that by parameterizing the velocity field via neural networks, NODE flows can achieve arbitrary accuracy in terms of the Wasserstein distance and KL-divergence. Furthermore, given a desired accuracy $\varepsilon > 0$, we give upper bounds on the required network depth, width, and size in terms of ε . Since the objective functional J contains first-order derivatives, we shall consider the approximation theory using $ReLU^2$ networks developed in Marzouk et al. (2024).

We first recall the definition of $ReLU^2$ networks.

Definition 6.1. Denote $\sigma_2(x) := \max\{0, x\}^2$ and let $d_1, d_2 \ge 1$. Then, the class of ReLU² networks mapping from $[0, 1]^{d_1}$ to \mathbb{R}^{d_2} , with depth L, width W, sparsity S, and bound B on the network weights, is denoted

$$\begin{split} \Phi^{d_1,d_2}(L,W,S,B) &= \Big\{ \big(W^{(L)}\sigma_2(\cdot) + b^{(L)} \big) \circ \dots \circ \big(W^{(1)}\sigma_2(\cdot) + b^{(1)} \big) : W^{(L)} \in \mathbb{R}^{1 \times W}, \\ b^{(L)} \in \mathbb{R}^{d_2}, W^{(1)} \in \mathbb{R}^{W \times d_1}, b^{(1)} \in \mathbb{R}^W, W^{(l)} \in \mathbb{R}^{W \times W}, b^{(l)} \in \mathbb{R}^W (1 < l < L), \\ \sum_{l=1}^L \big(\|W^{(l)}\|_0 + \|b^{(l)}\|_0 \big) \le S, \max_{1 \le l \le L} \big(\|W^{(l)}\|_{\infty,\infty} \vee \|b^{(l)}\|_{\infty} \big) \le B \Big\}. \end{split}$$

Remark 6.2. Since the representation of a function via neural networks is not unique in general, the statement "g is a ReLU² NN of depth L, width W, sparsity S, norm bound B" merely implies the existence of such a network satisfying the above properties. Possibly, for some other \tilde{L} , \tilde{W} , \tilde{S} and \tilde{B} , it may additionally hold that "g is a ReLU² NN of depth \tilde{L} , width \tilde{W} , sparsity \tilde{S} , and norm bound \tilde{B} ".

6.1 Wasserstein distance

As noted before, approximation in Wasserstein distance only requires C^0 , rather than C^1 , control of the velocity fields. In terms of the Wasserstein distance, we have the following result, which is a consequence of our regularity analysis of the velocity field in Theorem 4.12, Corollary 5.2 and neural network approximation results as e.g. presented in Marzouk et al. (2024):

Proposition 6.3. Let $k \ge 1$, $p \in [1, \infty]$, and let ρ , π be two probability distributions on $[0, 1]^d$ with Lebesgue densities in $C^k([0, 1]^d)$ satisfying Assumption 2.2. Then there exist constants $C_{d,k}$ and C'_{d,k,L_1,L_2} such that for every $\varepsilon \in (0, 1]$ there exists a ReLU² neural network $g \in \Phi^{d+1,d}(L, W, S, B)$ with

$$L \le C_{d,k}, \quad W \le C'_{d,k,L_1,L_2} \varepsilon^{-\frac{d+1}{k}}, \quad S \le C'_{d,k,L_1,L_2} \varepsilon^{-\frac{d+1}{k}}, \quad B \le C'_{d,k,L_1,L_2} \varepsilon^{-\frac{1}{k}}$$

such that for another constant C_{d,k,L_1,L_2} , we have

$$W_p(\rho, X_g(\cdot, 1)_{\sharp}\pi) \le C\varepsilon.$$

Proof. According to Theorem 4.12 there exists $f^{\Delta} \in C^k([0,1]^d \times [0,1])$ such that $X_{f^{\Delta}}(\cdot,1)_{\sharp}\pi = \rho$ and $\|f^{\Delta}\|_{C^k([0,1]^d \times [0,1])}$ only depends on L_1, L_2, k, d . We can extend f^{Δ} to some $\tilde{f^{\Delta}} \in C^k(\mathbb{R}^d \times [0,1])$ with compact support and such that

$$\|f^{\Delta}\|_{C^{k}(\mathbb{R}^{d}\times[0,1])} \leq C\|f^{\Delta}\|_{C^{k}(\Omega_{[0,1]})}$$

for some C solely depending on d (see for example Step 1 of the proof of (Marzouk et al., 2024, Theorem 33)). Since \tilde{f}^{Δ} has compact support and belongs to $C^1(\mathbb{R}^d \times [0,1])$, there exists $L < \infty$ such that $x \mapsto \tilde{f}^{\Delta}(x,t) : \mathbb{R}^d \to \mathbb{R}^d$ has Lipschitz constant L for all $t \in [0,1]$. Again, L solely depends on $\|f^{\Delta}\|_{C^1([0,1]^d \times [0,1])}$ and thus on L_1, L_2, k, d . Next, let $M = 1 + \exp(L)$.

According to Marzouk et al. (2024, Theorem 16), there exists a ReLU² neural network g satisfying the bounds (6.3) and

$$\|f^{\Delta}(x) - g(x)\|_{C^{0}([-M,M]^{d} \times [0,1])} \le \varepsilon.$$
(6.1)

Fix $x \in [0, 1]^d$. By (5.1) and (6.1), we have for all $t \in [0, 1]$

$$\|X_{\tilde{f}^{\Delta}}(x,t) - X_g(x,t)\| \le \varepsilon \exp(L) \le \exp(L)$$
(6.2)

and thus $||X_g(x,t)|| \leq 1 + \exp(L) = M$. Hence $X_g(x,1) \in \mathbb{R}^d$ is well-defined since the trajectory $t \mapsto X_g(x,t), t \in [0,1]$, remains within $[-M,M]^d \times [0,1]$. Finally, the first inequality in (6.2) and an application of Corollary 5.2 concludes the proof.

Next we discuss convergence of the objective J defined in (2.4), for the Wasserstein distance; specifically

$$J_{\mathrm{W}}(f) := W_2(\rho, X_f(\cdot, 1)_{\sharp}\pi)^2 + R(f)$$

where

$$R(f) = \int_0^1 \|\nabla_X f(X_f(x,t)) f(X_f(x,t),t) + \partial_t f(X_f(x,t),t)\|^2 \, \mathrm{d}t.$$
(6.3)

Since the regularization term R(f) contains first order derivatives of f, it will not suffice to have uniform approximation in f, but rather we'll additionally need uniform approximation of the derivatives of f. We also point out that by Theorem 3.12, $\inf_{f \in V} J_W(f) = 0$.

Theorem 6.4. Let $k \ge 2$, $p \in [1, \infty]$ and let ρ , π be two probability distributions on $[0, 1]^d$ with Lebesgue densities in $C^k([0, 1]^d)$ satisfying Assumption 2.2.

Then there exist constants $C_{d,k}$ and C'_{d,k,L_1,L_2} such that for every $\varepsilon \in (0,1]$ there exists a ReLU² neural network $g \in \Phi^{d+1,d}(L,W,S,B)$ with

$$L \le C_{d,k}, \quad W \le C'_{d,k,L_1,L_2} \varepsilon^{-\frac{d+1}{k-1}}, \quad S \le C'_{d,k,L_1,L_2} \varepsilon^{-\frac{d+1}{k-1}}, \quad B \le C'_{d,k,L_1,L_2} \varepsilon^{-\frac{1}{k-1}}$$
(6.4)

and such that for another constant C_{d,k,L_1,L_2} , we have

$$J_{\mathrm{W}}(g) \leq C_{d,k,L_1,L_2} \varepsilon^2.$$

Proof. According to Theorem 4.12, there exists $f^{\Delta} \in C^k([0,1]^d \times [0,1])$ such that $X_{f^{\Delta}}(\cdot,1)_{\sharp}\pi = \rho$ (i.e. $W_2(\rho, X_{f^{\Delta}}(\cdot,1)_{\sharp}\pi) = 0$), f^{Δ} realizes straight line trajectories (i.e. $R(f^{\Delta}) = 0$) and $||f^{\Delta}||_{C^k([0,1]^d \times [0,1])}$ only depends on L_1 , L_2 , k, d. In particular f^{Δ} can be extended to an element of \mathcal{V} in (2.7) such that $J_{\mathrm{W}}(f^{\Delta}) = 0$. Hence it suffices to show existence of g such that $J_{\mathrm{W}}(g) \leq C\varepsilon^2$.

We can extend f^{Δ} to some $\tilde{f^{\Delta}} \in C^k(\mathbb{R}^d \times [0,1])$ with compact support and such that

$$\|f^{\Delta}\|_{C^{k}(\mathbb{R}^{d}\times[0,1])} \leq C\|f^{\Delta}\|_{C^{k}([0,1]^{d}\times[0,1])}$$

for some C solely depending on d (see for example Step 1 of the proof of (Marzouk et al., 2024, Theorem 33)) Since \tilde{f}^{Δ} has compact support and belongs to C^k , $k \ge 2$, there exists $L < \infty$ such that the three maps

$$x \mapsto f^{\Delta}(x,t), \qquad x \mapsto \partial_t f^{\Delta}(x,t), \qquad x \mapsto \nabla_x f^{\Delta}(x,t),$$

have Lipschitz constant L for all $t \in [0, 1]$ and all $x \in \mathbb{R}^d$. Again, L solely depends on $||f^{\Delta}||_{C^k([0,1]^d \times [0,1])}$ and thus on L_1, L_2, k, d .

Next, let $M > \exp(L)$ be so large that $[0,1]^d \subseteq [-M + \exp(L), M - \exp(L)]^d$. According to (Marzouk et al., 2024, Theorem 16), there exists a ReLU² neural network g satisfying the bounds (6.4) and

$$\|\tilde{f}^{\Delta}(x) - g(x)\|_{C^{1}([-M,M]^{d} \times [0,1])} \le \varepsilon.$$
(6.5)

As in the proof of Proposition 6.3, we conclude that $X_q(x,1) \in \mathbb{R}^d$ is well-defined and

$$\|X_{\tilde{f}^{\Delta}}(x,t) - X_g(x,t)\| \le \varepsilon \exp(L) \qquad \forall x \in [0,1]^d, \ t \in [0,1].$$

Since the trajectories $t \mapsto X_{\tilde{f}^{\Delta}}(x,t)$ remain in $[0,1]^d \times [0,1]$ according to Proposition 4.9, we conclude that

$$X_g(x,t) \in [-M,M]^d \qquad \forall x \in [0,1]^d, \ t \in [0,1].$$
 (6.6)

Moreover, as in the proof of Proposition 6.3 holds

$$W_2(\rho, X_g(\cdot, 1)_{\sharp}\pi)^2 \le C\varepsilon^2.$$

To bound $J_W(g)$, it remains to treat the term R(g) in (6.3). We have

$$\begin{aligned} \|\nabla_X \tilde{f}^{\Delta}(X_{\tilde{f}^{\Delta}}(x,t)) - \nabla_X g(X_g(x,t))\| \leq & \|\nabla_X \tilde{f}^{\Delta}(X_{\tilde{f}^{\Delta}}(x,t)) - \nabla_X \tilde{f}^{\Delta}(X_g(x,t))\| \\ &+ \|\nabla_X \tilde{f}^{\Delta}(X_g(x,t)) - \nabla_X g(X_g(x,t))\|. \end{aligned}$$

Since $\nabla_X \tilde{f^{\Delta}}$ has Lipschitz constant L, The first term on the right-hand side is bounded by $L \exp(L)\varepsilon$. Due to $X_g(x,t) \in [0,1]^d \subseteq [-M,M]^d$ by (6.6), we can use (6.5) to bound the second term which gives

$$\|\nabla_X \tilde{f^{\Delta}}(X_{\tilde{f}^{\Delta}}(x,t)) - \nabla_X g(X_g(x,t))\| \le L \exp(L)\varepsilon + \varepsilon \qquad \forall x \in [0,1]^d, \ t \in [0,1]^d$$

Similarly we obtain

$$\|\partial_t f^{\Delta}(X_{\tilde{f}^{\Delta}}(x,t)) - \partial_t g(X_g(x,t))\| \le L \exp(L)\varepsilon + \varepsilon \qquad \forall x \in [0,1]^d, \ t \in [0,1]$$

and

$$\|f^{\Delta}(X_{\tilde{f}^{\Delta}}(x,t)) - g(X_g(x,t))\| \le L \exp(L)\varepsilon + \varepsilon \qquad \forall x \in [0,1]^d, \ t \in [0,1].$$

Therefore

$$\begin{split} R(g) &= |R(f^{\Delta}) - R(g)| \leq \int_0^1 \|\nabla_X \tilde{f}(X_{\tilde{f}^{\Delta}}(x,t)) \tilde{f^{\Delta}}(X_{\tilde{f}^{\Delta}}(x,t),t) + \partial_t \tilde{f}(X_{\tilde{f}^{\Delta}}(x,t),t) \\ &\quad - \nabla_X g(X_g(x,t)) g(X_g(x,t),t) - \partial_t g(X_g(x,t),t) \|^2 \, \mathrm{d}t \\ &\leq C \int_0^1 \varepsilon^2 \, \mathrm{d}t, \end{split}$$

where C only depends on L.

6.2 KL-divergence

Now we discuss convergence of the objective J defined in (2.4) for case where \mathcal{D} is the KL divergence, i.e.,

$$J_{\mathrm{KL}}(f) := \mathrm{KL}(X_f(\cdot, 1)_{\sharp}\pi || \rho) + R(f).$$

Note again that by Theorem 3.12, we have $\inf_{f \in \mathcal{V}} J_{KL}(f) = 0$.

For the KL-divergence to be well-defined, we need to enforce the constraint that the ODE flow map is a diffeomorphism onto $[0, 1]^d$. Therefore, we shall use the neural network based ansatz space introduced in Marzouk et al. (2024). To this end define $\chi_d(x_1, \ldots x_d) : D \to D$ via

$$\chi_d(x_1, \dots, x_d) = [x_1(1-x_1), \dots, x_d(1-x_d)]^T.$$

Letting \otimes be the coordinate-wise (Hadamard) multiplication of two vectors, for any velocity field $f : [0,1]^d \times [0,1] \to \mathbb{R}^d$, $f \otimes \chi_d$ then yields a vector field with vanishing normal components at the boundary of $[0,1]^d$ at any time $t \in [0,1]$. We define our neural network ansatz space as

$$\mathcal{F}_{NN}(L, W, S, B, r) := \{f^{NN}(x_1, \dots, x_d, t) \otimes \chi_d(x_1, \dots, x_d) : f^{NN} \in \Phi^{d+1, d}(L, W, S, B), \|f\|_{W^{2,\infty}(\Omega)} \le r\}.$$
(6.7)

Proposition 6.5. Let $k \ge 2$ and ρ, π two probability distributions satisfying Assumptions 2.1 and 2.2 with $\Omega_0 = \Omega_1 = [0, 1]^d$.

Then there exists constants $C_{d,k}$ and C'_{d,k,L_1,L_2} such that for every $\varepsilon \in (0,1]$, there exists a $ReLU^2$ network g in the ansatz space $\mathcal{F}_{NN}(L,W,S,B,r)$ with parameters satisfying

$$L \le C_{d,k}, \quad W \le C'_{d,k,L_1,L_2} \varepsilon^{-\frac{d+1}{k-1}}, \quad S \le C'_{d,k,L_1,L_2} \varepsilon^{-\frac{d+1}{k-1}}, \quad B \le C'_{d,k,L_1,L_2} \varepsilon^{-\frac{1}{k-1}}, \quad r \le C'_{d,k,L_1,L_2} \varepsilon^{-\frac{d+1}{k-1}},$$

such that for another constant C_{d,k,L_1,L_2} , we have

$$\operatorname{KL}(X_g(\cdot, 1)_{\sharp}\pi, \rho) \le C'_{d,k,L_1,L_2}\varepsilon^2.$$

Proof. According to Theorem 4.12, there exists $f^{\Delta} \in C^k([0,1]^d \times [0,1])$ such that $X_{f^{\Delta}}(\cdot,1)_{\sharp}\pi = \rho$ (i.e. $\mathcal{D}_{KL}(\rho, X_{f^{\Delta}}(\cdot,1)_{\sharp}\pi) = 0$), f^{Δ} realizes straight line trajectories (i.e. $R(f^{\Delta}) = 0$) and $||f^{\Delta}||_{C^k([0,1]^d \times [0,1])}$ only depends on L_1, L_2, k, d . By Step 1 of (Marzouk et al., 2024, Theorem 20), for all $N \geq 1$, there exists $g \in \mathcal{F}_{NN}(L, W, S, B, r)$ with

$$L \le C_{d,k}, \quad W \le N, \quad S \le N, \quad B \le C_{d,k} \| f^{\Delta} \|_{C^k([0,1]^d \times [0,1])} + N^{\frac{1}{d+1}}, \quad r \le C_{d,k,L_1,L_2}$$

such that $||f^{\Delta} - g||_{C^{1}([0,1]^{d} \times [0,1])} \leq C_{d,k,L_{1},L_{2}}N^{-\frac{k-1}{d+1}}$, where $C_{d,k}$ and $\tilde{C}_{d,k,L_{1},L_{2}}$ are constants depending on d, k and d, k, L_{1}, L_{2} respectively. Letting $\tilde{C}_{d,k,L_{1},L_{2}}N^{-\frac{k-1}{d+1}} = \varepsilon$, we solve for N to get $N = C_{d,k,L_{1},L_{2}}\varepsilon^{-\frac{d+1}{k-1}}$ for some $C_{d,k,L_{1},L_{2}}$. By Theorem 5.3, we then have $\mathcal{D}_{KL}(X_{g}(\cdot,1),\rho) \leq C_{d,k,L_{1},L_{2}}\varepsilon^{2}$.

Theorem 6.6. Let $k \ge 2$ and ρ, π two probability distributions satisfying Assumptions 2.1 and 2.2 with $\Omega_0 = \Omega_1 = [0, 1]^d$.

Then there exist constants $C_{d,k}$ and C'_{d,k,L_1,L_2} such that for every $\varepsilon \in (0,1]$, there exists a ReLU² network g in the ansatz space $\mathcal{F}_{NN}(L,W,S,B,r)$ with parameters satisfying

$$L \le C_{d,k}, \quad W \le C'_{d,k,L_1,L_2} \varepsilon^{-\frac{d+1}{k-1}}, \quad S \le C'_{d,k,L_1,L_2} \varepsilon^{-\frac{d+1}{k-1}}, \quad B \le C'_{d,k,L_1,L_2} \varepsilon^{-\frac{1}{k-1}}, \quad r \le C'_{d,k,L_1,L_2} \varepsilon^{-\frac{1}{k-1}},$$

such that for another constant C_{d,k,L_1,L_2} , we have

$$J(g) \le C_{d,k,L_1,L_2} \varepsilon^2.$$

Proof. The proof follows by the same arguments as the proof of Theorem 6.4. The error in the KL-divergence part of the objective follows from Proposition 6.5 and the error in minimal-energy regularization has already been treated in Theorem 6.4.

Remark 6.7. We comment here that the estimates we obtained above are L^{∞} in nature, thus we are able to derive distribution error estimates in other metrics as well, such as Hellinger, chi-square and total variation.

7 Discussion and future work

Our work is a crucial first step towards establishing a theoretical framework for sampling and distribution learning through ODE flow maps. In particular, the approximation results in this work can be viewed as quantifying the *bias* term in the classical context of statistical learning theory. In our parallel work Marzouk et al. (2024), we explore the *variance* term by analyzing the statistical complexity of the function class represented by ODE flow maps whose velocity fields come from a bounded neural network class.

While our work establishes a theoretical framework for analyzing ODE-based models, several important questions still remain open. First, we only consider distributions supported on bounded domains because the Lipschitz constant of straight-line ansatz considered in this work and Marzouk et al. (2024) can be uncontrollable when the distributions are not lower bounded. It will be interesting to see how our theories could be extended to the case of unbounded domains. Second, the approximation errors obtained in this work depend on d + 1 and k - 1, because we considered the velocity field as a generic function on \mathbb{R}^{d+1} and the approximation error metric considered is in C^1 for distributional stability results. This approximation error leads to nonparametric convergence rate of $n^{-\frac{2(k-1)}{d+1+2(k-1)}}$ in Marzouk et al. (2024). We note this statistical rate obtained is suboptimal, compared to the minimax optimal rate of learning k-smooth densities on d-dimensional domains (which is $n^{-\frac{2k}{d+2k}}$). Therefore, an important question to ask is whether neural-ODE based models can achieve the same minimax optimal statistical rates, as classical density estimators such as wavelets or kernel-based methods. We leave these open questions to future studies.

Acknowledgments and disclosure of funding

ZR and YM acknowledge support from the US Air Force Office of Scientific Research (AFOSR) MURI, Analysis and Synthesis of Rare Events, award number FA9550-20-1-0397, and from the US Department of Energy (DOE), Office of Advanced Scientific Computing Research, under grants DE-SC0021226 and DE-SC0023187. ZR also acknowledges support from a US National Science Foundation Graduate Research Fellowship.

Comments on training A

Training algorithm A.1

Here we present pseudocode for a notional training algorithm, i.e., an algorithm to learn the velocity field fof a neural ODE in the setting of problem P1 (see Section 2.2), when the information divergence used is the KL-divergence.

Recall the training objective we consider consists of two parts: one is the KL divergence from the source distribution to the pushforward of the target by the time-one flow map of the ODE; the other is the integration of (1.2) along the trajectories of the particles.

Given a distribution π for the initial condition x, the KL divergence from the source distribution to $\pi_{f,t}$ at time t = 1 can be written as

$$\mathcal{D}_{\mathrm{KL}}(\pi_{f,1} || \rho) = \mathcal{D}_{\mathrm{KL}}(X_f(\cdot, 1)_{\sharp} \pi || \rho) = \mathcal{D}_{\mathrm{KL}}(\pi || X_f^{-1}(\cdot, 1)_{\sharp} \rho)$$
$$= \mathbb{E}_{x \sim \pi} \left[\log \pi(x) - \log \rho(X_f(x, 1)) - \log \det \nabla_x X_f(x, 1) \right].$$

where $X_f^{-1}(\cdot, 1)$ denotes the inverse of $x \mapsto X_f(x, 1)$. The log determinant term above can be computed from the instantaneous change of variables formula (e.g., Chen et al. (2018)): $\log \det \nabla_x X(x, 1) = -\int_0^1 \operatorname{tr} \left(\nabla_X f(X(x,t),t) \right) dt$. The combined training objective becomes:

$$J(f) = \mathbb{E}_{x \sim \pi} \left[\log \pi(x) - \log \rho(X(x,1)) + \int_0^1 \operatorname{tr} \left(\nabla_X f(X(x,t),t) \right) dt + \lambda R(x,1) \right]$$
(A.1)

where λ controls the impact of penalization.

To evaluate the optimization objective, we need the ability to compute the matrix $\nabla_X f(X(x,t),t)$, as its trace appears in the change-of-variables term and the entire matrix appears in the regularization term. Also, we need the ability to compute $\partial_t f(X(x,t),t)$. We can assemble these two terms into a full Jacobian matrix, which we denote by $\nabla_{X,t} f(X(x,t),t)$; in practice, this is the Jacobian of a neural network with respect to all of its inputs. With the discretize-then-optimize approach of Onken et al. (2021), we can compute this matrix exactly via automatic differentiation. For details, see Appendix A.2.

In practice, we only have access to finite samples from the target measure, so we replace the population risk in objective (A.1) with an empirical risk based on this sample. Moreover, since $\log \pi(x)$ is independent of the velocity field f, it can be ignored in the optimization procedure. Hence, we arrive at the following empirical risk minimization problem:

$$J_{\text{ERM}}(f) = \frac{1}{N} \sum_{i=1}^{N} \left(-\log \rho(X(x_i, 1)) + \int_0^1 \operatorname{tr} \left(\nabla_X f(X(x_i, t), t) \right) dt + \lambda R(x_i, 1) \right) \,. \tag{ERM}$$

Putting everything together, we have Algorithm 1 in Appendix A.

Algorithm 1 Neural ODE training, problem P1

- 1: Input: sample $\mathcal{X} = \{x_i\}_{i=1}^N$ from target measure π , parameterized neural network $f(x, t; \theta)$, regularization parameter λ , source measure ρ .
- 2: Initialize θ ,
- 3: while θ not converged **do**
- Sample minibatch $\{x_i\}$ of size m from \mathfrak{X} 4:
- Set $x_i(0) = x_i$, $l_i(0) = r_i(0) = 0$ 5:
- Solve the following ODE system up to time t = 16:

$$\frac{dx_j}{dt} = f(x_j, t; \theta)$$
$$\frac{dl_j}{dt} = -\operatorname{tr}(\nabla_x f(x_j, t; \theta))$$
$$\frac{dr_j}{dt} = |\nabla_x f(x_j, t; \theta) f(x_j, t; \theta) + \partial_t f(x_j, t)|^2$$

- Compute the loss $L(\theta) = \frac{1}{m} \sum_{j=1}^{m} -\log \rho(x_j(1)) l_j(1) + \lambda r_j(1)$ Use automatic differentiation to backpropagate and update θ 7:
- 8:

9: end while

A.2 Exact computation of the Jacobian

Training neural ODEs consists of minimizing the (regularized) loss over the network weights subject to the ODE constraint. The adjoint-based methods in Chen et al. (2018), Grathwohl et al. (2019), and Finlay et al. (2020) can be viewed as an optimize-then-discretize approach: another continuous-time ODE (the *adjoint* equation) provides exact gradients with respect to network weights. Both the forward and adjoint equations are then discretized, checkpointing is typically employed to reduce memory requirements, and some care is required to ensure consistency of gradients. Alternatively, a discretize-then-optimize approach is proposed in Gholaminejad et al. (2019) and Onken et al. (2021), where one first discretizes the forward dynamics and computes gradients backwards in time via automatic differentiation. Since the training objective proposed in our work involves the entire Jacobian matrix of the velocity field, it is natural to use the discretize-thenoptimize approach, which allows exact computation of the Jacobian. As in Onken et al. (2021), the velocity field can be implemented as a ResNet and we can compute the Jacobian recursively.

Let s = (x, t) be the new variable formed by appending the time variable to the space variable. We then have the following recursive relation in a M-layer ResNet, where the $\{u_i\}$ are the outputs from intermediate layers:

$$u_{0} = \sigma(K_{0}s + b_{0})$$

$$u_{1} = u_{0} + h\sigma(K_{1}u_{0} + b_{1})$$

:

$$u_{M} = u_{M-1} + h\sigma(K_{M}u_{M-1} + b_{M})$$

Taking the gradient with respect to variable s, we have $\nabla_s u_i^T = \nabla_s u_{i-1} + h\sigma'(K_i u_{i-1} + b_i)K_i^T \nabla_s u_{i-1}$.

Therefore, we have the following update rule for the Jacobian:

$$J \leftarrow J + h\sigma'(K_i u_{i-1} + b_i)K_i^T J.$$

The network parameters K_i and b_i are to be learned. Since we use a discretized version of the velocity field in the implementation, these parameters can be updated through automatic differentiation.

B Knothe–Rosenblatt construction of triangular transport maps

Given probability measures ρ and π , the Knothe–Rosenblatt transport is, under appropriate conditions, the unique triangular monotone transport T such that $T_{\sharp}\pi = \rho$. In this section, we describe the explicit Knothe–Rosenblatt construction of triangular transport maps, as presented in Santambrogio (2015). Let $d \in \mathbb{N}$ be the dimension. For simplicity of presentation, we assume that π and ρ are supported on the hypercube $[0, 1]^d$. Let μ be a base measure (for example the Lebesgue measure) and assume that $\frac{d\pi}{d\mu} = \pi(x) \in C^0([0, 1]^d, \mathbb{R}^+)$ and $\frac{d\rho}{d\mu} = \rho(x) \in C^0([0, 1]^d, \mathbb{R}^+)$ are the corresponding densities. Assume also that the densities $\rho(x)$ and $\pi(x)$ are uniformly bounded from below by a positive constant.

For a continuous density function $f \in \{\rho, \pi\}$, we define the following auxiliary functions for $x \in [0, 1]^k$, $k \leq d$:

$$\hat{f}_{k}(x) = \int_{[0,1]^{d-k}} f(x, t_{k+1}, \dots, t_{d}) d\mu((t_{j})_{j=k+1}^{d})$$

$$f_{k}(x) = \frac{\hat{f}_{k}(x)}{\hat{f}_{k-1}(x_{[k-1]})}.$$
(B.1)

Hence $f_k(x_{[k-1]}, \cdot)$ is the marginal density of the variable x_k conditioned on $x_{[k-1]} = (x_1, \ldots, x_{k-1}) \in [0, 1]^{k-1}$.

Then, we define the corresponding CDFs:

$$F_{\pi,k}(x_{[k-1]}, x_k) = \int_0^{x_k} \pi_k(x_{[k-1]}, t_k) d\mu(t_k)$$

$$F_{\rho,k}(x_{[k-1]}, x_k) = \int_0^{x_k} \rho_k(x_{[k-1]}, t_k) d\mu(t_k),$$
(B.2)

which are well-defined for $x \in [0,1]^k$ and $k \in \{1,\ldots,d\}$. Note that these are interpreted as functions of the last variable x_k with $x_{[k-1]}$ fixed. In particular, we let $F_{\rho,k}(x_{[k-1]}, \cdot)^{-1}$ be the inverse of the map $x_k \to F_{\rho,k}(x_{[k-1]}, x_k)$

For $x \in [0, 1]^d$, the Knothe–Rosenblatt map is constructed recursively in the following way. First, define

$$T_1(x_1) = F_{\rho,1}^{-1} \circ F_{\pi,1}(x_1),$$

and for k > 1, define

$$T_k(x_{[k-1]}, \cdot) = F_{\rho,k}(T_1(x_1), \dots, T_{k-1}(x_{[k-1]}), \cdot)^{-1} \circ F_{\pi,k}(x_{[k-1]}, \cdot).$$
(B.3)

Then the map

$$T(x_1, \dots, x_d) = \left(T_1(x_1), T_2(x_{[2]}), \dots, T_d(x_{[d]})\right)$$

is the triangular Knothe–Rosenblatt transport $T : [0, 1]^d \to [0, 1]^d$, for which we have the following theorem:

Theorem B.1. The triangular Knothe–Rosenblatt map satisfies $T_{\sharp}\pi = \rho$ and det $\nabla T(x)\rho(T(x)) = \pi(x)$, $\forall x \in [0,1]^d$.

We comment that regularity assumptions for Theorem B.1 can be relaxed. For a more detailed discussion, see Bogachev et al. (2007).

C Auxiliary results

In this appendix, we collect statements and proofs that are required for the proofs of the main theorems.

First, we present two technical results about domains.

Lemma C.1. (uniform-cone characterization of convex domains) Let $\Omega \subset \mathbb{R}^d$ be a bounded, convex, and open domain. Then Ω is a Lipschitz domain.

Proof of Lemma C.1. Without loss of generality, we may assume that $0 \in \Omega$. Since Ω is bounded and open, there exist r, R > 0 such that $B(0, r) \subset \Omega \subset B(0, R)$, where B(0, r) and B(0, R) denote balls of radius r and R respectively.

Then, we can cover the surface of the ball of radius R by overlapping d-1 dimensional balls of radius ϵ such that the boundary of each such ball, $B_{d-1}(0,\epsilon)$, is completely covered by the adjacent balls. If \vec{n} denotes the unit vector emanating from the origin in the direction of the center of such a ball, then $U = \{t\vec{n} + y : t \ge 0, y \in B_{d-1}(0,\epsilon)\}$ is the cylinder of radius ϵ whose intersection with B(0,R) is the boundary of this ball.

Since the surface of B(0, R) can be covered by finitely many such d-1 dimensional balls, we can find a finite collection of such cylinders $\{U_j\}_{j=1}^J$ so that their union cover Ω . From this construction of $\{U_j\}_{j=1}^J$, the first property in the definition of Lipschitz domain is clearly satisfied.

To verify the second property, note that for each j, the coordinate system is simply the map that transforms the cylinder U_j to align with the direction of e_d , where e_d is the last vector of the standard basis of \mathbb{R}^d . For any $x \in \partial \Omega \cap U_j$, the cone defined by the convex closure of $\{x\} \cup B(0, r)$ is contained in the closure of Ω and the head angle α of the cone satisfies $\sin(\frac{\alpha}{2}) \geq \frac{r}{R}$, and thus the boundary is a Lipschitz function. \Box

The image of a Lipschitz domain under a sufficiently regular map remains a Lipschitz domain:

Theorem C.2 ((Hofmann et al., 2007, Theorem 4.1)). Assume $\Omega \subset \mathbb{R}^d$ is a bounded Lipschitz domain and \mathcal{O} is an open neighborhood of $\overline{\Omega}$ and $f : \mathcal{O} \to \mathbb{R}^n$ is a C^1 -diffeomorphism onto its image. Then, $\widetilde{\Omega} = f(\Omega)$ is also a Lipschitz domain.

Next, we we shall state the following regularity result about the regularity of optimal transport map from Panaretos and Zemel (2020), which is used in the proof of Theorem 3.12.

Theorem C.3. Fix open sets $\Omega_0, \Omega_1 \subset \mathbb{R}^d$ with Ω_1 convex and absolutely continuous measure μ, ν with finite second moment and bounded, strictly positive densities f, g respectively such that $\mu(\Omega_0) = \nu(\Omega_1) = 1$. Let ϕ be such that $\nabla \phi_{\sharp} \mu = \nu$. If Ω_0 and Ω_1 are bounded and f, g bounded from below, then ϕ is strictly convex and of class $C^{1,\alpha}(\Omega_0)$ for some $\alpha > 0$. In addition, if $f, g \in C^{k,\alpha}$, then $\phi \in C^{k+2,\alpha}(\Omega_0)$.

Then, we shall need the following results about composition and the inverse of a function in Section 4.

Theorem C.4 (Faá di Bruno). Let $k \in \mathbb{N}$. Let X, Y and Z be three Banach spaces, and let $F \in C^k(X, Y)$ and $G \in C^k(Y, Z)$. Then for all $0 \le n \le k$ and with $T_n := \{ \boldsymbol{\alpha} \in \mathbb{N}^n : \sum_{j=1}^n j\alpha_j = n \}$, for all $x, h \in X$ the nth derivative $[D^n(G \circ F)](x)(h^n) \in Z$ of $G \circ F$ at x evaluated at $h^n \in X^n$ equals

$$\sum_{\boldsymbol{\alpha}\in T_n} \frac{n!}{\boldsymbol{\alpha}!} [D^{|\boldsymbol{\alpha}|}G](F(x)) \left(\underbrace{\frac{[DF(x)](h)}{1!}, \dots, \frac{[DF(x)](h)}{1!}}_{\alpha_1 \text{ times}}, \dots, \underbrace{\frac{[D^nF(x)](h^n)}{n!}, \dots, \frac{[D^nF(x)](h^n)}{n!}}_{\alpha_n \text{ times}}\right).$$

Proof. Without loss of generality assume $F(0) = 0 \in Y$. Using Taylor's theorem (in Banach spaces) for G

$$G(F(x)) = \sum_{r=1}^{k} \frac{[D^r G](0)}{r!} (\underbrace{F(x), \dots, F(x)}_{r \text{ times}}) + o(\|F(x)\|_Y^k) \quad \text{as } \|F(x)\|_Y \to 0$$

Taylor expanding F(x) around $0 \in X$ then implies

$$G(F(x)) = \sum_{r=1}^{k} \frac{[D^r G](0)}{r!} (S_r(x)) + o(||F(x)||_Y^k)$$

where, using the notation x^{α_j} to denote $(x, \ldots, x) \in X^{\alpha_j}$,

$$S_r(x) = \left(\sum_{\alpha_1=1}^k \frac{[D^{\alpha_1}F](0)}{\alpha_1!}(x^{\alpha_1}) + o(\|x\|_X^k), \dots, \sum_{\alpha_r=1}^k \frac{[D^{\alpha_r}F](0)}{n!}(x^{\alpha_r}) + o(\|x\|_X^k)\right) \in Y^r.$$

Note that $F \in C^1$ and $F(0) = 0 \in Y$ implies $o(||F(x)||^k) = o(||x||^k)$ as $x \to 0$. Using multilinearity of the differential operators we thus find

$$G(F(x)) = \sum_{r=1}^{k} \sum_{\{\boldsymbol{\alpha} \in \mathbb{N}^r : |\boldsymbol{\alpha}| \le k\}} \frac{[D^r G](0)}{r!} \left(\frac{[D^{\alpha_1} F](0)(x^{\alpha_1})}{\alpha_1!}, \dots, \frac{[D^{\alpha_r} F](0)(x^{\alpha_r})}{\alpha_r!} \right) + o(||x||^k)$$

as $x \to 0$.

On the other hand, we can Taylor expand $G \circ F$. That is,

$$G(F(x)) = \sum_{r=1}^{k} \frac{[D^r(G \circ F)(0)]}{r!} (x^r) + o(||x||^k) \quad \text{as } x \to 0.$$

Comparing the powers of x, we get for $n \leq k$

$$D^{n}(G \circ F)(0)(x^{n}) = n! \sum_{r=1}^{k} \sum_{\{\alpha \in \mathbb{N}^{r} : |\alpha|=n\}} \frac{[D^{r}G](0)}{r!} \left(\frac{[D^{\alpha_{1}}F](0)(x^{\alpha_{1}})}{\alpha_{1}!}, \dots, \frac{[D^{\alpha_{r}}F](0)(x^{\alpha_{r}})}{\alpha_{r}!}\right)$$

To show that the expression for $D^n(G \circ F)(0)(x^n)$ is equivalent to the one given by Faá di Bruno's formula in Theorem 4.1, we make the following observation: the summation $\sum_{\{\alpha \in \mathbb{N}^r : |\alpha|=n\}}$ is over the partition of a set of n elements into r subsets each of α_i elements such that $r \ge i \ge 1, \alpha_i \ge 1$. If we let set $T_n = \{\alpha' = (\alpha'_1, \ldots, \alpha'_n) : \sum_{j=1}^n j\alpha'_j = n\}$ as in Theorem 4.1, each α'_j can be interpreted as the number of subsets with j elements and the total number of subsets is given by $|\alpha'|$. Another observation we make is that the summation in Theorem 4.1 takes ordered tuples while the summation in the above expression takes

unordered tuple $(\alpha_1, ..., \alpha_r)$. If $|\alpha'| = r$, the number of ways to arrange a tuple of r elements such that α'_1 of the elements are same (of value 1), α'_2 of the elements are the same (of value 2),..., and α'_n of the elements are same (of value n), is given by $\frac{r!}{\alpha'_1!...\alpha'_n!} = \frac{r!}{\alpha'!}$.

Therefore, by regrouping the summation, we obtain:

$$D^{n}(G \circ F)(0)(x^{n}) = n! \sum_{r=1}^{k} \sum_{\{\alpha \in \mathbb{N}^{r} : |\alpha| = n\}} \frac{[D^{r}G](0)}{r!} \left(\frac{[D^{\alpha_{1}}F](0)(x^{\alpha_{1}})}{\alpha_{1}!}, \dots, \frac{[D^{\alpha_{r}}F](0)(x^{\alpha_{r}})}{\alpha_{r}!} \right)$$

$$= \sum_{\alpha' \in T_{n}} \frac{n!}{r!} \frac{r!}{\alpha'!} [D^{|\alpha'|}G](0) \left(\underbrace{\frac{[DF](0)(x)}{1!}, \dots, \frac{[DF](0)(x)}{1!}}_{\alpha'_{1} \text{ times}}, \dots, \underbrace{\frac{[D^{n}F](0)(x^{n})}{n!}, \dots, \frac{[D^{n}F](0)(x^{n})}{\alpha'_{n} \text{ times}}}_{\alpha'_{n} \text{ times}} \right)$$

$$= \sum_{\alpha' \in T_{n}} \frac{n!}{\alpha'!} [D^{|\alpha'|}G](0) \left(\underbrace{\frac{[DF](0)(x)}{1!}, \dots, \frac{[DF](0)(x)}{1!}}_{\alpha'_{1} \text{ times}}, \dots, \underbrace{\frac{[D^{n}F](0)(x^{n})}{n!}, \dots, \frac{[D^{n}F](0)(x^{n})}{n!}}_{\alpha'_{n} \text{ times}} \right)$$

Theorem C.5 (Inverse function theorem). Let $k \ge 1$, let X, Y be two Banach spaces, and let $F \in C^k(X, Y)$. At every $x \in X$ for which $DF(x) \in L^1(X, Y)$ is an isomorphism, there exists an open neighbourhood $O \subseteq Y$ of F(x) and a function $G \in C^k(O, X)$ such that F(G(y)) = y for all $y \in O$.

Moreover, for every $n \leq k$ there exists a continuous function $C_n : \mathbb{R}^{n+1}_+ \to \mathbb{R}_+$ (independent of F, G, O) such that for y = F(x) with x as above

$$\|D^{n}G(y)\|_{\mathcal{L}^{n}_{\rm sym}(Y;X)} \leq C_{n}(\|[DF(x)]^{-1}\|_{\mathcal{L}^{1}_{\rm sym}(Y;X)}, \|DF(x)\|_{\mathcal{L}^{1}_{\rm sym}(X;Y)}, \dots, \|D^{n}F(x)\|_{\mathcal{L}^{n}_{\rm sym}(X;Y)}).$$
(C.1)

Proof. The stated local invertibility of F holds by the inverse function theorem in Banach spaces, see for instance (Deimling, 1985, Cor. 15.1).

Next, if F(G(y)) = y in a neighbourhood of y = F(x), then by Theorem 4.1, for $2 \le n \le k$

$$0 = D^n(F \circ G)(y) = \sum_{\boldsymbol{\alpha} \in T_n} \frac{n!}{\boldsymbol{\alpha}!} [D^{|\boldsymbol{\alpha}|}F](G(y)) \prod_{m=1}^n \left(\frac{[D^m G](y)}{m!}\right)^{\alpha_m},$$
(C.2)

where $T_n = \{ \boldsymbol{\alpha} \in \mathbb{N}^n : \sum_{j=1}^n j\alpha_j = n \}$. The only multiindex with $\alpha_n \neq 0$ is $\boldsymbol{\alpha} = (0, \dots, 0, 1)$. Solving (C.2) for $D^n G(y)$, we obtain with $\overline{T}_n \coloneqq \{ \boldsymbol{\alpha} \in \mathbb{N}^{n-1} : \sum_{j=1}^{n-1} j\alpha_j = n \}$ and y = F(x),

$$D^{n}G(y) = -([DF](G(y)))^{-1} \left(\sum_{\alpha \in \bar{T}_{n}} \frac{n!}{\alpha!} [D^{|\alpha|}F](G(y)) \left(\prod_{m=1}^{n-1} \left(\frac{[D^{m}G](y)}{m!} \right)^{\alpha_{m}} \right) \right).$$
(C.3)

Since G(y) = x, for n = 1, we have

$$||DG(y)||_{\mathcal{L}^{1}(Y;X)} = ||[DF(x)]^{-1}||_{\mathcal{L}^{1}(Y;X)}$$

and thus (C.1) holds with

$$C_1(\|[DF(x)]^{-1}\|_{\mathcal{L}^1(Y;X)}, \|DF(x)\|_{\mathcal{L}^1(X;Y)}) \coloneqq \|[DF(x)]^{-1}\|_{\mathcal{L}^1(Y;X)}.$$

Since the right-hand side of (C.3) only depends on G through $D^m G$ with $m \le n-1$ and on F through $D^{|\alpha|}F$ with $|\alpha| \le n$, an induction argument implies the existence of $C_n : \mathbb{R}^{n+1}_+ \to \mathbb{R}_+$ as in (C.1) for every $n \ge 2$.

A technical lemma that upper bounds terms in Faá di Bruno's formula:

Lemma C.6. For every $n \in \mathbb{N}$ holds $\sum_{\alpha \in T_n} \frac{n!}{\alpha!} \prod_{j=1}^n \frac{1}{(j!)^{\alpha_j}} \leq n^n$.

Proof. The sum $\sum_{\{\alpha \in T_n : |\alpha|=k\}} \frac{n!}{\alpha!} \prod_{j=1}^n \frac{1}{(j!)^{\alpha_j}}$ is equal to the so-called Stirling number of the second kind, S_n^k (Constantine and Savits (1996)). Therefore, all we need to do is to upper bound $\sum_{k=1}^n S_n^k$.

Note S_n^k denote the number of ways to distribute *n* distinct items into *k* non-distinct boxes such that each box contains at least one item. Then, $k!S_n^k$ is the number of ways to distribute *n* distinct items into *k* distinct boxes such that none of the boxes are empty. The total number of ways to distribute *n* distinct items into *n* distinct boxes is given by n^n (without the restriction that the boxes are nonempty). Therefore, we have

$$\sum_{k=1}^{n} \binom{n}{k} k! S_n^k = n^n,$$

and it follows that $\sum_{k=1}^{n} S_n^k \leq n^n$.

We also collect several auxiliary results about norm of matrices and their inverses.

Lemma C.7. Suppose M is a block triangular matrix:

$$M = \begin{pmatrix} A & B \\ 0 & D \end{pmatrix}$$

where A, D are invertible. Then,

$$||M^{-1}||_2 \le ||D^{-1}||_2 + ||A^{-1}BD^{-1}||_2 + ||A^{-1}||_2.$$

Proof. Since A, D are invertible, the inverse of M is

$$M^{-1} = \begin{pmatrix} A^{-1} & -A^{-1}BD^{-1} \\ 0 & D^{-1} \end{pmatrix}$$

The claim follows from the triangle inequality.

Lemma C.8. Let $A \in \mathbb{R}^{d \times d}$ be regular. Then $||A^{-1}||_2 \leq \frac{||A||_2^{d-1}}{|\det(A)|}$

Proof. Denote the singular values of A by $\sigma_1 \ge \sigma_2 \ge \cdots \ge \sigma_d > 0$. Then

$$\frac{\|A\|_2^{d-1}}{|\det(A)|} = \frac{\sigma_1^{d-1}}{\prod_{i=1}^d \sigma_i} = \left(\prod_{i=1}^{d-1} \frac{\sigma_1}{\sigma_i}\right) \frac{1}{\sigma_1} \ge \frac{1}{\sigma_1} = \|A^{-1}\|_2.$$

38

Lemma C.9. Let $T \in C^1(\Omega_0, \Omega_1)$ be a monotonic triangular map. Moreover, assume $\nabla T(x)$ has positive eigenvalues $\lambda_j(x)$ for $j = 1, \dots, d$ and all $x \in \Omega_0$. Then for all $t \in [0, 1]$ and all $x \in \Omega_0$

$$\|(\nabla_x T_t(x))^{-1}\|_2 = \|((1-t)I + t\nabla_x T(x))^{-1}\|_2 \le \frac{\max\{1, \|T\|_{C^1(\Omega_0)}\}^{d-1}}{\min\{1, \inf_{x \in \Omega_0} \det dT(x)\}}\}.$$

Proof. By Lemma C.8, for every $(x, t) \in \Omega_0 \times [0, 1]$

$$\|(\nabla_x T_t(x))^{-1}\|_2 \le \frac{\|\nabla_x T_t(x)\|_2^{d-1}}{|\det(T_t(x))|}$$

We have $\|\nabla T_t(x)\|_2 \leq (1-t) + t \|\nabla T(x)\|_2 \leq \max\{1, \|T\|_{C^1(\Omega_0)}\}$. For $s \in [0,1]$, $t \in (0,1)$ set $g_t(s) := \log(1-t+e^s)$. This function is convex in $s \in [0,1]$. Thus

$$g_t\left(\frac{1}{d}\sum_{i=1}^d \log(t\lambda_i(x))\right) \le \frac{1}{d}\sum_{i=1}^d g_t(\log(t\lambda_i(x))).$$

and therefore

$$\sum_{i=1}^{d} \log(1-t+t\lambda_i(x)) \ge d\log\left(1-t+t\left(\prod_{i=1}^{d}\lambda_i(x)\right)^{1/d}\right)$$
$$= d\log(1-t+t\det(\nabla T(x))^{1/d}).$$

Taking the exponential on both sides, we conclude that

$$\det((1-t)I + t\nabla T(x)) \ge (1-t + t\det(\nabla T(x))^{1/d})^d \ge \min\{1, \det dT(x)\}.$$

Finally, Lemma C.8 gives the result.

Finally, a technical lemma about the upper bounds on the C^k norm of the reciprocal of a C^k function bounded away from zero.

Lemma C.10. Let $k \in \mathbb{N}, k \geq 1$ and $f \in C^k(D)$ for domain $D \subset \mathbb{R}^d$ such that $\inf_{x \in D} f(x) > C_2$ for some constant $C_2 > 0$. Assume further $||f||_{C^k(D)} \leq C_1$ for another constant $C_1 > 0$. Then, it holds that

$$\|\frac{1}{f}\|_{C^k(D)} \le C \frac{C_1^k}{C_2^{k+1}}$$

for some constant C that depends on k but independent of f, C_1, C_2 .

Proof. We proceed as in (Zech and Marzouk, 2022a, Lemma C.4 (iii)). Let $n \in \mathbb{N}$ be an integer such that $0 \le n \le k$. We shall show by induction that $D^n(\frac{1}{f}) = \frac{p_n}{f^{n+1}}$ for some function p_n such that $||p_n||_{C^{k-n}(D)} \le C(||f||_{C^k(D)})^n$.

When n = 1, it holds that $D(\frac{1}{f}) = \frac{-Df}{f^2}$ and thus $||D(\frac{1}{f})||_{C^{k-1}(D)} \leq \frac{C_1}{C_2^2}$. Assume the induction hypothesis holds, for n + 1, we have

$$D^{n+1}(\frac{1}{f}) = D(\frac{p_n}{f^{n+1}}) = \frac{f^{n+1}Dp_n + p_n(n+1)f^nDf}{f^{2n+2}} = \frac{fDp_n + (n+1)p_nDf}{f^{n+2}} := \frac{p_{n+1}}{f^{n+2}}$$

Then it holds that

$$||p_{n+1}||_{C^{k-n-1}(D)} = ||fDp_n + (n+1)p_nDf||_{C^{k-n-1}(D)}$$

$$\leq C(n+2)||f||_{C^k(D)}||p_n||_{C^{k-n}}$$

$$\leq C(n+2)(||f||_{C^k(D)})^{n+1}.$$

Therefore, we have $\|\frac{1}{f}\|_{C^k(D)} \leq C \frac{C_1^k}{C_2^{k+1}}$ for some constant C that depends on k but independent of f, C_1, C_2 .

References

- Michael S. Albergo and Eric Vanden-Eijnden. Building normalizing flows with stochastic interpolants. In *The Eleventh International Conference on Learning Representations, ICLR 2023, Kigali, Rwanda, May 1-5, 2023.* OpenReview.net, 2023. URL https://openreview.net/forum?id=li7qeBbCR1t.
- Michael S. Albergo, Nicholas M. Boffi, and Eric Vanden-Eijnden. Stochastic interpolants: A unifying framework for flows and diffusions. *CoRR*, abs/2303.08797, 2023. doi: 10.48550/ARXIV.2303.08797. URL https://doi.org/10.48550/arXiv.2303.08797.
- Antonio Álvarez-López, Borjan Geshkovski, and Domènec Ruiz-Balet. Constructive approximate transport maps with normalizing flows. *arXiv preprint arXiv:2412.19366*, 2024.
- V.I. Arnold and R.A. Silverman. *Ordinary Differential Equations*. London, 1978. ISBN 9780262510189. URL https://books.google.com/books?id=5NumQgAACAAJ.
- Ricardo Baptista, Youssef Marzouk, and Olivier Zahm. On the representation and learning of monotone triangular transport maps. *Foundations of Computational Mathematics*, Nov 2023. ISSN 1615-3383. doi: 10.1007/s10208-023-09630-x. URL https://doi.org/10.1007/s10208-023-09630-x.
- Ricardo Baptista, Bamdad Hosseini, Nikola B Kovachki, Youssef M Marzouk, and Amir Sagiv. An approximation theory framework for measure-transport sampling algorithms. *Mathematics of Computation*, 2024.
- Jens Behrmann, Will Grathwohl, Ricky T. Q. Chen, David Duvenaud, and Jörn-Henrik Jacobsen. Invertible residual networks. In Kamalika Chaudhuri and Ruslan Salakhutdinov, editors, *Proceedings of the 36th International Conference on Machine Learning, ICML 2019, 9-15 June 2019, Long Beach, California, USA*, volume 97 of *Proceedings of Machine Learning Research*, pages 573–582. PMLR, 2019. URL http://proceedings.mlr.press/v97/behrmann19a.html.
- Jean-David Benamou and Yann Brenier. A computational fluid mechanics solution to the mongekantorovich mass transfer problem. *Numerische Mathematik*, 84(3):375–393, 2000. doi: 10.1007/ s002110050002. URL https://doi.org/10.1007/s002110050002.
- Joe Benton, George Deligiannidis, and Arnaud Doucet. Error bounds for flow matching methods. *Trans. Mach. Learn. Res.*, 2024, 2024. URL https://openreview.net/forum?id=uqQPyWFDhY.
- Vladimir Bogachev, Alexander Kolesnikov, and Kirill Medvedev. Triangular transformations of measures. *Sbornik: Mathematics*, 196:309, 10 2007. doi: 10.1070/SM2005v196n03ABEH000882.

- Yann Brenier. Polar factorization and monotone rearrangement of vector-valued functions. Communications on Pure and Applied Mathematics, 44:375–417, 1991.
- Soo Bong Chae. *Holomorphy and calculus in normed spaces*, volume 92 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker, Inc., New York, 1985. ISBN 0-8247-7231-8. With an appendix by Angus E. Taylor.
- Tian Qi Chen, Jesse Bettencourt, and David Duvenaud. Neural ordinary differential equations. *CoRR*, abs/1806.07366, 2018. URL http://arxiv.org/abs/1806.07366.
- G. Constantine and T. H. Savits. A multivariate faa di bruno formula with applications. *Transactions of the American Mathematical Society*, 348:503–520, 1996.
- Klaus Deimling. *Nonlinear functional analysis*. Springer-Verlag, Berlin, 1985. ISBN 3-540-13928-1. doi: 10.1007/978-3-662-00547-7. URL https://doi.org/10.1007/978-3-662-00547-7.
- Laurent Dinh, David Krueger, and Yoshua Bengio. NICE: non-linear independent components estimation. In Yoshua Bengio and Yann LeCun, editors, 3rd International Conference on Learning Representations, ICLR 2015, San Diego, CA, USA, May 7-9, 2015, Workshop Track Proceedings, 2015. URL http://arxiv.org/abs/1410.8516.
- Chris Finlay, Jörn-Henrik Jacobsen, Levon Nurbekyan, and Adam M. Oberman. How to train your neural ODE: the world of jacobian and kinetic regularization. In *Proceedings of the 37th International Conference on Machine Learning, ICML 2020, 13-18 July 2020, Virtual Event*, volume 119 of *Proceedings of Machine Learning Research*, pages 3154–3164. PMLR, 2020. URL http://proceedings.mlr.press/v119/finlay20a.html.
- Yuan Gao, Jian Huang, Yuling Jiao, and Shurong Zheng. Convergence of continuous normalizing flows for learning probability distributions. *CoRR*, abs/2404.00551, 2024. doi: 10.48550/ARXIV.2404.00551. URL https://doi.org/10.48550/arXiv.2404.00551.
- Amir Gholaminejad, Kurt Keutzer, and George Biros. ANODE: unconditionally accurate memoryefficient gradients for neural odes. In Sarit Kraus, editor, *Proceedings of the Twenty-Eighth International Joint Conference on Artificial Intelligence, IJCAI 2019, Macao, China, August 10-16, 2019*, pages 730–736. ijcai.org, 2019. doi: 10.24963/ijcai.2019/103. URL https://doi.org/10.24963/ijcai.2019/103.
- Will Grathwohl, Ricky T. Q. Chen, Jesse Bettencourt, Ilya Sutskever, and David Duvenaud. FFJORD: freeform continuous dynamics for scalable reversible generative models. In 7th International Conference on Learning Representations, ICLR 2019, New Orleans, LA, USA, May 6-9, 2019. OpenReview.net, 2019. URL https://openreview.net/forum?id=rJxgknCcK7.
- S. Hofmann, M. Mitrea, and Michael E. Taylor. Geometric and transformational properties of lipschitz domains, semmes-kenig-toro domains, and other classes of finite perimeter domains. *The Journal of Geometric Analysis*, 17:593–647, 2007.
- Chin-Wei Huang, David Krueger, Alexandre Lacoste, and Aaron C. Courville. Neural autoregressive flows. In Jennifer G. Dy and Andreas Krause, editors, *Proceedings of the 35th International Conference on Machine Learning, ICML 2018, Stockholmsmässan, Stockholm, Sweden, July 10-15, 2018,*

volume 80 of *Proceedings of Machine Learning Research*, pages 2083-2092. PMLR, 2018. URL http://proceedings.mlr.press/v80/huang18d.html.

- Chin-Wei Huang, Ricky T. Q. Chen, Christos Tsirigotis, and Aaron Courville. Convex potential flows: Universal probability distributions with optimal transport and convex optimization. In *International Conference on Learning Representations*, 2021. URL https://openreview.net/forum?id=te7PVH1sPxJ.
- Daniel Zhengyu Huang, Jiaoyang Huang, and Zhengjiang Lin. Convergence analysis of probability flow ODE for score-based generative models. *CoRR*, abs/2404.09730, 2024. doi: 10.48550/ARXIV.2404.09730. URL https://doi.org/10.48550/arXiv.2404.09730.
- Diederik P. Kingma, Tim Salimans, Rafal Józefowicz, Xi Chen, Ilya Sutskever, and Max Welling. Improving variational autoencoders with inverse autoregressive flow. In Daniel D. Lee, Masashi Sugiyama, Ulrike von Luxburg, Isabelle Guyon, and Roman Garnett, editors, Advances in Neural Information Processing Systems 29: Annual Conference on Neural Information Processing Systems 2016, December 5-10, 2016, Barcelona, Spain, pages 4736–4744, 2016. URL https://proceedings.neurips.cc/paper/2016/hash/ddeebdeefdb7e7e7a697e1c3e3d8ef54-F
- Durk P Kingma and Prafulla Dhariwal. Glow: Generative flow with invertible 1x1 convolutions. In S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett, editors, Advances in Neural Information Processing Systems, volume 31. Curran Associates, Inc., 2018. URL https://proceedings.neurips.cc/paper_files/paper/2018/file/d139db6a236200b21cc7f7
- Ivan Kobyzev, Simon Prince, and Marcus Brubaker. Normalizing flows: An introduction and review of current methods. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, page 1–1, 2020. ISSN 1939-3539. doi: 10.1109/tpami.2020.2992934. URL http://dx.doi.org/10.1109/TPAMI.2020.2992934.
- Zhifeng Kong and Kamalika Chaudhuri. The expressive power of a class of normalizing flow models. In Silvia Chiappa and Roberto Calandra, editors, *The 23rd International Conference on Artificial Intelligence and Statistics, AISTATS 2020, 26-28 August 2020, Online [Palermo, Sicily, Italy]*, volume 108 of *Proceedings of Machine Learning Research*, pages 3599–3609. PMLR, 2020. URL http://proceedings.mlr.press/v108/kong20a.html.
- Gen Li, Yuting Wei, Yuejie Chi, and Yuxin Chen. A sharp convergence theory for the probability flow odes of diffusion models. *CoRR*, abs/2408.02320, 2024. doi: 10.48550/ARXIV.2408.02320. URL https://doi.org/10.48550/arXiv.2408.02320.
- Qianxiao Li, Ting Lin, and Zuowei Shen. Deep learning via dynamical systems: An approximation perspective. *CoRR*, abs/1912.10382, 2019. URL http://arxiv.org/abs/1912.10382.
- Yaron Lipman, Ricky T. Q. Chen, Heli Ben-Hamu, Maximilian Nickel, and Matthew Le. Flow matching for generative modeling. In *The Eleventh International Conference on Learning Representations, ICLR 2023, Kigali, Rwanda, May 1-5, 2023.* OpenReview.net, 2023. URL https://openreview.net/forum?id=PqvMRDCJT9t.

- Xingchao Liu, Chengyue Gong, and Qiang Liu. Flow straight and fast: Learning to generate and transfer data with rectified flow. In *The Eleventh International Conference on Learning Representations, ICLR 2023, Kigali, Rwanda, May 1-5, 2023.* OpenReview.net, 2023. URL https://openreview.net/forum?id=XVjTT1nw5z.
- Yiping Lu, Aoxiao Zhong, Quanzheng Li, and Bin Dong. Beyond finite layer neural networks: Bridging deep architectures and numerical differential equations. In 6th International Conference on Learning Representations, ICLR 2018, Vancouver, BC, Canada, April 30 - May 3, 2018, Workshop Track Proceedings. OpenReview.net, 2018. URL https://openreview.net/forum?id=BlLatDVUM.
- Youssef Marzouk, Tarek Moselhy, Matthew Parno. and Alessio Spantini. Sammeasure transport: introduction. Handbook Uncertainty pling via An of Quan-2016. tification, page 1-41, doi: 10.1007/978-3-319-11259-6 23-1. URL http://dx.doi.org/10.1007/978-3-319-11259-6_23-1.
- Youssef Marzouk, Zhi Ren, Sven Wang, and Jakob Zech. Distribution learning via neural differential equations: a nonparametric statistical perspective. *Journal of Machine Learning Research*, 25:1–61, 2024. arXiv:2309.01043.
- Tarek A. Youssef M. Marzouk. Moselhy and Bayesian inference with op-2012. timal maps. Journal of Computational Physics, 231(23):7815-7850, https://doi.org/10.1016/j.jcp.2012.07.022. ISSN 0021-9991. doi: URL https://www.sciencedirect.com/science/article/pii/S0021999112003956.
- Derek Onken, Samy Wu Fung, Xingjian Li, and Lars Ruthotto. Ot-flow: Fast and accurate continuous normalizing flows via optimal transport. In Thirty-Fifth AAAI Conference on Artificial Intelligence, AAAI 2021, Thirty-Third Conference on Innovative Applications of Artificial Intelligence, IAAI 2021, The Eleventh Symposium on Educational Advances in Artificial Intelligence, EAAI 2021, Virtual Event, February 2-9, 2021, pages 9223–9232. AAAI Press, 2021. URL https://ojs.aaai.org/index.php/AAAI/article/view/17113.
- Victor M. Panaretos and Yoav Zemel. An Invitation to **Statistics** in URL Wasserstein Space. Springer. 2020. ISBN 978-3-030-38437-1. https://link.springer.com/content/pdf/10.1007%2F978-3-030-38438-8.pdf.
- Danilo Jimenez Rezende and Shakir Mohamed. Variational inference with normalizing flows. In Francis R. Bach and David M. Blei, editors, *Proceedings of the 32nd International Conference on Machine Learning, ICML 2015, Lille, France, 6-11 July 2015,* volume 37 of *JMLR Workshop and Conference Proceedings*, pages 1530–1538. JMLR.org, 2015. URL http://proceedings.mlr.press/v37/rezende15.html.
- Domènec Ruiz-Balet and Enrique Zuazua. Neural ODE control for classification, approximation, and transport. *SIAM Rev.*, 65(3):735–773, 2023. doi: 10.1137/21M1411433. URL https://doi.org/10.1137/21m1411433.
- Lars Ruthotto and Eldad Haber. Deep neural networks motivated by partial differential equations. J. Math. Imaging Vis., 62(3):352–364, 2020. doi: 10.1007/s10851-019-00903-1. URL https://doi.org/10.1007/s10851-019-00903-1.

- Filippo Santambrogio. Optimal transport for applied mathematicians. calculus of variations, pdes and modeling. 2015. URL https://www.math.u-psud.fr/~filippo/OTAM-cvgmt.pdf.
- Yang Song, Jascha Sohl-Dickstein, Diederik P. Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-based generative modeling through stochastic differential equations. In 9th International Conference on Learning Representations, ICLR 2021, Virtual Event, Austria, May 3-7, 2021. OpenReview.net, 2021. URL https://openreview.net/forum?id=PxTIG12RRHS.
- Sho Sonoda and Noboru Murata. Transport analysis of infinitely deep neural network. J. Mach. Learn. Res., 20:2:1–2:52, 2019. URL http://jmlr.org/papers/v20/16-243.html.
- Michael D. Spivak. Calculus on Manifolds: A Modern Approach to Classical Theorems of Advanced Calculus. Harper Collins Publishers, 1965. ISBN 0805390219. URL https://www.maa.org/press/maa-reviews/calculus-on-manifolds-a-modern-approach-to
- Elias M. Stein. Singular Integrals and Differentiability Properties of Functions (PMS-30). Princeton University Press, 1970. ISBN 9780691080796. URL http://www.jstor.org/stable/j.ctt1bpmb07.
- Esteban G. Tabak and Eric Vanden-Eijnden. Density estimation by dual ascent of the log-likelihood. *Communications in Mathematical Sciences*, 8(1):217 233, 2010.
- Takeshi Teshima, Isao Ishikawa, Koichi Tojo, Kenta Oono, Masahiro Ikeda, and Masashi Sugiyama. Coupling-based invertible neural networks are universal diffeomorphism approximators. In Hugo Larochelle, Marc'Aurelio Ranzato, Raia Hadsell, Maria-Florina Balcan, and Hsuan-Tien Lin, editors, Advances in Neural Information Processing Systems 33: Annual Conference on Neural Information Processing Systems 2020, NeurIPS 2020, December 6-12, 2020, virtual, 2020a. URL https://proceedings.neurips.cc/paper/2020/hash/2290a7385ed77cc5592dc2153229f082-F
- Takeshi Teshima, Koichi Tojo, Masahiro Ikeda, Isao Ishikawa, and Kenta Oono. Universal approximation property of neural ordinary differential equations. *CoRR*, abs/2012.02414, 2020b. URL https://arxiv.org/abs/2012.02414.
- Belinda Tzen and Maxim Raginsky. Theoretical guarantees for sampling and inference in generative models with latent diffusions. In Alina Beygelzimer and Daniel Hsu, editors, *Conference on Learning Theory, COLT 2019, 25-28 June 2019, Phoenix, AZ, USA*, volume 99 of *Proceedings of Machine Learning Research*, pages 3084–3114. PMLR, 2019. URL http://proceedings.mlr.press/v99/tzen19a.html.
- Cedric Villani. Optimal Transport: Old and New. Springer, 2008. ISBN 9783540710493.
- Zheyu Oliver Wang, Ricardo Baptista, Youssef Marzouk, Lars Ruthotto, and Deepanshu Verma. Efficient neural network approaches for conditional optimal transport with applications in bayesian inference, 2023.
- Dmitry Yarotsky. Error bounds for approximations with deep relu networks. *Neural Networks*, 94:103–114, 2017. doi: 10.1016/j.neunet.2017.07.002. URL https://doi.org/10.1016/j.neunet.2017.07.002.

- Dmitry Yarotsky and Anton Zhevnerchuk. The phase diagram of approximation rates for deep neural networks. In Hugo Larochelle, Marc'Aurelio Ranzato, Raia Hadsell, Maria-Florina Balcan, and Hsuan-Tien Lin, editors, *Advances in Neural Information Processing Systems 33: Annual Conference on Neural Information Processing Systems 2020, NeurIPS 2020, December 6-12, 2020, virtual, 2020.* URL https://proceedings.neurips.cc/paper/2020/hash/979a3f14bae523dc5101c52120c535e9-F
- Jakob Zech and Youssef Marzouk. Sparse approximation of triangular transports, part i: The finite-dimensional case. *Constructive Approximation*, 2022a. doi: https://doi.org/10.1007/s00365-022-09569-2.
- Jakob Zech and Youssef Marzouk. Sparse approximation of triangular transports, part ii: The infinite-dimensional case. *Constructive Approximation*, 2022b. doi: https://doi.org/10.1007/s00365-022-09570-9.