Rotational beta expansions and Schmidt games

Hajime Kaneko^{*1}, Jonathan Caalim^{$\dagger 2$}, and Nathaniel Nollen^{$\ddagger 2$}

¹University of Tsukuba, 1-1-1 Tennodai Tsukuba, Ibaraki 305-8577, Japan ²University of the Philippines - Diliman, Quezon City, Philippines 1101

Abstract

We consider rotational beta expansions in dimensions 1, 2 and 4 and view them as expansions on real numbers, complex numbers, and quaternions, respectively. We give sufficient conditions on the parameters $\alpha, \beta \in (0, 1)$ so that particular cylinder sets arising from the expansions are winning or losing Schmidt (α, β) -game.

Keywords: numeration systems, beta expansions, complex expansions, quaternions, Schmidt game

1 Introduction

Let $\alpha, \beta \in (0, 1)$. The Schmidt (α, β) -game on a complete metric space (X, λ) is a game with two players, Alice and Bob, and the following recursive rules [10]:

- 1. At the start, Bob chooses an initial radius $\rho = \rho_0 > 0$ and an initial center $x_0 \in X$. Set $B_0 := \overline{B(x_0, \rho)}$. Meanwhile, Alice chooses a *target set* $S \subseteq X$.
- 2. Bob and Alice play alternately. For $n \in \mathbb{N}$, set $\rho_n := (\alpha \beta)^n \rho$ and for
 - (a) Alice's *n*th turn: Alice chooses $y_n \in X$ such that

$$\lambda(x_{n-1}, y_n) + \alpha \rho_{n-1} \le \rho_{n-1},$$

(b) Bob's *n*th turn: Bob chooses $x_n \in X$ such that

$$\lambda(y_n, x_n) + \rho_n \le \alpha \rho_{n-1}.$$

^{*}kanekoha@math.tsukuba.ac.jp

[†]jcaalim@math.upd.edu.ph

 $^{^{\}ddagger}$ nnollen@math.upd.edu.ph

Let $B_n := \overline{B(x_n, \rho_n)}$ and $A_n := \overline{B(y_n, \alpha \rho_{n-1})}$. Then we have

$$B_0 \supseteq A_1 \supseteq B_1 \supseteq A_2 \supseteq \cdots$$
.

The *outcome* of the game is the unique point ω such that

$$\{\omega\} = \left(\bigcap_{n=1}^{\infty} A_n\right) \cap \left(\bigcap_{n=0}^{\infty} B_n\right).$$

Alice wins the game if $\omega \in S$. Otherwise, Bob wins.

Given α, β and ρ , the set S is called (α, β, ρ) -winning if Alice has a strategy to win the game regardless of Bob's play. If S is (α, β, ρ) -winning for all $\rho > 0$, we say that S is (α, β) -winning. If S is (α, β) -winning for any β , we say that S is α -winning. A set S is winning if S is α -winning for some α .

From [10], we know that if S is winning, then S is dense. In particular, if $X = \mathbb{R}^m$, then the Hausdorff dimension of S is m. Winning sets also exhibit large intersection properties. Indeed, let α be a real number with $0 < \alpha < 1$ and let S_n (n = 1, 2, ...) be α -winning sets. Then the intersection $\bigcap_{n\geq 1}S_n$ is also α -winning. Lemma 7.13 of [2] provides a more flexible result as follows. Let $P_j = \{m_j + d_jn \mid n = 0, 1, ...\}$ (j = 1, 2, ...) be an arithmetic progression of positive integers where m_j is the first term and d_j is the common difference. Assume that $\bigcup_{j\geq 1}P_j$ is a disjoint union partitioning N. Let α, β, ρ be positive real numbers with $\alpha < 1/3$ and $\beta < 1$. Set $\beta_j := \beta(\alpha\beta)^{-1+d_j}$ and $\rho_j := \rho(\alpha\beta)^{-1+m_j}$. If S_j is an $(\alpha, \beta_j, \rho_j)$ winning set for each $j \in \mathbb{N}$, then $\bigcap_{j\geq 1}S_j$ is an (α, β, ρ) -winning set. In particular, the intersection $\bigcap_{j\geq 1}S_j$ is uncountable. This gives one of the motivations to search for (α, β, ρ) winning sets.

Schmidt game is a useful tool to study the "denseness" of cylinder sets related to numeration systems. In the literature, there are several researches on the *b*-ary expansions $(1 < b \in \mathbb{N})$ and Schmidt games. In 1982, Freiling [6] showed that the set of real numbers containing the digit 0 or 5 in base 6 is (1/2, 1/2)-winning but (1/2, 1/3)-losing. In 2002, Dremov [5] proved that the set of real numbers containing the digit 0 or 3 in base 4 is (1/2, 1/2)-losing. In 2024, Neckrasov and Zhan [8] showed that set of real numbers where 0 appears at least half of the time in the *n*-tails ($n \in \mathbb{N}$) of its binary expansion is (α, β)-losing for $\alpha > \beta$. Given $b \in \mathbb{N} > 1$, Zanger-Tishler and Kalia [12] gave a sufficient condition on (α, β) so that the set of real numbers with 0 or b - 1 in their *b*-ary expansion is either (α, β) -winning or (α, β) -losing.

In this article, we apply Schmidt games to rotational beta expansions [1]. Let $m \in \mathbb{N}$. Let $\beta > 1$ and let M be an element of the special orthogonal group SO(m). Let \mathcal{L} be a lattice on \mathbb{R}^m with fundamental domain \mathcal{X} . Define the rotational beta transformation $\mathbb{T} = \mathbb{T}_{\beta M, \mathcal{L}, \mathcal{X}} : \mathcal{X} \to \mathcal{X}$ by

$$\mathbb{T}(z) = \beta M z - d(z),$$

where d(z) is the unique element of \mathcal{L} such that $\beta Mz - d(z) \in \mathcal{X}$. The rotational beta expansion of $z \in \mathcal{X}$ is the sequence $d(z) := d_1 d_2 \dots$ where the *n*th digit is $d_n = d_n(z) := d(\mathbb{T}^{n-1}(z))$. We have

$$z = (\beta M)^{-1} d_1 + (\beta M)^{-2} d_2 + \cdots$$

The digit set of the rotational beta expansion is $\mathcal{D} := \{d_1(z) \in \mathcal{L} : z \in \mathcal{X}\}$. An ordered *n*-tuple $d_1 d_2 \cdots d_n \in \mathcal{D}^n$ is called an admissible *n*-block (or, simply, block) if it is a finite subsequence of d(z) for some $z \in \mathcal{X}$. For an admissible block Ω , we define the set

$$\mathcal{C}[\Omega] = \mathcal{C}_{(\beta M, \mathcal{L}, \mathcal{X})}[\Omega] := \{ z \in \mathcal{X} \mid \Omega \text{ is a subblock of } d(z) \}.$$

In this paper, we study the winnability of $C[\Omega]$ with respect to Schmidt games for rotational beta expansions in dimensions m = 1, 2 and 4. The rotational beta expansion may be interpreted as: the classical beta expansion [9] if m = 1; a numeration system on \mathbb{C} with a complex number base if m = 2; and an expansion over the real quaternions with respect to a quaternion radix [4]. In Section 2, we look at the winnability of $C[\Omega]$ for m = 1. Note that we can show α -winnability (see Theorem 2.7) under certain assumptions, which may be considered reasonable. For comparison, see [7, Theorem 1.7], where Langeveld and Samuel investigated the α -winnability of the set badly approximable numbers x (i.e. the orbit of x under the (generalized) beta transformation avoids a particular given point) with respect to (generalized) beta expansions. In Section 3, we give sufficient conditions for winnability of $C[\Omega]$ where m = 2. This provides the first result of the winnability of rotational beta expansion. In Section 4, we treat the case where m = 4.

2 Real expansions

Rényi [9] introduced the so-called beta expansion on real numbers with real base b > 1. This generalizes the *b*-ary expansions on real numbers where the radix *b* is a natural number. Given a fixed $b \in \mathbb{R}$ with b > 1, the *j*th digit $d_j = d_j(x)$ of a real number $x \in [0, 1)$ is the unique integer $d \in \{0, 1, \ldots, s_b\} =: \mathcal{D}$ where

$$s_b = \begin{cases} b - 1, & \text{if } b \in \mathbb{N} \\ \lfloor b \rfloor, & \text{if } b \notin \mathbb{N} \end{cases}$$

such that

$$b^{j}\left(x-\sum_{k=1}^{j}d_{k}b^{-k}\right)\in\left[0,1\right)$$

We call the sequence $d(x) := d_1 d_2 \cdots$ the *b*-expansion of *x*.

Let $n \in \mathbb{N}$. We identify an *n*-block $a_1 a_2 \cdots a_n \in \mathcal{D}^n$ with the sequence $a_1 a_2 \cdots a_n 000 \cdots \in \mathcal{D}^{\mathbb{N}}$. For $a = a_1 a_2 \cdots \in \mathcal{D}^{\mathbb{N}}$, we define

$$a(b) := \frac{a_1}{b} + \frac{a_2}{b^2} + \cdots$$

Given $a = a_1 a_2 \cdots, b = b_1 b_2 \cdots \in \mathcal{D}^{\mathbb{N}}$, we say that a is lexicographically less than b and write

$$a <_{lex} b$$

if there exists $k \in \mathbb{N}$ such that $a_j = b_j$ when j < k and $a_k < b_k$. We write a $\leq_{\text{lex}} \mathbb{b}$ if a $<_{\text{lex}} \mathbb{b}$ or a = b. It is known that if a, $\mathbb{b} \in \mathcal{D}^{\mathbb{N}}$ are admissible, then a $<_{\text{lex}} \mathbb{b}$ if and only

if a(b) < b(b). Denote the set of admissible *n*-blocks in \mathcal{D}^n by \mathcal{D}^n_A . We can arrange the elements of \mathcal{D}^n (or \mathcal{D}^n_A) in an increasing manner with respect to the lexicographic order:

$$e_1 <_{\text{lex}} e_2 <_{\text{lex}} \cdots <_{\text{lex}} e_N.$$

We say that a and b are consecutive with respect to $<_{\text{lex}}$ if $(b, a) = (e_{j+1}, e_j)$ or $(b, a) = (e_j, e_{j+1})$ for some j. We give the following results on admissible *n*-blocks.

Lemma 2.1. Let a and b be consecutive blocks of \mathcal{D}^n_A with respect to $<_{\text{lex}}$. If a $<_{\text{lex}}$ b, then $b(b) - a(b) \le b^{-n}$.

Proof. Let $a = a_1 a_2 \cdots a_n$. Suppose $b(b) - a(b) > b^{-n}$. Let $z = a(b) + b^{-n}$ and $c = c_1 c_2 \cdots \in \mathcal{D}^{\mathbb{N}}$ be the *b*-expansion of *z*. Clearly, $a <_{\text{lex}} c$ since a(b) < z. Moreover, $c(b) = a(b) + b^{-n} < b(b)$ and so, $c <_{\text{lex}} b$. Since *a* and *b* are consecutive admissible *n*-blocks and $c <_{\text{lex}} b$, then

$$c_1c_2\cdots c_n=a_1a_2\cdots a_n.$$

Thus,

$$a(b) + b^{-n} = c(b) = \frac{a_1}{b} + \dots + \frac{a_n}{b^n} + \frac{c_{n+1}}{b^{n+1}} + \frac{c_{n+2}}{b^{n+2}} + \dots$$

Since $a(b) = \frac{a_1}{b} + \dots + \frac{a_n}{b^n}$, we have

$$1 = \frac{c_{n+1}}{b} + \frac{c_{n+2}}{b^2} + \cdots,$$

which is a contradiction as the admissibility criterion implies that $c_{n+1}c_{n+2}\cdots(b) < 1$. Therefore, $a <_{lex} b \leq_{lex} c$ and $b(b) - a(b) \leq c(b) - a(b) = z - a(b) = b^{-n}$.

Let $i \in \mathbb{N} \cup \{\infty\}$ be the length of the *b*-expansion of $b - \lfloor b \rfloor$, that is,

 $i = i_b := \max\{j \in \mathbb{N} : \text{ the } j\text{ th digit of the } b\text{-expansion of } b - \lfloor b \rfloor \text{ is nonzero}\}$

Also, we let $K = K_b$ be the maximal length of zero blocks (blocks containing only the digit zero) among the first *i* digits of the *b*-expansion of $b - \lfloor b \rfloor$.

For $d \in \mathcal{D}$ and $k \in \mathbb{N}$, let $V_k(b; d) := \{x \in [0, 1) : d_k(x) = d\}$. We follow [12] to study the winnability of the cylinder set

$$C_b[d] := \{x \in [0,1) : d \text{ appears in } d(x)\} = \bigcup_{k=1}^{\infty} V_k(b;d)$$

with respect to the Schmidt (α, β) -game.

For simplicity, we write $V_k(b; 0) = V_k$. Now, $V_1 = [0, 1/b)$. For any $j \in \{0, 1, \dots, \lfloor b \rfloor - 1\}$, we see that $j1 \in \mathcal{D}^2_A$. Thus, we have

$$V_{2} = \begin{cases} \bigcup_{j=0}^{b-1} \left[\frac{j}{b}, \frac{j}{b} + \frac{1}{b^{2}} \right) & \text{if } b \in \mathbb{N} \\ \bigcup_{\substack{\lfloor b \rfloor - 1 \\ j = 0}}^{\lfloor b \rfloor - 1} \left[\frac{j}{b}, \frac{j}{b} + \frac{1}{b^{2}} \right) \cup \left[\frac{\lfloor b \rfloor}{b}, \min\left\{ 1, \frac{\lfloor b \rfloor}{b} + \frac{1}{b^{2}} \right\} \right) & \text{if } b \notin \mathbb{N}. \end{cases}$$

For $j \in \mathbb{N}$, let $\phi_j = (j + \sqrt{j^2 + 4})/2$ be the *j*th metallic mean (that is, the golden, silver, bronze means, etc.). Note that, if $b \notin \mathbb{N}$,

$$\frac{\lfloor b \rfloor}{b} + \frac{1}{b^2} \le 1 \iff \lfloor b \rfloor < \phi_{\lfloor b \rfloor} \le b < \lfloor b \rfloor + 1.$$

For the remainder of this section, we let $c = c_1 c_2 \cdots = \lim_{\varepsilon \to 0^+} d(1 - \varepsilon)$. Let d' be the minimal digit of $d(1 - \varepsilon)$ and d an integer with $0 \le d \le d'$. In general, for $k \in \mathbb{N}$ with $k \ge 2$,

$$V_k(b;d) = \bigcup_{\mathbf{a}\in\mathcal{D}_A^{k-1}} \Delta(\mathbf{a}d) = \bigcup_{\mathbf{a}\in\mathcal{D}_A^{k-1}} [\mathbf{a}d(b), u_{\mathbf{a},d}),$$

where $\Delta(ad)$ is the set of real numbers in [0, 1) whose first k digits are given by ad, and $u_{a,d}$ is the the supremum of $\Delta(ad)$. Note that $u_{a,d} \in (ad(b), \min\{a(d+1)(b), 1\}]$.

Example 2.2. Let $b = (1 + \sqrt{5})/2$. Then $\mathcal{D}_A^3 = \{000, 001, 010, 100, 101\}$ and

• $\Delta(0000) = [0, b^{-4})$ • $\Delta(0010) = [b^{-3}, b^{-2})$

•
$$\Delta(0100) = [b^{-2}, b^{-2} + b^{-4})$$
 • $\Delta(1010) = [b^{-1} + b^{-3}, 1)$

Let

$$E_{k-1,d} := \{ a = a_1 a_2 \cdots a_{k-1} \in \mathcal{D}_A^{k-1} \mid a_j a_{j+1} \cdots a_{k-1} (d+1)(b) > 1 \text{ for some } j < k \}.$$

Proposition 2.3. Let $2 \le k \in \mathbb{N}$ and $a \in \mathcal{D}_A^{k-1}$. The following are equivalent:

- (1) $u_{\mathbf{a},d} = \mathbf{a}(d+1)(b)$, i.e., $\Delta(\mathbf{a}d)$ has length b^{-k} .
- (2) $a \notin E_{k-1,d}$.

Proof. Let $a = a_1 \cdots a_{k-1} \in \mathcal{D}_A^{k-1}$.

 $(1) \Longrightarrow (2)$ Assume $u_{a,d} = a(d+1)(b)$ and $a \in E_{k-1,d}$. Then $a_j \cdots a_{k-1}(d+1)(b) > 1$ for some j < k. If j = 1, then $u_{a,d} = a_1 \cdots a_{k-1}(d+1)(b) > 1$, a contradiction. So j > 1. Let $x = u_{a,d} - \varepsilon$ for some small $\varepsilon > 0$. Then $x \in \Delta(ad)$ and so d(x) has the form

$$\mathbf{d}(x) = a_1 \cdots a_{k-1} dx_1 x_2 \cdots$$

However, $a_i \cdots a_{k-1} dx_1 x_2 \cdots (b) \ge 1$, a contradiction.

 $(2) \Longrightarrow (1)$ Suppose a $\notin E_{k-1,d}$. Let $z \in [ad(b), a(d+1)(b))$. It is enough to show that the first k digits of z are given by ad. Observe that $a(d+1)(b) \leq 1$ and $a_2 \cdots a_{k-1}d(b) \leq bz - a_1 < a_2 \cdots a_{k-1}(d+1)(b) \leq 1$ since a $\notin E_{k-1,d}$. So $T(z) = bz - a_1$ and the first digit of z is a_1 . Similarly, $a_3 \cdots a_{k-1}d(b) \leq bT(z) - a_2 < a_3 \cdots a_{k-1}(d+1)(b) \leq 1$ since a $\notin E_{k-1,d}$ and the second digit of z is a_2 . By induction, the first k digits of z is given by ad. This completes the proof.

By Proposition 2.3, we may determine the length of an interval of V_k using the elements of E_{k-1} . Recall that if u, v, w are finite words such that w = uv, we say that u is a prefix of w and v is a suffix of w. In other words, w starts with u and ends with v. We also adopt this definition of prefixes and suffixes if w is an infinite word.

Proposition 2.4. Let $k \ge 2$ and $a \in E_{k-1,d}$. Then a has a suffix with positive length that is also a prefix of c.

Proof. Let $a = a_1 \cdots a_{k-1} \in E_{k-1,d}$. Then $a_j \cdots a_{k-1}(d+1)(b) > 1$ for some j < k. So a(d+1) is not admissible. This implies $c <_{\text{lex}} a_\ell \cdots a_{k-1}(d+1)$ for some $\ell < k$. However, $a_\ell \cdots a_{k-1}d <_{\text{lex}} c <_{\text{lex}} a_\ell \cdots a_{k-1}(d+1)$, where $a_\ell \cdots a_{k-1}d$ is identified with $a_\ell \cdots a_{k-1}d00\cdots$. This implies the first $k - \ell$ digits of c is given by $a_\ell \cdots a_{k-1}$.

Now, we consider consecutive elements of \mathcal{D}_A^{k-1} which are also elements of E_{k-1} .

Lemma 2.5. Let $k \ge 2$. Let a and b be consecutive elements of \mathcal{D}_A^{k-1} such that a $<_{\text{lex}}$ b. If a, b $\in E_{k-1,d}$, then b has the form

$$\mathbb{b} = b_1 \cdots b_{k'} \underbrace{00 \cdots 0}_{k-k'-1}$$

for some k' < k - 1 where $b_{k'} \neq 0$ and

$$a = b_1 \cdots b_{k'-1} (b_{k'} - 1) c_1 \cdots c_{k-k'-1}.$$

Moreover, the longest prefix of c that is also a suffix of a is $c_1 \cdots c_{k-k'-1}$ starting from the (k'+1)th digit.

Proof. Let a and b be consecutive elements of \mathcal{D}_A^{k-1} such that a $<_{\text{lex}}$ b and a, b $\in E_{k-1,d}$. Let $\mathbb{b} = b_1 \cdots b_{k-1}$.

Suppose $b_{k-1} \neq 0$. Then $a = b_1 \cdots b_{k-2}(b_{k-1} - 1)$. Since $a \in E_{k-1,d}$ and b is admissible, there exists j < k such that

$$b_j \cdots b_{k-2}(b_{k-1} - 1)(d+1)(b) > 1 > b_j \cdots b_{k-1}d(b)$$

$$\geq b_j \cdots b_{k-2}(b_{k-1} - 1)(d+1)(b).$$

We have a contradiction. Hence, $b_{k-1} = 0$.

Let $k' = \max\{1 \le j \le k-2 : b_j \ne 0\}$. In this case, $a = b_1 \cdots b_{k'-1}(b_{k'}-1)c_1 \cdots c_{k'-k-1}$. Indeed, if $a' \in \mathcal{D}^{k-1}$ such that $b_1 \cdots b_{k'-1}(b_{k'}-1)c_1 \cdots c_{k'-k-1} <_{\text{lex}} a'$, then either $b \le_{\text{lex}} a'$ or a' is not admissible.

Now, suppose $j \leq k'$ such that $b_j \cdots b_{k'-1}(b_{k'}-1)c_1 \cdots c_{k'-k-1}$ is a suffix of a that is also a prefix of c. However,

$$b_j \cdots b_{k'-1} b_{k'} 0^{\infty} <_{\text{lex}} c = b_j \cdots b_{k'-1} (b_{k'} - 1) c_1 c_2 \cdots$$

a contradiction.

Proposition 2.6. Let $a_1 <_{\text{lex}} a_2 <_{\text{lex}} \cdots <_{\text{lex}} a_N$ be consecutive elements of \mathcal{D}_A^{k-1} that are also elements of $E_{k-1,d}$. Then $N \leq K+2$.

Proof. For each j, denote $a_j = a_1^{(j)} \cdots a_{k-1}^{(j)}$. By Lemma 2.5,

$$a_2 = a_1^{(2)} \cdots a_{k_2}^{(2)} \underbrace{00 \cdots 0}_{k-k_2-1}$$
 for some $k_2 < k-1$ where $a_{k_2}^{(2)} \neq 0$

and $a_1 = a_1^{(2)} \cdots a_{k_2-1}^{(2)} (a_{k_2}^{(2)} - 1) c_1 c_2 \cdots$. Hence, the $(k_2 + 1)$ th digit of a_1 is $c_1 \neq 0$. Also,

$$a_3 = a_1^{(3)} \cdots a_{k_3}^{(3)} \underbrace{0 \cdots 0}_{k-k_3-1}$$
 for some $k_3 < k-1$ where $a_{k_3}^{(3)} \neq 0$.

Moreover, by Lemma 2.5, $a_2 = a_1^{(3)} \cdots a_{k_3-1}^{(3)} (a_{k_3} - 1)c_1 \cdots c_{k-k_3-1}$. So the $(k_3 + 1)$ th digit of a_2 is $c_1 \neq 0$ which implies $k_2 \geq k_3 + 1 > k_3$. Letting $k_j = \max\{1 \leq \ell \leq k - 1 : a_{\ell}^{(j)} \neq 0\}$ for each j yields a sequence

$$k-1 \ge k_1 > k_2 > \cdots > k_N.$$

So $k_N \leq k - N$.

Moreover, a_N ends with a zero block of length $k - k_N - 1$. Since K is the maximum length of zero blocks of $d(b - \lfloor b \rfloor)$, then c has at most K + 1 consecutive zeros. By Proposition 2.4, $k - k_N - 1 \leq K + 1$. Thus, $k - K - 2 \leq k_N \leq k - N$. Therefore, $N \leq K + 2$.

Hence, $V_k(b;d)$ is composed of intervals of length at most b^{-k} such that if an interval of $V_k(b;d)$ has length less than b^{-k} , then by Proposition 2.6, there is $j \leq K + 2$ such that the *j*th interval of $V_k(b;d)$ to the left of the current interval has length b^{-k} . Also, by Lemma 2.1, the consecutive centers of the intervals of $V_k(b;d)$ are at most $b^{-(k-1)}$ units apart.

We have the following result.

Theorem 2.7. Suppose $K < \infty$. Let $d \in \mathcal{D}$ such that $d \leq d'$ where d' is the minimal digit of $\lim_{\varepsilon \to 0^+} d(1 - \varepsilon)$ and d(x) is the *b*-expansion of $x \in [0, 1)$.

1. Let $\alpha, \beta \in (0, 1)$ such that $\log_b(\alpha\beta) \notin \mathbb{Q}$. Then $C_b[d]$ is (α, β) -winning if $\beta > A_b(\alpha)$ where

$$A_b(\alpha) := \frac{(2(Kb+2b)+1)\alpha - 1}{\alpha[(4(Kb+2b)-1) - \alpha(2(Kb+2b) - 1)]}.$$
(2.1)

2. Suppose that

$$\alpha < \frac{1}{2Kb^2 + 4b^2 + 1}$$

Then $C_b[d]$ is α -winning.

Remark. Denoting by γ_1 and γ_2 the numerator and denominator of the right-hand side of (2.1), respectively, we see that $\gamma_2 > 0$ and $\gamma_2 > \gamma_1$. In particular, we have $A_b(\alpha) < 1$. Note that Theorem 2.7 is flexible. In fact, if $\alpha < (2Kb + 4b + 1)^{-1}$, then $C_b[0]$ is (α, β) -winning for any $\beta \in (0, 1)$ with $\log_b(\alpha\beta) \notin \mathbb{Q}$.



Figure 1: Centers $a_{n+1}, b_{n+1}, a_{n+2}, b_{n+2}$ of the balls $A_{n+1}, B_{n+1}, A_{n+2}$ and B_{n+2}

Proof. (a) Suppose $\beta > A_b(\alpha)$. Observe that

$$2(Kb+2b)\alpha - \frac{4(Kb+2b)\alpha\beta(1-\alpha)}{1-\alpha\beta} < 1-\alpha$$

$$\iff 2(Kb+2b)\alpha + 2(Kb+2b)\alpha^2\beta - 4(Kb+2b)\alpha\beta < 1-\alpha\beta - \alpha + \alpha^2\beta$$

$$\iff 2(Kb+2b)\alpha - 1 + \alpha < \beta(-2(Kb+2b)\alpha^2 + 4(Kb+2b)\alpha - \alpha + \alpha^2)$$

$$\iff \beta > A_b(\alpha).$$

Since $\log_b(\alpha\beta)$ is irrational, there exist $n, k \in \mathbb{N}$ such that

$$2(Kb+2b)\alpha - \frac{4(Kb+2b)\alpha\beta(1-\alpha)}{1-\alpha\beta} < \frac{K+2}{\rho(\alpha\beta)^n b^{k-1}} < 1-\alpha$$

$$(2.2)$$

where $\rho > 0$ is the initial radius of the (α, β) -game. Hence,

(1)
$$\frac{K+2}{b^{k-1}} < \rho(\alpha\beta)^n (1-\alpha)$$

(2)
$$2\rho(\alpha\beta)^n \alpha - \frac{4\rho(\alpha\beta)^{n+1}(1-\alpha)}{1-\alpha\beta} < \frac{1}{b^k}.$$

For $j \in \mathbb{N}$, denote the center of Alice's and Bob's *j*-th balls A_j and B_j by a_j and b_j , respectively. For the center a_{n+1} , Alice must choose so that $|a_{n+1} - b_n| \leq \rho(\alpha\beta)^n(1-\alpha)$. Let *a* be the center of an interval of $V_k(b;d)$ that is closest to b_n . If such interval is of length b^{-k} , Alice chooses $a_{n+1} = a$. In this case, $|a_{n+1} - b_n| \leq b^{-(k-1)}$. If such interval is of length less than b^{-k} , Alice chooses $a_{n+1} = a'$, where *a'* is the center of an interval of $V_k(b;d)$ nearest to *a* such that the length of the interval of $V_k(b;d)$ centered at *a'* is b^{-k} . In this case, $|a_{n+1} - b_n| \leq (K+2)b^{-(k-1)}$ by Proposition 2.6. Hence, in any case, $|a_{n+1} - b_n| \leq (b^K + 1)b^{-(k-1)} < \rho(\alpha\beta)^n(1-\alpha)$ by (1).

Alice wins if she can force the situation where $A_{n+m} \subseteq V_k(b;d)$ for some $m \in \mathbb{N}$. For avoiding such a situation, Bob's best strategy is to move away as far as possible from a_{n+1} at each turn. See Figure 1. We may assume that Bob chooses the center b_{n+m} of the ball B_{n+m} to be farthest from the right of a_{n+m} , i.e.,

$$b_{n+m} = a_{n+m} + \rho \alpha (\alpha \beta)^{n+m-1} (1-\beta).$$

To counter, Alice chooses a_{n+m+1} to be the leftmost possible center, that is, the center of the ball A_{n+m+1} is

$$a_{n+m+1} = b_{n+m} - \rho(\alpha\beta)^{n+m}(1-\alpha)$$

= $a_{n+m} + \rho(\alpha\beta)^{n+m-1}[\alpha - 2\alpha\beta + \alpha^2\beta].$

Alice and Bob play optimally in this manner. We have

$$b_{n+1} = a_{n+1} + \rho \alpha (\alpha \beta)^n (1 - \beta)$$

$$a_{n+2} = a_{n+1} + \rho (\alpha \beta)^n (\alpha - 2\alpha \beta + \alpha^2 \beta)$$

$$b_{n+2} = a_{n+1} + \rho (\alpha \beta)^n [\alpha - 2\alpha \beta + 2\alpha^2 \beta - \alpha^2 \beta^2]$$

$$a_{n+3} = a_{n+1} + \rho (\alpha \beta)^n [\alpha - 2\alpha \beta + 2\alpha^2 \beta - 2\alpha^2 \beta^2 + \alpha^3 \beta^2]$$

In general,

$$a_{n+m+1} = a_{n+1} + \rho(\alpha\beta)^n \left[\alpha - 2(1-\alpha) \sum_{j=1}^m (\alpha\beta)^j - \alpha(\alpha\beta)^m \right]$$

So, the farthest point c_{n+m+1} of A_{n+m+1} from a_{n+1} is its rightmost point, i.e.,

$$c_{n+m+1} := a_{n+m+1} + \alpha \rho(\alpha \beta)^{n+m} = a_{n+1} + \rho(\alpha \beta)^n \left[\alpha - 2(1-\alpha) \sum_{j=1}^m (\alpha \beta)^j \right].$$

Computing for the distance between c_{n+m+1} and a_{n+1} , we have

$$|c_{n+m+1} - a_{n+1}| = \rho(\alpha\beta)^n \alpha - \frac{2\rho(\alpha\beta)^{n+1}(1-\alpha)[1-(\alpha\beta)^m]}{1-\alpha\beta}$$

As $m \to \infty$, the distance $|c_{n+m+1} - a_{n+1}|$ decreases towards

$$\rho(\alpha\beta)^n \alpha - \frac{2\rho(\alpha\beta)^{n+1}(1-\alpha)}{1-\alpha\beta}$$

By (2), there exists $m \in \mathbb{N}$ such that $|c_{n+m+1} - a| < 1/(2b^k)$, which implies that A_{n+m+1} is contained in an interval of V_k . Therefore, $A_{n+m+1} \subseteq V_k(b; d)$.

(b) It suffices to show that there exists $n, k \in \mathbb{N}$ satisfying (2.2). By the remark after Theorem 2.7, we may assume that

$$\frac{\log(\alpha^{-1}\beta^{-1})}{\log b} = \frac{t_2}{t_1},$$

where t_1, t_2 are coprime positive integers. Putting $\theta := b^{1/t_1}$, we have $\alpha\beta = \theta^{-t_2}$ and $b = \theta^{t_1}$. Since t_1, t_2 are coprime, we see that

$$\{(\alpha\beta)^n b^{k-1} \mid n, k \in \mathbb{N}\} = \{\theta^m \mid m \in \mathbb{Z}\}.$$

By assumption, we have $1 - \alpha > 2b(Kb + 2b)\alpha$. From $b \ge \theta$, we get that

$$2(Kb+2b)\alpha - \frac{4(Kb+2b)\alpha\beta(1-\alpha)}{1-\alpha\beta} < \theta(1-\alpha).$$

Therefore, there exist $n, k \in \mathbb{N}$ satisfying (2.2).



Figure 2: Graph of $\beta = A_b(\alpha)$ for $b = \phi_1$ (blue), ϕ_2 (red), ϕ_{10} (green). As $b \to \infty$, the graph of $\beta = A_b(\alpha)$ tends to $\beta = 1/(2 - \alpha)$ (black).

3 Complex expansions

We begin this section by reviewing some literature on the dynamics of the digits of rotational beta expansions in dimension 2. In [3, Proposition 4.1], a criterion that distinguishes admissible blocks from non-admissible ones was provided for a family of four-fold rotational beta expansions in \mathbb{R}^2 by relating the expansions to the so-called *beta Cantor series expansions*. In [1], necessary and sufficient conditions for the symbolic dynamical system associated to a rotational beta transformation to be sofic were given. In [11], an admissibility criterion for a family of rotational beta expansions in dimension 2 called the *zeta expansions* was given. In these zeta expansions, the digits are all rational integers. In general, determining the admissible blocks of rotational beta expansions in dimension 2 is a nontrivial problem.

In this section, we consider an expansion on \mathbb{C} with complex base ξ such that $|\xi| > 1$. Let \mathcal{L} be a lattice in \mathbb{C} and $\mathcal{X} \subseteq \mathbb{C}$ be a fundamental domain for \mathcal{L} . Then an analog definition for $(\xi, \mathcal{L}, \mathcal{X})$ -expansion (or simply ξ -expansion) is obtained from its corresponding rotational beta expansion.

Example 3.1. Consider $\xi = (1 + \sqrt{5})\hat{i}/2$ and the lattice $\mathcal{L} = \mathbb{Z}[\hat{i}]$ in \mathbb{C} with fundamental domain $\mathcal{X} = \{a + b\hat{i} : a, b \in [0, 1)\}$. Let $z = (5 - \sqrt{5})\hat{i}/10 \in \mathcal{X}$. Then $\xi z = -\sqrt{5}/5$ and $-\sqrt{5}/5 + 1 \in \mathcal{X}$. Hence, the first digit of z is -1 and $\mathbb{T}_{\xi,\mathcal{L},\mathcal{X}}(z) = (5 - \sqrt{5})/5$. Next, $\xi(\mathbb{T}_{\xi,\mathcal{L},\mathcal{X}}(z)) = 2\sqrt{5}\hat{i}/5 \in \mathcal{X}$. Thus, $\mathbb{T}^2_{\xi,\mathcal{L},\mathcal{X}}(z) = 2\sqrt{5}\hat{i}/5$ and the second digit of z is 0. Finally, $\xi(\mathbb{T}^2_{\xi,\mathcal{L},\mathcal{X}}(z)) + 2 = \mathbb{T}_{\xi,\mathcal{L},\mathcal{X}}(z)$. This means that the third digit of z is -2. Therefore, the $(\xi,\mathcal{L},\mathcal{X})$ -expansion of z is the periodic expansion $d(z) = (-1)\overline{0(-2)}$. Indeed,

$$z = \frac{-1}{\xi} + \frac{-2}{\xi^3} + \frac{-2}{\xi^5} + \frac{-2}{\xi^7} + \cdots$$

Example 3.2. Let $\xi = re^{\hat{i}\theta}$ where r > 1 and $\theta \in [0, 2\pi)$. Let $\mathcal{L} = \mathbb{Z}[\hat{i}]$ and $\mathcal{X} = \{a + b\hat{i} : -1/2 \le a, b < 1/2\}$. Let $z = x + y\hat{i} \in \mathcal{X}$. Then the first digit d(z) of the $(\xi, \mathcal{L}, \mathcal{X})$ -expansion of z is

$$d(z) = \left\lfloor \operatorname{Re}(\xi z) + \frac{1}{2} \right\rfloor + \left\lfloor \operatorname{Im}(\xi z) + \frac{1}{2} \right\rfloor \hat{\imath}$$

$$= \left\lfloor rx\cos(\theta) - ry\sin(\theta) + \frac{1}{2} \right\rfloor + \left\lfloor rx\sin(\theta) + ry\cos(\theta) + \frac{1}{2} \right\rfloor \hat{\imath}$$

where $\operatorname{Re}(\zeta)$ and $\operatorname{Im}(\zeta)$ are the real and imaginary parts of $\zeta \in \mathbb{C}$, respectively.

Analogous to the one-dimensional case, for d in the digit set \mathcal{D} and $k \in \mathbb{N}$, we let $V_k(\xi, d) := \{z \in \mathcal{X} : d_k(z) = d\}$ and

$$C_{\xi}[d] := \{z \in \mathcal{X} : d \text{ appears in } d(z)\} = \bigcup_{k=1}^{\infty} V_k(\xi, d).$$

We aim to give sufficient conditions so that the set $C_{\xi}[0]$ is (α, β) -winning.

3.1 Property (C_k)

When $0 \in \mathcal{D}$, we consider the scenario where $V_k(\xi; 0)$ can be expressed as union of translates of "scaled-down" copies of \mathcal{X} . This leads to a sufficient condition for which $C_{\xi}[0]$ is winning (see Theorem 3.12). To proceed, for $k \in \mathbb{N}$, we consider the property (C_k) for a ξ -expansion.

Definition 3.3. Let $k \in \mathbb{N}$. We say that the ξ -expansion has property (C_k) if

$$\xi^{-k}\mathcal{X} + \xi^{-(k-1)}a_{k-1} + \dots + \xi^{-1}a_1 \subseteq \mathcal{X}$$

for any admissible block $a_1 a_2 \cdots a_{k-1} \in \mathcal{D}^{k-1}$ with $(C_1) : \xi^{-1} \mathcal{X} \subseteq \mathcal{X}$.

We have the following result.

Proposition 3.4. Let $k \in \mathbb{N}$ and $a = a_1 a_2 \cdots a_k \in \mathcal{D}^k$ be an admissible block. If (C_n) holds for $n \in \{1, 2, \ldots, k+1\}$, then (a, 0) is admissible. Moreover, $V_{k+1}(\xi; 0)$ can be partitioned into disjoint squares as

$$V_{k+1}(\xi;0) = \bigcup_{a_1 a_2 \cdots a_k \in \mathcal{D}^k \text{ is admissible}} \xi^{-(k+1)} \mathcal{X} + \sum_{j=1}^k \xi^{-j} a_j.$$

Proof. We show the case when k = 1. Suppose (C_1) and (C_2) hold. By (C_1) , we have $\mathcal{X} \supseteq \xi^{-1} \mathcal{X} \supseteq \xi^{-2} \mathcal{X} \supseteq \cdots$. Let $a \in \mathcal{D}$. The set of points in \mathcal{X} with first digit a and second digit 0 is $S = (\xi^{-2} \mathcal{X} + \xi^{-1}a) \cap (\xi^{-1} \mathcal{X} + \xi^{-1}a) \cap \mathcal{X}$. By (C_1) ,

$$\xi^{-2}\mathcal{X} + \xi^{-1}a = \xi^{-1}(\xi^{-1}\mathcal{X}) + \xi^{-1}a \subseteq \xi^{-1}\mathcal{X} + \xi^{-1}a.$$

Also, by (C_2) , $\xi^{-2}\mathcal{X} + \xi^{-1}a \subseteq \mathcal{X}$. Therefore, $S = \xi^{-2}\mathcal{X} + \xi^{-1}a \neq \emptyset$. Hence, the block (a, 0) is admissible and

$$V_2(\xi;0) = \bigcup_{a \in \mathcal{D}} \xi^{-2} \mathcal{X} + \xi^{-1} a$$

The cases $k \geq 2$ are similar.

For the remainder of Section 3, we only consider $(\xi, \mathcal{L}, \mathcal{X})$ -expansions where $\mathcal{L} = \mathbb{Z}[\hat{i}]$ and $\mathcal{X} = \{a + b\hat{i} \mid a, b \in [-1/2, 1/2)\}$. For $\theta \in [0, 2\pi]$, define

$$c(\theta) := \max\{|\cos(\theta)|, |\sin(\theta)|\}$$

$$s(\theta) := \min\{|\cos(\theta)|, |\sin(\theta)|\}.$$

Note that θ and $\theta + \pi k/2$ ($k \in \mathbb{Z}$) produce the same results except for a few pathological cases (see remark after Theorem 3.11). Moreover, without loss of generality, we restrict $\theta \in [0, \pi/4]$ as we can switch the roles of $\cos(\theta)$ and $\sin(\theta)$ for $\theta \in (\pi/4, \pi/2]$. Thus, $c(\theta) = \cos(\theta)$ and $s(\theta) = \sin(\theta)$.

3.2 Square digit sets

Let $\xi = re^{\hat{i}\theta}$ where $\theta \in [0, \pi/4]$ and r > 1. Let $c = \cos(\theta)$ and $s = \sin(\theta)$. Let $\mathcal{L} = \mathbb{Z}[\hat{i}]$ and $\mathcal{X} = \{a + b\hat{i} : a, b \in [-1/2, 1/2)\}$ Note that the digit set \mathcal{D} is the smallest subset \mathcal{D}' of \mathcal{L} for which $\bigcup_{d \in \mathcal{D}'} (\overline{\mathcal{X}} + d)$ covers $\xi \overline{\mathcal{X}}$. Let $\mathbb{D} = \mathbb{D}_{(\xi, \mathcal{L}, \mathcal{X})} := \bigcup_{d \in \mathcal{D}} (\overline{\mathcal{X}} + d)$. Then \mathbb{D} is a contiguous

collection of squares that covers $\xi \overline{\mathcal{X}}$. We consider the simplest case where \mathbb{D} is a rectangle.

Proposition 3.5. If \mathbb{D} is a rectangle, then \mathbb{D} is a square.

Proof. Suppose \mathbb{D} is a rectangle. Since $0 \leq \theta \leq \pi/4$, it follows that ξP and P lie on the same quadrant in \mathbb{R}^2 if P is a corner point of $\overline{\mathcal{X}}$. Moreover, $\xi(1+\hat{\imath})/2, \xi(1-\hat{\imath})/2, \xi(-1+\hat{\imath})/2$ and $\xi(-1-\hat{\imath})/2$ are the topmost, rightmost, leftmost, and bottommost points of $\xi \overline{\mathcal{X}}$, respectively. Hence, for any admissible digit $a + b\hat{\imath}$ with $a, b \in \mathbb{Z}$, we have

$$\left\lfloor \frac{1 - r(c+s)}{2} \right\rfloor \le a, b \le \left\lfloor \frac{1 + r(c+s)}{2} \right\rfloor =: N.$$

Then

$$N \le \frac{1 + r(c+s)}{2} < N+1$$

-N < $\frac{1 - r(c+s)}{2} \le -N+1.$

Note that (1 - rc - rs)/2 = -N + 1 if and only if r(c + s) = 2N - 1. If r(c + s) is not an odd integer, then

$$\left\lfloor \frac{1 - r(c+s)}{2} \right\rfloor = -N.$$

In this case, $\mathcal{D} = \{a + b\hat{i} : a, b \in \mathbb{Z} \text{ such that } |a|, |b| \leq N\}$ and therefore, \mathbb{D} is a square. On the other hand, suppose r(c + s) = 2N - 1. Let $z = x + \hat{i}y \in \mathcal{X}$ be arbitrarily close to $(1 + \hat{i})/2$. Then $\text{Im}(d_1(z)) = \lfloor r(sx + cy) + 1/2 \rfloor$. However,

$$r(sx + cy) + \frac{1}{2} < \frac{r(c+s) + 1}{2} = N.$$

Hence,

$$\max_{d \in \mathcal{D}} \operatorname{Re}(d) = -\min_{d \in \mathcal{D}} \operatorname{Re}(d) = N - 1$$

$$\max_{d \in \mathcal{D}} \operatorname{Im}(d) = -\min_{d \in \mathcal{D}} \operatorname{Im}(d) = N - 1.$$

Thus, there is no admissible digit whose imaginary part is N. In other words, the corner points of $\xi \overline{\mathcal{X}}$ are on the boundaries (sides) of some adjacent translates of $\overline{\mathcal{X}}$. Hence, $\max_{d \in \mathcal{D}} |\operatorname{Re}(d)| = \max_{d \in \mathcal{D}} |\operatorname{Im}(d)| = N - 1$. See Figure 3.



Figure 3: $(re^{\hat{\imath}\theta})\overline{\mathcal{X}}$ inscribed in \mathbb{D} where $(r,\theta,N) = \left(\frac{5}{c+s} - \frac{1}{2}, \frac{\pi}{20}, 2\right)$ (left) and $(r,\theta,N) = \left(\frac{5}{c+s}, \frac{\pi}{20}, 3\right)$ (right)

We say that ξ has a square digit set of size $N \in \mathbb{N}$ if

$$\mathcal{D} = \{ a + b\hat{i} \mid a, b \in \mathbb{Z} \text{ such that } |a|, |b| \le N \}.$$

Note that, if $\xi = re^{i\theta}$ has a square digit set of size N, then

$$N = \left\lceil \frac{r(c+s)+1}{2} \right\rceil - 1.$$

The following result provides a criterion to determine if ξ has a square digit set.

Proposition 3.6. The base ξ has a square digit set of size $N \in \mathbb{N}$ if and only if

$$(2N-1)(c+s) < r \le \frac{2N+1}{c+s} =: u_N(\theta).$$

Proof. Note that ξ has a square digit set of size N if and only if $|\operatorname{Re}(a)|, |\operatorname{Im}(a)| \leq N$ for all $a \in \mathcal{D}$ and $N + N\hat{i} \in \mathcal{D}$. In fact, if $N + N\hat{i} \in \mathcal{D}$, then for any $\varepsilon_1, \varepsilon_2 \in \{1, -1\}$, we have $\varepsilon_1 N + \varepsilon_2 N\hat{i} \in \mathcal{D}$ by the proof of Proposition 3.5. We have

$$|\operatorname{Re}(a)|, |\operatorname{Im}(a)| \le N \iff \left\lceil \frac{r(c+s)+1}{2} \right\rceil - 1 \le N \iff r \le u_N(\theta).$$

For the rest of the proof, we assume that $r \leq u_N(\theta)$. It remains to show that $N + N\hat{i} \in \mathcal{D}$ if and only if r > (2N - 1)(c + s).

Now, $N + N\hat{i} \in \mathcal{D}$ if and only if there exist $x, y \in [-1/2, 1/2)$ such that

$$\left\lfloor rcx - rsy + \frac{1}{2} \right\rfloor = \left\lfloor rsx + rcy + \frac{1}{2} \right\rfloor = N.$$

Equivalently, (x, y) lies on the strips S_1 and S_2 given by

$$S_{1}: \frac{c}{s}x - \frac{2N+1}{2rs} < y \le \frac{c}{s}x - \frac{2N-1}{2rs}$$
$$S_{2}: \frac{2N-1}{2rc} - \frac{s}{c}x \le y < \frac{2N+1}{2rc} - \frac{s}{c}x.$$

Define the boundary lines of S_1 and S_2 as follows:

$$\ell_1: y = \frac{c}{s}x - \frac{2N+1}{2rs}$$
 and $\ell_2: y = \frac{c}{s}x - \frac{2N-1}{2rs}$
 $\ell_3: y = \frac{2N-1}{2rc} - \frac{s}{c}x$ and $\ell_4: y = \frac{2N+1}{2rc} - \frac{s}{c}x$.

Then $N + N\hat{i} \in \mathcal{D}$ if and only if the interior of $S_1 \cap S_2 \cap \overline{\mathcal{X}}$ is non-empty. Indeed, if $x + y\hat{i}$ is an interior point of $S_1 \cap S_2 \cap \overline{\mathcal{X}}$, then the first digit of $x + y\hat{i}$ is $N + N\hat{i}$.

Assuming $r \leq u_N(\theta)$, we show that the interior of $S_1 \cap S_2 \cap \overline{\mathcal{X}}$ is non-empty if and only if (2N-1)(c+s) < r. We divide the proof into 3 cases. When $\theta = \pi/4$, we only consider **CASES 2** and **3** as **CASE 1** is not applicable.

CASE 1: Suppose r > (2N-1)/(c-s). Let $\varepsilon > 0$ be small so that $z = (1-\varepsilon)(1+i)/2 \in \mathcal{X}$ is arbitrarily close to (1+i)/2. If $d_1(z) = a + bi$, then

$$a = \left\lfloor \frac{(1-\varepsilon)(rc-rs)+1}{2} \right\rfloor$$
 and $b = \left\lfloor \frac{(1-\varepsilon)(rc+rs)+1}{2} \right\rfloor$

Moreover, $a \leq b \leq N$. For some t > 0,

$$r = \frac{2N - 1 + t}{c - s}.$$

Choosing ε sufficiently small, we have

$$\frac{(1-\varepsilon)(rc-rs)+1}{2} = \frac{(1-\varepsilon)(2N-1+t)+1}{2} = N + \frac{t-(2N-1+t)\varepsilon}{2} > N.$$

Hence, $N \leq a \leq b \leq N$. Thus, a = b = N. This means that $N + N\hat{i} \in \mathcal{D}$.

CASE 2: Suppose $(2N-1)/(c+s) < r \le (2N-1)/(c-s)$. Then the corner point $(1+\hat{i})/2$ of $\overline{\mathcal{X}}$ is on the strip S_2 . Moreover, the corner point $(-1+\hat{i})/2$ of $\overline{\mathcal{X}}$ is below or on ℓ_3 since $r \le (2N-1)/(c-s)$. Likewise, $(1-\hat{i})/2$ is below ℓ_3 .

Let A and B be the intersection points of the line ℓ_3 with the line y = 1/2 and the line x = 1/2, respectively. Then

$$A = \frac{2N - 1 - rc}{2rs} + \frac{1}{2}\hat{\imath}$$
 and $B = \frac{1}{2} + \frac{2N - 1 - rs}{2rc}\hat{\imath}.$

Note that $S_1 \cap S_2 \cap \overline{\mathcal{X}}$ has an interior point if and only if A is above ℓ_1 and B is below ℓ_2 . See Figure 4.



Figure 4: Position of A and of B relative to the strips S_1 and S_2

Observe that

A is above
$$\ell_1 \iff \frac{c}{s} \cdot \frac{2N - 1 - rc}{2rs} - \frac{2N + 1}{2rs} \le \frac{1}{2} \iff r \ge 2N(c - s) - (c + s).$$

Since $0 \le \theta \le \pi/4$, we have

$$2N(c-s) - (c+s) = \frac{2N(c^2 - s^2) - (c+s)^2}{c+s} \le \frac{2N - (1+2cs)}{c+s} \le \frac{2N - 1}{c+s} < r.$$

So, A is always above ℓ_1 . It remains to determine when B is below ℓ_2 . We have

$$B \text{ is below } \ell_2 \iff \frac{2N - 1 - rs}{2rc} < \frac{c}{2s} - \frac{2N - 1}{2rs}$$
$$\iff (2N - 1)(c + s) < r.$$

Therefore, $S_1 \cap S_2 \cap \overline{\mathcal{X}}$ has an interior point if and only if r > (2N-1)(c+s).

CASE 3: Suppose $1 < r \le (2N - 1)/(c + s)$. Observe that

$$r \leq \frac{2N-1}{c+s} \Longleftrightarrow \frac{c+s}{2c} \leq \frac{2N-1}{2rc} \Longleftrightarrow \frac{1}{2} \leq \frac{2N-1}{rc} - \frac{1}{2} \cdot \frac{s}{c}.$$

This means that the corner point $(1 + \hat{\imath})/2$ of $\overline{\mathcal{X}}$ is below or on ℓ_3 (see Figure 5). In this case, $S_2 \cap \overline{\mathcal{X}}$, and consequently, $S_1 \cap S_2 \cap \overline{\mathcal{X}}$, has no interior point.



Figure 5: $(1+\hat{\imath})/2$ is below or on ℓ_3

Given $\theta \in [0, \pi/4]$, there exists r such that $\xi = re^{i\theta}$ has a square digit set of size $1, 2, \ldots, [(sc+1)/(2sc)] - 1$ since (sc+1)/(2sc) > 1 and

$$(2N-1)(c+s) < \frac{2N+1}{c+s} \iff N < \frac{sc+1}{2sc}.$$

Theorem 3.7. Let $\theta \in [0, \pi/4]$ and $K(\theta) := \lceil (sc+1)/(2sc) \rceil - 1$. Then $\xi = re^{i\theta}$ has a square digit set if and only if

$$r \in \bigcup_{N=1}^{K(\theta)} \left((2N-1)(c+s), u_N(\theta) \right].$$

Condition (C_k) and square digit sets 3.3

Let \mathcal{A} be the set of corner points of $\overline{\mathcal{X}}$. Then $\xi^{-2}P + a$ is a corner point of $\xi^{-2}\overline{\mathcal{X}} + a$ if and only if $P \in \mathcal{A}$ for any $a \in \mathcal{D}$. By convexity of $\overline{\mathcal{X}}$,

$$(C_k) \text{ holds } \iff \xi^{-k}\overline{\mathcal{X}} + \sum_{j=1}^{k-1} \xi^{-j} a_j \subseteq \mathcal{X}$$
$$\iff \xi^{-k}P + \sum_{j=1}^{k-1} \xi^{-j} a_j \subseteq \mathcal{X} \text{ for all } P \in \mathcal{A},$$

where the sums are taken over all admissible blocks $(a_1, a_2, \ldots, a_{k-1}) \in \mathcal{D}^{k-1}$. For $j \in \{1, 2, \ldots, k\}$, let $c^{(j)} := \cos(j\theta)$ and $s^{(j)} := \sin(j\theta)$. For $P \in \mathcal{A}$, the real and imaginary parts of $\xi^{-k}P$ are among the numbers

$$\frac{c^{(k)} + s^{(k)}}{2r^k}, -\frac{c^{(k)} + s^{(k)}}{2r^k}, \frac{c^{(k)} - s^{(k)}}{2r^k}, \text{ and } \frac{s^{(k)} + c^{(k)}}{2r^k}$$

Thus,

$$\max\{\operatorname{Re}(\xi^{-k}P)\} = \max\{\operatorname{Im}(\xi^{-k}P)\} = \frac{|c^{(k)}| + |s^{(k)}|}{2r^k}$$
$$\min\{\operatorname{Re}(\xi^{-k}P)\} = \min\{\operatorname{Im}(\xi^{-k}P)\} = -\frac{|c^{(k)}| + |s^{(k)}|}{2r^k}.$$

Let ξ have a square digit set of size N. If $a = a_1 + a_2 \hat{i} \in \mathcal{D}$, then

$$\xi^{-j}a = \frac{c^{(j)}a_1 + s^{(j)}a_2}{r^j} + \frac{c^{(j)}a_2 - s^{(j)}a_1}{r^j}\hat{\imath}.$$

So,

$$-\frac{N(|c^{(j)}|+|s^{(j)}|)}{r^{j}} \le \operatorname{Re}(\xi^{-j}a), \operatorname{Im}(\xi^{-j}a) \le \frac{N(|c^{(j)}|+|s^{(j)}|)}{r^{j}}.$$

Therefore, the maximum of $\operatorname{Re}\left(\xi^{-k}P + \sum_{j=1}^{n-1}\xi^{-j}a_j\right)$ and $\operatorname{Im}\left(\xi^{-k}P + \sum_{j=1}^{n-1}\xi^{-j}a_j\right)$ for $P \in \mathcal{A}$ and admissible block $(a_1, a_2, \ldots, a_{k-1}) \in \mathcal{D}^{k-1}$ is

$$\frac{|c^{(k)}| + |s^{(k)}|}{2r^k} + N\sum_{j=1}^{k-1} \frac{|c^{(j)}| + |s^{(j)}|}{r^j}$$

while the minimum is

$$-\frac{|c^{(k)}| + |s^{(k)}|}{2r^k} - N\sum_{j=1}^{k-1} \frac{|c^{(j)}| + |s^{(j)}|}{r^j}$$

For example, if $P = (1+\hat{\imath})/2$, then $\operatorname{Re}(\xi^{-2}P + \xi^{-1}(N+N\hat{\imath})) = (c^{(2)} + s^{(2)})/(2r^2) + N(c+N(2r^2))/(2r^2) + N(c+N(2r^2))/(2r^2)$ s)/r.

Let
$$f_N^{(k)}(r) := r^k - 2N \sum_{j=1}^{k-1} r^{k-j} (|c^{(j)}| + |s^{(j)}|) - (|c^{(k)}| + |s^{(k)}|)$$
. Then
 $f_N^{(k)}(r) > 0 \Longrightarrow (C_k).$

Note that $f_N^{(k)}$ has a unique positive root, which we call $v_N^{(k)} := v_N^{(k)}(\theta)$, by the Descartes' rule of signs If k = 2, we have

$$v_N^{(2)} = N(c+s) + \sqrt{N^2(c+s)^2 + c^{(2)} + s^{(2)}}.$$

Then, for ξ with a square digit set of size N, (C_k) holds if $r > v_N^{(k)}$. Observe that $f_N^{(k)}(1) < 0$ and so, $v_N^{(k)} > 1$. Also, if $t = v_N^{(k-1)}$, then

$$\begin{split} f_N^{(k)}(t) &= t^k - 2N \sum_{j=1}^{k-2} t^{k-j} (|c^{(j)}| + |s^{(j)}|) - 2Nt(|c^{(k-1)}| + |s^{(k-1)}|) - |c^{(k)}| - |s^{(k)}| \\ &< t^k - 2N \sum_{j=1}^{k-2} t^{k-j}(|c^{(j)}| + |s^{(j)}|) - 2Nt(|c^{(k-1)}| + |s^{(k-1)}|) \\ &< t^k - 2N \sum_{j=1}^{k-2} t^{k-j}(|c^{(j)}| + |s^{(j)}|) - (|c^{(k-1)}| + |s^{(k-1)}|) = t f_N^{(k-1)}(t) = 0. \end{split}$$

Thus, $\left\{v_N^{(k)}\right\}_{k\geq 2}$ is a strictly increasing sequence. Moreover,

$$v_N^{(k)} > v_N^{(2)} > 2N(c+s) > (2N-1)(c+s)$$

By Theorem 3.6, we have the following result.

Theorem 3.8. Let $\theta \in [0, \pi/4]$ and r > 1. Let $N \in \mathbb{N}$. Then $\xi = re^{i\theta}$ has a square digit set of size N and (C_n) holds for the ξ -expansion for $n \in \{1, 2, \dots, k\}$ if $v_N^{(k)}(\theta) < r < u_N(\theta)$.

For any $N \in \mathbb{N}$, observe that $v_N^{(1)} = c + s < u_N(\theta)$. Hence, (C_1) holds for any $re^{\hat{i}\theta}$ expansion with a square digit set where r > c + s. Since $c + s \le \sqrt{2}$, (C_1) holds whenever $r > \sqrt{2}$. Note that the digit set need not be square for (C_1) to hold since (C_1) does not depend on \mathcal{D} .

Proposition 3.9. Let $\xi = re^{i\theta}$ where $\theta \in [0, \pi/4]$ and r > 1. Then (C_1) holds if and only if $r > \cos(\theta) + \sin(\theta)$. In particular, if ξ has a square digit set, then (C_1) holds.

Fix $\theta \in [0, \pi/4]$. Does there exist $v_N^{(k)} < r < u_N$ such that $\xi = re^{i\theta}$ does not have a square digit set but (C_k) holds? The answer is negative. In other words, if (C_k) , then the digit set of the expansion is necessarily square.

Theorem 3.10. Let $\theta \in [0, \pi/4]$ and $v_N^{(k)}(\theta) < r < u_N(\theta)$. If (C_k) holds for some $k \in \mathbb{N}$ with $k \geq 2$, then $\xi = re^{i\theta}$ has a square digit set.

Proof. We show the proof for k = 2. Let $N = \lceil (rc + rs + 1)/2 \rceil - 1$. Then $r \leq u_N(\theta)$ and $|\operatorname{Re}(d)|, |\operatorname{Im}(d)| \leq N$ for any $d \in \mathcal{D}$.

Suppose $r > v_N^{(2)}$ but $\xi = re^{i\theta}$ does not have a square digit set. Then (C_2) holds. If r > (2N-1)/(c-s), then r > (2N-1)(c+s). By Proposition 3.6, ξ has a square digit set. So, $r \le (2N-1)/(c-s)$.

Let $\varepsilon > 0$ be sufficiently small. Let $x = 1/2 - \varepsilon$ and $y = (rc - 2N + 1)/(2rc) - c\varepsilon/s$. Then $x + y\hat{i} \in \mathcal{X}$ since $r \leq (2N - 1)/(c - s)$. Its first digit is $N + M\hat{i}$ as

$$N = \left\lfloor rcx - rsy + \frac{1}{2} \right\rfloor$$
$$M := \left\lfloor rsy + rcx + \frac{1}{2} \right\rfloor = \left\lfloor \frac{r - 2Nc + c + s}{2s} - \frac{r\varepsilon}{s} \right\rfloor$$

Since $\varepsilon > 0$ is small, we have

$$\frac{r-2Nc+c+s}{2s} \le M+1.$$

So, $r \leq 2Nc + 2Ms - c + s$.

Consider $z = (1 - \varepsilon)(1 + \hat{\imath})/2 \in \mathcal{X}$. By (C_2) , $\xi^{-2}z + \xi^{-1}(N + M\hat{\imath}) \in \mathcal{X}$. Then $\xi^{-2}z + \xi^{-1}(N + M\hat{\imath})$ has real part

$$\frac{c^{(2)} + s^{(2)}}{r^2} \left(\frac{1}{2} - \varepsilon\right) + \frac{Nc + Ms}{r} < \frac{1}{2}.$$

Since $\varepsilon > 0$ is small,

$$\frac{c^{(2)} + s^{(2)}}{2r^2} + \frac{Nc + Ms}{r} \le \frac{1}{2}.$$

Hence, $r^2 - 2(Nc + Ms)r - (c^{(2)} + s^{(2)}) \ge 0$. Then

$$(Nc + Ms) + \sqrt{(Nc + Ms)^2 + c^{(2)} + s^{(2)}} \le r \le 2(Nc + Ms) - (c - s).$$

Note that $M \geq -N$. We have

$$\begin{split} &\sqrt{(Nc + Ms)^2 + c^{(2)} + s^{(2)}} \le (Nc + Ms) - (c - s) \\ \implies c^{(2)} + s^{(2)} \le -2Nc(c - s) - 2Ms(c - s) + 1 - s^{(2)} \\ \implies 0 \le -2Nc(c - s) - 2Ms(c - s) + 1 - c^{(2)} - 2s^{(2)} \\ \implies 0 \le -2N(c - s)(c + s) + 1 - c^{(2)} - 2s^{(2)} = -2Nc^{(2)} + 1 - c^{(2)} - 2s^{(2)} \\ \implies 0 \le -3c^{(2)} - 2s^{(2)} + 1 \le -1 \end{split}$$

We have a contradiction. This completes the proof.

In what follows, we focus on condition (C_2) . Note that when $\theta \in [0, \pi/4]$, we get that (C_2) holds if and only if $v_N^{(2)}(\theta) < r < u_N(\theta)$ since $c^{(2)}, s^{(2)} \ge 0$. Given $\theta \in [0, \pi/4]$, there exists r such that (C_2) holds for the $\xi = re^{i\theta}$ (with square digit set of size $N \in \mathbb{N}$) if and only if

$$v_N^{(2)}(\theta) < u_N(\theta)$$

$$\iff N^2(c+s)^4 + (c^{(2)} + s^{(2)})(c+s)^2 < N^2(1-s^{(2)})^2 + 2N(1-s^{(2)}) + 1$$

$$\iff 0 > F(N) := 4s^{(2)}N^2 - 2(1-s^{(2)})N + [(c^{(2)} + s^{(2)})(c+s)^2 - 1].$$

Observe that 0 > F(x) for some x > 0 if and only if F has positive discriminant, that is,

$$\Delta := 4(1 - s^{(2)})^2 - 16s^{(2)}[(c^{(2)} + s^{(2)})(c + s)^2 - 1] > 0.$$

This happens if and only if $0 \le \theta < \gamma_1$ where $\gamma_1 \approx 0.12988$ is a particular constant. Suppose $0 \le \theta < \gamma_1$. Then there exists r such that $\xi = re^{i\theta}$ has a square digit set of size $N \in \mathbb{N}$ and (C_2) holds if and only if $L_- < N < L_+$ where

$$L_{\pm} := \frac{2(1 - s^{(2)}) \pm \sqrt{\Delta}}{8s^{(2)}}$$

are the roots of F. Note that $L_{-} < 1$. Also, $L_{+} > 1$ if and only if $0 \le \theta < \gamma_{2}$ where $0.1249 \approx \gamma_{2} = 2 \tan^{-1}(\delta)$ and $\delta \approx 0.625$ is the smallest positive root of $x^{8} + 16x^{7} + 30x^{4} - 16x + 1$. Therefore, if $0 < \theta < \gamma_{2}$, then there exists r such that $\xi = re^{i\theta}$ has a square digit set of size $N \in \mathbb{N}$ and (C_{2}) holds if and only if $N \in \{1, 2, \ldots, \lceil L_{+} \rceil - 1\}$.

We also have that if $\theta = 0$, then $v_N^{(2)}(\theta) = N + \sqrt{N^2 + 1} < 2N + 1 = u_N(\theta)$. Thus, for any $N \in \mathbb{N}$, there exists r such that $\xi = re^{i\theta}$ has a square digit set and (C_2) holds.

We have the following result.

Theorem 3.11. Let $\theta \in [0, \pi/4)$. Let

$$\mathscr{G}(\theta) := \left\{ r > 1 \mid (C_2) \text{ holds for } \xi = r e^{\hat{\imath} \theta} \right\}.$$

1. If $\theta \in [\gamma_2, \pi/4)$, then $\mathscr{G}(\theta) = \varnothing$.

2. If
$$\theta \in (0, \gamma_2)$$
, then $\mathscr{G}(\theta) = \bigcup_{N=1}^{\lceil L_+ \rceil - 1} (v(N, \theta), u(N, \theta)) \}.$

3. If
$$\theta = 0$$
, then $\mathscr{G}(\theta) = \bigcup_{N=1}^{\infty} (N + \sqrt{N^2 + 1}, 2N + 1].$

Remark. If $\theta \in [0, \pi/2)$ and $\pi/2 \leq \theta' \leq 2\pi$ such that $\theta' = \theta + k\pi/2$ for some $k \in \mathbb{Z}$, we obtain parallel results for $\xi = re^{i\theta}$ and $\xi' = re^{i\theta'}$, provided their digit sets are the same, since since $\xi \overline{\mathcal{X}} = \xi' \overline{\mathcal{X}}$. Note that the digit sets are different if and only if $r = u_N(\theta)$ and $\theta' \in [\pi/2, 3\pi/2)$. In such a case, we take $N = \lfloor (rc+rs+1)/2 \rfloor$, instead of $\lceil (rc+rs+1)/2 \rceil - 1$.

3.4 Schmidt game on $C_{\xi}[0]$

Let $\xi = re^{i\theta}$ and $N, k \in \mathbb{N}$ with $k \geq 2$ such that $v_N^{(k)}(\theta) \leq r \leq u_N(\theta)$. Then ξ has a square digit set and (C_n) holds for $n \in \{1, 2, \ldots, k\}$. Recall from Proposition 3.4 that if $(a_1, \ldots, a_{k-1}) \in \mathcal{D}^{(k-1)}$ is an admissible sequence, then $(a_1, \ldots, a_{k-1}, 0) \in \mathcal{D}^k$ is also admissible.

We introduce a notion of consecutive points in $\mathcal{D} = \{a + b\hat{i} : a, b \in \mathbb{Z}, |a|, |b| \leq N\}$. Write the $(2N+1)^2$ digits as

$$w_{s(2N+1)+t} = (-1)^{s+1}(N+1-t) + \hat{i}(-N+s),$$

where $s \in \{0, 1, ..., 2N\}, t \in \{1, 2, ..., 2N + 1\}$. Then $|w_{\ell+1} - w_{\ell}| = 1$ for any $\ell \in \{1, 2, ..., (2N + 1)^2 - 1\}$. See Figure 6.



Figure 6: A snake-like ordering on square digit sets of sizes N = 1 and N = 2

This defines a lexicographic ordering for \mathcal{D}^k , which, in turns, gives an ordering of points of the form

$$a_1\xi^{-1} + a_2\xi^{-2} + \dots + a_k\xi^{-k}$$

This translates to an order of the centers of the squares that make up V_k . Moreover, the distance between consecutive centers with respect to this order is at most $\sqrt{2}/r^{k-1}$. In summary, by Proposition 3.4, $V_k(\xi; 0)$ is composed of (rotated) squares with

- area $1/r^k$
- radius of biggest circle contained in a square is $1/(2r^k)$
- if the first k-1 digits coincide, gaps of consecutive centers are at most $\sqrt{2}/r^{k-1}$ wrt to the given order.

We have the following analog of Theorem 2.7.

Theorem 3.12. Let $\xi = re^{i\theta}$, $\mathcal{L} = \mathbb{Z}[\hat{i}]$ and $\mathcal{X} \cong [-1/2, 1/2)^2$. Let $N, k \in \mathbb{N}$ and suppose $v_N^{(k)}(\theta) < r < u_N^{(k)}(\theta)$. Let $0 < \alpha, \beta < 1$ and $\rho > 0$. Suppose

$$\beta > F_r(\alpha) := \frac{(2\sqrt{2r+1})\alpha - 1}{\alpha[(1 - 2\sqrt{2r})\alpha + (4\sqrt{2r-1})]}$$

If, either

1. $(2 - \alpha)\beta < 1$ and ρ satisfies

$$\frac{\log_r \rho + \log_r \left(2\alpha - \frac{4\alpha\beta(1-\alpha)}{1-\alpha\beta}\right) + k}{\log_r (\alpha^{-1}\beta^{-1})} \le n < \frac{\log_r \rho + \log_r \left(\frac{1-\alpha}{\sqrt{2}r}\right) + k}{\log_r (\alpha^{-1}\beta^{-1})}$$
(3.1)

for some $n \in \mathbb{N}$; or,

2. $(2-\alpha)\beta \ge 1$ and ρ is sufficiently large such that

$$1 < \frac{\log_r \rho + \log_r \left(\frac{1-\alpha}{\sqrt{2r}}\right) + k}{\log_r \left(\alpha^{-1}\beta^{-1}\right)},\tag{3.2}$$

then

 $C_{\xi}[0] := \{ z \in \mathcal{X} : \text{ the } \xi \text{-expansion of } z \text{ has digit } 0 \}$

is (α, β, ρ) -winning.

Proof. Note that $\beta > F_r(\alpha)$ if and only if

$$2\alpha - \frac{4\alpha\beta(1-\alpha)}{1-\alpha\beta} < \frac{1-\alpha}{\sqrt{2}r}.$$

Suppose that $\beta > F_r(\alpha)$ and $(2 - \alpha)\beta < 1$. Then (3.1) holds if and only if

$$2\alpha - \frac{4\alpha\beta(1-\alpha)}{1-\alpha\beta} \le \frac{1}{\rho(\alpha\beta)^n r^k} < \frac{1-\alpha}{\sqrt{2}r}.$$
(3.3)

Meanwhile, if $(2 - \alpha)\beta \ge 1$, then (3.3) with n = 1 follows from the fact that $(2 - \alpha)\beta \ge 1$ if and only if

$$2\alpha - \frac{4\alpha\beta(1-\alpha)}{1-\alpha\beta} \le 0$$

for any positive real number $\alpha, \beta < 1$. It follows that

(1)
$$\frac{\sqrt{2}}{r^{k-1}} < \rho(\alpha\beta)^n (1-\alpha)$$

(2)
$$\rho(\alpha\beta)^n \alpha - \frac{2\rho(\alpha\beta)^{n+1}(1-\alpha)}{1-\alpha\beta} \le \frac{1}{2r^k}.$$

By (1), Alice can choose a_{n+1} to be the center of one of the translates of \mathcal{X} that make up $V_k(\xi; 0)$. Bob's best strategy is to move away from a_{n+1} along a common direction, that is, for some unit vector \vec{v} .

$$b_{n+m} = a_{n+m} + \rho \alpha (\alpha \beta)^{n+m-1} (1-\beta) \vec{v}.$$

Then Alice responds with

$$a_{n+m+1} = b_{n+m} - \rho(\alpha\beta)^{n+m} (1-\alpha)\vec{v}.$$

Similar to the proof of Theorem 2.7, the outcome ω is at most $1/(2r^k)$ away from a_{n+1} . This means that $\omega \in V_k(\xi; 0) \subseteq C_{\xi}[0]$.

Remark. Let $N, k \in \mathbb{N}$ satisfy $v_N^{(k)}(\theta) < r < u_N^{(k)}(\theta)$. For any $0 < \alpha, \beta < 1$ with $\beta > F_r(\alpha)$ and $(2 - \alpha)\beta < 1$, we see that there exists $\rho > 0$ such that (3.1) holds for some $n \in \mathbb{N}$. Indeed, (3.1) holds if and only if

$$\left\lceil \frac{\log_r \rho}{\log_r (\alpha^{-1} \beta^{-1})} + \Phi \right\rceil < \left\lceil \frac{\log_r \rho}{\log_r (\alpha^{-1} \beta^{-1})} + \Psi \right\rceil \text{ and } 2 \le \left\lceil \frac{\log_r \rho}{\log_r (\alpha^{-1} \beta^{-1})} + \Psi \right\rceil,$$

where Φ and Ψ are positive real numbers independent of ρ with $\Phi < \Psi$.

Example 3.13. Let $\xi = re^{i\theta}$ where r = 9/2 and $0 \le \theta \le 0.064$. Then $v_2^{(2)}(\theta) < r < 5 = u_2(\theta)$. So, (C2) holds for the $(\xi, \mathcal{L}, \mathcal{X})$ -expansion where $\mathcal{L} = \mathbb{Z}[\hat{i}]$ and $\mathcal{X} = \{a + b\hat{i} : -1/2 \le a, b < 1/2\}$. Let $\rho = 2$, $\alpha = 3/5$ and

$$1 > \beta > F_{9/2}(3/5) = \frac{8495 - 180\sqrt{2}}{11901} \approx 0.6924.$$

Then we satisfy the conditions of Theorem 3.12 with n = 1 and k = 2. Hence, $C_{\xi}[0]$ is (α, β, ρ) -winning.

4 Quaternion expansions

The skew field (i.e. noncommutative division ring) \mathbb{H} of real quaternions is a unital associative algebra over \mathbb{R} with basis $\{1, \hat{i}, \hat{j}, \hat{k}\}$ where $\hat{i}, \hat{j}, \hat{k}$ are *imaginary* units satisfying

$$\hat{i}^2 = \hat{j}^2 = \hat{k}^2 = \hat{i}\hat{j}\hat{k} = -1.$$

A real quaternion $q \in \mathbb{H}$ is written uniquely as $q = a + b\hat{i} + c\hat{j} + d\hat{k}$, where $a, b, c, d \in \mathbb{R}$. We denote by \mathbb{H}_H and \mathbb{H}_L the rings of Hurwitz quaternions and Lipschitz quaternions, respectively:

$$\mathbb{H}_{H} := \{ a + b\hat{i} + c\hat{j} + d\hat{k} \mid a, b, c, d \in \mathbb{Z} \text{ or } a, b, c, d \in \mathbb{Z} + 1/2 \}$$
$$\mathbb{H}_{L} := \{ a + b\hat{i} + c\hat{j} + d\hat{k} \mid a, b, c, d \in \mathbb{Z} \}.$$

In this section, we consider a 4-dimensional rotational beta expansion where the matrix parameter M is isoclinic, that is, $M \in SO(4)$ and there exists a unit quaternion p such that Mx = px for all $x \in \mathbb{R}^4$ when we view \mathbb{R}^4 as the set \mathbb{H} of real quaternions. It has been shown in [4] that the expansion in this setting corresponds to an expansion on the set \mathbb{H} of real quaternions.

Let $q \in \mathbb{H}$ with |q| > 1. Let \mathcal{L} be a point lattice on \mathbb{H} with a fundamental domain \mathcal{X} such that $0 \in \mathcal{L} \cap \mathcal{X}$. Then

$$\mathbb{H} = \bigcup_{d \in \mathcal{L}} (\mathcal{X} + d).$$

For $z \in \mathcal{X}$, there exists a unique $d(z) \in \mathcal{L}$ such that $qz - d(z) \in \mathcal{X}$. We define the $(q, \mathcal{L}, \mathcal{X})$ -transformation $\mathbb{T} = \mathbb{T}_{q,\mathcal{L},\mathcal{X}} : \mathcal{X} \to \mathcal{X}$ by

$$\mathbb{T}(z) = qz - d(z).$$

The $(q, \mathcal{L}, \mathcal{X})$ -expansion of $z \in \mathcal{X}$ is $z = q^{-1}d_1 + q^{-2}d_2 + \cdots$, where the *j*th digit d_j of the expansion is given by $d(\mathbb{T}^{j-1}(z)) \in \mathcal{L}$. As before, we write $d(z) = d_q(z) := d_1d_2\cdots$ and $\mathcal{D} = \mathcal{D}(q, \mathcal{L}, \mathcal{X}) := \{d(z) \mid z \in \mathcal{X}\}$ for the digit set of the expansion.

We give some families of q-expansions.

Example 4.1 (Rotational beta expansion). Let $\{\eta_1, \eta_2, \eta_3, \eta_4\}$ be an \mathbb{R} -basis of \mathbb{H} . Then $\mathcal{L} := \mathbb{Z}\eta_1 + \mathbb{Z}\eta_2 + \mathbb{Z}\eta_3 + \mathbb{Z}\eta_4$ is a lattice of \mathbb{H} with fundamental domain $\mathcal{X} := \{t_1\eta_1 + t_2\eta_2 + t_3\eta_3 + t_4\eta_4 \mid t_1, t_2, t_3, t_4 \in [0, 1)\}$. Let $q \in \mathbb{H}$ with $|q| = \beta > 1$ and $q/\beta := \beta^{-1}q = a + b\hat{i} + c\hat{j} + d\hat{k}$ where $a, b, c, d \in \mathbb{R}$. If

$$M = \begin{bmatrix} a & -b & -c & -d \\ b & a & -d & c \\ c & d & a & -b \\ d & -c & b & a \end{bmatrix} \in SO(4)$$
(*)

(i.e., M is isoclinic), then $\beta M x = qx$ for any $x \in \mathbb{H}$ when x and qx are viewed as elements of \mathbb{R}^4 . Hence, the $(q, \mathcal{L}, \mathcal{X})$ -expansion (see [4]) corresponds to the 4-dimensional rotational beta expansion with parameters β and M.

Conversely, if M has the form (\star) where $a, b, c, d \in \mathbb{R}$ such that $a^2 + b^2 + c^2 + d^2 = 1$ and $\beta > 1$, then the rotational beta expansion with parameter (β, M) corresponds to the $(q, \mathcal{L}, \mathcal{X})$ -expansion where $q = \beta(a + b\hat{i} + c\hat{j} + d\hat{k}) \in \mathbb{H}$.

For instance, let $(\eta_1, \eta_2, \eta_3, \eta_4) = (1, \hat{i}, \hat{j}, \hat{k})$. Let $q = (1 + \sqrt{5})\hat{i}/2$. Then the q-expansion of $(1 + \hat{j})/2$ is given by the purely periodic expansion

$$0 (-2 - 2\hat{j}) (\hat{i} + \hat{k}) (-1 - \hat{j}) (\hat{i} + \hat{k}) (-1 - \hat{j}).$$

Example 4.2 (Zeta expansion). Let $\zeta \in \mathbb{H} \setminus \mathbb{R}$ with $|\zeta| > 1$. Let $\eta \in \mathbb{H}$ such that $\operatorname{Re}(\eta) = 0$ with $|\eta| = 1$ and $\eta \cdot \zeta = 0$ where \cdot is the usual dot product on \mathbb{R}^4 . Let $\varepsilon \in [0, 1)$. Consider the lattice

$$\mathcal{L} = \mathbb{Z} + \mathbb{Z}\overline{\zeta} + \mathbb{Z}\eta + \mathbb{Z}\overline{\zeta}\eta$$

with fundamental domain

$$\mathcal{X} = \{a_1 + a_2\overline{\zeta} + a_3\eta + a_4\overline{\zeta}\eta \mid -\varepsilon \le a_1, a_2, a_3, a_4 < 1 - \varepsilon\}.$$

Then the digit set \mathcal{D} of the ζ -expansion is contained in $\mathbb{Z} + \mathbb{Z}\eta$ (see [4]). This expansion extends ζ -expansion on complex numbers [11] to \mathbb{H} .

Example 4.3 (Symmetric q-expansion). Let $q \in \mathbb{H}$ with |q| > 1. For $1 \leq i \leq 4$, let $\varepsilon_i > 0$. Let $\mathcal{X} = \{a_1 + a_2\hat{i} + a_3\hat{j} + a_4\hat{k} \mid a_i \in [-\varepsilon_i, \varepsilon_i)\}$ be a fundamental domain of a point lattice \mathcal{L} be in \mathbb{H} such that $0 \in \mathcal{L}$. We say that the q-expansion is $(\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4)$ -symmetric. If $\varepsilon_1 = \cdots = \varepsilon_4 = \varepsilon$, then the q-expansion is ε -symmetric. For instance, if $\mathcal{L} = \mathbb{H}_L$ with fundamental domain $\mathcal{X} \cong [-1/2, 1/2)^2$, then the q-expansion is 1/2-symmetric. If $\mathcal{L} = \mathbb{H}_H$ and $\mathcal{X} = \Lambda$, then the q-expansion is (1/2, 1/2, 1/2, 1/4)-symmetric.

4.1 *q*-expansions and Schmidt game

Let us consider the $(q, \mathcal{L}, \mathcal{X})$ -expansion on \mathbb{H} where $q \in \mathbb{H}$ with $|q| > 1, \mathcal{L}$ is a point lattice in \mathbb{H} with a bounded fundamental domain \mathcal{X} . For an admissible block $\Omega = (a_1, a_2, \ldots, a_n)$, define the set

 $\mathcal{C}[\Omega] = \mathcal{C}_{(q,\mathcal{L},\mathcal{X})}[\Omega] := \{ z \in \mathcal{X} \mid \Omega \text{ appears in the expansion } d_q(z) \}.$

4.1.1 (α, β) -losing

In this section, we give values of $(\alpha, \beta) \in (0, 1)^2$ for which $\mathcal{C}[\Omega]$ is not (α, β) -winning with respect to the Schmidt game played in \mathcal{X} . This means that Bob can employ a strategy so that Ω is not a block of the outcome ω . In [12], such a set is called (α, β) -losing. Note that, for admissible blocks Ω and Ω' such that Ω is a sub-block of Ω' , if $\mathcal{C}[\Omega]$ is (α, β) -losing, then $\mathcal{C}[\Omega']$ is (α, β) -losing.

Theorem 4.4. There exists a positive constant $C_{\mathcal{X}}$ satisfying the following: Let $q \in \mathbb{H}$ with |q| > 1 and let $\Omega = d_1 d_2 \cdots d_n \in \mathcal{D}^n$ be a q-admissible word of length n such that

$$C_{\Omega} := C_{\mathcal{X}} + C_{\mathcal{X}} \left| \sum_{j=1}^{n} q^{n-j} d_j \right| < |q|^n.$$

Then, for any $(\alpha, \beta) \in (0, 1)^2$ with

$$C_{\Omega}|q|^{-n} \leq \alpha \text{ and } \alpha\beta \leq |q|^{-n},$$

we have $\mathcal{C}_q[\Omega]$ is (α, β) -losing.

Remark. Note that $C_q[\Omega]$ is (α, β) -losing if there exsits $\rho > 0$ such that $C_q[\Omega]$ is not (α, β) winning. More precisely, we can prove that $C_q[\Omega]$ is not (α, β) -winning for any $\rho < \rho_0(\mathcal{X})$, where $\rho_0(\mathcal{X})$ is a positive constant depending only on \mathcal{X} .

Proof. We show the case where n = 1 and denote $\Omega = d \in \mathcal{D}$. Let $\xi \neq 0$ be an interior point of \mathcal{X} . Let $\rho > 0$ such that $|\xi| > 2\rho$ and $\overline{B(\xi, \rho)} \subseteq \mathcal{X}$. Let $D := \sup_{z \in \mathcal{X}} |\xi - z|$ and

$$M := \sup_{\substack{z \in \mathcal{X} \\ \text{Let}}} |z| > 0.$$

$$C_{\mathcal{X}} := \max\left\{1 + \frac{D}{\rho}, \frac{M}{|\xi| - 2\rho}, \frac{1}{|\xi| - 2\rho}\right\}.$$

Note that

$$C_d \ge \max\left\{1 + \frac{D}{\rho}, \frac{M + |d|}{|\xi| - 2\rho}\right\}.$$

Assume that $C_d/|q| \leq \alpha$ and $\alpha\beta \leq 1/|q|$. Denote by a_k and b_k the centers of Alice's and Bob's kth ball, respectively. Let Bob choose the initial ball $B_0 = \overline{B(\xi, \rho)}$, i.e., $b_0 = \xi$. Then Alice chooses the ball $A_1 = \overline{B(a_1, \rho\alpha)} \subseteq B_0$. For $k, m \in \mathbb{N}$, let $a_k^{(m)}$ be the *m*th digit in the *q*-expansion of a_k .

We claim that $a_1^{(1)} \neq d$. Suppose otherwise. Then $qa_1 - d \in \mathcal{X}$. So, $|qa_1 - d| \leq M$ and

$$|a_1| \le \frac{M+|d|}{|q|} = (|\xi|-2\rho)\frac{M+|d|}{(|\xi|-2\rho)|q|} \le (|\xi|-2\rho)\frac{C_d}{|q|} \le (|\xi|-2\rho)\alpha.$$

Let $z \in A_1$. Then $|z| \leq |a_1| + \rho \alpha \leq (|\xi| - \rho)\alpha < |\xi| - \rho$. However, since $z \in A_1 \subseteq B_0 = \overline{B(\xi, \rho)}$, we have $|\xi| - |z| \leq |z - \xi| \leq \rho$. Contradiction.

Now, Bob chooses the ball $B_1 = B(q^{-1}(\xi + a_1^{(1)}), \alpha\beta\rho)$. First, we show that $B_1 \subseteq A_1$. Let $z \in B_1$. Then $|z - q^{-1}(\xi + a_1^{(1)})| \leq \alpha\beta\rho \leq \rho/|q|$. Note that since $qa_1 - a_1^{(1)} \in \mathcal{X}$, then $|qa_1 - a_1^{(1)} - \xi| \leq D$. Observe that

$$\begin{aligned} |z - a_1| &\leq |z - q^{-1}(\xi + a_1^{(1)})| + |a_1 - q^{-1}(\xi + a_1^{(1)})| \\ &\leq \frac{\rho}{|q|} + \frac{|qa_1 - a_1^{(1)} - \xi|}{|q|} \\ &\leq \frac{\rho}{|q|} + \frac{D}{|q|} = \frac{\rho}{|q|} \left(1 + \frac{D}{\rho}\right). \end{aligned}$$

So, $|z - a_1| \leq \rho C_d / |q| \leq \rho \alpha$. Then $B_1 \subseteq A_1$. Moreover, if $z \in B_1$, then $|qz - a_1^{(1)} - \xi| \leq \rho \alpha \beta |q| \leq \rho$. That is, $qz - a_1^{(1)} \in \overline{B(\xi, \rho)} \subseteq \mathcal{X}$. So, the first digit of any point in B_1 is $a_1^{(1)} \neq d$.

We explain Bob's strategy. Suppose Alice plays $A_k = B(a_k, \rho \alpha^k \beta^{k-1})$ for her kth turn for $k \in \mathbb{N}$. Bob responds with the ball

$$B_k = \overline{B(q^{-1}a_1^{(1)} + q^{-2}a_2^{(2)} + \dots + q^{-k}a_k^{(k)} + q^{-k}\xi, \alpha^k\beta^k\rho)}.$$

Note that by this strategy, a_k and a_{k+1} have the same first k digits in their q-expansions and $a_i^{(j)} \neq d$ for any $j \in \{1, \ldots, k\}$.

Therefore, the outcome ω has q-expansion $d(\omega) = a_1^{(1)} a_2^{(2)} \cdots$ where d does not appear. In other words, $C_q[d]$ is (α, β) -losing.

In the case where the length of Ω is general positive integer, we can show Theorem 4.4 in the same way as above.

We note that $C_{\mathcal{X}}$ depends only on the region \mathcal{X} . If |q| is sufficiently large compared to $\sum_{j=1}^{n} |q|^{n-j} |d_j|$, then we see $C_{\Omega} < |q|^n$. In this setting, we can choose $(\alpha, \beta) \in (0, 1)^2$ satisfying

the conditions of Theorem 4.4 because $C_{\Omega}|q|^{-n} < 1$.

In the proof above for n = 1, we see that, in any step, we can replace d by another digit d' where $|d'| \leq |d|$. Hence, we have the following corollary.

Corollary 4.5. Let $0 \leq t \in \mathbb{R}$. Then there exists $C_t > 0$ satisfying the following: If $q \in \mathbb{H}$ with |q| > 1 and $\alpha, \beta \in (0, 1)$ such that $C_t\beta \leq \alpha\beta|q| \leq 1$, then the set $\{z \in \mathcal{X} : d \text{ appears in } d_q(z) \text{ where } |d| \leq t\}$ is (α, β) -losing.

Moreover, in the proof of Theorem 4.4 for n = 1, we want to make C_d as small as possible. Hence, we want Bob to choose ξ and ρ so that $\overline{B(\xi, \rho)}$ is the biggest closed ball contained in \mathcal{X} . Likewise, we want to minimize D and maximize ρ and $|\xi| - 2\rho$.

In Examples 4.6 and 4.7, we choose ρ such that

$$1 + \frac{D}{\rho} = \frac{M + |d|}{|\xi| - 2\rho}$$

in order to control the value of C_d and provide a lower bound.

Example 4.6. In Example 4.1, we take $\{\eta_1, \eta_2, \eta_3, \eta_4\} = \{1, \hat{i}, \hat{j}, \hat{k}\}$. Let $\xi = (1 + \hat{i} + \hat{j} + \hat{k})/2 \in \mathcal{X}$. Note that $|\xi| = 1$, $\sup_{z \in \mathcal{X}} |z - \xi| = 1$ and $\sup_{z \in \mathcal{X}} |z| = 2$. For $d \in \mathcal{D}$, we take

$$\rho := \frac{\sqrt{|d|^2 + 6|d| + 17} - |d| - 3}{4} \le \frac{\sqrt{17} - 3}{4} < \frac{1}{2}$$

Then $|\xi| > 2\rho$ and $\overline{B(\xi,\rho)} \subseteq \mathcal{X}$. Also, $C_d = 1 + 1/\rho \ge 1 + 4/(\sqrt{17} - 3) \approx 4.56$.

Example 4.7. In Example 4.2, suppose $\operatorname{Re}(\zeta) \geq 0$ and $\varepsilon \neq 1/2$. Then $M = \sup_{z \in \mathcal{X}} |z| = \sqrt{2}|1+\overline{\zeta}| \max\{1-\varepsilon,\varepsilon\}$. Take $\xi = (1/2-\varepsilon)(1+\overline{\zeta}+\eta+\overline{\zeta}\eta)$. Then $|\xi| = \sqrt{2}|1/2-\varepsilon||1+\overline{\zeta}|$. The largest ball that fits inside \mathcal{X} is $B(\xi, 1/2)$. Moreover, $D = \sup_{z \in \mathcal{X}} |z-\xi| = \sqrt{2}|1+\overline{\zeta}|/2$, which is minimized by our choice of ξ . For $d \in \mathcal{D}$, we take $\rho < 1$ to be the positive solution to $1 + D/\rho = (M + |d|)/(|\xi| - 2\rho)$. Note that ρ decreases as |d| increases and $\rho \leq \sqrt{13 - 3\sqrt{17}}|1+\overline{\zeta}|/4 \approx 0.7942|1+\overline{\zeta}|$. Therefore, $C_d = 1 + D/\rho \geq 4.5616$.

Example 4.8. Let $\mathcal{L} = \mathbb{H}_H$ and $\mathcal{X} = \{a_1 + a_2\hat{i} + a_3\hat{j} + a_4\hat{k} : 0 \le a_1, a_2, a_3 < 1, 0 \le a_4 < 1/2\}$. Take $\xi = (2 + 2\hat{i} + 2\hat{j} + \hat{k})/4$. Then $\sup_{z \in \mathcal{X}} |z| = \sqrt{13}/2$ and $\sup_{z \in \mathcal{X}} |z - \xi| = \sqrt{13}/4$. Take $\rho = 1/4$. Then $C_d = \max\{1 + \sqrt{13}, (4|d| + 2\sqrt{13})/(\sqrt{13} - 2)\}$. If d = 0, then $C_d = 1 + \sqrt{13}$. If $d \ne 0$, then $C_d = (4|d| + 2\sqrt{13})/(\sqrt{13} - 2) \ge 2(17 + 4\sqrt{13})/9 \approx 6.98$.

Example 4.9. Consider the ε -symmetric q-expansion in Example 4.3. We modify the proof of Theorem 4.4 slightly as the optimal choice for ξ so that $B(\xi, \rho)$ is contained in \mathcal{X} is zero and, in this case, $|\xi| = 0$. Now, $M = \sup_{z \in \mathcal{X}} |z| = 2\varepsilon$. Let $0 < \tau < \varepsilon/2$ and $\xi = \tau(1+\hat{\imath}+\hat{\jmath}+\hat{k}) \in \mathcal{X}$. Thus, $D = \sup_{z \in \mathcal{X}} |z - \xi| = 2(\varepsilon + \tau)$. Take $\rho = \tau$. Then $\overline{B(\xi, \rho)} \subseteq \mathcal{X}$. Redefine

$$C_d = \max\{3 + 2\varepsilon/\tau, (2\varepsilon + |d|)/(2\tau)\}.$$

If $|d| \leq 6\tau + 2\varepsilon$, then $C_d = 3 + 2\varepsilon/\tau > 7$. Otherwise, $C_d = (2\varepsilon + |d|)/(2\tau)$.

Assume that $C_d/|q| \leq \alpha$. Let Bob start with the ball $B_0 = \overline{B(\xi, \rho)} = \overline{B(\xi, \tau)}$. By calculations, $|z| \geq \tau$ for $z \in B_0$.

Using this configuration, Bob can use similar strategy as in the proof of Theorem 4.4 for n = 1 so that $C_q[d]$ is (α, β) -losing.

Now, the interval $[C_{\Omega}/|q|^n, 1)$ is the set of α 's satisfying the hypothesis of Theorem 4.4 for some β . Note that in most cases, $C_{\Omega} = M_n$. For large |q|,

$$\frac{M_{n-1}/|q|^{n-1}}{M_n/|q|^n} = |q| \frac{M + \left|\sum_{j=1}^{n-1} q^{n-1-j} d_j\right|}{M + \left|\sum_{j=1}^n q^{n-j} d_j\right|} \gg 1.$$

This implies that, when |q| is large, the set of α 's for which $C_q[\Omega]$ is α -losing gets bigger as the length of Ω increases. This agrees with our expectations since $C_q[\Omega]$ gets smaller when Ω has more digits. For an admissible block Ω of length $|\Omega|$, we define

$$\widetilde{C}_{\Omega} := \inf\{C > 0 : \mathcal{C}_q[\Omega] \text{ is } (\alpha, \beta) \text{-losing if } C\beta \leq \alpha\beta |q|^{|\Omega|} \leq 1\}.$$

We remark that \widetilde{C}_{Ω} complements the idea of the winning dimension of $C_q[\Omega]$. Computing \widetilde{C}_{Ω} may not be trivial. Indeed, the constant C_{Ω} given in the proof of Theorem 4.4 may be improved. We give an example.

Example 4.10. Let $q \in \mathbb{H}$ with |q| > 1. Consider the lattice $\mathcal{L} = \mathbb{H}_H$ with fundamental domain $\Delta = \{a + b\hat{i} + c\hat{j} + d\hat{k} \mid 0 \leq a, b, c < 1, 0 \leq d < 1/2\}$. Note that $\sup_{z \in \Delta} |z| = \sqrt{13}/2$. Let $C \geq 2$. Suppose $C/|q| \leq \alpha$ and $\alpha\beta \leq 1/|q|$. Let Bob starts with the ball $B_0 = B(\xi, 1/k)$ where $k \in \mathbb{N}$ and $\xi = (2 + 2\hat{i} + 2\hat{j} + \hat{k})/4$. It can be shown that

$$\min_{z \in B_0} |z| = \frac{\sqrt{13}k - 4}{4k}.$$

Alice then chooses a ball $A_1 = B(a, \alpha/k)$ for some $a \in B_0$. Let $z \in A_1$ such that $|z| \leq |a| - \alpha/k$. Suppose that the first digit of a is 0. Then $qa \in \Delta$ and so, $|a| \leq \frac{\sqrt{13}}{2|q|} \leq \frac{\sqrt{13}}{2C}\alpha$. Since $C \geq 2$, then

$$|z| \le \left(\frac{\sqrt{13}}{2C} - \frac{1}{k}\right) \alpha < \frac{\sqrt{13}}{2C} - \frac{1}{k} \le \frac{\sqrt{13}}{4} - \frac{1}{k} \le |z|.$$

We have a contradiction. Thus, the first digit of a is not 0. In other words, if $2\beta < \alpha\beta |q| \leq 1$, then $C_q[0]$ is (α, β) -losing.

Given $d \in \mathcal{D}$, finding (α, β) such that $C_q[d]$ is (α, β) -winning is relatively difficult since the cylinder sets are not so easy to compute. However, using Theorem 2.7, we have the following.

Theorem 4.11. Let $1 < q \in \mathbb{R}$, $\mathcal{L} = \mathbb{H}_L$ and $\mathcal{X} \cong [0,1)^4$. Suppose $K_q < \infty$. If $a_i \in \{0, 1, 2, \ldots, \lfloor q \rfloor\}$ such that $a_i \leq d'$ where d' is the minimal digit of $\lim_{\varepsilon \to 0^+} d(1-\varepsilon)$ for $1 \leq i \leq 4$ and $\alpha, \beta \in (0,1)$ such that $\beta > A_{2q}(\alpha)$ with $\log_q(\alpha\beta) \notin \mathbb{Q}$, then $\mathcal{C}_q[a_1 + a_2\hat{i} + a_3\hat{j} + a_4\hat{k}]$ is (α, β) -winning.

Proof. Let x_i be the center of Bob's *i*th ball B_i and $y_i = a_i^{(1)} + a_i^{(2)}\hat{i} + a_i^{(3)}\hat{j} + a_i^{(4)}\hat{k}$ be the center of Alice's *i*th ball A_i .

Since $\beta > A_{2q}(\alpha)$, similar to the proof of Theorem 2.7, there exists $n, k \in \mathbb{N}$ such that for j = 1, 2, 3, 4, Alice can force $a_{n+1}^{(j)}$ to be a center of some interval of $V_k(q; a_j)$. In particular, $(K+2)/b^{k-1} < \rho(\alpha\beta)^n(1-\alpha)/2$ and the consecutive centers of the intervals of $V_k(q; a_j)$ are at most $q^{-(k-1)}$ units apart. Let $b_n^{(j)}$ be the *j*th component of x_n and $a^{(j)}$ be the center of an interval of $V_k(q; a_j)$ that is closest to $b_n^{(j)}$. If the length of the interval of $V_k(q; a_j)$ centered at $a^{(j)}$ is q^{-k} , then Alice chooses $a_{n+1}^{(j)} = a^{(j)}$. Otherwise, Alice chooses $a_{n+1}^{(j)}$ to be the nearest

center of an interval of $V_k(q; a_j)$ with length q^{-k} . In any case, $|a_{n+1}^{(j)} - b_n^{(j)}| \leq (K+2)q^{-(k-1)} < \rho(\alpha\beta)^n (1-\alpha)/2$. Thus,

$$|y_{n+1} - x_n| = \sqrt{\sum_{j=1}^4 |a_{n+1}^{(j)} - b_n^{(j)}|^2} < \sqrt{\sum_{j=1}^4 \left(\frac{\rho(\alpha\beta)^n (1-\alpha)}{2}\right)^2} = \rho(\alpha\beta)^n (1-\alpha).$$

We consider the case where Bob plays optimally. Now, Bob wants to move away from y_k as far as possible since the *j*th component of y_k has the digit a_j in its *q*-expansion. In particular, he wants to concentrate his movement on one of the components, since moving away from a component is enough for him to win. He also uses the same direction in his moves to maximize the distance.

WLOG, suppose that Bob chooses to move along the first component towards the positive direction. In this case, we have $x_k = y_k + \rho \alpha (\alpha \beta)^{k-1} (1-\beta)$. In response, Alice chooses,

$$y_{k+1} = x_k - \rho(\alpha\beta)^k (1-\alpha).$$

Hence, the game reduces to a 1-dimensional Schmidt game since the other 3 components do not move. $\hfill \Box$

5 Acknowledgements

The first author's work was supported by the JSPS KAKENHI Grant Number 24K06641. The third author acknowledges the Office of the Chancellor of the University of the Philippines Diliman, through the Office of the Vice Chancellor for Research and Development, for funding support through the Thesis and Dissertation Grant (242405 TND).

References

- [1] S. AKIYAMA AND J. CAALIM, Rotational beta expansion: ergodicity and soficness, Journal of the Mathematical Society of Japan, 69 (1) (2017), pp. 397–415.
- [2] Y. BUGEAUD, Distribution Modulo One and Diophantine Approximation, Cambridge Tracts in Mathematics, Cambridge University Press, 2012.
- [3] J. CAALIM AND S. DEMEGILLO, Beta Cantor series expansion and admissible sequences, Acta Polytech, 60 (3) (2020), pp. 214–224.
- [4] J. CAALIM AND N. NOLLEN, *Periodic quaternion expansion in Pisot bases*, Osaka Journal of Mathematics, to appear (2025).
- [5] V. DREMOV, On domains of (α, β) -winnability, Doklady Akademii Nauk, 384 (2002), pp. 304–307.
- [6] C. FREILING, An answer to a question of Schmidt on (α, β) games, Journal of Number Theory, 15 (2) (1982), pp. 226–228.

- [7] N. LANGEVELD AND T. SAMUEL, Intermediate β -shifts as greedy β -shifts with a hole, Acta Mathematica Hungarica, 170 (2023), pp. 269–301.
- [8] V. NECKRASOV AND E. ZHAN, On Nontrivial Winning and Losing Parameters of Schmidt games, arXiv preprint arXiv:2401.00614, (2024).
- [9] A. RÉNYI, Representations for real numbers and their ergodic properties, Acta Mathematica Academiae Scientiarum Hungaricae, 8 (1957), pp. 477–493.
- [10] W. M. SCHMIDT, On badly approximable numbers and certain games, Transactions of the American Mathematical Society, 123 (1966), pp. 178–199.
- [11] P. SURER, Representations for complex numbers with integer digits, Research in Number Theory, 6 (2020), p. 47.
- [12] M. ZANGER-TISHLER AND S. KALIA, On the Winning and Losing Parameters of Schmidt's game, 2013.