Derived representations of quantum character varieties

M. $Faitg^1$

¹Institut de Mathématiques de Toulouse, Université Paul Sabatier, 118 route de Narbonne, F-31062 Toulouse, France.

E-mail adress: matthieu.faitg@math.univ-toulouse.fr

Abstract

Quantum moduli algebras $\mathcal{L}_{g,n}^{inv}(H)$ were introduced by Alekseev–Grosse–Schomerus and Buffenoir–Roche in the context of quantization of character varieties of surfaces and exist for any quasitriangular Hopf algebra H. In this paper we construct representations of $\mathcal{L}_{g,n}^{inv}(H)$ on cohomology spaces $\operatorname{Ext}_{H}^{m}(X, M)$ for all $m \geq 0$, where X is any H-module and M is any $\mathcal{L}_{g,n}(H)$ -module endowed with a compatible H-module structure. As a corollary and under suitable assumptions on H, we obtain projective representations of mapping class groups of surfaces on such Ext spaces. This recovers the projective representations obtained in [LMSS23] from Lyubashenko theory, when the category $\mathcal{C} = H$ -mod is used in their construction. Other topological applications are matrix-valued invariants of knots in thickened surfaces and representations of skein algebras on Ext spaces.

1 Introduction

1.1 Context

The basic algebraic inputs in quantum topology are ribbon tensor categories, from which one constructs invariants of links, 3-manifolds or more generally TQFTs. Some assumptions are generally made on the category C one wants to use. In the famous Reshetikhin–Turaev construction [RT91, Tur94], one assumes that C is finite, semisimple (or is truncated to its semisimple part) and its braiding is non-degenerate. While non-degeneracy and usually also finiteness are widely accepted as reasonable assumptions, many works have focused on the creation of topological invariants based on non-semisimple categories, like representations of mapping class groups [Lyu95], link invariants [GKP13], invariants of 3-manifolds [CGP14] (where finiteness is also weakened) and TQFTs [DGGPR22], to cite a few.

An aspect which is peculiar to the non-semisimple situation, and which has been much less explored from a topological perspective, is the existence of non-zero Ext spaces $\text{Ext}_{\mathcal{C}}^m(-,-)$ for m > 0. It is natural to wonder how these cohomology spaces interact with the topological invariants defined from \mathcal{C} . Eventually one would like to obtain "derived TQFTs" with values in a category of complexes, or whose state spaces are cohomology spaces. Pioneering works in this direction are [LMSS18, LMSS23, SW21a, SW21b], where derived modular functors are constructed.

The present paper was inspired by the work of Lentner, Mierach, Schweigert and Sommerhäuser [LMSS18, LMSS23] who proved that the Lyubashenko construction [Lyu95] can be extended to obtain projective representations of mapping class groups of surfaces on certain Ext spaces. Let us briefly recall their main result; this will give a more precise idea of what is a "derived topological invariant" and will facilitate the comparison with our results explained below.

Let $\Sigma_{g,n}^{\circ}$ be the compact oriented surface of genus g with n + 1 boundary circles, one of which is distinguished. Let \mathcal{C} be a ribbon finite tensor category which is factorizable (*i.e.* non-degeneracy of the braiding). Label the boundary circles of $\Sigma_{g,n}^{\circ}$ by any objects $X, X_1, \ldots, X_n \in \mathcal{C}$ and denote the resulting "colored surface" by $\Sigma_{g,X_1,\ldots,X_n}^X$:

To such a colored surface, associate the vector space

 $Z(\Sigma_{g,X_1,\ldots,X_n}^X) := \operatorname{Hom}_{\mathcal{C}} \left(X \otimes X_1 \otimes \ldots \otimes X_n, L^{\otimes g} \right)$

where L is the coend $\int^{X \in \mathcal{C}} X^* \otimes X$, an important object in \mathcal{C} . Denote by $MCG(\Sigma_{g,n}^{\circ})$ the mapping class group which fixes the boundary pointwise. Using a sophisticated system of axiomatic rules, Lyubashenko [Lyu95, §4.4] defined a projective representation

$$Z_X : \mathrm{MCG}(\Sigma_{g,n}^{\circ}) \to \mathrm{PGL}(Z(\Sigma_{g,X_1,\dots,X_n}^X)).$$
(1)

The values of Z_X are defined on a set of generators of $MCG(\Sigma_g^{\circ})$ by explicit dinatural transformations which factor through the coend $L^{\otimes g}$. Moreover the choice $X = \mathbf{1}$ (the unit object of \mathcal{C}) is special: Z_1 descends to a projective representation of the mapping class group of the surface $\Sigma_{g,n}$ where the distinguished boundary component has been closed.

The idea of [LMSS23] is to replace the space $Z(\Sigma_{q,X_1,\ldots,X_n}^X)$ by its derived version:

$$Z^{m}(\Sigma^{X}_{g,X_{1},\ldots,X_{n}}) = \operatorname{Ext}^{m}_{\mathcal{C}}(X \otimes X_{1} \otimes \ldots \otimes X_{n}, L^{\otimes g}) \quad \text{for all } m \geq 0.$$

One can imagine that the distinguished boundary circle of $\Sigma_{g,n}^{\circ}$ is now colored by a projective resolution $0 \leftarrow X \leftarrow P_0 \leftarrow P_1 \leftarrow \ldots$ in \mathcal{C} , which by application of the functor $Z(\Sigma_{g,X_1,\ldots,X_n}^{-}) = \operatorname{Hom}_{\mathcal{C}}(-\otimes X_1 \otimes \ldots \otimes X_n, L^{\otimes g})$ gives the cochain complex

$$0 \longrightarrow Z(\Sigma_{g,X_1,\dots,X_n}^{P_0}) \longrightarrow Z(\Sigma_{g,X_1,\dots,X_n}^{P_1}) \longrightarrow \dots$$

whose cohomology is $Z^m(\Sigma_{g,X_1,\ldots,X_n}^X)$. It is not hard to see that the differential of this complex commute with the representations (1) on each cochain space, resulting in a projective representation of $MCG(\Sigma_{g,n}^{\circ})$ on the cohomology spaces:

$$Z_X^m : \mathrm{MCG}(\Sigma_{g,n}^\circ) \to \mathrm{PGL}\left(Z^m(\Sigma_{g,X_1,\dots,X_n}^X)\right), \quad Z_X^m(f)\big([s]\big) = \big[Z_{P_m}(s)\big]$$
(2)

for all cocycle $s \in Z(\Sigma_{g,X_1,\ldots,X_n}^{P_m})$, where [] denotes cohomology classes. The hard part, which is the main result of [LMSS23], is to prove that for $X = \mathbf{1}$ the morphism $Z_{\mathbf{1}}^m$ descends to a projective representation of $MCG(\Sigma_{g,n})$. Thanks to the classical fact that $Ext^0_{\mathcal{C}}(-,-) =$ $Hom_{\mathcal{C}}(-,-)$, the case m = 0 recovers Lyubashenko's construction.

Let H be a ribbon Hopf algebra. In this paper we show that a certain quantization $\mathcal{L}_{g,n}^{inv}(H)$ of the character variety of $\Sigma_{g,n}^{\circ}$ has representations on cohomology spaces $V^m(\Sigma_{g,X_1,\dots,X_n}^X)$ which are a variant of $Z^m(\Sigma_{g,X_1,\dots,X_n}^X)$ in the case $\mathcal{C} = H$ -mod. For surfaces $\Sigma_{g,0}^{\circ}$ and under certain assumptions on H, it is possible to associate a unique up to scalar element $\widehat{f} \in \mathcal{L}_{g,0}^{inv}(H)$ to each mapping class $f \in \text{MCG}(\Sigma_{g,0}^{\circ})$. By representing \widehat{f} on $V^m(\Sigma_g^X)$ we recover the projective representation of [LMSS23]. The fact that we have a representation of the whole algebra $\mathcal{L}_{g,n}^{inv}(H)$ and not just of the mapping class group permits further results. First, we define isotopy invariants of knots in surfaces with values in $\text{End}_{\Bbbk}(V^m(\Sigma_{g,X_1,\dots,X_n}^X))$ which obey a version of the claim that "the choice $X = \mathbf{1}$ is like closing the distinguished boundary circle". Second, we obtain representations of skein algebras on the spaces $V^m(\Sigma_{g,X_1,\dots,X_n}^X)$.

1.2 Summary of results

In order to explain our results precisely, we begin with a short description of the quantization of character varieties of surfaces that we use here and which is due to Alekseev–Grosse–Schomerus [AGS95] and Buffenoir–Roche [BR95]. For an algebraic Lie group G and an oriented surface Σ , the affine variety of representations (resp. characters) of $\pi_1(\Sigma)$ with values in G is

$$\operatorname{Rep}_{G}(\Sigma) = \operatorname{Hom}_{\operatorname{Grp}}(\pi_{1}(\Sigma), G) \qquad (\operatorname{resp. } \operatorname{Ch}_{G}(\Sigma) = \operatorname{Rep}_{G}(\Sigma) /\!\!/ G)$$

where G acts by conjugation on $\operatorname{Rep}_G(\Sigma)$. By definition $\mathcal{O}[\operatorname{Ch}_G(\Sigma)] = \mathcal{O}[\operatorname{Rep}_G(\Sigma)]^{G-\operatorname{inv}}$, where \mathcal{O} denotes the algebra of regular functions. Assume that $\Sigma = \Sigma_{g,n}^{\circ}$.¹ Since $\pi_1(\Sigma_{g,n}^{\circ})$ is a free group on 2g + n generators, we have $\mathcal{O}[\operatorname{Rep}_G(\Sigma_{g,n}^{\circ})] \cong \mathcal{O}(G)^{\otimes(2g+n)}$. The idea of the quantization is to consider the vector space $\mathcal{O}_q[\operatorname{Rep}_G(\Sigma_{g,n}^{\circ})] = \mathcal{O}_q(G)^{\otimes(2g+n)}$, where $\mathcal{O}_q(G)$ is the standard quantized algebra of functions on G, and to endow it with a "twisted product" which involves the R-matrix of $U_q(\mathfrak{g})$ where $\mathfrak{g} = \operatorname{Lie}(G)$. A crucial feature of the construction is that $\mathcal{O}_q[\operatorname{Rep}_G(\Sigma_{g,n}^{\circ})]$ is an $U_q(\mathfrak{g})$ -module-algebra under a certain coadjoint action. This permits to define the subalgebra $\mathcal{O}_q[\operatorname{Ch}_G(\Sigma_{g,n}^{\circ})] := \mathcal{O}_q[\operatorname{Rep}_G(\Sigma_{g,n}^{\circ})]^{U_q(\mathfrak{g})-\operatorname{inv}}$, called quantum moduli algebra or quantum character variety.

Being purely algebraic, this construction can be applied to any quasitriangular Hopf algebra H instead of $U_q(\mathfrak{g})$, with the restricted dual H° instead of $\mathcal{O}_q(G)$. In this way, see §4.1 for the precise definitions, one gets a (right) H-module-algebra $\mathcal{L}_{g,n}(H)$ which is $(H^\circ)^{\otimes(2g+n)}$ as a vector space and which has a subalgebra

$$\mathcal{L}_{q,n}^{\mathrm{inv}}(H) := \mathcal{L}_{q,n}(H)^{H-\mathrm{inv}}$$

of invariant elements under the coadjoint action of H^2 . In particular, for finite-dimensional H, $\mathcal{L}_{0,1}(H)$ is a variant of the coend $L = \int^{X \in H \text{-mod}} X^* \otimes X$ used in Lyubashenko theory.

It is well-known that $\mathcal{L}_{g,n}(H)$ has a representation on the spaces

$$(H^{\circ})^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n \tag{3}$$

for all *H*-modules X_1, \ldots, X_n . Moreover, we prove in §4.2 that there is a *H*-module structure on the spaces (3) such that the action of $\mathcal{L}_{g,n}(H)$ is *H*-linear. This is called a *H*-equivariant module, see Definition 2.1.

Let k be the ground field of the ribbon Hopf algebra H. Take another H-module X and consider the k-vector spaces

$$V(\Sigma_{g,X_1,\dots,X_n}^X) = \operatorname{Hom}_H \left(X, (H_{\operatorname{coad}}^\circ)^{\otimes g} \otimes X_1 \otimes \dots \otimes X_n \right)$$
$$V^m(\Sigma_{g,X_1,\dots,X_n}^X) = \operatorname{Ext}_H^m \left(X, (H_{\operatorname{coad}}^\circ)^{\otimes g} \otimes X_1 \otimes \dots \otimes X_n \right) \quad \text{for all } m \ge 0$$

where coad is a coadjoint action of H on H° . These are variants of the Lyubashenko spaces $Z(\Sigma_{g,X_1,\ldots,X_n}^X)$ and of their derived version.³ The H-equivariance of (3) easily implies that $\mathcal{L}_{g,n}^{\mathrm{inv}}(H)$ has a representation on $V(\Sigma_{g,X_1,\ldots,X_n}^X)$ for all X, given by acting on the target. Thus if $0 \leftarrow X \leftarrow P_0 \leftarrow P_1 \leftarrow \ldots$ is a projective resolution of H-modules, we have a representation of $\mathcal{L}_{g,n}^{\mathrm{inv}}(H)$ on each of the cochain spaces in the complex

$$0 \longrightarrow V(\Sigma_{g,X_1,\dots,X_n}^{P_0}) \longrightarrow V(\Sigma_{g,X_1,\dots,X_n}^{P_1}) \longrightarrow \dots$$

which are readily seen to commute with the differential. Hence:

¹Because of the use of stated skein algebras to be explained below, it is more convenient to assume that $\Sigma_{g,n}^{\circ}$ is the oriented surface of genus with *n* punctures and one boundary circle, instead of n + 1 boundary circles as in §1.1. This is what $\Sigma_{g,n}^{\circ}$ means in the sequel.

²The relation with the previous notations is thus $\mathcal{O}_q[\operatorname{Rep}_G(\Sigma_{g,n}^\circ)] = \mathcal{L}_{g,n}(U_q(\mathfrak{g}))$ and $\mathcal{O}_q[\operatorname{Ch}_G(\Sigma_{g,n}^\circ)] = \mathcal{L}_{g,n}(U_q(\mathfrak{g}))$, modulo the use of $\mathcal{O}_q(G)$ instead of the whole $U_q(\mathfrak{g})^\circ$.

³One might think that $V(\Sigma_{g,X_1,...,X_n}^X)$ is related to $Z(\Sigma_{g,X_1,...,X_n}^X)$ from §1.1 simply by duality, but actually this is more subtle because we use a *right H*-action. See §5.2.3 for an explicit isomorphism in the case n = 0.

Theorem 1. (Thm. 4.12) For all *H*-modules X, X_1, \ldots, X_n and all $m \ge 0$ there is a representation of $\mathcal{L}_{g,n}^{inv}(H)$ on $V^m(\Sigma_{g,X_1,\ldots,X_n}^X)$.

The arguments which lead to Theorem 1 are completely general and elementary: in §2.1 we construct representations of any *H*-module-algebra *A* on $\operatorname{Ext}_{H}^{m}(X, M)$ where *X* is a *H*-module and *M* is a *H*-equivariant *A*-module. In particular, if there exists a special morphism $\mu : H \to A$ called a quantum moment map then any *A*-module is *H*-equivariant, see §2.2.

Before stating our first topological application of Theorem 1 we need a few preliminary material. Given an element $\varphi \in \mathcal{L}_{0,1}^{\text{inv}}(H)$, *i.e.* a linear form $\varphi \in H^{\circ}$ which satisfies $\varphi(xy) = \varphi(yS^2(x))$, one can define a map

$$\mathbb{W}^{\varphi}: \left\{ \substack{\text{isotopy classes} \\ \text{of oriented framed}} \text{ knots in } \Sigma_{g,n}^{\circ} \times [0,1] \right\} \to \mathcal{L}_{g,n}^{\text{inv}}(H).$$
(4)

They appear in [BR96, BFK98a, Fai20b] under the name of Wilson loop maps. In §5.1, under certain assumptions on H, we give yet another construction of \mathbb{W}^{φ} which is based on an algebra isomorphism between $\mathcal{L}_{g,n}(H)$ and the stated skein algebra $\mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$, where $\Sigma_{g,n}^{\circ,\bullet}$ is $\Sigma_{g,n}^{\circ}$ with a puncture on the boundary. The main defining properties of $\mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$ are recalled in §5.1.1 from [BFR23, §6.1]. It is a generalization to any ribbon Hopf algebra H of the stated skein algebras defined in [Lê18, CL22] and [LS21], which correspond to the cases $H = U_q(\mathfrak{sl}_2)$ and $H = U_q(\mathfrak{sl}_n)$. The isomorphism $\mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet}) \cong \mathcal{L}_{g,n}(H)$ was established in [BFR23, §6.2] building upon [Fai20b] and is quickly recalled in §5.1.3. Taking advantage of the topological nature of stated skein algebras, we note in §5.1.2-§5.1.3 that there is a natural definition of knot maps

$$\mathfrak{i}_K : \mathcal{L}_{0,1}(H) \to \mathcal{S}_H^{\mathrm{st}}(\Sigma_{q,n}^{\circ,\bullet}) \cong \mathcal{L}_{g,n}(H)$$

where $K \subset \Sigma_{g,n}^{\circ,\bullet} \times [0,1]$ is any oriented framed knot which has a basepoint in $\partial(\Sigma_{g,n}^{\circ,\bullet}) \times [0,1]$. When \mathfrak{i}_K is restricted to $\mathcal{L}_{0,1}^{\mathrm{inv}}(H)$ the basepoint can be forgotten and we define $\mathbb{W}^{\varphi}(K) = \mathfrak{i}_K(\varphi)$.

Combining \mathbb{W}^{φ} with the representation in Theorem 1, we obtain a map

$$\mathcal{I}_X^{\varphi,m}: \left\{ \underset{\text{of oriented framed}}{\text{isotopy classes}} \text{ knots in } \Sigma_{g,n}^{\circ} \times [0,1] \right\} \xrightarrow{\mathbb{W}^{\varphi}} \mathcal{L}_{g,n}^{\text{inv}}(H) \xrightarrow{\text{rep.}} \text{End}_{\Bbbk} \left(V^m(\Sigma_{g,X_1,\dots,X_n}^X) \right).$$

for any $\varphi \in \mathcal{L}_{0,1}^{\text{inv}}(H)$, $m \geq 0$ and *H*-module *X*. This map is an isotopy invariant of knots and allows us to give a precise statement of the idea that the distinguished boundary component of $\Sigma_{q,n}^{\circ}$ "disappears" if we color it by $X = \mathbb{k}$, the trivial *H*-module:

Theorem 2. (Thm. 5.8) Assume that H and the modules X_1, \ldots, X_n are finite-dimensional. If two knots $K_1, K_2 \subset \Sigma_{g,n}^{\circ} \times [0,1]$ become isotopic in $\Sigma_{g,n} \times [0,1]$, then $\mathcal{I}_{\Bbbk}^{\varphi,m}(K_1) = \mathcal{I}_{\Bbbk}^{\varphi,m}(K_2)$.

Said differently, $\mathcal{I}_{\Bbbk}^{\varphi,m}$ descends to a (matrix-valued) isotopy invariant of knots in $\Sigma_{g,n} \times [0, 1]$. This is a nice interaction between topology and homological algebra, because the boundary component of $\Sigma_{g,n}^{\circ}$ disappears only when passing to the cohomology spaces. For n = 0 we obtain an isotopy invariant of knots in closed surfaces which in the case m = 0 was studied in [Fai20b, §6]. It makes no doubt that Theorem 2 can be extended to links, see Remark 5.10.

The second topological application of Theorem 1 is the construction of projective representations of mapping class groups. There is a natural action of the mapping class group on the character variety. This action survives quantization: there is an action of $MCG(\Sigma_{g,n}^{\circ})$ on $\mathcal{L}_{g,n}(H)$ by algebra automorphisms. The previous proof of this fact [AS96a, Fai20a] was rather involved. In §5.2.1 we note that $MCG(\Sigma_{g,n}^{\circ})$ obviously acts on the stated skein algebra $\mathcal{S}_{H}^{st}(\Sigma_{g,n}^{\circ,\bullet})$ and thus on $\mathcal{L}_{g,n}(H)$ thanks to the isomorphism between them; we then prove that it equals the "old" action defined by means of explicit formulas. Moreover the subalgebra $\mathcal{L}_{g,n}^{inv}(H)$ is stable under this action. We then restrict to n = 0, that is to surfaces $\Sigma_{g,0}^{\circ}$, and assume that the ribbon Hopf k-algebra H is finite-dimensional and factorizable. It is known that for such H there is an algebra isomorphism $\mathcal{L}_{g,0}(H) \cong \operatorname{End}_{\Bbbk}((H^*)^{\otimes g})$ ([Fai20a], reviewed in Appendix A). In other words $\mathcal{L}_{g,0}(H)$ is a matrix algebra and hence any algebra automorphism is by conjugation. Since $\operatorname{MCG}(\Sigma_{g,0}^{\circ})$ acts on $\mathcal{L}_{g,0}(H)$ by algebra automorphisms, we obtain a unique-up-to-scalar conjugation element $\widehat{f} \in \mathcal{L}_{g,0}(H)$ for each mapping class $f \in \operatorname{MCG}(\Sigma_{g,0}^{\circ})$. Actually $\widehat{f} \in \mathcal{L}_{g,0}^{\operatorname{inv}}(H)$ (Lemma 5.15) and Theorem 1 gives in particular a projective representation

$$\mathcal{R}_X^m : \operatorname{MCG}(\Sigma_{g,0}^\circ) \longrightarrow \mathcal{L}_{g,0}^{\operatorname{inv}}(H)^{\times}/\Bbbk^{\times} \longrightarrow \operatorname{PGL}\left(V^m(\Sigma_g^X)\right)$$

$$f \longmapsto \widehat{f} \longmapsto \operatorname{rep. of} \widehat{f}$$

for all *H*-module X and $m \ge 0$.

For a Dehn twist τ_c about a simple closed curve c there is a simple formula for $\hat{\tau}_c$ in terms of the knot map \mathbf{i}_c (Prop. 5.19). This allows us to deduce the following result from Theorem 2, thanks to the presentation of $MCG(\Sigma_{g,0})$ as a quotient of $MCG(\Sigma_{g,0}^{\circ})$ [Waj83]:

Theorem 3. (Thm. 5.16) Let \Bbbk be the trivial *H*-module. For all $m \ge 0$, \mathcal{R}^m_{\Bbbk} descends to a projective representation of $MCG(\Sigma_{g,0})$.

It should be noted that the definition of \mathcal{R}_X^m and the proof of Theorem 3 do not rely on explicit formulas for the representation of generators of the mapping class group. It is nevertheless interesting to have such formulas and we give them in §5.2.3 for the Lickorish generators. Moreover we prove that \mathcal{R}_X^m is equivalent to the representation Z_X^m of [LMSS23] recalled in §1.1 above.

For m = 0, Theorem 3 recovers the projective representations of [AS96a] (which was for semisimple H) and of [Fai20a] (non-semisimple H). The approach with stated skein algebras and knot maps presented here gives a less computational construction of these representations. Moreover, we use new notations (taken from [BFR23]) that should be easier to understand than the matrix formalism employed in previous papers.

Let $S_H(\Sigma_{g,n}^{\circ})$ be the skein algebra spanned by *H*-colored oriented framed links in $\Sigma_{g,n}^{\circ} \times [0, 1]$, modulo the linear relations given by the Reshetikhin–Turaev tangle invariant for *H*-mod. For $H = U_q(\mathfrak{sl}_2)$ this is the Kauffman bracket skein algebra $S_q(\Sigma_{g,n}^{\circ})$. In §5.3 we obtain, as another corollary of Theorem 1, representations of the skein algebra $S_H(\Sigma_{g,n}^{\circ})$ on the spaces $V^m(\Sigma_{g,X_1,\ldots,X_n}^{X})$. This is simply due to the existence of an algebra morphism $W : S_H(\Sigma_{g,n}^{\circ}) \to \mathcal{L}_{g,n}^{\mathrm{inv}}(H)$ which is a variant of (4) and also called Wilson loop map. In particular, for the Kauffman algebra $\mathcal{S}_{\epsilon}(\Sigma_{g,n}^{\circ})$ with q specialized to a root of unity $\epsilon \in \mathbb{C}$, we use a finite-dimensional quotient of $U_{\epsilon^2}(\mathfrak{sl}_2)$, known as *small quantum group* and denoted by $\overline{U}_{\epsilon^2}(\mathfrak{sl}_2)$, to define representations of $\mathcal{S}_{\epsilon}(\Sigma_{g,n}^{\circ})$ on the finite-dimensional spaces $\operatorname{Ext}_{\overline{U}_{\epsilon^2}}^m(X, (\overline{U}_{\epsilon^2}^*)^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n)$ for all $X, X_1, \ldots, X_n \in \overline{U}_{\epsilon^2}(\mathfrak{sl}_2)$ -mod and $m \geq 0$.

Acknowledgements. I thank S. Baseilhac and P. Roche for discussions and encouragements. Part of this work was done when the author was supported by the ANR LabEx CIMI within the French State funding "Investissements d'Avenir".

2 Equivariant modules over *H*-module-algebras

This section contains elementary remarks on representations of H-module-algebras, where H is a Hopf algebra. The key notion is that of H-equivariant module over a H-module-algebra A. We observe that any such module M gives rise to representations of the invariant subalgebra A^{inv} on the spaces $\text{Ext}_{H}^{m}(X, M)$ for any H-module X ("derived representations", §2.1). The derived representations of quantum character varieties constructed in §4 will appear as a corollary of this general fact and of the braided tensor product operation for H-equivariant modules which will be recalled in §3 below. Subsection 2.2 is independent and discusses a situation where any A-module is automatically H-equivariant.

2.1 Derived representations for *H*-equivariant modules

Let $H = (H, \cdot, 1, \Delta, \varepsilon, S)$ be a Hopf algebra with invertible antipode over a field k. We use Sweedler's notation for the coproduct and its iterations:

$$\Delta(h) = h_{(1)} \otimes h_{(2)}, \quad (\Delta \otimes \mathrm{id}) \circ \Delta(h) = h_{(1)} \otimes h_{(2)} \otimes h_{(3)}, \quad etc.$$

Denote by Mod-H the category of (not necessarily finite-dimensional) right modules over H, which is monoidal. We spell out the definitions above in this case. To avoid confusion, in all this paper, the action on a *left* module over a given associative algebra A will be denoted by a dot (\cdot) while the action on a *right* H-module will be denoted by a black square (\bullet).

A (right) *H*-module-algebra is an algebra object $A \in Mod-H$. More explicitly, it is a vector space A endowed with structures of associative algebra and of right *H*-module such that

$$\forall x, y \in A, \ \forall h \in H, \quad (xy) \bullet h = (x \bullet h_{(1)})(y \bullet h_{(2)}), \tag{5}$$

$$\forall h \in H, \quad 1_A \bullet h = \varepsilon(h) 1_A. \tag{6}$$

We use the following name for A-modules objects in Mod-H (taken from [VV10, §1.4] except that here we use right H-modules instead of left ones):

Definition 2.1. A H-equivariant A-module (M, \cdot, \bullet) is a left A-module (M, \cdot) which is also a right H-module (M, \bullet) and such that

$$\forall a \in A, \ \forall v \in M, \ \forall h \in H, \quad (a \cdot v) \bullet h = (a \bullet h_{(1)}) \cdot (v \bullet h_{(2)}). \tag{7}$$

We denote by $(A-Mod)_H$ the category of such modules; its morphisms are linear maps which commutes with both the actions of A and H.

Remark 2.2. In terms of representations, a *H*-equivariant module is a triple (M, ρ, D) where $\rho: A \to \operatorname{End}_{\Bbbk}(M)$ and $\mathsf{D}: H \to \operatorname{End}_{\Bbbk}(M)^{\operatorname{op}}$ are morphisms of algebras such that

$$\forall a \in A, \ \forall h \in H, \quad \rho(a \bullet h) = \mathsf{D}(h_{(1)}) \circ \rho(a) \circ \mathsf{D}\big(S^{-1}(h_{(2)})\big). \tag{8}$$

The relation with the notations in Def. 2.1 is of course $\rho(a)(v) = a \cdot v$ and $\mathsf{D}(h)(v) = v \cdot h$. The superscript ^{op} indicates that D is an anti-morphism (because M is a right H-module).

For any $X, M \in \text{Mod-}H$ let $\text{Hom}_H(X, M)$ be the subspace of linear maps $X \to M$ which commute with the right action of H. In particular when $X = \Bbbk$ is the trivial H-module defined by the counit $\varepsilon : H \to \Bbbk$ we get

$$\operatorname{Hom}_{H}(\mathbb{k}, M) \cong M^{\operatorname{inv}} = \left\{ v \in M \, \middle| \, \forall \, h \in H, \, v \bullet h = \varepsilon(h) v \right\}$$

i.e. the subspace of *H*-invariant elements in *M*. Note that A^{inv} is a subalgebra of *A*.

Lemma 2.3. 1. Any *H*-equivariant *A*-module is an (A^{inv}, H) -bimodule, i.e. the left action of A^{inv} and the right action of *H* commute.

2. For any right H-module X there is a functor

$$\mathcal{V}_X : (A\operatorname{-Mod})_H \to A^{\operatorname{inv}}\operatorname{-Mod}, \quad M \mapsto \operatorname{Hom}_H(X, M).$$

where the action of $a \in A^{inv}$ on $f \in Hom_H(X, M)$ is by acting on the target:

$$\forall x \in X, \quad (a \cdot f)(x) = a \cdot f(x). \tag{9}$$

3. In particular for any $M \in (A-Mod)_H$ the subspace M^{inv} is stable under the action of A^{inv} .

Proof. 1. Obvious from (7).

2. We just need to show that the subspace $\operatorname{Hom}_H(X, M) \subset \operatorname{Hom}_{\Bbbk}(X, M)$ is stable under the action (9), *i.e.* that $a \cdot f$ is *H*-linear. Using item 1 and *H*-linearity of *f*, we have for all $a \in A^{\operatorname{inv}}$:

$$\left((a \cdot f)(x)\right) \bullet h \stackrel{(9)}{=} \left(a \cdot f(x)\right) \bullet h = a \cdot \left(f(x) \bullet h\right) = a \cdot f(x \bullet h) \stackrel{(9)}{=} (a \cdot f)(x \bullet h).$$

3. Immediate consequence of item 1. It is also a consequence of item 2 with $X = \Bbbk$.

We now establish the "derived version" of Lemma 2.3. Let $X, Y \in Mod-H$ and take a projective resolution of X (in Mod-H):

$$0 \longleftarrow X \xleftarrow{d_0} P_0 \xleftarrow{d_1} P_1 \xleftarrow{d_2} \dots$$
(10)

Then the cohomology of the complex of vector spaces

$$\operatorname{Hom}_{H}(P_{0},Y) \xrightarrow{d_{1}^{*}} \operatorname{Hom}_{H}(P_{1},Y) \xrightarrow{d_{2}^{*}} \operatorname{Hom}_{H}(P_{2},Y) \xrightarrow{d_{3}^{*}} \dots$$
(11)

(where d_n^* means pullback) is denoted by $\operatorname{Ext}_H^{\bullet}(X, Y)$, *i.e.* $\operatorname{Ext}_H^0(X, Y) = \ker(d_1^*)$ and $\operatorname{Ext}_H^n(X, Y) = \ker(d_{n+1}^*)/\operatorname{im}(d_n^*)$ for all $n \ge 1$. In other words:

- An element in $\operatorname{Ext}_{H}^{n}(X, Y)$ is a cohomology class [f] where $f \in \operatorname{Hom}_{H}(P_{n}, Y)$ satisfies $f \circ d_{n+1} = 0$, *i.e.* f is a cocycle of degree n in the complex (11).
- Two cohomology classes [f], [f'] are equal if and only if $f f' = g \circ d_n$ for some $g \in$ Hom_H (P_{n-1}, Y) , *i.e.* f - f' is a coboundary of degree n in the complex (11).

Two different projective resolutions of X yield isomorphic Ext spaces, see e.g. [ML75, \S III.6].

Recall that $(A-Mod)_H$ denotes the category of *H*-equivariant *A*-modules (Def. 2.1)

Theorem 2.4. For all $n \in \mathbb{N}$ and any $X \in Mod$ -H there is a functor

$$\mathcal{V}_X^n : (A\operatorname{-Mod})_H \to A^{\operatorname{inv}}\operatorname{-Mod}, \quad M \mapsto \operatorname{Ext}_H^n(X, M).$$

The action of $a \in A^{\text{inv}}$ on a cohomology class $[f] \in \text{Ext}^n_H(X, M)$ is given by $a \cdot [f] = [a \cdot f]$, where $a \cdot f$ is the action on the target M of the cocycle f as defined in (9).

Proof. Take $M \in (A\operatorname{-Mod})_H$ with representation morphism $\rho_M : A \to \operatorname{End}_{\Bbbk}(M)$. Recall the projective resolution (10). By Lemma 2.3 the representation of A^{inv} on $\operatorname{Hom}_H(P_n, M)$, which we denote here by ρ_n , is given by pushforward of ρ_M , *i.e.* $\rho_n(a)(f) = \rho_M(a)_*(f) = \rho_M(a) \circ f$. On the other hand the differential of the complex $\operatorname{Hom}_H(P_{\bullet}, M)$ is given by pullback:

Hence the family of morphisms $(\rho_n(a)_*)_{n\in\mathbb{N}}$ commutes the differential and thus each $\rho_n(a)$ descends to an endomorphism $\bar{\rho}_n(a)$ of $\operatorname{Ext}^n_H(X, M)$ given by $\bar{\rho}_n(a)([f]) = [\rho_n(a)(f)] = [\rho_M(a) \circ f]$. It is clear that $\bar{\rho}_n(ab) = \bar{\rho}_n(a) \circ \bar{\rho}_n(b)$ since this is true for ρ_n . If $\varphi : M \to W$ is a morphism of *H*-equivariant *A*-modules we define $\mathcal{V}^n_X(\varphi)([f]) = [\varphi \circ f]$ for all $[f] \in \operatorname{Ext}^n_H(X, M)$. It is readily seen that $\mathcal{V}^n_X(\varphi)$ is A^{inv} -linear, by *A*-linearity of φ . This defines \mathcal{V}^n_X as a functor.

When the Hopf algebra H is finite-dimensional one might prefer to work within the category mod-H of *finite-dimensional* modules. All of the above remains true with this restriction.

2.2Quantum moment maps and derived representations

Let A be a right H-module-algebra, with right H-action \bullet : $A \otimes H \to A$. The following definition is found in [Lu93, Def. 1.2] and $[VV10, \S1.5]$:

Definition 2.5. A quantum moment map (QMM) is a morphism of algebras $\mu: H \to A$ such that

$$\forall a \in A, \ \forall h \in H, \quad a \,\mu(h) = \mu(h_{(1)}) \left(a \bullet h_{(2)} \right). \tag{12}$$

Until the end of this subsection we assume that a QMM μ is given on A. The property (12) is equivalent to

$$\forall a \in A, \ \forall h \in H, \quad a \bullet h = \mu(S(h_{(1)}))a\mu(h_{(2)}) \tag{13}$$

and in this form we see that the right H-module structure on A is entirely determined by μ . If we put $a = \mu(x)$ in (13) we find in particular

$$\forall x, h \in H, \quad \mu(x) \bullet h = \mu(\operatorname{ad}^{r}(h)(x))$$
(14)

which means that μ is H-linear when H is endowed with the right adjoint action defined by

$$\forall x, h \in H, \quad \mathrm{ad}^r(h)(x) = S(h_{(1)})xh_{(2)}.$$
 (15)

Recall that A^{inv} denotes the subalgebra of *H*-invariant elements in *A*. By (13) it is clear that

$$A^{\text{inv}} = \left\{ a \in A \mid \forall h \in H, \ \mu(h)a = a\mu(h) \right\}.$$

$$(16)$$

In other words A^{inv} is the centralizer of $\mu(H)$.

Having a quantum moment map on A implies that any A-module is automatically Hequivariant $[VV10, \S1.5]$:

Lemma 2.6. Let (M, \cdot) be any A-module. Define

$$\forall v \in M, \ \forall h \in H, \quad v \bullet_{\mu} h = \mu \big(S(h) \big) \cdot v.$$
(17)

Then $(M, \cdot, \bullet_{\mu})$ becomes a H-equivariant A-module. As a result there is a functor

A-Mod \rightarrow (A-Mod)_H, $(M, \cdot) \mapsto (M, \cdot, \bullet_{\mu}).$

Proof. Condition (7) is easily checked:

$$(a \cdot v) \bullet_{\mu} h = \mu (S(h)) a \cdot v = \mu (S(h_{(1)})) a \mu (h_{(2)}) \mu (S(h_{(3)})) \cdot v = (a \bullet h_{(1)}) \cdot (v \bullet_{\mu} h_{(2)})$$

the second equality is a trick and the third equality uses (13).

where the second equality is a trick and the third equality uses (13).

Warning. It is important to distinguish the algebra A and the regular module $_AA$. Indeed the right *H*-action on the algebra *A* is $a \cdot h = \mu(S(h_{(1)}))a\mu(h_{(2)})$ by (13) while the right *H*-action on the A-module _AA provided by Lemma 2.6 is $a \bullet_{\mu} h = \mu(S(h))a$.

Let X be any H-module. If we compose the functors in Lemmas 2.6 and 2.3 we get a functor

$$\mathcal{V}_X : A \operatorname{-Mod} \to A^{\operatorname{inv}} \operatorname{-Mod}, \quad M \mapsto \operatorname{Hom}_H(X, M_\mu).$$

where M_{μ} is M endowed with the H-equivariant structure (17) and the action of A^{inv} on $\operatorname{Hom}_H(X, M_\mu)$ is by acting on the target as in (9). In particular for any $M \in A$ -Mod the subspace

$$M_{\mu}^{\mathrm{inv}} = \mathrm{Hom}_{H}(\mathbb{k}, M_{\mu}) = \left\{ v \in M \mid \forall h \in H, \ \mu(h) \cdot v = \varepsilon(h)v \right\}$$

is stable under the action of A^{inv} . More generally if we compose the functor in Lemma 2.6 with the functor \mathcal{V}_X^n in Theorem 2.4 we get

Corollary 2.7. For all $n \in \mathbb{N}$ and any $X \in Mod$ -H there is a functor

 $\widetilde{\mathcal{V}}_X^n : A\operatorname{-Mod} \to A^{\operatorname{inv}}\operatorname{-Mod}, \quad M \mapsto \operatorname{Ext}^n_H(X, M_\mu).$

The action of $a \in A^{\text{inv}}$ on a cohomology class $[f] \in \text{Ext}^n_H(X, M_\mu)$ is given by $a \cdot [f] = [a \cdot f]$ where $a \cdot f$ denotes the action (9) defined by acting on the target M of the cocycle f.

3 Reminder on braided tensor product

If A_1 , A_2 are associative k-algebras then the vector space $A_1 \otimes_k A_2$ has a canonical algebra structure defined by $(a_1 \otimes a_2)(a'_1 \otimes a'_2) = (a_1a'_1) \otimes (a_2a'_2)$; this is known as *tensor product of* algebras. If M_i is an A_i -module for i = 1, 2 then the vector space $M_1 \otimes_k M_2$ has a canonical structure of $(A_1 \otimes_k A_2)$ -module defined by $(a_1 \otimes a_2)(v_1 \otimes v_2) = (a_1 \cdot v_1) \otimes (a_2 \cdot v_2)$; this is known as *external product of modules* and denoted by $M_1 \boxtimes M_2$. The well-known definitions below generalize these basic facts to other ambient braided monoidal categories than Vect_k .

Let C be a monoidal category, say strict for simplicity. We assume that C is equipped with a *braiding*, which is a natural isomorphism

$$c = \left(c_{X,Y} : X \otimes Y \xrightarrow{\sim} Y \otimes X\right)_{X,Y \in \mathcal{C}}$$

satisfying the usual axioms, see e.g. [Kas95, [XIII.1]. In this case algebras and modules in C interact nicely with the monoidal product:

Lemma 3.1. Let A_1 , A_2 be algebra objects in C, with products $m_i \in \operatorname{Hom}_{\mathcal{C}}(A_i \otimes A_i, A_i)$ and units $\eta_i \in \operatorname{Hom}_{\mathcal{C}}(\mathbf{1}, A_i)$.

1. Define

$$m_{12}: (A_1 \otimes A_2) \otimes (A_1 \otimes A_2) \xrightarrow{\operatorname{id}_{A_1} \otimes c_{A_2,A_1} \otimes \operatorname{id}_{A_2}} A_1 \otimes A_1 \otimes A_2 \otimes A_2 \xrightarrow{m_1 \otimes m_2} A_1 \otimes A_2$$
$$\eta_{12}: \mathbf{1} = \mathbf{1} \otimes \mathbf{1} \xrightarrow{\eta_1 \otimes \eta_2} A_1 \otimes A_2.$$

Then $(A_1 \otimes A_2, m_{12}, \eta_{12})$ is an algebra object in \mathcal{C} , which we denote by $A_1 \otimes A_2$. 2. For i = 1, 2 let M_i be an A_i -module object in \mathcal{C} , with action $\rho_i \in \text{Hom}_{\mathcal{C}}(A_i \otimes M_i, M_i)$. Define

$$\rho_{12}: (A_1 \otimes A_2) \otimes (M_1 \otimes M_2) \xrightarrow{\operatorname{id}_{A_1} \otimes c_{A_2,M_1} \otimes \operatorname{id}_{M_2}} A_1 \otimes M_1 \otimes A_2 \otimes M_2 \xrightarrow{\rho_1 \otimes \rho_2} M_1 \otimes M_2.$$

Then $(M_1 \otimes M_2, \rho_{12})$ is an $(A_1 \otimes A_2)$ -module object in \mathcal{C} , which we denote by $M_1 \otimes M_2$.

Proof. Straightforward computations for both items. The first item is [Maj92, Lem. 9.2.12], where $A_1 \otimes A_2$ is called the *braided tensor product* of A_1 and A_2 .

Let $H = (H, \cdot, 1, \Delta, \varepsilon, S, R)$ is a *quasitriangular* Hopf algebra over a field k, with invertible antipode. We write the *R*-matrix $R \in H^{\otimes 2}$ as

$$R = R_{[1]} \otimes R_{[2]}$$

which is a shorter notation for $R = \sum_{i} R_{[1],i} \otimes R_{[2],i}$. Here are properties of R for future reference [CP94, Kas95]:

Quasi-cocommutativity: $\forall h \in H, \quad R_{[1]}h_{(1)} \otimes R_{[2]}h_{(2)} = h_{(2)}R_{[1]} \otimes h_{(1)}R_{[2]},$ (18)

Quasitriangularity: $R_{1} \otimes R_{[1](2)} \otimes R_{[2]} = R_{[1]}^1 \otimes R_{[1]}^2 \otimes R_{[2]}^1 R_{[2]}^2$,

 $R_{[1]} \otimes R_{[2](1)} \otimes R_{2} = R^1_{[1]} R^2_{[1]} \otimes R^2_{[2]} \otimes R^1_{[2]},$ (20)

(19)

Yang–Baxter equation:
$$R_{[1]}^1 R_{[1]}^2 \otimes R_{[2]}^1 R_{[1]}^3 \otimes R_{[2]}^2 R_{[2]}^3 = R_{[1]}^2 R_{[1]}^1 \otimes R_{[1]}^3 R_{[2]}^1 \otimes R_{[2]}^3 R_{[2]}^2$$
 (21)

Inverse:
$$S(R_{[1]}) \otimes R_{[2]} = R_{[1]} \otimes S^{-1}(R_{[2]}) = R^{-1},$$
 (22)

Consequence of previous:
$$S(R_{[1]}) \otimes S(R_{[2]}) = R_{[1]} \otimes R_{[2]},$$
 (23)

"Skew-inverse":
$$R_{[1]}^{1}R_{[1]}^{2} \otimes S(R_{[2]}^{2})R_{[2]}^{1} = R_{[1]}^{1}S^{-1}(R_{[1]}^{2}) \otimes R_{[2]}^{2}R_{[2]}^{1} = 1 \otimes 1,$$
 (24)

Unitality for counit: $\varepsilon(R_{[1]}) R_{[2]} = R_{[1]} \varepsilon(R_{[2]}) = 1$ (25)

where $R^1 = R^1_{[1]} \otimes R^1_{[2]}$, $R^2 = R^2_{[1]} \otimes R^2_{[2]}$ and $R^3 = R^3_{[1]} \otimes R^3_{[2]}$ denote three copies of R.

Recall that Mod-H denotes the category of right modules over H. The R-matrix induces a braiding on Mod-H given by

 $\forall x \in X, \ \forall y \in Y, \quad c_{X,Y}(x \otimes y) = (y \bullet R_{[1]}) \otimes (x \bullet R_{[2]}).$ (26)

For this choice of ambient category, Lemma 3.1 translates as follows:

Corollary 3.2. Let A_1 and A_2 be right *H*-module-algebras.

1. There is a right H-module-algebra structure on the vector space $A_1 \otimes_{\Bbbk} A_2$, denoted by $A_1 \otimes A_2$, whose product and right H-module structure are given by

$$(a_1 \widetilde{\otimes} a_2)(a_1' \widetilde{\otimes} a_2') = a_1(a_1' \bullet R_{[1]}) \widetilde{\otimes} (a_2 \bullet R_{[2]})a_2',$$

$$(27)$$

$$(a_1 \widetilde{\otimes} a_2) \bullet h = (a_1 \bullet h_{(1)}) \widetilde{\otimes} (a_2 \bullet h_{(2)}).$$
(28)

2. For i = 1, 2, let M_i be a *H*-equivariant A_i -module. There is an $(A_1 \otimes A_2)$ -module structure on the vector space $M_1 \otimes_{\Bbbk} M_2$, denoted by $M_1 \boxtimes M_2$ and given by

$$(a_1 \widetilde{\otimes} a_2) \cdot (v_1 \widetilde{\boxtimes} v_2) = a_1 \cdot (v_1 \bullet R_{[1]}) \widetilde{\boxtimes} (a_2 \bullet R_{[2]}) \cdot v_2,$$

$$(29)$$

$$(v_1 \,\widetilde{\boxtimes} \, v_2) \bullet h = (v_1 \bullet h_{(1)}) \,\widetilde{\boxtimes} \, (v_2 \bullet h_{(2)}). \tag{30}$$

In all the sequel, tensor products are understood to be over k. Note that $a_1 \otimes a_2$ is just a notation for $a_1 \otimes a_2 \in A_1 \otimes A_2$ which allows us to insist on the use of the braided product in $A_1 \otimes A_2$. Similarly $v_1 \boxtimes v_2$ is just a notation for $v_1 \otimes v_2 \in M_1 \otimes M_2$.

As a side remark, quantum moment maps (Def. 2.5) behave well with respect to the braided tensor product of H-module-algebras, see Remark C.3 in Appendix C.

Let A_1, \ldots, A_l be a family of right *H*-module-algebras and let M_j be a *H*-equivariant A_j module for each j. For further use we write down the generalization of formulas (27)–(30) in this situation. It is very convenient to introduce the linear maps $\mathbf{e}_i : A_i \to A_1 \otimes \ldots \otimes A_l$ defined by

$$\mathbf{e}_1(a_1) = a_1 \widetilde{\otimes} \mathbf{1}_{A_2} \widetilde{\otimes} \dots \widetilde{\otimes} \mathbf{1}_{A_l}, \quad \mathbf{e}_2(a_2) = \mathbf{1}_{A_1} \widetilde{\otimes} a_2 \widetilde{\otimes} \mathbf{1}_{A_3} \widetilde{\otimes} \dots \widetilde{\otimes} \mathbf{1}_{A_l}, \quad etc$$
(31)

for all $a_i \in A_i$. The maps \mathfrak{e}_i are morphisms of *H*-module-algebras which moreover satisfy

$$\mathbf{e}_1(a_1)\mathbf{e}_2(a_2)\dots\mathbf{e}_l(a_l) = a_1 \,\widetilde{\otimes}\, a_2 \,\widetilde{\otimes}\,\dots\,\widetilde{\otimes}\, a_l \tag{32}$$

and
$$\mathbf{e}_j(a_j)\mathbf{e}_i(a_i) = \mathbf{e}_i(a_i \bullet R_{[1]})\mathbf{e}_j(a_j \bullet R_{[2]})$$
 if $i < j$ (33)

because of (6) and (25). It is clear that these two properties are enough to compute any product in $A_1 \otimes \ldots \otimes A_l$. The action of $A_1 \otimes \ldots \otimes A_l$ on $M_1 \otimes \ldots \otimes M_l$ is entirely described by

$$\mathbf{e}_{i}(a_{i}) \cdot (v_{1} \,\widetilde{\boxtimes} \, \dots \,\widetilde{\boxtimes} \, v_{l})$$

$$= v_{1} \bullet R_{1} \,\widetilde{\boxtimes} \, \dots \,\widetilde{\boxtimes} \, v_{i-1} \bullet R_{[1](i-1)} \,\widetilde{\boxtimes} \, (a_{i} \bullet R_{[2]}) \cdot v_{i} \,\widetilde{\boxtimes} \, v_{i+1} \,\widetilde{\boxtimes} \, \dots \,\widetilde{\boxtimes} \, v_{l}.$$

$$(34)$$

where the subscripts $(1), \ldots, (i-1)$ indicate iterated coproduct on $R_{[1]}$.

Keeping these notations, we record a few further facts which will be useful in Appendix A. For all i = 1, ..., l consider the associated representation morphisms $\rho_i : A_i \to \operatorname{End}_{\Bbbk}(M_i)$ and $\mathsf{D}_i : H \to \operatorname{End}_{\Bbbk}(M_i)^{\operatorname{op}}$ as in Remark 2.2. By (34), the representation morphism ρ of $A_1 \otimes \ldots \otimes A_l$ on the vector space $M_1 \otimes \ldots \otimes M_l$ is given by

$$\rho: A_1 \stackrel{\sim}{\otimes} \dots \stackrel{\sim}{\otimes} A_l \to \operatorname{End}_{\Bbbk}(M_1) \otimes \dots \otimes \operatorname{End}_{\Bbbk}(M_l)$$

$$\mathfrak{e}_i(a_i) \mapsto \mathsf{D}_1(R_{1}) \otimes \dots \otimes \mathsf{D}_{i-1}(R_{[1](i-1)}) \otimes \rho_i(a_i \bullet R_{[2]}) \otimes \operatorname{id}_{M_{i+1}} \otimes \dots \otimes \operatorname{id}_{M_l}.$$
(35)

The target of ρ is of course the usual (*i.e.* unbraided) tensor product of algebras. We can promote ρ to a morphism of right *H*-module-algebras if we define a right *H*-action on $\operatorname{End}_{\Bbbk}(M_1) \otimes \ldots \otimes \operatorname{End}_{\Bbbk}(M_l)$ by

$$(E_1 \otimes \ldots \otimes E_l) \bullet h = \bigotimes_{i=1}^l \left(\mathsf{D}_i(h_{(i)}) \circ E_i \circ \mathsf{D}_i(S^{-1}(h_{(2l+1-i)})) \right).$$
(36)

Lemma 3.3. Assume that the representation morphisms $\rho_i : A_i \to \operatorname{End}_{\Bbbk}(M_i)$ are injective (resp. surjective) for all *i*. Then ρ defined in (35) is injective (resp. surjective) as well.

Proof. It suffices to prove this for l = 2 because this case implies the result for all l by induction. Note first that:

$$\rho(a_1 \stackrel{\sim}{\otimes} a_2) \stackrel{(32)}{=} \rho(\mathfrak{e}_1(a_1)) \circ \rho(\mathfrak{e}_2(a_2)) \stackrel{(35)}{=} \rho_1(a_1) \circ \mathsf{D}_1(R_{[1]}) \otimes \rho_2(a_2 \bullet R_{[2]})$$
$$\stackrel{(8)}{=} \rho_1(a_1) \circ \mathsf{D}_1(R_{[1]}) \otimes \mathsf{D}_2(R_{[2](1)}) \circ \rho_2(a_2) \circ \mathsf{D}_2(S^{-1}(R_{2}))$$

Hence, using (24) and the fact that D_1 , D_2 and S are anti-morphism of algebras, we can rewrite this as

$$\rho_1(a_1) \otimes \rho_2(a_2) = \left(\mathrm{id}_{M_1} \otimes \mathsf{D}_2(R_{[2](1)}) \right) \circ \rho(a_1 \,\widetilde{\otimes} \, a_2) \circ \left(\mathsf{D}_1(S^{-1}(R_{[1]})) \otimes \mathsf{D}_2(S^{-1}(R_{2})) \right)$$

where \circ is here the product in $\operatorname{End}_{\Bbbk}(M_1) \otimes \operatorname{End}_{\Bbbk}(M_2)$. It follows that if $\rho(a_1^j \otimes a_2^j) = 0$, with implicit summation on j, then $\rho_1(a_1^j) \otimes \rho_2(a_2^j) = 0$. If ρ_1 and ρ_2 are injective, then $\rho_1 \otimes \rho_2$ is also injective and thus $a_1^j \otimes a_2^j = 0$. Surjectivity is proven along the same lines. \Box

4 Derived representations of moduli algebras

Let $H = (H, \cdot, 1, \Delta, \varepsilon, S, R)$ be a quasitriangular Hopf algebra over a field k and recall from §3 that we write $R = R_{[1]} \otimes R_{[2]}$. We introduce some more notations which will be intensively used in the sequel.

The Drinfeld element and its inverse are

$$u = S(R_{[2]})R_{[1]}, \qquad u^{-1} = S^{-2}(R_{[2]})R_{[1]}.$$
 (37)

Its relevance comes from the fact that

$$\forall h \in H, \quad S^2(h) = uhu^{-1}. \tag{38}$$

Let *H*-mod be the category of *finite-dimensional left H*-modules. For any $X \in H$ -mod, $\sigma \in X^*$ and $x \in X$ the linear form

$$_X \phi_x^{\sigma} : H \to \mathbb{k}, \quad h \mapsto \sigma(h \cdot x)$$
 (39)

is called a *matrix coefficient* of X. The subspace $H^{\circ} \subset H^{*}$ spanned by matrix coefficients of objects in *H*-mod is called *restricted dual* of *H*. When *H* is finite-dimensional one has $H^{\circ} = H^{*}$ because any linear form $\varphi \in H$ is the matrix coefficient $_{H}\phi_{1}^{\varphi}$ of the regular module *H*. The vector space H° has the structure of a Hopf algebra which is dual to that of *H*. Its product \star is defined by

$${}_X\phi_x^{\sigma} \star {}_Y\phi_y^{\tau} = {}_{X\otimes Y}\phi_{x\otimes y}^{\sigma\otimes \tau} \qquad i.e. \quad \forall h \in H, \quad (\varphi \star \psi)(h) = \varphi(h_{(1)})\psi(h_{(2)}) \tag{40}$$

while its coproduct is defined on matrix coefficients by

$$\Delta({}_X\phi^{\sigma}_x) = {}_X\phi^{\sigma}_{x_i} \otimes {}_X\phi^{x^i}_x \tag{41}$$

with implicit summation on i, where (x_i) is a basis of X and (x^i) is its dual basis. It is readily seen that

$$\forall \varphi \in H^{\circ}, \ \forall h, h' \in H, \quad \varphi_{(1)}(h) \varphi_{(2)}(h') = \varphi(hh').$$

$$(42)$$

The unit is $1_{H^{\circ}} = \varepsilon = {}_{\Bbbk} \phi_1^1$, namely the counit of H. The counit $\varepsilon_{H^{\circ}}$ (resp. antipode $S_{H^{\circ}}$) is defined by $\varepsilon_{H^{\circ}}(\varphi) = \varphi(1_H)$ (resp. $S_{H^{\circ}}(\varphi) = \varphi \circ S$).

The left and right coregular actions $\triangleright, \triangleleft$ of H on H^* are defined by

$$\forall h, h' \in H, \ \forall \varphi \in H^*, \quad (h \triangleright \varphi)(h') = \varphi(h'h), \quad (\varphi \triangleleft h)(h') = \varphi(hh').$$
(43)

The subspace $H^{\circ} \subset H^*$ is stable by these actions; actually, even the subspace spanned by the matrix coefficients of a given *H*-module *X* is stable:

$$\forall h \in H, \quad h \triangleright_X \phi_x^{\sigma} = {}_X \phi_{h \cdot x}^{\sigma}, \quad {}_X \phi_x^{\sigma} \triangleleft h = {}_X \phi_x^{\sigma(h \cdot ?)} \tag{44}$$

where $\sigma(h \cdot ?) \in X^*$ is the linear form $x' \mapsto \sigma(h \cdot x')$. Note that $h \triangleright \varphi = \varphi_{(2)}(h) \varphi_{(1)}$ and $\varphi \triangleleft h = \varphi_{(1)}(h) \varphi_{(2)}$. Also

$$h \triangleright (\varphi \star \psi) = (h_{(1)} \triangleright \varphi) \star (h_{(2)} \triangleright \psi), \quad (\varphi \star \psi) \triangleleft h = (\varphi \triangleleft h_{(1)}) \star (\psi \triangleleft h_{(2)})$$
(45)

for all $h \in H$ and $\varphi, \psi \in H^{\circ}$.

4.1 Definition of $\mathcal{L}_{g,n}(H)$ and $\mathcal{L}_{g,n}^{inv}(H)$

Our notations are those of [BFR23, §3.1, §4.1] where a construction of $\mathcal{L}_{g,n}(H)$ using twists is provided. Here we just recall the resulting definitions.

We first treat the cases (g, n) = (0, 1) and (g, n) = (1, 0) which are the building blocks of the general definition.

Definition 4.1. The right H-module-algebra $\mathcal{L}_{0,1}(H)$ is the vector space H° endowed with the product

$$\varphi\psi = \left(R_{[2]}^2 S(R_{[2]}^1) \triangleright \varphi\right) \star \left(R_{[1]}^2 \triangleright \psi \triangleleft R_{[1]}^1\right) \tag{46}$$

and with the right coadjoint action of H:

$$\forall h \in H, \ \forall \varphi \in H^{\circ}, \quad \operatorname{coad}^{r}(h)(\varphi) = S(h_{(2)}) \triangleright \varphi \triangleleft h_{(1)} = \varphi(h_{(1)}?S(h_{(2)})). \tag{47}$$

Here R^1, R^2 denote two copies of the *R*-matrix $R \in H \otimes H$, \star is defined in (40) and $\triangleright, \triangleleft$ are defined in (43). The product in $\mathcal{L}_{0,1}(H)$ is associative (straightforward computation using (21)) and has unit element ε , the counit of *H*. The module-algebra axioms are easily checked.

Definition 4.1 is a version of Majid's transmutation of H° [Maj92, Th. 7.4.1 and Ex. 9.4.10] (but here we work with *H*-modules instead of H° -comodules), also called "braided dual". For finite-dimensional *H* it is a variant of the Lyubashenko coend $\int^{X \in H \text{-mod}} X^* \otimes X$.

For later use we recall that $\mathcal{L}_{0,1}(H)$ is a Hopf algebra in Mod-H with the coproduct $\Delta_{H^{\circ}}$ defined in (41), but now viewed as a morphism of right H-module-algebras $\mathcal{L}_{0,1}(H) \rightarrow \mathcal{L}_{0,1}(H) \otimes \mathcal{L}_{0,1}(H)$. The antipode is given by

$$S_{\mathcal{L}_{0,1}}: \mathcal{L}_{0,1}(H) \to \mathcal{L}_{0,1}(H), \quad \varphi \mapsto S_{H^{\circ}}\big(S(R_{[1]}) \triangleright \varphi \triangleleft R_{[2]}u^{-1}\big) = \varphi\big(R_{[2]}u^{-1}S(?)S(R_{[1]})\big) \quad (48)$$

where $S_{H^{\circ}} = S^*$ is the antipode on H° and u is the Drinfeld element (37). The counit is the same as in H° , *i.e.* $\varphi \mapsto \varphi(1_H)$. By [Maj92, eq. (9.39)] the antipode of a Hopf algebra in a braided category is an antimorphism, but where one uses the braiding to switch the factors. If we apply this to $\mathcal{L}_{0,1}(H)$ in the category Mod-H with the braiding (26), it means that

$$S_{\mathcal{L}_{0,1}}(\varphi\psi) = S_{\mathcal{L}_{0,1}}(\text{coad}^{r}(R_{[1]})(\psi)) S_{\mathcal{L}_{0,1}}(\text{coad}^{r}(R_{[2]})(\varphi)).$$
(49)

One can also check directly this equality from the definitions, but the computation is involved.

The subalgebra of *H*-invariant elements in $\mathcal{L}_{0,1}(H)$ is

$$\mathcal{L}_{0,1}^{\mathrm{inv}}(H) = \left\{ \varphi \in \mathcal{L}_{0,1}(H) \, \big| \, \forall h \in H, \, \operatorname{coad}^{r}(h)(\varphi) = \varepsilon(h)\varphi \right\} \\ = \left\{ \varphi \in \mathcal{L}_{0,1}(H) \, \big| \, \forall x, y \in H, \, \varphi(xy) = \varphi(yS^{2}(x)) \right\}.$$
(50)

Example 4.2. For a finite-dimensional H-module X let $qTr_X(h) = Tr(\rho_X(uh))$ where $\rho_X : H \to End_k(X)$ is the representation morphism, u is the Drinfeld element (37) and Tr is the trace. It follows from (38) and the cyclicity of the trace that $qTr_X \in \mathcal{L}_{0,1}^{inv}(H)$. When H-mod is semisimple these are all the elements of $\mathcal{L}_{0,1}^{inv}(H)$, see Lemma B.1. If H has a ribbon element v one usually prefers to use the pivotal element $g = uv^{-1}$ instead of u for defining qTr_X .

Lemma 4.3. For all $\varphi \in \mathcal{L}_{0,1}(H)$ and $\psi \in \mathcal{L}_{0,1}^{inv}(H)$ we have $\varphi \psi = \varphi \star \psi = \psi \varphi$. In particular $\mathcal{L}_{0,1}^{inv}(H) \subset \mathcal{Z}(\mathcal{L}_{0,1}(H))$, where \mathcal{Z} denotes the center.

Proof. For all $\varphi, \psi \in \mathcal{L}_{0,1}(H)$ we can rewrite the product (46) as

$$\varphi\psi \stackrel{(23)}{=} \left(S(R_{[2]}^1 R_{[2]}^2) \triangleright \varphi\right) \star \left(S(R_{[1]}^2) \triangleright \psi \triangleleft R_{[1]}^1\right) \stackrel{(19)}{=} \left(S(R_{[2]}) \triangleright \varphi\right) \star \operatorname{coad}^r(R_{[1]})(\psi)$$

and also as

$$\psi\varphi \stackrel{(18)}{=} \left(\varphi \triangleleft R_{[1]}^1 R_{[1]}^2\right) \star \left(S(R_{[2]}^1) \triangleright \psi \triangleleft R_{[2]}^2\right) \stackrel{(19)}{=} \left(\varphi \triangleleft R_{[1]}\right) \star \operatorname{coad}^r(R_{[2]})(\psi).$$

If $\psi \in \mathcal{L}_{0,1}^{\text{inv}}(H)$ the *R*-matrices disappear because $(\varepsilon \otimes \text{id})(R) = (\text{id} \otimes \varepsilon)(R) = 1$.

The inclusion in Lemma 4.3 is actually an equality for $H = U_q(\mathfrak{g})$ [BR19, Prop. 6.19] and for H finite-dimensional and factorizable [Fai19a, Th. 3.7].

Let $\mathfrak{i}_b, \mathfrak{i}_a: H^\circ \to H^\circ \otimes H^\circ$ be the linear embeddings given by $\mathfrak{i}_b(\varphi) = \varphi \otimes \varepsilon, \mathfrak{i}_a(\varphi) = \varepsilon \otimes \varphi.$

Definition 4.4. The right *H*-module-algebra $\mathcal{L}_{1,0}(H)$ is the vector space $H^{\circ} \otimes H^{\circ}$ and its structure is uniquely characterized by the following conditions:

- the linear maps i_b, i_a are morphisms of H-module-algebras $\mathcal{L}_{0,1}(H) \to \mathcal{L}_{1,0}(H)$,
- for all $\beta, \alpha \in \mathcal{L}_{0,1}(H)$ it holds

$$\mathbf{i}_b(\beta)\mathbf{i}_a(\alpha) = \beta \otimes \alpha,\tag{51}$$

$$\mathbf{i}_{a}(\alpha)\,\mathbf{i}_{b}(\beta) = \mathbf{i}_{b}\left(R_{[2]}^{4}R_{[1]}^{3} \triangleright \beta \triangleleft R_{[1]}^{1}R_{[1]}^{2}\right)\,\mathbf{i}_{a}\left(R_{[2]}^{3}S(R_{[2]}^{1}) \triangleright \alpha \triangleleft R_{[2]}^{2}R_{[1]}^{4}\right) \tag{52}$$

where R^1, \ldots, R^4 denote four copies of the *R*-matrix.

See [BFR23, §3.1] for more details. The unit element is $\varepsilon \otimes \varepsilon$. As in the case of $\mathcal{L}_{0,1}(H)$, we denote by coad^r the right action of H on $\mathcal{L}_{1,0}(H)$. Due to (51) and (5) we see that it is just the diagonal action of (47):

$$\forall h \in H, \ \forall \beta \otimes \alpha \in \mathcal{L}_{1,0}(H), \quad \text{coad}^r(h)(\beta \otimes \alpha) = \text{coad}^r(h_{(1)})(\beta) \otimes \text{coad}^r(h_{(2)})(\alpha).$$
(53)

Said differently, $\mathcal{L}_{1,0}(H)$ is $\mathcal{L}_{0,1}(H) \otimes \mathcal{L}_{0,1}(H)$ as a right *H*-module.

For the general definition we use the braided tensor product $\tilde{\otimes}$ in Mod-*H* (recalled in §3) of many copies of the *H*-module-algebras $\mathcal{L}_{1,0}(H)$ and $\mathcal{L}_{0,1}(H)$:

Definition 4.5.
$$\mathcal{L}_{g,n}(H) = \underbrace{\mathcal{L}_{1,0}(H) \otimes \ldots \otimes \mathcal{L}_{1,0}(H)}_{g \ copies} \otimes \underbrace{\mathcal{L}_{0,1}(H) \otimes \ldots \otimes \mathcal{L}_{0,1}(H)}_{n \ copies}$$

As a vector space, $\mathcal{L}_{g,n}(H)$ is $(H^{\circ})^{\otimes(2g+n)}$. By construction it is a right *H*-module-algebra for the diagonal action of (47), which we denote again by coad^r :

$$\operatorname{coad}^{r}(h)(\varphi_{1}\otimes\ldots\otimes\varphi_{2g+n})=\operatorname{coad}^{r}(h_{(1)})(\varphi_{1})\otimes\ldots\otimes\operatorname{coad}^{r}(h_{(2g+n)})(\varphi_{2g+n}).$$
(54)

Said differently, $\mathcal{L}_{g,n}(H)$ is $\mathcal{L}_{0,1}(H)^{\otimes (2g+n)}$ as a right *H*-module.

In order to describe the product in $\mathcal{L}_{g,n}(H)$ it is convenient to introduce some notations. Let $b_1, a_1, \ldots, b_g, a_g, m_{g+1}, \ldots, m_{g+n}$ be formal symbols. Later they will be interpreted as generators for the fundamental group of the oriented surface of genus g with n punctures and one boundary component. Label each copy of $\mathcal{L}_{0,1}(H)$ in $\mathcal{L}_{g,n}(H)$ by a symbol as follows:

$$\mathcal{L}_{g,n}(H) = \underbrace{\left(\mathcal{L}_{1,0}(H) \otimes \mathcal{L}_{0,1}(H)\right)}_{b_1} \otimes \mathcal{L}_{0,1}(H)\right)}_{b_1} \otimes \ldots \otimes \underbrace{\left(\mathcal{L}_{1,0}(H) \otimes \mathcal{L}_{0,1}(H)\right)}_{b_g} \otimes \mathcal{L}_{0,1}(H)\right)}_{b_g} \otimes \mathcal{L}_{0,1}(H) \otimes \ldots \otimes \mathcal{L}_{0,1}(H).$$

This gives H-linear embeddings⁴

$$\mathfrak{i}_{b_i}, \,\mathfrak{i}_{a_i}, \,\mathfrak{i}_{m_{g+j}}: \mathcal{L}_{0,1}(H) \to \mathcal{L}_{g,n}(H) \qquad (1 \le i \le g, \ 1 \le j \le n).$$

$$(55)$$

For instance $\mathbf{i}_{b_1}(\varphi) = \varphi \otimes \varepsilon^{\otimes (2g+n-1)}$, where ε is the counit of H. The following conditions [BFR23, Prop. 4.4] entirely characterize the H-module-algebra structure in $\mathcal{L}_{g,n}(H)$:

Proposition 4.6. 1. For all $1 \leq i \leq g$ and $1 \leq j \leq n$ the embeddings \mathfrak{i}_{b_i} , \mathfrak{i}_{a_i} , $\mathfrak{i}_{m_{g+j}} : \mathcal{L}_{0,1}(H) \rightarrow \mathcal{L}_{g,n}(H)$ are morphisms of H-module-algebras.

2. For all $\beta_1, \alpha_1, \ldots, \beta_g, \alpha_g, \gamma_1, \ldots, \gamma_n \in \mathcal{L}_{0,1}(H)$ we have

$$\mathbf{i}_{b_1}(\beta_1) \, \mathbf{i}_{a_1}(\alpha_1) \dots \, \mathbf{i}_{b_g}(\beta_g) \, \mathbf{i}_{a_g}(\alpha_g) \, \mathbf{i}_{m_{g+1}}(\gamma_1) \dots \, \mathbf{i}_{m_{g+n}}(\gamma_n) \\ = \beta_1 \otimes \alpha_1 \otimes \dots \otimes \beta_g \otimes \alpha_g \otimes \gamma_1 \otimes \dots \otimes \gamma_n.$$

3. For all $1 \leq i \leq g$ and all $\varphi, \psi \in \mathcal{L}_{0,1}(H)$ we have

$$\mathfrak{i}_{a_i}(\varphi)\,\mathfrak{i}_{b_i}(\psi) = \mathfrak{i}_{b_i}\big(R^4_{[2]}R^3_{[1]} \triangleright \psi \triangleleft R^1_{[1]}R^2_{[1]}\big)\,\mathfrak{i}_{a_i}\big(R^3_{[2]}S(R^1_{[2]}) \triangleright \varphi \triangleleft R^2_{[2]}R^4_{[1]}\big).$$

4. Define $\operatorname{ind}(b_i) = \operatorname{ind}(a_i) = i$ and $\operatorname{ind}(m_{g+j}) = g + j$ and let s, t be symbols⁵ with $\operatorname{ind}(t) > \operatorname{ind}(s)$. For all $\varphi, \psi \in \mathcal{L}_{0,1}(H)$ and $1 \leq i < j \leq g + n$ we have

$$\begin{split} \mathbf{i}_{t}(\varphi)\,\mathbf{i}_{s}(\psi) &= \mathbf{i}_{s}\left(\mathrm{coad}^{r}(R_{[1]})(\psi)\right)\,\mathbf{i}_{t}\left(\mathrm{coad}^{r}(R_{[2]})(\varphi)\right) \\ &= \mathbf{i}_{s}\left(S(R_{[1]}^{3}R_{[1]}^{4}) \triangleright \psi \triangleleft R_{[1]}^{1}R_{[1]}^{2}\right)\,\mathbf{i}_{t}\left(S(R_{[2]}^{1}R_{[2]}^{3}) \triangleright \varphi \triangleleft R_{[2]}^{2}R_{[2]}^{4}\right). \end{split}$$

In the two items above R^1, \ldots, R^4 are four copies of the *R*-matrix.

Proof. Note first that by (31) we have embeddings of algebras

$$\mathfrak{e}_1, \ldots, \mathfrak{e}_g : \mathcal{L}_{1,0}(H) \to \mathcal{L}_{g,n}(H) \quad \text{and} \quad \mathfrak{e}_{g+1}, \ldots, \mathfrak{e}_{g+n} : \mathcal{L}_{0,1}(H) \to \mathcal{L}_{g,n}(H).$$

By definition, $\mathbf{i}_{b_i} = \mathbf{e}_i \circ \mathbf{i}_b$, $\mathbf{i}_{a_i} = \mathbf{e}_i \circ \mathbf{i}_a$ and $\mathbf{i}_{m_{g+j}} = \mathbf{e}_{g+j}$, where $\mathbf{i}_b, \mathbf{i}_a : \mathcal{L}_{0,1}(H) \to \mathcal{L}_{1,0}(H)$ are morphisms of algebras by Definition 4.4. Thus item 1 is proven. For item 2 we note that $\mathbf{i}_{b_i}(\beta_i)\mathbf{i}_{a_i}(\alpha_i) = \mathbf{e}_i(\beta_i \otimes \alpha_i)$ and use (32). Item 3 is obtained by applying \mathbf{e}_i to (52). Finally item 4 comes from (33) and (19)-(20).

In any H-module-algebra the subspace of elements which are H-invariant is a subalgebra, so we can make the following definition:

Definition 4.7. The moduli algebra, or quantum character variety, is

$$\mathcal{L}_{g,n}^{\text{inv}}(H) = \left\{ x \in \mathcal{L}_{g,n}(H) \, \middle| \, \forall \, h \in H, \, \text{coad}^r(h)(x) = \varepsilon(h)x \right\}$$

with coad^r from (54) and ε is the counit of H.

⁴These embeddings were denoted by $\mathfrak{i}_{B(i)}$, $\mathfrak{i}_{A(i)}$, $\mathfrak{i}_{M(g+j)}$ in [BFR23]. This slight change of notation will be more convenient for later purpose.

⁵That is elements in the list $b_1, a_1, \ldots, b_g, a_g, m_{g+1}, \ldots, m_{g+n}$.

4.2 *H*-equivariant $\mathcal{L}_{q,n}(H)$ -modules and derived representations

Since the seminal work by Alekseev [Ale94], it is known that $\mathcal{L}_{g,n}(H)$ has representations on the vector spaces $(H^{\circ})^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n$ where each X_i is a *H*-module. Our purpose here is to put a *H*-equivariant structure (Def. 2.1) on these spaces and to deduce derived representations for $\mathcal{L}_{g,n}^{inv}(H)$ thanks to Theorem 2.4.

For this we first define *H*-equivariant representations of $\mathcal{L}_{0,1}(H)$ on any *H*-module *X* and of $\mathcal{L}_{1,0}(H)$ on H° and then assemble them thanks to the braided product of equivariant modules (Cor. 3.2). This will result in a representation of $\mathcal{L}_{g,n}(H)$ on $(H^{\circ})^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n$ which is *H*-equivariant and is equivalent to the one obtained from the Alekseev morphism, see App. A.

Let ad^r be the right adjoint action (15) of H on itself; then (H, ad^r) is a right H-modulealgebra. Consider the linear map

$$\Phi_{0,1}: \mathcal{L}_{0,1}(H) \to H, \quad \varphi \mapsto (\varphi \otimes \mathrm{id}_H)(RR')$$
(56)

where $R' = R_{[2]} \otimes R_{[1]}$ is the flip of R. It is well-known and not difficult to check that $\Phi_{0,1}$ is a morphism of H-module-algebras $(\mathcal{L}_{0,1}(H), \operatorname{coad}^r) \to (H, \operatorname{ad}^r)$.

Lemma 4.8. Let (X, \cdot) be any left H-module and define

$$\forall \varphi \in \mathcal{L}_{0,1}(H), \ \forall h \in H, \ \forall x \in X, \quad \varphi \cdot x = \Phi_{0,1}(\varphi) \cdot x, \quad x \bullet h = S(h) \cdot x.$$

With this structure, X is a H-equivariant $\mathcal{L}_{0,1}(H)$ -module.

Proof. The *H*-equivariance condition (7) follows from the *H*-linearity of $\Phi_{0,1}$:

$$\operatorname{coad}^{r}(h_{(1)})(\varphi) \cdot (x \bullet h_{(2)}) = \Phi_{0,1} \left(\operatorname{coad}^{r}(h_{(1)})(\varphi) \right) S(h_{(2)}) \cdot x = \operatorname{ad}^{r}(h_{(1)}) \left(\Phi_{0,1}(\varphi) \right) S(h_{(2)}) \cdot x \\ = S(h_{(1)}) \Phi_{0,1}(\varphi) h_{(2)} S(h_{(3)}) \cdot x = S(h) \Phi_{0,1}(\varphi) \cdot x = (\varphi \cdot x) \bullet h. \quad \Box$$

Next we define a $\mathcal{L}_{1,0}(H)$ -module structure on the restricted dual H° by means of an induced representation. This strategy was already used in [AS96a, §7.2]; here we promote it to a H-equivariant representation. The resulting module is cyclic, generated by the counit ε :

Lemma 4.9. There is a structure of H-equivariant $\mathcal{L}_{1,0}(H)$ -module on H° given by

$$\mathbf{i}_{b}(\varphi) \cdot \psi = \varphi \psi, \quad \mathbf{i}_{a}(\varphi) \cdot \psi = \Phi_{0,1}(\operatorname{coad}^{r}(R_{[2]})(\varphi)) \triangleright \psi \triangleleft R_{[1]},$$
$$\psi \bullet h = \operatorname{coad}^{r}(h)(\psi)$$

for all $\varphi \in \mathcal{L}_{0,1}(H)$, $\psi \in H^{\circ}$, $h \in H$ and where $\varphi \psi$ is the product (46), coad^r is defined in (47).

Proof. Let $\mathcal{A} = \{ \mathfrak{i}_a(\varphi) | \varphi \in \mathcal{L}_{0,1}(H) \}$, which is a subalgebra of $\mathcal{L}_{1,0}(H)$ isomorphic to $\mathcal{L}_{0,1}(H)$. Lemma 4.8 gives a representation of $\mathcal{L}_{0,1}(H)$ on the ground field \Bbbk given by $\varphi \cdot 1_{\Bbbk} = \varphi(1_H) 1_{\Bbbk}$ due to (25). Consider the induced representation $\mathcal{L}_{1,0}(H) \otimes_{\mathcal{A}} \Bbbk$. Recall from (51) that any element in $\mathcal{L}_{1,0}(H)$ can be written as a linear combination of products of the form $\mathfrak{i}_b(\beta)\mathfrak{i}_a(\alpha)$ with $\alpha, \beta \in \mathcal{L}_{0,1}(H)$. Since

$$\mathbf{i}_b(\beta)\,\mathbf{i}_a(\alpha)\otimes_{\mathcal{A}}\mathbf{1}_{\Bbbk} = \mathbf{i}_b(\beta)\otimes_{\mathcal{A}}\alpha\cdot\mathbf{1}_{\Bbbk} = \alpha(\mathbf{1}_H)\,\mathbf{i}_b(\beta)\otimes_{\mathcal{A}}\mathbf{1}_{\Bbbk} \tag{57}$$

we have an isomorphism of vector spaces

$$\mathcal{L}_{1,0}(H) \otimes_{\mathcal{A}} \Bbbk \xrightarrow{\sim} H^{\circ}, \quad \mathfrak{i}_{b}(\psi) \otimes_{\mathcal{A}} 1_{\Bbbk} \mapsto \psi.$$
(58)

Let us compute the resulting representation of $\mathcal{L}_{1,0}(H)$ on H° . We have

$$\mathfrak{i}_b(\varphi)\cdot\psi\cong\mathfrak{i}_b(\varphi)\mathfrak{i}_b(\psi)\otimes_{\mathcal{A}}1_\Bbbk=\mathfrak{i}_b(\varphi\psi)\otimes_{\mathcal{A}}1_\Bbbk\cong\varphi\psi$$

for all $\varphi \in \mathcal{L}_{0,1}(H)$ and $\psi \in H^{\circ}$, where \cong denotes the identification (58). This shows in particular that this representation is cyclic, because $\psi = \mathbf{i}_b(\psi) \cdot \varepsilon$ for all $\psi \in H^{\circ}$, where ε is the counit of H. For the action of $\mathbf{i}_a(\varphi)$, note that (57) is rewritten as

$$\mathbf{i}_b(\beta)\,\mathbf{i}_a(\alpha)\cdot\boldsymbol{\varepsilon} = (\beta\otimes\alpha)\cdot\boldsymbol{\varepsilon} = \alpha(\mathbf{1}_H)\,\beta\tag{59}$$

Hence

$$\begin{aligned} \mathbf{i}_{a}(\varphi) \cdot \psi &= \mathbf{i}_{a}(\varphi) \, \mathbf{i}_{b}(\psi) \cdot \varepsilon \stackrel{(52)}{=} \mathbf{i}_{b} \left(R_{[2]}^{4} R_{[1]}^{3} \triangleright \psi \triangleleft R_{[1]}^{1} R_{[1]}^{2} \right) \, \mathbf{i}_{a} \left(R_{[2]}^{3} S(R_{[2]}^{1}) \triangleright \varphi \triangleleft R_{[2]}^{2} R_{[1]}^{4} \right) \cdot \varepsilon \\ &= \varphi \left(R_{[2]}^{2} R_{[1]}^{4} R_{[2]}^{3} S(R_{[2]}^{1}) \right) R_{[2]}^{4} R_{[1]}^{3} \triangleright \psi \triangleleft R_{[1]}^{1} R_{[1]}^{2} \stackrel{(20)}{=} \varphi \left(R_{[2](1)}^{1} R_{[1]}^{4} R_{[2]}^{3} S(R_{2}^{1}) \right) R_{[2]}^{4} R_{[1]}^{3} \triangleright \psi \triangleleft R_{[1]}^{1} \end{aligned}$$

which is the announced formula, by definitions of coad^r in (47) and $\Phi_{0,1}$ in (56). To prove the *H*-equivariant property, note first that for all $\beta \otimes \alpha \in \mathcal{L}_{1,0}(H)$

$$\operatorname{coad}^{r}(h)(\beta \otimes \alpha) \cdot \varepsilon \stackrel{(53)}{=} (\operatorname{coad}^{r}(h_{(1)})(\beta) \otimes \operatorname{coad}^{r}(h_{(2)})(\alpha)) \cdot \varepsilon \stackrel{(59),(47)}{=} \alpha(S(h_{(2)})h_{(3)}) \operatorname{coad}^{r}(h_{(1)})(\beta)$$
$$= \alpha(1_{H}) \operatorname{coad}^{r}(h)(\beta) \stackrel{(59)}{=} \operatorname{coad}^{r}(h)((\beta \otimes \alpha) \cdot \varepsilon).$$

It follows that $\operatorname{coad}^r(h)(x) \cdot \varepsilon = \operatorname{coad}^r(h)(x \cdot \varepsilon)$ for all $x \in \mathcal{L}_{1,0}(H)$. Since $\mathcal{L}_{1,0}(H)$ is *H*-modulealgebra under coad^r , we deduce that condition (7) holds as follows:

$$\operatorname{coad}^{r}(h_{(1)})(x) \cdot \operatorname{coad}^{r}(h_{(2)})(\psi) = \operatorname{coad}^{r}(h_{(1)})(x) \,\mathfrak{i}_{b}\left(\operatorname{coad}^{r}(h_{(2)})(\psi)\right) \cdot \varepsilon$$
$$= \operatorname{coad}^{r}(h)\left(x \,\mathfrak{i}_{b}(\psi)\right) \cdot \varepsilon = \operatorname{coad}^{r}(h)\left(x \,\mathfrak{i}_{b}(\psi) \cdot \varepsilon\right) = \operatorname{coad}^{r}(h)(x \cdot \psi)$$

for all $x \in \mathcal{L}_{1,0}(H)$, $\psi \in H^{\circ}$ and $h \in H$.

Recall that by definition $\mathcal{L}_{g,n}(H) = \mathcal{L}_{1,0}(H)^{\widetilde{\otimes} g} \widetilde{\otimes} \mathcal{L}_{0,1}(H)^{\widetilde{\otimes} n}$. Combining Lemmas 4.8 and 4.9 thanks to item 2 in Corollary 3.2, we obtain the *H*-equivariant $\mathcal{L}_{g,n}(H)$ -modules

$$(H^{\circ})^{\widetilde{\boxtimes} g} \widetilde{\boxtimes} X_1 \widetilde{\boxtimes} \dots \widetilde{\boxtimes} X_n \tag{60}$$

where X_1, \ldots, X_n are any left *H*-modules endowed with the *H*-equivariant structure from Lemma 4.8. As a vector space this is just $(H^\circ)^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n$. We write elements in the module (60) as $\psi_1 \otimes \ldots \otimes \psi_g \otimes x_1 \otimes \ldots \otimes x_n$ instead of the heavy notation $\psi_1 \boxtimes \ldots \boxtimes \psi_g \boxtimes x_1 \boxtimes \ldots \boxtimes x_n$.

In order to describe explicitly the structure on (60) we use the embeddings $\mathfrak{i}_{b_i}, \mathfrak{i}_{a_i}, \mathfrak{i}_{m_{g+j}} : \mathcal{L}_{0,1}(H) \to \mathcal{L}_{g,n}(H)$ from (55), which by Prop. 4.6 are the main tool to compute in $\mathcal{L}_{g,n}(H)$.

Proposition 4.10. The right H-module structure on (60) is

$$\begin{pmatrix} \psi_1 \otimes \ldots \otimes \psi_g \otimes x_1 \otimes \ldots \otimes x_n \end{pmatrix} \bullet h = \operatorname{coad}^r(h_{(1)})(\psi_1) \otimes \ldots \otimes \operatorname{coad}^r(h_{(g)})(\psi_g) \otimes S(h_{(1)}) \cdot x_1 \otimes \ldots \otimes S(h_{(g+n)}) \cdot x_n.$$

$$(61)$$

The left $\mathcal{L}_{g,n}(H)$ -module structure on (60) is given by

$$\begin{split} \mathbf{i}_{b_i}(\varphi) \cdot \left(\psi_1 \otimes \ldots \otimes \psi_g \otimes x_1 \otimes \ldots \otimes x_n\right) \\ &= \operatorname{coad}^r(R_{1})(\psi_1) \otimes \ldots \otimes \operatorname{coad}^r(R_{[1](i-1)})(\psi_{i-1}) \otimes \operatorname{coad}^r(R_{[2]})(\varphi)\psi_i \\ &\otimes \psi_{i+1} \otimes \ldots \otimes \psi_g \otimes x_1 \otimes \ldots \otimes x_n, \\ \\ &\mathbf{i}_{a_i}(\varphi) \cdot \left(\psi_1 \otimes \ldots \otimes \psi_g \otimes x_1 \otimes \ldots \otimes x_n\right) \\ &= \operatorname{coad}^r(R_{1}^1)(\psi_1) \otimes \ldots \otimes \operatorname{coad}^r(R_{[1](i-1)}^1)(\psi_{i-1}) \otimes \Phi_{0,1}\left(\operatorname{coad}^r(R_{[2]}^1R_{[2]}^2)(\varphi)\right) \triangleright \psi_i \triangleleft R_{[1]}^2 \\ &\otimes \psi_{i+1} \otimes \ldots \otimes \psi_g \otimes x_1 \otimes \ldots \otimes x_n. \end{split}$$

$$\begin{aligned} \mathbf{i}_{m_{g+j}}(\varphi) \cdot \left(\psi_1 \otimes \ldots \otimes \psi_g \otimes x_1 \otimes \ldots \otimes x_n\right) \\ &= \operatorname{coad}^r(R_{1})(\psi_1) \otimes \ldots \otimes \operatorname{coad}^r(R_{[1](g)})(\psi_g) \otimes S(R_{[1](g+1)}) \cdot x_1 \\ &\otimes \ldots \otimes S(R_{[1](g+j-1)}) \cdot x_{j-1} \otimes \Phi_{0,1}\left(\operatorname{coad}^r(R_{[2]})(\varphi)\right) \cdot x_j \otimes x_{j+1} \otimes \ldots \otimes x_n. \end{aligned}$$

for all $\varphi \in \mathcal{L}_{0,1}(H)$, with $1 \leq i \leq g$, $1 \leq j \leq n$, where coad^r is defined in (47) and $\operatorname{coad}^r(R_{[2]})(\varphi)\psi_i$ means product in the sense of (46).

Proof. The formula for the right *H*-action directly follows from (30) and the *H*-equivariant structures we put on H° and the X_i 's. The formulas for the action of $\mathcal{L}_{g,n}(H)$ directly follow from (34) and Lemma 4.9.

Remark 4.11. When $\varphi \in \mathcal{L}_{0,1}^{inv}(H)$, the formulas in Proposition 4.10 simplify dramatically:

$$\begin{aligned} \mathbf{i}_{b_i}(\varphi) \cdot \left(\bigotimes_k \psi_k \otimes \bigotimes_l x_l\right) &= \psi_1 \otimes \ldots \otimes (\psi_i \star \varphi) \otimes \ldots \otimes \psi_g \otimes \bigotimes_l x_l, \\ \mathbf{i}_{a_i}(\varphi) \cdot \left(\bigotimes_k \psi_k \otimes \bigotimes_l x_l\right) &= \psi_1 \otimes \ldots \otimes \left(\Phi_{0,1}(\varphi) \triangleright \psi_i\right) \otimes \ldots \otimes \psi_g \otimes \bigotimes_l x_l, \\ \mathbf{i}_{m_{g+j}}(\varphi) \cdot \left(\bigotimes_k \psi_k \otimes \bigotimes_l x_l\right) &= \bigotimes_k \psi_k \otimes x_1 \otimes \ldots \otimes \left(\Phi_{0,1}(\varphi) \cdot x_i\right) \otimes \ldots \otimes x_n, \end{aligned}$$

because of (25) and Lemma 4.3, where \star is the usual product (40) in H° .

For any left *H*-modules X, X_1, \ldots, X_n let $\Sigma_{g,X_1,\ldots,X_n}^X$ be the oriented surface of genus with n punctures labelled by X_1, \ldots, X_n and one boundary circle labelled by X^{6} .

To such a labelled surface, we associate the vector space

$$V(\Sigma_{g,X_1,\dots,X_n}^X) = \operatorname{Hom}_H(X, (H^\circ)^{\otimes g} \otimes X_1 \otimes \dots \otimes X_n)$$
(62)

where $(H^{\circ})^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n$ has the right *H*-module structure (61) and *X* is viewed as a right *H*-module thanks to the antipode: $x \cdot h = S(h) \cdot x$ for all $x \in X$ and $h \in H$ (or equivalently one can directly take $X \in \text{Mod-}H$). In particular when $X = \Bbbk$, the trivial *H*-module, $V(\Sigma_{g,X_1,\ldots,X_n}^{\Bbbk})$ is identified with the subspace of *H*-invariant elements in $(H^{\circ})^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n$ under (61).

Recall the subalgebra $\mathcal{L}_{g,n}^{\text{inv}}(H)$ of *H*-invariant elements under the right action coad^{*r*} (Def. 4.7). By Lemma 2.3 applied to the *H*-equivariant module $(H^{\circ})^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n$ there is a representation of $\mathcal{L}_{g,n}^{\text{inv}}(H)$ on the space $V(\Sigma_{g,X_1,\ldots,X_n}^X)$ given by acting on the target:

$$\forall a \in \mathcal{L}_{g,n}^{\text{inv}}(H), \ \forall f \in V(\Sigma_{g,X_1,\dots,X_n}^X), \ \forall x \in X, \quad (a \cdot f)(x) = a \cdot f(x).$$
(63)

In order to obtain the "derived" generalization of this representation, consider a projective resolution $0 \leftarrow X \leftarrow P_0 \leftarrow P_1 \leftarrow \ldots$ of X in Mod-H. Using the functor $V(\Sigma_{g,X_1,\ldots,X_n}^-) =$ Hom_H $(-,(H^\circ)^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n)$ we build the complex

$$0 \longrightarrow V(\Sigma_{g,X_1,\dots,X_n}^{P_0}) \longrightarrow V(\Sigma_{g,X_1,\dots,X_n}^{P_1}) \longrightarrow \dots$$

We denote the m-th cohomology space of this complex by

$$V^{m}(\Sigma_{g,X_{1},\dots,X_{n}}^{X}) = \operatorname{Ext}_{H}^{m}(X,(H^{\circ})^{\otimes g} \otimes X_{1} \otimes \dots \otimes X_{n}).$$
(64)

Note that $V^0(\Sigma^X_{g,X_1,\dots,X_n}) \cong V(\Sigma^X_{g,X_1,\dots,X_n})$ due to the general fact that $\operatorname{Ext}^0(M,N) \cong \operatorname{Hom}(M,N)$.

⁶The only reason we switch to n punctures and one boundary circle instead of n + 1 boundary circles (as in §1.1) is because it will be more convenient to have only one boundary circle when using stated skein algebras in §5.1.

Theorem 4.12. For all *H*-modules X, X_1, \ldots, X_n and all $m \ge 0$ there is a representation of $\mathcal{L}_{g,n}^{inv}(H)$ on $V^m(\Sigma_{g,X_1,\ldots,X_n}^X)$ given as follows:

$$\forall a \in \mathcal{L}_{g,n}^{\text{inv}}(H), \quad \forall [f] \in V^m(\Sigma_{g,X_1,\dots,X_n}^X), \qquad a \cdot [f] = [a \cdot f]$$

where [f] is the cohomology class of a cocycle $f \in V(\Sigma_{g,X_1,\ldots,X_n}^{P_m})$ and $a \cdot f$ is defined in (63).

Proof. Particular case of Theorem 2.4.

5 Topological applications

Assumption: $H = (H, \cdot, 1, \Delta, \varepsilon, S, R, v)$ is a ribbon Hopf algebra with invertible antipode over a field k. In the course of the text more assumptions on H will be required.

We denote the ribbon element of H by v. Recall from e.g. [CP94, §4.2.C], [Kas95, §XIV.6] that it is a central and invertible element such that

$$\Delta(v) = (R'R)^{-1}(v \otimes v), \quad \varepsilon(v) = 1, \quad S(v) = v, \quad v^2 = uS(u)$$

where u is the Drinfeld element (37). The *pivotal element* of H is

$$g = uv^{-1}. (65)$$

It satisfies

$$\Delta(g) = g \otimes g, \quad \text{and} \quad \forall h \in H, \ S^2(h) = ghg^{-1}.$$
(66)

The category *H*-mod of *finite-dimensional left H*-modules is ribbon, with braiding given by $c_{X,Y}(x \otimes y) = R_{[2]} \cdot y \otimes R_{[1]} \cdot x$ and twist $\theta_X : X \to X$ given by $\theta_X(x) = v^{-1} \cdot x$. We recall that for any $X \in H$ -mod the *H*-module structure on X^* is given by $(h \cdot \sigma)(x) = \sigma(S(h) \cdot x)$ for all $h \in H, \sigma \in X^*, x \in X$.

5.1 Knot maps and boundary closure

Let $\Sigma_{g,n}$ be the oriented surface of genus g with n punctures and $\Sigma_{g,n}^{\circ}$ be the surface obtained by cutting an open disk from $\Sigma_{g,n}$. Recall that the notation $\Sigma_{g,X_1,\ldots,X_n}^X$ means that the boundary circle is colored by X while the punctures are colored by X_1,\ldots,X_n . It is natural to expect that the choice $X = \Bbbk$ (the trivial H-module) has something to do with the surface where this boundary component has been closed. In this section we give a precise statement supporting this idea. Namely, to each isotopy class of knot $K \subset \Sigma_{g,n}^{\circ} \times [0,1]$, we associate an element $\mathbb{W}^{\varphi}(K) \in \mathcal{L}_{g,n}^{\mathrm{inv}}(H)$ and we show that the representation of $\mathbb{W}^{\varphi}(K)$ on the space $V^m(\Sigma_{g,X_1,\ldots,X_n}^{\Bbbk})$ defined in Theorem 4.12 only depends of the isotopy class of K viewed in $\Sigma_{g,n} \times [0,1]$. The superscript φ is any element in $\mathcal{L}_{0,1}^{\mathrm{inv}}(H)$

The elements $\mathbb{W}^{\varphi}(K)$ were first defined in [BR96] under the name Wilson loops. They were re-defined in [BFK98a] (in a different formalism) through an axiomatic system of rules. Finally in [Fai20b] they were recovered as a particular case of a holonomy operation for tangles in $\Sigma_{q,n}^{\circ} \times [0, 1]$.

It was proved in [BFR23] that, under some conditions on H, there is an isomorphism between $\mathcal{L}_{g,n}(H)$ and a so-called stated skein algebra $\mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$. This allows for an even more natural construction of \mathbb{W}^{φ} which is presented here. The point is to first define knot maps $\mathfrak{i}_{K} : \mathcal{L}_{0,1}^{\mathrm{inv}}(H) \to \mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet}) \cong \mathcal{L}_{g,n}(H)$ and then to let $\mathbb{W}^{\varphi}(K) = \mathfrak{i}_{K}(\varphi)$. This approach also greatly simplifies the construction of mapping class group representations in §5.2.

5.1.1 Stated skein algebras

Let $\Sigma_{g,n}^{\circ,\bullet}$ be the surface obtained by removing a point on the circle $\partial(\Sigma_{g,n}^{\circ})$. The extra puncture • is relevant only when considering tangles whose endpoints lie in $\partial(\Sigma_{g,n}^{\circ,\bullet}) \times [0,1]$, as we will shortly recall. As a result we indicate the puncture • only when it is relevant to do so; for instance there is no reason to write $\pi_1(\Sigma_{g,n}^{\circ,\bullet})$ instead of $\pi_1(\Sigma_{g,n}^{\circ})$.

Here is a picture of the surface $\Sigma_{g,n}^{\circ,\bullet}$. We fix some basepoint on $\partial(\Sigma_{g,n}^{\circ,\bullet})$, not to be confused with the puncture \bullet , for the fundamental group $\pi_1(\Sigma_{g,n}^{\circ})$ and fix generators $b_1, a_1, \ldots, b_g, a_g, m_{g+1}, \ldots, m_{g+n}$ of this (free) group:



We also put a preferred direction on the boundary arc $\partial(\Sigma_{g,n}^{\circ,\bullet})$ and introduce a simple loop $\partial_{g,n} \in \pi_1(\Sigma_{g,n}^{\circ})$. Note that the orientation of $\partial_{g,n}$ is opposed to the preferred direction on $\partial(\Sigma_{g,n}^{\circ,\bullet})$. By retracting $\partial(\Sigma_{g,n}^{\circ,\bullet})$ to a tubular neighborhood of these loops we get another picture for $\Sigma_{g,n}^{\circ,\bullet}$, which will be more convenient in the sequel:



With this view it is immediate that

$$\partial_{g,n} = b_1 a_1^{-1} b_1^{-1} a_1 b_2 a_2^{-1} b_2^{-1} a_2 \dots b_g a_g^{-1} b_g^{-1} a_g m_{g+1} m_{g+2} \dots m_{g+n}.$$
(68)

Let $S_H^{st}(\Sigma_{g,n}^{\circ,\bullet})$ the stated skein algebra of the bordered punctured surface $\Sigma_{g,n}^{\circ,\bullet}$. The definition of this algebra is explained in great detail in [BFR23, §6.1].⁷ Let us just recall that one considers the k-vector space spanned by isotopy classes of *H*-colored oriented stated ribbon graphs $G \subset \Sigma_{g,n}^{\circ,\bullet} \times [0,1]$ such that $\partial G \subset (\partial \Sigma_{g,n}^{\circ,\bullet}) \times [0,1]$ and the points of ∂G have strictly increasing heights when one follows the preferred direction on $\partial(\Sigma_{g,n}^{\circ,\bullet})$ fixed above. A state is a label of each end-point of the graph *G* by a vector in the *H*-module coloring the strand to which that point belong (or in its dual depending on the orientation). Then $S_H^{st}(\Sigma_{g,n}^{\circ,\bullet})$ is the quotient of this vector space by *skein relations* obtained by applying the Reshetikhin–Turaev functor [RT90] near the boundary. Examples of important stated skein relations are given in Figure 1. Hopefully the reader can grasp the definition from these examples; otherwise look at [BFR23, §6.1].

The product $G_1 * G_2$ of two stated ribbon graphs G_1, G_2 is obtained by putting G_1 (resp. G_2) in $\sum_{g,n}^{\circ,\bullet} \times [0, \frac{1}{2}[$ (resp. in $\sum_{g,n}^{\circ,\bullet} \times]\frac{1}{2}, 1]$) by isotopy without changing the states and then taking their union. This turns $\mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$ into an algebra.

There is moreover a right *H*-module-algebra structure on $\mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$ obtained by acting on states: let *G* be a stated ribbon graph and $p_1, \ldots, p_l \in \partial(\Sigma_{g,n}^{\circ,\bullet}) \times [0,1]$ be its boundary points,

⁷F. Costantino, J. Korinman and T. Lê are working on the definition and properties of stated skein algebras for surfaces with arbitrary many boundary components and arbitrary many punctures on each of these components.



Figure 1: Examples of stated skein relations. X and Y are finite-dimensional left *H*-modules, $x \in X, \varphi \in X^*, y \in Y, \psi \in Y^*, f \in \text{Hom}_H(X, Y)$. (x_i) is a basis of X, (x^i) is its dual basis. In (i) and (ii), f^* means the transpose of f. In (vii) and (viii) we use implicit summation on the index *i*.

ordered by increasing height. Let s_i be the state at p_i and $X_i \in H$ -mod be the color of the strand in G which contains p_i . For all $h \in H$, we define $G \bullet h$ to be G with the new states $s_1 \bullet h_{(1)} \otimes \ldots \otimes s_l \bullet h_{(l)}$, where

$$s_{i} \bullet h_{(i)} = \begin{cases} S(h_{(i)}) \cdot s_{i} & \text{if } s_{i} \in X_{i} \text{ (i.e. the strand goes upwards at } p_{i}) \\ s_{i}(h_{(i)} \cdot ?) & \text{if } s_{i} \in X_{i}^{*} \text{ (i.e. the strand goes downwards at } p_{i}) \end{cases}$$
(69)

with $s_i(h_{(i)} ?) \in X_i^*$ the linear form defined by $x_i \mapsto s_i(h_{(i)} \cdot x_i)$. Here is an example with three boundary points labelled by states $s_1 \in X_1^*$, $s_2 \in X_2$, $s_3 \in X_3$:

The arrow on the boundary in (70) indicates that the heights of the boundary points labelled by s_1, s_2, s_3 form a strictly increasing sequence in [0, 1] along this direction.

We now introduce the ingredients which will later enter in the definition of *knot maps*.

Definition 5.1. 1. A knot $K \subset \Sigma_{g,n}^{\circ} \times [0,1]$ is an oriented and framed embedding of the circle S^1 , up to isotopy.

2. A based knot $\mathbf{K} \subset \Sigma_{g,n}^{\circ,\bullet} \times [0,1]$ is an oriented and framed embedding of the circle S^1 such that $\mathbf{K} \cap \left(\partial(\Sigma_{g,n}^{\circ,\bullet}) \times [0,1]\right)$ is reduced to a point (the basepoint of the knot). We consider these knots up to isotopies such that the basepoint remains in $\partial(\Sigma_{g,n}^{\circ,\bullet}) \times [0,1]$.

3. We say that a based knot is positively oriented if its orientation near its basepoint looks like
when we draw it on the surface (67); otherwise we say it is negatively oriented.
4. From a based knot one can obtain a knot, by forgetting the conditions regarding the basepoint.

Note that a (based) knot is *not* allowed to contain coupons. Here is an example of a positively oriented and a negatively oriented based knot in $\Sigma_{1,0}^{\circ,\bullet} \times [0,1]$:



When consider as usual knots, \mathbf{K}_{ba} and \mathbf{L}_{ba} become isotopic up to orientation. Simple (*i.e.* nonself intersecting) loops in $\pi_1(\Sigma_{g,n}^\circ)$ have a basepoint in $\partial \Sigma_{g,n}^\circ$ by definition and can be embedded in $\Sigma_{g,n}^{\circ,\bullet} \times \{t\}$ for an arbitrary $t \in [0, 1]$; they are thus particular cases of based knots. This is illustrated in the example above: the based knot \mathbf{L}_{ba} is the simple loop $(ba)^{-1}$.

For a based knot $\mathbf{K} \subset \Sigma_{g,n}^{\circ,\bullet} \times [0,1], X \in H$ -mod, $x \in X$ and $\sigma \in X^*$ let ${}_X\mathbf{K}_x^{\sigma} \in \mathcal{S}_H^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$ be defined by:

• If K is positively oriented then



which means that the stated skein follows exactly \mathbf{K} in $\Sigma_{g,n}^{\circ,\bullet} \times [0,1]$ according to its orientation, except in this small neighbourhood of the basepoint where \mathbf{K} is represented in dashed line.

• If K is negatively oriented then



where the picture has the same meaning as in the previous item.

We see from (i) and (ii) in Fig. 1 that these elements obey dinaturality:

$$\forall f \in \operatorname{Hom}_{H}(X, Y), \ \forall x \in X, \ \forall \tau \in Y^{*}, \quad {}_{Y}\mathbf{K}_{f(x)}^{\tau} = {}_{X}\mathbf{K}_{x}^{f^{*}(\tau)}$$
(74)

where f^* is the transpose of f, because the coupon colored by f as in Fig. 1 can slide along **K**. If **K** is positively oriented then we moreover have

where in the first picture \mathbf{K}^{-1} is \mathbf{K} with the opposite orientation and $\langle -, x \rangle : X^* \to \mathbb{k}$ is defined by $\sigma \mapsto \sigma(x)$, while in the second picture the strands colored by X and Y are parallel and follow K closely. These facts are proved as in [BFR23, Proof of Lem. 6.4]. Note that if \mathbf{K} is positively oriented then \mathbf{K}^{-1} is negatively oriented and (xi) in Fig. 1, the 1st equality in (75) and (65) yield

$${}_{X}(\mathbf{K}^{-1})_{x}^{\sigma} = {}_{X^{*}}\mathbf{K}_{\sigma(R_{[2]}u^{-1}\cdot?)}^{\langle -,S(R_{[1]})\cdot x \rangle}.$$
(76)

Here are further basic properties which will be useful in the next subsection:

Lemma 5.2. If H is finite-dimensional then for any based knot K we have

$$_{H}\mathbf{K}_{h}^{\varphi} = {}_{H}\mathbf{K}_{1}^{h \triangleright \varphi}, \qquad {}_{X}\mathbf{K}_{x}^{\sigma} = {}_{H}\mathbf{K}_{1}^{\sigma(? \cdot x)}$$

where H is considered as an H-module by left multiplication and $\sigma(?\cdot x)$ is the matrix coefficient $_X\phi_x^{\sigma}: h \mapsto \sigma(h \cdot x).$

Proof. For the first equality consider $r_h : H \to H$, $a \mapsto ah$. It is a *H*-linear endomorphism of the regular representation and thus by (74) we get ${}_{H}\mathbf{K}_{h}^{\varphi} = {}_{H}\mathbf{K}_{r_{h}(1)}^{\varphi} = {}_{H}\mathbf{K}_{1}^{r_{h}^{*}(\varphi)} = {}_{H}\mathbf{K}_{1}^{h \mapsto \varphi}$. For the second equality, since the left *H*-module *X* is finite-dimensional it is in particular finitely generated and thus there exists a *H*-linear surjection $p : H^{l} \to X$ for some $l \ge 1$. Let $\iota_{s} : H \to H^{l}$ be the canonical injections and let $(h_{1}, \ldots, h_{l}) = \sum_{s} \iota_{s}(h_{s}) \in H^{l}$ be a preimage of *x* by *p*. Using (74) and the first equality in the current lemma we find

$${}_{X}\mathbf{K}_{x}^{\sigma} = \sum_{s=1}^{l} {}_{X}\mathbf{K}_{p \circ \iota_{s}(h_{s})}^{\sigma} = \sum_{s=1}^{l} {}_{H}\mathbf{K}_{h_{s}}^{(p \circ \iota_{s})^{*}(\sigma)} = \sum_{s=1}^{l} {}_{H}\mathbf{K}_{1}^{h_{s} \triangleright ((p \circ \iota_{s})^{*}(\sigma))}.$$

But note that for all $h \in H$ we have

$$\sum_{s=1}^{l} \langle h_s \triangleright (p \circ \iota_s)^*(\sigma), h \rangle = \sum_{s=1}^{l} \langle \sigma, (p \circ \iota_s)(hh_s) \rangle = \sum_{s=1}^{l} \langle \sigma, h \cdot (p \circ \iota_s)(h_s) \rangle = \sigma(h \cdot x).$$

5.1.2 Knot maps for stated skein algebras

For $X \in H$ -mod let $C(X) \subset H^{\circ}$ be the subspace of matrix coefficients of X, *i.e.* $_X \phi_x^{\sigma} : h \mapsto \sigma(h \cdot x)$, for all $\sigma \in X^*$ and $x \in X$. By definition there is a surjection

$$X^* \otimes X \twoheadrightarrow C(X), \quad \sigma \otimes x \mapsto {}_X \phi_x^{\sigma}.$$
 (77)

If we define the right *H*-action $(\sigma \otimes x) \cdot h = \sigma(h_{(1)} \cdot ?) \otimes S(h_{(2)}) \cdot x$ then this surjection becomes a *H*-linear map $(X^* \otimes X, \cdot) \to (C(X), \operatorname{coad}^r)$. We recall that $\mathcal{L}_{0,1}(H) = H^\circ = \sum_{X \in H \operatorname{-mod}} C(X)$ as vector spaces. For any based knot $\mathbf{K} \subset \Sigma_{g,n}^{\circ, \bullet} \times [0, 1]$ (Def. 5.1), we would like to use the stated skein ${}_X\mathbf{K}_x^\sigma$ from (72)–(73) to define a *based knot map* given by

$$\mathbf{j}_{\mathbf{K}}: \mathcal{L}_{0,1}(H) \to \mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet}), \quad {}_{X}\phi_{x}^{\sigma} \mapsto {}_{X}\mathbf{K}_{x}^{\sigma}.$$
(78)

A priori this is not well-defined: for example, elements in the kernel of (77) might give non-zero stated skeins. Nevertheless, the map $j_{\mathbf{K}}$ makes sense for two relevant choices for H:

- If H is $U_q(\mathfrak{g})$, the quantum group over $\mathbb{C}(q)$, then it is customary to use $\mathcal{O}_q(G)$ instead of the whole H° . By definition $\mathcal{O}_q(G)$ is the subspace of matrix coefficients of the type 1 highest weight $U_q(\mathfrak{g})$ -modules V_μ for all $\mu \in P_+$, where P_+ is the set of dominant weights of \mathfrak{g} . We have $\mathcal{O}_q(G) = \bigoplus_{\mu \in P_+} C(V_\mu)$ and $C(V_\mu) \cong V_\mu^* \otimes V_\mu$ [VY20, §3.10].
- If *H* is finite-dimensional then the second equality in Lemma 5.2 shows that ${}_{X}\mathbf{K}_{x}^{\sigma}$ actually only depends on the matrix coefficient ${}_{X}\phi_{x}^{\sigma}$ and not on the tensor $\sigma \otimes x$.

In the sequel of this subsection we assume that H satisfies one of these two items.

Here are the first basic properties of the based knot maps $j_{\mathbf{K}}$:

Lemma 5.3. 1. For all based knot **K** the linear map $\mathbf{j}_{\mathbf{K}}$ is *H*-linear with respect to the right actions coad^r on $\mathcal{L}_{0,1}(H)$ from (47) and \cdot on $\mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$ from (69). 2. Let **K** be a positively oriented based knot and \mathbf{K}^{-1} be **K** with the opposite orientation (thus

2. Let **K** be a positively oriented based knot and \mathbf{K}^{-1} be **K** with the opposite orientation (thus negatively oriented). Then

$$\forall \varphi \in \mathcal{L}_{0,1}(H), \quad \mathfrak{j}_{\mathbf{K}^{-1}}(\varphi) = \mathfrak{j}_{\mathbf{K}}(S_{\mathcal{L}_{0,1}}(\varphi))$$

where $S_{\mathcal{L}_{0,1}}$ is defined in (48).

Proof. 1. If $H = U_q(\mathfrak{g})$ then this follows from the *H*-equivariance of (77). If *H* is finitedimensional then by item 1 in Lemma 5.2 we have for all $\varphi \in \mathcal{L}_{0,1}(H)$

$$\mathfrak{j}_{\mathbf{K}}(\varphi) \bullet h = \mathfrak{j}_{\mathbf{K}}({}_{H}\phi_{1}^{\varphi}) \bullet h = {}_{H}\mathbf{K}_{1}^{\varphi} \bullet h \stackrel{(69)}{=} {}_{H}\mathbf{K}_{S(h_{(2)})}^{\varphi(h_{(1)}?)} = {}_{H}\mathbf{K}_{1}^{\varphi(h_{(1)}?S(h_{(2)}))} \stackrel{(47)}{=} \mathfrak{j}_{\mathbf{K}}(\operatorname{coad}^{r}(h)(\varphi)).$$

2. It is a consequence of (76):

$$\mathfrak{j}_{\mathbf{K}^{-1}}(_{X}\phi_{x}^{\sigma}) = {}_{X^{*}}\mathbf{K}_{\sigma(R_{[2]}u^{-1}\cdot?)}^{\langle -,S(R_{[1]})\cdot x\rangle} = \mathfrak{j}_{\mathbf{K}}\Big({}_{X^{*}}\phi_{\sigma(R_{[2]}u^{-1}\cdot?)}^{\langle -,S(R_{[1]})\cdot x\rangle}\Big)\,.$$

Now observe that for all $h \in H$,

$$\left\langle {}_{X^*}\phi^{\langle -,S(R_{[1]})\cdot x\rangle}_{\sigma(R_{[2]}u^{-1}\cdot?)},h\right\rangle = \left\langle h\cdot\sigma(R_{[2]}u^{-1}\cdot?),S(R_{[1]})\cdot x\right\rangle = \left\langle \sigma(R_{[2]}u^{-1}\cdot?),S(h)S(R_{[1]})\cdot x\right\rangle$$
$$= \sigma\left(R_{[2]}u^{-1}S(h)S(R_{[1]})\cdot x\right) = \left\langle S_{H^\circ}\left(S(R_{[1]})\triangleright_X\phi^{\sigma}_x \triangleleft R_{[2]}u^{-1}\right),h\right\rangle = \left\langle S_{\mathcal{L}_{0,1}}(x\phi^{\sigma}_x),h\right\rangle. \qquad \Box$$

From a based knot $\mathbf{K} \subset \Sigma_{g,n}^{\circ,\bullet} \times [0,1]$ one obtains a knot K by forgetting that the basepoint of \mathbf{K} is constrained to be attached to $\partial(\Sigma_{g,n}^{\circ,\bullet}) \times [0,1]$ (item 4 in Def. 5.1). Recall that $\mathcal{L}_{0,1}^{\mathrm{inv}}(H)$ is the subspace of invariant elements in $\mathcal{L}_{0,1}(H)$ under the right H-action coad^r; see (50).

Proposition 5.4. Let \mathbf{K}_1 and \mathbf{K}_2 be based knots in $\Sigma_{g,n}^{\circ,\bullet} \times [0,1]$ which reduce to the same knot when we forget that they are based. If $\varphi \in \mathcal{L}_{0,1}^{\mathrm{inv}}(H)$ then $\mathbf{j}_{\mathbf{K}_1}(\varphi) = \mathbf{j}_{\mathbf{K}_2}(\varphi)$. As a result there is a well-defined linear map $\mathbf{j}_K : \mathcal{L}_{0,1}^{\mathrm{inv}}(H) \to \mathcal{S}_H^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$ for all knot K.

Proof. The first claim is proven in Appendix B.1. For the second claim, choose any based knot **K** such that **K** reduces to *K* when we forget its basedpoint. Then it suffices to put $\mathbf{j}_K(\varphi) = \mathbf{j}_{\mathbf{K}}(\varphi)$ and the first claim ensures that this value is independent of the choice of **K**. \Box

We recall that a loop $\gamma \in \pi_1(\Sigma_{g,n}^{\circ})$ is called *simple* if it can be represented by a curve without self-intersections. Being a particular case of a based knot, we have the map \mathfrak{j}_{γ} . Recall the algebra structure on $\mathcal{L}_{0,1}(H)$ from Definition 4.1. Also recall the product * on $\mathcal{S}_H^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$, which is given by stacking.

Proposition 5.5. Let $\gamma \in \pi_1(\Sigma_{q,n}^\circ)$ be a simple loop.

1. If γ is positively oriented then \mathfrak{j}_{γ} is a morphism of right H-module-algebras $\mathcal{L}_{0,1}(H) \rightarrow \mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{q,n}^{\circ,\bullet})$.

2. If γ is negatively oriented then \mathfrak{j}_{γ} is a morphism of right H-modules $\mathcal{L}_{0,1}(H) \to \mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$ which satisfies

 $\forall \varphi, \psi \in \mathcal{L}_{0,1}(H), \quad \mathfrak{j}_{\gamma}(\varphi\psi) = \mathfrak{j}_{\gamma}(\operatorname{coad}^{r}(R_{[1]})(\psi)) * \mathfrak{j}_{\gamma}(\operatorname{coad}^{r}(R_{[2]})(\varphi)).$

Proof. The *H*-linearity was already proven for any based knot in item 1 of Lemma 5.3. 1. We check the algebra morphism property as follows:

$$\begin{split} \mathfrak{j}_{\gamma}(x\phi_{x}^{\sigma})*\mathfrak{j}_{\gamma}(y\phi_{y}^{\tau}) &= \overbrace{X}^{\sigma}\overbrace{\gamma} \underbrace{\gamma}_{Y} \underbrace{\gamma}_{Y} \underbrace{\gamma}_{\gamma} \underbrace{\gamma}_{Y} \underbrace{\gamma}_{\gamma} \underbrace{\gamma}_{Y} \underbrace{\gamma}_{Y} \underbrace{\gamma}_{\gamma} \underbrace{\gamma}_{Y} \underbrace{\gamma}_{Y} \underbrace{\gamma}_{\gamma} \underbrace{\gamma}_{Y} \underbrace{\gamma}_{Y} \underbrace{\gamma}_{\gamma} \underbrace{\gamma}_{Y} \underbrace{\gamma}_{Y} \underbrace{\gamma}_{Y} \underbrace{\gamma}_{\gamma} \underbrace{\gamma}_{Y} \underbrace{\gamma}$$

In the pictures γ is drawn in dashed line and we only represent a neighborhood of its basepoint. Outside of this, the strands colored by X and Y are parallel to γ and all three stay in a same layer $\sum_{g,n}^{\circ,\bullet} \times \{t\}$. For the second equality we used (x) and (xii) in Fig. 1, for the third we used the 2nd formula in (75), for the fourth we used the definition of \star in (40), for the fifth we used (44) and the last is (46).

2. Follows immediately from item 2 in Lemma 5.3, (49) and the previous item. \Box

To conclude this subsection, we note that there is an obvious action of the mapping class group on the stated skein algebra. Indeed, let $MCG(\Sigma_{g,n}^{\circ})$ be the mapping class group of $\Sigma_{g,n}^{\circ}$, *i.e.* the group of isotopy classes of diffeomorphisms $\Sigma_{g,n}^{\circ} \to \Sigma_{g,n}^{\circ}$ which fix pointwise both the punctures and the boundary. A mapping class $f \in MCG(\Sigma_{g,n}^{\circ})$ acts on an isotopy class of stated ribbon graph $G \subset \Sigma_{g,n}^{\circ,\bullet} \times [0,1]$ simply by $f(G) = (f \times id_{[0,1]})(G)$ and without changing the states. Let us linearly extend this formula to formal linear combinations of stated ribbon graphs. Then the skein relations are compatible with the resulting action, since they occur only a small neighborhood of $\partial(\Sigma_{g,n}^{\circ,\bullet}) \times [0,1]$. As a result for all $f \in MCG(\Sigma_{g,n}^{\circ})$ we get a linear map

$$\widetilde{f}: \mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet}) \to \mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet}), \quad G \mapsto (f \times \mathrm{id}_{[0,1]})(G)$$
(79)

whose inverse is $\widetilde{f^{-1}}$. It is clear that \widetilde{f} is an algebra automorphism, because the product in $\mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$ is given by stacking in the height component [0,1] which is not modified by \widetilde{f} . From these remarks we get a group morphism

$$\operatorname{MCG}(\Sigma_{g,n}^{\circ}) \to \operatorname{Aut}_{\operatorname{alg}}(\mathcal{S}_{H}^{\operatorname{st}}(\Sigma_{g,n}^{\circ,\bullet})), \quad f \mapsto \widetilde{f}.$$

We note as a side remark that \tilde{f} is also compatible with the right *H*-module structure on $\mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{q,n}^{\circ,\bullet})$ from (69), and is thus an automorphism of *H*-module-algebra.

An element $f \in MCG(\Sigma_{g,n}^{\circ})$ transforms a based knot **K** into the based knot $f(\mathbf{K}) = (f \times id_{[0,1]})(\mathbf{K})$ because f fixes the boundary. Moreover f preserves the orientation of **K** and hence it is immediate from the definitions that

$$\forall f \in \mathrm{MCG}(\Sigma_{g,n}^{\circ}), \quad \widetilde{f} \circ \mathfrak{j}_{\mathbf{K}} = \mathfrak{j}_{f(\mathbf{K})}.$$
(80)

5.1.3 Knots maps for $\mathcal{L}_{g,n}(H)$ and boundary closure

In [BFR23, §6.2] a morphism of algebras

$$\operatorname{hol}^{\operatorname{st}} : \mathcal{S}_{H}^{\operatorname{st}}(\Sigma_{g,n}^{\circ,\bullet}) \to \mathcal{L}_{g,n}(H)$$

was defined by building upon results of [Fai20b]; the notation stands for "stated holonomy". Under the following assumption it is actually an *isomorphism*:

Assumption: As in §5.1.2 we assume that $H = U_q(\mathfrak{g})$ with q generic or that H is finite dimensional. More generally one can take any ribbon Hopf algebra H such that the knot maps (78) are well-defined.

We give a property which characterizes holst and is enough for our purposes. Recall that a simple loop in $\pi_1(\Sigma_{g,n}^{\circ})$ can be seen as a based knot in $\Sigma_{g,n}^{\circ,\bullet} \times [0,1]$ (Def. 5.1). Consider the resulting knot map $\mathfrak{j}_s: \mathcal{L}_{0,1}(H) \to \mathcal{S}_H^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$ for any generator $s \in \{b_1, a_1, \ldots, b_g, a_g, m_{g+1}, \ldots, m_{g+n}\}$ of $\pi_1(\Sigma_{g,n}^{\circ})$ fixed in (67). Also recall the canonical embeddings $\mathfrak{i}_s: \mathcal{L}_{0,1}(H) \to \mathcal{L}_{g,n}(H)$ from (55). Then for each such s there is a commutative diagram

In particular we deduce that holst is an isomorphism of *H*-module-algebras when $\mathcal{S}_{H}^{st}(\Sigma_{g,n}^{\circ,\bullet})$ is endowed with the right *H*-action (69), because \mathfrak{i}_{s} and \mathfrak{j}_{s} are *H*-linear.

For any based knot $\mathbf{K} \subset \Sigma_{g,n}^{\circ,\bullet} \times [0,1]$ define a linear map

$$\mathfrak{i}_{\mathbf{K}} = \operatorname{hol}^{\operatorname{st}} \circ \mathfrak{j}_{\mathbf{K}} : \mathcal{L}_{0,1}(H) \to \mathcal{L}_{g,n}(H)$$
(82)

where $\mathfrak{j}_{\mathbf{K}}$ is the map from (78). Obviously:

Corollary 5.6. The statements of Lemma 5.3 and Propositions 5.4 and 5.5 remain true if we replace everywhere j by i.

In particular, by Proposition 5.4, for all (unbased) knot $K \subset \Sigma_{q,n}^{\circ} \times [0,1]$ there is a knot map

$$\mathfrak{i}_K : \mathcal{L}_{0,1}^{\mathrm{inv}}(H) \to \mathcal{L}_{g,n}^{\mathrm{inv}}(H), \quad \varphi \mapsto \mathfrak{i}_K(\varphi)$$
(83)

where **K** is any based knot which is isotopic to K when we consider it as an unbased knot. Example 5.7. In $\Sigma_{0,2}^{\circ} \times [0,1]$ let



(the puncture on the boundary is relevant only when working with based knots). Then \mathbf{K} reduces to K when considered as an unbased knot and we have



The first equality is by definition of $j_{\mathbf{K}}$ from (72) and (78), the second equality uses (vii), (xi) and (xii) in Fig. 1, the third equality uses the first identity (75)and the last equality uses the

second identity in (75). To obtain a simpler expression, let us evaluate this on an element $h \otimes k \in H \otimes H$:

$$\sigma(h_{(1)}S(R_{[1]}^1)g \cdot x_i) \Big\langle h_{(2)} \cdot x^i (R_{[2]}^2 g^{-1} \cdot ?), S(R_{[1]}^2) R_{[2]}^1 \cdot x_j \Big\rangle x^j (k \cdot x)$$

= $\sigma(h_{(1)}S(R_{[1]}^1)g R_{[2]}^2 g^{-1}S(h_{(2)})S(R_{[1]}^2) R_{[2]}^1 k \cdot x) = \sigma(h_{(1)}S(R_{[2]}^2 R_{[1]}^1)S(h_{(2)}) R_{[1]}^2 R_{[2]}^1 k \cdot x)$

where the second equality uses (66) and (23). It follows that for all $\varphi \in \mathcal{L}_{0,1}(H)$, the element $\mathfrak{i}_{\mathbf{K}}(\varphi) \in \mathcal{L}_{0,2}(H) = H^{\circ} \otimes H^{\circ} \subset (H \otimes H)^{*}$ is characterized by

$$\forall h, k \in H, \quad \left< \mathbf{i}_{\mathbf{K}}(\varphi), h \otimes k \right> = \varphi \left(h_{(1)} S(R_{[2]}^2 R_{[1]}^1) S(h_{(2)}) R_{[1]}^2 R_{[2]}^1 k \right).$$

For more complicated knots/surfaces it is faster to use the technique \dot{a} la Hennings to evaluate $i_{\mathbf{K}}(\varphi)$ [Fai20b, §4.3]. The following picture illustrates this technique:



Note that in [Fai20b, §4.3] there was an extra pivotal element g in hen_K because we used symmetric linear forms (*i.e.* $\varphi(xy) = \varphi(yx)$) instead of elements of $\mathcal{L}_{0,1}^{\text{inv}}(H)$, see (50) and (66).

Now let us change our point of view on the knot map (83): we fix φ while we allow K to vary. More precisely if Σ is an oriented surface denote by $\text{Knots}(\Sigma)$ the set of isotopy classes of (unbased) oriented framed knots in $\Sigma \times [0, 1]$. Then for any $\varphi \in \mathcal{L}_{0,1}^{\text{inv}}(H)$ we have a map

$$\mathbb{W}^{\varphi}$$
: Knots $(\Sigma_{g,n}^{\circ}) \to \mathcal{L}_{g,n}^{\text{inv}}(H), \quad K \mapsto \mathfrak{i}_{K}(\varphi).$ (84)

The inclusion $\Sigma_{g,n}^{\circ} \hookrightarrow \Sigma_{g,n}$ induces a surjection

$$\pi : \operatorname{Knots}(\Sigma_{q,n}^{\circ}) \twoheadrightarrow \operatorname{Knots}(\Sigma_{g,n}).$$

Recall from §4.2 the spaces $V(\Sigma_{g,X_1,\ldots,X_n}^X)$ for all $X, X_1, \ldots, X_n \in H$ -Mod, their derived version $V^m(\Sigma_{g,X_1,\ldots,X_n}^X)$ and the representations of $\mathcal{L}_{g,n}^{inv}(H)$ on these spaces. In the next result we take $X = \mathbb{k}$, the ground field on which H acts through the counit $\varepsilon : H \to \mathbb{k}$.

Theorem 5.8. Let H be a finite-dimensional ribbon Hopf algebra or a quantum group $U_q(\mathfrak{g})$ over $\mathbb{C}(q)$. Let $\varphi \in \mathcal{L}_{0,1}^{\mathrm{inv}}(H)$.

1. We have a commutative diagram

$$\operatorname{Knots}(\Sigma_{g,n}^{\circ}) \xrightarrow{\mathbb{W}^{\varphi}} \mathcal{L}_{g,n}^{\operatorname{inv}}(H)$$

$$\downarrow^{\operatorname{rep. from (63)}}_{\operatorname{Knots}(\Sigma_{g,n}) \xrightarrow{\exists !}} \operatorname{End}_{\Bbbk} \left(V(\Sigma_{g,X_{1},\ldots,X_{n}}^{\Bbbk}) \right)$$

2. Assume that H and X_1, \ldots, X_n are all finite-dimensional. We have a commutative diagram

In item 1, $V(\Sigma_{g,X_1,\ldots,X_n}^{\Bbbk})$ is just the subspace of invariant elements in $(H^{\circ})^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n$ for the right action of H in (61).

In order to prove the theorem, note that two knots K_1, K_2 satisfy $\pi(K_1) = \pi(K_2)$ if and only if they are related by a finite sequence of the following transformation taking place in a small neighborhood of the boundary annulus $\partial(\Sigma_{q,n}^{\circ}) \times [0,1]$:

$$\bigcirc \longleftrightarrow \bigcirc (85)$$

On the left picture the strand represent any portion of a knot which is brought near the boundary by isotopy. We need to find how the transformation (85) interacts with the knot maps i_K . Consider two based knots \mathbf{K} and \mathbf{K}_{∂} which look as follows in a neighborhood of $\partial(\Sigma_{g,n}^{\circ,\bullet}) \times [0,1]$:



Thanks to isotopy we can always assume that the unbased knots in (85) have this orientation in the neighborhood of the boundary.

Lemma 5.9. For all $\varphi \in \mathcal{L}_{0,1}(H)$,

$$\mathfrak{i}_{\mathbf{K}_{\partial}}(\varphi) = \mathfrak{i}_{\mathbf{K}}(\varphi_{(1)}) \,\mathfrak{i}_{\partial_{g,n}}(S_{\mathcal{L}_{0,1}}(\varphi_{(2)}))$$

with the coproduct (41), the antipode (48) and $\partial_{g,n}$ is the boundary loop (68).

Proof. It is a simple computation in $\mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{q,n}^{\circ,\bullet})$. Since \mathbf{K}_{∂} is negatively oriented,

The first equality is from the definition of $\mathbf{j}_{\mathbf{K}_{\partial}}$ in (73), the second equality uses isotopy and naturality of the twist, the third equality uses (vii) in Fig. 1 and the last equality is by the definition of $\mathbf{j}_{\mathbf{K}}$, $\mathbf{j}_{\partial g,n}$ in (72), (73) and the definition of the product in $\mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$. The result now follows from the formula (41) which gives the coproduct on matrix coefficients, item 2 in Lemma 5.3 and finally by applying holst, recall the definition of $\mathbf{i}_{\mathbf{K}}$ in (82).

The use of the resolution (95) and of the comparison theorem in the proof below were inspired by [LMSS23, p. 50], although they are used to prove a different result there.

Proof of Theorem 5.8. In both items, for $K_1, K_2 \in \text{Knots}(\Sigma_{g,n}^\circ)$ such that $\pi(K_1) = \pi(K_2)$, we want to show that the representations of $\mathbf{i}_{K_1}(\varphi)$ and $\mathbf{i}_{K_2}(\varphi)$ are equal. It suffices to show this for $\mathbf{i}_{\mathbf{K}}(\varphi)$ and $\mathbf{i}_{\mathbf{K}_{\partial}}(\varphi)$, where \mathbf{K} and \mathbf{K}_{∂} are based knots which look like (86) in a neighborhood of the boundary. Write

$$M = (H^{\circ})^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n$$

equipped with the *H*-equivariant structure from Prop. 4.10, so that $V(\Sigma_{g,X_1,\ldots,X_n}^{\Bbbk}) = \operatorname{Hom}_H(\Bbbk, M) \cong M^{\operatorname{inv}}$ and $V^m(\Sigma_{g,X_1,\ldots,X_n}^{\Bbbk}) = \operatorname{Ext}_H^m(\Bbbk, M)$. Let $\varphi \in \mathcal{L}_{0,1}^{\operatorname{inv}}(H)$ and consider the element

$$s_i \otimes t_i = \mathfrak{i}_{\mathbf{K}}(\varphi_{(1)}) \otimes \Phi_{0,1}(S_{\mathcal{L}_{0,1}}(\varphi_{(2)})) \in \mathcal{L}_{g,n}(H) \otimes H$$

with implicit summation on i, where $\Phi_{0,1}$ has been introduced in (56). According to Lemma 5.9 and Proposition C.1 we have

$$\mathbf{i}_{\mathbf{K}_{\partial}}(\varphi) \cdot w = \mathbf{i}_{\mathbf{K}}(\varphi_{(1)}) \, \mathbf{i}_{\partial g, n} \big(S_{\mathcal{L}_{0,1}}(\varphi_{(2)}) \big) \cdot w = s_i \cdot \big(w \bullet S^{-1}(t_i) \big) \tag{87}$$

for all $w \in M$. Also note that

$$s_i \,\varepsilon(t_i) = \mathfrak{i}_{\mathbf{K}}(\varphi) \tag{88}$$

because $\varepsilon_H \circ \Phi_{0,1} = \varepsilon_{H^*}$ and $\varepsilon_{H^*} \circ S_{\mathcal{L}_{0,1}} = \varepsilon_{H^*}$. *Proof of item 1.* If $w \in M^{\text{inv}}$ then by (87) and (88), $\mathbf{i}_{\mathbf{K}_{\partial}}(\varphi) \cdot w = s_i \cdot (w \varepsilon(S^{-1}(t_i))) = \mathbf{i}_{\mathbf{K}}(\varphi)$. *Proof of item 2.* Since $\varphi \in \mathcal{L}_{0,1}^{\text{inv}}(H)$, it is easily seen from (42) and (50) that

$$\forall h \in H, \ \operatorname{coad}^{r}(h)(\varphi_{(1)}) \otimes \varphi_{(2)} = \varphi_{(1)} \otimes \operatorname{coad}^{r}(S(h))(\varphi_{(2)})$$

By *H*-linearity of the maps $i_{\mathbf{K}}$, $S_{\mathcal{L}_{0,1}}$ and $\Phi_{0,1}$ we deduce that

$$\forall h \in H, \quad \operatorname{coad}^{r}(h)(s_{i}) \otimes t_{i} = s_{i} \otimes \operatorname{ad}^{r}(S(h))(t_{i})$$
(89)

where ad^r is the right adjoint action (15). Recall that for all $P \in \operatorname{mod} H$ the representation (63) of $\mathcal{L}_{g,n}^{\operatorname{inv}}(H)$ on $\operatorname{Hom}_H(P, M)$ is given by acting on the target. Using Lemma 5.9 and Proposition C.1 we get

$$\left\langle \mathbf{i}_{\mathbf{K}_{\partial}}(\varphi) \cdot f, p \right\rangle = \mathbf{i}_{\mathbf{K}}(\varphi) \cdot f(p) \stackrel{(87)}{=} s_i \cdot \left(f(p) \bullet S^{-1}(t_i) \right) = s_i \cdot f\left(p \bullet S^{-1}(t_i) \right)$$
(90)

for all $f \in \operatorname{Hom}_{H}(P, M)$ and $p \in P$. Endow the dual vector space M^{*} with the right H-action defined by

$$\forall \xi \in M^*, \ \forall h \in H, \ \forall w \in M, \quad \langle \xi \bullet h, w \rangle = \xi \big(w \bullet S^{-1}(h) \big). \tag{91}$$

By finite-dimensionality of M, there is an isomorphism

$$J_P : \operatorname{Hom}_H(P, M) \xrightarrow{\sim} \operatorname{Hom}_H(M^* \otimes P, \Bbbk), \quad \langle J_P(f), \xi \otimes p \rangle = \xi(f(p))$$

and the family $J = (J_P)_{P \in \text{mod}-H}$ is a natural isomorphism. Through J_P we can transport the representation of $\mathcal{L}_{g,n}^{\text{inv}}(H)$ on $\text{Hom}_H(P, M)$ to a representation on $\text{Hom}_H(M^* \otimes P, \Bbbk)$. For $a \in \mathcal{L}_{g,n}^{\text{inv}}(H)$ and $F \in \text{Hom}_H(M^* \otimes P, \Bbbk)$ it is defined by

$$\forall \xi \otimes p \in M^* \otimes P, \quad \langle a \cdot F, \xi \otimes p \rangle = F(\xi(a \cdot ?) \otimes p)$$
(92)

where $\xi(a \cdot ?)$ is the linear form $w \mapsto \xi(a \cdot w)$. It follows from (90) and (92) that

$$\forall F \in \operatorname{Hom}_{H}(M^{*} \otimes P, \Bbbk), \qquad \mathfrak{i}_{\mathbf{K}_{\partial}}(\varphi) \cdot F = F \circ T_{P}^{\partial} \quad \text{and} \quad \mathfrak{i}_{\mathbf{K}}(\varphi) \cdot F = F \circ T_{P}.$$
(93)

where T_P^{∂} and T_P are the endomorphisms of $M^* \otimes P$ defined by

$$T_P^{\partial}(\xi \otimes p) = \xi(s_i \cdot ?) \otimes (p \bullet S^{-1}(t_i)), \quad T_P(\xi \otimes p) = \xi(\mathbf{i}_{\mathbf{K}}(\varphi) \cdot ?) \otimes p.$$

A key property is that these endomorphisms are *H*-linear. Indeed,

$$T_{P}^{\partial}(\xi \otimes p) \bullet h \stackrel{(91)}{=} \xi \left(s_{i} \cdot (? \bullet S^{-1}(h_{(1)})) \otimes \left(p \bullet S^{-1}(t_{i})h_{(2)} \right) \right)$$

= $\xi \left(s_{i} \cdot (? \bullet S^{-1}(h_{(1)})) \otimes \left(p \bullet h_{(3)}S^{-1} \left(\operatorname{ad}^{r}(h_{(2)})(t_{i}) \right) \right) \right)$
 $\stackrel{(89)}{=} \xi \left(\operatorname{coad}^{r}(S^{-1}(h_{(2)}))(s_{i}) \cdot (? \bullet S^{-1}(h_{(1)})) \otimes \left(p \bullet h_{(3)}S^{-1}(t_{i}) \right) \right)$
 $\stackrel{(7)}{=} \xi \left((s_{i} \cdot ?) \bullet S^{-1}(h_{(1)}) \right) \otimes \left(p \bullet h_{(2)}S^{-1}(t_{i}) \right) = T_{P}^{\partial} ((\xi \otimes p) \bullet h)$

while *H*-linearity of T_P is clear because $\mathfrak{i}_{\mathbf{K}}(\varphi) \in \mathcal{L}_{g,n}^{\mathrm{inv}}(H)$ by *H*-linearity of $\mathfrak{i}_{\mathbf{K}}$ and hence its action on *M* commutes with • by item 1 in Lemma 2.3. The following naturality property is readily seen: for all $d \in \operatorname{Hom}_H(P, Q)$ we have commutative diagrams

$$\begin{array}{lll}
M^* \otimes Q \stackrel{\operatorname{id}_{M^*} \otimes d}{\longleftarrow} M^* \otimes P & M^* \otimes Q \stackrel{\operatorname{id}_{M^*} \otimes d}{\longleftarrow} M^* \otimes P & (94) \\
T^{\partial}_{Q} \downarrow & \downarrow T^{\partial}_{P} & T_{Q} \downarrow & \downarrow T_{P} \\
M^* \otimes Q \stackrel{\operatorname{id}_{M^*} \otimes d}{\longleftarrow} M^* \otimes P & M^* \otimes Q \stackrel{\operatorname{id}_{M^*} \otimes d}{\longleftarrow} M^* \otimes P
\end{array}$$

Now let $0 \leftarrow \Bbbk \leftarrow \stackrel{d_0}{\longleftarrow} P_0 \leftarrow \stackrel{d_1}{\longleftarrow} P_1 \leftarrow \stackrel{d_2}{\longleftarrow} \dots$ be a projective resolution of \Bbbk in mod-H. For $P, X \in \text{mod-}H$, it is well-known that if P is projective then so is $X \otimes P$. Moreover the functor $M^* \otimes -$ is clearly exact (because it is exact in Vect_{\Bbbk}). Thus

$$0 \longleftarrow M^* \xleftarrow{\operatorname{id}_{M^*} \otimes d_0} M^* \otimes P_0 \xleftarrow{\operatorname{id}_{M^*} \otimes d_1} M^* \otimes P_1 \xleftarrow{\operatorname{id}_{M^*} \otimes d_2} \dots$$
(95)

is a projective resolution of M^* in mod-H. This observation together with the naturality of J yields the isomorphism

$$\overline{J}^m : \operatorname{Ext}_H^m(\Bbbk, M) \xrightarrow{\sim} \operatorname{Ext}_H^m(M^*, \Bbbk), \quad [f] \mapsto \left[J_{P_m}(f) \right]$$
(96)

for all $m \ge 0$. Through \overline{J}^m , the representation from Th. 4.12 translates of course to $a \cdot [F] = [a \cdot F]$ for all $a \in \mathcal{L}_{g,n}^{\text{inv}}(H)$ with $F \in \text{Hom}_H(M^* \otimes P_m, \Bbbk)$ and $a \cdot F$ defined in (92).

We are finally ready to prove item 2. By (94), the ordered collections of *H*-morphisms $T_{\bullet}^{\partial} = (T_{\Bbbk}^{\partial}, T_{P_0}^{\partial}, T_{P_1}^{\partial}, \ldots)$ and $T_{\bullet} = (T_{\Bbbk}, T_{P_0}, T_{P_1}, \ldots)$ both are endomorphisms of the complex (95). Note from (88) that

$$T^{\partial}_{\Bbbk}(\xi \otimes 1) = \xi(s_i \cdot ?) \otimes \left(1 \bullet S^{-1}(t_i)\right) = \xi(s_i \cdot ?) \otimes \varepsilon \left(S^{-1}(t_i)\right) = T_{\Bbbk}(\xi \otimes 1)$$

for all $\xi \in M^*$, so that $T_{\Bbbk}^{\partial} = T_{\Bbbk}$. Hence the endomorphisms of complexes T_{\bullet}^{∂} and T_{\bullet} both lift T_{\Bbbk} along the projective resolution (95). By the comparison theorem, see e.g. [ML75, Th. 6.1], it follows that T_{\bullet}^{∂} and T_{\bullet} are chain-homotopic. Applying the pullback functor $\operatorname{Hom}_{H}(-, \Bbbk)$ to T_{\bullet}^{∂} , T_{\bullet} and to the homotopy between them, we conclude that the endomorphisms of complexes

$$\operatorname{Hom}_{H}(M^{*} \otimes P_{\bullet}, \Bbbk) \to \operatorname{Hom}_{H}(M^{*} \otimes P_{\bullet}, \Bbbk), \quad F \mapsto F \circ T_{P_{\bullet}}^{\partial} \quad \text{and} \quad F \mapsto F \circ T_{P_{\bullet}}$$

are chain-homotopic as well. They thus yield the same maps on cohomology, *i.e.* on $\operatorname{Ext}_{H}^{\bullet}(M^*, \Bbbk)$. Hence for all cohomology class $[F] \in \operatorname{Ext}_{H}^{m}(M^*, \Bbbk)$ coming from a cocycle $F \in \operatorname{Hom}_{H}(M^* \otimes P_m, \Bbbk)$ we have

$$\mathbf{i}_{\mathbf{K}_{\partial}}(\varphi) \cdot [F] = [\mathbf{i}_{\mathbf{K}_{\partial}}(\varphi) \cdot F] \stackrel{(93)}{=} [F \circ T_{P_{m}}^{\partial}] = [F \circ T_{P_{m}}] \stackrel{(93)}{=} [\mathbf{i}_{\mathbf{K}}(\varphi) \cdot F] = \mathbf{i}_{\mathbf{K}}(\varphi) \cdot [F]$$

which is the desired result through the isomorphism (96).

Remark 5.10. The maps \mathbb{W}^{φ} defined in (84) can be generalized to a map \mathbb{W} on the set of framed and oriented links in $\Sigma_{g,n}^{\circ} \times [0,1]$, which are colored by elements in $\mathcal{L}_{0,1}^{\mathrm{inv}}(H)$ *i.e.* each connected component of the link is labelled by such an element; see [Fai20b, Def. 4.13]⁸. It is a generalization in the sense that $\mathbb{W}^{\varphi}(K) = \mathbb{W}(K^{\varphi})$ where K^{φ} is K colored by φ . It is possible to define \mathbb{W} by means of stated skein algebras as we did for knots. It is highly probable that Theorem 5.8 remains true when generalized to $\mathcal{L}_{0,1}^{\mathrm{inv}}(H)$ -colored links, with \mathbb{W}^{φ} replaced by \mathbb{W} . We have restricted ourselves to knots to avoid technicalities induced by having multiple basepoints (one for each connected component of the link).

⁸In [Fai20b] the links are colored by symmetric linear forms, *i.e.* $\varphi(xy) = \varphi(yx)$. These are in bijection with $\mathcal{L}_{0,1}^{\text{inv}}(H)$ via $\varphi \mapsto \varphi \triangleleft g$ where g is the pivotal element, because of (50) and (66).

5.2 Projective representations of mapping class groups

The mapping class group of an oriented surface Σ is the group of orientation-preserving homeomorphisms of Σ considered up to isotopy; we denote it by MCG(Σ). A reference is [FM12].

Recall that $\Sigma_{g,0}$ denotes the closed oriented surface of genus g and $\Sigma_{g,0}^{\circ}$ is $\Sigma_{g,0}$ with one boundary circle; in this section we mainly work with such surfaces. Our goal is to obtain projective representations of the mapping class group $MCG(\Sigma_{g,0}^{\circ})$ on the derived spaces $V^m(\Sigma_g^X)$ as a corollary of the representations of $\mathcal{L}_{g,0}^{inv}(H)$ on these spaces (Thm. 4.12). Moreover we prove as a corollary of Theorem 5.8 that the choice $X = \Bbbk$ "closes the boundary" and gives projective representations of $MCG(\Sigma_{g,0})$. These representations are shown to be equivalent to those from [LMSS23] and generalise our previous work [Fai20a] where we constructed a projective representation of $MCG(\Sigma_{g,0}^{\circ})$ on $(H^*)^{\otimes g} \cong V^0(\Sigma_g^H)$ and of $MCG(\Sigma_{g,0})$ on $Inv((H^*)^{\otimes g}) \cong V^0(\Sigma_g^k)$.

The starting point of the construction is an action of $MCG(\Sigma_{g,n}^{\circ})$ on $\mathcal{L}_{g,n}(H)$ by algebra automorphisms. While in previous papers [AS96a, Fai20a] this action was rather painful to define (see Remark 5.14), the isomorphism $\mathcal{L}_{g,n}(H) \cong \mathcal{S}_{H}^{st}(\Sigma_{g,n}^{\circ,\bullet})$ makes its existence obvious.

5.2.1 Mapping class group action on $\mathcal{L}_{q,n}(H)$

Assume that the ribbon Hopf algebra H is finite-dimensional or is a quantum group $U_q(\mathfrak{g})$ with q generic. Recall from (79) the action of the group $MCG(\Sigma_{g,n}^{\circ})$ on $\mathcal{S}_H^{st}(\Sigma_{g,n}^{\circ,\bullet})$ by algebra automorphisms. Thanks to the isomorphism $\mathcal{L}_{g,n}(H) \cong \mathcal{S}_H^{st}(\Sigma_{g,n}^{\circ,\bullet})$ we obtain a group morphism

$$\operatorname{MCG}(\Sigma_{g,n}^{\circ}) \to \operatorname{Aut}_{\operatorname{alg}}(\mathcal{L}_{g,n}(H)), \quad f \mapsto \widetilde{f}$$

$$\tag{97}$$

which actually takes values in the subgroup of *H*-module-algebra automorphisms, as already noted below (79). Recall the canonical embeddings $\mathbf{i}_{b_i}, \mathbf{i}_{a_i}, \mathbf{i}_{m_{g+j}} : \mathcal{L}_{0,1}(H) \to \mathcal{L}_{g,n}(H)$ from (55); they are particular cases of the based knots maps (82). Using the compatibility (80) between the mapping class group action and the based knots maps, we see that \tilde{f} is described as follows:

Corollary 5.11. Let $f \in MCG(\Sigma_{g,n}^{\circ})$. Then for any based knot \mathbf{K} we have $\tilde{f} \circ \mathfrak{i}_{\mathbf{K}} = \mathfrak{i}_{f(\mathbf{K})}$. In particular

$$\forall \varphi \in \mathcal{L}_{0,1}(H), \quad \widetilde{f}(\mathfrak{i}_{b_i}(\varphi)) = \mathfrak{i}_{f(b_i)}(\varphi), \quad \widetilde{f}(\mathfrak{i}_{a_i}(\varphi)) = \mathfrak{i}_{f(a_i)}(\varphi), \quad \widetilde{f}(\mathfrak{i}_{m_{g+j}}(\varphi)) = \mathfrak{i}_{f(m_{g+j})}(\varphi)$$

for all $1 \leq i \leq g$, $1 \leq j \leq n$ and these values entirely define \tilde{f} by item 2 in Prop. 4.6.

In order to have explicit formulas for \tilde{f} , one would like to express each $\mathbf{i}_{f(b_i)}$, $\mathbf{i}_{f(a_i)}$ and $\mathbf{i}_{f(m_{g+j})}$ in terms of the "standard" embeddings $\mathbf{i}_{b_1}, \mathbf{i}_{a_1}, \ldots, \mathbf{i}_{b_g}, \mathbf{i}_{a_g}, \mathbf{i}_{m_{g+1}}, \ldots, \mathbf{i}_{m_{g+n}}$ used to describe $\mathcal{L}_{g,n}(H)$ in Prop. 4.6. This relies on a remarkably simple formula:

Proposition 5.12. Let $\gamma \in \pi_1(\Sigma_{g,n}^\circ)$ be a simple loop and $\gamma = \gamma_1 \dots \gamma_l$ be its decomposition in terms of the generators $a_i^{\pm 1}$, $b_i^{\pm 1}$, $m_{g+j}^{\pm 1}$ fixed in (67). Then

$$\forall \varphi \in \mathcal{L}_{0,1}(H), \quad \mathfrak{i}_{\gamma}(\varphi) = \varphi_{(1)}(v^{N(\gamma)}) \mathfrak{i}_{\gamma_1}(\varphi_{(2)}) \mathfrak{i}_{\gamma_2}(\varphi_{(3)}) \dots \mathfrak{i}_{\gamma_l}(\varphi_{(l+1)})$$

where $\varphi_{(1)} \otimes \ldots \otimes \varphi_{(l+1)}$ is iteration of the coproduct (41), $v \in H$ is the ribbon element and $N(\gamma) \in \mathbb{Z}$ is defined in Def. B.2 of Appendix B.2.

Proof. The proof is contained in Appendix B.2. Here we just note that $\pi_1(\Sigma_{g,n}^{\circ})$ is a free group generated by the loops a_i , b_i , m_{g+j} . Hence the decomposition $\gamma = \gamma_1 \dots \gamma_l$ is almost unique: the only possible variations are of the form ss^{-1} and $s^{-1}s$ for some generator s. They can be cancelled without changing the claimed formula for \mathbf{i}_{γ} because by item 2 in Lemma 5.3 and item 1 in Proposition 5.5 we have $\mathbf{i}_s(\varphi_{(1)})\mathbf{i}_{s^{-1}}(\varphi_{(2)}) = \mathbf{i}_s(\varphi_{(1)}S_{\mathcal{L}_{0,1}}(\varphi_{(2)})) = \mathbf{i}_s(\varepsilon) = \mathbf{1}_{S_H^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})}$ and similarly for $s^{-1}s$. This example shows how to use Cor. 5.11 and Prop. 5.12 to describe explicitly f:

Example 5.13. Let τ_{a_1} be the Dehn twist about the of the loop $a_1 \in \pi_1(\Sigma_{g,n}^\circ)$ [FM12, §3.1]. Precisely, one must consider the free homotopy class of a_1 , *i.e.* forget that it is attached to the basepoint. Then we have $\tau_{a_1}(a_i) = a_i$ for all i, $\tau_{a_1}(b_1) = b_1a_1$, $\tau_{a_1}(b_j) = b_j$ if j > 1 and $\tau_{a_1}(m_{g+k}) = m_{g+k}$ for all k. Hence $\tilde{\tau_{a_1}} : \mathcal{L}_{g,n}(H) \to \mathcal{L}_{g,n}(H)$ is given by

$$\begin{split} \widetilde{\tau_{a_1}}\big(\mathfrak{i}_{b_1}(\varphi)\big) &= \mathfrak{i}_{b_1a_1}(\varphi) = \varphi_{(1)}(v^{-1})\,\mathfrak{i}_{b_1}(\varphi_{(2)})\,\mathfrak{i}_{a_1}(\varphi_{(3)}), \quad \widetilde{\tau_{a_1}}\big(\mathfrak{i}_{a_1}(\varphi)\big) = \mathfrak{i}_{a_1}(\varphi), \\ \widetilde{\tau_{a_1}}\big(\mathfrak{i}_{b_j}(\varphi)\big) &= \mathfrak{i}_{b_j}(\varphi), \quad \widetilde{\tau_{a_1}}\big(\mathfrak{i}_{a_j}(\varphi)\big) = \mathfrak{i}_{a_j}(\varphi), \quad \widetilde{\tau_{a_1}}\big(\mathfrak{i}_{m_{g+k}}(\varphi)\big) = \mathfrak{i}_{m_{g+k}}(\varphi) \end{split}$$

for all $\varphi \in H^{\circ}$, $2 \leq j \leq g$ and $1 \leq k \leq n$.

Remark 5.14. Proposition 5.12 and Corollary 5.11 were definitions of i_{γ} and \tilde{f} in [Fai19b, §5.3.3] and [Fai20a, §5.2], building upon the ideas of [AS96a, §9.1]. Actually in my papers this was only for surfaces $\Sigma_{g,0}^{\circ}$. The automorphisms \tilde{f} were first defined for a generating set of MCG($\Sigma_{g,0}^{\circ}$); then long computations were required to check that they were indeed morphisms of algebras and that (97) was indeed a group morphism. This was then used to prove that i_{γ} is a morphism of algebras, by transforming γ into a_1 thanks to the mapping class group. Clearly the approach based on stated skein algebras presented here is much more natural. Note that these papers use a matrix formalism which makes their formulas look different from the ones here; the relation with the present notations is explained in [BFR23, §3.1, §4.1].

5.2.2 Construction of the representation

Recall that a quasitriangular Hopf algebra H is called *factorizable* if the morphism $\Phi_{0,1}$ from (56) is bijective [RS88].

Assumption: From now on, we assume that H is finite-dimensional, ribbon and factorizable. We recall that \Bbbk denotes the ground field of H.

It is proven in Proposition A.1 that, under these assumptions on H, there is an isomorphism of algebras

$$\mathcal{L}_{g,n}(H) \cong \operatorname{End}_{\Bbbk}(H^*)^{\otimes g} \otimes H^{\otimes n}$$
, and in particular $\mathcal{L}_{g,0}(H) \cong \operatorname{End}_{\Bbbk}((H^*)^{\otimes g})$.

Hence $\mathcal{L}_{g,0}(H)$ is isomorphic to a matrix algebra. By the Skolem–Noether theorem, any automorphism of $\mathcal{L}_{g,0}(H)$ is the conjugation by some invertible element unique up to scalar. In other words there is an isomorphism of groups

$$\operatorname{Aut}_{\operatorname{Alg}}(\mathcal{L}_{g,0}(H)) \cong \mathcal{L}_{g,0}(H)^{\times}/\mathbb{k}^{\times}$$
(98)

where the superscript \times means subset of invertible elements.

Recall the action of $MCG(\Sigma_{g,n}^{\circ})$ on $\mathcal{L}_{g,n}(H)$ by algebra automorphisms, described in Corollary 5.11 and which comes from the isomorphism between $\mathcal{L}_{g,n}(H)$ and the stated skein algebra. Combined with (98), it gives a map

$$\operatorname{MCG}(\Sigma_{g,0}^{\circ}) \longrightarrow \operatorname{Aut}_{\operatorname{Alg}}(\mathcal{L}_{g,0}(H)) \xrightarrow{\sim} \mathcal{L}_{g,0}(H)^{\times}/\mathbb{k}^{\times}
 f \longrightarrow \widetilde{f} \longrightarrow \widehat{f}
 \tag{99}$$

where the unique up to scalar invertible element \hat{f} is defined by

$$\forall x \in \mathcal{L}_{g,0}(H), \quad \tilde{f}(x) = \hat{f} x \, \hat{f}^{-1}.$$
(100)

Lemma 5.15. For all $f \in MCG(\Sigma_{g,0}^{\circ})$ we have $\widehat{f} \in \mathcal{L}_{g,0}^{inv}(H)$, i.e. the element \widehat{f} is invariant for the *H*-module structure coad^r from (54).

Proof. Since H is factorizable we have the quantum moment map $\mu_{g,0} = \mathfrak{i}_{\partial_{g,0}} \circ \Phi_{0,1}^{-1} : H \to \mathcal{L}_{g,0}(H)$ where $\partial_{g,0}$ is the boundary loop (68), see Appendix C. Thus by (16) it suffices to check that \widehat{f} commutes with $\mu_{g,0}(h)$ for all $h \in H$:

$$\widehat{f}\,\mu_{g,0}(h) = \widetilde{f}\left(\mu_{g,0}(h)\right)\,\widehat{f} = \widetilde{f}\circ\mathfrak{i}_{\partial_{g,0}}\left(\Phi_{0,1}^{-1}(h)\right)\,\widehat{f} = \mathfrak{i}_{f(\partial_{g,0})}\left(\Phi_{0,1}^{-1}(h)\right)\,\widehat{f} = \mathfrak{i}_{\partial_{g,0}}\left(\Phi_{0,1}^{-1}(h)\right)\,\widehat{f} = \mu_{g,0}(h)\,\widehat{f}$$

where the first equality uses the defining property of \hat{f} , the third uses Corollary 5.11 and the fourth uses that the mapping class group fixes pointwise $\partial_{g,0}$ by definition.

Lemma 5.15 allows us to combine the group morphism (99) with the representations of $\mathcal{L}_{g,0}^{\text{inv}}(H)$ on $V(\Sigma_g^X) = \text{Hom}_H(X, (H^*)^{\otimes g})$ and $V^m(\Sigma_g^X) = \text{Ext}_H^m(X, (H^*)^{\otimes g})$ from §4.2. Hence, for all $X \in \text{mod-}H$, we get projective representations:

$$\mathcal{R}_X : \mathrm{MCG}(\Sigma_{g,0}^{\circ}) \longrightarrow \mathrm{Aut}_{\mathrm{Alg}}(\mathcal{L}_{g,0}(H)) \xrightarrow{\sim} \mathcal{L}_{g,0}(H)^{\times}/\Bbbk^{\times} \xrightarrow{(63)} \mathrm{PGL}(V(\Sigma_g^X)), \qquad (101)$$

$$\mathcal{R}_X^m : \mathrm{MCG}(\Sigma_{g,0}^{\circ}) \longrightarrow \mathrm{Aut}_{\mathrm{Alg}}(\mathcal{L}_{g,0}(H)) \xrightarrow{\sim} \mathcal{L}_{g,0}(H)^{\times}/\Bbbk^{\times} \xrightarrow{\mathrm{Th.\,4.12}} \mathrm{PGL}(V^m(\Sigma_g^X)).$$

More explicitly

$$\forall f \in \mathrm{MCG}(\Sigma_{g,0}^{\circ}), \ \forall s \in V(\Sigma_g^X), \quad \mathcal{R}_X(f)(s) = \widehat{f} \cdot s \tag{102}$$

where $\widehat{f} \cdot s$ is given by acting on the target of s, *i.e.* on $(H^*)^{\otimes g}$, thanks to the representation of $\mathcal{L}_{g,0}(H)$ on $(H^*)^{\otimes g}$ in Prop. 3.2. If $0 \leftarrow X \leftarrow P_0 \leftarrow P_1 \leftarrow \ldots$ is a projective resolution in mod-H, then for any cohomology class $[s] \in V^m(\Sigma_g^X)$ of a cocycle $s \in V(\Sigma_g^{P_m})$ we have

$$\forall f \in \mathrm{MCG}(\Sigma_{g,0}^{\circ}), \quad \mathcal{R}_X^m(f)([s]) = [\mathcal{R}_{P_m}(f)(s)] = [\widehat{f} \cdot s].$$
(103)

Note that $V^0(\Sigma_q^X) \cong V(\Sigma_q^X)$ and the projective representation \mathcal{R}_X^0 is equivalent to \mathcal{R}_X .

The natural group morphism $MCG(\Sigma_{g,0}^{\circ}) \to MCG(\Sigma_{g,0})$ is surjective [FM12, §4.2.5] and thus $MCG(\Sigma_{g,0})$ is a quotient of $MCG(\Sigma_{g,0}^{\circ})$. For a special choice of X in (101) we obtain a projective representation of this quotient:

Theorem 5.16. Let \Bbbk be the trivial *H*-module. Then \mathcal{R}^m_{\Bbbk} descends to a projective representation of $MCG(\Sigma_{g,0})$ on $V^m(\Sigma^k_g)$, for all $m \ge 0$. In particular this is true for the projective representation \mathcal{R}_k on $V(\Sigma^k_g)$.

The rest of this section is dedicated to the proof of this theorem.

Lemma 5.17. Let $\rho : \text{MCG}(\Sigma_{g,0}^{\circ}) \to G$ be a group morphism. Consider the curves α_g and δ defined in Fig. 2 on p.35 below. If $\rho(\tau_{\alpha g}) = \rho(\tau_{\delta})$ then ρ factors through $\text{MCG}(\Sigma_{g,0})$.

Proof. This follows from Wajnryb's presentation of $MCG(\Sigma_{g,0})$ and $MCG(\Sigma_{g,0})$ ([Waj83], also given in [FM12, Th. 5.3]). In this presentation one uses the Humphries generators, which are the Dehn twists about the curves $\alpha_1, \alpha_2, \beta_i$ $(1 \le i \le g)$ and γ_j $(1 \le j \le g - 1)$ in Fig. 2 below. For commodity write $\gamma_0 = \alpha_1$ and let

$$f = (\tau_{\beta_g} \tau_{\gamma_{g-1}})(\tau_{\beta_{g-1}} \tau_{\gamma_{g-2}}) \dots (\tau_{\beta_1} \tau_{\gamma_0})(\tau_{\gamma_0} \tau_{\beta_1})(\tau_{\gamma_1} \tau_{\beta_2}) \dots (\tau_{\gamma_{g-1}} \tau_{\beta_g}).$$

Then the presentation for $MCG(\Sigma_{g,0})$ is obtained by adding the relation $f\tau_{\delta}f^{-1} = \tau_{\delta}$ to the presentation of $MCG(\Sigma_{g,0}^{\circ})$.⁹ Using drawings, it is not hard to see that $f(\delta) = \alpha_g$ and thus

$$\rho(f)\rho(\tau_{\delta})\rho(f)^{-1} = \rho(f\tau_{\delta}f^{-1}) = \rho(\tau_{f(\delta)}) = \rho(\tau_{\alpha_g})$$

where the second equality is a classical fact on Dehn twists [FM12, §3.3]. Hence by assumption on ρ we have $\rho(f)\rho(\tau_{\delta})\rho(f)^{-1} = \rho(\tau_{\delta})$, which shows that ρ is compatible with the extra relation.

⁹Actually to have a presentation in the formal sense one must express τ_{δ} in terms of the Humphries generators, see [FM12, p. 127].

We want to apply Lemma 5.17 with $\rho = \mathcal{R}^m_{\Bbbk}$. This is based on an auxilliary result (Prop. 5.19) whose statement and proof require some preliminaries.

Recall that a *left integral* for H is a non-zero linear form $\lambda \in H^*$ such that

$$\forall h \in H, \quad \lambda(h_{(2)})h_{(1)} = \lambda(h)1_H. \tag{104}$$

Such a λ is known to be unique up to non-zero scalar and satisfies

$$\forall x, y \in H, \quad \lambda(xy) = \lambda(yS^2(x)), \qquad i.e. \ \lambda \in \mathcal{L}_{0,1}^{\text{inv}}(H)$$
(105)

see e.g. [Rad11]. A consequence of the factorizability assumption on H is the following formula

$$\lambda \circ S^{-1} = \lambda \triangleleft g^{-2} \tag{106}$$

where g is the pivotal element (65); for a proof see e.g. [Fai19a, Lem. 5.9]. Here is a consequence of these facts:

Lemma 5.18. Let $S_{\mathcal{L}_{0,1}}$ be the antipode (48) on $\mathcal{L}_{0,1}(H)$. We have

$$S_{\mathcal{L}_{0,1}}(\lambda) = \lambda, \qquad S_{\mathcal{L}_{0,1}}(v \triangleright \lambda) = v \triangleright \lambda.$$

Proof. For all $h \in H$,

$$\langle S_{\mathcal{L}_{0,1}}(\lambda), h \rangle = \lambda \left(R_{[2]} u^{-1} S(h) S(R_{[1]}) \right) \stackrel{(105)}{=} \lambda \left(S^{-1}(R_{[1]}) R_{[2]} u^{-1} S(h) \right)$$

Note that

$$S^{-1}(R_{[1]})R_{[2]}u^{-1} = S^{-1}\left(S(R_{[2]})R_{[1]}\right)u^{-1} \stackrel{(37)}{=} S^{-1}(u)u^{-1} \stackrel{(65)}{=} S^{-1}(gv)g^{-1}v^{-1} = g^{-2}$$

because g is grouplike (whence $S(g) = g^{-1}$) and v is central and satisfies S(v) = v by definition. We conclude that $\langle S_{\mathcal{L}_{0,1}}(\lambda), h \rangle = \lambda(g^{-2}S(h)) = \lambda(h)$ thanks to (106). For the second claim, simply note that $S_{\mathcal{L}_{0,1}}(v \triangleright \lambda) = v \triangleright S_{\mathcal{L}_{0,1}}(\lambda)$ because v is central and satisfies S(v) = v. \Box

Let $c \,\subset\, \Sigma_{g,0}^{\circ}$ be a closed curve considered up to homotopy. As opposed to elements in $\pi_1(\Sigma_{g,0}^{\circ})$, c is not assumed to be attached to a basepoint or to have an orientation. The closed curve c is called simple if its homotopy class contains a representative without self-intersections. In this case we have the *Dehn twist* $\tau_c \in \mathrm{MCG}(\Sigma_{g,0}^{\circ})$, see e.g. [FM12, §3.1]. We now give a formula for the element $\hat{\tau}_c \in \mathcal{L}_{g,n}^{\mathrm{inv}}(H)$ defined by (100). Note that a simple

We now give a formula for the element $\hat{\tau}_c \in \mathcal{L}_{g,n}^{\mathrm{inv}}(H)$ defined by (100). Note that a simple closed curve c endowed with an orientation can be seen as a knot in $\Sigma_{g,0}^{\circ} \times [0,1]$ and thus by (83) we have the knot map $\mathfrak{i}_c : \mathcal{L}_{0,1}^{\mathrm{inv}}(H) \to \mathcal{L}_{g,n}^{\mathrm{inv}}(H)$. Also recall that c is called non-separating if $\Sigma_{g,0}^{\circ} \setminus c$ is connected.

Proposition 5.19. Let $c \in \Sigma_{g,0}^{\circ}$ be a non-separating simple closed curve endowed with an arbitrary orientation and denote by τ_c the Dehn twist about c. Then

$$\widehat{\tau}_c = \mathfrak{i}_c(v \triangleright \lambda).$$

A related formula was given in [AS96a, §9.2] for semisimple H; it was generalized to any H in [Fai20a, Lem. 5.3] and proved through the isomorphism between $\mathcal{L}_{1,0}(H)$ and the Heisenberg double (see App. A). Here is a proof entirely in $\mathcal{L}_{g,n}(H)$ and within the current notations:

Proof. Note that the element $\mathbf{i}_c(v \triangleright \lambda)$ is independent of the orientation on c, thanks to item 2 in Lemma 5.3 and Lemma 5.18. We first prove the result in the case where c is the simple closed curve α_1 in Fig. 2 below. The generator $a_1 \in \pi_1(\Sigma_{g,n}^\circ)$ defined in (67) can be choosen as a

based knot representing α_1 , so that by definition (83) we have $\mathfrak{i}_{\alpha_1}(v \triangleright \lambda) = \mathfrak{i}_{a_1}(v \triangleright \lambda)$. According to Example 5.13 we must show that for all $\varphi \in \mathcal{L}_{0,1}(H)$:

$$\mathbf{i}_{a_1}(v \triangleright \lambda)\mathbf{i}_{a_1}(\varphi) = \mathbf{i}_{a_1}(\varphi)\mathbf{i}_{a_1}(v \triangleright \lambda),\tag{107}$$

$$\mathbf{i}_{a_1}(v \triangleright \lambda)\mathbf{i}_{b_1}(\varphi) = \varphi_{(1)}(v^{-1})\mathbf{i}_{b_1}(\varphi_{(2)})\mathbf{i}_{a_1}(\varphi_{(3)})\mathbf{i}_{a_1}(v \triangleright \lambda),$$
(108)

 $\mathbf{i}_{a_1}(v \triangleright \lambda)\mathbf{i}_{a_j}(\varphi) = \mathbf{i}_{a_j}(\varphi)\mathbf{i}_{a_1}(v \triangleright \lambda), \quad \mathbf{i}_{a_1}(v \triangleright \lambda)\mathbf{i}_{b_j}(\varphi) = \mathbf{i}_{b_j}(\varphi)\mathbf{i}_{a_1}(v \triangleright \lambda) \quad \text{for } j \ge 2.$ (109)

(107) is immediate from Lemma 4.3 and (105). The two equalities in (109) are also immediate from item 4 in Prop. 4.6 and (105). For (108) note that $\varphi(v \triangleright \lambda) = \varphi \star (v \triangleright \lambda)$ for all $\varphi \in \mathcal{L}_{0,1}(H)$, due to (105) and Lemma 4.3. But

$$\langle \varphi \star (v \triangleright \lambda), h \rangle = \varphi(h_{(1)})\lambda(vh_{(2)}) = \varphi\big(S(v_{(1)})v_{(2)}h_{(1)}\big)\lambda\big(v_{(3)}h_{(2)}\big) \stackrel{(104)}{=} \langle \varphi\big(S(v_{(1)})\big)(\lambda \triangleleft v_{(2)}), h \rangle.$$
(110)

Using this fact we get that the right-hand side of (108) equals

$$\varphi_{(1)}(v^{-1}) \, \mathfrak{i}_{b_1}(\varphi_{(2)}) \mathfrak{i}_{a_1} \big(\varphi_{(3)}(v \triangleright \lambda) \big) \stackrel{(110)}{=} \varphi_{(1)}(v^{-1}) \, \mathfrak{i}_{b_1}(\varphi_{(2)}) \varphi_{(3)} \big(S(v_{(1)}) \big) \mathfrak{i}_{a_1}(\lambda \triangleleft v_{(2)})$$

$$\stackrel{(42)}{=} \mathfrak{i}_{b_1} \big(v^{-1} S(v_{(1)}) \triangleright \varphi \big) \mathfrak{i}_{a_1}(\lambda \triangleleft v_{(2)})$$

where we also used the centrality of v. The exchange relation in item 3 of Prop. 4.6 allows us to check that this result agrees with the left-hand side of (108):

$$\begin{aligned} \mathbf{i}_{a_{1}}(v \triangleright \lambda)\mathbf{i}_{b_{1}}(\varphi) &= \mathbf{i}_{b_{1}}\left(R_{[2]}^{4}R_{[1]}^{3} \triangleright \varphi \triangleleft R_{[1]}^{1}R_{[1]}^{2}\right)\mathbf{i}_{a_{1}}\left(vR_{[2]}^{3}S(R_{[2]}^{1}) \triangleright \lambda \triangleleft R_{[2]}^{2}R_{[1]}^{4}\right) \\ \stackrel{(105)}{=} \mathbf{i}_{b_{1}}\left(R_{[2]}^{4}R_{[1]}^{3} \triangleright \varphi \triangleleft R_{[1]}^{1}R_{[1]}^{2}\right)\mathbf{i}_{a_{1}}\left(\lambda \triangleleft vS^{-2}(R_{[2]}^{3})S^{-1}(R_{[2]}^{1})R_{[2]}^{2}R_{[1]}^{4}\right) \\ \stackrel{(22)}{=} \mathbf{i}_{b_{1}}\left(R_{[2]}^{4}R_{[1]}^{3} \triangleright \varphi\right)\mathbf{i}_{a_{1}}\left(\lambda \triangleleft vS^{-2}(R_{[2]}^{3})R_{[1]}^{4}\right) \\ \stackrel{(23)}{=} \mathbf{i}_{b_{1}}\left(S\left(S(R_{[1]}^{3})S^{-1}(R_{[2]}^{4})\right) \triangleright \varphi\right)\mathbf{i}_{a_{1}}\left(\lambda \triangleleft vR_{[2]}^{3}R_{[1]}^{4}\right) \\ &= \mathbf{i}_{b_{1}}\left(v^{-1}S(v_{(1)}) \triangleright \varphi\right)\mathbf{i}_{a_{1}}(\lambda \triangleleft v_{(2)}) \end{aligned}$$

where for the last equality we used (22) and the axiom $\Delta(v) = (R'R)^{-1}(v \otimes v)$ of the ribbon element, also recall (22) for R^{-1} . Now let c be as in the statement. By the change of coordinate principle [FM12, §1.3.1], there exists $f \in \text{MCG}(\Sigma_{g,0}^{\circ})$ such that $f(\alpha_1) = c$. Then $\tau_c = f \circ \tau_{\alpha_1} \circ f^{-1}$ [FM12, Fact 3.7] and $\mathfrak{i}_c = \tilde{f} \circ \mathfrak{i}_{\alpha_1}$ (Cor. 5.11). Hence, using the result for α_1 just established, we find

$$\widehat{\tau_c} = \widehat{f} \, \widehat{\tau_{\alpha_1}} \, \widehat{f}^{-1} \stackrel{(100)}{=} \widetilde{f}(\widehat{\tau_{\alpha_1}}) = \widetilde{f}(\mathfrak{i}_{\alpha_1}(v \triangleright \lambda)) = \mathfrak{i}_c(v \triangleright \lambda). \qquad \Box$$

Proof of Theorem 5.16. By Lemma 5.17 it suffices to show that $\mathcal{R}^m_{\Bbbk}(\tau_{\alpha g}) = \mathcal{R}^m_{\Bbbk}(\tau_{\delta})$. Recall that simple closed curves in Σ can be seen as knots in $\Sigma \times [0, 1]$. Note that the simple closed curves α_g and δ become isotopic under the injection $\Sigma^{\circ}_{g,0} \hookrightarrow \Sigma_{g,0}$, thus the same is true if we look at them as knots in $\Sigma^{\circ}_{g,0} \times [0, 1]$. As a result Thm. 5.8 applies to the associated knot maps $\mathfrak{i}_{\alpha g}, \mathfrak{i}_{\delta} : \mathcal{L}^{\mathrm{inv}}_{0,1}(H) \to \mathcal{L}^{\mathrm{inv}}_{g,n}(H)$ and we get

$$\mathcal{R}^{m}_{\Bbbk}(\tau_{\alpha g})\big([s]\big) \stackrel{(103)}{=} \widehat{\tau_{\alpha g}} \cdot [s] = \mathfrak{i}_{\alpha g}(v \triangleright \lambda) \cdot [s] = \mathfrak{i}_{\delta}(v \triangleright \lambda) \cdot [s] = \widehat{\tau_{\delta}} \cdot [s] \stackrel{(103)}{=} \mathcal{R}^{m}_{\Bbbk}(\tau_{\delta})\big([s]\big).$$

for all $[s] \in V^m(\Sigma_g^{\Bbbk})$, where the second and fourth equalities use Prop. 5.19 while the third equality uses Thm. 5.8.

5.2.3 Explicit formulas and equivalence with the LMSS representations

Note that $V(\Sigma_g^H) = \operatorname{Hom}_H(H, (H^*)^{\otimes g}) \cong (H^*)^{\otimes g}$, where H is considered as a module over itself (regular representation). Through this identification, \mathcal{R}_H from (101) becomes a projective representation on $(H^*)^{\otimes g}$ which we simply denote by \mathcal{R} :

$$\forall f \in \mathrm{MCG}(\Sigma_{g,0}^{\circ}), \ \forall \Psi \in (H^*)^{\otimes g}, \quad \mathcal{R}(f)(\Psi) = \widehat{f} \cdot \Psi$$

where \cdot is the representation of $\mathcal{L}_{g,0}(H)$ on $(H^*)^{\otimes g}$ (§4.2) and \widehat{f} is defined in (100). As a side remark, $\mathcal{R}(f)$ commutes with the right action (61) of H on $(H^*)^{\otimes g}$; this is due to Lemma 5.15 and item 1 in Lemma 2.3. The representation \mathcal{R} encapsulates the action of $\mathrm{MCG}(\Sigma_{g,0}^{\circ})$ on the spaces $V(\Sigma_g^X)$ and their derived version $V^m(\Sigma_g^X)$ for all $X \in \mathrm{mod}$ -H. Indeed (102) is rewritten as

$$\forall f \in \mathrm{MCG}(\Sigma_{g,0}^{\circ}), \ \forall s \in V(\Sigma_g^X), \quad \mathcal{R}_X(f)(s) = \mathcal{R}(f) \circ s.$$
(111)

Similarly (103) is rewritten as

$$\forall f \in \mathrm{MCG}(\Sigma_{g,0}^{\circ}), \ \forall [s] \in V^m(\Sigma_g^X), \quad \mathcal{R}_X^m(f)([s]) = [\mathcal{R}(f) \circ s].$$
(112)

Let us then give the value of \mathcal{R} on the *Lickorish generators* of $MCG(\Sigma_{g,0}^{\circ})$, which are the Dehn twists about the curves α_i , β_i , γ_j represented in Fig. 2.



Figure 2: Some simple closed curves on the surface $\sum_{a,0}^{\circ}$.

Proposition 5.20. The representations of the Lickorish generators through \mathcal{R} are given by

 $\mathcal{R}(\tau_{\alpha_i})(\psi_1 \otimes \ldots \otimes \psi_g) = \psi_1 \otimes \ldots \otimes \psi_{i-1} \otimes (v^{-1} \triangleright \psi_i) \otimes \psi_{i+1} \otimes \ldots \otimes \psi_g,$ $\mathcal{R}(\tau_{\beta_i})(\psi_1 \otimes \ldots \otimes \psi_g) = \psi_i \big(S(v_{(1)}) \big) \psi_1 \otimes \ldots \otimes \psi_{i-1} \otimes (\lambda \triangleleft v_{(2)}) \otimes \psi_{i+1} \otimes \ldots \otimes \psi_g,$ $\mathcal{R}(\tau_{\gamma_j})(\psi_1 \otimes \ldots \otimes \psi_g) = \psi_1 \otimes \ldots \otimes \psi_{j-1} \otimes \big(S(v_{(1)}^{-1}) \triangleright \psi_j \big) \otimes \big(\psi_{j+1} \triangleleft v_{(2)}^{-1} \big) \otimes \psi_{j+2} \otimes \ldots \otimes \psi_g$

for all $\psi_1 \otimes \ldots \otimes \psi_g \in (H^*)^{\otimes g}$, where v is the ribbon element and λ is the left integral of H normalized by $\lambda(v) = 1$.

Proof. We already computed these formulas in [Fai20a, Th. 5.12], modulo the fact that there we used another representation of $\mathcal{L}_{g,0}(H)$ on $(H^*)^{\otimes g}$, namely the Alekseev representation explained in appendix A. This gives another projective representation \mathcal{R}' of $MCG(\Sigma_{g,0}^{\circ})$ on $(H^*)^{\otimes g}$ defined by $\mathcal{R}'(f) = \Phi'_{g,0}(\widehat{f})$ with $\Phi'_{g,0} : \mathcal{L}_{g,0}(H) \to End_{\Bbbk}((H^*)^{\otimes g})$ from Prop. A.1. Within the present notations, the resulting formulas are¹⁰

$$\mathcal{R}'(\tau_{\alpha_i})(\psi_1 \otimes \ldots \otimes \psi_g) = \psi_1 \otimes \ldots \otimes \psi_{i-1} \otimes (v^{-1} \triangleright \psi_i) \otimes \psi_{i+1} \otimes \ldots \otimes \psi_g,$$

$$\mathcal{R}'(\tau_{\beta_i})(\psi_1 \otimes \ldots \otimes \psi_g) = \psi_1 \otimes \ldots \otimes \psi_{i-1} \otimes \left[v^{-1} \triangleright \left((\lambda \triangleleft g^{-1}v) \star (v \triangleright \psi_i)\right)\right] \otimes \psi_{i+1} \otimes \ldots \otimes \psi_g,$$

$$\mathcal{R}'(\tau_{\gamma_j})(\psi_1 \otimes \ldots \otimes \psi_g) = \psi_1 \otimes \ldots \otimes \psi_{j-1} \otimes \left(R^1_{[2]}v^{-1}_{(2)}R^2_{[2]} \triangleright \psi_j \triangleleft S^{-1}(R^2_{[1]})R^1_{[1]}\right)$$

$$\otimes \left(R^3_{[2]}R^4_{[2]} \triangleright \psi_{j+1} \triangleleft S^{-1}(R^4_{[1]})S^{-1}(v^{-1}_{(1)})R^3_{[1]}\right) \otimes \psi_{j+2} \otimes \ldots \otimes \psi_g$$

From (124) below we have an isomorphism $F^{\otimes g} : (H^*)^{\otimes g} \to (H^*)^{\otimes g}$ from the representation of §4.2 (used to build \mathcal{R}) to the Alekseev representation (used to build \mathcal{R}'), where $F(\psi) = R_{[2]} \triangleright \psi \triangleleft R_{[1]}$. Hence

$$F^{\otimes g}\big(\mathcal{R}(f)(\psi_1\otimes\ldots\otimes\psi_g)\big)=\mathcal{R}'(f)\big(F(\psi_1)\otimes\ldots\otimes F(\psi_g)\big).$$

¹⁰In [Fai19b], α_i was denoted by a_j , β_j by b_j and γ_j by d_{j+1} . Note the shift in this last case.

The formula for $\mathcal{R}(\tau_{\alpha_i})$ immediately follows from this equality by centrality of v. The formula for $\mathcal{R}(\tau_{\gamma_j})$ is easily seen as well, using that S(v) = v. The formula for $\mathcal{R}(\tau_{\beta_i})$ can also be obtained from the isomorphism $F^{\otimes g}$, but it is a non-trivial exercise which requires involved computations. It is much faster to prove it directly. Due to Proposition 5.19, the fact (105) that $v \triangleright \lambda \in \mathcal{L}_{0,1}^{inv}(H)$ and Remark 4.11 we have

$$\mathcal{R}(\tau_{\beta_i})(\psi_1 \otimes \ldots \otimes \psi_g) = \mathfrak{i}_{b_i}(v \triangleright \lambda) \cdot (\psi_1 \otimes \ldots \otimes \psi_g) = \psi_1 \otimes \ldots \otimes (\psi_i \star (v \triangleright \lambda)) \otimes \ldots \otimes \psi_g.$$

In (110) we already noted that $\psi \star (v \triangleright \lambda) = \psi (S(v_{(1)})) \lambda \triangleleft v_{(2)}$, hence the result.

Remark 5.21. Recall the morphism $\Phi_{0,1}$ from (56). One can show that $\Phi_{0,1}(v \triangleright \lambda) = v^{-1}$ due to the property $\Delta(v) = (R'R)^{-1}(v \otimes v)$. Using this fact together with Proposition 5.19 and Remark 4.11 gives a direct proof of the formula for $\mathcal{R}(\tau_{\alpha_i})$, *i.e.* without using \mathcal{R}' . The formula for $\mathcal{R}(\tau_{\gamma_i})$ is much harder to obtain directly (as it was with \mathcal{R}').

We finish by discussing the relation of our construction with that of [Lyu95] and [LMSS23]. As already recalled in §1.1, for any finite ribbon factorizable category \mathcal{C} , the Lyubashenko construction [Lyu95] produces a projective representation Z_X of $MCG(\Sigma_{g,0}^{\circ})$ on the space $Z(\Sigma_g^X) = Hom_{\mathcal{C}}(X, L^{\otimes g})$ for all $X \in \mathcal{C}$, where $L = \int^{X \in \mathcal{C}} X^* \otimes X$. It is given by pushforward:

$$\forall f \in \mathrm{MCG}(\Sigma_{g,0}^{\circ}), \ \forall s \in \mathrm{Hom}_{\mathcal{C}}(X, L^{\otimes g}), \quad Z_X(f)(s) = Z(f) \circ s$$
(113)

where $Z(f) \in \operatorname{Aut}_{\mathcal{C}}(L^{\otimes g})$ is defined on the Lickorish generators by means of certain dinatural transformations which factorize through $L^{\otimes g}$.

Take $\mathcal{C} = H$ -mod. Then L is H^* endowed with the left coadjoint action: $\operatorname{coad}^l(h)(\varphi) = h_{(2)} \triangleright \varphi \triangleleft S(h_{(1)}) = \varphi(S(h_{(1)})?h_{(2)})$ for all $h \in H$ and $\varphi \in H^*$. Recall that any right H-module (X, \bullet) is automatically a left module (and conversely) through the formula $h \cdot x = x \bullet S^{-1}(h)$. For the vector space $(H^*)^{\otimes g}$ endowed with the right H-action (61) this gives the left H-action

$$h \cdot (\psi_1 \otimes \ldots \otimes \psi_g) = \left(h_{(2g-1)} \triangleright \psi_1 \triangleleft S^{-1}(h_{(2g)})\right) \otimes \ldots \otimes \left(h_{(1)} \triangleright \psi_g \triangleleft S^{-1}(h_{(2)})\right).$$

Denote by L'_q this left *H*-module structure on $(H^*)^{\otimes g}$. Then

$$V(\Sigma_g^X) \stackrel{(62)}{=} \operatorname{Hom}_{\operatorname{mod-}H}(X, (H^*)^{\otimes g}) = \operatorname{Hom}_{H\operatorname{-mod}}(X, L'_g)$$

for any H-module X. The map

$$I: L^{\otimes g} \to L'_g, \quad \psi_1 \otimes \ldots \otimes \psi_g \mapsto (\psi_g \circ S) \otimes \ldots \otimes (\psi_2 \circ S) \otimes (\psi_1 \circ S)$$

is easily seen to be an isomorphism of *H*-modules, where $L^{\otimes g}$ is of course equipped with the diagonal action given by coproduct. As a result the pushforward $I_* : Z(\Sigma_g^X) \cong V(\Sigma_g^X)$ is an isomorphism of vector spaces for all X. We proved in [Fai20a, §6.2]¹¹ that

$$\forall f \in \mathrm{MCG}(\Sigma_{q,0}^{\circ}), \quad I \circ Z(f) = \mathcal{R}(f) \circ I.$$

It immediately follows from (111) and (113) that I_* is an intertwiner between Lyubashenko's projective representation Z_X and the projective representation \mathcal{R}_X defined in (101).

We also explained in §1.1 that in [LMSS23] the projective representation Z_X^m of $MCG(\Sigma_{g,0}^{\circ})$ on $Z^m(\Sigma_g^X) = \operatorname{Ext}_{H-\mathrm{mod}}^m(X, L^{\otimes g})$ is given by

$$\forall f \in \mathrm{MCG}(\Sigma_{g,0}^{\circ}), \ \forall [s] \in \mathrm{Ext}_{H-\mathrm{mod}}^{m}(X, L^{\otimes g}), \quad Z_{X}^{m}(f)([s]) = \left[Z(f) \circ s\right]$$
(114)

for all $m \ge 0$. It immediately follows from (112) and (114) that the linear map $[s] \mapsto [I \circ s]$ is an intertwiner between the projective representation Z_X^m of [LMSS23] and the projective representation \mathcal{R}_X^m defined in (101).

If X is the trivial H-module k then they all descend to equivalent projective representations of $MCG(\Sigma_{q,0})$.

¹¹The intertwiner I has a simpler expression than the intertwiner used in [Fai20a] because here we use \mathcal{R} while in [Fai20a] we used \mathcal{R}' , where \mathcal{R}' was recalled in the proof of Prop. 5.20.

5.3 Derived representations of skein algebras

We first explain how Theorem 4.12 yields derived representations for skein algebras defined from any ribbon Hopf algebra H by means of the Reshetikhin–Turaev tangle invariant for H-mod. We then discuss the case of the Kauffman bracket skein algebra at roots of unity, which is different in nature since it is defined by specialization.

5.3.1 General skein algebras

Let H be a ribbon Hopf algebra over a field k. The *skein algebra* $S_H(\Sigma)$ of an oriented surface Σ is the k-vector space spanned by isotopy classes of H-colored framed and oriented links in $\Sigma \times [0, 1]$, modulo so-called skein relations which are obtained by applying the Reshetikhin–Turaev invariant of tangles inside small cubes in $\Sigma \times [0, 1]$; see e.g. [BFR23, §6.2] for a complete definition.

Recall the stated skein algebra $\mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{a,n}^{\circ,\bullet})$ from §5.1.1. There is a natural map

 $\{H\text{-colored, framed, oriented links in } \Sigma_{g,n}^{\circ} \times [0,1]\} \to \mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$

because a link in $\Sigma_{g,n}^{\circ} \times [0,1]$ can be seen as a ribbon graph in $\Sigma_{g,n}^{\circ,\bullet} \times [0,1]$ without boundary points. By very definition of the right action of H on $\mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$ in (69), we see that this map actually takes values in the subalgebra of invariant elements $\mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})^{\mathrm{inv}}$. It is moreover compatible with the skein relations in $\mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$ [BFR23, Lem. 6.7] so it descends to an algebra morphism $\mathcal{S}_{H}(\Sigma_{g,n}^{\circ,\bullet}) \to \mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})^{\mathrm{inv}}$. This morphism is not surjective when H-mod is not semisimple. More surprisingly, it might a priori be not injective; the point is that there are more skein relations in $\mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$ than in $\mathcal{S}_{H}(\Sigma_{g,n}^{\circ,\bullet})$ because of the possibility to interact with the boundary. On the other hand when H-mod is semisimple then $\mathcal{S}_{H}(\Sigma_{g,n}^{\circ,\bullet}) \to \mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})^{\mathrm{inv}}$ is an isomorphism [BFR23, Cor. 6.12].

For any ribbon Hopf algebra H, there is a morphism of H-module-algebras holst : $\mathcal{S}_{H}^{st}(\Sigma_{g,n}^{\circ,\bullet}) \to \mathcal{L}_{g,n}(H)$ [BFR23, §6.2]; the morphism holst was already used in §5.1.3 under certain assumptions on H which we do not need here. From these remarks we have an algebra morphism

$$W: \mathcal{S}_H(\Sigma_{g,n}^{\circ}) \to \mathcal{S}_H^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})^{\mathrm{inv}} \xrightarrow{\mathrm{hol}^{\mathrm{st}}} \mathcal{L}_{g,n}^{\mathrm{inv}}(H)$$

called Wilson loop map. By Theorem 4.12, its existence implies:

Corollary 5.22. For all $m \ge 0$ there is a representation of $\mathcal{S}_H(\Sigma_{g,n}^\circ)$ on the derived spaces $V^m(\Sigma_{g,X_1,\ldots,X_n}^X)$.

The skein algebra $\mathcal{S}_H(\Sigma_{g,n})$ is the quotient of $\mathcal{S}_H(\Sigma_{g,n}^\circ)$ by the relation (85). Because of Theorem 5.8, when H is finite-dimensional, it is natural to expect that the choice $X = \Bbbk$ (trivial H-module) in Corollary 5.22 closes the boundary component:

More precisely this would immediately follow from the generalization of Thm. 5.8 suggested in Rmk. 5.10, because a *H*-colored link is a particular case of a $\mathcal{L}_{0,1}^{\text{inv}}(H)$ -colored link, where all colors are of the form qTr_X (see Example 4.2). In particular this would give representations of the skein algebra of the closed surface $\Sigma_{g,0}$.

5.3.2 Kauffman bracket skein algebras at roots of unity

Let q be a formal variable and $U_{q^2} = U_{q^2}(\mathfrak{sl}_2)$ be the quantum envelopping algebra of $\mathfrak{sl}_2(\mathbb{C})$, considered over the ground ring $\mathbb{C}[q^{\pm 1/2}]$.¹² It is generated by $E, F, K^{\pm 1}$ modulo the well-known relations, see e.g. in [Kas95, Def. VI.1.1]. It has a Hopf structure which can be found e.g. in [Kas95, §VII.1]. For $H = U_{q^2}$, we use $\mathcal{O}_{q^2}(\mathrm{SL}_2)$ instead of the whole $H^{\circ,13}$ Although the Hopf algebra $U_{q^2}(\mathfrak{sl}_2)$ is not quasitriangular in the usual sense, there is the well-known formula for R whose action is well defined on $X \otimes Y$ for all finite-dimensional U_{q^2} -modules X, Y; see e.g. [CP94, Prop 6.4.8]. This is enough to define $\mathcal{L}_{g,n}(U_{q^2})$ as the $\mathbb{C}[q^{\pm 1/2}]$ -module $\mathcal{O}_{q^2}(\mathrm{SL}_2)^{\otimes(2g+n)}$ endowed with the product described in Prop. 4.6.

Here is a more concrete description of $\mathcal{L}_{g,n}(U_{q^2})$. Let $\rho : U_{q^2} \to M_2(\mathbb{C}[q^{\pm 1/2}])$ be the 2dimensional representation of U_{q^2} given by

$$\rho(E) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \ \rho(F) = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \ \rho(K) = \begin{pmatrix} q^2 & 0 \\ 0 & q^{-2} \end{pmatrix} \text{ and let } \rho(h) = \begin{pmatrix} \phi_0^0(h) & \phi_1^0(h) \\ \phi_0^1(h) & \phi_1^1(h) \end{pmatrix}$$
(115)

for all $h \in U_{q^2}$. This defines the matrix coefficients $\phi_0^0, \phi_1^0, \phi_0^1, \phi_1^1 \in \mathcal{O}_{q^2}(\mathrm{SL}_2)$ of ρ . A more popular notation is $\phi_0^0 = a, \phi_1^0 = b, \phi_0^1 = c, \phi_1^1 = d$. Then $\mathcal{L}_{g,n}(U_{q^2})$ is the $\mathbb{C}[q^{\pm 1/2}]$ -algebra generated by

$$\mathfrak{i}_{b_i}(\phi_t^s), \quad \mathfrak{i}_{a_i}(\phi_t^s), \quad \mathfrak{i}_{m_{g+j}}(\phi_t^s) \quad \text{with } 1 \leq i \leq g, \ 1 \leq j \leq n, \ 0 \leq s,t \leq 1$$

modulo the relations that can be found e.g. in [Fai20b, Lem. 5.1].¹⁴

The right action coad^r of U_{q^2} on $\mathcal{L}_{0,1}(U_{q^2})$ is easily computed: thanks to the module-algebra property, it suffices to determine the action of the generators $E, F, K \in U_{q^2}$ on the generators $\phi_t^s \in \mathcal{L}_{0,1}(U_{q^2})$. Since the embeddings \mathbf{i}_{b_i} , \mathbf{i}_{a_i} , $\mathbf{i}_{m_{g+j}} : \mathcal{L}_{0,1}(U_{q^2}) \to \mathcal{L}_{g,n}(U_{q^2})$ are morphisms of module-algebras, this also entirely determines the action of U_{q^2} on $\mathcal{L}_{q,n}(U_{q^2})$. For instance:

$$\begin{pmatrix} \operatorname{coad}^{r}(E)(\phi_{0}^{0}) & \operatorname{coad}^{r}(E)(\phi_{1}^{0}) \\ \operatorname{coad}^{r}(E)(\phi_{0}^{1}) & \operatorname{coad}^{r}(E)(\phi_{1}^{1}) \end{pmatrix} = - \begin{pmatrix} \phi_{0}^{0} & \phi_{1}^{0} \\ \phi_{0}^{1} & \phi_{1}^{1} \end{pmatrix} \rho(EK^{-1}) + \rho(E) \begin{pmatrix} \phi_{0}^{0} & \phi_{1}^{0} \\ \phi_{0}^{0} & \phi_{1}^{1} \end{pmatrix} \rho(K^{-1}) = \begin{pmatrix} q^{-2}\phi_{0}^{1} & q^{2}(\phi_{1}^{1} - \phi_{0}^{0}) \\ 0 & -q^{2}\phi_{0}^{1} \end{pmatrix} .$$

Recall that we denote by $\mathcal{L}_{g,n}^{\text{inv}}(U_{q^2})$ the subalgebra of invariant elements for this action.

Denote by $S_q^{\text{st}}(\Sigma)$ the Kauffman bracket *stated* skein algebra of a punctured bordered oriented surface Σ . It is the $\mathbb{C}[q^{\pm 1/2}]$ -module freely spanned by the isotopy classes of framed (unoriented, uncolored) tangles in $\Sigma \times [0, 1]$ whose endpoints in $\partial(\Sigma) \times [0, 1]$ are labelled by states + or -, modulo the Kauffman bracket skein relations and additional boundary relations as defined in [Lê18, CL22]. The only difference with these papers here is that we use the opposite product.¹⁵.

Remark 5.23. This particular definition of stated skein algebra is of course equivalent to the one given in §5.1.1 if one takes $H = U_{q^2}$ there. The states -, + have to be interpreted as the basis vectors (1,0), (0,1) of the representation ρ in (115). All strands can be colored by ρ because every U_{q^2} -module appears as a direct summand of some tensor power $\rho^{\otimes k}$, see e.g. [Kas95, §VII.7]. Finally the orientation of the strands can be forgotten because ρ is self-dual, modulo the choice of a non-standard pivotal element [Fai20b, §5] (*i.e.* we do not use g = K).

¹²We choose this ground ring because the stated skein algebra is defined over it [Lê18, CL22], as we will recall later. One can even take \mathbb{Z} instead of \mathbb{C} .

 $^{{}^{13}}U_{q^2}(\mathfrak{sl}_2)^{\circ}$ is generated by the matrix coefficients of the irreducible highest modules V_k^{\pm} defined e.g. in [Kas95, VI.3.5]. For $\mathcal{O}_{q^2}(\mathrm{SL}_2)$ we only take matrix coefficients of the type 1 modules V_k^{\pm} .

¹⁴The change of notations with [Fai20b] is $B(i) = \begin{pmatrix} \mathbf{i}_{b_i}(\phi_0^0) & \mathbf{i}_{b_i}(\phi_1^0) \\ \mathbf{i}_{b_i}(\phi_0^1) & \mathbf{i}_{b_i}(\phi_1^1) \end{pmatrix}$ and similarly for A(i) and M(g+j). ¹⁵In our convention, $L_1 * L_2$ means that put L_1 below L_2 in $\Sigma \times [0, 1]$.

Denote by $S_q(\Sigma)$ the usual Kauffman bracket skein algebra of Σ , still over the ground ring $\mathbb{C}[q^{\pm 1/2}]$. In [Fai20b, §5] we proved that there is a commutative diagram of $\mathbb{C}[q^{\pm 1/2}]$ -algebras

$$\mathcal{S}_{q}^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet}) \xrightarrow{\cong} \mathcal{L}_{g,n}(U_{q^{2}}) \tag{116}$$

$$\int_{\mathfrak{S}_{q}} \int_{\mathbb{C}_{g,n}^{\circ}} \int_{W}^{\cong} \mathcal{L}_{g,n}^{\mathrm{inv}}(U_{q^{2}})$$

where $S_q(\Sigma_{g,n}^{\circ})$ is sent to the subalgebra of links in $S_q^{\text{st}}(\Sigma_{g,n}^{\circ,\bullet})$. The isomorphism W is explicit: if γ is a simple closed curve in $S_q(\Sigma_{g,n}^{\circ})$ we have $W(\gamma) = \mathbf{i}_{\gamma}(q\text{Tr}_{\rho}) = -\mathbf{i}_{\gamma}(q^2\phi_0^0 + q^{-2}\phi_1^1)$, with $q\text{Tr}_{\rho}$ from Example 4.2 and $\mathbf{i}_{\gamma} : \mathcal{L}_{0,1}(U_{q^2}) \to \mathcal{L}_{g,n}(U_{q^2})$ can be computed through Proposition 5.12. We note that J. Korinman generalized diagram (116) to surfaces with arbitrary many boundary circles and arbitrary many punctures on each boundary circle [Kor23].

Now let ϵ be a root of unity, say of order 4p with $p \geq 2$ to follow the choices in [Fai20b, §7]. Being $\mathbb{C}[q^{\pm 1/2}]$ -modules, the algebras above can be specialized to $q = \epsilon$ (with an arbitrary choice of $\epsilon^{1/2}$) and we denote them by U_{ϵ^2} , $\mathcal{L}_{g,n}(U_{q^2})_{\epsilon^2}$, $\mathcal{L}_{g,n}^{\text{inv}}(U_{q^2})_{\epsilon^2}$ and $\mathcal{S}_{\epsilon}(\Sigma_{g,n}^{\circ})$; they are \mathbb{C} -algebras. We want to construct "derived representations" of $\mathcal{S}_{\epsilon}(\Sigma_{g,n}^{\circ})$. Note that Theorem 4.12 cannot be applied to $\mathcal{L}_{g,n}(U_{q^2})_{\epsilon^2}$ because this algebra is not of the form $\mathcal{L}_{g,n}(H)$. Instead the idea is to use a finite-dimensional quotient which has this form.

Recall that the small quantum group is the following $2p^3$ -dimensional quotient of U_{ϵ^2} :

$$\overline{U}_{\epsilon^2} = \overline{U}_{\epsilon^2}(\mathfrak{sl}_2) = U_{\epsilon}/\langle E^p, F^p, K^{2p} - 1 \rangle.$$

The Hopf algebra $\overline{U}_{\epsilon^2}$ is not quasitriangular; instead there is an *R*-matrix in the extension of $\overline{U}_{\epsilon^2}$ by a square root of *K* [FGST06a, §4.1]. Despite this small technical issue one can still define the algebra $\mathcal{L}_{g,n}(\overline{U}_{\epsilon^2})$: it is $(\overline{U}_{\epsilon^2})^{\otimes(2g+n)}$ as a \mathbb{C} -vector space and is endowed with the product of §4.1 [Fai19a, §6]. Let $\overline{\rho}: \overline{U}_{\epsilon^2} \to M_2(\mathbb{C})$ be the representation defined as ρ in (115) but with q replaced by ϵ and consider the associated matrix coefficients $\overline{\phi}_t^s$ with $0 \leq s, t \leq 1$. Then $\mathcal{L}_{g,n}(\overline{U}_{\epsilon^2})$ is generated by the elements $\mathbf{i}_{b_i}(\overline{\phi}_t^s)$, $\mathbf{i}_{a_i}(\overline{\phi}_t^s)$ and $\mathbf{i}_{m_{g+j}}(\overline{\phi}_t^s)$ with $1 \leq i \leq g, 1 \leq j \leq n$, $0 \leq s, t \leq 1$ modulo the same relations as $\mathcal{L}_{g,n}(U_q)_{\epsilon^2}$ together with the extra relations

$$\mathbf{i}_s(\overline{\phi}_1^0)^p = 0, \quad \mathbf{i}_s(\overline{\phi}_1^0)^p = 0, \quad \mathbf{i}_s(\overline{\phi}_1^1)^{2p} = 1$$

for all $s \in \{b_1, a_1, \ldots, b_g, a_g, m_{g+1}, \ldots, m_{g+n}\}$. As a result there are projections pr : $U_{\epsilon^2} \twoheadrightarrow \overline{U}_{\epsilon^2}$ and $\operatorname{pr}_{g,n} : \mathcal{L}_{g,n}(U_q)_{\epsilon^2} \twoheadrightarrow \mathcal{L}_{g,n}(\overline{U}_{\epsilon^2})$ which readily satisfy

$$\forall h \in U_{\epsilon^2}, \ \forall a \in \mathcal{L}_{g,n}(U_q)_{\epsilon^2}, \quad \mathrm{pr}_{g,n}\big(\mathrm{coad}^r(h)(a)\big) = \mathrm{coad}^r\big(\mathrm{pr}(h)\big)\big(\mathrm{pr}_{g,n}(a)\big).$$

It follows that $\operatorname{pr}_{g,n}$ restricts to a morphism $\mathcal{L}_{g,n}^{\operatorname{inv}}(U_q)_{\epsilon^2} \to \mathcal{L}_{g,n}^{\operatorname{inv}}(\overline{U}_{\epsilon^2})$, which as a side remark is no longer surjective. Composing with W in (116) specialized to ϵ , we obtain a morphism of \mathbb{C} -algebras

$$\overline{W}: \mathcal{S}_{\epsilon}(\Sigma_{g,n}^{\circ}) \xrightarrow{W_{\epsilon}} \mathcal{L}_{g,n}^{\mathrm{inv}}(U_q)_{\epsilon^2} \xrightarrow{\mathrm{pr}_{g,n}} \mathcal{L}_{g,n}^{\mathrm{inv}}(\overline{U}_{\epsilon^2})$$

and Theorem 4.12 gives:

Corollary 5.24. For any $\overline{U}_{\epsilon^2}$ -modules X, X_1, \ldots, X_n and all $m \ge 0$, there is a representation of the Kauffman bracket skein algebra $\mathcal{S}_{\epsilon}(\Sigma_{g,n}^{\circ})$ on $\operatorname{Ext}_{\overline{U}_{\epsilon^2}}^m (X, (\overline{U}_{\epsilon^2}^*)^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n)$.

This is less abstract than what it might look. Take the example of $\Sigma_{1,0}^{\circ}$. It is known that $\mathcal{S}_{\epsilon}(\Sigma_{1,0}^{\circ})$ is generated by the simple loops α, β in Fig. 2 [BP99]. We have $\overline{W}(\alpha) = \mathfrak{i}_a(\mathrm{qTr}_{\overline{\rho}})$ and $\overline{W}(\beta) = \mathfrak{i}_b(\mathrm{qTr}_{\overline{\rho}})$, where $\mathrm{qTr}_{\overline{\rho}} = -\epsilon^2 \overline{\phi}_0^0 - \epsilon^{-2} \overline{\phi}_1^1 \in \mathcal{L}_{0,1}^{\mathrm{inv}}(\overline{U}_{\epsilon^2})$. A straightforward computation

reveals that $\Phi_{0,1}(qTr_{\overline{\rho}}) = \Omega$ where $\Omega = (\epsilon^2 - \epsilon^{-2})FE + \epsilon^2 K + \epsilon^{-2}K^{-1} \in \mathcal{Z}(\overline{U}_{\epsilon^2})$ is the Casimir element. By Remark 4.11, the action on any $\psi \in \overline{U}_{\epsilon^2}^*$ is

$$\overline{W}(\alpha) \cdot \psi = \Omega \triangleright \psi, \quad \overline{W}(\beta) \cdot \psi = \psi \star \operatorname{qTr}_{\overline{\rho}}.$$

Endow $\overline{U}_{\epsilon^2}^*$ with the right coadjoint action (53). Then for any $P \in \text{mod}-\overline{U}_{\epsilon^2}$, the action of $L \in \mathcal{S}_{\epsilon}(\Sigma_{1,0}^{\circ})$ on $f \in \text{Hom}_{\overline{U}_{\epsilon^2}}(P, \overline{U}_{\epsilon^2}^*)$ is by acting on the target $\overline{U}_{\epsilon^2}^*$, *i.e.* $(L \cdot f)(x) = \overline{W}(L) \cdot f(x)$. Finally, for a projective resolution $0 \leftarrow X \leftarrow P_0 \leftarrow P_1 \leftarrow \dots$ in $\text{mod}-\overline{U}_{\epsilon^2}$, the action of L on a cohomology class $[f] \in \text{Ext}_{\overline{U}_{\epsilon^2}}^m(X, \overline{U}_{\epsilon^2}^*)$ represented by a cocycle $f \in \text{Hom}_{\overline{U}_{\epsilon^2}}(P_m, \overline{U}_{\epsilon^2}^*)$ is given by $L \cdot [f] = [L \cdot f]$. Projective resolutions of simple $\overline{U}_{\epsilon^2}$ -modules are given in [FGST06b, §3.1].

It is very likely that if X is the trivial module \mathbb{C} then Corollary 5.24 descends to a representation of $\mathcal{S}_{\epsilon}(\Sigma_{g,0})$. For m = 0, *i.e.* for the representation on $\operatorname{Hom}_{\overline{U}_{\epsilon^2}}(\mathbb{C}, (\overline{U}_{\epsilon^2})^{\otimes g})$, this is true and was proved in [Fai20b, §7].¹⁶

Remark 5.25. 1. In the forthcoming paper [BFR], building upon techniques in [BR24], we generalize some of the arguments above to any simple complex Lie algebra \mathfrak{g} , thus working with $U_q(\mathfrak{g})$ instead of $U_q(\mathfrak{sl}_2)$. More precisely, we show that there is an algebra surjection $\mathcal{L}_{g,n}(U_q(\mathfrak{g}))_{\zeta} \to \mathcal{L}_{g,n}(u_{\zeta}(\mathfrak{g}))$ which restricts to a morphism $\mathcal{L}_{g,n}^{inv}(U_q(\mathfrak{g}))_{\zeta} \to \mathcal{L}_{g,n}^{inv}(u_{\zeta}(\mathfrak{g}))$, where $\zeta \in \mathbb{C}$ is a root of a unity whose order has suitable arithmetic properties and $u_{\zeta}(\mathfrak{g})$ is a small quantum group, finite-dimensional quotient of $U_{\zeta}(\mathfrak{g})$.

2. Let $S_q(\Sigma, \mathfrak{g})$ be the skein algebra whose relations come from the Reshetikhin–Turaev tangle invariant for $U_q(\mathfrak{g})$. At least for $\mathfrak{g} = \mathfrak{sl}_N(\mathbb{C})$, there is a complete list of skein relations with coefficients in $\mathbb{C}[q^{\pm 1}]$ [Sik05, LS21], and hence a specialization $S_{\zeta}(\Sigma, \mathfrak{g})$ of $S_q(\Sigma, \mathfrak{g})$ at the root of unity ζ is possible. For $\Sigma = \Sigma_{g,n}^{\circ}$, the morphism $\mathcal{L}_{g,n}^{\mathrm{inv}}(U_q(\mathfrak{g}))_{\zeta} \to \mathcal{L}_{g,n}^{\mathrm{inv}}(u_{\zeta}(\mathfrak{g}))$ yields representations of $S_{\zeta}(\Sigma, \mathfrak{g})$ on Ext spaces over $u_{\zeta}(\mathfrak{g})$ as in Corollary 5.24.

A Alekseev morphism and representations of $\mathcal{L}_{g,n}(H)$

Let H be a quasitriangular Hopf algebra over a field k, with invertible antipode and R-matrix $R = R_{[1]} \otimes R_{[2]}$. In this appendix we provide a short construction of a morphism $\Phi'_{g,n} : \mathcal{L}_{g,n}(H) \to \operatorname{End}_{\Bbbk}(H^{\circ})^{\otimes g} \otimes H^{\otimes n}$ which is an adaptation of [Ale94] and was used in [Fai20a]. An important property, which was a key-fact for the construction of mapping class group representations in §5.2.2, is that $\Phi'_{g,n}$ is an isomorphism when H is finite-dimensional and factorizable. The morphism $\Phi'_{g,n}$ also yields representations of $\mathcal{L}_{g,n}(H)$ on $(H^{\circ})^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n$ for any H-modules X_i and we will see that they are isomorphic to those of §4.2.

Recall first that the *Heisenberg double* $\mathcal{H}(H^{\circ}) = H^{\circ} \# H$ is the vector space $H^{\circ} \otimes H$ endowed with the (associative) product

$$(\alpha \# a)(\beta \# b) = \alpha \star (a_{(1)} \triangleright \beta) \# a_{(2)}b$$

where \triangleright is the left coregular action (43) and \star is the product (40) which is dual to Δ . Its unit element is $\varepsilon \# 1_H$ where ε is the counit of H. It is customary to write # instead of \otimes because $\mathcal{H}(H^\circ)$ is a smash product [Mon93, §4.1.10]. In [BFR23, Lem. 3.6] it is proved that $\mathcal{H}(H^\circ)$ is a right H-module-algebra when endowed with the following action:

$$(\alpha \# a) \bullet h = S(h_{(2)}) \triangleright \alpha \triangleleft h_{(3)} \# S(h_{(1)})ah_{(4)}.$$

There is a well-known representation of $\mathcal{H}(H^{\circ})$ on H° given by

$$\forall (\alpha \# a) \in \mathcal{H}(H^{\circ}), \ \forall \psi \in H^{\circ}, \quad (\alpha \# a) \cdot \psi = \alpha \star (a \triangleright \psi).$$
(117)

¹⁶Actually in that paper we considered a slightly different representation which is a right representation, but the computations can be easily reproduced in the present setting.

We can make it H-equivariant (Def. 2.1) by letting

$$\forall \psi \in H^{\circ}, \ \forall h \in H, \quad \psi \bullet h = S(h_{(1)}) \triangleright \psi \triangleleft h_{(2)}.$$
(118)

Now recall the morphisms $i_b, i_a : \mathcal{L}_{0,1}(H) \to \mathcal{L}_{1,0}(H)$ from Def. 4.4 and consider the linear map

$$\Phi_{1,0} : \mathcal{L}_{1,0}(H) \to \mathcal{H}(H^{\circ})$$

$$\mathbf{i}_{b}(\varphi) \mapsto \left(R^{1}_{[2]}R^{2}_{[2]} \triangleright \varphi \triangleleft R^{3}_{[1]}R^{1}_{[1]}\right) \# R^{3}_{[2]}R^{2}_{[1]}$$

$$\mathbf{i}_{a}(\varphi) \mapsto \varepsilon \# \Phi_{0,1}(\varphi)$$
(119)

for all $\varphi \in \mathcal{L}_{0,1}(H)$, where $\Phi_{0,1} : \mathcal{L}_{0,1}(H) \to H$ is defined in (56) and \mathbb{R}^1 , \mathbb{R}^2 , \mathbb{R}^3 are three copies of the *R*-matrix. It is known that $\Phi_{1,0}$ is a morphism of algebras [Ale94] which is moreover a *H*-linear map $(\mathcal{L}_{1,0}(H), \operatorname{coad}^r) \to (\mathcal{H}(H^\circ), \bullet)$ [BFR23, Prop. 3.7]. As a result, by pullback of (117) and (118), we get a structure of *H*-equivariant $\mathcal{L}_{1,0}(H)$ -module on H° :

$$\mathbf{i}_{b}(\varphi) \cdot \psi = \left(R^{1}_{[2]}R^{2}_{[2]} \triangleright \varphi \triangleleft R^{3}_{[1]}R^{1}_{[1]}\right) \star \left(R^{3}_{[2]}R^{2}_{[1]} \triangleright \psi\right), \quad \mathbf{i}_{a}(\varphi) \cdot \psi = \Phi_{0,1}(\varphi) \triangleright \psi, \\
\psi \bullet h = S(h_{(1)}) \triangleright \psi \triangleleft h_{(2)}$$
(120)

for all $\varphi \in \mathcal{L}_{0,1}(H)$, $\psi \in H^{\circ}$ and $h \in H$. Introduce the following notations for the corresponding representation morphisms:

$$\rho_{1,0}: \mathcal{L}_{1,0}(H) \to \operatorname{End}_{\Bbbk}(H^{\circ}), \quad \rho_{1,0}(\beta \otimes \alpha)(\psi) = \mathfrak{i}_{b}(\beta)\mathfrak{i}_{a}(\alpha) \cdot \psi,$$
$$\mathsf{D}: H \to \operatorname{End}_{\Bbbk}(H^{\circ})^{\operatorname{op}}, \quad \mathsf{D}(h)(\psi) = S(h_{(1)}) \triangleright \psi \triangleleft h_{(2)}.$$

Let $\mathfrak{e}_1, \ldots, \mathfrak{e}_g : \mathcal{L}_{1,0}(H) \to \mathcal{L}_{g,n}(H)$ and $\mathfrak{e}_{g+1}, \ldots, \mathfrak{e}_{g+n} : \mathcal{L}_{0,1}(H) \to \mathcal{L}_{g,n}(H)$ be the tensor-wise embeddings in $\mathcal{L}_{g,n}(H) = \mathcal{L}_{1,0}(H)^{\widetilde{\otimes} g} \widetilde{\otimes} \mathcal{L}_{0,1}(H)^{\widetilde{\otimes} n}$, defined as in (31). With these notations:

Proposition A.1. 1. There is a morphism of algebras

$$\Phi'_{g,n}: \mathcal{L}_{g,n}(H) \to \operatorname{End}_{\Bbbk}(H^{\circ})^{\otimes g} \otimes H^{\otimes n}$$

(where the target is equipped with the usual tensor-wise product) defined by

$$\Phi_{g,n}'(\mathfrak{e}_{i}(x)) = \mathsf{D}(R_{1}) \otimes \ldots \otimes \mathsf{D}(R_{[1](i-1)}) \otimes \rho_{1,0}(\operatorname{coad}^{r}(R_{[2]})(x)) \otimes \operatorname{id}_{H^{\circ}}^{\otimes (g-i)} \otimes 1_{H}^{\otimes n}$$

for all $1 \leq i \leq g$, $x \in \mathcal{L}_{1,0}(H)$, with coad^r from (53), and

$$\Phi_{g,n}'(\mathfrak{e}_{g+j}(\varphi)) = \mathsf{D}(R_{1}) \otimes \ldots \otimes \mathsf{D}(R_{[1](g)}) \otimes S(R_{[1](g+1)}) \otimes \ldots \otimes S(R_{[1](g+j-1)}) \\ \otimes \Phi_{0,1}(\operatorname{coad}^{r}(R_{[2]})(\varphi)) \otimes 1_{H}^{\otimes n-j}$$

for all $1 \leq j \leq g$, $\varphi \in \mathcal{L}_{0,1}(H)$, with coad^r from (47) and $\Phi_{0,1}$ from (56). 2. If we endow $\operatorname{End}_{\Bbbk}(H^{\circ})^{\otimes g} \otimes H^{\otimes n}$ with the following structure of right H-module-algebra

$$\begin{pmatrix} E_1 \otimes \ldots \otimes E_g \otimes k_1 \otimes \ldots \otimes k_n \end{pmatrix} \bullet h = \left(\mathsf{D}(h_{(1)}) \circ E_1 \circ \mathsf{D}(S^{-1}(h_{(2g+2n)})) \right) \otimes \ldots \otimes \left(\mathsf{D}(h_{(g)}) \circ E_g \circ \mathsf{D}(S^{-1}(h_{(g+2n+1)})) \right) \\ \otimes S(h_{(g+1)}) k_1 h_{(g+2n)} \otimes \ldots \otimes S(h_{(g+n)}) k_n h_{(g+n+1)}$$

then $\Phi'_{g,n}$ becomes a morphism of H-module-algebras. 3. If the Hopf algebra H is finite-dimensional and factorizable, then $\Phi'_{g,n}$ is an isomorphism. *Proof.* 1. By Lemma 4.8 there is a *H*-equivariant representation of $\mathcal{L}_{0,1}(H)$ on *H*; the corresponding representation morphisms are

$$\rho_{0,1} : \mathcal{L}_{0,1}(H) \to \operatorname{End}_{\Bbbk}(H), \quad \rho_{0,1}(\varphi)(k) = \Phi_{0,1}(\varphi)k,$$

$$\mathsf{D}' : H \to \operatorname{End}_{\Bbbk}(H)^{\operatorname{op}}, \quad \mathsf{D}'(h)(k) = S(h)k.$$
(121)

Combining it with the *H*-equivariant representation (120) of $\mathcal{L}_{1,0}(H)$ on H° , item 2 in Corollary 3.2 gives a representation of $\mathcal{L}_{g,n}(H) = \mathcal{L}_{1,0}(H)^{\widetilde{\otimes} g} \widetilde{\otimes} \mathcal{L}_{0,1}(H)^{\widetilde{\otimes} n}$ on $(H^{\circ})^{\widetilde{\otimes} g} \widetilde{\otimes} H^{\widetilde{\otimes} n}$. By (35) the associated representation morphism $\rho_{g,n} : \mathcal{L}_{g,n}(H) \to \operatorname{End}_{\Bbbk}(H^{\circ})^{\otimes g} \otimes \operatorname{End}_{\Bbbk}(H)^{\otimes n}$ is given by

$$\rho_{g,n}(\mathfrak{e}_i(x)) = \mathsf{D}(R_{1}) \otimes \ldots \otimes \mathsf{D}(R_{[1](i-1)}) \otimes \rho_{1,0}(\operatorname{coad}(R_{[2]})(x)) \otimes \operatorname{id}_{H^\circ}^{\otimes (g-i)} \otimes 1_H^{\otimes i}$$

for all $1 \leq i \leq g$ and

$$\rho_{g,n}(\mathfrak{e}_{g+j}(\varphi)) = \mathsf{D}(R_{1}) \otimes \ldots \otimes \mathsf{D}(R_{[1](g)}) \otimes \mathsf{D}'(S(R_{[1](g+1)})) \otimes \ldots \otimes \mathsf{D}'(S(R_{[1](g+j-1)}))$$
$$\otimes \rho_{0,1}(\operatorname{coad}^r(R_{[2]})(\varphi)) \otimes 1_H^{\otimes n-j}$$

for all $1 \leq j \leq n$. Now note that because of the particular form of $\rho_{0,1}$ and D' (based on left multiplication in H), $\rho_{g,n}$ actually takes values in $\operatorname{End}_{\Bbbk}(H^{\circ})^{\otimes g} \otimes \operatorname{End}(H_H)^{\otimes n}$, where $\operatorname{End}(H_H)$ is the subspace of linear maps $f: H \to H$ such that f(hk) = f(h)k for all $h, k \in H$. Using the isomorphism of algebras $\operatorname{End}(H_H) \xrightarrow{\sim} H$ given by $f \mapsto f(1_H)$, we define

$$\Phi'_{g,n} : \mathcal{L}_{g,n}(H) \xrightarrow{\rho_{g,n}} \operatorname{End}_{\Bbbk}(H^{\circ})^{\otimes g} \otimes \operatorname{End}(H_{H})^{\otimes n} \xrightarrow{\sim} \operatorname{End}_{\Bbbk}(H^{\circ})^{\otimes g} \otimes H^{\otimes n}$$
(122)

which is a morphism of algebras as a composition of such. The claimed formulas for $\Phi'_{g,n}$ immediately follow.

2. Note from (121) that if $f \in \text{End}(H_H)$ then for all $h, h' \in H$ we have $(\mathsf{D}'(h) \circ f \circ \mathsf{D}'(S^{-1}(h')))(1_H) = S(h)f(h') = S(h)f(1_H)h'$. Now the result follows from (36) and (122).

3. This is [Fai20a, Prop. 3.5]. Since the notations used there look quite different, here are a few details. By assumption $\Phi_{0,1}$ is an isomorphism and thus $\rho_{0,1}$ is injective. Recall that $\rho_{1,0}$ is the composition $\mathcal{L}_{1,0}(H) \xrightarrow{\Phi_{1,0}} \mathcal{H}(H^*) \xrightarrow{(117)} \operatorname{End}_{\Bbbk}(H^*)$. It is not difficult to see that $\Phi_{1,0}$ is an isomorphism under our assumptions on H [Fai19a, Th. 4.8]. Moreover, it is a remarkable fact that the representation of $\mathcal{H}(H^\circ)$ on H° is faithful [Mon93, Lem. 9.4.2] for all H. Hence, when H is finite-dimensional, the arrow $\mathcal{H}(H^*) \to \operatorname{End}_{\Bbbk}(H^*)$ is an isomorphism by equality of dimensions. As a result the composition $\rho_{1,0}$ is an isomorphism. It follows that $\rho_{g,n}$ is injective by Lemma 3.3 and thus $\Phi'_{g,n}$ is injective as well because of the factorization (122). To conclude, it suffices to note that the source and target of $\Phi'_{g,n}$ have the same dimension, namely $\dim(H)^{2g+n}$.

Remark A.2. 1. In [Fai20a] we used an isomorphism which has $\mathcal{H}(H^{\circ})^{\otimes g} \otimes H^{\otimes n}$ for target. When H is finite-dimensional it is known that $\mathcal{H}(H^{\circ}) \cong \operatorname{End}_{\Bbbk}(H^{\circ})$ [Mon93, Lem. 9.4.2] and the morphism $\Phi'_{g,n}$ above coincide exactly with the one in [Fai20a] through this identification.

2. Our definition of $\Phi'_{g,n}$ is especially suited for finite-dimensional H. See [BFR23, §5] for the definition of a morphism $\Phi_{g,n}$ which is more suited for the infinite-dimensional case, because in this case its target is smaller. Also see [BFR23, Rmk. 5.5] for a comparison on the two different conventions to combine $\Phi_{0,1}$ and $\Phi_{1,0}$ in order to define $\Phi'_{g,n}$ and $\Phi_{g,n}$.

Recall the braided external product $\widetilde{\boxtimes}$ of *H*-equivariant modules from Cor.3.2. Let H_{old}° (resp. H_{new}°) be the representation of $\mathcal{L}_{1,0}(H)$ defined in (120) (resp. in Lemma 4.9) and let X_1, \ldots, X_n be left *H*-modules. On the one hand, the morphism $\Phi'_{g,n}$ gives a representation of $\mathcal{L}_{g,n}(H)$ on the vector space $(H^{\circ})^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n$; by the proof of Proposition A.1, this representation is exactly $(H_{\text{old}}^{\circ})^{\widetilde{\boxtimes} g} \widetilde{\boxtimes} X_1 \widetilde{\boxtimes} \ldots \widetilde{\boxtimes} X_n$. On the other hand, the representation of $\mathcal{L}_{g,n}(H)$ constructed in §4.2 is $(H_{\text{new}}^{\circ})^{\widetilde{\boxtimes} g} \widetilde{\boxtimes} X_1 \widetilde{\boxtimes} \ldots \widetilde{\boxtimes} X_n$ by definition. The linear map

$$F: H^{\circ}_{\text{new}} \to H^{\circ}_{\text{old}}, \quad \psi \mapsto R_{[2]} \triangleright \psi \triangleleft R_{[1]}$$
(123)

is easily seen to be an isomorphism of H-equivariant $\mathcal{L}_{1,0}(H)$ -modules. As a result

$$F^{\otimes g} \otimes \operatorname{id}_{X_1} \otimes \ldots \otimes \operatorname{id}_{X_n} : (H^{\circ}_{\operatorname{new}})^{\widetilde{\boxtimes} g} \widetilde{\boxtimes} X_1 \widetilde{\boxtimes} \ldots \widetilde{\boxtimes} X_n \to (H^{\circ}_{\operatorname{old}})^{\widetilde{\boxtimes} g} \widetilde{\boxtimes} X_1 \widetilde{\boxtimes} \ldots \widetilde{\boxtimes} X_n$$
(124)

is an isomorphism of *H*-equivariant $\mathcal{L}_{g,n}(H)$ -modules.

B Proofs of results on knot maps

We recall that in these results we assume that $H = U_q(\mathfrak{g})$ over $\mathbb{C}(q)$ or that H is finitedimensional. The ground field of H is denoted by \Bbbk .

B.1 Proof of Prop. 5.4

Assume first that $H = U_q(\mathfrak{g})$ over $\Bbbk = \mathbb{C}(q)$. Recall that $\mathcal{L}_{0,1}(\mathfrak{g})$ is the vector space $\mathcal{O}_q(G)$ endowed with the product (46) and the right action coad^r from (47). Denote by P_+ the set of dominant weights of \mathfrak{g} and for each $\mu \in P_+$ let V_{μ} be the irreducible $U_q(\mathfrak{g})$ -module with highest weight μ . The following lemma is well-known and generalized to $\mathcal{L}_{0,n}(\mathfrak{g})$ in [BR19, Prop. 6.21].

Lemma B.1. $\mathcal{L}_{0,1}^{\text{inv}}(\mathfrak{g}) = \text{Span}_{\mathbb{C}(q)} \{ q \text{Tr}_{\mu} \mid \mu \in P_+ \}, \text{ where } q \text{Tr}_{\mu} = q \text{Tr}_{V_{\mu}} \text{ is defined in Ex. 4.2.}$

Proof. We have the decomposition of right $U_q(\mathfrak{g})$ -modules $\mathcal{L}_{0,1}(\mathfrak{g}) \cong \bigoplus_{\mu \in P_+} (C(V_\mu), \operatorname{coad}^r)$ where $C(V_\mu)$ is the subspace of matrix coefficients of V_μ . There is an isomorphism of $\mathbb{C}(q)$ -vector spaces

$$\operatorname{End}_{\mathbb{C}(q)}(V_{\mu}) \to C(V_{\mu}), \quad f \mapsto \sum_{i} w^{i}_{\mu} (2g \cdot f(w_{\mu,i}))$$

where $(w_{\mu,i})$ is a basis of V_{μ} with dual basis (w_{μ}^{i}) and g is the pivotal element. It is easy to see that it establishes an isomorphism between $\operatorname{End}_{U_{q}(\mathfrak{g})}(V_{\mu})$ and $C(V_{\mu})^{\operatorname{inv}}$. Since $\operatorname{End}_{U_{q}(\mathfrak{g})}(V_{\mu}) \cong$ $\mathbb{C}(q)$ we find that, up to scalar, the unique element in $C(V_{\mu})^{\operatorname{inv}}$ is $\sum_{i} w_{\mu}^{i} (?g \cdot w_{\mu,i}) = \operatorname{qTr}_{\mu}$. \Box

Now we are ready to prove Prop. 5.4 for $H = U_q(\mathfrak{g})$. For a knot $K \subset \Sigma_{g,n}^{\circ} \times [0,1]$, choose any based knot **K** which reduces to K when considered as an usual knot. This is done by choosing any point in K, isotoping it to $\partial(\Sigma_{g,n}^{\circ,\bullet}) \times [0,1]$ and considering it as a basepoint. If **K** is positively oriented, we get from (viii) in Fig. 1



which is clearly independent of the particular choice of \mathbf{K} . The argument is similar if \mathbf{K} is negatively oriented.

Now assume that H is finite-dimensional. The strategy of proof in this case is a bit different because an element invariant for \cdot in $X^* \otimes X$ does not necessarily comes from an H-endomorphism of X. Let K be a knot in $\sum_{g,n}^{\circ} \times [0,1]$ and choose two arbitrary points $p_1, p_2 \in K$. Using isotopy, we assume that these two points are very close to $\partial \sum_{g,n}^{\circ, \bullet} \times [0,1]$. Then by another small isotopy, we can put p_1 (resp. p_2) on the boundary and consider it as the basepoint of a based knot \mathbf{K}_1 (resp. \mathbf{K}_2). By well-chosen isotopies, we can always arrange the following configuration:

$$K = \underbrace{p_1 \qquad p_2}_{p_2} \qquad \mathbf{K}_1 = \underbrace{\mathbf{K}_2}_{p_2} = \underbrace{\mathbf{K}_2}_{p$$

The dashed lines schematize the knot K outside of this small neighborhood of the boundary. Then Proposition 5.4 will clearly follow from the following equality:

$$\forall \varphi \in \mathcal{L}_{0,1}^{\mathrm{inv}}(H), \quad \mathfrak{j}_{\mathbf{K}_1}(\varphi) = \mathfrak{j}_{\mathbf{K}_2}(\varphi)$$

To prove it, assume first that \mathbf{K}_1 is positively oriented; then so is \mathbf{K}_2 . We have

where (h_i) is a basis of H with dual basis (h^i) , the third equality uses (viii) in Fig. 1 and the last equality is obtained by dinaturality as the first claim in Lemma 5.2. Note using (50) and (66) that for all $x, y \in H$,

$$\langle \varphi(?gh_i) \otimes h^i, x \otimes y \rangle = \varphi(xgy) = \varphi(yS^2(x)g) = \varphi(ygx) = \langle h^i \otimes \varphi(?gh_i), x \otimes y \rangle.$$

Hence

Now if \mathbf{K}_1 is negatively oriented then so is \mathbf{K}_2 . Let \mathbf{K}_1^{-1} and \mathbf{K}_2^{-1} be the based knots with opposite orientations (then positively oriented). By item 2 in Lemma 5.3 and the previous computation,

$$\mathfrak{j}_{\mathbf{K}_1}(\varphi) = \mathfrak{j}_{\mathbf{K}_1^{-1}} \big(S_{\mathcal{L}_{0,1}}(\varphi) \big) = \mathfrak{j}_{\mathbf{K}_2^{-1}} \big(S_{\mathcal{L}_{0,1}}(\varphi) \big) = \mathfrak{j}_{\mathbf{K}_2}(\varphi)$$

for all $\varphi \in \mathcal{L}_{0,1}^{\mathrm{inv}}(H)$. We used that $S_{\mathcal{L}_{0,1}}$ is a *H*-linear endomorphism of $\mathcal{L}_{0,1}(H)$ and thus $S_{\mathcal{L}_{0,1}}(\varphi) \in \mathcal{L}_{0,1}^{\mathrm{inv}}(H)$.

B.2 Proof of Prop. 5.12

Let $\gamma \in \pi_1(\Sigma_{g,n}^{\circ})$. Fix a particular drawing D_{γ} of γ on the surface (67). We allow only drawings which do not contain self-intersections and such a drawing is made of oriented caps (\cap) , cups (\cup) , two lines ending to the base point and half-circles in handles (the punctured half-disks in (67) being also considered as handles). Let:

- $\vec{N}_{\cap}(D_{\gamma})$ be the number of caps oriented from left to right in the picture D_{γ} .
- $\stackrel{\leftarrow}{N}_{\cap}(D_{\gamma})$ be the number of caps oriented from right to left in the picture D_{γ} .
- $\vec{N}_{\cup}(D_{\gamma})$ be the number of half-circles oriented from left to right in handles and cups oriented from left to right in the picture D_{γ} .
- $\stackrel{\leftarrow}{N}_{\cup}(D_{\gamma})$ be the number of half-circles oriented from right to left in handles and cups oriented from right to left in the picture D_{γ} .

Here is an example in $\Sigma_{1,1}^{\circ,\bullet}$, with a particular drawing D_{γ} of the loop $\gamma = b_1 a_1^{-1} = b_1 a_1^{-1} m_2 m_2^{-1}$

Then we have $\overrightarrow{N}_{\cap}(D_{\gamma}) = 3$, $\overleftarrow{N}_{\cap}(D_{\gamma}) = 1$, $\overrightarrow{N}_{\cup}(D_{\gamma}) = 3$ and $\overleftarrow{N}_{\cup}(D_{\gamma}) = 2$.

We have an important relation:

$$\overleftarrow{N}_{\cup}(D_{\gamma}) + \overrightarrow{N}_{\cup}(D_{\gamma}) = \overleftarrow{N}_{\cap}(D_{\gamma}) + \overrightarrow{N}_{\cap}(D_{\gamma}) + 1.$$
(126)

Indeed, when we follow D_{γ} along its orientation starting from the basepoint, we necessarily first meet a cup or a half-circle in a handle. After we necessarily meet a cap, and then again a cup or a half-circle *etc*, which gives a sequence [CUP, cap], [CUP, cap], ..., [CUP, cap], CUP where CUP means cup or half-circle. The sequence necessarily ends by a CUP, in order to return to the basepoint. Thus any CUP is paired with a cap as indicated by square brackets in the sequence above, except for the last one, whence the "+1" term in the formula.

Now define

$$\overleftarrow{N}(\gamma) = \overleftarrow{N}_{\cup}(D_{\gamma}) - \overleftarrow{N}_{\cap}(D_{\gamma}), \qquad \overrightarrow{N}(\gamma) = \overrightarrow{N}_{\cup}(D_{\gamma}) - \overrightarrow{N}_{\cap}(D_{\gamma})$$
(127)

where D_{γ} is some drawing of γ . These quantities are independent of the particular drawing, whence the notation. This is because the only possible variations in a drawing are zig-zags made of cups-caps and cc^{-1} where c is a half-circle in a handle, as in (125).

Definition B.2. Recall item 3 in Definition 5.1 about positive orientation of based knots (which applies in particular to simple loops). For any simple loop $\gamma \in \pi_1(\Sigma_{q,n}^\circ)$ we let

$$N(\gamma) = \begin{cases} \overleftarrow{N}(\gamma) & \text{if } \gamma \text{ is positively oriented,} \\ \overleftarrow{-N}(\gamma^{-1}) & \text{if } \gamma \text{ is negatively oriented.} \end{cases}$$

Lemma B.3. 1. If γ is a negatively oriented simple loop then $N(\gamma) = \overleftarrow{N}(\gamma) - 1$. 2. For any simple loop γ we have $N(\gamma) = -N(\gamma^{-1})$.

Proof. 1. By (126) we have $\overleftarrow{N}(\gamma) + \overrightarrow{N}(\gamma) = 1$. Note also that $\overleftarrow{N}(\gamma^{-1}) = \overrightarrow{N}(\gamma)$ for all γ , which follows immediately from the definitions. Hence $N(\gamma) = -\overrightarrow{N}(\gamma^{-1}) = -\overrightarrow{N}(\gamma) = \overleftarrow{N}(\gamma) - 1$. 2. Immediate by definition.

We now give a recursion formula for $N(\gamma)$, based on a factorization $\gamma = \gamma' \gamma''$ where γ', γ'' are *non-intersecting* simple loops. Such a factorization is obtained by bringing a cap of γ near the basepoint without creating self-intersections and then turning the cap into a concatenation of two loops. Near the basepoint, there are 6 possible configurations for a positively oriented simple loop:



Of course this is schematic and the dashed parts of the loops γ' and γ'' never intersect. Similarly there are 6 possible configurations for a negatively oriented simple loop:



With these notations, define

$$P(\gamma',\gamma'') = \begin{cases} -1 & \text{in cases } 4^+, 1^-, 5^-, \\ 0 & \text{in cases } 2^+, 3^+, 6^+, 2^-, 3^-, 6^-, \\ 1 & \text{in cases } 1^+, 5^+, 4^-. \end{cases}$$
(128)

Lemma B.4. For any simple loop $\gamma \in \pi_1(\Sigma_{q,n}^\circ)$ we have

$$N(\gamma) = N(\gamma') + N(\gamma'') + P(\gamma', \gamma'').$$

Proof. Note that in any case we have $\overset{\leftarrow}{N_{\cup}}(\gamma) = \overset{\leftarrow}{N_{\cup}}(\gamma') + \overset{\leftarrow}{N_{\cup}}(\gamma'')$. The computation of $\overset{\leftarrow}{N_{\cap}}(\gamma)$ is a case-by-case analysis. Here are two examples, the others are left to the reader. In case 4^+ it holds $\overset{\leftarrow}{N_{\cap}}(\gamma) = \overset{\leftarrow}{N_{\cap}}(\gamma') + \overset{\leftarrow}{N_{\cap}}(\gamma'') + 1$; hence since γ' and γ'' are positively oriented we get $N(\gamma) = \overset{\leftarrow}{N}(\gamma) = \overset{\leftarrow}{N}(\gamma') + \overset{\leftarrow}{N}(\gamma'') - 1 = N(\gamma') + N(\gamma'') - 1$ by Def. B.2 and (127). In case 2^+ it holds $\overset{\leftarrow}{N_{\cap}}(\gamma) = \overset{\leftarrow}{N_{\cap}}(\gamma') + \overset{\leftarrow}{N_{\cap}}(\gamma'') + 1$; hence since γ' is negatively oriented and γ'' is positively oriented we get $N(\gamma) = \overset{\leftarrow}{N}(\gamma) = \overset{\leftarrow}{N}(\gamma') + \overset{\leftarrow}{N}(\gamma'') - 1 = N(\gamma') + N(\gamma'')$ by Def. B.2, (127) and item 1 in Lemma B.3. Also the result for negatively oriented loops can be deduced from oriented loops. For instance if $\gamma = \gamma'\gamma''$ is in case 1^- then $\gamma^{-1} = \gamma''^{-1}\gamma'^{-1}$ is in case 1^+ and thus $N(\gamma) = -N(\gamma^{-1}) = -N(\gamma''^{-1}) - N(\gamma'^{-1}) - 1 = N(\gamma') + N(\gamma'') - 1$ by item 2 in Lemma B.3.

Now we give a recursion formula for the loop maps $i_{\gamma} : \mathcal{L}_{0,1}(H) \to \mathcal{L}_{g,n}(H)$, which are particular cases of the based knot maps defined in (82).

Proposition B.5. Let $\gamma \in \pi_1(\Sigma_{g,n}^\circ)$ be a simple loop and take a decomposition $\gamma = \gamma' \gamma''$ where γ', γ'' are two non-intersecting simple loops. Then

$$\mathbf{i}_{\gamma}(\varphi) = \varphi_{(1)} \left(v^{P(\gamma',\gamma'')} \right) \mathbf{i}_{\gamma'}(\varphi_{(2)}) \, \mathbf{i}_{\gamma''}(\varphi_{(3)})$$

for all $\varphi \in \mathcal{L}_{0,1}(H)$, with $P(\gamma', \gamma'') \in \{-1, 0, 1\}$ from (128) and $\varphi_{(1)} \otimes \varphi_{(2)} \otimes \varphi_{(3)}$ is iteration of the coproduct (42).

Proof. The proof is a case-by-case verification for each possible configuration of γ' and γ'' listed above. In each case, one first compute j_{γ} in terms of $j_{\gamma'}$, $j_{\gamma''}$ thanks to the stated skein relations and gets the desired result by applying the morphism holst. We give details in two cases, the other cases are left to the reader.

In case 2^+ , γ is positively oriented so \mathfrak{j}_{γ} is defined by (72) and we have:



where the third equality uses naturality of the twist and (vii) in Fig. 1. Note that γ' is negatively oriented while γ'' is positively oriented, so $j_{\gamma'}$ is defined by (73) while $j_{\gamma''}$ is defined by (72) and the last term is equal to $j_{\gamma'}(_X\phi_{x_i}^{\sigma})j_{\gamma''}(_X\phi_x^{x^i})$ by definition of the product in $\mathcal{S}_H^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$. Applying holst to this equality gives the result, because $\Delta(_X\phi_x^{\sigma}) = _X\phi_{x_i}^{\sigma} \otimes _X\phi_x^{x^i}$ by (41). In case 4⁻, γ is negatively oriented so j_{γ} is defined by (73) and we have:



For the third equality the extra twist is compensated by the ribbon element v inserted in x^i . Note that γ' , γ'' are negatively oriented so $\mathbf{j}_{\gamma'}$, $\mathbf{j}_{\gamma''}$ are defined by (73) and the last term is equal to $\mathbf{j}_{\gamma'}(_X \phi_{x_i}^{\sigma})\mathbf{j}_{\gamma''}(_X \phi_x^{x^i(v\cdot?)})$ by definition of the product in $\mathcal{S}_H^{\mathrm{st}}(\Sigma_{g,n}^{\circ,\bullet})$. Applying holst to this equality and using (41) and (44) we get $\mathbf{i}_{\gamma}(\varphi) = \mathbf{i}_{\gamma'}(\varphi_{(1)})\mathbf{i}_{\gamma''}(\varphi_{(2)} \triangleleft v) = \varphi_{(2)}(v)\mathbf{i}_{\gamma'}(\varphi_{(1)})\mathbf{i}_{\gamma''}(\varphi_{(3)})$ for all $\varphi \in \mathcal{L}_{0,1}(H)$. Since v is central it holds $\varphi_{(2)}(v)\varphi_{(1)} = \varphi_{(1)}(v)\varphi_{(2)}$ by (42) and we are done.

Proof of Proposition 5.12. Recall that $\pi_1(\Sigma_{g,n}^{\circ})$ is a free group on the generators a_i , b_i , m_{g+j} with $1 \leq i \leq g$ and $1 \leq j \leq n$. Assume that the word $\gamma_1 \ldots \gamma_l$ in these generators and their inverses which represents γ is reduced, *i.e.* does not contain factors of the form ss^{-1} . We will prove the theorem by induction on the length l of the word $\gamma_1 \ldots \gamma_l$. For length 1 it is tautological. Assume that the formula is true for all simple loops which can be represented by a word of length < l for some l. Take a factorization $\gamma = \gamma' \gamma''$ where γ' , γ'' are non-intersecting simple loops. Then $\gamma' = \gamma_1 \ldots \gamma_k$ and $\gamma'' = \gamma_{k+1} \ldots \gamma_l$ for some 1 < k < l. Using Proposition B.5 and the induction hypothesis we get

$$\mathbf{i}_{\gamma}(\varphi) = \varphi_{(1)} \left(v^{P(\gamma',\gamma'')} \right) \mathbf{i}_{\gamma'}(\varphi_{(2)}) \mathbf{i}_{\gamma''}(\varphi_{(3)}) = \varphi_{(1)} \left(v^{P(\gamma',\gamma'')} \right) \varphi_{(2)} \left(v^{N(\gamma')} \right) \mathbf{i}_{\gamma_1}(\varphi_{(3)}) \dots \mathbf{i}_{\gamma_k}(\varphi_{(k+2)}) \varphi_{(k+3)} \left(v^{N(\gamma'')} \right) \mathbf{i}_{\gamma_{k+1}}(\varphi_{(k+4)}) \dots \mathbf{i}_{\gamma_l}(\varphi_{(l+3)}).$$

Since v is central, (42) implies $\varphi_{(2)}(v)\varphi_{(1)} = \varphi_{(1)}(v)\varphi_{(2)}$. Hence

$$\mathbf{i}_{\gamma}(\varphi) = \varphi_{(1)} \left(v^{N(\gamma') + N(\gamma'') + P(\gamma', \gamma'')} \right) \mathbf{i}_{\gamma_1}(\varphi_{(2)}) \dots \mathbf{i}_{\gamma_k}(\varphi_{(k+1)}) \mathbf{i}_{\gamma_{k+1}}(\varphi_{(k+2)}) \dots \mathbf{i}_{\gamma_l}(\varphi_{(l+1)})$$

and Lemma B.4 finishes the proof.

C On the quantum moment map for $\mathcal{L}_{q,n}(H)$

Let H be a ribbon Hopf algebra over a field k. The main result of this appendix is Prop. C.1, which is a key ingredient in the proof of Theorem 5.8. We also recall that when H is factorizable $\mathcal{L}_{g,n}(H)$ has a quantum moment map $\mu_{g,n}$ and then Prop. C.1 can be reformulated in terms of $\mu_{g,n}$ (Cor. C.2). The QMM $\mu_{g,n}$ is a "fusion" of many copies of $\mu_{1,0}$ and $\mu_{0,1}$; such a fusion operation for QMMs is discussed in general in Remark C.3.

We will need the following "braided coproduct" on H

$$\underline{\Delta}: \quad H \to H \otimes H \\ h \mapsto \operatorname{ad}^{r}(R_{[1]})(h_{(1)}) \otimes S(R_{[2]})h_{(2)} = R_{[1]}^{1}h_{(1)}R_{[1]}^{2} \otimes S(R_{[2]}^{2})R_{[2]}^{1}h_{(2)}$$
(129)

where ad^{r} is the right adjoint action of H on itself (15) and the equality follows from (19) and (23). Note using (20) and (18) that it can also be written as

$$\underline{\Delta}(h) = h_{(2)} R_{[1]} \otimes \operatorname{ad}^{r}(R_{[2]})(h_{(1)}).$$
(130)

It is known [Maj92, Th. 7.4.2]¹⁷ that the linear map $\underline{\Delta}$ is a morphism of *H*-module-algebras $(H, \mathrm{ad}^r) \to (H, \mathrm{ad}^r) \widetilde{\otimes} (H, \mathrm{ad}^r)$ and is coassociative: $(\underline{\Delta} \otimes \mathrm{id}_H) \circ \underline{\Delta} = (\mathrm{id}_H \otimes \underline{\Delta}) \circ \underline{\Delta}$. For completeness, we note that the antipode $\widetilde{S} : (H, \mathrm{ad}^r) \to (H, \mathrm{ad}^r)$ is given by

$$\widetilde{S}(h) = S\left(\operatorname{ad}^{r}(R_{[1]})(h)\right)R_{[2]} = S\left(R_{[2]}u^{-1}hS(R_{[1]})\right)$$

where the second equality follows from the properties of *R*-matrices and *u* is the Drinfeld element recalled in (37). Then $\underline{H} = (H, \cdot, 1_H, \underline{\Delta}, \varepsilon_H, \underline{S})$ is a Hopf algebra object in Mod-*H*, [Maj92, Ex. 9.4.9].

Recall that $\mathcal{L}_{0,1}(H)$ is a Hopf algebra object in Mod-H with product (46), coproduct (41) and antipode (48). Consider the map $\Phi_{0,1} : \mathcal{L}_{0,1}(H) \to \underline{H}$ defined by $\varphi \mapsto \varphi(R^1_{[1]}R^2_{[2]})R^1_{[2]}R^2_{[1]}$ and already used in §4.2. Then $\Phi_{0,1}$ is a morphism of Hopf algebras in Mod-H, meaning that

$$\Phi_{0,1} \circ \operatorname{coad}^{r}(h) = \operatorname{ad}^{r}(h) \circ \Phi_{0,1}, \qquad \Phi_{0,1}(\varphi\psi) = \Phi_{0,1}(\varphi)\Phi_{0,1}(\psi),
(\Phi_{0,1} \otimes \Phi_{0,1}) \circ \Delta_{H^{\circ}} = \underline{\Delta} \circ \Phi_{0,1}, \qquad \Phi_{0,1} \circ S_{\mathcal{L}_{0,1}} = \underline{S} \circ \Phi_{0,1}.$$
(131)

All these equalities follow from the properties of *R*-matrices recalled in §3. A variant of the map $\Phi_{0,1}$ appears in [Maj92, eq. (9.48)].

Recall the boundary loop $\partial_{g,n} = \partial(\Sigma_{g,n}^{\circ})$ defined in (68), which is positively oriented. We have seen in Prop. 5.5 and Cor. 5.6 that for any positively oriented simple loop $\gamma \in \pi_1(\Sigma_{g,n}^{\circ})$ we have a *H*-module-algebra morphism $\mathbf{i}_{\gamma} : \mathcal{L}_{0,1}(H) \to \mathcal{L}_{g,n}(H)$. In particular, by Proposition 5.12, the morphism $\mathbf{i}_{\partial_{1,0}}$ is given by

$$\mathfrak{i}_{\partial_{1,0}}(\varphi) = \varphi_{(1)}(v^2)\mathfrak{i}_b(\varphi_{(2)})\mathfrak{i}_a\big(S_{\mathcal{L}_{0,1}}(\varphi_{(3)})\big)\mathfrak{i}_b\big(S_{\mathcal{L}_{0,1}}(\varphi_{(4)})\big)\mathfrak{i}_a(\varphi_{(5)})$$

and the morphism $i_{\partial q,n}$ is given by

$$\mathbf{i}_{\partial g,n}(\varphi) = \mathbf{i}_{\partial_{1,0}}(\varphi_{(1)}) \widetilde{\otimes} \dots \widetilde{\otimes} \mathbf{i}_{\partial_{1,0}}(\varphi_{(g)}) \widetilde{\otimes} \varphi_{(g+1)} \widetilde{\otimes} \dots \widetilde{\otimes} \varphi_{(g+n)}.$$
(132)

where we use that $\mathcal{L}_{g,n}(H) = \mathcal{L}_{1,0}(H)^{\tilde{\otimes} g} \tilde{\otimes} \mathcal{L}_{0,1}(H)^{\tilde{\otimes} n}$. When H does not satisfies the assumptions of §5.1.3 we cannot use the based knot maps; then formula (132) must be taken as a definition and remains a morphism of H-module-algebras [BFR23, Proof of Lem. 7.12]. From [BFR23, Lem. 7.13] we have the following exchange relation

$$\forall \varphi, \psi \in \mathcal{L}_{0,1}(H), \quad \mathfrak{i}_{\partial g,n} \left(R^1_{[2]} \triangleright \varphi \triangleleft R^2_{[1]} \right) \mathfrak{i}_s \left(\psi \triangleleft R^2_{[2]} R^1_{[1]} \right) = \mathfrak{i}_s \left(R^2_{[2]} R^1_{[1]} \triangleright \psi \right) \mathfrak{i}_{\partial g,n} \left(R^1_{[2]} \triangleright \varphi \triangleleft R^2_{[1]} \right).$$

where $s \in \{b_1, a_1, \ldots, b_g, a_g, m_{g+1}, \ldots, m_{g+n}\}$ is any generator of $\pi_1(\Sigma_{g,n}^\circ)$ and R^1 , R^2 are two copies of the *R*-matrix. The properties of the *R*-matrix (see §3) allow us to rewrite this as

$$\forall \varphi, \psi \in \mathcal{L}_{0,1}(H), \quad \mathfrak{i}_{\partial g,n}(\varphi)\mathfrak{i}_s(\psi) = \mathfrak{i}_s \left(\operatorname{coad}^r \left(S^{-1}(R^1_{[2]}R^2_{[1]}) \right)(\psi) \right) \mathfrak{i}_{\partial g,n} \left(R^2_{[2]} \triangleright \varphi \triangleleft R^1_{[1]} \right). \tag{133}$$

Proposition C.1. Recall the *H*-equivariant $\mathcal{L}_{g,n}(H)$ -module structure on $(H^{\circ})^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n$ from Prop. 4.10 and in particular its right *H*-action •. We have

$$\mathfrak{i}_{\partial g,n}(\varphi) \cdot w = w \bullet S^{-1}(\Phi_{0,1}(\varphi))$$

for all $\varphi \in \mathcal{L}_{0,1}(H)$ and $w \in (H^{\circ})^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n$, with $\Phi_{0,1}$ from (56).

Proof. • *Case* (g, n) = (0, 1): We have $i_{\partial_{0,1}}(\varphi) = \varphi$ and the result is just the definition of the representation of $\mathcal{L}_{0,1}(H)$ on a *H*-module in Lemma 4.8.

¹⁷The only difference with Majid's book is that here we use the right adjoint action of H on itself instead of the left one, which explains the difference in the formulas.

• Case (g, n) = (1, 0): Here it is helpful to use stated skein algebras. To do so we assume that H satisfies the assumptions of §5.1.3; the result is true for any ribbon H but algebraic computations without stated skein algebras are involved. We have $\mathcal{L}_{1,0}(H) \cong \mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{1,0}^{\circ,\bullet})$ as algebras and $H^{\circ} = \mathcal{L}_{0,1}(H) \cong \mathcal{S}_{H}^{\mathrm{st}}(\Sigma_{0,1}^{\circ,\bullet})$ as vector spaces. Under these identifications, the representation of $\mathcal{L}_{1,0}(H)$ on H° in Lemma 4.9 becomes



where G is any ribbon graph and the two strands in the handles of $\Sigma_{1,0}^{\circ,\bullet}$ actually represent a bunch of parallel strands. Because of skein relations it is sufficient to prove this formula for elements of the form $\mathbf{j}_b(X_1\phi_{x_1}^{\sigma_1})\mathbf{j}_a(X_2\phi_{x_2}^{\sigma_2})$ in $\mathcal{S}_H^{\mathrm{st}}(\Sigma_{1,0}^{\circ,\bullet})$, *i.e.* when G is the identity graph (4 parallel strands). For such elements it is easy to use the characterization of the isomorphism $\mathcal{L}_{1,0}(H) \cong \mathcal{S}_H^{\mathrm{st}}(\Sigma_{1,0}^{\circ,\bullet})$ from (81) in order to identify the diagrams above with the formulas in Lemma 4.9; details are left to the reader. In particular:



which means that $\mathbf{i}_{\partial_{1,0}}(\varphi) \cdot \varepsilon = \varphi(1_H)\varepsilon$ for all $\varphi \in \mathcal{L}_{0,1}(H)$, since the empty diagram corresponds to the counit $\varepsilon = {}_{\mathbb{k}} \phi_1^{\mathrm{id}}$. The result now easily follows from Lemma 4.9, (133) and the definition of $\Phi_{0,1}$:

$$\begin{split} \mathbf{i}_{\partial_{1,0}}(\varphi) \cdot \psi &= \mathbf{i}_{\partial_{g,n}}(\varphi) \mathbf{i}_{b}(\psi) \cdot \varepsilon = \mathbf{i}_{b} \Big(\operatorname{coad}^{r} \left(S^{-1}(R^{1}_{[2]}R^{2}_{[1]}) \right)(\psi) \Big) \mathbf{i}_{\partial_{1,0}} \left(R^{2}_{[2]} \triangleright \varphi \triangleleft R^{1}_{[1]} \right) \cdot \varepsilon \\ &= \mathbf{i}_{b} \Big(\operatorname{coad}^{r} \left(S^{-1}(\Phi_{0,1}(\varphi)) \right)(\psi) \Big) \cdot \varepsilon = \operatorname{coad}^{r} \left(S^{-1}(\Phi_{0,1}(\varphi)) \right)(\psi) \end{split}$$

for all $\psi \in H^{\circ}$. We have $\bullet = \operatorname{coad}^{r}$ in this case by (61), so we are done.

• General case: It is divided in three steps: first prove the result for $\mathcal{L}_{g,0}(H)$ by induction on g thanks to the decomposition $\mathcal{L}_{g+1,0}(H) = \mathcal{L}_{g,0}(H) \otimes \mathcal{L}_{1,0}(H)$, then do the same for $\mathcal{L}_{0,n}(H)$ by induction on n, and finally deduce the result in general from the decomposition $\mathcal{L}_{g,n}(H) = \mathcal{L}_{g,0}(H) \otimes \mathcal{L}_{0,n}(H)$. We give details only for the first step, the others are based on completely similar computations. Assume that the Proposition is true for $\mathcal{L}_{g,0}(H)$. We use the notation $\underline{\Delta}(h) = h_{(\underline{1})} \otimes h_{(\underline{2})}$ for the coproduct (129). Then for all $\varphi \in \mathcal{L}_{0,1}(H)$ and $w \boxtimes \psi \in (H^{\circ})^{\boxtimes g} \boxtimes H^{\circ}$,

$$\begin{split} & \mathfrak{i}_{\partial_{g+1,0}}(\varphi) \cdot \left(w \,\widetilde{\boxtimes} \,\psi\right) \stackrel{(132)}{=} \left(\mathfrak{i}_{\partial_{g,0}}(\varphi_{(1)}) \,\widetilde{\otimes} \,\mathfrak{i}_{\partial_{1,0}}(\varphi_{(2)})\right) \cdot \left(w \,\widetilde{\boxtimes} \,\psi\right) \\ & \stackrel{(29)}{=} \left(\mathfrak{i}_{\partial_{g,0}}(\varphi_{(1)}) \cdot \left(w \bullet R_{[1]}\right)\right) \,\widetilde{\boxtimes} \left(\mathfrak{i}_{\partial_{1,0}}\left(\operatorname{coad}^{r}(R_{[2]})(\varphi_{(2)})\right) \cdot \psi\right) \\ & = \left(w \bullet R_{[1]}S^{-1}\left(\Phi_{0,1}(\varphi_{(1)})\right)\right) \,\widetilde{\boxtimes} \left(\operatorname{coad}^{r}\left[S^{-1}\left(\Phi_{0,1}\left(\operatorname{coad}^{r}(R_{[2]})(\varphi_{(2)})\right)\right)\right](\psi)\right) \\ & \stackrel{(131)}{=} \left(w \bullet R_{[1]}S^{-1}\left(\Phi_{0,1}(\varphi_{(1)})\right)\right) \,\widetilde{\boxtimes} \left(\operatorname{coad}^{r}\left[S^{-1}\left(\operatorname{ad}^{r}(R_{[2]})\left(\Phi_{0,1}(\varphi_{(2)})\right)\right)\right](\psi)\right) \\ & \stackrel{(131)}{=} \left(w \bullet R_{[1]}S^{-1}\left(\Phi_{0,1}(\varphi_{(1)})\right)\right) \,\widetilde{\boxtimes} \left(\operatorname{coad}^{r}\left[S^{-1}\left(\operatorname{ad}^{r}(R_{[2]})\left(\Phi_{0,1}(\varphi_{(2)})\right)\right)\right](\psi)\right) \end{split}$$

$$\stackrel{(130)}{=} \left(w \bullet R_{[1]}^1 S^{-1} \left(\Phi_{0,1}(\varphi)_{(2)} R_{[1]}^2 \right) \right) \widetilde{\boxtimes} \left(\operatorname{coad}^r \left[S^{-1} \left(\operatorname{ad}^r (R_{[2]}^2 R_{[2]}^1) \left(\Phi_{0,1}(\varphi)_{(1)} \right) \right) \right] (\psi) \right)$$

$$\stackrel{(24)}{=} \left(w \bullet S^{-1} \left(\Phi_{0,1}(\varphi)_{(2)} \right) \right) \widetilde{\boxtimes} \left(\operatorname{coad}^r \left[S^{-1} \left(\Phi_{0,1}(\varphi)_{(1)} \right) \right] (\psi) \right) \stackrel{(61)}{=} \left(w \widetilde{\boxtimes} \psi \right) \bullet S^{-1} \left(\Phi_{0,1}(\varphi) \right)$$

where the unnumbered equality uses the induction hypothesis and the case (g, n) = (1, 0) established previously.

Now assume that H is factorizable, i.e. that $\Phi_{0,1}$ is an isomorphism, and define

$$\mu_{g,n}: H \xrightarrow{\Phi_{0,1}^{-1}} \mathcal{L}_{0,1}(H) \xrightarrow{i_{\partial_{g,n}}} \mathcal{L}_{g,n}(H).$$

In particular $\mu_{0,1} = \Phi_{0,1}^{-1}$. The morphism of algebras $\mu_{g,n}$ is a quantum moment map (QMM, see Def. 2.5). This property was obtained in [Jor14] for $H = U_q(\mathfrak{gl}_n)$ and generalized to any H in [BFR23, §7.2]; the proof of the QMM axiom uses the equality (133). Due to (132) and the third equality in (131) we have the factorization

$$\forall h \in H, \quad \mu_{g,n}(h) = \mu_{1,0}(h_{(\underline{1})}) \widetilde{\otimes} \dots \widetilde{\otimes} \mu_{1,0}(h_{(\underline{g})}) \widetilde{\otimes} \mu_{0,1}(h_{(\underline{g+1})}) \widetilde{\otimes} \dots \widetilde{\otimes} \mu_{0,1}(h_{(\underline{g+n})})$$
(135)

where the underlined subscripts are a Sweedler notation for the braided coproduct (129).

The existence of the QMM $\mu_{g,n}$ implies that $any \mathcal{L}_{g,n}(H)$ -module (W, \cdot) has automatically a *H*-equivariant structure, defined by $w \cdot_{\mu g,n} h = \mu_{g,n}(S(h)) \cdot w$ (Lemma 2.6). This applies in particular to the modules $(H^{\circ})^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n$ constructed in §4.2, which are already endowed with the *ad hoc H*-equivariant structure • defined in (61). These two *H*-equivariant structures agree:

Corollary C.2. For all $h \in H$ and $w \in (H^{\circ})^{\otimes g} \otimes X_1 \otimes \ldots \otimes X_n$, we have $w \bullet_{\mu_{g,n}} h = w \bullet h$.

Proof. By Proposition C.1,
$$w \bullet_{\mu g,n} h = \mu_{g,n} (S(h)) \cdot w = \mathfrak{i}_{\partial g,n} (\Phi_{0,1}^{-1}(S(h))) \cdot w = w \bullet h.$$

Remark C.3. The formula (135) shows that the QMM $\mu_{g,n}$ is a "fusion" of copies of the QMMs $\mu_{1,0}$ and $\mu_{0,1}$. Such a fusion operation is general, as we now explain. Here we only assume that H is quasitriangular. Let A_1 and A_2 be right H-module-algebras and assume that A_i has a QMM μ_i . Then the algebra morphism

$$\mu_{12}: H \xrightarrow{\Delta} H \stackrel{\sim}{\otimes} H \xrightarrow{\mu_1 \otimes \mu_2} A_1 \stackrel{\sim}{\otimes} A_2$$

is a QMM for $A_1 \otimes A_2$, where we use the coproduct (129). Let us prove the QMM axiom (12). It is more convenient to check it separately on elements of the form $1_{A_1} \otimes a_2$ and $a_1 \otimes 1_{A_2}$. For $1_{A_1} \otimes a_2$, note first from (129) and (14) that

$$\mu_{12}(h) = \mu_1(h_{(1)}) \bullet R_{[1]} \widetilde{\otimes} \mu_2 \big(S(R_{[2]}) h_{(2)} \big)$$
(136)

for all $h \in H$. Hence we have

$$\begin{aligned} &(1_{A_{1}} \otimes a_{2})\mu_{12}(h) = \mu_{1}(h_{(1)}) \bullet R_{[1]}^{1}R_{[1]}^{2} \otimes (a_{2} \bullet R_{[2]}^{2})\mu_{2}\left(S(R_{[2]}^{1})h_{(2)}\right) & \text{by (136), (27)} \\ &= \mu_{1}(h_{(1)}) \bullet R_{[1]}^{1}R_{[1]}^{2} \otimes \mu_{2}\left(S(R_{[2]}^{1})_{(1)}h_{(2)}\right)\left(a_{2} \bullet R_{[2]}^{2}S(R_{[2]}^{1})_{(2)}h_{(3)}\right) & \text{by (12)} \\ &= \mu_{1}(h_{(1)}) \bullet R_{[1]}^{1}R_{[1]}^{3}R_{[1]}^{2} \otimes \mu_{2}\left(S(R_{[2]}^{1})h_{(2)}\right)\left(a_{2} \bullet R_{[2]}^{2}S(R_{[2]}^{3})h_{(3)}\right) & \text{by (20)} \\ &= \mu_{1}(h_{(1)}) \bullet R_{[1]}^{1} \otimes \mu_{2}\left(S(R_{[2]}^{1})h_{(2)}\right)\left(a_{2} \bullet h_{(3)}\right) & \text{by (24)} \\ &= \mu_{12}(h_{(1)})\left(1_{A_{1}} \otimes (a_{2} \bullet h_{(2)})\right) & \text{by (136), (27), (6), (25)} \\ &= \mu_{12}(h_{(1)})\left((1_{A_{1}} \otimes a_{2}) \bullet h_{(2)})\right) & \text{by (28), (6).} \end{aligned}$$

The proof for $a_1 \otimes 1_{A_2}$ is similar and left to the reader.

Recall that if M_i is an A_i -module for i = 1, 2 then it is automatically a H-equivariant module with the right action \bullet_{μ_i} (Lemma 2.6) and we can form the $(A_1 \otimes A_2)$ -module $M_1 \otimes M_2$ as in Corollary 3.2. Now $M_1 \otimes M_2$ has two H-equivariant structures. The first is the diagonal action (30), given by $(v_1 \otimes v_2) \bullet h = (v_1 \bullet_{\mu_1} h_{(1)}) \otimes (v_2 \bullet_{\mu_2} h_{(2)}) = \mu_1(S(h_{(1)}) \cdot v_1 \otimes \mu_2(S(h_{(2)})) \cdot v_2$. The second comes directly from the QMM μ_{12} and is thus given by $(v_1 \otimes v_2) \bullet_{\mu_{12}} h = \mu_{12}(S(h)) \cdot (v_1 \otimes v_2)$. These two H-actions are equal:

$$\mu_{12}(S(h)) \cdot (v_1 \,\widetilde{\boxtimes} \, v_2) = \left[\mu_1 \left(S(h_{(1)}) R_{[1]} \right) \widetilde{\otimes} \, \mu_2(S(h_{(2)})) \bullet R_{[2]} \right] \cdot (v_1 \,\widetilde{\boxtimes} \, v_2)$$
 by (130) and (14)

$$= \mu_1 \left(S(h_{(1)}) R_{[1]}^1 \right) \cdot \left(v_1 \bullet_{\mu_1} R_{[1]}^2 \right) \widetilde{\boxtimes} \left(\mu_2(S(h_{(2)})) \bullet R_{[2]}^1 R_{[2]}^2 \right) \cdot v_2$$
 by (29)

$$= \mu_1 \left(S(h_{(1)}) R_{[1]}^1 S(R_{[1]}^2) \right) \cdot v_1 \,\widetilde{\boxtimes} \left(\mu_2(S(h_{(2)})) \bullet R_{[2]}^1 R_{[2]}^2 \right) \cdot v_2$$
 by (17)

$$= \mu_1(S(h_{(1)})) \cdot v_1 \,\widetilde{\boxtimes} \, \mu_2(S(h_{(2)})) \cdot v_2$$
 by (22).

This can be seen as a generalization of Corollary C.2, which in Proposition C.1 is proven from the building blocks $\mu_{0,1}$ and $\mu_{1,0}$.

References

- [Ale94] A. Yu. Alekseev, Integrability in the Hamiltonian Chern-Simons theory, Alg. i Anal. 6 (2) (1994) 53-66.
- [AGS95] A. Y. Alekseev, H. Grosse, V. Schomerus, Combinatorial quantization of the Hamiltonian Chern-Simons theory I, Comm. Math. Phys. 172 (1995), 317–358.
- [AS96a] A. Yu. Alekseev, V. Schomerus, Representation theory of Chern-Simons observables, Duke Math. J. 85 (1996), 447–510.
- [BR19] S. Baseilhac, P. Roche, Unrestricted Quantum Moduli Algebras, I: The Case of Punctured Spheres, SIGMA 18, 025, 78 pages (2022).
- [BR24] S. Baseilhac, P. Roche, Unrestricted Quantum Moduli Algebras, II: Noetherianity and simple fraction rings at roots of 1, SIGMA 20, 047, 70 pages (2024).
- [BFR23] S. Baseilhac, M. Faitg, P. Roche, Unrestricted Quantum Moduli Algebras, III: Surfaces or arbitrary genus and skein algebras, arXiv:2302.00396 (2023).
- [BFR] S. Baseilhac, M. Faitg, P. Roche, On the structure and representations of quantum graph algebras at roots of unity, in preparation.
- [BR95] E. Buffenoir, P. Roche, Two dimensional lattice gauge theory based on a quantum group, Comm. Math. Phys. 170 (1995), 669–698.
- [BR96] E. Buffenoir, P. Roche, Link invariants and combinatorial quantization of hamiltonian Chern Simons theory, Comm. Math. Phys. 181 (1996), 331–365.
- [BP99] D. Bullock, J.H. Przytycki, Multiplicative structure of Kauffman bracket skein modules quantizations, Proc. Amer. Math. Soc 128(3), 923–931 (1999).
- [BFK98a] D. Bullock, C. Frohman, J. Kania-Bartoszynska, Topological Interpretations of Lattice Gauge Field Theory, Comm. Math. Phys. 198 (1998), 47–81.
- [CP94] V. Chari, A. Pressley, A Guide To Quantum Groups, Cambridge Univ. Press (1994).

- [CGP14] F. Costantino, N. Geer, B. Patureau-Mirand, Quantum invariants of 3-manifolds via link surgery presentations and non-semi-simple categories, J. Topology7(4), 1005–1053 (2014).
- [CL22] F. Costantino, T. T. Q. Lê, Stated skein algebras of surfaces, J. Eur. Math. Soc. 24 (2022), 4063–4142.
- [DGGPR22] M. De Renzi, A.M. Gainutdinov, N. Geer, B. Patureau-Mirand, I. Runkel, 3-Dimensional TQFTs from non-semisimple modular categories, Selecta Math. New Ser. 28, article no. 42 (2022).
- [Fai19a] M. Faitg, Modular group representations in combinatorial quantization with nonsemisimple Hopf algebras, SIGMA 15 (2019), 077, 39 pages.
- [Fai19b] M. Faitg, Mapping class groups, skein algebras and combinatorial quantization, PhD thesis, Université de Montpellier, arXiv:1910.04110.
- [Fai20a] M. Faitg, Projective representations of mapping class groups in combinatorial quantization, Comm. Math. Phys. 377 (2020), 161–198.
- [Fai20b] M. Faitg, Holonomy and (stated) skein algebras in combinatorial quantization, Quantum Topol. (2024), published online first. DOI 10.4171/QT/176
- [FM12] B. Farb, D. Margalit, A primer on mapping class groups, Oxford University Press, 2012.
- [FGST06a] B.L. Feigin, A.M. Gainutdinov, A.M. Semikhatov, I.Yu. Tipunin, Modular Group Representations and Fusion in Logarithmic Conformal Field Theories and in the Quantum Group Center, Comm. Math. Phys. 265 (2006), 47–93.
- [FGST06b] B.L. Feigin, A.M. Gainutdinov, A.M. Semikhatov, I.Yu. Tipunin, Kazhdan-Lusztig correspondence for the representation category of the triplet W-algebra in logarithmic CFT, Theor. Math. Phys. 148, 1210–1235 (2006)
- [GKP13] N. Geer, J. Kujawa, B. Patureau-Mirand, Ambidextrous objects and trace functions for nonsemisimple categories, Proc. Amer. Math. Soc. 141(9), 2963–2978 (2013).
- [Jor14] D. Jordan, Quantized multiplicative quiver varieties, Adv. Math. 250 (2014), 420–466.
- [Kas95] C. Kassel, Quantum Groups, Graduate Texts in Mathematics 155, Springer, 1995.
- [Kor23] J. Korinman, Finite presentations for stated skein algebras and lattice gauge field theory, Algebr. Geom. Topol. 23:3, 1249–1302 (2023).
- [Lê18] T. T. Q. Lê, Triangular decomposition of skein algebras, Quantum Topol. 9 (2018), 591–632.
- [LS21] T. T. Q. Lê, A. S. Sikora, Stated SL(n)-skein modules and algebras, arXiv:2201.00045.
- [LMSS18] S. Lentner, S.N. Mierach, C. Schweigert, Y. Sommerh äuser, Hochschild cohomology and the modular group, J. Algebra 507, 400–420 (2018).
- [LMSS23] S. Lentner, S. N. Mierach, C. Schweigert, Y. Sommerhäuser, Hochschild Cohomology, Modular Tensor Categories, and Mapping Class Groups I, Springer Briefs in Mathematical Physics 44 (2023).

- [Lu93] J.-H. Lu, Moment maps at the quantum level, Comm. Math. Phys. 157(2) (1993) 389– 404.
- [Lyu95] V. Lyubashenko, Invariants of 3-manifolds and projective representations of mapping class groups via quantum groups at roots of unity, Commun. Math. Phys. 172, 467–516 (1995).
- [ML75] S. Mac Lane, *Homology*, Classics in Mathematics, Springer, 1975.
- [Maj92] S. Majid, Foundations of Quantum Group Theory, Cambridge University Press (1995).
- [Mon93] S. Montgomery, Hopf algebras and their actions on rings, CBMS Regional Conference Series in Mathematics, Vol. 82, Amer. Math. Soc., Providence, RI, 1993.
- [Rad11] D.E. Radford, *Hopf algebras*, Series on Knots and Everything **49**, World Scientific, 2011.
- [RS88] N. Reshetikhin, M.A. Semenov-Tian-Shansky, Quantum R-matrices and factorization problems, J. Geom. Phys. 5(4), 533–550 (1988).
- [RT90] N. Yu. Reshetikhin, V. G. Turaev, Ribbon Graphs and Their Invariants Derived from Quantum Groups, Comm. Math. Phys 127 (1990), 1–26.
- [RT91] N. Reshetikhin, V. G. Turaev, Invariants of 3-manifolds via link polynomials and quantum groups, Invent. Math. 103, 547–597 (1991).
- [Sik05] A. S. Sikora, Skein theory for SU(n)-quantum invariants, Algebr. Geom. Topol. 5 (2005), 865–897.
- [SW21a] C. Schweigert, L. Woike, The Hochschild complex of a finite tensor category, Algebr. Geom. Topol. 21, 3689–3734 (2021).
- [SW21b] C. Schweigert, L. Woike, Homotopy coherent mapping class group actions and excision for Hochschild complexes of modular categories, Adv. Math. 386, 107814 (2021).
- [Tur94] V. G. Turaev, Quantum invariants of knots and 3-manifolds, De Gruyter Studies in Mathematics 18, 1994.
- [VV10] M. Varagnolo, E. Vasserot, Double affine Hecke algebras at roots of unity, Representation Theory 14 (2010), 510–600.
- [VY20] C. Voigt, R. Yuncken, Complex Semisimple Quantum Groups and Representation Theory, Lecture Notes in Mathematics 2264, Springer, 2020.
- [Waj83] B. Wajnryb, A simple presentation for the mapping class group of an orientable surface, Israel J. Math. 45, 157–174 (1983).