

Energy needed to propel a tiny spacecraft to Proxima Centauri, and, an unstated assumption in Einstein's 1905 paper

C. J. Umrigar^{a1} and Tyler A. Anderson^{b1}

¹*Laboratory of Atomic and Solid State Physics,
Cornell University, Ithaca, NY 14853, USA.*

The Breakthrough Starshot project aims to send a tiny 2 gram spacecraft to Proxima Centauri propelled by a light sail and powerful Earth-based lasers. We provide two derivations of the laser energy required to propel the spacecraft and give the reader the opportunity to decide which one is correct before providing the answer. In the second part of this paper we point out that one of the formulae in Einstein's amazing 1905 paper is correct only in certain limits, but Einstein fails to mention that. This has caused some confusion in the Breakthrough Starshot literature.

^a CyrusUmrigar@cornell.edu

^b taa65@cornell.edu

I. INTRODUCTION

The prospects for space travel capture the human imagination and are the subject of much scientific and engineering endeavor, as well as creative science fiction. The two Voyager spacecraft, launched in 1977, are the fastest human-made objects currently travelling in interstellar space at approximately 62,000 km/hour or $6 \times 10^{-5}c$, where c is the speed of light in vacuum. In Table I we show the distances of some well-known astronomical objects and the time it would take the Voyager spacecraft to get there. If one limits oneself to missions that can occur within a human lifetime, it is apparent that while the entire solar system is accessible using current technology, the closest galaxy is not accessible even taking into account any reasonable projections of improvements to technology. (One may be tempted to think that because of relativistic time-dilation, one could travel arbitrarily far in a human lifetime, but in fact the energy requirements to approach the speed of light are enormous.) Travel to the closest star, Proxima Centauri, is in-between – it takes too long at Voyager speeds, but would take only 21 years or 8.5 years of Earth-frame time if one could have spacecraft with a speed of $0.2 c$ or $0.5 c$ respectively.

TABLE I. Distances to various astronomical objects in light travel and Voyager travel units.

Object	Distance	Time on Voyager
Moon	1.3 light seconds	6 hours
Sun	8 light minutes	93 days
Pluto	5.3 light hours	10 years
Closest star (Proxima Centauri)	4.24 light years	71,000 years
Closest galaxy (Canis Major Dwarf)	25,000 light years	420,000,000 years

The outline for the rest of the paper is as follows. In Section II we dispel any notion that advances in chemical rocket technology could get us there. In Section III we point out that photon rockets are not a viable solution because there is no known technology that can convert a sufficiently large fraction of rest mass into photons. In Section IV we discuss the Breakthrough Starshot project, which aims to propel a tiny spacecraft weighing about 2 grams using the pressure exerted by laser light on a light sail. Major technical problems would have to be overcome for this project to succeed. Instead, here we focus on the basic question of how much laser energy is needed because that is a trickier question than first meets the eye. We provide two derivations, which give two different expressions for the required energy that differ greatly at relativistic speeds. We encourage the reader to ponder which is the correct result and why, before reading the explanation given at the end of the section. In Section V we derive the expression for the energy of light reflected from a moving perfect mirror and find that the expression derived by Einstein in his famous 1905 paper is correct only in limiting cases. Next, we discuss some confusion in the Breakthrough Starshot literature related to this expression.

II. CHEMICAL ROCKETS ARE NOT THE ANSWER

The first question that may occur to the reader is whether improvements to conventional chemical rocket technology could propel spacecraft to a good fraction of the speed of light. The answer is a resounding no!

Chemical rockets eject hot gas, obtained by burning fuel, in one direction to propel the remainder of the rocket in the opposite direction. The rocket equation relates the mass of the entire rocket M (most of which is comprised of the fuel), to the mass of the payload m (the part of the rocket that travels to the final destination), the velocity of the exhaust gases v_{exh} , and the desired eventual speed of the payload v . A typical value for v_{exh} is $3 \text{ km/s} = 10^{-5} c$. Using $v = 0.5 c$, the rocket equation gives

$$M = m e^{\frac{v}{v_{\text{exh}}}} = m e^{\frac{0.5c}{10^{-5}c}} = m e^{50000} \approx m 10^{21715} \quad (\text{for } v = 0.5c). \quad (1)$$

This is truly an astronomically large mass, M , even for a tiny payload! The reason is that early in the flight fuel is being burned to accelerate not just the payload but also the rest of the fuel. In fact, the Voyager spacecrafts achieved even their rather modest speeds not by just using chemical rocket propulsion, but by also using gravity assists from Jupiter, Saturn and Uranus.

III. PHOTON ROCKETS ARE NOT THE ANSWER

Since M/m is exponential in v/v_{exh} , an enormous reduction in the mass, M , can be achieved by increasing v_{exh} . Let's carry this to the logical extreme, as has been proposed before¹, and say that instead of emitting exhaust gases, the rocket emits a collimated beam of photons. Let M_0 be the initial rest mass of the rocket, m_0 be the rest mass of the payload and E_{photon} be the energy of the emitted photons. Then, from energy and momentum conservation we have

$$M_0 c^2 = \gamma m_0 c^2 + E_{\text{photon}} \quad (\text{Energy cons.}) \quad (2)$$

$$\vec{p}_{\text{photon}} = -\vec{p}_{\text{rocket}} \quad (\text{Momentum cons.}) \quad (3)$$

$$\text{so } p_{\text{photon}} = p_{\text{rocket}} = \gamma m_0 v \equiv \gamma m_0 c \beta \quad (4)$$

Since the energy of photons is c times their momentum, $E_{\text{photon}} = \gamma m_0 c^2 \beta$. Plugging this back into the energy conservation equation, we have

$$M_0 c^2 = \gamma m_0 c^2 + \gamma m_0 c^2 \beta \quad (5)$$

$$M_0 = \gamma(1 + \beta)m_0 = \sqrt{\frac{1 + \beta}{1 - \beta}}m_0 \quad (6)$$

$$\rightarrow \begin{cases} (1 + \beta)m_0, & \text{when } \beta \rightarrow 0 (\gamma \rightarrow 1) \\ 2\gamma m_0, & \text{when } \beta \rightarrow 1 (\gamma \gg 1) \\ \sqrt{3}m_0 = 1.73m_0, & \text{when } \beta \rightarrow 1/2 \end{cases} \quad (7)$$

This is an enormous improvement compared to a chemical rocket, but is still not feasible with current technology for the following reason. We have assumed that mass can be converted into a collimated beam of photons with 100% efficiency but there is no way to do this, aside from matter-antimatter annihilation. There is no known way to make and store this quantity of antimatter. Nuclear reactors are not the answer – a fission reactor converts at most only $\sim 0.15\%$ of its rest mass into heat, and, further, the efficiency for converting heat into laser light is small.

IV. LIGHT SAILS AND THE BREAKTHROUGH STARSHOT PROJECT

Another idea is to leave the energy source that propels the spacecraft on Earth, or, in Earth orbit. This can be achieved by using the pressure of laser light shining on a light sail to propel the spacecraft. The goal of the Breakthrough Starshot project² is to propel a tiny spacecraft equipped with a light sail, weighing only about 2 grams, with an intense light beam from a phased array of lasers to something like $0.2c$. This project has \$100 million in seed funding from science philanthropist Yuri Milner, and many distinguished scientists and engineers are associated with the project². Needless to say, some very tough technical problems need to be overcome:

1. Deploying a sail that is sufficiently light, with a sufficiently high area and a high reflectivity. It is helpful to have a reflectivity close to one, but it is essential that the absorptivity is very close to zero so that the sail does not vaporize. Further the sail must maintain its shape and orientation while it is being propelled.
2. Making the camera and the communication devices really tiny, so that the total rest mass of the spacecraft is just 2 grams.
3. Having powerful enough lasers with a sufficiently small angular spread.

Here, we do not concern ourselves with these technical problems or the likelihood that they will ever be surmounted. Instead we address the basic question of how much laser light energy is needed to propel the spacecraft to velocity v .

We use standard relativistic notation, namely,

$$\beta = \frac{v}{c}, \quad (8)$$

$$\gamma = \frac{1}{\sqrt{1 - \beta^2}}. \quad (9)$$

We now give 2 derivations that give different expressions for the energy needed and then give the reader the opportunity to think about which is right and relevant, which is wrong or not relevant, and why.

A. Derivation 1

We find the energy, E_I , of the laser light needed to accelerate the spaceship with rest mass m_0 from an initial velocity β_i to a final velocity β_f . Denote the energy of the reflected light by E_R .

$$\gamma_i m_0 c^2 + E_I = \gamma_f m_0 c^2 + E_R \quad (\text{energy cons.}) \quad (10)$$

$$\gamma_i m_0 c \beta_i + \frac{E_I}{c} = \gamma_f m_0 c \beta_f - \frac{E_R}{c} \quad (\text{momentum cons.}) \quad (11)$$

Adding Eq. 10 to c times Eq. 11, we eliminate E_R and get

$$\begin{aligned} \gamma_i(1 + \beta_i)m_0c^2 + 2E_I &= \gamma_f(1 + \beta_f)m_0c^2 \\ \text{i.e., } E_I &= \frac{1}{2}(\gamma_f(1 + \beta_f) - \gamma_i(1 + \beta_i))m_0c^2 \\ &= \boxed{\frac{1}{2} \left(\sqrt{\frac{1 + \beta_f}{1 - \beta_f}} - \sqrt{\frac{1 + \beta_i}{1 - \beta_i}} \right) m_0c^2} \end{aligned} \quad (12)$$

If the initial velocity, $\beta_i = 0$, then

$$E_I = \frac{1}{2} \left(\sqrt{\frac{1 + \beta_f}{1 - \beta_f}} - 1 \right) m_0c^2 \rightarrow \begin{cases} \frac{\beta_f}{2} m_0c^2, & \text{when } \beta_f \rightarrow 0 \\ \frac{(\sqrt{3}-1)}{2} m_0c^2 \approx 0.37 m_0c^2, & \text{when } \beta_f = \frac{1}{2} \\ (\gamma_f - \frac{1}{2}) m_0c^2, & \text{when } \gamma_f \gg 1 (\beta_f \rightarrow 1). \end{cases} \quad (13)$$

Note that the final velocity, β_f , depends only on the total incident energy, E_I , and not on the energy schedule.

At small β_f , the required energy reduces to the nonrelativistic expression,

$$E_I = \frac{\beta_f}{2} m_0 c^2, \quad (14)$$

as it must. That this is the nonrelativistic expression can be seen as follows. In the small β_f limit, the reflected light momentum, $-\frac{E_R}{c}$, is equal in magnitude to the incident light momentum, $\frac{E_I}{c}$, so that the final momentum of the craft, $m_0 v$, must equal $2E_I/c$. So, $E_I = \frac{c}{2} m_0 v = \frac{\beta_f}{2} m_0 c^2$.

As an aside, we note that, instead of solving for E_I as a function of β_i and β_f , as in Eq. 12, we can solve for β_f as a function of β_i and E_I to obtain

$$\beta_f = \frac{\beta_i + 2\sqrt{1 - \beta_i^2} r + 2(1 - \beta_i)r^2}{1 + 2\sqrt{1 - \beta_i^2} r + 2(1 - \beta_i)r^2}, \quad (15)$$

where $r = E_I/(m_0 c^2)$. Eq. 13 is the same as Eq. 17 of Ref. 3 but our derivation is simpler. Eq. 15 is much simpler than Eq. 3 of Ref. 3, but they are equivalent as can be checked by plugging in arbitrary numerical values for β_i and r .

B. Derivation 2

We now give the second derivation, which consists of the following steps:

1. Relate the laser power to the force on the spacecraft.
2. Relate the force to the acceleration.
3. Integrate over time to relate the laser energy to the spacecraft velocity.

1) Relate the laser power to the force on the spacecraft

If photons are emitted at intervals of T , the photons are spaced cT apart in the Earth frame. In the Earth frame photons are moving with speed $c - v$ relative to the spacecraft, so in the Earth frame the photons are received by the spacecraft

$$T_r = \frac{cT}{(c - v)} = \frac{T}{1 - \beta} \quad (16)$$

apart in time.

Now, in the Earth frame, the spacecraft receives photons of the same energy that the laser emits, but the reflected photons have a lower energy. To figure that out, following the argument in

Einstein's 1905 paper⁴, transform to the spacecraft frame and back. In the spacecraft frame the energy of the received photons is Doppler shifted to

$$E'_I = E_I \sqrt{\frac{1-\beta}{1+\beta}} \quad (17)$$

and the reflected photons have very nearly the same energy because the spacecraft mass is about 10^{33} times greater than the relativistic mass of an optical photon. (The rest mass of the photon is of course zero, but the relativistic mass is not.) In the Earth frame, these reflected photons will be further Doppler shifted, so the energy in the Earth frame of the reflected photons is

$$E_R = E'_I \sqrt{\frac{1-\beta}{1+\beta}} = E_I \left(\frac{1-\beta}{1+\beta} \right) \quad (18)$$

Now, the force on the spacecraft equals the rate of change of its momentum, i.e., $F = \frac{dp}{dT} = \frac{\Delta p}{T_r}$, the magnitude of the momentum of a photon is its energy divided by c , and, the laser power is the rate at which it is emitting energy, i.e., $P = \frac{dE_I}{dT} = \frac{E_I}{T}$. Hence, using Eqs. 16 and 18, we get

$$F = \frac{\Delta p}{T_r} = \frac{E_I + E_R}{cT_r} = \frac{1}{c} \frac{1-\beta}{T} E_I \left(1 + \frac{1-\beta}{1+\beta} \right) = \frac{2}{c} \left(\frac{1-\beta}{1+\beta} \right) P$$

i.e. $\boxed{P = \frac{c}{2} \left(\frac{1+\beta}{1-\beta} \right) F}$ (relate force to power) (19)

2) Relate the force to the acceleration

Next, we derive the relativistic expression relating force and acceleration.

$$\begin{aligned} F &= \frac{d(\gamma m_0 v)}{dt} = \frac{d(\gamma m_0 c \beta)}{dt} = m_0 c \left(\frac{d(\gamma \beta)}{dt} \right) \\ &= m_0 c \left(\frac{d\gamma}{d\beta} \frac{d\beta}{dt} \beta + \gamma \frac{d\beta}{dt} \right) = m_0 c \left(\frac{d\gamma}{d\beta} \beta + \gamma \right) \frac{d\beta}{dt} \\ &= m_0 c \gamma (\beta^2 \gamma^2 + 1) \frac{d\beta}{dt} = m_0 c \gamma (\gamma^2) \frac{d\beta}{dt} \\ &= \gamma^3 m_0 c \frac{d\beta}{dt}. \end{aligned} \quad (20)$$

The relativistic force is γ^3 times larger than the nonrelativistic expression.

3) Integrate over time to relate the laser energy to the spacecraft velocity

Using Eqs. 19 and 20:

$$\begin{aligned}
 E_I &= \int_0^T P dt = \int_0^T \frac{c}{2} \left(\frac{1+\beta}{1-\beta} \right) \gamma^3 m_0 c \frac{d\beta}{dt} dt = \frac{m_0 c^2}{2} \int_0^{\beta_f} d\beta \left(\frac{1+\beta}{1-\beta} \right) \frac{1}{(1-\beta^2)^{3/2}} \\
 &= \boxed{\left(\frac{(2-\beta_f)\sqrt{1-\beta_f^2}}{3(1-\beta_f)^2} - \frac{2}{3} \right) \frac{m_0 c^2}{2}} \tag{21}
 \end{aligned}$$

If we divide Eq. 21 by the power of the laser, we obtain an expression for the time as a function of β_f that has also been published several times in the literature⁵⁻⁸ (see e.g. Eq. 11 of Ref. 6).

C. Summary of results from derivations 1 and 2

We have provided two derivations of the laser energy, E_I , needed to accelerate the spacecraft from zero velocity to β (we now drop the subscript, f , to simplify the notation), and arrived at two different expressions (Eqs. 13 and 21), which we repeat below for easy comparison:

Derivation 1 Result:

$$\boxed{E_I = \left(\sqrt{\frac{1+\beta}{1-\beta}} - 1 \right) \frac{m_0 c^2}{2}} \rightarrow \begin{cases} \frac{\beta}{2} m_0 c^2, & \text{when } \beta \rightarrow 0 \\ (\gamma - \frac{1}{2}) m_0 c^2, & \text{when } \gamma \gg 1 (\beta \rightarrow 1) \end{cases} \tag{22}$$

Derivation 2 Result:

$$\boxed{E_I = \left(\frac{(2-\beta)\sqrt{1-\beta^2}}{3(1-\beta)^2} - \frac{2}{3} \right) \frac{m_0 c^2}{2}} \rightarrow \begin{cases} \frac{\beta}{2} m_0 c^2, & \text{when } \beta \rightarrow 0 \\ \frac{4}{3} \gamma^3 m_0 c^2, & \text{when } \gamma \gg 1 (\beta \rightarrow 1) \end{cases} \tag{23}$$

We encourage the reader to pause and think about which is correct and relevant, and which is incorrect or not relevant. We note that both have the correct nonrelativistic limit, so this cannot be the basis for deciding. On the other hand, in the highly relativistic limit ($\gamma \rightarrow \infty$) they are dramatically different – the first scales as γ , whereas the second scales as γ^3 .

D. Resolution of the seeming contradiction

The two expressions answer two different questions. Eq. 22 answers the question: how much energy, E_I , has been received by the spacecraft when its velocity is β . Eq. 23 answers the question: how




FIG. 1. Comparison of the energy expressions in Eq. 22 and 23 and the nonrelativistic expression for the energy needed to achieve velocity β .

much energy, E_I , has been emitted by the laser when the velocity of the spacecraft is β , assuming that the laser emits with constant power. As the spacecraft moves away from Earth, a progressively larger fraction of the light emitted by the laser is in transit between Earth and the spacecraft and that light has not yet imparted any momentum to the spacecraft. This is why the energy in Eq. 23 is so much larger than that in Eq. 22 for large γ .

Which expression is relevant to the actual experiment? The laser beam has a small but nonzero angular spread, θ , so part of the beam will miss the light sail once the spacecraft is at a distance $d > \frac{L}{\theta}$, where L is the linear dimension of the light sail. Hence it makes sense to shine the laser during only the very early part of the mission, so that all of the laser light is captured by the sail.⁹ The spacecraft will reach its terminal velocity once this energy reaches it. Hence it is the first expression, Eq. 22, that is the relevant one, and that is fortunate since it gives the smaller energy estimate. Eq. 23 gives an energy that is 45% too large at $\beta = 0.5$, and 4.6 times too large at $\beta = 0.9$. Fig. IV D compares the energies of Eqs. 22 and 23 as well as the nonrelativistic expression.

E. Energy estimate for $\beta = 0.5$

We now estimate the energy needed to accelerate a spacecraft, with rest mass $m_0 = 2$ grams, to $\beta = 0.5$. From Eq. 22,

$$\begin{aligned} E_I &= \frac{1}{2} \left(\sqrt{\frac{1+\beta}{1-\beta}} - 1 \right) m_0 c^2 \\ &= 0.37 \times 0.002 \text{ kg} \times (3 \times 10^8 \text{ m/s})^2 = 7 \times 10^{13} \text{ J} = 19 \text{ Gigawatt hours} \end{aligned} \quad (24)$$

A typical power plant produces 1 Gigawatt of power. Assuming that the propulsion occurs during the first hour of flight, this would require harnessing the power of 19 power plants, but only for an hour. Since conversion of electric power to laser light is not 100% efficient, in practice more energy would be required.

V. ENERGY OF LIGHT REFLECTED BY A MOVING PERFECT MIRROR

We now find the expression for the energy reflected back by a moving perfect mirror. Again, let E_I and E_R be the energy of incident and the reflected light respectively, m_0 be the rest mass of the mirror, β_i and β_f be its initial and final velocities, and, $\hat{\mathbf{n}}$ and $-\hat{\mathbf{n}}$ be the direction of the incident and reflected light. We again use conservation of energy and momentum, but use of the 4-vector formalism simplifies the derivation. Equating 4-momenta before and after:

$$\frac{E_I}{c}(1, \hat{\mathbf{n}}) + \gamma_i m_0 c(1, \beta_i \hat{\mathbf{n}}) = \frac{E_R}{c}(1, -\hat{\mathbf{n}}) + \gamma_f m_0 c(1, \beta_f \hat{\mathbf{n}}) \quad (25)$$

$$\text{i.e., } \frac{E_I}{c}(1, \hat{\mathbf{n}}) - \frac{E_R}{c}(1, -\hat{\mathbf{n}}) + \gamma_i m_0 c(1, \beta_i \hat{\mathbf{n}}) = \gamma_f m_0 c(1, \beta_f \hat{\mathbf{n}}) \quad (26)$$

Squaring the 4-momenta we obtain,

$$(m_0 c)^2 - 4 \frac{E_I E_R}{c^2} + 2E_I \gamma_i m_0 (1 - \beta_i) - 2E_R \gamma_i m_0 (1 + \beta_i) = (m_0 c)^2 \quad (27)$$

$$\text{i.e., } \left((1 + \beta_i) \gamma_i + 2 \frac{E_I}{m_0 c^2} \right) E_R = (1 - \beta_i) \gamma_i E_I \quad (28)$$

$$\text{i.e., } \boxed{E_R = \frac{(1 - \beta_i)}{1 + \beta_i + 2r/\gamma_i} E_I}, \quad \text{where } r = \frac{E_I}{m_0 c^2} \quad (29)$$

Note that the frequency of the reflected photons changes over time as the mirror speeds up; the reflected energy, E_R , is the total reflected energy after energy E_I has been shone on the mirror which had initial velocity β_i . For a single photon and an electron as the target, this is the Compton

scattering formula, except that we allow the initial velocity, β_i , to be nonzero, and restrict the scattering angle to be 180° .

Einstein, in Section 8 of his first paper on special relativity⁴ in 1905, uses the argument presented in the section leading up to Eq. 18 to derive the expression

$$\nu_R = \frac{1 - \beta_i}{1 + \beta_i} \nu_I. \quad (30)$$

The more general expression of Eq. 29 reduces to Eq. 30 when either $m_0 \rightarrow \infty$, or, if β_i is interpreted as the instantaneous velocity and one considers classical electrodynamics so that the energy of photons is not quantized and the energy received, E_I , can be infinitesimal. On the other hand for quantized photons, Eq. 30 is never exactly correct though the correction term, r , is 10^{-33} for optical photons and a 2 gram spacecraft. Interestingly, just three months prior to his special relativity paper, Einstein had published his paper on the photoelectric effect¹⁰ (for which he was awarded the Nobel prize); nevertheless in his paper on special relativity he considers classical electrodynamics only.

Did Einstein realize that he was making an assumption when he derived Eq 30? Surprisingly, it appears he did not. He is rather emphatic in his paper:

“All problems in the optics of moving bodies can be solved by the method here employed. What is essential is, that the electric and magnetic force of the light which is influenced by a moving body, be transformed into a system of co-ordinates at rest relatively to the body. By this means all problems in the optics of moving bodies will be reduced to a series of problems in the optics of stationary bodies.”

A. The Breakthrough Starshot Project and Einstein’s Formula

There has been some confusion in the Breakthrough Starshot literature, in part caused by Einstein’s failure to state his assumptions when deriving Eq. 30. Kipping in his paper³ attributed the large (several percent) larger time that Lubin and coworkers found for the time required for the spacecraft to reach a given velocity to Lubin et al. using Eq. 30 rather than Eq. 29. In reality, that difference is due to Lubin computing a quantity that is not very relevant to the actual space mission, as explained in Section IVD. Later, Kipping published an erratum¹¹ in which he computed the same quantity as Lubin and found a much smaller difference from Lubin et al., which he again attributed to Lubin et al. using Eq. 30 rather than Eq. 29. However, even this much smaller difference is much larger than the actual difference, which is of $\mathcal{O}(r) \approx 10^{-33}$ as shown in the Supplementary Material. It is worth noting that the difference between Eq. 29 and Eq. 30 is large if the energy E_I incident

on the sail since the time that the sail had velocity β_i is sufficiently large that $r = E_I/(m_0c^2)$ is $\mathcal{O}(1)$. On the other hand, in our Derivation 2, which is equivalent to that of Lubin et al., β is the instantaneous velocity that is integrated over as it changes, and the difference between using Eq. 30 (the classical electrodynamics expression) and Eq. 29 (which is exact even for quantized photons) is $\mathcal{O}(r)$, where $r = h\nu/(m_0c^2) \approx 10^{-33}$ for optical photons and $m_0 \approx 2$ gram.

VI. CONCLUSIONS

In the first portion of this paper, we discussed the laser energy needed to propel a tiny spacecraft to Proxima Centauri. We derived two formulae and explained why the one most commonly presented in the literature is not relevant to the problem. In the second half of the paper, we discuss the energy reflected by a moving perfect mirror and point out that the formula in Einstein’s 1905 paper is correct only in two limiting cases. There has been some confusion in the Breakthrough Starshot literature, in part due to Einstein’s failure to point out the assumptions in his derivation, that we have cleared up.

-
- ¹ A. P. French, *Special Relativity* (The MIT Introductory Physics Series, W. W. Norton & Co., 1968).
- ² “Internet Investor and Science Philanthropist Yuri Milner & Physicist Stephen Hawking Announce Breakthrough Starshot Project to Develop 100 Million Mile per Hour Mission to the Stars within a Generation,” <https://breakthroughinitiatives.org/news/4>.
- ³ David Kipping, “Relativistic Light Sails,” *The Astronautical Journal* **153**, 277 (2017).
- ⁴ Albert Einstein, “Zur Elektrodynamik bewegter Körper (On the Electrodynamics of Moving Bodies),” *Annalen der Physik* **17**, 891 (1905).
- ⁵ Philip Lubin, “A Roadmap to Interstellar Flight,” arXiv preprint arXiv:1604.01356v7 (2016).
- ⁶ Neeraj Kulkarni, Philip Lubin, and Qicheng Zhang, “Relativistic Spacecraft Propelled by Directed Energy,” *The Astronautical Journal* **155**, 155 (2018).
- ⁷ P. Lubin and W. Hettel, “The Path to Interstellar Flight,” *Acta Futura* **12**, 9 (2020).
- ⁸ Philip Lubin, *The path to interstellar flight* (World Scientific Series on Emerging Technologies, Vol. 2, 2022).
- ⁹ Continuing to shine the laser beyond this point is wasteful because, for typical parameters, it results in only about a 10% increase in the terminal velocity of the spacecraft.
- ¹⁰ Albert Einstein, “Über einen die Erzeugung und Verwandlung des Lichtes betreffenden heuristischen Gesichtspunkt (Heuristic Point of View about the Creation and Conversion of Light),” *Annalen der Physik* **17**, 132 (1905).
- ¹¹ David Kipping, “Erratum: Relativistic Light Sails,” *The Astronomical Journal* **155**, 103 (2018).

Supplementary Information: Energy Needed to Propel a Tiny Spacecraft to Proxima Centauri, and, An Unstated Assumption in Einstein's 1905 Paper

C. J. Umrigar^{a1} and Tyler A. Anderson^{b1}

¹*Laboratory of Atomic and Solid State Physics,
Cornell University, Ithaca, NY 14853, USA.*

I. EXACT FORMULA FOR THE TIME NEEDED TO REACH β

In derivation 2 of the paper we find a formula for the total energy emitted by the laser at the instant the craft reaches a particular β . Dividing this expression by the power P of the laser we obtain an expression for the time in the Earth frame when the craft reaches a speed β :

$$t = \frac{m_0 c^2}{2P} \left(\frac{(2 - \beta)\sqrt{1 - \beta^2}}{3(1 - \beta)^2} - \frac{2}{3} \right). \quad (1)$$

This formula is strictly correct only in the continuum (classical electrodynamics) limit where the discrete nature of the photons emitted by the laser is ignored. We now derive an exact expression which is correct even outside the continuum limit and show that the correction to Eq. 1 is extremely small - on the order of 10^{-33} for Breakthrough Starshot. Although the beginning of our analysis resembles one already given by Kipping in Ref. 1, we make no approximations and arrive at a very different conclusion.

Suppose the laser emits photons of energy $E_\gamma = r m_0 c^2$ (recall that m_0 is the mass of the craft) at intervals of T so that $P = \frac{r m_0 c^2}{T}$. Let t_n be the time when n photons have been received by the craft and let β_n be the speed of the craft immediately after the n th photon has been received. We have already shown in the paper that, for a craft moving at β , the time between consecutive photon collisions with the sail is just

$$T_r = \frac{T}{1 - \beta}, \quad (2)$$

so that we may find the cumulative time t_n by summing over all these times:

$$t_n = \sum_{m=0}^{n-1} \frac{T}{1 - \beta_m}. \quad (3)$$

^a CyrusUmrigar@cornell.edu

^b taa65@cornell.edu

The total energy incident on the craft immediately after t_n is just

$$E_n = nE_\gamma = nrm_0c^2. \quad (4)$$

Using the formula from derivation 1 of the paper, we can relate the total energy incident on the craft E_n (and thus n) to β_n :

$$nr = \frac{E_n}{m_0c^2} = \frac{1}{2} \left(\sqrt{\frac{1+\beta_n}{1-\beta_n}} - 1 \right). \quad (5)$$

This expression can then be inverted to find β_n as a function of n :

$$\beta_n = \frac{(2nr+1)^2 - 1}{(2nr+1)^2 + 1}, \quad (6)$$

so that

$$\frac{1}{1-\beta_n} = \frac{(2nr+1)^2 + 1}{2}. \quad (7)$$

Substituting this expression into Eq. 3 and using the elementary formulae

$$\sum_{m=0}^{n-1} 1 = n,$$

$$\sum_{m=0}^{n-1} m = \frac{n(n-1)}{2},$$

$$\sum_{m=0}^{n-1} m^2 = \frac{n(n-1)(2n-1)}{6}$$

gives

$$\begin{aligned} t_n &= T \sum_{m=0}^{n-1} (1 + 2nr + 2(nr)^2) \\ &= T \left(n + rn(n-1) + \frac{1}{3}r^2n(n-1)(2n-1) \right). \end{aligned} \quad (8)$$

Rearranging and using $\frac{T}{r} = \frac{m_0c^2}{P}$ we find

$$t_n = \frac{m_0c^2}{P} \left(nr + (nr)^2 + \frac{2}{3}(nr)^3 - nr^2 - n^2r^3 + \frac{1}{3}nr^3 \right). \quad (9)$$

In the continuum limit $r \rightarrow 0$ but $n \rightarrow \infty$ in such a way that the product nr remains finite. Thus,

we expect the first three terms in Eq. 9 to agree with Eq. 1 while the remaining three terms represent a correction that goes to zero in the continuum limit. We now show that this is indeed the case by using Eq. 5 to write these three terms in terms of β_n :

$$nr = \frac{1}{2} \sqrt{\frac{1+\beta_n}{1-\beta_n}} - \frac{1}{2} \quad (10)$$

$$(nr)^2 = \frac{1}{4} \left(\frac{1+\beta_n}{1-\beta_n} \right) - \frac{1}{2} \sqrt{\frac{1+\beta_n}{1-\beta_n}} + \frac{1}{4} \quad (11)$$

$$\frac{2}{3}(nr)^3 = \frac{1}{12} \left(\sqrt{\frac{1+\beta_n}{1-\beta_n}} \right)^3 - \frac{1}{4} \left(\frac{1+\beta_n}{1-\beta_n} \right) + \frac{1}{4} \sqrt{\frac{1+\beta_n}{1-\beta_n}} - \frac{1}{12}. \quad (12)$$

Adding these three equations together we see that there are a number of immediate cancellations (indicated by terms of the same color), and we are left with

$$nr + (nr)^2 + \frac{2}{3}(nr)^3 = \frac{1}{12} \left(\sqrt{\frac{1+\beta_n}{1-\beta_n}} \right)^3 + \frac{1}{4} \sqrt{\frac{1+\beta_n}{1-\beta_n}} - \frac{1}{3} \quad (13)$$

$$= \frac{1}{6} \frac{(2-\beta_n)\sqrt{1-\beta_n^2}}{(1-\beta_n)^2} - \frac{1}{3}. \quad (14)$$

Substituting the above expression into Eq. 9 we finally obtain

$$t_n = \frac{m_0 c^2}{2P} \left(\frac{(2-\beta_n)\sqrt{1-\beta_n^2}}{3(1-\beta_n)^2} - \frac{2}{3} \right) + \delta t_n \quad (15)$$

where

$$\delta t_n = -\frac{m_0 c^2 nr}{P} \left(r + \left(n - \frac{1}{3} \right) r^2 \right). \quad (16)$$

Note that this correction is negative, but that is because the times t_n that we have chosen are immediately after the n th photon has struck. If instead, we had defined t_n to be the time just before the n th photon has struck, then the opposite sign would be obtained. What is important is that since nr is $\mathcal{O}(1)$ and r is $\mathcal{O}(10^{-33})$, δt_n is $\mathcal{O}(10^{-33})$ which is totally negligible.

¹ David Kipping, ‘‘Erratum: Relativistic Light Sails,’’ *The Astronomical Journal* **155**, 103 (2018).