Conference Proceedings for BCVSPIN 2024: Particle Physics and Cosmology in the Himalayas Kathmandu, Nepal, December 9-13, 2024

Exploring generalized Starobinsky Model of Inflation: Observational Constraints

Saisandri Saini¹, Akhilesh Nautiyal²

¹Department of Physics, Malaviya National Insitute of Technology, Jaipur, India ²Department of Physics, Malaviya National Insitute of Technology, Jaipur, India

E-mail: 2019rpy9082@mnit.ac.in

E-mail: akhilesh.phy@mnit.ac.in

Abstract. We examine the power-law Starobinsky model, a generalized version of the Starobinsky inflation model, characterized by a power-law correction to Einstein gravity. Employing the f(R) formalism, the scalar and tensor power spectra were numerically computed as functions of the dimensionless parameters M and β . A Markov Chain Monte Carlo (MCMC) analysis was conducted using Planck-2018, BICEP3 and BAO observational data, yielding precise constraints on $\beta = 1.987^{+0.013}_{-0.016}$, 95% C. L. and $\log_{10} M = -4.72^{+0.21}_{-0.20}$. The derived scalar spectral index $n_s = 0.9676^{+0.0069}_{-0.0068}$ and tensor-to-scalar ratio $r = 0.0074^{+0.0061}_{-0.0044}$ lie within the bounds set by Planck observations. We analyze a general reheating scenario while keeping the number of e-folds during inflation, N_{pivot} , fixed. The analysis confirms that deviations for high-energy physics and supergravity-based inflationary models.

1 Introduction

The inflationary paradigm [1] provides a robust framework for addressing the shortcomings of the standard Big Bang cosmology, such as the horizon and flatness problems [2], while simultaneously offering a mechanism for the origin of cosmic microwave background (CMB) anisotropies and large-scale structure [3–5]. Among various inflationary scenarios [6], the Starobinsky R^2 model [7] stands out as one of the most compelling in analysis of recent Planck 2018 results [8]. It relies solely on a curvature-squared term within the f(R) framework, without requiring the introduction of additional scalar fields.

Despite its success, generalizations of the Starobinsky model have gained attention, particularly those incorporating power-law corrections to the curvature term [9, 10]. Such extensions not only allow deviations from the canonical R^2 form but also provide a fertile ground for connecting inflationary physics with high-energy frameworks like supergravity [11, 12].

In this work, we focus on the power-law Starobinsky model, where the action includes a term proportional to R^{β} . This generalization introduces two free parameters, M and β , which are subject to observational constraints. By numerically evaluating the scalar and tensor perturbations, we perform a Bayesian parameter estimation using the latest CMB and BAO data to assess the observational viability of this model. The derived constraints on β offer new insights into the parameter space beyond the Starobinsky R^2 limit, highlighting the potential for deviations and their implications for inflationary dynamics and reheating scenarios. The analyses in [13–16] using the slow-roll approximation also found a deviation from $\beta = 2$. In [17], derived constraints on the inflaton potential parameters M and β from equation 3 using Planck-2018 data, in combination with BICEP3 [18], (BAO), (DES) and Pantheon observations. A general reheating scenario is considered, along with variations in the number of e-folds during inflation, N_{pivot} . However, in this study, we have set $N_{pivot} = 50$ as a fixed value.

2 The Generalized Starobinsky Model of Inflation

2.1 Overview of the Model

The generalized Starobinsky model introduces a power-law correction to the f(R) framework of gravity, extending the well-known R^2 inflationary model. This approach explores deviations from the canonical Starobinsky potential, offering a broader parameter space for analyzing inflationary dynamics and connecting with high-energy physics scenarios. The action for the model in the Jordan frame is given by [13]:

$$S_J = \frac{-M_{Pl}^2}{2} \int \sqrt{-g} \left(R + \frac{1}{6M^2} \frac{R^\beta}{M_{Pl}^{2\beta-2}} \right) d^4x.$$
(1)

where $M_{Pl}^2 = (8\pi G)^{-1}$, g is the determinant of the metric $g_{\mu\nu}$, R is the Ricci Scalar and M is a dimensionless real parameter.

For $\beta = 2$, the model reduces to the original Starobinsky R^2 inflationary model.

2.2 Transformation to the Einstein Frame

The formulation in the Jordan frame can be re-expressed in the Einstein frame by a conformal transformation. This allows the action to take the standard Einstein-Hilbert form with an additional scalar field χ . The transformed action in the Einstein frame is:

$$S_E = \int d^4x \sqrt{-\tilde{g}} \left(\frac{-M_{Pl}^2}{2} \tilde{R} + \frac{1}{2} \tilde{g}^{\mu\nu} \partial_\mu \chi \partial_\nu \chi + U(\chi) \right), \tag{2}$$

Where \tilde{g} and \tilde{R} are the metric and Ricci scalar in the Einstein frame. $U(\chi)$ is the scalar potential derived from the original f(R) function.

The scalar field potential in the Einstein frame for the generalized Starobinsky model is given as:

$$U(\chi) = \left(\frac{\beta - 1}{2}\right) \left(\frac{6M^2}{\beta^\beta}\right)^{\frac{1}{\beta - 1}} \exp\left[\frac{2\chi}{\sqrt{6}} \left(\frac{2 - \beta}{\beta - 1}\right)\right] \times \left(1 - \exp\left(\frac{-2\chi}{\sqrt{6}}\right)\right)^{\frac{\beta}{\beta - 1}}.$$
 (3)

2.3 Background and Perturbation Equations

In the Einstein frame, the equations governing the background dynamics during inflation are: **Friedmann equations:**

$$H^{2} = \frac{1}{3M_{Pl}^{2}} \left[\frac{1}{2} \dot{\chi}^{2} + V(\chi) \right].$$
(4)

$$\dot{H} = -\frac{1}{2M_{Pl}^2} \tag{5}$$

Scalar Field Equation:

$$\ddot{\chi} + 3H\dot{\chi} + \frac{dV(\chi)}{d\chi} = 0.$$
(6)

Here, the dot represents the differentiation with respect to cosmic time.

During inflation, perturbations in the metric and scalar field give rise to observable quantities such as the scalar power spectrum P_s and and the tensor-to-scalar ratio r. These perturbations are described by the Mukhanov-Sasaki equation for scalar perturbations and the tensor perturbation

equation [19,20] . Mukhanov-Sasaki Equation (Scalar Perturbations):

$$\frac{d^2 u_k}{d\tau^2} + \left(k^2 - \frac{1}{z}\frac{d^2 z}{d\tau^2}\right)u_k = 0,\tag{7}$$

where $z = \frac{1}{H} \frac{d\chi}{d\tau}$ encodes the background dynamics, τ denotes the conformal time. k is the comoving wavenumber. The scalar power spectrum is expressed as:

$$\mathcal{P}_{\mathcal{R}}(k) = \frac{k^3}{2\pi^2} \left| \frac{u_k}{z} \right|^2,\tag{8}$$

Tensor Perturbations: Similarly the mode equation for tensor perturbations generated during inflation is given as

$$\frac{dv_k}{d\tau^2} + \left(k^2 - \frac{1}{a}\frac{da''}{d\tau^2}\right)v_k = 0,\tag{9}$$

and the primordial tensor power spectrum is given as

$$\mathcal{P}_t(k) = \frac{4}{\pi^2} \frac{k^3}{M_{Pl}^2} \left| \frac{v_k}{a} \right|^2.$$
(10)

The scalar spectral index n_s and the tensor spectral index n_t are computed numerically from the power spectra by applying their respective definitions [21].

$$n_s = 1 + \frac{d\ln \mathcal{P}_{\mathcal{R}}}{d\ln k},\tag{11}$$

$$n_t = \frac{d\ln \mathcal{P}_t}{d\ln k},\tag{12}$$

The tensor-to-scalar ratio r is defined by [21]

$$r = \frac{\mathcal{P}_t}{\mathcal{P}_{\mathcal{R}}}.$$
(13)

The background and perturbation equations are solved numerically using the e-folding number , $N = \ln a$ as the independent variable. The initial conditions are set deep in the slow-roll regime, where the scalar field's velocity is small, and the perturbation modes are initialized in the Bunch-Davies vacuum state. The Planck CMB observations impose constrains on n_s and r. However, in this study n_s and r are treated as derived quantities, while the primary parameters constrained directly by the CMB data are those of the inflaton potential (3), namely M and β .

3 Observational Results and Parameter Constraints

In this study, the parameters of the power-law Starobinsky model were constrained using data from Planck-2018, BICEP3 and baryon acoustic oscillations (BAO). The analysis was conducted using a numerical framework to solve the background and perturbation equations of the model without relying on the slow-roll approximation. To explore the parameter space of the model, the publicly available ModeCode [22] was adapted to incorporate the power-law Starobinsky potential. This numerical tool computed the scalar and tensor power spectra, which were then input into CAMB [23] to generate the angular power spectra for the cosmic microwave background (CMB) anisotropy and polarization. CosmoMC [24] was employed for the Markov Chain Monte Carlo (MCMC) analysis, enabling robust parameter estimation.

Table 1 presents the constraints derived for the parameters of potential 3, including the efoldings N_{pivot} and the derived parameters r and n_s . The priors for the model parameters were set same as [17] as follows:

$$\log_{10} M \in [-6.5, -3.0], \quad \beta \in [1.90, 2.07].$$
(14)

Parameter	68% limits	95% limits	99% limits
N_{pivot}	$48.7^{+2.9}_{-7.8}$	49^{+10}_{-8}	49^{+10}_{-8}
$\log_{10} M$	$-4.72^{+0.11}_{-0.14}$	$-4.72^{+0.21}_{-0.20}$	$-4.72^{+0.27}_{-0.23}$
β	$1.987^{+0.011}_{-0.0048}$	$1.987^{+0.013}_{-0.016}$	$1.987^{+0.014}_{-0.022}$
n_s	0.9676 ± 0.0035	$0.9676^{+0.0069}_{-0.0068}$	$0.9676^{+0.0090}_{-0.0089}$
r	$0.0074_{-0.0039}^{+0.0016}$	$0.0074^{+0.0061}_{-0.0044}$	$0.0074_{-0.0048}^{+0.0086}$

Table 1: Constraints on parameters of potential, r and n_s using Planck-2018, BICEP/Keck (BK15) and BAO observations.



Figure 1: Marginalized joint two-dimensional 68% C.L. and 95% C.L. constraints on parameters of potential (a) and N_{pivot} (b) using Planck-2018, BICEP/Keck (BK15) and BAO data

The Table 1 clearly indicates that the most suitable value of β is

$$\beta = 1.987^{+0.013}_{-0.016}, \ 95\% C. L., \tag{15}$$

The triangular plots 1 display the posterior distributions and correlation contours between different cosmological parameters. Fig. 1(a) indicates a strong correlation between model parameters $(\log_{10} M, \beta)$ and N_{pivot} also to each other. Fig.1(b) the spectral index n_s and the tensor-to-scalar ratio r display a strong correlation. As observed in Table 1, the best-fit values of the scalar spectral index n_s and the tensor-to-scalar ratio r, derived from the optimal potential parameters of the power-law Starobinsky model, remain consistent with the Planck limits.

In this study, we investigate power law Starobinsky inflation in the context of Planck-2018, BICEP3 [18] CMB data. We consider the inflaton potential (3) for power-law Starobinsky inflation in the Einstein frame and numerically compute the scalar and tensor power spectra using MODE-CODE. Utilizing these results, we conduct an MCMC analysis with COSMOMC to constrain the inflaton potential parameters β and M, as well as the number of e-folds at the pivot scale, N_{pivot} . To account for deviations from Starobinsky inflation, we vary β within the range 1.9 to 2.07. Based on Planck-2018 data, we find that $\beta = 1.987^{+0.013}_{-0.016}$, 95% confidence level ($\beta = 1.966^{+0.032}_{-0.042}$, in [17]), indicating that current CMB and LSS observations favor a slight deviation from the standard Starobinsky inflationary scenario.

References

[1] A. H. Guth, Phys. Rev. **D23**, 347 (1981).

- [2] A. D. Linde, Phys. Lett. B 108, 389-393 (1982) doi:10.1016/0370-2693(82)91219-9
- [3] V. F. Mukhanov and G. V. Chibisov, JETP Lett. 33, 532 (1981).
- [4] A. A. Starobinsky, Phys. Lett. **B117**, 175 (1982).
- [5] A. H. Guth and S.-Y. Pi, Phys. Rev. D32, 1899 (1985).
- [6] J. Martin, C. Ringeval and V. Vennin, Phys. Dark Univ. 5-6, 75-235 (2014) doi:10.1016/j.dark.2014.01.003 [arXiv:1303.3787 [astro-ph.CO]].
- [7] A. A. Starobinsky, Phys. Lett. B **91**, 99-102 (1980) doi:10.1016/0370-2693(80)90670-X
- [8] Y. Akrami et al. [Planck], Astron. Astrophys. 641, A10 (2020) doi:10.1051/0004-6361/201833887 [arXiv:1807.06211 [astro-ph.CO]].
- [9] V. Muller, H. J. Schmidt and A. A. Starobinsky, Class. Quant. Grav. 7, 1163-1168 (1990) doi:10.1088/0264-9381/7/7/012
- [10] S. Gottlober, V. Muller, H. J. Schmidt and A. A. Starobinsky, Int. J. Mod. Phys. D 1, 257-279 (1992) doi:10.1142/S0218271892000136
- [11] J. Ellis, D. V. Nanopoulos and K. A. Olive, JCAP 10, 009 (2013) doi:10.1088/1475-7516/2013/10/009 [arXiv:1307.3537 [hep-th]].
- [12] J. Ellis, D. V. Nanopoulos, K. A. Olive and S. Verner, JHEP 03, 099 (2019) doi:10.1007/JHEP03(2019)099 [arXiv:1812.02192 [hep-th]].
- [13] G. K. Chakravarty and S. Mohanty, Phys. Lett. B 746, 242-247 (2015) doi:10.1016/j.physletb.2015.04.056 [arXiv:1405.1321 [hep-ph]].
- [14] H. Motohashi, Phys. Rev. D 91, 064016 (2015) doi:10.1103/PhysRevD.91.064016 [arXiv:1411.2972 [astro-ph.CO]].
- [15] S. D. Odintsov and V. K. Oikonomou, Int. J. Mod. Phys. D 32, no.01, 2250135 (2023) doi:10.1142/S0218271822501358 [arXiv:2210.11351 [gr-qc]].
- [16] S. Meza, D. Altamirano, M. Z. Mughal and C. Rojas, Int. J. Mod. Phys. D 30, no.08, 2150062 (2021) doi:10.1142/S0218271821500620 [arXiv:2104.01139 [gr-qc]].
- [17] S. Saini and A. Nautiyal, Phys. Rev. D 108, no.12, 123505 (2023) doi:10.1103/PhysRevD.108.123505 [arXiv:2305.00682 [astro-ph.CO]].
- [18] P. A. R. Ade *et al.* [BICEP and Keck], Phys. Rev. Lett. **127**, no.15, 151301 (2021) doi:10.1103/PhysRevLett.127.151301 [arXiv:2110.00483 [astro-ph.CO]].
- [19] V. F. Mukhanov, Sov. Phys. JETP 67, 1297-1302 (1988)
- [20] M. Sasaki, Prog. Theor. Phys. 76, 1036 (1986) doi:10.1143/PTP.76.1036
- [21] B. A. Bassett, S. Tsujikawa and D. Wands, Rev. Mod. Phys. 78, 537-589 (2006) doi:10.1103/RevModPhys.78.537 [arXiv:astro-ph/0507632 [astro-ph]].
- [22] M. J. Mortonson, H. V. Peiris and R. Easther, Phys. Rev. D 83, 043505 (2011) doi:10.1103/PhysRevD.83.043505 [arXiv:1007.4205 [astro-ph.CO]].
- [23] A. Lewis, A. Challinor and A. Lasenby, Astrophys. J. 538, 473-476 (2000) doi:10.1086/309179
 [arXiv:astro-ph/9911177 [astro-ph]].
- [24] A. Lewis and S. Bridle, Phys. Rev. D 66, 103511 (2002) doi:10.1103/PhysRevD.66.103511 [arXiv:astro-ph/0205436 [astro-ph]].