

Correspondence between Myrzakulov $F(R, Q)$ gravity and Tsallis cosmology

Andreas Lymp^a Gulgassyl Nugmanova^b Almira Sergazina^b

^a*Department of Physics, University of Patras, 26500 Patras, Greece*

^b*Eurasian National University, Astana, 010008, Kazakhstan*

E-mail: alymp@upatras.gr

ABSTRACT: We investigate the correspondence between Myrzakulov $F(R, Q)$ gravity and Tsallis cosmology. The former is a modified gravity that uses both curvature and non-metricity, while the latter is a modified cosmology arising from the gravity-thermodynamics conjecture, employing Tsallis entropy instead of the Bekenstein-Hawking one. By appropriately identifying the functional dependencies and the model parameters, we demonstrate that both frameworks can give identical background evolution, reproducing the standard cosmological sequence of matter and dark energy domination. However, their perturbation behavior exhibits differences, since the growth of density fluctuations and the effective Newton constant deviate between the two scenarios, indicating that perturbative observables, such as structure formation and weak-lensing ones, could serve as distinguishing factors between them.

KEYWORDS: Metric-affine gravity, gravity-thermodynamics conjecture, Tsallis cosmology

Contents

1	Introduction	1
2	Myrzakulov $F(R, Q)$ Cosmology	2
3	Tsallis cosmology	5
4	Correspondence between Myrzakulov $F(R, Q)$ gravity and Tsallis cosmology	7
4.1	Background evolution	7
4.2	Perturbation evolution	9
5	Conclusions	10

1 Introduction

The Standard Model of Cosmology, which consists of Λ -Cold Dark Matter (Λ CDM) within general relativity, has proven to be very effective in explaining the evolution of the universe, both in terms of background dynamics and perturbation theory [1]. Despite its successes, certain theoretical challenges, such as the cosmological constant problem [2], the non-renormalizability of general relativity [3], and possible observational tensions [4], have motivated the exploration of various extensions and alternative approaches. Broadly speaking, these extensions fall into two main categories. The first category retains general relativity as the fundamental gravitational framework but introduces additional components, such as the dark energy sector [5, 6]. The second category involves modifications of gravity itself, where general relativity emerges as a limiting case, capable of driving cosmic acceleration and richer cosmological behavior [7–9].

One approach to construct modified gravity theories involves modifying the Einstein-Hilbert action by introducing additional terms, leading to various extensions such as $f(R)$ gravity [10–12] or Gauss-Bonnet and $f(G)$ gravity [13–16]. Alternatively, modifications can arise within the torsional gravitational framework, giving rise to theories such as $f(T)$ gravity [17–23], as well as its extensions like $f(T, T_G)$ gravity [24] and $f(T, B)$ gravity [25]. Moreover, there can be other formulations based on nonmetricity and the symmetric teleparallel connection, leading to $f(Q)$ gravity [26, 27] and its variants such as $f(Q, C)$ gravity [28].

Within this geometrical framework, Myrzakulov $F(R, Q)$ gravity incorporates both curvature and nonmetricity to describe gravitational interactions [29]. By employing a generalized connection that simultaneously accounts for both geometric properties, this theory establishes a more intricate gravitational structure, leading to a richer cosmological

behavior. This enhanced framework allows for interesting phenomenology, and has led to many applications [29–44].

On the other hand, there is another approach to account for the theoretical and observational puzzles, namely to apply the gravity-thermodynamics conjecture [45–47]. Specifically, it has been demonstrated that the cosmological Friedmann equations can be reformulated as the first law of thermodynamics when treating the universe as a thermodynamic system bounded by the apparent horizon [48–58]. Apart from the basic formulation, using the standard Bekenstein-Hawking entropy, this approach has been applied to extended entropies too, such as Tsallis, Barrow, Kaniadakis etc entropies, and proves to exhibit rich cosmological phenomenology [59–79].

In this paper we examine the Correspondence between Myrzakulov $F(R, Q)$ gravity and Tsallis cosmology. In particular, we are interested in studying under which connection choices one obtains equivalence of the two frameworks. The paper is structured as follows: In Section 2, we briefly review Myrzakulov $F(R, Q)$ gravity and cosmology, while in Section 3 we present the modified Friedmann equations in Tsallis cosmology. Then, in Section 4 we reconstruct the involved functions and parameters in order to make the two theories coincide at the background level, and additionally we study the perturbation evolution showing that the two theories deviate. Finally, we summarize our results in Section 5.

2 Myrzakulov $F(R, Q)$ Cosmology

The central idea of this modified gravity is the modification of the underlying connection. In particular, we introduce a connection [41]

$$\Gamma^\rho{}_{\mu\nu} = \hat{\Gamma}^\rho{}_{\mu\nu} + L^\rho{}_{\mu\nu}, \quad (2.1)$$

with $\hat{\Gamma}^\rho{}_{\mu\nu}$ the Levi-Civita connection and $L^\rho{}_{\mu\nu}$ the disformation tensor given by

$$L^\rho{}_{\mu\nu} = \frac{1}{2}g^{\rho\lambda}(-Q_{\mu\nu\lambda} - Q_{\nu\mu\lambda} + Q_{\lambda\mu\nu}), \quad (2.2)$$

and where $Q_{\rho\mu\nu} = \nabla_\rho g_{\mu\nu}$ is the nonmetricity tensor.

As it is known, the Ricci curvature tensor $R_{\mu\nu}$ in terms of the general connection is written as [35, 80, 81]

$$R_{\mu\nu} = \partial_\lambda \Gamma^\lambda{}_{\mu\nu} - \partial_\mu \Gamma^\lambda{}_{\lambda\nu} + \Gamma^\lambda{}_{\lambda\alpha} \Gamma^\alpha{}_{\mu\nu} - \Gamma^\lambda{}_{\mu\alpha} \Gamma^\alpha{}_{\lambda\nu}, \quad (2.3)$$

which can be expressed in terms of the Levi-Civita connection as

$$R_{\mu\nu} = \hat{R}_{\mu\nu} + \partial_\lambda L^\lambda{}_{\mu\nu} - \partial_\mu L^\lambda{}_{\lambda\nu} + \hat{\Gamma}^\lambda{}_{\lambda\alpha} L^\alpha{}_{\mu\nu} + \hat{\Gamma}^\alpha{}_{\mu\nu} L^\lambda{}_{\lambda\alpha} - \hat{\Gamma}^\lambda{}_{\mu\alpha} L^\alpha{}_{\lambda\nu} - \hat{\Gamma}^\alpha{}_{\lambda\nu} L^\lambda{}_{\mu\alpha} + L^\lambda{}_{\lambda\alpha} L^\alpha{}_{\mu\nu} - L^\lambda{}_{\mu\alpha} L^\alpha{}_{\lambda\nu}, \quad (2.4)$$

where by $\hat{R}_{\mu\nu}$ we mark the Ricci curvature tensor with respect to Levi-Civita connection. Hence, the Ricci scalar $R = g^{\mu\nu} R_{\mu\nu}$ with respect to the general connection can be re-written as

$$R = \hat{R} + u, \quad (2.5)$$

where u is a function depending on the metric, its first and second derivatives, and the connection and its first derivative.

In the same lines, the nonmetricity tensor with respect to general connection can be expressed as

$$Q_{\rho\mu\nu} = \partial_\rho g_{\mu\nu} - \Gamma_{\mu\rho}^\lambda g_{\lambda\nu} - \Gamma_{\nu\rho}^\lambda g_{\lambda\mu}, \quad (2.6)$$

which can be re-written as

$$Q_{\rho\mu\nu} = \check{Q}_{\rho\mu\nu} + (-L_{\mu\rho}^\lambda g_{\lambda\nu} - L_{\nu\rho}^\lambda g_{\lambda\mu}), \quad (2.7)$$

where $\check{Q}_{\rho\mu\nu}$ is the nonmetricity scalar calculated with the symmetric teleparallel connection. Thus, we can express the nonmetricity scalar $Q = -g^{\mu\nu} (L_{\beta\mu}^\alpha L_{\nu\alpha}^\beta - L_{\beta\alpha}^\alpha L_{\mu\nu}^\beta)$ as

$$Q = \check{Q} + w, \quad (2.8)$$

where w is a function depending on the metric and its first derivative and on the connection.

One can now proceed to introducing the action of the theory as [41]

$$S = \frac{1}{16\pi G} \int [F(R, Q) + L_m] \sqrt{-g} d^4x, \quad (2.9)$$

where g is the metric determinant, G is the Newton constant, and L_m is the matter Lagrangian. Thus, performing variation in terms of the metric yields [41]

$$-\frac{1}{2}g_{\mu\nu}F + F_R R_{(\mu\nu)} + F_Q L_{(\mu\nu)} + \hat{\nabla}_\lambda (F_Q J^\lambda_{(\mu\nu)}) + g_{\mu\nu} \hat{\nabla}_\lambda (F_Q \zeta^\lambda) = 8\pi G T_{\mu\nu}, \quad (2.10)$$

with $F_R = \frac{\partial F}{\partial R}$, $F_Q = \frac{\partial F}{\partial Q}$ and where as usual the matter energy-momentum tensor is defined as $T_{\mu\nu} \equiv -\frac{2}{\sqrt{-g}} \frac{\delta(\sqrt{-g} \mathcal{L}_m)}{\delta g^{\mu\nu}}$. In the above expressions we have introduced the quantities

$$L_{\mu\nu} = \frac{1}{4} \left[(Q_{\mu\alpha\beta} - 2Q_{\alpha\beta\mu}) Q_\nu^{\alpha\beta} + (Q_\mu + 2\tilde{Q}_\mu) Q_\nu + (2Q_{\mu\nu\alpha} - Q_{\alpha\mu\nu}) Q^\alpha \right] \quad (2.11)$$

$$J^\lambda_{\mu\nu} = \sqrt{-g} \left(\frac{1}{4} Q^\lambda_{\mu\nu} - \frac{1}{2} Q_{\mu\nu}{}^\lambda \right), \quad (2.12)$$

where $Q_\lambda = Q_\lambda{}^\alpha{}_\alpha$ and $\tilde{Q}^\lambda = Q_\alpha{}^{\lambda\alpha}$, while $\hat{\nabla}_\lambda = \left(-\frac{1}{2}Q^\lambda + \tilde{Q}^\lambda \right) - \nabla_\lambda$, with ∇_λ the covariant derivative associated with the general connection and $\hat{\nabla}_\lambda$ the covariant derivative corresponding to the Levi-Civita connection. Additionally, the connection field equations are [41]

$$P_\lambda{}^{\mu\nu}(F_R) + F_Q \left[2Q^{[\nu\mu]}{}_\lambda - Q_\lambda{}^{\mu\nu} + (\tilde{Q}^\nu - Q^\nu) \delta_\lambda^\mu + Q_\lambda g^{\mu\nu} + \frac{1}{2} Q^\mu \delta_\lambda^\nu \right] = 0, \quad (2.13)$$

where $P_\lambda{}^{\mu\nu}(F_R)$ is the Palatini-like tensor given by

$$P_\lambda{}^{\mu\nu}(F_R) := -\frac{\nabla_\lambda (\sqrt{-g} F_R g^{\mu\nu})}{\sqrt{-g}} + \frac{\nabla_\alpha (\sqrt{-g} F_R g^{\mu\alpha} \delta_\lambda^\nu)}{\sqrt{-g}}. \quad (2.14)$$

We proceed to the cosmological application. We choose a flat Friedmann-Robertson-Walker (FRW) metric, namely

$$ds^2 = -dt^2 + a^2(t) \delta_{ij} dx^i dx^j, \quad (2.15)$$

where $a(t)$ is the scale factor. In this case, the Ricci scalar calculated in terms of the Levi-Civita connection is

$$\hat{R} = 6 \left(\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} \right), \quad (2.16)$$

while the nonmetricity scalar calculated in terms of the symmetric teleparallel connection is

$$\check{Q} = 6 \frac{\dot{a}^2}{a^2}. \quad (2.17)$$

In order to extract the field equations we follow the mini-superspace approach [30] and we set $u = u(a, \dot{a}, \ddot{a})$ and $w = w(a, \dot{a})$. Moreover, for the matter Lagrangian we set $L_m = -\rho_m(a)$, with ρ_m the matter energy density considered to correspond to a perfect fluid [82]. Thus, the Lagrangian of the action (2.9) becomes

$$\begin{aligned} \mathcal{L}(a, R, Q, \dot{a}, \dot{R}, \dot{Q}) &= a^3(F - RF_R - QF_Q) + 6a\dot{a}^2(F_R + F_Q) \\ &+ 6a^2\dot{a}\dot{F}_R + a^3(uF_R + wF_Q) + 16\pi G a^3 L_m. \end{aligned} \quad (2.18)$$

Hence, varying the above Lagrangian provides the field equations

$$\begin{aligned} -\frac{1}{2}(F - RF_R - QF_Q) + 3H^2(F_R + F_Q) - \frac{1}{2}[(u - \dot{a}u_{\dot{a}})F_R + (w - \dot{a}w_{\dot{a}})F_Q] \\ + 3H(\dot{R}F_{RR} + \dot{Q}F_{RQ}) = 8\pi G \rho_m \end{aligned} \quad (2.19)$$

and

$$\begin{aligned} -\frac{1}{2}(F - RF_R - QF_Q) + (2\dot{H} + 3H^2)(F_R + F_Q) - \frac{1}{2}(u + \frac{1}{3}au_a - \dot{a}u_{\dot{a}} - \frac{1}{3}a\dot{u}_{\dot{a}})F_R \\ - \frac{1}{2}(w + \frac{1}{3}aw_a - \dot{a}w_{\dot{a}} - \frac{1}{3}a\dot{w}_{\dot{a}})F_Q + 2H(\dot{F}_R + \dot{F}_Q) + \frac{1}{6}a(u_{\dot{a}}\dot{F}_R + w_{\dot{a}}\dot{F}_Q) + \ddot{F}_R = -8\pi G p_m, \end{aligned} \quad (2.20)$$

with $H = \frac{\dot{a}}{a}$ the Hubble parameter and p_m the matter pressure.

In this paper we will investigate the simplest model, namely

$$F(R, Q) = R + \lambda Q + \tilde{\Lambda}. \quad (2.21)$$

Inserting this ansatz into the FRW field equations we finally extract the modified Friedmann equations as

$$3H^2 = 8\pi G (\rho_m + \rho_{DE}) \quad (2.22)$$

$$2\dot{H} + 3H^2 = -8\pi G (p_m + p_{DE}), \quad (2.23)$$

where we have defined an effective dark energy sector, with energy density and pressure respectively given by

$$\rho_{DE} = \frac{1}{8\pi G} \left[\frac{1}{2}(u - \dot{a}u_{\dot{a}}) + \frac{\lambda}{2}(w - \dot{a}w_{\dot{a}}) - 3\lambda H^2 + \frac{\tilde{\Lambda}}{2} \right] \quad (2.24)$$

$$p_{DE} = -\frac{1}{8\pi G} \left[-\frac{1}{2}\left(u + \frac{1}{3}au_a - \dot{a}u_{\dot{a}} - \frac{1}{3}a\dot{u}_{\dot{a}}\right) - \frac{\lambda}{2}\left(w + \frac{1}{3}aw_a - \dot{a}w_{\dot{a}} - \frac{1}{3}a\dot{w}_{\dot{a}}\right) - \frac{\tilde{\Lambda}}{2} + \lambda(2\dot{H} + 3H^2) \right], \quad (2.25)$$

where the subscripts a, \dot{a}, \ddot{a} mark the partial derivatives with respect to them. As we can see, assuming that the matter sector is conserved, we obtain the effective dark energy conservation as

$$\dot{\rho}_{DE} + 3H(\rho_{DE} + p_{DE}) = 0. \quad (2.26)$$

Finally, we can introduce the equation-of-state parameter of the effective dark energy as $w_{DE} \equiv p_{DE}/\rho_{DE}$.

3 Tsallis cosmology

In this section we present Tsallis cosmology, namely the modified Friedmann equations that arise in the gravity-thermodynamics framework if one uses the extended Tsallis entropy instead of the standard Bekenstein-Hawking one.

Tsallis entropy is a generalization of the standard Boltzmann-Gibbs entropy, introduced by Tsallis in 1988 as a framework for extending statistical mechanics to non-extensive systems [83–86]. It is defined as

$$S_T = k_B \frac{1 - \sum_i p_i^q}{q - 1}, \quad (3.1)$$

where q is the entropic index that characterizes the degree of non-additivity, p_i represents the probability of the system's microstates, and k_B is the Boltzmann constant. In the limit $q \rightarrow 1$, Tsallis entropy recovers the classical Boltzmann-Gibbs entropy. Tsallis entropy is relevant in systems exhibiting long-range interactions, correlations, or fractal-like phase space structures, where the standard thermodynamic formalism fails. In the case of a black hole, the nonextensive Tsallis entropy can be written in compact form as [87]

$$S_T = \frac{\tilde{\alpha}}{4G} A^\delta, \quad (3.2)$$

with $A \propto L^2$ the area of the system, having characteristic length L , and $\tilde{\alpha}$ and δ the parameters. As one can see, for $\delta = 1$ and $\tilde{\alpha} = (4G)^{-1}$ (in units where $\hbar = k_B = c = 1$), we obtain the usual additive entropy.

Let us now apply the gravity-thermodynamics conjecture. This formulation states that the first law of thermodynamics can be applied to the cosmological horizon [45–47]. The

most commonly considered horizon is the apparent horizon [48, 49, 88], which is defined as:

$$r_A = \frac{1}{\sqrt{H^2 + \frac{k}{a^2}}}. \quad (3.3)$$

To describe the thermodynamic properties of the horizon, one uses the temperature formula analogous to that of black holes, replacing the black hole event horizon with the apparent horizon [89]:

$$T_h = \frac{1}{2\pi r_A}. \quad (3.4)$$

Similarly, the entropy associated with the apparent horizon will be the standard Bekenstein-Hawking one, namely

$$S_h = \frac{1}{4G}A. \quad (3.5)$$

By considering the energy flux across the apparent horizon, given by $\delta Q = -dE = A(\rho_m + p_m)Hr_A dt$, and employing the first law of thermodynamics $-dE = TdS$, alongside (3.4) and (3.5), we derive [49]:

$$-4\pi G(\rho_m + p_m) = \dot{H} - \frac{k}{a^2}, \quad (3.6)$$

$$\frac{8\pi G}{3}(\rho_m) = H^2 + \frac{k}{a^2} - \frac{\Lambda}{3}, \quad (3.7)$$

with the integration constant giving rise to the cosmological constant.

The gravity-thermodynamics conjecture has been applied in the case of various extended entropies [59–79]. If one uses Tsallis entropy (3.2) instead of the standard one, then he finds [59]

$$H^2 = \frac{8\pi G}{3}(\rho_m + \rho_{DE}) \quad (3.8)$$

$$\dot{H} = -4\pi G(\rho_m + p_m + \rho_{DE} + p_{DE}), \quad (3.9)$$

where we have defined the effective dark energy density and pressure as

$$\rho_{DE} = \frac{1}{8\pi G} \left\{ \Lambda + 3H^2 \left[1 - \alpha \frac{\delta}{2-\delta} H^{2(1-\delta)} \right] \right\}, \quad (3.10)$$

$$p_{DE} = \frac{1}{8\pi G} \left\{ \Lambda + 2\dot{H} \left[1 - \alpha \delta H^{2(1-\delta)} \right] + 3H^2 \left[1 - \alpha \frac{\delta}{2-\delta} H^{2(1-\delta)} \right] \right\}, \quad (3.11)$$

where $\alpha \equiv (4\pi)^{\delta-1} \tilde{\alpha}$. Lastly, the dark-energy equation-of-state parameter reads as

$$w_{DE} \equiv \frac{p_{DE}}{\rho_{DE}} = -1 - \frac{2\dot{H} \left[1 - \alpha \delta H^{2(1-\delta)} \right]}{\Lambda + 3H^2 \left[1 - \frac{\alpha \delta}{2-\delta} H^{2(1-\delta)} \right]}. \quad (3.12)$$

As expected, for $\delta = 1$ all above equations reduce to the ones of Λ CDM scenario.

4 Correspondence between Myrzakulov $F(R, Q)$ gravity and Tsallis cosmology

In this section, we investigate the conditions under which the Myrzakulov $F(R, Q)$ gravity framework can be mapped to Tsallis cosmology. We first study the background behavior, and then we proceed to the examination of perturbations.

4.1 Background evolution

To make the two theories coincide at the background level, we appropriately choose the functions $u(a, \dot{a}, \ddot{a})$ and $w(a, \dot{a})$ in such a way that the modified Friedmann equations in both scenarios become identical. Observing the forms (2.24), (2.25) as well as (3.10), (3.11) we impose the ansatz

$$u(a, \dot{a}, \ddot{a}) = \epsilon \left(\frac{\dot{a}}{a} \right)^\zeta + \eta \left(\frac{\dot{a}}{a} \right)^\theta, \quad (4.1)$$

$$w(a, \dot{a}) = \xi \left(\frac{\dot{a}}{a} \right). \quad (4.2)$$

Hence, we can see that we obtain a correspondence between the two theories if we make the identifications $\lambda = 3$, $\tilde{\Lambda} = 2\Lambda$, $\eta = -12$, $\theta = 2$, $\xi = 10$, $\zeta = 4 - 2\delta$ and $\epsilon = \frac{6\alpha\delta}{(2-\delta)(3-2\delta)}$. This correspondence may establish a connection between metric-affine modifications of gravity and non-additive entropy formalisms in cosmology.

Let us examine the cosmological evolution in more detail. As the independent variable we use the redshift z , defined as $1+z = 1/a$ (we set the present scale factor to 1). Moreover, we introduce the matter and dark energy density parameters as

$$\Omega_m = \frac{8\pi G}{3H^2} \rho_m \quad (4.3)$$

$$\Omega_{DE} = \frac{8\pi G}{3H^2} \rho_{DE}. \quad (4.4)$$

Without loss of generality we will focus on dust matter, namely we take $p_m = 0$. In this case the conservation equation gives $\rho_m = \frac{\rho_{m0}}{a^3}$, with ρ_{m0} the current value of the matter energy (from now on the subscript ‘‘0’’ marks the present value of a quantity). Thus, for dust matter we have $\Omega_m = \Omega_{m0} H_0^2 / (a^3 H^2)$. Moreover, we set the value of $\tilde{\Lambda}$ (and thus of Λ too) in order for the Friedmann equation (2.22) applied at present time to give $\Omega_{m0} \approx 0.31$ in agreement with observations [90]. Additionally, without loss of generality we set $\alpha = 1$, namely to its standard value.

In the upper panel of Fig. 1 we present the evolution of the density parameters $\Omega_{DE}(z)$ and $\Omega_m(z) = 1 - \Omega_{DE}(z)$, for Myrzakulov cosmology and Tsallis cosmology, under the aforementioned identifications. In the lower panel we show the dark-energy equation-of-state parameter. Note that we extend the evolution into the future too. As expected, there is no distinction between the two scenarios, since they were reconstructed to coincide at the background level. As we can see, we obtain the sequence of matter and dark energy epochs, while in the future the universe results to the de-Sitter state. Concerning the dark-energy equation-of-state parameter we can see that it may enter in the phantom regime, experiencing the phantom-divide crossing.




Figure 1. Upper graph: The dark-energy density parameter Ω_{DE} (black-solid) and the matter density parameter Ω_m (red-dashed), as a function of the redshift z , for Tsallis cosmology with $\delta = 1.1$ and $\alpha = 1$ in units where $H_0 = 1$, and for Myrzakulov gravity under the identifications mention in the text and for $\epsilon = \frac{6\alpha\delta}{(2-\delta)(3-2\delta)} = 9.2$, in units where $H_0 = 1$. Lower graph: The corresponding dark-energy equation-of-state parameter w_{DE} . In both graphs we have imposed $\Omega_m(z = 0) = \Omega_{m0} \approx 0.31$.

Additionally, in Fig. 2 we present the evolution of the dark-energy equation-of-state parameter for various values of the model parameters. Similarly to the previous figure, since w_{DE} is a background-related quantity the two models coincide.




Figure 2. The dark-energy equation-of-state parameter w_{DE} as a function of the redshift z , for Tsallis cosmology with $\alpha = 1$ in units where $H_0 = 1$, and various values of the parameter δ , and for Myrzakulov gravity for various parameters of $\epsilon = \frac{6\alpha\delta}{(2-\delta)(3-2\delta)}$. For each curve we set $\Omega_m(z = 0) = \Omega_{m0} \approx 0.31$ at present.

4.2 Perturbation evolution

Let us first review briefly the behavior of Tsallis cosmology at the perturbative level. We introduce the matter overdensity $\delta_m = \delta\rho_m/\rho_m$, we focus on dust matter and we set $\alpha = 1$ for convenience. Additionally, we assume that the dark-energy sector does not cluster. In this case we have [65]

$$\delta_m'' + \frac{2(4-2\delta) - (9-6\delta+8\pi G\Lambda H^{2\delta-4})}{(4-2\delta)(1+z)}\delta_m' + \frac{3^{\frac{1}{\delta-2}} \left[(1-2\delta)\rho_m^{\frac{1}{2-\delta}} - 9(1-\delta)\Lambda\rho_m^{\frac{\delta-1}{2-\delta}} \right] 8\pi G}{2(2-\delta)^2 H^2(1+z)^2}\delta_m = 0, \quad (4.5)$$

where primes denote derivatives with respect to the redshift. As one can see, we obtain a different friction term as well as an effective Newton constant comparing to standard cosmology. In the case $\delta = 1$ one obtains the standard result, which in the case of matter domination ($\Omega_m \approx 1$) yields

$$\delta_m'' + \frac{1}{2(1+z)}\delta_m' - \frac{3}{2(1+z)^2}\delta_m = 0, \quad (4.6)$$

as expected.

Let us now study the perturbations in Myrzakulov $F(R, Q)$ gravity. Although the full analysis lies beyond the scope of the present work, we can obtain approximate expressions working within the mini-superspace framework. We will focus on the model where we have obtained a correspondence at the background level, and we assume non-clustering dark energy. We work in the Newtonian gauge and we write the scalar perturbations of the metric as

$$ds^2 = -(1+2\Phi)dt^2 + a^2(1-2\Psi)\delta_{ij}dx^i dx^j. \quad (4.7)$$

Inserting this into (2.18), and imposing the aforementioned identifications that make the background behavior of the two theories to coincide, we find

$$\delta_m'' + \frac{2(4-2\delta) - (9-6\delta+8\pi G\Lambda H^{2\delta-4})}{(4-2\delta)(1+z)}\delta_m' + \frac{G \left(1 + \lambda \frac{w_a + w_{\dot{a}}}{2} + \frac{u_a + u_{\dot{a}}}{2} \right)}{(1+\lambda)(1+z)^2}\delta_m = 0. \quad (4.8)$$

As we observe, although the friction term between the two theories is the same, the last term, related to the effective Newton constant is different. This is an important result since it implies that the growth of density perturbations and the matter power spectrum will be different in the two theories, and therefore confronting them with perturbation-related datasets, such as σ_8 and weak-lensing ones, will lead to different results. Hence, such an analysis will be useful in order to distinguish the two theories.

Let us proceed to a numerical example. We solve (4.5) for Tsallis cosmology and (4.8) for Myrzakulov gravity, under the aforementioned identifications in which the two models coincide. Then, can calculate the quantity $f\sigma_8(z) = f(z)\sigma(z)$, with $f(z) := -\frac{d\ln\delta_m(z)}{d\ln z}$ and $\sigma(z) := \sigma_8 \frac{\delta_m(z)}{\delta_m(0)}$ [4].

In Fig. 3 we present the evolution of $f\sigma_8$ for Λ CDM scenario, as well as for Tsallis cosmology and Myrzakulov gravity, for the aforementioned identifications in which the two theories coincide at the background level. In this case the curves do not coincide anymore, and thus the behavior of perturbations can be used to distinguish the two theories.




Figure 3. The behavior of $f\sigma_8$ for Myrzakulov gravity (black solid) and Tsallis cosmology (red-dashed), under the aforementioned identifications and for $\delta = 1.1$ and $\alpha = 1$, and $\epsilon = \frac{6\alpha\delta}{(2-\delta)(3-2\delta)} = 9.2$, in units where $H_0 = 1$, in which the two theories coincide at the background level. The curves do not coincide and thus the behavior of perturbations can be used to distinguish the two theories.

5 Conclusions

Myrzakulov $F(R, Q)$ gravity is a modified gravitational framework that extends general relativity by incorporating both curvature and nonmetricity. By introducing an additional function of the Ricci scalar R and the nonmetricity scalar Q , this theory provides a richer geometrical structure that allows for deviations from Einstein's theory, and in particular in the context of cosmology. The inclusion of a non-special connection modifies the gravitational dynamics and affects the evolution of the universe, leading to novel phenomenological implications. Myrzakulov gravity has been studied in various contexts, including inflation, late-time acceleration, and structure formation, demonstrating its potential as an alternative to the standard cosmological model.

On the other hand, Tsallis cosmology emerges from the gravity-thermodynamics conjecture, where the modified Friedmann equations arise from the first law of thermodynamics when employing Tsallis entropy instead of the usual Bekenstein-Hawking entropy. The Tsallis entropy formalism introduces a non-additive generalization of standard thermodynamic entropy, characterized by a parameter δ that encapsulates deviations from extensive statistical mechanics. Applying this approach to the cosmological setting results in modified equations that can describe an evolving dark energy sector. Hence, it provides a thermodynamic interpretation of cosmic acceleration and can lead to interesting cosmological phenomenology.

In this work, we examined the correspondence between Myrzakulov $F(R, Q)$ gravity and Tsallis cosmology, focusing on their equivalence at the background level. By appropriately identifying the functional dependencies of the connection-related functions in Myrzakulov gravity, we reconstructed the effective dark energy sector of Tsallis cosmology. Our

analysis demonstrated that both frameworks can give identical background evolution for specific function and parameter choices, reproducing the standard cosmological sequence of matter and dark energy domination. However, despite the fact that the two theories coincide at the background level, their perturbation behavior exhibits differences. In particular, the growth of density fluctuations and the effective Newton constant deviate between the two scenarios, indicating that perturbative observables, such as structure formation and weak-lensing ones, could serve as distinguishing factors between them.

Future research directions could study several extensions of this correspondence. A more detailed investigation into the stability of perturbations in Myrzakulov gravity, particularly within the full metric-affine formalism, could offer more information into the viability of the theory. Additionally, observational constraints from large-scale structure data, such as galaxy clustering and weak-lensing surveys, could be employed to quantify deviations from standard predictions. The extension of this correspondence to include other entropy-based modifications, such as Barrow and Kaniadakis entropies, may also provide a broader understanding of the connection between gravity and thermodynamics. These projects are left for future investigation.

References

- [1] P. J. E. Peebles and B. Ratra, *Rev. Mod. Phys.* **75**, 559-606 (2003) [arXiv:astro-ph/0207347 [astro-ph]].
- [2] S. Weinberg, *Rev. Mod. Phys.* **61**, 1-23 (1989).
- [3] A. Addazi, *et al.* *Prog. Part. Nucl. Phys.* **125**, 103948 (2022) [arXiv:2111.05659 [hep-ph]].
- [4] E. Abdalla, *et al.* *JHEAp* **34**, 49-211 (2022) [arXiv:2203.06142 [astro-ph.CO]].
- [5] E. J. Copeland, M. Sami and S. Tsujikawa, *Int. J. Mod. Phys. D* **15**, 1753-1936 (2006) [arXiv:hep-th/0603057 [hep-th]].
- [6] Y. F. Cai, E. N. Saridakis, M. R. Setare and J. Q. Xia, *Phys. Rept.* **493**, 1-60 (2010) [arXiv:0909.2776 [hep-th]].
- [7] E. N. Saridakis *et al.* [CANTATA], Springer (2021), [arXiv:2105.12582 [gr-qc]].
- [8] S. Capozziello and M. De Laurentis, *Phys. Rept.* **509**, 167-321 (2011) [arXiv:1108.6266 [gr-qc]].
- [9] Y. F. Cai, S. Capozziello, M. De Laurentis and E. N. Saridakis, *Rept. Prog. Phys.* **79**, no.10, 106901 (2016) [arXiv:1511.07586 [gr-qc]].
- [10] A. De Felice and S. Tsujikawa, *Living Rev. Rel.* **13**, 3 (2010) [arXiv:1002.4928 [gr-qc]].
- [11] S. Nojiri and S. D. Odintsov, *Phys. Rept.* **505**, 59-144 (2011) [arXiv:1011.0544 [gr-qc]].
- [12] A. A. Starobinsky, *JETP Lett.* **86**, 157-163 (2007) [arXiv:0706.2041 [astro-ph]].
- [13] S. Nojiri and S. D. Odintsov, *Phys. Lett. B* **631**, 1-6 (2005) [arXiv:hep-th/0508049].
- [14] A. De Felice and S. Tsujikawa, *Phys. Lett. B* **675**, 1-8 (2009) [arXiv:0810.5712 [hep-th]].
- [15] G. B. Zhao, R. G. Crittenden, L. Pogosian and L. Samushia, *Phys. Rev. Lett.* **109**, 171301 (2012) [arXiv:1207.3804 [astro-ph.CO]].
- [16] M. F. Shamir, *Eur. Phys. J. C* **80**, no.12, 1102 (2020) [arXiv:2012.00558 [gr-qc]].

- [17] G. R. Bengochea and R. Ferraro, Phys. Rev. D **79**, 124019 (2009) [arXiv:0812.1205 [astro-ph]].
- [18] E. V. Linder, Phys. Rev. D **81**, 127301 (2010) [arXiv:1005.3039 [astro-ph.CO]].
- [19] S. H. Chen, J. B. Dent, S. Dutta and E. N. Saridakis, Phys. Rev. D **83**, 023508 (2011) [arXiv:1008.1250 [astro-ph.CO]].
- [20] N. Tamanini and C. G. Boehmer, Phys. Rev. D **86**, 044009 (2012) [arXiv:1204.4593 [gr-qc]].
- [21] G. R. Bengochea, Phys. Lett. B **695**, 405-411 (2011) [arXiv:1008.3188 [astro-ph.CO]].
- [22] D. Liu and M. J. Reboucas, Phys. Rev. D **86**, 083515 (2012) [arXiv:1207.1503 [astro-ph.CO]].
- [23] M. H. Daouda, M. E. Rodrigues and M. J. S. Houndjo, Eur. Phys. J. C **72**, 1890 (2012) [arXiv:1202.1147 [gr-qc]].
- [24] G. Kofinas and E. N. Saridakis, Phys. Rev. D **90**, no.8, 084044 (2014) [arXiv:1404.2249 [gr-qc]].
- [25] S. Bahamonde, C. G. Böhrer and M. Wright, Phys. Rev. D **92**, no.10, 104042 (2015) [arXiv:1508.05120 [gr-qc]].
- [26] J. Beltrán Jiménez, L. Heisenberg and T. Koivisto, Phys. Rev. D **98**, no.4, 044048 (2018) [arXiv:1710.03116 [gr-qc]].
- [27] L. Heisenberg, Phys. Rept. **1066**, 1-78 (2024) [arXiv:2309.15958 [gr-qc]].
- [28] A. De, T. H. Loo and E. N. Saridakis, JCAP **03**, 050 (2024) [arXiv:2308.00652 [gr-qc]].
- [29] R. Myrzakulov, Eur. Phys. J. C **72**, 2203 (2012) [arXiv:1207.1039 [gr-qc]].
- [30] E. N. Saridakis, S. Myrzakul, K. Myrzakulov and K. Yerzhanov, Phys. Rev. D **102**, no.2, 023525 (2020) [arXiv:1912.03882 [gr-qc]].
- [31] F. K. Anagnostopoulos, S. Basilakos and E. N. Saridakis, Phys. Rev. D **103**, no.10, 104013 (2021) [arXiv:2012.06524 [gr-qc]].
- [32] T. Myrzakul, K. Yesmakhanova, N. Myrzakulov, S. Myrzakul, K. Myrzakulov, K. Yerzhanov, R. Myrzakulov and G. Nugmanova, [arXiv:2101.05318 [gr-qc]].
- [33] D. Iosifidis, N. Myrzakulov and R. Myrzakulov, Universe **7**, no.8, 262 (2021) [arXiv:2106.05083 [gr-qc]].
- [34] N. Myrzakulov, R. Myrzakulov and L. Ravera, Symmetry **13**, no.10, 1855 (2021) [arXiv:2108.00957 [gr-qc]].
- [35] T. Harko, N. Myrzakulov, R. Myrzakulov and S. Shahidi, Phys. Dark Univ. **34**, 100886 (2021) [arXiv:2110.00358 [gr-qc]].
- [36] G. Papagiannopoulos, S. Basilakos and E. N. Saridakis, Phys. Rev. D **106**, no.10, 103512 (2022) [arXiv:2202.10871 [gr-qc]].
- [37] S. Kazempour and A. R. Akbarieh, Astropart. Phys. **165**, 103060 (2025) [arXiv:2309.09230 [gr-qc]].
- [38] E. H. M. Zahran, A. Bekir, R. A. Ibrahim and R. Myrzakulov, AIMS Math. **9**, no.3, 6145-6160 (2024)
- [39] D. C. Maurya and R. Myrzakulov, Eur. Phys. J. C **84**, no.5, 534 (2024) [arXiv:2401.00686 [gr-qc]].

- [40] D. C. Maurya and R. Myrzakulov, *Eur. Phys. J. C* **84**, no.6, 625 (2024) [arXiv:2402.02123 [gr-qc]].
- [41] D. C. Maurya, K. Yesmakhanova, R. Myrzakulov and G. Nugmanova, *Chin. Phys. C* **48**, no.12, 125101 (2024) [arXiv:2403.11604 [gr-qc]].
- [42] D. C. Maurya, K. Yesmakhanova, R. Myrzakulov and G. Nugmanova, *Phys. Scripta* **99**, no.10, 105014 (2024) [arXiv:2404.09698 [gr-qc]].
- [43] D. Momeni and R. Myrzakulov, [arXiv:2412.04524 [gr-qc]].
- [44] A. Lymperis, G. Nugmanova and A. Sergazina, [arXiv:2501.18524 [gr-qc]].
- [45] T. Jacobson, *Phys. Rev. Lett.* **75**, 1260-1263 (1995) [arXiv:gr-qc/9504004 [gr-qc]].
- [46] T. Padmanabhan, *Phys. Rept.* **406**, 49-125 (2005) [arXiv:gr-qc/0311036 [gr-qc]].
- [47] T. Padmanabhan, *Rept. Prog. Phys.* **73**, 046901 (2010) [arXiv:0911.5004 [gr-qc]].
- [48] A. V. Frolov and L. Kofman, *JCAP* **05**, 009 (2003) [arXiv:hep-th/0212327 [hep-th]].
- [49] R. G. Cai and S. P. Kim, *JHEP* **02**, 050 (2005) [arXiv:hep-th/0501055 [hep-th]].
- [50] M. Akbar and R. G. Cai, *Phys. Rev. D* **75**, 084003 (2007) [arXiv:hep-th/0609128 [hep-th]].
- [51] A. Paranjape, S. Sarkar and T. Padmanabhan, *Phys. Rev. D* **74**, 104015 (2006) [arXiv:hep-th/0607240 [hep-th]].
- [52] M. Akbar and R. G. Cai, *Phys. Lett. B* **635**, 7-10 (2006) [arXiv:hep-th/0602156 [hep-th]].
- [53] A. Sheykhi, B. Wang and R. G. Cai, *Nucl. Phys. B* **779**, 1-12 (2007) [arXiv:hep-th/0701198 [hep-th]].
- [54] R. G. Cai and N. Ohta, *Phys. Rev. D* **81**, 084061 (2010) [arXiv:0910.2307 [hep-th]].
- [55] M. Wang, J. Jing, C. Ding and S. Chen, *Phys. Rev. D* **81**, 083006 (2010) [arXiv:0912.4832 [gr-qc]].
- [56] Y. Gim, W. Kim and S. H. Yi, *JHEP* **07**, 002 (2014) [arXiv:1403.4704 [hep-th]].
- [57] Z. Y. Fan and H. Lu, *Phys. Rev. D* **91**, no.6, 064009 (2015) [arXiv:1501.00006 [hep-th]].
- [58] S. A. Tsilioukas, N. Petropoulos and E. N. Saridakis, [arXiv:2412.21146 [gr-qc]].
- [59] A. Lymperis and E. N. Saridakis, *Eur. Phys. J. C* **78**, no.12, 993 (2018) [arXiv:1806.04614 [gr-qc]].
- [60] C. Arias, F. Diaz, R. Olea and P. Sundell, *JHEP* **04**, 124 (2020) [arXiv:1906.05310 [hep-th]].
- [61] S. Nojiri, S. D. Odintsov and E. N. Saridakis, *Eur. Phys. J. C* **79**, no.3, 242 (2019) [arXiv:1903.03098 [gr-qc]].
- [62] C. Q. Geng, Y. T. Hsu, J. R. Lu and L. Yin, *Eur. Phys. J. C* **80**, no.1, 21 (2020) [arXiv:1911.06046 [astro-ph.CO]].
- [63] E. N. Saridakis, *JCAP* **07**, 031 (2020) [arXiv:2006.01105 [gr-qc]].
- [64] A. Lymperis, S. Basilakos and E. N. Saridakis, *Eur. Phys. J. C* **81**, no.11, 1037 (2021) [arXiv:2108.12366 [gr-qc]].
- [65] A. Sheykhi and B. Farsi, *Eur. Phys. J. C* **82**, no.12, 1111 (2022) [arXiv:2205.04138 [gr-qc]].
- [66] S. Di Gennaro and Y. C. Ong, *Universe* **8**, no.10, 541 (2022) [arXiv:2205.09311 [gr-qc]].
- [67] G. G. Luciano and J. Gine, *Phys. Lett. B* **833**, 137352 (2022) [arXiv:2204.02723 [gr-qc]].

- [68] Z. Çoker, Ö. Ökcü and E. Aydiner, EPL **143**, no.5, 59001 (2023) [arXiv:2308.10212 [gr-qc]].
- [69] S. Basilakos, A. Lymperis, M. Petronikolou and E. N. Saridakis, Eur. Phys. J. C **84**, no.3, 297 (2024) [arXiv:2308.01200 [gr-qc]].
- [70] A. Salehi, [arXiv:2309.15956 [gr-qc]].
- [71] P. Jizba and G. Lambiase, Entropy **25**, no.11, 1495 (2023) [arXiv:2310.19045 [gr-qc]].
- [72] P. Jizba, G. Lambiase, G. G. Luciano and L. Mastrototaro, Eur. Phys. J. C **84**, no.10, 1076 (2024) [arXiv:2403.09797 [gr-qc]].
- [73] E. Ebrahimi and A. Sheykhi, Phys. Dark Univ. **45**, 101518 (2024) [arXiv:2405.13096 [gr-qc]].
- [74] U. K. Tyagi, S. Haridasu and S. Basak, Phys. Rev. D **110**, no.6, 063503 (2024) [arXiv:2406.07446 [astro-ph.CO]].
- [75] Ö. Ökcü and E. Aydiner, Gen. Rel. Grav. **56**, no.7, 87 (2024) [arXiv:2407.14685 [gr-qc]].
- [76] M. Petronikolou, S. Basilakos, A. Lymperis and E. Saridakis, PoS **CORFU2023**, 174 (2024)
- [77] M. L. Mendoza-Martínez, A. Cervantes-Contreras, J. J. Trejo-Alonso and A. Hernandez-Almada, Eur. Phys. J. C **84**, no.7, 717 (2024) [arXiv:2404.18346 [astro-ph.CO]].
- [78] M. Yarahmadi and A. Salehi, Eur. Phys. J. C **84**, no.4, 443 (2024) [arXiv:2501.07860 [astro-ph.CO]].
- [79] S. A. Tsilioukas, M. Petronikolou, S. Basilakos and E. N. Saridakis, [arXiv:2501.15927 [gr-qc]].
- [80] S. Capozziello, V. De Falco and C. Ferrara, Eur. Phys. J. C **82**, no.10, 865 (2022) [arXiv:2208.03011 [gr-qc]].
- [81] K. Shimada, K. Aoki and K. i. Maeda, Phys. Rev. D **99**, no.10, 104020 (2019) [arXiv:1812.03420 [gr-qc]].
- [82] A. Paliathanasis, Class. Quant. Grav. **33**, no.7, 075012 (2016) [arXiv:1512.03239 [gr-qc]].
- [83] C. Tsallis, J. Statist. Phys. **52**, 479-487 (1988)
- [84] M. L. Lyra and C. Tsallis, Phys. Rev. Lett. **80**, 53-56 (1998) [arXiv:cond-mat/9709226 [cond-mat.stat-mech]].
- [85] C. Tsallis, R. S. Mendes and A. R. Plastino, Physica A **261**, 534 (1998)
- [86] G. Wilk and Z. Włodarczyk, Phys. Rev. Lett. **84**, 2770 (2000) [arXiv:hep-ph/9908459 [hep-ph]].
- [87] C. Tsallis and L. J. L. Cirto, Eur. Phys. J. C **73**, 2487 (2013) [arXiv:1202.2154 [cond-mat.stat-mech]].
- [88] R. G. Cai, L. M. Cao and Y. P. Hu, Class. Quant. Grav. **26**, 155018 (2009) [arXiv:0809.1554 [hep-th]].
- [89] G. W. Gibbons and S. W. Hawking, Phys. Rev. D **15**, 2738-2751 (1977).
- [90] N. Aghanim *et al.* [Planck Collaboration], Astron. Astrophys. **641**, A6 (2020) [arXiv:1807.06209 [astro-ph.CO]].