A-PRIORI ESTIMATES FOR GENERALIZED KORTEWEG-DE VRIES EQUATIONS IN $H^{-1}(\mathbb{R})$

MIHAELA IFRIM AND THIERRY LAURENS

ABSTRACT. We prove local-in-time a-priori estimates in $H^{-1}(\mathbb{R})$ for a family of generalized Korteweg–de Vries equations. This is the first estimate for any non-integrable perturbation of the KdV equation that matches the regularity of the sharp well-posedness theory for KdV. In particular, we show that our analysis applies to models for long waves in a shallow channel of water with an uneven bottom.

The proof of our main result is based upon a bootstrap argument for the renormalized perturbation determinant coupled with a local smoothing norm.

Contents

1. Introduction	1
1.1. Acknowledgments	6
2. Preliminaries	7
3. The local smoothing norm	10
4. The local smoothing estimate	21
5. The a-priori estimates	27
References	30

1. INTRODUCTION

The Korteweg–de Vries equation

(KdV) $\partial_t u = -u'' + 6uu'$

(where $u' = \partial_x u$) was originally derived as a model for the propagation of waves in a shallow channel of water [42]. However, over the next century, (KdV) has proved to be a fundamental model for nonlinear dispersive waves, with applications to a wide variety of physical systems spanning many fields of science (see, for example, [14]).

Mathematically, (KdV) has also played a central role in our understanding of dispersive equations. In particular, the (KdV) equation was the first discovered example of a completely integrable PDE, a feature that has been heavily exploited in some works. This discovery sparked a pursuit into the well-posedness and dispersive behavior for this system over the next 50 years, and generated a long list of cutting-edge techniques [2, 5, 9, 12, 25, 27, 34, 36, 37, 40, 41, 59, 60, 62], some of which rely on complete integrability, and some of which do not. On the integrable side this effort culminated in global well-posedness for initial data in H^{-1} on the line and the circle [33, 38], a result that is sharp in the class of H^s spaces for both geometries [50, 51]. Although (KdV) lied at the center of this effort, it also served as a testing ground for the introduction of new tools and innovative techniques. After their introduction, many of these methods (both integrable and non-integrable) were later adapted to broader classes of dispersive equations. However, this important step has yet to be achieved for the completely integrable methods used to prove the sharp well-posedness results [33, 38].

One physically important class of examples are the KdV equations with variable bottom. As (KdV) has proved to be an effective model for the propagation of waves in a shallow channel of water, many authors have introduced generalizations to describe an uneven bottom [15, 22, 29, 31, 44, 49, 54, 55, 61, 63, 64]. For concreteness, consider the model

(1.1)
$$\partial_t u = -b^5 u''' + 6uu' - 4bu' - 6b'u,$$

which was derived and rigorously justified in [29]. Here, b(x) is defined in terms of a function c(x) which describes the bottom of the channel:

$$b(x) := \sqrt{1 - c(x)}.$$

When c(x) is a small Schwartz function, the equation (gKdV) closely resembles (KdV). Nevertheless, our understanding of the well-posedness problem is comparatively lacking.

After a change of variables (see (5.15) below), the equation (1.1) can be put in the form

(gKdV)
$$\partial_t u = -u''' + 6uu' + (a_1u')' + a_2u^2 + a_3u' + a_4u$$

for certain coefficients a_j . In this work, we will assume that the coefficients $a_j(t, x)$ are given functions that are sufficiently regular and localized in space, and study the corresponding solutions $u : [-T, T] \times \mathbb{R} \to \mathbb{R}$.

Our main contribution is the following local-in-time a-priori estimate in H^{-1} :

Theorem 1.1 (A-priori estimate). There exists $\epsilon > 0$ so that, if the coefficients are given smooth functions that satisfy the decay bounds uniformly for $|t| \leq T$ and $x \in \mathbb{R}$:

(1.2)
$$|a_j(t,x)| + |\partial_x a_j(t,x)| \le \epsilon (1+x^2)^{-1}$$
 for $j = 1, 2, 3,$

(1.3)
$$|a_4(t,x)| + |\partial_x a_4(t,x)| \lesssim (1+x^2)^{-1},$$

then for any A > 0 there exist constants C, T > 0 so that for any smooth solution u to (gKdV) in [-T, T] whose initial data satisfies

(1.4)
$$||u(0)||_{H^{-1}} \le A$$
,

satisfies the uniform bound

(1.5)
$$\sup_{|t| \le T} \|u(t)\|_{H^{-1}} \le C$$

Remark 1.2. The lifespan T for the bounds in the Theorem is more accurately identified in the proof as

$$T \gtrsim (1+A)^{-4}.$$

On the other hand one might expect that C should be comparable to A; however this is not the case because in the proof of the Theorem we work with a weighted form of the H^{-1} norm. So instead we get the weaker bound

$$C \lesssim A(1+A)^2.$$

Remark 1.3. The assumption that the coefficients a_j are smooth in the above theorem is purely qualitative, and only serves to ensure that we can talk about smooth solutions to (gKdV). Alternatively, if we want to only assume that the coefficients satisfy (1.2) and (1.3), then the conclusion of the theorem would remain valid for rougher solutions, e.g. in L^2 , once local well-posedness is known.

We do not claim that the conditions (1.2)-(1.3) in this theorem are sharp. In fact, our proof will require slightly weaker hypotheses (see (4.1)-(4.4) below for details). The main thrust of this work is to match the sharp regularity of the well-posedness theory for (KdV), and so any hypotheses on the coefficients that work constitutes significant progress. In particular, (1.5) rules out strong forms of instantaneous norm inflation in H^{-1} .

Applying our general result to the equation (1.1), we obtain:

Corollary 1.4. There exists a constant $\epsilon > 0$ so that, if $c : \mathbb{R} \to \mathbb{R}$ is a smooth function that satisfies the small pointwise bounds

$$|\partial_x^j c(x)| \le \epsilon (1+x^2)^{-1}$$
 for $j = 0, 1, \dots, 4$,

then for any A > 0 there exist constants $\epsilon, T > 0$ so that for any smooth solution u to (1.1) in [-T, T] whose initial data satisfies

$$||u(0)||_{H^{-1}} \le A,$$

satisfies the uniform bound

$$\sup_{|t| \le T} \|u(t)\|_{H^{-1}} \le C.$$

We contend that for each of the numerous applications of (KdV), our a-priori estimate (1.5) could also be employed to describe small physical imperfections using similar methods.

Another main way in which (gKdV) arises naturally is in the study of localized perturbations of a background solution to (KdV): if we take our solution u = q + V to be a given background wave V plus a perturbation q, then the equation for q often takes the form (gKdV). For example, if V(t, x) solves (KdV) then q solves

(1.6)
$$\partial_t q = -q''' + 6qq' + 6Vq' + 6V'q,$$

and if V(x) is a fixed background profile then a forcing term -V''' + 6VV' is added to the RHS above. Through this lens, the special case (1.6) of gKdV equations have been of great interest in the literature.

The first phase of results for (1.6) addressed the construction of solutions using the inverse scattering transform. Two particularly common choices for V are profiles that are periodic [18–20, 43], which describe localized defects to periodic wave trains, and step-like [8, 10, 11, 32], which arise in the study of bore propagation and rarefaction waves. Later, the authors of [16, 17] established a general framework that encompasses both of these cases, and even a mixture of the two as $x \to +\infty$ and $x \to -\infty$. Although the main thrust is to prove existence, a key step in many of these works is establishing persistence of regularity for solutions, much like (1.5).

While many of these results work at high regularity, these methods for existence have been adapted to classes of one-sided step-like initial data [24, 57, 58] and even to one-sided step-like elements of $H_{\text{loc}}^{-1}(\mathbb{R})$ [23]. Despite the lack of assumptions as $x \to -\infty$ (the direction in which radiation propagates), these low-regularity results require rapid decay at $x \to +\infty$ and global boundedness from below.

An alternative approach was introduced in [48], which proves global existence and uniqueness for initial data that satisfies u(0, x) = o(|x|) as $x \to \pm \infty$. Here, the background wave V is evolved according to the inviscid Burger's equation $\partial_t V = -6VV'$ using the method of characteristics, and then solutions to the equation for q are constructed using a family of discretized approximate equations. Existence and uniqueness for q is then established in a weighted H^3 space.

Inspired by the theory of tidal bores and rarefaction waves (as well as kink solutions to KdV with higher-power nonlinearities), authors began turning their attention to the well-posedness problem for step-like initial data. The first phase of results employed BBM and parabolic regularizations [3, 28, 65], which involve adding a term to the LHS of (1.6) that eases the proof of well-posedness $(-\varepsilon \partial_t \partial_x^2 q \text{ and } -\varepsilon \partial_x^2 q \text{ for } \varepsilon > 0$, respectively) and then sending $\varepsilon \to 0$. The former was introduced by Bona and Smith [4] in the case $V \equiv 0$ and leads to the Benjamin–Bona–Mahony equation, for which the method is named. These approaches culminated in local well-posedness for initial data $q \in H^s$ for $s > \frac{3}{2}$, and was later advanced to s > 1 in [21] through the incorporation of Strichartz estimates.

Recently, the work [53] extends local well-posedness to the range $s > \frac{1}{2}$ using a synthesis of much more modern tools for well-posedness. In addition to the cases of step-like and periodic background waves V, this result also applies to more general choices of V with bounded asymptotics as $x \to \pm \infty$, and even more general nonlinearities in (KdV). Nevertheless, it only applies to (gKdV) in the special case where $a_1 \equiv 0$ and a_2 is constant.

Around the same time, the second author proved that (KdV) is globally well-posed for initial perturbations $q \in H^{-1}$ in [46], provided that the background wave V is a suitable solution to (KdV). These conditions on V do include regularity but do not impose any assumptions on spatial asymptotics. In particular, this result applies to the important cases of smooth periodic [46] and step-like [45] initial data.

All of these works for (1.6) focus on the coefficients a_3 and a_4 in (gKdV). The introduction of the coefficient a_1 already significantly changes the analysis. The local well-posedness of this equation for initial data in H^s with $s > \frac{3}{2}$ and suitable coefficients was demonstrated in [30] using energy methods. Recently, local well-posedness was extended to the range $s > \frac{1}{2}$ in [52].

In the most general case, the optimal well-posedness results that cover the a_2 term in (gKdV) are [1, 13, 26] to the best of our knowledge. These works establish well-posedness for initial data at a considerably high regularity, but they apply to very general families of nonlinear equations with a KdV-type dispersion.

Let us now turn to our methods for proving the priori estimate (1.5). At the center of our analysis lies the renormalized perturbation determinant $\alpha(\kappa, u)$ (see (2.9) below for details). The perturbation determinant is a conserved quantity for (KdV) originating from scattering theory, and is rooted in the complete integrability of this system. However, the authors of [39] (and independently [56]) discovered a renormalization α of this quantity that satisfies

(1.7)
$$\kappa \alpha(\kappa, u) \approx \int \frac{|\widehat{u}(\xi)|^2}{\xi^2 + 4\kappa^2} d\xi =: \|u\|_{H_{\kappa}^{-1}}^2 \text{ for all } \kappa \gg \|u\|_{H^{-1}}^2$$

and is still conserved, and used this to prove a global-in-time a-priori estimate for solutions to (KdV) in H^s spaces for $-1 \le s < 1$. At the same time, conserved energies in H^s spaces

were independently constructed in [41] for a full range of Sobolev exponents $s \ge -1$. As it turned out, these energies are also connected to the perturbation determinant, but with a different choice of renormalization. See also the earlier work [47] where similar bounds were proved at slightly higher regularity $s > -\frac{4}{5}$.

The same quantity α was then used as a starting point in [38] to prove that (KdV) is well-posed in H^{-1} using the method of commuting flows.

Of course, introducing the coefficients a_j in (gKdV) breaks the conservation of α , along with all of the other conservation laws of (KdV). In general, the value of α can grow in time, and consequently we cannot expect a global-in-time estimate. Instead, in order to establish (1.5), we prove that α must remain finite for a short period of time using a bootstrap/continuity argument.

However, we immediately encounter an obstacle. As an illustrative example, consider the L^2 norm; if u is a Schwartz solution of (KdV), then the L^2 norm of u(t) is constant in time. On the other hand, if u solves (gKdV), then the L^2 norm evolves according to

(1.8)
$$\partial_t \int \frac{1}{2} u^2 \, dx = \int -a_1 (\partial_x u)^2 + a_2 u^3 - \frac{1}{2} (\partial_x a_3) u^2 + a_4 u^2 \, dx.$$

How can we bound the right-hand side solely in terms of $||u||_{L^2}$ in order to close the argument?

Our strategy in this paper is to use local smoothing: due to the dispersive nature of the equation, we expect a gain in the regularity of solutions locally in space on average in time. For (KdV), this effect was discovered by Kato [35], who proved that L^2 solutions are in fact in H^1 locally in space at almost every time. In particular, this gives us a way to make sense of the RHS of (1.8) even for L^2 solutions u.

For the a-priori estimate (1.5), we encounter an analogous loss of derivatives phenomenon in computing the time evolution of α . The local smoothing effect attendant to this problem is to show that an H^{-1} solution of (gKdV) is in L^2 locally in space and time:

Theorem 1.5 (Local smoothing estimate). There exists $\epsilon > 0$ so that, if the smooth coefficients a_j satisfy the bounds in (1.2)–(1.3) uniformly for $|t| \leq T$ and $x \in \mathbb{R}$, then for any A > 0 there exist constants C, T > 0 so that for any smooth solution u to (gKdV) in [-T, T] whose initial data satisfies

(1.9)
$$||u(0)||_{H^{-1}} \le A,$$

satisfies the local energy bound

(1.10)
$$\sup_{x_0 \in \mathbb{R}} \int_{-T}^{T} \int_{x_0 - 1}^{x_0 + 1} |u(t, x)|^2 \, dx \, dt \le C.$$

We will prove (1.5) and (1.10) simultaneously, using a bootstrap argument in terms of both α and a local smoothing norm.

In the case where the coefficients $a_j \equiv 0$ vanish, an analogous local smoothing estimate was first proved in [7]; see also the earlier work in [47], where similar local smoothing estimates are proved for H^s solutions with $s > -\frac{4}{5}$. However, our proof is more closely related to the alternative approach in [38, Th. 1.2] using the renormalized perturbation determinant. This argument is made possible by the discovery of a density $\rho(x; \kappa, u)$ for α (see (2.9) below) not known during the earlier work [39]. Moreover, if u solves (KdV), then ρ satisfies the density-flux equation (or 'microscopic conservation law')

(1.11)
$$\partial_t \rho + \partial_x j = 0$$

for a certain current $j(x; \kappa, u)$ (see (2.12)). Integrating this equation in space yields the conservation of α . Alternatively, first multiplying by a smooth step function and then integrating in space leads to a local smoothing estimate; this is basis of the short proof presented in [38].

Turning our attention back to (gKdV), we again encounter obstacles. The presence of the coefficients a_i breaks the conservation law (1.11), and instead we have

(1.12)
$$\partial_t \rho + \partial_x j = d\rho|_u [(a_1 u')' + a_2 u^2 + a_3 u' + a_4 u].$$

(Here, $d\rho|_u$ denotes the functional derivative of ρ at u; see (2.14) for details.) All of terms on the RHS of (1.12) are new, and must that must be controlled in terms of α and our local smoothing norm.

Many of our estimates are in the spirit of [6], which proves that the fifth-order KdV equation is globally well-posed in H^{-1} . In order to prove their result, the authors had to prove an analogous local smoothing estimate for H^{-1} solutions of their system and a family of approximate equations. Together, the estimates in [6, 38] provide upper bounds on the density ρ and current j in terms of the local smoothing and H^{-1} norms, the latter of which can then be controlled by α using (1.7). For example, when $\kappa \geq 1$ we may bound

(1.13)
$$\|u\|_{H_{\kappa}^{-1}}^{2} = \int \frac{|\widehat{u}(\xi)|^{2}}{\xi^{2} + 4\kappa^{2}} d\xi \leq \|u\|_{H^{-1}}^{2}$$

and use this to construct α for all $||u||_{H^{-1}}^2 \ll 1$. For the (KdV) and fifth-order KdV equations, this small-data assumption does not pose a problem; one can use the scaling symmetry to make the initial data arbitrarily small, and then the exact conservation of α implies that solutions remain small globally in time.

By comparison, (gKdV) does not posses a scaling symmetry in general, and we expect that the H^{-1} norm of solutions can be growing in time without bound. This in turn forces us to choose $\kappa \gg ||u||_{H^{-1}}^2$ large, which introduces a large implicit constant in (1.13). Consequently, the estimates in [6, 38] are insufficient, even in estimating the LHS of (1.12). In order to close our bootstrap argument, we must establish new estimates for the density ρ and current j using the H_{κ}^{-1} norm, rather than H^{-1} , as this is the quantity whose growth we can efficiently estimate using (1.7). In comparison to the estimate (1.13) used throughout [6, 38], this requires a much more careful estimation of high frequency contributions.

This paper is organized as follows. We begin in Section 2 by reviewing the material developed in [6, 38] that we will need, including the key players α , ρ , and j mentioned previously as well as the estimates they satisfy. We then proceed in Section 3 by introducing our local smoothing norm and developing the new estimates required for our bootstrap argument.

We employ these ingredients in Sections 4 and 5 to prove more precise versions of our local smoothing and a-priori estimates, Theorem 4.1 and Proposition 5.1, respectively. We then combine these two estimates with a bootstrap argument, and present the full result in Theorem 5.2. Finally, we conclude Section 5 by describing how the statements of Theorems 1.1 and 1.5 and Corollary 1.4 represent special cases of Theorem 5.2.

1.1. Acknowledgments. While working on this project the first author was supported by the CAREER grant DMS-1845037, by the NSF grant DMS-2348908, by the Miller Foundation and by a Simons Fellowship.

2. Preliminaries

Here we recall some of the key objects from [38] which are used in order to prove that (KdV) is well-posed in H^{-1} , and which will serve us well in obtaining the a-priori bounds for our nonintegrable model (gKdV) displayed in Theorem 1.1 and Theorem 1.5. Our starting point is represented by the self-adjoint Lax operator associated to the KdV flow,

$$L_u = -\partial_x^2 + u.$$

Formally, the spectrum of L_u is conserved along the KdV flow. More precisely, $L_{u(t)}$ and $L_{u(s)}$ are unitarily equivalent operators. In general L_u is not invertible, which is one reason it is convenient to replace it with $L_u + \kappa^2$; given $u \in H^{-1}$ the operator $L_u + \kappa^2$ is invertible if $\kappa > 0$ is large enough. One may construct conserved quantities for KdV by looking at the trace of functions of $L_u + \kappa^2$, and in particular the trace of its renormalized logarithm, which can be also viewed as a function of κ . As shown in [38], this trace is closely related to the diagonal of the kernel of $L_u + \kappa^2$.

The first step in the analysis is to construct the diagonal of the integral kernel G(x, y) for the resolvent

(2.1)
$$R_u := (-\partial_x^2 + u + \kappa^2)^{-1}$$

of the Lax operator associated to a state u. With this operator in mind, it will be convenient to work with the norms

$$\|f\|_{H^s_{\kappa}}^2 := \int |\widehat{f}(\xi)|^2 (\xi^2 + 4\kappa^2)^s \, d\xi$$

where our convention for the Fourier transform is

$$\widehat{f}(\xi) = \frac{1}{\sqrt{2\pi}} \int e^{-i\xi x} f(x) \, dx \quad \text{so that} \quad \langle f, g \rangle = \int \overline{f}(x) g(x) \, dx = \langle \widehat{f}, \widehat{g} \rangle.$$

In the definition of H^s_{κ} , the constant 4 is simply included in order to make (2.4) an equality. One should see these spaces as versions of the traditional inhomogeneous Sobolev spaces H^s but adapted to the frequency scale κ rather than the unit frequency. Topologically these are the same as the classical Sobolev spaces H^s , but with the implicit constants in the norm equivalence depending on κ . In particular, we have the elementary estimates

$$\|fg\|_{H^{\pm 1}_{\kappa}} \lesssim \|f\|_{H^{1}} \, \|g\|_{H^{\pm 1}_{\kappa}} \quad \text{and} \quad \|fg\|_{H^{\pm 1}_{\kappa}} \lesssim \|f\|_{W^{1,\infty}} \, \|g\|_{H^{\pm 1}_{\kappa}}$$

uniformly for $\kappa \geq 1$. Notice that the results for H_{κ}^{-1} follow from those for H_{κ}^{1} by duality. We will also frequently use the embedding

$$\|f\|_{L^{\infty}} \lesssim \kappa^{-\frac{1}{2}} \|f\|_{H^{1}_{\kappa}}$$
 uniformly for $\kappa \ge 1$,

which implies the algebra bound

$$\|fg\|_{H^1_{\kappa}} \lesssim \kappa^{-\frac{1}{2}} \|f\|_{H^1_{\kappa}} \|g\|_{H^1_{\kappa}} \quad \text{uniformly for } \kappa \ge 1.$$

The resolvent R_0 associated to $u \equiv 0$,

(2.2)
$$R_0(\kappa) := (-\partial_x^2 + \kappa^2)^{-1},$$

plays an important role in what follows, and in particular has the mapping property

$$R_0: H^s_\kappa \to H^{s+2}_\kappa, \quad s \in \mathbb{R}.$$

This property is also shared by R_u for $u \in H^{-1}$ for κ large enough, but only for restricted range of s. One can formally use R_0 to obtain an expansion for R_u :

(2.3)
$$R_u = \sum_{\ell=0}^{\infty} (-1)^{\ell} (R_0 u)^{\ell} R_0,$$

which is justified for $u \in H^{-1}$ for κ large enough.

Our first bound involving R_0 is as follows:

Lemma 2.1 ([39]). For $\kappa > 0$, we have

(2.4)
$$\|\sqrt{R_0}f\sqrt{R_0}\|_{\mathfrak{I}_2} = \kappa^{-\frac{1}{2}} \|f\|_{H_{\kappa}^{-1}}$$

Here, \mathfrak{I}_2 denotes the class of Hilbert–Schmidt operators on $L^2(\mathbb{R})$. Such operators are automatically continuous due to the inequality

$$\|A\|_{\mathfrak{I}_{\infty}} \le \|A\|_{\mathfrak{I}_{2}} = \sqrt{\operatorname{tr}\{A^{*}A\}},$$

where \mathfrak{I}_{∞} denotes the operator norm. Moreover, the space of such operators forms a twosided ideal in the space of bounded operators:

$$\|ABC\|_{\mathfrak{I}_{2}} \leq \|A\|_{\mathfrak{I}_{\infty}} \|B\|_{\mathfrak{I}_{2}} \|C\|_{\mathfrak{I}_{\infty}}$$

Lastly, the product of two Hilbert–Schmidt operators is trace class, and we have

$$|\operatorname{tr}(AB)| \le ||AB||_{\mathfrak{I}_1} \le ||A||_{\mathfrak{I}_2} ||B||_{\mathfrak{I}_2}$$

Here, \mathfrak{I}_1 denotes the trace class; this consists of operators whose singular values are ℓ^1 -summable. In this way, the second inequality above is simply an application of the Cauchy–Schwarz inequality.

The Green's function is the integral kernel for the resolvent of the Lax operator R_u , which may be expressed as

$$G(x,y) = \langle \delta_x, (-\partial_x^2 + u + \kappa^2)^{-1} \delta_y \rangle$$

As explained earlier, its restriction to the diagonal plays a key role in our analysis.

Proposition 2.2 (Diagonal Green's function [38]). There exists a constant C > 0 so that the following statements are true for any R > 0:

(1) For each $||u||_{H^{-1}_{\epsilon}} \leq R$, the diagonal Green's function

$$g(x;\kappa,u) := \langle \delta_x, (-\partial_x^2 + u + \kappa^2)^{-1} \delta_x \rangle$$

exists for all $x \in \mathbb{R}$ and $\kappa \ge 1 + CR^2$.

(2) The mappings

(2.5)
$$u \mapsto g - \frac{1}{2\kappa} \quad and \quad u \mapsto \frac{1}{g} - 2\kappa$$

are real-analytic functionals from $\{u : ||u||_{H^{-1}} \leq R\}$ into H^1 for all $\kappa \geq 1 + CR^2$. (3) We have the estimates

(2.6)
$$||g(\kappa, u) - \frac{1}{2\kappa}||_{H^1_{\kappa}} \lesssim \kappa^{-1} ||u||_{H^{-1}_{\kappa}},$$

(2.7)
$$\left\|\frac{1}{g(\kappa,u)} - 2\kappa\right\|_{H^1_{\kappa}} \lesssim \kappa \|u\|_{H^{-1}_{\kappa}}$$

uniformly for $\kappa \geq 1 + CR^2$.

As a consequence of the expansion of R_u in (2.3), for g we have a similar expansion

(2.8)
$$g = \frac{1}{2\kappa} + h_1 + h_2 + \dots, \qquad h_\ell(x) = (-1)^\ell \langle \delta_x, (R_0 u)^\ell R_0 \delta_x \rangle.$$

In particular, using the integral kernel $\langle \delta_x, R_0(\kappa) \delta_y \rangle = \frac{1}{2\kappa} e^{-\kappa |x-y|}$ for the free resolvent, we find that

$$h_1 = -\kappa^{-1} R_0(2\kappa) u.$$

As it turns out, it is the function 1/q that is linearly related to the logarithmic renormalized trace of the resolvent. A consequence of the proposition is that the integral of 1/q diverges. Even after removing the constant, its integral still turns out to diverge for u in any H^s space. To rectify this, one also needs to remove the linear term in u, which formally integrates to a multiple of $\int u \, dx$; this is a conserved quantity for the KdV flow, and thus harmless. A similar renormalization was also implemented in [41] for the logarithm of the transmission coefficient. Taking these renormalizations into account, we can define the conserved quantity which controls the H^{-1} norm of u:

(2.9)
$$\alpha(\kappa, u) := \int \rho \, dx, \qquad \rho(x; \kappa, u) := -\frac{1}{2g(x; \kappa, u)} + \kappa + 2\kappa R_0(2\kappa)u$$

The formula for α is the trace of the integral kernel $-1/2G(x, y; \kappa, u)$ with the first two terms of its Taylor series about $u \equiv 0$ removed.

Proposition 2.3 (Introducing α [38]). There exists a constant C > 0 so that the following statements are true for any R > 0:

- (1) The quantities $\rho(x; \kappa, u)$ and $\alpha(\kappa, u)$ defined in (2.9) are finite and nonnegative for
- all $x \in \mathbb{R}$, $||u||_{H_{\kappa}^{-1}} \leq R$, and $\kappa \geq 1 + CR^2$. (2) The mapping $u \mapsto \alpha$ is a real analytic functional on $\{u : ||u||_{H_{\kappa}^{-1}} \leq R\}$ for all $\kappa \geq 1 + CR^2.$
- (3) We have

(2.10)
$$\frac{1}{4}\kappa^{-1} \|u\|_{H_{\kappa}^{-1}}^{2} \le \alpha(\kappa, u) \le \kappa^{-1} \|u\|_{H_{\kappa}^{-1}}^{2}$$

for all $||u||_{H^{-1}} \leq R$ and $\kappa \geq 1 + CR^2$.

The authors of [38] also proved that under the (KdV) flow, we have the following densityflux equation

(2.11)
$$\partial_t \rho + j' = 0,$$

where $\rho = \rho(x; \kappa, u)$ is defined in (2.9) and

(2.12)
$$j(x;\kappa,u) := \frac{1}{g}(4\kappa^3 g - 2\kappa^2 + u) + 2\kappa R_0(2\kappa)(u'' - 3u^2).$$

We see in particular by integrating (2.11) in space that α is conserved under the (KdV) flow.

By comparison, the coefficients a_i in (gKdV) break the density-flux equation (2.11). Instead, we obtain an additional term as follows:

(2.13)
$$\partial_t \rho + j' = d\rho|_u \big[(a_1 u')' + a_2 u^2 + a_3 u' + a_4 u \big],$$

where

(2.14)
$$d\rho|_{u}(f) = \frac{d}{ds}\rho(x;\kappa,u+sf)\Big|_{\substack{s=0\\9}} = \frac{1}{2g^{2}}dg|_{u}(f) + 2\kappa R_{0}(2\kappa)f$$

is the functional derivative of ρ at u, and

(2.15)
$$dg|_{u}(f) = -\int G(x,y)f(y)G(y,x)\,dy.$$

The identity (2.13) lies at the heart of both of the proofs of our a-priori and local smoothing estimates.

In order to leverage local smoothing, we will frequently need to commute slowly varying weights past R_0 , as in the following estimates.

/ \

Lemma 2.4. If $w : \mathbb{R} \to (0, \infty)$ satisfies

(2.16)
$$|w'(x)| + |w''(x)| \lesssim w(x) \quad and \quad \frac{w(y)}{w(x)} \lesssim e^{|x-y|/2}$$

uniformly for $x, y \in \mathbb{R}$, then

$$(2.17) \|wR_0\frac{1}{w}\|_{L^p\to L^p} \lesssim \kappa^{-2}, \|w\partial R_0\frac{1}{w}\|_{L^p\to L^p} \lesssim \kappa^{-1}, \|wR_0\frac{1}{w}\|_{H^{-1}_\kappa\to H^1_\kappa} \lesssim 1$$

uniformly for $\kappa \geq 1$, for any $1 \leq p \leq \infty$.

The proof is elementary; see, for example, [6, Lem. 2.7].

We will use this lemma in particular for the functions ψ^{ℓ} , $\ell = 1, 2, 3$, where

$$\psi(x) = \operatorname{sech}(\frac{x}{6})$$

The constant 6 is simply included to ensure that ψ^{ℓ} satisfies (2.16) for $\ell = 1, 2, 3$. (In turn, the constant 2 in (2.16) is based on the integral kernel $\langle \delta_x, R_0 \delta_y \rangle = \frac{1}{2\kappa} e^{-\kappa |x-y|}$ and the condition $\kappa \geq 1$.)

Over the course of our analysis, we will also encounter the 'double commutator'

$$R_0\psi^2 R_0 - \psi R_0^2 \psi = [[R_0, \psi], R_0 \psi].$$

In order to bound this operator, the naive estimate (2.17) unfortunately does not yield enough decay as $\kappa \to \infty$. To this end, we record the following operator estimates, which capture the double commutator structure:

Lemma 2.5. We have

(2.18)
$$\left\| \frac{1}{\psi} (R_0 \psi^2 R_0 - \psi R_0^2 \psi) \frac{1}{\psi} \right\|_{H^{-2}_{\kappa} \to H^2_{\kappa}} \lesssim \kappa^{-2},$$

(2.19)
$$\left\|\frac{1}{\psi}(R_0\partial\psi^2 R_0\partial - \psi R_0^2\partial^2\psi)\frac{1}{\psi}\right\|_{H_{\kappa}^{-2} \to H_{\kappa}^2} \lesssim 1$$

uniformly for $\kappa \geq 1$.

In comparison to (2.17), the proofs of (2.18) and (2.19) exhibit extra cancellation by computing the double commutator in a symmetric way. For example, (2.18) follows from [6, Lem. 2.10], and (2.19) can be verified using a similar argument.

3. The local smoothing norm

Here, we will record some useful estimates for the following local smoothing norm:

(3.1)
$$\|u\|_{\mathrm{LS}_{\kappa}} = \sup_{x_0 \in \mathbb{R}} \|\psi_{x_0} u'\|_{L^2_t H^{-1}_{\kappa}([-T,T] \times \mathbb{R})^2}$$

where T > 0 and

$$\psi(x) = \operatorname{sech}(\frac{x}{6}), \qquad \psi_{x_0}(x) = \psi(x - x_0).$$

We begin with the following elementary observations:

Lemma 3.1. Given a Schwartz function $\phi(x)$, we have

(3.2)
$$\|\phi u\|_{L^2_t H^{-2}_\kappa} \lesssim \kappa^{-1} \|u\|_{L^2_t H^{-1}_\kappa},$$

(3.3)
$$\|(\phi u)'\|_{L^2_t H^{-2}_\kappa} + \|\phi u'\|_{L^2_t H^{-2}_\kappa} \lesssim \kappa^{-1} \big(\|u\|_{\mathrm{LS}_\kappa} + \|u\|_{L^2_t H^{-1}_\kappa} \big),$$

(3.4)
$$\|(\phi u)''\|_{L^2_t H^{-2}_\kappa} + \|\phi u''\|_{L^2_t H^{-2}_\kappa} \lesssim \|u\|_{\mathrm{LS}_\kappa} + \kappa^{-1} \|u\|_{L^2_t H^{-1}_\kappa},$$

uniformly for $\kappa \geq 1$ and T > 0. (All space-time norms are taken over $[-T, T] \times \mathbb{R}$.)

Proof. For the estimate (3.2) we simply write

$$\|\phi u\|_{L^2_t H^{-2}_{\kappa}} \lesssim \|u\|_{L^2_t H^{-2}_{\kappa}} \lesssim \kappa^{-1} \|u\|_{L^2_t H^{-1}_{\kappa}}.$$

For the estimate (3.3) we use Leibnitz rule $(\phi u)' = \phi' u + \phi u'$. For the first term we use (3.2). For the second we write

$$\int \psi_z^2(x) \, dz = c$$

for some constant c, so that

$$\phi(x) = \frac{1}{c} \int \phi(x) \psi_z^2(x) \, dz$$

Then, it follows that

$$\|\phi u'\|_{H^{-1}_{\kappa}} \leq \frac{1}{c} \int \|\phi \psi_z^2 u'\|_{H^{-1}_{\kappa}} dz \lesssim \int \|\phi \psi_z\|_{H^4} \|\psi_z u'\|_{H^{-1}_{\kappa}} dz.$$

Integrating in z and noting that

$$\int \|\phi\psi_z\|_{H^4} \ dz \lesssim_{\phi} 1,$$

we obtain

(3.5)

 $\|\phi u'\|_{H^{-1}_{\kappa}} \lesssim \|u\|_{\mathrm{LS}_{\kappa}},$

and the estimate
$$(3.3)$$
 follows, as

$$\|\phi u'\|_{H^{-2}_{\kappa}} \lesssim \kappa^{-1} \|\phi u'\|_{H^{-1}_{\kappa}}$$

For the estimate (3.4) we simply write

$$(\phi u)'' = (\phi u')' + \phi' u' + \phi'' u_{\pm}$$

and we use twice (3.5) and once (3.2).

The operator estimate (3.7) below will play an important role in our analysis. In comparison to (2.4), we now bound the operator norm (rather than Hilbert–Schmidt) and only require a local norm of u to do so.

Lemma 3.2. We have

(3.6)
$$||fu'||_{L^2_t H^{-1}_\kappa} \lesssim \kappa^{-\frac{1}{2}} ||f||_{H^1_\kappa} ||u||_{\mathrm{LS}_\kappa} \text{ for all } f \in H^1(\mathbb{R}),$$

uniformly for $\kappa \geq 1$. As a consequence,

(3.7)
$$\left\|\sqrt{R_0}u'\sqrt{R_0}\right\|_{L^2_t\mathfrak{I}_\infty} \lesssim \kappa^{-\frac{1}{2}} \|u\|_{\mathrm{LS}_\kappa}.$$

Proof. We argue by duality. Let $h \in L^2_t H^1_{\kappa}$. Fix a smooth partition of unity

$$\sum_{j \in \mathbb{Z}} \chi_j^2 \equiv 1, \quad \text{with} \quad \chi_j \equiv 1 \text{ on } [j, j+1] \quad \text{and} \quad \text{supp } \chi_j \subset [j-1, j+2].$$

Then

$$\langle h, fu' \rangle = \sum_{j \in \mathbb{Z}} \int \widetilde{\chi}_j^2 u' \cdot \chi_j^2 f \overline{h} \, dx$$

where $\tilde{\chi}_j^2 = \chi_{j-1}^2 + \chi_j^2 + \chi_{j+1}^2$ is a fattened bump function. We estimate

$$\begin{split} \int_{-T}^{T} |\langle h, fu' \rangle| \, dt &\lesssim \sum_{j} \|\widetilde{\chi}_{j}^{2}u'\|_{L_{t}^{2}H_{\kappa}^{-1}} \left\|\chi_{j}^{2}f\overline{h}\right\|_{L_{t}^{2}H_{\kappa}^{1}} \\ &\lesssim \kappa^{-\frac{1}{2}} \sum_{j} \|u\|_{\mathrm{LS}_{\kappa}} \|\chi_{j}f\|_{H_{\kappa}^{1}} \|\chi_{j}h\|_{L_{t}^{2}H_{\kappa}^{1}} \\ &\lesssim \kappa^{-\frac{1}{2}} \|u\|_{\mathrm{LS}_{\kappa}} \left(\sum_{j} \|\chi_{j}f\|_{H_{\kappa}^{1}}^{2}\right)^{\frac{1}{2}} \left(\sum_{j} \|\chi_{j}h\|_{L_{t}^{2}H_{\kappa}^{1}}^{2}\right)^{\frac{1}{2}} \\ &\lesssim \kappa^{-\frac{1}{2}} \|u\|_{\mathrm{LS}_{\kappa}} \|f\|_{H_{\kappa}^{1}} \|h\|_{L_{t}^{2}H_{\kappa}^{1}} \,. \end{split}$$

Taking a supremum over $||h||_{L^2_t H^1_r} \leq 1$, the estimate (3.6) follows.

The second estimate (3.7) now follows immediately:

$$\left\|\sqrt{R_0}u'\sqrt{R_0}\right\|_{L^2_t\mathfrak{I}_{\infty}} \lesssim \kappa^{-\frac{1}{2}} \left\|\sqrt{R_0}\right\|_{H^{-1}_{\kappa} \to L^2} \|u\|_{\mathrm{LS}_{\kappa}} \left\|\sqrt{R_0}\right\|_{L^2 \to H^1_{\kappa}} \lesssim \kappa^{-\frac{1}{2}} \|u\|_{\mathrm{LS}_{\kappa}}.$$

Our next lemma provides an estimate for the diagonal Green's function analogous to (2.6) that captures the gain of regularity we expect from our local smoothing norm.

Lemma 3.3. There exists a constant C > 0 so that for any Schwartz function $\phi(x)$, we have

(3.8) $\|\phi g'\|_{L^2_t H^1_\kappa} \lesssim_{\phi} \kappa^{-1} \|u\|_{\mathrm{LS}_\kappa}$

uniformly for

(3.9)
$$\kappa \ge 1 + C \|u\|_{L_t^{\infty} H_{\kappa}^{-1}}^2.$$

Proof. Using a 'continuous partition of unity' as in the proof of (3.3), it suffices to prove (3.8) in the special case where $\phi = \psi$. We argue by duality. For $f \in L_t^2 H_{\kappa}^{-1}$, we can use the expansion we have for g in (2.8) to write

$$\int_{-T}^{T} \left| \int f\psi g' \, dx \right| dt \leq \sum_{\ell \geq 1} \int_{-T}^{T} \left| \operatorname{tr} \left\{ f\psi [\partial, (R_0 u)^{\ell} R_0] \right\} \right| dt.$$

When $\ell = 1$, there is only one copy of u on which the derivative ∂ may fall. In this case, we write

$$f\psi R_0 u' = f[\psi R_0 \frac{1}{\psi}]\psi u'$$

and note that the operator in square brackets is bounded $H_{\kappa}^{-1} \to H_{\kappa}^{1}$ by (2.17). Therefore

$$\left| \operatorname{tr} \left\{ f \psi R_0 u' R_0 \right\} \right| \lesssim \left\| \sqrt{R_0} f \sqrt{R_0} \right\|_{\mathfrak{I}_2} \left\| \sqrt{R_0} \psi u' \sqrt{R_0} \right\|_{\mathfrak{I}_2} \lesssim \kappa^{-1} \left\| f \right\|_{H_{\kappa}^{-1}} \left\| \psi u' \right\|_{H_{\kappa}^{-1}},$$

and thus

$$\int_{-T}^{T} \left| \operatorname{tr} \left\{ f \psi R_0 u' R_0 \right\} \right| dt \lesssim \kappa^{-1} \left\| f \right\|_{L^2_t H^{-1}_\kappa} \left\| u \right\|_{\mathrm{LS}_\kappa}$$

Next, we turn to the case where $\ell \geq 2$. We distribute the derivative ∂ using the product rule $[\partial, AB] = [\partial, A]B + A[\partial, B]$. For the one factor of u', we estimate its contribution in operator norm using (3.7). The requirement $\ell \geq 2$ then guarantees that there is at least one other factor of u, which together with f provides us with two terms we can put in Hilbert–Schmidt norm:

$$\begin{split} \sum_{\ell \ge 2} \int_{-T}^{T} \left| \operatorname{tr} \left\{ f \psi[\partial, (R_{0}u)^{\ell}R_{0}] \right\} \right| dt \\ & \le \sum_{\ell \ge 2} \ell \left\| \sqrt{R_{0}} f \psi \sqrt{R_{0}} \right\|_{L_{t}^{2}\mathfrak{I}_{2}} \left\| \sqrt{R_{0}}u' \sqrt{R_{0}} \right\|_{L_{t}^{2}\mathfrak{I}_{\infty}} \left\| \sqrt{R_{0}}u \sqrt{R_{0}} \right\|_{L_{t}^{\infty}\mathfrak{I}_{2}}^{\ell-1} \\ & \lesssim \sum_{\ell \ge 2} \ell \kappa^{-1} \left\| f \right\|_{L_{t}^{2}H_{\kappa}^{-1}} \left\| u \right\|_{\mathrm{LS}_{\kappa}} \left(\kappa^{-\frac{1}{2}} \left\| u \right\|_{L_{t}^{\infty}H_{\kappa}^{-1}} \right)^{\ell-1} \\ & \lesssim \kappa^{-1} \left\| f \right\|_{L_{t}^{2}H_{\kappa}^{-1}} \left\| u \right\|_{\mathrm{LS}_{\kappa}}. \end{split}$$

Altogether, we conclude

$$\int_{-T}^{T} \left| \int f \psi g' \, dx \right| dt \lesssim \kappa^{-1} \, \|f\|_{L^{2}_{t} H^{-1}_{\kappa}} \, \|u\|_{\mathrm{LS}_{\kappa}} \, .$$

Taking a supremum over $||f||_{L^2_t H^{-1}_{\kappa}} \leq 1$, the estimate (3.8) follows.

The next few results provide estimates for the terms h_{ℓ} appearing in the series (2.8) for g. We will start with h_1 , which follows easily from (3.2)–(3.4).

Lemma 3.4. We have

(3.10)
$$\|\psi h_1\|_{L^2_{t,x}} \lesssim \kappa^{-2} \|u\|_{L^2_t H^{-1}_\kappa},$$

(3.11)
$$\|\psi h'_1\|_{L^2_{t,x}} \lesssim \kappa^{-2} \|u\|_{\mathrm{LS}_{\kappa}},$$

(3.12)
$$\|\psi h_1''\|_{L^2_{t,x}} \lesssim \kappa^{-1} \|u\|_{\mathrm{LS}_{\kappa}} + \kappa^{-2} \|u\|_{L^2_t H^{-1}_{\kappa}}$$

uniformly for $\kappa \geq 1$ and T > 0.

Proof. Recall that $h_1 = -\kappa^{-1} R_0(2\kappa) u$. Using (2.17), we see that

$$\|\psi h_1'\|_{L^2_{t,x}} \lesssim \kappa^{-1} \|\psi h_1'\|_{L^2_t H^1_\kappa} = \kappa^{-2} \|\psi R_0(2\kappa) \frac{1}{\psi} \cdot \psi u'\|_{L^2_t H^1_\kappa} \lesssim \kappa^{-2} \|u\|_{\mathrm{LS}_\kappa}.$$

This proves (3.11). For (3.10) and (3.12), we also use (3.4) and (3.2):

$$\|\psi h_1\|_{L^2_{t,x}} \lesssim \kappa^{-1} \|\psi u\|_{L^2_t H^{-2}_{\kappa}} \lesssim \kappa^{-2} \|u\|_{L^2_t H^{-1}_{\kappa}},$$

$$\|\psi h_1''\|_{L^2_{t,x}} \lesssim \kappa^{-1} \|\psi u''\|_{L^2_t H^{-2}_{\kappa}} \lesssim \kappa^{-1} (\|u\|_{\mathrm{LS}_{\kappa}} + \kappa^{-1} \|u\|_{L^2_t H^{-1}_{\kappa}}).$$

The following lemma will be useful in estimating all of the cubic and higher order terms of j (see (4.16) for details). The proof is rather involved, because we need to be efficient in our estimation. Specifically, we need to get a factor on the RHS of (3.13) that is $o(\kappa^{-5})$. (Otherwise, we would not be able to start at $\ell = 3$ in (4.16).)

Lemma 3.5. Given a Schwartz function $\phi(x)$, we have

(3.13)
$$\sum_{\ell \ge 3} \|\phi h_\ell\|_{L^1_{t,x}} \lesssim (T\kappa^2)^{\frac{3}{4}} \kappa^{-\frac{11}{2}} \|u\|_{L^{\infty}_t H^{-1}_{\kappa}} \left(\|u\|^2_{L^{\infty}_t H^{-1}_{\kappa}} + \|u\|^2_{\mathrm{LS}_{\kappa}} \right)$$

uniformly for $T \leq \kappa^{-2}$ and κ satisfying (3.9).

Proof. Let us begin with (3.13) for the case $\ell = 3$. We argue by duality: for $f \in L^{\infty}([-T, T] \times \mathbb{R})$, we have

(3.14)
$$\int_{-T}^{T} \int_{-\infty}^{\infty} f \phi h_{\ell} \, dx \, dt = (-1)^{\ell} \int_{-T}^{T} \operatorname{tr} \left\{ \sqrt{R_0} f \phi(R_0 u)^{\ell} \sqrt{R_0} \right\} dt.$$

Using a continuous partition of unity argument as in the proof of (3.4), it suffices to prove (3.13) in the special case when $\phi = \psi^3$. Write

$$\sqrt{R_0} f \psi^3 (R_0 u)^3 \sqrt{R_0} = \sqrt{R_0} f [\psi^3 R_0 \frac{1}{\psi^3}] \psi u [\psi^2 R_0 \frac{1}{\psi^2}] \psi u [\psi R_0 \frac{1}{\psi}] \psi u \sqrt{R_0}.$$

By (2.17), each operator in square brackets above is bounded $H_{\kappa}^{-1} \to H_{\kappa}^{1}$. This allows us to write

$$\sqrt{R_0} f \psi^3 (R_0 u)^3 \sqrt{R_0} = (\sqrt{R_0} f \sqrt{R_0}) A_3 (\sqrt{R_0} \psi u \sqrt{R_0}) A_2 (\sqrt{R_0} \psi u \sqrt{R_0}) A_1 (\sqrt{R_0} \psi u \sqrt{R_0}) A_2 (\sqrt{R_0} \psi u \sqrt{R_0}) A_$$

for operators A_j with $||A_j||_{\mathfrak{I}_{\infty}} \lesssim 1$. We seek to estimate the \mathfrak{I}_1 norm of the product, for which it suffices to bound two of the factors in round brackets in \mathfrak{I}_{∞} and two in \mathfrak{I}_2 . The factor containing f is directly estimated in \mathfrak{I}_{∞} by

(3.15)
$$\|\sqrt{R_0}f\sqrt{R_0}\|_{\mathfrak{I}_{\infty}} \lesssim \kappa^{-2}\|f\|_{L^{\infty}}.$$

It remains to consider the three remaining factors. To estimate them, we will use two bounds. On one hand for the Hilbert–Schmidt norm we use (2.4) which gives

(3.16)
$$\|\sqrt{R_0}w\sqrt{R_0}\|_{\mathfrak{I}_2} = \kappa^{-\frac{1}{2}}\|w\|_{H_{\kappa}^{-1}}.$$

On the other hand for the operator norm we combine (3.15) for the low frequencies with (3.16) for the high frequencies and then use an interpolation inequality,

$$\left\|\sqrt{R_0}w\sqrt{R_0}\right\|_{\mathfrak{I}_{\infty}} \lesssim \kappa^{-\frac{1}{2}} \|w_{>\kappa}\|_{H_{\kappa}^{-1}} + \kappa^{-2} \|w_{<\kappa}\|_{L^{\infty}} \lesssim \kappa^{-\frac{1}{2}} \|w_{>\kappa}\|_{H_{\kappa}^{-1}} + \kappa^{-2} \|w_{<\kappa}\|_{L^2}^{\frac{1}{2}} \|w_{<\kappa}'\|_{L^2}^{\frac{1}{2}},$$

which we can shorten to

(3.17)
$$\|\sqrt{R_0}w\sqrt{R_0}\|_{\mathfrak{I}_{\infty}} \lesssim \kappa^{-\frac{1}{2}} \|w\|_{H_{\kappa}^{-1}}^{\frac{1}{2}} \|w'\|_{H_{\kappa}^{-2}}^{\frac{1}{2}}$$

Applying these bounds to $h = \psi u$, we obtain

$$\begin{aligned} \left\|\sqrt{R_0}f\psi^3(R_0u)^3\sqrt{R_0}\right\|_{\mathfrak{I}_1} &\lesssim \kappa^{-2-\frac{3}{2}}\|h\|_{H_{\kappa}^{-1}}^{\frac{5}{2}}\|h'\|_{H_{\kappa}^{-2}}^{\frac{1}{2}}\|f\|_{L^{\infty}} \\ &\lesssim \kappa^{-2-\frac{3}{2}}\|h\|_{H_{\kappa}^{-1}}(c^3\|h'\|_{H_{\kappa}^{-2}}^2 + c^{-1}\|h\|_{H_{\kappa}^{-1}}^2)\|f\|_{L^{\infty}} \end{aligned}$$

with an arbitrary positive constant c. Now we integrate in time,

$$\left\|\sqrt{R_0}f\psi^3(R_0u)^3\sqrt{R_0}\right\|_{L^1_t\mathfrak{I}_1} \lesssim \kappa^{-2-\frac{3}{2}} \|h\|_{L^\infty_tH^{-1}_\kappa} (c^3\|h'\|_{L^2_tH^{-2}_\kappa}^2 + c^{-1}\|h\|_{L^2_tH^{-1}_\kappa}^2) \|f\|_{L^\infty_{t,x}}.$$

Now we use (3.3),

$$\begin{split} \left\| \sqrt{R_0} f \psi^3(R_0 u)^3 \sqrt{R_0} \right\|_{L^1_t \mathfrak{I}_1} \\ &\lesssim \kappa^{-2 - \frac{3}{2}} \| u \|_{L^\infty_t H^{-1}_\kappa} \left(c^3 \kappa^{-2} \| u \|_{\mathrm{LS}_\kappa}^2 + (c^{-1} + c^3 \kappa^{-2}) \| u \|_{L^2_t H^{-1}_\kappa}^2 \right) \| f \|_{L^\infty_{t,x}} \\ &\lesssim \kappa^{-4 - \frac{3}{2}} \| u \|_{L^\infty_t H^{-1}_\kappa} \left(c^3 \| u \|_{\mathrm{LS}_\kappa}^2 + T (c^{-1} \kappa^2 + c^3) \| u \|_{L^\infty_t H^{-1}_\kappa}^2 \right) \| f \|_{L^\infty_{t,x}}. \end{split}$$

Finally we make a good choice for c in order to balance the constants,

(3.18)
$$c = T^{\frac{1}{4}} \kappa^{\frac{1}{2}}.$$

This gives

 $(3.19) \|\sqrt{R_0} f \psi^3(R_0 u)^3 \sqrt{R_0} \|_{L^1_t \mathfrak{I}_1} \lesssim \kappa^{-\frac{11}{2}} (T\kappa^2)^{\frac{3}{4}} \|u\|_{L^{\infty}_t H^{-1}_{\kappa}} \left(\|u\|^2_{L^{\infty}_t H^{-1}_{\kappa}} + \|u\|^2_{\mathrm{LS}_{\kappa}} \right) \|f\|_{L^{\infty}_{t,x}},$ which suffices for the case $\ell = 3$.

Next, we turn to the terms with $\ell \geq 4$. Using (3.19), we estimate

$$\begin{split} \left\| \sqrt{R_0} f \psi^3(R_0 u)^{\ell} \sqrt{R_0} \right\|_{L^1_t \mathfrak{I}_1} \\ & \leq \left\| \sqrt{R_0} f \psi^3(R_0 u)^3 \sqrt{R_0} \right\|_{L^1_t \mathfrak{I}_1} \left\| \sqrt{R_0} u \sqrt{R_0} \right\|_{L^\infty_t \mathfrak{I}_2}^{\ell-3} \\ & \lesssim \kappa^{-\frac{11}{2}} (T\kappa^2)^{\frac{3}{4}} \left\| f \right\|_{L^\infty_{t,x}} \left\| u \right\|_{L^\infty_t H^{-1}_\kappa} \left(\left\| u \right\|_{L^\infty_t H^{-1}_\kappa}^2 + \left\| u \right\|_{\mathrm{LS}_\kappa}^2 \right) \left(\kappa^{-\frac{1}{2}} \left\| u \right\|_{L^\infty_t H^{-1}_\kappa} \right)^{\ell-3}. \end{split}$$

Choosing C larger if necessary, the condition (3.9) implies

$$\kappa^{-\frac{1}{2}} \|u\|_{L^{\infty}_{t}H^{-1}_{\kappa}} \le \frac{1}{2}$$

Summing over $\ell \geq 3$ this relation will make our geometric sum convergent. We now can take the supremum over $\|f\|_{L^{\infty}_{t,r}} \leq 1$ and conclude the proof of (3.13).

Using a similar argument, we also obtain estimates in $L_t^2 H_{\kappa}^1$ (instead of $L_{t,x}^1$) using only one copy of the local smoothing norm:

Lemma 3.6. Given a Schwartz function $\phi(x)$, we have

(3.20)
$$\begin{aligned} \left\| \phi(g - \frac{1}{2\kappa} - h_1) \right\|_{L^2_t H^1_\kappa} &\leq \sum_{\ell \geq 2} \|\phi h_\ell\|_{L^2_t H^1_\kappa} \\ &\lesssim \kappa^{-\frac{5}{2}} (T\kappa^2)^{\frac{1}{4}} \|u\|_{L^\infty_t H^{-1}_\kappa} \left(\|u\|_{L^\infty_t H^{-1}_\kappa} + \|u\|_{\mathrm{LS}_\kappa} \right) \end{aligned}$$

uniformly for $T \leq \kappa^{-2}$ and κ satisfying (3.9).

Proof. Using a continuous partition of unity argument as in the proof of (3.4), it suffices to prove (3.20) in the special case $\phi = \psi^2$. We use the expansion (2.8) for g, which shows that

$$g - \frac{1}{2\kappa} - h_1 = \sum_{\ell \ge 2} h_\ell.$$

We argue by duality. For $f \in L^2_t H^{-1}_{\kappa}([-T,T] \times \mathbb{R})$, we have

$$|\langle \phi(g - \frac{1}{2\kappa} - h_1), f \rangle| \lesssim \sum_{\ell \ge 2} \|\sqrt{R_0} f \psi^2(R_0 u)^2 \sqrt{R_0}\|_{L^1_t \mathfrak{I}_1}.$$

It remains to estimate the terms on the right. We begin with the case $\ell = 2$, where we write

$$\sqrt{R_0} f \psi^2 (R_0 u)^2 \sqrt{R_0} = \sqrt{R_0} f [\psi^2 R_0 \frac{1}{\psi^2}] \psi u [\psi R_0 \frac{1}{\psi}] \psi u \sqrt{R_0}.$$

By (2.17), each operator in square brackets above is bounded $H_{\kappa}^{-1} \to H_{\kappa}^{1}$. This allows us to write

$$\sqrt{R_0}f\psi^2(R_0u)^2\sqrt{R_0} = \left(\sqrt{R_0}f\sqrt{R_0}\right)A_2\left(\sqrt{R_0}\psi u\sqrt{R_0}\right)A_1\left(\sqrt{R_0}\psi u\sqrt{R_0}\right)$$

for operators A_j with $||A_j||_{\mathfrak{I}_{\infty}} \lesssim 1$.

Now we use (3.16) for the f contribution and for one of the ψu factors, and (3.17) for the other ψu factor. This yields

$$\begin{aligned} \left\|\sqrt{R_0}f\psi^2(R_0u)^2\sqrt{R_0}\right\|_{\mathfrak{I}_1} &\lesssim \kappa^{-\frac{3}{2}}\|f\|_{H_{\kappa}^{-1}}\|\psi u\|_{H_{\kappa}^{-1}}^{\frac{3}{2}}\|(\psi u)'\|_{H_{\kappa}^{-2}}^{\frac{1}{2}} \\ &\lesssim \kappa^{-\frac{3}{2}}\|f\|_{H_{\kappa}^{-1}}\|\psi u\|_{H_{\kappa}^{-1}}\left(c^{-1}\|\psi u\|_{H_{\kappa}^{-1}}+c\|(\psi u)'\|_{H_{\kappa}^{-2}}\right)\end{aligned}$$

Finally, we integrate in time and use (3.3) to arrive at

$$\begin{split} \left\|\sqrt{R_0}f\psi^2(R_0u)^2\sqrt{R_0}\right\|_{L^1_t\mathfrak{I}_1} \\ \lesssim \kappa^{-\frac{3}{2}}\|f\|_{L^2_tH^{-1}_\kappa}\|\psi u\|_{L^\infty_tH^{-1}_\kappa} \Big(\kappa^{-1}c\|u\|_{\mathrm{LS}_\kappa} + (c^{-1}+\kappa^{-1}c)\|u\|_{L^2_tH^{-1}_\kappa}\Big). \end{split}$$

Then by applying Hölder's inequality in time, we obtain

$$\left\|\sqrt{R_0}f\psi^2(R_0u)^2\sqrt{R_0}\right\|_{L^1_t\mathfrak{I}_1} \lesssim \kappa^{-\frac{5}{2}}\|f\|_{L^2_tH^{-1}_\kappa}\|\psi u\|_{L^\infty_tH^{-1}_\kappa}\left(c\|u\|_{\mathrm{LS}_\kappa} + T^{\frac{1}{2}}(\kappa c^{-1} + c)\|u\|_{L^\infty_tH^{-1}_\kappa}\right)$$

Optimizing the choice of c as in (3.18) this yields

$$\left\|\sqrt{R_0}f\psi^2(R_0u)^2\sqrt{R_0}\right\|_{L^1_t\mathfrak{I}_1} \lesssim \kappa^{-2}T^{\frac{1}{4}}\|f\|_{L^2_tH^{-1}_\kappa}\|u\|_{L^\infty_tH^{-1}_\kappa}\left(\|u\|_{L^\infty_tH^{-1}_\kappa}+\|u\|_{\mathrm{LS}_\kappa}\right),$$

which suffices for $\ell = 2$.

For $\ell \geq 3$, we estimate

$$\begin{aligned} \left\|\sqrt{R_0}f\psi^2(R_0u)^\ell\sqrt{R_0}\right\|_{L^1_t\mathfrak{I}_1} &\leq \left\|\sqrt{R_0}f\psi^2(R_0u)^2\sqrt{R_0}\right\|_{L^1_t\mathfrak{I}_1}\left\|\sqrt{R_0}u\sqrt{R_0}\right\|_{L^\infty_t\mathfrak{I}_2}^{\ell-2} \\ &\lesssim \kappa^{-2}T^{\frac{1}{4}}\left\|f\right\|_{L^2_tH^{-1}_\kappa}\left\|u\right\|_{L^\infty_tH^{-1}_\kappa}\left(\left\|u\right\|_{L^\infty_tH^{-1}_\kappa}+\left\|u\right\|_{\mathrm{LS}_\kappa}\right)\left(\kappa^{-\frac{1}{2}}\left\|u\right\|_{L^\infty_tH^{-1}_\kappa}\right)^{\ell-2}.\end{aligned}$$

Summing over $\ell \geq 2$ (exactly like in the proof of the previous lemma) and taking a supremum over $\|f\|_{L^2_t H^{-1}_{\kappa}} \leq 1$ yields the second inequality in (3.20).

Altogether, the previous lemmas provide us with the control we need over the functional $u \mapsto g$. However, the other key functional in our analysis is $u \mapsto \frac{1}{g}$; indeed, this is what appears in the definition (2.9) of ρ . The following lemma provides us with an estimate for the quadratic and higher order terms of $\frac{1}{g}$, using the tools that we have already developed.

Lemma 3.7. Given a Schwartz function $\phi(x)$, we have

(3.21)
$$\left\| \phi \left(\frac{1}{g} - 2\kappa + \frac{4\kappa^2 h_1}{1 + 2\kappa h_1} \right) \right\|_{L^2_{t,x}} \lesssim \kappa^{-\frac{3}{2}} (T\kappa^2)^{\frac{1}{4}} \| u \|_{L^{\infty}_{t} H^{-1}_{\kappa}} \left(\| u \|_{L^{\infty}_{t} H^{-1}_{\kappa}} + \| u \|_{\mathrm{LS}_{\kappa}} \right)$$

uniformly for $T \leq \kappa^{-2}$ and κ as in (3.9).

Proof. Recall that $h_1 = -\kappa^{-1} R_0(2\kappa) u$. Choosing C larger if necessary, the condition (3.9) implies

$$\|2\kappa h_1\|_{L^{\infty}_x} = \|2R_0(2\kappa)u\|_{L^{\infty}_x} \lesssim \kappa^{-\frac{1}{2}} \|u\|_{H^{-1}_{\kappa}} < 1.$$

Hence,

(3.22)
$$\left\|\frac{1}{1+2\kappa h_1}\right\|_{L^{\infty}_{t,x}} + \left\|\frac{2\kappa h_1}{1+2\kappa h_1}\right\|_{L^{\infty}_{t,x}} \lesssim 1$$

uniformly for κ satisfying (3.9).

Now, we use the identity

$$\frac{1}{g} - 2\kappa + \frac{4\kappa^2 h_1}{1 + 2\kappa h_1} = -\frac{2\kappa}{g} \left(1 - \frac{2\kappa h_1}{1 + 2\kappa h_1} \right) \left(g - \frac{1}{2\kappa} - h_1 \right)$$

together with the estimates (2.7) and (3.20) to bound

$$\begin{split} \|\phi\big(\frac{1}{g} - 2\kappa + \frac{4\kappa^{2}h_{1}}{1 + 2\kappa h_{1}}\big)\|_{L^{2}_{t,x}} &\lesssim \kappa \|\frac{1}{g}\|_{L^{\infty}_{t,x}} \|\phi\big(g - \frac{1}{2\kappa} - h_{1}\big)\|_{L^{2}_{t,x}} \\ &\lesssim \kappa \|\psi\big(g - \frac{1}{2\kappa} - h_{1}\big)\|_{L^{2}_{t}H^{1}_{\kappa}} \\ &\lesssim \kappa^{-1}T^{\frac{1}{4}} \|u\|_{L^{\infty}_{t}H^{-1}_{\kappa}} \left(\|u\|_{L^{\infty}_{t}H^{-1}_{\kappa}} + \|u\|_{\mathrm{LS}_{\kappa}}\right). \end{split}$$

Before proceeding, we will record one more estimate for the product h_1h_2 , similar in spirit to the preceding analysis. (This will be useful in estimating (4.15).)

Lemma 3.8. Given a Schwartz function $\phi(x)$, we have

(3.23)
$$\|\phi h_1 h_2\|_{L^1_{t,x}} \lesssim \kappa^{-\frac{13}{2}} (T\kappa^2)^{\frac{3}{4}} \|u\|_{L^{\infty}_t H^{-1}_{\kappa}} \left(\|u\|^2_{L^{\infty}_t H^{-1}_{\kappa}} + \|u\|^2_{\mathrm{LS}_{\kappa}} \right)$$

uniformly for $T \leq \kappa^{-2}$ and $\kappa \geq 1$.

Proof. Using a continuous partition of unity argument as in the proof of (3.4), it suffices to prove (3.23) in the special cases where $\phi = \psi^3$.

We argue by duality: for $f \in L^{\infty}([-T,T] \times \mathbb{R})$, we have

$$\int_{-T}^{T} \int_{-\infty}^{\infty} f\psi^{3}h_{1}h_{2} \, dx \, dt = \int_{-T}^{T} \operatorname{tr} \left\{ \sqrt{R_{0}} f\psi^{3}h_{1}(R_{0}u)^{2} \sqrt{R_{0}} \right\} dt.$$

By (2.17), we may write

$$\sqrt{R_0} f \psi^3 h_1 (R_0 u)^2 \sqrt{R_0} = \sqrt{R_0} f \psi h_1 \sqrt{R_0} A_2 \sqrt{R_0} \psi u \sqrt{R_0} A_1 \sqrt{R_0} \psi u \sqrt{R_0}$$

for operators A_j with $||A_j||_{\mathfrak{I}_{\infty}} \lesssim 1$.

Now we use the bounds (3.16) and (3.17) to estimate

$$\left\|\sqrt{R_0}f\psi^3h_1(R_0u)^2\sqrt{R_0}\right\|_{\mathfrak{I}_1} \lesssim \kappa^{-\frac{3}{2}}\|f\psi h_1\|_{H_{\kappa}^{-1}}\|\psi u\|_{H_{\kappa}^{-1}}^{\frac{3}{2}}\|(\psi u)'\|_{H_{\kappa}^{-2}}^{\frac{1}{2}}.$$

As $h_1 = -\kappa^{-1} R_0(2\kappa) u$, the first factor on the right is estimated as

 $\|f\psi h_1\|_{H^{-1}_{\kappa}} \lesssim \kappa^{-1} \|f\|_{L^{\infty}} \|\psi h_1\|_{L^2} \lesssim \kappa^{-3} \|f\|_{L^{\infty}} \|\psi R_0 u\|_{H^1_{\kappa}} \lesssim \kappa^{-3} \|f\|_{L^{\infty}} \|\psi u\|_{H^{-1}_{\kappa}}.$ Hence we obtain the fixed time bound

$$\begin{aligned} \left\|\sqrt{R_0}f\psi^3h_1(R_0u)^2\sqrt{R_0}\right\|_{\mathfrak{I}_1} &\lesssim \kappa^{-\frac{9}{2}}\|f\|_{L^{\infty}}\|\psi u\|_{H^{-1}_{\kappa}}^{\frac{5}{2}}\|(\psi u)'\|_{H^{-2}_{\kappa}}^{\frac{1}{2}}\\ &\lesssim \kappa^{-\frac{9}{2}}\|f\|_{L^{\infty}}\|\psi u\|_{H^{-1}_{\kappa}}(c^3\|(\psi u)'\|_{H^{-2}_{\kappa}}^2 + c^{-1}\|\psi u\|_{H^{-1}_{\kappa}}^2). \end{aligned}$$

Integrating in time and using (3.3) this gives

$$\begin{split} \left\| \sqrt{R_0} f \psi^3 h_1(R_0 u)^2 \sqrt{R_0} \right\|_{L^1_t \mathfrak{I}_1} \\ \lesssim \kappa^{-\frac{13}{2}} \| f \|_{L^\infty_{t,x}} \| \psi u \|_{L^\infty_t H^{-1}_\kappa} (c^3 \| u \|_{\mathrm{LS}_\kappa}^2 + T(c^{-1} \kappa^2 + c^3) \| \psi u \|_{L^\infty_t H^{-1}_\kappa}^2). \end{split}$$

Balancing c as in (3.18) we arrive at

$$\left\|\sqrt{R_0}f\psi^3h_1(R_0u)^2\sqrt{R_0}\right\|_{L^1_t\mathfrak{I}_1} \lesssim \kappa^{-\frac{13}{2}}(T\kappa^2)^{\frac{3}{4}}\|f\|_{L^\infty_{t,x}}\|\psi u\|_{L^\infty_t H^{-1}_\kappa}\left(\|u\|^2_{L^\infty_t H^{-1}_\kappa} + \|u\|^2_{\mathrm{LS}_\kappa}\right)$$
needed.

as needed.

We conclude this section with two estimates for the functional derivative of ρ , which appears on RHS(2.13). First, we have the following estimate that does not make use of local smoothing:

Lemma 3.9. Given T > 0, we have

(3.24)
$$\|d\rho|_{u}(f)\|_{L^{1}_{t,x}} \lesssim \kappa^{-1} \|u\|_{L^{\infty}_{t}H^{-1}_{\kappa}} \|f\|_{L^{1}_{t}H^{-1}_{\kappa}}$$

uniformly for κ satisfying (3.9).

Proof. We argue by duality: let $h \in L^{\infty}([-T,T] \times \mathbb{R})$. Expanding the series (2.3) in the definition (2.15) of dg, we obtain

$$\int_{-T}^{T} \int h \frac{1}{g^2} dg|_u(f) \, dx \, dt = \sum_{\ell, m \ge 0} \int_{-T}^{T} (-1)^{\ell+m+1} \operatorname{tr} \left\{ \frac{h}{g^2} (R_0 u)^{\ell} R_0 f(R_0 u)^m R_0 \right\} dt$$

We will insert this into the formula (2.14) for $d\rho$.

For $\ell = m = 0$, we have

$$-\operatorname{tr}\left\{\frac{h}{g^{2}}R_{0}fR_{0}\right\} = -\int \frac{h}{g^{2}}\kappa^{-1}R_{0}(2\kappa)f\,dx$$
$$= -\int h\,4\kappa R_{0}(2\kappa)f\,dx - \int h\left(\frac{1}{g^{2}} - 4\kappa^{2}\right)\kappa^{-1}R_{0}(2\kappa)f\,dx.$$

The first term on the RHS above is canceled out by the contribution of $2\kappa R_0(2\kappa)f$ in (2.14). For the second term, we use (2.7) to estimate

$$\int_{-T}^{T} \int \left| h \left(\frac{1}{g^2} - 4\kappa^2 \right) \kappa^{-1} R_0(2\kappa) f \right| dx dt$$

$$\lesssim \|h\|_{L^{\infty}_{t,x}} \|\frac{1}{g} - 2\kappa\|_{L^{\infty}_t L^2_x} \left(\|\frac{1}{g}\|_{L^{\infty}_{t,x}} + 2\kappa \right) \kappa^{-1} \|R_0(2\kappa) f\|_{L^1_t L^2_x}$$

$$\lesssim \kappa^{-1} \|h\|_{L^{\infty}_{t,x}} \|u\|_{L^{\infty}_t H^{-1}_\kappa} \|f\|_{L^1_t H^{-1}_\kappa}.$$

So it only remains to deal with the terms for which $\ell + m \ge 1$. In this case, there are at least two operators we can put in Hilbert–Schmidt norm:

$$\sum_{\ell+m\geq 1} \left\| \frac{h}{g^2} (R_0 u)^{\ell} R_0 f(R_0 u)^m R_0 \right\|_{L^1_t \mathfrak{I}_1} \leq \sum_{\ell+m\geq 1} \left\| \sqrt{R_0} \frac{h}{g^2} \sqrt{R_0} \right\|_{L^\infty_t \mathfrak{I}_\infty} \left\| \sqrt{R_0} f\sqrt{R_0} \right\|_{L^1_t \mathfrak{I}_2} \left\| \sqrt{R_0} u\sqrt{R_0} \right\|_{L^\infty_t \mathfrak{I}_2}^{\ell+m}$$
18

$$\lesssim \kappa^{-\frac{5}{2}} \sum_{\ell+m \ge 1} \left\| \frac{h}{g^2} \right\|_{L^{\infty}_{t,x}} \|f\|_{L^1_t H^{-1}_{\kappa}} \left(\kappa^{-\frac{1}{2}} \|u\|_{L^{\infty}_t H^{-1}_{\kappa}} \right)^{\ell+m} \\ \lesssim \kappa^{-1} \|u\|_{L^{\infty}_t H^{-1}_{\kappa}} \|h\|_{L^{\infty}_{t,x}} \|f\|_{L^1_t H^{-1}_{\kappa}}.$$

Altogether, we have

$$\int_{-T}^{T} \int \left| h \, d\rho \right|_{u}(f) \left| \, dx \, dt \lesssim \kappa^{-1} \, \|u\|_{L_{t}^{\infty} H_{\kappa}^{-1}} \, \|h\|_{L_{t,x}^{\infty}} \, \|f\|_{L_{t}^{1} H_{\kappa}^{-1}} \, .$$

Taking a supremum over $||h||_{L^{\infty}_{t,x}} \leq 1$, we obtain (3.24).

We can also handle one derivative inside $d\rho$, provided that the input is localized in space and we may estimate one copy of u in local smoothing norm:

Lemma 3.10. We have

(3.25)
$$\left| \int_{-T}^{T} \int \phi_{x_0} \, d\rho |_u (\psi^2 f)' \, dx \, dt \right| \lesssim \kappa^{-1} \, \|\psi f\|_{L^2_t H^{-1}_\kappa} \left(\, \|u\|_{L^\infty_t H^{-1}_\kappa} + \|u\|_{\mathrm{LS}\kappa} \, \right)$$

uniformly for $x_0 \in \mathbb{R}$, $T \le \kappa^{-2}$, and κ satisfying (3.9), where ϕ is defined by (3.26) $\phi_{x_0}(x) = \phi(x - x_0) = 6 \tanh(\frac{x - x_0}{\epsilon})$ so that $\phi'_{x_0} = \psi^2_{x_0}$.

(3.26)
$$\phi_{x_0}(x) = \phi(x - x_0) = 6 \tanh(\frac{x - x_0}{6})$$
 so that $\phi'_{x_0} = \psi^2_{x_0}$

Proof. In the following, we will set $\phi = \phi_{x_0}$ for simplicity. First, we use (2.14) to write

(3.27)
$$2\int \phi \, d\rho|_{u}(\psi^{2}f)' \, dx = \int \phi \left\{ \frac{1}{g^{2}} dg|_{u}(\psi^{2}f)' + 4\kappa R_{0}(2\kappa)(\psi^{2}f)' \right\} \, dx$$
$$= \int \frac{\phi}{g^{2}} \left\{ dg|_{u}(\psi^{2}f)' + \kappa^{-1}R_{0}(2\kappa)(\psi^{2}f)' \right\} \, dx$$

(3.28)
$$-\kappa^{-1} \int \phi\left(\frac{1}{g} + 2\kappa\right) \left(\frac{1}{g} - 2\kappa + \frac{4\kappa^2 h_1}{1 + 2\kappa h_1}\right) R_0(2\kappa) (\psi^2 f)' dx$$

(3.29)
$$+ \kappa^{-1} \int \phi \left(\frac{1}{g} + 2\kappa\right) \frac{4\kappa^2 h_1}{1 + 2\kappa h_1} R_0(2\kappa) (\psi^2 f)' \, dx.$$

We will estimate the contributions of (3.27)-(3.29) individually.

Let us start with (3.28). By (2.7), (3.21), and (2.17) we have

$$\kappa^{-1} \int_{-T}^{T} \left| \int \phi \left(\frac{1}{g} + 2\kappa \right) \left(\frac{1}{g} - 2\kappa + \frac{4\kappa^{2}h_{1}}{1 + 2\kappa h_{1}} \right) R_{0}(2\kappa) (\psi^{2}f)' dx \right| dt$$

$$\lesssim \kappa^{-1} \|\phi\|_{L^{\infty}} \left\| \frac{1}{g} + 2\kappa \right\|_{L^{\infty}_{t,x}} \left\| \psi \left(\frac{1}{g} - 2\kappa + \frac{4\kappa^{2}h_{1}}{1 + 2\kappa h_{1}} \right) \right\|_{L^{2}_{t,x}} \left\| \frac{1}{\psi} R_{0} \partial \psi \right\|_{H^{-1}_{\kappa} \to L^{2}} \|\psi f\|_{L^{2}_{t}H^{-1}_{\kappa}}$$

$$\lesssim \kappa^{-\frac{3}{2}} (T\kappa^{2})^{\frac{1}{4}} \|u\|_{L^{\infty}_{t}H^{-1}_{\kappa}} \left(\|u\|_{L^{\infty}_{t}H^{-1}_{\kappa}} + \|u\|_{\mathrm{LS}_{\kappa}} \right) \|\psi f\|_{L^{2}_{t}H^{-1}_{\kappa}}.$$

Next, we turn to (3.29). Integrating by parts and using (2.17), (2.7), (3.11), (3.22), (3.8), and (3.10), we have

$$\begin{split} \kappa^{-1} \int_{-T}^{T} \left| \int \phi \left(\frac{1}{g} + 2\kappa \right) \frac{4\kappa^{2}h_{1}}{1 + 2\kappa h_{1}} R_{0}(2\kappa) (\psi^{2}f)' \, dx \right| \, dt \\ \lesssim \kappa^{-1} \left\| \psi f \right\|_{L_{t}^{2}H_{\kappa}^{-1}} \left\| \psi R_{0} \frac{1}{\psi} \right\|_{L^{2} \to H_{\kappa}^{1}} \left\| \psi \left[\phi \left(\frac{1}{g} + 2\kappa \right) \frac{4\kappa^{2}h_{1}}{1 + 2\kappa h_{1}} \right]' \right\|_{L_{t,x}^{2}} \\ \lesssim \kappa^{-2} \left\| \psi f \right\|_{L_{t}^{2}H_{\kappa}^{-1}} \left\{ \left\| \frac{1}{g} + 2\kappa \right\|_{L_{t,x}^{\infty}} \left\| \psi h_{1}' \right\|_{L_{t,x}^{2}} \left\| \frac{4\kappa^{2}}{1 + 2\kappa h_{1}} \right\|_{L_{t,x}^{\infty}} \left(1 + \left\| \frac{2\kappa h_{1}}{1 + 2\kappa h_{1}} \right\|_{L_{t,x}^{\infty}} \right) \right\|_{H_{t,x}^{2}} \end{split}$$

$$+ \|\psi g'\|_{L^{2}_{t,x}} \left\|\frac{1}{g}\right\|^{2}_{L^{\infty}_{t,x}} \left\|\frac{4\kappa^{2}h_{1}}{1+2\kappa h_{1}}\right\|_{L^{\infty}_{t,x}} + \left\|\frac{1}{g} + 2\kappa\right\|_{L^{\infty}_{t,x}} \left\|\frac{4\kappa^{2}}{1+2\kappa h_{1}}\right\|_{L^{\infty}_{t,x}} \|\psi h_{1}\|_{L^{2}_{t,x}} \right\} \\ \lesssim \kappa^{-1} \|\psi f\|_{L^{2}_{t}H^{-1}_{\kappa}} \left(\|u\|_{L^{\infty}_{t}H^{-1}_{\kappa}} + \|u\|_{\mathrm{LS}_{\kappa}}\right).$$

It remains to estimate (3.27). We write

$$\int \frac{\phi}{g^2} \Big\{ dg|_u (\psi^2 f)' + \kappa^{-1} R_0(2\kappa) (\psi^2 f)' \Big\} dx$$

= $\sum_{\ell+m \ge 1} (-1)^{\ell+m+1} \operatorname{tr} \Big\{ \frac{\phi}{g^2} (R_0 u)^\ell R_0 \psi^2 [\partial, \psi^2 f] (R_0 u)^m R_0 \Big\}$
= $\sum_{\ell+m \ge 1} (-1)^{\ell+m} \operatorname{tr} \Big\{ \psi^2 f \Big[\partial, (R_0 u)^\ell R_0 \frac{\phi}{g^2} (R_0 u)^m R_0 \Big] \Big\}.$

In the last equality, we cycled the trace (i.e. tr(AB) = tr(BA)). Now, we will distribute the derivative ∂ using the product rule $[\partial, AB] = [\partial, A]B + A[\partial, B]$.

First, let us consider the terms with $\ell + m = 1$. When the derivative lands on u, we use (2.17) and (2.7) to estimate:

$$\begin{split} \int_{-T}^{T} \left| \operatorname{tr} \left\{ \psi^{2} f R_{0} u' R_{0} \frac{\phi}{g^{2}} R_{0} \right\} \right| + \left| \operatorname{tr} \left\{ \psi^{2} f R_{0} \frac{\phi}{g^{2}} R_{0} u' R_{0} \right\} \right| dt \\ & \lesssim \left\| \sqrt{R_{0}} \psi f \sqrt{R_{0}} \right\|_{L_{t}^{2} \mathfrak{I}_{2}} \left\| \sqrt{R_{0}} \psi u' \sqrt{R_{0}} \right\|_{L_{t}^{2} \mathfrak{I}_{2}} \left\| \sqrt{R_{0}} \frac{\phi}{g^{2}} \sqrt{R_{0}} \right\|_{L_{t}^{\infty} \mathfrak{I}_{\infty}} \\ & \lesssim \kappa^{-1} \left\| \psi f \right\|_{L_{t}^{2} H_{\kappa}^{-1}} \left\| u \right\|_{\mathrm{LS}_{\kappa}}. \end{split}$$

When the derivative lands on $\frac{\phi}{g^2}$, we use (2.4), (2.7), and (3.9) to estimate

(3.30)
$$\begin{aligned} \|\sqrt{R_0}(\frac{\phi}{g^2})'\sqrt{R_0}\|_{L^2_t\mathfrak{I}_\infty} &\lesssim \kappa^{-\frac{3}{2}}T^{\frac{1}{2}} \|(\frac{\phi}{g^2})'\|_{L^\infty_t L^2_x} \lesssim \kappa^{-\frac{3}{2}}T^{\frac{1}{2}} \left(\left\|\frac{1}{g}\right\|^2_{L^\infty_{t,x}} + \left\|\frac{1}{g}\right\|^3_{L^\infty_{t,x}} \|g'\|_{L^2_{t,x}}\right) \\ &\lesssim \kappa^{\frac{1}{2}}T^{\frac{1}{2}} \left(1 + \|u\|_{L^\infty_t H^{-1}_\kappa}\right) \lesssim 1. \end{aligned}$$

This yields

$$\int_{-T}^{T} \left| \operatorname{tr} \left\{ \psi^{2} f R_{0} u R_{0} \left(\frac{\phi}{g^{2}} \right)' R_{0} \right\} \right| + \left| \operatorname{tr} \left\{ \psi^{2} f R_{0} \left(\frac{\phi}{g^{2}} \right)' R_{0} u R_{0} \right\} \right| dt$$

$$\lesssim \left\| \sqrt{R_{0}} \psi^{2} f \sqrt{R_{0}} \right\|_{L_{t}^{2} \mathfrak{I}_{2}} \left\| \sqrt{R_{0}} u \sqrt{R_{0}} \right\|_{L_{t}^{\infty} \mathfrak{I}_{2}} \left\| \sqrt{R_{0}} \left(\frac{\phi}{g^{2}} \right)' \sqrt{R_{0}} \right\|_{L_{t}^{2} \mathfrak{I}_{\infty}}$$

$$\lesssim \kappa^{-1} \left\| \psi f \right\|_{L_{t}^{2} H_{\kappa}^{-1}} \left\| u \right\|_{L_{t}^{\infty} H_{\kappa}^{-1}}.$$

Lastly, we turn to the terms with $\ell + m \geq 2$. When the derivative lands on u, we use (3.7). When the derivative lands on $\frac{\phi}{g^2}$, we use (3.30). As $\ell + m \geq 2$, there are always at least two terms that we can put in Hilbert–Schmidt norm:

$$\int_{-T}^{T} \left| \operatorname{tr} \left\{ \psi^{2} f \left[\partial, (R_{0}u)^{\ell} R_{0} \frac{\phi}{g^{2}} (R_{0}u)^{m} R_{0} \right] \right\} \right| dt \\
\lesssim \left\| \sqrt{R_{0}} \psi^{2} f \sqrt{R_{0}} \right\|_{L_{t}^{2} \mathfrak{I}_{2}} \left\{ \left\| \sqrt{R_{0}} \left(\frac{\phi}{g^{2}} \right)' \sqrt{R_{0}} \right\|_{L_{t}^{2} \mathfrak{I}_{\infty}} \left\| \sqrt{R_{0}} u \sqrt{R_{0}} \right\|_{L_{t}^{\infty} \mathfrak{I}_{2}} \right. \\
\left. + (\ell + m) \left\| \sqrt{R_{0}} u' \sqrt{R_{0}} \right\|_{L_{t}^{2} \mathfrak{I}_{\infty}} \left\| \sqrt{R_{0}} \frac{\phi}{g^{2}} \sqrt{R_{0}} \right\|_{L_{t}^{\infty} \mathfrak{I}_{\infty}} \left\| \sqrt{R_{0}} u \sqrt{R_{0}} \right\|_{L_{t}^{\infty} \mathfrak{I}_{2}}^{\ell + m - 1} \right\} \\
\lesssim \kappa^{-1} \left\| \psi f \right\|_{L_{t}^{2} H_{\kappa}^{-1}} \left(\left\| u \right\|_{L_{t}^{\infty} H_{\kappa}^{-1}} + \left\| u \right\|_{\mathrm{LS}_{\kappa}} \right) (\ell + m + 1) \left(\kappa^{-\frac{1}{2}} \left\| u \right\|_{L_{t}^{\infty} H_{\kappa}^{-1}} \right)^{\ell + m - 1}.$$

Summing over $\ell + m \geq 2$ then yields

$$\sum_{\ell+m\geq 2} \int_{-T}^{T} \left| \operatorname{tr} \left\{ \psi^2 f \left[\partial, (R_0 u)^{\ell} R_0 \frac{\phi}{g^2} (R_0 u)^m R_0 \right] \right\} \right| dt \lesssim \kappa^{-1} \|\psi f\|_{L^2_t H^{-1}_{\kappa}} \left(\|u\|_{L^{\infty}_t H^{-1}_{\kappa}} + \|u\|_{\mathrm{LS}_{\kappa}} \right).$$

This was the final term that we needed to estimate, and thus concludes the proof of the lemma. \square

4. The local smoothing estimate

In this section, we will prove our main local smoothing estimate for solutions to (gKdV):

Theorem 4.1. Let $\kappa \geq 1$, and u a solution to (gKdV) which satisfies the bound (3.9) uniformly in a time interval [0,T] with $T \leq \kappa^{-2}$. For some $\varepsilon > 0$ assume that the coefficients a_1, a_2, a_3, a_4 obey the following bounds in [0, T], uniformly in $z \in \mathbb{R}$:

(4.1)
$$\int \|\psi_z a_1\|_{L^{\infty}_t H^1_x} dz \leq \varepsilon \quad or \quad \int \|\psi_z a_1\|_{L^{\infty}_t W^{1,\infty}_x} dz \leq \varepsilon,$$

(4.2)
$$\int \|\psi_z a_2\|_{L^{\infty}_{t,x}} \, dz \le \varepsilon_z$$

(4.3)
$$\int \|\psi_z a_3\|_{L^2_t H^1_x} dz \le \varepsilon \quad or \quad \int \|\psi_z a_3\|_{L^2_t W^{1,\infty}_x} dz \le \varepsilon,$$

(4.4)
$$||a_4||_{L^{\infty}_t H^1_x} < \infty \quad or \quad ||a_4||_{L^{\infty}_t W^{1,\infty}_x} < \infty$$

Then the solution u(t) to (gKdV) satisfies the local energy estimate

(4.5)
$$\|u\|_{\mathrm{LS}_{\kappa}}^{2} \leq 2 \|u\|_{C_{t}H_{\kappa}^{-1}}^{2} + C(\varepsilon + (T\kappa^{2})^{\frac{1}{4}} + \kappa^{-2}) \left(\|u\|_{L_{t}^{\infty}H_{\kappa}^{-1}}^{2} + \|u\|_{\mathrm{LS}_{\kappa}}^{2} \right)$$

For the definition of ψ_{x_0} and LS_{κ} , see (3.1). The rest of the section is devoted to the proof of the theorem.

Proof of Theorem 4.1. Recall that solutions u(t) to (gKdV) obey the approximate conservation law (2.13). In order to prove the local smoothing estimate, we will multiply (2.13) by the function ϕ defined in (3.26) and integrate in space and time:

$$\int_{-T}^{T} \int \psi_{x_0}^2 j(t,x) \, dx \, dt = \int \phi_{x_0} [\rho(T,x) - \rho(-T,x)] \, dx$$
$$- \int_{-T}^{T} \int \phi_{x_0} \, d\rho |_u [(a_1 u')' + a_2 u^2 + a_3 u' + a_4 u] \, dx \, dt.$$

Here we seek to identify the left hand side, respectively the first term on the right, with the left hand side, respectively the first term on the right, in (4.5) modulo acceptable errors.

We begin with the first term on the right, which is easiest. Recall that $\rho(t, x) \ge 0$, and so $\|\rho\|_{L^1} = \alpha(\kappa, u)$. For the first term on the RHS, we use (2.10) to estimate

(4.6)
$$\kappa \left| \int \phi_{x_0} [\rho(T, x) - \rho(-T, x)] \, dx \right| \le 2 \, \|u\|_{C_t H_{\kappa}^{-1}}^2$$

We next consider the left hand side, for which we will show that

(4.7)
$$\kappa \int_{-T}^{T} \int \psi_{x_0}^2 j(t,x) = -\frac{3}{2} \left\| \psi_{x_0} u' \right\|_{L_t^2 H_\kappa^{-1}}^2 dx \, dt + Err_0,$$

where for brevity Err_0 is an expression satisfying the bound

$$|Err_0| \lesssim ((T\kappa^2)^{\frac{1}{4}} + \kappa^{-2}) \big(\|u\|_{L^{\infty}_t H^{-1}_{\kappa}}^2 + \|u\|_{\mathrm{LS}_{\kappa}}^2 \big).$$

The last task is then to estimate the second term on the right, and show that

(4.8)
$$\kappa \int_{-T}^{T} \int \phi_{x_0} d\rho |_u \left[(a_1 u')' + a_2 u^2 + a_3 u' + a_4 u \right] dx \, dt = Err_1,$$

where

$$|Err_1| \lesssim (\varepsilon + \kappa^{-2}) (\|u\|_{L^{\infty}_t H^{-1}_{\kappa}}^2 + \|u\|_{\mathrm{LS}_{\kappa}}^2).$$

The desired estimate (4.5) is obtained by combining (4.6), (4.7) and (4.8) followed by taking the supremum over $x_0 \in \mathbb{R}$. It remains to prove the bounds (4.7) and (4.8).

We begin with the proof of (4.7). To leading order, j is quadratic in u. Specifically, if we insert the series (2.8) for g, then the quadratic terms of j are:

$$j_2 = 8\kappa^4 h_2 - h_1(16\kappa^5 h_1 + 4\kappa^2 u) - 6\kappa R_0(2\kappa)[u^2].$$

It is then natural to expect the leading contribution to come from j_2 , while the cubic and higher contributions coming from $j - j_2$ to be perturbative. We do this in the next two lemmas. In the first lemma we examine the contribution of j_2 , and show that this generates the local smoothing norm of u:

Lemma 4.2. Assume that $\kappa \geq 1$ and u satisfies (3.9) uniformly in [0, T]. Then we have

$$\left\|\psi_{x_0}u'\right\|_{L^2_t H^{-1}_{\kappa}}^2 = -\frac{2}{3}\kappa \int_{-T}^{T} \int_{-\infty}^{\infty} \psi_{x_0}^2 j_2 \, dx \, dt + O\left(\kappa^{-2} \left(\left\|u\right\|_{L^{\infty}_t H^{-1}_{\kappa}}^2 + \left\|u\right\|_{\mathrm{LS}_{\kappa}}^2\right)\right)$$

uniformly for $x_0 \in \mathbb{R}$.

Proof. We need κj_2 to be O(1), so there must be some cancellation for the terms that are $O(\kappa)$ and higher. In order to exhibit this cancellation, we use the identities (cf. [6, Lem. 2.5])

(4.9)
$$16\kappa^{5}h_{2} = 3u^{2} - 3\kappa^{2}(h_{1}'')^{2} - 20\kappa^{4}\left[(h_{1}')^{2} - (h_{1}^{2})''\right] + 4\kappa^{4}\partial^{2}R_{0}(2\kappa)\left[(h_{1}')^{2} + 2(h_{1}^{2})''\right],$$
$$16\kappa^{5}h_{1} + 4\kappa^{2}u = 4\kappa^{3}h_{1}''$$

to write

$$-2\kappa j_2 = 3\kappa^2 (h_1'')^2 + 12\kappa^4 (h_1')^2 - 3\left[u^2 - 4\kappa^2 R_0(2\kappa)u^2\right] - 4\kappa^4 \partial^2 R_0(2\kappa) \left[(h_1')^2 + 2(h_1^2)''\right].$$

We multiply this by $\psi_{x_0}^2$ and integrate in space and time. Working from left to right, we claim that

(4.10)
$$\int_{-T}^{T} \langle \psi^{2}, 3\kappa^{2}(h_{1}'')^{2} \rangle dt = 3 \int_{-T}^{T} \langle (\psi u')', R_{0}^{2}(\psi u')' \rangle dt + O\left(\kappa^{-2} \|u\|_{\mathrm{LS}_{\kappa}}^{2}\right),$$

(4.11)
$$\int_{-T}^{T} \langle \psi^{2}, 12\kappa^{4}(h_{1}')^{2} \rangle dt = 3 \int_{-T}^{T} \langle \psi u', R_{0}\psi u' \rangle - \langle (\psi u')', R_{0}^{2}(\psi u')' \rangle dt$$
$$+ O\left(\kappa^{-2} \|u\|_{\mathrm{LS}_{\kappa}}^{2}\right),$$

(4.12)
$$\int_{-T}^{T} \langle \psi^2, (\text{rest of } -2\kappa j_2) \rangle \, dt = O\left(\kappa^{-2} \left(\|u\|_{L^\infty_t H^{-1}_\kappa}^2 + \|u\|_{\mathrm{LS}_\kappa}^2 \right) \right)$$

uniformly for $x_0 \in \mathbb{R}$, where $\psi = \psi_{x_0}$ and $R_0 = R_0(2\kappa)$. Adding these together, this would yield

$$\int_{-T}^{T} -2\kappa \langle \psi^2, j_2 \rangle \, dt = 3 \int_{-T}^{T} \langle \psi u', R_0 \psi u' \rangle \, dt + O\left(\kappa^{-\frac{1}{2}} \left(\|u\|_{L_t^{\infty} H_{\kappa}^{-1}}^2 + \|u\|_{\mathrm{LS}_{\kappa}}^2 \right) \right).$$

The first term on the RHS is exactly $\|\psi u'\|_{L^2_t H^{-1}_{\kappa}}^2$, and so this would finish the proof.

Let us start with (4.10). As $h_1 = -\kappa^{-1} R_0(2\kappa) u$, we have

$$\kappa^2 \langle \psi^2, (h_1'')^2 \rangle = \langle \psi R_0 u'', \psi R_0 u'' \rangle$$

It remains to replace $\psi R_0 \partial$ by $R_0 \partial \psi$ above: by (2.19) we have

$$\begin{split} \int_{-T}^{T} \left| \langle \psi R_0 u'', \psi R_0 u'' \rangle - \langle R_0(\psi u')', R_0(\psi u')' \rangle \right| dt \\ &= \int_{-T}^{T} \left| \langle \psi u', \frac{1}{\psi} \left(R_0 \partial \psi^2 R_0 \partial - \psi R_0^2 \partial^2 \psi \right) \frac{1}{\psi} \cdot \psi u' \rangle \right| dt \\ &\lesssim \| \psi u' \|_{L_t^2 H_\kappa^{-2}}^2 \\ &\lesssim \kappa^{-2} \| u \|_{\mathrm{LS}_\kappa}^2 \,. \end{split}$$

This proves (4.10).

The proof of (4.11) proceeds in a similar way. We write

$$\psi^2, 12\kappa^4(h_1')^2 \rangle = 12\kappa^2 \langle \psi R_0 u', \psi R_0 u' \rangle.$$

First, we use (2.18) to replace ψR_0 by $R_0 \psi$ above:

$$12\kappa^{2} \int_{-T}^{T} \left| \langle \psi R_{0}u', \psi R_{0}u' \rangle - \langle R_{0}\psi u', R_{0}\psi u' \rangle \right| dt$$

$$= 12\kappa^{2} \int_{-T}^{T} \left| \langle \psi u', \frac{1}{\psi} \left(R_{0}\psi^{2}R_{0} - \psi R_{0}^{2}\psi \right) \frac{1}{\psi} \cdot \psi u' \rangle \right| dt$$

$$\lesssim \left\| \psi u' \right\|_{L_{t}^{2}H_{\kappa}^{-2}}^{2}$$

$$\lesssim \kappa^{-2} \left\| u \right\|_{\mathrm{LS}_{\kappa}}^{2}.$$

Next we use the identity $4\kappa^2 R_0 = 1 + \partial^2 R_0$ to write

$$12\kappa^2 \langle R_0 \psi u', R_0 \psi u' \rangle = 3 \langle \psi u', R_0 \psi u' \rangle - 3 \langle (\psi u')', R_0^2 (\psi u')' \rangle,$$

and (4.11) follows.

Lastly, we turn to (4.12). There are two terms remaining in $-2\kappa j_2$ which we want to show make a negligible contribution. For the first term, we use the identity $1 - 4\kappa^2 R_0 = -\partial^2 R_0$ to write

$$\langle \psi^2, 3[u^2 - 4\kappa^2 R_0 u^2] \rangle = 3\langle \psi^2, R_0(u^2)'' \rangle = 3\langle \frac{1}{\psi^2} R_0(\psi^2)'', \psi^2 u^2 \rangle.$$

Therefore, by (2.17) and (3.2)–(3.4) we have for $T \leq \kappa^{-2}$

$$\int_{-T}^{T} \left| \langle \psi^2, 3 \left[u^2 - 4\kappa^2 R_0 u^2 \right] \rangle \right| dt \leq 3 \left\| \frac{1}{\psi^2} R_0(\psi^2)'' \right\|_{L^{\infty}_x} \| \psi u \|_{L^2_{t,x}}^2$$
$$\lesssim \kappa^{-2} \left\| (-\partial^2 + 4\kappa^2)(\psi u) \right\|_{L^2_t H^{-2}_\kappa}^2$$

$$\lesssim \kappa^{-2} (\|u\|_{L^{\infty}_{t}H^{-1}_{\kappa}}^{2} + \|u\|_{\mathrm{LS}_{\kappa}}^{2}).$$

For the last term of $-2\kappa j_2$, we write

$$\langle \psi^2, 4\kappa^4 \partial^2 R_0[(h_1')^2 + 2(h_1^2)''] \rangle = 4\kappa^2 \langle R_0(\psi^2)'', (R_0u')^2 \rangle + 4\kappa^2 \langle R_0(\psi^2)^{(4)}, (R_0u)^2 \rangle,$$

in order to estimate

$$\begin{split} &\int_{-T}^{T} \left| \langle \psi^{2}, 4\kappa^{4} \partial^{2} R_{0}[(h_{1}')^{2} + 2(h_{1}^{2})''] \rangle \right| dt \\ &\lesssim \kappa^{2} \Big\{ \left\| \frac{1}{\psi^{2}} R_{0}(\psi^{2})'' \right\|_{L_{x}^{\infty}} \left\| \psi R_{0} \frac{1}{\psi} \cdot \psi u' \right\|_{L_{t,x}^{2}}^{2} + \left\| \frac{1}{\psi^{2}} R_{0}(\psi^{2})^{(4)} \right\|_{L_{x}^{\infty}} \left\| \psi R_{0} \frac{1}{\psi} \cdot \psi u \right\|_{L_{t,x}^{2}}^{2} \Big\} \\ &\lesssim \kappa^{2} \Big\{ \kappa^{-2} \left\| \psi u' \right\|_{L_{t}^{2} H_{\kappa}^{-2}}^{2} + \kappa^{-2} \left\| \psi u \right\|_{L_{t}^{2} H_{\kappa}^{-2}}^{2} \Big\} \lesssim \kappa^{-2} \Big(\left\| u \right\|_{L_{t}^{\infty} H_{\kappa}^{-1}}^{2} + \left\| u \right\|_{\mathrm{LS}_{\kappa}}^{2} \Big). \end{split}$$

Altogether, this proves (4.12).

In the next lemma we show that the cubic and higher order terms of j make a negligible contribution:

Lemma 4.3. Assume that $\kappa \geq 1$ and u satisfies (3.9) uniformly in [0,T] with $T \leq \kappa^{-2}$. Then we have

(4.13)
$$\left| \int_{-T}^{T} \int_{-\infty}^{\infty} \psi_{x_0}^2(j-j_2) \, dx \, dt \right| \lesssim \kappa^{-1} (T\kappa^2)^{\frac{1}{4}} \left(\|u\|_{L^{\infty}_t H^{-1}_{\kappa}}^2 + \|u\|_{\mathrm{LS}_{\kappa}}^2 \right)$$

uniformly in $x_0 \in \mathbb{R}$.

Proof. In order to exhibit cancellation in $j - j_2$, we expand g as the series (2.8) in the expression (2.12) for j:

(4.14)
$$j - j_2 = \frac{1}{g} (4\kappa^3 h_1 + u) + 2\kappa R_0 (2\kappa) u'' + h_1 (16\kappa^5 h_1 + 4\kappa^2 u)$$

$$(4.15) \qquad \qquad + \left(\frac{1}{g} - 2\kappa\right)4\kappa^3h_2$$

(4.16)
$$+4\kappa^3 \frac{1}{g} \sum_{\ell \ge 3} h_\ell.$$

We will estimate the contributions (4.14)–(4.16) one at a time.

Let us start with (4.14). Using the identity (4.9), we write

$$(4.14) = \kappa \left(\frac{1}{g} - 2\kappa + 4\kappa^2 h_1\right) h_1'' = \kappa \left(\frac{1}{g} - 2\kappa + \frac{4\kappa^2 h_1}{1 + 2\kappa h_1}\right) h_1'' + \frac{8\kappa^4}{1 + 2\kappa h_1} h_1^2 h_1''.$$

For the first term on the RHS, we use (3.21) and (3.12) to bound:

$$\begin{split} \left| \int_{-T}^{T} \int_{-\infty}^{\infty} \psi^{2} \kappa \left(\frac{1}{g} - 2\kappa + \frac{4\kappa^{2}h_{1}}{1 + 2\kappa h_{1}} \right) h_{1}'' \, dx \, dt \right| \\ \lesssim \kappa \left\| \psi \left(\frac{1}{g} - 2\kappa + \frac{4\kappa^{2}h_{1}}{1 + 2\kappa h_{1}} \right) \right\|_{L^{2}_{t,x}} \left\| \psi h_{1}'' \right\|_{L^{2}_{t,x}} \\ \lesssim \kappa^{-\frac{3}{2}} (T\kappa^{2})^{\frac{1}{4}} \left\| u \right\|_{L^{\infty}_{t}H^{-1}_{\kappa}} \left(\left\| u \right\|_{L^{\infty}_{t}H^{-1}_{\kappa}}^{2} + \left\| u \right\|_{\mathrm{LS}_{\kappa}}^{2} \right). \end{split}$$

For the second term, we use (3.22), (3.10), and (3.12) to bound:

$$\begin{split} \left| \int_{-T}^{T} \int_{-\infty}^{\infty} \psi^{2} \frac{8\kappa^{4}}{1+2\kappa h_{1}} h_{1}^{2} h_{1}^{\prime \prime} \, dx \, dt \right| &\lesssim \kappa^{\frac{7}{2}} \left\| \frac{1}{1+2\kappa h_{1}} \right\|_{L_{t,x}^{\infty}} \|h_{1}\|_{L_{t}^{\infty} H_{\kappa}^{1}} \|\psi h_{1}\|_{L_{t,x}^{2}} \, \|\psi h_{1}^{\prime \prime}\|_{L_{t,x}^{2}} \\ &\lesssim \kappa^{-\frac{1}{2}} T^{\frac{1}{2}} \, \|u\|_{L_{t}^{\infty} H_{\kappa}^{-1}}^{2} \left(\|u\|_{L_{t}^{\infty} H_{\kappa}^{-1}} + \|u\|_{\mathrm{LS}_{\kappa}} \right) \\ &\lesssim \kappa^{-\frac{3}{2}} (T\kappa^{2})^{\frac{1}{2}} \, \|u\|_{L_{t}^{\infty} H_{\kappa}^{-1}}^{-1} \left(\|u\|_{L_{t}^{\infty} H_{\kappa}^{-1}}^{2} + \|u\|_{\mathrm{LS}_{\kappa}}^{2} \right). \end{split}$$

Next, we turn to (4.15). We write

$$(4.15) = 4\kappa^3 \left(\frac{1}{g} - 2\kappa + \frac{4\kappa^2 h_1}{1 + 2\kappa h_1}\right) h_2 - \frac{16\kappa^5}{1 + 2\kappa h_1} h_1 h_2.$$

For the first term on the RHS, we use (3.21) and (3.20) to estimate

$$\begin{split} \left| \int_{-T}^{T} \int_{-\infty}^{\infty} \psi^{2} 4\kappa^{3} \left(\frac{1}{g} - 2\kappa + \frac{4\kappa^{2}h_{1}}{1 + 2\kappa h_{1}} \right) h_{2} \, dx \, dt \right| \\ &\lesssim \kappa^{3} \left\| \psi \left(\frac{1}{g} - 2\kappa + \frac{4\kappa^{2}h_{1}}{1 + 2\kappa h_{1}} \right) \right\|_{L^{2}_{t,x}} \|\psi h_{2}\|_{L^{2}_{t,x}} \\ &\lesssim \kappa^{-2} (T\kappa^{2})^{\frac{1}{2}} \|u\|_{L^{\infty}_{t}H^{-1}_{\kappa}}^{2} \left(\|u\|_{L^{\infty}_{t}H^{-1}_{\kappa}}^{2} + \|u\|_{\mathrm{LS}_{\kappa}}^{2} \right) \\ &\lesssim \kappa^{-\frac{3}{2}} (T\kappa^{2})^{\frac{1}{2}} \|u\|_{L^{\infty}_{t}H^{-1}_{\kappa}}^{2} \left(\|u\|_{L^{\infty}_{t}H^{-1}_{\kappa}}^{2} + \|u\|_{\mathrm{LS}_{\kappa}}^{2} \right). \end{split}$$

For the second term, we use (3.23) to bound

$$\left| \int_{-T}^{T} \int_{-\infty}^{\infty} \psi^{2} \frac{16\kappa^{5}}{1+2\kappa h_{1}} h_{1}h_{2} \, dx \, dt \right| \lesssim \kappa^{5} \left\| \frac{1}{1+2\kappa h_{1}} \right\|_{L^{\infty}_{t,x}} \left\| \psi^{2}h_{1}h_{2} \right\|_{L^{1}_{t,x}} \\ \lesssim \kappa^{-\frac{3}{2}} (T\kappa^{2})^{\frac{3}{4}} \left\| u \right\|_{L^{\infty}_{t}H^{-1}_{\kappa}} \left(\left\| u \right\|_{L^{\infty}_{t}H^{-1}_{\kappa}}^{2} + \left\| u \right\|_{\mathrm{LS}_{\kappa}}^{2} \right)$$

Finally, we turn to (4.16). Using (3.13), we estimate

$$\left| \int_{-T}^{T} \int_{-\infty}^{\infty} \psi^{2}(4.16) \, dx \, dt \right| \lesssim \kappa^{3} \left\| \frac{1}{g} \right\|_{L_{t,x}^{\infty}} \sum_{\ell \ge 3} \left\| \psi^{2} h_{\ell} \right\|_{L_{t,x}^{1}} \\ \lesssim \kappa^{-\frac{3}{2}} (T\kappa^{2})^{\frac{3}{4}} \left\| u \right\|_{L_{t}^{\infty} H_{\kappa}^{-1}} \left(\left\| u \right\|_{L_{t}^{\infty} H_{\kappa}^{-1}}^{2} + \left\| u \right\|_{\mathrm{LS}_{\kappa}}^{2} \right).$$

Altogether, we obtain

$$\left| \int_{-T}^{T} \int_{-\infty}^{\infty} \psi_{x_{0}}^{2} (j - j_{2}) \, dx \, dt \right| \lesssim \kappa^{-\frac{3}{2}} (T \kappa^{2})^{\frac{1}{4}} \, \|u\|_{L_{t}^{\infty} H_{\kappa}^{-1}} \left(\, \|u\|_{L_{t}^{\infty} H_{\kappa}^{-1}}^{2} + \|u\|_{\mathrm{LS}_{\kappa}}^{2} \right).$$

This implies (4.13) by (3.9), and thus finishes the proof of the lemma.

The last step in the proof of the theorem is to prove the estimate (4.8), which shows that the contributions of the source terms in the density flux relation (2.13) are perturbative:

Lemma 4.4. Assume that $\kappa \geq 1$ and u satisfies (3.9) uniformly in [0,T], and that the coefficients a_1, a_2, a_3, a_4 in (gKdV) satisfy (4.1)-(4.4). Then

$$(4.17) \left| \int_{-T}^{T} \int \phi_{x_0} d\rho |_u \left[(a_1 u')' + a_2 u^2 + a_3 u' + a_4 u \right] dx dt \right| \lesssim \kappa^{-1} (\varepsilon + \kappa^{-2}) \left(\|u\|_{L^{\infty}_t H^{-1}_{\kappa}}^2 + \|u\|_{\mathrm{LS}_{\kappa}}^2 \right)$$
uniformly in $x_0 \in \mathbb{R}$

uniformly in $x_0 \in \mathbb{R}$.

Proof. We begin with the contribution of $(a_1u')'$. Write

(4.18)
$$a_1(t,x) = c \int a_1(t,x) \psi_z^3(x) \, dz$$

for a constant c. By (4.1) we have

$$\int \left\| \psi_z^2 a_1 u' \right\|_{L^2_t H^{-1}_{\kappa}} dz \lesssim \int \left\| \psi_z a_1 \right\|_{L^{\infty}_t X} \left\| \psi_z u' \right\|_{L^2_t H^{-1}_{\kappa}} dz \lesssim \varepsilon \left\| u \right\|_{\mathrm{LS}_{\kappa}},$$

where $X = H^1$ or $X = W^{1,\infty}$. Therefore, by (3.25) we have

$$\left| \int_{-T}^{T} \int \phi \, d\rho |_{u}(a_{1}u')' \, dx \, dt \right| \lesssim \varepsilon \kappa^{-1} \left(\left\| u \right\|_{L_{t}^{\infty} H_{\kappa}^{-1}} + \left\| u \right\|_{\mathrm{LS}_{\kappa}} \right) \left\| u \right\|_{\mathrm{LS}_{\kappa}},$$

and this is an acceptable contribution to (4.17).

For the contributions of a_2u^2 , a_3u' , and a_4u , we use (3.24) to estimate

(4.19)
$$\left| \int_{-T}^{T} \int \phi \, d\rho |_{u} (a_{2}u^{2} + a_{3}u' + a_{4}u) \, dx \, dt \right| \\ \lesssim \kappa^{-1} \, \|u\|_{L_{t}^{\infty}H_{\kappa}^{-1}} \left(\|a_{2}u^{2}\|_{L_{t}^{1}H_{\kappa}^{-1}} + \|a_{3}u'\|_{L_{t}^{1}H_{\kappa}^{-1}} + \|a_{4}u\|_{L_{t}^{1}H_{\kappa}^{-1}} \right).$$

For a_4u , we use (4.4) and (3.9) to bound

$$\|a_4 u\|_{L^1_t H^{-1}_{\kappa}} \lesssim T \|a_4\|_{L^{\infty}_t X} \|u\|_{L^{\infty}_t H^{-1}_{\kappa}} \lesssim \kappa^{-2} \|u\|_{L^{\infty}_t H^{-1}_{\kappa}}$$

where $X = H^1$ or $X = W^{1,\infty}$.

For a_3u' , we use (4.18) and (4.3) to bound

$$\|a_{3}u'\|_{L_{t}^{1}H_{\kappa}^{-1}} \lesssim \int \|\psi_{z}^{2}a_{3}u'\|_{L_{t}^{1}H_{\kappa}^{-1}} dz \lesssim \|u\|_{\mathrm{LS}_{\kappa}} \int \|\psi_{z}a_{3}\|_{L_{t}^{2}X} dz \lesssim \varepsilon \|u\|_{\mathrm{LS}_{\kappa}},$$

where $X = H^1$ or $X = W^{1,\infty}$. Finally, for a_2u^2 , we use (3.2)–(3.4) to bound

(4.20)
$$\|\psi_{z}u\|_{L^{2}_{t,x}} \lesssim \|u\|_{\mathrm{LS}_{\kappa}} + \kappa \|u\|_{L^{2}_{t}H^{-1}_{\kappa}} \lesssim \|u\|_{\mathrm{LS}_{\kappa}} + \|u\|_{L^{\infty}_{t}H^{-1}_{\kappa}}$$

provided that $T \leq \kappa^{-2}$. Using (4.18), the embedding $L^1 \hookrightarrow H_{\kappa}^{-1}$, and (4.2), we see that

$$\begin{aligned} \|a_{2}u^{2}\|_{L_{t}^{1}H_{\kappa}^{-1}} &\lesssim \int \|\psi_{z}^{3}a_{2}u^{2}\|_{L_{t}^{1}H_{\kappa}^{-1}} \, dz \lesssim \kappa^{-\frac{1}{2}} \int \|\psi_{z}^{3}a_{2}u^{2}\|_{L_{t,x}^{1}} \, dz \\ &\lesssim \kappa^{-\frac{1}{2}} \int \|\psi_{z}u\|_{L_{t,x}^{2}}^{2} \, \|\psi_{z}a_{2}\|_{L_{t,x}^{\infty}} \, dz \lesssim \varepsilon \kappa^{-\frac{1}{2}} \big(\|u\|_{L_{t}^{\infty}H_{\kappa}^{-1}}^{2} + \|u\|_{\mathrm{LS}_{\kappa}}^{2} \big). \end{aligned}$$

Returning to (4.19), we conclude

$$\left| \int_{-T}^{T} \int \phi \, d\rho |_{u} (a_{2}u^{2} + a_{3}u' + a_{4}u) \, dx \, dt \right| \\ \lesssim \kappa^{-1} (\varepsilon + \kappa^{-2}) \left(\left\| u \right\|_{L_{t}^{\infty} H_{\kappa}^{-1}}^{2} + \left\| u \right\|_{\mathrm{LS}_{\kappa}}^{2} \right) + \varepsilon \kappa^{-\frac{3}{2}} \left\| u \right\|_{L_{t}^{\infty} H_{\kappa}^{-1}} \left(\left\| u \right\|_{\mathrm{LS}_{\kappa}}^{2} + \left\| u \right\|_{L_{t}^{\infty} H_{\kappa}^{-1}}^{2} \right).$$

This implies (4.17) by (3.9), and thus completes the proof.

The last three lemmas complete the proof of Theorem 4.1.

5. The A-priori estimates

In this section, we will prove the energy estimate for solutions to (gKdV), and use it to conclude the proof of our main results in Theorem 1.1 and Theorem 1.5.

Proposition 5.1. For any $\varepsilon \in (0, 1]$, if the coefficients a_1, a_2, a_3, a_4 of (gKdV) satisfy (4.1)–(4.4), then any solution u(t) to (gKdV) satisfies

(5.1)
$$\|u\|_{C_t H_{\kappa}^{-1}}^2 \lesssim \|u(0)\|_{H_{\kappa}^{-1}}^2 + (\varepsilon + T) \left(\|u\|_{L_t^{\infty} H_{\kappa}^{-1}}^2 + \|u\|_{\mathrm{LS}_{\kappa}}^2\right)$$

uniformly for $T \leq \kappa^{-2}$ and κ satisfying (3.9) in [0, T].

Proof. Integrating (2.13) in space and using (2.14), we obtain

$$\partial_t \alpha = \int \left[\frac{1}{2g^2} dg |_u + 2\kappa R_0(2\kappa) \right] \left[(a_1 u')' + a_2 u^2 + a_3 u' + a_4 u \right] dy.$$

Next, we use (2.15) and the identity

$$\int \frac{G(x,y)G(y,x)}{2g(y)^2} \, dy = g(x)$$

(see [38, Lem. 2.5] for a proof) to write

$$\partial_t \alpha = -\int (g - \frac{1}{2\kappa}) \left[(a_1 u')' + a_2 u^2 + a_3 u' + a_4 u \right] dx$$

By (2.10) and the fundamental theorem of calculus, this yields

(5.2)
$$\|u\|_{C_t H_{\kappa}^{-1}}^2 \lesssim \|u(0)\|_{H_{\kappa}^{-1}}^2 + \kappa \int_{-T}^T \left| \int (g - \frac{1}{2\kappa}) \left[(a_1 u')' + a_2 u^2 + a_3 u' + a_4 u \right] dx \right| dt.$$

We will successively estimate the contribution of each of the terms inside the integral.

We begin with the contribution of a_4 . Using (2.6) and (4.4), we have

$$\begin{split} \int_{-T}^{T} \left| \int (g - \frac{1}{2\kappa}) a_4 u \, dx \right| dt &\leq T \left\| g - \frac{1}{2\kappa} \right\|_{L_t^{\infty} H_{\kappa}^1} \| a_4 u \|_{L_t^{\infty} H_{\kappa}^{-1}} \\ &\lesssim \kappa^{-1} T \| a_4 \|_{L_t^{\infty} X} \| u \|_{L_t^{\infty} H_{\kappa}^{-1}}^2 \\ &\lesssim \kappa^{-1} T \| u \|_{L_t^{\infty} H_{\kappa}^{-1}}^2, \end{split}$$

where $X = H^1$ or $X = W^{1,\infty}$.

Next, we address a_3 . Using the continuous partition of unity (4.18) and (4.3), we have

$$\begin{split} \int_{-T}^{T} \left| \int (g - \frac{1}{2\kappa}) a_{3} u' \, dx \right| dt &\leq \left\| g - \frac{1}{2\kappa} \right\|_{L_{t}^{\infty} H_{\kappa}^{1}} \left\| a_{3} u' \right\|_{L_{t}^{1} H_{\kappa}^{-1}} \\ &\lesssim \kappa^{-1} \left\| u \right\|_{L_{t}^{\infty} H_{\kappa}^{-1}} \int \left\| \psi_{z} a_{3} \right\|_{L_{t}^{2} X} \left\| \psi_{z} u' \right\|_{L_{t}^{2} H_{\kappa}^{-1}} \, dz \\ &\lesssim \varepsilon \kappa^{-1} \left\| u \right\|_{L_{t}^{\infty} H_{\kappa}^{-1}} \left\| u \right\|_{\mathrm{LS}_{\kappa}}. \end{split}$$

For a_2 , we use the embedding $H^1_{\kappa} \hookrightarrow L^{\infty}$, (4.18), and (4.20) to bound

$$\int_{-T}^{T} \left| \int (g - \frac{1}{2\kappa}) a_2 u^2 \, dx \right| dt \le \left\| g - \frac{1}{2\kappa} \right\|_{L^{\infty}_{t,x}} \left\| b u^2 \right\|_{L^{1}_{t,x}}$$
27

$$\lesssim \kappa^{-\frac{3}{2}} \|u\|_{L_{t}^{\infty} H_{\kappa}^{-1}} \int \|\psi_{z} a_{2}\|_{L_{t,x}^{\infty}} \|\psi_{z} u\|_{L_{t,x}^{2}}^{2} dz$$

$$\lesssim \varepsilon \kappa^{-\frac{3}{2}} \|u\|_{L_{t}^{\infty} H_{\kappa}^{-1}} \left(\|u\|_{\mathrm{LS}_{\kappa}}^{2} + \|u\|_{L_{t}^{\infty} H_{\kappa}^{-1}}^{2} \right)$$

$$\lesssim \varepsilon \kappa^{-1} \left(\|u\|_{\mathrm{LS}_{\kappa}}^{2} + \|u\|_{L_{t}^{\infty} H_{\kappa}^{-1}}^{2} \right),$$

using (3.9) at the last step.

Lastly, we turn to the contribution of a_1 . We integrate by parts once in space, and then we use (4.18), (3.8), and (4.1) to estimate:

$$\int_{-T}^{T} \left| \int (g - \frac{1}{2\kappa}) (a_1 u')' \, dx \right| dt \leq \int \|\psi_z g'\|_{L^2_t H^1_\kappa} \|\psi_z u'\|_{L^2_t H^{-1}_\kappa} \|\psi_z a_1\|_{L^\infty_t X} \, dz$$
$$\lesssim \varepsilon \kappa^{-1} \|u\|_{\mathrm{LS}_\kappa}^2 \, .$$

Altogether, returning to (5.2), we obtain

$$\|u\|_{C_{t}H_{\kappa}^{-1}}^{2} \lesssim \|u(0)\|_{H_{\kappa}^{-1}}^{2} + (\varepsilon + T) \left(\|u\|_{L_{t}^{\infty}H_{\kappa}^{-1}}^{2} + \|u\|_{\mathrm{LS}_{\kappa}}^{2} \right).$$

At this point we are prepared to complete the proofs of Theorems 1.1 and 1.5, by combining Propositions 4.1 and 5.1 and a continuity argument. We restate the results together in a stronger form:

Theorem 5.2. There exist $\epsilon_0, c_0 > 0$ so that given coefficients a_1, \ldots, a_4 in (gKdV) satisfying (4.1)–(4.4) with $\varepsilon = \varepsilon_0$ and $R \ge 1$ the solutions u(t) to (gKdV) and initial data u_0 satisfying

(5.3)
$$||u_0||_{H^{-1}} \le R$$

exist up to time

(5.4)
$$T_0 = c_0 R^{-4}$$

and satisfy in $[0, T_0]$

(5.5)
$$\|u\|_{L^{\infty}_{t}H^{-1}_{\kappa}} + \|u\|_{\mathrm{LS}_{\kappa}} \lesssim R, \qquad \kappa = R^{2}.$$

Proof. For any $\kappa \gg R^2$, the condition (3.9) holds at t = 0 and thus on some time interval [0, T]. Let $T \leq \kappa^{-2}$ be any time for which the solution exists in [0, T] and satisfies (3.9) uniformly in [0, T]. Then both Propositions 4.1 and 5.1 apply in [0, T].

Consider the quantity

$$B_T := \|u\|_{C_t H_{\kappa}^{-1}([-T,T] \times \mathbb{R})}^2 + \frac{1}{4} \|u\|_{\mathrm{LS}_{\kappa}([-T,T] \times \mathbb{R})}^2$$

Adding (4.5) and (5.1), we see that there exists a universal constant $C \ge 1$ so that

(5.6)
$$B_T \le CR^2 + C(\varepsilon_0 + (T\kappa^2)^{\frac{1}{4}} + \kappa^{-2})B_T.$$

Assuming

(5.7) $\epsilon_0 \ll 1, \quad T \ll \kappa^{-2}, \quad \kappa \gg 1,$

we conclude that

$$(5.8) B_T \le 2CR^2$$

Now we fix our parameters in order, beginning with $\epsilon_0 \ll 1$ so that the first relation (5.7) holds, then

$$\kappa = 1 + 4CR^2,$$

so that the third relation in (5.7) holds, and finally

$$T_0 = c\kappa^{-2}, \qquad c \ll 1$$

so that the second relation of (5.7) holds.

Finally, we run a standard continuity argument. We let $T \in (0, T_0]$ be maximal so that (3.9) holds in [0, T]. Then by the above reasoning shows that (5.8) holds in [0, T]. In particular, given the choice of κ , this implies that (3.9) holds strictly at time T. But this contradicts the maximality of T unless $T = T_0$. We conclude that $T = T_0$, and thus (5.8) holds in $[0, T_0]$. Hence the conclusion of the theorem follows.

Next, we will show that the assumptions (1.2)-(1.3) on the coefficients a_1, \ldots, a_4 in Theorems 1.1 and 1.5 provide an example of when our hypotheses (4.1)-(4.4) are satisfied.

Lemma 5.3. Given $\varepsilon, T \in (0, 1]$, there exists a constant $\delta \in (0, 1]$ so that, if the coefficients a_1, \ldots, a_4 satisfy

(5.9)
$$|a_j(t,x)| + |\partial_x a_j(t,x)| \le \delta (1+x^2)^{-1} \text{ for } j = 1,2,3,$$

(5.10)
$$|a_4(t,x)| + |\partial_x a_4(t,x)| \lesssim (1+x^2)^{-1}$$

uniformly for $|t| \leq T$ and $x \in \mathbb{R}$, then (4.1)–(4.4) are satisfied for ε .

Proof. We partition ψ as follows:

$$\psi^2(x) = \operatorname{sech}^2 x \le \sum_{n \ge 0} \operatorname{sech}^2(n) \mathbf{1}_{\{n \le |x| < n+1\}}(x).$$

This yields

$$\int |\psi_z(x)a_j(x)|^2 dx \le \sum_{n\ge 0} \operatorname{sech}^2(n) \int_{n\le |x-z|< n+1} |a_1(x)|^2 dx \lesssim \delta^2 \langle z \rangle^{-4},$$

and so

$$\int \|\psi_z a_j\|_{L^2_x} dz \lesssim \delta \int \langle z \rangle^{-2} dz \lesssim \delta.$$

Using the same estimates for $\partial_x a_j$, we obtain

$$\int \left\|\psi_z a_j\right\|_{H^1_x} dz \lesssim \delta \quad \text{for } j = 1, 2, 3, 4.$$

For j = 4, this shows that (4.4) is satisfied. For j = 1, 2, 3, this demonstrates that we may choose $\delta \leq 1$ sufficiently small so that (4.1)–(4.3) hold.

Lastly, we show that our result applies to the model (1.1) for the propagation of waves over a variable channel bottom.

Proof of Corollary 1.4. If the function $c : \mathbb{R} \to \mathbb{R}$ that describes the channel bottom is smooth and satisfies $\|c\|_{L^{\infty}} < 1$, then the function $y : \mathbb{R} \to \mathbb{R}$ given by

$$y(x) = \int_0^x \frac{1}{b^{\frac{5}{3}}(\xi)} d\xi = \int_0^x \frac{1}{[1 - c(\xi)]^{\frac{5}{6}}} d\xi$$

is well-defined and satisfies

(5.11)
$$|y'(x)| \approx 1$$
 uniformly for $x \in \mathbb{R}$.

Consequently,

(5.12)
$$\int |f(y)|^2 dy \approx \int |(f \circ y)(x)|^2 dx$$
 and $\int |f'(y)|^2 dy \approx \int |(f \circ y)'(x)|^2 dx$,

and so

(5.13)
$$\|f \circ y\|_{H^1_{\kappa}} \approx \|f\|_{H^1_{\kappa}} \quad \text{uniformly for } \kappa \ge 1.$$

Moreover, notice that $y^{-1} : \mathbb{R} \to \mathbb{R}$ exists by the inverse function theorem, and also satisfies (5.11)–(5.13). A straightforward duality argument then shows

(5.14)
$$\|g \circ y\|_{H^{-1}_{\kappa}} \approx \|g\|_{H^{-1}_{\kappa}} \quad \text{uniformly for } \kappa \ge 1.$$

Making the change of variables

(5.15)
$$u(t,x) = b^{\frac{5}{3}}(x)v(t,y(x) - 4t),$$

we find that if u(t, x) solves (1.1) then v(t, y) solves (gKdV) with coefficients

$$\begin{aligned} a_1(t,y) &= 0, \\ a_2(t,y) &= 10b^{\frac{2}{3}}b' \circ y^{-1}, \\ a_3(t,y) &= \left[\frac{5}{9}b^{\frac{4}{3}}(b')^2 - \frac{10}{3}b^{\frac{7}{3}}b'' + 4(1-b^{-\frac{2}{3}})\right] \circ y^{-1}, \\ a_4(t,y) &= \left[\frac{10}{3}b^2(b')^3 - \frac{10}{3}b^3b'b'' - \frac{5}{3}b^4b''' - \frac{38}{3}b'\right] \circ y^{-1}. \end{aligned}$$

Clearly, we may choose $\eta \in (0, 1]$ sufficiently small so that

$$|\partial_x^j c(x)| \le \eta (1+x^2)^{-1}$$
 for $j = 0, 1, \dots, 4$

imply that the coefficients a_1, \ldots, a_4 above satisfy (5.9)–(5.10). Note that the 4 in (5.15) contributes the constant term 4 in a_3 , which is needed to ensure that this coefficient vanishes as $y \to \pm \infty$.

This demonstrates that v satisfies the a-priori estimate (5.5). This in turn implies that u satisfies (5.5) as well, by (5.14).

References

- T. Akhunov, D. M. Ambrose, and J. D. Wright, Well-posedness of fully nonlinear KdV-type evolution equations, Nonlinearity 32 (2019), no. 8, 2914–2954. MR3987807
- J. Bona and R. Scott, Solutions of the Korteweg-de Vries equation in fractional order Sobolev spaces, Duke Math. J. 43 (1976), no. 1, 87–99. MR393887
- [3] J. L. Bona, S. V. Rajopadhye, and M. E. Schonbek, Models for propagation of bores. I. Two-dimensional theory, Differential Integral Equations 7 (1994), no. 3-4, 699–734. MR1270099
- [4] J. L. Bona and R. Smith, The initial-value problem for the Korteweg-de Vries equation, Philos. Trans. Roy. Soc. London Ser. A 278 (1975), no. 1287, 555–601. MR385355
- [5] J. Bourgain, Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. II. The KdV-equation, Geom. Funct. Anal. 3 (1993), no. 3, 209–262. MR1215780
- [6] B. Bringmann, R. Killip, and M. Visan, Global well-posedness for the fifth-order KdV equation in H⁻¹(ℝ), Ann. PDE 7 (2021), no. 2, Paper No. 21, 46. MR4304314
- T. Buckmaster and H. Koch, The Korteweg-de Vries equation at H⁻¹ regularity, Ann. Inst. H. Poincaré C Anal. Non Linéaire 32 (2015), no. 5, 1071–1098. MR3400442
- [8] V. Buslaev and V. Fomin, An inverse scattering problem for the one-dimensional Schrödinger equation on the entire axis, Vestnik Leningrad. Univ. 17 (1962), no. 1, 56–64. MR0139371

- M. Christ, J. Colliander, and T. Tao, Asymptotics, frequency modulation, and low regularity ill-posedness for canonical defocusing equations, Amer. J. Math. 125 (2003), no. 6, 1235–1293. MR2018661
- [10] A. Cohen, Solutions of the Korteweg-de Vries equation with steplike initial profile, Comm. Partial Differential Equations 9 (1984), no. 8, 751–806. MR748367
- [11] A. Cohen and T. Kappeler, Solutions to the Korteweg-de Vries equation with initial profile in $L_1^1(\mathbf{R}) \cap L_N^1(\mathbf{R}^+)$, SIAM J. Math. Anal. 18 (1987), no. 4, 991–1025. MR892486
- [12] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao, Sharp global well-posedness for KdV and modified KdV on ℝ and T, J. Amer. Math. Soc. 16 (2003), no. 3, 705–749. MR1969209
- [13] W. Craig, T. Kappeler, and W. Strauss, Gain of regularity for equations of KdV type, Ann. Inst. H. Poincaré C Anal. Non Linéaire 9 (1992), no. 2, 147–186. MR1160847
- [14] D. G. Crighton, Applications of KdV, 1995, pp. 39–67. KdV '95 (Amsterdam, 1995). MR1329553
- [15] M. W Dingemans, Water wave propagation over uneven bottoms, World Scientific Publishing Company, 1997.
- [16] I. Egorova, K. Grunert, and G. Teschl, On the Cauchy problem for the Korteweg-de Vries equation with steplike finite-gap initial data. I. Schwartz-type perturbations, Nonlinearity 22 (2009), no. 6, 1431–1457. MR2507328
- [17] I. Egorova and G. Teschl, On the Cauchy problem for the Korteweg-de Vries equation with steplike finitegap initial data II. Perturbations with finite moments, J. Anal. Math. 115 (2011), 71–101. MR2855034
- [18] V. D. Ermakova, The asymptotics of the solution of the Cauchy problem for the Korteweg-de Vries equation with nondecreasing initial data of special type, Dokl. Akad. Nauk Ukrain. SSR Ser. A 7 (1982), 3-6, 88. MR671494
- [19] _____, The inverse scattering problem on the whole axis for the Schrödinger equation with nondecreasing potential of special form, Vestnik Khar'kov, Univ 230 (1982), 50–60.
- [20] N. E. Firsova, Solution of the Cauchy problem for the Korteweg-de Vries equation with initial data that are the sum of a periodic and a rapidly decreasing function, Mat. Sb. (N.S.) 135(177) (1988), no. 2, 261–268, 272. MR937811
- [21] C. Gallo, Korteweg-de Vries and Benjamin-Ono equations on Zhidkov spaces, Adv. Differential Equations 10 (2005), no. 3, 277–308. MR2123133
- [22] R. Grimshaw and S. R. Pudjaprasetya, Hamiltonian formulation for solitary waves propagating on a variable background, 1999, pp. 89–98. Applications of solitons and symmetries. MR1697970
- [23] S. Grudsky, C. Remling, and A. Rybkin, The inverse scattering transform for the KdV equation with step-like singular Miura initial profiles, J. Math. Phys. 56 (2015), no. 9, 091505, 14. MR3395045
- [24] S. Grudsky and A. Rybkin, On positive type initial profiles for the KDV equation, Proc. Amer. Math. Soc. 142 (2014), no. 6, 2079–2086. MR3182026
- [25] Z. Guo, Global well-posedness of Korteweg-de Vries equation in $H^{-3/4}(\mathbb{R})$, J. Math. Pures Appl. (9) **91** (2009), no. 6, 583–597. MR2531556
- [26] B. Harrop-Griffiths, Large data local well-posedness for a class of KdV-type equations, Trans. Amer. Math. Soc. 367 (2015), no. 2, 755–773. MR3280026
- [27] M. Ifrim, H. Koch, and D. Tataru, Dispersive decay of small data solutions for the KdV equation, Ann. Sci. Éc. Norm. Supér. (4) 56 (2023), no. 6, 1709–1746. MR4706572
- [28] R. Iorio, F. Linares, and M. Scialom, KdV and BO equations with bore-like data, Differential Integral Equations 11 (1998), no. 6, 895–915. MR1659252
- [29] S. Israwi, Variable depth KdV equations and generalizations to more nonlinear regimes, M2AN Math. Model. Numer. Anal. 44 (2010), no. 2, 347–370. MR2655953
- [30] S. Israwi and R. Talhouk, Local well-posedness of a nonlinear KdV-type equation, C. R. Math. Acad. Sci. Paris 351 (2013), no. 23-24, 895–899. MR3133600
- [31] R. S. Johnson, On the development of a solitary wave moving over an uneven bottom, Proc. Cambridge Philos. Soc. 73 (1973), 183–203. MR312817
- [32] T. Kappeler, Solutions of the Korteweg-de Vries equation with steplike initial data, J. Differential Equations 63 (1986), no. 3, 306–331. MR848272
- [33] T. Kappeler and P. Topalov, Global wellposedness of KdV in H⁻¹(T, ℝ), Duke Math. J. 135 (2006), no. 2, 327–360. MR2267286

- [34] T. Kato, Quasi-linear equations of evolution, with applications to partial differential equations, Spectral theory and differential equations (Proc. Sympos., Dundee, 1974; dedicated to Konrad Jörgens), 1975, pp. 25–70. Lecture Notes in Math., Vol. 448. MR0407477
- [35] _____, On the cauchy problem for the (generalized) Korteweg-de Vries equation, Studies in applied mathematics 8 (1983), 93-128.
- [36] C. E. Kenig, G. Ponce, and L. Vega, Well-posedness of the initial value problem for the Korteweg-de Vries equation, J. Amer. Math. Soc. 4 (1991), no. 2, 323–347. MR1086966
- [37] _____, A bilinear estimate with applications to the KdV equation, J. Amer. Math. Soc. 9 (1996), no. 2, 573–603. MR1329387
- [38] R. Killip and M. Vişan, KdV is well-posed in H^{-1} , Ann. of Math. (2) **190** (2019), no. 1, 249–305. MR3990604
- [39] R. Killip, M. Vişan, and X. Zhang, Low regularity conservation laws for integrable PDE, Geom. Funct. Anal. 28 (2018), no. 4, 1062–1090. MR3820439
- [40] N. Kishimoto, Well-posedness of the Cauchy problem for the Korteweg-de Vries equation at the critical regularity, Differential Integral Equations 22 (2009), no. 5-6, 447–464. MR2501679
- [41] H. Koch and D. Tataru, Conserved energies for the cubic nonlinear Schrödinger equation in one dimension, Duke Math. J. 167 (2018), no. 17, 3207–3313. MR3874652
- [42] D. J. Korteweg and G. de Vries, On the change of form of long waves advancing in a rectangular canal, and on a new type of long stationary waves, Philos. Mag. (5) 39 (1895), no. 240, 422–443. MR3363408
- [43] E. A. Kuznetsov and A. V. Mikhaĭlov, Stability of stationary waves in nonlinear weakly dispersive media,
 Ž. Èksper. Teoret. Fiz. 67 (1974), no. 5, 1717–1727. MR0387847
- [44] D. Lannes, The water waves problem, Mathematical Surveys and Monographs, vol. 188, American Mathematical Society, Providence, RI, 2013. Mathematical analysis and asymptotics. MR3060183
- [45] T. Laurens, KdV on an incoming tide, Nonlinearity 35 (2022), no. 1, 343–387. MR4356987
- [46] _____, Global well-posedness for $H^{-1}(\mathbb{R})$ perturbations of KdV with exotic spatial asymptotics, Comm. Math. Phys. **397** (2023), no. 3, 1387–1439. MR4541923
- [47] B. Liu, A priori bounds for KdV equation below $H^{-\frac{3}{4}}$, J. Funct. Anal. **268** (2015), no. 3, 501–554. MR3292346
- [48] A. Menikoff, The existence of unbounded solutions of the Korteweg-de Vries equation, Comm. Pure Appl. Math. 25 (1972), 407–432. MR303129
- [49] J. W. Miles, On the Korteweg-de Vries equation for a gradually varying channel, J. Fluid Mech. 91 (1979), no. 1, 181–190. MR532306
- [50] L. Molinet, A note on ill posedness for the KdV equation, Differential Integral Equations 24 (2011), no. 7-8, 759–765. MR2830706
- [51] _____, Sharp ill-posedness results for the KdV and mKdV equations on the torus, Adv. Math. 230 (2012), no. 4-6, 1895–1930. MR2927357
- [52] L. Molinet, R. Talhouk, and I. Zaiter, On well-posedness for some Korteweg-de Vries type equations with variable coefficients, J. Evol. Equ. 23 (2023), no. 3, Paper No. 52, 37. MR4612353
- [53] J. M. Palacios, Local well-posedness for the gKdV equation on the background of a bounded function, Rev. Mat. Iberoam. 39 (2023), no. 1, 341–396. MR4571609
- [54] S. R. Pudjaprasetya and E. van Groesen, Unidirectional waves over slowly varying bottom. II. Quasihomogeneous approximation of distorting waves, Wave Motion 23 (1996), no. 1, 23–38. MR1388509
- [55] S. R. Pudjaprasetya, E. Van Groesen, and E. Soewono, The splitting of solitary waves running over a shallower water, Wave Motion 29 (1999), no. 4, 375–389. MR1686790
- [56] A. Rybkin, Regularized perturbation determinants and KdV conservation laws for irregular initial profiles, Topics in operator theory. Volume 2. Systems and mathematical physics, 2010, pp. 427–444. MR2683250
- [57] _____, The Hirota τ -function and well-posedness of the KdV equation with an arbitrary step-like initial profile decaying on the right half line, Nonlinearity **24** (2011), no. 10, 2953–2990. MR2842104
- [58] _____, KdV equation beyond standard assumptions on initial data, Phys. D 365 (2018), 1–11. MR3739918
- [59] J. C. Saut and R. Temam, Remarks on the Korteweg-de Vries equation, Israel J. Math. 24 (1976), no. 1, 78–87. MR454425

- [60] R. Temam, Sur un problème non linéaire, J. Math. Pures Appl. (9) 48 (1969), 159–172. MR261183
- [61] B. Tian and Y.-T. Gao, Variable-coefficient balancing-act method and variable-coefficient KdV equation from fluid dynamics and plasma physics, Eur. Phys. J. B 22 (2001), 351–360.
- [62] M. Tsutsumi and T. Mukasa, Parabolic regularizations for the generalized Korteweg-de Vries equation, Funkcial. Ekvac. 14 (1971), 89–110. MR312097
- [63] E. van Groesen and S. R. Pudjaprasetya, Uni-directional waves over slowly varying bottom. I. Derivation of a KdV-type of equation, Wave Motion 18 (1993), no. 4, 345–370. MR1258081
- [64] S. B. Yoon and P. L.-F. Liu, A note on Hamiltonian for long water waves in varying depth, Wave Motion 20 (1994), no. 4, 359–370. MR1307821
- [65] P. Zhidkov, Korteweg-de Vries and nonlinear Schrödinger equations: qualitative theory, Lecture Notes in Mathematics, vol. 1756, Springer-Verlag, Berlin, 2001. MR1831831

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, MADISON, WI 53706, USA *Email address*: ifrim@math.wisc.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, MADISON, WI 53706, USA *Email address*: laurens@math.wisc.edu