REFINED REGULARITY FOR NONLOCAL ELLIPTIC EQUATIONS AND APPLICATIONS

WENXIONG CHEN, CONGMING LI, LEYUN WU, AND ZHOUPING XIN

ABSTRACT. In this paper, we establish refined regularity estimates for nonnegative solutions to the fractional Poisson equation

$$(-\Delta)^{s}u(x) = f(x), \ x \in B_{1}(0).$$

Specifically, we have derived Hölder, Schauder, and Ln-Lipschitz regularity estimates for any nonnegative solution u, provided that only the local L^{∞} norm of u is bounded. These estimates stand in sharp contrast to the existing results where the global L^{∞} norm of u is required. Our findings indicate that the local values of the solution u and f are sufficient to control the local values of higher order derivatives of u. Notably, this makes it possible to establish a priori estimates in unbounded domains by using blowing up and re-scaling argument.

As applications, we derive singularity and decay estimates for solutions to some super-linear nonlocal problems in unbounded domains, and in particular, we obtain a priori estimates for a family of fractional Lane-Emden type equations in \mathbb{R}^n . This is achieved by adopting a different method using auxiliary functions, which is applicable to both local and nonlocal problems.

1. INTRODUCTION AND MAIN RESULTS

In recent years, there has been a surge of interest in employing the fractional Laplacian to model a diverse array of physical phenomena characterized by long-range interactions. Notably, it finds applications in modeling anomalous diffusion, quasi-geostrophic flows, turbulence, water waves, molecular dynamics, and other phenomena (refer to [2, 9, 15, 31] and the associated references). Specifically, the fractional Laplacian serves as the infinitesimal generator of a stable Lévy diffusion process (refer to [1]), playing a crucial role in describing anomalous diffusions observed in various contexts such as plasmas, flame propagation, chemical reactions in liquids, and population dynamics.

The fractional Laplacian can be defined as

$$(-\Delta)^{s}u(x) = C_{n,s}PV \int_{\mathbb{R}^{n}} \frac{u(x) - u(y)}{|x - y|^{n+2s}} dy$$

= $C_{n,s}PV \lim_{\varepsilon \to 0} \int_{\mathbb{R}^{n} \setminus B_{\varepsilon}(x)} \frac{u(x) - u(y)}{|x - y|^{n+2s}} dy,$ (1.1)

where s is any real number between 0 and 1, PV stands for the Cauchy principal value, $B_{\varepsilon}(x)$ is the ball of radius ε centered at x, and $C_{n,s}$ is a dimensional constant that depends on n and s, precisely given by

$$C_{n,s} = \left(\int_{\mathbb{R}^n} \frac{1 - \cos(\zeta_1)}{|\zeta|^{n+2s}} d\zeta\right)^{-1}$$

To ensure the integrability of the right hand side of (1.1), we require that $u \in C^{1,1}_{loc}(\mathbb{R}^n) \cap \mathcal{L}_{2s}$, where

$$\mathcal{L}_{2s} = \left\{ u \in L^1_{loc} \left| \int_{\mathbb{R}^n} \frac{|u(x)|}{1+|x|^{n+2s}} dx < \infty \right. \right\}$$

²⁰¹⁰ Mathematics Subject Classification. 35R11, 35B45, 35B65.

Key words and phrases. Schauder estimate, a priori estimates.

endowed naturally with the norm

$$||u||_{\mathcal{L}_{2s}} := \int_{\mathbb{R}^n} \frac{|u(x)|}{1+|x|^{n+2s}} dx.$$

The nonlocal nature of the fractional Laplacian makes it difficult to derive the regularity estimates by using the well-known traditional approaches for local differential operators. As a remedy, Caffarelli and Silvestre [5] introduced the extension method which turns nonlocal equations into local ones in higher dimensions, so that a series of existing approaches on local equations can be applied, and thus it has become a powerful tool in investigating equations involving the fractional Laplacian. However, one sometimes needs to assume extra conditions such as $s \ge \frac{1}{2}$ in proving symmetry of the fractional Lane-Emden equation ([3]) which may not be necessary if one approaches the problem directly.

To address these challenges and explore further aspects of nonlocal equations involving the fractional Laplacian and other nonlocal elliptic operators, various powerful direct methods have been developed. These include the direct method of moving planes and of moving spheres ([10-12, 16, 30, 32], sliding methods ([28, 29]), the method of scaling spheres ([17]), and others ([30]). These methods have been widely applied to study nonlocal problems and a series of fruitful results have been obtained.

In this paper, by direct elaborate analysis instead of extension, we establish Hölder regularity, Schauder regularity, Ln-Lipschitz regularity and C^k regularity for nonnegative classical solutions to the fractional Poisson equation

$$(-\Delta)^s u(x) = f(x), \ x \in B_1(0).$$
 (1.2)

As important applications, we derive a priori estimates concerning possible singularities and decay for nonnegative solutions to nonlocal problems involving more general nonlinearities in an unbounded domain Ω :

$$-\Delta)^{s}u(x) = b(x)|\nabla u|^{q}(x) + f(x, u(x)), \ x \in \Omega,$$
(1.3)

where $0 < q < \frac{2sp}{2s+p-1}$. It is noteworthy that the following interior Hölder regularity and Schauder regularity for solutions to (1.2) with $f \in L^{\infty}(B_1(0))$ and $f \in C^{\alpha}(\overline{B}_1(0))$ for all $0 < \varepsilon < 2s$, have been proved

$$\|u\|_{C^{[2s-\varepsilon]},\{2s-\varepsilon\}}(B_{1/2}(0)) \le C(\|f\|_{L^{\infty}(B_{1}(0))} + \|u\|_{L^{\infty}(\mathbb{R}^{n})}),$$
(1.4)

and

$$\|u\|_{C^{[2s+\alpha]},\{2s+\alpha\}}(B_{1/2}(0)) \le C\left(\|f\|_{C^{\alpha}(\bar{B}_{1}(0))} + \|u\|_{L^{\infty}(\mathbb{R}^{n})}\right),\tag{1.5}$$

where $2s + \alpha \notin \mathbb{N}$, $[2s + \alpha]$ and $\{2s + \alpha\}$ are the integer part and the fraction part of $2s + \alpha$ respectively, see [11, 18, 26, 27] and also [6–8] for fully nonlinear nonlocal equations. Additionally, later results indicated that the right hand side of (1.5) can be relaxed to

$$C\left(\|f\|_{C^{\alpha}(\bar{B}_{1}(0))}+\|u\|_{\mathcal{L}_{2s}}\right)$$

For more details, please refer to [11, Theorem 12.2.4] and [22, Theorem 1.3].

However, probably due to the non-local nature of the fractional Laplacian, it appears that these results are considerably weaker than those established for the classical Poisson equation

$$-\Delta u(x) = f(x), \ x \in B_1(0),$$
 (1.6)

for which one usually needs only $||u||_{L^{\infty}(B_1(0))}$ and $||f||_{C^{\alpha}(\bar{B}_1(0))}$ to bound $||u||_{C^{2,\alpha}(B_{1/2}(0))}$.

More importantly, the known a priori estimates, such as (1.4) and (1.5) with its improvements are insufficient for applying the powerful blowing-up and re-scaling arguments to analyze the qualitative properties of solutions, in particular, to establish a priori estimates for fractional equations on unbounded domains due to the requirement on the bonded-ness of global norms of u.

As an illustration, let Ω be an unbounded domain in \mathbb{R}^n , and let us compare a simple problem of local nature

$$\begin{cases} -\Delta u(x) = u^p(x), & x \in \Omega\\ u(x) = 0, & x \in \partial \Omega \end{cases}$$
(1.7)

with it nonlocal counter part

$$\begin{cases} (-\Delta)^s u(x) = u^p(x), & x \in \Omega\\ u(x) = 0, & x \in \mathbb{R}^n \setminus \Omega, \end{cases}$$
(1.8)

where 0 < s < 1, and p is a subcritical exponent.

To obtain an a priori estimate of positive solutions, one would usually apply a typical blowing up and re-scaling argument briefly as follows.

If the solutions are not uniformly bounded, then there exist a sequence of solutions $\{u_k\}$ and a sequence of points $\{x^k\} \in \Omega$, such that $u_k(x^k) \to \infty$. Notice that since Ω is unbounded, this $u_k(x^k)$ may not be the maximum of u_k in Ω . Upon re-scaling, the new sequence of functions v_k are bounded and satisfy the same equation on certain domains B_k .

For the equation in (1.7) of local nature, the bounded-ness of $\{v_k\}$ on B_k is sufficient to guarantee the bound on its higher norms, say $C^{2,\alpha}$ norm. This leads to the convergence of $\{v_k\}$ to a function v, which is a bounded solution of the same equation as (1.7) either in the whole space or in a half space. By the known results on the nonexistence of solutions in these two spaces, one derives a contradiction and hence obtains the a priori estimate.

Nevertheless, when dealing with nonlocal problem (1.8), the currently available regularity estimates, such as (1.4) and (1.5), indicate that the bounded-ness of $\{v_k\}$ on B_k are not sufficient to guarantee the bound on its higher norms, they also require $\{v_k\}$ be bounded across the entire space. Such a condition cannot be met because there is no information on the behavior of v_k beyond B_k during the re-scaling process. This posses a substantial difficulty on employing blowing and re-scaling arguments for nonlocal equations on unbounded domains.

The above arguments lead to the following natural question:

Can one use only $||u||_{L^{\infty}(B_1(0))}$ and $||f||_{L^{\infty}(B_1(0))} (||f||_{C^{\alpha}(\bar{B}_1(0))})$ to bound $||u||_{C^{[2s-\varepsilon]}, \{2s-\varepsilon\}}(B_{1/2}(0))$ ($||u||_{C^{[2s+\alpha]}, \{2s-\alpha\}}(B_{1/2}(0))$) in (1.4) ((1.5)) under certain conditions?

This is one of the main problems we study in this paper (see Theorem 1.1 and Theorem 1.3 below). By careful analysis, we answer this question affirmatively for nonnegative solutions and conclude that

$$\|u\|_{C^{[2s-\varepsilon]},\{2s-\varepsilon\}}(B_{1/2}(0)) \le C(\|f\|_{L^{\infty}(B_{1}(0))} + \|u\|_{L^{\infty}(B_{1}(0))})$$

for all $0 < \varepsilon < 2s$, and

$$\|u\|_{C^{[2s+\alpha]},\{2s+\alpha\}}(B_{1/2}(0)) \le C\left(\|f\|_{C^{\alpha}(\bar{B}_{1}(0))} + \|u\|_{L^{\infty}(B_{1}(0))}\right)$$
(1.9)

for $2s + \alpha \notin \mathbb{N}$.

We now proceed to outline the main ingredients of our analysis and present our main results. To derive the Schauder estimates for non-negative solutions of the fractional Poisson equation (1.2), we split the solution u into two parts: the s-harmonic function in $B_1(0)$

$$h(x) := u(x) - C_{n,s} \int_{B_1(0)} \frac{f(y)}{|x-y|^{n-2s}} dy$$

and the remaining term

$$w(x) := C_{n,s} \int_{B_1(0)} \frac{f(y)}{|x-y|^{n-2s}} dy.$$

For w(x), one expects that $||w||_{C^{[2s+\alpha]}, \{2s+\alpha\}}(B_{1/2}(0))}$ can be controlled by $||f||_{C^{\alpha}(B_1(0))}$. While for the s-harmonic function h(x), it has been proved in [11] (see also [22]) that $h(x) \in C^{\infty}(B_1(0))$ and can be represented by

$$h(x) = \int_{|y|>1} P(x,y)h(y)dy, \ \forall \ |x| < 1,$$

where

$$P(x,y) = \begin{cases} \frac{\Gamma(n/2)\sin(\pi s)}{\pi^{\frac{n}{2}+1}} \left(\frac{1-|x|^2}{|y|^2-1}\right)^s \frac{1}{|x-y|^n}, & |y| > 1, \\ 0, & |y| \le 1 \end{cases}$$

is the so-called Poisson kernel.

By the definition of P(x, y) and some elaborate calculations, we find that the higher order derivatives of P(x, y) can be controlled by itself. Then, $\|D^k h\|_{L^{\infty}(B_{1/2}(0))}$ can be bounded by $\|f\|_{L^{\infty}(B_1(0))}$ and $\|u\|_{L^{\infty}(B_1(0))}$ due to the fact that h = u - w. This leads to better Schauder estimates for nonnegative solutions, and our observations here may provide some useful insights on the research of local regularity for nonlocal operators.

Thus we have shown that for $2s + \alpha \notin \mathbb{N}$,

$$\|u\|_{C^{[2s+\alpha]},\{2s+\alpha\}}(B_{1/2}(0)) \le C\left(\|f\|_{C^{\alpha}(\bar{B}_{1}(0))} + \|u\|_{L^{\infty}(B_{1}(0))}\right)$$

While for $2s + \alpha \in \mathbb{N}$, two versions of estimates will be obtained:

i) for $f \in C^{\alpha}$, $\|u\|_{C^{2s+\alpha-1,\ln L}(B_{1/2}(0))}$ can be controlled by $\|u\|_{L^{\infty}(B_1(0))}$ and $\|f\|_{C^{\alpha}(\bar{B}_1(0))}$, which is the optimal Ln–Lipschitz regularity.

ii) for $f \in C^{\alpha,Dini}$, $\|u\|_{C^{2s+\alpha}(B_{1/2}(0))}$ can be estimated in terms of $\|u\|_{L^{\infty}(B_{1}(0))}$ and $\|f\|_{C^{\alpha,Dini}(\bar{B}_{1}(0))}$.

Roughly speaking, the above regularity results imply that the local values of the solution u and non-homogeneous term f are sufficient to control the higher order derivatives of u. These newly derived estimates are both novel and desirable.

Based on the above refined regularity estimates, we are able to establish singularity and decay estimates for solutions to equation (1.3) (see Theorem 1.8). To this end, one usually used the well-known "doubling lemma" ([25, Lemma 5.1]) combining with the re-scaling and blowing-up to derive such kind of estimates. For example, Poláčik, Quittner and Souplet in [25] established decay and singularity for the classical integer order Lane-Emden equation

$$-\Delta u(x) = u^p(x)$$

by using this method.

In order to circumvent the use of "the doubling lemma" and make the proof more direct, we adopted an alternative approach involving the construction of auxiliary functions. The idea is that if an estimate (in terms of the distance to $\partial\Omega$) such as (1.16) fails, then by the contradiction argument, there exist sequences Ω_k , u_k , $x_k \in \Omega_k$ such that

$$u_k(x_k) \left(1 + dist(x_k, \partial \Omega_k)^{\frac{-2s}{p-1}}\right)^{-1} > k \to \infty \quad \text{as} \quad k \to \infty,$$

see (4.2). Then for each k, we construct an auxiliary function $S_k(x)$ and find its maximum point denoted as a_k in a neighborhood of x_k . After appropriate rescaling (related to a_k), we can guarantee that the rescaled sequence $\{v_k\}$ is bounded in a neighborhood of a_k . In order to pass to the limit to obtain a bounded solution of a limiting problem in the whole space \mathbb{R}^n to derive a contradiction with a Liouville theorem for the fractional Lane-Emden equation in \mathbb{R}^n , we use the refined regularity results established in Theorem 1.1 and Theorem 1.3 below. Compared with previous results for fractional equations (see [23]), we do not use "the doubling lemma" and do not need to assume $u \in \dot{H}^s(\mathbb{R}^n)$.

This new method enables us not only to obtain singularity and decay estimates for fractional order elliptic problems without using the doubling lemma, but also to deal effectively with both integer order elliptic and parabolic problems, as well as for fractional parabolic problems. As other by-products, we derive the boundedness of nonnegative solutions for a family of fractional Lane-Emden type equations in the whole space \mathbb{R}^n and thus simplifies the problem when we study such kind of equations. These results can be extended to a family of fractional Lane-Emden type systems. It may shed some light on the fractional Lane-Emden conjecture which states that if

$$\frac{1}{p+1} + \frac{1}{q+1} > \frac{n-2s}{n}$$

then there are no nontrivial positive classical solutions of

$$\begin{cases} (-\Delta)^s u(x) = v^p(x), & x \in \mathbb{R}^n, \\ (-\Delta)^s v(x) = u^q(x), & x \in \mathbb{R}^n. \end{cases}$$

We now present the main results in this paper in details.

We start with some standard notations. $C^{k,\ln L}(\bar{\Omega})$ is the so called "Ln-Lipshitz" spaces with

$$\|f\|_{C^{k,\ln L}(\bar{\Omega})} = \|f\|_{C^{k}(\bar{\Omega})} + \sum_{|\beta|=k} \sup_{x,y\in\Omega, x\neq y} \frac{|D^{\beta}f(x) - D^{\beta}f(y)|}{|x-y||\ln\min(|x-y|,1/2)|} < \infty.$$

A function f is said to be Dini continuous (in Ω) if ω_f satisfies the Dini condition

$$I_{\omega_f} := \int_0^{\operatorname{diam}(\Omega)} \frac{\omega_f(r)}{r} dt < +\infty,$$

where

$$\omega_f(r) := \sup\{|f(x) - f(y)| : x, y \in \Omega, \ |x - y| \le r\}.$$

Its norm is defined by

$$||f||_{C^{Dini}(\bar{\Omega})} = ||f||_{C^0(\bar{\Omega})} + I_{\omega_f}$$

In addition, we say that f is $C^{\alpha,Dini}$ (in Ω) for $0 < \alpha < 1$ if ω_f satisfies

$$I_{\omega_f}^{\alpha} := \int_0^{\operatorname{diam}(\Omega)} \frac{\omega_f(r)}{r^{1+\alpha}} dr < +\infty$$

with the norm

$$||f||_{C^{\alpha,Dini}(\bar{\Omega})} = ||f||_{C^0(\bar{\Omega})} + I^{\alpha}_{\omega_f}$$

It is obvious that Dini continuity is a stronger condition than continuity, and there exists a positive constant C such that

$$\|f\|_{C^{\alpha}(\bar{\Omega})} \le C \|f\|_{C^{\alpha,Dini}(\bar{\Omega})}.$$

The first main result is the Hölder regularity for fractional Poisson equations.

Theorem 1.1. Assume that $u \in C^{1,1}_{loc}(\mathbb{R}^n) \cap \mathcal{L}_{2s}$ is a nonnegative solution of

$$(-\Delta)^s u(x) = f(x), \ x \in B_1(0),$$
 (1.10)

where $f(x) \in L^{\infty}(B_1(0))$. Then there exist some positive constants C_1 and C_2 such that

$$\|u\|_{C^{[2s],\{2s\}}(B_{1/2}(0))} \le C_1 \left(\|f\|_{L^{\infty}(B_1(0))} + \|u\|_{L^{\infty}(B_1(0))} \right), \ if s \neq \frac{1}{2}, \tag{1.11}$$

and

$$\|u\|_{C^{0,\ln L}(B_{1/2}(0))} \le C_2 \left(\|f\|_{L^{\infty}(B_1(0))} + \|u\|_{L^{\infty}(B_1(0))} \right), \ if s = \frac{1}{2}, \tag{1.12}$$

where [2s] and $\{2s\}$ denote the integer part and the fraction part of 2s respectively.

Remark 1.2. Under the assumptions of Theorem 1.1, for any 0 < s < 1 and for all $0 < \varepsilon < 2s$, it follows from (1.11) and (1.12) that

$$\|u\|_{C^{[2s-\varepsilon]},\{2s-\varepsilon\}}(B_{1/2}(0)) \le C\left(\|f\|_{L^{\infty}(B_{1}(0))} + \|u\|_{L^{\infty}(B_{1}(0))}\right).$$

If $f(x) \in C^{\alpha}(\overline{B}_1(0))$ $(0 < \alpha < 1)$, then the following Schauder regularity for $2s + \alpha \notin \mathbb{N}$ and Ln-Lipschitz regularity for $2s + \alpha \in \mathbb{N}$ hold.

Theorem 1.3. Assume that $u \in C^{1,1}_{loc}(\mathbb{R}^n) \cap \mathcal{L}_{2s}$ is a nonnegative solution of (1.10), and $f(x) \in C^{\alpha}(\bar{B}_1(0))$ $(0 < \alpha < 1)$.

If $2s + \alpha \notin \mathbb{N}$, then there exists a constant $C_3 > 0$ such that

$$\|u\|_{C^{[2s+\alpha]},\{2s+\alpha\}}(B_{1/2}(0)) \le C_3\left(\|f\|_{C^{\alpha}(\bar{B}_1(0))} + \|u\|_{L^{\infty}(B_1(0))}\right),\tag{1.13}$$

where $[2s + \alpha]$ and $\{2s + \alpha\}$ are the integer part and the fraction part of $2s + \alpha$ respectively.

If $2s + \alpha \in \mathbb{N}$, then there exists a constant $C_4 > 0$ such that

$$\|u\|_{C^{2s+\alpha-1,\ln L}(B_{1/2}(0))} \le C_4 \left(\|f\|_{C^{\alpha}(\bar{B}_1(0))} + \|u\|_{L^{\infty}(B_1(0))} \right).$$
(1.14)

Remark 1.4. It should be noted that $||u||_{L^{\infty}(B_1(0))}$ can be replaced by $\inf_{B_{3/4}(0)} u$ in (1.11), (1.12),

(1.13) and (1.14). In fact, first, it will follow from (2.7) that $||u||_{L^{\infty}(B_1(0))}$ can be replaced by $||u||_{L^{\infty}(B_{1/2}(0))}$. Second, since $h + ||w||_{L^{\infty}(B_1(0))} + 1 > 0$ and $(-\Delta)^s(h + ||w||_{L^{\infty}(B_1(0))} + 1) = 0$ in $B_1(0)$, then for any $x, y \in B_{3/4}(0)$, it holds that

$$h(x) + \|w\|_{L^{\infty}(B_{1}(0))} + 1 \leq \frac{(9/16 - |x|^{2})^{s}(3/4 + |y|)^{n}}{(9/16 - |y|^{2})^{s}(3/4 - |x|)^{n}} (h(y) + \|w\|_{L^{\infty}(B_{1}(0))} + 1)$$

$$\leq C (h(y) + \|w\|_{L^{\infty}(B_{1}(0))} + 1),$$

which implies

$$\sup_{B_{3/4}(0)} h \le C \inf_{B_{3/4}(0)} h$$

It follows that

$$\begin{split} \|u\|_{L^{\infty}(B_{1/2}(0))} &\leq \|h\|_{L^{\infty}(B_{3/4}(0))} + \|w\|_{L^{\infty}(B_{1}(0))} \\ &\leq C \inf_{B_{3/4}(0)} h + \|w\|_{L^{\infty}(B_{1}(0))} \\ &\leq C \inf_{B_{3/4}(0)} u + 2\|w\|_{L^{\infty}(B_{1}(0))} \\ &\leq C \inf_{B_{3/4}(0)} u + 2\|f\|_{L^{\infty}(B_{1}(0))}. \end{split}$$

Therefore, the conclusion is true.

Remark 1.5. The estimates here are established by a direct method. It worth noting that Jin, Li and Xiong ([20, Theorem 2.11]) addressed a similar problem by exploying an extension method and derived

$$\|u\|_{C^{[2s+\alpha]},\{2s+\alpha\}}(B_{1/2}(0)) \le C\left(\|f\|_{C^{\alpha}(\bar{B}_{1}(0))} + \inf_{B_{3/4}(0)} u\right)$$
(1.15)

for $2s + \alpha \notin \mathbb{N}$. As noted in the above Remark 1.4, this estimate is essentially equivalent to our result in (1.13). However, besides their basic condition that $u \in \mathcal{L}_{2s}$, due to the use of the extension method, they imposed the additional assumption that $u \in \dot{H}^s(\mathbb{R}^n)$. Clearly, this is a significantly stronger requirement than our condition that $u \in \mathcal{L}_{2s}$. There are many \mathcal{L}_{2s} functions that do not belong to $\dot{H}^s(\mathbb{R}^n)$, for instance,

$$u(x) = |x| \text{ for } s > \frac{1}{2} \text{ and } u(x) = (x_n + 2)^{\gamma}_+ \text{ for any } 2s > \gamma > s - \frac{n}{2}$$

are in \mathcal{L}_{2s} , however $||u||_{\dot{H}^{s}(\mathbb{R}^{n})} = \infty$.

Remark 1.6. If there is no sign assumption on u, then the conclusions in Theorem 1.1 and Theorem 1.3 hold with $||u||_{L^{\infty}(B_1(0))}$ replaced by $||u||_{L^{\infty}(\mathbb{R}^n)}$, which have been proved in the above mentioned references [6–8, 11, 18, 22, 26, 27] and so on.

Our next result concerns the $C^{2s+\alpha}$ regularity for nonnegative solutions of the fractional Poisson equation (1.10) under the condition $f(x) \in C^{\alpha,Dini}(\bar{B}_1(0))$ ($0 < \alpha < 1$), which seems to be optimal.

Theorem 1.7. Assume that $u \in C_{loc}^{1,1}(\mathbb{R}^n) \cap \mathcal{L}_{2s}$ is a nonnegative solution of (1.6). Let $f(x) \in C^{\alpha,Dini}(\bar{B}_1(0))$ $(0 < \alpha < 1)$. If $2s + \alpha \in \mathbb{N}$, then there exists a constant $C_5 > 0$ such that

$$\|u\|_{C^{2s+\alpha}(B_{1/2}(0))} \le C_5 \left(\|f\|_{C^{\alpha,Dini}(\bar{B}_1(0))} + \|u\|_{L^{\infty}(B_1(0))} \right).$$

We now state the singularity and decay estimates for solutions to (1.3) derived by constructing an auxiliary function combining with rescaling, for which the following conditions will be assumed:

(H1) $b(x) : \mathbb{R}^n \to \mathbb{R}$ is uniformly bounded and uniformly Hölder continuous.

(H2) $f(x,t): \Omega \times [0,\infty) \to \mathbb{R}$ is uniformly Hölder continuous with respect to x and continuous with respect to t.

(H3) $f(x,t) \leq C_0(1+t^p)$ uniformly for all x in Ω , and

$$\lim_{t \to \infty} \frac{f(x,t)}{t^p} = K(x), \ 1$$

where $K(x) \in (0, \infty)$ is uniformly continuous and $\lim_{|x|\to\infty} K(x) = \overline{C} \in (0, \infty)$.

Theorem 1.8. Let Ω be an arbitrary domain of \mathbb{R}^n and $u \in C^{1,1}_{loc}(\Omega) \cap \mathcal{L}_{2s}$ be a nonnegative solution of (1.3) with b(x) = 0. Suppose that the assumptions (H2) and (H3) hold. Then there exists a positive constant C_6 such that

$$u(x) \le C_6 \left(1 + dist^{-\frac{2s}{p-1}}(x, \partial\Omega) \right), \ x \in \Omega.$$
(1.16)

More precisely, if $\Omega \neq \mathbb{R}^n$, we have

$$u(x) \le C_6 dist^{-\frac{2s}{p-1}}(x, \partial\Omega), \ x \in \Omega.$$
(1.17)

In particular, if Ω is the whole space \mathbb{R}^n , then

$$u(x) \le C_6, \ x \in \mathbb{R}^n; \tag{1.18}$$

if $\Omega \neq \mathbb{R}^n$ is an exterior domain, i.e. $\Omega \supset \{x \in \mathbb{R}^n \mid |x| > R\}$, then

$$u(x) \le C_6 |x|^{-\frac{2s}{p-1}}, \ |x| \ge 2R;$$
 (1.19)

and if Ω is a punctured ball, i.e. $\Omega = B_R(0) \setminus \{0\}$ for some R > 0, then

$$u(x) \le C_6 |x|^{-\frac{2s}{p-1}}, \ 0 < |x| < \frac{R}{2}.$$
 (1.20)

Remark 1.9. As a typical example, (1.18) implies that all the nonnegative solutions to

$$(-\Delta)^s u(x) = u^p(x), \ x \in \mathbb{R}^n$$

are bounded in the subcritical case when 1 . Therefore, in this case, the nonexistence of nontrivial nonnegative bounded solutions implies the nonexistence of nontrivial nonnegative solutions, bounded or not. This is also true for the Lane-Emden system under certain conditions, which may provide new insights and ideas for completely solving the Lane-Emden conjecture. (1.17), (1.19) and (1.20) provide a priori estimates for possible singularities of local solutions to equation (1.3) and also decay estimates in the case of exterior domains.

If (1.3) contains $|\nabla u|$, we also derive similar singularity and decay estimates. In this case, we need to assume $s > \frac{1}{2}$ due to the appearance of the gradient term.

Theorem 1.10. Let Ω be an arbitrary domain in \mathbb{R}^n and $u \in C^{1,1}_{loc}(\Omega) \cap \mathcal{L}_{2s}$ is a nonnegative solution of (1.3). Suppose that the assumptions (H1), (H2) and (H3) hold. If $s > \frac{1}{2}$, then there exists a positive constant C_7 such that

$$u(x) + |\nabla u|^{\frac{2s}{2s+p-1}}(x) \le C_7 \left(1 + dist^{-\frac{2s}{p-1}}(x, \partial\Omega)\right), \ x \in \Omega.$$

More precisely, if $\Omega \neq \mathbb{R}^n$, we have

$$u(x) + |\nabla u|^{\frac{2s}{2s+p-1}}(x) \le C_7 \operatorname{dist}^{-\frac{2s}{p-1}}(x,\partial\Omega), \ x \in \Omega.$$

In particular, if Ω is the whole space, i.e. $\Omega = \mathbb{R}^n$, then

$$u(x) + |\nabla u|^{\frac{2s}{2s+p-1}}(x) \le C_7, \ x \in \mathbb{R}^n;$$

if $\Omega \neq \mathbb{R}^n$ is an exterior domain, i.e. $\Omega \supset \{x \in \mathbb{R}^n \mid |x| > R\}$, then

$$u(x) + |\nabla u|^{\frac{2s}{2s+p-1}}(x) \le C_7 |x|^{-\frac{2s}{p-1}}, \ |x| \ge 2R;$$

and if Ω is a punctured ball, i.e. $\Omega = B_R(0) \setminus \{0\}$ for some R > 0, then

$$u(x) + |\nabla u|^{\frac{2s}{2s+p-1}}(x) \le C_7 |x|^{-\frac{2s}{p-1}}, \ 0 < |x| \le \frac{R}{2}.$$

Remark 1.11. In the previous works on L^{∞} -estimates for solutions to fractional subcritical Lane-Emden equations (see [11, Theorem 9.1.1]), it is often assumed that Ω is bounded and each nonnegative solution is bounded in $\overline{\Omega}$. Then $\|u\|_{L^{\infty}(\Omega)}$ can be bounded by a uniform positive constant independent of u. While Theorems 1.8 and 1.10 give L^{∞} estimate for nonnegative solutions to the fractional subcritical Lane-Emden equation in \mathbb{R}^n , where there is no requirement on the boundedness of solutions in advance since \mathbb{R}^n is unbounded.

The rest of the paper is organized as follows. In Section 2, the interior Hölder continuity of nonnegative solutions to the factional Poisson equation is obtained, which proves Theorem 1.1. In Section 3, according to $2s + \alpha \notin \mathbb{N}$ and $2s + \alpha \in \mathbb{N}$, we classify completely the regularity for nonnegative solutions to the fractional Poisson equation under the condition $f \in C^{\alpha}(\bar{B}_1(0))$, including Schauder regularity if $2s + \alpha \notin \mathbb{N}$ and Ln-Lipschitz regularity if $2s + \alpha \in \mathbb{N}$ (Theorem 1.3). In addition, we prove the $C^{2s+\alpha}$ regularity under the condition $f \in C^{\alpha,Dini}(\bar{B}_1(0))$ and $2s + \alpha \in \mathbb{N}$ (Theorem 1.3). Section 4 is devoted to the proofs of singularity and decay estimates without gradient term (Theorem 1.8), and singularity and decay estimates with gradient term (Theorem 1.10).

2. Hölder regularity

In what follows, we will use C and C_i $(i = 1, 2, \dots)$ to denote generic positive constants, whose values may differ from line to line.

In this section, we show the Hölder regularity for any given nonnegative solution u to (1.10) by splitting u into two parts:

$$u(x) = \left(u(x) - C_{n,s} \int_{B_1(0)} \frac{f(y)}{|x - y|^{n - 2s}} dy\right) + C_{n,s} \int_{B_1(0)} \frac{f(y)}{|x - y|^{n - 2s}} dy$$

:= $h(x) + w(x)$.

It can be proved that the first part

$$h(x) = u(x) - C_{n,s} \int_{B_1(0)} \frac{f(y)}{|x - y|^{n - 2s}} dy$$
(2.1)

is a s-harmonic function with

$$(-\Delta)^{s}h(x) = 0, \ x \in B_1(0).$$

The second part is denoted as

$$w(x) = C_{n,s} \int_{B_1(0)} \frac{f(y)}{|x - y|^{n - 2s}} dy.$$

It is well-known that w(x) is called the Newtonian potential of f if s = 1 and n > 2. In this context, w(x) is briefly called the potential of f in the case that 0 < s < 1 and n > 2s. The following maximum principle will be useded in the grave f

The following maximum principle will be needed in the proof.

Lemma 2.1. [26, Proposition 2.17] Let Ω be a bounded domain in \mathbb{R}^n . Assume that u is a lower-semicontinuous function in $\overline{\Omega}$ and satisfies

$$\begin{cases} (-\Delta)^s u(x) \ge 0, & x \in \Omega\\ u(x) \ge 0, & x \in \mathbb{R}^n \backslash \Omega. \end{cases}$$

in the sense of distribution. Then $u(x) \ge 0$ in \mathbb{R}^n .

To derive the Hölder regularity for nonnegative solutions to the fractional Poisson equation (1.10), we first estimate the s-harmonic function $h(x) = u(x) - C_{n,s} \int_{B_1(0)} \frac{f(y)}{|x-y|^{n-2s}} dy$.

Lemma 2.2. Let the assumptions in Theorem 1.1 hold and h(x) is the s-harmonic function defined in (2.1). Then there exists a positive constant \tilde{C}_k such that for any $k = 0, 1, \cdots$, it holds that

$$\|D^k h\|_{L^{\infty}(B_{1/2}(0))} \le \tilde{C}_k \left(\|f\|_{L^{\infty}(B_1(0))} + \|u\|_{L^{\infty}(B_1(0))} \right)$$

Proof. Since $h(x) \in \mathcal{L}_{2s}$, by Theorem 4.1.2 in [11] or Lemma 4.1 in [21], it holds that $h(x) \in C^{\infty}(B_1(0))$. Then it follows from Theorem 2.10 in [4] that h(x) in $B_1(0)$ can be expressed in terms of an integral on $\mathbb{R}^n \setminus B_1(0)$ consisting of a Poisson kernel and h itself:

$$h(x) = \int_{|y|>1} P(x,y)h(y)dy, \ \forall |x|<1,$$
(2.2)

where

$$P(x,y) = \begin{cases} \frac{\Gamma(n/2)\sin(\pi s)}{\pi^{\frac{n}{2}+1}} \left(\frac{1-|x|^2}{|y|^2-1}\right)^s \frac{1}{|x-y|^n}, & |y| > 1, \\ 0, & |y| \le 1 \end{cases}$$

is the so-called Poisson kernel.

Direct calculations, for any $|x| < \frac{1}{2}$ and |y| > 1, imply

$$\frac{\partial P}{\partial x_i}(x,y) = \left(\frac{-2sx_i}{1-|x|^2} + \frac{-n(x_i-y_i)}{|x-y|^2}\right)P(x,y).$$

Then for any $|x| < \frac{1}{2}$ and |y| > 1, it holds that

$$\left|\frac{\partial P}{\partial x_i}(x,y)\right| \le \left(\frac{2s|x|}{1-|x|^2} + \frac{n}{|x-y|}\right) P(x,y) \le C_1 P(x,y).$$

$$(2.3)$$

For the second order derivatives of h(x), for any $|x| < \frac{1}{2}$ and |y| > 1, and each $i, j = 1, \dots, n$, one has

$$\begin{aligned} \frac{\partial^2 P}{\partial x_i \partial x_j}(x,y) = & \frac{\partial}{\partial x_j} \left(\frac{-2sx_i}{1-|x|^2} + \frac{-n(x_i-y_i)}{|x-y|^2} \right) P(x,y) \\ & + \left(\frac{-2sx_i}{1-|x|^2} + \frac{-n(x_i-y_i)}{|x-y|^2} \right) \frac{\partial P}{\partial x_j}(x,y). \end{aligned}$$

Then the smoothness of the function $\frac{-2sx_i}{1-|x|^2} + \frac{-n(x_i-y_i)}{|x-y|^2}$ and (2.3) imply

$$\begin{split} \left| \frac{\partial^2 P}{\partial x_i \partial x_j}(x, y) \right| &\leq \left| \frac{\partial}{\partial x_j} \left(\frac{-2sx_i}{1 - |x|^2} + \frac{-n(x_i - y_i)}{|x - y|^2} \right) \right| P(x, y) \\ &+ \left| \frac{-2sx_i}{1 - |x|^2} + \frac{-n(x_i - y_i)}{|x - y|^2} \right| \frac{\partial P}{\partial x_j}(x, y) \\ &\leq C_2 P(x, y). \end{split}$$

By recursion, for any $k = 1, 2, \cdots$, we conclude that there exists a constant $C_k > 0$ such that

$$\left| D^k P(x,y) \right| \le C_k P(x,y), \ |x| < \frac{1}{2} \text{ and } |y| > 1.$$
 (2.4)

Due to (2.2) and (2.4), one can obtain

$$\begin{aligned} |D^{k}h(x)| &= \left| \int_{|y|>1} D^{k}P(x,y)h(y)dy \right| \\ &= \left| \int_{|y|>1} D^{k}P(x,y)(u(y) - w(y))dy \right| \\ &\leq \int_{|y|>1} \left| D^{k}P(x,y) \right| (u(y) + |w(y)|)dy \\ &\leq C_{k} \left(\int_{|y|>1} P(x,y)u(y)dy + \int_{|y|>1} P(x,y)|w(y)|dy \right) \\ &:= C_{k}(I_{1} + I_{2}), \ x \in B_{1/2}(0), \end{aligned}$$

$$(2.5)$$

where the fact that u is nonnegative has been used.

Note that

$$|w(x)| = C_{n,s} \left| \int_{B_1(0)} \frac{f(y)}{|x-y|^{n-2s}} dy \right| \le C ||f||_{L^{\infty}(B_1(0))}, \ x \in \mathbb{R}^n \setminus B_1(0),$$

hence

$$I_{2} = \int_{|y|>1} P(x,y)|w(y)|dy \le C \|f\|_{L^{\infty}(B_{1}(0))} \int_{|y|>1} P(x,y)dy \le C \|f\|_{L^{\infty}(B_{1}(0))}.$$
(2.6)

To estimate I_1 , we define

$$v(x) = \begin{cases} \int_{|y|>1} P(x,y)u(y)dy, & x \in B_1(0), \\ u(x), & x \in \mathbb{R}^n \setminus B_1(0). \end{cases}$$

It follows from the definition of P(x, y) and $u \in \mathcal{L}_{2s}$ that $v \in \mathcal{L}_{2s}$.

By virtue of [11, Theorem 4.1.1], we obtain that v(x) satisfies

$$\begin{cases} (-\Delta)^s v(x) = 0, & x \in B_1(0), \\ v(x) = u(x), & x \in \mathbb{R}^n \backslash B_1(0). \end{cases}$$

Recall that the function

$$g_0(x) = \frac{2^{-2s}\Gamma(n/2)}{\Gamma((n+2s)/2)\Gamma(1+s)} (1 - |x|_+^2)^s$$

satisfies

$$(-\Delta)^s g_0(x) = 1.$$

 Set

$$g(x) = ||f||_{L^{\infty}(B_1(0))} g_0(x), \ x \in \mathbb{R}^n.$$

Since $v - (u + g) \in \mathcal{L}_{2s}$ and satisfies

$$\begin{cases} (-\Delta)^s (u+g-v)(x) = f(x) + \|f\|_{L^{\infty}(B_1(0))} \ge 0, & x \in B_1(0), \\ u(x) + g(x) - v(x) = 0, & x \in \mathbb{R}^n \setminus B_1(0), \end{cases}$$

Lemma 2.1 implies that

$$v(x) \le u(x) + g(x), \ x \in \mathbb{R}^n.$$

It follows that

$$I_1 = \int_{|y|>1} P(x,y)u(y)dy \le u(x) + g(x) \le ||u||_{L^{\infty}(B_1(0))} + C||f||_{L^{\infty}(B_1(0))}, \ x \in B_{1/2}(0).$$
(2.7)

Collecting estimates (2.5)–(2.7) shows that for any $k = 0, 1, \cdots$, there exists a positive constant \tilde{C}_k such that

$$\|D^k h\|_{L^{\infty}(B_{1/2}(0))} \leq \tilde{C}_k \left(\|f\|_{L^{\infty}(B_1(0))} + \|u\|_{L^{\infty}(B_1(0))} \right).$$

This completes the proof of Lemma 2.2.

Denote the fundamental solution of the fractional Laplacian $(-\Delta)^s$ in \mathbb{R}^n by

$$\Gamma(x,y) := \frac{C_{n,s}}{|x-y|^{n-2s}}, \ n \ge 3.$$

Then

$$w(x) = \int_{B_1(0)} \Gamma(x, y) f(y) dy.$$

Lemma 2.3. If $f \in L^{\infty}(B_1(0))$ and 2s > 1, or $f \in C^{\alpha}(\overline{B}_1(0))$ $(0 < \alpha < 1)$ and $2s + \alpha > 1$, then $w \in C^1(B_1(0))$ satisfies

$$\partial_i w(x) = \int_{B_1(0)} \partial_i \Gamma(x, y) \left(f(x) - f(y) \right) dy + f(x) \int_{\partial B_1(0)} \Gamma(x, y) \nu_i dS_y, \ i = 1, \cdots, n,$$

where ν_i is the *i*-th component of the unit outward normal vector of $\partial B_1(0)$.

The proof of the lemma is standard which is similar to the case s = 1 in [24].

Proof of Theorem 1.1. Due to Lemma 2.2, it suffices to show

$$\|w\|_{C^{[2s],\{2s\}}(B_{1/2}(0))} \le C\left(\|f\|_{L^{\infty}(B_{1}(0))} + \|u\|_{L^{\infty}(B_{1}(0))}\right), \text{ if } s \neq \frac{1}{2},$$

and

$$||w||_{C^{0,\ln L}(B_{1/2}(0))} \le C \left(||f||_{L^{\infty}(B_1(0))} + ||u||_{L^{\infty}(B_1(0))} \right), \text{ if } s = \frac{1}{2}.$$

For any 0 < s < 1, it is clear that

$$\|w\|_{L^{\infty}(B_{1/2}(0))} = C_{n,s} \sup_{B_{1/2}(0)} \int_{B_{1}(0)} \frac{f(y)}{|x-y|^{n-2s}} dy \le C \|f\|_{L^{\infty}(B_{1}(0))}.$$
(2.8)

1

For any $x, \bar{x} \in B_{1/2}(0)$, let $\zeta = \frac{x+\bar{x}}{2}$ be the midpoint of x and \bar{x} , and $\delta = |x - \bar{x}|$ be the distance between x and \bar{x} . Then

$$\begin{split} |w(x) - w(\bar{x})| = C_{n,s} \left| \int_{B_1(0)} \left(\frac{1}{|x - y|^{n - 2s}} - \frac{1}{|\bar{x} - y|^{n - 2s}} \right) f(y) dy \right| \\ \leq C_{n,s} ||f||_{L^{\infty}(B_1(0))} \int_{B_2(\zeta)} \left| \frac{1}{|x - y|^{n - 2s}} - \frac{1}{|\bar{x} - y|^{n - 2s}} \right| dy \\ \leq C_{n,s} ||f||_{L^{\infty}(B_1(0))} \left(\int_{B_{\delta}(\zeta)} \left| \frac{1}{|x - y|^{n - 2s}} - \frac{1}{|\bar{x} - y|^{n - 2s}} \right| dy \\ + \int_{B_2(\zeta) \setminus B_{\delta}(\zeta)} \left| \frac{1}{|x - y|^{n - 2s}} - \frac{1}{|\bar{x} - y|^{n - 2s}} \right| dy \right). \end{split}$$
(2.9)

Since $B_{\delta}(\zeta) \subset B_{\frac{3}{2}\delta}(x)$, if $s \neq \frac{1}{2}$, one has

$$\int_{B_{\delta}(\zeta)} \frac{1}{|x-y|^{n-2s}} dy \le \int_{B_{\frac{3}{2}\delta}(x)} \frac{1}{|x-y|^{n-2s}} dy \le C \int_{0}^{\frac{3}{2}\delta} r^{2s-1} dr \le C\delta^{2s}.$$

Thus for any $s \in (0, 1)$, it holds that

$$\int_{B_{\delta}(\zeta)} \left| \frac{1}{|x-y|^{n-2s}} - \frac{1}{|\bar{x}-y|^{n-2s}} \right| dy \le \int_{B_{\delta}(\zeta)} \frac{1}{|x-y|^{n-2s}} dy + \int_{B_{\delta}(\zeta)} \frac{1}{|\bar{x}-y|^{n-2s}} dy \le C\delta^{2s}.$$
(2.10)

In addition, noting that

$$|\xi - y| \ge |\zeta - y| - |\zeta - \xi| \ge \frac{|\zeta - y|}{2}, \ y \in B_2(\zeta) \setminus B_\delta(\zeta),$$

by the mean value theorem, if $0 < s < \frac{1}{2}$, one can get

$$\int_{B_{2}(\zeta)\setminus B_{\delta}(\zeta)} \left| \frac{1}{|x-y|^{n-2s}} - \frac{1}{|\bar{x}-y|^{n-2s}} \right| dy$$

$$\leq C \int_{B_{2}(\zeta)\setminus B_{\delta}(\zeta)} \frac{|x-\bar{x}|}{|\xi-y|^{n-2s+1}} dy$$

$$\leq C|x-\bar{x}| \int_{B_{2}(\zeta)\setminus B_{\delta}(\zeta)} \frac{2}{|\zeta-y|^{n-2s+1}} dy$$

$$\leq C|x-\bar{x}| \int_{\delta}^{2} r^{2s-2} dr$$

$$\leq C|x-\bar{x}|^{2s},$$
(2.11)

where ξ lies between x and \bar{x} . If $s = \frac{1}{2}$, a similar argument as (2.11) yields

$$\int_{B_{2}(\zeta)\setminus B_{\delta}(\zeta)} \left| \frac{1}{|x-y|^{n-2s}} - \frac{1}{|\bar{x}-y|^{n-2s}} \right| dy$$

$$\leq C \int_{B_{2}(\zeta)\setminus B_{\delta}(\zeta)} \frac{|x-\bar{x}|}{|\xi-y|^{n}} dy$$

$$\leq C|x-\bar{x}| \int_{B_{2}(\zeta)\setminus B_{\delta}(\zeta)} \frac{2}{|\zeta-y|^{n}} dy$$

$$\leq C|x-\bar{x}| \left| \ln\min(|x-\bar{x}|, 1/2) \right|.$$
(2.12)

It follows from (2.9)–(2.12) that for any $x, \bar{x} \in B_{1/2}(0)$,

$$|w(x) - w(\bar{x})| \le C|x - \bar{x}|^{2s} ||f||_{L^{\infty}(B_1(0))}, \quad \text{if } 0 < s < \frac{1}{2}, \tag{2.13}$$

and

$$|w(x) - w(\bar{x})| \le C|x - \bar{x}| \left| \ln \min(|x - \bar{x}|, 1/2) \right| ||f||_{L^{\infty}(B_1(0))}, \quad \text{if } s = \frac{1}{2}.$$

$$(2.14)$$

Next we estimate $[\partial_i w]_{C^{2s-1}(B_{1/2}(0))}$, $i = 1, \dots, n$, by showing that

$$|\partial_i w(x) - \partial_i w(\bar{x})| \le C |x - \bar{x}|^{2s-1} ||f||_{L^{\infty}(B_1(0))}, \quad \text{if } \frac{1}{2} < s < 1$$
(2.15)

for any $x, \bar{x} \in B_{1/2}(0)$.

If $\frac{1}{2} < s < 1$, by Lemma 2.3, for any $x \in B_{1/2}(0)$ and $i = 1, \dots, n$, one has

$$\begin{split} &|\partial_{i}w(x)| \\ &= \left| \int_{B_{1}(0)} \partial_{i}\Gamma(x-y) \left(f(y) - f(x) \right) dy + f(x) \int_{\partial B_{1}(0)} \Gamma(x-y)\nu_{i} dS_{y} \right| \\ &\leq \left| \int_{B_{1}(0)} \partial_{i}\Gamma(x-y) \left(f(y) - f(x) \right) dy \right| + \left| f(x) \int_{\partial B_{1}(0)} \Gamma(x-y)\nu_{i} dS_{y} \right| \\ &\leq C \|f\|_{L^{\infty}(B_{1}(0))} \int_{B_{1}(0)} \frac{1}{|x-y|^{n-2s+1}} dy + C \|f\|_{L^{\infty}(B_{1}(0))} \\ &\leq C \|f\|_{L^{\infty}(B_{1}(0))}, \end{split}$$

which implies that

$$||Dw||_{L^{\infty}(B_{1/2}(0))} \le C||f||_{L^{\infty}(B_{1}(0))}, \text{ if } \frac{1}{2} < s < 1.$$
 (2.16)

If $\frac{1}{2} < s < 1$, by virtue of Lemma 2.3, for any $x, \bar{x} \in B_{1/2}(0)$, it holds that

$$\begin{split} \partial_i w(x) &- \partial_i w(\bar{x}) \\ &= \int_{B_1(0)} \partial_i \Gamma(x-y) \left(f(y) - f(x) \right) dy - \int_{B_1(0)} \partial_i \Gamma(\bar{x}-y) \left(f(y) - f(\bar{x}) \right) dy \\ &+ f(x) \int_{\partial B_1(0)} \Gamma(x-y) \nu_i dS_y - f(\bar{x}) \int_{\partial B_1(0)} \Gamma(\bar{x}-y) \nu_i dS_y \\ &:= I_1 + I_2, \end{split}$$

where

$$I_{1} = \int_{B_{1}(0)} \partial_{i} \Gamma(x-y) \left(f(y) - f(x)\right) dy - \int_{B_{1}(0)} \partial_{i} \Gamma(\bar{x}-y) \left(f(y) - f(\bar{x})\right) dy$$

and

$$I_2 = f(x) \int_{\partial B_1(0)} \Gamma(x-y)\nu_i dS_y - f(\bar{x}) \int_{\partial B_1(0)} \Gamma(\bar{x}-y)\nu_i dS_y.$$

For I_1 , one can evaluate the integral on $B_{\delta}(\zeta)$ and $B_1(0) \setminus B_{\delta}(\zeta)$ separately, where $\delta = |x - \bar{x}|$ and $\zeta = \frac{x + \bar{x}}{2}$. For the integral I_1 on $B_{\delta}(\zeta)$, one has by direct estimates that

$$\left| \int_{B_{\delta}(\zeta)} \partial_{i} \Gamma(x-y) \left(f(y) - f(x) \right) dy \right| + \left| \int_{B_{\delta}(\zeta)} \partial_{i} \Gamma(x-y) \left(f(y) - f(\bar{x}) \right) dy \right|$$

$$\leq C \|f\|_{L^{\infty}(B_{1}(0))} \int_{B_{\delta}(\zeta)} \frac{1}{|x-y|^{n-2s+1}} dy$$

$$\leq C \|f\|_{L^{\infty}(B_{1}(0))} |x-\bar{x}|^{2s-1}.$$

$$(2.17)$$

The integral I_1 on $B_1(0) \setminus B_{\delta}(\zeta)$ can be estimated as follows

$$\int_{B_{1}(0)\setminus B_{\delta}(\zeta)} \partial_{i}\Gamma(x-y) \left(f(y)-f(x)\right) dy - \int_{B_{1}(0)\setminus B_{\delta}(\zeta)} \partial_{i}\Gamma(\bar{x}-y) \left(f(y)-f(\bar{x})\right) dy$$

$$= \int_{B_{1}(0)\setminus B_{\delta}(\zeta)} \left(\partial_{i}\Gamma(x-y)-\partial_{i}\Gamma(\bar{x}-y)\right) \left(f(y)-f(x)\right) dy$$

$$+ \int_{B_{1}(0)\setminus B_{\delta}(\zeta)} \partial_{i}\Gamma(\bar{x}-y) \left(f(\bar{x})-f(x)\right) dy$$

$$:= I_{11} + I_{12}.$$
(2.18)

The mean value theorem and the fact that $f \in L^{\infty}(B_1(0))$ give

$$|I_{11}| = \left| \int_{B_1(0)\setminus B_{\delta}(\zeta)} \left(\partial_i \Gamma(x-y) - \partial_i \Gamma(\bar{x}-y) \right) (f(y) - f(x)) \, dy \right|$$

$$\leq C \|f\|_{L^{\infty}(B_1(0))} \int_{B_1(0)\setminus B_{\delta}(\zeta)} \frac{|x-\bar{x}|}{|\xi-y|^{n-2s+2}} dy$$

$$\leq C \|f\|_{L^{\infty}(B_1(0))} |x-\bar{x}| \int_{B_1(0)\setminus B_{\delta}(\zeta)} \frac{1}{|\zeta-y|^{n-2s+2}} dy$$

$$\leq C \|f\|_{L^{\infty}(B_1(0))} |x-\bar{x}|^{2s-1},$$
(2.19)

where ξ is a point lying between x and \bar{x} , and one has used the fact that

$$|x-y| \le |x-\zeta| + |\zeta-y| \le 2|\zeta-y|, y \in B_1(0) \setminus B_{\delta}(\zeta),$$

and

$$|\xi - y| \ge |y - \zeta| - |\zeta - \xi| \ge |y - \zeta| - \frac{|y - \zeta|}{2} = \frac{|y - \zeta|}{2}, \ y \in B_1(0) \setminus B_{\delta}(\zeta).$$

For I_{12} , one has by the divergence theorem that

$$I_{12} = (f(\bar{x}) - f(x)) \int_{\partial B_1(0) \cup \partial B_\delta(\zeta)} \Gamma(\bar{x} - y) \nu_i dS_y,$$

which together with I_2 gives

$$\begin{aligned} |I_{12} + I_2| \\ &= \left| \left(f(\bar{x}) - f(x) \right) \int_{\partial B_1(0) \cup \partial B_{\delta}(\zeta)} \Gamma(\bar{x} - y) \nu_i dS_y \right. \\ &+ f(x) \int_{\partial B_1(0)} \Gamma(x - y) \nu_i dS_y - f(\bar{x}) \int_{\partial B_1(0)} \Gamma(\bar{x} - y) \nu_i dS_y \right| \\ &= \left| \left(f(\bar{x}) - f(x) \right) \int_{\partial B_{\delta}(\zeta)} \Gamma(\bar{x} - y) \nu_i dS_y \right. \end{aligned}$$

$$\left. + f(x) \int_{\partial B_1(0)} \left(\Gamma(x - y) - \Gamma(\bar{x} - y) \right) \nu_i dS_y \right| \\ &\leq C[f]_{L^{\infty}(B_1(0))} |x - \bar{x}|^{2s} + |f(x)| \int_{\partial B_1(0)} \frac{|x - \bar{x}|}{|\xi - y|^{n - 2s + 1}} dS_y \\ &\leq C \|f\|_{L^{\infty}(B_1(0))} |x - \bar{x}|^{2s - 1}, \end{aligned}$$

$$(2.20)$$

where ξ lies between x and \bar{x} .

Thus (2.15) follows from (2.17)-(2.20).

By virtue of Lemma 2.2, (2.8), (2.16), (2.13), (2.14) and (2.15), we obtain (1.11) and (1.12). This completes the proof of Theorem 1.1. \Box

3. Schauder Regularity, LN-Lipschitz Regularity and $C^{2s+\alpha}$ regularity

In this section, we show the Schauder regularity and Ln-Lipschitz regularity for nonnegative solutions to (1.10). To this end, we need to estimate $w(x) := \int_{B_1(0)} \Gamma(x, y) f(y) dy$ with $\Gamma(x, y) := \frac{C_{n,s}}{|x-y|^{n-2s}}, n \ge 3.$

Assume that $f \in C^{\alpha}(\bar{B}_1(0))$, which may be extended to be 0 outside of $B_1(0)$.

Lemma 3.1. Let w(x) be the potential of f in $B_1(0)$. Then

$$\|w\|_{C^{[2s+\alpha]},\{2s+\alpha\}}(B_{1/2}(0)) \le C \|f\|_{C^{\alpha}(\bar{B}_{1}(0))}, \quad \text{if } 2s+\alpha \notin \mathbb{N},$$

$$(3.1)$$

and

$$\|w\|_{C^{2s+\alpha-1,\ln L}(B_{1/2}(0))} \le C \|f\|_{C^{\alpha}(\bar{B}_{1}(0))}, \quad \text{if } 2s+\alpha \in \mathbb{N},$$
(3.2)

where $[2s + \alpha]$ and $\{2s + \alpha\}$ are the integer part and the fractional part of $2s + \alpha$ respectively.

Proof. If $2s + \alpha \notin \mathbb{N}$, by a similar argument as Theorem 12.2.2 in [11], one can derive (3.1). Details of the proof are omitted here.

If $2s + \alpha \in \mathbb{N}$, to estimate $||w||_{C^{2s+\alpha-1,\ln L}(B_{1/2}(0))}$, we consider two possible cases: $2s + \alpha = 1$ and $2s + \alpha = 2$.

Case i. $2s + \alpha = 1$.

In this case, we show that

$$\|w\|_{C^{0,\ln L}(B_{1/2}(0))} \le C \|f\|_{C^{\alpha}(\bar{B}_{1}(0))}.$$
(3.3)

By (2.8), it suffices to show that for any $x, \bar{x} \in B_{1/2}(0)$,

$$|w(x) - w(\bar{x})| \le C[f]_{C^{\alpha}(\bar{B}_{1}(0))} |x - \bar{x}| |\ln\min(|x - \bar{x}|, 1/2)|,$$

where C is a constant independent of x and \bar{x} .

Set $\zeta = \frac{x+\bar{x}}{2}$. Then $x, \bar{x} \in B_1(0) \subset B_2(\zeta)$. By symmetry,

$$\int_{B_2(\zeta)} \left(\frac{1}{|x-y|^{n-2s}} - \frac{1}{|\bar{x}-y|^{n-2s}} \right) dy = 0,$$

it thus follows that

$$w(x) - w(\bar{x}) = C_{n,s} \int_{B_1(0)} \left(\frac{1}{|x - y|^{n - 2s}} - \frac{1}{|\bar{x} - y|^{n - 2s}} \right) f(y) dy$$

= $C_{n,s} \int_{B_2(\zeta)} \left(\frac{1}{|x - y|^{n - 2s}} - \frac{1}{|\bar{x} - y|^{n - 2s}} \right) f(y) dy$
= $C_{n,s} \int_{B_2(\zeta)} \left(\frac{1}{|x - y|^{n - 2s}} - \frac{1}{|\bar{x} - y|^{n - 2s}} \right) (f(y) - f(x)) dy,$
(3.4)

due to the fact supp $f \subset B_1(0)$. Denote $\delta = |x - \bar{x}|$. We will estimate the integrand in (3.4) on $B_{\delta}(\zeta)$ and $B_2(\zeta) \setminus B_{\delta}(\zeta)$ respectively. By virtue of $f \in C^{\alpha}(\bar{B}_1(0)), B_{\delta}(\zeta) \subset B_{\frac{3}{2}\delta}(x)$ or $B_{\delta}(\zeta) \subset B_{\frac{3}{2}\delta}(\bar{x})$, one has

$$\begin{split} & \left| \int_{B_{\delta}(\zeta)} \frac{1}{|x-y|^{n-2s}} (f(y) - f(x)) dy \right| \\ \leq & C[f]_{C^{\alpha}(\bar{B}_{1}(0))} \int_{B_{\delta}(\zeta)} \frac{1}{|x-y|^{n-1}} dy \\ \leq & C[f]_{C^{\alpha}(\bar{B}_{1}(0))} \int_{B_{\frac{3}{2}\delta}(x)} \frac{1}{|x-y|^{n-1}} dy \\ \leq & C[f]_{C^{\alpha}(\bar{B}_{1}(0))} |x-\bar{x}|. \end{split}$$

and

$$\begin{split} & \left| \int_{B_{\delta}(\zeta)} \frac{1}{|\bar{x} - y|^{n-2s}} (f(y) - f(x)) dy \right| \\ \leq & C[f]_{C^{\alpha}(\bar{B}_{1}(0))} \int_{B_{\delta}(\zeta)} \frac{|x - y|^{\alpha}}{|\bar{x} - y|^{n-2s}} dy \\ \leq & C[f]_{C^{\alpha}(\bar{B}_{1}(0))} \int_{B_{\delta}(\zeta)} \frac{|x - \bar{x}|^{\alpha} + |\bar{x} - y|^{\alpha}}{|\bar{x} - y|^{n-2s}} dy \\ \leq & C[f]_{C^{\alpha}(\bar{B}_{1}(0))} \left(|x - \bar{x}|^{\alpha} \int_{B_{\frac{3}{2}\delta}(\bar{x})} \frac{1}{|\bar{x} - y|^{n-2s}} dy + \int_{B_{\frac{3}{2}\delta}(\bar{x})} \frac{1}{|\bar{x} - y|^{n-1}} dy \right) \\ \leq & C[f]_{C^{\alpha}(\bar{B}_{1}(0))} |x - \bar{x}|. \end{split}$$

Hence

$$\left| \int_{B_{\delta}(\zeta)} \left(\frac{1}{|x-y|^{n-2s}} - \frac{1}{|\bar{x}-y|^{n-2s}} \right) (f(y) - f(x)) dy \right| \le C[f]_{C^{\alpha}(\bar{B}_{1}(0))} |x-\bar{x}|.$$
(3.5)

For any $y \in B_2(\zeta) \setminus B_{\delta}(\zeta)$, since

$$|x - y| \le |x - \zeta| + |\zeta - y| \le 2|\zeta - y|.$$

Then the mean value theorem implies that

$$\begin{aligned} \left| \int_{B_{2}(\zeta)\setminus B_{\delta}(\zeta)} \left(\frac{1}{|x-y|^{n-2s}} - \frac{1}{|\bar{x}-y|^{n-2s}} \right) (f(y) - f(x)) dy \right| \\ \leq C \int_{B_{2}(\zeta)\setminus B_{\delta}(\zeta)} \frac{|x-\bar{x}|}{|\xi-y|^{n-2s+1}} [f]_{C^{\alpha}(B_{1}(0))} |x-y|^{\alpha} dy \\ \leq C [f]_{C^{\alpha}(\bar{B}_{1}(0))} |x-\bar{x}| \int_{B_{2}(\zeta)\setminus B_{\delta}(\zeta)} \frac{(2|\zeta-y|)^{\alpha}}{|\xi-y|^{n-2s+1}} dy \\ \leq C [f]_{C^{\alpha}(\bar{B}_{1}(0))} |x-\bar{x}| \int_{B_{2}(\zeta)\setminus B_{\delta}(\zeta)} \frac{1}{|\zeta-y|^{n}} dy \\ \leq C [f]_{C^{\alpha}(\bar{B}_{1}(0))} |x-\bar{x}| \left| \ln \min(|x-\bar{x}|, 1/2) \right|, \end{aligned}$$
(3.6)

where ξ is a point lying between x and \bar{x} , and one has used the fact that

$$|\xi - y| \ge |y - \zeta| - |\zeta - \xi| \ge |y - \zeta| - \frac{|y - \zeta|}{2} = \frac{|y - \zeta|}{2}, \ y \in B_2(\zeta) \setminus B_\delta(\zeta).$$

Collecting estimates (3.4)-(3.6) yields

$$|w(x) - w(\bar{x})| \le C[f]_{C^{\alpha}(\bar{B}_{1}(0))} |x - \bar{x}| |\ln\min(|x - \bar{x}|, 1/2)|,$$

which shows that

$$||w||_{C^{0,\ln L}(B_{1/2}(0))} \le C||f||_{C^{\alpha}(\bar{B}_{1}(0))}.$$

Case ii. $2s + \alpha = 2$.

In this case, we will estimate $\|\partial_i w(x)\|_{L^{\infty}(B_{1/2}(0))}$ and $[w]_{C^{1,\ln L}(B_{1/2}(0))}$ to obtain

$$\|w\|_{C^{1,\ln L}(B_{1/2}(0))} \le C \|f\|_{C^{\alpha}(\bar{B}_{1}(0))} |x - \bar{x}|| \ln \min(|x - \bar{x}|, 1/2)|.$$
(3.7)

By Lemma 2.3, for any $x \in B_{1/2}(0)$, it holds that $|\partial_{-w}(x)|$

$$\begin{aligned} &|\partial_{i}w(x)| \\ &= \left| \int_{B_{1}(0)} \partial_{i}\Gamma(x-y) \left(f(y) - f(x)\right) dy + f(x) \int_{\partial B_{1}(0)} \Gamma(x-y)\nu_{i} dS_{y} \right| \\ &\leq \left| \int_{B_{1}(0)} \partial_{i}\Gamma(x-y) \left(f(y) - f(x)\right) dy \right| + \left| f(x) \int_{\partial B_{1}(0)} \Gamma(x-y)\nu_{i} dS_{y} \right| \\ &\leq C[f]_{C^{\alpha}(\bar{B}_{1}(0))} \int_{B_{1}(0)} \frac{1}{|x-y|^{n-1}} dy + C ||f||_{L^{\infty}(B_{1}(0))} \\ &\leq C ||f||_{C^{\alpha}(\bar{B}_{1}(0))}. \end{aligned}$$
(3.8)

Next we estimate $[w]_{C^{1,\ln L}(B_{1/2}(0))}$ by showing that

$$|\partial_i w(x) - \partial_i w(\bar{x})| \le C ||f||_{C^{\alpha}(\bar{B}_1(0))} |x - \bar{x}|| \ln \min(|x - \bar{x}|, 1/2)|, \ \forall x, \bar{x} \in B_{1/2}(0).$$

By virtue of Lemma 2.3, for any $x, \bar{x} \in B_{1/2}(0)$, one has

$$\partial_i w(x) - \partial_i w(\bar{x}) := I_1 + I_2,$$

where

$$I_{1} = \int_{B_{1}(0)} \partial_{i} \Gamma(x-y) \left(f(y) - f(x)\right) dy - \int_{B_{1}(0)} \partial_{i} \Gamma(\bar{x}-y) \left(f(y) - f(\bar{x})\right) dy$$

and

$$I_2 = f(x) \int_{\partial B_1(0)} \Gamma(x-y)\nu_i dS_y - f(\bar{x}) \int_{\partial B_1(0)} \Gamma(\bar{x}-y)\nu_i dS_y.$$

It can be estimated in a similar way as for the Case i by evaluating the integrands in $B_{\delta}(\zeta)$ and $B_1(0) \setminus B_{\delta}(\zeta)$ separately, where $\delta = |x - \bar{x}|$ and $\zeta = \frac{x + \bar{x}}{2}$. On $B_{\delta}(\zeta)$, note that

$$\begin{split} & \left| \int_{B_{\delta}(\zeta)} \partial_{i} \Gamma(x-y) \left(f(y) - f(x) \right) dy \right| \\ \leq & C[f]_{C^{\alpha}(\bar{B}_{1}(0))} \int_{B_{\delta}(\zeta)} \frac{1}{|x-y|^{n-1}} dy \\ \leq & C[f]_{C^{\alpha}(\bar{B}_{1}(0))} |x-\bar{x}|, \end{split}$$

and

$$\left| \int_{B_{\delta}(\zeta)} \partial_i \Gamma(\bar{x} - y) \left(f(y) - f(\bar{x}) \right) dy \right| \le C[f]_{C^{\alpha}(\bar{B}_1(0))} |x - \bar{x}|.$$

Then

$$\left| \int_{B_{\delta}(\zeta)} \partial_{i} \Gamma(x-y) \left(f(y) - f(x) \right) dy - \int_{B_{\delta}(\zeta)} \partial_{i} \Gamma(\bar{x}-y) \left(f(y) - f(\bar{x}) \right) dy \right|$$

$$\leq C[f]_{C^{\alpha}(\bar{B}_{1}(0))} |x-\bar{x}|.$$

$$(3.9)$$

On $B_1(0) \setminus B_{\delta}(\zeta)$, one has

$$\int_{B_{1}(0)\setminus B_{\delta}(\zeta)} \partial_{i}\Gamma(x-y) \left(f(y)-f(x)\right) dy - \int_{B_{1}(0)\setminus B_{\delta}(\zeta)} \partial_{i}\Gamma(\bar{x}-y) \left(f(y)-f(\bar{x})\right) dy$$

$$= \int_{B_{1}(0)\setminus B_{\delta}(\zeta)} \left(\partial_{i}\Gamma(x-y)-\partial_{i}\Gamma(\bar{x}-y)\right) \left(f(y)-f(x)\right) dy$$

$$+ \int_{B_{1}(0)\setminus B_{\delta}(\zeta)} \partial_{i}\Gamma(\bar{x}-y) \left(f(\bar{x})-f(x)\right) dy$$

$$:= I_{11} + I_{12}.$$
(3.10)

By the mean value theorem and the fact that $f \in C^{\alpha}(\bar{B}_1(0))$, one can get

$$\begin{aligned} |I_{11}| &= \left| \int_{B_1(0)\setminus B_{\delta}(\zeta)} \left(\partial_i \Gamma(x-y) - \partial_i \Gamma(\bar{x}-y) \right) (f(y) - f(x)) \, dy \right| \\ &\leq \int_{B_1(0)\setminus B_{\delta}(\zeta)} \frac{|x-\bar{x}|}{|\xi-y|^{n-2s+2}} [f]_{C^{\alpha}(B_1(0))} |x-y|^{\alpha} dy \\ &\leq C[f]_{C^{\alpha}(\bar{B}_1(0))} |x-\bar{x}| \int_{B_1(0)\setminus B_{\delta}(\zeta)} \frac{|\zeta-y|^{\alpha}}{|\zeta-y|^{n-2s+2}} dy \\ &\leq C[f]_{C^{\alpha}(\bar{B}_1(0))} |x-\bar{x}| \left| \ln \min(|x-\bar{x}|, 1/2) \right|, \end{aligned}$$
(3.11)

where ξ is a point lying between x and \bar{x} , and one has used the fact that

$$|x-y| \le |x-\zeta| + |\zeta-y| \le 2|\zeta-y|, y \in B_1(0) \setminus B_{\delta}(\zeta),$$

and

$$|\xi - y| \ge |y - \zeta| - |\zeta - \xi| \ge |y - \zeta| - \frac{|y - \zeta|}{2} = \frac{|y - \zeta|}{2}, \ y \in B_1(0) \setminus B_{\delta}(\zeta).$$

For the integral I_{12} , the divergence theorem yields

$$I_{12} = (f(\bar{x}) - f(x)) \int_{\partial B_1(0) \cup \partial B_\delta(\zeta)} \Gamma(\bar{x} - y) \nu_i dS_y,$$

which, combined with I_2 , gives

$$\begin{aligned} |I_{12} + I_2| \\ &= \left| \left(f(\bar{x}) - f(x) \right) \int_{\partial B_1(0) \cup \partial B_{\delta}(\zeta)} \Gamma(\bar{x} - y) \nu_i dS_y \\ &+ f(x) \int_{\partial B_1(0)} \Gamma(x - y) \nu_i dS_y - f(\bar{x}) \int_{\partial B_1(0)} \Gamma(\bar{x} - y) \nu_i dS_y \right| \\ &= \left| \left(f(\bar{x}) - f(x) \right) \int_{\partial B_{\delta}(\zeta)} \Gamma(\bar{x} - y) \nu_i dS_y \\ &+ f(x) \int_{\partial B_1(0)} \left(\Gamma(x - y) - \Gamma(\bar{x} - y) \right) \nu_i dS_y \right| \\ &\leq C[f]_{C^{\alpha}(\bar{B}_1(0))} |x - \bar{x}| + |f(x)| \int_{\partial B_1(0)} \frac{|x - \bar{x}|}{|\xi - y|^{n - 2s + 1}} dS_y \\ &\leq C \|f\|_{C^{\alpha}(\bar{B}_1(0))} |x - \bar{x}|, \end{aligned}$$
(3.12)

where ξ lies between x and \bar{x} .

It follows from (3.8)–(3.12) that

$$||w||_{C^{1,\ln L}(B_{1/2}(0))} \le C||f||_{C^{\alpha}(\bar{B}_{1}(0))}|x-\bar{x}||\ln\min(|x-\bar{x}|,1/2)|.$$

Combining the two cases leads to (3.2) by (3.3) and (3.7). This completes the proof of Lemma 3.1.

Proof of Theorem 1.3. To show the Schauder regularity for nonnegative solutions to the fractional Poisson equation (1.10) in the case $f \in C^{\alpha}(\bar{B}_1(0))$ and $2s + \alpha \notin \mathbb{N}$, one needs only to combine the estimate for w(x) in (3.1) and with the estimate for h(x) in Lemma 2.2.

Then if $2s + \alpha \notin \mathbb{N}$, it holds that

$$\begin{aligned} &\|u\|_{C^{[2s+\alpha]},\{2s+\alpha\}}(B_{1/2}(0))\\ \leq &\|w\|_{C^{[2s+\alpha]},\{2s+\alpha\}}(B_{1/2}(0)) + \|h\|_{C^{[2s+\alpha]},\{2s+\alpha\}}(B_{1/2}(0))\\ \leq &\|w\|_{C^{[2s+\alpha]},\{2s+\alpha\}}(B_{1/2}(0)) + \|h\|_{C^{[2s+\alpha]},\{2s+\alpha\}}(B_{1/2}(0))\\ \leq &C\left(\|f\|_{C^{\alpha}(\bar{B}_{1}(0))} + \|u\|_{L^{\infty}(B_{1}(0))}\right),\end{aligned}$$

which shows (1.13).

For the Ln-Lipschitz regularity of nonnegative solutions to the fractional Poisson equation (1.10) in the case $f \in C^{\alpha}(\bar{B}_1(0))$ and $2s + \alpha \in \mathbb{N}$, it suffices to combine the estimate for w(x) in (3.2) and with the estimate for h(x) in Lemma 2.2.

Therefore, if $2s + \alpha \in \mathbb{N}$, it follows from (3.2) and Lemma 2.2 that

$$\begin{aligned} &\|u\|_{C^{2s+\alpha-1,\ln L}(B_{1/2}(0))} \\ \leq &\|w\|_{C^{2s+\alpha-1,\ln L}(B_{1/2}(0))} + \|h\|_{C^{2s+\alpha-1,\ln L}(B_{1/2}(0))} \\ \leq &\|w\|_{C^{2s+\alpha-1,\ln L}(B_{1/2}(0))} + \|D^{2s+\alpha}h\|_{L^{\infty}(B_{1/2}(0))} \\ \leq &C\left(\|f\|_{C^{\alpha}(\bar{B}_{1}(0))} + \|u\|_{L^{\infty}(B_{1}(0))}\right). \end{aligned}$$

Therefore, (1.14) holds. This completes the proof of Theorem 1.3.

Proof of Theorem 1.7. If $2s + \alpha \in \mathbb{N}$ and $f \in C^{\alpha,Dini}(\bar{B}_1(0))$, we can show the $C^{2s+\alpha}$ regularity for nonnegative solutions to the fractional Poisson equation (1.10) by considering two cases: $2s + \alpha = 1$ and $2s + \alpha = 2$.

Case 1. $2s + \alpha = 1$.

Then we claim that

$$\|u\|_{C^{1}(B_{1/2}(0))} \le C\left(\|f\|_{C^{\alpha,Dini}(\bar{B}_{1}(0))} + \|u\|_{L^{\infty}(B_{1}(0))}\right)$$

Indeed, by Lemma 2.2, it suffices to prove

$$\|w\|_{C^{1}(B_{1/2}(0))} \le C \|f\|_{C^{\alpha,Dini}(\bar{B}_{1}(0))}.$$
(3.13)

Let ζ be the midpoint of x and \bar{x} , then $x, \bar{x} \in B_1(0) \subset B_2(\zeta)$. Denote $\delta = |x - \bar{x}|$. It follows from (3.4) that

$$\begin{split} w(x) - w(\bar{x}) \\ = & C_{n,s} \int_{B_2(\zeta)} \left(\frac{1}{|x - y|^{n - 2s}} - \frac{1}{|\bar{x} - y|^{n - 2s}} \right) (f(y) - f(x)) dy \\ = & C_{n,s} \int_{B_{\delta}(\zeta)} \left(\frac{1}{|x - y|^{n - 2s}} - \frac{1}{|\bar{x} - y|^{n - 2s}} \right) (f(y) - f(x)) dy \\ &+ & C_{n,s} \int_{B_2(\zeta) \setminus B_{\delta}(\zeta)} \left(\frac{1}{|x - y|^{n - 2s}} - \frac{1}{|\bar{x} - y|^{n - 2s}} \right) (f(y) - f(x)) dy \\ := & C_{n,s} (I_1 + I_2). \end{split}$$
(3.14)

 I_1 and I_2 can be treated respectively as follows. By virtue of $f \in C^{\alpha,Dini}(\bar{B}_1(0))$ and $2s + \alpha = 1$, one can get

$$\begin{split} & \left| \int_{B_{\delta}(\zeta)} \frac{1}{|x-y|^{n-2s}} \left(f(y) - f(x) \right) dy \right| \\ & \leq \left| \int_{B_{3\delta/2}(x)} \frac{1}{|x-y|^{n-2s-\alpha}} \|f\|_{C^{\alpha}(\bar{B}_{1}(0))} dy \right| \\ & \leq C \delta \|f\|_{C^{\alpha}(\bar{B}_{1}(0))} \\ & \leq C |x-\bar{x}| \|f\|_{C^{\alpha,Dini}(\bar{B}_{1}(0))}, \end{split}$$

and

$$\begin{split} & \left| \int_{B_{\delta}(\zeta)} \frac{1}{|\bar{x} - y|^{n-2s}} \left(f(y) - f(x) \right) dy \right| \\ \leq & \|f\|_{C^{\alpha}(\bar{B}_{1}(0))} \int_{B_{\delta}(\zeta)} \frac{|x - y|^{\alpha}}{|\bar{x} - y|^{n-2s}} dy \\ \leq & \|f\|_{C^{\alpha}(\bar{B}_{1}(0))} \int_{B_{\delta}(\zeta)} \frac{|x - \bar{x}|^{\alpha} + |\bar{x} - y|^{\alpha}}{|\bar{x} - y|^{n-2s}} dy \\ \leq & \|f\|_{C^{\alpha}(\bar{B}_{1}(0))} \left(|x - \bar{x}|^{\alpha} \int_{B_{3\delta/2}(\bar{x})} \frac{1}{|\bar{x} - y|^{n-2s}} dy + \int_{B_{3\delta/2}(\bar{x})} \frac{1}{|\bar{x} - y|^{n-1}} dy \right) \\ \leq & C |x - \bar{x}| \|f\|_{C^{\alpha}(\bar{B}_{1}(0))} \\ \leq & C |x - \bar{x}| \|f\|_{C^{\alpha}(\bar{B}_{1}(0))}. \end{split}$$

Hence,

$$|I_1| \le C|x - \bar{x}| \|f\|_{C^{\alpha,Dini}(\bar{B}_1(0))}.$$
(3.15)

For I_2 , since

$$|x-y| \le |x-\zeta| + |\zeta-y| \le 2|\zeta-y|, \ x \in B_{1/2}(0), \ y \in B_2(\zeta) \setminus B_{\delta}(\zeta),$$

one can use the mean value theorem to derive

$$\begin{aligned} |I_{2}| &= \left| \int_{B_{2}(\zeta) \setminus B_{\delta}(\zeta)} \left(\frac{1}{|x-y|^{n-2s}} - \frac{1}{|\bar{x}-y|^{n-2s}} \right) (f(y) - f(x)) dy \right| \\ &\leq C \int_{B_{2}(\zeta) \setminus B_{\delta}(\zeta)} \frac{|x-\bar{x}|}{|\xi-y|^{n-2s+1}} \omega_{f}(|x-y|) dy \\ &\leq C|x-\bar{x}| \int_{B_{2}(\zeta) \setminus B_{\delta}(\zeta)} \frac{\omega_{f}(2|\zeta-y|)}{|\xi-y|^{n-2s+1}} dy \\ &\leq C|x-\bar{x}| \int_{B_{2}(\zeta) \setminus B_{\delta}(\zeta)} \frac{\omega_{f}(2|\zeta-y|)}{|\zeta-y|^{n-2s+1}} dy \\ &\leq C|x-\bar{x}| \int_{\delta}^{2} \frac{\omega_{f}(2r)}{r^{1+\alpha}} dr \\ &\leq C|x-\bar{x}| ||f||_{C^{\alpha,Dini}(\bar{B}_{1}(0))}, \end{aligned}$$
(3.16)

where ξ is a point lying between x and \bar{x} , and one has used the fact that

$$|\xi - y| \ge |y - \zeta| - |\zeta - \xi| \ge |y - \zeta| - \frac{|y - \zeta|}{2} = \frac{|y - \zeta|}{2}, \ y \in B_2(\zeta) \setminus B_\delta(\zeta).$$

Collecting (3.14)-(3.16) yields (3.13).

Case 2. $2s + \alpha = 2$.

Then it holds that

$$\|u\|_{C^2(B_{1/2}(0))} \le C\left(\|f\|_{C^{\alpha,Dini}(\bar{B}_1(0))} + \|u\|_{L^{\infty}(B_1(0))}\right),$$

which follows from Lemma 2.2 and the following estimate

$$\|w\|_{C^2(B_{1/2}(0))} \le C \|f\|_{C^{\alpha,Dini}(\bar{B}_1(0))}.$$
(3.17)

To prove (3.17), by Lemma 2.3, for any $x \in B_{1/2}(0)$, one has

$$\begin{aligned} |\partial_{i}w(x)| \\ &= \left| \int_{B_{1}(0)} \partial_{i}\Gamma(x-y) \left(f(y) - f(x)\right) dy + f(x) \int_{\partial B_{1}(0)} \Gamma(x-y)\nu_{i} dS_{y} \right| \\ &\leq \left| \int_{B_{1}(0)} \partial_{i}\Gamma(x-y)\omega_{f}(|x-y|) dy \right| + \left| f(x) \int_{\partial B_{1}(0)} \Gamma(x-y)\nu_{i} dS_{y} \right| \\ &\leq C \int_{B_{1}(0)} \frac{\omega_{f}(|x-y|)}{|x-y|^{n-2s+1}} dy + C \|f\|_{L^{\infty}(B_{1}(0))} \\ &\leq C \int_{0}^{1} \frac{\omega_{f}(r)}{r^{1+\alpha}} dy + C \|f\|_{L^{\infty}(B_{1}(0))} \\ &\leq C \|f\|_{C^{\alpha,Dini}(\bar{B}_{1}(0))}. \end{aligned}$$
(3.18)

Next, the second order derivatives of w can be estimated as

$$|\partial_i w(x) - \partial_i w(\bar{x})| \le C |x - \bar{x}| \|f\|_{C^{\alpha, Dini}(\bar{B}_1(0))}, \ \forall x, \bar{x} \in B_{1/2}(0).$$
(3.19)

To this end, by Lemma 2.3, for any $x, \bar{x} \in B_{1/2}(0)$, one has

$$\partial_i w(x) - \partial_i w(\bar{x}) := J_1 + J_2,$$

where

$$J_{1} = \int_{B_{1}(0)} \partial_{i} \Gamma(x-y) \left(f(y) - f(x)\right) dy - \int_{B_{1}(0)} \partial_{i} \Gamma(\bar{x}-y) \left(f(y) - f(\bar{x})\right) dy$$

and

$$J_2 = f(x) \int_{\partial B_1(0)} \Gamma(x-y)\nu_i dS_y - f(\bar{x}) \int_{\partial B_1(0)} \Gamma(\bar{x}-y)\nu_i dS_y.$$

Decompose J_1 into integrals on $B_{\delta}(\zeta)$ and $B_1(0) \setminus B_{\delta}(\zeta)$ and estimate each one as follows. First, it holds

$$\left| \int_{B_{\delta}(\zeta)} \partial_{i} \Gamma(x-y) \left(f(y) - f(x) \right) dy \right|$$

$$\leq C \int_{B_{3\delta/2}(x)} \frac{1}{|x-y|^{n-2s+1}} \omega_{f}(|x-y|) dy$$

$$\leq C|x-\bar{x}| \int_{0}^{3\delta/2} \frac{\omega_{f}(r)}{r^{1+\alpha}} dr$$

$$\leq C|x-\bar{x}| ||f||_{C^{\alpha,Dini}(\bar{B}_{1}(0))},$$
(3.20)

similarly, one has

$$\left| \int_{B_{\delta}(\zeta)} \partial_{i} \Gamma(\bar{x} - y) \left(f(y) - f(\bar{x}) \right) dy \right| \leq C |x - \bar{x}| \|f\|_{C^{\alpha, Dini}(\bar{B}_{1}(0))}.$$
(3.21)

Next, note that

$$\int_{B_{1}(0)\setminus B_{\delta}(\zeta)} \partial_{i}\Gamma(x-y) \left(f(y)-f(x)\right) dy - \int_{B_{1}(0)\setminus B_{\delta}(\zeta)} \partial_{i}\Gamma(\bar{x}-y) \left(f(y)-f(\bar{x})\right) dy$$

$$= \int_{B_{1}(0)\setminus B_{\delta}(\zeta)} \left(\partial_{i}\Gamma(x-y)-\partial_{i}\Gamma(\bar{x}-y)\right) \left(f(y)-f(x)\right) dy$$

$$+ \int_{B_{1}(0)\setminus B_{\delta}(\zeta)} \partial_{i}\Gamma(\bar{x}-y) \left(f(\bar{x})-f(x)\right) dy$$

$$:= J_{11} + J_{12}.$$
(3.22)

It follows from the mean value theorem and $f \in C^{\alpha,Dini}(\bar{B}_1(0))$ that

$$\begin{aligned} J_{11} &= \left| \int_{B_1(0) \setminus B_{\delta}(\zeta)} \left(\partial_i \Gamma(x-y) - \partial_i \Gamma(\bar{x}-y) \right) \left(f(y) - f(x) \right) dy \right| \\ &\leq \int_{B_1(0) \setminus B_{\delta}(\zeta)} \frac{|x-\bar{x}|}{|\xi-y|^{n-2s+2}} \omega_f(|x-y|) dy \\ &\leq C|x-\bar{x}| \int_{B_1(0) \setminus B_{\delta}(\zeta)} \frac{\omega_f(2|\zeta-y|)}{|\zeta-y|^{n-2s+2}} dy \\ &\leq C|x-\bar{x}| \int_{\delta}^2 \frac{\omega_f(2r)}{r^{1+\alpha}} dr \\ &\leq C|x-\bar{x}| \|f\|_{C^{\alpha,Dini}(\bar{B}_1(0))}, \end{aligned}$$

$$(3.23)$$

where ξ is a point lying between x and \bar{x} , and one has used the facts that

$$|x-y| \le |x-\zeta| + |\zeta-y| \le 2|\zeta-y|, \ y \in B_1(0) \setminus B_{\delta}(\zeta),$$

and

$$|\xi - y| \ge |y - \zeta| - |\zeta - \xi| \ge |y - \zeta| - \frac{|y - \zeta|}{2} = \frac{|y - \zeta|}{2}, \ y \in B_1(0) \setminus B_\delta(\zeta).$$

For the integral J_{12} , the divergence theorem implies

$$J_{12} = (f(\bar{x}) - f(x)) \int_{\partial B_1(0) \cup \partial B_\delta(\zeta)} \Gamma(\bar{x} - y) \nu_i dS_y.$$

Since

$$|f(x) - f(\bar{x})| \le C|x - \bar{x}| ||f||_{C^{\alpha}(\bar{B}_{1}(0))} \le C|x - \bar{x}| ||f||_{C^{\alpha,Dini}(\bar{B}_{1}(0))},$$

we can estimate J_{12} together J_2 to obtain

$$\begin{aligned} |J_{12} + J_2| \\ &= \left| (f(\bar{x}) - f(x)) \int_{\partial B_1(0) \cup \partial B_{\delta}(\zeta)} \Gamma(\bar{x} - y) \nu_i dS_y \right. \\ &+ f(x) \int_{\partial B_1(0)} \Gamma(x - y) \nu_i dS_y - f(\bar{x}) \int_{\partial B_1(0)} \Gamma(\bar{x} - y) \nu_i dS_y \right| \\ &= \left| (f(\bar{x}) - f(x)) \int_{\partial B_{\delta}(\zeta)} \Gamma(\bar{x} - y) \nu_i dS_y \right. \\ &+ f(x) \int_{\partial B_1(0)} (\Gamma(x - y) - \Gamma(\bar{x} - y)) \nu_i dS_y \right| \\ &\leq C |x - \bar{x}|^{\alpha} ||f||_{C^{\alpha}(\bar{B}_1(0))} |x - \bar{x}|^{2s - 1} + |f(x)| \int_{\partial B_1(0)} \frac{|x - \bar{x}|}{|\xi - y|^{n - 2s + 1}} dS_y \\ &\leq C |x - \bar{x}| ||f||_{C^{\alpha, Dini}(\bar{B}_1(0))}, \end{aligned}$$

$$(3.24)$$

where ξ lies between x and \bar{x} , and one has used the fact $2s + \alpha = 2$.

Collecting (3.20)–(3.24) leads to (3.19), which together with (3.18) gives (3.17). This completes the proof of Theorem 1.7.

4. A priori estimates

Proof of Theorem 1.8. By the assumptions, u sloves the semi-linear equation

$$(-\Delta)^s u(x) = f(x, u(x)), \ x \in \Omega,$$

and we will show the following a priori estimates

 $u(x) \leq C(1 + dist^{-\frac{2s}{p-1}}(x,\partial\Omega)), \ x \in \Omega$

by a contradiction argument.

Assume otherwise. Then, there exist sequences of Ω_k , u_k , $x_k \in \Omega_k$, such that u_k solves

$$\begin{cases} (-\Delta)^s u_k(x) = f(x, u_k(x)), & x \in \Omega_k, \\ u_k(x) = 0, & x \in \mathbb{R}^n \setminus \Omega_k, \end{cases}$$
(4.1)

and

$$\frac{1}{2}(u_k(x_k))^{\frac{p-1}{2s}}d_k \ge \frac{1}{2}(u_k(x_k))^{\frac{p-1}{2s}}(1+d_k^{-1})^{-1} > k \to \infty \text{ as } k \to \infty,$$
(4.2)

where $d_k = dist(x_k, \partial \Omega_k)$.

Define

$$r_k := 2ku_k^{-\frac{p-1}{2s}}(x_k)$$

Then (4.2) implies that $r_k < d_k$ and thus $B_{r_k}(x_k) \subset \Omega_k$.

Define a function

$$S_k(x) = u_k(x)(r_k - |x - x_k|)^{\frac{2s}{p-1}}, \ x \in B_{r_k}(x_k)$$

Then there exists $a_k \in B_{r_k}(x_k)$ such that

$$S_k(a_k) = \max_{B_{r_k}(x_k)} S_k(x).$$

It follows that

$$u_k(x) \le u_k(a_k) \frac{(r_k - |a_k - x_k|)^{\frac{2s}{p-1}}}{(r_k - |x - x_k|)^{\frac{2s}{p-1}}}, \ x \in B_{r_k}(x_k).$$

$$(4.3)$$

 Set

$$\lambda_k = (u_k(a_k))^{-\frac{p-1}{2s}}$$

Taking $x = x_k$ in (4.3) yields

$$\frac{u_k(x_k)}{u_k(a_k)} \le \frac{(r_k - |a_k - x_k|)^{\frac{2s}{p-1}}}{r_k^{\frac{2s}{p-1}}},\tag{4.4}$$

which implies that

$$u_k(x_k) \le u_k(a_k)$$

Using the definition of r_k and λ_k , one gets from (4.4) that

$$2k\lambda_k \le r_k - |a_k - x_k|. \tag{4.5}$$

In addition, one can get

$$|x_k - |a_k - x_k| \le 2(r_k - |x - x_k|), \ x \in B_{k\lambda_k}(a_k).$$
(4.6)

Indeed, for any $x \in B_{k\lambda_k}(a_k)$, one has

$$|x - x_k| \le |x - a_k| + |a_k - x_k| \le k\lambda_k + r_k - 2k\lambda_k \le r_k$$

due to (4.5). It follows that $x \in B_{r_k}(x_k)$ and thus $B_{k\lambda_k}(a_k) \subset B_{r_k}(x_k)$. Now for any $x \in B_{k\lambda_k}(a_k)$, using (4.5), one can get

$$2(r_{k} - |x - x_{k}|) \ge 2(r_{k} - (|x_{k} - a_{k}| + |x - a_{k}|))$$

$$\ge 2(r_{k} - (|x_{k} - a_{k}| + k\lambda_{k}))$$

$$\ge r_{k} - |x_{k} - a_{k}| + 2k\lambda_{k} - 2k\lambda_{k}$$

$$\ge r_{k} - |x_{k} - a_{k}|,$$

therefore, (4.6) follows.

The combination of (4.3) with (4.6) yields

$$u_k(x) \le u_k(a_k) 2^{\frac{2s}{p-1}}, \ x \in B_{k\lambda_k}(a_k).$$
 (4.7)

We now rescale the solutions as

$$v_k(x) = \frac{1}{u_k(a_k)} u_k(\lambda_k x + a_k), \ x \in B_k(0).$$

By (4.1), it is easy to check that the function $v_k(x)$ solves

$$(-\Delta)^{s} v_{k}(x) = \lambda_{k}^{\frac{2sp}{p-1}} f(\lambda_{k} x + a_{k}, \lambda_{k}^{-\frac{2s}{p-1}} v_{k}(x)) := F_{k}(x, v_{k}(x)), \ x \in B_{k}(0).$$
(4.8)

Moreover, it follows from the definition of v_k and (4.7) that

$$v_k(0) = 1 \tag{4.9}$$

and

$$v_k(x) \le 2^{\frac{2s}{p-1}}, \ x \in B_k(0).$$
 (4.10)

Then by (4.10) and the assumptions (H1)-(H3), one can obtain

$$F_k(x, v_k(x)) = \lambda_k^{\frac{2sp}{p-1}} f(\lambda_k x + a_k, \lambda_k^{-\frac{2s}{p-1}} v_k(x))$$

$$\leq C_0 \lambda_k^{\frac{2sp}{p-1}} \left(1 + \lambda_k^{-\frac{2sp}{p-1}} v_k^p(x) \right)$$

$$\leq C_0 \lambda_k^{\frac{2sp}{p-1}} + C_0 v_k^p(x)$$

$$\leq C_1.$$

Applying Theorem 1.1 to (4.8) yields

$$||v_k||_{C^{[2s-\varepsilon]},\{2s-\varepsilon\}}(B_{3k/4}(0)) \le C_2$$

for all $0 < \varepsilon < 2s$. It follows that $F_k(x, v_k(x))$ is Hölder continuous in $B_{3k/4}(0)$. Then by virtue of Theorem 1.3, we conclude that there exist some positive constants $\beta \in (0, 1)$ and C_3 independent of k such that

$$\|v_k\|_{C^{[2s+\beta]},\{2s+\beta\}}(B_{k/2}(0)) \le C_3.$$
(4.11)

Due to the above uniform regularity estimates, it then follows from the Ascoli-Arzelá theorem that there exists a converging subsequence of $\{v_k\}$ (stilled denoted by $\{v_k\}$) that converges pointwisely in \mathbb{R}^n to a function v(x). Taking limit in (4.9) yields

$$v(0) = 1. (4.12)$$

Moreover, v_k converges in $C_{loc}^{[2s+\epsilon],\{2s+\epsilon\}}(\mathbb{R}^n)$ and hence $(-\Delta)^s v_k$ converges point-wisely in \mathbb{R}^n . Applying a result in [19, Theorem 1.1] we obtain

$$\lim_{k \to \infty} (-\Delta)^s v_k(x) = (-\Delta)^s v(x) - b, \forall x \in \mathbb{R}^n$$
(4.13)

with

$$b = C_{n,s} \lim_{R \to \infty} \lim_{k \to \infty} \int_{B_R^c} \frac{v_k(x)}{|x|^{n+2s}} dx \ge 0$$

In addition, for any fixed $x \in \mathbb{R}^n$, if $\{\lambda_k x + a_k\}$ is bounded, we assume that $\lambda_k x + a_k \to \bar{x}$ by extracting a further subsequence. Therefore, the assumption (H3) implies

$$\lambda_{k}^{\frac{2sp}{p-1}} f(\lambda_{k}x + a_{k}, \lambda_{k}^{-\frac{2s}{p-1}} v_{k}(x))$$

$$= (v_{k}(x))^{p} \frac{f(\lambda_{k}x + a_{k}, \lambda_{k}^{-\frac{2s}{p-1}} v_{k}(x))}{\left(\lambda_{k}^{-\frac{2s}{p-1}} v_{k}(x)\right)^{p}}$$

$$\to K(\bar{x})v^{p}(x) \quad \text{as} \quad k \to \infty,$$

$$(4.14)$$

which is still valid in the case $|\lambda_k x + a_k| \to \infty$ with $|\bar{x}| = \infty$ by (H3).

Consequently, it follows from (4.13) and (4.14) that v is a solution of

$$(-\Delta)^s v(x) = K(\bar{x})v^p(x) + b, \ x \in \mathbb{R}^n$$
(4.15)

with $b \ge 0$.

To proceed with the proof, we show that b equals to 0. Let

$$F_R(x) = \begin{cases} K(\bar{x})v^p(x) + b, & x \in B_R(0) \\ 0, & x \in B_R^C(0) \end{cases}$$
$$v_R(x) = C_{n,s} \int_{\mathbb{R}^n} \frac{F_R(y)}{|x - y|^{n - 2s}} dy.$$

Then it is easy to see that

$$\begin{cases} (-\Delta)^s v_R(x) = F_R(x), & x \in \mathbb{R}^n \\ \lim_{|x| \to \infty} v_R(x) = 0. \end{cases}$$

Denote

$$w_R(x) = v(x) - v_R(x), \ x \in \mathbb{R}^n.$$

Then

$$\begin{cases} (-\Delta)^s w_R(x) \ge 0, & x \in \mathbb{R}^n, \\ \lim_{|x| \to \infty} w_R(x) \ge 0. \end{cases}$$

By the maximum principle, we have

$$w_R(x) \ge 0, \ x \in \mathbb{R}^n.$$

Now let $R \to \infty$, we arrive at

$$v(x) \ge \int_{\mathbb{R}^n} \frac{C_{n,s}}{|x-y|^{n-2s}} \left(K(\bar{x}) v^p(y) + b \right) dy, \ x \in \mathbb{R}^n$$

If b > 0, then the integral on the right hand side diverges and $v(x) = \infty$. This contradicts the boundedness of v. Therefore, we must have b = 0 and

$$(-\Delta)^s v(x) = K(\bar{x})v^p(x), \ x \in \mathbb{R}^n$$

By virtue of (4.12) and the maximum principle, one gets that

$$v(x) > 0, \ \forall x \in \mathbb{R}^n.$$

On the other hand, by the Liouville theorem in [10, Theorem 4], (4.15) has no positive solution if 1 . This is a contradiction. Therefore, (1.16) and (1.17) are true. Therefore, the proof of Theorem 1.8 is completed.

Proof of Theorem 1.10. In this case, Ω is an unbounded domain in \mathbb{R}^n , and u solves the quasi-linear equation

$$(-\Delta)^s u(x) = b(x) |\nabla u|^q(x) + f(x, u(x)), \ x \in \Omega$$

for $s > \frac{1}{2}$.

We show also that

$$u(x) + |\nabla u|^{\frac{2s}{2s+p-1}}(x) \le C\left(1 + dist^{-\frac{2s}{p-1}}(x,\partial\Omega)\right), \ x \in \Omega$$
(4.16)

by a contradiction argument.

If (4.16) is not valid, then there exist sequences of $\Omega_k \subset \mathbb{R}^n$, u_k , $x_k \in \Omega_k$ such that u_k solves

$$(-\Delta)^s u_k(x) = b(x) |\nabla u_k|^q(x) + f(x, u_k(x)), \quad x \in \Omega_k,$$

and

$$\frac{1}{2}M_k(x_k)d_k \ge \frac{1}{2}M_k(x_k)(1+d_k^{-1})^{-1} > k \to \infty \text{ as } k \to \infty,$$

where $d_k = dist(x_k, \partial \Omega_k)$ and

$$M_k(x) := u_k^{\frac{p-1}{2s}}(x) + |\nabla u_k|^{\frac{p-1}{2s+p-1}}(x).$$

Define

$$r_k := 2kM_k^{-1}(x_k).$$

It follows that $r_k < d_k$ and therefore $B_{r_k}(x_k) \subset \Omega_k$.

To start the rescaling procedure and obtain upper bounds for solutions, we consider the following function

$$S_k(x) = (M_k(x)(r_k - |x - x_k|))^{\frac{2s}{p-1}}, \ x \in B_{r_k}(x_k).$$

Then there exists a point $a_k \in B_{r_k}(x_k)$ such that

$$S_k(a_k) = \max_{B_{r_k}(x_k)} S_k(x)$$

It follows that

$$M_k(x) \le M_k(a_k) \frac{r_k - |a_k - x_k|}{r_k - |x - x_k|}, \ x \in B_{r_k}(x_k).$$

$$(4.17)$$

Denote

$$\lambda_k = (M_k(a_k))^{-1}.$$

Taking $x = x_k$ in (4.17) gives

$$\frac{M_k(x_k)}{M_k(a_k)} \le \frac{r_k - |a_k - x_k|}{r_k},\tag{4.18}$$

which implies that

 $M_k(x_k) \le M_k(a_k).$

It follows from the definition of r_k , λ_k and (4.18) that

$$2k\lambda_k \le r_k - |a_k - x_k|. \tag{4.19}$$

In addition, using (4.19) and by a similar argument as in deriving (4.6), one gets

$$|x_k - |a_k - x_k| \le 2(r_k - |x - x_k|), \ x \in B_{k\lambda_k}(a_k) \subset B_{r_k}(x_k).$$
(4.20)

Combining (4.17) with (4.20) yields

$$M_k(x) \le 2M_k(a_k), \ x \in B_{k\lambda_k}(a_k).$$

$$(4.21)$$

Rescale the solutions sequence as

$$v_k(x) = \frac{1}{(M_k(a_k))^{\frac{2s}{p-1}}} u_k(\lambda_k x + a_k), \ x \in B_k(0).$$

Then $v_k(x)$ satisfies the following equation

$$(-\Delta)^{s} v_{k}(x) = \lambda_{k}^{\frac{2sp-(2s+p-1)q}{p-1}} b(\lambda_{k}x + a_{k}) |\nabla v_{k}|^{q}(x) + \lambda_{k}^{\frac{2sp}{p-1}} f(\lambda_{k}x + a_{k}, \lambda_{k}^{-\frac{2s}{p-1}} v_{k}(x))$$

$$:= F_{k}(x, v_{k}(x), \nabla v_{k}(x)), \ x \in B_{k}(0).$$

$$(4.22)$$

Moreover, by the definition of λ_k and (4.21), one has

$$\left(v_k^{\frac{p-1}{2s}} + |\nabla v_k|^{\frac{p-1}{2s+p-1}}\right)(0) = \lambda_k M_k(a_k) = 1$$
(4.23)

and

$$\left(v_k^{\frac{p-1}{2s}} + |\nabla v_k|^{\frac{p-1}{2s+p-1}}\right)(x) \le 2, \ |x| \le k.$$
(4.24)

It follows from (4.24) that v_k and $|\nabla v_k|$ are uniformly bounded in $B_k(0)$. Then for k large enough, one can get

$$\lambda_k^{\frac{2sp-(2s+p-1)q}{p-1}} b(\lambda_k x + a_k) |\nabla v_k|^q(x) \le C_8, \ x \in B_k(0),$$

and

$$\lambda_k^{\frac{2sp}{p-1}} f(\lambda_k x + a_k, \lambda_k^{-\frac{2s}{p-1}} v_k(x))$$

$$\leq C_0 \lambda_k^{\frac{2sp}{p-1}} \left(1 + \lambda_k^{-\frac{2sp}{p-1}} (v_k(x))^p \right)$$

$$\leq C_0 \lambda_k^{\frac{2sp}{p-1}} + C_0 (v_k(x))^p$$

$$\leq C_9,$$

where one has used the assumptions (H1)-(H3) and the fact $0 < q < \frac{2sp}{2s+p-1}$. As a result, there exists a constant $C_{10} > 0$ independent of k such that

$$0 \le F_k(x, v_k(x), \nabla v_k(x)) \le C_{10}, \ x \in B_k(0).$$

It follows from Theorem 1.1 that there exists a positive constant C_{11} independent of k such that

$$\|v_k\|_{C^{[2s-\varepsilon]},\{2s-\varepsilon\}}(B_{3k/4}(0)) \le C_{11}$$

for all $0 < \varepsilon < 2s$. Therefore $F_k(x, v_k(x), \nabla v_k(x))$ in (4.22) is Hölder continuous for $x \in B_{3k/4}(0)$ due to $s > \frac{1}{2}$. By virtue of Theorem 1.3, there exist some positive constants $\beta \in (0, 1)$ and C_{12} which are independent of k such that

$$\|v_k\|_{C^{[2s+\beta]},\{2s+\beta\}}(B_{k/2}(0)) \le C_{12}.$$
(4.25)

Due to the above uniform regularity estimates, it follows from the Ascoli-Arzelá theorem that there exists a converging subsequence of $\{v_k\}$ (still denoted by $\{v_k\}$) that converges pointwisely in \mathbb{R}^n to a function v(x), and

$$v_k \to v$$
 in $C^1_{loc}(\mathbb{R}^n)$ as $k \to \infty$.

Taking limit in (4.23), we obtain

$$v^{\frac{p-1}{2s}}(0) + |\nabla v|^{\frac{p-1}{2s+p-1}}(0) = 1.$$
(4.26)

(4.25) and a similar argument as deriving (4.13) leads to

$$\lim_{k \to \infty} (-\Delta)^s v_k(x) = (-\Delta)^s v(x) - b, \ x \in \mathbb{R}^n$$
(4.27)

for some $b \ge 0$.

To derive a contradiction, one needs to analyze the limit equation satisfied by v.

Since $0 < q < \frac{2sp}{2s+p-1}$, v_k and $|\nabla v_k|$ are uniformly bounded in $B_k(0)$, then

$$\lambda_k^{\frac{2sp-(2s+p-1)q}{p-1}}b(\lambda_k x + a_k)|\nabla v_k|^q(x) \to 0 \text{ as } k \to \infty,$$

In addition, for any fixed $x \in \mathbb{R}^n$, if $\{\lambda_k x + a_k\}$ is bounded, we assume that $\lambda_k x + a_k \to \bar{x}$ by extracting a further subsequence. Therefore, the assumption (H3) implies

$$\lambda_k^{\frac{2sp}{p-1}} f(\lambda_k x + a_k, \lambda_k^{-\frac{2s}{p-1}} v_k(x))$$

$$= (v_k(x))^p \frac{f(\lambda_k x + a_k, \lambda_k^{-\frac{2s}{p-1}} v_k(x))}{\left(\lambda_k^{-\frac{2s}{p-1}} v_k(x)\right)^p}$$

$$\to K(\bar{x}) v^p(x) \quad \text{as} \quad k \to \infty,$$

$$(4.28)$$

which is still valid in the case $|\lambda_k x + a_k| \to \infty$ with $|\bar{x}| = \infty$ by (H3).

Consequently, combining (4.27) and (4.28) shows that v is a solution of

$$(-\Delta)^s v(x) = K(\bar{x})v^p(x) + b, \ x \in \mathbb{R}^n.$$

Similar to the previous argument, one can show that b = 0, and hence

$$(-\Delta)^{s}v(x) = K(\bar{x})v^{p}(x), \ x \in \mathbb{R}^{n}.$$
(4.29)

By virtue of (4.26) and the maximum principle, one can deduce that

$$v(x) > 0, \ \forall x \in \mathbb{R}^n$$

On the other hand, by the Liouville theorem in [10, Theorem 4], (4.29) has no positive solution. This is a contradiction. Therefore, (4.16) holds and thus the proof of Theorem 1.10 is completed.

Acknowledgements. This research is supported in part by the Zhang Ge Ru Foundation, Hong Kong RGC Earmarked Research Grants CUHK-14301421, CUHK-14300917, CUHK-14302819 and CUHK-14300819. Z. Xin is also supported by the key project of NSFC (Grants No. 12131010 and 11931013). C. Li is partially supported by NSFC (Grant No. 12031012). W. Chen is partially supported by MPS Simons Foundation 847690 and NSFC (Grant No. 12071229). L. Wu is partially supported by NSFC (Grant No. 12401133).

Date availability statement: Data will be made available on reasonable request.

Conflict of interest statement: There is no conflict of interest.

References

- J. Bertoin, Lévy Processes, Cambridge Tracts in Mathematics, 121 Cambridge University Press, Cambridge, 1996.
- [2] J. P. Bouchard, A. Georges, Anomalous diffusion in disordered media, Statistical mechanics, models and physical applications, Physics reports, 195 (1990).
- [3] C. Brandle, E. Colorado, A. de Pablo, U. Sanchez, A concave-convex elliptic problem involving the fractional Laplacian, Proc. Roy. Soc. Edinburgh Sect. A, 143 (2013), 39–71.
- [4] C. Bucur, Some observations on the Green function for the ball in the fractional Laplace framework, Commun. Pure Appl. Anal., 15 (2) (2016), 657-699.
- [5] L. Caffarelli, L. Silvestre, An extension problem related to the fractional Laplacian, Comm. Partial Differential Equations, 32 (7-9) (2007), 1245-1260.
- [6] L. Caffarelli, L. Silvestre, Regularity theory for fully nonlinear integro-differential equations, Comm. Pure Appl. Math., 62 (2009), 597-638.
- [7] L. Caffarelli, L. Silvestre, Regularity results for nonlocal equations by approximation, Arch. Ration. Mech. Anal., 200 (2011), 59-88.
- [8] L. Caffarelli, L. Silvestre, The Evans-Krylov theorem for nonlocal fully nonlinear equations, Ann. of Math., 174 (2) (2011), 1163-1187.
- [9] L. Caffarelli, L. Vasseur, Drift diffusion equations with fractional diffusion and the quasigeostrophic equation, Ann. Math., 3 (2010), 1903-1930.
- [10] W. Chen, C. Li, Y. Li, A direct method of moving planes for the fractional Laplacian, Adv. Math., 308 (2017), 404-437.
- [11] W. Chen, Y. Li, P. Ma, The fractional Laplacian, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2020.
- [12] W. Chen, Y. Li, R. Zhang, A direct method of moving spheres on fractional order equations, J. Funct. Anal., 272(2017) 4133-4157.
- [13] W. Chen, L. Wu, Liouville theorems for fractional parabolic equations, Adv. Nonlinear Stud., 21 (4) (2021), 939-958.
- [14] W. Chen, L. Wu, Uniform a priori estimates for solutions of higher critical order fractional equations. Calc. Var. & PDEs, 60 (2021), no. 3, Paper No. 102.

- [15] P. Constantin, Euler equations, Navier-Stokes equations and turbulence, in Mathematical Foundation of Turbulent Viscous Flows, Vol. 1871 of Lecture Notes in Math. Springer, Berlin, 2006.
- [16] W. Dai, G. Qin, Classification of nonnegative classical solutions to third-order equations, Adv. Math., 328 (2018), 822-857.
- [17] W. Dai, G. Qin, Liouville type theorems for fractional and higher order Hénon-Hardy type equations via the method of scaling spheres, arXiv:1810.02752, 2018.
- [18] S. Dipierro, N. Soave, E. Valdinoci, On fractional elliptic equations in Lipschitz sets and epigraphs: regularity, monotonicity and rigidity results, Math. Ann., 369 (3-4), (2017), 1283-1326.
- [19] X. Du, T. Jin, J. Xiong, H. Yang, Blow up limits of the fractional Laplacian and their applications to the fractional Nirenberg problem, Proc. Amer. Math. Soc., 151 (11) (2023), 4693-4701.
- [20] T. Jin, Y. Li, J. Xiong, On a fractional Nirenberg problem, part I: blow up analysis and compactness of solutions, J. Eur. Math. Soc. (JEMS), 16 (6) (2014), 1111-1171.
- [21] C. Li, C. Liu, Z. Wu, H. Xu, Non-negative solutions to fractional Laplace equations with isolated singularity, Adv. Math., 373 (2020), 107329, 38 pp.
- [22] C. Li, L. Wu, Pointwise regularity for fractional equations, J. Differential Equations, 302 (2021), 1-36.
- [23] Y. Li, J. Bao, Fractional Hardy-Hénon equations on exterior domains, J. Differential Equations, 266 (2-3) (2019), 1153-1175.
- [24] D. Gilbarg, N. S. Trudinger, Elliptic partial differential equations of second order, Second edition. 224. Springer-Verlag, Berlin, 1983. xiii+513 pp.
- [25] P. Poláčik, P. Quittner, P. Souplet, Singularity and decay estimates in superlinear problems via Liouville-type theorems. I. Elliptic equations and systems, Duke Math. J., 139 (3) (2007), 555-579.
- [26] L. Silvestre, Regularity of the obstacle problem for a fractional power of the Laplace operator, Comm. Pure Appl. Math., 60 (1) (2007), 67-112.
- [27] X. Ros-Oton, J. Serra, Regularity theory for general stable operators, J. Differential Equations, 260 (12) (2016), 8675-8715.
- [28] L. Wu, Sliding methods for the higher order fractional Laplacians, Fract. Calc. Appl. Anal., 24 (3) (2021), 923-949.
- [29] L. Wu, W. Chen, The sliding methods for the fractional p-Laplacian, Adv. Math., 361 (2020), 106933, 26 pp.
- [30] L. Wu, W. Chen, Radial Symmetry of Ancient solutions to nonlocal parabolic equations, Adv. Math, 408(2022) 108607.
- [31] V. Tarasov, G. Zaslasvky, Fractional dynamics of systems with long range interaction, Comm. Nonl. Sci. Numer. Simul., 11 (2006), 885-889.
- [32] R. Zhuo, C. Li, Classification of anti-symmetric solutions to nonlinear fractional Laplace equations, Calc. Var. Partial Differential Equations, 61 (1), 2022, Paper No. 17, 23pp.

WENXIONG CHEN DEPARTMENT OF MATHEMATICAL SCIENCES YESHIVA UNIVERSITY NEW YORK, NY, 10033, USA *Email address*: wchen@yu.edu

CONGMING LI SCHOOL OF MATHEMATICAL SCIENCES CMA-SHANGHAI, SHANGHAI JIAO TONG UNIVERSITY SHANGHAI, 200240, P. R. CHINA *Email address*: congming.li@sjtu.edu.cn

LEYUN WU SCHOOL OF MATHEMATICS SOUTH CHINA UNIVERSITY OF TECHNOLOGY GUANGZHOU, 510640, P. R. CHINA AND THE INSTITUTE OF MATHMATICAL SCIENCES & DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, N.T., HONG KONG, P. R. CHINA *Email address*: leyunwu@scut.edu.cn

ZHOUPING XIN THE INSTITUTE OF MATHMATICAL SCIENCES & DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, N.T., HONG KONG, P. R. CHINA *Email address*: zpxin@ims.cuhk.edu.hk