# Multivalued forbidden numbers of two-rowed configurations – the missing cases

Wallace Peaslee<sup>\*</sup> University of Cambridge Department of Applied Mathematics and Theoretical Physics Cambridge, United Kingdom

Attila Sali<sup>†</sup> HUN-REN Alfréd Rényi Institute of Mathematics and Department of Computer Science, BUTE Budapest, Hungary

Jun Yan<sup>‡</sup>

University of Warwick Mathematics Institute Coventry, United Kingdom

#### Abstract

The present paper considers extremal combinatorics questions in the language of matrices. An s-matrix is a matrix with entries in  $\{0, 1, \ldots, s-1\}$ . An s-matrix is simple if it has no repeated columns. A matrix F is a configuration in a matrix A, denoted  $F \prec A$ , if it is a row/column permutation of a submatrix of A. Avoid(m, s, F) is the set of m-rowed, simple s-matrices not containing a configuration of F and forb $(m, s, F) = \max\{|A|: A \in \text{Avoid}(m, s, F)\}$ . Dillon and Sali initiated the systematic study of forb(m, s, F) for 2-matrices F, and computed forb(m, s, F) for all 2-rowed F when s > 3. In this paper we tackle the remaining cases when s = 3. In particular, we determine the asymptotics of forb $(m, 3, p \cdot K_2) - \text{forb}(m, 3, p \cdot I_2)$  for p > 3, where  $K_2$  is the  $2 \times 4$  simple 2-matrix and  $I_2$  is the  $2 \times 2$  identity matrix, as well as the exact values of forb(m, 3, F) for many 2-rowed 2-matrices F.

Keywords: Forbidden configurations, (0,1)- matrices, s-matrices, extremal set theory

# 1 Introduction

Several questions of extremal combinatorics can be conveniently expressed in the language of (0, 1)matrices. An matrix A is said to be *simple* if it contains no repeated columns. There is a natural correspondence between columns of an *m*-rowed (0, 1)-matrix and subsets of [m]. We consider an extremal set theory problem in matrix terminology as follows. Let |A| be the number of columns of A. For a given matrix F, we say F is a *configuration* in A, denoted  $F \prec A$ , if there is a submatrix

<sup>\*</sup>Supported by a UK Engineering and Physical Sciences Research Council Doctoral Training Partnership grant.

<sup>&</sup>lt;sup>†</sup>Research was partially supported by the National Research, Development and Innovation Office (NKFIH) grants K–132696 and SNN-135643.

<sup>&</sup>lt;sup> $\ddagger$ </sup>Supported by the Warwick Mathematics Institute Centre for Doctoral Training, and by funding from the UK EPSRC (Grant number: EP/W523793/1).

of A which can be obtained by permuting the rows and columns of F. Define

Avoid $(m, F) = \{A \colon A \text{ is an } m \text{-rowed } (0, 1) \text{-matrix and } F \not\prec A\},\$ 

$$forb(m, F) = \max\{|A| : A \in Avoid(m, F)\}$$

We say a matrix  $A \in Avoid(m, F)$  is *extremal* if |A| = forb(m, F), and let

$$\operatorname{ext}(m, F) = \{A \in \operatorname{Avoid}(m, F) \colon |A| = \operatorname{forb}(m, F)\}.$$

The foremost theorem in forbidden configurations is Theorem 1.1 below. Here,  $K_k$  denotes the  $k \times 2^k$  matrix containing all possible 0, 1-columns.

**Theorem 1.1.** [Sau72, She72, VC71]

forb
$$(m, K_k) = \binom{m}{k-1} + \binom{m}{k-2} + \dots + \binom{m}{0}.$$

Further interesting results and research problems can be found in the survey paper [Ans13]. However, our main interest in the present paper is the generalization of these concepts to s-matrices, which have entries in  $\{0, 1, \ldots, s-1\}$ . Such matrices can be thought of as s-coloured set systems or as representations of collections of functions from a given finite set into  $\{0, 1, \ldots, s-1\}$ . The concepts of simple matrices and configurations naturally extend. For a given collection  $\mathcal{F}$  of matrices, we denote by Avoid $(m, s, \mathcal{F})$  the collection of m-rowed, simple s-matrices that avoid every matrix  $F \in \mathcal{F}$ . The main extremal function in the study of forbidden configurations is

$$forb(m, s, \mathcal{F}) = \max\{|A| \colon A \in Avoid(m, s, \mathcal{F})\},\$$

which we call the *forbidden number*, and we similarly define

$$ext(m, s, F) = \{A \in Avoid(m, s, F) \colon |A| = forb(m, s, F)\}.$$

Alon [Alo83] and Steele [Ste78] gave a generalization of Theorem 1.1 to s-matrices, with the bound given being an exponential function of m. This is a special case of the following more general phenomenon proved by Füredi and Sali [FS12]. A matrix is called an (i, j)-matrix if its entries are all from the set  $\{i, j\}$ .

**Theorem 1.2.** Let  $\mathcal{F}$  be a family of s-matrices. If for every pair of distinct  $i, j \in \{0, 1, \ldots, s-1\}$  there is an (i, j)-matrix in  $\mathcal{F}$ , then forb $(m, s, \mathcal{F}) = O(m^k)$  for some positive integer k. If  $\mathcal{F}$  has no (i, j)-matrix for some distinct  $i, j \in \{0, 1, \ldots, s-1\}$ , then forb $(m, s, \mathcal{F}) = \Omega(2^m)$ .

Several papers [ADLS23, AL14, ELS20] considered special sets of forbidden configurations for which the forbidden numbers are of polynomial magnitudes. The systematic investigation of exact exponential bounds when forbidding (0, 1)-configurations was started in [DS21]. They provided exact formulas for the forbidden numbers forb(m, s, F) of many (0, 1)-matrices F, including all 2-rowed matrices when s > 3. Along the way, they exposed some interesting qualitative differences between the cases s = 2, s = 3, and s > 3.

Some important matrices include  $I_k$ , the  $k \times k$  identity matrix and  $K_k$ , the  $k \times 2^k$  matrix of all possible (0,1)-columns on k rows. We use the following notation for general 2-rowed configurations:

$$F(a, b, c, d) = \begin{bmatrix} a & b & c & d \\ 00 \cdots 0 & 11 \cdots 1 & 00 \cdots 0 & 11 \cdots 1 \\ 00 \cdots 0 & 00 \cdots 0 & 11 \cdots 1 & 11 \cdots 1 \end{bmatrix}$$

For example,  $F(1,1,1,1) = K_2$ . Let  $p \cdot F$  denote the concatenation of p copies of F, so that in particular  $p \cdot K_2$  and F(p, p, p, p) are the same as configurations, while  $p \cdot I_2$  is the same as F(0, p, p, 0). When s > 3, the forbidden numbers forb(m, s, F(a, b, c, d)) for all 2-rowed configurations were determined exactly in [DS21]. Under mild conditions similar to Lemma 1.3 below, it was shown that forb(m, s, F(a, b, c, d)) = forb(m, s, F(a, p, p, d)), where  $p = \min\{b, c\}$ . In particular, it turned out that forb $(m, s, p \cdot K_2) = \text{forb}(m, s, p \cdot I_2)$ , which implies that forb $(m, s, p \cdot K_2) = \text{forb}(m, s, F(a, p, p, d))$  if s > 3 and  $a, d \leq p$ .

For general 2-rowed configurations F(a, b, c, d), the case when s = 3 is harder than when s > 3. In particular, it was shown in [DS21] that  $\operatorname{forb}(m, 3, p \cdot K_2) > \operatorname{forb}(m, 3, p \cdot I_2)$  if  $p \ge 3$ , exhibiting a different behaviour to the s > 3 case. More generally,  $\operatorname{forb}(m, 3, F(a, b, c, d))$  were determined in [DS21] only when  $\max\{a, d\} \ge \min\{b, c\}$ . Our goal in this paper is to study the missing cases when  $\max\{a, d\} < \min\{b, c\}$ .

The following important reduction lemma was proved in [DS21].

**Lemma 1.3.** [DS21] Let  $p = \min\{b, c\}$ . If  $2^{m-2} \ge (\max\{a, b, c, d\} - 1)m^2$  and  $p \ge 1$ , then

forb(m, 3, F(a, b, c, d)) = forb(m, 3, F(a, p, p, d)).

Note that by Lemma 1.3, we may assume without loss of generality that b = c = p provided that m is large. It was also proved in [DS21] that  $\operatorname{forb}(m, 3, F(p, p, p, p)) = \operatorname{forb}(m, 3, p \cdot K_2) = 2^m + m2^{m-1} + (p-1)\binom{m}{2}$ . In what follows, we investigate the forbidden numbers  $\operatorname{forb}(m, 3, F(a, p, p, d))$  for various parameters a, d < p.

In Section 2, we begin by introducing some terminologies and notations. Then, we prove first a generalization of a lemma in [DS21], which roughly says that any matrix in Avoid(m, 3, F(a, p, p, d)) that contains many columns must satisfy some stability-type structural properties. In Section 2.1, we determine the difference between forb $(m, 3, p \cdot K_2)$  and forb $(m, 3, p \cdot I_2)$  asymptotically. Throughout the paper we let  $r = \lceil \log_2(p-1) \rceil$ , so  $2^{r-1} < p-1 \le 2^r$ . Our first, maybe surprising, observation is that the difference above does not depend on m.

**Proposition 1.4.** For all  $p \ge 2$ , forb $(m, 3, p \cdot K_2)$  – forb $(m, 3, p \cdot I_2)$  is independent of m as long as  $m \ge 2r + 2$ .

This allows us to define  $g_p$  to be the common value of  $\operatorname{forb}(m, 3, p \cdot K_2) - \operatorname{forb}(m, 3, p \cdot I_2)$  for  $m \geq 2r + 2$ , and we prove the following asymptotic result for  $g_p$ .

## Theorem 1.5.

$$g_p \sim \frac{1}{2} p(\log_2 p)^2.$$

In Section 2.2, we compute forb(m, 3, F(a, p, p, d)) for all  $p \leq 4$  and  $max\{a, d\} < p$ , which all turn out to be equal to  $forb(m, 3, p \cdot I_2)$ .

This no longer holds for larger values of p, but in Section 2.3 we have further exact results for many cases. Let

$$N_r = \binom{\left\lceil \frac{r}{2} \right\rceil + 1}{2} + \binom{\left\lfloor \frac{r}{2} \right\rfloor + 1}{2}$$

The next theorem determines forb(m, 3, F(p-q, p, p, p-q)) for a wide range of values p and q.

**Theorem 1.6.** For all  $m \ge 2N_r + 2$  and  $0 \le q \le p - 1$ ,

$$forb(m, 3, F(p-q, p, p, p-q)) = forb(m, 3, p \cdot K_2) - qN_r$$

if one of the following holds

- $p \ge 2^{r-1} + 2q + 1$ ,
- $p \leq 2^{r-1} + 2q$ ,  $qN_r \leq 2^{r-3}$  and  $\frac{1}{2}(\lfloor \frac{r}{2} \rfloor + 1)(p 1 2^{r-1}) \geq q$ .

In particular, this is true for all fixed constants q and all p sufficiently large.

With a more careful analysis of the proof of Theorem 1.6, we prove the following result.

Corollary 1.7. For all  $m \ge 2N_r + 2$  and  $p \ge 2$ ,

 $forb(m, 3, F(p-1, p, p, p-1)) = forb(m, 3, p \cdot K_2) - N_r.$ 

Determining the exact forbidden number in the remaining cases when q is relatively large might be difficult, given that the exact value of forb $(m, 3, p \cdot I_2)$  is not known. In particular, the following result shows that the exact formula in Theorem 1.6 may not hold if those conditions are not satisfied.

**Proposition 1.8.** If  $m \ge 2N_r + 2$ ,  $0 \le q \le p - 1$  and  $d := q - (\lceil \frac{r}{2} \rceil + 1)(p - 1 - 2^{r-1}) > 0$ , then

$$\operatorname{forb}(m, 3, F(p-q, p, p, p-q)) \ge \operatorname{forb}(m, 3, p \cdot K_2) - qN_r + \left\lfloor \frac{r}{2} \right\rfloor d.$$

We also have the following more general lower bound.

**Proposition 1.9.** For every  $0 \le q_0, q_1 \le p-1$ , every  $0 \le r_1, r_2 \le r$  satisfying  $r_1 + r_2 = r$ , and every  $m \ge 2\binom{r_1+1}{2} + 2\binom{r_2+1}{2} + 2$ , we have

forb
$$(m, 3, F(p - q_0, p, p, p - q_1)) \ge$$
 forb $(m, 3, p \cdot K_2) - \min\{q_0, q_1\} \binom{r_1 + 1}{2} - \max\{q_0, q_1\} \binom{r_2 + 1}{2}$ 

For fixed  $p, q_0, q_1$ , we may optimize the right-hand side by viewing  $r_1$  as the variable and using  $r_2 = r - r_1$ . That is, assuming  $q_0 \leq q_1$ , we minimize  $q_0 \binom{r_1+1}{2} + q_1 \binom{r-r_1+1}{2}$ . Easy calculation shows that if  $q_1 = \alpha q_0$  for some  $\alpha \geq 1$ , then the minimum is achieved when  $r_1$  is the closest integer to  $\frac{(2r+1)\alpha-1}{2(\alpha+1)}$ , as long as this fraction is at most r, which is equivalent to  $\alpha \leq 2r+1$ . Otherwise,  $r_1 = r$  is optimal. Note that if  $q_0 = q_1$  and so  $\alpha = 1$ , then the minimum is achieved when  $r_1 = \lceil \frac{r}{2} \rceil, r_2 = \lfloor \frac{r}{2} \rfloor$ . The resulting lower bound in this case is tight in many cases by Theorem 1.6. We suspect that a more involved version of this proof could potentially show that after optimization, the general lower bound for forb $(m, 3, F(p - q_0, p, p, p - q_1))$  given by Proposition 1.9 is also tight in several cases.

# 2 Results

Recall that from Lemma 1.3, we may assume that b = c, as long as m is sufficiently large. From now on, we will assume  $b = c = p \ge 3$ ,  $0 \le a, d \le p - 1$  unless stated otherwise. Generalizing some ideas from [DS21], we now make some definitions to help better understand the structure of matrices in Avoid(m, 3, F(a, p, p, d)).

**Definition 2.1.** Let  $A \in \operatorname{Avoid}(m, 3, F(a, p, p, d))$ . To each pair (i, j) with  $1 \le i < j \le m$ , associate a vector  $\begin{bmatrix} x \\ y \end{bmatrix} \in \left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$  so that if  $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$  or  $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$  then rows i, j contain at most p-1 copies of  $\begin{bmatrix} x \\ y \end{bmatrix}, if \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$  then rows i, j contain at most a-1 copies of  $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$  and if  $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$  then rows i, j contain at most d-1 copies of  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ . Such a vector must exist as  $F(a, p, p, d) \not\prec A$ , and if multiple choices are possible, we pick one arbitrarily. In all cases, we say a column v of A has a mark at the pair (i, j) if  $v_i = x$  and  $v_j = y$ .

Let B be the matrix consisting of all columns of A with no mark, and let C be the matrix consisting of all columns of A with at least one mark. We call  $A = \begin{bmatrix} B \mid C \end{bmatrix}$  the standard decomposition of A.

We also associate to A a directed graph T on [m] as follows: for each pair  $1 \le i < j \le m$ , if  $\begin{bmatrix} 0\\1 \end{bmatrix}$  is associated with (i, j), add the directed edge ij; if  $\begin{bmatrix} 1\\0 \end{bmatrix}$  is associated with (i, j), add the directed edge ij.

*ji*; otherwise do nothing. Let N denote the number of pairs (i, j) that are associated with either  $\begin{bmatrix} 0\\0 \end{bmatrix}$ 

or  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ . We say that these pairs form 0-non-edges and 1-non-edges in T, respectively.

Using the notations above, we have some immediate observations. It follows from the definition that B is in Avoid $(m, 3, K_2)$  so  $|B| \leq 2^m + m2^{m-1}$ . We also have  $|C| \leq (p-1)\binom{m}{2} - N + N(\max\{a, d\} - 1) = (p-1)\binom{m}{2} - N(p - \max\{a, d\})$ , since every column in C contains at least one mark.

The following key lemma, generalisaing one from [DS21], shows that if |B| is large, then the associated directed graph T is transitive.

# **Lemma 2.2.** If $|B| > 2^m + m2^{m-1} - 2^{m-3}$ , then T is transitive.

*Proof.* Assume T is not transitive, then there exists a triple (i, j, k) of distinct vertices such that ij, jk are edges of T but either ki is an edge of T or there is no edge between k and i. For a column v in B, define its support to be the set of row indices i such that  $v_i$  is either 0 or 1. For any set  $X \subset [m]$  of rows of B, let B(X) be the matrix consisting of columns of B with support X. Then its restriction  $B(X)|_X$  of B(X) to the rows in X is simple and is in Avoid $(|X|, 2, K_2)$ . Thus, if  $\{i, j, k\} \not\subset X$ , then  $|B(X)| = |B(X)|_X| \leq \operatorname{forb}(|X|, 2, K_2) = |X| + 1$  by Theorem 1.1.

If, however,  $\{i, j, k\} \subset X$ , then consider  $B(X)|_{\{i, j, k\}}$ . If ij, jk, ki are all edges of T, the only

possible columns in  $B(X)|_{\{i,j,k\}}$  are  $\begin{bmatrix} 0\\0\\0 \end{bmatrix}$  and  $\begin{bmatrix} 1\\1\\1 \end{bmatrix}$ . So  $B(X)|_{X\setminus\{i,j\}}$  is simple and avoids  $K_2$ . Thus  $|B(X)| = |B(X)|_{X\setminus\{i,j\}}| \leq \operatorname{forb}(|X|-2,2,K_2) = |X|-1$ . If ij,jk are edges of T but there is no edge between i and k, there are two cases. If rows i,k are associated with  $\begin{bmatrix} 0\\0 \end{bmatrix}$ , then any column in  $B(X)|_{\{i,j,k\}}$  has a 1 in entry i so  $B(X)|_{X\setminus\{i\}}$  is simple and avoids  $K_2$ . If rows i,k are associated with  $\begin{bmatrix} 1\\1 \end{bmatrix}$ , then any column in  $B(X)|_{\{i,j,k\}}$  has a 0 in entry k so  $B(X)|_{X\setminus\{k\}}$  is simple and avoids  $K_3$ .

 $K_2$ . In both cases, we have  $|B(X)| \le \text{forb}(|X| - 1, 2, K_2) = |X|$ .

Therefore, if ij, jk, ki are all edges of T, by counting columns in B according to their support, we have

$$\begin{split} |B| &\leq \sum_{\substack{X \subset [m] \\ \{i,j,k\} \not \in X}} (|X|+1) + \sum_{\substack{X \subset [m] \\ \{i,j,k\} \subset X}} (|X|-1) \\ &= \sum_{X \subset [m]} (|X|+1) - 2 \sum_{\substack{X \subset [m] \\ \{i,j,k\} \subset X}} 1 \\ &= \sum_{j=0}^m \binom{m}{j} (j+1) - 2 \sum_{j=0}^{m-3} \binom{m-3}{j} = 2^m + m2^{m-1} - 2^{m-2} \end{split}$$

A similar calculation shows that  $|B| \leq 2^m + m2^{m-1} - 2^{m-3}$  if ij, jk are edges of T but neither ik nor ki is. Both contradict the assumption that  $|B| > 2^m + m2^{m-1} - 2^{m-3}$ , which completes the proof.

Note that if  $|A| > 2^m + m2^{m-1} - 2^{m-3} + (p-1)\binom{m}{2}$ , then as  $|C| \le (p-1)\binom{m}{2}$ , we must have  $|B| > 2^m + m2^{m-1} - 2^{m-3}$ . Therefore, by Lemma 2.2, T is transitive, and hence the directed comparability graph of a finite poset P. Thus, by taking a linear extension of P, we can rearrange

the rows of A so that no pair  $1 \le i < j \le m$  is associated with  $\begin{bmatrix} 1\\ 0 \end{bmatrix}$ . In particular, unless (i, j) is associated with  $\begin{bmatrix} 0\\ 0 \end{bmatrix}$  or  $\begin{bmatrix} 1\\ 1 \end{bmatrix}$ , there is no occurrence of 0 above 1 in rows i, j in B.

# **2.1** Asymptotics of $\operatorname{forb}(m, 3, p \cdot K_2) - \operatorname{forb}(m, 3, p \cdot I_2)$

By [DS21], if  $2^{m-2} \ge p-1$ , then forb $(m, 3, p \cdot K_2) = 2^m + m2^{m-1} + (p-1)\binom{m}{2}$ , but the exact value of forb $(m, 3, p \cdot I_2)$  is not known for general p. In this subsection, we determine asymptotically the difference between forb $(m, 3, p \cdot K_2)$  and forb $(m, 3, p \cdot I_2)$ .

We first give a lower bound construction that uses only 0-mark and 1-mark columns.

Lemma 2.3. forb $(m, 3, p \cdot I_2) \ge 2^m + m2^{m-1} + (p-1)\left(\binom{m}{2} - \binom{r+1}{2}\right) + (r-1)2^r + 1.$ 

*Proof.* Let every pair i < j of rows be associated with  $\begin{bmatrix} 0\\1 \end{bmatrix}$ , so a mark in a column corresponds exactly to an instance of 0 above 1.

If a column v contains a unique mark at (i, j), then  $v_{\ell} \in \{1, 2\}$  for all  $1 \leq \ell < i$ ,  $v_{\ell} = 2$  for all  $i < \ell < j$ , and  $v_{\ell} \in \{0, 2\}$  for  $j < \ell \leq m$ . Thus, there are exactly  $2^{m-1+i-j}$  such columns, and we can include  $\min\{p-1, 2^{m-1+i-j}\}$  of them. Let  $r = \lceil \log_2(p-1) \rceil$ , then  $2^{m-1+i-j} < p-1$  if and only if  $i \in [r]$  and  $m-r+i \leq j \leq m$ . Therefore, the number of 1-mark columns we can include is

$$(p-1)\left(\binom{m}{2} - \binom{r+1}{2}\right) + \sum_{i=1}^{r} \sum_{j=m-r+i}^{m} 2^{m-1+i-j} = (p-1)\left(\binom{m}{2} - \binom{r+1}{2}\right) + (r-1)2^r + 1.$$

Since there are  $2^m + m2^{m-1}$  columns with no 0 above 1, and we can always include all of them, the result follows.

If  $A \in \operatorname{ext}(m, 3, p \cdot I_2)$ , let  $A = \begin{bmatrix} B \mid C \end{bmatrix}$  be the standard decomposition. Since  $|A| \ge 2^m + m2^{m-1} + (p-1)\left(\binom{m}{2} - \binom{r+1}{2}\right) + (r-1)2^r + 1$  by Lemma 2.3, and  $|C| \le (p-1)\binom{m}{2}$  from definition, we have that  $|B| > 2^m + m2^{m-1} - 2^{m-3}$  as long as  $m \ge 2r + 2$ . Hence, by Lemma 2.2, we can suitably permute the rows of A so that every pair i < j of rows is associated with  $\begin{bmatrix} 0\\1 \end{bmatrix}$ . Note that in this case the directed graph T is actually a tournament as there are no non-edges.

Motivated by Lemma 2.3 above, we say a pair of rows i < j is abundant if there are at least p-1 possible columns with exactly one mark at (i, j), or equivalently if  $2^{m-1+i-j} \ge p-1$ . A pair of rows i < j is called *scarce* if it is not abundant. It follows that if  $r = \lceil \log_2 (p-1) \rceil$ , then there are  $\binom{r+1}{2}$  scarce pairs of rows, which are exactly (i, j) for  $i \in [r]$  and  $m - r + i \le j \le m$ . The next lemma shows that we can always assume the maximum number of 1-mark columns are present in A.

**Lemma 2.4.** There exists  $A \in ext(m, 3, p \cdot I_2)$  containing  $2^m + m2^{m-1}$  0-mark columns and  $(p-1)\left\binom{m}{2} - \binom{r+1}{2}\right) + (r-1)2^r + 1$  1-mark columns. More precisely, for every abundant pair (i, j) of rows, A contains exactly p-1 columns with their only mark at (i, j), and for every scarce pair (i, j) of rows, A contains all columns with their only mark at (i, j).

Proof. We can always take all  $2^m + m2^{m-1}$  columns with no mark. Let  $A \in \text{ext}(m, 3, p \cdot I_2)$  maximize the number of 1-mark columns. Assume for some pair i < j of rows, A does not contain  $\min\{2^{m-1+i-j}, p-1\}$  columns with their unique marks at (i, j). Then there exists a column v with its only mark at (i, j) that is not in A. There are two cases. If A contains less than  $\min\{2^{m-1+i-j}, p-1\}$  marks at (i, j), [A | v] is then in Avoid $(m, 3, p \cdot I_2)$  with more columns than A, contradiction. If A contains exactly  $\min\{2^{m-1+i-j}, p-1\}$  marks at (i, j), then A contains a column u with at least two marks including a mark at (i, j). Remove column u and add column v to get a new matrix A'. Then A' is in  $\exp(m, 3, p \cdot I_2)$  with one more 1-mark column than A, contradiction. Thus, for all pair i < j

of rows, A contains  $\min\{2^{m-1+i-j}, p-1\}$  columns with their unique marks at (i, j), which add up to  $(p-1)\left(\binom{m}{2} - \binom{r+1}{2}\right) + (r-1)2^r + 1$  1-mark columns in total.

We can now prove Proposition 1.4, which shows that  $forb(m, 3, p \cdot K_2) - forb(m, 3, p \cdot I_2)$  is the same constant for all  $m \ge 2r + 2$ .

Proof of Proposition 1.4. Since forb $(m, 3, p \cdot K_2) = 2^m + m2^{m-1} + (p-1)\binom{m}{2}$  by [DS21], it suffices to show that for all  $m \ge 2r+2$ 

$$forb(m+1,3,p \cdot I_2) - forb(m,3,p \cdot I_2) = forb(m+1,3,p \cdot K_2) - forb(m,3,p \cdot K_2)$$
$$= 2^{m-1}(m+4) + (p-1)m.$$

First we prove that  $\operatorname{forb}(m+1,3,p \cdot I_2) - \operatorname{forb}(m,3,p \cdot I_2) \geq 2^{m-1}(m+4) + (p-1)m$ . Let  $A \in \operatorname{ext}(m,3,p \cdot I_2)$ . From above and by Lemma 2.2 we can assume A contains at most p-1 instances of 0 above 1 in each pair i < j of rows. A must contain all  $2^m + m2^{m-1}$  columns with no mark. Let S denote the number of columns in A with at least one mark.

Consider the following matrix A' with m + 1 rows. First take all  $2^{m+1} + (m + 1)2^m$  columns with no 0 above 1. Then, fix an index  $r + 1 \leq k \leq m + 1 - r$ , which exists as  $m \geq 2r + 2$ . For every column v in A with at least one mark, define a new column v' with m + 1 rows by  $v'_i = v_i$ if i < k,  $v'_k = 2$  and  $v'_i = v_{i-1}$  if  $k < i \leq m + 1$ . The new columns v' obtained in this way are all distinct and have marks only on row set  $\{1, 2, \ldots, k - 1, k + 1, \ldots m + 1\}$  corresponding to the marks in the original columns. Add them all to A'. Moreover, for each  $1 \leq j < k$ , the pair (j, k)is abundant so we can find p - 1 columns with their unique marks at (j, k). Similarly, for each  $k < j \leq m + 1$ , the pair (k, j) is abundant so we can find p - 1 columns with their unique marks at (k, j). Add all such (p - 1)m columns to A'. Then for any two rows i < j of A', there are at most p - 1 columns with 0 above 1 by construction, so  $A' \in \text{Avoid}(m + 1, 3, p \cdot I_2)$ , which implies forb $(m + 1, 3, p \cdot I_2) \geq |A'| = 2^{m+1} + (m + 1)2^m + S + (p - 1)m$ . Therefore,

forb
$$(m + 1, 3, p \cdot I_2)$$
 - forb $(m, 3, p \cdot I_2)$   

$$\geq (2^{m+1} + (m+1)2^m + S + (p-1)m) - (2^m + m2^{m-1} + S)$$

$$= 2^{m-1}(m+4) + (p-1)m.$$

Now we prove that  $\operatorname{forb}(m+1,3,p\cdot I_2) - \operatorname{forb}(m,3,p\cdot I_2) \leq 2^{m-1}(m+4) + (p-1)m$ . Let  $A \in \operatorname{ext}(m+1,3,p\cdot I_2)$ . We can assume that A has all  $2^{m+1} + (m+1)2^m$  columns with no mark. By Lemma 2.4, we can also assume A has the maximum number  $(p-1)\left\binom{m+1}{2} - \binom{r+1}{2} + (r-1)2^r + 1$  of 1-mark columns.

Fix a row index k with  $r+1 \le k \le m+1-r$ . Let S be the number of columns in A with at least two marks and let v be any one of them. For any abundant pair (i, j), every column with a mark at (i, j) is a 1-mark column by Lemma 2.4, so all marks in v correspond to scarce pairs. In particular, if v has a mark at (i, j), then  $i \le r < k < m+2-r \le j$ , which implies that  $v_k = 2$ , as otherwise v will have a mark at abundant pair (i, k) or (k, j). It follows that all columns in A with at least two marks remain distinct and do not lose any mark after removing row k. Now we construct an m-rowed matrix A' by taking all  $2^m + m2^{m-1}$  columns with no mark,  $(p-1)(\binom{m}{2} - \binom{r+1}{2}) + (r-1)2^r + 1$ columns with exactly one mark, as well as all columns obtained by removing the row k entries from all columns in A with at least two marks. Then  $A' \in Avoid(m, 3, p \cdot I_2)$ , so we have

forb
$$(m, 3, p \cdot I_2) \ge |A'| = (2^m + m2^{m-1}) + (p-1)\left(\binom{m}{2} - \binom{r+1}{2}\right) + (r-1)2^r + 1 + S$$
  
= forb $(m+1, 3, p \cdot I_2) - 2^{m-1}(m+4) - (p-1)m$ .

This completes the proof.

This allows us to define  $g_p$  to be the common value of  $\operatorname{forb}(m, 3, p \cdot K_2) - \operatorname{forb}(m, 3, p \cdot I_2)$  for all  $m \geq 2r + 2$ . By [DS21], we have  $g_1 = g_2 = 0, g_3 = 1, g_4 = 2$  and  $g_5 = 5$ . In the remaining part of this section, we prove Theorem 1.5, which establishes the asymptotic formula  $g_p \sim \frac{1}{2}p(\log_2 p)^2$ . As will be seen later, it follows from Lemma 2.3 that  $g_p \leq \frac{1}{2}p(\log_2 p)^2 + o(p(\log_2 p)^2)$ . To prove the reverse inequality, we use the following two results.

**Lemma 2.5.** Let  $m \ge 2r+2$ . Suppose  $b, c \ge 1$  and  $b+c \le r+1$ . Then there are at most  $2^r(r-b-c+1)+2^{b+c-2} \le r2^r$  m-rowed 0, 1, 2-columns with exactly bc marks formed by c 0's above b 1's, all corresponding to scarce pairs of rows.

*Proof.* Let the positions of the c 0's be  $i_1 < \cdots < i_c$  and the positions of the b 1's be  $j_1 < \cdots < j_b$ . All bc marks correspond to scarce pairs if and only if  $j_1 - i_c \ge m - r$ . Moreover, any such column v satisfies  $v_k = 2$  for all  $i_1 < k < j_b$  that is not one of  $i_1, \ldots, i_c$  or  $j_1, \ldots, j_b$ . Therefore, if  $b, c \ge 2$ , then the number of such columns is

$$\begin{split} &\sum_{i_c=c}^{r-b+1} \sum_{j_1=m+i_c=r}^{m-b+1} \sum_{i_1=1}^{i_c-c+1} \binom{i_c-i_1-1}{c-2} \sum_{j_b=j_1+b-1}^{m} \binom{j_b-j_1-1}{b-2} 2^{i_1-1+m-j_b} \\ &= \sum_{i_c=c}^{r-b+1} \sum_{j_1=m+i_c=r}^{m-b+1} \sum_{i=c-2}^{i_c-2} \binom{i}{c-2} \sum_{j=b-2}^{m-j_1-1} \binom{j}{b-2} 2^{m-3+i_c-j_1-i-j} \\ &= \sum_{i_c=c}^{r-b+1} \sum_{j_1=m+i_c=r}^{m-b+1} 2^{m-3+i_c-j_1} \left( \sum_{i=c-2}^{i_c-2} \binom{i}{c-2} 2^{-i} \right) \left( \sum_{j=b-2}^{m-j_1-1} \binom{j}{b-2} 2^{-j} \right) \\ &\leq \sum_{i_c=c}^{r-b+1} \sum_{j_1=m+i_c=r}^{m-b+1} 2^{m-3+i_c-j_1} \left( \sum_{i=c-2}^{\infty} \binom{i}{c-2} 2^{-i} \right) \left( \sum_{j=b-2}^{\infty} \binom{j}{b-2} 2^{-j} \right) \\ &= \sum_{i_c=c}^{r-b+1} \sum_{j_1=m+i_c=r}^{m-b+1} 2^{m-3+i_c-j_1} \cdot 2 \cdot 2 = \sum_{i_c=c}^{r-b+1} 2^{i_c} \sum_{j_1=m+i_c=r}^{m-b+1} 2^{m-1-j_1} \\ &= \sum_{i_c=c}^{r-b+1} 2^{i_c} (2^{r-i_c}-2^{b-2}) = \sum_{i_c=c}^{r-b+1} (2^r-2^{i_c+b-2}) \\ &= 2^r (r-b-c+1) + 2^{b+c-2} \leq r2^r, \end{split}$$

as required. The cases when b = 1 or c = 1 can be verified with similar calculations.

**Proposition 2.6.** Let  $k \ge 2$  be fixed, then we have  $g_p \ge \frac{(p-1)k}{k+1} \binom{r+1}{2} - r2^r \sum_{i=1}^k \tau(i)$ , where  $\tau(n)$  is the number of divisors of n.

Proof. Let  $A \in \text{ext}(m, 3, p \cdot I_2)$ . As before we can assume A contains at most p-1 instances of 0 above 1 in every pair of rows, and contains the maximum number of 1-mark columns. For  $j \ge 0$ , let  $a_j$  be the number of j-mark columns in A. Note that if  $j \ge 2$  and v is a j-mark column, then every mark in v corresponds to a scarce pair of rows. From before, we have  $a_0 = 2^m + m2^{m-1}$  and  $a_1 = (p-1)\left(\binom{m}{2} - \binom{r+1}{2}\right) + (r-1)2^r + 1$ . Since there are at most p-1 marks at each pair of rows, we have  $\sum_{i=1}^{\infty} ia_i \le (p-1)\binom{m}{2}$ . Let  $\sum_{i=1}^{\infty} ia_i = (p-1)\binom{m}{2} - S$ .

Let  $m \ge 2r+2$ ,  $2 \le i \le k$ , and consider an *m*-rowed 0, 1, 2-column *v* with exactly *i* marks, all of which are scarce. Since  $m \ge 2r+2$ , there does not exist x < y < x' < y' such that both (x, y)and (x', y') are scarce. So there exists *b*, *c* with bc = i such that all *i* marks of *v* are formed as *c* 0's above *b* 1's. Thus Lemma 2.5 shows there are at most  $\tau(i)r2^r$  such columns, and so  $a_i \le \tau(i)r2^r$ . We have

$$\sum_{i=k+1}^{\infty} a_i \le \sum_{i=k+1}^{\infty} \frac{1}{k+1} i a_i = \frac{1}{k+1} \left( (p-1) \binom{m}{2} - S - \sum_{i=1}^{k} i a_i \right)$$

and therefore

$$\sum_{i=1}^{\infty} a_i = \sum_{i=1}^{k} a_i + \sum_{i=k+1}^{\infty} a_i \le \frac{1}{k+1} \left( (p-1)\binom{m}{2} - S \right) + \sum_{i=1}^{k} \left( 1 - \frac{i}{k+1} \right) a_i.$$

It follows that

$$g_{p} = \operatorname{forb}(m, 3, p \cdot K_{2}) - \operatorname{forb}(m, 3, p \cdot I_{2}) = \operatorname{forb}(m, 3, p \cdot K_{2}) - |A|$$

$$= (p-1)\binom{m}{2} - \sum_{i=1}^{\infty} a_{i}$$

$$\geq \frac{(p-1)k}{k+1}\binom{r+1}{2} + \frac{S}{k+1} - \frac{k}{k+1}((r-1)2^{r}+1) - \sum_{i=2}^{k} \left(1 - \frac{i}{k+1}\right)a_{i}$$

$$\geq \frac{(p-1)k}{k+1}\binom{r+1}{2} - r2^{r} - \sum_{i=2}^{k} a_{i}$$

$$\geq \frac{(p-1)k}{k+1}\binom{r+1}{2} - r2^{r} \sum_{i=1}^{k} \tau(i).$$

Proof of Theorem 1.5. By Lemma 2.3,  $forb(m, 3, p \cdot I_2) \ge 2^m + m2^{m-1} + (p-1)\binom{m}{2} - \binom{r+1}{2} + (r-1)2^r + 1$ . Hence,

$$g_p \le \operatorname{forb}(m, 3, p \cdot K_2) - 2^m - m2^{m-1} - (p-1)\left(\binom{m}{2} - \binom{r+1}{2}\right) - (r-1)2^r - 1$$
$$= (p-1)\binom{r+1}{2} - (r-1)2^r - 1 = \frac{1}{2}p(\log_2 p)^2 + o(p(\log_2 p)^2).$$

On the other hand, Proposition 2.6 implies that for every fixed  $k \ge 2$ ,

$$g_p \ge \frac{(p-1)k}{k+1} \binom{r+1}{2} - r2^r \sum_{i=1}^k \tau(i) = \left(1 - \frac{1}{k+1}\right) \frac{1}{2} p(\log_2 p)^2 + o(p(\log_2 p)^2).$$

Combining these two bounds and letting  $k \to \infty$ , we get  $g_p \sim \frac{1}{2} p (\log_2 p)^2$ .

## **2.2** $p \le 4$

In this subsection, we compute the exact values of  $\operatorname{forb}(m, 3, F(a, b, c, d))$  when  $\max\{a, d\} < \min\{b, c\} = p \le 4$ . If p = 1, 2, then  $\operatorname{forb}(m, 3, p \cdot I_2) = \operatorname{forb}(m, 3, p \cdot K_2)$  by [DS21]. Then, for large m, Lemma 1.3 implies that  $\operatorname{forb}(m, 3, p \cdot I_2) \le \operatorname{forb}(m, 3, F(a, b, c, d)) = \operatorname{forb}(m, 3, F(a, p, p, d)) \le \operatorname{forb}(m, 3, p \cdot K_2)$ , so  $\operatorname{forb}(m, 3, F(a, b, c, d)) = \operatorname{forb}(m, 3, p \cdot K_2) = \operatorname{forb}(m, 3, p \cdot K_2) = 2^m + m2^{m-1} + (p-1)\binom{m}{2}$ .

Next, we treat the cases when p = 3 and p = 4.

**Lemma 2.7.** For  $p \ge 3$ , let  $1 \le k \le \min\{p, g_p\}$ . If  $\min\{b, c\} = p$ ,  $\max\{a, d\} \le p - k$ ,  $m \ge 2r + 2$  and  $2^{m-2} \ge (\max\{b, c\} - 1)m^2$ , then

$$forb(m, 3, F(a, b, c, d)) \le forb(m, 3, p \cdot I_2) + g_p - k = forb(m, 3, p \cdot K_2) - k.$$

Proof. The conditions ensure that Lemma 1.3 applies, so we can assume that b = c = p, and that forb $(m, 3, p \cdot I_2) = \operatorname{forb}(m, 3, p \cdot K_2) - g_p$  by Proposition 1.4. Let  $A \in \operatorname{Avoid}(m, 3, F(a, p, p, d))$  and let  $A = \begin{bmatrix} B \mid C \end{bmatrix}$  be a standard decomposition. Recall from the definition that N is the number of pairs of rows that are associated with either  $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$  or  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ . If N = 0, then  $A \in \operatorname{Avoid}(m, 3, p \cdot I_2)$  so  $|A| \leq \operatorname{forb}(m, 3, p \cdot I_2) = \operatorname{forb}(m, 3, p \cdot K_2) - g_p$ . If  $N \geq 1$ , then  $|A| \leq |B| + |C| \leq 2^m + m2^{m-1} + (p-1)\binom{m}{2} - N(p-(p-k)) \leq 2^m + m2^{m-1} + (p-1)\binom{m}{2} - k = \operatorname{forb}(m, 3, p \cdot K_2) - k$ .

Corollary 2.8. If  $\min\{b, c\} = 3$ ,  $\max\{a, d\} \le 2$ ,  $m \ge 4$  and  $2^{m-2} \ge (\max\{b, c\} - 1)m^2$ , then

 $forb(m, 3, F(a, b, c, d)) = forb(m, 3, 3 \cdot I_2) = forb(m, 3, 3 \cdot K_2) - 1.$ 

*Proof.* Set p = 3, k = 1 in Lemma 2.7 and note that  $g_3 = 1$ .

**Proposition 2.9.** If  $\min\{b, c\} = 4$ ,  $\max\{a, d\} \le 3$ ,  $m \ge 6$  and  $2^{m-2} \ge (\max\{b, c\} - 1)m^2$ , then

 $forb(m, 3, F(a, b, c, d)) = forb(m, 3, 4 \cdot I_2) = forb(m, 3, 4 \cdot K_2) - 2.$ 

*Proof.* The conditions ensure that Lemma 1.3 applies so we can assume that b = c = 4. We have that  $4 \cdot I_2 \prec F(a, 4, 4, d)$  and  $g_4 = 2$  by [DS21], so  $2^m + m2^{m-1} + 3\binom{m}{2} - 2 = \operatorname{forb}(m, 3, 4 \cdot I_2) \leq \operatorname{forb}(m, 3, F(a, 4, 4, d))$ . It suffices to show the reverse inequality.

Let  $A \in \operatorname{Avoid}(m, 3, F(a, 4, 4, d))$ , and let  $A = \lfloor B \mid C \rfloor$  be a standard decomposition. If N = 0, then  $A \in \operatorname{Avoid}(m, 3, 4 \cdot I_2)$  so  $|A| \leq \operatorname{forb}(m, 3, 4 \cdot I_2)$ . If  $N \geq 2$ , then  $|A| \leq |B| + |C| \leq (2^m + m2^{m-1}) + (3\binom{m}{2} - N) \leq \operatorname{forb}(m, 3, 4 \cdot I_2)$ .

If N = 1,  $|A| \leq |B| + |C| \leq (2^m + m2^{m-1}) + (3\binom{m}{2} - 1) = \operatorname{forb}(m, 3, 4 \cdot I_2) + 1$ . Assume for a contradiction that equality holds. Then  $|B| = 2^m + m2^{m-1}$ , so T is transitive by Lemma 2.2. Let i' < j' be the unique pair of rows associated to  $\begin{bmatrix} 0\\0 \end{bmatrix}$  or  $\begin{bmatrix} 1\\1 \end{bmatrix}$ . Without loss of generality we can assume it is associated with  $\begin{bmatrix} 0\\0 \end{bmatrix}$ . We can rearrange the rows of A so that there is no 0

above 1 in B for every pair of rows except possibly for rows i', j'. For equality to hold we also need  $|C| = 3(\binom{m}{2} - 1) + 2 = 3\binom{m}{2} - 1$ . This means that every column in C has exactly one mark, and for every i < j with  $(i, j) \neq (i', j')$  there are 3 columns in C with a mark at (i, j), and there are 2 columns in C with a mark at (i', j'). There are four cases.

If 1 < i' < j' < m, then for a column v to have exactly one mark at (1, m), we need  $v_k = 2$  for all  $2 \le k \le m - 1$ , which leaves only 1 choice for such v, contradiction. If i' = 1, j' = m, then for a column v to have exactly one mark at (1, m - 1), we need  $v_k = 2$  for all  $2 \le k \le m - 2$ . So there are only 2 choices for such v, which contradicts the above. If i' = 1, j' < m, then for a column v to have exactly one mark at (1, m - 1), we need  $v_k = 2$  for all  $2 \le k \le m - 2$ . So there are only 2 choices for such v, which contradicts the above. If i' = 1, j' < m, then for a column v to have exactly one mark at (1, m), we need  $v_k = 2$  for all  $2 \le k \le m - 1$  except that  $v_{j'}$  could be either 1 or 2, which gives only 2 choice for such v, contradiction. If i' > 1, j' = m, then for a column v to have exactly one mark at (1, m), we need  $v_k = 2$  for all  $2 \le k \le m - 1$  except that  $v_{i'}$  could be either 0 or 2, which gives only 2 choice for such v, contradiction. In all cases we have a contradiction so equality cannot hold and  $|A| \le \text{forb}(m, 3, 4 \cdot I_2)$ . Thus  $\text{forb}(m, 3, F(a, 4, 4, d)) \le \text{forb}(m, 3, 4 \cdot I_2)$ , and this completes the proof.

Up to now, we have shown that for  $p \leq 4$ , forb(m, 3, F(a, b, c, d)) =forb $(m, 3, p \cdot I_2) =$ forb $(m, 3, p \cdot I_2) =$ forb $(m, 3, p \cdot I_2) - g_p$  as long as max $\{a, d\} mia<math>\{b, c\}$  and Lemma 1.3 applies. This need not be true for larger p as will be seen in the next subsection.

## **2.3** F(a, p, p, d)

Throughout this section we assume that  $a, d \leq p$  and  $A \in ext(m, 3, F(a, p, p, d))$ .



## **2.3.1** Structure of $A \in ext(m, 3, F(a, p, p, d))$

Let  $A = \begin{bmatrix} B \mid C \end{bmatrix}$  be a standard decomposition of A as defined in Definition 2.1. Note that  $|A| \ge forb(m, 3, p \cdot I_2) = forb(m, 3, p \cdot K_2) - g_p = 2^m + m2^{m-1} + (p-1)\binom{m}{2} - g_p$  and  $|C| \le (p-1)\binom{m}{2}$ . Thus, for all  $m \ge 2N_r + 2$ , we have  $|B| > 2^m + m2^{m-1} - 2^{m-3}$ , so by Lemma 2.2 we may assume that no pair i < j of rows is associated with  $\begin{bmatrix} 1\\ 0 \end{bmatrix}$ .

**Lemma 2.10.** Let  $1 \le i < j < k \le m$ .

- *if j*, *k* forms a 0-non-edge, then one of *i*, *j* and *i*, *k* also forms a 0-non-edge,
- *if i*, *j* forms a 1-non-edge, then one of *i*, *k* and *j*, *k* also forms a 1-non-edge.

*Proof.* The proof is similar to the one of Lemma 2.2. First assume that i < j < k and j, k forms a 0-non-edge, and neither i, j nor i, k form a 0-non-edge. For any subset  $X \subset [m]$  of rows, let B(X) be the matrix consisting of columns of B with support X. Note that  $B(X)|_X$  is simple and it is in Avoid $(|X|, 2, K_2)$ , hence  $|B(X)|_X| \leq |X| + 1$  if  $\{i, j, k\} \not\subseteq X$ . For  $\{i, j, k\} \subseteq X$  we show below that  $|B(X)|_X| \leq |X|$ , so then a similar calculation as in Lemma 2.2 shows  $|B| \leq 2^m + m2^{m-1} - 2^{m-3}$ , a contradiction.

- (i) Pair (i, j) is assigned  $\begin{bmatrix} 0\\1 \end{bmatrix}$ .
  - (ia) Pair (i, k) is assigned  $\begin{bmatrix} 0\\1 \end{bmatrix}$ . In this case entries in row i of B(X) must all be 1, since i = 0 would imply both j = 0 and k = 0, which contradicts the assumption that j, k forms a 0-non-edge. Therefore,  $B(X)|_{X \setminus \{i\}}$  is simple.
  - (ib) Pair (i,k) is assigned  $\begin{bmatrix} 1\\1 \end{bmatrix}$ . In this case the only possible columns of  $B(X)|_{\{i,j,k\}}$  are  $i \mid 0 \quad 1$   $j \mid 0 \quad 1$ , hence  $B(X)|_{X \setminus \{i\}}$  is simple.  $k \mid 1 \quad 0$

(ii) Pair 
$$(i, j)$$
 is assigned  $\begin{bmatrix} 1\\1 \end{bmatrix}$ 

- (iia) Pair (i, k) is assigned  $\begin{bmatrix} 0\\1 \end{bmatrix}$ . Now, the only possible columns of  $B(X)|_{\{i,j,k\}}$  are  $\begin{array}{c}i\\j\\k\\0\end{array}$  hence  $B(X)|_{X\setminus\{i\}}$  is simple.
- (iib) Pair (i, k) is assigned  $\begin{bmatrix} 1\\1 \end{bmatrix}$ . Here, row *i* of B(X) cannot contain any entry 1, hence  $B(X)|_{X \setminus \{i\}}$  is simple.
- In all cases above, it follows that  $|B(X)| = |B(X)|_{X \setminus \{i\}}| \le \operatorname{forb}(|X| 1, 2, K_2) = |X|$ , as required. The case when i < j < k and i, j forms a 1-non-edge is similar.

**Lemma 2.11.** If i < j < k and i, k forms a non-edge, then one of i, j and j, k also forms a non-edge.

*Proof.* If ij and jk are both directed edges, then ik would also be a directed edge by the transitivity provided by Lemma 2.2, a contradiction.

**Lemma 2.12.** If there are N non-edges, then all the 0-non-edges are in the first N + 1 rows, and all the 1-non-edges are in the last N + 1 rows.

*Proof.* Let k be the largest number such that there exists j < k with j, k forming a 0-non-edge. Then either 1, j or 1, k is also a 0-non-edge by Lemma 2.10. If 1, k is a 0-non-edge, then by Lemma 2.11, for every  $1 < \ell < k$ , either 1,  $\ell$  or  $\ell, k$  is a non-edge. This means that the total number of non-edges is at least k - 1. If instead 1, j is a 0-non-edge, then similarly for 1 < i < j, one of 1, i and i, j is a non-edge, and for every  $j < \ell < k$ , one of  $j, \ell$  and  $\ell, k$  is a non-edge. Together with the non-edges 1, j and j, k we get at least (j - 1) + (k - j) = k - 1 non-edges. In both cases, we conclude that  $N \ge k - 1$ , so  $k \le N + 1$ . The 1-non-edges case is similar.

#### 2.3.2 Lower bounds

Similar to Section 2.1, we say a pair (i, j) of rows with i < j is abundant if  $2^{i-1+m-j} \ge p-1$ , and scarce otherwise. Equivalently, if  $r = \lceil \log_2(p-1) \rceil$ , then (i, j) is abundant if  $i-1+m-j \ge r$  and scarce otherwise.

**Lemma 2.13.** Suppose  $m \ge \max\{2N+2, 2r+2\}$  and there are N non-edges. Then

- for every scare pair (i, m + 1 j), if  $a_i$  is the number of 0-non-edges (i, k) with i < k, and  $b_j$  is the number of 1-non-edges (k, m + 1 j) with k < m + 1 j, then the number of columns with their unique marks at (i, m + 1 j) is  $2^{i+j-2+a_i+b_j}$ .
- for every abundant pair (i, j) that is not a non-edge, there are at least p 1 1-mark columns, each of whose unique mark is at (i, j).
- for every non-edge (i, j), there are at least p 1 1-mark columns, each of whose unique mark is at (i, j).

*Proof.* Since  $m \ge 2N + 2$ , the first and last N + 1 rows of A are disjoint. By Lemma 2.12, all 0-non-edges are within the first N + 1 rows, and all 1-non-edges are within the last N + 1 rows.

If (i, m + 1 - j) is a scarce pair, then  $i - 1 + m - (m + 1 - j) \leq r - 1$ , so  $(m + 1 - j) - i \geq m - r \geq \frac{1}{2}m \geq N + 1$ . In particular, (i, m + 1 - j) is not a non-edge. Let  $i < k_1 < k_2 < \ldots < k_{a_i}$  be such that  $(i, k_t)$  is a 0-non-edge for all  $t \in [a_i]$ , and let  $\ell_1 < \ell_2 < \ldots < \ell_{b_j} < m + 1 - j$  be such that  $(\ell_s, m + 1 - j)$  is a 1-non-edge for all  $s \in [b_j]$ . By Lemma 2.12,  $k_{a_i} \leq N + 1 < m - N \leq \ell_1$ . If v is a 1-mark column with its unique mark at (i, m + 1 - j), then we must have  $v_i = 0, v_{m+1-j} = 1$  and  $v_z = 2$  for all  $z \in \{i + 1, i + 2, \ldots, m + 1 - j\} \setminus \{k_1, k_2, \ldots, k_{a_i}, \ell_1, \ell_2, \ldots, \ell_{b_j}\}$ , while  $v_z$  can be 1 or 2 for all z < i and all  $z \in \{k_1, k_2, \ldots, k_{a_i}\}$ , and  $v_z$  can be 0 or 2 for all z > m + 1 - j and all  $z \in \{\ell_1, \ell_2, \ldots, \ell_{b_j}\}$ . Moreover, every column v of this form does contain exactly one mark at (i, m + 1 - j), so there are  $2^{i+j-2+a_i+b_j}$  1-mark columns with their unique mark at (i, m + 1 - j).

If (i, j) is abundant and not a non-edge, then  $i - 1 + m - j \ge r$ . It follows that we can find  $0 \le r_1 \le i - 1$  and  $0 \le r_2 \le m - j$ , such that  $r_1 + r_2 = r$ . Consider the columns v of the form  $v_i = 0, v_j = 1, v_z = 1$  or 2 for all  $z \le r_1, v_z = 0$  or 2 for all  $z \ge m - r_2 + 1$ , and  $v_z = 2$  for all other z. There are  $2^{r_1+r_2} = 2^r \ge p - 1$  such columns, and one can verify that each of these contains a unique mark at (i, j).

If (i, j) is a non-edge, without loss of generality a 0-non-edge, consider the columns v of the form  $v_i = v_j = 0$ ,  $v_z = 0$  or 2 for all  $z \ge m - r + 1$ , and  $v_z = 2$  for all other z. There are  $2^r \ge p - 1$  such columns, and one can verify that each of these contains a unique mark at (i, j).

**Lemma 2.14.** If there exist  $r_1, r_2 \ge 0$  with  $r_1 + r_2 \le m - 2$ , such that the 0-non-edges are exactly (i, j) with  $1 \le i < j \le r_1 + 1$  and the 1-non-edges are exactly (i, j) with  $m - r_2 \le i < j \le m$ , then there are exactly  $2^m + m2^{m-1}$  columns with no mark.

*Proof.* If  $v_z = 1$  or 2 for all  $z \le m - r_2 - 1$  and  $v_z = 0$  or 2 for all  $z \ge m - r_2$ , then v contains no mark and there are  $2^m$  such columns.

If v is not of this form and has no mark, then  $v_z = 0$  for some  $z \le m - r_2 - 1$ , or  $v_z = 1$  for some  $z \ge m - r_2$ . In the first case, let  $i \le m - r_2 - 1$  be minimal such that  $v_i = 0$ . If  $i \le r_1 + 1$ , then  $v_z = 1$ 

must be 1 or 2 for all other  $z \in [r_1 + 1]$  and  $v_z$  must be 0 or 2 for all  $z > r_1 + 1$ , and all of these  $2^{m-1}$  columns have no mark. If  $r_1 + 1 < i \leq m - r_2 - 1$ , then  $v_z$  must be 1 or 2 for all other z < iand  $v_z$  must be 0 or 2 for all z > i, and all of these  $2^{m-1}$  columns have no mark. If we are not in the first case, then  $v_z = 1$  or 2 for all  $z \le m - r_2 - 1$ . Let  $i \ge m - r_2$  be minimal such that  $v_i = 1$ . Then  $v_z$  must be 0 or 2 for all other  $z \ge m - r_2$ , and all of these  $2^{m-1}$  columns have no mark. In total, there are  $m2^{m-1}$  such columns with no mark. 

Recall that  $N_r = {\binom{\lfloor \frac{r}{2} \rfloor + 1}{2}} + {\binom{\lceil \frac{r}{2} \rceil + 1}{2}}$ . We say that a scarce pair (i, m + 1 - j) is *fixed*, if  $2^{i+j-2+a_i+b_j} \ge p-1$ , or equivalently  $i+j-2+a_i+b_j \ge r$ , using the notation of Lemma 2.13.

*Proof of Proposition 1.9.* Since  $forb(m, 3, F(p - q_0, p, p, p - q_1)) = forb(m, 3, F(p - q_1, p, p, p - q_0))$ by taking 0,1-complements, we may without loss of generality assume  $q_0 \leq q_1$ . Consider  $\binom{r_1+1}{2}$ 0-non-edges, which are (i, j) for all  $1 \le i < j \le r_1 + 1$ , and  $\binom{r_2+1}{2}$  1-non-edges, which are (i, j) for all  $m - r_2 \leq i < j \leq m$ . Using  $r_1 + r_2 = r$ , one can check that all  $\binom{r+1}{2}$  scarce pairs are fixed. By Lemma 2.14, there are  $2^m + m2^{m-1}$  columns with no mark. Since all scarce pairs are fixed, for each pair (i, j) that is not a non-edge, by Lemma 2.13, we can find p-1 columns, each of whose unique mark is at (i, j). Also by Lemma 2.13, for each 0-non-edge (i, j), we can find  $p - q_0 - 1$  columns with their unique marks at (i, j), and for each 1-non-edge (i, j), we can find  $p - q_1 - 1$  columns with their unique marks at (i, j). The matrix formed by all these distinct columns avoids  $F(p-q_0, p, p, p-q_1)$ as a configuration, so

forb
$$(m, 3, F(p - q_0, p, p, p - q_1)) \ge 2^m + m2^{m-1} + (p-1)\binom{m}{2} - q_0\binom{r_1 + 1}{2} - q_1\binom{r_2 + 1}{2}$$
  
= forb $(m, 3, p \cdot K_2) - q_0\binom{r_1 + 1}{2} - q_1\binom{r_2 + 1}{2}$ ,  
equired.

as required.

Corollary 2.15. For every  $0 \le q \le p-1$  and every  $m \ge 2N_r + 2$ ,

$$\operatorname{forb}(m, 3, F(p-q, p, p, p-q)) \ge \operatorname{forb}(m, 3, p \cdot K_2) - qN_r.$$

*Proof.* Apply Proposition 1.9 with  $q_0 = q_1 = q$  and  $r_1 = \left\lceil \frac{r}{2} \right\rceil$ ,  $r_2 = \left\lfloor \frac{r}{2} \right\rfloor$ .

Let us call a pair (i, m + 1 - j) of rows weakly fixed if, using the notation of Lemma 2.13, we have  $a_i + b_j = r + 1 - i - j$ .

Proof of Proposition 1.8. From assumption,  $d = q - (\lceil \frac{r}{2} \rceil + 1)(p - 1 - 2^{r-1}) > 0$ . We may assume  $r \geq 2$  so  $\left\lfloor \frac{r}{2} \right\rfloor > 0$ , as otherwise the result follows directly from Corollary 2.15. Consider  $\binom{\left\lceil \frac{r}{2} \right\rceil + 1}{2}$ 0-non-edges, which are (i, j) for all  $1 \le i < j \le \left\lceil \frac{r}{2} \right\rceil + 1$ , and  $\binom{\left\lfloor \frac{r}{2} \right\rfloor}{2} = \binom{\left\lfloor \frac{r}{2} \right\rfloor + 1}{2} - \lfloor \frac{r}{2} \rfloor$  1-non-edges, which are (i, j) for all  $m - \lfloor \frac{r}{2} \rfloor + 1 \le i < j \le m$ . This corresponds to  $r_1 = \lceil \frac{r}{2} \rceil$  and  $r_2 = \lfloor \frac{r}{2} \rfloor - 1$  in the language of Proposition 1.9, though as  $r_1 + r_2 = r - 1 < r$ , not every scarce pair is fixed. It can be checked that the scarce pairs (i, m+1-j) are weakly fixed for all  $i \in \lfloor \frac{r}{2} \rfloor + 1$  and  $j \in \left\lfloor \left\lfloor \frac{r}{2} \right\rfloor\right]$ , while all other scarce pairs are fixed. This means that we used  $N_r - \left\lfloor \frac{r}{2} \right\rfloor$  non-edges to fix  $\binom{r+1}{2} - \lfloor \frac{r}{2} \rfloor (\lceil \frac{r}{2} \rceil + 1)$  scarce pairs, and leave the remaining  $\lfloor \frac{r}{2} \rfloor (\lceil \frac{r}{2} \rceil + 1)$  scarce pairs weakly fixed. Therefore, forb(m, 3, F(p-q, p, p, p-q)) is at least the number of columns with 0 or 1 mark, which by Lemma 2.13 and Lemma 2.14 is

$$2^{m} + m2^{m-1} + (p-1)\binom{m}{2} - q\left(N_{r} - \lfloor \frac{r}{2} \rfloor\right) - \lfloor \frac{r}{2} \rfloor \left( \lceil \frac{r}{2} \rceil + 1 \right) (p-1-2^{r-1})$$
  
= forb(m, 3, p \cdot K\_{2}) - qN\_{r} + \begin{bmatrix} \frac{r}{2} \\ \frac{r}{2} \end{bmatrix} \left( q - \left( \lceil \frac{r}{2} \rceil + 1 \right) (p-1-2^{r-1}) \right)

$$= \operatorname{forb}(m, 3, p \cdot K_2) - qN_r + \left\lfloor \frac{r}{2} \right\rfloor d,$$

as claimed.

## 2.3.3 Upper bounds

**Lemma 2.16.** If  $m \ge 2N_r + 2$ , and N non-edges are used to fix M scarce pairs, then

$$\max(M-N) = \binom{r+1}{2} - N_r.$$

*Proof.* Since  $M \leq \binom{r+1}{2}$ , we may assume that  $N \leq N_r$ , so that  $m \geq 2N+2$ . For every  $i \in [r]$ , let  $a_i$  be the number of 0-non-edges (i,k) with i < k, and let  $b_i$  be the number of 1-non-edges (k, m+1-i) with k < m+1-i. By definition, for every  $i \in [r]$  and  $j \in [r+1-i]$ , the scarce pair (i, m+1-j) is fixed if and only if  $2^{i+j-2+a_i+b_j} \geq p-1$ , which happens if and only if  $a_i + b_j \geq r+2-i-j$ .

Consider the construction in the proof of Proposition 1.9 with  $r_1 = \lceil \frac{r}{2} \rceil$  and  $r_2 = \lfloor \frac{r}{2} \rfloor$ . Here,  $N_r = \binom{\lceil \frac{r}{2} \rceil + 1}{2} + \binom{\lfloor \frac{r}{2} \rfloor + 1}{2}$  non-edges are used to fix all  $\binom{r+1}{2}$  scarce pairs, so M - N can achieve the value  $\binom{r+1}{2} - N_r$ .

To show  $\binom{r+1}{2} - N_r$  is the best possible, consider the following equivalent problem on the triangular array  $T_r$  of size r, whose (i, j) entry is  $T_r(i, j) = r + 2 - i - j$  for all  $i \in [r]$  and  $j \in [r + 1 - i]$ .

For every  $i \in [r]$ , a row operation on row *i* reduces every entry in row *i* by 1, and a column operation on column *i* reduces every entry in column *i* by 1. If we perform  $a_i$  row operations on row *i* and  $b_i$ column operations on column *i* for all  $i \in [r]$ , then after all of these operations, the (i, j) entry of  $T_r$ is non-positive if and only if  $a_i + b_j \ge r + 2 - i - j$ , and so if and only if the scarce pair (i, m + 1 - j)is fixed. Therefore, it suffices to show that if N row or column operations turn M entries in  $T_r$ non-positive, then  $M - N \le {r+1 \choose 2} - N_r$ .

To this end, we use induction on r to show  $M - N \leq \binom{r+1}{2} - N_r = \lfloor \frac{r^2}{4} \rfloor$ . The r = 1, 2 cases can be checked easily, so we assume  $r \geq 3$ .

For each  $i \in [r]$ , let  $a_i$  and  $b_i$  be the number of operations performed on row i and column i, respectively. Observe that we may assume  $a_r = 0$ , as otherwise reducing  $a_r$  by 1 and increasing  $b_1$  by 1 does not decrease M and leaves N unchanged. Similarly, we may assume  $b_r = 0$ .

Let  $M_1$  be the number of non-positive integers in the first row or column of  $T_r$  after N operations, and let  $M' = M - M_1$ . Let  $N' = \sum_{i=2}^{r-1} a_i + \sum_{i=2}^{r-1} b_j$ , so that  $N - N' = a_1 + a_r + b_1 + b_r = a_1 + b_1$ . From the induction hypothesis,  $M' - N' \leq \left\lfloor \frac{(r-2)^2}{4} \right\rfloor$ , so we are done if  $M_1 - a_1 - b_1 \leq \left\lfloor \frac{r^2}{4} \right\rfloor - \left\lfloor \frac{(r-2)^2}{4} \right\rfloor = r - 1$ . If the top left entry of  $T_r$ , which is r, is turned non-positive, then  $a_1 + b_1 \geq r$ , and so  $M_1 - a_1 - b_1 \leq 2r - 1 - r = r - 1$  and we are done.

If the top left entry of  $T_r$  remains positive, let x be the number of non-positive entries in row 1 and let y be the number of non-positive entries in column 1, so that  $x + y = M_1$ . If  $x \le a_1$ , then  $y - b_1 \le y \le r - 1$ , so  $M_1 - a_1 - b_1 = x - a_1 + y - b_1 \le r - 1$  and we are done. Similarly, we are done if  $y \le b_1$ , so we assume neither is true.

For each  $2 \leq i \leq r$ , the (1,i) entry of  $T_r$  is non-positive after the operations if and only if  $a_1 + b_i \geq r + 1 - i$ . Since x of these entries are non-positive, we have  $\sum_{i=2}^r b_i \geq \sum_{i=r-x+1}^r \max\{r + 1 - i - a_1, 0\} = \sum_{i=1}^x \max\{i - a_1, 0\} = \sum_{k=1}^{x-a_1} k = \binom{x-a_1+1}{2}$ . Similarly, we have  $\sum_{i=2}^r a_i \geq \binom{y-b_1+1}{2}$ .

Like above, we may assume  $M_1 - a_1 - b_1 = x - a_1 + y - b_1 \ge r$ , as otherwise we are done. By convexity,  $N \ge \sum_{i=2}^r a_i + \sum_{i=2}^r b_i \ge \binom{\lfloor \frac{r+2}{2} \rfloor}{2} + \binom{\lceil \frac{r+2}{2} \rceil}{2} = N_r$ , so  $M - N \le \binom{r+1}{2} - N_r = \lfloor \frac{r^2}{4} \rfloor$ , as required.

**Lemma 2.17.** If  $m \ge 2N_r + 2$  and  $p \ge 2^{r-1} + 2q + 1$ , then

$$forb(m, 3, F(p-q, p, p, p-q)) = forb(m, 3, p \cdot K_2) - qN_r.$$

Proof. Suppose N non-edges are used to fix M scarce pairs. Let  $\mathcal{P}$  be the set of scarce pairs (i, m+1-j) that remained unfixed. For every  $(i, m+1-j) \in \mathcal{P}$ , let the (i, j) entry of  $T_r$  after the N corresponding row or columns operations be  $a_{i,j}$ , so that by Lemma 2.13 there are  $n_{i,j} := 2^{r-a_{i,j}}$  possible 1-mark columns with their unique marks at (i, m+1-j). Since (i, m+1-j) is not fixed,  $a_{i,j} \geq 1$  and so  $p-1-n_{i,j} \geq p-1-2^{r-1} \geq 2q$ .

It follows that the maximum number of 1-mark columns plus columns with at least two marks is at most

$$(p-1)\left(\binom{m}{2} - \left(\binom{r+1}{2} - M\right)\right) - qN + \sum_{(i,m+1-j)\in\mathcal{P}} n_{i,j} + \frac{1}{2} \sum_{(i,m+1-j)\in\mathcal{P}} (p-1-n_{i,j})$$

$$= (p-1)\binom{m}{2} - qN - \frac{1}{2} \sum_{(i,m+1-j)\in\mathcal{P}} (p-1-n_{i,j})$$

$$\le (p-1)\binom{m}{2} - qN - q\left(\binom{r+1}{2} - M\right)$$

$$\le (p-1)\binom{m}{2} - qN_r,$$

where the last inequality follows from Lemma 2.16.

There are at most  $2^m + m2^{m-1}$  columns with no mark, as they together form a matrix in Avoid $(m, 3, K_2)$ . Hence, the total number of columns is at most  $2^m + m2^{m-1} + (p-1)\binom{m}{2} - qN_r =$ forb $(m, 3, p \cdot K_2) - qN_r$ . This combined with Corollary 2.15 completes the proof.

**Lemma 2.18.** Let  $m \ge 2N_r + 2$ ,  $qN_r \le 2^{r-3}$ , and  $A \in Avoid(m, 3, F(p-q, p, p, p-q))$ . Suppose there is at least one scarce pair that is neither fixed nor weakly fixed, then A has at most forb $(m, 3, p \cdot K_2) - qN_r$  columns.

*Proof.* Let  $\mathcal{P}_1$  be the set of scarce pairs that are weakly fixed and let  $\mathcal{P}_2$  be the set of scarce pairs that are neither fixed nor weakly fixed. For every  $(i, m+1-j) \in \mathcal{P} := \mathcal{P}_1 \cup \mathcal{P}_2$ , let the (i, j) entry of  $T_r$  after the N operations be  $a_{i,j}$ , so  $a_{i,j} = 1$  if  $(i, m+1-j) \in \mathcal{P}_1$ , and  $a_{i,j} \geq 2$  if  $(i, m+1-j) \in \mathcal{P}_2$ . From Lemma 2.13, there are  $n_{i,j} := 2^{r-a_{i,j}}$  possible 1-mark columns with their unique marks at (i, m+1-j).

Therefore, the maximum number of columns is at most, like in the proof of Lemma 2.17,

$$2^{m} + m2^{m-1} + (p-1)\binom{m}{2} - qN - \frac{1}{2} \sum_{(i,j)\in\mathcal{P}} (p-1-n_{i,j})$$
  
$$< 2^{m} + m2^{m-1} + (p-1)\binom{m}{2} - qN - \frac{1}{2} \cdot 2^{r-2}$$
  
$$\leq 2^{m} + m2^{m-1} + (p-1)\binom{m}{2} - qN_{r}$$
  
$$= \operatorname{forb}(m, 3, p \cdot K_{2}) - qN_{r},$$

as required, where the first inequality uses  $\emptyset \neq \mathcal{P}_2 \subset \mathcal{P}$  and  $p-1-n_{i,j} \geq p-1-2^{r-2} > 2^{r-2}$  for all  $(i, m+1-j) \in \mathcal{P}_2$ , while the second inequality holds because  $\frac{1}{2} \cdot 2^{r-2} \geq qN_r$ .

**Lemma 2.19.** If all scarce pairs are fixed or weakly fixed by N non-edges, then  $N_r - N \leq \left\lceil \frac{r}{2} \right\rceil$ .

Proof. In the language of Lemma 2.16, all entries in the triangular array  $T_r$  are at most 1 after N row or column operations, and we want to show  $N \ge N_r - \lceil \frac{r}{2} \rceil$ . We may assume that all operations are performed on the first r-1 rows and columns, as the entries in the last row and the last column are already at most 1. Now, apply this set of operations on the first r-1 rows and columns of  $T_r$  to the triangular array  $T_{r-1}$  instead, and note that they turn all  $\binom{r}{2}$  entries in  $T_{r-1}$  non-positive. Thus, by Lemma 2.16 we have  $\binom{r}{2} - N \le \binom{r}{2} - N_{r-1}$ , and so  $N \ge N_{r-1} = N_r - \lceil \frac{r}{2} \rceil$ .

For  $i \in [r]$  and  $j \in [r+1-i]$ , we say the (i, j) entry of  $T_r$  is fixed (or weakly fixed) if the corresponding scarce pair (i, m+1-j) is fixed (or weakly fixed).

**Lemma 2.20.** If all scarce pairs are fixed or weakly fixed by  $N_r - t$  non-edges for some  $0 \le t \le \left\lceil \frac{r}{2} \right\rceil$ , then there are at least  $t(\left| \frac{r}{2} \right| + 1)$  weakly fixed pairs.

*Proof.* We use induction on r and work equivalently in the triangular array  $T_r$ . The r = 1, 2 cases are easy to verify, so we assume  $r \ge 3$ .

For every  $1 \le i \le \left\lceil \frac{r+1}{2} \right\rceil$ , let  $a_i$  be the number of operations performed on row i of  $T_r$  and let  $b_i$  be the number of operations performed on column i of  $T_r$ . For every  $1 \le i \le \left\lceil \frac{r+1}{2} \right\rceil$ , since the pair (1, m + 1 - i) is fixed or weakly fixed, the (1, i) entry of  $T_r$ , which was r - i + 1 at the beginning, is now at most 1. Thus,  $r - i + 1 - a_1 - b_i \le 1$ , and so  $a_1 + b_i \ge r - i$ . In particular,  $a_1 + b_1 \ge r - 1$ . Moreover, if the (1, i) entry of  $T_r$  for every  $1 \le i \le \left\lceil \frac{r+1}{2} \right\rceil$ , we see that  $a_i + b_i \ge r - i + 1$ . Similarly, by looking at the (i, 1) entry of  $T_r$  for every  $1 \le i \le \left\lceil \frac{r+1}{2} \right\rceil$ , we see that  $a_i + b_1 \ge r - i$ , and  $a_i + b_1 \ge r - i + 1$  if this entry is non-positive.

Consider the subtriangle of size r-2 obtained by removing the first row and column from  $T_r$ . Let  $N_{r-2} - S$  be the number of operations performed in this subtriangle. Then  $r-1 \leq a_1 + b_1 = N_r - t - (N_{r-2} - S)$ , so  $S \geq r - 1 - N_r + N_{r-2} + t = t - 1$ . Also, since all entries in this subtriangle are fixed or weakly fixed, we have  $N_{r-2} - S \geq N_{r-2} - \left\lceil \frac{r-2}{2} \right\rceil = N_{r-2} - \left\lceil \frac{r}{2} \right\rceil + 1$  by Lemma 2.19. It follows that  $S \leq \left\lceil \frac{r}{2} \right\rceil - 1$ , and we can apply the induction hypothesis to see that there are at least  $S(\lfloor \frac{r-2}{2} \rfloor + 1) = S \lfloor \frac{r}{2} \rfloor$  weakly fixed entries in this subtriangle.

If  $S \ge t+1$ , then as  $S \le \lfloor \frac{r}{2} \rfloor - 1$  from above, we have  $t \le \lfloor \frac{r}{2} \rfloor - 2$ . From the induction hypothesis, the subtriangle contains at least  $(t+1) \lfloor \frac{r}{2} \rfloor \ge t(\lfloor \frac{r}{2} \rfloor + 1)$  weakly fixed entries, and hence so does the original triangle, as required. Otherwise, S = t + x for x = -1 or 0. Then, we have  $a_1 + b_1 = N_r - t - (N_{r-2} - S) = r + x$ . Therefore,

$$\sum_{i=2}^{\left\lceil \frac{r+1}{2} \right\rceil} (a_i + b_1) + \sum_{i=2}^{\left\lceil \frac{r+1}{2} \right\rceil} (a_1 + b_i) \le \left\lceil \frac{r-1}{2} \right\rceil (r+x) + (N_{r-2} - S)$$

$$= \begin{cases} \frac{3}{4}r^2 + \frac{1}{2}(x-1)r - x - t, & \text{if } r \text{ is even,} \\ \frac{3}{4}r^2 + \frac{1}{2}(x-2)r + \frac{1}{4}(1 - 6x - 4t), & \text{if } r \text{ is odd.} \end{cases}$$

On the other hand,

$$\sum_{i=2}^{\left\lceil \frac{r+1}{2} \right\rceil} (r-i+1) + \sum_{i=2}^{\left\lceil \frac{r+1}{2} \right\rceil} (r-i+1) = \begin{cases} \frac{3}{4}r^2 - \frac{1}{2}r, & \text{if } r \text{ is even,} \\ \frac{3}{4}r^2 - r + \frac{1}{4}, & \text{if } r \text{ is odd.} \end{cases}$$

Therefore, if r is even, then at least

$$\frac{3}{4}r^2 - \frac{1}{2}r - \left(\frac{3}{4}r^2 + \frac{1}{2}(x-1)r - x - t\right) = x + t - \frac{1}{2}xr$$

of the inequalities  $a_i + b_1 \ge r - i + 1$  for  $2 \le i \le \frac{r}{2} + 1$  and  $a_1 + b_i \ge r - i + 1$  for  $2 \le i \le \frac{r}{2} + 1$  must fail to hold, which leads to  $x + t - \frac{1}{2}xr$  weakly fixed entries in the first row or column. If x = -1,

this is  $\frac{1}{2}r + t - 1$ , which along with the weakly fixed (1, 1) entry and the at least  $(t - 1)\frac{r}{2}$  weakly fixed entries in the subtriangle guaranteed by the induction step above gives the required  $t(\frac{r}{2} + 1)$  weakly fixed entries overall. If x = 0, this is t, and there are at least  $t\frac{r}{2}$  weakly fixed entries in the subtriangle by induction, so we have the required  $t(\frac{r}{2} + 1)$  overall as well.

If r is odd, then at least

$$\frac{3}{4}r^2 - r + \frac{1}{4} - \left(\frac{3}{4}r^2 + \frac{1}{2}(x-2)r + \frac{1}{4}(1-6x-4t)\right) = \frac{3}{2}x + t - \frac{1}{2}xr$$

of the inequalities  $a_i + b_1 \ge r - i + 1$  for  $2 \le i \le \left\lceil \frac{r+1}{2} \right\rceil$  and  $a_1 + b_i \ge r - i + 1$  for  $2 \le i \le \left\lceil \frac{r+1}{2} \right\rceil$  must fail to hold, leading to  $\frac{3}{2}x + t - \frac{1}{2}xr$  weakly fixed entries in the first row or column. If x = -1, this is  $\frac{1}{2}r + t - \frac{3}{2} = \left\lfloor \frac{r}{2} \right\rfloor + t - 1$ , which along with the weakly fixed (1, 1) entry and the at least  $(t-1) \lfloor \frac{r}{2} \rfloor$ weakly fixed entries in the subtriangle guaranteed by induction gives the required  $t(\lfloor \frac{r}{2} \rfloor + 1)$  weakly fixed entries overall. If x = 0, this is t, and there are at least  $t \lfloor \frac{r}{2} \rfloor$  weakly fixed entries in the subtriangle by induction, so we have the required  $t(\lfloor \frac{r}{2} \rfloor + 1)$  overall as well, finishing the proof.  $\Box$ 

Proof of Theorem 1.6. We have already seen in Lemma 2.17 that equality holds if  $p \ge 2^{r-1} + 2q + 1$ . If  $p \le 2^{r-1} + 2q$  and  $qN_r \le 2^{r-3}$ , then by Lemma 2.18, the number of columns is at most forb $(m, 3, p \cdot K_2) - qN_r$  if at least one scarce pair is neither fixed nor weakly fixed. Now assume all scarce pairs fixed or weakly fixed and  $\frac{1}{2}(\lfloor \frac{r}{2} \rfloor + 1)(p - 1 - 2^{r-1}) \ge q$ .

Suppose there are  $N = N_r - t$  non-edges, and they fix M pairs, then by Lemma 2.19 we have  $t \leq \lfloor \frac{r}{2} \rfloor$ . Using the notation in the proof of Lemma 2.18, the number of columns with at least 1 mark is at most

$$(p-1)\binom{m}{2} - qN - \frac{1}{2} \sum_{(i,m+1-j)\in\mathcal{P}_1} (p-1-n_{i,j})$$
$$= (p-1)\binom{m}{2} - qN - \frac{1}{2}|\mathcal{P}_1|(p-1-2^{r-1})$$
$$= (p-1)\binom{m}{2} - qN_r + qt - \frac{1}{2}|\mathcal{P}_1|(p-1-2^{r-1}).$$

If  $t \leq 0$ , then this is at most  $(p-1)\binom{m}{2} - qN_r$ . If  $1 \leq t \leq \lceil \frac{r}{2} \rceil$ , then Lemma 2.20 implies that  $|\mathcal{P}_1| \geq t \left( \lfloor \frac{r}{2} \rfloor + 1 \right)$ , so this is at most

$$(p-1)\binom{m}{2} - qN_r + qt - \frac{1}{2}t\left(\left\lfloor\frac{r}{2}\right\rfloor + 1\right)(p-1-2^{r-1}) \le (p-1)\binom{m}{2} - qN_r.$$

Thus, in both cases, the maximum number of columns is at most

$$2^{m} + m2^{m-1} + (p-1)\binom{m}{2} - qN_r = \text{forb}(m, 3, p \cdot K_2) - qN_r.$$

Together with Corollary 2.15, this completes the proof.

Proof of Corollary 1.7. If p = 2, then  $N_r = N_0 = 0$ . From [DS21],  $g_2 = 0$ , so we have forb $(m, 3, 2 \cdot K_2) = \operatorname{forb}(m, 3, 2 \cdot I_2)$  for  $m \ge 2$  by Proposition 1.4. Since forb $(m, 3, 2 \cdot I_2) \le \operatorname{forb}(m, 3, F(1, 2, 2, 1)) \le \operatorname{forb}(m, 3, 2 \cdot K_2)$ , we have forb $(m, 3, F(1, 2, 2, 1)) = \operatorname{forb}(m, 3, 2 \cdot K_2) = \operatorname{forb}(m, 3, 2 \cdot K_2) - N_0$ . If p = 3, then  $N_r = N_1 = 1$ . By Corollary 2.8, noting that we do not need to apply the Lemma 1.3, for  $m \ge 4$  we have forb $(m, 3, F(2, 3, 3, 2)) = \operatorname{forb}(m, 3, 3 \cdot K_2) - 1 = \operatorname{forb}(m, 3, 3 \cdot K_2) - N_1$ . If p = 4, then  $N_r = N_2 = 2$ . By Proposition 2.9, again without the need for Lemma 1.3, for  $m \ge 6$  we have forb $(m, 3, F(3, 4, 4, 3)) = \operatorname{forb}(m, 3, 4 \cdot K_2) - 2 = \operatorname{forb}(m, 3, 4 \cdot K_2) - N_2$ . Therefore, the statement holds for p = 2, 3, 4.

For  $p \geq 5$ , we have  $r \geq 2$ . It follows from Corollary 2.15 that  $\operatorname{forb}(m, 3, F(p-1, p, p, p-1)) \geq \operatorname{forb}(m, 3, p \cdot K_2) - N_r$ , so we just need to prove the reverse inequality. If all scarce pairs are fixed or weakly fixed, then as  $\frac{1}{2}(\lfloor \frac{r}{2} \rfloor + 1)(p-1-2^{r-1}) \geq 1$ , the proof of Theorem 1.6 in the case when all scarce pairs are fixed or weakly fixed shows the maximum number of columns is  $\operatorname{forb}(m, 3, p \cdot K_2) - N_r$ .

Therefore, we just need to consider the case when at least one scarce pair is neither fixed nor weakly fixed. Assume that N non-edges are used and  $\mathcal{P}$  is the set of scarce pairs (i, m + 1 - j) that are not fixed. Then, as in the proof of Lemma 2.18, the maximum possible number of columns is

$$2^{m} + m2^{m-1} + (p-1)\binom{m}{2} - N - \frac{1}{2} \sum_{(i,m+1-j)\in\mathcal{P}} (p-1-n_{i,j}).$$

This is at most for  $(m, 3, p \cdot K_2) - N_r$  if and only if

$$N + \frac{1}{2} \sum_{(i,m+1-j)\in\mathcal{P}} (p-1-n_{i,j}) \ge N_r.$$
(1)

Let  $w_f$  denote the number of weakly fixed pairs and  $n_f$  denote the number of neither fixed nor weakly fixed ones, so that  $w_f + n_f = |\mathcal{P}|$ . If (i, m + 1 - j) is neither fixed nor weakly fixed, then  $p - 1 - n_{i,j} \ge p - 1 - 2^{r-2} \ge 2^{r-2} + 1$ , so

$$N + \frac{1}{2} \sum_{(i,j)\in\mathcal{P}} (p-1-n_{i,j}) \ge N + \frac{1}{2} w_f + (2^{r-3} + \frac{1}{2}) n_f.$$
<sup>(2)</sup>

We will prove that for all r,

$$N + \frac{1}{2}w_f + (2^{r-3} + \frac{1}{2})n_f \ge N_r,$$
(3)

which then implies (1) and finishes the proof.

If  $r \ge 7$ , then  $2^{r-3} \ge N_r$ , and as  $n_f \ge 1$ , (3) holds. The values of  $N_r$  for  $2 \le r \le 6$  are as follows.

If r = 2, then we are done if  $N \ge 1$  as  $n_f \ge 1$ . If N = 0, then  $w_f = 2$  and (3) holds as well.

For r = 3, we need  $N + \frac{1}{2}w_f + \frac{3}{2}n_f \ge 4$ . If  $N \ge 3$ , then the inequality holds as  $n_f \ge 1$ . If N = 2, (3) holds as at least one of the three diagonal entries is weakly fixed, while  $N \le 1$  implies  $n_f \ge 2$ and  $w_f \ge 2$ , so (3) still holds.

For r = 4, we need  $N + \frac{1}{2}w_f + \frac{5}{2}n_f \ge 6$ , which holds if  $N \ge 4$  as  $n_f \ge 1$ . If N = 3, it holds as  $w_f \ge 1$ . If N = 1, 2, then at least two of the three entries in the upper left corner of  $T_4 = \begin{pmatrix} 4 & 3 \\ 3 & 1 \end{pmatrix}$  are not at least weakly fixed, so  $n_f \ge 2$  and (3) holds. If N = 0,  $n_f = 6$  and (3) holds.

For r = 5, we need  $N + \frac{1}{2}w_f + \frac{9}{2}n_f \ge 9$ , which holds if  $N \ge 5$ . If N = 4, then (3) holds as  $w_f \ge 1$ . If  $N \le 3$ , then  $n_f \ge 2$  and (3) holds, as at least two of the three upper left corner entries of  $T_5 - \frac{5}{4} - \frac{4}{4}$  are not at least weakly fixed.

Finally for r = 6, we need  $N + \frac{1}{2}w_f + \frac{17}{2}n_f \ge 12$ , which holds if  $N \ge 4$ . If  $N \le 3$ , then  $n_f \ge 3$  and (3) holds as all three entries of the upper left corner of  $T_6 \begin{bmatrix} 6 & 5 \\ 5 \end{bmatrix}$  are not at least weakly fixed. This proves (3) for all  $r \ge 2$  and finishes the proof.

# References

- [ADLS23] Richard P Anstee, Jeffrey Dawson, Linyuan Lu, and Attila Sali. Multivalued matrices and forbidden configurations. *Discrete Mathematics*, 346(9):113492, 2023.
- [AL14] Richard Anstee and Linyuan Lu. Unavoidable multicoloured families of configurations, 2014. arXiv:1409.4123.
- [Alo83] Noga Alon. On the density of sets of vectors. Discrete Mathematics, 46(2):199–202, 1983.
- [Ans13] Richard Anstee. A survey of forbidden configuration results. *The Electronic Journal of Combinatorics*, 20(1), 2013.
- [DS21] Travis Dillon and Attila Sali. Exponential multivalued forbidden configurations. Discrete Mathematics & Theoretical Computer Science, 23(Combinatorics), 2021.
- [ELS20] Keaton Ellis, Baian Liu, and Attila Sali. Multi-symbol forbidden configurations. Discrete Applied Mathematics, 276:24 – 36, 2020.
- [FS12] Zoltán Füredi and Attila Sali. Optimal multivalued shattering. SIAM Journal on Discrete Mathematics, 26:737–744, 2012.
- [Sau72] Norbert Sauer. On the density of families of sets. Journal of Combinatorial Theory, Series A, 13:145–147, 1972.
- [She72] Saharon Shelah. A combinatorial problem: Stability and order for models and theories in infinitary language. *Pacific Journal of Mathematics*, 41:247–261, 1972.
- [Ste78] J Michael Steele. Existence of submatrices with all possible columns. Journal of Combinatorial Theory, Series A, 24(1):84–88, 1978.
- [VC71] Vladimir Vapnik and Alexey Chervonenkis. On the uniform convergence of relative frequencies of events to their probabilities. Theory of Probability and Its Applications, 16:264–280, 1971.