

A Family of Algebraic Operations Extending the Turaev Cobracket

Toyo TANIGUCHI *

Abstract

We introduce a family of maps parametrised by certain ribbon graphs. It is based on a connection in non-commutative geometry and contains the double divergence as a special case. Applying the construction to the case of the group algebra of the fundamental group of a compact connected oriented surface with boundary, we obtain an algebraic generalisation of the Turaev cobracket. If the connection is flat, they define classes in the Lie algebra cohomology of the space of derivations. In the case of the free associative algebra, we show that they are canonically identified with the standard generators of the cohomology ring of the matrix Lie algebra \mathfrak{gl}_n .

2020 Mathematics Subject Classification: 16D20, 17B56, 17B65, 53C05, 57K20, 58B34.

Keywords: non-commutative geometry, divergence maps, flat connections, ribbon graph, loop operations.

1. INTRODUCTION

The Turaev cobracket is a *loop operation* introduced in [Tur91] together with its quantisation, and gives a Lie bialgebra structure on the vector space spanned by the homotopy set of free loops on a connected compact oriented surface. This Lie bialgebra is a topological counterpart of the Batalin-Vilkovisky structure on the moduli space of flat principal super Lie group bundles; the Lie algebra part is classically known by Goldman [Gol86], and the extension to the whole BV structure is done by Alekseev, Naef, Pulmann, and Ševera [ANPŠ24].

If the surface has only one boundary component, the associated graded of this Lie bialgebra with a suitable filtration is isomorphic to Schedler's necklace Lie bialgebra on the cyclic quotient $|T(W)|$ of the free associative algebra over a vector space W ; see Section 6 of [KK16] for the precise statement and original references. This algebraic operation also appears in the framework of operad theory. In [MW15], Merkulov–Willwacher introduced a prop of ribbon graphs with a canonical representation on $|T(W)|$ where W is equipped with a skew-symmetric pairing. In particular, Theorem 4.2.3 in [MW15] shows that a representation over this prop is automatically an involutive Lie bialgebra, and it can be shown that their bialgebra structure on $|T(W)|$ coincides with Schedler's one.

In this paper, we will define a family of algebraic operations on the space of free loops extending the Turaev cobracket. We shall briefly see the construction in a special case. First, we define a series of maps called *k-divergence*

$$\mathrm{Div}_k^\nabla : \mathrm{Der}_{\mathbb{K}}(A)^{\otimes k} \rightarrow |A|^{\otimes 2}$$

associated with a \mathbb{K} -algebra A , an A -bimodule M , an action of $\mathrm{Der}_{\mathbb{K}}(A)$ on M and a non-commutative version of a connection (or a covariant derivative) ∇ on M . In some situations, $|A|$ is endowed with a Lie bracket and a Lie algebra homomorphism $\psi : |A| \rightarrow \mathrm{Der}_{\mathbb{K}}(A)$. The pull-back

$$\mathrm{Div}_k^\nabla \circ \psi^{\otimes k} : |A|^{\otimes k} \rightarrow |A|^{\otimes 2}$$

is denoted by $\delta_k^{\psi, \nabla}$.

*Graduate School of Mathematical Sciences, The University of Tokyo. 3-8-1, Komaba, Meguro-ku, Tokyo, 153-8914, Japan.
E-mail: toyo(at)ms.u-tokyo.ac.jp

For $k = 1$, the map Div_1^∇ is an analogue of the usual divergence; see the author's previous paper [Tan24b] for more details. Its pull-back $\delta_1^{\psi, \nabla}$ has a very interesting interaction with loop operations. The result of Alekseev–Kawazumi–Kuno–Naef [AKKN23] and the author [Tan24b] state that, in the case of group algebra $A = \mathbb{K}\pi$ of the fundamental group $\pi = \pi_1(\Sigma)$ of a connected oriented compact surface Σ with non-empty boundary, we can choose M , ∇ and ψ suitably so that $-\delta_1^{\psi, \nabla}$ coincides the framed version of the Turaev cobracket in [AKKN23]. In fact, the Lie algebra map ψ is given by the Kawazumi–Kuno action σ introduced in [KK14], which is another loop operation. See Remark 2.7 for a more precise statement. For $k = 2$, some values of $\delta_2^{\sigma, \nabla}$ are computed in Example 2.8.

The original motivation for the construction of k -divergence was to craft another loop operation such that multiple intersections are resolved at the same time. Our rather algebraic definition of the operations $\delta_k^{\psi, \nabla}$ lacks topological viewpoint: even in the case of the group algebra $\mathbb{K}\pi$, each operation is not mapping-class-group equivariant as opposed to the case of the Turaev cobracket, but instead, the framework is applicable to an arbitrary algebra. See Remark 3.6 for the discussion.

In the case of $A = T(W)$, the free associative algebra over a \mathbb{K} -vector space W , the ribbon graph operations reappear. Suppose we are given a *skew-symmetric* pairing

$$\langle \cdot, \cdot \rangle : W \otimes W \rightarrow \mathbb{K},$$

from which we obtain a Lie algebra homomorphism $\text{Ham}_{\langle \cdot, \cdot \rangle} : |T(W)| \rightarrow \text{Der}_{\mathbb{K}}(T(W))$. Here, $|T(W)|$ is equipped with the necklace bracket.

Theorem (Theorem 3.4). *Let ∇_W be the canonical flat connection given in Definition 3.1 on the space of non-commutative 1-forms on $T(W)$. Then, the map*

$$(-1)^k \delta_k^{\text{Ham}_{\langle \cdot, \cdot \rangle}, \nabla_W} : |T(W)|^{\otimes k} \rightarrow |T(W)|^{\otimes 2}$$

coincides with the operation associated with the unique bivalent connected ribbon graph with k vertices and k edges.

Recall that the divergence map on an oriented Riemannian manifold in the usual differential geometry is a 1-cocycle on the Lie algebra of vector fields. The following is a corresponding result for k -divergences. Let A be any \mathbb{K} -algebra and $\text{alt} : \wedge^k \text{Der}_{\mathbb{K}}(A) \rightarrow \text{Der}_{\mathbb{K}}(A)^{\otimes k}$ the alternisation map.

Theorem (Theorem 4.13). *If the connection ∇ is flat, the map $\text{Div}_k^\nabla \circ \text{alt}$ is a Lie algebra k -cocycle on $\text{Der}_{\mathbb{K}}(A)$. If k is even, the cocycle vanishes.*

Along the way, we develop a theory of non-commutative de Rham cohomology with vector bundle coefficients, which is eventually applied to give a proof of the theorem above.

In the important case of $A = T(W)$, we have an explicit description of the cocycle for k odd. Recall that the cohomology ring $H_{\text{CE}}^\bullet(\mathfrak{gl}(W), \mathbb{K})$ is isomorphic to the exterior algebra with a single degree- k generator for each k odd, $1 \leq k \leq 2 \dim(W) - 1$; for the proof, see Theorem 2.1.6 of [Fuk86] for example.

Theorem (Theorem 4.15). *Let $\text{Der}_{\mathbb{K}}(T(W))^{(\geq 0)}$ be the space of derivations with non-negative degree. Then, we have a canonical isomorphism*

$$H_{\text{CE}}^\bullet(\mathfrak{gl}(W), \mathbb{K}) \cong H_{\text{CE}}^\bullet(\text{Der}_{\mathbb{K}}(T(W))^{(\geq 0)}, |T(W)|),$$

under which the cohomology class of $(-1)^k \text{Div}_k^{\nabla_W} \circ \text{alt}$ for k odd is sent to the standard generator of $H_{\text{CE}}^\bullet(\mathfrak{gl}(W), \mathbb{K})$ of degree k .

Organisation of the paper. In Section 2, We define k -divergences and calculate some values in the case of group algebra. Section 3 is occupied with the proof of Theorem 3.4. Section 4 is devoted to the general theory of non-commutative de Rham cohomology with vector bundle coefficients, which is used in the proof of Theorem 4.13. In Appendix A, we define the infinitesimal action of a derivation to a connection and discuss the relation with the associated divergence for further analysis of divergence maps.

Acknowledgements. The author thanks Sergei Merkulov for the heart-warming discussion and for sharing their draft on ribbon graphs and loop operations, and Nariya Kawazumi for many comments on the draft.

Conventions. \mathbb{K} is a field of characteristic zero. Unadorned tensor products are always over \mathbb{K} . We use Sweedler-type notation: an element of the tensor product $X \otimes X$ is denoted as $x = x' \otimes x''$.

2. DIVERGENCE MAPS AND A FAMILY OF OPERATIONS

Let B be a unital associative \mathbb{K} -algebra. Recall some notions introduced in [Tan24b]. Also, see [Gin05] for a general theory of non-commutative differential geometry.

Definition 2.1.

- The *trace space* of B is defined by $|B| = B/[B, B]$, which is the quotient by the \mathbb{K} -vector space generated by commutators. This is merely the cyclic quotient, not the abelianisation.
- The space of *non-commutative 1-forms* $\Omega^1 B$ on B is a B -bimodule defined by

$$\Omega^1 B = B^e \{ db : b \in B \} / \langle d(bb') = (db)b' + b(db'), d \text{ is } \mathbb{K}\text{-linear} \rangle.$$

The space of p -forms is defined by $\Omega^p B = \underbrace{\Omega^1 B \otimes_B \cdots \otimes_B \Omega^1 B}_{p \text{ times}}$.

- For $f \in \text{Der}_{\mathbb{K}}(B)$, the *Lie derivative* $L_f : \Omega^1 B \rightarrow \Omega^1 B$ is defined by

$$L_f(bdb') = f(b)db' + bdf(b') \quad \text{for } b, b' \in B.$$

- A *derivation action* on a B -module M is a triple $(\mathfrak{d}, \varphi, \rho)$ where \mathfrak{d} is a Lie algebra over \mathbb{K} and

$$\varphi : \mathfrak{d} \rightarrow \text{Der}_{\mathbb{K}}(B) \quad \text{and} \quad \rho : \mathfrak{d} \rightarrow \text{End}_{\mathbb{K}}(M)$$

are Lie algebra homomorphisms satisfying the compatibility:

$$\rho(f)(bm) = \varphi(f)(b)m + b\rho(f)(m) \quad \text{for } f \in \mathfrak{d}, b \in B \text{ and } m \in M.$$

- A *connection* on M is a \mathbb{K} -linear map $\nabla : M \rightarrow \Omega^1 B \otimes_B M$ satisfying the Leibniz rule:

$$\nabla(bm) = db \otimes m + b\nabla(m) \quad \text{for } b \in B \text{ and } m \in M.$$

This is extended to the degree +1 derivation by the following:

$$\begin{aligned} \nabla : \Omega^p B \otimes_B M &\rightarrow \Omega^{p+1} B \otimes_B M \\ \omega \otimes m &\mapsto d\omega \otimes m + (-1)^p \omega \nabla(m), \end{aligned}$$

and the connection ∇ is *flat* if $\nabla^2 : M \rightarrow \Omega^2 B \otimes_B M$ vanishes.

Definition 2.2. Let $(\mathfrak{d}, \varphi, \rho)$ be a derivation action on M , and ∇ a connection on M . With these data given, we define a map c_{∇} as

$$c_{\nabla} : \mathfrak{d} \rightarrow \text{End}_B(M) : f \mapsto (i_{\varphi(f)} \otimes \text{id}_M) \circ \nabla - \rho(f).$$

Note that it takes its value in the space of B -linear maps by the compatibility. The property of this map is discussed in Section 4.

Definition 2.3. Suppose M is dualisable (i.e., finitely generated and projective). For $k \geq 1$, we define the *k -divergence map* $\text{Div}_k^{\nabla} : \mathfrak{d}^{\otimes k} \rightarrow |B|$ associated with a derivation action $(\mathfrak{d}, \varphi, \rho)$ and a connection ∇ by the following: for $f_1, \dots, f_k \in \mathfrak{d}$,

$$\text{Div}_k^{\nabla}(f_1, \dots, f_k) = \text{Tr}(c_{\nabla}(f_1) \circ \cdots \circ c_{\nabla}(f_k)).$$

We denote the pullback by a Lie algebra homomorphism $\psi : \mathfrak{g} \rightarrow \mathfrak{d}$ for some Lie algebra \mathfrak{g} by $\delta_k^{\psi, \nabla} = \text{Div}_k^{\nabla} \circ \psi^{\otimes k}$.

Proposition 2.4. Div_k^∇ is cyclically symmetric in its arguments: $\text{Div}_k^\nabla(f_1, f_2, \dots, f_k) = \text{Div}_k^\nabla(f_2, \dots, f_k, f_1)$.

Proof. This follows from the property of the trace map. \square

Remark 2.5. The case of $k = 1$ is simply called the *divergence map*, and its details and applications are discussed in [Tan24b]. Note that the divergence here differs from [Tan24b] by a sign, which is purely conventional.

Definition 2.6. For a \mathbb{K} -algebra A , the default derivation action is defined by $B = A^e$, $M = \Omega^1 A$, $\mathfrak{d} = \text{Der}_{\mathbb{K}}(A)$,

$$\begin{aligned} \varphi: \text{Der}_{\mathbb{K}}(A) &\rightarrow \text{Der}_{\mathbb{K}}(A^e) &: f &\mapsto f \otimes \text{id} + \text{id} \otimes f, \text{ and} \\ \rho: \text{Der}_{\mathbb{K}}(A) &\rightarrow \text{End}_{\mathbb{K}}(\Omega^1 A) &: f &\mapsto L_f, \end{aligned}$$

and we will use this action unless otherwise specified.

Remark 2.7. Let $\Sigma = \Sigma_{g,n+1}$, an oriented connected compact surface of genus g with $n+1$ boundary components. Put $\pi = \pi_1(\Sigma, *)$, the fundamental group of Σ with a base point $* \in \partial\Sigma$, and consider the group algebra $A = \mathbb{K}\pi$. Then, the Kawazumi-Kuno action $\sigma: |\mathbb{K}\pi| \rightarrow \text{Der}_{\mathbb{K}}(\mathbb{K}\pi)$ is defined by the following formula: for a free loop α and a based loop β , both generically immersed, we set

$$\sigma(\alpha)(\beta) = \sum_{p \in \alpha \cap \beta} \text{sign}(p; \alpha, \beta) \alpha * _p \beta$$

where $\text{sign}(p; \alpha, \beta)$ is a local intersection number with respect to the orientation of Σ , and $\alpha * _p \beta$ is a curve obtained by surgery at the intersection p . This is a Lie algebra homomorphism with $|\mathbb{K}\pi|$ endowed with the structure of the Goldman Lie algebra. See [AKKN23] for the details.

Now take a free generating system $\mathcal{C} = (\alpha_i, \beta_i, \gamma_j)_{1 \leq i \leq g, 1 \leq j \leq n}$ of π so that α_i and β_i form a genus pair, γ_j is a boundary loop and

$$\alpha_1 \beta_1 \alpha_1^{-1} \beta_1^{-1} \dots \alpha_g \beta_g \alpha_g^{-1} \beta_g^{-1} \gamma_1 \dots \gamma_n$$

represents a boundary loop based at $* \in \partial\Sigma$. The flat connection $\nabla_{\mathcal{C}}$ on $\Omega^1 \mathbb{K}\pi$ is defined by $\nabla_{\mathcal{C}}((dc)c^{-1}) = 0$ for all $c \in \mathcal{C}$. Then, by the result of Alekseev–Kawazumi–Kuno–Naef [AKKN23] and the author [Tan24b], the map $-\delta_1^{\sigma, \nabla_{\mathcal{C}}}: |\mathbb{K}\pi| \rightarrow |\mathbb{K}\pi|^{\otimes 2}$ coincides with the framed version of the Turaev cobracket. From this viewpoint, the series of maps $\delta_k^{\sigma, \nabla_{\mathcal{C}}}$ is regarded as an algebraic extension of the Turaev cobracket.

Table 1: Some values of $\delta_2^{\sigma, \nabla_{\mathcal{C}}}$.

x	y	$\delta_2^{\sigma, \nabla_{\mathcal{C}}}(x, y)$	(x, y) disjoint?
s^m	any element	0	Y
st	st	$3stst \otimes 1 - 2tsst \otimes 1 - 2st \otimes st + 1 \otimes stst$	N
st	tu	$tus \otimes t - tuts \otimes 1 + tust \otimes 1 - tu \otimes st - st \otimes tu + t \otimes t^{-1}sttu$	N
st	tu^{-1}	$tu^{-1}s \otimes t - tu^{-1}ts \otimes 1 + tu^{-1}st \otimes 1 - tu^{-1} \otimes st - st \otimes tu^{-1} + t \otimes t^{-1}sttu^{-1}$	Y
su	$u^{-1}tuv$	$t[u, s] \otimes v + t \otimes tuv u^{-1} t^{-1} [s, u] + tuv u^{-1} [s, u] \otimes 1$	Y
sv	tu	$2tu[s, v] \otimes 1 + t \otimes tut^{-1}[s, v] + ut[v, s] \otimes 1 + t \otimes [v, s]u + 1 \otimes [s, v]tu$	Y
tu	$stuvu^{-1}t^{-1}$	0	Y
a	b	$-ab \otimes 1$	N
a	bc	$-acb \otimes 1$	N
a	bcd	$-acdb \otimes 1$	N
ab	ab	$-2ab \otimes ab + 2abab \otimes 1$	Y
ab	bc	$-bcba \otimes 1 + cbba \otimes 1 - ab \otimes bc$	N
a	$aba^{-1}b^{-1}$	$-2ba \otimes b^{-1} + b \otimes b^{-1}a + baa \otimes a^{-1}b^{-1}$	Y

Example 2.8. Some values of $\delta_2^{\sigma, \nabla^c}$ are listed in Table 1. We put $a = \alpha_1, b = \beta_1, c = \alpha_2, \dots, s = \gamma_1, t = \gamma_2$ and so on, and omit the vertical bar $|$ for the trace space for the sake of legibility. $[x, y] = xy - yx$ is the usual commutator.

It yields a non-zero value for some pair (x, y) of disjoint loops, so $\delta_2^{\sigma, \nabla^c}$ cannot be written as a composite of a combination of the Goldman bracket and the Turaev cobracket. There are hints of topological features, such as disjointness, reflected in the values (e.g., the sum of coefficients or the number of terms), but none are particularly clear. For similar arguments involving cyclic words, the reader may want to compare it with a combinatorial description of the Goldman–Turaev Lie bialgebra [Cha04] and Counting Intersection Theorem for the TWG Lie algebra [CK21].

The original motivation to define the maps Div_k^∇ was to eliminate the choice of a connection in reformulating the divergence Div_1^∇ . Since we take the trace of c_∇ , the off-diagonal entries of the connection form of ∇ do not contribute to Div_1^∇ at all, and the author expected that the conditions on Div_k^∇ , to which they certainly contribute, will restrict the choice of a connection. Therefore, we ask the following:

Question 2.9. *Is there a (flat) connection ∇ on $\Omega^1 \mathbb{K}\pi$ that makes the operation $\delta_k^{\sigma, \nabla}$ (anti-)symmetric in its values?*

Remark 2.10. In the case of the tensor algebra $A = T(W)$, we have a canonical flat connection ∇_W explained below, and a similar statement holds, as we will see in the next section.

3. PROPS OF RIBBON GRAPHS

In this section, we will provide an explicit expression of Div_k^∇ in the case of tensor algebra and see a relation to operations associated with ribbon graphs introduced in [MW15] by Merkulov and Willwacher.

Definition 3.1. Let W be a finite-dimensional \mathbb{K} -vector space and $T(W)$ the tensor algebra on W . The canonical flat connection ∇_W on $\Omega^1 T(W)$ is defined by $\nabla_W(dw) = 0$ for all $w \in W$.

Definition 3.2. Let $\langle \cdot, \cdot \rangle: W \otimes W \rightarrow \mathbb{K}$ be a pairing. We extend the pairing to $T(W) \otimes T(W) \rightarrow T(W) \otimes T(W)$ by, for $u_i, w_j \in W$,

$$\langle u_1 \cdots u_r, w_1 \cdots w_s \rangle = \sum_{\substack{1 \leq i \leq r \\ 1 \leq j \leq s}} \langle u_i, w_j \rangle w_1 \cdots w_{j-1} u_{i+1} \cdots u_r \otimes u_1 \cdots u_{i-1} w_{j+1} \cdots w_s.$$

The associated *Hamiltonian flow* is defined by

$$\text{Ham}_{\langle \cdot, \cdot \rangle}: |T(W)| \rightarrow \text{End}_{\mathbb{K}}(T(W)) : a \mapsto \langle a, \cdot \rangle.$$

We can show that the image is contained in $\text{Der}_{\mathbb{K}}(T(W))$, and it is a Lie algebra morphism if the pairing is skew-symmetric.

Remark 3.3. The map $\langle \cdot, \cdot \rangle: T(W) \otimes T(W) \rightarrow T(W) \otimes T(W)$ satisfies the axiom of a *double bracket* introduced in [vdB08] by Van den Bergh. In general, a double bracket $\Pi: A \otimes A \rightarrow A \otimes A$ induces the associated hamiltonial flow $\text{Ham}_{\Pi}: |A| \rightarrow \text{Der}_{\mathbb{K}}(A)$. See [Tan24a] for the details.

The following result supports the validity of the construction in the last section:

Theorem 3.4. *Suppose that the pairing $\langle \cdot, \cdot \rangle: W \otimes W \rightarrow \mathbb{K}$ is skew-symmetric. Then, the map*

$$(-1)^k \delta_k^{\text{Ham}_{\langle \cdot, \cdot \rangle}, \nabla_W} : |T(W)|^{\otimes k} \rightarrow |T(W)|^{\otimes 2}$$

coincides with the operation given by the ribbon graph L_k in Figure 1 for each k .

The explanation and the proof of the proposition occupy the rest of the section.

Before the proof, we recall the definition of a ribbon graph and the construction of the associated operations. The notations below are aligned with the ones in [MW15].




Figure 1: The ribbon graph $L_k \in \mathcal{R}_{2,k}^k$. The labelling of the edges is suppressed, as any choice of these gives the same element in $\mathcal{R}Gra_1(2, k)$.

Definition 3.5.

- A *ribbon graph* is a tuple $\Gamma = (H, \tau_1, \tau_0, I)$ where H is a non-empty finite set of *half edges*, $\tau_1: H \rightarrow H$ is an involution (i.e., $\tau_1^2 = \text{id}$) with no fixing points, $\tau_0: H \rightarrow H$ is a bijection, and I is a (possibly empty) finite set of *isolated vertices*. The set of *vertices*, *edges* and *boundary components* are defined by

$$V(\Gamma) = (H/\tau_0) \sqcup I, \quad E(\Gamma) = H/\tau_1, \quad \text{and} \quad B(\Gamma) = H/(\tau_1^{-1} \circ \tau_0) \sqcup I.$$

- For a vertex $v \in V(\Gamma) \setminus I$, let $H(v) \subset H$ be the orbit of τ_0 representing v . This set is equipped with a natural cyclic order induced by τ_0 . The *valency* of the vertex is the cardinality of the set $H(v)$. If $v \in I$, the valency is defined to be zero.
- A *direction* of an edge $e = \{h, \tau_1(h)\} \in E(\Gamma)$ is a total order on the set $\{h, \tau_1(h)\}$. A *directed* ribbon graph is a ribbon graph with a choice of direction for each edge.
- A *labelling* of a ribbon graph Γ with n vertices, ℓ edges and m boundary components is a tuple of bijections

$$V(\Gamma) \rightarrow \{1, \dots, n\}, \quad E(\Gamma) \rightarrow \{1, \dots, \ell\} \quad \text{and} \quad B(\Gamma) \rightarrow \{\bar{1}, \dots, \bar{m}\}.$$

Boundary components are conventionally labelled with barred numbers \bar{k} .

- $\mathcal{R}_{m,n}^\ell$ is the set of (isomorphism classes of) all such directed labelled ribbon graphs.

A ribbon graph is depicted as a planar graph as in Figure 1, with the cyclic order at each vertex specified in counterclockwise rotation.

Finally, the prop $\mathcal{R}Gra_1$ is defined as follows: as a \mathbb{K} -vector space,

$$\mathcal{R}Gra_1(m, n) = \bigoplus_{\ell \geq 0} \mathbb{K} \mathcal{R}_{m,n}^\ell / (S_\ell \times S_2^{\times \ell})$$

for $m, n \geq 1$, where the symmetric group S_ℓ acts on it by the re-numbering of edges, and each copy of S_2 acts by the corresponding edge flip with the factor (-1) for each transposition. The prop structure is given by a composition of ribbon graphs, which is an analogue of surface glueing; we will not use the explicit formula below and refer to Section 4.2 of [MW15] for the detail.

The prop $\mathcal{R}Gra_1$ has a canonical representation on the space of cyclic words $|T(W)|$ over a vector space W with a skew-symmetric pairing (Theorem 4.2.2 of [MW15]), and we shall briefly recall it. With a ribbon graph $\Gamma \in \mathcal{R}_{m,n}^\ell$ is associated a \mathbb{K} -linear map $\Gamma: |T(W)|^{\otimes n} \rightarrow |T(W)|^{\otimes m}$ by the following procedure. To compute $\Gamma(\mathbf{w}_1, \dots, \mathbf{w}_n)$ for cyclic words $\mathbf{w}_i = w_1^{(i)} \cdots w_{r_i}^{(i)}$ with $w_j^{(i)} \in W$:

- (1) Take a cyclic-order preserving injection $q_i: H(i) \rightarrow \{1, \dots, r_i\}$ for each $i \in V(\Gamma)$. Here each vertex v is identified with its label i , and the set $\{1, \dots, r_i\}$ is equipped with the cyclic order $1 \leq 2 \leq \dots \leq r_i \leq 1$. This gives a cyclic arrangement of $w_j^{(i)}$'s around the vertex i with $w_{q_i(h)}^{(i)}$ on the half-edge h and other letters *in-between*. If not possible (i.e., if $\#H(i) > r_i$), the value of $\Gamma(\mathbf{w}_1, \dots, \mathbf{w}_n)$ is zero.

- (2) For each directed edge $e = \{h, \tau_1(h) : h \leq \tau_1(h)\}$ from a vertex i to the other i' in this direction, take a pairing $\langle w_{q_i(h)}^{(i)}, w_{q_{i'}(\tau_1(h))}^{(i')} \rangle$. Denote by λ the product of these scalars over all edges. The letters used in the pairing are thus *consumed*.
- (3) Obtain the cyclic words \mathbf{w}'_k by collecting the surviving letters $w_j^{(i)}$ along the k -th boundary of Γ .
- (4) $\Gamma(\mathbf{w}_1, \dots, \mathbf{w}_n)$ is the sum of $\lambda \cdot \mathbf{w}'_1 \otimes \dots \otimes \mathbf{w}'_m$ for all possible combination of injections $q = (q_i)_{1 \leq i \leq n}$.

Then, this only depends on the class in $\mathcal{RGr}_1(m, n)$ by the skew-symmetry of the pairing.

Moreover, there is also a canonical map $\text{Lie}^\diamond \rightarrow \mathcal{RGr}_1$ from the prop governing involutive Lie bialgebras given by

$$(\text{Lie bracket}) \mapsto \begin{array}{c} 1 \longrightarrow 2 \\ \downarrow \quad \uparrow \\ \bar{1} \end{array} \quad \text{and} \quad (\text{Lie cobracket}) \mapsto \bar{2} \begin{array}{c} \circlearrowleft \\ \bar{1} \end{array} \cdot 1 = L_1.$$

For the proof, see Theorem 4.2.3 of [MW15]. Therefore, an \mathcal{RGr}_1 -algebra can be considered as an enhancement of an involutive Lie bialgebra, and $|T(W)|$ is automatically endowed with an involutive Lie bialgebra structure. In general, for each integer d , we can define a \mathbb{Z} -graded prop \mathcal{RGr}_d with the canonical representation on a graded vector space with graded-symmetric pairing.

Proof of Proposition 3.4. The proof is done by direct computation. Take $\mathbf{w}_1, \dots, \mathbf{w}_k \in |T(W)|$ of the form $\mathbf{w}_i = w_1^{(i)} \dots w_{r_i}^{(i)}$ with $w_j^{(i)} \in W$. For $w \in W$, we have

$$\begin{aligned} \text{Ham}_{\langle \cdot, \cdot \rangle}(\mathbf{w}_i)(w) &= \sum_{1 \leq s_i \leq r_i} \langle w_{s_i}^{(i)}, w \rangle w_{s_i+1}^{(i)} \dots w_{s_i-1}^{(i)}, \\ c_{\nabla W}(\text{Ham}_{\langle \cdot, \cdot \rangle}(\mathbf{w}_i))(dw) &= -d \left(\sum_{1 \leq s_i \leq r_i} \langle w_{s_i}^{(i)}, w \rangle w_{s_i+1}^{(i)} \dots w_{s_i-1}^{(i)} \right) \\ &= - \sum_{\substack{1 \leq s_i \leq r_i \\ 1 \leq t_i \leq r_i \\ s_i \neq t_i}} \langle w_{s_i}^{(i)}, w \rangle w_{s_i+1}^{(i)} \dots w_{t_i-1}^{(i)} dw_{t_i}^{(i)} w_{t_i+1}^{(i)} \dots w_{s_i-1}^{(i)}, \end{aligned}$$

$$\begin{aligned} &c_{\nabla W}(\text{Ham}_{\langle \cdot, \cdot \rangle}(\mathbf{w}_1)) \circ \dots \circ c_{\nabla W}(\text{Ham}_{\langle \cdot, \cdot \rangle}(\mathbf{w}_k))(dw) \\ &= c_{\nabla W}(\text{Ham}_{\langle \cdot, \cdot \rangle}(\mathbf{w}_1)) \circ \dots \circ c_{\nabla W}(\text{Ham}_{\langle \cdot, \cdot \rangle}(\mathbf{w}_{k-1})) \left(- \sum_{\substack{1 \leq s_i \leq r_i \\ 1 \leq t_i \leq r_i \\ s_i \neq t_i}} \langle w_{s_i}^{(k)}, w \rangle w_{s_i+1}^{(k)} \dots w_{t_i-1}^{(k)} dw_{t_i}^{(k)} w_{t_i+1}^{(k)} \dots w_{s_i-1}^{(k)} \right) \\ &= \dots (\text{repeatedly applied}) \dots \\ &= \sum_{1 \leq i \leq k} \sum_{\substack{1 \leq s_i \leq r_i \\ 1 \leq t_i \leq r_i \\ s_i \neq t_i}} \langle w_{s_i}^{(1)}, w_{t_i}^{(2)} \rangle \dots \langle w_{s_i}^{(k)}, w \rangle (-1)^k \\ &\quad w_{s_k+1}^{(k)} \dots w_{t_k-1}^{(k)} \dots w_{s_2+1}^{(2)} \dots w_{t_2-1}^{(2)} w_{s_1+1}^{(1)} \dots w_{t_1-1}^{(1)} dw_{t_1}^{(1)} w_{t_1+1}^{(1)} \dots w_{s_1-1}^{(1)} w_{t_2+1}^{(2)} \dots w_{s_2-1}^{(2)} \dots w_{t_k+1}^{(k)} \dots w_{s_k-1}^{(k)}, \end{aligned}$$

and, by taking the trace, we have

$$\begin{aligned} (-1)^k \delta_k^{\text{Ham}_{\langle \cdot, \cdot \rangle}, \nabla W}(\mathbf{w}_1, \dots, \mathbf{w}_k) &= \sum_{1 \leq i \leq k} \sum_{\substack{1 \leq s_i \leq r_i \\ 1 \leq t_i \leq r_i \\ s_i \neq t_i}} \langle w_{s_i}^{(1)}, w_{t_i}^{(2)} \rangle \dots \langle w_{s_i}^{(k)}, w_{t_i}^{(1)} \rangle \\ &\quad |w_{s_k+1}^{(k)} \dots w_{t_k-1}^{(k)} \dots w_{s_2+1}^{(2)} \dots w_{t_2-1}^{(2)} w_{s_1+1}^{(1)} \dots w_{t_1-1}^{(1)}| \\ &\quad \otimes |w_{t_1+1}^{(1)} \dots w_{s_1-1}^{(1)} w_{t_2+1}^{(2)} \dots w_{s_2-1}^{(2)} \dots w_{t_k+1}^{(k)} \dots w_{s_k-1}^{(k)}|. \end{aligned}$$

which is exactly equal to $L_k(\mathbf{w}_1, \dots, \mathbf{w}_k)$ by the definition of the ribbon graph operation. \square

Remark 3.6. Having seen this, the original motivation (in addition to giving a constraint to the choice of connections) was to craft a family of loop operations, each corresponding to an element of $\mathcal{R}Gra_1$ being an extension of the Goldman–Turaev Lie bialgebra. The operations given by the graphs

$$\begin{array}{c} 1 \longrightarrow 2 \\ \downarrow \quad \uparrow \\ \text{1} \end{array} \quad \text{and} \quad \begin{array}{c} \bar{2} \\ \circlearrowleft \\ \text{1} \end{array}$$

are the associated graded of the Goldman bracket and the Turaev cobracket, respectively, for the surface with only one boundary component. For the reference, see the survey [KK16] by Kawazumi–Kuno, for example.

There were already some attempts by several people (and the author is, again, thankful to Sergei Merkulov for sharing their past attempts on it), but all were unfortunately unsuccessful; operations given by the smoothing–intersections type argument are not well-defined when two or more intersections are resolved at the same time. In contrast, our definition of the operations $\delta_k^{\psi, \nabla}$ are purely algebraic with a lack of topological viewpoint. The interpretation of these operations regarding the topology of curves would be an interesting topic.

Another result in their paper is that there is a map of props $\mathcal{H}oLieb_{0,1} \rightarrow \mathcal{R}Gra_1$ (Theorem 4.3.3 of [MW15]). It would also be interesting to investigate the $\mathcal{H}oLieb_{0,1}$ -relations in the case of the group algebra $\mathbb{K}\pi$.

4. NON-COMMUTATIVE DE RHAM COHOMOLOGY AND VECTOR BUNDLES

The quantity c_{∇} defined in the section 2 was essential for the construction. To see what it really is, we will develop a theory on a non-commutative version of de Rham cohomology with a vector bundle coefficient. We will relate the space of endomorphism-valued non-commutative differential forms to the Chevalley–Eilenberg (CE) complex over a derivation Lie algebra in Theorem 4.8 and show that the alternated version $\text{Div}_k^{\nabla} \circ \text{alt}$ is a k -cocycle in the CE complex.

Let B be a unital associative \mathbb{K} -algebra.

Definition 4.1. The *graded trace space* $\text{DR}^{\bullet}B = \Omega^{\bullet}B / [\Omega^{\bullet}B, \Omega^{\bullet}B]$ of $\Omega^{\bullet}B$ is the space of *de Rham differential forms*. An element of $\text{DR}^n B$ is of the form $|b_0 db_1 \cdots db_n|$ with the relation generated by

$$|b_0 db_1 \cdots db_n| = (-1)^{n-1} |db_2 \cdots db_n b_0 db_1|.$$

Being a graded derivation, the exterior derivative d descends to $\text{DR}^{\bullet}B$.

Definition 4.2. We define the space of *endomorphism-valued differential forms* by

$$\Omega^{\bullet}(B, \text{End } M) = \text{Hom}_B(M, \Omega^{\bullet}B \otimes_B M).$$

For a connection ∇ on a B -module M , the space $\Omega^{\bullet}(B, \text{End } M)$ is endowed with a degree +1 map D_{∇} given by the covariant derivative:

$$D_{\nabla} : \Omega^{\bullet}(B, \text{End } M) \rightarrow \Omega^{\bullet+1}(B, \text{End } M) : \xi \mapsto [\nabla, \xi] := \nabla \circ \xi - (-1)^{|\xi|} (\text{id} \otimes \xi) \circ \nabla.$$

The space $\Omega^{\bullet}(B, \text{End } M)$ is a (curved) differential graded Lie algebra with the graded commutator $[\xi, \eta] = (\text{id} \otimes \xi) \circ \eta - (-1)^{|\xi||\eta|} (\text{id} \otimes \eta) \circ \xi$, and $D_{\nabla}^2 = 0$ holds if ∇ is flat.

Remark 4.3. The space $\Omega^{\bullet}(B, \text{End } M)$ is not equal to the B -tensor product of $\Omega^{\bullet}B$ and $\text{End}_B(M)$: the latter does not even have a B -module structure unless B is commutative. Also, since we use the Koszul sign convention, we have $(\text{id} \otimes \xi) \circ \eta = (-1)^{|\xi||\eta|} \eta' \otimes (\xi \circ \eta'')$.

Definition 4.4. Let M be B -dualisable. The *trace map* on $\Omega^{\bullet}(B, \text{End } M)$ with values in the non-commutative de Rham complex $\text{DR}^{\bullet}B$ is defined by

$$\text{Tr} : \Omega^{\bullet}(B, \text{End } M) \rightarrow \text{DR}^{\bullet}B : \theta \otimes \omega \otimes m \mapsto |\theta(m)\omega|$$

using the natural surjection $|\Omega^{\bullet}B| \twoheadrightarrow \text{DR}^{\bullet}B$.

Proposition 4.5. *The trace map*

$$\mathrm{Tr} : (\Omega^\bullet(B, \mathrm{End} M), D_\nabla) \rightarrow (\mathrm{DR}^\bullet B, d)$$

is a (curved) differential graded Lie algebra homomorphism. Here $\mathrm{DR}^\bullet B$ is regarded as an abelian Lie algebra.

Proof. Putting $\nabla = \nabla' \otimes \nabla'' : M \rightarrow \Omega^1 B \otimes_B M$, one has

$$\begin{aligned} D_\nabla(\theta \otimes \omega \otimes m) &= [\nabla, \theta \otimes \omega \otimes m] \\ &= d\theta(\cdot)\omega \otimes m + \theta(\cdot)d\omega \otimes m + (-1)^{|\omega|}\theta(\cdot)\omega\nabla'(m) \otimes \nabla''(m) \\ &\quad - (-1)^{|\omega|+|\omega|}\nabla'(\cdot)\theta(\nabla''(\cdot))\omega \otimes m, \end{aligned}$$

so that

$$\begin{aligned} \mathrm{Tr} D_\nabla(\theta \otimes \omega \otimes m) &= |d\theta(m)\omega + \theta(m)d\omega + (-1)^{|\omega|}\theta(\nabla''(m))\omega\nabla'(m) - \nabla'(m)\theta(\nabla''(m))\omega| \\ &= d|\theta(m)\omega| = d\mathrm{Tr}(\theta \otimes \omega \otimes m), \end{aligned}$$

which shows that Tr intertwines the differentials. The proof that the trace map is a Lie algebra homomorphism is analogous to the proof that the trace map is cyclic in its argument. \square

Since de Rham cohomology is closely related to Chevalley–Eilenberg cohomology in the usual differential geometry, we shall now relate these in a non-commutative setting.

Definition 4.6.

- For a Lie algebra \mathfrak{g} over \mathbb{K} and a \mathfrak{g} -module V , $C_{\mathrm{CE}}^n(\mathfrak{g}, V) = \mathrm{Hom}_{\mathbb{K}}(\wedge^n \mathfrak{g}, V)$ is the Chevalley–Eilenberg complex with the differential

$$\begin{aligned} (d_{\mathrm{CE}}\psi)(x_0, \dots, x_n) &= \sum_{0 \leq i \leq n} (-1)^i x_i \cdot \psi(x_0, \dots, \check{x}_i, \dots, x_n) \\ &\quad + \sum_{0 \leq i < j \leq n} (-1)^{i+j} \psi([x_i, x_j], x_0, \dots, \check{x}_i, \dots, \check{x}_j, \dots, x_n). \end{aligned}$$

If V is an associative \mathbb{K} -algebra with a \mathfrak{g} -equivariant multiplication, the complex is a differential graded algebra with the shuffle product.

- For a derivation action $(\mathfrak{d}, \varphi, \rho)$ on a dualisable B -module M , the space $\mathrm{End}_B(M)$ is regarded as a \mathfrak{d} -module by $f \cdot \mu = [\rho(f), \mu]$ for $f \in \mathfrak{d}$ and $\mu \in \mathrm{End}_B(M)$.
- The maps

$$\iota : \Omega^\bullet(B, \mathrm{End} M) \rightarrow C_{\mathrm{CE}}^\bullet(\mathfrak{d}, \mathrm{End}_B(M)) \quad \text{and} \quad \iota : \mathrm{DR}^\bullet B \rightarrow C_{\mathrm{CE}}^\bullet(\mathfrak{d}, |B|)$$

are given by, for $f_i \in \mathfrak{d}$ and $\alpha \in \Omega^n(B, \mathrm{End} M)$ or $\mathrm{DR}^n B$,

$$\iota(\alpha)(f_1 \wedge \dots \wedge f_n) = i_{\varphi(f_n)} \dots i_{\varphi(f_1)} \alpha.$$

Here $i_f \otimes \mathrm{id}$ is briefly denoted by i_f .

Remark 4.7. The map c_∇ defined in Section 2 lives in $C_{\mathrm{CE}}^1(\mathfrak{d}, \mathrm{End}_B(M))$.

Theorem 4.8. *The following is a commutative diagram of curved complexes:*

$$\begin{array}{ccc} (\Omega^\bullet(B, \mathrm{End} M), D_\nabla) & \xrightarrow{\mathrm{Tr}} & (\mathrm{DR}^\bullet B, d) \\ \downarrow \iota & & \downarrow \iota \\ (C_{\mathrm{CE}}^\bullet(\mathfrak{d}, \mathrm{End}_B(M)), d_{\mathrm{CE}} + [c_\nabla, \cdot]) & \xrightarrow{\mathrm{Tr}} & (C_{\mathrm{CE}}^\bullet(\mathfrak{d}, |B|), d_{\mathrm{CE}}). \end{array}$$

We need some preparations for the proof.

Proposition 4.9. *Two ι 's are cochain maps.*

Proof. We only check the left one since the right one is easier. For $\theta \otimes \omega \otimes m \in M^* \otimes_B \Omega^n B \otimes_B M$, we have (compare with the computation in the proof of Proposition 4.5), denoting $i_{\varphi(f)}$ simply by i_f ,

$$\begin{aligned} & \iota(D_{\nabla}(\theta \otimes \omega \otimes m))(f_0 \wedge \cdots \wedge f_n) \\ &= i_{f_n} \cdots i_{f_0}(d\theta \otimes m + \theta d\omega \otimes m + (-1)^n \theta \omega \nabla m - (\theta \circ \nabla) \cdot \omega \otimes m) \\ &= \sum_{0 \leq j \leq n} (-1)^j (f_j \circ \theta) \cdot i_{f_n} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_0} \omega \cdot m + \theta \cdot i_{f_n} \cdots i_{f_0} d\omega \cdot m \\ & \quad + \sum_{0 \leq j \leq n} \left((-1)^{n+(n-j)} \theta \cdot i_{f_n} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_0} \omega \cdot i_{f_j} \nabla m - (-1)^j (\theta \circ i_{f_j} \nabla) \cdot i_{f_n} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_0} \omega \otimes m \right). \end{aligned}$$

On the other hand, we have

$$\begin{aligned} & ((d_{\text{CE}} + [c_{\nabla}, \cdot]) \circ \iota)(\theta \otimes \omega \otimes m)(f_0 \wedge \cdots \wedge f_n) \\ &= \sum_{0 \leq j \leq n} (-1)^j [f_j, \theta \cdot i_{f_n} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_0} \omega \cdot m] + \sum_{0 \leq j < k \leq n} (-1)^{j+k} \theta \cdot i_{f_n} \cdots i_{f_k} \check{i}_{f_k} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_0} i_{[f_j, f_k]} \omega \cdot m \\ & \quad + \sum_{0 \leq j \leq n} (-1)^j [c_{\nabla}(f_j), \theta \cdot i_{f_n} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_0} \omega \cdot m] \\ &= \sum_{0 \leq j \leq n} (-1)^j [i_{f_j} \nabla, \theta \cdot i_{f_n} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_0} \omega \cdot m] + \sum_{0 \leq j < k \leq n} (-1)^{j+k} \theta \cdot i_{f_n} \cdots i_{f_k} \check{i}_{f_k} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_0} i_{[f_j, f_k]} \omega \cdot m \\ &= \sum_{0 \leq j \leq n} (-1)^j \left((f_j \circ \theta) \cdot i_{f_n} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_0} \omega \cdot m + \theta \cdot L_{f_j}(i_{f_n} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_0} \omega) \cdot m \right. \\ & \quad \left. + \theta \cdot i_{f_n} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_0} \omega \cdot i_{f_j} \nabla m - (\theta \circ i_{f_j} \nabla) \cdot i_{f_n} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_0} \omega \cdot m \right) \\ & \quad + \sum_{0 \leq j < k \leq n} (-1)^{j+k} \theta \cdot i_{f_n} \cdots i_{f_k} \check{i}_{f_k} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_0} i_{[f_j, f_k]} \omega \cdot m. \end{aligned}$$

Finally, we are done if we prove, on $\Omega^n B$,

$$i_{f_n} \cdots i_{f_0} d = \sum_{0 \leq j \leq n} (-1)^j L_{f_j} i_{f_n} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_0} + \sum_{0 \leq j < k \leq n} (-1)^{j+k} i_{f_n} \cdots i_{f_k} \check{i}_{f_k} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_0} i_{[f_j, f_k]}.$$

We proceed by induction. If $n = 0$, this is equivalent to $i_{f_0} d = L_{f_0}$, which is true because $\Omega^0 B$ is simply equal to B . For $n \geq 1$, by the induction hypothesis,

$$\begin{aligned} i_{f_n} \cdots i_{f_0} d &= i_{f_n} \cdots i_{f_1} L_{f_0} - (i_{f_n} \cdots i_{f_1} d) i_{f_0} \\ &= L_{f_0} i_{f_n} \cdots i_{f_1} + \sum_{1 \leq k \leq n} i_{f_n} \cdots i_{[f_k, f_0]} \cdots i_{f_1} \\ & \quad - \left(\sum_{1 \leq j \leq n} (-1)^{j-1} L_{f_j} i_{f_n} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_1} + \sum_{1 \leq j < k \leq n} (-1)^{j+k} i_{f_n} \cdots i_{f_k} \check{i}_{f_k} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_1} i_{[f_j, f_k]} \right) i_{f_0} \\ &= \sum_{0 \leq j \leq n} (-1)^j L_{f_j} i_{f_n} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_0} + \sum_{0 \leq j < k \leq n} (-1)^{j+k} i_{f_n} \cdots i_{f_k} \check{i}_{f_k} \cdots i_{f_j} \check{i}_{f_j} \cdots i_{f_0} i_{[f_j, f_k]}. \end{aligned}$$

This completes the proof. \square

Proof of Theorem 4.8. The square is commutative since contractions and trace maps are mutually compatible. Furthermore, since M is dualisable, $\text{Tr} : \text{End}_B(M) \rightarrow |B|$ is automatically a $\text{Der}_{\mathbb{K}}(B)$ -module map; hence is also a \mathfrak{d} -module map. Therefore, Tr commutes with d_{CE} . For any ψ , we have

$$\text{Tr}(d_{\text{CE}}(\psi) + [c_{\nabla}, \psi]) = \text{Tr}(d_{\text{CE}}(\psi)) = d_{\text{CE}} \text{Tr}(\psi).$$

This shows that Tr in the bottom row is a cochain map. Finally, the proposition above ensures that ι 's are cochain maps. \square

The following is the key property of the map c_{∇} .

Proposition 4.10. *If ∇ is flat, c_∇ is a Maurer–Cartan element with respect to the differential d_{CE} . Therefore, $d_{\text{CE}} + [c_\nabla, \cdot]$ is also a differential.*

Proof. Let $R = \nabla^2$ be the curvature. For a general connection ∇ , we have

$$(d_{\text{CE}}c_\nabla)(f \wedge g) + [c_\nabla(f), c_\nabla(g)] = \iota(R)(f \wedge g).$$

The proof is essentially the same as Proposition 4.11 in [Tan24b]. By the definition of the shuffle product, $[c_\nabla(f), c_\nabla(g)]$ is equal to $(c_\nabla \circ c_\nabla)(f \wedge g)$; this implies $d_{\text{CE}}c_\nabla + c_\nabla \circ c_\nabla = \iota(R)$, which is zero if ∇ is flat. \square

We will now look at the map Div_k^∇ with these tools available. To apply the above, we only consider the alternated version $\text{Div}_k^\nabla \circ \text{alt}$, where alt is defined by

$$\text{alt}(f_1 \wedge \dots \wedge f_k) = \sum_{s \in S_k} (-1)^s f_{s(1)} \otimes \dots \otimes f_{s(k)}.$$

Lemma 4.11. *The map $\text{Div}_k^\nabla \circ \text{alt}: \wedge^k \mathfrak{d} \rightarrow |B|$ is equal to $\text{Tr}(c_\nabla^k) \in C_{\text{CE}}^k(\mathfrak{d}, |B|)$.*

Proof. This follows from the definition of Div_k^∇ . \square

Corollary 4.12. *For k even, $\text{Div}_k^\nabla \circ \text{alt} = \text{Tr}(c_\nabla^k)$ is zero.*

Proof. By the cyclic symmetry of the trace, we have

$$\text{Tr}(c_\nabla^k) = \text{Tr}(c_\nabla \circ c_\nabla^{k-1}) = (-1)^{k-1} \text{Tr}(c_\nabla^{k-1} \circ c_\nabla) = -\text{Tr}(c_\nabla^k).$$

This completes the proof. \square

Theorem 4.13. *If ∇ is flat, the map $\text{Div}_k^\nabla \circ \text{alt} \in C_{\text{CE}}^k(\mathfrak{d}, |B|)$ is a Lie algebra k -cocycle. If k is even, the cocycle vanishes.*

Proof. If k is even, the map $\text{Div}_k^\nabla \circ \text{alt}$ itself is zero. Suppose k is odd. Since $(d_{\text{CE}} + [c_\nabla, \cdot])(c_\nabla) = c_\nabla^2$ and $d_{\text{CE}} + [c_\nabla, \cdot]$ is a degree-1 derivation of the Chevalley-Eilenberg complex, which is a differential graded algebra, we have

$$(d_{\text{CE}} + [c_\nabla, \cdot])(c_\nabla^k) = \sum_{1 \leq i \leq k} (-1)^{i-1} c_\nabla^{(i-1)+2+(k-i)} = c_\nabla^{k+1}.$$

Taking the trace, we obtain $d_{\text{CE}}(\text{Tr}(c_\nabla^k)) = \text{Tr}(c_\nabla^{k+1})$ in $C_{\text{CE}}^{k+1}(\mathfrak{d}, |B|)$, which is zero by the corollary above. \square

Remark 4.14. By the theorem above, the pullback $\delta_k^{\psi, \nabla}$ is also a k -cocycle. In the case of the tensor algebra $T(W)$, an alternative proof of the map $\delta_k^{\text{Ham}(\cdot, \cdot), \nabla_W}$ in Proposition 3.4 being a Lie algebra cocycle for $k \equiv 1 \pmod{4}$ is available in the framework of the ribbon graph complex by Merkulov and Willwacher [MW15]. First, using the notation there, the ribbon graph complex is given as the deformation complex

$$\text{RGC}_{1,1} := \text{Def}(\text{Lieb} \xrightarrow{s_1} \text{RGr}_1)$$

associated with the properad morphism s_1 given by

$$(\text{Lie bracket}) \mapsto \begin{array}{c} 1 & \xrightarrow{2} & 2 \\ \bullet & & \bullet \\ \downarrow & & \downarrow \\ & 1 & \end{array} \quad \text{and} \quad (\text{Lie cobracket}) \mapsto 0.$$

This complex is automatically a differential graded Lie algebra with differential induced by s_1 , and it is naturally isomorphic to the space of all infinite series of directed labelled ribbon graphs with their vertex- and boundary-labelling skew-symmetrised. Furthermore, the differential is pictorially given by the vertex expansion, which increases the number of edges by exactly one and preserves bivalent vertices. In addition, the element corresponding to the graph L_k is non-zero if and only if $k \equiv 1 \pmod{4}$. From these facts, it is readily seen that the differential of L_k is always zero.

Finally, the functorial property of the deformation complex gives the push-out map

$$\mathrm{RGC}_{1,1} = \mathrm{Def}(\mathcal{L}ieb \xrightarrow{s_1} \mathcal{R}Gra_1) \rightarrow \mathrm{Def}(\mathcal{L}ieb \rightarrow \mathcal{E}nd_{|T(W)|}) = \hat{C}_{\mathrm{CE}}^\bullet(|T(W)|, \wedge |T(W)|),$$

which is a differential graded Lie algebra morphism. Here, the CE complex in the last term is suitably completed. Under this morphism, the element L_k is sent to $\mathrm{alt} \circ \delta_k^{\mathrm{Ham}(\cdot, \cdot), \nabla_W}$, which ensures its cocycle property. When $k \equiv 1 \pmod{4}$, L_k is already skew-symmetric and therefore is sent to $\delta_k^{\mathrm{Ham}(\cdot, \cdot), \nabla_W}$.

We refer to [MV09] for a detailed exposition on deformation complexes.

In the case of $A = T(W)$, the cocycles $\mathrm{Tr}(c_{\nabla_W}^k)$ are the essential ones in the following sense:

Theorem 4.15. *Let $D = \mathrm{Der}_{\mathbb{K}}(T(W))^{(\geq 0)}$ be the space of derivations with non-negative degree, and put $V = |T(W)|$, which is a D -module. Then, we have a canonical isomorphism*

$$H_{\mathrm{CE}}^\bullet(\mathfrak{gl}(W), \mathbb{K}) \cong H_{\mathrm{CE}}^\bullet(D, V),$$

under which the cohomology class of $(-1)^k \mathrm{Tr}(c_{\nabla_W}^k)$ for k odd is sent to the standard generator of the cohomology ring $H_{\mathrm{CE}}^\bullet(\mathfrak{gl}(W), \mathbb{K})$ of degree k .

We need the following lemma. The statement is standard, but we repeat the argument for clarity.

Lemma 4.16. *Let $\iota: S \hookrightarrow R$ and $\pi: R \rightarrow S$ be \mathbb{K} -algebra maps with $\pi \circ \iota = \mathrm{id}$. Put $I = \mathrm{Ker} \pi$. Then, we have an adjunction*

$$F: R\text{-Mod} \rightleftarrows S\text{-Mod}: G$$

given by $F(V) = \iota^*(V/IV)$ and $G(N) = \pi^*N$.

Proof. We construct the universal maps $\eta: V \rightarrow GF(V)$ and $\varepsilon: FG(N) \rightarrow N$. First, we have $GF(V) = (\iota \circ \pi)^*(V/IV)$, so we define $\eta: V \rightarrow (\iota \circ \pi)^*(V/IV)$ to be the natural projection. This is an R -module map: for $r \in R$, it is uniquely written as $r = \iota(s) + z$ with $s \in S$ and $z \in I$, so that, for $v \in V$,

$$\eta(rv) = \eta(\iota(s)v + zv) = [\iota(s)v] = \iota(s)\eta(v) = (\iota \circ \pi)(\iota(s) + z)\eta(v) = (\iota \circ \pi)(r)\eta(v)$$

holds in V/IV . Next, we have $FG(N) = \iota^*((\pi^*N)/(I\pi^*N)) = N$ since $I\pi^*N = 0$ and $\pi \circ \iota = \mathrm{id}$. We define ε to be the identity map. Then, we have $\varepsilon \circ F(\eta) = \mathrm{id}$ and $\eta \circ G(\varepsilon) = \mathrm{id}$ due to the isomorphism $FG \cong \mathrm{id}$. \square

Corollary 4.17. *Let \mathfrak{g} be a non-negatively graded \mathbb{K} -Lie algebra, and denote by $\mathfrak{g}^{(0)}$ the degree-0 part. For a $\mathfrak{g}^{(0)}$ -module N , we have $H_{\mathrm{CE}}^\bullet(\mathfrak{g}^{(0)}, N) \cong H_{\mathrm{CE}}^\bullet(\mathfrak{g}, \pi^*N)$.*

Proof. Set $R = U\mathfrak{g}$, $S = U\mathfrak{g}^{(0)}$, ι the natural inclusion, π the natural projection and $V = \mathbb{K}$: the trivial $U\mathfrak{g}$ -module. By the lemma above, we have an isomorphism of functors

$$\mathrm{Hom}_{U\mathfrak{g}^{(0)}}(\mathbb{K}, -) \cong \mathrm{Hom}_{U\mathfrak{g}}(\mathbb{K}, \pi^*(-)).$$

because $\iota^*(\mathbb{K}/I\mathbb{K})$ is just the trivial $U\mathfrak{g}^{(0)}$ -module \mathbb{K} . Therefore, their derived functors are isomorphic. Note that the right-hand side is a composition $\mathrm{Hom}_{U\mathfrak{g}}(\mathbb{K}, -) \circ \pi^*$, and π^* is exact by definition. This implies the derived functor of the right-hand side is just $\mathbb{R}\mathrm{Hom}_{U\mathfrak{g}}(\mathbb{K}, -) \circ \pi^*$. Now we have

$$H_{\mathrm{CE}}^\bullet(\mathfrak{g}^{(0)}, N) = \mathrm{Ext}_{U\mathfrak{g}^{(0)}}^\bullet(\mathbb{K}, N) \cong \mathrm{Ext}_{U\mathfrak{g}}^\bullet(\mathbb{K}, \pi^*N) = H_{\mathrm{CE}}^\bullet(\mathfrak{g}, \pi^*N).$$

This completes the proof. \square

Proof of Theorem 4.15. We have a natural map of complexes

$$j: C_{\mathrm{CE}}^\bullet(\mathfrak{gl}(W), \mathbb{K}) \hookrightarrow C_{\mathrm{CE}}^\bullet(D, V)$$

by the identifications $\mathfrak{gl}(W) = D^{(0)} \subset D$ and $\mathbb{K} = V^{(0)} \subset V$. In addition, the complex $C_{\mathrm{CE}}^\bullet(D, V)$ is naturally \mathbb{Z} -graded and the differential d_{CE} has degree 0.

Now, we set up a spectral sequence. Define a decreasing filtration on C^\bullet by, for $p \in \mathbb{Z}$,

$$F^p C^n = \bigoplus_{m \geq p} \text{Hom}_{\mathbb{K}}((\wedge^n D)^{(m)}, V).$$

We can check $\bigcap_p F^p C^n = 0$ and $d_{\text{CE}}(F^p C^n) \subset F^p C^{n+1}$ by using the fact that D is non-negatively graded. The E_0 -page is given by

$$E_0^{p,q} = F^p C^{p+q} / F^{p+1} C^{p+q} \cong \text{Hom}_{\mathbb{K}}((\wedge^{p+q} D)^{(p)}, V).$$

We compute the induced differential $d_0^{p,q} : E_0^{p,q} \rightarrow E_0^{p,q+1}$. For $\psi \in E_0^{p,q}$ and $f_0, \dots, f_{p+q} \in D$ with $\sum_{0 \leq i \leq p+q} \deg(f_i) = p$, we have

$$\begin{aligned} d_0^{p,q}(\psi)(f_0, \dots, f_{p+q}) &= \sum_{0 \leq i \leq p+q} (-1)^i f_i \cdot \psi(f_0, \dots, \check{f}_i, \dots, f_{p+q}) \\ &\quad + \sum_{0 \leq i < j \leq p+q} (-1)^{i+j} \psi([f_i, f_j], f_0, \dots, \check{f}_i, \dots, \check{f}_j, \dots, f_{p+q}). \end{aligned}$$

If $\deg(f_{i_0}) > 0$, the term $f_{i_0} \cdot \psi(f_0, \dots, \check{f}_{i_0}, \dots, f_{p+q})$ is zero since we have $\sum_{\substack{0 \leq i \leq p+q \\ i \neq i_0}} \deg(f_i) < p$. This is re-stated as that the action of f_i with $\deg(f_i) > 0$ on the space V looks trivial. More precisely, we have

$$\left(\bigoplus_{p+q=\bullet} E_0^{p,q}, d_0 \right) \cong (C_{\text{CE}}^\bullet(D, \pi^* \iota^* V), d_{\text{CE}})$$

as complexes. By Corollary 4.17, we obtain $\bigoplus_{p+q=\bullet} E_1^{p,q} \cong H_{\text{CE}}^\bullet(D, \pi^* \iota^* V) \cong H_{\text{CE}}^\bullet(D^{(0)}, \iota^* V)$.

The centre of $D^{(0)} = \mathfrak{gl}(W)$ is isomorphic to \mathbb{K} and spanned by the Euler operator eu , which is a unique derivation on $T(W)$ specified by $\text{eu}(w) = w$ for all $w \in W$. It is acting on V by multiplication-by-degree. Therefore, the ‘‘centre kills’’ argument shows $H_{\text{CE}}^\bullet(D^{(0)}, \iota^* V) \cong H_{\text{CE}}^\bullet(D^{(0)}, (\iota^* V)^{(0)}) = H_{\text{CE}}^\bullet(\mathfrak{gl}(W), \mathbb{K})$. The map j induces an isomorphism on E_1 . The only surviving terms in the E_1 page are $E_1^{0,q}$, so the spectral sequence converges, and j is a quasi-isomorphism.

Next, the result of Fuks (Theorem 2.1.6 of [Fuk86]) states that the cohomology ring $H_{\text{CE}}^\bullet(\mathfrak{gl}(W), \mathbb{K})$ is isomorphic to the exterior algebra generated by φ_k for $1 \leq k \leq 2 \dim(W) - 1$ and k odd, where

$$\varphi_k : \wedge^k \mathfrak{gl}(W) \rightarrow \mathbb{K} : A_1 \wedge \dots \wedge A_k \rightarrow \sum_{s \in S_k} (-1)^s \text{Tr}(A_{s(1)} \dots A_{s(k)}).$$

For $f \in D^{(0)} = \mathfrak{gl}(W)$, we have

$$(c_{\nabla_W}(f))(dw) = i_f \nabla_W(dw) - L_f(dw) = -df(w).$$

Since f is just a matrix over \mathbb{K} , the endomorphism $c_{\nabla_W}(f) \in \text{End}_{T(W)^e} \Omega^1 T(W)$ is represented by the matrix $-f$. Therefore, we have

$$\text{Tr}(c_{\nabla_W}^k)(f_1 \wedge \dots \wedge f_k) = (-1)^k \varphi_k(f_1 \wedge \dots \wedge f_k)$$

for $f_i \in D^{(0)}$, which implies $j(\varphi_k) = (-1)^k \text{Tr}(c_{\nabla_W}^k)$. \square

For many classical results on the commutative counterpart, see Chapter 2 of the book [Fuk86] by Fuks. For the cohomology of the derivation Lie algebra on a free algebra over an (augmented) operad with various other coefficients, see the paper [Dot23] by Dotsenko.

APPENDIX A. THE INFINITESIMAL ACTION AND DIVERGENCES

This section is devoted to the further study of the interaction between the map c_{∇} and a derivation action. Let B be a \mathbb{K} -algebra and M a left B -module. Recall the following in [Tan24b], Section 7:

Definition A.1. A *homological connection* on M is a triple $\nabla = (P_\bullet, \partial_\bullet, \nabla_\bullet)$ where $P = (P_\bullet, \partial_\bullet)$ is a projective resolution of M and ∇_i is a connection on P_i . A homological connection ∇ is said to be flat if each ∇_i is flat.

Now, we introduce the infinitesimal action.

Definition A.2. Let $(\mathfrak{d}, \varphi, \rho)$ be a derivation action on M . For $f \in \mathfrak{d}$, we define the adjoint action of f on ∇ as follows. Firstly, choose a lift $\lambda[f]$ of $\rho(f): M \rightarrow M$ to the projective resolution P once and for all, so that $\lambda[f]$ is a $\varphi(f)$ -derivation and the association $f \mapsto \lambda[f]$ is \mathbb{K} -linear. We define the adjoint action by

$$\text{ad}_f \nabla = \{(L_{\varphi(f)} \otimes \text{id} + \text{id} \otimes \lambda[f]_i) \circ \nabla_i - \nabla_i \circ \lambda[f]_i: P_i \rightarrow \Omega^1 B \otimes_B P_i\}_{i \geq 0}.$$

Each map is an element of $\Omega^1(B, \text{End } P_i)$.

Proposition A.3. $\text{Tr}(\text{ad}_f \nabla) \in |B|$ is independent of the choice of a lift.

Proof. If we take another lift $\lambda'[f]$, the difference is B -linearly null-homotopic: $\lambda[f] - \lambda'[f] = [\partial, h]$ for some $h \in \text{End}_B^{(1)}(P)$. Then, we have

$$\begin{aligned} & \text{Tr}\{(L_{\varphi(f)} \otimes \text{id} + \text{id} \otimes \lambda[f]) \circ \nabla - \nabla \circ \lambda[f]\} - \text{Tr}\{(L_{\varphi(f)} \otimes \text{id} + \text{id} \otimes \lambda'[f]) \circ \nabla - \nabla \circ \lambda'[f]\} \\ &= \text{Tr}([\partial, h], \nabla) \\ &= \text{Tr}([\partial, \nabla], h) + [\partial, [h, \nabla]]. \end{aligned}$$

Since $[\partial, \nabla]$ and $[h, \nabla]$ are B -linear, $\text{Tr}([\partial, \nabla], h) + [\partial, [h, \nabla]]$ vanishes due to the cyclic invariance of the trace. \square

Remark A.4. Note that ∇ is not B -linear, so we cannot deduce $\text{Tr}([\partial, h], \nabla) = 0$ immediately.

Now, we see the relation between the adjoint action and the divergence.

Proposition A.5. Let $R = \nabla^2 \in \Omega^2(M, \text{End } P)$ be the curvature. Then, we have $\text{ad}_f \nabla = D_\nabla(c_\nabla(f)) + i_{\varphi(f)} R$ in $\Omega^1(B, \text{End } P)$ for $f \in \mathfrak{d}$.

Proof. What we have to show is the equality

$$(L_{\varphi(f)} \otimes \text{id} + \text{id} \otimes \lambda[f]) \circ \nabla - \nabla \circ \lambda[f] = [\nabla, i_{\varphi(f)} \nabla - \lambda[f]] + i_{\varphi(f)} R.$$

The terms involving $\lambda[f]$ can be cancelled out, and the remaining is $(L_{\varphi(f)} \otimes \text{id}) \circ \nabla = [\nabla, i_{\varphi(f)} \nabla] + i_{\varphi(f)} R$. Denoting $\varphi(f)$ by f , we have, for $p \in P$,

$$\begin{aligned} & ((L_f \otimes \text{id}) \circ \nabla - [\nabla, i_f \nabla])(p) \\ &= L_f(\nabla'(p)) \otimes \nabla''(p) - \nabla(i_f \nabla(p)) + (\text{id} \otimes i_f \nabla)(\nabla(p)) \\ &= L_f(\nabla'(p)) \otimes \nabla''(p) - \nabla(i_f \nabla'(p) \cdot \nabla''(p)) + (\text{id} \otimes i_f \nabla)(\nabla'(p) \otimes \nabla''(p)) \\ &= L_f(\nabla'(p)) \otimes \nabla''(p) - di_f(\nabla'(p)) \cdot \nabla''(p) - i_f \nabla'(p) \cdot \nabla(\nabla''(p)) + \nabla'(p) \cdot i_f \nabla(\nabla''(p)) \\ &= i_f d(\nabla'(p)) \cdot \nabla''(p) - i_f(\nabla'(p) \cdot \nabla(\nabla''(p))) \\ &= i_f(\nabla^2(p)) \\ &= i_f(R(p)). \end{aligned}$$

This completes the proof. \square

Corollary A.6. If ∇ is trace-flat, i.e., $\text{Tr}(R) = 0$, we have $\text{Tr}(\text{ad}_f \nabla) = d(\text{Div}^{(\nabla, \varphi, \rho)}(f))$ in $\text{DR}^1 B$ for $f \in \mathfrak{d}$.

Proof. This follows from the above proposition that Tr is a cochain map. \square

Remark A.7. For an oriented Riemannian manifold X , the divergence $\text{div}_\mu(\xi)$ of a vector field ξ on X is a change of a fixed volume form μ by the Lie derivative: $\text{div}_\mu(\xi)\mu = L_\xi(\mu)$. This is recovered from our definition of the divergence by setting $\mathbb{K} = \mathbb{R}$, $B = C^\infty(X)$, $M = \wedge^{\text{top}} T^*X$, $\mathfrak{d} = \text{Der}(C^\infty(X))$,

$$\varphi = \text{id}: \mathfrak{d} \rightarrow \text{Der}_{\mathbb{K}}(B),$$

$$\begin{aligned}\rho &= L: \mathfrak{d} \rightarrow \text{End}_{\mathbb{K}}(M), \text{ and} \\ \nabla &= \nabla_{\mu}: M \rightarrow \Omega^1 B \otimes_B M: \mu \mapsto 0.\end{aligned}$$

Here, the flat connection ∇_{μ} should be regarded as an incarnation of the volume form μ . The Proposition above says that, under the assumption of the flatness, the infinitesimal change of the connection (conversely thought of as a non-commutative volume form) by a vector field f roughly gives the divergence as one may expect.

The map c_{∇} also naturally emerges from the infinitesimal adjoint action.

Proposition A.8. *For any homological connection ∇ and $f \in \mathfrak{d}$, $\iota(\text{ad}_f \nabla)$ is equal to*

$$\text{ad}_f(c_{\nabla}) := [\lambda[f], c_{\nabla}] - c_{\nabla} \circ \text{ad}_f$$

in $C_{\text{CE}}^1(\mathfrak{d}, \text{End}_B(P))$ up to homotopy term.

Proof. For $g \in \mathfrak{d}$, we have

$$\begin{aligned}\iota(\text{ad}_f \nabla)(g) &= i_g((L_f \otimes \text{id} + \text{id} \otimes \lambda[f]) \circ \nabla - \nabla \circ \lambda[f]) \\ &= (i_g L_f \otimes \text{id} + i_g \otimes \lambda[f]) \circ \nabla - i_g \nabla \circ \lambda[f].\end{aligned}$$

On the other hand, we have

$$\begin{aligned}([\lambda[f], c_{\nabla}] - c_{\nabla} \circ \text{ad}_f)(g) &= [\lambda[f], i_g \nabla - \lambda[g]] - (i_{[f,g]} \nabla - \lambda[[f, g]]) \\ &= [\lambda[f], i_g \nabla] - i_{[f,g]} \nabla + (\lambda[[f, g]] - [\lambda[f], \lambda[g]]).\end{aligned}$$

The last two terms $\lambda[[f, g]] - [\lambda[f], \lambda[g]]$ is equal to $[\partial, h]$ for some $h: \mathfrak{d} \rightarrow \text{End}_B^{(-1)}(P)$ since it is the difference of the lifts of the same map $\rho([f, g]) = [\rho(f), \rho(g)]$. Therefore,

$$\begin{aligned}(\iota(\text{ad}_f \nabla) - \text{ad}_f(c_{\nabla}))(g) &= (i_g L_f \otimes \text{id} + i_g \otimes \lambda[f]) \circ \nabla - i_g \nabla \circ \lambda[f] - [\lambda[f], i_g \nabla] + i_{[f,g]} \nabla - [\partial, h] \\ &= (L_f i_g \otimes \text{id} + i_g \otimes \lambda[f]) \circ \nabla - \lambda[f] \circ i_g \nabla - [\partial, h] \\ &= -[\partial, h].\end{aligned}$$

The last equality comes from the assumption that $\lambda[f]$ is a $\varphi(f)$ -derivation. □

REFERENCES

- [AKKN23] Anton Alekseev, Nariya Kawazumi, Yusuke Kuno, and Florian Naef. The Goldman-Turaev Lie bialgebra and the Kashiwara-Vergne problem in higher genera. 2023. [arXiv:1804.09566v3](#). 1, 2, 7
- [ANPŠ24] Anton Alekseev, Florian Naef, Ján Pulmann, and Pavol Ševera. Batalin-Vilkovisky structures on moduli spaces of flat connections. *Advances in Mathematics*, 443:109580, 2024. [arXiv:2210.08944](#), [doi:10.1016/j.aim.2024.109580](#). 1
- [Cha04] Moira Chas. Combinatorial Lie bialgebras of curves on surfaces. *Topology*, 43(3):543–568, 2004. [arXiv:math/0105178](#). 2, 8
- [CK21] Moira Chas and Arpan Kabiraj. The Lie bracket of undirected closed curves on a surface. *Transactions of the American Mathematical Society*, 375(4):2365–2386, 2021. [arXiv:1910.08991](#), [doi:10.1090/tran/8541](#). 2, 8
- [Dot23] Vladimir Dotsenko. Stable homology of Lie algebras of derivations and homotopy invariants of wheeled operads. 2023. [arXiv:2311.18594](#). 4
- [Fuk86] Dmitry Borisovich Fuks. *Cohomology of Infinite-Dimensional Lie Algebras*. Springer US, 1986. [doi:10.1007/978-1-4684-8765-7](#). 1, 4
- [Gin05] Victor Ginzburg. Lectures on noncommutative geometry. 2005. [arXiv:math/0506603](#). 2
- [Gol86] William M. Goldman. Invariant functions on Lie groups and Hamiltonian flows of surface group representations. *Inventiones mathematicae*, 85(2):263–302, 1986. [doi:10.1007/BF01389091](#). 1
- [KK14] Nariya Kawazumi and Yusuke Kuno. The logarithms of Dehn twists. *Quantum Topology*, 5(3):347–423, 2014. [arXiv:1008.5017v1](#), [doi:10.4171/QT/54](#). 1

- [KK16] Nariya Kawazumi and Yusuke Kuno. *The Goldman–Turaev Lie bialgebra and the Johnson homomorphisms*, volume V of *Handbook of Teichmüller Theory*, pages 97–165. EMS Press, 2016. [arXiv:1304.1885](#), [doi:10.4171/160](#). 1, 3.6
- [MV09] Sergei Merkulov and Bruno Vallette. Deformation theory of representations of prop(erad)s II. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 2009(636), 2009. [arXiv:0707.0889v4](#), [doi:10.1515/crelle.2009.084](#). 4.14
- [MW15] Sergei Merkulov and Thomas Willwacher. Props of ribbon graphs, involutive Lie bialgebras and moduli spaces of curves. 2015. [arXiv:1511.07808](#). 1, 3, 3, 3, 3, 3.6, 4.14
- [Tan24a] Toyo Taniguchi. Modular vector fields in non-commutative geometry. 2024. [arXiv:2410.24064](#). 3.3
- [Tan24b] Toyo Taniguchi. Non-commutative divergence and the Turaev cobracket. *To appear in Algebraic & Geometric Topology*, 2024. [arXiv:2403.16566](#). 1, 2, 2.5, 2.7, 4, A
- [Tur91] Vladimir G. Turaev. Skein quantization of Poisson algebras of loops on surfaces. *Annales scientifiques de l'École Normale Supérieure*, Ser. 4, 24(6):635–704, 1991. [doi:10.24033/asens.1639](#). 1
- [vdB08] Michel van den Bergh. Double Poisson algebras. *Transactions of the American Mathematical Society*, 360(11):5711–5769, 2008. [arXiv:math/0410528](#). 3.3