# DEFINITIONS AND EXAMPLES OF ALGEBRAIC MORAVA K-THEORIES

NOBUAKI YAGITA

ABSTRACT. Algebraic Morava K-theories are defined by Sechin, Vishik, and others [Vi], [Ge-La-Pe-Se] as quotients of algebraic cobordism theories, which are oriented theories but not cohomology theories. On the other hand the author had defined the Morava theories [Ya7] as (the double degree) generalized cohomology theories. We compare these Morava-K-theories.

# 1. INTRODUCTION

In the  $\mathbb{A}^1$ -stable homotopy category, over a field k with ch(k) = 0, Voevodsky [Vo1] defined the space (spectrum) MGL(k) which defines the algebraic (motivic) cobordism

$$MGL^{*,*'}(X) \cong Mor(X, \mathbb{T}^{*,*'} \land MGL(k))$$

where Mor(-, -) is the group of morphisms in this category.

Soon later, Levine-Morel [Le-Mo1,2] defined an another algebraic cobordism  $\Omega^*(X)$  for a smooth X such that it is a universal oriented (for the formal groups laws). In particular, Levine shows [Le]

$$\Omega^*(X) \cong MGL^{2*,*}(X)$$
 for smooth X.

We will extend the above formula for the Morava K-theory, and compute their examples.

We first recall the Brown-Peterson cohomology (for details see §2 below or [No]) such that

$$\Omega^*(pt.)_{(p)} \supset BP^* = \mathbb{Z}_{(p)}[v_1, v_2, ...] \quad |v_i| = -2p^i + 2.$$

(For a topological cohomology theory  $A^*(X)$ , the notation  $A^*$  means its coefficients  $A^*(pt)$ , but for the algebraic case it means  $A^{2*,*}(pt)$ , here.)

Next recall that the (topological) Morava K-theory  $K(n)^*(X)$  is a generalized cohomology theory with the coefficient  $K(n)^* = \mathbb{Z}/p[v_n, v_n^{-1}]$ . In §2, we can construct [Ya7] the algebraic Morava K-theory  $AK(n)^{**'}(X)$  with  $deg(v_n) = (-2p^n + 2, -p^n + 1)$ .

<sup>2020</sup> Mathematics Subject Classification. 14C15, 14P99, 55N20.

Key words and phrases. algebraic cobordism, Morava K-theory.

We also recall the (topological) cobordism theory  $P(n)^*(X)$  in §3 and its algebraic version  $AP(n)^{*,*'}(X)$  such that  $P(n)^* = \mathbb{Z}/p[v_n, v_{n+1}, ...]$ . We define

$$AK(n)^{*,*'}(X) = K(n)^* \otimes_{BP^*} AP(n)^{*,*'}(X).$$

Let us write the ideal  $I_n = (p, v_1, ..., v_{n-1}) \subset BP^*$  so that  $P(n)^* \cong BP^*/I_n$ . Then we have (in §3)

**Lemma 1.1.** For a smooth X, we have

$$\Omega^*(X)/I_n \cong \Omega(X)^* \otimes_{BP^*} P(n)^* \cong AP(n)^{2*,*}(X),$$
$$\Omega^*(X) \otimes_{BP^*} K(n)^* \cong AK(n)^{2*,*}(X).$$

On the other side, Sechin , Vishik and others [Vl], [Ge-La-Pe-Se] define the (algebraic)  $K(n)^*(X)$  by directly  $\Omega^*(X) \otimes_{BP^*} K(n)^*$  (write it here by  $OK(n)^*(X)$ ).

To compute  $\Omega^*(X)$ , or  $\Omega(X)/I_n$  directly is the rather difficult. In this paper, we first compute  $ABP^{*,*'}(X)$ ,  $AP(n)^{*,*'}(X)$  by using the Atiyah-Hirzebruch (type) spectral sequences. Next we compute  $OP(n)^*(X)$  and the last  $OK(n)^*(X)$ .

As examples for Lemma 1.1, we consider the some cases X = BG classifying spaces BG of algebraic groups in §5. In §6, we consider the cases X = G algebraic groups themselves. In §7, we see

**Theorem 1.2.** Let G be an algebraic group, corresponding a simply connected Lie group which has a p-torsion. Then the algebraic group G is not homotopy nilpotent in  $\mathbb{A}^1$ -homotopy category.

In §8, we compute  $AP(n)^{*,*'}(\tilde{\chi}_V)$  when X is the reduced Ceck complex of the norm variety V. In §9, we study the K(n)-theory of the norm variety over  $k = \mathbb{R}$ .

## 2. Algebraic BP-theories

At first, we recall the algebraic  $MU^*$ -theory  $AMU^{*,*'}(X)$  from [Ya7].

For a topological space X, recall that  $MU^*(X)$  is the complex cobordism theory defined in the usual (topological) spaces and

$$MU^* = MU^*(pt.) \cong \mathbb{Z}[x_1, x_2, ...] \quad |x_i| = -2i.$$

Here each  $x_i$  is represented by a sum of hypersurfaces of  $dim(x_i) = 2i$  in some product of complex projective spaces ([Ha],[Ra]). Let  $MGL^{*,*'}(-)$ be the motivic cobordism theory defined by Voevodsky [Vo1]. Let us write by AMU the spectrum  $MGL_{(p)}$  in the stable  $\mathbb{A}^1$ -homotopy category representing this motivic cobordism theory (localized at p), i.e.,

$$MGL^{*,*'}(-)_{(p)} = AMU^{*,*'}(-).$$

Here note that  $AMU^{2*,*}(pt.) \cong MU^{2*}_{(p)}$ . It is not isomorphic to  $AMU^{*,*'}(pt)$  in general, while  $AMU^{*,*'}(X)$  is an  $MU^*_{(p)}$ -algebra.

Given a regular sequence  $S_n = (s_1, ..., s_n)$  with  $s_i \in MU^*_{(p)}$ , we can inductively construct the AMU-module spectrum by the cofibering of spectra ([Ya7,9], [Ra])

(2.1) 
$$\mathbb{T}^{-1/2|s_i|} \wedge AMU(S_{i-1}) \xrightarrow{\times s_i} AMU(S_{i-1}) \to AMU(S_i)$$

where  $\mathbb{T} = \mathbb{A}/(\mathbb{A} - \{0\})$  is the Tate object. For the realization map  $t_{\mathbb{C}}$  induced from  $k \subset \mathbb{C}$ , it is also immediate that  $t_{\mathbb{C}}(AMU(S_n)) \cong MU(S_n)$  with  $MU(S_n)^* = MU^*/(S_n)$ . Therefore  $t_{\mathbb{C}}$  induces the natural map

$$t_{\mathbb{C}}$$
 :  $AMU(S)^{*,*'}(X) \to MU(S)^{*}(X).$ 

Recall that the Brown-Peterson cohomology theory  $BP^*(-)$  with the coefficient  $BP^* \cong \mathbb{Z}_{(p)}[v_1, v_2...]$  by identifying  $v_i = x_{p^i-1}$ . (So  $|v_i| = -2(p^i - 1)$ .) We can construct spectra (in the stable  $\mathbb{A}^1$ -homotopy category)

$$ABP = AMU(x_i | i \neq p^j - 1)$$

such that  $t_{\mathbb{C}}(ABP) \cong BP$ . For  $S = (v_{i_1}, ..., v_{i_n})$ , let us write

$$ABP(S) = AMU(S \cup \{x_i | i \neq p^j - 1\})$$

so that  $t_{\mathbb{C}}(ABP(S)) = BP(S)$  with  $BP(S)^* = BP^*/(S)$ .

In particular, let  $AH\mathbb{Z} = ABP(v_1, v_2, ...)$  so that  $AH\mathbb{Z}^{2*,*}(pt.) \cong \mathbb{Z}_{(p)}$ . In the  $\mathbb{A}^1$ -stable homotopy category, Hopkins-Morel showed that

$$AH\mathbb{Z} \cong H_{\mathbb{Z}}$$
, *i.e.*,  $AH\mathbb{Z}^{*,*'}(X) \cong H^{*,*'}(X,\mathbb{Z}_{(p)})$ 

the (usual) motivic cohomology. Using this result, we can construct the motivic Atiyah-Hirzebruch (type) spectral sequence.

**Theorem 2.1.** ([Ya7,9]) Let Ah = ABP(S) for  $S = (v_{i_1}, v_{i_2}, ...)$ , and recall

$$h^{2*''} = BP^{2*''}/(S) \cong Ah^{2*'',*''}(pt)$$

Then there is AHss (the Atiyah-Hirzebruch spectral sequence)

$$E(Ah)_{2}^{(*,*',2*'')} = H^{*,*'}(X;h^{2*''}) \Longrightarrow Ah^{*+2*'',*'+*''}(X)$$

with the differential  $d_{2r+1}: E_{2r+1}^{(*,*',2*'')} \to E_{2r+1}^{(*+2r+1,*'-r,2*''-2r)}.$ 

Note that  $E_2^{*,*',2*''} \cong H^{*,*'}(X.Ah^{2*'',*''}(pt))$ . The coefficient part is the only difference place from the usual AHss.

Note that the cohomology  $H^{m,n}(X, h^{2n'})$  here is the usual motivic cohomology with (constant) coefficients in the abelian group  $h^{2n'}$ .

**Remark.** We do *not* asume the existence of the natural map  $AE_r \to E_r$  of spectral sequences for algebraic to topological. (while there is a map  $Ah^{*,*'}(X) \to h^*(X)$ )).

Here we recall some important properties of the motivic cohomology. When X is smooth, we know the (p - localized) Chow ring is

> (2.2)  $CH^*(X) \cong H^{2*,*}(X; \mathbb{Z}_{(p)}),$ (2.3)  $H^{*,*'}(X; \mathbb{Z}_{(p)}) \cong 0 \text{ for } 2*' < *.$

Hence if X is smooth, then  $E_r^{m,n,2n'} \cong 0$  for m > 2n.

Let  $S \subset R = (v_{j_1}, ...)$ . Then the induced map  $ABP(S) \to ABP(R)$ of spectra induces the  $BP^*$ -module map of AHss :  $E(ABP(S))_r^{*,*',*''} \to E(ABP(R))_r^{*,*',*''}$ . In general,  $ABP(S)^{*,*'}(X) \not\cong ABP^{*,*'}(X)/(S)$ . However, from the above maps and dimensional reason (2.3) with the differential (see also §4 below)

$$d_{2r+1}: E_{2r+1}^{(2*,*,0)} \to E_{2r+1}^{(2*+2r+1,*-r,-2r)} = 0,$$

we see that all elements in  $E_2^{2^{*,*,0}} \cong H^{2^{*,*}}(X) \otimes 1 \cong CH^*(X)$  are infinite cycles  $d_r = 0$  for all r > 0. Hence, for a smooth X (taking  $S = \emptyset, R = S$ ) we have the surjection and isomorphisms

(2.4) 
$$BP^{2*}/(S) \otimes CH^*(X) \twoheadrightarrow ABP(S)^{2*,*}(X)$$
  
(2.5)  $ABP(S)^{2*,*}(X) \otimes_{BP^*} \mathbb{Z}_{(p)} \cong H^{2*,*}(X) \cong CH^*(X)$ 

In this paper, a connective oriented theory  $h^{2*}(X)$  means  $ABP(S)^{2*,*}(X)$  as above. We mainly consider the connective oriented theory  $ABP^{2*,*}(X)$ . Hereafter, we write it simply

$$\Omega^*(X) = ABP^{2*,*}(X) \cong MGL^{2*,*}(X) \otimes_{MU^*} BP^*.$$

Hence from (2.5),  $\Omega^*(X) \otimes_{\Omega^*} \mathbb{Z}_{(p)} \cong CH^*(X)$  for a smooth X.

# 3. Algebraic Morava K theories

By the arguments in previous section we can define the algebraic versions  $Ak(n)^{*,*'}(X)$ ,  $AK(n)^{*,*'}(X)$  and  $AP(n)^{*,*'}(X)$ .

Recall the invariant ideal  $I_n = (p, v_1, ..., v_{n-1}) \subset BP^*$ . Define  $P(n) = BP(I_n)$  so that  $P(n)^* = BP^*/(I_n) \cong \mathbb{Z}/p[v_n, v_{n+1}, ...]$ . (Since  $I_n$  is invariant ideal,  $P(n)^*(X)$  has the Landweber-Novikov operations, but  $K(n)^*(X)$  below does not.)

The usual Morava K-theory is defined from the following (Conner-Floyd type) formula

(3.1) 
$$K(n)^*(X) = K(n)^* \otimes_{P(n)^*} P(n)^*(X).$$

so that  $K(n)^* = \mathbb{Z}/p[v_n, v_n^{-1}]$ . (We also know  $K(n)^*(X) \cong k(n)^*(X)[v_n^{-1}]$ , but most cases,  $k(n)^*(X)$  seems not easier to compute than  $P(n)^*(X)$ .) Remark that for general,

$$K(n)^*(X) \not\cong K(n)^* \otimes_{P(s)^*} P(s)^*(X) \quad for \ s \neq n.$$

We will study the algebraic version of these theorems.

Levine-Morel define algebraic cobordism  $\Omega^*(X)$  as the universal theory of the (some oriented) theories having the formal group laws. In particular, Levine (essentially) shows

$$\Omega^*(X) \cong ABP^{2**}(X)$$
 when X smooth.

Moreover, Sechin, Vishik and others define the oriented theories

(3,2) 
$$OBP(S)^*(X) = \Omega^*(X)/(S)$$

e.g. 
$$OP(n)^{*}(X) = \Omega^{*}(X)/I_{n}, \quad OK(n)^{*}(X) = \Omega^{*}(X) \otimes_{BP^{*}} K(n)^{*}$$

These oriented theories are not represented by the motivic cohomology theories in general. But we have the following lemma.

**Lemma 3.1.** Let X be snooth. Then we have the ring isomorphism

$$OP(n)^{2*}(X) = \Omega^*(X) / I_n \cong AP(n)^{2*,*}(X).$$

*Proof.* From (2.1) in the preceding section, we consider the exact sequence for the motivic cohomology

$$AP(n)^{2*'-1,*'}(X) \xrightarrow{\delta} AP(n-1)^{2*',*'}(X) \xrightarrow{v_{n-1}} AP(n-1)^{2*,*}(X) \to AP(n)^{2*,*}(X) \xrightarrow{\delta} AP(n-1)^{2*+1,*}(X) = 0$$

The last term in the above sequence follows from the dimensional reason (2.3). By induction, we have

$$AP(n)^{2*,*}(X) \cong AP(n-1)^{2*,*}(X)/(v_{n-1})$$
$$\cong OP(n-1)^{*}(X)/(v_{n-1}) = OP(n)^{*}(X).$$

**Remark.** It seems unknown that  $ABP(S)^{*,*'}(X)$  has a good ring structure for  $S \neq \emptyset$ . However  $AP(n)^{*,*'}(X)$  has when  $k = \mathbb{C}$  (Theorem 7.4 in [Ya7]).

For other regular sequence S in §2 in (2.1), by the same arguments, we have

**Lemma 3.2.** Let X be smooth. Then we have a  $BP^*$ -module isomorrphism

$$OBP(S)^{2*}(X) = \Omega^{*}(X)/(S) \cong ABP(S)^{2*,*}(X).$$

Next we recall the *truncated BP*-theory. Suppose that we fix generators  $v_1, v_2, ...,$  (and some finiteness conditions,) let us write  $J_{n+1} = (v_{n+1}, v_{n+2}, ...) \subset BP^*$  for the fixed generators (note  $J_n$  is not an invariant (under the Landweber-Novikov operations) ideal. Define the truncated BP-theory

 $BP\langle n \rangle^*(X) = BP(J_{n+1})^*(X)$ 

so that  $BP\langle n \rangle^* \cong BP^*/J_{n*1} \cong \mathbb{Z}_{(p)}[v_1, ..., v_n]$ . Similarly, we can define  $ABP\langle n \rangle^{*,*'}(X)$ . Moreover we have from the preceding lemma

**Lemma 3.3.** Let X be snooth. Then we have  $BP^*$ -module isomorphisms

$$OBP\langle n \rangle^{2*}(X) = \Omega^*(X)/J_{n+1} \cong ABP\langle n \rangle^{2*,*}(X),$$
$$Ok(n)^{2*}(X) = \Omega^*(X)/(I_n, J_{n+1}) \cong Ak(n)^{2*,*}(X).$$

## Remark.

The truncated theory  $ABP\langle n \rangle^{*,*'}(X)$  is decided by the choice of  $v_i \in J_{n+1}$  but not decided by n. Hence when we consider  $k(n)^*(X)$ , we fix some typical generators of  $BP^*$ .

## 4. MILNOR OPERATIONS AND WEIGHT DEGREE

Voevodsky showed that there exists the Milnor operation  $Q_i$  in the motivic cohomology  $H^{*,*'}(X; \mathbb{Z}/p)$  ([Vo1,3]). For an object  $\chi$  in  $\mathbb{A}^1$  homotopy category, we have

$$Q_i: H^{*,*'}(\chi; \mathbb{Z}/p) \to H^{*+2p^i - 1, *' + p^i - 1}(\chi; \mathbb{Z}/p)$$

which is compatible with the usual Milnor operation on  $H^*(t_{\mathbb{C}}(\chi); \mathbb{Z}/p)$ for the realization map  $t_{\mathbb{C}}$ . (Here the topological operation is defined  $Q_0 = \beta$  Bockstein operation and  $Q_{i+1} = Q_i P^{p^i} - P^{p^i}Q_i$ .) This operation  $Q_i$  can be extended on  $H^{*,*'}(M; \mathbb{Z}/p)$  for a motive M in M(X) (Lemma 7.1 in [Ya10]).

The relation  $Q_n$  to the exact sequence (2.1) for  $k(n)^*(X)$  is giving

$$(4.1) \quad k(n)^*(X) \xrightarrow{v_n} k(n)^*(X) \xrightarrow{r} H^*(X; \mathbb{Z}/p) \xrightarrow{\delta} k(n)^*(X) \to \dots$$

Here we can see  $Q_n = r\delta$ .

For  $0 \neq x \in H^{*,*'}(X; \mathbb{Z}/p)$  (or cohomology operation), let us write \* = |x|, w(x) = 2 \*' - \*, d(x) = \* - \*' so that  $0 \leq d(x) \leq dim(X)$  and  $w(x) \geq 0$  when X is smooth. We also note

$$|\tau| = 0, \ w(\tau) = 2, \quad |Q_i| = 2p^i - 1, \ w(Q_i) = -1$$

here  $0 \neq \tau \in H^{0,1}(Speck); \mathbb{Z}/p).$ 

For the differential  $d_r$  in AHss, we have  $w(d_r) = -1$ . Then the arguments for (2.4-2.5) are immediate. In fact some differencial  $d_r$  is represented as  $v_n \otimes Q_n$ .

 $\mathbf{6}$ 

We also note that (for  $h^*(-)$  in §2)

(4.2) 
$$Ah^{2**}(X) \cong Ah^{*',*''}(X)/\{x \in Ah^{*',*''}(X) | w(x) \ge 1\}.$$

# 5. Examples ; classifying spaces X = BG.

We give here the most simple but a non-trivial example, ([Ya5,8]) for Lemma 3.1. Let  $k = \bar{k} \subset \mathbb{C}$ , X = BG and  $G = \mathbb{Z}/p$ . Hence

$$H^{*,*'}(Spec(k);\mathbb{Z}/p)\cong\mathbb{Z}/p[\tau].$$

At first, we recall the topological cases. Then (for p odd)

$$H^*(BG) \cong \mathbb{Z}[y]/(py), \quad H^*(BG; \mathbb{Z}/p) \cong \mathbb{Z}/p[y] \otimes \Lambda(x),$$

where |x| = 1, |y| = 2 and  $x^2 = 0$ . (For p = 2,  $x^2 = y$ .) By the AHss, we have

$$E_2^{*,*'} \cong H^*(BG) \otimes BP^{*'} \cong BP^*[y]/(py) \Longrightarrow BP^*(X).$$

The  $E_2$ -term is generated by even degrees, and hence generates the  $E_{\infty}$ -term, That is (the graded ring)  $grBP^*(BG) \cong BP^*[y]/(py)$ .

More precisely, we can see

$$BP^*(BG) \cong BP^*[y]/([p](y)) \quad |y| = 2.$$

(For easy of notations we simply write by  $BP^*[y]$  the formal power series  $BP^*[[y]]$  in this paper.) Here [p](y) the *p*-th product of  $BP^*$ -formal group law, in particular

$$[p](y) = py + v_1y^p + v_2y^{p^2} + \dots \ mod(I_{\infty}^2).$$

Since  $y = c_1$  the first Chern class, we have the surjection

$$\Omega^*(X) \twoheadrightarrow BP^*(BG) \cong BP^*[y]/([p]y)$$

Next we consider the case P(s) for  $s \ge 1$ . The AHss is written

$$E_2^{*,*'} \cong H^*(BG; \mathbb{Z}/p) \otimes P(s)^{*'} \Longrightarrow P(s)^*(BG).$$

It is known the first non-zero differencial is

$$d_{2p^s-1}(x) = v_s Q_s(x) = v_s y^{p^s}.$$

Hence we see

$$grP(s)^*(X) \cong E_{2p^s} \cong P(s)^*[y]/(v_s y^{p^s}).$$

In fact, the right hand side is generated by even degree elements, and it is isomorphic to the infinite term. In particular,  $P(n)^{odd}(X) = 0$ .

The motivic  $(k = \bar{k})$  cohomology is written as

$$H^{*,*'}(BG:\mathbb{Z}/p)\cong H^*(BG;\mathbb{Z}/p)\otimes\mathbb{Z}/p[\tau],\quad \tau\in H^{0,1}(pt.;\mathbb{Z}/p).$$

Here for p = 2,  $x^2 = y\tau$  [Vo1-2]. We can get the same result for the algebraic case  $AP(n)^{odd,*'}(BG) = 0$ . We consider the exact sequence

$$0 = AP(1)^{2*-1,*}(X) \to ABP^{2*,*}(X) \xrightarrow{p} ABP^{2*,*}(X) \to AP(1)^{2*,*}(X) \to ABP^{2*+1,*}(X) = 0.$$

From the result of  $AP(1)^{*,*'}(X)$  and the fact  $\Omega^*(X)$  has no *p*-divisible element, we can see  $\Omega^*(X)/p \cong AP(1)^{2**}(X)$ . Thus we have

 $\Omega^*(X) \cong BP^*(X) \cong BP^*[y]/([p](y)).$ 

Hence we can apply the main lemma.

**Lemma 5.1.** When X = BG,  $G = \mathbb{Z}/p$ , we have  $OP(n)^*(X) = \Omega^*/I_n \cong AP(n)^{2*,*}(X).$ 

Thus, we can write

$$AP(n)^{2*,*}(BG) \cong \Omega^*(BG)/I_n \cong BP^*[y]/([p](y), I_n).$$
  
$$\cong \mathbb{Z}/p[v_n, v_{n+1}, \dots][y]/(v_n y^{p^n} + v_{n+1} y^{p^{n+1}} + \dots)$$
  
$$\cong P(n)^{2*}[y]/(v_n y^{p^n} + v_{n+1} y^{p^{n+1}} + \dots),$$

which has no  $v_n$ -torsion.

**Corollary 5.2.** Let X=BG for  $G=\mathbb{Z}/p$ . Then

$$AK(n)^{2*,*}(BG) \cong \Omega^*(BG) \otimes_{BP^*} K(n)^*$$
$$\cong K(n)^*[y]/(v_n y^{p^n}) \cong K(n)^*[y]/(y^{p^n}).$$

Hence the corresponding formal group law is the Honda group law (identifying  $v_n = 1$ ).

The similar (but more weak) facts hold for other groups. (Theorem 12.1, Proposition 12.4 in [Ya8])

**Lemma 5.3.** Let X = BG and G be  $\mathbb{Z}/p$ , the orthogonal groups  $O_m$ ,  $SO_m$  (or their products). Then we have  $AP(1)^{odd.*'}(X) = 0$  and

$$AP(1)^{*'',*'}(X) \cong AP(1)^{2*,*}(X) \otimes \mathbb{Z}/p[\tau]$$
  
where  $AP(1)^{2*,*}(X) \cong P(1)^{2*}(X) \cong BP^{2*}(X)/p$ 

**Example.** We consider the case X = BG for  $G = SO_3$  and p = 2. The cohomology is (given by the Stiefel-Whitney classes  $w_2, w_3$  and Chern classes  $c_2, c_3$ )

 $H^*(X; \mathbb{Z}/2) \cong \mathbb{Z}/2[w_2, w_3].$  with  $Q_0 w_2 = w_3.$ 

Hence  $H^*(X) \cong (\mathbb{Z}\{1\} \oplus \mathbb{Z}/2[c_3]^+) \otimes \mathbb{Z}[c_2]$  with  $c_i = w_i^2$ .

By using  $Q_1(w_3) = w_3^2 = c_3$  and the AHss, we have

 $grBP^*(X) \cong (BP^*\{1\} \oplus BP^*[c_3]^+/(2,v_1)) \otimes \mathbb{Z}[c_2]$ 

Here note that in  $BP^*(X)$ , the element  $v_1c_3 \neq 0$  (it is  $v_2c_2^2 + ...$ , this fact is shown considering the restriction to  $BP^*(B\mathbb{Z}/2)$ ).

Hence we see  $BP^*(X)/2 \cong P(1)^*(X)$ . (For the details of the AHss converging  $P(1)^*(X)$ , see §12 in [Ya8].)

## 6. Lie groups

Let  $k = \overline{k} \subset \mathbb{C}$ . In this section, we study for X = G; Lie groups themselves. For G exceptional groups and  $p \neq 2$  are studied in [Ya1,2,3]. Hereafter let X = C = SO. Then

Hereafter let  $X = G = SO_7$ , Then

$$H^*(G; \mathbb{Z}/2) \cong \Lambda(x_3, x_5, y_6), \quad x_3^2 = y_6$$
$$H^*(G) \cong \Lambda(x_3) \otimes (\mathbb{Z}\{1, x_5y_6\} \oplus \mathbb{Z}/2\{y_6\}).$$

Here the suffix *i* of  $x_i$ ,  $y_6$  means its degree and  $x_5y_6$  is an element in  $H^*(G)$  such that its image in  $H^*(G; \mathbb{Z}/2)$  is the same name.

We consider AHss

$$E_2 \cong H^*(G) \otimes BP^* \Longrightarrow BP^*(G).$$

The differencial is given by  $d_3(x_3) = v_1Q_1x_3 = v_1y_6$ . Hence infinitive term is written as

$$E_{\infty} \cong BP^*A \oplus BP^*B/2 \oplus BP^*C/(2, v_1)$$

with  $A = \mathbb{Z}\{1, 2x_3, x_5y_6, x_3x_5x_6\}, \quad B = \mathbb{Z}/2\{x_3x_6\}, \quad C = \mathbb{Z}/2\{y_6\}.$ Here  $2y_3$  and  $x_iy_6$  are elements in  $E_{\infty}$  induced from the same named elements in  $E_2$ .

Then we can compute more exactly [Ya1,2]

$$BP^{*}(G) \cong BP^{*}\{1, 2x_{3}, x_{3}x_{5}x_{6}\}$$
  

$$\oplus BP^{*}\{x_{3}y_{6}, x_{5}y_{6}\}/(2x_{3}y_{6} - v_{1}x_{5}y_{6})$$
  

$$\oplus BP^{*}/(2, v_{1})\{y_{6}\}.$$

Similarly, we can compute (for example using  $P(2)^*(X) \cong P(2)^* \otimes H^*(X; \mathbb{Z}/2)$ )

$$P(1)^*(G) \cong (P(1)^*\{1, x_3y_6\} \oplus P(2)^*\{y_6\}) \otimes \Lambda(x_5).$$

Indeed, take  $x_5 \in P(1)^5(X)$  with  $\delta(x_5) = y_6$ . The element  $2x_3 \in BP^*(X)$  goes (by the reduction u) to  $v_1x_5 \in P(1)^*(X)$  i.e.,  $u(2x_3) + v_1x_5 = 0$ , recall  $Q_1x_3 = Q_0x_5$ .

By Kac [Ya6], we knew  $(CH^*(X)/2 = \mathbb{Z}/2\{1, y_6\})$ 

$$\Omega^*(G)/I_{\infty} \cong CH^*(G)/2 \cong \mathbb{Z}/2\{1\} \oplus \mathbb{Z}/2\{y_6\}.$$

Moreoer we know  $v_1y_6 = 0$  from  $Q_1x_3 = y_6$ . Hence we see  $w(y_6) = 0$  but  $w(x_ix_j) \ge 1$  for the other generators. Thus we get

$$ABP^{2*,*}(G) \cong \Omega^*(G) \cong BP^*\{1\} \oplus P(2)^*\{y_6\}.$$

Since  $k = \bar{k}$ , we see  $H^*_{et}(Spec(k); \mathbb{Z}/2) \cong \mathbb{Z}/2[\tau]$ . Hence each  $\tau^i$  is a permanent cycle in the above AHss, and we get

**Theorem 6.1.** Let  $G = SO_7$  Then

$$AP(1)^{*,*'}(G) \cong \Omega^*(G) \otimes \mathbb{Z}/2[\tau],$$
  
where  $\Omega^*(G)/2 \cong P(1)^*\{1\} \oplus P(2)^*\{y_6\}.$ 

Corollary 6.2. Let  $G = SO_7$ . Then

$$AK(s)^{2*,*}(G) \cong \begin{cases} K(1)^*\{1\} & for \ s = 1\\ K(2)^*\{1, y_6\} & for \ s \ge 2 \end{cases}$$

Quite recently, Gelthauser- Lavrenov- Petrov-Sechin [Ge-La-Pe-Se] study the Morava K-theories for all  $SO_m$ .

# 7. Homotopy Nilpotency

Let X = G be a simply connected Lie group. For ease of arguments, let us assume p odd in this section. By Borel, its mod(p) cohomology is a tensor product of truncated polynomial and an exterior algebra.

Moreover when it has *p*-torsion. there is an surjective map [Ya4]

(7.1) 
$$H^*(X; \mathbb{Z}/p) \twoheadrightarrow A = \mathbb{Z}/p[y]/(y^p) \otimes \Lambda(x, x')$$
  
where  $|x| = 3$ ,  $P^1(x) = x'$ ,  $Q_1x = Q_0x' = y$ ,

(Hence |y| = 2p + 2, |x'| = 2p + 1)

We also see, by dimensional reason, that for X in (7.1)

(7.2)  $K(2)^*(X) \twoheadrightarrow K(2)^* \otimes A$ 

We consider the homology theory (the  $\mathbb{Z}/p$ -dual of the cohomology theory). There is the injection (but *not* a *ring* map)

$$K(2)_* \otimes B \subset K(2)_*(X) \text{ where } B = \mathbb{Z}/p[y]/(y^p) \otimes \Lambda(z, z'),$$

with y (resp. z.z') is the dual of y (resp. x, x')

We want to study the Pontryagin ring structure of  $K(2)_*(X)$ , i.e., the product is induced from that of the Lie group G. Let us write the adjoint

$$ad(y)(z) = [y, z] = yz - zy.$$

(Here the product is the Ponriyagin product.) The definition of homotopy nilpotents implies as all Pontriyagin products are nilpotents. In particular

$$ad^{i}(y)(z) = 0$$
 for large *i*.

**Lemma 7.1.** ([Ya4] Theorem 1.5) Let G in the group (7.1). Then G is not homotopy nilpotent.

$$ad^{p-1}(y)(z) = -v_2 z \neq 0 \in K(2)_*(G).$$

We know

$$x \in H^{3,3}(G; \mathbb{Z}/p) \cong H^3_{et}(G; \mathbb{Z}/p) \cong H^3(X(\mathbb{C}); \mathbb{Z}/p).$$

Hence x (resp. z) exists in  $AK(2)^{*,*'}(X)$  (resp. $AK(2)_{*,*'}(X)$ ) By the compatible of  $t_{\mathbb{C}}$  and K2)\* we have that X is not homotopy nilpotent.

**Lemma 7.2.** Let G be an algebraic group, corresponding a simply connected Lie group which has a p-torsion. Then G is not homotopy nilpotent in  $\mathbb{A}^1$ -homotopy category.

Examples. Let  $G = F_4, p = 3$  (or  $G = E_8, p = 5$ ). Then we have  $K(2)_*(G) \cong K(2)_*[y]/(y^p) \otimes C \otimes C'$ where  $\begin{cases} C = \Lambda(z, ad^1(y)(z), ...ad^{p-2}(y)(z)) \\ C' = \Lambda(z', ad^1(y)(z'), ...ad^{p-2}(y)(z')). \\ ad^{p-1}(y)(z) = -v_2 z. \quad ad^{p-1}(y)(z') = -v_2 z'. \end{cases}$ 8. Cêck COMPLEX FOR  $k \neq \bar{k}$ 

Suppose ch(k) = 0 but we do not assume  $k = \overline{k}$  in this section. (We do not assume X smooth.) The following lemmas also hold for the algebraic theories. Let us write  $Q(n) = \Lambda(Q_0, ..., Q_n)$  for the Milnor operation  $Q_i$ .

**Lemma 8.1.** Let  $P(1)^*(X) \cong P(n)^* \otimes B_n$ , for some  $\mathbb{Z}/p$ -module  $B_n$ , (*i.e.* it is an  $I_n$ -torsion and  $P(n)^*$ -free module ). Then

$$P(s)^{*}(X) \cong P(n)^{*} \otimes Q(s-1) \otimes B_{s}$$
 for  $s \leq n$   
where  $Q_{1}...Q_{n-1}B_{n} = B_{1}$ , (i.e.,  $B_{s} = Q_{1}^{-1}...Q_{s}^{-1}B_{1}$ ). Hence

$$K(s)^*(X) \cong \begin{cases} 0 \quad s < n \\ K(n)^* \otimes Q(n-1) \otimes B_n \quad s = n \end{cases}$$

*Proof.* Let s < n. By induction, we assume

$$P(s)^*(X) \cong P(n)^* \otimes Q(s-1)B_s.$$

We consider the exact sequence (2.1)

$$P(s)^*(X) \xrightarrow{v_s} P(s)^*(X) \xrightarrow{r} P(s+1)^*(X) \xrightarrow{\delta} P(s)^{*+1}(X)...$$

Here we assume  $P(s)^*(X)$  is a  $P(n)^*$ -module by induction. That is  $v_s = 0$ . Hence we have

$$P(s+1)^*(X) \cong \mathbb{Z}/p\{r, \delta^{-1}\}P(s)^*(X)$$
  
$$\cong \mathbb{Z}/p\{r, \delta^{-1}\} \otimes P(n)^* \otimes Q(s-1)B_s \quad by \ induction$$
  
$$\cong \mathbb{Z}/p\{1, Q_s^{-1}\} \otimes P(n)^* \otimes Q(s-1)B_s \quad by \ Q_s = r\delta$$
  
$$\cong P(n)^* \otimes Q(s-1) \otimes \mathbb{Z}/p\{B_s, B_{s+1}\} \quad by \ B_s = Q_s B_{s+1}.$$

$$\cong P(n)^* \otimes Q(s) \otimes B_{s+1}$$

Taking s = n - 1, we get the result.

**Corollary 8.2.** By the same assumption as the above lemma. we see that

$$H^*(X;\mathbb{Z}/p)\cong Q(n-1)\otimes B_n$$

*Proof.* We study the exact sequence in the proof of the preceding lemma, but for  $s \ge n+1$ . Since the map  $v_s$  on  $P(s)^*(X) \cong P(s)^*B_n$  is injective, we have

$$P(s+1)^*(X) \cong P(s+1)^* \otimes B_n.$$

We get the corollary (for the algebraic case from the Hopkins and Morel theorem  $\lim_{s\to\infty} AP(s)^{*,*'}(X) \cong H^{*,*'}(X;\mathbb{Z}/p)$ .)

**Lemma 8.3.** Suppose that as a Q(n-1)-module

$$H^*(X;\mathbb{Z}/p)\cong Q(n-1)\otimes B_n$$

Then we have  $P(1)^*(X) \cong P(n)^* \otimes B_1$ .

*Proof.* We consider AHss

$$E_2 \cong H^*(X; \mathbb{Z}/p) \otimes P(1)^* \cong Q(n-1)B_n \otimes P(1)^* \Longrightarrow P(1)^*(X).$$

By the naturality, the first non-zero differential is given

$$d_{2p-1}(B_n) = v_1 \otimes Q_1 B_n.$$
  
Hence  $Ker(d_{2p-1})$  is  $P(1)^* \otimes Q_1 B_n$  from  $(Q_1)^2 = 0$ . Moreover we see  
 $E_{2p} \cong P(1)^* \otimes Q_1 B_n / (v_1 Q_1 B_n).$ 

By induction and the assumption of degree and  $d_{2p^i-1}(x) = v_i \otimes Q_i x$ , we can compute

$$E_{\infty} \cong P(1)^* / (v_1, v_2, ..., v_{n-1}) \otimes Q_1 Q_2 ... Q_{n-1}) B_n$$

which is isomorphic to  $P(n)^* \otimes B_1$ . (Note that  $Q_1 \dots Q_{n-1} B_n \in P(1)^*(X)$ ,)

Corollary 8.4. Let  $P(1)^*(X) \cong P(n)^* \otimes B_n$ . Then  $K(n)^*(X) \cong K(n)^* \otimes B_n$ , and  $K(s)^*(X) \cong 0$  s < n.

*Proof.* We have the corollary from  $K(s)^* \otimes P(n)^*_{BP^*} \otimes B_s = 0.$ 

## Example 1. V(n).

In the (topological) stable homotopy category, the space X = V(n) is called the Smith-Toda space (spectrum) if

$$P(1)^*(X) \cong P(n+1)^* \quad (or \ H^*(X; \mathbb{Z}/p) \cong Q(n)).$$

12

If V(n) exists then we can define a (non-zero) stable homotopy element of a sphere

$$S^0 \to V(n) \xrightarrow{v_n} V(n) \xrightarrow{qu.} S^N \text{ some } N$$

This is thought as an important problem in the stable homotopy groups of spheres. However V(n) does not exist in general. For example V(3) exists if and only if  $p \ge 7$ .

**Example 2** ; norm variety for  $\bar{k} \neq k$ .

Let  $\chi_X$  be the  $\hat{C}ech$  complex, which is defined as  $(\chi_X)^n = X^{n+1}$  (see details [Vo1,2,4]). Let  $\tilde{\chi}_X$  be defined from the cofiber sequence

$$\tilde{\chi}_X \to \chi_X \to Spec(k)$$

in the stable  $\mathbb{A}^1$ -homotopy category. Voevodsky defined the motivic cohomology  $H^{*,*'}(\chi; \mathbb{Z}/p)$  for all objects  $\chi$  in the stable  $\mathbb{A}^1$ -homotopy category.

It is known that the ideal

$$I(X) = \pi_* BP^*(X) \subset BP^*(pt.) = BP^* \quad for \ \pi : X \to pt.$$

is an invariant ideal (e.g. Lemma 5.3 in [Ya7]).

The following two lemmas are known.

**Lemma 8.5** (Vo1). Let  $p: X \to Spec(k)$  be the projection and  $t_{\mathbb{C}}(X) = v \in BP^*$ . Let Ah = ABP(S) for some regular sequence S. Then

$$Ah^{*,*'}(\tilde{\chi}_X)$$
 is  $v-torsion$ .

*Proof.* Let us write  $\chi_X$  by  $\chi$  simply. We consider the maps

$$Ah^{*,*'}(\chi) \xrightarrow{p^*} Ah^{*,*'}(\chi \times X) \xrightarrow{p_*} Ah^{d,d'}(\chi)$$

so that  $p_*p^*(x) = vx$ . The lemma is proved from  $Ah^{**'}(\tilde{\chi}) = 0$  because

$$Ah^{*,*'}(\chi \times X) \cong Ah^{*,*'}(\chi).$$

**Lemma 8.6.** (Lemma 6.4, 6.5 in [Ya10]) If  $I_n \subset I(X)$ , then  $ABP^{*,*'}(\tilde{\chi}_X)$  is  $I_n$ -torsion, and  $H^{*,*'}(\tilde{\chi}_X; \mathbb{Z}/p)$  is Q(n-1)-free.

Now we recall the norm variety. Given a pure symbol a in the mod p Milnor K-theory  $K_n^M(k)/p$ , by Rost, we can construct the norm variety  $V_a$  such that

$$\pi_*([V_a]) = v_{n-1}, \quad a|_{k(V_a)} = 0 \in K_n^M(k(V_a))/p$$

where  $[V_a] = 1 \in \Omega^0(V_a), \pi : V_a \to pt$  is the projection and  $k(V_a)$  is the function field of  $V_a$  over k. Note  $I(V_a) = I_n$ .

**Theorem 8.7.** (For p = 2, [Ya4]) Let  $0 \neq a = (a_0, ..., a_n) \in K_n^M(k)/p$ . Then there is a  $K_*^M(k) \otimes Q(n-1)$ -modules isomorphism

 $H^{*,*'}(\tilde{\chi}_a; \mathbb{Z}/p) \cong K^M_*(k)/(Ker(a)) \otimes Q(n-1) \otimes \mathbb{Z}/p[\xi_a]\{a'\}$ where  $\xi_a = Q_{n-1}...Q_0(a')$  and deg(a') = (n, n-1).

Thus from Lemma 8.3 and Lemma 3.1, we see

Corollary 8.8. We have

$$AK(s)^{*,*'}(\tilde{\chi}_a) \cong \begin{cases} K(n)^* \otimes H^{*,*'}(\tilde{\chi}_a; \mathbb{Z}/p) & s = n \\ 0 & 1 \le s < n. \end{cases}$$

///

**Remark.** For details of etale cohomologies of  $\chi_a$  and  $M_a$  are studied in §8,9 in [Ya 10]. For an example

$$H^{*,*'}(M_a; \mathbb{Z}/2) \cong H^{*,*'}(\chi_a; \mathbb{Z}/p) \quad for * < 2(p^{n-2}-1)/(p-1).$$

# 9. The motivic cohomology of quadrics over $\mathbb{R}$ with coefficients $\mathbb{Z}/2$

Let X be a smooth variety over the field  $\mathbb{R}$  of real numbers, and we consider the cohomologies of  $\mathbb{Z}/2$  coefficients. In this paper the mod(2) étale cohomology means the motivic cohomology of the same first and the second degrees  $H^*_{\acute{e}t}(X;\mathbb{Z}/2) \cong H^{*,*}(X:\mathbb{Z}/2)$ .

It is well known ([Vo1], [Vo2])

$$H^*_{\acute{e}t}(Spec(\mathbb{C});\mathbb{Z}/2)\cong\mathbb{Z}/2, \quad H^{*,*'}(Spec(\mathbb{C});\mathbb{Z}/2)\cong\mathbb{Z}/2[\tau],$$

 $H^*_{\acute{e}t}(Spec(\mathbb{R}); \mathbb{Z}/2) \cong \mathbb{Z}/2[\rho], \quad H^{*,*'}(Spec(\mathbb{R}); \mathbb{Z}/2) \cong \mathbb{Z}/2[\tau, \rho]$ 

where  $0 \neq \tau \in H^{0,1}(Spec(\mathbb{R}); \mathbb{Z}/2) \cong \mathbb{Z}/2$  and where  $\rho = -1 \in \mathbb{R}^*/(\mathbb{R}^*)^2 \cong K^M_{\mathfrak{k}}(\mathbb{R})/2 \cong H^1_{\mathfrak{k}}(Spec(\mathbb{R}); \mathbb{Z})$ 

$$\rho = -1 \in \mathbb{R}^*/(\mathbb{R}^*)^2 \cong K_1^M(\mathbb{R})/2 \cong H^1_{\acute{e}t}(Spec(\mathbb{R}); \mathbb{Z}/2).$$

We recall the cycle map from the Chow ring to the étale cohomology

$$cl/2: CH^*(X)/2 \to H^{2*}_{\acute{e}t}(X; \mathbb{Z}/2).$$

This map is also written as  $H^{2*,*}(X;\mathbb{Z}/2) \xrightarrow{\times \tau^*} H^{2*,2*}(X;\mathbb{Z}/2)$ .

Let  $X = Q^d$  be an anisotropic quadric of dimension  $2^n - 1$  (i.e. the norm variety for  $\rho^{n+1} \in K^M_{n+1}(\mathbb{R})/2$ ). Then we have the Rost motive  $M \subset Q^d$  [Ro]. It is known (the remark in page 575 in [Ya2])

$$H^*_{\acute{e}t}(M; \mathbb{Z}/2) \cong \mathbb{Z}/2[\rho]/(\rho^{2^{n+1}-1}) \cong \mathbb{Z}/2\{1, \rho, \rho^2, ..., \rho^{2^{n+1}-2}\}$$

The Chow ring is also known [Ro]

$$CH^*(M)/2 \cong \mathbb{Z}/2\{1, c_0, c_1, \dots, c_{n-1}\}, \quad cl(c_i) = \rho^{2^{n+1}-2^{i+1}}.$$

The cycle map cl/2 is injective. The elements  $c_i$  is also written as

$$c_i = \rho^{2^{n+1}-2^{i+1}} \tau^{-2^n+2^i}$$
 in  $CH^*(M)/2 \subset H^{2*}_{\acute{e}t}(M:\mathbb{Z}/2)[\tau^{-1}]$ 

The mod(2) motivic cohomology is known (Theorem 5.3 in [Ya9]).

**Theorem 9.1.** (Theorem 5.3 in [Ya9], [Ya3]) The cohomology  $H^{*,*'}(M_n; \mathbb{Z}/2)$  is isomorphic to the  $\mathbb{Z}/2[\rho, \tau]$ -subalgebra of

$$\mathbb{Z}/2[\rho, \tau, \tau^{-1}]/(\rho^{2^{n+1}-1})$$

generated by  $a = \rho^{n+1}$ ,  $a' = a\tau^{-1}$ , and elements in  $\Lambda(Q_0, ..., Q_{n-1})\{a'\}$ . **Lemma 9.2.** We have  $Q_0(\tau^{-1}) = \rho\tau^{-2}$ . Hence  $Q_0(a') = \rho a\tau^{-2}$ , while  $Q_0(a) = 0$ . Similarly, we see  $Q_1(\tau^{-2}) = \rho^3 \tau^{-4}$ , and  $Q_1(a') = \rho^6 \tau^{-3}$ .

*Proof.* We see the first equation from

$$0 = Q_0(1) = Q_0(\tau\tau^{-1}) = \rho\tau^{-1} + \tau Q_0(\tau^{-1}).$$

**Example.**  $X = M_2$ . Recall that  $\tau : H^{*,*'}(X; \mathbb{Z}/2) \to H^{*,*'+1}(X; \mathbb{Z}/2)$  is injective. The motivic cohomology is given as (Example 5.15 in [Ya9])

$$H^{*,*'}(X; \mathbb{Z}/2) \cong \mathbb{Z}/2[\tau] \{1, \rho, \rho^2, a' = \rho^3 \tau^{-1}, Q_0 a' = \rho^4 \tau^{-2}, \rho Q_0 a' = \rho^5 \tau^{-2}, Q_1 a' = \rho^6 \tau^{-3} \}.$$

We consider AHss for s = 1 or 2

$$E_2 \cong H^{*,*'}(X; \mathbb{Z}/2) \otimes P(s)^* \Longrightarrow AP(s)^{*,*'}(X).$$

We first consider the case s = 2. The first degree of the differential are (see Theorem 2.1)

 $-|v_2| + 1 > 6 = first \ degree \ of H^{2*,*}(X:\mathbb{Z}/2).$ 

Hence all differentials (to  $CH^*(X)/2$ ) are zero.

Next, we consider the case s = 1 We note

$$d_3(a') = Q_1 a' = \rho^6 \tau^{-3}.$$

Then we see

**Theorem 9.3.** *Let*  $X = M_2$ *.* 

$$AP(s)^{2*,*}(X) \cong \begin{cases} P(1)^* \{1, Q_0 a'\} \oplus P(2)^* \{Q_1 a'\} & s = 1\\ P(s)^* \{1, Q_0 a', Q_1 a'\} & s \ge 2. \end{cases}$$

Recall  $Q^3$  is the norm variety  $M_2 \subset Q^3$ . We have the decomposition of the motives  $M(Q^3) \cong M_2 \oplus M_1 \otimes \mathbb{T}$  (see §8 in [Ya9]). The Chow ring is written

 $CH^*(Q^3) \cong \mathbb{Z}_2\{1, h, h^2, h^3\} \oplus \mathbb{Z}/2\{c\}.$ Here  $h \in CH^1(\mathbb{T}), h^2 \in CH^2(M_1), h^3 = Q_1a'$  and  $c = Q_0a'$ . Hence **Corollary 9.4.** For the norm variety  $Q^3$  over  $\mathbb{R}$ , we have

$$AP(1)^{2*,*}(Q^3) \cong P(1)^*\{1, h, h^2, Q_0(a')\} \oplus P(2)^*\{h^3\}$$
$$AK(1)^{2*,*}(Q^3) \cong K(1)^*\{1, h, h^2, Q_0(a')\}.$$

Remark that (Lemma 6.2 in [Ya9])) we know  $Q_1(\tau^2) = \rho^3$ . However this element does note related elements of w(x) = 0 (In fact  $w(\tau) = 2$ ).

## Theorem 9.5. Let $X = M_n$ .

$$AP(s)^{2*,*}(X) \begin{cases} \supset P(n)^* \{Q_1 ... Q_{n-1} a'\} & s = n-1 \\ \cong P(s)^* \otimes CH^*(X)^*\} & s \ge n \end{cases}$$

Recall  $a = \rho^{n+1}$ .  $a' = a\tau^{-1}$ . So w(a') = n - 1. In fact, we have

$$CH^*(X)/2 \cong H^{2*,*}(X; \mathbb{Z}/2) \cong \mathbb{Z}/2\{Q_0...\hat{Q}_i...Q_{n-1}(a')|0 \le i \le n-1\}.$$

#### References

- [Ge-La-Pe-Se] N.Geldhauser, A.Lavrenov, V.Petrov, P.Sechin. Morava J-invariant ArXiv. 2024.14099v1 [Math.KT]
- [Ha] M.Hazewinkel. Formal groups and applications. *Pure and applied Math.* 78., *Academic Press, Inc.* (1978), 573pp.
- [Le] M. Levine. Comparison of cobordism theories. J. Algebra **322** (2009), 3291-3317.
- [Le-Mo1] M. Levine and F. Morel. Cobordisme algébrique I. C. R. Acad. Sci. Paris 332 (2001), 723-728.
- [Le-Mo2] M. Levine and F. Morel. Cobordisme algébrique II. C. R. Acad. Sci. Paris 332 (2001), 815-820.
- [Mi] J.Milnor. On cobordism ring  $\Omega_*$  and a complex analogue, I. Amer. J. Math. 82 (1960), 505-521.
- [Mo-Vo] F. Morel and V. Voevodsky. A<sup>1</sup>-homotopy theory of schemes. *IHES Publ.* Math. 90 (2001), 45-143.
- [No] P. Novikov The methods of algebraic topology from the view point of cobordism theory. *Math. USSR. Izv.* **1** (1967), 827-913.
- [Ra] D.Ravenel. Complex cobordism and stable homotopy groups of spheres. Pure and Applied Mathematics, 121. Academic Press (1986).
- [Qu] D. Quillen. Elementary proofs of some results of cobordism theory using Steenrod operations. Adv. Math. 7 (1971), 29-56.
- [To1] B. Totaro. Torsion algebraic cycles and complex cobordism. J. Amer. Math. Soc. 10 (1997), 467–493.
- [To2] B. Totaro. The Chow ring of classifying spaces. Proc. of Symposia in Pure Math. "Algebraic K-theory" (1997:University of Washington, Seattle) 67 (1999), 248-281.
- [Vi] A.Vishik. Isotropic and numerical equivalence for the Cow rings and Morava K-theories Invent. Math. 237 (2024), 770-808.
- [Vo1] V. Voevodsky. The Milnor conjecture. www.math.uiuc.edu/K-theory/0176 (1996).

- [Vo2] V. Voevodsky (Noted by Weibel). Voevodsky's Seattle lectures : K-theory and motivic cohomology Proc. of Symposia in Pure Math. "Algebraic Ktheory" (1997:University of Washington, Seattle) 67 (1999), 283-303.
- [Vo3] V. Voevodsky. Motivic cohomology with Z/2 coefficient. Publ. Math. IHES 98 (2003), 59-104.
- [Vo4] V.Voevodsky. Reduced power operations in motivic cohomology. Publ.Math. IHES 98 (2003),1-57.
- [Vo6] V.Voevodsky. On motivic cohomology with Z/l-coefficients. Ann. Math. 174 (2011) 401-438.
- [Ya1] N. Yagita. Brown-Peterson cohomology of exceptional Lie group J. Pure and Applied Algebra 17 (1980), 223-23223-226.
- [Ya2] N. Yagita. On the mod odd prime Brown-Peterson cohomology groups of exceptional Lie groups. J. Math. Soc. Japan 34 (1982) 294-305.
- [Ya2] N. Yagita. On relations between Brown-Peterson cohomology and the ordinary mod p cohomology theory.. Kodai Math. J. 7 (1984), 273-285. /////
- [Ya4] N. Yagita. Pontrojagin rings of the Morava K-theory for finite H-spaces. newblock J. Math. Kyoto Univ. c. 36 (1996), 447-452.
- [Ya5] N. Yagita. Examples for the mod p motivic cohomology of classifying spaces. Trans. Amer. Math. Soc. 335 (2003) 4427-4450.
- [Ya6] N. Yagita. Algebraic cobordism of simply connected Lie groups. Math.Proc. Camb.Phil.Soc. 139 (2005), 243-260.
- [Ya7] N. Yagita. Applications of Atiyah-Hirzebruch spectral sequence for motivic cobordism. Proc. London Math. Soc. 90 (2005) 783-816.
- [Ya8] N. Yagita. Coniveau filtrations of cohomology of groups. Proc. London Math. Soc. 101 (2010), 179-206.
- [Ya9] N. Yagita, Motivic cohomology of quadrics and the coniveau spectral sequence. J. K-theory 6 (2010) 547-589.
- [Ya10] N. Yagita. Algebraic BP-theory and norm varieties. Hokkaido Math. J. 41 (2012), 275-316.

[N. YAGITA] DEPARTMENT OF MATHEMATICS, FACULTY OF EDUCATION, IBARAKI UNIVERSITY, MITO, IBARAKI, JAPAN

Email address: nobuaki.yagita.math@vc.ibaraki.ac.jp