# arXiv:2502.04930v1 [math.CT] 7 Feb 2025

# A 2-dimensional torsion theory on symmetric monoidal categories

Mariano Messora

#### Abstract

In this paper we describe a homotopy torsion theory in the category of small symmetric monoidal categories. Thanks to the use of natural isomorphisms as basis for the nullhomotopy structure, this homotopy torsion theory enjoys some interesting 2-dimensional properties which may be the starting point for a definition of "2-dimensional torsion theory".

As torsion objects we take symmetric 2-groups, thus generalising a known pointed torsion theory in the category of commutative monoids where abelian groups play the part of torsion objects. In the last part of the paper we carry out an analogous generalisation for the classical torsion theory in the category of abelian groups given by torsion and torsion-free groups.

## Contents

1	Introduction	<b>2</b>
<b>2</b>	Preliminaries	3
3	Purely monoidal categories	6
4	Homotopy torsion theories	12
<b>5</b>	A nullhomotopy structure $\Theta$ for monoidal categories	15
6	A $\Theta$ -torsion theory for symmetric monoidal categories	16

7	2-dimensional aspects	19
8	Homotopy invariants	20
9	Homotopy torsion theories and pretorsion theories	23
10	A remark on the symmetric setting	25
11	Another 2-dimensional torsion theory	27

# 1 Introduction

Torsion theories were originally introduced in the context of abelian categories ([7]) as a generalisation of a well-known property of the category of abelian groups: this category includes two special classes of objects—torsion and torsion-free groups—and these classes enjoy the following key features.

- 1. Any group homomorphism from a torsion group to a torsion-free group is necessarily zero;
- 2. any abelian group A sits in the middle of a short exact sequence  $T \rightarrow A \rightarrow F$  of abelian groups, where T is a torsion group and F is a torsion-free group.

More recently, different authors have studied wider generalisations of this notion, both in pointed and in non-pointed categories (see, for example, [5, 6, 18, 13, 11, 9]). The objective of the present work is to introduce and study a torsion theory in a 2-dimensional context. One can already find a very interesting case of a torsion theory developed in the 2-category **Cat** of small categories in the recent work by Borceux, Campanini, Gran and Tholen ([4]); in that work, however, due to the use of the formalism of pretorsion theories, the 2-category **Cat** is essentially treated as a 1-category, and the 2-dimensional aspects are not fully taken into account. Our goal is instead to construct a torsion theory in a 2-category in a way such that the 2-dimensional structure is extensively incorporated into the theory. We achieve this goal by applying the formalism of *homotopy torsion theories*, recently introduced in [17].

Our starting point is a pointed torsion theory on the (1-)category CMon of commutative monoids, which was first introduced in [8]. Torsion objects

are given by abelian groups, seen as commutative monoids in which every element is invertible. As torsion free objects we have those monoids in which the only invertible element is the unit. In literature, monoids with this particular property have been variously called *pure*, *sharp* or *reduced*, and we adopt the first name. In [8] it is proven that abelian groups and pure commutative monoids form indeed a torsion theory on CMon.

The main idea in this paper is to generalise this torsion theory, moving from the 1-dimensional context of commutative monoids to the 2-dimensional one of symmetric monoidal categories. However, this generalisation is not straightforward. In the first place, the 1-category of symmetric monoidal categories (with symmetric monoidal functors) is *not* a pointed category, and therefore we do not have zero morphisms and short exact sequences in the traditional sense. Second, we know that symmetric monoidal categories actually form (with symmetric monoidal functors and monoidal natural transformations) a 2-category, and, as mentioned above, we would like to take this richer structure into account.

The framework of homotopy torsion theories allows us to address both the above issues. In this setting, we prove that symmetric 2-groups are the torsion objects of a homotopy torsion theory in the category of symmetric monoidal categories, using particular natural isomorphisms as nullhomotopies. We also show that, thanks to the use of 2-cells as nullhomotopies, this homotopy torsion theory exhibits some remarkable properties in dimension 2 which may be the basis for a notion of "2-dimensional torsion theory" in future work. We conclude the paper by carrying out an analogous generalisation for the classical torsion theory on abelian groups given by torsion and torsion-free groups, thus describing another homotopy torsion theory which enjoys the same 2-dimensional properties.

# 2 Preliminaries

In the present section we will recall some fundamental elements we will need for the new torsion theory that we want to construct.

#### 2.1 Monoidal categories and monoidal functors

Since the core idea is to move from the 1-dimensional context of monoids to the 2-dimensional context of monoidal categories, we first fix the notations and the conventions we will use for monoidal categories. For the basic theory of monoidal categories we refer to [16] or [3]. See also [15] and [14] for coherence issues. We denote the full structure of a monoidal category by

$$(\mathfrak{M}, \otimes_{\mathfrak{M}}, I_{\mathfrak{M}}, a_{\mathfrak{M}}, \ell_{\mathfrak{M}}, r_{\mathfrak{M}}),$$

where  $\mathcal{M}$  is the underlying category,  $\otimes_{\mathcal{M}}$  is the tensor,  $I_{\mathcal{M}}$  is the tensor unit,  $a_{\mathcal{M}}$ ,  $\ell_{\mathcal{M}}$ ,  $r_{\mathcal{M}}$  are the associator, the left and the right unitors respectively. We will often simply write  $(\mathcal{M}, \otimes, I, a, \ell, r)$ , or even just  $\mathcal{M}$ . When  $\mathcal{M}$  is symmetric, we will denote the symmetric braiding by  $b_{\mathcal{M}}$  (or simply b).

For a monoidal functor

$$F: \mathcal{M} \to \mathcal{N},$$

we denote by  $e_F$  and  $m_F$  (or e and m) the maps representing the coherence with the unit and the tensor product respectively, and we always assume they are isomorphisms (so monoidal functors for us will actually be strong monoidal functors).

We denote by MonCat the 2-category of small monoidal categories, monoidal functors and monoidal natural transformations, and by SymMonCat the 2-category of small symmetric monoidal categories, symmetric monoidal functors and monoidal natural transformations. Since these 2-categories are both strict, sometimes we will implicitly regard them as ordinary 1-categories, and no confusion should arise.

In what follows, the symmetry b, the structure maps a,  $\ell$  and r of monoidal categories, and the coherence maps e and m of monoidal functors will often not be explicitly written down.

#### 2.2 2-groups

If we are to generalise the torsion theory considered in the introduction, and we want to replace commutative monoids with symmetric monoidal categories, then abelian groups should be replaced by symmetric 2-groups.

Even though the notion of (weak) 2-group has appeared in literature before, it is in Hoàng Xuân Sính's doctoral thesis ([12]) that we find a first systematic study of the concept, under the name of "gr-catégories" (see also [1]). Let us recall here the definition of 2-group (weak 2-group in [2]).

**Definition 2.1.** In a monoidal category we say that an object A is weakly invertible if there exists an object B (called a weak inverse of A) such that

$$A \otimes B \cong B \otimes A \cong I.$$

A 2-group is a monoidal category  $\mathcal{G}$  such that

- every morphism in G is invertible (i.e. G is a groupoid);
- every object in G is weakly invertible.

A symmetric 2-group is a 2-group which is symmetric as a monoidal category. We denote by 2-Grp (respectively, Sym2-Grp) the 2-category of small 2groups (respectively, small symmetric 2-groups), monoidal functors (respectively, symmetric monoidal functors) and monoidal natural transformations.

We state here the following basic property that will be useful later (cf. for example [2]).

**Proposition 2.2.** In a 2-group  $\mathcal{G}$ , for any object A, the endofunctors  $A \otimes -$  and  $- \otimes A$  of  $\mathcal{G}$  are equivalences (with inverses  $B \otimes -$  and  $- \otimes B$  respectively, for any weak inverse B of A). Furthermore, it is always possible to find for each object A an object B and isomorphisms

$$h\colon I \to A \otimes B, \qquad k\colon B \otimes A \to I,$$

such that (A, B, h, k) is an adjoint equivalence, i.e. it makes the following compositions identities

$$A \longrightarrow I \otimes A \xrightarrow{h \otimes h} (A \otimes B) \otimes A \longrightarrow A \otimes (B \otimes A) \xrightarrow{A \otimes k} A \otimes I \longrightarrow A,$$
$$B \longrightarrow B \otimes I \xrightarrow{B \otimes h} B \otimes (A \otimes B) \longrightarrow (B \otimes A) \otimes B \xrightarrow{k \otimes B} I \otimes B \longrightarrow B$$

(where the unnamed arrows are components of the unitors, of the associator or of their inverses). Clearly  $(B, A, k^{-1}, h^{-1})$  is an adjoint equivalence as well.

There is a meaningful way to assign a 2-group to any monoidal category: by considering its weakly invertible objects and isomorphisms between them. We recall this in the following definition.

**Definition 2.3.** Given any monoidal category  $\mathcal{M}$ , the *Picard 2-group* of  $\mathcal{M}$ , which we denote by  $\Gamma(\mathcal{M})$ , is the subcategory of  $\mathcal{M}$  consisting of all the weakly invertible objects of  $\mathcal{M}$  and isomorphisms between them. This inherits the monoidal structure of  $\mathcal{M}$ , and is therefore a 2-group.

The Picard 2-group of a monoidal category satisfies a well-known property which we report here for future reference.

Remark 2.4. In this remark we consider the 2-categories 2-Grp, Sym2-Grp, MonCat and SymMonCat, but we regard them as 1-categories, because for the time being we are only interested in the 1-dimensional properties. 2-Grp is a coreflective subcategory of MonCat, and Sym2-Grp is a coreflective subcategory of SymMonCat. In both cases the coreflector is given by extending to 1-cells the mapping  $\mathcal{M} \mapsto \Gamma(\mathcal{M})$ , and the counit of the adjunction has components given by the inclusions  $\Gamma(\mathcal{M}) \hookrightarrow \mathcal{M}$ , for any (symmetric) monoidal category  $\mathcal{M}$ .

In the next section we want to define particular monoidal categories which have properties that are somehow diametrically opposite to those of 2-groups.

# **3** Purely monoidal categories

In this part we introduce and study the notion of "purely monoidal category", which is meant to provide a 2-dimensional equivalent of the notion of pure monoid (see the introduction). Consider the following equivalent statements about monoidal categories.

**Proposition 3.1.** Let  $\mathcal{M}$  be a monoidal category, and recall that  $\Gamma(\mathcal{M})$  is its Picard 2-group (see Definition 2.3). The following are equivalent.

- (1)  $\Gamma(\mathcal{M})$  is monoidally equivalent to 1, the monoidal category with just one object  $I_1$  and one arrow.
- (2) All weakly invertible objects of  $\mathcal{M}$  are isomorphic to  $I_{\mathcal{M}}$ , and the only automorphism of  $I_{\mathcal{M}}$  is the identity.
- (3) For any pair of weakly invertible objects, there exists exactly one isomorphism between them.

(We recall that a monoidal equivalence is simply an equivalence in the 2-category MonCat.)

*Proof.* The equivalence of (2) and (3) is obvious by the fact that  $I_{\mathcal{M}}$  is weakly invertible itself.

Now call  $\mathcal{G}$  the Picard 2-group of  $\mathcal{M}$ , and call F the only (monoidal) functor from  $\mathcal{G}$  to 1. If (1) holds, F has an inverse  $G: \mathbf{1} \to \mathcal{G}$  with a monoidal natural isomorphism  $\eta: GF \Rightarrow \operatorname{id}_{\mathcal{G}}$ . For any A in  $\mathcal{G}$ , the composition  $e_G \circ \eta_A^{-1}$ gives an isomorphism between A and  $I_{\mathcal{M}}$  (where  $e_G$  is the coherence map of G described in Section 2.1). By the naturality of  $\eta$ , this is the only map in  $\mathcal{G}$  from A to  $I_{\mathcal{M}}$ . Vice versa, if (3) holds, call  $G: \mathbf{1} \to \mathcal{G}$  the functor defined by  $GI_1 = I_{\mathcal{M}}$ . Plainly, this is a monoidal functor with the obvious coherence maps, and clearly  $FG = \operatorname{id}_1$ . For every object A in  $\mathcal{G}$ , the unique isomorphism between A and  $I_{\mathcal{M}}$  provides an isomorphism between GFA and A. The fact that these isomorphisms are the components of a monoidal natural isomorphism from GF to  $\operatorname{id}_{\mathcal{M}}$  follows from the uniqueness hypothesis in (3).

**Remark 3.2.** Observe that Proposition 3.1 can be specialised to the category SymMonCat simply by noticing that, if  $\mathcal{M}$  is a symmetric monoidal category and  $\Gamma(\mathcal{M})$  is equivalent to 1 in MonCat, then  $\Gamma(\mathcal{M})$  is equivalent to 1 in SymMonCat.

**Definition 3.3.** We say that a monoidal category  $\mathcal{M}$  is *pure* (or that  $\mathcal{M}$  is a *purely monoidal category*) if any (and hence all) of the properties in the above proposition hold. When a purely monoidal category  $\mathcal{M}$  is symmetric as a monoidal category then we say that it is a *symmetric purely monoidal category*. We denote by Pur (respectively, SymPur) the 2-category of small symmetric monoidal categories (respectively, small symmetric purely monoidal categories), monoidal functors (respectively, symmetric monoidal functors) and monoidal natural transformations.

**Example 3.4.** Cartesian and cocartesian monoidal categories are symmetric purely monoidal categories.

In fact, suppose  $\mathcal{C}$  is a category with finite products, and that for some objects A and B we have an isomorphism  $A \times B \cong 1$ , where 1 is the terminal object of  $\mathcal{C}$ . We show that A and B are also terminal.

Since  $A \times B \cong 1$ , we have that  $A \times B$  is itself terminal. Let then t be the unique map with domain A and codomain  $A \times B$ , and call  $t_1: A \to A$  and  $t_2: A \to B$  its components, so that  $t = (t_1, t_2): A \to A \times B$ . By uniqueness, we must have  $(t_1, t_2) = (\mathrm{id}_A, t_2): A \to A \times B$ , and so  $t_1 = \mathrm{id}_A$ . Consider then the product projection  $p_1: A \times B \to A$ . We have

•  $t \circ p_1 = id_{A \times B}$ , since  $A \times B$  is terminal;

•  $p_1 \circ t = t_1 = \mathrm{id}_A$ .

We conclude that  $p_1$  is an isomorphism and so A is terminal. An identical proof applies to B.

We have therefore shown that all weakly invertible objects in a cartesian monoidal category are terminal, and therefore uniquely isomorphic to the unit.

We introduce now an important property of symmetric purely monoidal categories (the proof relies however on some results we will introduce later on in this paper).

**Proposition 3.5.** The (1-)category of symmetric purely monoidal categories and symmetric monoidal functors between them is a reflective subcategory of the category SymMonCat of symmetric monoidal categories and symmetric monoidal functors.

*Proof.* In this proof we will explicitly exhibit a way to associate to any symmetric monoidal category  $\mathcal{M}$  a symmetric purely monoidal category  $\Pi(\mathcal{M})$  and a symmetric monoidal functor  $Q_{\mathcal{M}} \colon \mathcal{M} \to \Pi(\mathcal{M})$ , since we will soon need this construction. However, we will not prove here that this yields in fact a reflection, as this will be a simple consequence of a later result (see Proposition 6.2 and Corollary 6.4).

The construction we present here is based on a work by Vitale ([19]), where the existence of a certain notion of cokernel for morphisms between 2-groups is proven. We report here (an adaptation of) the construction of the cokernel in [19] applied to the inclusion of the Picard 2-group into a symmetric monoidal category, and we check that it gives rise to a purely monoidal category.

Let us first fix a symmetric monoidal category  $\mathcal{M}$  and then proceed by steps. Recall that  $\Gamma(\mathcal{M})$  is the Picard 2-group of  $\mathcal{M}$  (see Definition 2.3).

- 1. The first step is to define a bicategory  $\mathcal{B}(\mathcal{M})$  as follows.
  - The objects of  $\mathcal{B}(\mathcal{M})$  are the same as those of  $\mathcal{M}$ .
  - Given two objects X and Y in  $\mathcal{M}$ , a 1-arrow in  $\mathcal{B}(\mathcal{M})$  from X to Y is given by a pair (A, f), where A is an object of  $\Gamma(\mathcal{M})$ , and  $f: X \to Y \otimes A$  is an arrow in  $\mathcal{M}$ .


• Given two 1-arrows in  $\mathcal{B}(\mathcal{M})$ 

$$X \xrightarrow{(A,f)} Y \xrightarrow{(B,g)} Z$$

the composition is given by  $(B \otimes A, h)$ , where h is given by the following composition of arrows in  $\mathcal{M}$ .

$$X \xrightarrow{f} Y \otimes A \xrightarrow{g \otimes A} (Z \otimes B) \otimes A \xrightarrow{a_{Z,B,A}} Z \otimes (B \otimes A)$$

- The 1-identity on an object X in  $\mathcal{B}(\mathcal{M})$  is given by  $(I, r_X^{-1})$ .
- Given parallel 1-arrows (A, f),  $(A', f'): X \to Y$ , we define a 2arrow  $\alpha: (A, f) \Rightarrow (A', f')$  to be just an arrow  $\alpha: A \to A'$  in  $\Gamma(\mathcal{M})$ such that the following triangle commutes in  $\mathcal{M}$ .



- Vertical composition and 2-identities are just composition and identities in  $\Gamma(\mathcal{M})$ , and horizontal composition is just the tensor product in  $\Gamma(\mathcal{M})$ .
- The unitors and associators of  $\mathcal{B}(\mathcal{M})$  (as a bicategory) are given by the unitors and associators of  $\mathcal{M}$  (as a monoidal category).

This makes  $\mathcal{B}(\mathcal{M})$  into a bicategory where every 2-arrow is invertible (being an invertible arrow in  $\Gamma(\mathcal{M})$ ).

- Now that we have B(M), we define Π(M) as the classifying category of B(M). Explicitly, we have that Π(M) is a category having
  - the same objects as  $\mathcal{B}(\mathcal{M})$  (and therefore the same objects as  $\mathcal{M}$ );
  - as arrows, 2-isomorphism classes of 1-arrows of  $\mathcal{B}(\mathcal{M})$ .

We denote the 2-isomorphism class of a 1-arrow (A, f) in  $\mathcal{B}(\mathcal{M})$  by [A, f]. By definition, given objects X and Y in  $\mathcal{M}$ , and 1-arrows  $(A, f), (A', f'): X \to Y$  in  $\mathcal{B}(\mathcal{M})$ , we have that these determine equal arrows  $[A, f] = [A', f']: X \to Y$  in  $\Pi(\mathcal{M})$  if and only if there exists an isomorphism  $\alpha: A \to A'$  in  $\mathcal{M}$  such that the above triangle 3.6 commutes.

3. We have a functor

$$Q_{\mathcal{M}} \colon \mathcal{M} \to \Pi(\mathcal{M})$$

sending each object of  $\mathcal{M}$  to itself in  $\Pi(\mathcal{M})$ , and sending a morphism  $f: X \to Y$  in  $\mathcal{M}$  to  $[I, r_Y^{-1} \circ f]$ , where r is the left unitor:

$$X \xrightarrow{f} Y \xrightarrow{r_Y^{-1}} Y \otimes I.$$

- 4. We now introduce a symmetric monoidal structure on  $\Pi(\mathcal{M})$  and observe that  $Q_{\mathcal{M}}$  is a symmetric monoidal functor.
  - The tensor product of objects is the same as in  $\mathcal{M}$ .
  - Given two arrows  $[A, f]: X \to Y$  and  $[A', f']: X' \to Y'$  in  $\Pi(\mathfrak{M})$ , their tensor product is given by  $[A \otimes A', g]$ , where g is the following composition of maps in  $\mathfrak{M}$ .

$$\begin{array}{c} X \otimes X' \xrightarrow{f \otimes f'} & (Y \otimes A) \otimes (Y' \otimes A') \\ & \checkmark \\ Y \otimes \left( (A \otimes Y') \otimes A' \right) \xrightarrow{Y \otimes (b_{A,Y'} \otimes A')} & Y \otimes \left( (Y' \otimes A) \otimes A' \right) \\ & \checkmark \\ & \swarrow \\ & (Y \otimes Y') \otimes (A \otimes A') \end{array}$$

(We recall that b is the symmetry, and the unnamed winding arrows are compositions of the appropriate components of the associator of  $\mathcal{M}$ .)

• The associator, the unitors and the braiding of  $\Pi(\mathcal{M})$  are just the image under  $Q_{\mathcal{M}}$  of the corresponding maps of  $\mathcal{M}$ .

With this additional structure, we have made  $\Pi(\mathcal{M})$  into a symmetric monoidal category. The symmetry of  $\mathcal{M}$  (and not just the braiding) is used to prove the naturality of the braiding of  $\Pi(\mathcal{M})$ .

5. Now we just need to check that  $\Pi(\mathcal{M})$  is pure. We use a property that we state and prove separately in Lemma 3.7 below, as we will need it again later.

In order to verify that  $\Pi(\mathcal{M})$  is pure we prove condition (2) of Proposition 3.1. Consider a weakly invertible object A of  $\mathcal{M}$ . Then  $[A, \ell_A^{-1}]$ is a morphism from A to I in  $\Pi(\mathcal{M})$ , and it is also an isomorphism by Lemma 3.7 (we recall that  $\ell$  is the left unitor).

Now, suppose

 $[A, f], [B, g] \colon I \to I$ 

are isomorphisms in  $\Pi(\mathcal{M})$ . Then, again by Lemma 3.7, f and g are isomorphisms in  $\mathcal{M}$ . Call  $\phi = \ell_B \circ g \circ f^{-1} \circ \ell_A^{-1}$ . By naturality of  $\ell$  we have  $(I \otimes \phi) \circ f = g$ , yielding [A, f] = [B, g] (by definition).

**Lemma 3.7.** Consider a symmetric monoidal category  $\mathcal{M}$  and the associated functor  $Q_{\mathcal{M}} \colon \mathcal{M} \to \Pi(\mathcal{M})$  described in Proposition 3.5. Then a morphism  $[A, f] \colon X \to Y$  in  $\Pi(\mathcal{M})$  is an isomorphism if and only if f is an isomorphism in  $\mathcal{M}$ . In particular,  $Q_{\mathcal{M}}$  reflects isomorphisms.


*Proof.* As reported in [19], if f is an isomorphism in  $\mathcal{M}$ , and we consider an adjoint equivalence (A, B, h, k) (see Proposition 2.2), then an inverse of [A, f] in  $\Pi(\mathcal{M})$  is [B, g], where g is the composition of the following maps.

$$Y \xrightarrow{r_Y} Y \otimes I \xrightarrow{Y \otimes h} Y \otimes (A \otimes B) \xrightarrow{}_{a_{Y,A,B}^{-1}} (Y \otimes A) \otimes B \xrightarrow{}_{f^{-1} \otimes B} X \otimes B$$

Vice versa, suppose [A, f] is an isomorphism in  $\Pi(\mathcal{M})$ , with inverse [B, g]. Then, starting from the fact that  $[B, g] \circ [A, f] = \operatorname{id}_X$  in  $\Pi(\mathcal{M})$  and spelling out the definition of composition and identity in  $\Pi(\mathcal{M})$ , we get that in  $\mathcal{M}$ the map f has a left inverse and the map  $g \otimes A$  has a right inverse. Since  $- \otimes A$  is an equivalence (see Proposition 2.2), and equivalences preserve and reflect split monomorphisms and split epimorphisms, we conclude that g has a right inverse as well. Conversely, applying the same reasoning to the reverse composition  $[A, f] \circ [B, g] = \operatorname{id}_Y$ , we conclude that g has a left inverse and fhas a right inverse. Since f has both left and right inverses, it must be an isomorphism.

Finally, to see that  $Q_{\mathcal{M}}$  reflects isomorphisms consider an arrow  $f: X \to Y$  in  $\mathcal{M}$ , and suppose that  $Q_{\mathcal{M}}(f) = [I, r_Y^{-1} \circ f]$  is an isomorphism in  $\Pi(\mathcal{M})$ . Then, by the first part of this corollary,  $r_Y^{-1} \circ f$  is an isomorphism in  $\mathcal{M}$ , and therefore f also is.  $\Box$  We conclude the section with another useful property about the construction of Proposition 3.5. This property is exactly the content of Lemma 2.2 in [19], except for the fact that the lemma in [19] is given in the context of 2-groups; however, it is easily proven to work in our context as well.

**Lemma 3.8.** Let  $\mathcal{M}$  be a symmetric monoidal category, and  $Q_{\mathcal{M}} \colon \mathcal{M} \to \Pi(\mathcal{M})$  the associated functor described in Proposition 3.5. Then, for any arrow  $[A, f] \colon X \to Y$  in  $\Pi(\mathcal{M})$ , the following diagram is commutative,



where  $\zeta_A = [A, \ell_A^{-1}] \colon A \to I.$ 

# 4 Homotopy torsion theories

As mentioned in the introduction we want to set our new torsion theory in the framework of homotopy torsion theories. Therefore in this section we want to recall the essential notions we are going to need (refer to [17] for more details).

We begin with the basic definitions of nullhomotopy structure and category with nullhomotopies, which were first introduced (in a slightly different form) by Grandis in [10].

**Definition 4.1.** Let  $\mathcal{C}$  be any category. A nullhomotopy structure  $\Theta$  on  $\mathcal{C}$  is given by the following data.

- 1) For every arrow f in  $\mathfrak{C}$ , a set  $\Theta(f)$ , called the set of nullhomotopies on f;
- 2) for any triple of composable arrows  $\cdot \xrightarrow{p} \cdot \xrightarrow{f} \cdot \xrightarrow{q} \cdot$ , a function

$$q \bullet - \bullet p \colon \Theta(f) \to \Theta(q \circ f \circ p)$$

such that the following axioms are satisfied.

a) Given  $f: X \to Y$  and  $\vartheta \in \Theta(f)$ , we have that

$$\operatorname{id}_Y \bullet \vartheta \bullet \operatorname{id}_X = \vartheta;$$

b) given a quintuple of composable maps  $\cdot \xrightarrow{p'} \cdot \xrightarrow{p} \cdot \xrightarrow{f} \cdot \xrightarrow{q} \cdot \xrightarrow{q'} \cdot$ , and given  $\vartheta \in \Theta(f)$ , then we have that

$$(q' \circ q) \bullet \vartheta \bullet (p' \circ p) = q' \bullet (q \bullet \vartheta \bullet p) \bullet p'.$$

A category with nullhomotopies is given by a pair  $(\mathcal{C}, \Theta)$ , where  $\mathcal{C}$  is a category and  $\Theta$  is a nullhomotopy structure on  $\mathcal{C}$ .

Given a category with nullhomotopies  $(\mathcal{C}, \Theta)$ , we say that a morphism fin  $\mathcal{C}$  is  $\Theta$ -trivial (or nullhomotopic) if  $\Theta(f)$  is non-empty. We say instead that an object in  $\mathcal{C}$  is  $\Theta$ -trivial if its corresponding identity morphism is  $\Theta$ -trivial. Finally, given two objects X and Y in  $\mathcal{C}$ , we say that X is  $\Theta$ -orthogonal to Y (in symbols,  $X \perp_{\Theta} Y$ ) if for all arrows  $f: X \to Y$  we have that  $\Theta(f)$  is a singleton.

Notation 4.2. In this section, to visually represent a nullhomotopy  $\vartheta$  on a map f, we will use the notation




as if  $\vartheta$  was a 2-cell in a 2-category from f to a phantom 0-arrow (of course, in general, we may not be in a 2-category and we may not have a 0-arrow).

Moreover, in light of Axioms a) and b), given maps  $f: X \to Y$  and  $g: Y \to Z$ , and nullhomotopies  $\phi \in \Theta(f)$  and  $\psi \in \Theta(g)$ , we write  $g \bullet \phi$  for  $g \bullet \phi \bullet \operatorname{id}_X$  and  $\psi \bullet f$  for  $\operatorname{id}_Z \bullet \psi \bullet f$ .


Next, we introduce the natural notion of kernel and cokernel in this context, i.e. the notion of (strong) homotopy kernel and (strong) homotopy cokernel.

**Definition 4.3.** Let  $(\mathcal{C}, \Theta)$  be a category with nullhomotopies, and let  $f: X \to Y$  be an arrow in  $\mathcal{C}$ . A homotopy kernel of f with respect to  $\Theta$  (or a  $\Theta$ -kernel of f) is a triple  $(K, k, \vartheta)$ , where  $k: K \to X$  is a map in  $\mathcal{C}$  and  $\vartheta \in \Theta(f \circ k)$ , with the property that given any other triple  $(H, h, \phi)$ , with  $h: H \to X$  in  $\mathcal{C}$ 

and  $\phi \in \Theta(f \circ h)$ , there exists a unique map  $h' \colon H \to K$  such that  $k \circ h' = h$ and  $\vartheta \bullet h' = \phi$ .



We say that  $(K, k, \vartheta)$  is strong if, additionally, for every map  $a: A \to K$  and every nullhomotopy  $\phi \in \Theta(k \circ a)$  such that  $f \bullet \phi = \vartheta \bullet a$ , there exists a unique nullhomotopy  $\alpha \in \Theta(a)$  such that  $k \bullet \alpha = \phi$ .



We will sometimes just write K or k to denote the (strong)  $\Theta$ -kernel  $(K, k, \vartheta)$ . Of course (strong) homotopy cokernels are defined dually.

A homotopy exact sequence in  $(\mathcal{C}, \Theta)$  (or a  $\Theta$ -exact sequence) will be given by a pair of composable morphisms  $X \xrightarrow{f} Y \xrightarrow{g} Z$  and a nullhomotopy  $\vartheta \in \Theta(g \circ f)$  such that  $(X, f, \vartheta)$  is the  $\Theta$ -kernel of g and  $(Z, g, \vartheta)$  is the  $\Theta$ cokernel of f.

We now have all the elements to define homotopy torsion theories.

**Definition 4.4.** Let  $(\mathcal{C}, \Theta)$  be a category with nullhomotopies. A homotopy torsion theory on  $(\mathcal{C}, \Theta)$  (or a  $\Theta$ -torsion theory on  $\mathcal{C}$ ) is given by a pair  $(\mathcal{T}, \mathcal{F})$  of full and replete subcategories of  $\mathcal{C}$  such that

- 1. given objects T in  $\mathfrak{T}$  and F in  $\mathfrak{F}$ , then T is  $\Theta$ -orthogonal to F (see Definition 4.1);
- 2. for every object X in C there exists a  $\Theta$ -exact sequence

$$T_X \xrightarrow{\vartheta_X \Uparrow} X \xrightarrow{f_X} F_X$$

(see Definition 4.3), with  $T_X$  in  $\mathfrak{T}$  and  $F_X$  in  $\mathfrak{F}$ .

# 5 A nullhomotopy structure $\Theta$ for monoidal categories

Now let us go back to monoidal categories. In Sections 2 and 3 we considered two special classes of (symmetric) monoidal categories (namely 2-groups and purely monoidal categories) which could play the role of the torsion and torsion-free objects in the new torsion theory we are trying to develop. However, in order to construct a torsion theory, we need get ourselves a notion of trivial map and of exact sequence, and of course we make use of the machinery just described in Section 4: we want to endow the category of (symmetric) monoidal categories with a nullhomotopy structure, i.e., we want to define a set of nullhomotopies on any given (symmetric) monoidal functor.

We know that between any two monoidal categories we always have the constant functor at the unit object of the codomain. This is not a zero (1-)morphism (since MonCat does not have a zero object), however we will use it as a "reference trivial morphism", meaning that we will consider trivial any functor naturally isomorphic to it, and use these natural isomorphims as nullhomotopies. More precisely, we give the following definition.

**Definition 5.1.** Given any monoidal functor  $F: \mathcal{M} \to \mathcal{N}$ , define the set

 $\Theta(F) = \{\phi \colon F \Rightarrow \Delta I_{\mathcal{N}} \mid \phi \text{ is a monoidal natural isomorphism} \}$ 

(where  $\Delta I_{\mathcal{N}}$  is the constant functor at  $I_{\mathcal{N}}$ ). Given monoidal functors

$$\mathcal{M}' \xrightarrow{P} \mathcal{M} \xrightarrow{F} \mathcal{N} \xrightarrow{Q} \mathcal{N}'$$

and given  $\phi \in \Theta(F)$ , we define the composition

$$Q \bullet \phi \bullet P \in \Theta(QFP),$$

having components

$$(Q \bullet \phi \bullet P)_X = e_Q^{-1} \circ Q(\phi_{PX})$$

for all X in  $\mathcal{M}'$ , where we recall that  $e_Q$  is the coherence map of Q defined in Section 2.1 (in other words,  $Q \bullet \phi \bullet P$  is the natural transformation obtained by vertically composing the horizontal composition  $\mathrm{id}_Q \circ \phi \circ \mathrm{id}_P$  with the constant natural transformation  $e_Q^{-1} \colon \Delta(QI_N) \Rightarrow \Delta I_{\mathcal{N}'} \colon \mathcal{M}' \to \mathcal{N}'$ ). **Remark 5.2.** It is easy to verify that the definition of  $\Theta$  we have just given makes (MonCat,  $\Theta$ ) into a category with nullhomotopies. Clearly, if we restrict the definition of  $\Theta$  to symmetric monoidal functors, we get a nullhomotopy structure on SymMonCat (which we still call  $\Theta$ ), making (SymMonCat,  $\Theta$ ) a category with nullhomotopies as well.

From now on, unless otherwise specified,  $\Theta$  will always denote the specific nullhomotopy structure of Definition 5.1.

# 6 A Θ-torsion theory for symmetric monoidal categories

Thus far we have laid out the ingredients for our 2-dimensional generalisation of the torsion theory for commutative monoids given in the introduction in the following way.

Commutative monoids	$\Rightarrow$	Symmetric monoidal categories
Trivial (=constant) maps	$\Rightarrow$	$\Theta$ -trivial (=naturally isomorphic to constant) functors
Abelian groups	$\implies$	Symmetric 2-groups
Commutative pure monoids	$\Rightarrow$	Symmetric purely monoidal categories

We now need to check that these ingredients do produce the desired result. We do so in Proposition 6.2, after proving the following preparatory lemma.

**Lemma 6.1.** Let  $F: \mathcal{M} \to \mathcal{N}$  be a (symmetric) monoidal functor between (symmetric) monoidal categories. If  $\mathcal{N}$  is pure, then  $\Theta(F)$  has at most one element. If in addition  $\mathcal{M}$  is a 2-group, then  $\Theta(F)$  has exactly one element.

Proof. Suppose  $\mathcal{N}$  is pure. If there exists a nullhomotopy  $\phi \in \Theta(F)$ , then  $\phi$  is unique because its components are isomorphisms between weakly invertible objects (in fact, for each object X in  $\mathcal{M}$  the map  $\phi_X$  is an isomorphism between FX and  $I_{\mathcal{N}}$ , and since  $I_{\mathcal{N}}$  is weakly invertible, FX must also be), and there exists exactly one isomorphism between every given pair of weakly invertible objects in a purely monoidal category. If we know that  $\mathcal{M}$  is a 2-group, then for every object X in  $\mathcal{M}$  we have that FX is weakly invertible in  $\mathcal{N}$  (as X is weakly invertible in  $\mathcal{M}$ ), and so there is a unique isomorphism  $\phi_X \colon FX \to I_{\mathcal{N}}$ . The maps  $\phi_X$  with X in  $\mathcal{M}$ give rise to a monoidal natural isomorphism between F and  $\Delta I_{\mathcal{N}}$  (naturality and monoidality follow from the uniqueness of isomorphisms between weakly invertible objects in  $\mathcal{N}$ ).

#### **Proposition 6.2.** (Sym2-Grp, SymPur) is a $\Theta$ -torsion theory on SymMonCat.

Proof. We need to verify that Axioms 1 and 2 of Definition 4.4 are satisfied.

Axiom 1 ( $\Theta$ -orthogonality of symmetric 2-groups and symmetric purely monoidal categories) is part of Lemma 6.1. To prove Axiom 2, fix a symmetric monoidal category  $\mathcal{M}$  and consider the following sequence:


(6.3) 
$$\Gamma(\mathcal{M}) \xrightarrow{J_{\mathcal{M}}} \mathcal{M} \xrightarrow{Q_{\mathcal{M}}} \Pi(\mathcal{M}),$$

where  $J_{\mathcal{M}} \colon \Gamma(\mathcal{M}) \to \mathcal{M}$  is the inclusion of the Picard 2-group of  $\mathcal{M}$  in  $\mathcal{M}$  itself (see Definition 2.3), and  $Q_{\mathcal{M}} \colon \mathcal{M} \to \Pi(\mathcal{M})$  is the functor constructed in the proof of Proposition 3.5. We know  $\Gamma(\mathcal{M})$  is a symmetric 2-group and that  $\Pi(\mathcal{M})$  is a symmetric purely monoidal category. Therefore we just need to check that 6.3 is a  $\Theta$ -exact sequence (see Definition 4.3). Call  $\zeta_{\mathcal{M}}$  the unique nullhomotopy on  $Q_{\mathcal{M}} \circ J_{\mathcal{M}}$ . By uniqueness, its components are given, for any object A in  $\Gamma(\mathcal{M})$ , by

$$(\zeta_{\mathcal{M}})_A \equiv \zeta_A = [A, \ell_A^{-1}] \colon A \to I.$$

Let us now give an outline of the proofs of the required universal properties for the homotopy exactness of 6.3.

First, let us check that  $(\Gamma(\mathcal{M}), J_{\mathcal{M}}, \zeta_{\mathcal{M}})$  is the  $\Theta$ -kernel of  $Q_{\mathcal{M}}$ . Suppose we are in the following situation,



where  $F: \mathcal{L} \to \mathcal{M}$  is a symmetric monoidal functor,  $\phi \in \Theta(Q_{\mathcal{M}} \circ F)$  and the unnamed arrows are constant functors at the units of the respective codomains. We need to prove that there exists a unique symmetric monoidal functor  $F': \mathcal{L} \to \Gamma(\mathcal{M})$  such that  $J_{\mathcal{M}} \circ F' = F$  and  $\zeta_{\mathcal{M}} \bullet F' = \phi$ . We actually just need to prove that for all object X in  $\mathcal{L}$  we have that FX is weakly invertible, and that for all morphisms f in  $\mathcal{L}$  we have that Ff is an isomorphism, so that F lifts to the Picard 2-group of its codomain. Uniqueness then follows from the fact that  $J_{\mathcal{M}}$  is an inclusion, and the condition on the nullhomotopy is trivial because there can exist at most one nullhomotopy on any functor into a pure monoidal category by Lemma 6.1. So let X be an object in  $\mathcal{L}$ . Then  $\phi_X$  gives us an isomorphism between  $Q_{\mathcal{M}}(F(X)) = F(X)$ and I in  $\Pi(\mathcal{M})$ . This isomorphism will be given by some  $[A, u]: FX \to I$ , with A a weakly invertible object in  $\mathcal{M}$  and  $u: FX \to I \otimes A$  an arrow in  $\mathcal{M}$ . By Lemma 3.7, u is actually an isomorphism in  $\mathcal{M}$ , yielding


$$FX \cong I \otimes A \cong A$$

with A weakly invertible. We conclude that FX is weakly invertible in  $\mathcal{M}$ . If  $f: X \to Y$  is a morphism in  $\mathcal{L}$ , then, the naturality of  $\phi$  gives us

$$\phi_Y \circ Q_{\mathcal{M}}(F(f)) = \phi_X$$


We conclude that  $Q_{\mathcal{M}}(F(f))$  is an isomorphism. Since, by Lemma 3.7,  $Q_{\mathcal{M}}$  reflects isomorphisms, we are done.

Next, we check that  $(\Pi(\mathcal{M}), Q_{\mathcal{M}}, \zeta_{\mathcal{M}})$  is the  $\Theta$ -cokernel of  $J_{\mathcal{M}}$ . Consider the following situation.



Here  $G: \mathcal{M} \to \mathcal{N}$  is a symmetric monoidal functor with a nullhomotopy  $\psi \in \Theta(G \circ J_{\mathcal{M}})$ ; the unnamed arrows are the constant functor at the appropriate unit objects. We need to show that there exists a unique symmetric monoidal functor  $G': \Pi(\mathcal{M}) \to \mathcal{N}$  such that  $G' \circ Q_{\mathcal{M}} = G$  and  $G' \bullet \zeta_{\mathcal{M}} = \psi$ . The first condition determines G' uniquely on objects: given any X in  $\mathcal{M}$  we must have

 $G'X = G'(Q_{\mathfrak{M}}(X)) = GX$ . To determine the action of G' on arrows we rely on Lemma 3.8. By the lemma, we have that for any arrow  $[A, f]: X \to Y$  in  $\Pi(\mathfrak{M}), G'([A, f])$  must necessarily be the composition of the following arrows in  $\mathfrak{N}$  (recall the definition of the coherence map  $m_G$  in Section 2.1).




(Use also that  $(m_{G'})_{X,Y}$  must be equal to  $(m_G)_{X,Y}$  for all X and Y in  $\mathcal{M}$ , by the fact that  $G = G' \circ Q_{\mathcal{M}}$  and by definition of composition of monoidal functors.) It is now routine to check that this defines G' uniquely as a symmetric monoidal functor satisfying the required properties.

Corollary 6.4. In this corollary, we regard the 2-categories Sym2-Grp, SymPur and SymMonCat just as 1-categories. By Corollary 7.8 in [17] we have that Sym2-Grp is a coreflective subcategory of SymMonCat and SymPur is a reflective subcategory of SymMonCat, thus confirming Remark 2.4 and completing the proof of Proposition 3.5. We denote by  $\Gamma$  the right adjoint to the inclusion Sym2-Grp  $\hookrightarrow$  SymMonCat, and by  $\Pi$  the left adjoint to the inclusion SymPur  $\hookrightarrow$  SymMonCat (in fact they are obtained as an extension to 1-cells of the constructions  $\Gamma$  and  $\Pi$  described in Definition 2.3 and Proposition 3.5 respectively).

## 7 2-dimensional aspects

We would now like to highlight a few properties that give the homotopy torsion theory just defined a true 2-dimensional flavour.

1. Nullhomotopies on a given symmetric monoidal functor  $F: \mathcal{M} \to \mathcal{N}$  are given by monoidal natural isomorphisms between F and the constant functor  $\Delta I_{\mathcal{N}}$ ; this nullhomotopy structure is the key ingredient which allows us to incorporate the 2-dimensional structure of SymMonCat into the theory. One should also remark that even if the category SymMonCat is not pointed as a 1-category, the one-object-one-arrow category 1 is a 2-terminal and bi-initial object in SymMonCat. 2. The Θ-kernels and Θ-cokernels that we have described in Proposition 6.2 satisfy an additional universal property at level 2 that actually makes them bipullbacks and bipushouts respectively. This comes as no surprise, since this 2-dimensional universal property (or a variant of it) is actually required by definition by Vitale in [19], in the context of symmetric 2-groups. To be precise, in the 2-category SymMonCat of symmetric monoidal categories, we find that for any symmetric monoidal category M the following diagram



is both a bipullback and a bipushout. The proofs of the universal properties are essentially given in [19] (even if kernels and cokernels are not explicitly described as bipullbacks and bipushouts in [19]).

3. The adjunctions considered in Corollary 6.4 can actually be promoted to 2-adjunctions by appropriately extending the functors  $\Gamma$  and  $\Pi$  to 2-cells (the key fact needed to prove this property being that the components along weakly invertible elements of a monoidal natural transformation are always isomorphisms).

These properties (which are also shared by the torsion theory to be introduced in Section 11) might be the basis for a definition of "2-dimensional torsion theory".


### 8 Homotopy invariants

In this section we discuss how the "1-dimensional" torsion theory on CMon described in the introduction is related to the homotopy torsion theory we described on SymMonCat. We denote by  $\gamma$  and  $\varpi$  the torsion and torsion-free functors associated to the torsion theory in CMon, so that for a commutative monoid M we have the following exact presentation.

$$\gamma(M) \xrightarrow{j_M} M \xrightarrow{q_M} \varpi(M)$$

(In particular  $\gamma(M)$  is the Picard group of M, i.e. the subgroup of invertible elements of M.)


Consider the functor  $K: \mathsf{Cat} \to \mathsf{Set}$  taking each (small) category to the set of isomorphism classes of its objects. If X is an object in a category, we denote by  $[\![X]\!]$  the isomorphism class containing X. The functor K can be specialised to functors from MonCat to Mon and from SymMonCat to CMon (we call them again K), making the following diagram commute



(the horizontal arrows are forgetful functors). For a monoidal category  $\mathcal{M}$  and objects X, Y in  $\mathcal{M}$ , the product of  $\llbracket X \rrbracket$  and  $\llbracket Y \rrbracket$  in  $K\mathcal{M}$  is given by  $\llbracket X \rrbracket \cdot \llbracket Y \rrbracket = \llbracket X \otimes_{\mathcal{M}} Y \rrbracket$ , and the unit is  $\llbracket I_{\mathcal{M}} \rrbracket$ . Clearly, if we started off with a symmetric monoidal category we would end up with a commutative monoid.

It is easy to check that the functor K "commutes" with the functors  $\Gamma$  and  $\Pi$ , meaning that we have natural isomorphisms  $K \circ \Gamma \cong \gamma \circ K$  and  $K \circ \Pi \cong \varpi \circ K$ . Hence, given a symmetric monoidal category  $\mathfrak{M}$ , we have three "homotopy invariants",  $K\Gamma \mathfrak{M} \cong \gamma K \mathfrak{M}$ ,  $K \mathfrak{M}$ , and  $K\Pi \mathfrak{M} \cong \varpi K \mathfrak{M}$ , which give rise to an exact sequence of commutative monoids, as made precise in the following proposition.

**Proposition 8.1.** Let  $\mathcal{M}$  be a symmetric monoidal category. Then we have the following isomorphism of short exact sequences of monoids.




If we call  $\operatorname{Pic}(\mathcal{M}) = K\Gamma\mathcal{M} \cong \gamma K\mathcal{M}$  (this is the standard Picard group of  $\mathcal{M}$ ), and  $\operatorname{Pure}(\mathcal{M}) = K\Pi\mathcal{M} \cong \varpi K\mathcal{M}$ , then the short exact sequence of commutative monoids

 $\operatorname{Pic}(\mathcal{M}) \longrightarrow K\mathcal{M} \longrightarrow \operatorname{Pure}(\mathcal{M})$ 

is a "homotopy invariant" of  $\mathcal{M}$ , in the sense that symmetric monoidal equivalences induce isomorphisms between the corresponding exact sequences.


We can also consider the following pairs of functors.



- $\pi_0$  takes each monoidal category  $\mathcal{M}$  to the monoid  $\pi_0(\mathcal{M})$  having as elements the connected components of  $\mathcal{M}$ . For objects X and Y, if we denote by [X] and [Y] the connected components containing X and Y, then the product is given by  $[X] \cdot [Y] = [X \otimes_{\mathcal{M}} Y]$ , and the unit is given by  $[I_{\mathcal{M}}]$ .
- $\iota_0$  takes each monoid M to the discrete monoidal category  $\iota_0(M)$  having the elements of M as objects and tensor product and unit given by the product and unit of M.
- $\pi_1$  takes a monoidal category  $\mathcal{M}$  to the monoid  $\pi_1(\mathcal{M})$ , consisting of the endomorphisms of  $I_{\mathcal{M}}$ , and with product given by composition and unit obviously given by  $\mathrm{id}_{I_{\mathcal{M}}}$ . This is always a commutative monoid.
- $\iota_1$  takes a commutative monoid M to the symmetric monoidal category  $\iota_1(M)$  having exactly one object whose endomorphisms are given by the elements of M. Composition and tensor product in  $\iota_1 M$  are both simply given by the product in M.

Applying  $\pi_0$  and  $\pi_1$  to  $\Theta$ -exact sequence of the form 6.3 does not yield in general exact sequences in CMon. However the following holds.

**Proposition 8.2.** For any commutative monoid M denote by  $\mathcal{M}_k$  the symmetric monoidal category  $\iota_k M$ , for  $k \in \{0, 1\}$ . We have the following natural isomorphisms of short exact sequences, for  $k \in \{0, 1\}$ .



In other words, the short exact sequence associated to M in the torsion theory on CMon can be obtained in two ways from homotopy exact sequences

arising in the homotopy torsion theory on SymMonCat. Furthermore, for any commutative monoid M, we have that M is an abelian group if and only if  $\iota_0(M)$  is a 2-group, if and only if  $\iota_1(M)$  is a 2-group; similarly, M is a pure monoid if and only if  $\iota_0(M)$  is a purely monoidal category, if and only if  $\iota_1(M)$  is a purely monoidal category, if and only if  $\iota_1(M)$  is a purely monoidal category.

# 9 Homotopy torsion theories and pretorsion theories

In this section we want to compare the homotopy torsion theory that we have described with another notion of non-pointed torsion theory existing in literature.

Let us start by considering the subcategory Z of  $\Theta$ -trivial monoidal categories, i.e. of monoidal categories whose identity functor is  $\Theta$ -trivial. We want to study a few of its properties.

First of all, reasoning as in the proof of Proposition 3.1, it is easy to see that a monoidal category  $\mathcal{Z}$  is in Z if and only if for any pair of objects X and Y in Z there exists exactly one morphism from X to Y. Therefore, we have that every category in Z is automatically symmetric with a unique possible symmetry. The following remark is also obvious.

**Remark 9.1.** For any monoidal category  $\mathcal{M}$  and any  $\Theta$ -trivial monoidal category  $\mathcal{Z}$ , we have  $\mathcal{M} \perp_{\Theta} \mathcal{Z}$  and  $\mathcal{Z} \perp_{\Theta} \mathcal{M}$  (see Definition 4.1).

Next, we want to study  $\Theta$ -kernels and  $\Theta$ -cokernels of the identity morphisms in MonCat and SymMonCat. We do so in the following proposition.

**Proposition 9.2.** Identity morphisms in MonCat and SymMonCat have strong  $\Theta$ -kernels and strong  $\Theta$ -cokernels that lie in Z.

*Proof.* Let us fix a monoidal category  $\mathcal{M}$ ; we construct the  $\Theta$ -kernel and  $\Theta$ -cokernel of  $\mathrm{id}_{\mathcal{M}}$ .

The construction of the kernel is based on the one we find in [19] for symmetric 2-groups. Consider the monoidal category  $S(\mathcal{M})$  defined as follows.

- $\operatorname{obj}(S(\mathcal{M})) = \{(X, x) \mid X \in \operatorname{obj}(\mathcal{M}), x \colon X \to I_{\mathcal{M}} \text{ isomorphism}\};$
- for (X, x) and (Y, y) in  $obj(S(\mathcal{M}))$ , set

 $\operatorname{Hom}_{S(\mathcal{M})}((X, x), (Y, y)) = \{f \colon X \to Y \mid f \circ x = y\} = \{y \circ x^{-1}\};$ 

- the identity morphism on (X, x) is just  $id_X$  in  $\mathcal{M}$ ;
- for (X, x) and (Y, y) in  $obj(S(\mathcal{M}))$ , define their tensor product as

$$(X, x) \otimes_{S(\mathcal{M})} (Y, y) = (X \otimes_{\mathcal{M}} Y, i_{\mathcal{M}} \circ (x \otimes_{\mathcal{M}} y)),$$

where  $i_{\mathcal{M}} = (\ell_{\mathcal{M}})_{I_{\mathcal{M}}} = (r_{\mathcal{M}})_{I_{\mathcal{M}}};$ 

- given morphisms in  $S(\mathcal{M})$ , their tensor product in  $S(\mathcal{M})$  is just their tensor product in  $\mathcal{M}$ ;
- the unit element in  $S(\mathcal{M})$  is  $(I_{\mathcal{M}}, \mathrm{id}_{I_{\mathcal{M}}})$ ;
- the associators and unitors in  $S(\mathcal{M})$  are given by the corresponding maps in  $\mathcal{M}$ .

The monoidal category just defined is clearly  $\Theta$ -trivial (all hom-sets are singletons). We have a monoidal functor  $\epsilon_{\mathfrak{M}} \colon S(\mathfrak{M}) \to \mathfrak{M}$ , sending any morphism  $f \colon (X, x) \to (Y, y)$  in  $S(\mathfrak{M})$  to  $f \colon X \to Y$  in  $\mathfrak{M}$  (and this functor is symmetric when  $\mathfrak{M}$  is symmetric). Since  $S(\mathfrak{M})$  is trivial, by 9.1 there exists a unique nullhomotopy—call it  $\kappa$ —on  $\epsilon_{\mathfrak{M}}$ . One can easily prove that the triple  $(S(\mathfrak{M}), \epsilon_{\mathfrak{M}}, \kappa)$  is a  $\Theta$ -kernel of id\_{\mathfrak{M}}. Since  $S(\mathfrak{M})$  is trivial, one also shows (using 9.1) that this kernel is strong.

The construction of the  $\Theta$ -cokernel of  $\mathrm{id}_{\mathcal{M}}$  is much simpler. Call  $R(\mathcal{M})$  the unique monoidal category having the same objects and unit as  $\mathcal{M}$ , exactly one morphism between any pair of objects, and tensor product on objects equal to that of  $\mathcal{M}$ . Clearly  $R(\mathcal{M})$  is  $\Theta$ -trivial. There exists a unique functor  $\eta_{\mathcal{M}} \colon \mathcal{M} \to R(\mathcal{M})$  acting as the identity on objects; this functor is automatically monoidal (and symmetric when  $\mathcal{M}$  is symmetric). Again, since  $R(\mathcal{M})$  is trivial, by 9.1 there exists a unique nullhomtopy—call it  $\lambda$ —on  $\eta_{\mathcal{M}}$ . Then, it is easy to prove that the triple  $(R(\mathcal{M}), \eta_{\mathcal{M}}, \lambda)$  is a strong  $\Theta$ -cokernel of  $\mathrm{id}_{\mathcal{M}}$ .  $\Box$ 

In light of Remark 9.1 and Proposition 9.2, we can use Proposition 6.5 in [17] to deduce the following result.

**Proposition 9.3.** The category Z of monoidal categories that are monoidally equivalent to the terminal category 1 is a reflective and coreflective subcategory of the category MonCat of monoidal categories and (strong) monoidal functors, and of the category SymMonCat of symmetric monoidal categories and (strong) symmetric monoidal functors. For each monoidal category  $\mathcal{M}$ , we have that the morphisms  $\eta_{\mathcal{M}} \colon \mathcal{M} \to R(\mathcal{M})$  and  $\epsilon_{\mathcal{M}} \colon S(\mathcal{M}) \to \mathcal{M}$  described in 9.2 are, respectively, the reflection and the coreflection of  $\mathcal{M}$  into Z. In particular, we can also conclude that the  $\Theta$ -kernel of any monoidal functor  $F: \mathcal{M} \to \mathcal{N}$  can be obtained as the (1-categorical) pullback of F along  $\epsilon_{\mathcal{N}}$ , and its  $\Theta$ -cokernel as the pushout of F along  $\eta_{\mathcal{M}}$ . Notice however that  $\Theta$ -kernels and  $\Theta$ -cokernels are in general different notions from those of Zprekernel and Z-precokernel as defined in [9], and (Sym2-Grp, SymPur) is not a pretorsion theory according again to [9]. In fact, the morphisms defining prekernels are always monomorphisms (see Proposition 2.1 in [9]), and dually the morphims defining precokernels are always epimorphisms. However, that is not true for  $\Theta$ -kernels and  $\Theta$ -cokernels, and, in particular, even the  $Q_{\mathcal{M}}$ 's appearing in the  $\Theta$ -exact presentations given for the  $\Theta$ -torsion theory (Sym2-Grp, SymPur) are not always epimorphisms (instead, the  $J_{\mathcal{M}}$ 's happen to always be monomorphisms). Check out the following example.

**Example 9.4.** Let  $C = \{1, -1\}$  be the 2-element (multiplicative) group, and let  $\mathcal{M} = \iota_0(C)$  and  $\mathcal{N} = \iota_1(C)$  (see Section 8). Notice that there exists a unique (monoidal) functor H from  $\mathcal{M}$  to  $\mathcal{N}$ .

Since  $\mathcal{M}$  is a 2-group, the  $\Theta$ -presentation of  $\mathcal{M}$  is given by

$$\Gamma(\mathcal{M}) = \mathcal{M} \xrightarrow{\mathrm{id}_{\mathcal{M}}} \mathcal{M} \xrightarrow{Q_{\mathcal{M}} = \eta_{\mathcal{M}}} \Pi(\mathcal{M}) = S(\mathcal{M}).$$

Call  $\Pi(\mathcal{M}) = S(\mathcal{M})$  just  $\mathcal{P}$ , and  $Q_{\mathcal{M}} = \eta_{\mathcal{M}}$  just P. The category  $\mathcal{P}$  has just two objects and one map between any pair of objects. There are exactly two functors from  $\mathcal{P}$  to  $\mathcal{N}$ : one—call it F—sending the two non-identity isomorphisms to 1 in  $\mathcal{N}$ , and the other—call it G—sending the two nonidentity isomorphisms to -1 in  $\mathcal{N}$ . We then have that  $F \circ P = G \circ P = H$ and so P is not an epimorphism in SymMonCat.

## 10 A remark on the symmetric setting

We want to show that neither the pointed torsion theory recalled in the introduction, nor the homotopy torsion theory introduced in the subsequent sections, can be generalised to the non-commutative or non-symmetric settings. More precisely, we show that we cannot have a pointed torsion theory on the category Mon of not necessarily commutative monoids with groups as torsion objects; similarly, not necessarily symmetric 2-groups cannot be the torsion part of a homotopy torsion theory on the 2-category MonCat of not necessarily symmetric monoidal categories (with the usual nullhomotopy structure of Definition 5.1).

First of all, we know that a necessary condition for the category Grp of groups to be the torsion part of some pointed torsion theory on Mon, is that it is a normal monocoreflective subcategory of Mon (see for example [13]). As a subcategory of Mon, Grp is indeed monocoreflective, with the coreflector sending any monoid M to the group  $\gamma(M)$  of its invertible elements, and counit of the coreflection along M being simply the inclusion  $j_M: \gamma(M) \hookrightarrow$ M. However this map is not in general a normal monomorphism. To see that, consider a cokernel  $q_M \colon M \to \varpi(M)$  of  $j_M$  in Mon. If  $j_M$  was a normal monomorphism, it would be the kernel of its cokernel, which is not the case in general. In fact, consider the monoid  $M = \operatorname{End}_{\mathsf{Set}}(X)$  of the endomorphisms in Set of some *infinite* set X, with product given by composition (therefore  $\gamma(M)$  is the set of bijective endomorphisms of X). Write X as  $X = A \sqcup B$ , with A and B disjoint subsets of X having the same cardinality as X, and choose bijections  $\alpha: A \to X$  and  $\beta: B \to X$ . Define then two elements f and q of M as follows. The function  $f: X \to X$  is simply defined as  $f(x) = \alpha^{-1}(x)$  for all  $x \in X$ . To define  $g: X \to X$ , fix an element  $a_0$  in A, and set

$$g(x) = \begin{cases} a_0 & x \in A, \\ \beta(x) & x \in B. \end{cases}$$

Clearly  $g \circ f \in M$  is the constant function at  $a_0$  and it is not invertible (so  $g \circ f \notin \gamma(M)$ ). However, if we define the invertible function  $\gamma(M) \ni u \colon X \to X$  by

$$u(x) = \begin{cases} \beta^{-1}(\alpha(x)) & x \in A, \\ \alpha^{-1}(\beta(x)) & x \in B \end{cases}$$

(so that u "swaps" A and B), we find that  $g \circ u \circ f = id_X$ . We therefore have

$$1_{\varpi(M)} = q_M(\mathrm{id}_X) = q_M(g \circ u \circ f) = q_M(g) \circ q_M(u) \circ q_M(f) = q_M(g \circ f)$$

(where we used that  $q_M(u) = \mathrm{id}_X$  since  $u \in \gamma M$ ). If  $j_M$  was the kernel of  $q_M$ , then we would have that  $g \circ f$  would be in  $\Gamma M$ , which is not true. We conclude that Grp is not a torsion subcategory of Mon.

By this last fact, we can also conclude that the category of (not necessarily symmetric) 2-groups is not the torsion part of any  $\Theta$ -torsion theory on MonCat (with the nullhomotopy structure  $\Theta$  on MonCat of Definition 5.1). In fact, if it were the case, for any monoidal category  $\mathcal{M}$  we would have a  $\Theta$ -exact sequence in MonCat of the form

(10.1) 
$$\Gamma(\mathcal{M}) \longrightarrow \mathcal{M} \xrightarrow{P} \mathcal{P}_{P}$$

for some monoidal category  $\mathcal{P}$ , and some monoidal functor P. The fact that on the left of this sequence we find the inclusion of the Picard 2-group of  $\mathcal{M}$ in  $\mathcal{M}$  is due to this inclusion being the counit of the coreflection of 2-Grp in MonCat (see Section 2.2), and the torsion part in a homotopy torsion theory is always given by the coreflection of the torsion objects (cf. [17]). Now take  $\mathcal{M} = \iota_0(M)$ , with  $M = \operatorname{End}_{\mathsf{Set}}(X)$  the monoid defined above, and  $\iota_0$ defined in Section 8. Applying K to 10.1 and using the universal property of  $\Theta$ -kernels, one can easily show that  $j_M : \gamma(M) \to M$  would need to be the kernel in Mon of  $KP : M \to K\mathcal{P}$  (K as defined in Section 8), which contradicts the fact that  $j_M$  is not a normal monomorphism in Mon.

## 11 Another 2-dimensional torsion theory

In this work we have produced a homotopy torsion theory in the 2-category of symmetric monoidal categories starting from a classical torsion theory in the 1-category of commutative monoids. In this last section we want to carry out a similar process, but this time we start from the classical torsion theory in the category of abelian groups given by torsion and torsion-free groups.

Naturally we want to work in the 2-category Sym2-Grp of symmetric 2groups as a 2-dimensional generalisation of abelian groups, and we keep the same nullhomotopy structure  $\Theta$  of Definition 5.1, though restricted to symmetric monoidal functors between symmetric 2-groups; we keep calling this restriction  $\Theta$ .

Let us start by introducing some preliminary definitions.

**Definition 11.1.** Given an object X in a symmetric 2-group  $\mathcal{G}$ , we say that X is a torsion object if there exists a positive integer n such that  $X^{\otimes n} \cong I$  (where the tensor power  $X^{\otimes n}$  is defined recursively by  $X^{\otimes 1} = X$  and  $X^{\otimes (m+1)} = X \otimes X^{\otimes m}$ ).

We say that a symmetric 2-group  $\mathcal{G}$  is torsion if all of its objects are torsion; we say that a symmetric 2-group  $\mathcal{G}$  is torsion-free if for any torsion object X in  $\mathcal{G}$  there exists a unique isomorphism from X to I in  $\mathcal{G}$ .

We claim in the following proposition that torsion and torsion-free symmetric 2-groups form a  $\Theta$ -torsion theory of the category of symmetric 2-groups.

**Proposition 11.2.** Torsion and torsion-free symmetric 2-groups (as defined in Definition 11.1) form a homotopy torsion theory on the category Sym2-Grp

of symmetric 2-groups with respect to the nullhomotopy structure  $\Theta$  defined at the start of the section.

*Proof.* We outline here the key elements of the proof

- (1) Since the image by a symmetric monoidal functor of a torsion object is plainly again a torsion object, it is obvious by the definitions of torsion and torsion-free symmetric 2-group that  $\Theta(F)$  is a singleton for any symmetric monoidal functor F from a torsion symmetric 2-group to a torsion-free symmetric 2-group.
- (2) Actually, since an object which is isomorphic to the unit object is automatically a torsion object, then in a torsion-free symmetric 2-group there is at most one map from any object to the unit object. We conclude that  $\Theta(F)$  is either the empty set or a singleton for any functor F to a torsion-free symmetric 2-group.
- (3) For any symmetric 2-group  $\mathcal{G}$ , we can consider the full subcategory  $T(\mathcal{G})$  of its torsion objects. Clearly  $T(\mathcal{G})$  is a torsion symmetric 2-group, and will give us the torsion part of  $\mathcal{G}$ .
- (4) Consider the inclusion T<sub>g</sub>: T(G) → G. Since we work in the category of symmetric 2-groups, we can directly apply the construction of cokernels from [19] to get the Θ-cokernel F<sub>g</sub>: G → Φ(G) of T<sub>g</sub> in Sym2-Grp. This construction is essentially the same we used in Proposition 3.5 and is fully described in [19]. We briefly recall here the main features we are going to need.
  - The category  $\Phi(\mathcal{G})$  has the same objects as  $\mathcal{G}$ .
  - A morphism from X to Y in Φ(𝔅) is an equivalence class [A, f], where A is an object in T(𝔅) and f: X → Y ⊗ A is a morphism in 𝔅; we have [A, f] = [B, g]: X → Y when there exists a morphism ψ: A → B in 𝔅 such that g = (Y ⊗ ψ) ∘ f.
  - The tensor product of objects in  $\Phi(\mathcal{G})$  is the same as in  $\mathcal{G}$ .
- (5) We have thus obtained a sequence

(11.3) 
$$T(\mathfrak{G}) \xrightarrow{T_{\mathfrak{G}}} \mathfrak{G} \xrightarrow{F_{\mathfrak{G}}} \Phi(\mathfrak{G}),$$

where  $F_{\mathfrak{g}}$  is a  $\Theta$ -cokernel of  $T_{\mathfrak{g}}$ . To complete the proof we just need to show that  $\Phi(\mathfrak{g})$  is torsion-free and that  $T_{\mathfrak{g}}$  is the  $\Theta$ -kernel of  $F_{\mathfrak{g}}$ . We rely on the following property (Item (6)).

- (6) If an object X is torsion in Φ(G) then it is torsion in G. In fact, if X is torsion in Φ(G), then we have a positive integer n and a morphism [A, f]: X<sup>⊗n</sup> → I in Φ(G), for some map f in G and some object A in T(G). By definition of the arrows in Φ(G), this tells us that X<sup>⊗n</sup> ≅ I ⊗ A ≅ A in G. But since A is torsion in G, there exists a positive integer m such that A<sup>⊗m</sup> ≅ I in G. Therefore we have that X<sup>⊗(nm)</sup> ≅ (X<sup>⊗n</sup>)<sup>⊗m</sup> ≅ A<sup>⊗m</sup> ≅ I, and so X is torsion in G.
- (7) To check that  $\Phi(\mathfrak{G})$  is torsion-free, suppose that X is a torsion object in  $\Phi(\mathfrak{G})$ . By Item (6), X is torsion in  $\mathfrak{G}$ , and so we can consider the isomorphism  $[X, \ell_X^{-1}]: X \to I$  in  $\Phi(\mathfrak{G})$ . Given two isomorphisms  $[A, f], [B, g]: X \to I$  in  $\Phi(\mathfrak{G})$ , we have  $g = (I \otimes \psi) \circ f$ , with  $\psi = \ell_B \circ g \circ f^{-1} \circ \ell_A^{-1}$ , and therefore [A, f] = [B, g].
- (8) To check that  $T_{\mathfrak{G}}$  is the  $\Theta$ -kernel of  $F_{\mathfrak{G}}$ , consider a 2-group  $\mathfrak{H}$  and symmetric monoidal functor  $F: \mathfrak{H} \to \mathfrak{G}$  such that  $F_{\mathfrak{G}} \circ F$  is  $\Theta$ -trivial. We just need to check that for all X in  $\mathfrak{H}$  we have that FX is torsion in  $\mathfrak{G}$ , i.e. that F actually takes values in the full subcategory  $T(\mathfrak{G})$  of  $\mathfrak{G}$  (in fact, the condition on nullhomotopies is automatically satisfied thanks to Item (2), and uniqueness is ensured by the fact that  $T_{\mathfrak{G}}$  is an inclusion). If  $F_{\mathfrak{G}} \circ F$  is  $\Theta$ -trivial, then for all X in  $\mathfrak{H}$ , we have that  $FX \cong I$  in  $\Phi(\mathfrak{G})$ . But then by Item (6) we conclude that FX is torsion in  $\mathfrak{G}$  and we are done.  $\Box$

The torsion theory just introduced shares the 2-dimensional properties listed in Section 7. It is also well-behaved with respect to the functors  $\pi_0$  and  $\pi_1$  (see Section 8), meaning that by applying these functors to the  $\Theta$ -exact sequence 11.3 we obtain exact sequences of abelian groups corresponding to two different torsion theories. In particular, applying  $\pi_0$  we get the classical presentation of the abelian group  $\pi_0(\mathfrak{G})$  as an extension of its (classical) torsion-free part by its torsion subgroup; applying  $\pi_1$  we get instead the trivial exact sequence  $\pi_1(\mathfrak{G}) = \pi_1(\mathfrak{G}) \longrightarrow 0$ .

# References

- J. C. Baez, Hoàng Xuân Sính's thesis: Categorifying Group Theory, Thang Long Journal of Science: Mathematics and Mathematical Sciences 2 (8) (2023).
- [2] J. C. Baez, A. D. Lauda, Higher-Dimensional Algebra V: 2-Groups, Theory and Applications of Categories 12 (14) (2004).
- [3] F. Borceux, Handbook of Categorical Algebra, Vol. 2, Cambridge University Press, 1994.
- [4] F. Borceux, F. Campanini, M. Gran, W. Tholen, Groupoids and skeletal categories form a pretorsion theory in Cat, Advances in Mathematics 426 (2023).
- [5] D. Bourn, M. Gran, Torsion theories in homological categories, Journal of Algebra 305 (1) (2006) 18–47.
- [6] M. M. Clementino, D. Dikranjan, W. Tholen, Torsion theories and radicals in normal categories, Journal of Algebra 305 (1) (2006) 98–129.
- [7] S. D. Dickson, A torsion theory for Abelian categories, Trans. Amer. Math. Soc. (1966).
- [8] A. Facchini, Commutative Monoids, Noncommutative Rings and Modules, Springer International Publishing, 2021.
- [9] A. Facchini, C. Finocchiaro, M. Gran, Pretorsion theories in general categories, Journal of Pure and Applied Algebra 225 (2) (2021).
- [10] M. Grandis, Simplicial Homotopical Algebra and Satellites, Applied Categorical Structures 5 (1997).
- [11] M. Grandis, G. Janelidze, From torsion theories to closure operators and factorization systems, Categories and General Algebraic Structures with Application 12 (2020).
- [12] Hoàng Xuân Sính, Gr-catégories, Ph.D. thesis, Université Paris VII (1973).

- [13] G. Janelidze, W. Tholen, Characterization of torsion theories in general categories, Contemp. Math. 431 (2007).
- [14] A. Joyal, R. Street, Braided Tensor Categories, Advances in Mathematics 102 (1) (1993).
- [15] G. M. Kelly, On MacLane's conditions for coherence of natural associativities, commutativities, etc., Journal of Algebra 1 (4) (1964).
- [16] S. Mac Lane, Categories for the Working Mathematician, 2nd Edition, Springer New York, 1978.
- [17] S. Mantovani, M. Messora, E. M. Vitale, Homotopy torsion theories, Journal of Pure and Applied Algebra 228 (12) (2024).
- [18] J. Rosický, W. Tholen, Factorization, Fibration and Torsion, Journal of Homotopy and Related Structures 2 (2008).
- [19] E. M. Vitale, A Picard–Brauer exact sequence of categorical groups, Journal of Pure and Applied Algebra 175 (1) (2002).