Duality of Selmer groups of an abelian variety over a number field

SAIKAT BISWAS

ABSTRACT. Let A be an abelian variety defined over a number field K and let A^{\vee} be the dual abelian variety. For an odd prime p, we consider two Selmer groups attached to A[p] and relate the orders of these groups along with those of their corresponding duals to the order of the component groups of A^{\vee} at primes v.

1. INTRODUCTION

In order to motivate the main result of this paper, we begin with a quick review of *generalized Selmer groups*, fixing notations along the way. Let K be a number field with Galois group $G_K := \operatorname{Gal}(\overline{K}/K)$, for a separable closure \overline{K} of K. For every prime v of K, let K_v be the completion of K at v with Galois group $G_v := G_{K_v}$. When v is finite, we denote by $K_v^{\rm nr}$ the maximal unramified extension of K_v whose Galois group is identified with the *inertia group* $I_v \subset G_v$ at v. The Galois group of the finite residue field k_v of K_v is given by the quotient group G_v/I_v . Let M be a discrete G_K -module of finite cardinality, μ_{∞} be the G_K -module of roots of unity in \overline{K} , and $M^* := \operatorname{Hom}_{\mathbb{Z}}(M, \mu_{\infty})$ the *Cartier dual* of *M*. We shall also denote by $\widehat{M} := \operatorname{Hom}_{\mathbb{Z}}(M, \mathbb{Q}/\mathbb{Z})$ the *Pontryagin dual* of M. There is a continuous discrete action of G_K on M^* arising out of the corresponding action of G_K on M as well as on \overline{K}^{\times} . We consider a Selmer structure \mathcal{L} for M, i.e. a family of subgroups $H^1_{\mathcal{L}}(K_v, M) \subset H^1(K_v, M)$ for every prime v such that $H^1_{\mathcal{L}}(K_v, M) = H^{\tilde{1}}(G_v/I_v, M^{I_v})$ for all but finitely many v. The corresponding Selmer structure \mathcal{L}^* for M^* is the family of subgroups $H^1_{\mathcal{L}^*}(K_v, M^*) \subset H^1(K_v, M^*)$, where $H^1_{\mathcal{L}^*}(K_v, M^*)$ is the orthogonal complement of $H^1_{\mathcal{L}}(K_v, M)$ under the local Tate pairing

 $H^1(K_v, M) \times H^1(K_v, M^*) \to \mathbb{Q}/\mathbb{Z}$

²⁰¹⁰ Mathematics Subject Classification. Primary 11G10; Secondary 11S25, 12G05.

Key words and phrases. Selmer groups, Tamagawa numbers, abelian varieties.

The generalized Selmer group $H^1_{\mathcal{L}}(K, M)$ of M is the subgroup of all classes in $H^1(K, M)$ whose image under the restriction map

$$\operatorname{Res}_v : H^1(K, M) \to H^1(K_v, M)$$

lies in $H^1_{\mathcal{L}}(K_v, M)$ for every v. The Selmer group $H^1_{\mathcal{L}^*}(K, M^*)$ of M^* is defined analogously, and is called the *dual Selmer group* attached to $H^1_{\mathcal{L}}(K, M)$. We have the following important result due to Wiles [Wil95], suggested by a result of Greenberg [Gre89], which allows for the comparison of the order of a Selmer group to that of its dual.

Theorem 1.1. The groups $H^1_{\mathcal{L}}(K, M)$ and $H^1_{\mathcal{L}^*}(K, M^*)$ are finite, and we have

$$\frac{\#H^1_{\mathcal{L}}(K,M)}{\#H^1_{\mathcal{L}^*}(K,M^*)} = \frac{\#H^0(K,M)}{\#H^0(K,M^*)} \prod_v \frac{\#H^1_{\mathcal{L}}(K_v,M)}{\#H^0(K_v,M)}$$

Since $\#H^1(G_v/I_v, M^{I_v}) = \#H^0(K_v, M)$ [DDT95, §2.3], all but finitely many terms in the product on the right hand side are 1 so that the product makes sense. The formulation of Theorem 1.1 as given above appears in [DDT95] which is slightly different from that presented in [Wil95], but the two are equivalent. Following a terminology suggested by Darmon in [Dar04], we refer to the term on the right side of Theorem 1.1 as the *Euler characteristic* attached to $H^1_{\mathcal{L}}(K, M)$ and denote it by $\chi_{\mathcal{L}}(K, M)$. Thus, we can write

(1.1)
$$\chi_{\mathcal{L}}(K,M) = \frac{\#H^0(K,M)}{\#H^0(K,M^*)} \prod_v \frac{\#H^1_{\mathcal{L}}(K_v,M)}{\#H^0(K_v,M)}$$

If $\mathcal{L}_1, \mathcal{L}_2$ are Selmer structures for M, we say that $\mathcal{L}_1 \subseteq \mathcal{L}_2$ if $H^1_{\mathcal{L}_1}(K_v, M) \subseteq H^1_{\mathcal{L}_2}(K_v, M)$ for every v. In this case, it can be shown that $H^1_{\mathcal{L}_1}(K, M) \subseteq H^1_{\mathcal{L}_2}(K, M)$ and that $\mathcal{L}_1^* \supseteq \mathcal{L}_2^*$. The next corollary follows immediately from (1.1) above.

Corollary 1.2. Suppose that $\mathcal{L}_1 \subseteq \mathcal{L}_2$ are Selmer structures for M. Then we have

(1.2)
$$\frac{\chi_{\mathcal{L}_2}(K,M)}{\chi_{\mathcal{L}_1}(K,M)} = \prod_v \frac{\# H^1_{\mathcal{L}_2}(K_v,M)}{\# H^1_{\mathcal{L}_1}(K_v,M)}$$

In particular, the comparison of the Euler characteristics attached to two different Selmer groups is a purely local calculation. We can also rewrite (1.2), in view of Theorem 1.1, as

(1.3)
$$\frac{\#H^1_{\mathcal{L}_2}(K,M)}{\#H^1_{\mathcal{L}_2^*}(K,M^*)} = \chi_{\mathcal{L}_1}(K,M) \cdot \prod_v \frac{\#H^1_{\mathcal{L}_2}(K_v,M)}{\#H^1_{\mathcal{L}_1}(K_v,M)}$$

 $\mathbf{2}$

The goal of this paper is to carry out the computation involved in Corollary 1.2 for the case when, for an odd prime p, M = A[p] is the p-torsion subgroup of an abelian variety A defined over K. Let A^{\vee} be the dual abelian variety. Then it can be shown that $M^* = A^{\vee}[p]$. Let S be any finite set of primes of K containing all archimedean primes, all primes above p, and all primes of bad reduction for A. We consider three Selmer structures for A[p]. The first, denoted by $\operatorname{Sel}_p(A/K_v)$, is defined by $\operatorname{im}(\delta_v)$, where δ_v is the injection

$$A(K_v)/pA(K_v) \hookrightarrow H^1(K_v, A[p])$$

arising out of the Kummer exact sequence. The second, denoted by $H^1_{(S)}(K_v, A[p])$, is defined by ker(φ), where φ is the composition

$$H^1(K_v, A[p]) \to H^1(K_v, A)[p] \to H^1(K_v^{\mathrm{nr}}, A)[p]$$

The third, denoted by $H^1_{[S]}(K_v, A[p])$, is defined by $\delta_v(\kappa_v)$, where κ_v is the kernel of the map

$$A(K_v)/pA(K_v) \to \Phi_{A,v}(k_v)/p\Phi_{A,v}(k_v)$$

Here, $\Phi_{A,v}(k_v)$ is the arithmetic component group of A at v.

The respective Selmer groups, all subgroups of $H^1(K, A[p])$, are denoted by $\operatorname{Sel}_p(A/K)$, $H^1_{(S)}(K, A[p])$, and $H^1_{[S]}(K, A[p])$. The corresponding Selmer groups for $A^{\vee}[p]$ are defined similarly. Our main result in this paper is the following theorem.

Theorem 1.3 (Main Theorem). Under the notations introduced above, we have (K = A[-1])

$$\frac{\chi_{(S)}(K, A[p])}{\chi_{\operatorname{Sel}_p}(K, A[p])} = \# \bigoplus_{v \in S} \Phi_{A^{\vee}, v}(k_v)[p]$$

or, equivalently

$$\frac{\#H^1_{(S)}(K, A[p])}{\#H^1_{[S]}(K, A^{\vee}[p])} = \frac{\#\operatorname{Sel}_p(A/K)}{\#\operatorname{Sel}_p(A^{\vee}/K)} \cdot \#\bigoplus_{v \in S} \Phi_{A^{\vee}, v}(k_v)[p]$$

The local terms on the right hand side of Theorem 1.3 can be computed explicitly by Tate's algorithm [Sil94, IV.§9] when A is an elliptic curve, by the monodromy pairing [Gro73, §10] when A is a semistable abelian variety, and by the intersection pairing [BL99, §1] when A is the Jacobian of a curve.

Corollary 1.4. Suppose that A is an elliptic curve defined over K and p is a prime of good reduction for A. Then

$$\frac{\#H^1_{(S)}(K, A[p])}{\#H^1_{[S]}(K, A[p])} = \# \bigoplus_{v \in S} \Phi_{A,v}(k_v)[p]$$

and

$$\# \operatorname{Sel}_{p}(A/K) < \# H^{1}_{[S]}(K, A[p]) \# \bigoplus_{v \in S} \Phi_{A,v}(k_{v})[p]$$

Proof. In this case, $A^{\vee} \cong A$ and $\chi_{\operatorname{sel}_p}(K, A[p]) = 1$ [Dar04, Prop. 10.15]. The first equality then follows immediately from Theorem 1.3. To prove the second inequality, we show in Lemma 3.1 below that there are inclusions

$$H^1_{[S]}(K, A[p]) \subset \operatorname{Sel}_p(A/K) \subset H^1_{(S)}(K, A[p])$$

which in turn imply that

$$#H^{1}_{[S]}(K, A[p]) < # \operatorname{Sel}_{p}(A/K) < #H^{1}_{(S)}(K, A[p])$$

The desired inequality now follows by dividing throughout by $\#H^1_{[S]}(K, A[p])$ and using the first equality. \Box

In Section 2, we investigate the different Selmer structures on A[p] and the corresponding dual Selmer structures on $A^{\vee}[p]$. We relate the order of these structures to each other using Tate local duality as our key ingredient. In Section 3, we consider the corresponding Selmer groups on A[p] as well as on $A^{\vee}[p]$ and, using Corollary 1.2, prove our main result. In Section 3, we present an alternative proof of our main theorem by deriving it from a refinement of a well known dual exact sequence due to Cassels.

2. Local Computations

We collect some definitions and results from [Bis15]. Let $\Phi_{A,v}$ be the k_v -group scheme of connected components of the Neron model of A, and let $\Phi_{A,v}(k_v)$ the arithmetic component group of A at v. Let

$$c_{A,v} = \#\Phi_{A,v}(k_v)$$

be the Tamagawa number of A at v. We write

$$c(A/K) = \prod_{v} c_{A,v}$$

Equivalently, we can also define $\Phi_{A,v}(k_v)$ by the exactness of the sequence

(2.1)
$$1 \to A_0(K_v) \to A(K_v) \to \Phi_{A,v}(k_v) \to 1$$

Here, $A_0(K_v)$ is the subgroup of *universal norms* of $A(K_v)$ from K_v^{nr} defined by

$$A_0(K_v) := \operatorname{Nm}(A(K_v^{\operatorname{nr}})) = \bigcap_{L_i} N_{L_i/K_v}(A(L_i))$$

4

where, the intersection is over all finite, unramified extensions L_i of K_v and N_{L_i/K_v} is the norm map from L_i to K_v . The group $TT(A/K_v)$ of *Tamagawa torsors* of A/K_v is defined by the exactness of the sequence

(2.2)
$$1 \to \operatorname{TT}(A/K_v) \to H^1(K_v, A) \xrightarrow{\operatorname{res}_v} H^1(K_v^{\operatorname{nr}}, A)$$

In particular, $TT(A/K_v)$ is trivial when A has good reduction at v. There is a perfect duality of finite groups

(2.3)
$$\operatorname{TT}(A/K_v) \times \Phi_{A^{\vee},v}(k_v) \to \mathbb{Q}/\mathbb{Z}$$

Let p be an odd prime. For every prime v, the Kummer exact sequence for A/K_v is

$$(2.4) \quad 1 \to A(K_v)/pA(K_v) \xrightarrow{\delta_v} H^1(K_v, A[p]) \xrightarrow{\pi_v} H^1(K_v, A)[p] \to 1$$

Let

$$\operatorname{Sel}_p(A/K_v) := \operatorname{im}(\delta_v) \subset H^1(K_v, A[p])$$

Since δ_v is injective, it follows immediately that $\# \operatorname{Sel}_p(A/K_v) = \#A(K_v)/pA(K_v)$. Furthermore, for $v \notin S$, A has good reduction at v and $\operatorname{Sel}_p(A/K_v)$ is the subgroup of all classes in $H^1(K_v, A[p])$ that are unramified at v, i.e. their restriction to $H^1(K_v^{\operatorname{nr}}, A[p])$ is trivial [HS00, Prop.C.4.6]. In other words, $\operatorname{Sel}_p(A/K_v) = H^1(K_v^{\operatorname{nr}}/K_v, A(K_v^{\operatorname{nr}})[p])$ for all $v \notin S$. We have thus proved

Lemma 2.1. The family of subgroups $\operatorname{Sel}_p(A/K_v) \subset H^1(K_v, A[p])$ for all v form a Selmer structure for A. Furthermore, we have

$$\#\operatorname{Sel}_p(A/K_v) = \#A(K_v)/pA(K_v)$$

The Kummer exact sequence for A^{\vee}/K_v is (2.5)

$$1 \to A^{\vee}(K_v)/pA^{\vee}(K_v) \xrightarrow{\delta_v^{\vee}} H^1(K_v, A^{\vee}[p]) \xrightarrow{\pi_v^{\vee}} H^1(K_v, A^{\vee})[p] \to 1$$

We note that $A^{\vee}[p]$ is the Cartier dual of A[p] [Mum70, §15]. This also follows from the non-degenerate Weil pairing

$$A[p] \times A^{\vee}[p] \to \mu_p$$

There is an isomorphism [Mil02, Cor I.3.4] of compact groups

$$\widehat{A(K_v)} \cong H^1(K_v, A^{\vee})$$

and an isomorphism [Mil02, Cor I.2.3] of finite groups

$$H^1(\widetilde{K_v, A}[p]) \cong H^1(K_v, A^{\vee}[p])$$

that allow us to identify the exact sequence (2.5) as the Pontryagin dual of the exact sequence (2.4). In particular,

$$\operatorname{Sel}_p(A^{\vee}/K_v) = \operatorname{im}(\delta_v^{\vee}) \subset H^1(K_v, A^{\vee}[p])$$

is the orthogonal complement of

$$\operatorname{Sel}_p(A/K_v) = \operatorname{im}(\delta_v) \subset H^1(K_v, A[p])$$

under the local Tate pairing

(2.6)
$$H^1(K_v, A[p]) \times H^1(K_v, A^{\vee}[p]) \to \mathbb{Q}/\mathbb{Z}$$

It follows from Lemma 2.1 that

Lemma 2.2. The family of subgroups $\operatorname{Sel}_p(A^{\vee}/K_v) \subset H^1(K_v, A^{\vee}[p])$ for all v form a Selmer structure for A^{\vee} .

Let

$$H^1_{(S)}(K_v, A[p]) := \ker \varphi_v$$

where φ_v is the composition

Theorem 2.3. There is an exact sequence

(2.7)
$$1 \to A(K_v)/pA(K_v) \xrightarrow{\delta_v} H^1_{(S)}(K_v, A[p]) \to \mathrm{TT}(A/K_v)[p] \to 1$$

Proof. The pair of maps π_v , res^{*p*}_{*v*} yields the exact sequence

 $1 \to \ker(\pi_v) \to \ker(\varphi_v) \to \ker(\operatorname{res}_v^p) \to \operatorname{coker}(\pi_v)$

From (2.4), we find that $\ker(\pi_v) = \operatorname{im}(\delta_v) \cong A(K_v)/pA(K_v)$, and $\operatorname{coker}(\pi_v) = 1$ since π_v is surjective. From (2.2), we get that $\ker(\operatorname{res}_v^p) \cong \operatorname{TT}(A/K_v)[p]$. The desired exact sequence now follows.

Corollary 2.4. The family of subgroups $H^1_{(S)}(K_v, A[p]) \subset H^1(K_v, A[p])$ for all v form a Selmer structure for A. Furthermore, we have

(2.8)
$$\frac{\#H^1_{(S)}(K_v, A[p])}{\#\operatorname{Sel}_p(A/K_v)} = \#\Phi_{A^{\vee}, v}(k_v)[p]$$

Proof. As noted above, A has good reduction at all primes $v \notin S$ so that $TT(A/K_v)$ is trivial at these primes. Thus, we have

$$H^1_{(S)}(K_v, A[p]) = \ker(\varphi_v) \cong \operatorname{im}(\delta_v) = \operatorname{Sel}_p(A/K_v)$$

which shows that the subgroups $H^1_{(S)}(K_v, A[p])$ form a Selmer structure for A. Eq.(2.8) follows directly from Eq.(2.7) and Eq.(2.3), noting that $\Phi_{A^{\vee},v}(k_v)$ is finite. Remark 2.5. We note that

$$\#\Phi_{A^{\vee},v}(k_v) = c_{A^{\vee},v} = c_{A,v} = \#\Phi_{A,v}(k_v)$$

by [Lor11, Prop 4.3]. It follows that

$$c(A/K) = \prod_{v} c_{A,v} = \prod_{v} c_{A^{\vee},v} = c(A^{\vee}/K)$$

Let κ_v be defined by the exactness of the sequence

(2.9)
$$1 \to \kappa_v \to A(K_v)/pA(K_v) \xrightarrow{\alpha_v} \Phi_{A,v}(k_v)/p\Phi_{A,v}(k_v) \to 1$$

Let

$$H^1_{[S]}(K_v, A[p]) := \delta_v(\kappa_v) \subset H^1(K_v, A[p])$$

The following lemma is evident.

Lemma 2.6. We have

(2.10)
$$\#H^1_{[S]}(K_v, A[p]) = \#\kappa_v = \#[A(K_v) \cap p\Phi_{A,v}(k_v)]/pA(K_v)$$

Theorem 2.7. $H^1_{[S]}(K_v, A^{\vee}[p])$ and $H^1_{(S)}(K_v, A[p])$ are orthogonal complements of each other under the local Tate pairing (2.6).

Proof. Consider the commutative diagram

The top row is the exact sequence (2.5), while the bottom row is the dual of the exact sequence (2.7). The right vertical map is an identity, while the left vertical map α_v^{\vee} follows from (2.9), whence it is surjective. It follows that the middle vertical map β_v , induced by the two boundary maps, is surjective and that

$$\ker(\beta_v) \cong \delta_v^{\vee}(\ker(\alpha_v^{\vee})) = H^1_{[S]}(K_v, A^{\vee}[p])$$

Corollary 2.8. The family of subgroups $H^1_{[S]}(K_v, A^{\vee}[p]) \subset H^1(K_v, A^{\vee}[p])$ for all v form a Selmer structure for A^{\vee} . Furthermore,

$$\frac{\#\operatorname{Sel}_p(A^{\vee}/K_v)}{\#H^1_{[S]}(K_v, A^{\vee}[p])} = \#\Phi_{A^{\vee},v}(k_v)[p]$$

Lastly, we take note of the following lemma whose proof follows directly from definition.

 $\overline{7}$

Lemma 2.9. The aforementioned Selmer structures have inclusions

$$H^1_{[S]}(K_v, A[p]) \subset \operatorname{Sel}_p(A/K_v) \subset H^1_{(S)}(K_v, A[p])$$

for all $v \in S$.

3. GLOBAL COMPUTATIONS

Let K_S be the maximal extension of K unramified outside S and let $G_S = \text{Gal}(K_S/K)$. For each $v \in S$, let w be some prime of K_S above v, and let $K_{S,w}$ be the union of the completions at w of the finite extensions of K contained in K_S . The composition

$$G_{K_v} \twoheadrightarrow \operatorname{Gal}(K_{S,w}/K_v) \hookrightarrow G_S$$

induces the restriction map

(3.1)
$$\operatorname{Res}_{v} : H^{1}(G_{S}, A[p]) \longrightarrow H^{1}(K_{v}, A[p])$$

which in turn give the homomorphism

(3.2)
$$\beta_S : H^1(G_S, A[p]) \longrightarrow \bigoplus_{v \in S} H^1(K_v, A[p])$$

For each of the Selmer structures introduced in the previous section, we define the corresponding Selmer groups by

(3.3)
$$\operatorname{Sel}_p(A/K) := \beta_S^{-1} \left(\bigoplus_{v \in S} \operatorname{Sel}_p(A/K_v) \right) \subset H^1(G_S, A[p])$$

(3.4)
$$H^{1}_{(S)}(K, A[p]) := \beta_{S}^{-1} \left(\bigoplus_{v \in S} H^{1}_{(S)}(K_{v}, A[p]) \right) \subset H^{1}(G_{S}, A[p])$$

(3.5)
$$H^1_{[S]}(K, A[p]) := \beta_S^{-1} \left(\bigoplus_{v \in S} H^1_{[S]}(K_v, A[p]) \right) \subset H^1(G_S, A[p])$$

Since $H^1(G_S, A[p])$ is finite [Mil02, Cor. I.4.15], it follows that each of the Selmer groups above are also finite. It also follows from Lemma 2.9 above that

Lemma 3.1. There are proper inclusions

$$H^1_{[S]}(K, A[p]) \subset \operatorname{Sel}_p(A/K) \subset H^1_{(S)}(K, A[p])$$

The following proposition will be used in the next section.

Proposition 3.2. There are exact sequences

$$1 \to \operatorname{Sel}_{p}(A/K) \to H^{1}(G_{S}, A[p]) \xrightarrow{\lambda_{S}} \bigoplus_{v \in S} H^{1}(K_{v}, A)[p]$$
$$1 \to \operatorname{Sel}_{p}(A/K) \to H^{1}_{(S)}(K, A[p]) \xrightarrow{\rho_{S}} \bigoplus_{v \in S} \operatorname{TT}(A/K_{v})[p]$$
$$1 \to H^{1}_{[S]}(K, A[p]) \to \operatorname{Sel}_{p}(A/K) \xrightarrow{\eta_{S}} \bigoplus_{v \in S} \Phi_{A,v}(k_{v})/p\Phi_{A,v}(k_{v})$$

Proof. Lemmas 2.9 and 3.1 imply the exact sequences

$$1 \to \operatorname{Sel}_{p}(A/K) \to H^{1}(G_{S}, A[p]) \to \bigoplus_{v \in S} H^{1}(K_{v}, A[p]) / \operatorname{im}(\delta_{v})$$
$$1 \to \operatorname{Sel}_{p}(A/K) \to H^{1}_{(S)}(K, A[p]) \to \bigoplus_{v \in S} H^{1}_{(S)}(K_{v}, A[p]) / \operatorname{im}(\delta_{v})$$
$$1 \to H^{1}_{[S]}(K, A[p]) \to \operatorname{Sel}_{p}(A/K) \to \bigoplus_{v \in S} \operatorname{im}(\delta_{v}) / \delta_{v}(\kappa_{v})$$

The proposition now follows from Eqs (2.4), (2.7), and (2.9).

Furthermore, we find that the dual Selmer groups attached to each of these aforementioned Selmer groups can be identified as

(3.6)
$$\operatorname{Sel}_{p^*}(A^{\vee}/K) = \operatorname{Sel}_p(A^{\vee}/K)$$

(3.7)
$$H^{1}_{(S)^{*}}(K, A^{\vee}[p]) = H^{1}_{[S]}(K, A^{\vee}[p])$$

The Euler characteristics attached to the Selmer groups $\operatorname{Sel}_p(A/K)$ and $H^1_{(S)}(K, A[p])$ can be given, using Theorem 1.1, as

(3.8)
$$\chi_{\operatorname{Sel}_p}(K, A[p]) = \frac{\#\operatorname{Sel}_p(A/K)}{\#\operatorname{Sel}_p(A^{\vee}/K)}$$

(3.9)
$$\chi_{(S)}(K, A[p]) = \frac{\#H^1_{(S)}(K, A[p])}{\#H^1_{[S]}(K, A^{\vee}[p])}$$

Using Eqs (1.2) and (2.8), we obtain

$$\frac{\chi_{(S)}(K, A[p])}{\chi_{\mathrm{Sel}_p}(K, A[p])} = \prod_v \frac{\#H^1_{(S)}(K_v, A[p])}{\#\operatorname{Sel}_p(A/K_v)} = \#\bigoplus_{v\in S} \Phi_{A^{\vee}, v}(k_v)[p]$$

which proves Theorem 1.3.

Remark 3.3. $\operatorname{Sel}_p(A/K)$ is usually referred to as the classical *p*-Selmer group of *A*. On the other hand, we may refer to $H^1_{(S)}(K, A[p])$ as the relaxed Selmer group of *A* at *S* and to $H^1_{[S]}(K, A[p])$ as the restricted Selmer group of *A* at *S*. We note that these notations and terminologies (along with the construction of *S*) have been used differently in [Dar04].

4. A DUAL EXACT SEQUENCE

We present an alternative proof of Theorem 1.3 in this section. The following result is often referred to as the Cassels-Poitou-Tate sequence. The original version of this theorem appeared in [Cas64] which involved elliptic curves.

Theorem 4.1. There is an exact sequence

The results obtained in the last section can be used to establish the following refinement to Theorem 4.1. Note that Theorem 1.3 is an immediate consequence of Theorem 4.2.

Theorem 4.2. There is an exact sequence of finite groups

Proof. Consider the following exact sequences from Proposition 3.2

$$1 \to \operatorname{Sel}_{p}(A/K) \to H^{1}_{(S)}(K, A[p]) \xrightarrow{\rho_{S}} \bigoplus_{v \in S} \operatorname{TT}(A/K_{v})[p]$$
$$1 \to H^{1}_{[S]}(K, A^{\vee}[p]) \to \operatorname{Sel}_{p}(A^{\vee}/K) \xrightarrow{\eta_{S}^{\vee}} \bigoplus_{v \in S} \Phi_{A^{\vee}, v}(k_{v})/p \Phi_{A^{\vee}, v}(k_{v})$$

It suffices to prove that the images of ρ_S and η_S^{\vee} are orthogonal complements of each other under the sum of the local pairings (2.3). Equivalently, we prove that the sequence

$$H^1_{(S)}(K, A[p]) \xrightarrow{\rho_S} \bigoplus_{v \in S} \operatorname{TT}(A/K_v)[p] \xrightarrow{\eta_S^{\vee}} \operatorname{Sel}_{p(A^{\vee}/K)}$$

is exact at the middle term, i.e. $\operatorname{im} \rho_S = \ker \widehat{\eta_S^{\vee}}$. The commutative diagram



yields the exact sequence

$$1 \to \ker \lambda_S \to \ker \delta_S \to \ker (\oplus \operatorname{res}_v^p) \xrightarrow{\phi} \operatorname{coker} \lambda_S \to \operatorname{coker} \delta_S$$

part of which we identify as

$$H^1_{(S)}(K, A[p]) \xrightarrow{\rho_S} \bigoplus_{v \in S} \operatorname{TT}(A/K_v)[p] \xrightarrow{\phi} \operatorname{coker} \lambda_S$$

Hence, we have im $\rho_S = \ker \phi$. On the other hand, Theorem 4.1 yields an inclusion

$$\operatorname{coker} \lambda_S \hookrightarrow \widehat{\operatorname{Sel}_p(A^\vee/K)}$$

The map $\widehat{\eta}_S$ is the composition

$$\bigoplus_{v \in S} \operatorname{TT}(A/K_v)[p] \xrightarrow{\phi} \operatorname{coker} \lambda_S \hookrightarrow \operatorname{Sel}_p(A^{\vee}/K)$$

which implies that ker $\phi = \ker \widehat{\eta_S}$. Hence, im $\rho_S = \ker \widehat{\eta_S}$.

Remark 4.3. More generally, if $\mathcal{L}_1 \subseteq \mathcal{L}_2$ are Selmer structures for M then there are exact sequences

$$1 \longrightarrow H^1_{\mathcal{L}_1}(K, M) \longrightarrow H^1_{\mathcal{L}_2}(K, M) \longrightarrow \bigoplus_v H^1_{\mathcal{L}_2}(K_v, M) / H^1_{\mathcal{L}_1}(K_v, M)$$

$$1 \longrightarrow H^1_{\mathcal{L}^*_2}(K, M^*) \longrightarrow H^1_{\mathcal{L}^*_1}(K, M^*) \longrightarrow \bigoplus_v H^1_{\mathcal{L}^*_1}(K_v, M^*) / H^1_{\mathcal{L}^*_2}(K_v, M^*)$$

such that the images of the two right-hand maps are orthogonal complements of each other under the local Tate pairing [Rub00, Thm. 1.7.3].

Consequently, there is a canonical exact sequence of finite groups

$$1 \longrightarrow H^{1}_{\mathcal{L}_{1}}(K, M) \longrightarrow H^{1}_{\mathcal{L}_{2}}(K, M) \longrightarrow \bigoplus_{v} H^{1}_{\mathcal{L}_{2}}(K_{v}, M) / H^{1}_{\mathcal{L}_{1}}(K_{v}, M)$$

$$1 \longleftarrow H^{1}_{\mathcal{L}_{2}^{*}}(\widehat{K, M^{*}}) \longleftarrow H^{1}_{\mathcal{L}_{1}^{*}}(\widehat{K, M^{*}})$$

Note that Eq.(1.3) follows immediately from this sequence.

References

- [Bis15] S. Biswas, Tamagawa torsors of an Abelian variety, J. Ramanujan Math. Soc. 30, No. 1 (2015), 67–81.
- [BL99] S. Bosch, Q. Liu, Rational points of group of components of a Néron Model, Manuscripta Math. 98 (1999), no. 3, 275–293.
- [Cas64] J.W.S Cassels, The dual exact sequence, J. Reine Angew. Math. 216 (1964), 150–158.
- [Dar04] H. Darmon, Rational Points on Modular Elliptic Curves, CBMS Regional Conference Series in Mathematics, 101, American Mathematical Society, Providence, RI, 2004.
- [DDT95] H. Darmon, F. Diamond, R. Taylor, *Fermat's Last Theorem*, in: Elliptic Curves, Modular Forms & Fermat's Last Theorem (Hong Kong 1993), 2–140, International Press, Cambridge, MA, 1997.
- [Gre89] R. Greenberg, Iwasawa theory for p-adic representations, in: Algebraic Number Theory, in honor of K. Iwasawa, Adv. Studies in Pure Math. 17, Academic Press, San Diego, 1989, pp. 97–137.
- [Gro73] A. Grothendieck, Modèles de Néron et monodromie, LNM 288, Seminaire de Géometrie 7, Expose IX, Springer-Verlag, 1973.
- [HS00] M. Hindry, J. Silverman, *Diophantine Geometry*, Springer, 2000.
- [Lor11] D. Lorenzini, Torsion and Tamagawa numbers, Ann. Inst. Fourier, 61 no. 5 (2011), 1995–2037.
- [Mil02] J. S. Milne, Arithmetic Duality Theorems, Book Surge, 2002.
- [Mum70] D. Mumford, Abelian Varieties, Oxford University Press: Oxford, 1970.
- [Rub00] K. Rubin, *Euler systems*, Annals of Math. Studies 147, Princeton University Press, 2000.
- [Sil94] J. Silverman, Advanced topics in the arithmetic of elliptic curves, GTM 151, Springer, 1994.
- [Wil95] A. Wiles, Modular elliptic curves and Fermat's Last Theorem, Annals of Math. 141 (1995), 443–551.

DEPARTMENT OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TEXAS AT DALLAS, RICHARDSON, TX 75080-3021.

Email address: saikat.biswas@utdallas.edu