FINEST POSITROID SUBDIVISIONS FROM MAXIMAL WEAKLY SEPARATED COLLECTIONS

Gleb A. Koshevoy, Fang Li, Lujun Zhang *

Contents

1	Intr	oduction and notations	2
2	Posi	tive Tropical Grassmannian and The First Theorem	3
	2.1	Preliminaries on positive tropical Grassmannian and positroid subdivisions	3
	2.2	Blade arrangement and weighted blade arrangement	4
	2.3	Construction of positive tropical Grassmannian by weighted blade arrangement	6
	2.4	The first theorem	12
3	Bou	ndary Maps on Reduced Plabic Graphs and The Second Theorem	13
	3.1	The boundary maps on plabic graphs	14
	3.2	Some lemmas for preparations	15
	3.3	Proof of the second theorem and a corollary	19
4	The	Conclusions and the Further Work	22

ABSTRACT

We study cell decomposition of positive tropical Grassmannian Trop⁺Gr_{k,n} following an approach by Early in [1]. Specifically, we deal with positroid subdivision of hypersimplex induced by translated blades from any maximal weakly separated collection. One of our main results gives a necessary and sufficient condition on a maximal weakly separated collection to form a positroid subdivision of a hypersimplex corresponding to a simplicial cone in Trop⁺Gr_{k,n}. For k = 2 our condition says that any weakly separated collection of two-elements sets gives such a simplicial cone, and all cones are of such a form.

Then our second result shows that the maximality of any weakly separated collection is preserved under the boundary map, which affirmatively answers a question by Early in [1]. The main tool in proving this theorem is the plabic graph proposed by Postnikov [2]. As a corollary, we find that all those positroid subdivisions are the finest. Thus, the flip of two maximal weakly separated collections corresponds to a pair of adjacent maximal cones in positive tropical Grassmannian.

Keywords Positive tropical Grassmannian · Positroid subdivision · Weak separation · Plabic graph · Cluster algebra

arXiv:2502.05033v1 [math.RT] 7 Feb 2025

^{*}The corresponding author

1 Introduction and notations

The positive tropical Grassmannian Trop⁺Gr_{k,n} was proposed by Speyer and Williams ([3]) in 2005. It is defined as the space of realizable positive tropical linear spaces. Recently, there are many researches on this object with other areas of math and physics ([4], [5], [6]). Among them, the relation with the positroid subdivisions of hypersimplex $\Delta_{k,n}$ is the main focus in this paper. From the results in [4] [7], we know that Trop⁺Gr_{k,n} parameterizes all the regular positroid subdivisions of $\Delta_{k,n}$. To be more precise, maximal cones in Trop⁺Gr_{k,n} parameterize the finest positroid subdivisions of $\Delta_{k,n}$ while rays parameterize the coarsest positroid subdivisions.

More specifically, Regular matroid sudivisions of a hypersimplex $\Delta_{k,n}$ is an implementation of $\Delta_{k,n}$ as a polyhedral complex of matroid polytopes. Such subdivisions arise as affinity areas of M-convex functions by Murota [8] with the domain $\Delta_{k,n}$. which generalize valued matroids by Dress and Wentzel ([9]) over the tropical ring.

M-convex function has a local descriptions which read as follows: the restriction of an M-convex function with the domain $\Delta_{k,n}$ on an octahedron with vertices *Lab*, *Lcd*, *Lac*, *Lbd*, *Lad*, *Lbc*, $\binom{L \in [n]}{k-2}$, a < b < c < d, $\{a, b, c, d\} \cap L = \emptyset$, has at most two affinity areas of full dimension. M-convex functions are stable under convolution but not under the summation.

We are interested in a subclass of tropical Plücker functions (TP functions for short), which are also called positive tropical vectors. Such a subclass is less complicated. Namely, the local characterisation of such functions reads as follows; there are three of four possibilities for M-convex functions. Namely, either the whole octahedron, or the halves with the vertices containing the non-separated diagonal with endpoints *Lac*, *Lbd*.

The space of TP functions on $\Delta_{k,n}$ is of dimension k(n - k) + 1. Namely the restriction of a TP-function to the vertices labelling by sets of a maximal weakly separated collection W is a bijection indeed (see Appendix A to [10]). The point is that weakly separated collections corresponds to seeds of the cluster algebra of regular functions on Grassmaniann.

Early in [1] first made a connection between weakly separated collections and positroid subdivisions of $\Delta_{k,n}$ by moving standard blades to vertices of hypersimplex. A standard blade β is the union of codimension 1 faces of a complete fan in hyperplane $\sum_{i=1}^{n} x_i = 0$. By moving β to a vertex e_J of $\Delta_{k,n}$, we get the translated blade β_J which induces a multi-split of $\Delta_{k,n}$. Recall that a k-split ($k \ge 2$) is polyhedral subdivision into k-maximal cells which intersect in an inner face of codimension k - 1. The following theorem by Herrmann guarantees that the positroid subdivision induced by β_J corresponds to a ray in Trop⁺Gr_{k,n}.

Theorem 1.1. ([11]) A k-split is a coarsest regular subdivision.

In [6] the notion of blade arrangement is extended to weighted blade arrangement which is a \mathbb{R} -linear combination of translated blades. This generalisation allows to implement the tropical Grassmannian by weighted blade arrangements that satisfy compatibility condition and positivity condition under the boundary map ∂_L (Definition 2.26). Here we use a slightly different setting in defining this collection of weighted blade arrangements $\mathcal{Z}_{k,n}$, and prove that $\mathcal{Z}_{k,n}$ is exactly Trop⁺Gr_{k,n} (See Lemma 2.27). Let $\overline{\mathcal{Z}_{k,n}}$ be the quotient fan of $\mathcal{Z}_{k,n}$ modulo the *n*-dimensional linearity space $\mathcal{F}_{k,n} := \operatorname{span}{\beta_I \mid I}$ is frozen}. When k = 2, the polyhedron cone $\mathcal{P}_W := {\sum_{J \in W} \mathbb{R}_{\geq} \beta_J}$ from a nonfrozen maximal weakly separated collection *W* is maximal simplicial in $\overline{\mathcal{Z}_{2,n}}$. However, when k > 2, this is not always true. Thus we provide a sufficient and necessary condition for \mathcal{P}_W being simplicial of maximal dimension in $\overline{\mathcal{Z}_{k,n}}$.

Theorem 1.2. (Theorem 2.32) Define

$$\mathcal{S} := \{W' \subseteq \binom{[n]}{k} \mid \partial_L(W') \text{ contains a nonfrozen maximal } w - \text{ collection in } \binom{[n] \setminus L}{2} \forall L \subseteq [n] \text{ with } |L| = k - 2\}$$

as a poset of nonfrozen w-collections ordered by inclusion. Then for any nonfrozen maximal w-collection W, \mathcal{P}_W is maximal simplicial in $\overline{\mathcal{Z}_{k,n}}$ if and only if W is minimal in S.

In this paper, we mainly focus on those subdivisions induced by a maximal weakly separated collection W. For their characterization we affirmatively answer a question in [1] on stability of maximality of weak separation under the boundary map ∂_j .

Theorem 1.3. (*Theorem 3.1*) Let W be a maximal w-collection in $\binom{[n]}{k}$, then $\partial_j(W)$ is also a maximal w-collection in $\binom{[n]\setminus\{j\}}{k-1}$.

We complete the proof by using plabic graphs. The plabic graph is a combinatorial tool proposed by Postnikov [2] to study the totally nonnegative Grassmannian. Danilov et al [12] affirmatively answered to the Leclerc-Zelevinsky conjecture on purity of maximal weakly separated collections and proved that any two maximal weakly separated

collections can be connected by a sequence of flips. Later one, Oh et al [13] proved the same results using technique of plabic tilings, which are closely related to plabic graphs.

Combining with the weighted blade arrangement construction of positive tropical Grassmannian and the above theorem, we get the following corollary.

Corollary 1.4. (Corollary 3.13) The translated blades of a maximal weakly separated collection induce a finest regular positroid subdivison of $\Delta_{k,n}$.

This paper is organized as follows. We start by introducing a new construction of positive tropical Grassmannian from Nick Early in Section 2. Based on this construction, we prove our first theorem in Section 2.4. In Section 3, we introduce a boundary map on a reduced plabic graph and offer some lemmas for preparations. Then we give the proof of the second theorem and an example in Section 3.3.

We use the following notation throughout the paper. Let [n] denote $\{1, 2, \dots, n\}$ with cyclic order and $\binom{[n]}{k}$ denote the set of k-subsets of [n]. For any subset $L \subseteq [n]$, we endow $[n] \setminus L$ with a cyclic order inherited from [n]. For example, if we take n = 8 and $L = \{3, 4, 7\}$, the cyclic order of $[n] \setminus L = \{1, 2, 5, 6, 8\}$ is given by 1 < 2 < 5 < 6 < 8 and its cyclic rotations.

We write [a, b] for the closed cyclic interval from a to b and $(a, b) := [a, b] \setminus \{a, b\}$ for the open cyclic interval. If L is a subset of [n] and $a \in [n]$, we abbreviate $L \cup \{a\}$ by La. For any $I \in {[n] \choose k}$, we use $e_I := \sum_{i \in I} e_i$ and $x_I : \sum_{i \in I} x_i$ to denote the sum of unit vectors and coordinates. If no confusion, we write the set $I = \{i_1, i_2, \dots, i_k\}$ by $i_1 i_2 \cdots i_k$. For brevity, we refer to weakly separated collections as w-collections.

2 Positive Tropical Grassmannian and The First Theorem

2.1 Preliminaries on positive tropical Grassmannian and positroid subdivisions

It has been proved in [4] that the positive Dressian equals the positive tropical Grassmannian which implies that $Trop^+Gr_{k,n}$ is the intersection of tropical hypersurfaces determined by all the three-terms Plücker relations.

Definition 2.1. The positive tropical Grassmannian Trop⁺Gr_{k,n} is the set of points $p = (p_I)_{I \in \binom{[n]}{k}} \in \mathbb{R}^{\binom{n}{k}}$ such that

$$p_{Lac} + p_{Lbd} = \min \left\{ p_{Lab} + p_{Lcd}, \ p_{Lbc} + p_{Lad} \right\}$$

for any $1 \le a < b < c < d \le n$ and $L \in {[n] \setminus \{a, b, c, d\} \choose k-2}$ and p is called a positive tropical Plücker vector.

Definition 2.2. For $\mathcal{M} \subset {\binom{[n]}{k}}$, let

$$S_{\mathcal{M}} := \{ V \in Gr_{k,n}^{\geq 0}(\mathbb{R}) \mid p_I(V) > 0 \text{ if and only if } I \in \mathcal{M} \}.$$

If $S_M \neq \emptyset$, then \mathcal{M} is called a **positroid** and $S_{\mathcal{M}}$ a **positroid cell**. $P_{\mathcal{M}} := Conv\{e_J \mid J \in \mathcal{M}\}$ is the **positroid polytope** related to \mathcal{M} . A polyheral subdivision is called a **positroid subdivision** if every face in this subdivision is a positroid polytope.

If we take $p = (p_I)_{I \in \binom{[n]}{k}}$ as height function on vertices of $\Delta_{k,n}$ and project the lower faces of convex({(e_I, p_I)|I $\in \binom{[n]}{k}$ }) $\subset \mathbb{R}^{n+1}$ onto $\Delta_{k,n}$, then it induces a polyhedral subdivision \mathcal{D}_p of $\Delta_{k,n}$. We have the following statement.

Theorem 2.3. ([5], Theorem 9.12) $p \in \text{Trop}^+\text{Gr}_{k,n}$ if and only if \mathcal{D}_p is a positroid subdivision of $\Delta_{k,n}$.

Thus Trop⁺Gr_{k,n} is equipped with a secondary fan structure such that p and p' are in the same interior of a cone if and only if they induce the same positroid subdivision of $\Delta_{k,n}$. In Trop⁺Gr_{k,n}, a ray corresponds to a coarsest positroid subdivision while a maximal cone corresponds to a finest positroid subdivision. Besides, Trop⁺Gr_{k,n} has a Plücker fan structure as a subfan of Gröbner fan of Grassmannian $Gr_{k,n}$. The following theorem says that this two structures coincide.

Theorem 2.4. ([14], Theorem 14) The Plücker fan structure coincides with the secondary fan structure in Trop⁺Gr_{k,n}.

There is a direct corollary from the above theorem.

Corollary 2.5. Two regular positroid subdivisions \mathcal{D}_p and $\mathcal{D}_{p'}$ coincide on $\Delta_{k,n}$ if and only if they induce the same subdivision on the 3-skeleton (every octahedral face) of $\Delta_{k,n}$.

Theorem 2.6. Let p be a positive tropical Plücker vector. Then the following statements are equivalent

- (i) \mathcal{D}_p induces a finest subdivision of $\Delta_{k,n}$, that is, for any positive Plücker vector p' such that p + p' is positive Plücker, there holds $\mathcal{D}_p = \mathcal{D}_{p+p'}$.
- (ii) Every octohedral face in \mathcal{D}_p is subdivided in two pyramids, each of which contains the non-separated diagonal.

Proof. (ii) \Rightarrow (i) is directly by Corollary 2.5. We only prove the converse direction. Suppose for a finest subdivision \mathcal{D}_p , an octahedron with vertices *Lab*, *Lcd*, *Lac*, *Lbd*, *Lad*, *Lbc* is not subdivided. Since with vertices *Lab*, *Lcd*, *Lad*, *Lbc* are weakly separated, we can expand them to a maximal w-collection W. Define a Plücker function by assigning the values of vertices of W such that $p'_{Lab} + p'_{Lcd} \neq p'_{Lad} + p'_{Lbc}$. Then, for such a positive tropical Plücker vector p', the subdivision $\mathcal{D}_{p+p'}$ has the octahedron with vertices *Lab*, *Lcd*, *Lac*, *Lbd*, *Lad*, *Lbc*, subdivided in two halves. A contradiction.

2.2 Blade arrangement and weighted blade arrangement

We follow the notations from [1] [6]. In [6]] there is a construction of a bijection between $\text{Trop}^+\text{Gr}_{k,n}$ and weighted blade arrangements, which we will use.

Definition 2.7. ([1]) A decorated ordered set partition $((S_1)_{s_1}, \dots, (S_l)_{s_l})$ is an ordered set partition (S_1, \dots, S_l) of $\{1, \dots, n\}$ together with a list of nonnegative integers (s_1, \dots, s_l) such that $\sum_{j=1}^l s_j = k$. Additionally, It is said to be of type $\Delta_{k,n}$ if $|S_j| \ge 2$ and $1 \le s_j \le |S_j| - 1$ (for each $j = 1, \dots, l$), briefly written as $((S_1)_{s_1}, \dots, (S_l)_{s_l}) \in OSP(\Delta_{k,n})$. Denote by $[(S_1)_{s_1}, \dots, (S_l)_{s_l}]$ the polyhedral cone in $H_{k,n}$ formulated by the following facet inequalities

$$x_{S_1} \ge s_1$$

$$x_{S_1 \cup S_2} \ge s_1 + s_2$$

$$\vdots$$

$$x_{S_1 \cup \dots \cup S_{l-1}} \ge s_1 + \dots + s_{l-1}.$$
(1)

The l cones $[(S_1)_{s_1}, (S_2)_{s_2}, \dots, (S_l)_{s_l}], [(S_2)_{s_2}, (S_3)_{s_3}, \dots, (S_l)_{s_l}] \cdots [(S_l)_{s_l}, (S_1)_{s_1}, \dots, (S_{l-1})_{s_{l-1}}]$ form a complete simplicial fan in $H_{k,n}$ ([15]). The blade $(((S_1)_{s_1}, \dots, (S_l)_{s_l}))$ is the union of codimension 1 faces of this fan (here we use $\partial(\mathcal{P})$ to denote the boundary of the closed cone \mathcal{P}), that is

$$(((S_1)_{s_1},\cdots,(S_l)_{s_l})) = \bigcup_{j=1}^l \partial([(S_j)_{s_j},(S_{j+1})_{s_{j+1}}\cdots,(S_{j-1})_{s_{j-1}}]).$$
(2)

Besides, if $s_i = 0$ for some $j = 1, \dots, l$ then the subscript s_i will be omitted.

The standrad blade $\beta = ((1, 2, \dots, n))$ is characterized as follows

$$\beta = ((1, 2, \cdots, n)) = \bigcup_{j=1}^{n} \partial([j, j+1, \cdots, j-1]).$$

Denote by $\Pi_j := [j, j+1, \dots, j-1] = \left\{ t_1(e_1 - e_2) + t_2(e_2 - e_3) + \dots + t_j(e_j - e_{j+1}) + \dots + t_n(e_n - e_1) \right\} (j+1 := 1 \text{ if } j = n) \text{ and } \beta_J = ((1, 2, \dots, n))_{e_J} \text{ is the translation of } \beta \text{ from origin to the vertex } e_J \text{ of } \Delta_{k,n}. \text{ Given a piecewise-linear function } h(x) \text{ on hyperplane } H_{0,n} := \{x \in \mathbb{R}^n \mid \sum_{i=0}^n x_i = 0\} \text{ by}$

$$h(x) = \min\{L_1(x), L_2(x), \cdots, L_n(x)\}$$

where

$$L_i(x) = x_{i+1} + 2x_{i+2} + \dots + (n-i)x_n + x_1 + \dots + (n-1)x_{i-1}$$

Proposition 2.8. ([1]) The affine areas of h(x) are exactly $\Pi_1, \Pi_2, \dots, \Pi_n$, and

$$\min\{L_1(x), L_2(x), \cdots, L_n(x)\} = L_i(x)$$

for $x \in \Pi_i$. Besides, the standard blade β is the tropical hypersurface of h(x).

Now since $\Pi_1, \Pi_2, \dots, \Pi_n$ form a complete simplicial fan in $H_{0,n}$, then for any $x \in (H_{0,n} \cap \mathbb{Z}^n)$, there exists a unique maximal subset $\{i_1, i_2, \dots, i_l\} \subset [n]$ such that $x \in \bigcap_{j \in \{i_1, i_2, \dots, i_l\}} \Pi_j$. Moreover, it is not hard to check that $\{e_j - e_{j+1} \mid j \in \{i_1, i_2, \dots, i_l\}^c\}$ is a minimal system of generators for the monoid $(\bigcap_{j \in \{i_1, i_2, \dots, i_l\}} \Pi_j) \cap \mathbb{Z}^n$. Thus we have a unique **positive expression**

$$x = \sum_{j \in \{i_1, i_2, \cdots, i_l\}^c} t_j (e_j - e_{j+1})$$

where $t_j > 0$ for $j \notin \{i_1, i_2, \dots, i_l\}^c$. The set $\{i_1, i_2, \dots, i_l\}^c$ is called the **support** of *x*, denoted by Supp(x). In particular, let $x = e_J - e_I$ for any *k*-subsets $I \neq J$, the positive integer $\sum_{j \in \{i_1, i_2, \dots, i_l\}^c} t_j$ is called the **distance** from e_I to e_J , denoted by $d(e_I, e_J)$. Clearly, $d(e_I, e_J) + d(e_J, e_I) = n$ whenever *I*, *J* satisfies $|I \setminus J| = |J \setminus I| = 1$.

Lemma 2.9. Let $h(x - e_J)$ be the translation of h(x) from the origin to e_J , then

$$h(e_I - e_J) = -d(e_J, e_I)$$

Proof. We first take the unique positive expression $e_I - e_J = \sum_{j \in Supp(e_I - e_J)} t_j(e_j - e_{j+1})$. Without loss of generality, suppose $e_I - e_J \in \Pi_1$ (In other words, $t_n = 0$), then

$$h(e_{I} - e_{J}) = L_{1}(e_{I} - e_{J})$$

= $(e_{I} - e_{J}) \cdot (0, 1, 2, \dots, n-1)$
= $(\sum_{j \in Supp(e_{I} - e_{J})} t_{j}(e_{j} - e_{j+1})) \cdot (0, 1, 2, \dots, n-1)$
= $-\sum_{j \in Supp(e_{I} - e_{J})} t_{j}$

Theorem 2.10. ([1]) Let e_J be a vertex of $\Delta_{k,n}$, the translated blade $\beta_J = ((1, 2, \dots, n))_{e_J}$ induces a *l*-split of $\Delta_{k,n}$ such that

$$((1,2,\cdots,n))_{e_J} \cap \Delta_{k,n} = ((S_1)_{s_1},(S_2)_{s_2},\cdots,(S_l)_{s_l}) \cap \Delta_{k,n}$$

Where $((S_1)_{s_1}, (S_2)_{s_2}, \dots, (S_l)_{s_l}) \in OSP(\Delta_{k,n})$ satisfies that l is the number of cyclic intervals of J and $1 \in S_1$. The blade β_J induces a trivial subdivision of $\Delta_{k,n}$ if and only if J is an interval. In this case, J is called frozen while others are called nonfrozen.

Remark 2.11. The blade $((S_1)_{s_1}, (S_2)_{s_2}, \dots, (S_l)_{s_l})$ is completely determined by e_J . Let $J = \bigcup_{i=1}^l J_i$ be the union of cyclic intervals and (C_1, \dots, C_l) be the interlaced complement to intervals of J. Then $(S_1, S_2, \dots, S_l) = (C_1, J_1, C_2, J_2, \dots, C_l, J_l)$ where $S_i = C_i \cup J_i$ and $s_i = |J_i|$. Every translated blade β_J induce such a subdivision of $\Delta_{k,n}$, so we may abuse the notation in the following sections to represent such a subdivision by β_J .

Definition 2.12. A blade arrangement is a collection of translated blades $\{\beta_{J_1}, \beta_{J_2}, \dots, \beta_{J_l}\}$. A weighted blade arrangement is a formal \mathbb{R} -linear combination of some translated blades $\{\beta_{J_1}, \beta_{J_2}, \dots, \beta_{J_l}\}$

Definition 2.13. Two sets $I, J \in {\binom{[n]}{k}}$ are weakly separated if there do not exist $a, b, c, d \in [n]$ such that

- (i) a < b < c < d or d < a < b < c or c < d < a < b or b < c < d < a;
- (*ii*) $a, c \in I \setminus J$ and $b, d \in J \setminus I$.

A collection of k-sets is called a weakly separated collection if any two in it are weakly separated.

In [1], Nick Early treated the weak separation phenomena from the viewpoint of discrete geometry. We will see in Lemma 2.27 that this is equivalent to telling whether two rays $\mathbb{R}_{\geq 0}\beta_I$ and $\mathbb{R}_{\geq 0}\beta_J$ are in some maximal cone of the positive tropical Grassmannian Trop⁺Gr_{k,n}.

Theorem 2.14. ([1]) The refinement of subdivisions $\{\beta_{J_1}, \beta_{J_2}, \dots, \beta_{J_l}\}$ is a positroid subdivision of $\Delta_{k,n}$ if and only if $\{J_1, J_2, \dots, J_l\}$ is a weakly separated collection.

2.3 Construction of positive tropical Grassmannian by weighted blade arrangement

For any $j \in [n]$, the multi-split β_J over $\Delta_{k,n}$ may also induce a multi-split over the boundary $\partial_{(\{j\},1)}(\Delta_{k,n}) := \{x \in \Delta_{k,n} \mid x_j = 1\} \cong \Delta_{k-1,n-1}$. The following lemma illustrates the process

Lemma 2.15. ([1]) The multi-split over the boundary satisfies

$$\partial_{(\{j\},1)}(((1,2,\cdots,n))_{e_J} \cap \Delta_{k,n}) = ((1,2,\cdots,\widehat{j},\cdots,n))_{e_{J'}} \cap \partial_{(\{j\},1)}(\Delta_{k,n})$$

where $J = \{j_1, \dots, j_k\}$ and $J' = J \setminus \{j_{a+1}\}$ if j satisfies $j_a < j \le j_{a+1}$ (the indices are cyclic). We can also replace $\{j\}$ by a subset $L = \{a_1, \dots, a_t\} \subset [n]$ such that $1 \le t \le k-2$, then

$$\partial_{(L,1)} := \partial_{(\{a_1\},1)} \circ \partial_{(\{a_2\},1)} \circ \cdots \circ \partial_{(\{a_t\},1)}$$

does not depend on the order of $\{j_1, \dots, j_t\}$. Besides, the order in $[n] \setminus L$ is in inherited from $1 \le 2 \le \dots \le n$. So the boundary map on a k-subset can be defined as $\partial_{\{j\},1\}}(J) := J'$.

The boundary map on a collection of balde arrangements can be extended linearly to weighted blade arrangements. We will replace $\partial_{(\{j\},1)}$ by ∂_j in the following passage for simplicity. Let $\beta_J^{(L)}$ be the translated blade on $\partial_L(\Delta_{k,n})$ for a subset $L \subset [n]$ with $0 \le |L| \le k - 2$ and $J \subset {[n] \setminus L \choose k - |L|}$. The operator ∂_j can be defined on the linear span of $\beta_J^{(L)}$ as follows,

$$\partial_j(\beta_J^{(L)}) = \begin{cases} 0 & j \in L \\ \beta_{J \setminus \{j_{a+1}\}}^{(L \cup \{j\})} & j \notin L \end{cases}$$

where $J = \{j_1, \dots, J_{k-|L|}\}$ and $j_a < j \le j_{a+1}$ in the cyclic order of $[n] \setminus L$ inherited from [n].

Definition 2.16. ([6]) Let

$$\mathfrak{B}_{k,n}^{\bullet} := \bigoplus_{m=0}^{n-(k-2)} \left(\bigoplus_{L \in \binom{[n]}{m}} \mathfrak{B}_{k,n}^{(L)} \right),$$

where $\mathfrak{B}_{k,n}^{(L)}$ is a vector space of formal linear span of $\beta_J^{(L)}$

$$\mathfrak{B}_{k,n}^{(L)} := \operatorname{span}\left\{\beta_J^{(L)} : J \in \binom{[n] \setminus L}{k - |L|}\right\}$$

Let $\mathfrak{B}_{k,n} := \mathfrak{B}_{k,n}^{(\emptyset)}$ be the top component of $\mathfrak{B}_{k,n}^{\bullet}$. Then $(\mathfrak{B}_{k,n}^{\bullet}, \partial)$ is called the **hypersimplicial blade complex** where $\partial := \partial_1 + \partial_2 + \cdots + \partial_n$ is the sum of all the boundary maps.

Remark 2.17. $(\mathfrak{B}^{\bullet}_{k,n},\partial)$ is not a complex in the usual sense since $\partial^2 \neq 0$ although $\partial_j^2 = 0$ for any $j \in [n]$. In the following text, we will see that a weighted blade is a function with a positroidal decomposition of $\Delta_{k,n}$, however the sum of weighted blades is not necessary a function with positroidal decomposition of $\Delta_{k,n}$. Therefore, we need a subset $\mathbb{Z}_{k,n}$ of such sums, whose elements induce positroidal decomposition.

We consider a weighted blade arrangement \mathcal{L}_J denoted by

$$\mathcal{L}_J := \sum_{M \subset J^{\bullet}} (-1)^{1+|M|} \beta_{J_M},$$

where $C_{J^{\bullet}} = \left\{ e_{J_M} = e_J + \sum_{i=1}^t \delta_M(j_i)(e_{j_i-1} - e_{j_i}) \mid M \subset J^{\bullet} \right\}$ with $J^{\bullet} = \{j_1, \dots, j_t\}$ being the initial points of the cyclic intervals of J and δ_M being the characteristic function of subset $M \subset J^{\bullet}$ (that is, $\delta_M(j_i) = 1$ if $j_i \in J^{\bullet}$, otherwise $\delta_M(j_i) = 0$).

Proposition 2.18. We have the following relation

$$\sum_{I \in \binom{[n]}{k}} d(e_J, e_I) \mathcal{L}_I = n\beta_J - (\beta_{[1,k]} + \beta_{[2,k+1]} + \dots + \beta_{[n,k-1]})$$

where $d(e_J, e_I)$ is the smallest positive number of steps from e_J to e_I , where each step has to be one of the directions $e_1 - e_2, e_2 - e_3, \dots, e_n - e_1$.

Proof. Let $K_{\bullet} = \{k'_1, k'_2, \dots, k'_t\}$ be the end points of cyclic intervals of K for some $K \in \binom{[n]}{k}$ and $C_{K_{\bullet}} = \begin{cases} e_{K_M} = e_K - \sum_{i=1}^t \delta_M(k'_i)(e_{k'_i} - e_{k'_i+1}) \mid M \subset K_{\bullet} \end{cases}$. Then we calculate the coefficient of β_K on the left side of the equation. It is equal to

$$\sum_{M\subset K_{\bullet}}(-1)^{1+|M|}d(e_J,e_{K_M}).$$

Let $e_K - e_J = \sum_{j=1}^n t_j (e_j - e_{j+1})$ be the unique positive expression of $e_K - e_J$, so $d(e_J, e_K) = \sum_{j=1}^n t_j$ and $e_{K_M} - e_J = e_{K_M} - e_K + e_K - e_J$ $= -\sum_{i=1}^t \delta_M(k'_i)(e_{k'_i} - e_{k'_i+1}) + \sum_{i=1}^n t_j(e_j - e_{j+1}).$

Case 1: $K \neq J$. The unique positive expression of $e_{K_M} - e_J$ can be written as

$$e_{K_M} - e_J = \begin{cases} -\sum_{i=1}^t \delta_M(k'_i)(e_{k'_i} - e_{k'_i+1}) + \sum_{j=1}^n t_j(e_j - e_{j+1}) & M \setminus Supp(e_K - e_J) = \emptyset \\ \sum_{l=1}^n (e_l - e_{l+1}) - \sum_{i=1}^t \delta_M(k'_i)(e_{k'_i} - e_{k'_i+1}) + \sum_{j=1}^n t_j(e_j - e_{j+1}) & M \setminus Supp(e_K - e_J) \neq \emptyset \end{cases}$$

so the distance from e_J to e_{K_M} can be expressed by

$$d(e_J, e_{K_M}) = \begin{cases} d(e_J, e_K) - |M| & M \setminus Supp(e_K - e_J) = \emptyset \\ d(e_J, e_K) - |M| + n & M \setminus Supp(e_K - e_J) \neq \emptyset \end{cases}$$

We noticed that every $M \subset K_{\bullet}$ has a unique partition $M = M' \sqcup M''$ where $M' \subset K_{\bullet} \setminus Supp(e_K - e_J)$ and $M'' \subset K_{\bullet} \cap Supp(e_K - e_J)$. By Lemma 2.19, $K_{\bullet} \cap Supp(e_K - e_J) \neq \emptyset$, so we consider two subcases as follows

Subcase 1: $|K_{\bullet}| = 1$ (This is equivalent to that K is an interval). Then $K_{\bullet} \subset Supp(e_K - e_J)$, so the coefficient of β_K is

$$-d(e_J, e_K) + (d(e_J, e_K) - 1) = -1$$

Subcase 2: $|K_{\bullet}| \ge 2$. Then

$$\begin{split} \sum_{M \subset K_{\bullet}} (-1)^{1+|M|} d(e_J, e_{K_M}) &= (-1)^{1+|L|} \sum_{\substack{M \subset K_{\bullet} \\ M' \neq \emptyset}} (d(e_J, e_K) - |M| + n) + \sum_{\substack{M \subset K_{\bullet} \\ M = M''}} (-1)^{1+|M|} (d(e_J, e_K) - |M| + n) - \sum_{\substack{M \subset K_{\bullet} \\ M = M''}} (-1)^{1+|M|} \cdot n = 0 - 0 = 0 \end{split}$$

Case 2: K = J. In this case,

$$e_{K_M} - e_J = \sum_{l=1}^n (e_l - e_{l+1}) - \sum_{i=1}^t \delta_M(k_i')(e_{k_i'} - e_{k_i'+1})$$

is the unique positive expression of $e_{K_M} - e_J$ for $M \neq \emptyset$, so $d(e_{K_M}, e_J) = n - |M|$. Similarly, we consider two subcases.

Subcase 3: $|K_{\bullet}| = 1$. The coefficient of β_K is n - 1. Subcase 4: $|K_{\bullet}| \ge 2$. The coefficient is

$$\sum_{M \subset K_{\bullet}} (-1)^{1+|M|} d(e_J, e_{K_M}) = \sum_{\substack{M \subset K_{\bullet} \\ L \neq \emptyset}} (-1)^{1+|M|} (n - |M|) = n$$

Lemma 2.19. For any $K, J \subset {\binom{[n]}{k}}$ and $K \neq J$, we have $K_{\bullet} \cap Supp(e_K - e_J) \neq \emptyset$.

Proof. Take the unique positive expression $e_K - e_J = \sum_{l=1}^n t_l(e_l - e_{l+1})$, it is not hard to find that either $|t_l - t_{l+1}| = 0$ or 1. Since $K \neq J$, there exists some $a \in K \setminus J$ such that $t_a \neq 0$. Then we can take the minimal positive integer b such that $t_{a+b} = 0$ (Such b do exists since not all the t_l are positive). This implies that $t_{a+b-1} = 1$. So the coefficient of e_{a+b} is -1 which implies that $a + b \in J \setminus K$. Therefore there must exist some point of K_{\bullet} appearing in the interval [a, a + b].

Corollary 2.20. $(\mathcal{L}_J)_{J \in \binom{[n]}{k}}$ is a basis of linear space $\mathfrak{B}_{k,n} = \operatorname{span} \left\{ \beta_J : J \in \binom{[n]}{k} \right\}.$

Proof. The transition matrix from $(\beta_J)_{J \in \binom{[n]}{k}}$ to $(n\beta_J - (\beta_{[1,k]} + \beta_{[2,k+1]} + \dots + \beta_{[n,k-1]}))_{J \in \binom{[n]}{k}}$ is

(п	0	•••	0	0	0	•••	0)
	0	п	•••	0	0	0	•••	0
	÷	÷	·	÷	:	:	·	:
	0	0		n	0	0		0
-	-1	-1	•••	-1	<i>n</i> – 1	-1	• • •	-1
	-1 -1	-1 -1	· · · · · · ·	-1 -1	n - 1 - 1	$-1 \\ n - 1$	· · · · · · ·	-1 -1
	-1 -1 :	-1 -1 :	···· ···	-1 -1 :	n-1 -1 \vdots	-1 n-1 \vdots	···· ···	-1 -1 :

which is invertible for $n \ge 4$. So $(\mathcal{L}_J)_{J \in \binom{[n]}{k}}$ is a basis.

Remark 2.21. Similar to Definition 2.16, define $(\mathcal{L}_J^{(L)})_{J \in \binom{[n]\setminus L}{k-|L|}}$ to be the linear combination of $(\beta_J^{(L)})_{J \in \binom{[n]\setminus L}{k-|L|}}$ for any $L \subset [n]$ with $0 \leq |L| \leq k-2$. The cyclic intervals of J should be taken in the base set $[n]\setminus L$, where the cyclic order of $[n]\setminus L$ is inherited from [n] (This is explicitly defined in Definition 2.23). Thus by Corollary 2.20, $(\mathcal{L}_J^{(L)})_{J \in \binom{[n]\setminus L}{k-|L|}}$ is an \mathbb{R} -basis of $\mathfrak{B}_{k,n}^{(L)}$.

Example 2.22. *Take* k = 2, n = 4 and J = 13, we have

$$d(e_{13}, e_{12}) = d(e_{13}, e_{34}) = 1 \quad d(e_{13}, e_{13}) = 0$$

$$d(e_{13}, e_{14}) = d(e_{13}, e_{23}) = 3 \quad d(e_{13}, e_{24}) = 2$$

It is not hard to find that $d(e_J, -)$ satisfies the positive tropical Plücker relation $d(e_{13}, e_{13}) + d(e_{13}, e_{24}) = \min\{d(e_{13}, e_{12}) + d(e_{13}, e_{34}), d(e_{13}, e_{14}) + d(e_{13}, e_{23})\}$. The basis $(\mathcal{L}_J)_{J \in \binom{[4]}{J}}$ are

$$\begin{array}{ccc} \mathcal{L}_{12} = -\beta_{12} + \beta_{24} & \mathcal{L}_{14} = -\beta_{14} + \beta_{13} \\ \mathcal{L}_{23} = -\beta_{23} + \beta_{13} & \mathcal{L}_{34} = -\beta_{34} + \beta_{24} \\ \mathcal{L}_{24} = -\beta_{24} + \beta_{14} + \beta_{23} - \beta_{13} & \mathcal{L}_{13} = -\beta_{13} + \beta_{12} + \beta_{34} - \beta_{24} \end{array}$$

Thus

$$\sum_{I \in \binom{[4]}{2}} d(e_J, e_I) \mathcal{L}_I = (-\beta_{12} + \beta_{24}) + 3(-\beta_{14} + \beta_{13}) + 3(-\beta_{23} + \beta_{13}) + (-\beta_{24} + \beta_{14} + \beta_{23} - \beta_{13}) + (-\beta_{34} + \beta_{24}) = 4\beta_{13} - \beta_{12} - \beta_{23} - \beta_{34} - \beta_{14}.$$

Definition 2.23. Let $[n] \setminus L = \{l_1, l_2, \dots, l_{n-m}\}$ with $l_1 < l_2 < \dots < l_{n-m}$ for some $L \subset [n]$ with |L| = m $(0 \le m \le k-2)$. Then $l_I := \{l_i \mid i \in I\}$ is called frozen in $[n] \setminus L$ if I is a cyclic interval in [n-m], otherwise unfrozen where $I \subset [n-m]$ and |I| = k - m.

Proposition 2.24. ([6]) For any $j \in [n]$, the action of boundary map ∂_j on the basis $(\mathcal{L}_J^{(L)})_{J \in \binom{[n] \setminus L}{k-|L|}}$ can be described as follows

$$\partial_{j}(\mathcal{L}_{J}^{(L)}) = \begin{cases} \mathcal{L}_{J \setminus \{j\}}^{(L \cup \{j\})} & j \in J \\ 0 & j \notin J \end{cases}$$

Proof. To avoid more notations, we suppose that $L = \emptyset$. Let $J^{\bullet} = \{j_1, j_2, \dots, j_t\}$. Therefore, for any $j \in [n]$, there exists a unique pair $\{j_a, j_{a+1}\}$ such that $j_a \leq j < j_{a+1}$. Then

$$\partial_j(\mathcal{L}_J) = \sum_{M \subset J^\bullet} (-1)^{1 + |M|} \partial_j(\beta_{J_M})$$

If $j \notin J$, we have $\partial_j(\beta_{J_M}) = \partial_j(\beta_{J_M \cup \{j_{a+1}\}})$ for any $M \subset J^{\bullet} \setminus \{j_{a+1}\}$. Therefore,

$$\partial_j(\mathcal{L}_J) = \sum_{M \subset J^{\bullet} \setminus \{j_{a+1}\}} ((-1)^{1+|M|} + (-1)^{1+|M \cup \{j_{a+1}\}|}) \partial_j(\beta_{J_M}) = 0$$

Or if $j \in J$, then $\partial_j(\mathcal{L}_J) = \mathcal{L}_{J \setminus \{j\}}^{(\{j\})}$ can be vertified directly.

Corollary 2.25. ([6]) Given any $\mathbf{c} = (c_I)_{I \in \binom{[n]}{k}} \in \mathbb{R}^{\binom{[n]}{k}}$ and $\mathcal{L}(\mathbf{c}) := \sum_{J \in \binom{[n]}{k}} c_J \mathcal{L}_J$. $\partial_L(\mathcal{L}(\mathbf{c}))$ can be written as a

linear combination as

$$\partial_L(\mathcal{L}(\mathbf{c})) = \sum_{i,j \in [n-k+2]} \pi_{l_i l_j}^{(L)} \beta_{l_i l_j}^{(L)}$$

where |L| = k - 2 and $[n] \setminus L = \{l_1, l_2, \dots, l_{n-k+2}\}$. Choose any unfrozen pair $\{l_i, l_j\}$ (i.e. $|i - j| \ge 2$), the coefficient $\pi_{l_i l_j}^{(L)} = -(c_{Ll_i l_j} - c_{Ll_i l_{j+1}} - c_{Ll_{i+1} l_j} + c_{Ll_{i+1} l_{j+1}})$

Proof. The coefficient of $\beta_{l_i l_j}^{(L)}$ is from $\mathcal{L}_{l_i l_j}^{(L)}$, $\mathcal{L}_{l_{i+1} l_j}^{(L)}$, $\mathcal{L}_{l_i l_{j+1}}^{(L)}$ and $\mathcal{L}_{l_{i+1} l_{j+1}}^{(L)}$. Thus by Proposition 2.24, we get this relation.

Take any

$$\beta(\mathbf{c}) = \sum_{J \in \binom{[n]}{k}} c_I \beta_I \in \mathfrak{B}_{k,n}$$

where $\mathbf{c} = (c_I) \in \mathbb{R}^{\binom{n}{k}}$. Choose some $L \subset [n]$ with |L| = k - 2, then the image of $\beta(\mathbf{c})$ under the boundary map is

$$\partial_L(\beta(\mathbf{c})) = \sum_{\{i,j\} \subset [n] \setminus L \text{ nonfrozen}} \pi_{l_i l_j}^{(L)} \beta_{l_i l_j}^{(L)}$$

where $\pi_{l_i l_j}^{(L)} = \sum_{\substack{I \in \binom{[n]}{k} : \partial_L(\beta_I) = \beta_{l_i l_j}^{(L)} \\ 0 \text{ f } \beta_{l_i l_j}^{(L)} \text{ such that } \pi_{l_i l_i}^{(L)} \neq 0.} c_I$. Let the notation $Supp_L(\beta(\mathbf{c}))$ denote the support of $\beta(\mathbf{c})$ under $\partial_L i.e.$ the set

Definition 2.26. Denote by $Z_{k,n}$ the collection of elements $z \in \mathfrak{B}_{k,n}$ that satisfy

- (i) (Compatibility condition) Supp_L(z) induce a positroid subdivision of $\partial_L(\Delta_{k,n})$.
- (ii) (Positivity condition) $\pi_{l_i l_i}^{(L)} \ge 0$ for any nonfrozen $\{l_i, l_j\}$ in $[n] \setminus L$.

for any $L \subset [n]$ with |L| = k - 2.

The following important lemma shows an explicit relation between $Z_{k,n}$ and positive tropical Grassmannian Trop⁺Gr_{k,n}.

Lemma 2.27. ([6]) Let $\mathbf{c} = (c_I)_{I \in \binom{[n]}{k}} \in \mathbb{R}^{\binom{n}{k}}$. Then \mathbf{c} satisfies the positive tropical Plücker relation

 $c_{Ll_al_c} + c_{Ll_bl_d} = \min\{c_{Ll_al_b} + c_{Ll_cl_d}, c_{Ll_al_d} + c_{Ll_bl_c}\}$

for any cyclic order a < b < c < d in [n - k + 2] if and only if $\mathcal{L}(\mathbf{c}) := \sum_{I} c_{I} \mathcal{L}_{I} \in \mathbb{Z}_{k,n}$. Moreover, this induces a bijection between the positive tropical Grassmannian Trop⁺Gr_{k,n} and $\mathbb{Z}_{k,n}$.

Proof. Suppose that $\mathbf{c} \in \text{Trop}^+\text{Gr}_{k,n}$. In particular, we take a = i, b = i + 1, c = j, d = j + 1, then by Corollary 2.25

$$\pi_{l_i l_j}^{(L)} = -(c_{Ll_i l_j} - c_{Ll_i l_{j+1}} - c_{Ll_{i+1} l_j} + c_{Ll_{i+1} l_{j+1}}) \ge 0$$

for any nonfrozen pair $\{l_i, l_j\}$. It remains to show that $Supp_L(\mathcal{L}(\mathbf{c}))$ induce a positroid subdivision of $\partial(\Delta_{k,n})$. Since $\{l_i, l_j\}$ and $\{l_p, l_q\}$ are not weakly separated where $p \in [i + 1, j - 1]$ and $q \in [j + 1, i - 1]$, we have

$$\sum_{\substack{p \in [i+1,j-1] \\ q \in [j+1,i-1]}} \pi_{l_p l_q}^{(L)} = -(c_{Ll_i l_j} - c_{Ll_i l_{i+1}} - c_{Ll_j l_{j+1}} + c_{Ll_{i+1} l_{j+1}}).$$

It follows that min $\left\{ \pi_{l_i l_j}^{(L)}, \sum_{\substack{p \in [i+1,j-1] \\ q \in [j+1,i-1]}} \pi_{l_p l_q}^{(L)} \right\}$ is equal to

$$\min\{-(c_{Ll_il_j} - c_{Ll_il_{j+1}} - c_{Ll_{i+1}l_j} + c_{Ll_{i+1}l_{j+1}}), -(c_{Ll_il_j} - c_{Ll_il_{i+1}} - c_{Ll_jl_{j+1}} + c_{Ll_{i+1}l_{j+1}})\} = \min\{(c_{Ll_il_{j+1}} + c_{Ll_{i+1}l_j}), (c_{Ll_il_{i+1}} + c_{Ll_jl_{j+1}})\} - (c_{Ll_il_j} + c_{Ll_{i+1}l_{j+1}}) = 0.$$

Thus this proves that $\mathcal{L}(\mathbf{c})$ satisfies (i) in Definition 2.26.

Conversely, suppose that $\mathcal{L}(\mathbf{c}) \in \mathbb{Z}_{k,n}$. Given any a < b < c < d in [n - k + 2], we observed that any $\{l_p, l_q\}$ in A and $\{l_{p'}, l_{q'}\}$ in B are not weakly separated where

$$A := \{\{l_p, l_q\} \mid p \in [a, b-1], q \in [c, d-1]\}$$
$$B := \{\{l_{p'}, l_{q'}\} \mid p' \in [b, c-1], q' \in [d, a-1]\}$$

Therefore min $\left\{\sum_{\{l_p,l_q\}\in A} \pi_{l_pl_q}^{(L)}, \sum_{\{l_{p'},l_{q'}\}\in B} \pi_{l_{p'}l_{q'}}^{(L)}\right\} = 0$ by conditions (i) and (ii) in Definition 2.26. As explained on the left of Figure 1, $\pi_{l_pl_q}^{(L)}$ is presented by the labeled square with a number on each corner. The number on the corner (p,q) refers to the sign of $c_{Ll_pl_q}$ in the expansion of Corollary 2.25.

$$\begin{array}{c} (a,d) \\ (a,d-1) \\ \hline 1 & 0 \\ (a,d-1) \\ \hline 1 & 0 \\ (a,d-1) \\ \hline 1 & 0 \\ (a,d-1) \\ \hline 0 & 0 \\ (a,c+1) \\ \hline 0 & 0 \\ (a,c+1) \\ \hline 0 & 0 \\ (a,c+1) \\ \hline 0 & 0 \\ (a,c) \\ (a+1,c) \\ (b,c+1) \\ (b$$

Thus $\sum_{\{l_p, l_q\} \in A} \pi_{l_p l_q}^{(L)}$ can be presented on the right by splicing all the squares together. And we add up all the numbers overlapped at the lattice point. So

$$\begin{split} & \sum_{\{l_p, l_q\} \in A} \pi_{l_p l_q}^{(L)} = -(c_{Ll_a l_c} - c_{Ll_a l_d} - c_{Ll_b l_c} + c_{Ll_b l_d}) \\ & \sum_{\{l_{p'}, l_{q'}\} \in B} \pi_{l_{p'} l_{q'}}^{(L)} = -(c_{Ll_b l_d} - c_{Ll_a l_b} - c_{Ll_c l_d} + c_{Ll_a l_c}) \end{split}$$

Now it is not difficult to see that min $\begin{cases} \sum_{\{l_p, l_q\} \in A} \pi_{l_p l_q}^{(L)}, & \sum_{\{l_{p'}, l_{q'}\} \in B} \pi_{l_{p'} l_{q'}}^{(L)} \end{cases} = 0 \text{ is equivalent to } c_{Ll_a l_c} + c_{Ll_b l_d} = \min\{c_{Ll_a l_b} + c_{Ll_b l_c}\}. \end{cases}$

Corollary 2.28. The vector $\mathbf{d}_J := (d(e_J, e_I))_{I \in \binom{[n]}{k}}$ satisfies the positive tropical Plücker relations. Moreover, the subdivison $\mathcal{D}_{\mathbf{d}_I}$ is exactly that induced by translated blade $((1, 2, \dots, n))_{e_I}$.

Proof. Since $h(x - e_J) = \min\{L_1(x - e_J), L_2(x - e_J), \dots, L_n(x - e_J)\}\$ is concave on the hyperplane $\sum_{i=1}^n x_i = k$, we know that $-h(x - e_J)$ is convex. By Proposition 2.8, the intersections of all the affine areas of $-h(x - e_J)$ with $\Delta_{k,n}$ form the subdivision induced by $((1, 2, \dots, n))_{e_J}$. On the other hand, because of the convexity of $-h(x - e_J)$ and $-h(e_J, e_I) = d(e_J, e_I)$, the projection of any lower face of convex($\{(e_I, d(e_J, e_I))\}\)$ is exactly the intersection of some affine area of $-h(x - e_J)$ with $\Delta_{k,n}$.

Remark 2.29. From Corollary 2.28 together with the equation in Proposition 2.18

$$\sum_{I \in \binom{[n]}{k}} d(e_J, e_I) \mathcal{L}_I = n\beta_J - (\beta_{[1,k]} + \beta_{[2,k+1]} + \dots + \beta_{[n,k-1]}),$$

 β_J can be regarded as the positroid subdivision induced by $((1, 2, \dots, n))_{e_J}$. Also, we ignore the negative terms on the right since β_I induces a trivial subdivision of Δ when I is a cyclic interval. Through these observations, it is not hard to see that $\mathcal{Z}_{k,n}$ contains a linearity spcae $\mathcal{F}_{k,n} := \text{span}\{\beta_I \mid I \text{ is frozen}\}$.

Definition 2.30. Let $\overline{Z_{k,n}} := Z_{2,n}/\mathcal{F}_{k,n}$ be the quotient fan modulo the linearity space. Then evel element $x \in \overline{Z_{k,n}}$ can be uniquely written as a weighted blade arrangement of nonfrozen translated blades.

2.4 The first theorem

From now on we only consider all the weighted blade arrangements in $\overline{Z_{k,n}}$, that is, we will omit all the terms of frozen translated blades. Let us first see the case k = 2.

Proposition 2.31. All the maximal cones of $\overline{\mathbb{Z}_{k,n}}$ are simplicial with dimension n-3. And they are of the form $\mathcal{P}_W := \{\sum_{J \in W} \mathbb{R}_{\geq 0} \beta_J\}$ for some nonfrozen maximal w-collection W.

Proof. Any finest positroidal subdivision of $\Delta_{k,n}$ contains n-2 top dimensional cells (see [4]) and n-3 facets which split these cells (2-splits). Every translated blade to a non-frozen vertex induces a 2-split. Therefore we have weakly separated collection of n-3 vertices corresponding to these splits. But n-3 is a maximal cardinality of a weakly separated collection of two-elements sets. This implies that the cone spanned by a such collection of weighted blades forms a simplicial cone of $\overline{Z_{2,n}}$. Moreover, there is a bijection between maximal weakly separated sets of two elements sets and vertices of the associahedron (see Fomin and Zelevinsky [16]). Therefore all simplical cones of $\overline{Z_{2,n}}$ are of such a form.

From Proposition 2.31, we know that every maximal cone in $\overline{Z_{2,n}}$ is simplicial. However when $k \ge 3$, it is not the case. Here we provide a necessary and sufficient condition for $\mathcal{P}_W := \{\sum_{J \in W} \mathbb{R} \ge \beta_J\}$ being a maximal simplical cone in $\overline{Z_{k,n}}$.

Theorem 2.32. The following statements are equivalent

- (1) Let $W = \{J_1, \dots, J_l\} \subset {\binom{[n]}{k}}$ be a nonfrozen maximal w-collection and $W' \subseteq W$ then W' = W if and only if $\partial_L(W')$ contains a nonfrozen maximal w-collection in ${\binom{[n]\setminus L}{2}}$ for any $L \subseteq [n]$ with |L| = k 2. Equivalently, W is minimal in S defined in Theorem 1.2.
- (2) The positive linear expansion $\mathcal{P} = \{\mathbb{R}_{\geq 0}\beta_{J_1} + \cdots \otimes \mathbb{R}_{\geq 0}\beta_{J_l}\}$ forms a maximal simplicial cone in $\overline{\mathcal{Z}_{k,n}}$ where $W = \{J_1, J_2, \cdots, J_l\}$ is a nonfrozen maximal w-collection in $\binom{[n]}{k}$ for $2 \le k \le n-2$.

Proof. (1) \Rightarrow (2). It is obvious for k = 2 by Proposition 2.31. For $k \geq 3$, since every octahedral face $\{Lab, Lbc, Lcd, Lad, Lac, Lbd\}$ (a < b < c < d is cyclic in [n]) is contained in some face $\partial_L(\Delta_{k,n})$, by assumption, $\{\beta_{J'}^{(L)}|J' \in \partial_L(W)$ induce a finest positroid subdivision of $\partial_L(\Delta_{k,n}) \cong \Delta_{2,n-k+2}$. By Theorem 2.6, every octahedral face in $\Delta_{k,n}$ is subdivided. Thus we conclude that the regular positroid subdivision induced by $\beta_{J_1}, \dots, \beta_{J_l}$ is the finest. Let $\mathcal{P} := \{\mathbb{R}_{\geq}\beta_{J_1} + \mathbb{R}_{\geq}\beta_{J_2} + \dots \mathbb{R}_{\geq}\beta_{J_l}\}$ be the cone spanned by $\beta_{J_1}, \dots, \beta_{J_l}$, then \mathcal{P} is contained in some maximal cone \mathcal{M} of $\overline{Z}_{k,n}$ by the compatibility of $\{\beta_{J_1}, \beta_{J_2}, \dots, \beta_{J_l}\}$. We consider the face $\mathcal{P}' = \{\sum_{J'_i \in W'} \mathbb{R}_{\geq}\beta_{J'_i}\}$ of \mathcal{P} spanned by a proper subset $\{\beta_{J'_i} \mid J'_i \in W'\}$ of $\{\beta_{J_1}, \beta_{J_2}, \dots, \beta_{J_l}\}$. Now by assumption in (1), there exists a subset $L_0 \subseteq [n]$ with $|L_0| = k - 2$ such that $\partial_{L_0}(W')$ does not contain a nonfrozen maximal w-collection. Therefore the subdivision of $\partial_{L_0}(\Delta_{k,n}) \cong \Delta_{2,n-k+2}$ induced by $\partial_{L_0}(W')$ is not the finest which implies that there exists an octahedral face in $\partial_{L_0}(\Delta_{k,n})$ not subdivided. Again by Theorem 2.6, we know that $\{\beta_{J'_i}|J'_i \in W'\}$ does not induce a finest positroid subdivision of $\Delta_{k,n}$. This proves that the faces of \mathcal{P} do not intersect the interior of \mathcal{M} . On the other hand, it is showed in Theorem 4.1 of [3] and Example 4.20 of [6], there exists a parametrization (a surjection) from $\mathbb{R}^{(k-1)(n-k-1)}$ to $\overline{Z}_{k,n}$. Therefore $dim(\mathcal{M}) \leq (k-1)(n-k-1)$. Since $\{\beta_{J_1}, \beta_{J_2}, \dots, \beta_{J_l}\}$ are linear independent and the cardinality of W is (k-1)(n-k-1), we have $(k-1)(n-k-1) = dim(\mathcal{P}) \leq dim(\mathcal{M}) \leq (k-1)(n-k-1)$. Thus $\mathcal{P} = \mathcal{M}$ is maximal simplicial.

(2) \Rightarrow (1). Conversely we know that the positive tropical Grassmannian possess a secondary fan structure. Thus from the assumption that \mathcal{P} is maximal, we say that $\{\beta_J | J \in W\}$ induces a finest positroid subdivision of $\Delta_{k,n}$. So every octahedral face is subdivided by Theorem 2.6. Therefore $\partial_L(W)$ contains a maximal w-collection of $\binom{[n]\setminus L}{2}$. On the other hand, if we take any proper subset $W' \subseteq W$, since \mathcal{P} is simplicial, $\mathcal{P}' = \{\sum_{J'_i \in W'} \mathbb{R}_{\geq 0} \beta_{J'_i} | J'_i \in W'\}$ does not induce a finest positroid subdivision of $\Delta_{k,n}$. Then there exists some L_0 such that $\partial_{L_0}(W)$ does not contains a maximal w-collection of $\binom{[n]\setminus L}{2}$.

Example 2.33. In $\Delta_{3,6}$, we take four nonfrozen maximal w-collections {124, 125, 134, 145}. Take the boundary map,

$$\begin{split} \partial(\beta_{124}) &= \beta_{24}^{(1)} + \beta_{14}^{(2)} + \beta_{24}^{(5)} + \beta_{24}^{(6)} \\ \partial(\beta_{125}) &= \beta_{25}^{(1)} + \beta_{15}^{(2)} + \beta_{25}^{(6)} \\ \partial(\beta_{134}) &= \beta_{14}^{(2)} + \beta_{14}^{(3)} + \beta_{13}^{(4)} \\ \partial(\beta_{145}) &= \beta_{15}^{(2)} + \beta_{15}^{(3)} + \beta_{15}^{(4)} + \beta_{14}^{(5)} \end{split}$$

It can be checked that {124, 125, 134, 145} satisfy (1) of Theorem 2.32, thus rays β_{124} , β_{125} , β_{134} , β_{145} comprise a maximal simplicial cone in $\mathbb{Z}_{3,6}$.

Below is an example showing that the positive weighted blade arrangements from some nonfrozen maximal w-collection does not form a maximal simplicial cone in $\overline{Z_{k,n}}$.

Example 2.34. Still in $\Delta_{3,6}$, consider a nonfrozen maximal w-collection {135, 235, 145, 136}. Then

$$\begin{aligned} \partial(\beta_{135}) &= \beta_{35}^{(1)} + \beta_{15}^{(2)} + \beta_{15}^{(3)} + \beta_{13}^{(4)} + \beta_{13}^{(5)} + \beta_{35}^{(6)} \\ \partial(\beta_{235}) &= \beta_{35}^{(1)} + \beta_{35}^{(2)} + \beta_{25}^{(3)} + \beta_{35}^{(6)} \\ \partial(\beta_{145}) &= \beta_{15}^{(2)} + \beta_{15}^{(3)} + \beta_{15}^{(4)} + \beta_{14}^{(5)} \\ \partial(\beta_{136}) &= \beta_{36}^{(1)} + \beta_{13}^{(4)} + \beta_{13}^{(5)} + \beta_{13}^{(6)} \end{aligned}$$

Obviously $Supp(\beta_{135}) \subset Supp(\beta_{235}) \cup Supp(\beta_{145}) \cup Supp(\beta_{136})$. In fact the support of 3-split β_{135} is contained in the support of the 2-splits $\beta_{235}, \beta_{145}, \beta_{136}$, that is,

$$((16)_1, (23)_1, (45)_1) \cap \Delta_{3,6} \subset (\{x_{1236} = 2\} \cup \{x_{16} = 1\} \cup \{x_{1456} = 2\}) \cap \Delta_{3,6}.$$

And here we explicitly write the generators of this cone in standard coordinate $\{\mathcal{L}_I\}$ of $\mathbb{R}^{\binom{6}{3}}$ where $I \in \binom{[6]}{3}$.

$$\mathbb{R}_{\geq}\beta_{135} = \mathbb{R}_{\geq}(3\mathcal{L}_{123} + 2\mathcal{L}_{124} + \mathcal{L}_{125} + 6\mathcal{L}_{126} + \mathcal{L}_{134} + 5\mathcal{L}_{136} + 5\mathcal{L}_{145} + 4\mathcal{L}_{146} + 3\mathcal{L}_{156} + 6\mathcal{L}_{234} + 5\mathcal{L}_{235} + 4\mathcal{L}_{236} + 4\mathcal{L}_{245} + 3\mathcal{L}_{246} + 2\mathcal{L}_{256} + 3\mathcal{L}_{345} + 2\mathcal{L}_{346} + \mathcal{L}_{356} + 6\mathcal{L}_{456})$$

$$\mathbb{R}_{\geq}\beta_{235} = \mathbb{R}_{\geq}(4\mathcal{L}_{123} + 3\mathcal{L}_{124} + 2\mathcal{L}_{125} + 7\mathcal{L}_{126} + 2\mathcal{L}_{134} + 1\mathcal{L}_{135} + 6\mathcal{L}_{136} + 6\mathcal{L}_{145} + 5\mathcal{L}_{146} + 6\mathcal{L}_{156} + \mathcal{L}_{234} + 5\mathcal{L}_{236} + 5\mathcal{L}_{245} + 4\mathcal{L}_{246} + 3\mathcal{L}_{256} + 5\mathcal{L}_{345} + 3\mathcal{L}_{346} + 2\mathcal{L}_{356} + 7\mathcal{L}_{456})$$

$$\mathbb{R}_{\geq}\beta_{145} = \mathbb{R}_{\geq}(4\mathcal{L}_{123} + 3\mathcal{L}_{124} + 2\mathcal{L}_{125} + 7\mathcal{L}_{126} + 2\mathcal{L}_{134} + \mathcal{L}_{135} + 6\mathcal{L}_{136} + 5\mathcal{L}_{146} + 4\mathcal{L}_{156} + 7\mathcal{L}_{234} + 6\mathcal{L}_{235} + 5\mathcal{L}_{236} + 5\mathcal{L}_{245} + 4\mathcal{L}_{246} + 3\mathcal{L}_{256} + 4\mathcal{L}_{345} + 3\mathcal{L}_{346} + 2\mathcal{L}_{356} + 7\mathcal{L}_{456})$$

$$\mathbb{R}_{\geq}\beta_{136} = \mathbb{R}_{\geq}(4\mathcal{L}_{123} + 3\mathcal{L}_{124} + 2\mathcal{L}_{125} + 1\mathcal{L}_{126} + 2\mathcal{L}_{134} + \mathcal{L}_{135} + 6\mathcal{L}_{145} + 5\mathcal{L}_{146} + 4\mathcal{L}_{156} + 7\mathcal{L}_{234} + 6\mathcal{L}_{235} + 5\mathcal{L}_{236} + 5\mathcal{L}_{245} + 4\mathcal{L}_{246} + 4\mathcal{L}_{256} + 4\mathcal{L}_{345} + 3\mathcal{L}_{346} + 2\mathcal{L}_{356} + 7\mathcal{L}_{456})$$

The above equations hold modulo the linearity space $\operatorname{span}_{\mathbb{R}} \{ \sum_{i \in I} \mathcal{L}_I \mid i = 1, \cdots, 6 \}$ of $\operatorname{Trop}^+\operatorname{Gr}_{3,6}$.

3 Boundary Maps on Reduced Plabic Graphs and The Second Theorem

In this section, we introduce some basic properties for reduced plabic graphs in [13]. Besides, we replace the boundary map ∂_j on a maximal w-collection W by a series of manipulations on corresponding reduced plabic graph $\Sigma_0(W)$. Then we prove Theorem 3.1 by showing that the face labels of $\partial_j(\Sigma_0(W))$ is exactly $\partial_j(W)$ and $\partial_j(\Sigma_0(W))$ is a biparite reduced plabic graph. As a corollary, we prove that the translated blades from a maximal w-collection induce a finest positroid subdivision of $\Delta_{k,n}$. At last, we characterize the flip between two finest positroid subdivisions by a pair of adjacent maximal cones in $\overline{Z}_{k,n}$.

Theorem 3.1. Let W be a maximal w-collection in $\binom{[n]}{k}$, then $\partial_j(W)$ is also a maximal w-collection in $\binom{[n]\setminus\{j\}}{k-1}$.

The proof of this Theorem will be conducted in the end of this section by using plabic graphs.

3.1 The boundary maps on plabic graphs

Definition 3.2. A plabic graph is a planar graph G embedded in a closed disk **D** satisfying the following conditions

- 1. No edges cross each other.
- 2. Each internal vertex is colored black or white.
- 3. Each internal vertex is connected by a path to some boundary vertex.
- 4. The boundary vertices are labeled by $1, 2, \dots, n$ in clockwise order for some $n \in \mathbb{Z}_{>0}$.
- 5. Each boundary vertex is incident to exactly one internal vertex by an edge.

Definition 3.3. Two plabic graphs G and G' are said to be move equivalenct to each other if G and G' can be related to each other via a sequence of the following local moves (M1), (M2) and (M3):

(M1) (The square move) Change the colors of vertices of a quadrilateral face whose vertices are trivalent.



Figure 2: (M_1)

(M2) Contract an edge of connecting two internal vertices of the same color or split an internal vertex into two vertices of the same color joined by an edge.



Figure 3: (M_2)

(M3) Remove a bivalent vertex and merge the edge adjacent to it or insert a bivalent in the middle of an edge.



Definition 3.4. Let W be a maximal w-collection in $\binom{[n]}{k}$, and $\Sigma_0(W)$ be the biparite reduced plabic graph obtained from the duality of plabic tiling. Define $\Sigma(W)$ to be the equivalence class of $\Sigma_0(W)$ under moves (M2) such that every *vertex has degree at least 3. (Obviously* $\Sigma_0(W) \in \Sigma(W)$ *by the definition of plabic tiling)*

According to the construction of plabic tiling in [13] or combined tiling in [17], Every black point in $\Sigma_0(W)$ is labeled by a (k + 1)-set \mathcal{B} such that the faces adjacent to \mathcal{B} are all the K-sets $\{\mathcal{B} \setminus b_1, \mathcal{B} \setminus b_2, \cdots, \mathcal{B} \setminus b_s\}$ of W arranged clockwise, where $b_1 < b_2 < \cdots < b_s$ is cyclic ordered. Similarly, every white point is labeled by a (k - 1)-set Wwhile the adjacent faces are $\{W \cup w_1, W \cup w_2, \dots, W \cup w_r\}$ in the clockwise order, where $w_1 < w_2 < \dots < w_r$ is cyclic ordered.

Definition 3.5. For any $i \in [n]$, $<_i$ denotes a linear order on [n] as follows:

$$i <_i i + 1 <_i i + 1 \cdots <_i i - 1$$

and for any subset $S \subset [n]$, we use the notation $M_i^i(S)$ to denote the set of minimal l elements in S under $<_i$.

Definition 3.6. Without loss of generality, we take j = n. The plabic graph $\partial_n(\Sigma_0(W))$ is obtained from $\Sigma_0(W)$ through the following steps

- 1. Delete the vertex labeled n k on the boundary and the unique edge adjacent to it, but preserve the internal vertex of this edge.
- 2. Delete all the edge with vertices \mathcal{B} and \mathcal{W} such that $\mathcal{B}\setminus \mathcal{W} = M_2^n(\mathcal{B})$.
- 3. Relabel the boundary vertices $n, 1, 2, \dots, n-k-1$ by $1, 2, \dots, n-k$ clockwise.
- 4. Delete all the single points and bivalent points, use moves (M2) to get a biparite plabic graph.

We use the same notation as the Appendix of [10]. The collection $W_0 = I_n^k \cup S_n^k$, where I_n^k consists of the intervals of size k and S_n^k consists of the sets of size k represented as the union of two nonempty intervals $[1,m] \cup [l, l+k-m-1]$ with $l \ge m+2$ and $l+k-m-1 \le n-1$. To prove Theorem 3.1, we first prove a special case then we show that the maximality of $\partial_i(W_0)$ preserves under flips.

3.2 Some lemmas for preparations

Lemma 3.7. Let $W_0 = I_n^k \cup S_n^k$, then $\partial_n(\Sigma_0(W_0)) \cong \Sigma_0(\partial_n(W_0))$

Proof. We can see that $\partial_n(W_0)$ consists of cyclic intervals of size k-1 in [n-1] and the sets of size k-1 representable as $[2,m] \cup [l, l+k-m-1]$. Clearly $\partial_n(W_0)$ is a maximal w-collection in $\binom{[n-1]}{k-1}$. Let us consider the local area of every face F of $\partial_n(\Sigma_0(W_0))$, where the local area, denoted by L(F), consists of face F and faces that have common vertex or common edge with F. We classify the L(F) into four cases:

Case 1: As in Figure 5, the left is the local area of face $F = \{1\} \cup [l, k + l - 2]$ with $l \ge 4$. Since $\partial_n([l-1, k+l-2]) = \partial_n(\{1\} \cup [l, k+l-2]) = [l, k+l-2]$ and $\partial_n([l, k+l-1]) = \partial_n(\{1\} \cup [l+1, k+l-1]) = [l+1, k+l-1]$, we delete the two edges that separate these two pairs of faces respectively. Then follow the rest steps in Definition 3.6, we get the graph on the right.



Figure 5: Boundary map on local area of $F = \{1\} \cup [l, k + l - 2]$ with $l \ge 4$

Case 2: As in Figure 6, the left is the local area of face $F = \{1\} \cup [3, k+1]$. Since $\partial_n([2, k+l]) = \partial_n(\{1\} \cup [3, k+l]) = [3, k+l]$ and $\partial_n([3, k+2]) = \partial_n(\{1\} \cup [4, k+2]) = [4, k+2]$, the same as Case 1, we delete these edges to combine the adjacent faces into one face.



Figure 6: Boundary map on local area of $F = \{1\} \cup [3, k+l]$

Case 3: As in Figure 7, the left is the local area of $F = [1, m] \cup [l, k + l - m - 1]$ with $m \ge 2$ and $l \ge m + 3$. By definition, none of the adjacent faces in L(F) will be combined into one face. So we only need to change the label on each face while keep the graph unchanged.



Figure 7: Boundary map on local area of $F = [1, m] \cup [l, k + l - m - 1]$ with $m \ge 2$ and $l \ge m + 3$

Case 4: As in Figure 8, the left is the local area of face $F = [1, m] \cup [m + 2, k + 1]$ with $m \ge 3$. The only edge that need to be deleted is the edge that separates face $F_1 = \{1\} \cup [3, k + 1][2, k + 1]$ and face $F_2 = [2, k + 1]$. So F_1 and F_2 are united into $F'_{1,2}$ on the right.



Figure 8: Boundary map on local area of $F = [1, m] \cup [m + 2, k + l]$ with $m \ge 3$

Given any element $F' \in \partial_n(W_0)$, we observed that the local area of F' in $\Sigma_0(\partial_n(W_0))$ is exactly presented on the right of some case above. Since a plabic graph is a planar graph, the connection of these local areas must be unique. So $\partial_n(\Sigma_0(W_0)) \cong \Sigma_0(\partial_n(W_0))$.

Remark 3.8. We can extend the operations in Definition 3.6 to the equivalence class $\Sigma(W)$. Since every $G \in \Sigma(W)$ is obtained from $\Sigma_0(W)$ by a series of moves (M2), a black (white) point in $\Sigma_0(W)$ may be extended to a tree with black (white) vertives. We can assign the same label on every point of this tree as the point in $\Sigma_0(W)$. So we can still do the operations in Definition 3.6. It is illustrated in the following figure.



Figure 9: Labels of black points in G

For an extension Lemma 3.7 from W_0 to any maximal w-collection in $\binom{[n]}{k}$, we need that any two maximal w-collection can be joined by a sequence of flips (cluster mutations). Recall (Leclecr-Zelevinsky) that if a w-collection collection W contain five sets *Lab*, *Lcd*, *Lac*, *Lad*, *Lbc*, then the collection $W' = W \setminus \{Lac\} \cup \{Lbd\}$ is weakly separated. The transformation $W \to W'$ is called a **flip**.

Theorem 3.9. ([12],[13]) Any two maximal w-collection in $\binom{[n]}{k}$ can be joined by a sequence of flips.

Lemma 3.10. For any $G \in \Sigma(W)$, the following forbidden structure with $\mathcal{B}\backslash W = M_2^n(\mathcal{B})$ will not appear in G (*a*, *b*, *c* are cyclic ordered in $[n]\backslash L$).



Figure 10: A forbidden structure in plabic graph

Proof. Suppose that this structure appear in *G*, since $\mathcal{B}\setminus \mathcal{W} = \{Labc\}\setminus \{Lb\} = \{a, c\} = M_2^n(\mathcal{B})$, we have either $(a, c) \cap Lb = \emptyset$ or $(c, a) \cap Lb = \emptyset$. But *a*, *b*, *c* are cyclic ordered, so $(a, c) \cap Lb \neq \emptyset$. However, since every internal point in *G* has degree no less than three, there exists a face on the left of \mathcal{B} such that $F = \{Labc\}\setminus \{x\}$ where $x \in (c, a)$. There is a contradiction to the fact $(c, a) \cap Lb = \emptyset$.

Lemma 3.11. Let W be any maximal w-collection, then $\partial_n(\Sigma_0(W)) \cong \Sigma_0(\partial_n(W))$. Moreover, the induced subgraph (1-dimensional subcomplex) of points $\partial_n^{-1}|_W(\partial_n(J)) := \{I \in W \mid \partial_n(I) = \partial_n(J), J \in W\}$ is connected in the plabic tiling of W.

Proof. We start with $W_0 = I_n^k \cup S_n^k$, we have known that $\partial_n(\Sigma_0(W_0)) \cong \Sigma_0(\partial_n(W_0))$. We will show that this relation holds under flips. Let W be a maximal w-collection satisfying $\partial_n(\Sigma_0(W)) \cong \Sigma_0(\partial_n(W))$ and $W' = (W \setminus \{Lac\}) \cup \{Lbd\}$. We noticed that, for any fixed black point \mathcal{B} in $G \in \Sigma(W)$, there are at most one edge adjacednt to \mathcal{B} can be deleted under the boundary map ∂_n . Thus we only need to consider two cases up to symmetry. These two cases depend on how many edges are deleted in the local area formed by Lab, Lbc, Lcd, Lad, Lac(Lbd).

Case 1: As in figure 11, only one edge is deleted in step 2 in the local area formed by Lab, Lbc, Lcd, Lad, Lac. In this case, $M_2^n(Lacd) = \{a, d\}$ and $(d, a) \cap Lc = \emptyset$. So the edge that separates Lcd and Lac is deleted in step 2. The rest edges stay unchanged which implies that $M_2^n(Labc) \notin \{\{a, b\}, \{b, c\}, \{c, a\}\}$. Then we consider the local area formed by Lab, Lbc, Lcd, Lad, Lbd after a flip. From the assumption $M_2^n(Lacd) = \{a, d\}$ and the cyclic order of a, b, c, d, we obtain $M_2^n(Labd) = \{a, d\}$. Thus the edge that separates Lbd and Lab is deleted under ∂_n . The converse direction, that is considering the inverse flip $W' \longrightarrow W$ is similar.



Figure 11: Case 1

Case 2: As in figure 12, two edges are deleted in step 2 in the local area formed by Lab, Lbc, Lcd, Lad, Lac. This is equivalent to $M_2^n(Lacd) = \{a, d\}$ with $(d, a) \cap Lc = \emptyset$ and $M_2^n(Labc) = \{a, b\}$ with $(a, b) \cap Lc = \emptyset$. By Lemma 3.10, the upper right edge outside of the square region must connect to a white point or boundary vertex. So if we consider the boundary map after flip, that is on the local area formed by Lab, Lbc, Lcd, Lad, Lbd, then edges that separate faces Lbc, Lcd and faces Lac, Lab will be deleted. Conversely, it is also true as case 1.



Figure 12: Case 2

Case 3: No edge is deleted in step 2. So the flip $W \longrightarrow W'$ induces a flip $\partial_n(W) \longrightarrow \partial_n(W')$ as in figure 13.



Therefore, it is showed in the case 1 or 2 that when $\partial_n(W) = \partial_n(W')$, we have $\partial_n(\Sigma_0(W)) \cong \partial_n(\Sigma_0(W'))$. So by the assumption $\partial_n(\Sigma_0(W)) \cong \Sigma_0(\partial_n(W))$, we get $\partial_n(\Sigma_0(W')) \cong \Sigma_0(\partial_n(W'))$. This isomorphism is obvious in case 3. Besides, when $|\partial_n^{-1}|_W(\partial_n(J))| \ge 2$, we notice that $\partial_n^{-1}|_W(\partial_n(J))$ form a series of consecutive vertices of a white polygon labeled by $\partial_n(J)$ in the plabic tiling of W. So the induced subgraph is connected. \Box

3.3 Proof of the second theorem and a corollary

Proof of Theorem 3.1.

 $\partial_n(W_0)$ is a maximal w-collection as $W_0 = \mathcal{I}_n^k \cup \mathcal{S}_n^k$ by Lemma 3.7. Suppose that W is obtained from W_0 by a sequence of flips, then by Lemma 3.11, either $\Sigma_0(\partial_n(W_0)) \cong \Sigma_0(\partial_n(W))$ or $\Sigma_0(\partial_n(W))$ is obtained from $\Sigma_0(\partial_n(W_0))$ by a sequence of flips. Since $\Sigma_0(\partial_n(W_0))$ is reduced, then $\Sigma_0(\partial_n(W))$ is also reduced so $\partial_n(W)$ is a maximal w-collection.

Example 3.12. Take a maximal w-collection $W = \{127, 137, 136, 156, 167, 135, 145, 134, 123, 234, 345, 456, 567, 678, 178, 128\}$ in $\binom{[8]}{3}$, then by Definition 3.6, $\partial_8(\Sigma_0(W))$ is produced as showed in the following figure.



Figure 14: An example of producing reduced plabic graph $\partial_8(\Sigma_0(W))$

Since $\partial_8(W) = \{27, 37, 36, 35, 12, 23, 34, 45, 56, 67, 17\}$ is the collection of face labels on the reduced plabic graph $\partial_8(\Sigma_0(W))$, it is a maximal w-collection in $\binom{[7]}{2}$

Corollary 3.13. The translated blades $\{\beta_J | J \in W\}$ of a maximal w-collection W induce a finest regular positroid subdivison of $\Delta_{k,n}$.

Proof. Choose any $L \subset [n]$ with |L| = k - 2, then we use Theorem 3.1 repeatedly. Thus, $\partial_L(W)$ is a maximal w-collection in $\binom{[n]\setminus L}{2}$. By the same arguments as in the proof of Theorem 2.32, we obtain that every octahedral face of $\Delta_{k,n}$ is subdivided. So $\{\beta_J | J \in W\}$ induce a finest positroid subdivision.

Following Example 2.34, we give an nontrivial example for Corollary 3.13.

Example 3.14. Hypersimplex $\Delta_{3,6}$ is divided into six top dimensional positroid polytopes by blade arrangements $\beta_{135}, \beta_{235}, \beta_{145}, \beta_{136}$ as showed in Figure 16. For $i \in \{1, 2, 3, 4, 5, 6\}$, let \mathcal{M}_i be the series-parallel matroid (see [18]) obtained from the spanning tree of a series-parallel graph with labeled edges and let $P_{\mathcal{M}_i}$ be its positroid polytope. Since every series-parallel matroid does not contain an octahedral face ([18]) and the labeled graph does not contain the following forbidden substructure (Figure 15),



Figure 15: The forbidden structures in the labeled series-parallel graph with a < b < c < d

Then this implies that pyramids $\{Lab, Lbc, Lcd, Lda, Lac\}$ and $\{Lab, Lbc, Lcd, Lda, Lbd\}$ which do not contain non-separated diagonal $\{Lac, Lbd\}$ are not allowed in M_i . Thus this is a finest positroid subdivision.





 $P_{\mathcal{M}_1} = \{ x_{16} \ge 1 \ x_{1236} \ge 2, \ x_{1456} \ge 2 \} \cap \Delta_{3,6}$



 $P_{\mathcal{M}_3} = \{ x_{16} \ge 1 \ x_{1236} \le 2, \ x_{1456} \ge 2 \} \cap \Delta_{3,6}$



 $P_{\mathcal{M}_4} = \{ x_{16} \le 1 \ x_{1236} \ge 2, \ x_{1456} \le 2 \} \cap \Delta_{3,6}$

 $P_{\mathcal{M}_2} = \{ x_{16} \ge 1 \ x_{1236} \ge 2, \ x_{1456} \le 2 \} \cap \Delta_{3,6}$



 $P_{\mathcal{M}_5} = \{x_{16} \le 1 \ x_{1236} \le 2, \ x_{1456} \ge 2\} \cap \Delta_{3,6}$



 $P_{\mathcal{M}_6} = \{x_{16} \le 1 \ x_{1236} \le 2, \ x_{1456} \le 2\} \cap \Delta_{3,6}$

Figure 16: Finest positroid subdivision of $\Delta_{3,6}$ induced by $\beta_{135}, \beta_{235}, \beta_{145}, \beta_{136}$

Remark 3.15. From Corollary 3.13, we know that there exists unique maximal cone \mathcal{M}_W in $\overline{\mathcal{Z}_{k,n}}$ such that $\mathcal{P}_W \subseteq \mathcal{M}_W$ for every nonfrozen w-collection W. And in Theorem 2.32, we discussed when $\mathcal{P}_W = \mathcal{M}_W$. If two maximal w-collection W_1 and W_2 are connected by a flip i.e. $W_2 = W_1 \setminus \{Lac\} \cup \{Lbd\}$, then \mathcal{M}_{W_1} and \mathcal{M}_{W_2} are two adjacent maximal cones in $\overline{\mathcal{Z}_{k,n}}$ and $\mathcal{P}_{W_1} \cap \mathcal{P}_{W_2} \subseteq \mathcal{M}_{W_1} \cap \mathcal{M}_{W_2}$. We use the Figure 17 to present the flip between these objects.



Figure 17: The flip of a positroid subdivision induced by translated blades

4 The Conclusions and the Further Work

In this paper, we have shown another hypostasis of maximal weakly separated collection of \mathcal{W}_n^k , they label finest positroids subdivisions of $\Delta_{k,n}$, or the cones in $\overline{\mathcal{Z}_{k,n}}$.

Futhermore, each $W \in W_n^k$ is uniquely lifted to a maximal weakly separated collection in $2^{[n]}$ by adding interval sets of size > k and cointerval sets of size < k, and is a basis of TP- functions on $2^{[n]}$ satisfying, for any X and $\{i < j < k < l\} \cap X = \emptyset$,

$$F(Xj) + F(Xik) = \min(F(Xi) + F(Xjk), F(Xij) + F(Xk),$$

and

$$F(Xik) + F(Xjk) = \min(F(Xij) + F(Xkl), F(Xil) + F(Xjk).$$

The supermodular TP-functions, a subset of which cut out by inequalities $F(Xi) + F(Xj) \le F(Xij) + F(X)$ form a crystal $B(\infty)$ for SL_n [19]. A supermodular TP-functions is a (sup) support function to a MV polytope [20]. Let us define cones of supermodular TP-functions, two supermodular TP-functions F and G belong to the same cone if F + G is a TP-function (it is supermodular since supermodularity is stable under summation). A subdivision is finest if all octahedrons are subdivided in two halves. Kamnitzer in [20] considers cones in the set of MV-polytopes. MV-polytopes P and Q belong to the same cone if P + Q is an MV-polytope. For example, for SL_4 there are 13 such cones of maximal dimension, 12 of which are simplicial with 6 generators and one is not simplicial with 7 generators (Section 6 in [20]).

Note that each function defined on vertices on the Boolean cube can be extended as a convex or a concave function to the whole cube $[0, 1]^{[n]}$. Thus, we get two dissections of the cube by affinity areas of corresponding extensions. A supermodular TP-function being extended to a convex function on $[0, 1]^{[n]}$ is a (sup) support function to an MV-polytope. In such a case, the cube is dissected by Weyl chambers.

If we consider a concave extension of a submodular functions, we get a dissection of the cube into generalised polymatroids, since any supermodular TP-function is a M^{\ddagger} -function [8]. One can regard such generalised polymatroids as *generalised positroids*.

For any $k = 2, \dots, n-2$, the restriction of a supermodular TP-function F to the vertices of $\Delta_{k,n}$ is a positive tropical Plücker vector p. Hence the intersection of the generalised positroid subdivision obtained from the concave extension of F with the hyperplane $\sum_i x_i = k$ gives a subdivision of $\Delta_{k,n}$ into poistroids for p (finest subdivison if the subdivison for F is finest).

In a subsequent publication we plan to describe cones of finest genralised positroids subdivisions of unit cubes.

On the another hand, we can expand a positive tropical Plücker vector on $\Delta_{k,n}$ to a supermodular TP-function on the Boolean cube $2^{[n]}$ and, hence get (not uniquely) a generalised positroid subdivision of the cube. This extension gives us finest positroid subdivisions of all $\Delta_{k',n}$, for all $k' \neq k$, compatible with \mathcal{D}_p .

Acknowledgments

For this project, FL and LZ are supported by the National Natural Science Foundation of China (No.12071422, No.12131015)

References

- [1] Nick Early. From weakly separated collections to matroid subdivisions. Combinatorial Theory, 2019.
- [2] Alexander Postnikov. Total positivity, grassmannians, and networks, 2006.
- [3] David Speyer and Lauren Williams. The tropical totally positive Grassmannian. J. Algebraic Combin., 22(2):189–210, 2005.
- [4] David Speyer and Lauren K. Williams. The positive Dressian equals the positive tropical Grassmannian. *Trans. Amer. Math. Soc. Ser. B*, 8:330–353, 2021.
- [5] Tomasz Ł ukowski, Matteo Parisi, and Lauren K. Williams. The positive tropical Grassmannian, the hypersimplex, and the *m* = 2 amplituhedron. *Int. Math. Res. Not. IMRN*, (19):16778–16836, 2023.
- [6] Nick Early. Weighted blade arrangements and the positive tropical grassmannian, 2022.
- [7] Nima Arkani-Hamed, Thomas Lam, and Marcus Spradlin. Positive configuration space. *Communications in Mathematical Physics*, 384(2):909–954, April 2021.

- [8] Kazuo Murota. Discrete convex analysis. *Mathematical Programming*, 83:313–371, 1998.
- [9] Andreas W.M Dress and Walter Wenzel. Valuated matroids. Advances in Mathematics, 93(2):214–250, 1992.
- [10] Vladimir I. Danilov, Alexander V. Karzanov, and Gleb A. Koshevoy. Plücker environments, wiring and tiling diagrams, and weakly separated set-systems. *Advances in Mathematics*, 224(1):1–44, 2010.
- [11] Sven Herrmann. On the facets of the secondary polytope. *Journal of Combinatorial Theory, Series A*, 118(2):425–447, February 2011.
- [12] Vladimir I Danilov, Alexander V Karzanov, and Gleb A Koshevoy. On maximal weakly separated set-systems. *Journal of Algebraic Combinatorics*, 32:497–531, 2010.
- [13] Suho Oh, Alexander Postnikov, and David E. Speyer. Weak separation and plabic graphs. *Proceedings of the London Mathematical Society*, 110(3):721–754, February 2015.
- [14] Jorge Alberto Olarte, Marta Panizzut, and Benjamin Schröter. On local dressians of matroids, 2018.
- [15] Nick Early. Honeycomb tessellations and graded permutohedral blades, 2022.
- [16] Sergey Fomin and Andrei Zelevinsky. Y-systems and generalized associahedra. Annals of Mathematics, 158:977– 1018, 2001.
- [17] V. I. Danilov, A. V. Karzanov, and G. A. Koshevoy. Combined tilings and separated set-systems. Selecta Mathematica, 23(2):1175–1203, September 2016.
- [18] Theory of Matroids. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 1986.
- [19] V Danilov, A Karzanov, and G Koshevoy. Tropical plucker functions and kashiwara crystals. *Contemporary Mathematics*, 616:77–100, 2014.
- [20] Joel Kamnitzer. Mirković-vilonen cycles and polytopes. Annals of mathematics, pages 245–294, 2010.

Gleb A. Koshevoy, Institute for Information Transmission Problems, Russian Academy of Science of Moscow, Moscow 127051, Russia; koshevoyga@gmail.com

Fang Li, School of Mathmatical Sciences, Zhejiang University, Hangzhou 310058, China; fangli@zju.edu.cn

Lujun Zhang, School of Mathmatical Sciences, Zhejiang University, Hangzhou 310058, China;

12135007@zju.edu.cn