A NEW FAMILY OF SOFIC ONE-RELATOR GROUPS

FEDERICO BERLAI

ABSTRACT. We provide an infinite family of sofic one-relator groups that are not residually finite nor residually solvable. The proof is essentially different from the one in [1] as it does not use Magnus' decompositions.

Throughout the paper we use the notation $x^y := y^{-1}xy$ and $[x, y] := x^{-1}y^{-1}xy$.

In this short note, we focus on a family of one-relator groups introduced by Baumslag, Miller III and Troeger [4] in 2007. These one-relator groups generalize a classical example of Baumslag [3] from 1969, and they are not residually finite nor residually solvable. To introduce this family of one-relator groups, let $r, w \in F(a, b)$ be two elements that do not commute (more generally, one could consider a free group on more than two generators, but for the sake of simplicity we will focus on the two-generated case here). Define

(1)
$$G_{r,w} := \langle a, b \mid r = [r, r^w] \rangle = \langle a, b \mid r^2 = r^{r^w} \rangle.$$

In [4] it is proved that $G_{r,w}$ has the same finite quotients as the group $\langle a, b | r \rangle$, and that $G_{r,w}$ is not residually finite (because, for instance, the non-trivial element r of $G_{r,w}$ has trivial image in all finite quotients). Such groups are not residually solvable either, as the element r belongs to all derived subgroups of $G_{r,w}$. In the case r = a and w = b one recovers the classical Baumslag group

(2)
$$B := \langle a, b \mid a = [a, a^b] \rangle = \langle a, b \mid a^2 = (b^{-1}ab)^{-1}a(b^{-1}ab)] \rangle.$$

This group was considered by Gilbert Baumslag in 1969 to provide (yet another) example of a non residually finite one-relator group [3]. Bannon [1] (compare also [2]) noticed that, despite Bnot being residually finite nor residually solvable, it is a sofic group. Indeed, looking at Magnus' decomposition for such one-relator group one sees that B is an HNN-extension of a Baumslag-Solitar group BS(1, 2) amalgamating a cyclic subgroup, that is, $B = BS(1, 2)*\mathbb{Z}$. Such HNN extensions are known to be sofic [8, 11] because the base group is sofic and the amalgamated subgroup is amenable. This was the first example of a sofic one-relator group that is not residually finite nor residually solvable, with more examples considered in [2], applying the same strategy.

As noticed in [2], soficity of the groups G_{a,b^n} for n > 1 does not follow from the argument of [1, 2], because their Magnus' decomposition is $(BS(1,2) * F_{n-1}) *_{F_n}$, where (for n > 1) a nonamenable free group appears as amalgamated subgroup. For such HNN extensions there is no general result implying soficity. The definition of sofic group will not play a role here, hence we refer to [6, 12] for excellent introductions to these groups. We will just underline that it is an open problem to see if all groups are sofic [12, Open Question 3.8], or if all one-relator groups are sofic [12, Open Question 4.9].

In what follows, we will mostly focus on the case r = a and plug in exponents $l, k \in \mathbb{Z}$ into the relator in a fashion similar to Baumslag-Solitar groups, so that the relation reads $r^k = (r^l)^{r^w}$, to cover a slighly more general family of groups (one could go even further and consider relators as done in [5], but we will not do this here). Given two elements $r, w \in F(a, b)$ that do not commute and two non-zero integers l and k, we define

(3)
$$G_{r,w}(l,k) := \langle a, b, \dots \mid (r^l)^{r^w} = r^k \rangle.$$

As justified in the following section, many of the one-relator groups from Equation (3) are not residually finite nor residually solvable. Whenever l = 1 and k = 2, we simplify the notation and write directly $G_{r,w}$ instead of $G_{r,w}(1,2)$ (these are exactly the groups recalled in Equation (1)). We prove:

Theorem A. For all non-zero integers $n, l, k \in \mathbb{Z}$, the group $G_{a,b^n}(l,k)$ is sofic.

This settles the open problem raised in [2] (see [2, Theorem 2] and the discussion before that). A careful choice of the integers l and k immediately implies:

Corollary B. For all $n \in \mathbb{Z} \setminus \{0\}$ and $l \in \mathbb{Z} \setminus \{0, -1\}$ the group $G_{a,b^n}(l, l+1)$ is sofic but not residually finite nor residually solvable.

Theorem A can be used iteratively to generate more sofic one-relator groups that are not residually finite nor residually solvable:

Theorem C. For all non-zero integers $n, l, k \in \mathbb{Z}$ the one-relator group $G_{a,b^{-n}ab^n}(l,k)$ is sofic.

As in Corollary B, correctly choosing the integers l and k produces groups that are sofic but not residually finite nor residually solvable.

It is not evident if all groups among the one-relator groups introduced by Baumslag, Miller III and Troeger can be shown to be sofic using Magnus' decompositions and the argument used in this paper (or by other methods). For instance, is the group $G_{a,b^{-1}ab^2}$ sofic?

Acknowledgements. The author is supported by the Spanish Government, grant PID2020-117281GB-I00, partly by the European Regional Development Fund (ERDF), and the Basque Government, grant IT1483-22. He is indebted to Goulnara Arzhantseva for suggesting to work on this project during his PhD studies, back in 2013. He is very thankful to Marco Linton for insightful conversations on this topic and for pointing out a gap in a proof in a previous version of this note, and to an anonymous referee for suggesting a slightly shorter/cleaner proof of Theorem A.

1. Preliminary Lemmas

In this preliminary section we collect some observations that will be useful later, justifying our generalisation of the family of one-relator groups introduced by Baumslag, Miller III and Troeger [4]. In what follows, r and w are elements of the free group F(a, b) that do not commute.

We start by noticing that many of these groups are not residually finite. Indeed, if $l = \pm 1$ and $k = \pm 2$ (or viceversa), then $G_{r,w}(l,k)$ is not residually finite. In particular, if $(l,k) \in \{(1,2), (-1,-2), (2,1), (-2,-1)\}$, then the claim follows directly applying [4, Theorem 1]. If signs are mixed, that is if $(l,k) \in \{(1,-2), (-1,2), (2,-1), (-2,1)\}$, the claim essentially follows from [4, Theorem 1], again, because [4, Lemma 1] holds in this case: that is, if $x, y \in G$ are two elements of the same finite order and satisfy the relation $x^y = x^{-2}$, then the subgroup $\langle x, y \rangle$ of G is trivial. If l and k are different in absolute value and both different from ± 1 , then the group $G_{r,w}(l,k)$ is not residually finite by [10, Theorem B]. These remarks and Lemma 1.1, joint with Theorem A, already imply Corollary B.

The groups $G_{r,w}$ considered in [4] are always non-residually finite, for any pair of non-commuting elements r, w in a free groups. A full characterization of (non-) residual finiteness of the groups $G_{r,w}(l,m)$ is not given here, but it seems plausible to suppose that these groups also are never residually finite.

Many of these groups are also not residually solvable:

Lemma 1.1. If $k \neq 0 \neq l$ and $k = l \pm 1$, then the group $G_{r,w}(l,k)$ is not residually solvable.

Proof. Suppose that k = l+1. The defining relation $r^k = (r^l)^{r^w}$ can be rewritten as $r = r^{-l}(r^l)^{r^w}$, that is $r = [r^l, (r)^w]$ is an element of the derived subgroup. Consequently the elements r^w and r^l also lie in the derived subgroup. Inductively, r is an element of the intersection of all derived subgroups of $G_{r,w}(l,k)$. This element is not trivial by the same argument of [4, Lemma 1]; thus, $G_{r,w}(l,k)$ is not residually solvable.

If k = l - 1, then the same argument shows that r^{-1} is a non-trivial element in the intersection of all derived subgroups of $G_{r,w}(l,k)$.

Even if it will be of no use in this note, we record the following observations:

Lemma 1.2. Let $r, w \in F(a, b)$ be two non-commuting elements and $l, k \in \mathbb{Z}$. Then $G_{r,w}(l, k) \cong G_{r,w}(-l, -k)$ and $G_{r,w}(l, k) \cong G_{r^{-1},w}(k, l)$.

Proof. For the first isomorphism, notice that the relation $(r^l)^{r^w} = r^k$ implies that $((r^l)^{r^w})^{-1} = (r^k)^{-1}$, that is $(r^{-l})^{r^w} = r^{-k}$. Therefore

are group homomorphisms such that $\xi \circ \eta$ and $\eta \circ \xi$ are both identity homomorphisms. Therefore they are isomorphisms.

For the second isomorphism, notice that $(r^l)^{r^w} = r^k$ immediately implies that $r^l = (r^k)^{(r^{-1})^w}$, and the claim follows taking inverses.

Lemma 1.3. Let $w \in F(a,b)$ be a non-trivial element whose normal form starts and ends with non-trivial powers of b. Then $G_{a,a^nw}(l,k) \cong G_{a,w}(l,k) \cong G_{a,wa^n}(l,k)$.

Proof. This is easy to see by explicitly writing the relations of $G_{a,a^nw}(l,k)$ and of $G_{a,wa^n}(l,k)$ and reducing them to the relation of $G_{a,w}(l,k)$.

2. Proof of Theorem A

We may assume without loss of generality that n is positive, because the map defined on the generators as $a \mapsto a$ and $b \mapsto b^{-1}$ is an automorphism. For the case n = 1, note that $G := G_{a,b}(l,k)$ is the HNN extension

$$G_{a,b}(l,k) = \langle a,b \mid (a^b)^{-1}a^l a^b = a^k \rangle = \langle a_0, a_1, b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = \langle a_1,b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = \langle a_1,b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = \langle a_1,b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = \langle a_1,b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = \langle a_1,b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = \langle a_1,b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = \langle a_1,b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = \langle a_1,b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = \langle a_1,b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = \langle a_1,b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = \langle a_1,b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = \langle a_1,b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = \langle a_1,b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = \langle a_1,b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = \langle a_1,b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = \langle a_1,b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = \langle a_1,b \mid a_1^{-1}a_0^l a_1 = a_0^k, b^{-1}a_0b = a_1 \rangle = BS(l,k) *_{\mathbb{Z}} A_{a_1}(l,k) = A(l,k) = A(l,k)$$

The Baumslag-Solitar group BS(l, k) is residually solvable [9], hence sofic. Thus, $G_{a,b}(l, k)$ is sofic [8, 11]. Consider the the relator of the group of Equation (3). As the generator b has exponent-sum equal to zero in this relator, the group G admits a surjective homomorphism onto $\mathbb{Z} = \langle z \rangle$ defined by the assignments $a \mapsto e$ and $b \mapsto z$. The kernel K of this homomorphism is the normal closure in G of the element a, that is

$$(4) \quad K = \langle \langle a \rangle \rangle_G = \langle \{a_i\}_{i \in \mathbb{Z}} \rangle = \langle \{a_i\}_{i \in \mathbb{Z}} \mid \{a_{i+1}^{-1}a_i^l a_{i+1} = a_i^k\}_{i \in \mathbb{Z}} \rangle, \qquad a_i := b^{-i}ab^i \quad \forall i \in \mathbb{Z}.$$

This kernel is sofic, being a subgroup of the sofic group $G_{a,b}(l,k)$.

For the general case, consider $n \ge 2$ and the group $G_{a,b^n}(l,k)$. Also in this case the exponentsum of the generator b in the relator is equal to zero. Therefore $G_{a,b^n}(l,k)$ admits a surjective homomorphism onto \mathbb{Z} with kernel the normal closure of a in $G_{a,b^n}(l,k)$:

$$K_n := \langle \langle a \rangle \rangle = \langle \{a_i\}_{i \in \mathbb{Z}} \mid \{a_{i+n}^{-1} a_i^l a_{i+n} = a_i^k\}_{i \in \mathbb{Z}} \rangle.$$

Grouping the generators modulo n, we see that K_n is the free product of n copies of the group K appearing in Equation (4). Therefore K_n is sofic as well [7, Theorem 1], and thus $G_{a,b^n}(l,k)$ too, being sofic-by-amenable [6, Proposition 7.5.14]. Thus, the proof of Theorem A is completed.

3. Proof of Theorem C

We now deduce Theorem C building on the result and arguments of the previous section. Let

$$G := G_{a,b^{-n}ab^n}(l,k) = \langle a,b \mid \left((b^{-n}ab^n)^{-1}a(b^{-n}ab^n) \right)^{-1}a^l \left((b^{-n}ab^n)^{-1}a(b^{-n}ab^n) \right) = a^k \rangle.$$

As the generator b has exponent-sum equal to zero in the relator, G admits a surjective homomorphism onto $\mathbb{Z} = \langle z \rangle$ defined on the generators by $a \mapsto e$ and $b \mapsto z$. The kernel of this homomorphism is the normal closure of a in G: it is generated by the conjugates $a_i := b^{-i}ab^i$, where $i \in \mathbb{Z}$, subject to the relations $(a_{n+i}^{-1}a_ia_{n+i})^{-1}a_i^l(a_{n+i}^{-1}a_ia_{n+i}) = a_i^k$. Regrouping the generators modulo n, this kernel is isomorphic to the free product of n copies of

$$H := \langle \{g_i\}_{i \in \mathbb{Z}} \mid \{(g_{i+1}^{-1}g_ig_{i+1})^{-1}g_i^l(g_{i+1}^{-1}g_ig_{i+1}) = g_i^k\}_{i \in \mathbb{Z}} \rangle$$

= $\dots \underset{g_0 \mapsto g_0}{*} \langle g_0, g_1 \mid (g_0^l)^{g_0^{g_1}} = g_0^k \rangle \underset{g_1 \mapsto g_1}{*} \langle g_1, g_2 \mid (g_1^l)^{g_1^{g_2}} = g_1^k \rangle \underset{g_2 \mapsto g_2}{*} \dots$

The group G is locally sofic, and hence sofic [6, Proposition 7.5.5], being an iterated amalgamated product of $G_{a,b^n}(l,k)$ (which is sofic by Theorem A) amalgamated along cyclic (hence amenable) subgroups [8, 11]. Thus $K = H * \ldots * H$ is sofic as well [7, Theorem 1], and G too, being an extension of the sofic group K by the amenable quotient \mathbb{Z} .

FEDERICO BERLAI

References

- [1] J. P. Bannon, A non-residually solvable hyperlinear one-relator group. Proc. Amer. Math. Soc. 139 (2011), no. 4, 1409 1410;
- [2] J. P. Bannon, N. Noblett, A note on nonresidually solvable hyperlinear one-relator groups. Involve 3 (2010), no. 3, 345 - 347;
- [3] G. Baumslag, A non-cyclic one-relator group all of whose finite quotients are cyclic. J. Austral. Math. Soc. 10 (1969), 497 - 498;
- [4] G. Baumslag, C. F. Miller III, D. Troeger, Reflections on the residual finiteness of one-relator groups. Groups Geom. Dyn. 1 (2007), no. 3, 209 - 219;
- [5] A. Brunner, On a class of one-relator groups. Canadian J. Math. 32 (1980), no. 2, 414 420;
- [6] T. Ceccherini-Silberstein, M. Coornaert, Cellular automata and groups. Second edition. Springer Monographs in Mathematics. Springer, Cham, (2023);
- [7] G. Elek, E. Szabó, On sofic groups. J. Group Theory 9, (2006), 161 171;
- [8] G. Elek, E. Szabó, Sofic representations of amenable groups. Proc. Amer. Math. Soc. 139 (2011), no. 12, 4285 - 4291;
- [9] P. Kropholler, Baumslag-Solitar groups and some other groups of cohomological dimension two. Comment. Math. Helvetici 65 (1990), 547 - 558;
- [10] S. Meskin, Nonresidually finite one-relator groups. Trans. Amer. Math. Soc. 164 (1972), 105 114;
- [11] L. Păunescu, On sofic actions and equivalence relations. J. Funct. Anal. 261 (2011), no. 9, 2461 2485;
- [12] V. Pestov, Hyperlinear and sofic groups: a brief guide. Bull. Symbolic Logic 14 (2008), no. 4, 449 480.

(Federico Berlai) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF THE BASQUE COUNTRY UPV/EHU, BARRIO SARRIENA S/N, 48940 LEIOA, SPAIN

Email address, Federico Berlai: federico.berlai@ehu.eus