NOISE SENSITIVITY OF HIERARCHICAL FUNCTIONS AND DEEP LEARNING LOWER BOUNDS IN GENERAL PRODUCT MEASURES

RUPERT LI AND ELCHANAN MOSSEL

ABSTRACT. Recent works explore deep learning's success by examining functions or data with hierarchical structure. Complementarily, research on gradient descent performance for deep nets has shown that noise sensitivity of functions under independent and identically distributed (i.i.d.) Bernoulli inputs establishes learning complexity bounds. This paper aims to bridge these research streams by demonstrating that functions constructed through repeated composition of non-linear functions are noise sensitive under general product measures.

1. INTRODUCTION

Our goal is to bridge two lines of research in deep learning, exploring the role of hierarchical structure in understanding neural network success. The first line investigates how hierarchical properties in data generative processes or learned functions contribute to deep learning effectiveness. This hypothesis has been independently proposed by multiple researchers, including:

- Bruna and Mallat's wavelet scattering networks [6],
- Mhaskar, Liao, and Poggio's compositional function models [13],
- Patel, Nguyen, and Baraniuk's hierarchical rendering models [18].

Mossel [16] provided a rigorous attempt to establish computational hardness, though subsequent work by Koehler and Mossel [12] only demonstrated low-degree hardness. In terms of compositions of functions, Telgarsky [21] presents a notable result demonstrating an iteratively composed simple function that can be precisely expressed as a deep narrow network, yet requires super-polynomial computational complexity for small network depth.

The second research line examines stochastic gradient descent performance on fully connected deep networks with inputs drawn from the uniform distribution on the discrete cube and characterizes the learning complexity through Fourier expansion. One result in this line of work, Abbe, Bengio, Cornacchia, Kleinberg, Lotfi, Raghu, and Zhang [2], demonstrates that the noise-sensitivity of a function implies a lower bound on the number of iterations of gradient descent. These results build on fundamental results in [3, 1] that reveal that the structure of the "hierarchy" in non-vanishing Fourier coefficients is a critical factor determining gradient descent convergence speed. Note that this hierarchical structure in the Fourier domain differs from the earlier notion of function composition that is the key in the current paper.

In our main result we show that hierarchical functions, i.e., functions obtained by decompositions of non-linear functions on disjoint independent inputs, are noise sensitive. This implies a learning lower bound under gradient descent. It also immediately implies lowdegree lower bounds on such functions, see, e.g., [20] for background on low degree hardness.

RUPERT LI AND ELCHANAN MOSSEL

Our work builds upon a long line of work in discrete Fourier analysis. The fact that recursively composing the majority function on 3 bits leads to a noise sensitive function is folklore in the area, see e.g., [4, 5]. The results of [14] further study decompositions of monotone functions and construct such decompositions that are essentially "most" noise sensitive among monotone functions. We also note that recursive constructions, i.e., Sipser functions, provide the best results in circuit lower bounds for circuits with AND, OR and NOT gates [10].

The main contribution of the current paper is by showing that noise sensitivity of composition of non-linear functions on non-overlapping inputs is generic—the functions to be composed do not have to be identical, to be chosen carefully, or to be balanced or have any other property other than being somewhat uncorrelated with linear functions of their inputs. Our results hold for all product measures, both discrete and continuous.

1.1. Our Results.

1.1.1. Multilinear functions. To begin with we consider the case where all functions involved are multilinear. This includes in particular the case where all functions have binary inputs and outputs which is the classical case where analysis of Boolean functions and noise sensitivity are discussed. In this case we obtain the following result concerning a single multilinear function. Precise definitions are provided in Section 2, though we comment that d(f, Lin) quantifies the non-linearity of f.

Lemma 1.1. Suppose $\{(X_i, Y_i) : i \in [n]\}$ are mutually independent, where for some $0 \le \rho \le 1$, we have $\operatorname{Corr}(X_i, Y_i) \le \rho$ for all *i*. Suppose X_i and Y_i have the same first and second moments, *i.e.*, $\mathbb{E}[X_i] = \mathbb{E}[Y_i]$ and $\mathbb{E}[X_i^2] = \mathbb{E}[Y_i^2] < \infty$, for all $i \in [n]$. If $f : \mathbb{R}^n \to \mathbb{R}$ is a multilinear function satisfying $d(f, \operatorname{Lin}) \ge \varepsilon > 0$, then

(1.1)
$$\operatorname{Corr}(f(X), f(Y)) \le (1 - \varepsilon)\rho + \varepsilon \rho^2.$$

Remark 1.2. Such a Y can always be created by setting $Y_i = X_i$ with probability ρ and otherwise letting Y_i be independent and identically distributed as X_i , independently for all *i*; note that this is not in general the unique way to construct such a Y. Moreover, this construction yields X and Y that have the same marginal distribution, which does not need to be true in general for Lemma 1.1 to hold.

Applying Lemma 1.1 inductively yields the following result demonstrating exponential decay in noise stability for a hierarchical multilinear function, precisely defined in Section 2.3.

Theorem 1.3. Suppose $\{(X_i, Y_i) : i \in [n]\}$ are mutually independent, where for some $0 \leq \rho < 1$, we have $\operatorname{Corr}(X_i, Y_i) \leq \rho$ for all *i*. Suppose X_i and Y_i have the same first and second moments, *i.e.*, $\mathbb{E}[X_i] = \mathbb{E}[Y_i]$ and $\mathbb{E}[X_i^2] = \mathbb{E}[Y_i^2] < \infty$, for all $i \in [n]$. If $f : \mathbb{R}^n \to \mathbb{R}$ is a hierarchical multilinear function of non-linearity $\varepsilon > 0$ and depth *d*, then for any $\delta < \varepsilon$,

$$\operatorname{Corr}(f(X), f(Y)) \le C_{\rho,\varepsilon,\delta} (1-\delta)^d$$

for some constant $C_{\rho,\varepsilon,\delta} > 0$ depending only on ρ , ε , and δ .

As discussed in Remark 2.2, this bound is essentially tight, as it is impossible to have exponential rate smaller than $1-\varepsilon$. Remark 2.3 discusses how this can be improved to doubly exponential decay in depth in the case where each component function has 0 correlation with

 $\mathbf{2}$

linear functions, such as *t*-resilient functions from cryptography, which are further discussed in Remark 2.4, and the parity bit function as further discussed in Remark 2.5.

The result above immediately implies a low-degree bound on such a function—in particular we get that the correlation M of such a function f with a degree-D polynomial is bounded by

(1.2)
$$M \le C_{\rho,\varepsilon,\delta} \rho^{-D/2} (1-\delta)^{d/2};$$

see Lemma 2.6 below. In other words, we get a degree bound that is proportional to the circuit depth, i.e., in order to get $M = \Omega(1)$ correlation, one requires $D = \Omega(d)$. In the special case where $\varepsilon = 1$, we obtain a degree bound that is exponential in the depth, i.e., $M = \Omega(1)$ requires $D = \Omega(2^d)$; see Remark 2.3.

Similarly, the results of [2] imply that for functions as above, in their setting of fully connected neural networks and their initialization, the complexity of learning the function using their version of gradient descent is proportional exponential in the depth of the circuit. In the special case where all function are resilient, i.e., have 0 correlation with linear function, the lower bound becomes doubly exponential in the depth, i.e., exponential in the input size.

1.1.2. *General functions.* The results above leave something to be desired, as neural network activation functions, notably ReLU, are typically not multilinear in their inputs. However, some care is needed in this general case as the following examples show.

Example 1.4. Consider a hierarchical function with a binary tree structure, where at even levels we have component functions $f(x_1, x_2) = \cos(\pi x_1)$ and at odd levels $g(x_1, x_2) = \frac{\arccos(x_1)}{\pi}$, where each of the component variables takes values in [0, 1]. For example, such a function of depth 2 would be given by $g(f(x_1, x_2), f(x_3, x_4))$. We note that f and g are clearly non-linear, but any such function of even depth will output x_1 , which is as noise stable as possible, and there is no decay in correlation as depth increases.

Example 1.5. The phenomenon above does not require the functions to have continuous inputs. In fact, it suffices for the inputs and outputs of the functions to take on a small number of values, e.g., four. For example, we can consider the input distributions to be uniform over $\{\pm 1\}^{3^d}$, i.e., 3^d i.i.d. variables uniformly distributed on $\{-1, 1\}$. In our correlated setup for noise sensitivity, we consider two such random vectors $X = (X_1, \ldots, X_n)$ and $Y = (Y_1, \ldots, Y_n)$ coupled such that $\{(X_i, Y_i)\}_{i=1}^n$ are mutually independent, and $\operatorname{Corr}(X_i, Y_i) = \rho$ for all i.

In our construction, our functions can output at most four values, $\pm 10 \pm 1$, and our overall hierarchical function has a ternary tree structure. For a number $x = 10b_1 + b_2$ for $b_1, b_2 \in \{-1, 1\}$, let us write $B_1(x) = b_1$ and $B_2(x) = b_2$. We can then take the functions at the lowest layer (depth d) to be

$$f(x_1, x_2, x_3) = x_1 + 10 \operatorname{maj}(x_1, x_2, x_3),$$

where $\operatorname{maj}(x_1, x_2, x_3) = 2 \cdot 1_{\{x_1+x_2+x_3>0\}} - 1$ is the majority function, i.e., takes value 1 if a majority of inputs take value 1, and takes value -1 if a majority of inputs takes value -1. We let the remaining functions be

$$g(x_1, x_2, x_3) = B_2(x_1) + 10 \operatorname{maj}(B_1(x_1), B_1(x_2), B_1(x_3)),$$

RUPERT LI AND ELCHANAN MOSSEL

It is easy to see that all functions involved are far from linear, but the hierarchical function on 3^d inputs computes the first input plus 10 times the recursive majority of all 3^d inputs, and is therefore very noise stable as it has constant correlation with x_1 .

To account for potential non-linear relationships, we strengthen our notions of correlation in both our assumptions and our conclusions. The precise definitions are provided in Section 3, but loosely speaking, $\operatorname{Corr}(L^2(X), L^2(Y))$ denotes the maximum correlation between any two functions where one is a function of X and the other is a function of Y. The analog of Lemma 1.1 in this more general setting is the following lemma.

Lemma 1.6. Suppose $\{(X_i, Y_i) : i \in [n]\}$ are mutually independent, where for some $0 \le \rho \le 1$, we have $\operatorname{Corr}(L^2(X_i), L^2(Y_i)) \le \rho$ for all i. If $f : \mathbb{R}^n \to \mathbb{R}$ satisfies $\mathbb{E}[f(X)^2], \mathbb{E}[f(Y)^2] < \infty$ and

 $\operatorname{Corr}(L^{2}(f(X)), L^{2}(X_{1}) + \dots + L^{2}(X_{n})), \operatorname{Corr}(L^{2}(f(Y)), L^{2}(Y_{1}) + \dots + L^{2}(Y_{n})) \leq \sqrt{1 - \varepsilon}$

for some $\varepsilon > 0$, then

$$\operatorname{Corr}(L^2(f(X)), L^2(f(Y))) \le (1 - \varepsilon)\rho + \varepsilon \rho^2.$$

In Section 3.3, we justify why (1.3) is the natural analog of the condition $d(f, \text{Lin}) \ge \varepsilon$ in Lemma 2.6.

It is straightforward to check that the construction in Remark 1.2 still satisfies the stronger condition $\operatorname{Corr}(L^2(X_i), L^2(Y_i)) = \rho$: letting Z_i be an independent copy of X_i such that $Y_i = X_i$ with probability ρ and $Y_i = Z_i$ otherwise, we have, for $f \in L^2(X_i)$ and $g \in L^2(Y_i)$ normalized so that $\mathbb{E}[f] = \mathbb{E}[g] = 0$ and $\mathbb{E}[f^2] = \mathbb{E}[g^2] = 1$, by the Cauchy–Schwarz inequality

$$\operatorname{Corr}(f(X_i), g(Y_i)) = \rho \mathbb{E}[f(X_i)g(X_i)] + (1-\rho)\mathbb{E}[f(X_i)g(Z_i)] \le \rho \cdot 1 + (1-\rho) \cdot 0 = \rho,$$

where equality is achieved when f = g as functions $\mathbb{R} \to \mathbb{R}$.

Using Lemma 1.6, we obtain the following general version of Theorem 1.3.

Theorem 1.7. Suppose $\{(X_i, Y_i) : i \in [n]\}$ are mutually independent, where for some $0 \leq \rho < 1$, we have $\operatorname{Corr}(L^2(X_i), L^2(Y_i)) \leq \rho$ for all i. If $f : \mathbb{R}^n \to \mathbb{R}$ is a hierarchical function of depth d and non-separability $\varepsilon > 0$ in both $L^2(X)$ and $L^2(Y)$, then for any $\delta < \varepsilon$,

$$\operatorname{Corr}(L^2(f(X)), L^2(f(Y))) \le C_{\rho,\varepsilon,\delta}(1-\delta)^d$$

for some constant $C_{\rho,\varepsilon,\delta} > 0$ depending only on ρ , ε , and δ .

The result above immediately implies a low-degree bound on such a function, similar to (1.2). In particular, the correlation M of such a function f with a function of Efron–Stein degree at most D (see Section 3.2 for precise definitions), i.e., a sum of functions that each depend on at most D of the n variables, is bounded by

(1.4)
$$M \le C_{\rho,\varepsilon,\delta} \rho^{-D/2} (1-\delta)^{d/2};$$

see Lemma 3.4. In particular, for fixed D this bound decays exponentially in depth d. In particular, by picking suitable $\rho < 1$, we find that f has o(1) correlation with any function expressed as a linear combination of functions that each depend on at most O(d) of the variables. And in the special case where $\varepsilon = 1$, we obtain a doubly exponential bound for M, which implies a degree bound that is exponential in depth, rather than proportional;

see Remark 3.3. In this case, the correlation is o(1) for any function expressed as a linear combination of functions each depending on $o(2^d)$ variables.

We conjecture that the lower bounds of [2] should extend in the appropriate setup and leave this for future work.

1.2. Section Overview. In Section 2, we introduce the necessary definitions, and then prove Lemma 1.1, Theorem 1.3, and Lemma 2.6. Then in Section 3, we first introduce the necessary formalization of correlated probability spaces in Section 3.1, then introduce the Efron–Stein decomposition in Section 3.2 and its fundamental properties, especially in relation to Markov operators; finally, properly equipped, we prove Lemma 1.6 and Theorem 1.7.

2. Noise stability of compositions of multilinear functions

2.1. **Preliminary setup.** For any probability space (Ω, \mathbb{P}) , function $f \in L^2(\Omega, \mathbb{P})$, and subset $V \subseteq L^2(\Omega, \mathbb{P})$, define the (squared) distance between f and V by

$$d(f, V) = \inf_{g \in V} \frac{\mathbb{E}[(f-g)^2]}{\operatorname{Var}(f)}.$$

If f is almost surely constant, i.e., $\operatorname{Var}(f) = 0$, we define d(f, V) = 0. Similarly, throughout this paper we use the convention $\operatorname{Corr}(X, Y) = 0$ if either X or Y are almost surely constant. We now specialize to the case where \mathbb{P} is a product measure on product space Ω over laws of real-valued random variables X_1, X_2, \ldots, X_n . In the product measure, these random variables are mutually independent, and we additionally assume the random variables have finite second moment, i.e., $\mathbb{E}[X_i^2] < \infty$ for all $i \in [n]$. Beyond this, we make no further assumptions on the distributions of the random variables, e.g., some may be discrete, others continuous, or neither. We will consider the setting where $f \in L^2(\Omega, \mathbb{P})$ is multilinear, i.e., f is of the form

$$f(x_1,\ldots,x_n) = \sum_{S \subseteq [n]} a_S \prod_{i \in S} x_i$$

for some coefficients $a_S \in \mathbb{R}$. Note that any function of the above form is in $L^2(\Omega, \mathbb{P})$. In the Boolean case, i.e., when X_1, \ldots, X_n are each random variables supported on at most two points, note that all $f \in L^2(\Omega, \mathbb{P})$ are multilinear in X_1, \ldots, X_n .

Let Lin denote the subspace of $L^2(\Omega, \mathbb{P})$ spanned by linear functions, i.e.,

$$Lin = span(\{1, x_1, x_2, \dots, x_n\}).$$

For each $i \in [n]$, define the normalized random variable $\widetilde{X}_i = \frac{X_i - \mathbb{E}[X_i]}{\sqrt{\operatorname{Var}(X_i)}}$, which has mean 0 and variance 1, and similarly denote $\widetilde{x}_i = \frac{x_i - \mathbb{E}[X_i]}{\sqrt{\operatorname{Var}(X_i)}}$ for all $x_i \in \mathbb{R}$. Then for any multilinear f, we let $\widehat{f}(S) \in \mathbb{R}$ for $S \subseteq [n]$ denote the unique coefficients satisfying

$$f(x_1,\ldots,x_n) = \sum_{S \subseteq [n]} \widehat{f}(S) \prod_{i \in S} \widetilde{x}_i.$$

For each $S \subseteq [n]$, we let $\chi_S : \mathbb{R}^n \to \mathbb{R}$ denote the function given by

$$\chi_S(x_1,\ldots,x_n)=\prod_{i\in S}\widetilde{x}_i.$$

By independence of X_1, \ldots, X_n , we have $\{\chi_S : S \subseteq [n]\}$ is a set of orthonormal multilinear polynomials: for distinct $S, S' \subseteq [n]$, we have $\mathbb{E}[\chi_S \chi_{S'}] = 0$ and $\mathbb{E}[\chi_S^2] = 1$. Also note that $\mathbb{E}[\chi_S] = 1_{\{S = \emptyset\}}$. Hence, we have

$$\mathbb{E}[f^2] = \sum_{S \subseteq [n]} \widehat{f}(S)^2 \quad \text{and} \quad \operatorname{Var}(f) = \sum_{S \neq \emptyset} \widehat{f}(S)^2.$$

From this, it is easy to see that

$$d(f, \operatorname{Lin}) = \frac{\sum_{|S| \ge 2} \widehat{f}(S)^2}{\sum_{S \ne \emptyset} \widehat{f}(S)^2}.$$

For any $S \subseteq [n]$, we let X_S denote the random vector consisting of the X_i for $i \in S$, and similarly define Y_S , as well as x_S and y_S for $x, y \in \mathbb{R}^n$. For example, $\chi_S(x)$ depends only on x_S .

2.2. Main lemma. Lemma 1.1 shows that if we have some noisy observation Y_i of each X_i , namely $\operatorname{Corr}(X_i, Y_i) \leq \rho$ for all $i \in [n]$ for some constant $\rho < 1$, and f is far from linear, then $\operatorname{Corr}(f(X), f(Y)) < \rho$, with this correlation displaying exponential decay with respect to application of f. We now prove this lemma.

Proof of Lemma 1.1. Note that because X_1, \ldots, X_n are mutually independent, and Y_1, \ldots, Y_n are also mutually independent, we have Var(f(X)) = Var(f(Y)) as

$$\mathbb{E}[\chi_{S}(X)\chi_{S'}(X)] = \mathbb{E}[\chi_{S}(Y)\chi_{S'}(Y)] = 1_{\{S=S'\}}.$$

We similarly also have $\mathbb{E}[f(X)] = \mathbb{E}[f(Y)] = \hat{f}(0)$ as $\mathbb{E}[\chi_S(X)] = \mathbb{E}[\chi_S(Y)] = 1_{\{S=\emptyset\}}$. Thus, without loss of generality, we can rescale and shift f so that $\operatorname{Var}(f) = 1$ and $\mathbb{E}[f] = 0$. Then we expand

$$\operatorname{Corr}(f(X), f(Y)) = \mathbb{E}[f(X)f(Y)] = \sum_{S \neq \emptyset} \sum_{S' \neq \emptyset} \widehat{f}(S)\widehat{f}(S')\mathbb{E}[\chi_S(X)\chi_{S'}(Y)].$$

Note that if $S \neq S'$, we have $\mathbb{E}[\chi_S(X)\chi_{S'}(Y)] = 0$. Then either $S \not\subseteq S'$ or $S' \not\subseteq S$. Say $S \not\subseteq S'$. Then

$$\mathbb{E}[\chi_S(X)\chi_{S'}(Y)] = \mathbb{E}[\mathbb{E}[\chi_S(X)\chi_{S'}(Y)|(X_{S'},Y_{S'})]]$$
$$= \mathbb{E}[\mathbb{E}[\chi_S(X)|(X_{S'},Y_{S'})]\chi_{S'}(Y)]$$
$$= \mathbb{E}[0 \cdot \chi_{S'}(Y)]$$
$$= 0.$$

The case $S' \not\subseteq S$ follows symmetrically. If S = S', then by independence of the (X_i, Y_i) we have

$$\mathbb{E}[\chi_S(X)\chi_S(Y)] = \prod_{i \in S} \mathbb{E}[\widetilde{X}_i \widetilde{Y}_i] = \prod_{i \in S} \operatorname{Corr}(X_i, Y_i) \le \rho^{|S|}.$$

Thus,

$$\operatorname{Corr}(f(X), f(Y)) \le \sum_{S \neq \emptyset} \rho^{|S|} \widehat{f}(S)^2$$

Recalling that

$$\operatorname{Var}(f) = \sum_{S \neq \emptyset} \widehat{f}(S)^2 = 1$$

while

$$d(f, \operatorname{Lin}) = \sum_{|S| \ge 2} \widehat{f}(S)^2 \ge \varepsilon,$$

we obtain the bound

$$\operatorname{Corr}(f(X), f(Y)) \le (1 - \varepsilon)\rho + \varepsilon \rho^2,$$

as desired.

Note that this bound is sharp when the conditions in the statement of the lemma are tight, i.e., $\operatorname{Corr}(X_i, Y_i) = \rho$ for all *i*, and $d(f, \operatorname{Lin}) = \varepsilon$, and *f* is of degree 2.

Remark 2.1. In the Boolean setting, i.e., when the X_i and Y_i are all Boolean random variables, when $\operatorname{Corr}(X_i, Y_i) = \rho$ for all *i*, then the quantity $\operatorname{Corr}(f(X), f(Y))$ that we study in Lemma 1.1 coincides with the well-studied noise stability $\operatorname{Stab}_{\rho}(f)$. We refer readers to [17] for a comprehensive overview of the analysis of Boolean functions. As we wish to state results that apply for general settings beyond just the Boolean setting, we will not use the specific definition of noise stability $\operatorname{Stab}_{\rho}(f)$, though the spirit remains the same.

2.3. Hierarchical multilinear functions. We now consider functions consisting of a composition of multilinear functions in a hierarchical structure. Specifically, for any positive integers n and d, we define a function $f : \mathbb{R}^n \to \mathbb{R}$ to be a *hierarchical multilinear function* of non-linearity $\varepsilon > 0$ and depth d recursively as follows. A hierarchical multilinear function of non-linearity $\varepsilon > 0$ and depth 1 is simply a function satisfying the conditions of Lemma 1.1. Note that $d(f, \operatorname{Lin})$ does not depend on the underlying random variables X_1, \ldots, X_n . A hierarchical multilinear function of non-linearity $\varepsilon > 0$ and depth $d \ge 2$ is a function $f : \mathbb{R}^n \to \mathbb{R}$ of the form

$$f(x) = g(h_1(x_{\mathcal{P}_1}), \dots, h_m(x_{\mathcal{P}_m}))$$

for some partition \mathcal{P} of [n] with $m \geq 2$ parts and hierarchical multilinear functions g, h_1, \ldots, h_m of non-linearity ε , where g has depth 1 and each h_i has depth at least d-1. While this definition allows for the h_i to have depth greater than or equal to d, note that any depth-dhierarchical multilinear function of non-linearity $\varepsilon > 0$ must be a function of at least 2^d variables, so our recursive definition does terminate and is well-defined.

As previously mentioned, all functions on Boolean random variables are multilinear, so any function on Boolean random variables of this hierarchical form whose component functions all have Boolean outputs, i.e., take on two values, except for potentially the top component function, are hierarchical multilinear functions. Thus, our definition of a hierarchical multilinear function includes all hierarchical Boolean functions.

We now prove Theorem 1.3, which shows that $\operatorname{Corr}(f(X), f(Y))$ decays exponentially in depth d, for hierarchical multilinear functions of positive non-linearity.

Proof of Theorem 1.3. Let $g(x) = (1 - \varepsilon)x + \varepsilon x^2$. For positive integer $m \ge 1$, let g^m denote the composition of g a total of m times. Because X_i and Y_i have the same first and second moments, and because the (X_i, Y_i) are mutually independent across i, any multilinear function h will have $\mathbb{E}[h(X)] = \mathbb{E}[h(Y)]$ and $\mathbb{E}[h(X)^2] = \mathbb{E}[h(Y)^2]$, and thus we can repeatedly apply Lemma 1.1 for all component functions of our hierarchical multilinear function f, as the first and second moment conditions of Lemma 1.1 inductively hold. By repeatedly applying Lemma 1.1 throughout the hierarchical structure of f, and noting that g is an increasing function on [0, 1] with $0 \leq g(x) < x$ for all $0 \leq x < 1$, we have $\operatorname{Corr}(f(X), f(Y)) \leq g^d(\rho)$, so it suffices to show $g^d(\rho) \leq C_{\rho,\varepsilon,\delta} (1-\delta)^d$ for all $d \geq 1$, for some constant $C_{\rho,\varepsilon,\delta}$. Note that $\frac{g(x)}{x} = 1 - \varepsilon + \varepsilon x$, so there exists some $\alpha = \alpha_{\varepsilon,\delta} \in (0, 1)$ such that for all $x \leq \alpha$, we have $g(x) \leq (1-\delta)x$. Thus it suffices to show there exists some nonnegative integer $N = N_{\rho,\varepsilon,\delta}$ such that $g^N(\rho) \leq \alpha$. If $\rho \leq \alpha$, we simply take N = 0. Otherwise, note that for $x \leq \rho < 1$, we have

$$\frac{1-g(x)}{1-x} = 1 + \varepsilon - \varepsilon(1-x) \ge 1 + \varepsilon\rho > 1,$$

$$_{1+\varepsilon\rho} \left(\frac{1-\alpha}{1-\rho}\right).$$

so we may take $N \ge \log_{1+\varepsilon\rho} \left(\frac{1-\alpha}{1-\rho}\right)$.

Remark 2.2. We remark that this bound is essentially tight. Suppose all component functions defining our hierarchical multilinear function f have non-linearity exactly $\varepsilon > 0$, i.e., each component function g satisfies $d(g, \operatorname{Lin}) = \varepsilon$, and $\operatorname{Corr}(X_i, Y_i) = \rho$ for all i. With these tight conditions, following the proof of Lemma 1.1, it is straightforward to see that we could have proven $\operatorname{Corr}(f(X), f(Y)) \ge (1 - \varepsilon)\rho$ instead of (1.1). Inductively applying this bound instead of Lemma 1.1, we see that our hierarchical multilinear function of depth d has $\operatorname{Corr}(f(X), f(Y)) \ge (1 - \varepsilon)^d \rho$. Hence, it is impossible to have exponential rate smaller than $1 - \varepsilon$, and we have provided a bound with exponential rate α for all $\alpha > 1 - \varepsilon$.

Remark 2.3. Note that in the case $\varepsilon = 1$, i.e., f is a hierarchical multilinear function where each component function is completely uncorrelated with any linear function, Lemma 1.1 actually implies $\operatorname{Corr}(f(X), f(Y)) \leq \rho^{2^d}$. In this case, we have doubly exponential decay in depth, with decay rate depending on ρ , stronger than the singly exponential decay provided by Theorem 1.3, where the decay rate depends on ε .

Remark 2.4. In the literature of cryptography, a Boolean function is *t*-resilient if it is balanced and remains unpredictable even if an adversary knows up to t of the n input bits. Concretely, this means $\hat{f}(S) = 0$ for all $|S| \leq t$. We refer interested readers to [7, Chapter 8] for further background on Boolean functions in cryptography. Our results do not care about the balanced condition, i.e., $\hat{f}(\emptyset) = 0$, but aside from this, our case $\varepsilon = 1$ in the Boolean setting corresponds to 1-resilient functions. It is straightforward to see, by adapting the proof of Lemma 1.1, that for any positive integer t, if we have a hierarchical function whose components are all t-resilient Boolean functions, then we have $\operatorname{Corr}(f(X), f(Y)) \leq \rho^{(t+1)^d}$.

Remark 2.5. One example of such a component function is the parity function in the Boolean case, i.e., $\operatorname{Parity}(x) = x_1 \cdot x_2 \cdots x_n$, where $X_i \sim \operatorname{Unif}(\{\pm 1\})$ for all *i*. This function takes value 1 if an even number of bits are on, i.e., have value -1, and value -1 if an odd number of bits are on. In the coding theory literature, this function is more commonly written as

$$\operatorname{Parity}(x_1,\ldots,x_n)=x_1\oplus\cdots\oplus x_n,$$

where \oplus denotes the XOR (exclusive OR), and each $x_i \in \{0, 1\}$ is a bit. However, to fit this into our multilinear polynomial framework, we convert the additive group $\{0, 1\}$ modulo 2, i.e., $\mathbb{Z}/2\mathbb{Z}$, into the multiplicative group $\{1, -1\}$, which is a standard transformation in the literature of analysis of Boolean functions. Assuming $\operatorname{Corr}(X_i, Y_i) = \rho$ for all *i*, it is well-known, and can be seen by adapting the proof of Lemma 1.1, that $\operatorname{Corr}(\operatorname{Parity}(X), \operatorname{Parity}(Y)) = \rho^n$, i.e., exponential in n. Note that one can view the parity function as a hierarchical function of smaller component parity functions, and by expressing the *n*-bit parity function as a hierarchical function of depth $\lfloor \log_2 n \rfloor$ with components being 2-bit parity functions, we have, via the $\varepsilon = 1$ version of Theorem 1.3 discussed in this remark, that

$$\operatorname{Corr}(\operatorname{Parity}(X), \operatorname{Parity}(Y)) \le \rho^{2^{\lfloor \log_2 n \rfloor}} < \rho^{n/2},$$

so our result still demonstrates exponential decay in n, albeit with a different rate than the true rate.

For example, note that Theorem 1.3 shows that the noise stability of the Boolean function on 3^d bits consisting of recursively composing the majority function on 3 bits, which is not linear, decays exponentially in d. Thus, the recursive majority function is noise sensitive for logarithmic depth. Recall that the fact that the recursive majority function is noise sensitive is folklore; see, for example, [4, 5].

Theorem 1.3 implies low-degree bounds on hierarchical multilinear functions. To make this concrete, the following lemma shows how bounds on noise stability, that is, Corr(f(X), f(Y)), imply bounds on the maximum correlation between f and any multilinear function of degree at most D.

Lemma 2.6. Suppose $\{(X_i, Y_i) : i \in [n]\}$ are mutually independent, where for some $0 \leq \rho \leq 1$, we have $\operatorname{Corr}(X_i, Y_i) = \rho$ for all *i*. Suppose X_i and Y_i have the same first and second moments, *i.e.*, $\mathbb{E}[X_i] = \mathbb{E}[Y_i]$ and $\mathbb{E}[X_i^2] = \mathbb{E}[Y_i^2] < \infty$, for all $i \in [n]$. Let $f : \mathbb{R}^n \to \mathbb{R}$ be a multilinear function, and let M denote the maximum correlation between f and any multilinear function of degree at most D, *i.e.*,

$$M = \sup_{\substack{g: \mathbb{R}^n \to \mathbb{R} \text{ multilinear} \\ \deg(g) \le D}} \operatorname{Corr}(f(X), g(X)).$$

Then

$$M \le \rho^{-D/2} \sqrt{\operatorname{Corr}(f(X), f(Y))}.$$

Proof. We wish to bound

$$M = \sup_{\substack{g: \mathbb{R}^n \to \mathbb{R} \text{ multilinear} \\ \deg(g) \le D}} \operatorname{Corr}(f(X), g(X)).$$

For any such g, decompose f and g as

$$f = \sum_{S \subseteq [n]} \widehat{f}(S) \chi_S, \quad g = \sum_{S \subseteq [n]} \widehat{g}(S) \chi_S,$$

and without loss of generality rescale g so that Var(g) = 1, i.e.,

$$\sum_{S \neq \emptyset} \widehat{g}(S)^2 = 1$$

Following the proof of Lemma 1.1, we know

$$\operatorname{Corr}(f(X), f(Y)) = \frac{1}{\operatorname{Var}(f)} \sum_{S \neq \emptyset} \rho^{|S|} \widehat{f}(S)^2.$$

Similarly, i.e., via orthogonality of $\{\chi_S : S \subseteq [n]\}$, by the Cauchy–Schwarz inequality, we have

 $\sum \hat{c}(\alpha)^2$

$$\operatorname{Corr}(f(X), g(X)) = \frac{1}{\sqrt{\operatorname{Var}(f)}} \sum_{S \neq \emptyset} \widehat{f}(S) \widehat{g}(S) = \frac{1}{\sqrt{\operatorname{Var}(f)}} \sum_{1 \le |S| \le D} \widehat{f}(S) \widehat{g}(S) \le \frac{\sqrt{\sum_{1 \le |S| \le D} f(S)^2}}{\sqrt{\operatorname{Var}(f)}},$$

with equality if and only if

$$g = C + \frac{\sum_{1 \le |S| \le D} \widehat{f}(S) \chi_S}{\sqrt{\sum_{1 \le |S| \le D} \widehat{f}(S)^2}}$$

for some constant $C \in \mathbb{R}$. Thus,

$$M = \frac{1}{\sqrt{\operatorname{Var}(f)}} \sqrt{\sum_{1 \le |S| \le D} \widehat{f}(S)^2}.$$

Then we relate $\operatorname{Corr}(f(X), f(Y))$ with M via

$$\operatorname{Corr}(f(X), f(Y)) = \frac{1}{\operatorname{Var}(f)} \sum_{S \neq \emptyset} \rho^{|S|} \widehat{f}(S)^2$$
$$\geq \frac{1}{\operatorname{Var}(f)} \sum_{1 \leq |S| \leq D} \rho^{|S|} \widehat{f}(S)^2$$
$$\geq \frac{1}{\operatorname{Var}(f)} \sum_{1 \leq |S| \leq D} \rho^D \widehat{f}(S)^2$$
$$= \rho^D M^2,$$

which rearranges to our desired inequality.

Using Theorem 1.3, for any hierarchical multilinear function f of non-linearity $\varepsilon > 0$ and depth d, and for any $\delta < \varepsilon$ and $0 \le \rho < 1$, then applying Lemma 2.6 yields (1.2), i.e., the maximum correlation M between f and any multilinear function of degree at most D is bounded by

(2.1)
$$M \le C_{\rho,\varepsilon,\delta} \rho^{-D/2} (1-\delta)^{d/2},$$

for some constant $C_{\rho,\varepsilon,\delta} > 0$ depending only on $\rho, \varepsilon, \delta$. Note that this is because, as previously discussed, we can always construct a ρ -correlated Y satisfying the conditions of Theorem 1.3. In particular, for fixed D, our bound (1.2) decays exponentially in depth d.

3. Non-separable functions

3.1. Correlated probability spaces. If we hope to generalize Theorem 1.3 beyond multilinear functions, we will need to strengthen our initial assumption, i.e., that $\operatorname{Corr}(X_i, Y_i) \leq \rho$. Because $\operatorname{Corr}(X_i, Y_i)$ only captures linear relationships between X_i and Y_i , if our function is not multilinear, it has the potential to have higher correlation $\operatorname{Corr}(f(X), f(Y))$ than that of its components, i.e., $\operatorname{Corr}(X_i, Y_i)$, by utilizing these nonlinear relationships. Recall Examples 1.4 and 1.5, which illustrate how nonlinear relationships in a single variable can lead to noise stability that does not decay with depth. To this end, we broaden our concept of correlation via the following definition of correlation between probability spaces, sometimes referred to as the "Hirschfeld–Gebelein–Rényi maximal correlation" as it was first introduced by Hirschfeld [11] and Gebelein [9] and then studied by Rényi [19].

Definition 3.1. Given a probability measure \mathbb{P} defined on $\Omega = \prod_{i=1}^{k} \Omega_i$, we say that $\Omega_1, \ldots, \Omega_k$ are *correlated spaces*. Given two linear subspaces A and B of $L^2(\mathbb{P})$, we define the *correlation* between A and B by

$$\rho(A, B; \mathbb{P}) = \rho(A, B) = \sup \left\{ \operatorname{Corr}(f, g) : f \in A, g \in B \right\}$$

For $i, j \in [k]$, we define the *correlation* $\rho(\Omega_i, \Omega_j; \mathbb{P})$ by

$$\rho(\Omega_i, \Omega_j; \mathbb{P}) = \rho(\Omega_i, \Omega_j) = \rho(L^2(\Omega_i, \mathbb{P}), L^2(\Omega_j, \mathbb{P}); \mathbb{P}).$$

Because ρ is used as a constant in this paper, we will also use the notation $\operatorname{Corr}(\cdot)$ in place of $\rho(\cdot)$ for clarity.

For random vector $X : \Omega \to \mathbb{R}^n$, let $L^2(X)$ denote the linear subspace of $L^2(\mathbb{P})$ that is $\sigma(X)$ -measurable. Each $f \in L^2(X)$ can be identified with a function $\mathfrak{f} : \mathbb{R}^n \to \mathbb{R}$ satisfying $\mathbb{E}[\mathfrak{f}(X)^2] < \infty$, and we will frequently make no distinction between these two functions.

Equipped with this new definition, we will consider the condition that $\operatorname{Corr}(L^2(X_i), L^2(Y_i)) \leq \rho$, in place of the condition that $\operatorname{Corr}(X_i, Y_i) \leq \rho$.

We recall some theory from Mossel [15, §2.2] concerning noise correlation of functions. For two random variables $X, Y : \Omega \to \mathbb{R}$, if $f \in L^2(X)$ satisfies $\mathbb{E}[f(X)] = 0$ and $\mathbb{E}[f(X)^2] = 1$, then the maximizer of $|\mathbb{E}[f(X)g(Y)]|$ among $g \in L^2(Y)$ satisfying $\mathbb{E}[g(Y)^2] = 1$ is given by

$$g = \frac{\mathbb{E}[f(X)|Y]}{\|\mathbb{E}[f(X)|Y]\|_2} = \frac{\mathbb{E}[f(X)|Y]}{\sqrt{\mathbb{E}[f(X)|Y]^2}},$$

and

$$\left|\mathbb{E}[f(X)g(Y)]\right| = \left\|\mathbb{E}[f(X)|Y]\right\|_{2}$$

The function $\mathbb{E}[f(X)|Y] \in L^2(Y)$ is the *Markov operator* from $L^2(X)$ to $L^2(Y)$, and as such we will denote this g by $Tf \in L^2(Y)$. We have $\mathbb{E}[Tf] = 0$, so

(3.1)
$$\operatorname{Corr}(L^{2}(X), L^{2}(Y)) = \sup \left\{ \mathbb{E}[f(X)Tf(X)] : f \in L^{2}(X), \mathbb{E}[f] = 0, \mathbb{E}[f^{2}] = 1 \right\}$$

3.2. Efron–Stein decomposition. Returning to our setup in Section 2, where \mathbb{P} is a product measure on product space Ω over the laws of correlated pairs of random variables $(X_1, Y_1), \ldots, (X_n, Y_n)$, we now introduce the Efron–Stein decomposition as our analog to the χ_S that decomposed our multilinear polynomials.

Definition 3.2. Let $(\Omega_1, \mu_1), \ldots, (\Omega_n, \mu_n)$ be laws of correlated pairs of real-valued random variables $(X_1, Y_1), \ldots, (X_n, Y_n)$, and let $(\Omega, \mu) = \prod_{i=1}^n (\Omega_i, \mu_i)$. The *Efron–Stein decomposition* of $f \in L^2(X)$ is given by

$$f(x) = \sum_{S \subseteq [n]} f_S(x),$$

where each function f_S depends only on x_S and for all $S \not\subseteq S'$, we have $\mathbb{E}[f_S|X_{S'}] = 0$ almost surely. We can similarly define the Efron–Stein decomposition for any $g \in L^2(Y)$.

RUPERT LI AND ELCHANAN MOSSEL

It is well-known that the Efron–Stein decomposition exists and is unique [8]. A standard result (see, for example, [15, Proposition 2.11], which only proves the result in the discrete case but naturally holds in general) states that the Efron–Stein decomposition "commutes" with the Markov operator, i.e., $(Tf)_S = T(f_S) \in L^2(Y)$, where $T : L^2(X) \to L^2(Y)$ is the Markov operator, which is given by $T = \bigotimes_{i=1}^n T_i$ for Markov operators $T_i : L^2(X_i) \to L^2(Y_i)$. It thus is unambiguous to write Tf_S . Similarly generalizing the proof of [15, Proposition 2.12], if $\operatorname{Corr}(L^2(X_i), L^2(Y_i)) \leq \rho_i$ for some ρ_i for each i, then for all $S \subseteq [n]$,

(3.2)
$$\|Tf_S\|_2 \le \left(\prod_{i\in S} \rho_i\right) \|f_S\|_2.$$

3.3. Hierarchical non-separable functions. Recall the goal of this section is to generalize Theorem 1.3 beyond multilinear functions. Just as we had to broaden our assumption that $\operatorname{Corr}(X_i, Y_i) \leq \rho$ to $\operatorname{Corr}(L^2(X_i), L^2(Y_i)) \leq \rho$, we must also broaden our assumption that f is non-linear to account for non-linear relationships in a single variable. To this end, we replace the concept of non-linearity with non-separability, where recall an (additively) separable function $f : \mathbb{R}^n \to \mathbb{R}$ is a function of the form $f(x) = f_1(x_1) + \cdots + f_n(x_n)$ for $f_1, \ldots, f_n : \mathbb{R} \to \mathbb{R}$. Note that in the context of multilinear polynomials, separability and linearity are equivalent, so this definition of a separable function is consistent with our setup in Section 2. We can then define non-separability by the condition that

$$\operatorname{Corr}(L^2(f(X)), L^2(X_1) + \dots + L^2(X_n)) \le 1 - \varepsilon.$$

For any $g \in L^2(f(X))$ and $h_i \in L^2(X_i)$ for all $i \in [n]$, where we normalize $\mathbb{E}[g] = 0 = \mathbb{E}[h_i]$ for all i and $\mathbb{E}[g^2] = 1 = \mathbb{E}[(h_1 + \cdots + h_n)^2]$, we have by the defining properties of the Efron–Stein decomposition, where we view $G = g \circ f \in L^2(X)$,

$$\operatorname{Corr}(g, h_1 + \dots + h_n) = \mathbb{E}[g(f(X))(h_1(X_1) + \dots + h_n(X_n))]$$

= $\sum_{S \neq \emptyset} \mathbb{E}[G_S(h_1 + \dots + h_n)_S]$
= $\sum_{i=1}^n \mathbb{E}[G_{\{i\}}h_i]$
 $\leq \sum_{i=1}^n \|G_{\{i\}}\|_2 \|h_i\|$
 $\leq \sqrt{\sum_{i=1}^n \|G_{\{i\}}\|_2^2} \sqrt{\sum_{i=1}^n \|h_i\|_2^2}$
= $\sqrt{\sum_{i=1}^n \|G_{\{i\}}\|_2^2}.$

Note that by picking h_i appropriately, one can make both applications of the Cauchy–Schwarz inequality tight, and thus

(3.3)
$$\operatorname{Corr}(L^{2}(f(X)), L^{2}(X_{1}) + \dots + L^{2}(X_{n})) = \sup_{\substack{g \in L^{2}(f(X))\\ \mathbb{E}[g]=0, \mathbb{E}[g^{2}]=1}} \sqrt{\sum_{i=1}^{n} \|G_{\{i\}}\|_{2}^{2}}$$

Because of the square root on the right hand side, we will reparametrize ε so that our non-separability condition is

$$\operatorname{Corr}(L^2(f(X)), L^2(X_1) + \dots + L^2(X_n)) \le \sqrt{1 - \varepsilon}$$

By naturally extending our definition of d(f, V) to d(U, V) for $U, V \subseteq L^2(\Omega, \mathbb{P})$ by

$$d(U,V) = \inf_{f \in U, g \in V} \frac{\mathbb{E}[(f-g)^2]}{\operatorname{Var}(f)},$$

note that

(3.4)
$$d(L^{2}(f(X)), L^{2}(X_{1}) + \dots + L^{2}(X_{n})) = \inf_{g,h_{1},\dots,h_{n}} \frac{\sum_{S} \|G_{S} - (h_{1})_{S} - \dots - (h_{n})_{S})\|_{2}^{2}}{\sum_{S \neq \emptyset} \|G_{S}\|_{2}^{2}}$$
$$= \inf_{\substack{g \in L^{2}(f(X))\\ \mathbb{E}[g] = 0, \mathbb{E}[g^{2}] = 1}} \sum_{|S| \ge 2} \|G_{S}\|_{2}^{2},$$

where we have used the fact that $f \mapsto f_S$ is a linear operator, which follows from linearity of expectation. Assuming $\mathbb{E}[g] = 0$ and $\mathbb{E}[g^2] = 1$, we have $G_{\emptyset} \equiv 0$ and $\sum_{S \subseteq [n]} ||G_S||_2^2 = 1$, so we see that the extremizers in (3.3) and (3.4) are the same g, and (3.5)

$$\operatorname{Corr}(L^2(f(X)), L^2(X_1) + \dots + L^2(X_n)) \le \sqrt{1 - \varepsilon} \iff d(L^2(f(X)), L^2(X_1) + \dots + L^2(X_n)) \ge \varepsilon$$

While the latter condition is more consistent with our condition in Section 2, we will use the former condition to consistently use the language of correlated probability spaces. Of course, this choice is purely stylistic, as these two conditions are equivalent. We say a function $f \in L^2(X)$ satisfying this condition is ε -non-separable. Likewise, we can define a hierarchical function $f \in L^2(X)$ of non-separability $\varepsilon > 0$ and depth d analogously to how we defined hierarchical multilinear functions.

We are now ready to prove Lemma 1.6, our generalization of Lemma 1.1 to non-separable functions.

Proof of Lemma 1.6. Let Z = f(X) and W = f(Y). Pick arbitrary $g, h : \mathbb{R} \to \mathbb{R}$ such that $\mathbb{E}[g(Z)] = \mathbb{E}[h(W)] = 0$ and $\mathbb{E}[g(Z)^2] = \mathbb{E}[h(W)^2] = 1$. It suffices to show $\mathbb{E}[g(Z)h(W)] \leq (1-\varepsilon)\rho + \varepsilon\rho^2$. Let

$$G = g \circ f \in L^2(X)$$
 and $H = h \circ f \in L^2(Y).$

Note that for distinct $S, S' \subseteq [n]$, we have $\mathbb{E}[G_S(X)H_{S'}(Y)] = 0$. This is because we either have $S \not\subseteq S'$ or $S' \not\subseteq S$, and in the former case, by the defining properties of the Efron–Stein

decomposition and recalling the (X_i, Y_i) are mutually independent across i,

$$\mathbb{E}[G_S(X)H_{S'}(Y)] = \mathbb{E}[\mathbb{E}[G_S(X)H_{S'}(Y)|(X_{S'},Y_{S'})]]$$
$$= \mathbb{E}[\mathbb{E}[G_S(X)|(X_{S'},Y_{S'})]H_{S'}(Y)]$$
$$= \mathbb{E}[0 \cdot H_{S'}(Y)]$$
$$= 0.$$

Thus,

$$\begin{split} \mathbb{E}[g(Z)h(W)] &= \sum_{S \neq \emptyset} \mathbb{E}[G_S(X)H_S(Y)] \\ &= \sum_{S \neq \emptyset} \|G_S\|_2 \, \|H_S\|_2 \, \mathbb{E}\left[\frac{G_S(X)}{\|G_S\|_2} \frac{H_S(Y)}{\|H_S\|_2}\right] \\ &\leq \sum_{S \neq \emptyset} \|G_S\|_2 \, \|H_S\|_2 \, \mathbb{E}\left[\frac{G_S(X)}{\|G_S\|_2} T\left(\frac{G_S}{\|G_S\|_2}\right)\right] \\ &= \sum_{S \neq \emptyset} \|G_S\|_2 \, \|H_S\|_2 \, \|TG_S\|_2 \\ &\leq \sum_{S \neq \emptyset} \|G_S\|_2 \, \|H_S\|_2 \, \rho^{|S|}, \end{split}$$

where the first inequality uses the property that the Markov operator maximizes covariance as discussed in Section 3.1, and the second inequality follows from (3.2). Applying the Cauchy–Schwarz inequality and then our non-separability condition, we bound

$$\mathbb{E}[g(Z)h(W)] \leq \sqrt{\sum_{S \neq \emptyset} \|G_S\|_2^2 \rho^{|S|}} \sqrt{\sum_{S \neq \emptyset} \|H_S\|_2^2 \rho^{|S|}} \leq (1-\varepsilon)\rho + \varepsilon \rho^2,$$

which completes the proof.

Theorem 1.7 follows from the same proof as that of Theorem 1.3 using Lemma 1.6 in place of Lemma 1.1.

Remark 3.3. Again, we note that in the case $\varepsilon = 1$, i.e., f is a hierarchical function where each component function has zero correlation with any function of a single one of its component variables, Lemma 1.6 implies $\operatorname{Corr}(L^2(f(X)), L^2(f(Y))) \leq \rho^{2^d}$. So, analogously to Remark 2.3, in this case we have doubly exponential decay in depth with decay rate depending on ρ , stronger than the singly exponential decay provided by Theorem 1.7 with decay rate depending on ε .

Lastly, we state and prove the following general version of Lemma 2.6, which we use to justify (1.4).

Lemma 3.4. Suppose $\{(X_i, Y_i) : i \in [n]\}$ are mutually independent, where for some $0 \le \rho \le 1$, we have $\operatorname{Corr}(L^2(X_i), L^2(Y_i)) = \rho$ for all *i*. Suppose X and Y have the same marginal distributions. If $f : \mathbb{R}^n \to \mathbb{R}$ satisfies $\mathbb{E}[f(X)^2] < \infty$, then let M denote the maximum

correlation between $L^2(f)$ and any function of Efron–Stein degree at most D, i.e., $M = Corr(L^2(f(X)), \mathcal{G})$, where $\mathcal{G} = \sum_{|S| \leq D} L^2(X_S)$. Let

$$\gamma = \sup_{G,H} \sum_{S \neq \emptyset} \|G_S\|_2 \, \|H_S\|_2 \, \rho^{|S|},$$

where the supremum is taken over all $g, h : \mathbb{R} \to \mathbb{R}$ such that $\mathbb{E}[g(f(X))] = \mathbb{E}[h(f(Y))] = 0$ and $\mathbb{E}[g(f(X))^2] = \mathbb{E}[h(f(Y))^2] = 1$, and $G = g \circ f \in L^2(X)$ and $H = h \circ f \in L^2(Y)$. Then

$$M \le \rho^{-D/2} \sqrt{\gamma}.$$

Proof. Let Z = f(X) and W = f(Y). Using a slightly generalized but essentially similar argument as that which we used to show (3.3), we have

$$M = \sup_{\substack{g \in L^2(f(X)) \\ \mathbb{E}[g] = 0, \mathbb{E}[g^2] = 1}} \sqrt{\sum_{1 \le |S| \le D} \|G_S\|_2^2},$$

where $G = g \circ f \in L^2(X)$. It then suffices to show that for any such g,

$$\sqrt{\sum_{1 \le |S| \le D} \|G_S\|_2^2} \le \rho^{-D/2} \left(\sup_{\substack{h \in L^2(f(Y))\\ \mathbb{E}[h] = 0, \mathbb{E}[h^2] = 1}} \sum_{S \ne \emptyset} \|G_S\|_2 \|H_S\|_2 \rho^{|S|} \right)^{1/2},$$

where $H = h \circ f \in L^2(Y)$. Letting g and h be the same when viewed as functions $\mathbb{R} \to \mathbb{R}$, because X and Y have the same marginal distributions, we note that $H_S = G_S$ for all S, which achieves the equality case of the Cauchy–Schwarz inequality that implies

$$\sup_{\substack{h \in L^2(f(Y))\\ \mathbb{E}[h] = 0, \mathbb{E}[h^2] = 1}} \sum_{S \neq \emptyset} \|G_S\|_2 \|H_S\|_2 \rho^{|S|} = \sum_{S \neq \emptyset} \|G_S\|_2^2 \rho^{|S|}.$$

Hence, it suffices to show

$$\sum_{1 \le |S| \le D} \|G_S\|_2^2 \rho^D \le \sum_{S \ne \emptyset} \|G_S\|_2^2 \rho^{|S|},$$

which is clearly true as $0 \le \rho \le 1$.

To justify (1.4), note that for any $0 \le \rho < 1$, using Remark 1.2 we can construct Y satisfying the conditions of Lemma 3.4. Recall that Theorem 1.7 was proven by inductively applying Lemma 1.6. Using the fact that the proof of Lemma 1.6 implies the stronger technical statement

$$\sup_{G,H} \sum_{S \neq \emptyset} \|G_S\|_2 \|H_S\|_2 \rho^{|S|} \le (1-\varepsilon)\rho + \varepsilon \rho^2,$$

the proof of Theorem 1.7 implies that for a hierarchical function $f \in L^2(X)$ of non-separability $\varepsilon > 0$ and depth d, and for any $\delta < \varepsilon$ and $0 \le \rho < 1$ (where because our constructed Y has the same marginal distribution as X, we have f is also such a hierarchical function in $L^2(Y)$),

$$\sup_{G,H} \sum_{S \neq \emptyset} \|G_S\|_2 \|H_S\|_2 \rho^{|S|} \le C_{\rho,\varepsilon,\delta} (1-\delta)^d.$$

Thus, Lemma 3.4 implies

$$M \le C'_{\rho,\varepsilon,\delta} \,\rho^{-D/2} (1-\delta)^{d/2},$$

where $C'_{\rho,\varepsilon,\delta} = \sqrt{C_{\rho,\varepsilon,\delta}} > 0$, which justifies (1.4).

Acknowledgements

E.M. was partially supported by NSF DMS-2031883, Bush Faculty Fellowship ONR-N00014-20-1-2826, and Simons Investigator award (622132).

References

- E. Abbe, E. B. Adsera, and T. Misiakiewicz, The merged-staircase property: a necessary and nearly sufficient condition for sgd learning of sparse functions on two-layer neural networks, Conference on Learning Theory, PMLR, 2022, pp. 4782–4887.
- [2] E. Abbe, S. Bengio, E. Cornacchia, J. Kleinberg, A. Lotfi, M. Raghu, and C. Zhang, *Learning to reason with neural networks: Generalization, unseen data and boolean measures*, Advances in Neural Information Processing Systems 35 (2022), 2709–2722.
- [3] E. Abbe, E. Boix-Adsera, M. S. Brennan, G. Bresler, and D. Nagaraj, *The staircase property: How hierarchical structure can guide deep learning*, Advances in Neural Information Processing Systems 34 (2021), 26989–27002.
- [4] M. Ben-Or and N. Linial, Collective coin flipping, Randomness and Computation (S. Micali, ed.), Academic Press, New York, 1990.
- [5] I. Benjamini, G. Kalai, and O. Schramm, Noise sensitivity of boolean functions and applications to percolation, Inst. Hautes Études Sci. Publ. Math. 90 (1999), 5–43.
- [6] J. Bruna and S. Mallat, *Invariant scattering convolution networks*, IEEE transactions on pattern analysis and machine intelligence 35 (2013), no. 8, 1872–1886.
- [7] Y. Crama and P. L. Hammer (eds.), Boolean models and methods in mathematics, computer science, and engineering, Encyclopedia of Mathematics and its Applications, vol. 134, Cambridge University Press, Cambridge, 2010.
- [8] B. Efron and C. Stein, The jacknife estimate of variance, Annals of Statistics 9 (1996), no. 3, 586–596.
- [9] H. Gebelein, Das statistische Problem der Korrelation als Variations- und Eigenwertproblem und sein Zusammenhang mit der Ausgleichsrechnung, Z. Angew. Math. Mech. 21 (1941), 364–379.
- [10] J. Håstad, B. Rossman, R. A. Servedio, and L.-Y. Tan, An average-case depth hierarchy theorem for boolean circuits, Journal of the ACM (JACM) 64 (2017), no. 5, 35.
- [11] H. O. Hirschfeld, A connection between correlation and contingency, Mathematical Proceedings of the Cambridge Philosophical Society, vol. 31, Cambridge University Press, 1935, pp. 520–524.
- [12] F. Koehler and E. Mossel, Reconstruction on trees and low-degree polynomials, Advances in Neural Information Processing Systems 35 (2022), 18942–18954.
- [13] H. Mhaskar, Q. Liao, and T. Poggio, *Learning functions: when is deep better than shallow*, arXiv preprint arXiv:1603.00988 (2016).
- [14] E. Mossel and R. O'Donnell, On the noise sensitivity of monotone functions, Mathematics and computer science, II (Versailles, 2002) (Basel), Trends Math., Birkhäuser, 2002, pp. 481–495.
- [15] E. Mossel, Gaussian bounds for noise correlation of functions, Geom. Funct. Anal. 19 (2010), no. 6, 1713–1756.
- [16] _____, Deep learning and hierarchal generative models, arXiv preprint arXiv:1612.09057, 2016.
- [17] R. O'Donnell, Analysis of boolean functions, Cambridge University Press, 2014.
- [18] A. B. Patel, T. Nguyen, and R. G. Baraniuk, A probabilistic theory of deep learning, arXiv preprint arXiv:1504.00641 (2015).
- [19] A. Rényi, On measures of dependence, Acta Math. Acad. Sci. Hungar. 10 (1959), 441–451.
- [20] T. Schramm and A. S. Wein, Computational barriers to estimation from low-degree polynomials, The Annals of Statistics 50 (2022), no. 3, 1833–1858.

[21] M. Telgarsky, benefits of depth in neural networks, Conference on Learning Theory, 2016, arXiv preprint arXiv:1602.04485, pp. 1517–1539.

UNIVERSITY OF CAMBRIDGE, CAMBRIDGE, CB2 1TN, GBR *Email address:* rml61@cam.ac.uk

MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, USA $\mathit{Email}\ address: \texttt{elmos}\texttt{Qmit.edu}$