

# Stochastic internal habit formation and optimality

M. Aleandri<sup>\*1</sup>, A. Bondi<sup>†1</sup>, and F. Gozzi<sup>‡1</sup>

<sup>1</sup> LUISS University, DEF, 00197 viale Romania 32, Roma, Italy.

## Abstract

Growth models with internal habit formation have been studied in various settings under the assumption of deterministic dynamics. The purpose of this paper is to explore a stochastic version of the model in Carroll et al. [1997, 2000], one of the most influential on the subject. The goal is twofold: on one hand, to determine how far we can advance in the technical study of the model; on the other, to assess whether at least some of the deterministic outcomes remain valid in the stochastic setting. The resulting optimal control problem proves to be challenging, primarily due to the lack of concavity in the objective function. This feature is present in the model even in the deterministic case (see, e.g., Bambi and Gozzi [2020]). We develop an approach based on Dynamic Programming to establish several useful results, including the regularity of the solution to the corresponding HJB equation and a verification theorem. These results lay the groundwork for studying the model optimal paths and comparing them with the deterministic case.

**Keywords:** Habit formation; Optimal investment and consumption; Stochastic control; Dynamic programming; Viscosity solution; Hamilton–Jacobi–Bellman equation.

**Mathematics Subject Classification (2000):** 49L20; 49L12; 49L25; 49J55; 93E20;

**JEL Classification:** C32; C61; D91; E21; O40.

---

\*Email: [maleandri@luiss.it](mailto:maleandri@luiss.it).

†Email: [abondi@luiss.it](mailto:abondi@luiss.it).

‡Email: [fgozzi@luiss.it](mailto:fgozzi@luiss.it).

# 1 Introduction

In two highly influential works, Carroll et al. [1997, 2000] analyze the dynamics of an endogenous growth model incorporating internal habit formation. They demonstrate that this feature plays a key role in explaining how increased economic growth can lead to higher savings. This finding is particularly significant because it is based on a “multiplicative” habit formation model, as opposed to the “subtractive” form, which has several drawbacks. Among them, the “subtractive” form fosters addictive behavior by requiring consumption to consistently exceed the habit stock, potentially resulting in infinitely negative utility.

In Bambi and Gozzi [2020], the authors revisit the optimality of the solution found in Carroll et al. [2000] using a Dynamic Programming approach, differing from previous works that relied on the Maximum Principle. The advantage of Dynamic Programming lies in its ability to explore global dynamics and, more importantly, to establish sufficient optimality conditions without requiring concavity assumptions. Indeed, concavity does not hold in this model (see, e.g., Bambi and Gozzi [2020] for a discussion).

This paper investigates a stochastic version of the above model, incorporating multiplicative noise in the state equations. The aim is to conduct a thorough technical analysis of the model while also evaluating whether at least some of its original deterministic outcomes remain valid. We refer to, for instance, Angoshtari et al. [2022] for another stochastic model with habit formation in a different context.

The resulting optimal control problem proves to be highly challenging, for two main reasons: the lack of concavity in the objective function – which, as already mentioned, is one of the model’s key characteristics even in the deterministic case – and its singularity (i.e., the limit is  $-\infty$ ) when the consumption goes to 0.

We tackle this optimal control problem by developing the Dynamic Programming approach. In particular, we show that the corresponding value function, denoted by  $V$ , is a classical solution of the HJB equation in the first open quadrant  $\mathbb{R}_{++}^2$ , see Theorem 5.2. This is a nontrivial result, which is established in three main steps. More precisely, we first obtain some properties of  $V$  (see Section 4), then we prove that  $V$  is a viscosity solution of the HJB equation, and finally we apply a local regularization approach based on crucial results by Safonov [1989] and Crandall et al. [1992]. We also establish a verification theorem, see Theorem 6.2, which gives a sufficient condition for optimality.

These results provide a solid foundation for the study of the model optimal paths and their connections with the deterministic case. This will be the subject of subsequent research.

The paper is organized as follows. Section 2 introduces the model and the optimal control problem we are interested in. In Section 3, the corresponding HJB equation is

formally obtained, together with some properties of the maximum value Hamiltonian. Some important properties (such as monotonicity and continuity) of the value function  $V$  associated with the control problem are then investigated in Section 4. In Section 5 we prove that  $V$  is a classical solution of the HJB equation, see Theorem 5.2. This section is divided into two subsections: in the first (Subsection 5.1), we demonstrate that  $V$  is a viscosity solution, while in the second (Subsection 5.2) we prove the main Theorem 5.2. Finally, the verification theorem is established in Section 6. In Appendix A, we prove some technical results which are employed in the arguments of Subsection 5.2.

## 2 The model

Consider a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  endowed with a filtration  $\mathbb{F} = (\mathcal{F}_t)_t$  which satisfies the usual hypotheses. The model state variables are the processes  $k = (k_t)_{t \geq 0}$  and  $h = (h_t)_{t \geq 0}$ , representing the capital and the habit, respectively. Their evolution is described by the following controlled linear stochastic differential equations (henceforth, SDEs):

$$dk_t = (Bk_t - c_t)dt + \beta_1 k_t dW_{1,t}, \quad k_0 > 0, \quad (1)$$

$$dh_t = \rho(c_t - h_t)dt + \beta_2 h_t dW_{2,t}, \quad h_0 > 0. \quad (2)$$

Here,  $W_1, W_2$  are two independent Brownian motions,  $\beta_1, \beta_2 > 0$ ,  $\rho \in (0, 1)$  and  $B \geq 0$ . The positive initial conditions  $k_0$  and  $h_0$  are assumed to be deterministic. The control process  $c = (c_t)_{t \geq 0}$  represents the consumption; it is progressively measurable and (strictly) positive, with locally integrable trajectories. For any choice of  $c$  and  $h_0, k_0$ , the dynamics (1)-(2) admit unique solutions, up to indistinguishability. Their explicit expressions are given by

$$k_t = \mathcal{E}(Bt + \beta_1 W_{1,t}) \left( k_0 - \int_0^t \frac{c_s}{\mathcal{E}(Bs + \beta_1 W_{1,s})} ds \right), \quad t \geq 0, \quad (3)$$

$$h_t = \mathcal{E}(-\rho t + \beta_2 W_{2,t}) \left( h_0 + \rho \int_0^t \frac{c_s}{\mathcal{E}(-\rho s + \beta_2 W_{2,s})} ds \right), \quad t \geq 0, \quad (4)$$

where we denote by  $\mathcal{E}(X_t)$  the Doléans-Dade exponential of a continuous semimartingale  $X$  evaluated at time  $t$ . In the sequel, when the context requires to clarify the dependence of these solutions on the initial conditions and the control, we will denote by

$$h^{h_0, 0, c} \text{ the solution of (2) controlled by } c \text{ starting from } h_0 \text{ at time 0.} \quad (5)$$

An analogous notation will be used for the capital process  $k^{k_0, 0, c}$ .

For a fixed initial condition  $(k_0, h_0)$ , we denote by  $\mathcal{A}(k_0, h_0)$  the set of admissible controls.

It is determined by imposing the positivity state constraints

$$k_t > 0 \quad \text{and} \quad h_t > 0, \quad \text{for every } t \geq 0, \mathbb{P} - \text{a.s.}, \quad (6)$$

together with the following relation between  $c$  and  $k$ :

$$c \leq Rk, \quad \mathbb{P} \otimes dt - \text{a.e.}, \text{ for some constant } R > 0. \quad (7)$$

Summarizing,

$$\mathcal{A}(k_0, h_0) := \left\{ c: \Omega \times [0, \infty) \rightarrow \mathbb{R} \text{ progr. meas.} \begin{cases} c(\omega, \cdot) \in L^1_{\text{loc}}(0, \infty), \text{ for } \mathbb{P}\text{-a.e. } \omega \in \Omega, \\ c > 0, \text{ for } \mathbb{P} \otimes dt - \text{a.e.}, \text{ the solutions of} \\ (1)\text{-}(2) \text{ satisfy (6), and (7) holds} \end{cases} \right\}. \quad (8)$$

Given  $\theta > 0$ ,  $\sigma \in (1, \infty)$  and  $\gamma \in [0, 1)$ , the utility function of our control problem is defined by

$$\begin{aligned} J(k_0, h_0; c) &:= \mathbb{E} \left[ \int_0^\infty \frac{1}{1-\sigma} \left( \frac{c_t}{h_t^\gamma} \right)^{1-\sigma} e^{-\theta t} dt \right] \\ &= -\frac{1}{\sigma-1} \mathbb{E} \left[ \int_0^\infty \left( \frac{h_t^\gamma}{c_t} \right)^{\sigma-1} e^{-\theta t} dt \right] \leq 0, \quad k_0, h_0 > 0, c \in \mathcal{A}(k_0, h_0). \end{aligned} \quad (9)$$

From now on, we write  $\mathbb{R}_+$  for  $(0, \infty)$  and  $\mathbb{R}_{++}^2$  for  $(0, \infty)^2$ . The value function  $V: \mathbb{R}_{++}^2 \rightarrow [-\infty, 0]$  associated with  $J$  in (9) is

$$V(k_0, h_0) = \sup_{c \in \mathcal{A}(k_0, h_0)} J(k_0, h_0; c), \quad (k_0, h_0) \in \mathbb{R}_{++}^2. \quad (10)$$

We assume that the parameters of the model fulfill the following requirement (cf. [Freni et al., 2008, Assumption 2.4]).

**Assumption 1.** Suppose that  $R \geq B$ . Furthermore, the parameters  $(\beta_1, \beta_2, \gamma, \sigma, \theta)$  satisfy the following conditions:

(i) if  $\rho + \frac{1}{2}(\beta_2^2 - \beta_1^2) \leq 0$ , then

$$\theta > (\sigma - 1) \left( -\gamma\rho - \frac{1}{2}(\gamma\beta_2^2 - \beta_1^2) + \max \left\{ 2, \frac{1}{\gamma(\sigma - 1)} \right\} (\sigma - 1)(\beta_1^2 + \gamma^2\beta_2^2) \right); \quad (11)$$

(ii) if  $\rho + \frac{1}{2}(\beta_2^2 - \beta_1^2) > 0$ , then

$$\theta > (\sigma - 1) \left( \frac{1}{2}\beta_1^2(1 - \gamma) + \max \left\{ 2, \frac{1}{\gamma(\sigma - 1)} \right\} (\sigma - 1)(\beta_1^2 + \gamma^2\beta_2^2) \right). \quad (12)$$

This assumption is sufficient to ensure the existence of strategies with finite utility, as demonstrated in Proposition 4.2.

We conclude this section with a lemma showing that, for every initial point in  $\mathbb{R}_{++}^2$ , the set of admissible controls is nonempty. This is crucial to ensure the meaningfulness of our setting.

**Lemma 2.1.** *For every  $(k_0, h_0) \in \mathbb{R}_{++}^2$ , the set  $\mathcal{A}(k_0, h_0)$  of admissible controls is nonempty. In addition,  $\mathcal{A}(k_0, h_0)$  does not depend on  $h_0$ , that is,*

$$\mathcal{A}(k_0, h_0) = \mathcal{A}(k_0). \quad (13)$$

*Proof.* Fix  $(k_0, h_0) \in \mathbb{R}_{++}^2$ . Choosing  $c = \nu \mathcal{E}(\beta_1 W_1)$ , for some constant  $\nu > 0$  to be determined, (3) reads

$$k_t = e^{Bt} \mathcal{E}(\beta_1 W_{1,t}) \left( k_0 - \frac{\nu}{B} (1 - e^{-Bt}) \right), \quad t \geq 0. \quad (14)$$

Therefore, if we take  $\nu > 0$  such that  $\frac{\nu}{B} < k_0$ ,

$$k_t \geq \left( k_0 - \frac{\nu}{B} \right) \mathcal{E}(\beta_1 W_{1,t}) > 0, \quad t \geq 0,$$

hence the positivity constraint (6) on  $k$  is satisfied. If we further require that  $\frac{R}{\nu} (k_0 - \frac{\nu}{B}) > 1$ , then, considering also that  $c > 0$ ,

$$Rk_t \geq \frac{R}{\nu} \left( k_0 - \frac{\nu}{B} \right) c_t > c_t, \quad t \geq 0,$$

which proves that (7) is fulfilled.

The two previous conditions on  $\nu$  are satisfied by taking  $\nu < Bk_0$  sufficiently small. With this choice of  $\nu$ , we claim that  $c \in \mathcal{A}(k_0, h_0) \neq \emptyset$ . Indeed, from the explicit expression of the habit process  $h$  in (4), the positivity constraint (6) is always satisfied when the control process is positive. This also demonstrates that (13) holds. Thus, the proof is complete.  $\square$

### 3 HJB equation: Formal derivation

Given  $(k, h) \in \mathbb{R}_{++}^2$ ,  $P \in \mathbb{R}^2$  and  $Q \in \mathbb{R}^{2 \times 2}$ , the current value Hamiltonian is

$$H(k, h, P, Q; c) = a(k, h)^\top P + \frac{1}{2} \text{tr}(\Sigma(k, h)^\top Q \Sigma(k, h)) + \frac{1}{1 - \sigma} \left( \frac{h^\gamma}{c} \right)^{1 - \sigma}, \quad c \in (0, Rk].$$

Here,  $a(k, h) = \begin{bmatrix} Bk-c \\ \rho(c-h) \end{bmatrix}$  and  $\Sigma(k, h) = \begin{bmatrix} \beta_1 k & 0 \\ 0 & \beta_2 h \end{bmatrix}$ . Therefore, denoting by  $D^2g$  the Hessian matrix of a sufficiently smooth function  $g$ , the HJB equation associated with the value function  $V$  is, formally,

$$\theta V = \sup_{c \in (0, Rk]} H(k, h, \nabla V, D^2V; c). \quad (15)$$

More explicitly, (15) reads

$$\begin{aligned} \theta V(k, h) &= Bk \partial_k V(k, h) - \rho h \partial_h V(k, h) + \frac{1}{2} \text{tr}(\Sigma(k, h)^\top D^2V(k, h) \Sigma(k, h)) \\ &\quad + \sup_{c \in (0, Rk]} \left\{ -c \partial_k V(k, h) + \rho c \partial_h V(k, h) - \frac{1}{\sigma-1} \left( \frac{h^\gamma}{c} \right)^{\sigma-1} \right\} \\ &=: \mathcal{L}_{k,h} V + \sup_{c \in (0, Rk]} g(c; h, \partial_k V(k, h), \partial_h V(k, h)). \end{aligned} \quad (16)$$

We now focus on the supremum appearing in (16), writing  $\partial_k V$  (resp.,  $\partial_h V$ ) for  $\partial_k V(k, h)$  (resp.,  $\partial_h V(k, h)$ ) to make the notation shorter. When  $\partial_k V - \rho \partial_h V \leq 0$ , the mapping  $g(\cdot; h, \partial_k V, \partial_h V)$  is increasing in  $(0, Rk]$ , hence it achieves its maximum at  $c = Rk$ .

When  $\partial_k V - \rho \partial_h V > 0$ , the study of the first derivative sign entails that

$$g(\cdot; h, \partial_k V, \partial_h V) \text{ is increasing in } \left( 0, \left( \frac{h^{\gamma(\sigma-1)}}{\partial_k V - \rho \partial_h V} \right)^{\frac{1}{\sigma}} \right] \text{ and decreasing in } \left[ \left( \frac{h^{\gamma(\sigma-1)}}{\partial_k V - \rho \partial_h V} \right)^{\frac{1}{\sigma}}, \infty \right).$$

As a result, also in this case the supremum is in fact a maximum achieved at

$$c = \min \left\{ Rk, \left( \frac{h^{\gamma(\sigma-1)}}{\partial_k V - \rho \partial_h V} \right)^{\frac{1}{\sigma}} \right\}.$$

In particular, notice that

$$\left( \frac{h^{\gamma(\sigma-1)}}{\partial_k V - \rho \partial_h V} \right)^{\frac{1}{\sigma}} \leq Rk \quad \iff \quad \partial_k V - \rho \partial_h V \geq \frac{h^{\gamma(\sigma-1)}}{(Rk)^\sigma}.$$

Combining the two cases, it follows that

$$\begin{aligned} G(k, h, \partial_k V, \partial_h V) &= \sup_{c \in (0, Rk]} g(c; h, \partial_k V, \partial_h V) \\ &= \begin{cases} -Rk(\partial_k V - \rho \partial_h V) - \frac{1}{\sigma-1} \left( \frac{h^\gamma}{Rk} \right)^{\sigma-1}, & \partial_k V - \rho \partial_h V \leq h^{\gamma(\sigma-1)}(Rk)^{-\sigma}, \\ -\left( 1 + \frac{1}{\sigma-1} \right) [h^\gamma(\partial_k V - \rho \partial_h V)]^{1-\frac{1}{\sigma}}, & \partial_k V - \rho \partial_h V > h^{\gamma(\sigma-1)}(Rk)^{-\sigma}. \end{cases} \end{aligned} \quad (17)$$

Thus, the HJB equation in (16) can be rewritten as

$$\theta V(k, h) = \mathcal{L}_{k,h}V + G(k, h, \partial_k V(k, h), \partial_h V(k, h)) =: H_{\max}(k, h, \nabla V(k, h), D^2V(k, h)). \quad (18)$$

In the sequel, given a point  $P \in \mathbb{R}_{++}^2$  and positive constant  $r > 0$ , we denote by  $B_r(P)$  the open ball in  $\mathbb{R}^2$  centered at  $P$  of radius  $r$ , and by  $\overline{B}_r(P)$  its closure. For any  $n \in \mathbb{N}$ , let

$\mathcal{S}^n$  be the space of symmetric matrices in  $\mathbb{R}^{n \times n}$ .

Given  $A, B \in \mathcal{S}^n$ , the expression  $A \leq B$  means that  $B - A \in \mathcal{S}^n$  is positive semidefinite.

We conclude this section with two lemmas collecting some properties of the maximum value Hamiltonian  $H_{\max}$  in (18). In Subsection 5.2, such properties will be crucial to study the regularity of  $V$  via PDE techniques.

**Lemma 3.1.** *Let  $D \subset \mathbb{R}^2$  be an open and bounded set such that its closure  $\overline{D} \subset \mathbb{R}_{++}^2$ . Define the function*

$$\tilde{F} : \mathcal{S}^2 \times \mathbb{R}^2 \times \mathbb{R} \times \overline{D} \rightarrow \mathbb{R} \quad \text{by} \quad \tilde{F}(Q, p, v, (k, h)) = -\theta v + H_{\max}(k, h, p, Q). \quad (19)$$

*Then there exists a positive constant  $\bar{C} > 0$  such that, for every  $Q = (Q_{ij})_{i,j} \in \mathcal{S}^2$ ,  $(k, h) \in \overline{D}$ ,  $p = \begin{bmatrix} p_k \\ p_h \end{bmatrix}$ ,  $\bar{p} = \begin{bmatrix} \bar{p}_k \\ \bar{p}_h \end{bmatrix} \in \mathbb{R}^2$  and  $v, \bar{v} \in \mathbb{R}$ ,*

$$|\tilde{F}(Q, p, v, (k, h)) - \tilde{F}(Q, \bar{p}, \bar{v}, (k, h))| \leq \bar{C}(|p_k - \bar{p}_k| + |p_h - \bar{p}_h| + |v - \bar{v}|), \quad (20)$$

*and, for every  $\alpha \in (0, 1)$ ,*

$$\langle \tilde{F}(Q, p, v, \cdot) \rangle_D^\alpha \leq \bar{C} \left( 1 + \sum_{i,j=1}^2 |Q_{ij}| + |p_k| + |p_h| + |v| \right), \quad (21)$$

*where*

$$\langle f \rangle_D^\alpha = \sup_{\substack{(k,h) \in D \\ \delta > 0}} \delta^{-\alpha} \left( \sup_{D \cap B_\delta(k,h)} f - \inf_{D \cap B_\delta(k,h)} f \right), \quad \text{for a map } f : D \rightarrow \mathbb{R}.$$

**Lemma 3.2.** *Let  $D \subset \mathbb{R}_{++}^2$  be an open and bounded set such that its closure  $\overline{D} \subset \mathbb{R}_{++}^2$ . Define the function  $\tilde{F} : \mathcal{S}^2 \times \mathbb{R}^2 \times \mathbb{R} \times D \rightarrow \mathbb{R}$  as in Lemma 3.1. Then there is a constant  $\tilde{C} > 0$  such that*

$$\begin{aligned} \tilde{F}(Q, \alpha(k - \bar{k}, h - \bar{h}), v, (k, h)) - \tilde{F}(\bar{Q}, \alpha(k - \bar{k}, h - \bar{h}), v, (\bar{k}, \bar{h})) \\ \leq \tilde{C}(\alpha|(k - \bar{k}, h - \bar{h})|^2 + |(k - \bar{k}, h - \bar{h})|), \end{aligned} \quad (22)$$

for every  $\alpha > 0$ ,  $(k, h)$ ,  $(\bar{k}, \bar{h}) \in D$ ,  $v \in \mathbb{R}$  and  $Q, \bar{Q} \in \mathcal{S}^2$  satisfying

$$\begin{bmatrix} Q & 0 \\ 0 & -\bar{Q} \end{bmatrix} \leq 3\alpha \begin{bmatrix} I & -I \\ -I & I \end{bmatrix}. \quad (23)$$

The proofs of Lemmas 3.1 and 3.2 are postponed to Appendix A.

## 4 Properties of the Value function $V$

This section is devoted to the analysis of the value function  $V$  defined in (10). We will show, in particular, that  $V$  is finite valued and continuous, from which we deduce the dynamic programming principle in Proposition 4.5.

We start with a preliminary result concerning the homogeneity of  $V$  in  $\mathbb{R}_{++}^2$ .

**Lemma 4.1.** *For every  $\alpha > 0$ ,*

$$V(\alpha k_0, \alpha h_0) = \alpha^{(1-\gamma)(1-\sigma)} V(k_0, h_0), \quad (k_0, h_0) \in \mathbb{R}_{++}^2. \quad (24)$$

*Proof.* Fix  $\alpha > 0$  and an initial condition  $(k_0, h_0) \in \mathbb{R}_{++}^2$ . Recall that the definition of the sets  $\mathcal{A}(k_0, h_0)$  and  $\mathcal{A}(\alpha k_0, \alpha h_0)$  of admissible controls is given in (8). According to (13) in Lemma 2.1, these sets only depend on  $k_0$  and  $\alpha$ , i.e., they are independent of  $h_0$ . Moreover, by the explicit expression of the capital  $k$  in (3), it is straightforward to conclude that

$$\mathcal{A}(\alpha k_0, \alpha h_0) = \alpha \mathcal{A}(k_0, h_0).$$

Using the notation introduced in (5), by formula (4),

$$h_t^{\alpha h_0, 0, \alpha c} = \alpha h_t^{h_0, 0, c}, \quad t \geq 0, \mathbb{P} - \text{a.s.}, \text{ for every } c \in \mathcal{A}(\alpha k_0, \alpha h_0).$$

As a result, also using (9) and (10),

$$\begin{aligned} V(\alpha k_0, \alpha h_0) &= \sup_{c \in \mathcal{A}(\alpha k_0, \alpha h_0)} \mathbb{E} \left[ \int_0^\infty \frac{1}{1-\sigma} \left( \frac{(h_t^{\alpha h_0, 0, c})^\gamma}{c_t} \right)^{\sigma-1} e^{-\theta t} dt \right] \\ &= \sup_{c \in \mathcal{A}(k_0, h_0)} \mathbb{E} \left[ \int_0^\infty \frac{1}{1-\sigma} \left( \frac{(h_t^{\alpha h_0, 0, \alpha c})^\gamma}{\alpha c_t} \right)^{\sigma-1} e^{-\theta t} dt \right] \\ &= \sup_{c \in \mathcal{A}(k_0, h_0)} \mathbb{E} \left[ \int_0^\infty \frac{1}{1-\sigma} \left( \frac{(\alpha h_t^{h_0, 0, c})^\gamma}{\alpha c_t} \right)^{\sigma-1} e^{-\theta t} dt \right] = \alpha^{(1-\gamma)(1-\sigma)} V(k_0, h_0). \end{aligned}$$

The proof is then complete.  $\square$

Since the utility function  $J$  in (9) is nonpositive, by the definition (10),  $V \leq 0$  in  $\mathbb{R}_{++}^2$ , as well. Recalling our standing Assumption 1, the following proposition gives a lower bound for  $V$ . In particular, we deduce that  $V(k_0, h_0) \in (-\infty, 0]$  (i.e.,  $V(k_0, h_0)$  is finite) for every  $(k_0, h_0) \in \mathbb{R}_{++}^2$ .

**Proposition 4.2.** *There exists a positive constant  $C = C(\sigma, \gamma, \theta, \rho, \beta_1, \beta_2, B) > 0$  such that*

$$V(k_0, h_0) \geq -C \left( \left( \frac{h_0^\gamma}{k_0} \right)^{\sigma-1} + \frac{1}{k_0^{(1-\gamma)(\sigma-1)}} \right), \quad (k_0, h_0) \in \mathbb{R}_{++}^2. \quad (25)$$

*Proof.* Fix  $(k_0, h_0) \in \mathbb{R}_{++}^2$  and consider the control  $c_t = Bk_0\mathcal{E}(\beta_1 W_{1,t})$ ,  $t \geq 0$ . According to (14), the corresponding capital process  $k$  is

$$k_t = k_0\mathcal{E}(\beta_1 W_{1,t}) = \frac{c_t}{B} > 0, \quad t \geq 0.$$

Since  $B \leq R$  by Assumption 1, the constraint (7) is fulfilled. Thus, also recalling (13) in Lemma 2.1, we have  $c \in \mathcal{A}(k_0, h_0)$ . Using the explicit expressions of the habit process  $h$  in (4) and the Doléans-Dade exponential of a continuous semimartingale, by (10),

$$\begin{aligned} V(k_0, h_0) &\geq -\frac{(Bk_0)^{1-\sigma}}{\sigma-1} \mathbb{E} \left[ \int_0^\infty e^{-\theta t} e^{(1-\sigma) \left[ (\gamma\rho + \frac{1}{2}(\gamma\beta_2^2 - \beta_1^2))t + \beta_1 W_{1,t} - \gamma\beta_2 W_{2,t} \right]} \right. \\ &\quad \left. \times \left( h_0 + \rho Bk_0 \int_0^t e^{(\rho + \frac{1}{2}(\beta_2^2 - \beta_1^2))s + \beta_1 W_{1,s} - \beta_2 W_{2,s}} ds \right)^{\gamma(\sigma-1)} dt \right] \\ &=: -\frac{(Bk_0)^{1-\sigma}}{\sigma-1} \mathbb{E} \left[ \int_0^\infty \mathbf{I}_t \times \mathbf{II}_t dt \right]. \end{aligned}$$

By Tonelli's theorem and Hölder's inequality, it then follows that

$$V(k_0, h_0) \geq -\frac{(Bk_0)^{1-\sigma}}{\sigma-1} \int_0^\infty (\mathbb{E}[\mathbf{I}_t^p])^{\frac{1}{p}} (\mathbb{E}[\mathbf{II}_t^q])^{\frac{1}{q}} dt, \quad (26)$$

where we pick  $q > 2$  such that  $q\gamma(\sigma-1) > 1$ , and  $p > 1$  is its Hölder's conjugate, that is,  $p = q(q-1)^{-1}$ . In particular, notice that  $q > 2$  entails  $p < q$ . Since  $W_1$  and  $W_2$  are independent Brownian motions and, for every  $t > 0$ , the random variables  $e^{\beta_1 W_{1,t}}$  and  $e^{-\gamma\beta_2 W_{2,t}}$  have a log-normal distribution,

$$\mathbb{E}[\mathbf{I}_t^p] = \exp \left\{ p(1-\sigma) \left( \gamma\rho + \frac{1}{2}(\gamma\beta_2^2 - \beta_1^2) + \frac{\theta}{\sigma-1} + \frac{1}{2}p(1-\sigma)(\beta_1^2 + \gamma^2\beta_2^2) \right) t \right\}, \quad t \geq 0.$$

As for **II**, we compute, defining  $q^* = q\gamma(\sigma - 1) > 1$  and denoting by  $p^*$  its Hölder's conjugate, i.e.,  $p^* = q^*(q^* - 1)^{-1}$ ,

$$\begin{aligned}
\mathbb{E}[|\mathbf{II}_t|^q] &\leq 2^{q^*-1} \left( h_0^{q^*} + (\rho B k_0)^{q^*} \mathbb{E} \left[ \left( \int_0^t e^{(\rho + \frac{1}{2}(\beta_2^2 - \beta_1^2))s + \beta_1 W_{1,s} - \beta_2 W_{2,s}} ds \right)^{q^*} \right] \right) \\
&\leq 2^{q^*-1} \left( h_0^{q^*} + (\rho B k_0)^{q^*} \left( \int_0^t e^{p^*(\rho + \frac{1}{2}(\beta_2^2 - \beta_1^2))s} ds \right)^{\frac{q^*}{p^*}} \mathbb{E} \left[ \int_0^t e^{q^*(\beta_1 W_{1,s} - \beta_2 W_{2,s})} ds \right] \right) \\
&= 2^{q^*-1} \left( h_0^{q^*} + (\rho B k_0)^{q^*} \left( \int_0^t e^{p^*(\rho + \frac{1}{2}(\beta_2^2 - \beta_1^2))s} ds \right)^{\frac{q^*}{p^*}} \int_0^t e^{\frac{1}{2}(q^*)^2(\beta_1^2 + \beta_2^2)s} ds \right) \\
&\leq 2^{q^*-1} \left( h_0^{q^*} + C_1 k_0^{q^*} \left( \int_0^t e^{p^*(\rho + \frac{1}{2}(\beta_2^2 - \beta_1^2))s} ds \right)^{\frac{q^*}{p^*}} \exp \left\{ \frac{1}{2}(q^*)^2(\beta_1^2 + \beta_2^2)t \right\} \right), \quad t \geq 0,
\end{aligned}$$

for some constant  $C_1 = C_1(\rho, \beta_1, \beta_2, q^*, B) > 0$ . Here, in the second estimate we apply Hölder's inequality with exponents  $p^*$  and  $q^*$ , and in the equality, use the independence of  $W_1$  and  $W_2$ , together with Fubini's theorem and the properties of the log-normal distribution. Plugging the two previous estimates on **I** and **II** into (26) and recalling that  $q^*/q = \gamma(\sigma - 1)$ , using also that the power  $x^{\frac{1}{q}}$  is subadditive on  $[0, \infty)$ , we obtain

$$\begin{aligned}
V(k_0, h_0) &\geq -C_2 k_0^{1-\sigma} \int_0^\infty \exp \left\{ (1-\sigma) \left( \gamma\rho + \frac{1}{2}(\gamma\beta_2^2 - \beta_1^2) + \frac{\theta}{\sigma-1} + \frac{1}{2}p(1-\sigma)(\beta_1^2 + \gamma^2\beta_2^2) \right) t \right\} \\
&\quad \times \left( h_0^{\gamma(\sigma-1)} + k_0^{\gamma(\sigma-1)} \left( \int_0^t e^{p^*(\rho + \frac{1}{2}(\beta_2^2 - \beta_1^2))s} ds \right)^{\frac{\gamma(\sigma-1)}{p^*}} \exp \left\{ \frac{1}{2}q\gamma^2(\sigma-1)^2(\beta_1^2 + \beta_2^2)t \right\} \right) dt,
\end{aligned} \tag{27}$$

for a constant  $C_2 = C_2(\sigma, q, \rho, \beta_1, \beta_2, q^*, B) > 0$ .

From now on, we denote by  $C_i$ ,  $i = 3, 4, \dots$ , positive constant possibly dependent on  $(\sigma, q, \rho, \beta_1, \beta_2, p^*, q^*, B)$ . We now argue that the integral on the right-hand side of (27) converges thanks to Assumption 1. To do this, we distinguish two cases.

If  $\rho + \frac{1}{2}(\beta_2^2 - \beta_1^2) \leq 0$ , then

$$\int_0^t e^{p^*(\rho + \frac{1}{2}(\beta_2^2 - \beta_1^2))s} ds \leq \max\{1, t\}, \quad t \geq 0.$$

Hence, recalling that  $q > p$  and  $\gamma \in [0, 1)$ , we deduce from (27) that

$$V(k_0, h_0) \geq -C_3 \left( \left( \frac{h_0^\gamma}{k_0} \right)^{\sigma-1} + \frac{1}{k_0^{(1-\gamma)(\sigma-1)}} \right) \int_0^\infty \max\{1, t\} \exp\{(\sigma-1)e_1(q)t\} dt,$$

where  $e_1(q) := -\gamma\rho - \frac{1}{2}(\gamma\beta_2^2 - \beta_1^2) - \frac{\theta}{\sigma-1} + q(\sigma-1)(\beta_1^2 + \gamma^2\beta_2^2)$ .

Condition (11) in Assumption 1(i) enables us to find a  $q > \max\left\{2, \frac{1}{\gamma(\sigma-1)}\right\}$  such that  $e_1(q) < 0$ , whence (25).

In the case  $\rho + \frac{1}{2}(\beta_2^2 - \beta_1^2) > 0$ , using again the fact that  $q > p$  and  $\gamma \in [0, 1)$ , from (27) we obtain the estimate

$$V(k_0, h_0) \geq -C_4 \left( \left( \frac{h_0^\gamma}{k_0} \right)^{\sigma-1} + \frac{1}{k_0^{(1-\gamma)(\sigma-1)}} \right) \int_0^\infty \exp\{(\sigma-1)e_2(q)t\} dt,$$

where  $e_2(q) := \frac{1}{2}\beta_1^2(1-\gamma) - \frac{\theta}{\sigma-1} + q(\sigma-1)(\beta_1^2 + \gamma^2\beta_2^2)$ .

As before, Condition (12) in Assumption 1(ii) enables us to find a  $q > \max\left\{2, \frac{1}{\gamma(\sigma-1)}\right\}$  such that  $e_2(q) < 0$ , whence (25). The proof is now complete.  $\square$

The following proposition shows the monotonicity of the value function  $V$  in  $\mathbb{R}_{++}^2$  with respect to its variables.

**Proposition 4.3.** *The value function  $V$  is nondecreasing in the capital variable  $k$  and nonincreasing in the habit variable  $h$ .*

*Proof.* We start by studying the monotonicity of  $V$  with respect to  $k$ . Fix  $h_0 \in (0, \infty)$  and  $k_1, k_2 \in (0, \infty)$  such that  $k_1 < k_2$ . By the explicit expression of the capital process in (3),  $\mathcal{A}(k_1, h_0) \subset \mathcal{A}(k_2, h_0)$ . Since the utility function  $J$  in (9) does not depend on the capital  $k$ , we deduce that  $V(k_1, h_0) \leq V(k_2, h_0)$ .

As for the monotonicity in  $h$ , fix  $k_0 \in (0, \infty)$  and  $h_1, h_2 \in (0, \infty)$  such that  $h_1 < h_2$ . By (13) in Lemma 2.1,  $\mathcal{A}(k_0, h_1) = \mathcal{A}(k_0, h_2) = \mathcal{A}(k_0)$ . Moreover, from the explicit expression in (4), we have  $h^{h_1, 0, c} \leq h^{h_2, 0, c}$ ,  $\mathbb{P} \otimes dt$ -a.e., for every  $c \in \mathcal{A}(k_0)$ . As a result,

$$J(k_0, h_1; c) \geq J(k_0, h_2; c), \quad c \in \mathcal{A}(k_0).$$

Taking the supremum over  $\mathcal{A}(k_0)$  yields  $V(k_0, h_1) \geq V(k_0, h_2)$ , which completes the proof.  $\square$

Combining the homogeneity in Lemma 4.1 and the monotonicity in Proposition 4.3, under an additional requirement on the parameters  $\gamma$  and  $\sigma$  we now demonstrate the continuity of  $V$  in  $\mathbb{R}_{++}^2$ .

**Theorem 4.4.** *Suppose that*

$$\gamma(\sigma - 1) \leq 1. \quad (28)$$

*Then the value function  $V: \mathbb{R}_{++}^2 \rightarrow (-\infty, 0]$  is continuous.*

*Proof.* Fix an arbitrary point  $P_0 = (k_0, h_0) \in \mathbb{R}_{++}^2$ . Note that, by Lemma 4.1, the separate continuity of  $V(k_0, \cdot)$  at  $h_0$ , i.e.,

$$\lim_{h \rightarrow h_0} V(k_0, h) = V(k_0, h_0), \quad (29)$$

is sufficient to deduce the joint continuity of  $V$  at  $P_0$ . Indeed, given two sequences  $(h_n)_n, (k_n)_n \subset \mathbb{R}_+$  which converge to  $h_0$  and  $k_0$ , respectively, by (24),

$$V(k_n, h_n) = V\left(\frac{k_n}{k_0}k_0, \frac{k_n}{k_0}\frac{k_0}{k_n}h_n\right) = \left(\frac{k_n}{k_0}\right)^{(1-\gamma)(1-\sigma)} V\left(k_0, \frac{k_0}{k_n}h_n\right) \xrightarrow{n \rightarrow \infty} V(k_0, h_0),$$

where we use (29) to pass to the limit.

We then focus on (29). Since the value function  $V$  is nonincreasing with respect to  $h$  by Proposition 4.3,

$$\lim_{h \rightarrow h_0^-} V(k_0, h) \geq V(k_0, h_0) \geq \lim_{h \rightarrow h_0^+} V(k_0, h).$$

Thus, we only need to show the reversed chain of inequalities, starting from

$$\lim_{h \rightarrow h_0^+} V(k_0, h) \geq V(k_0, h_0). \quad (30)$$

Consider  $(h_n)_n \subset (h_0, \infty)$  such that  $\lim_{n \rightarrow \infty} h_n = h_0$ . For every  $\epsilon > 0$ , by (10), there exists an  $\epsilon$ -optimal control  $\bar{c} \in \mathcal{A}(k_0, h_0)$  satisfying

$$V(k_0, h_0) \leq \mathbb{E}\left[\int_0^\infty \frac{1}{1-\sigma} e^{-\theta t} \frac{1}{\bar{c}_t^{\sigma-1}} (h_t^{h_0, 0, \bar{c}})^{\gamma(\sigma-1)} dt\right] + \epsilon.$$

Since  $\bar{c} \in \mathcal{A}(k_0, h_n)$  by Lemma 2.1,

$$V(k_0, h_n) - V(k_0, h_0) \geq \frac{1}{1-\sigma} \mathbb{E}\left[\int_0^\infty \frac{e^{-\theta t}}{\bar{c}_t^{\sigma-1}} \left( (h_t^{h_n, 0, \bar{c}})^{\gamma(\sigma-1)} - (h_t^{h_0, 0, \bar{c}})^{\gamma(\sigma-1)} \right) dt\right] - \epsilon,$$

which holds for every  $n \in \mathbb{N}$ . The explicit expression in (4) implies that  $h^{h_n, 0, \bar{c}}$  converges to  $h^{h_0, 0, \bar{c}}$  locally uniformly in  $[0, \infty)$ ,  $\mathbb{P}$ -a.s. Therefore, also observing that  $h^{h_n, 0, \bar{c}} \geq h^{h_0, 0, \bar{c}}$ ,  $n \in \mathbb{N}$ ,  $\mathbb{P} \otimes dt$ -a.e., Fatou's lemma yields

$$\lim_{n \rightarrow \infty} V(k_0, h_n) \geq V(k_0, h_0) - \epsilon.$$

This estimate gives (30) as  $\epsilon$  is arbitrary.

Concerning  $\lim_{h \rightarrow h_0^-} V(k_0, h)$ , we suppose by contradiction that

$$\lim_{h \rightarrow h_0^-} V(k_0, h) > V(k_0, h_0). \quad (31)$$

By (10), we consider a sequence  $(h_n)_n \subset (0, h_0)$  which converges to  $h_0$  and corresponding admissible controls  $c_n \in \mathcal{A}(k_0, h_n)$  such that

$$J(k_0, h_n; c_n) > V(k_0, h_n) - \frac{1}{n}, \quad n \in \mathbb{N}, \quad \text{whence, by (31), } \lim_{n \rightarrow \infty} J(k_0, h_n; c_n) > V(k_0, h_0). \quad (32)$$

Note that (32) combined with (4) and (9) entails

$$\sup_{n \in \mathbb{N}} \mathbb{E} \left[ \int_0^\infty \frac{e^{-\theta t}}{c_{n,t}^{\sigma-1}} \mathcal{E}(-\rho t + \beta_2 W_{2,t})^{\gamma(1-\sigma)} \left( 1 + \int_0^t \frac{c_{n,s}}{\mathcal{E}(-\rho s + \beta_2 W_{2,s})} ds \right)^{\gamma(\sigma-1)} dt \right] < \infty. \quad (33)$$

Fix a generic  $n \in \mathbb{N}$ . Since  $h_n < h_0$ , the modified control  $\frac{h_n}{h_0} c_n$  belongs to  $\mathcal{A}(k_0, h_n)$ . We then consider  $J(k_0, h_n; \frac{h_n}{h_0} c_n)$  and aim to estimate the difference

$$\Delta_n := J(k_0, h_n; \frac{h_n}{h_0} c_n) - J(k_0, h_n; c_n).$$

In particular, by (9) we compute

$$\begin{aligned} |\Delta_n| &\leq \frac{1}{\sigma-1} \left( \left( \frac{h_0}{h_n} \right)^{\sigma-1} - 1 \right) \mathbb{E} \left[ \int_0^\infty \frac{e^{-\theta t}}{c_{n,t}^{\sigma-1}} \left( h_t^{h_n, 0, \frac{h_n}{h_0} c_n} \right)^{\gamma(\sigma-1)} dt \right] \\ &\quad + \frac{1}{\sigma-1} \mathbb{E} \left[ \int_0^\infty \frac{e^{-\theta t}}{c_{n,t}^{\sigma-1}} \left| \left( h_t^{h_n, 0, \frac{h_n}{h_0} c_n} \right)^{\gamma(\sigma-1)} - \left( h_t^{h_n, 0, c_n} \right)^{\gamma(\sigma-1)} \right| dt \right] =: \mathbf{I}_n + \mathbf{II}_n. \end{aligned} \quad (34)$$

By (4),

$$h_t^{h_n, 0, \frac{h_n}{h_0} c_n} = \frac{h_n}{h_0} h_t^{h_0, 0, c_n}, \quad t \geq 0, \quad (35)$$

whence, considering also that the sequence  $(J(k_0, h_0; c_n))_n$  is bounded by (33),

$$\mathbf{I}_n \leq \left( \left( \frac{h_0}{h_n} \right)^{\sigma-1} - 1 \right) \left( \frac{h_n}{h_0} \right)^{\gamma(\sigma-1)} \sup_{n \in \mathbb{N}} |J(k_0, h_0; c_n)| < \frac{1}{2n}, \quad \text{for } n \text{ sufficiently large.}$$

As for  $\mathbf{II}_n$ , notice that the power function  $x^{\gamma(\sigma-1)}$  is  $\gamma(\sigma-1)$ -Hölder continuous in  $[0, \infty)$  thanks to the assumption (28). Denoting by  $C_1$  its Hölder-continuity constant, by (4) and (33) we deduce that

$$\begin{aligned} \mathbf{II}_n &\leq \frac{C_1}{\sigma-1} \rho^{\gamma(\sigma-1)} \left( 1 - \frac{h_n}{h_0} \right)^{\gamma(\sigma-1)} \\ &\quad \times \mathbb{E} \left[ \int_0^\infty \frac{e^{-\theta t}}{c_{n,t}^{\sigma-1}} \mathcal{E}(-\rho t + \beta_2 W_{2,t})^{\gamma(\sigma-1)} \left( \int_0^t \frac{c_{n,s}}{\mathcal{E}(-\rho s + \beta_2 W_{2,s})} ds \right)^{\gamma(\sigma-1)} dt \right] \\ &\leq C_1 \rho^{\gamma(\sigma-1)} \left( 1 - \frac{h_n}{h_0} \right)^{\gamma(\sigma-1)} \sup_{n \in \mathbb{N}} |J(k_0, h_0; c_n)| \leq \frac{1}{2n}, \quad \text{for } n \text{ sufficiently large.} \end{aligned}$$

Plugging the estimates we have just established on  $\mathbf{I}_n$  and  $\mathbf{II}_n$  into (34) gives  $|\Delta_n| < \frac{1}{n}$ , for  $n$  sufficiently large. Thus, by (32),

$$\lim_{n \rightarrow \infty} J\left(k_0, h_n; \frac{h_n}{h_0} c_n\right) = \lim_{n \rightarrow \infty} J(k_0, h_n; c_n) + \lim_{n \rightarrow \infty} \Delta_n = \lim_{n \rightarrow \infty} J(k_0, h_n; c_n) > V(k_0, h_0).$$

At the same time, recalling (9) and (35),

$$J\left(k_0, h_n; \frac{h_n}{h_0} c_n\right) = \left( \frac{h_0}{h_n} \right)^{(1-\gamma)(\sigma-1)} J(k_0, h_0; c_n) \leq \left( \frac{h_0}{h_n} \right)^{(1-\gamma)(\sigma-1)} V(k_0, h_0),$$

whence  $\lim_{n \rightarrow \infty} J\left(k_0, h_n; \frac{h_n}{h_0} c_n\right) \leq V(k_0, h_0)$ . This is a contradiction demonstrating that (31) cannot hold. Consequently,  $V(k_0, \cdot)$  is also left-continuous at  $h_0$  and the proof is complete.  $\square$

Proposition 4.2 and Theorem 4.4 enable us to establish the dynamic programming principle, as shown in the next result. Its proof, which follows standard arguments (see, e.g., Fleming and Soner [2006], Gassiat et al. [2014], Ishii and Loreti [2002]) is omitted for brevity.

**Proposition 4.5.** *Suppose that (28) holds. Then the value function  $V$  satisfies the dynamic programming equation, that is, for every  $(k, h) \in \mathbb{R}_{++}^2$  and stopping time  $\tau$  (possibly depending on  $c \in \mathcal{A}(k, h)$ ), the following functional equation holds:*

$$V(k, h) = \sup_{c \in \mathcal{A}(k, h)} \mathbb{E} \left[ \int_0^\tau \frac{e^{-\theta t}}{1-\sigma} \left( \frac{h_t^\gamma}{c_t} \right)^{\sigma-1} dt + e^{-\theta \tau} V(k_\tau, h_\tau) \right]. \quad (36)$$

## 5 Solving the HJB equation

In this section, we show that the value function  $V$  solves the HJB equation (18) both in the viscosity and the classical sense. In particular, we demonstrate that  $V$  is a regular map in the class  $C^2(\mathbb{R}_{++}^2; \mathbb{R})$ .

### 5.1 $V$ is a viscosity solution of the HJB equation

We begin this subsection by recalling the definition of (continuous) *viscosity solutions* of the HJB equation (18).

**Definition 1.** Let  $D$  be an open set in  $\mathbb{R}_{++}^2$ . A continuous function  $v : D \rightarrow \mathbb{R}$  is a *viscosity subsolution* (resp., *viscosity supersolution*) of (18) in  $D$  if, for any  $\psi \in C^2(D; \mathbb{R})$  and any local maximum point  $(k_M, h_M) \in D$  (resp., local minimum point  $(k_m, h_m) \in D$ ) of  $v - \psi$ , the following inequality holds:

$$\begin{aligned} & \theta v(k_M, h_M) - H_{\max}(k_M, h_M, \nabla \psi(k_M, h_M), D^2 \psi(k_M, h_M)) \leq 0 \\ & \text{(resp., } \theta v(k_m, h_m) - H_{\max}(k_m, h_m, \nabla \psi(k_m, h_m), D^2 \psi(k_m, h_m)) \geq 0). \end{aligned}$$

A continuous function  $v : D \rightarrow \mathbb{R}$  is a *viscosity solution* of (18) in  $D$  if it is both a viscosity subsolution and a viscosity supersolution in  $D$ .

In the next theorem, adapting in a nontrivial way some standard arguments in, e.g., Choulli et al. [2003], Di Giacinto et al. [2011], we prove that the value function  $V$  defined in (10) solves (18) on  $\mathbb{R}_{++}^2$  in the viscosity sense. Recall that, given a point  $P \in \mathbb{R}_{++}^2$  and positive constant  $r > 0$ , we denote by  $B_r(P)$  the open ball in  $\mathbb{R}^2$  centered at  $P$  of radius  $r$ , and by  $\overline{B}_r(P)$  its closure.

**Theorem 5.1.** *Assume (28). Then the value function  $V$  is a viscosity solution of the HJB equation (18) in  $\mathbb{R}_{++}^2$ .*

*Proof.* We structure the proof in two steps. More specifically, in the first (resp., second) step, we demonstrate that  $V$  is a viscosity supersolution (resp., viscosity subsolution) of (18) on  $\mathbb{R}_{++}^2$ .

Step I. Consider  $\psi \in C^2(\mathbb{R}_{++}^2; \mathbb{R})$  and a local minimum point  $(k_m, h_m) \in \mathbb{R}_{++}^2$  for  $V - \psi$ . Without loss of generality, we suppose that  $V(k_m, h_m) = \psi(k_m, h_m)$ , hence there exists  $\delta > 0$  such that  $V(k, h) \geq \psi(k, h)$ ,  $(k, h) \in B_\delta(k_m, h_m) \subset \mathbb{R}_{++}^2$ .

Fix a constant  $\bar{c} \in (0, Rk_m)$ . Recalling the notation in (5), we define the stopping time

$$\bar{\tau} = \inf \left\{ t \geq 0 : k_t^{k_m, 0, \bar{c}} \leq \frac{\bar{c}}{R} \right\};$$

observe that  $\bar{\tau} > 0$ ,  $\mathbb{P}$ -a.s., by the continuity of  $k^{k_m, 0, \bar{c}}$ . We now employ  $\bar{\tau}$  to construct the following positive progressively measurable process  $c_\nu^*$ ,  $\nu > 0$ :

$$c_{\nu, t}^* = \begin{cases} \bar{c}, & t \leq \bar{\tau}, \\ \nu (\mathcal{E}(B\bar{\tau} + \beta_1 W_{1, \bar{\tau}}))^{-1} \mathcal{E}(\beta_1 W_{1, t}), & t > \bar{\tau}. \end{cases} \quad (37)$$

Notice that, choosing  $\nu$  sufficiently small,  $c_\nu^* \in \mathcal{A}(k_m, h_m)$ . Indeed, by the definition of  $\bar{\tau}$  and (3), on the event  $\{\tau < \infty\}$

$$\frac{\bar{c}}{R} (\mathcal{E}(B\bar{\tau} + \beta_1 W_{1, \bar{\tau}}))^{-1} = (\mathcal{E}(B\bar{\tau} + \beta_1 W_{1, \bar{\tau}}))^{-1} k_\tau^{k_m, 0, c_\nu^*} = k_m - \bar{c} \int_0^{\bar{\tau}} \frac{ds}{\mathcal{E}(Bs + \beta_1 W_{1, s})}.$$

Consequently, for every  $t > \bar{\tau}$ , using (37) we have

$$\begin{aligned} k_t^{k_m, 0, c_\nu^*} &= \mathcal{E}(Bt + \beta_1 W_{1, t}) \left( k_m - \bar{c} \int_0^{\bar{\tau}} \frac{ds}{\mathcal{E}(Bs + \beta_1 W_{1, s})} - \int_{\bar{\tau}}^t \frac{c_{\nu, s}^*}{\mathcal{E}(Bs + \beta_1 W_{1, s})} ds \right) \\ &= (\mathcal{E}(B\bar{\tau} + \beta_1 W_{1, \bar{\tau}}))^{-1} \mathcal{E}(Bt + \beta_1 W_{1, t}) \left( \frac{\bar{c}}{R} - \frac{\nu}{B} (e^{-B\bar{\tau}} - e^{-Bt}) \right), \end{aligned}$$

which shows that the constraints (6) and (7) are satisfied as  $\nu$  approaches 0.

We then fix  $\nu^* > 0$  such that  $c^* := c_{\nu^*}^* \in \mathcal{A}(k_m, h_m)$ , and define the stopping times

$$\tau_\delta = \inf \left\{ t \geq 0 : |k_t^{k_m, 0, c^*} - k_m|^2 + |h_t^{h_m, 0, c^*} - h_m|^2 \geq \delta^2 \right\}, \quad \bar{\tau}_{\delta, \epsilon} = \tau_\delta \wedge \epsilon \wedge \bar{\tau}.$$

Since trivially  $\mathbb{E}[\bar{\tau}_{\delta, \epsilon}] \leq \epsilon < \infty$ , we can apply Dynking's formula to the function  $(t, k, h) \mapsto e^{-\theta t} \psi(k, h)$  to obtain, denoting by  $h^*$  (resp.,  $k^*$ ) the process  $h^{h_m, 0, c^*}$  (resp.,  $k^{k_m, 0, c^*}$ ) to make the notation shorter,

$$\begin{aligned} \mathbb{E} \left[ e^{-\theta \bar{\tau}_{\delta, \epsilon}} \psi(k_{\bar{\tau}_{\delta, \epsilon}}^*, h_{\bar{\tau}_{\delta, \epsilon}}^*) \right] - \psi(k_m, h_m) &= \mathbb{E} \left[ \int_0^{\bar{\tau}_{\delta, \epsilon}} e^{-\theta t} \left( -\theta \psi(k_t^*, h_t^*) + (B(k_t^* - c_t^*)) \partial_k \psi(k_t^*, h_t^*) \right. \right. \\ &\quad \left. \left. + \rho(c_t^* - h_t^*) \partial_h \psi(k_t^*, h_t^*) + \frac{1}{2} \beta_1^2 (k_t^*)^2 \partial_{kk} \psi(k_t^*, h_t^*) + \frac{1}{2} \beta_2^2 (h_t^*)^2 \partial_{hh} \psi(k_t^*, h_t^*) \right) dt \right]. \quad (38) \end{aligned}$$

We observe that, since  $\bar{\tau}_{\delta, \epsilon} \leq \tau_\delta$ ,

$$e^{-\theta \bar{\tau}_{\delta, \epsilon}} V(k_{\bar{\tau}_{\delta, \epsilon}}^*, h_{\bar{\tau}_{\delta, \epsilon}}^*) - V(k_m, h_m) \geq e^{-\theta \bar{\tau}_{\delta, \epsilon}} \psi(k_{\bar{\tau}_{\delta, \epsilon}}^*, h_{\bar{\tau}_{\delta, \epsilon}}^*) - \psi(k_m, h_m), \quad \mathbb{P} - \text{a.s.}$$

Hence, by the dynamic programming principle in Proposition 4.5, we deduce from (38) that

$$\begin{aligned} 0 \geq \mathbb{E} \left[ \int_0^{\bar{\tau}_{\delta, \epsilon}} e^{-\theta t} \left( -\theta \psi(k_t^*, h_t^*) + \frac{1}{1 - \sigma} \left( \frac{(h_t^*)^\gamma}{\bar{c}} \right)^{\sigma-1} + (B(k_t^* - \bar{c})) \partial_k \psi(k_t^*, h_t^*) \right. \right. \\ \left. \left. + \rho(\bar{c} - h_t^*) \partial_h \psi(k_t^*, h_t^*) + \frac{1}{2} \beta_1^2 (k_t^*)^2 \partial_{kk} \psi(k_t^*, h_t^*) + \frac{1}{2} \beta_2^2 (h_t^*)^2 \partial_{hh} \psi(k_t^*, h_t^*) \right) dt \right], \end{aligned}$$

where we also use the fact that, by (37),  $c^* = \bar{c}$  in  $[0, \bar{\tau}_{\delta, \epsilon}]$ , as  $\bar{\tau}_{\delta, \epsilon} \leq \bar{\tau}$ . Notice that  $\tau_\delta > 0$ ,  $\mathbb{P}$ -a.s., by the continuity of  $h^*$  and  $k^*$ . Recalling that the same holds true for  $\bar{\tau}$ , too, for  $\mathbb{P}$ -a.e.  $\omega \in \Omega$ ,

$$\bar{\tau}_{\delta, \epsilon}(\omega) = \epsilon, \quad \text{when } \epsilon \text{ is sufficiently small.}$$

Therefore, dividing by  $\epsilon$  the previous estimate and taking the limit as  $\epsilon \rightarrow 0$ , by the fundamental theorem of calculus and dominated convergence,

$$\begin{aligned} 0 &\geq -\theta\psi(k_m, h_m) + \frac{1}{1-\sigma} \left( \frac{h_m^\gamma}{\bar{c}} \right)^{\sigma-1} + (B(k_m - \bar{c})\partial_k\psi(k_m, h_m) + \rho(\bar{c} - h_m))\partial_h\psi(k_m, h_m) \\ &\quad + \frac{1}{2}\beta_1^2 k_m^2 \partial_{kk}\psi(k_m, h_m) + \frac{1}{2}\beta_2^2 h_m^2 \partial_{hh}\psi(k_m, h_m) \\ &= -\theta\psi(k_m, h_m) + \mathcal{L}_{k_m, h_m}\psi + g(\bar{c}; h_m, \partial_k\psi(k_m, h_m), \partial_h\psi(k_m, h_m)), \end{aligned}$$

where the last equality employs the notation in (16). Since  $\bar{c}$  is chosen arbitrarily in  $(0, Rk_m)$  and  $g(\cdot; h_m, \partial_k\psi(k_m, h_m), \partial_h\psi(k_m, h_m))$  is continuous at  $Rk_m$ , by taking the sup for  $\bar{c} \in (0, Rk_m]$  we conclude that

$$\theta V(k_m, h_m) - H_{\max}(k_m, h_m, \nabla\psi(k_m, h_m), D^2\psi(k_m, h_m)) \geq 0,$$

hence  $V$  is a viscosity supersolution of (18).

*Step II.* Consider  $\psi \in C^2(\mathbb{R}_{++}^2; \mathbb{R})$  and a local maximum point  $(k_M, h_M) \in \mathbb{R}_{++}^2$  for  $V - \psi$ . Without loss of generality, we assume  $V(k_M, h_M) = \psi(k_M, h_M)$ , hence there exists a  $\delta > 0$  such that  $V(k, h) \leq \psi(k, h)$ ,  $(k, h) \in B_\delta(k_M, h_M) \subset \mathbb{R}_{++}^2$ . We suppose by contradiction that  $V$  is *not* a viscosity subsolution of (18); by Definition 1, this means that there exists  $\eta_1 > 0$  such that

$$\eta_1 < \theta V(k_M, h_M) - \mathcal{L}_{k_M, h_M}\psi - G(k_M, h_M, \partial_k\psi(k_M, h_M), \partial_h\psi(k_M, h_M)).$$

By the continuity of  $V$ , see Theorem 4.4, since  $G$  in (17) is jointly continuous in its arguments and  $\psi \in C^2$ , we can find an  $\epsilon \in (0, \delta)$  such that, for any  $(k, h) \in B_\epsilon(k_M, h_M)$ ,

$$\frac{\eta_1}{2} < \theta V(k, h) - \mathcal{L}_{k, h}\psi + c(\partial_k\psi(k, h) - \rho\partial_h\psi(k, h)) + \frac{1}{\sigma-1} \frac{h^{\gamma(\sigma-1)}}{c^{\sigma-1}}, \quad c \in (0, Rk]. \quad (39)$$

For any admissible control  $\bar{c} \in \mathcal{A}(k_M, h_M)$ , we define the stopping time

$$\tau_\epsilon = \inf \left\{ t \geq 0 : \left( k_t^{k_M, 0, \bar{c}}, h_t^{h_M, 0, \bar{c}} \right) \notin B_\epsilon(k_M, h_M) \right\};$$

notice that  $(k^{k_M, 0, \bar{c}}, h^{h_M, 0, \bar{c}}) \in B_\delta(k_M, h_M)$  in the random interval  $[0, \tau_\epsilon]$  as  $\epsilon < \delta$ . Using  $\tau_\epsilon$ , from (39) we compute, denoting by  $\bar{h}$  (resp.,  $\bar{k}$ ) the process  $h^{h_M, 0, \bar{c}}$  (resp.,  $k^{k_M, 0, \bar{c}}$ ) to shorten the notation,

$$\begin{aligned} & \frac{\eta_1}{2} \mathbb{E} \left[ \int_0^{\tau_\epsilon \wedge 1} e^{-\theta t} dt \right] - \mathbb{E} \left[ \int_0^{\tau_\epsilon \wedge 1} \frac{e^{-\theta t}}{\sigma - 1} \left( \frac{\bar{h}_t^\gamma}{\bar{c}_t} \right)^{\sigma-1} dt \right] \\ & < \mathbb{E} \left[ \int_0^{\tau_\epsilon \wedge 1} e^{-\theta t} \left( \theta \psi(\bar{k}_t, \bar{h}_t) - \mathcal{L}_{\bar{k}_t, \bar{h}_t} \psi + \bar{c}_t (\partial_k \psi(\bar{k}_t, \bar{h}_t) - \rho \partial_h \psi(\bar{k}_t, \bar{h}_t)) \right) dt \right]. \end{aligned} \quad (40)$$

Since  $\mathbb{E}[\tau_\epsilon \wedge 1] \leq 1 < \infty$ , an application Dynkin's formula as in Step I yields

$$\begin{aligned} & \mathbb{E} \left[ e^{-\theta(\tau_\epsilon \wedge 1)} \psi(\bar{k}_{\tau_\epsilon \wedge 1}, \bar{h}_{\tau_\epsilon \wedge 1}) \right] - \psi(k_M, h_M) \\ & = \mathbb{E} \left[ \int_0^{\tau_\epsilon \wedge 1} e^{-\theta t} \left( -\theta \psi(\bar{k}_t, \bar{h}_t) + \mathcal{L}_{\bar{k}_t, \bar{h}_t} \psi - \bar{c}_t (\partial_k \psi(\bar{k}_t, \bar{h}_t) - \rho \partial_h \psi(\bar{k}_t, \bar{h}_t)) \right) dt \right]. \end{aligned} \quad (41)$$

Therefore, combining (40) and (41),

$$V(k_M, h_M) \geq \mathbb{E} \left[ \int_0^{\tau_\epsilon \wedge 1} \frac{e^{-\theta t}}{1 - \sigma} \left( \frac{\bar{h}_t^\gamma}{\bar{c}_t} \right)^{\sigma-1} dt + e^{-\theta(\tau_\epsilon \wedge 1)} V(\bar{k}_{\tau_\epsilon \wedge 1}, \bar{h}_{\tau_\epsilon \wedge 1}) \right] \quad (42)$$

$$+ \frac{\eta_1}{2} \mathbb{E} \left[ \int_0^{\tau_\epsilon \wedge 1} e^{-\theta t} dt \right]. \quad (43)$$

We now focus on the last addend in (43). Inspired by [Choulli et al., 2003, Equation (2.49)], we define the  $C^2$ -function  $w: \bar{B}_\epsilon(k_M, h_M) \rightarrow (-\infty, 0]$  by

$$w(k, h) = \frac{1}{C} \left( (k - k_M)^2 + (h - h_M)^2 - \epsilon^2 \right), \quad (k, h) \in \bar{B}_\epsilon(k_M, h_M),$$

where  $C \in \mathbb{R}_+$  is the positive constant given by

$$\begin{aligned} C := & \sup_{(k, h) \in \bar{B}_\epsilon(k_M, h_M)} \sup_{c \in (0, Rk]} \left( 2B|k - c||k - k_M| + 2\rho|c - h||h - h_M| + \beta_1^2 k^2 + \beta_2^2 h^2 \right. \\ & \left. + \theta(|k - k_M|^2 + |h - h_M|^2 + \epsilon^2) \right). \end{aligned}$$

Thanks to this choice of  $C$ , employing again Dynkin's formula with the function  $(t, k, h) \mapsto$

$e^{-\theta t}w(k, h)$ , we infer that

$$\begin{aligned}
& \mathbb{E} \left[ e^{-\theta(\tau_\epsilon \wedge 1)} w(\bar{k}_{\tau_\epsilon \wedge 1}, \bar{h}_{\tau_\epsilon \wedge 1}) \right] - w(k_M, h_M) \\
&= \frac{1}{C} \mathbb{E} \left[ \int_0^{\tau_\epsilon \wedge 1} e^{-\theta t} \left( -\theta w(\bar{k}_t, \bar{h}_t) + 2 \left[ \frac{B\bar{k}_t - \bar{c}_t}{\rho(\bar{c}_t - \bar{h}_t)} \right]^\top \begin{bmatrix} \bar{k}_t - k_M \\ \bar{h}_t - h_M \end{bmatrix} + \beta_1^2 \bar{k}_t^2 + \beta_2^2 \bar{h}_t^2 \right) dt \right] \\
&\leq \mathbb{E} \left[ \int_0^{\tau_\epsilon \wedge 1} e^{-\theta t} dt \right]. \tag{44}
\end{aligned}$$

Since  $w \equiv 0$  on  $\partial B_\epsilon(k_M, h_M)$  and  $w \geq -\frac{1}{C}\epsilon^2$  in its domain, by the continuity of  $\bar{k}$  and  $\bar{h}$  we have

$$\begin{aligned}
\mathbb{E} \left[ e^{-\theta(\tau_\epsilon \wedge 1)} w(\bar{h}_{\tau_\epsilon \wedge 1}, \bar{h}_{\tau_\epsilon \wedge 1}) \right] &= \mathbb{E} \left[ e^{-\theta\tau_\epsilon} w(\bar{k}_{\tau_\epsilon}, \bar{h}_{\tau_\epsilon}) 1_{\{\tau_\epsilon \leq 1\}} \right] + \mathbb{E} \left[ e^{-\theta} w(\bar{k}_1, \bar{h}_1) 1_{\{\tau_\epsilon > 1\}} \right] \\
&\geq 0 - \frac{1}{C}\epsilon^2 e^{-\theta} \mathbb{P}(\tau_\epsilon > 1) \geq -\frac{1}{C}\epsilon^2 e^{-\theta}.
\end{aligned}$$

Consequently, noticing also that  $w(k_M, h_M) = -\frac{1}{C}\epsilon^2$ , (44) gives

$$\mathbb{E} \left[ \int_0^{\tau_\epsilon \wedge 1} e^{-\theta t} dt \right] \geq \frac{1}{C}\epsilon^2 (1 - e^{-\theta}) =: \eta_2 > 0.$$

Thus, (43) implies that

$$V(k_M, h_M) > \mathbb{E} \left[ \int_0^{\tau_\epsilon \wedge 1} \frac{e^{-\theta t}}{1 - \sigma} \left( \frac{\bar{h}_t^\gamma}{\bar{c}_t} \right)^{\sigma-1} dt + e^{-\theta(\tau_\epsilon \wedge 1)} V(\bar{k}_{\tau_\epsilon \wedge 1}, \bar{h}_{\tau_\epsilon \wedge 1}) \right] + \frac{\eta_1}{2} \eta_2,$$

We recall that  $\bar{c} \in \mathcal{A}(k_M, h_M)$  is arbitrary and observe that the constant  $\eta_2 > 0$  is independent of  $\bar{c}$ . Therefore, taking the supremum over  $\mathcal{A}(k_M, h_M)$  in the previous estimate, we conclude that

$$V(k_M, h_M) > \sup_{c \in \mathcal{A}(k_M, h_M)} \mathbb{E} \left[ \int_0^{\tau_\epsilon \wedge 1} \frac{e^{-\theta t}}{1 - \sigma} \left( \frac{(h_t^{h_M, 0, c})^\gamma}{c_t} \right)^{\sigma-1} dt + e^{-\theta(\tau_\epsilon \wedge 1)} V(\bar{k}_{\tau_\epsilon \wedge 1}, \bar{h}_{\tau_\epsilon \wedge 1}) \right].$$

This violates the dynamic programming principle in Proposition 4.5. It follows that  $V$  is a viscosity subsolution of (18), and the proof is complete.  $\square$

## 5.2 $V$ is a classical solution of the HJB equation

We recall the definition of classical solution of the HJB equation (18).

**Definition 2.** Let  $D$  be an open subset of  $\mathbb{R}_{++}^2$ . A function  $v \in C^2(D; \mathbb{R})$  is called a *classical solution* of (18) if

$$\theta v(k, h) = H_{\max}(k, h, \nabla v(k, h), D^2 v(k, h)), \quad (k, h) \in D.$$

In the next theorem, we combine the regularity properties of  $H_{\max}$  obtained in Section 3 with fundamental results in Crandall et al. [1992], Safonov [1989] to demonstrate that the value function  $V$  is a classical solution of (18) in  $\mathbb{R}_{++}^2$ . This also shows that  $V$  is a regular function of class  $C^2(\mathbb{R}_{++}^2; \mathbb{R})$ .

**Theorem 5.2.** *Suppose that (28) is fulfilled. Then the value function  $V$  satisfies the HJB equation (18) in  $\mathbb{R}_{++}^2$  in the classical sense.*

*Proof.* Fix a generic  $(k_0, h_0) \in \mathbb{R}_{++}^2$  and consider  $\delta_0 > 0$  such that  $\overline{B}_{\delta_0}(k_0, h_0) \subset \mathbb{R}_{++}^2$ . Thanks to Lemma 3.1 and Theorem 4.4, we can apply Theorem 1.1 in Safonov [1989] to state that, for some  $\alpha \in (0, 1)$ , the HJB equation (18) has a unique classical solution  $v$  in  $B_{\delta_0}(k_0, h_0)$  in the class

$$C^{2+\alpha}(B_{\delta_0}(k_0, h_0); \mathbb{R}) \cap C(\overline{B}_{\delta_0}(k_0, h_0); \mathbb{R})$$

such that  $v = V$  on  $\partial B_{\delta_0}(k_0, h_0)$ . In particular,  $v$  is also a (continuous) viscosity solution of (18) in  $B_{\delta_0}(k_0, h_0)$ . Therefore, by Theorem 3.3 in Crandall et al. [1992] – which can be applied by Lemma 3.2 and Theorem 5.1 – we deduce that

$$V = v \quad \text{in} \quad \overline{B}_{\delta_0}(k_0, h_0).$$

Since  $(k_0, h_0)$  is arbitrary, the proof is complete.  $\square$

*Remark 1.* From the proof of Theorem 5.2, it follows that, for every open bounded set  $D$  such that  $\overline{D} \subset \mathbb{R}_{++}^2$ , the value function  $V$  is of class  $C^{2+\alpha}(D; \mathbb{R})$ , for some  $\alpha \in (0, 1)$  possibly dependent on  $D$ .

## 6 Verification theorem

In this section, we provide a sufficient condition for optimality of controls using the HJB equation. We start with a preliminary lemma.

**Lemma 6.1.** *Fix  $(k_0, h_0) \in \mathbb{R}_{++}^2$ . Then, for every  $p \geq 1$ , there exists a constant  $C > 0$  dependent on  $p$  and the parameters of the model such that*

$$\mathbb{E} \left[ |(k_t, h_t)|^p \right] \leq C \left( 1 + t^{p-\frac{1}{2}} \right) |(h_0, k_0)|^p \exp \left\{ p \left( B + \frac{1}{2} \left( (2p-1)\beta_1^2 + 4p\beta_2^2 \right) t \right) \right\}, \quad t \geq 0, \quad (45)$$

for every admissible control  $c \in \mathcal{A}(k_0, h_0)$ .

*Proof.* Fix  $p \geq 1$ . Given  $(k_0, h_0) \in \mathbb{R}_{++}^2$  and  $c \in \mathcal{A}(k_0, h_0)$ , from the explicit expression of the capital process  $k$  in (3) we compute

$$\mathbb{E}[|k_t|^p] \leq k_0^p e^{p(Bt - \frac{1}{2}\beta_1^2 t)} \mathbb{E}\left[e^{p\beta_1 W_{1,t}}\right] = k_0^p \exp\left\{p\left(B + \frac{1}{2}\beta_1^2(p-1)\right)t\right\}, \quad t \geq 0. \quad (46)$$

Here, in the last equality we employ the fact that  $\exp\{W_{1,t}\}$  has a log-normal distribution, for every  $t \geq 0$ . As for the habit process  $h$  in (4), by the control constraint (7) we compute, using also Hölder's inequality,

$$\begin{aligned} \mathbb{E}[|h_t|^p] &\leq |h_0|^p \exp\left\{p\left(-\rho + \frac{1}{2}\beta_2^2(p-1)\right)t\right\} \\ &\quad + (k_0 R \rho)^p \mathbb{E}\left[\left(\mathcal{E}(-\rho t + \beta_2 W_{2,t})\right)^p \left(\int_0^t \frac{\mathcal{E}(Bs + \beta_1 W_{1,s})}{\mathcal{E}(-\rho s + \beta_2 W_{2,s})} ds\right)^p\right] \\ &\leq \exp\left\{p\left(-\rho + \frac{1}{2}\beta_2^2(p-1)\right)t\right\} \left(|h_0|^p + (k_0 R \rho)^p e^{\frac{1}{2}\beta_2^2 p^2 t} \times \mathbf{I}_t\right), \quad t \geq 0, \end{aligned} \quad (47)$$

where we define

$$\mathbf{I}_t = \mathbb{E}\left[\left(\int_0^t \exp\left\{\left(B + \rho - \frac{1}{2}(\beta_1^2 - \beta_2^2)\right)s + \beta_1 W_{1,s} - \beta_2 W_{2,s}\right\} ds\right)^{2p}\right]^{\frac{1}{2}}, \quad t \geq 0.$$

By Jensen's inequality, Fubini's theorem and the independence of  $W_1$  and  $W_2$ ,

$$\mathbf{I}_t^2 \leq t^{2p-1} \int_0^t \exp\left\{2p\left(B + \rho + \frac{1}{2}((2p-1)\beta_1^2 + (2p+1)\beta_2^2)\right)s\right\} ds.$$

Therefore, denoting by  $C > 0$  a constant possibly dependent on  $p$  and the parameters of the model, from (47) we deduce that, for every  $t \geq 0$ ,

$$\begin{aligned} \mathbb{E}[|h_t|^p] &\leq C \exp\left\{p\left(-\rho + \frac{1}{2}\beta_2^2(p-1)\right)t\right\} |(h_0, k_0)|^p \\ &\quad \times \left(1 + t^{p-\frac{1}{2}} \exp\left\{p\left(B + \rho + \frac{1}{2}((2p-1)\beta_1^2 + (3p+1)\beta_2^2)\right)t\right\}\right) \\ &\leq C \left(1 + t^{p-\frac{1}{2}}\right) |(h_0, k_0)|^p \exp\left\{p\left(B + \frac{1}{2}((2p-1)\beta_1^2 + 4p\beta_2^2)\right)t\right\}. \end{aligned}$$

Combining this estimate with (46) we obtain (45). The proof is then complete.  $\square$

We now impose the following assumptions on the model parameters and the growth of a map  $v$ .

**Assumption 2.** Given a twice-differentiable function  $v : \mathbb{R}_{++}^2 \rightarrow \mathbb{R}$ , suppose that  $v$  and  $\nabla v$  satisfy the polynomial growth condition

$$|v(k, h)| + |\nabla v(k, h)| \leq C(1 + |(k, h)|^p), \quad (k, h) \in \mathbb{R}_{++}^2,$$

for some constant  $C > 0$ . Here,  $p \geq 1$  and satisfies, together with the model parameters, the following inequality:

$$\theta > p \left( B + \frac{1}{2} \left( (2p-1)\beta_1^2 + 4p\beta_2^2 \right) \right).$$

**Theorem 6.2** (Smooth Verification, Sufficient Condition). *Let  $v : \mathbb{R}_{++}^2 \rightarrow \mathbb{R}$  be a classical solution of the HJB equation (18), see Definition 2, which satisfies Assumption 2. Then*

$$v(k, h) \geq V(k, h), \quad (k, h) \in \mathbb{R}_{++}^2, \quad (48)$$

where  $V$  is the value function in (10).

In addition, given  $(k_0, h_0) \in \mathbb{R}_{++}^2$ , suppose that  $c^*$  is an admissible control in  $\mathcal{A}(k_0, h_0)$  such that, denoting by  $k^*$  (resp.,  $h^*$ ) the process  $k^{k_0, 0, c^*}$  (resp.,  $h^{h_0, 0, c^*}$ ),

$$c_s^* \in \arg \max_{c \in (0, Rk_s^*]} g(c; h_s^*, \partial_k v(k_s^*, h_s^*), \partial_h v(k_s^*, h_s^*)), \quad \text{for a.e. } s \in [0, \infty), \mathbb{P}\text{-a.s.}, \quad (49)$$

where  $g$  is defined in (16). Then  $c^*$  is optimal at  $(k_0, h_0)$ , and

$$v(k_0, h_0) = V(k_0, h_0).$$

*Proof.* Fix  $(k_0, h_0) \in \mathbb{R}_{++}^2$  and consider, an admissible control  $c \in \mathcal{A}(k_0, h_0)$ . Denoting by  $k = k^{k_0, 0, c}$  and  $h = h^{h_0, 0, c}$  the solutions of (1) and (2), respectively, by Itô's formula we deduce that

$$\begin{aligned} e^{-\theta T} v(k_T, h_T) = & v(k_0, h_0) + \int_0^T e^{-\theta t} \left( -\theta v(k_t, h_t) + \partial_k v(k_t, h_t)(Bk_t - c_t) + \partial_h v(k_t, h_t)\rho(c_t - h_t) \right. \\ & \left. + \frac{1}{2} \text{tr}(\Sigma(k_t, h_t)^\top D^2 v(k_t, h_t) \Sigma(k_t, h_t)) \right) dt \\ & + \beta_1 \int_0^T e^{-\theta t} k_t \partial_k v(k_t, h_t) dW_{1,t} + \beta_2 \int_0^T e^{-\theta t} h_t \partial_h v(k_t, h_t) dW_{2,t}, \quad T > 0, \mathbb{P}\text{-a.s.} \end{aligned}$$

By Lemma 6.1 and Assumption 2, we can take the expected value in the previous equation.

This passage gives, by the martingale property of the stochastic integrals,

$$\begin{aligned} \mathbb{E} \left[ e^{-\theta T} v(k_T, h_T) \right] + \mathbb{E} \left[ \int_0^T \frac{e^{-\theta t}}{1-\sigma} \left( \frac{c_t}{h_t^\gamma} \right)^{1-\sigma} dt \right] &= v(k_0, h_0) + \mathbb{E} \left[ \int_0^T e^{-\theta t} \left( -\theta v(k_t, h_t) \right. \right. \\ &+ \partial_k v(k_t, h_t) (Bk_t - c_t) + \partial_h v(k_t, h_t) \rho(c_t - h_t) + \frac{1}{2} \text{tr}(\Sigma(k_t, h_t)^\top D^2 v(k_t, h_t) \Sigma(k_t, h_t)) \\ &\left. \left. + \frac{1}{1-\sigma} \left( \frac{c_t}{h_t^\gamma} \right)^{1-\sigma} \right) dt \right]. \end{aligned}$$

Notice that the two sides of the previous equality might not be finite. Since  $v$  is a classical solution of (18), we write

$$\begin{aligned} \mathbb{E} \left[ e^{-\theta T} v(k_T, h_T) \right] + \mathbb{E} \left[ \int_0^T \frac{e^{-\theta t}}{1-\sigma} \left( \frac{c_t}{h_t^\gamma} \right)^{1-\sigma} dt \right] &= v(k_0, h_0) \\ &+ \mathbb{E} \left[ \int_0^T e^{-\theta t} \left( g(c_t; h_t, \partial_k v(k_t, h_t), \partial_h v(k_t, h_t)) - G(k_t, h_t, \partial_k v(k_t, h_t), \partial_h v(k_t, h_t)) \right) dt \right]. \end{aligned} \tag{50}$$

Observe that, by definition of  $g$  and  $G$  (see (17)), the last term in the right-hand side of (50) is nonpositive. Moreover, by Assumption 2 and Lemma 6.1,

$$\lim_{T \rightarrow \infty} \mathbb{E} \left[ e^{-\theta T} v(k_T, h_T) \right] = 0.$$

Thus, by the monotone convergence theorem, we compute the limit as  $T \rightarrow \infty$  in (50) to infer that

$$\begin{aligned} \mathbb{E} \left[ \int_0^\infty \frac{e^{-\theta t}}{1-\sigma} \left( \frac{c_t}{h_t^\gamma} \right)^{1-\sigma} dt \right] &= v(k_0, h_0) \\ &+ \mathbb{E} \left[ \int_0^\infty e^{-\theta t} \left( g(c_t; h_t, \partial_k v(k_t, h_t), \partial_h v(k_t, h_t)) - G(k_t, h_t, \partial_k v(k_t, h_t), \partial_h v(k_t, h_t)) \right) dt \right]. \end{aligned} \tag{51}$$

In particular,

$$v(k_0, h_0) \geq \mathbb{E} \left[ \int_0^\infty \frac{e^{-\theta t}}{1-\sigma} \left( \frac{c_t}{h_t^\gamma} \right)^{1-\sigma} dt \right].$$

Since  $c$  is arbitrary, we can take the supremum over  $\mathcal{A}(k_0, h_0)$  in this estimate to establish (48), as  $(k_0, h_0)$  is a generic point in  $\mathbb{R}_{++}^2$ .

If we have an admissible control  $c^* \in \mathcal{A}(k_0, h_0)$  satisfying (49), then, by the definition of  $G$  in (17),

$$\mathbb{E} \left[ \int_0^\infty e^{-\theta t} \left( g(c_t^*; h_t^*, \partial_k v(k_t^*, h_t^*), \partial_h v(k_t^*, h_t^*)) - G(k_t^*, h_t^*, \partial_k v(k_t^*, h_t^*), \partial_h v(k_t^*, h_t^*)) \right) dt \right] = 0.$$

Consequently, by (48) (which we have just proved) and (51),

$$V(k_0, h_0) \geq \mathbb{E} \left[ \int_0^\infty \frac{e^{-\theta t}}{1 - \sigma} \left( \frac{c_t^*}{(h_t^*)^\gamma} \right)^{1-\sigma} dt \right] = v(k_0, h_0) \geq V(k_0, h_0),$$

which completes the proof.  $\square$

*Remark 2.* Equation (51) in the proof of Theorem 6.2 is the so-called *fundamental identity* for the optimal control problem. Note also that the conclusion (48) of Theorem 6.2 holds when  $v$  is only a classical *supersolution* of the HJB equation (18).

We conclude this section with the *closed loop equations (CLEs)*. As in Theorem 6.2, we consider a classical solution  $v: \mathbb{R}_{++}^2 \rightarrow \mathbb{R}$  of (18) that satisfies Assumption 2. Recalling the arguments leading to (17) in Section 3, we define the *feedback map*

$$\Phi_v: \mathbb{R}_{++}^2 \rightarrow \mathbb{R}_+ \quad \text{such that} \quad \Phi_v(k, h) = \arg \max_{c \in (0, Rk]} g\left(c; h, \partial_k v(k, h), \partial_h v(k, h)\right) \in \mathbb{R}_+.$$

The corresponding CLEs are given by

$$dk_t^{k_0, 0, \Phi_v} = (Bk_t^{k_0, 0, \Phi_v} - \Phi_v(k_t^{k_0, 0, \Phi_v}, h_t^{h_0, 0, \Phi_v}))dt + \beta_1 k_t^{k_0, 0, \Phi_v} dW_{1,t}, \quad k_0 > 0, \quad (52)$$

$$dh_t^{h_0, 0, \Phi_v} = \rho(\Phi_v(k_t^{k_0, 0, \Phi_v}, h_t^{h_0, 0, \Phi_v}) - h_t^{h_0, 0, \Phi_v})dt + \beta_2 h_t^{h_0, 0, \Phi_v} dW_{2,t}, \quad h_0 > 0. \quad (53)$$

By construction, the following corollary is a straightforward consequence of Theorem 6.2.

**Corollary 6.3.** *Given  $(k_0, h_0) \in \mathbb{R}_{++}^2$ , suppose that there exists a solution  $(k^{k_0, 0, \Phi_v}, h^{h_0, 0, \Phi_v})$  of the CLEs (52) and (53) satisfying the positivity constraint (6). Then the process  $c^{\Phi_v} = \Phi(k^{\Phi_v}, h^{\Phi_v})$  belongs to  $\mathcal{A}(k_0, h_0)$  and is an optimal control, that is,*

$$V(k_0, h_0) = \mathbb{E} \left[ \int_0^\infty \frac{e^{-\theta t}}{1 - \sigma} \left( \frac{c_t^{\Phi_v}}{(h_t^{h_0, 0, \Phi_v})^\gamma} \right)^{1-\sigma} dt \right].$$

## 7 Conclusions

We develop and study a growth model with internal habit formation which is a stochastic version of the well-known model in Carroll et al. [1997, 2000]. The corresponding optimal control problem is challenging to analyze, due to the lack of concavity and the singularity of the utility function. We prove crucial results regarding classical solutions of the associated HJB equation and sufficient conditions for optimality. The study of the model optimal paths and their comparison with findings in the deterministic case remains open and will be the subject of a future research project.

## Acknowledgements

Michele Aleandri is member of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

Michele Aleandri, Alessandro Bondi and Fausto Gozzi are supported by the Italian Ministry of University and Research (MIUR), in the framework of PRIN projects 2017FKHBA8 001 (The Time-Space Evolution of Economic Activities: Mathematical Models and Empirical Applications) and 20223PNJ8K (Impact of the Human Activities on the Environment and Economic Decision Making in a Heterogeneous Setting: Mathematical Models and Policy Implications).

## References

Bahman Angoshtari, Erhan Bayraktar, and Virginia R. Young. Optimal investment and consumption under a habit-formation constraint. *SIAM Journal on Financial Mathematics*, 13(1):321–352, 2022.

Mauro Bambi and Fausto Gozzi. Internal habits formation and optimality. *Journal of Mathematical Economics*, 91:165–172, 2020. ISSN 0304-4068. doi: <https://doi.org/10.1016/j.jmateco.2020.09.008>. URL <https://www.sciencedirect.com/science/article/pii/S0304406820301075>.

Christopher D. Carroll, Jody Overland, and David N. Weil. Comparison utility in a growth model. *Journal of Economic Growth*, 2(4):339 – 367, 1997. doi: [10.1023/A:1009740920294](https://doi.org/10.1023/A:1009740920294). URL <https://www.scopus.com/inward/record.uri?eid=2-s2.0-0002969888&doi=10.1023%2fa%3a1009>

- Christopher D. Carroll, Jody Overland, and David N. Weil. Saving and growth with habit formation. *American Economic Review*, 90(3):341–355, 06 2000. doi: 10.1257/aer.90.3.341. URL <https://www.aeaweb.org/articles?id=10.1257/aer.90.3.341>.
- Tahir Choulli, Michael Taksar, and Xun Yu Zhou. A diffusion model for optimal dividend distribution for a company with constraints on risk control. *SIAM Journal on Control and Optimization*, 41(6):1946–1979, 2003.
- Michael G Crandall, Hitoshi Ishii, and Pierre-Louis Lions. User’s guide to viscosity solutions of second order partial differential equations. *Bulletin of the American mathematical society*, 27(1):1–67, 1992.
- Marina Di Giacinto, Salvatore Federico, and Fausto Gozzi. Pension funds with a minimum guarantee: a stochastic control approach. *Finance and Stochastics*, 15:297–342, 2011.
- Wendell H Fleming and Halil Mete Soner. *Controlled Markov processes and viscosity solutions*, volume 25. Springer Science & Business Media, 2006.
- Giuseppe Freni, Fausto Gozzi, and Cristina Pignotti. Optimal strategies in linear multisector models: Value function and optimality conditions. *Journal of Mathematical Economics*, 44(1):55–86, 2008.
- Paul Gassiat, Fausto Gozzi, and Huy en Pham. Investment/consumption problem in illiquid markets with regime-switching. *SIAM Journal on Control and Optimization*, 52(3):1761–1786, 2014.
- Hitoshi Ishii and Paola Loreti. A class of stochastic optimal control problems with state constraint. *Indiana University Mathematics Journal*, pages 1167–1196, 2002.
- Mikhail V Safonov. On the classical solution of nonlinear elliptic equations of second order. *Mathematics of the USSR-Izvestiya*, 33(3):597, 1989.

## A Proof of Lemmas 3.1 and 3.2

*Proof of Lemma 3.1.* In the proof, we denote by  $C > 0$  a positive constant possibly depending on  $\overline{D} \subset \mathbb{R}_{++}^2$  and the parameters of the model allowed to change from line to line. Fix  $Q = (Q_{ij})_{i,j} \in \mathcal{S}^2$ ,  $(k, h) \in \overline{D}$ ,  $p = \begin{bmatrix} p_k \\ p_h \end{bmatrix}$ ,  $\bar{p} = \begin{bmatrix} \bar{p}_k \\ \bar{p}_h \end{bmatrix} \in \mathbb{R}^2$  and  $v, \bar{v} \in \mathbb{R}$ . To prove (20) we distinguish two cases: (i)  $p$  and  $\bar{p}$  satisfy the same inequality in (17); (ii)  $p$  satisfies the first inequality in (17) and  $\bar{p}$  the second.

Case (i). We consider two subcases:

$$(I) \quad p_k - \rho p_h \leq h^{\gamma(\sigma-1)}(Rk)^{-\sigma} \quad \text{and} \quad \bar{p}_k - \rho \bar{p}_h \leq h^{\gamma(\sigma-1)}(Rk)^{-\sigma};$$

$$(II) \quad p_k - \rho p_h > h^{\gamma(\sigma-1)}(Rk)^{-\sigma} \quad \text{and} \quad \bar{p}_k - \rho \bar{p}_h > h^{\gamma(\sigma-1)}(Rk)^{-\sigma}.$$

Notice that, by the continuity of  $G$  in (17), in Case (II) we can also consider the inequalities with  $\geq$  instead of  $>$ . Since  $D$  is bounded, straightforward computations based on (18) and (19) yield

$$\begin{aligned} & |\tilde{F}(Q, p, v, (k, h)) - \tilde{F}(Q, \bar{p}, \bar{v}, (k, h))| \leq \theta |v - \bar{v}| + C(|p_k - \bar{p}_k| + |p_h - \bar{p}_h|) \\ & + \begin{cases} Rk(|p_k - \bar{p}_k| + \rho|p_h - \bar{p}_h|), & \text{if (I) holds,} \\ (1 + \frac{1}{\sigma-1})h^{\gamma(1-\frac{1}{\sigma})} |(p_k - \rho p_h)^{1-\frac{1}{\sigma}} - (\bar{p}_k - \rho \bar{p}_h)^{1-\frac{1}{\sigma}}|, & \text{if (II) holds.} \end{cases} \end{aligned}$$

Considering also that the power  $x^{1-\frac{1}{\sigma}}$  is Lipschitz-continuous in  $(\epsilon, \infty)$ , for any  $\epsilon > 0$ , and that  $\bar{D} \subset \mathbb{R}_{++}^2$ , (20) follows.

Case (ii). Suppose that  $p_k - \rho p_h \leq h^{\gamma(\sigma-1)}(Rk)^{-\sigma}$  and  $\bar{p}_k - \rho \bar{p}_h > h^{\gamma(\sigma-1)}(Rk)^{-\sigma}$ . Since

$$\begin{aligned} 0 & < \max\{h^{\gamma(\sigma-1)}(Rk)^{-\sigma} - (p_k - \rho p_h), (\bar{p}_k - \rho \bar{p}_h) - h^{\gamma(\sigma-1)}(Rk)^{-\sigma}\} \\ & \leq (\bar{p}_k - \rho \bar{p}_h) - (p_k - \rho p_h) \leq C(|p_k - \bar{p}_k| + |p_h - \bar{p}_h|), \end{aligned}$$

(20) follows from Case (i).

As regards (21), by (17), (18), the sublinear growth of the power  $x^{1-\frac{1}{\sigma}}$  in  $\mathbb{R}_+$  and the fact that  $\bar{D} \subset \mathbb{R}_{++}^2$ , an argument similar to the one above for (20) entails that

$$C_1 = C \left( 1 + \sum_{i,j=1}^2 |Q_{ij}| + |p_k| + |p_h| + |v| \right)$$

is a Lipschitz-continuity constant for  $\tilde{F}(Q, p, v, \cdot)$  in  $D$ . Moreover,  $\tilde{F}(Q, p, v, \cdot)$  is bounded by  $C_1$  in  $D$ . Therefore,  $\tilde{F}(Q, p, v, \cdot)$  is  $\alpha$ -Hölder's continuous with constant  $2C_1$  in  $D$  for any  $\alpha \in (0, 1)$ , whence (21). This completes the proof.  $\square$

*Proof of Lemma 3.2.* As in the proof of Lemma 3.1, we denote by  $C > 0$  a positive constant possibly depending on  $D \subset \mathbb{R}_{++}^2$  and the parameters of the model allowed to change from line to line. Fix  $(k, h), (\bar{k}, \bar{h}) \in D$ ,  $v \in \mathbb{R}$  and  $Q, \bar{Q} \in \mathcal{S}^2$  such that (23) is fulfilled. Based on (17), we distinguish the two cases (A) and (B).

(A) One of the following holds:

$$(A_1) \quad \alpha(k - \bar{k} - \rho(h - \bar{h})) \leq \bar{h}^{\gamma(\sigma-1)}(R\bar{k})^{-\sigma} \quad \text{and} \quad \alpha(k - \bar{k} - \rho(h - \bar{h})) \leq h^{\gamma(\sigma-1)}(Rk)^{-\sigma};$$

$$(A_2) \quad \alpha(k - \bar{k} - \rho(h - \bar{h})) > \bar{h}^{\gamma(\sigma-1)}(R\bar{k})^{-\sigma} \quad \text{and} \quad \alpha(k - \bar{k} - \rho(h - \bar{h})) > h^{\gamma(\sigma-1)}(Rk)^{-\sigma}.$$

(B) One of the following holds:

$$(B_1) \quad \alpha(k - \bar{k} - \rho(h - \bar{h})) > \bar{h}^{\gamma(\sigma-1)}(R\bar{k})^{-\sigma} \quad \text{and} \quad \alpha(k - \bar{k} - \rho(h - \bar{h})) \leq h^{\gamma(\sigma-1)}(Rk)^{-\sigma};$$

$$(B_2) \quad \alpha(k - \bar{k} - \rho(h - \bar{h})) \leq \bar{h}^{\gamma(\sigma-1)}(R\bar{k})^{-\sigma} \quad \text{and} \quad \alpha(k - \bar{k} - \rho(h - \bar{h})) > h^{\gamma(\sigma-1)}(Rk)^{-\sigma}.$$

In Case (A), writing  $\Sigma$  (resp.,  $\bar{\Sigma}$ ) for the diagonal matrix  $\Sigma(k, h)$  (resp.,  $\Sigma(\bar{k}, \bar{h})$ ) to simplify the notation, we compute

$$\begin{aligned} & \tilde{F}(Q, \alpha(k - \bar{k}, h - \bar{h}), v, (k, h)) - \tilde{F}(\bar{Q}, \alpha(k - \bar{k}, h - \bar{h}), v, (\bar{k}, \bar{h})) \\ &= \alpha B(k - \bar{k})^2 - \rho \alpha (h - \bar{h})^2 + \frac{1}{2} \text{tr}(\Sigma^\top Q \Sigma - \bar{\Sigma}^\top \bar{Q} \bar{\Sigma}) \\ &+ \begin{cases} -\alpha R(k - \bar{k})^2 + \alpha R \rho (k - \bar{k})(h - \bar{h}) + \frac{1}{\sigma-1} \left( \left( \frac{\bar{h}^\gamma}{R\bar{k}} \right)^{\sigma-1} - \left( \frac{h^\gamma}{Rk} \right)^{\sigma-1} \right), & \text{if } (A_1), \\ \left( 1 + \frac{1}{\sigma-1} \right) (\alpha(k - \bar{k}) - \rho \alpha (h - \bar{h}))^{1-\frac{1}{\sigma}} \left( \bar{h}^{\gamma(1-\frac{1}{\sigma})} - h^{\gamma(1-\frac{1}{\sigma})} \right), & \text{if } (A_2). \end{cases} \end{aligned}$$

The boundedness of  $D$  and the fact that  $\bar{D} \subset \mathbb{R}_{++}^2$ , together with the trivial inequality  $2ab \leq a^2 + b^2$ ,  $(a, b) \in \mathbb{R}^2$ , yield (cf. Case (i) (I) in the proof of Lemma 3.1)

$$\begin{aligned} & \tilde{F}(Q, \alpha(k - \bar{k}, h - \bar{h}), v, (k, h)) - \tilde{F}(\bar{Q}, \alpha(k - \bar{k}, h - \bar{h}), v, (\bar{k}, \bar{h})) \\ & \leq C(\alpha|(k - \bar{k}, h - \bar{h})|^2 + |(k - \bar{k}, h - \bar{h})|) + \frac{1}{2} \text{tr}(\Sigma^\top Q \Sigma - \bar{\Sigma}^\top \bar{Q} \bar{\Sigma}). \end{aligned} \quad (54)$$

To estimate the trace in (54), we multiply (23) on the right by the positive semidefinite matrix

$$\begin{bmatrix} \Sigma^\top \Sigma & \bar{\Sigma}^\top \Sigma \\ \Sigma^\top \bar{\Sigma} & \bar{\Sigma}^\top \bar{\Sigma} \end{bmatrix} \in \mathcal{S}^4$$

and then take the trace. Since this passage does not alter the order in (23), we have

$$\text{tr}(\Sigma^\top Q \Sigma - \bar{\Sigma}^\top \bar{Q} \bar{\Sigma}) \leq 3\alpha \text{tr}((\Sigma^\top - \bar{\Sigma}^\top)(\Sigma - \bar{\Sigma})) \leq C\alpha|(k - \bar{k}, h - \bar{h})|^2.$$

Plugging this bound into (54) gives (22), completing the proof of Case (A).

Notice that, by the continuity of  $G$  in (17), in Case  $(A_2)$  we can also consider the inequalities with  $\geq$  instead of  $>$ . Thus, Case (B) can be readily deduced from Case (A), see also Case (ii) in the proof of Lemma 3.1. The proof is now complete.  $\square$