PRESENTATION OF FINITE REEDY CATEGORIES AS LOCALIZATIONS OF FINITE DIRECT CATEGORIES

GENKI SATO

ABSTRACT. In this paper, we present a construction from a Reedy category C of a direct category Down(C) and a functor $\text{Down}(C) \to C$, which exhibits C as an $(\infty, 1)$ -categorical localization of Down(C). This result refines previous constructions in the literature by ensuring finiteness of the direct category Down(C) whenever C is finite, which is not guaranteed by existing approaches. The finiteness property is useful when we want to embed the construction into the syntax of a (non-infinitary) logic: the author expects the construction may be used to develop a meta-theory of finitely truncated simplicial types for homotopy type theory.

Contents

1. Introduction	2
Acknowledgements	4
2. Preliminaries	4
2.1. Grossaries	4
2.2. Remarks on foundations	7
2.3. Direct and Reedy categories	8
2.4. The simplex category	11
2.5. Simplicial sets, quasicategories and $(\infty, 1)$ -categories	11
3. Several categories related to a Reedy category	17
4. The hom-posets of the categories of skew ladders	20
5. Reedy structure, directness and finiteness	24
6. Functors between the constructed categories	28
7. The proof of 1-localization	31
8. Shapes of $(\infty, 1)$ -diagrams used in the proof of $(\infty, 1)$ -localization	37
8.1. Some endofunctors on \mathbf{Set}_{Δ}	38
8.2. Some natural transformations between the endofunctors	42
8.3. Properties of the transformations 1: two universal localizations	45
8.4. Properties of the transformations 2: an inner anodyne map	50
8.5. Properties of the transformations 3: another inner anodyne map	55
8.6. Main lemma for $(\infty, 1)$ -localization	60
9. The proof of $(\infty, 1)$ -localization	62
10. Conclusion: proof of the main theorem	67
References	67

 $E\text{-}mail\ address:\ \texttt{gettaplacetogo} \texttt{@gmail.com}.$

²⁰²⁰ Mathematics Subject Classification. 18N55 .

Key words and phrases. Reedy category, direct category, finite category, $(\infty, 1)$ -localization, simplicial set, homotopy type theory.

2 FINITE REEDY CATEGORIES AS LOCALIZATIONS OF FINITE DIRECT CATEGORIES

1. INTRODUCTION

In this paper, we prove the following result:

Theorem 1.1. Let C be a Reedy category. Then there is a concrete construction (to be seen in Definitions 3.8 and 7.1) of a direct category Down(C) and a functor $\text{Down}(C) \to C$ which exhibits C as an $(\infty, 1)$ -categorical localization of Down(C). Furthermore, the category Down(C) is finite whenever C is.

Without the mention of finiteness, the proof of this result appears in literature in a stronger form:

Theorem 1.2 (Preceding result). Let X be any simplicial set. Then there is a well-founded partial order P and a functor $P \to X$, i.e., a simplicial map $N(P) \to X$ exhibiting X as an $(\infty, 1)$ -localization of P.

A direct proof of Theorem 1.2 may be seen, for example, in the Lurie's online textbook *Kerodon* as [16, Theorem 02MD]. The proof of a claim "equivalent to Theorem 1.2 up to homotopy" can also be found in an earlier literature: in [4, 3], Barwick and Kan state that the cofibrant objects of a model category of relative categories, which models all $(\infty, 1)$ -categories, are necessarily relative posets; therefore any $(\infty, 1)$ -category is a localization of a poset.

As you can see, the statement of Barwick and Kan does not refer to well-foundedness (or equivalently directness). Strictly speaking, Lurie's statement does not either. However, with a closer inspection of Lurie's proof, it is easily seen that the poset he has constructed is well-founded. I, the author, have not inspected Barwick and Kan's proof, but I roughly expect that the poset they have constructed is well-founded as well, as they use a modified nerve functor.

Although the preceding Theorem 1.2 is almost stronger than our Theorem 1.1, there is an improvement in the latter: finiteness. If C is a finite Reedy category, the category Down(C), which has C as a localization, is in fact a finite direct category. The preceding constructions do not seem to guarantee the finiteness of the poset P except in very limited cases. For example, Lurie's construction of P may be taken to be finite precisely when X is a finite simplicial set. If we wish to take X = N(C), the nerve of a 1-category, the simplicial set X is finite if and only if C is already finite and *direct*. Therefore, if you wish to get a finite direct category from Theorem 1.2, you need to start with a finite direct category, which renders Theorem 1.2 useless for the purpose.

On the other hand, our Theorem 1.1 provides a finite direct category from a finite Reedy category. Furthermore, by applying our construction and then Lurie's construction, we obtain the following corollary:

Corollary 1.3. Let C be a finite Reedy category. Then there is a finite partial order P and a functor $P \to C$ exhibiting C as an $(\infty, 1)$ -localization of P.

The author expects that the result of this paper may be useful when one wishes to embed the construction into the syntax of a (non-infinitary) logic. In particular, it may have applications to the meta-theory of homotopy type theory (HoTT) [22]. In HoTT, there is a meta-notion of finite direct presheaves (or finite inverse diagrams): for each finite direct category C, there is a well-defined notion of the type of presheaves on C [20, 13]. The result of this paper suggests that these notions can be extended to finite Reedy presheaves: if C is a finite Reedy category, a presheaf over C may be defined as a presheaf over Down(C) equipped with a point of an appropriate subcontractible space. A key example is that of finitely truncated simplicial types. For each n, there is a theory of n-truncated semisimplicial types; however, the theory of n-truncated simplicial types for an arbitrary meta-level natural number n has not yet been developed. Our result suggests that the theory of n-simplicial types can be developed in a straightforward way.

Kraus and Sattler have already worked along these lines in their extended abstract [13]. As a part of their work, they have proposed a sketch of a in-HoTT theory of diagrams over a fixed Reedy category with certain conditions via a "direct replacement". An important example of their work is a simple infinite direct replacement of the untruncated simplex category, which is far simpler than the present construction. This approach yields a definition of simplicial types in terms of countably infinite meta-series of type judgements in HoTT. However, in private communication, Kraus and the author have confirmed that truncated simplex categories do not satisfy the conditions proposed in [13], contrary to the claim in it. Our present construction applies to all Reedy categories, including truncated simplex categories.

Let us roughly outline the construction of Down(C). Given a finite Reedy category $C = (C, C_-, C_+)$, the construction of Down(C) begins with the direct subcategory C_+ . If we added morphisms from C_- to this, we would obtain C, but it would break directness. Instead, for each non-identity $f: x \to y$ in C_- , we would like to add a new object c_f and two morphisms, as in:

$$x \longrightarrow c_f \xleftarrow{\sim} y$$

If we make the morphism $c_f \leftarrow y$ a weak equivalence, we would get a "factorization" of f. If we make c_f have a higher degree than x and y, the directness or the finiteness would not be broken. However, since we need to ensure coherence, just adding these objects c_f and connecting morphisms is not enough. We actually need to add an arbitraily long composable chains in C_- as objects and we consider any series of morphisms in C_+ that connect these chains as morphisms; the resulting category is called $\int \mathbf{N}^{-,+}(C)$ in Section 3. However this does not guarantee finiteness, because this construction allows arbitrarily long chains of idenitities. These arbitrary long chains actually breaks the directness (it is merely Reedy). By quotienting out these idenities, we obtain a finite direct category Down(C), because C_- is finite inverse. This is the rough idea of the construction.

Let us now give an outline of the paper. In Section 2, we wrap up some terminology and some results in category and ∞ -category theory. In Section 3, we define the category Down(C) and some other related categories. In Section 4, we investigate the structure of hom-set of Down(C). From this, the directness and the finiteness of Down(C) will be shown in Section 5. In Section 6, we shall study some functors among Down(C) and other categories constructed in Section 3. This helps us to show that $\text{Down}(C) \to C$, among other C-valued functors, are 1localizations in Section 7. The functor $\text{Down}(C) \to C$ itself, called last, will also be constructed in Section 7. After that, Sections 8 and 9 will be devoted to the proof that $\text{Down}(C) \to C$ is an $(\infty, 1)$ -localization. In Section 8, we shall define some endofunctors on the category of simplicial sets and investigate their properties. These endofunctors will describe the shapes of the $(\infty, 1)$ -diagrams in the actual proof in Section 9 that $\text{Down}(C) \to C$ is an $(\infty, 1)$ -localization. Our main theorem, Theorem 1.1, will be proven in Section 10. The content of the theorem shall have been proven part by part in the previous sections, and the proof in Section 10 will be a concise summary of these results.

Throughout this paper except for Section 2, we shall fix a Reedy category $C = (C, C_{-}, C_{+})$.

4 FINITE REEDY CATEGORIES AS LOCALIZATIONS OF FINITE DIRECT CATEGORIES

Acknowledgements

I would like to thank my advisors, Toshitake Kohno and Nariya Kawazumi, for their invaluable guidance and encouragement. I am also grateful to Yohsuke Matsuzawa for organizing regular progress meetings, despite not sharing my mathematical background, and to Jun Yoshida and to Satoshi Sugiyama for their helpful conversations. I appreciate the great previous research by Nicolai Kraus and Christian Sattler, and I am grateful to Nicolai Kraus for the wounderful discussion on their work. I appreciate the insights provided by Andrej Bauer, Selgei Burkin, Simon Henry, Peter LeFanu Lamsdaine, Maxime Ramzi, Mike Shulman, Paul Taylor, and David White through MathOverflow. My thanks also go to my collegemates who have supported and delighted me, during my course at the University of Tokyo. Finally, although this paper was completed after my departure from the university, my doctoral research was supported by a JSPS Research Fellowship and the FMSP program at the University of Tokyo.

2. Preliminaries

In this section, we review some category-theoretic preliminaries. In Section 2.1, we list some standard category-theoretic terminology and notations, including the weak and the strict definitions of 1-localizations. In Section 2.2, we remark on the set-theoretic foundations of this paper. Since the first half of this paper is based on the language of constructive mathematics, we shortly discuss the possible formalization of the part. The subsection also includes the definition of well-founded relations, which is foundational for defining direct categories. In Section 2.3, we discuss the definitions of direct categories and Reedy categories. Those notions need explicit reconsideration under the constructive settings, we shall discuss them in detail. In Section 2.4, we recall the definition of our most important Reedy category, the simplex category Δ , and list some related notations and terminologies. Finally in Section 2.5, we shall cite some terminology, notation and results from $(\infty, 1)$ -category theory based on simplicial sets as models. This will be vital in Sections 8 and 9 for formulating and proving that Down(C) $\rightarrow C$ is an $(\infty, 1)$ -localization.

2.1. **Grossaries.** We first list some category-theoretic terminology and notations. This is long, but it is a standard glossary in category theory, so the readers may wish to skip this list. See, for instance, [2, 6, 17] for reference.

Definition 2.1 (Category-theoretic glossary).

- Let C be a category. Then we write Ob C for the collection of objects of C, $\operatorname{Hom}_C(x, y)$ for the collection of morphisms $x \to y$ in C, and Mor C for the collection of all morphisms of C. If $f: x \to y$ and $g: y \to z$ are morphisms in a category, their composition $x \to z$ will be denoted by $g \circ f$. The identity morphism of an object x in a category is written as id_x .
- If there is no danger of confusion, we may write $f \in C$ to mean $f \in Mor C$, and $x \in C$ to say $x \in Ob C$.
- A functor $F: C \to D$, as usual, stands for a covariant one: a pair of a function $F: \operatorname{Ob} C \to \operatorname{Ob} D$ and a family of functions $F: \operatorname{Hom}_C(x, y) \to \operatorname{Hom}_D(F(x), F(y))$ preserving identities and compositions. If $F: C \to D$ and $G: D \to E$ are functors, their composition $x \mapsto G(F(x))$ is denoted by $G \circ F: C \to E$. The identity functor of a category C is denoted by id_C .
- A functor $F: C \to D$ is *faithful* if, for any pair of objects x, y in C, the function $F: \operatorname{Hom}_C(x, y) \to \operatorname{Hom}_D(F(x), F(y))$ is injective. The functor is *full* if this function is surjective, and *fully faithful* if it is an isomorphism of sets.

- A category C' is said to be a *subcategory* of a category C, denoted by $C' \subseteq C$, if there is a faithful functor $C' \to C$ solely consisting of inclusions of subsets on objects and morphisms. A subcategory C' is said to be *full* if the functor is fully faithful; it is said to be *wide* if the functor is the identity on objects.
- A functor $F: C \to D$ is said to reflect identity if for any morphism $f: x \to y$ in C with F(x) = F(y) and $F(f) = id_{F(x)}$, we have x = y and $f = id_x$.
- A natural transformation $\alpha: F \Rightarrow G$ between functors $F, G: C \to D$, as always, consists of a morphism $\alpha_x: F(x) \to G(x)$ in D for each object x in C satisfying suitable commutativity condition.
- The vertical composition of natural transformations $\alpha \colon F \Rightarrow G$ and $\beta \colon G \Rightarrow H$, is denoted by $\beta \circ \alpha \colon F \Rightarrow H$. The identity natural transformation of a functor F is denoted by $\mathrm{id}_F \colon F \Rightarrow F$.
- The functor category D^C has functors $C \to D$ as objects, natural transformations between them as morphisms, and the vertical composition of natural transformations as composition.
- The left whiskering of a natural transformation $\alpha \colon F \Rightarrow G \colon C \to D$ by a functor $H \colon D \to D'$ is denoted by $H \triangleleft \alpha \colon H \circ F \Rightarrow H \circ G$.
- The right whiskering of a natural transformation $\alpha \colon F \Rightarrow G \colon C \to D$ by a functor $H \colon C' \to C$ is denoted by $\alpha \triangleright H \colon F \circ H \Rightarrow G \circ H$.
- Let C and D be a category. If $F: C' \to C$ is a functor, precomposition with F and the left whiskering by F define the precomposition functor $D^C \to D^{C'}$, denoted by $F^* = (-\circ F)$. Similarly for a functor $G: D \to D'$, the postcomposition functor $D^C \to D'^C$ is denoted by $G_* = (G \circ -)$.
- The dual of a category C will be denoted by C^{op} .
- The category $D^{C^{\text{op}}}$ of functors $C^{\text{op}} \to D$ may be denoted by D_C . The objects of this category may also be called *D*-valued presheaves over *C* (or contravariant functors from *C* to *D*, but we will not use this term).
- If x, y are objects in a category C and the hom-set $\text{Hom}_C(x, y)$ is a singleton, then we write $! = !_{xy} : x \to y$ for the unique morphism in $\text{Hom}_C(x, y)$.
- Let C and D be categories. The *join* $C \star D$ is the category whose objects are $Ob(C \star D) := Ob(C) \amalg Ob(D) = (\{0\} \times Ob(C)) \cup (\{1\} \times Ob(D))$ and whose morphisms are

 $\operatorname{Hom}_{C\star D}((i,x),(j,y)) = \begin{cases} \operatorname{Hom}_{C}(x,y) & \text{if } i = j = 0, \\ \operatorname{Hom}_{D}(x,y) & \text{if } i = j = 1, \\ \{!\} & \text{if } i = 0 \text{ and } j = 1, \\ \emptyset & \text{if } i = 1 \text{ and } j = 0. \end{cases}$

The composition in $C \star D$ is given by the compositions in C and D when possible, and by the unique possible choice when composing with !.

- We use the word *semicategory* for categories without the assumption of identities; in other words, a quiver with associative compositions. We can forget the structure of identities in a category to get a semicategory, and can freely adjoin identities to a semicategory to obtain a category.
- A finite category is a small category whose sets of objects its morphisms are both finite, i.e. bijective to some set of the form $\{0, 1, ..., n-1\}$.
- A preordered set (P, ≤) is canonically considered as a category by defining Ob P to be P regarded as a set and setting

$$\operatorname{Hom}_{P}(x, y) \coloneqq \begin{cases} \{ !_{xy} \} & \text{if } x \leq y, \\ \emptyset & \text{otherwise.} \end{cases}$$

6 FINITE REEDY CATEGORIES AS LOCALIZATIONS OF FINITE DIRECT CATEGORIES

- If a category C has every hom-set a subset of a singleton, then C is identified with the preordered set of its objects. In particular, the notation for joins of categories and of preordered sets are identified. Also, the notation Q^P for the category of functors $P \to Q$ is identified as the preordered set of order-preserving functions $P \to Q$.
- We employ some examples of *large categories*: the category of small categories **Cat**, that of sets **Set**, that of (reflexively) partially ordered sets **Poset**, and the category of functors from a small category to these three categories. The most important example of such functor categories in this paper is the category of simplicial sets \mathbf{Set}_{Δ} . The term "category," when used without qualification or reference to these examples, will always mean a *small* category.
- If C is a small category, the Yoneda embedding $C \to \mathbf{Set}_C$ is denoted by \mathbf{y} : $\mathbf{y}(c) = \operatorname{Hom}_C(-, c)$ for $c \in \operatorname{Ob} C$.
- The *colimit* of the *diagram* $F: D \to C$ is denoted by $\operatorname{colim}_{d \in D} F(d)$.
- As a specific case of colimits, if $F: \Lambda \to C$ and $G: \Lambda \to D$ are functors, the pointwise left Kan extension of G along F is denoted by $(\operatorname{Lan}_G F)(c) = \operatorname{colim}_{F(\lambda)\to c} G(\lambda)$.
- If C is a category and $F: C^{\text{op}} \to \mathbf{Cat}$ is a functor, the *Grothendieck construction* of F is denoted by $\int F \to C$. We will clarify the actual construction in its specialized form in Definition 3.1.
- We use some terminology from enriched category theory for the specific case of enrichment over **Poset**. When we endow a **Poset**-enrichment on a category, we do not introduce notation for hom-objects; instead, we simply endow hom-sets with order structures. A notable usage of the term from the theory is the change of enriching base by a lax monoidal functor.

We also need to revisit the weak and strict definitions of (1-)localizations of categories. For the strict definition, we refer the reader to [6, Section 5.2] and [7, Section I.1]. The definition of weak localizations may be found in [16, Definition 01M9].

Definition 2.2. Let $F: C \to D$ be a functor between categories, and let $W \subseteq$ Mor C. Consider the following conditions:

- (1) F sends morphisms in W to isomorphisms in D.
- (2) If a functor $G: C \to E$ sends morphisms in W to isomorphisms in E, then there exists a unique functor $H: D \to E$ such that $H \circ F = G$.
- (3) If a functor $G: C \to E$ sends morphisms in W to isomorphisms in E, then there exists a functor $H: D \to E$ and a natural isomorphism $\theta: H \circ F \Rightarrow G$.
- (4) If $H, H': D \to E$ are functors and $\theta: H \circ F \Rightarrow H' \circ F$ is a natural isomorphism, then there is a unique natural isomorphism $\hat{\theta}: H \Rightarrow H'$ such that $\hat{\theta} \triangleright F = \theta$.
- (5) For any category E, the precomposition functor $F^* \colon E^D \to E^C$ is fully faithful.

We say that F exhibits D as the strict (1-)localization of C at (or with respect to) W if it satisfies the conditions (1) and (2). We may instead say that the pair (D, F) is a strict (1-)localization of C at, or with respect to, W. We say F exhibits D as the weak (1-)localization of C at (or with respect to) W if it satisfies the conditions (1), (3) and (4), or equivalently the conditions (1), (3) and (5). In this case, we may instead say that the pair (D, F) is a weak (1-)localization of C at, or with respect to, W.

Remark 2.3. A strict (1-)localization is a weak (1-)localization. More specifically, a pair $(D, F: C \to D)$ of a category and a functor is a weak 1-localization of C at

 $W \subseteq \text{Mor } C$ if and only if there is a strict 1-localization (D', F') of C at W, an equivalence $H: D' \xrightarrow{\sim} D$ of categories, and a natural isomorphism $H \circ F' \cong F$. On the other hand, the pair (D, F) is a strict 1-localization of C at W if and only if it is a weak 1-localization of C at W and $F: \text{Ob } C \to \text{Ob } D$ is a bijection.

2.2. Remarks on foundations. Although we need to enter the realm of classical axiomatic set theory in Sections 8 and 9 to enable our discussion of $(\infty, 1)$ -category theory, we shall use the language of constructive mathematics in Sections 3–7. We only place restrictions on the use of non-constructive principles, and we do not additionally assume constructive principles that contradict classical mathematics.

If P is a proposition or a property, we say "P is decidable" to mean that "either P is true or P is false." If P is a property of some object, we may say "We have dichotomy of x into P and non-P" to mean that "x either satisfies P or does not."

In order to formalize our results in Sections 3–7, it should be sufficient to have, as the foundation, Aczel's Constructive Zermelo-Fraenkel (CZF) set theory [1]. CZF is an intuitionistic first-order theory with equality whose unique non-logical symbol is the membership binary relation \in . Its axioms (and axiom schemata) are Axioms of Extensionality, Pairing, Union, Empty Set and Infinity, and the Axiom Schemata of Δ_0 -Separation, Strong Collection, Subset Collection, and \in -Induction. Note that CZF does not include the dedicate notion of classes, and the large categories should be treated through the well-known trick of metatheoretically regarding properties of sets as classes.

In fact, since our discussion in Sections 3–7 is almost completely finitistic, the author strongly conjectures that the foundation for the formalization of these sections can be weakened to Intuitionistic Kripke-Platek set theory with the Axiom of Infinity (IKP ω) [14]. The theory IKP ω is axiomatized by Axioms of Extensionality, Pairing, Union, Empty Set and Infinity, as well as the Axiom Schemata of Δ_0 -Separation, Δ_0 -Collection, and \in -Induction. The author has not checked the details of the formalization, but the categories and functors constructed in Sections 3–7 may be built using finite-domain function sets, which are Δ_1 -constructible in IKP ω . Since IKP admits the Theorem Scheme of Strong Σ_1 -Collection, this provides a strong indication that the entire formalization can indeed be carried out in IKP ω .

The author furthermore conjectures that we may use Intuitionistic Kripke-Platke set theory without the Axiom of Infinity (IKP) by reformulating and omitting some of the results in Sections 5 and 7. Specifically, the author suspects that Sections 3, 4 and 6 remain valid if we allow large categories in our construction, and that the following results admit modified proofs:

- the categories ∫ N^{--,+}₊(C) from Definition 3.3 and Down(C) from Definition 3.8 are small categories;
- the two categories are direct (Corollary 5.5 and Proposition 5.7);
- the two categories are finite if C is finite (Lemma 5.8 and Corollary 5.9);
- the functor $\text{Down}(C) \to C$ is a 1-localization functor (Theorem 7.3).

However, without the Axiom of Infinity, the arguments cannot be executed as smoothly as in the present paper, since we then only have the *class* of natural numbers. Consequently the following categories are not necessarily *small*, meaning that we need to meta-theoretically treat them using classes:

- the simplex category Δ ;
- the category $\int \mathbf{N}(C)$ in Definition 3.1;
- the category $\int \mathbf{N}^{-,+}(C)$ in Definition 3.2;
- the category $\text{Down}_*(C)$ in Definition 3.8.

8 FINITE REEDY CATEGORIES AS LOCALIZATIONS OF FINITE DIRECT CATEGORIES

The natural definitions of Reedy structures on these large categories, their directness, or their localizations naturally involve univeral quantification over all classes satisfying certain conditions, which is not possible in this set theory. In the present paper, Proposition 5.3 and several from Section 7 show such properties of these categories. We might be able to reformulate them as meta-theorems or treat them with workarounds, or might need to give them up. The author has not investigated this further.

As mentioned at the beginning of this subsection, to formalize the results in Sections 8 and 9, we need the language of classical mathematics and a stronger set-theoretic foundation. In these sections, we shall use the language of $(\infty, 1)$ category theory to prove that $\text{Down}(C) \to C$ is an $(\infty, 1)$ -localization. Since the author is not aware of any constructive theory of $(\infty, 1)$ -categories, we shall not attempt to formalize these sections in a constructive setting. One possible choice of foundation for these sections is Zermelo-Fraenkel set theory with the Axiom of Choice (ZFC), but that does not provide an internalized notion of classes, and therefore some lemmas (especially those in Section 2.5) would have to be treated as meta-theorems. Alternatively, it can be more convenient to use a foundation that includes or allows a definition of classes, such as Morse-Kelley set theory (MK), or Tarski-Grothendieck set theory (TG). See Shulman's survey [21] for a discussion on this topic.

As we treat direct and Reedy categories, we shall need to use the notion of well-foundedness. We simply recall the definition here:

Definition 2.4 (Well-foundedness). Let X be a set and < be a binary relation on X. We say that < is a *well-founded relation* on X if X is the only subset Y of X satisfying the following property: for any $x \in X$, if $\{y \in X \mid y < x\} \subseteq Y$, then $x \in Y$. The set X equipped with such < is said to be a *well-founded set*.

However, the treatment of well-foundedness in this paper is abstract, and we shall only need the following facts for this paper:

- If X is a set and < is a well-founded relation on X, then there is no <-cycle: $x_0 < x_1 < \cdots < x_n < x_0$ for some $n \ge 0$.
- The set \mathbb{N} of natural numbers is well-founded under the usual order.
- Let $(X, <_X)$ and $(Y, <_Y)$ be sets with relations, and assume that the latter is well-founded. If there is a function $f: X \to Y$ such that $x_1 <_X x_2$ implies $f(x_1) <_Y f(x_2)$, then $<_X$ is well-founded.
- Let $(\Lambda, <)$ be a well-founded set, and $(X_{\lambda}, <_{\lambda})$ be a well-founded set for each $\lambda \in \Lambda$. Endow the disjoint union

$$\coprod_{\lambda \in \Lambda} X_{\lambda} = \{ (\lambda, x) \mid \lambda \in \Lambda, x \in X_{\lambda} \}$$

with the lexicographic relation, i.e., $(\lambda_1, x_1) < (\lambda_2, x_2)$ if and only if either $\lambda_1 < \lambda_2$ or $\lambda_1 = \lambda_2$ and $x_1 <_{\lambda_1} x_2$. Then the disjoint union is well-founded.

2.3. Direct and Reedy categories. In this subsection, we review the definitions of direct and Reedy categories constructively. Textbook account with classic foundation can be found, for example, in [9, Sections 5.1 and 5.2].

We first begin with the definition of direct categories. The constructive definition of direct categories suffers from ambiguity; the correct categorification of wellfounded sets would be direct *semicategories*, and it is not clear what is the vaild extension of the notion to categories with identities. The following definition is the one supported by Shulman's helpful answer [12] to my MathOverflow question:

Definition 2.5. Let C be a semicategory. For any $x, y \in Ob C$, let us write $x \prec_C y$ if there exists morphism $x \to y$ in C. We say that C is *direct* if \prec_C is a well-founded

relation on Ob C. A category is said to be *direct* if it is isomorphic to the one obtained by freely adjoining identities to a direct semicategory. A semicategory or a category is said to be *inverse* if its dual is direct.

This definition is good in that it allows us to construct a presheaf over any direct category by the most intuitive form of induction. The definition is also equivalent to the inductive definition by Shulman in [19]. Bartels, on the other hand, raised a remarkable opposition in comments to the same question: the definition above does not include the canonical reflexively ordered class of ordinals, even if we see the irreflexively ordered class of ordinals as well-founded.

The following lemma proves that this constructive definition is equivalent to that from the classical mathematics:

Lemma 2.6. Let C be a category. For any $x, y \in ObC$, let us write $x \prec y$ if there is a non-identity morphism $x \rightarrow y$. Assume that \prec is a well-founded relation on ObC. Then C is a direct category if and only if any morphism in C is either an identity or non-identity.

Proof. The "only if" part is clear. For the "if" part, assume the dichotomy of morphisms into identities and non-identities. We need to show that C is direct. Let the wide sub-semicategory C' consist of all objects in C and all non-identity morphisms in C. For well-definedness, we need to show that C' is closed under compositions. Let $f: x \to y$ and $g: y \to z$ be non-identity morphisms in C. If $g \circ f$ were an identity, then we would have $x \prec y \prec z = x$, which contradicts the well-foundedness of \prec . Therefore, $g \circ f$ is a non-identity, and C' is a semicategory. By the assumption of the lemma, C' is direct. It suffices to see that C is isomorphic to the category obtained by freely adjoining identities to C', which is immediate from our dichotomy assumption.

Remark 2.7. If C is a finite category, the following three conditions are equivalent:

- (1) C is a direct category.
- (2) C is an inverse category.
- (3) Any endomorphisms and isomorphisms of C are identities.

We now move on to the definition of Reedy categories.

Definition 2.8 (Reedy categories). Let C be a category. A *Reedy structure* on C is a pair (C_{-}, C_{+}) of wide subcategories of C satisfying the following conditions:

- (1) Every morphism in C can be uniquely factored as the composition of a morphism in C_{-} followed by one in C_{+} : for each morphism $f: x \to y$ in C, there is a unique triple (z, g, h) of $z \in Ob(C)$, $g: x \to z$ in C_{-} , and $h: z \to y$ in C_{+} with $f = h \circ g$.
- (2) Each of C_{-} and C_{+} is isomorphic to the category obtained by freely adjoining identities to some semicategory.
- (3) Define a relation <' on Ob(C) by setting x <' y if and only if there is a non-identity $x \to y$ in C_+ or a non-identity $x \leftarrow y$ in C_- . Then <' is a well-founded relation on Ob(C).

A Reedy category $C = (C, C_{-}, C_{+})$ is a category C equipped with a Reedy structure (C_{-}, C_{+}) on it.

The definition may look different from the usual definition (for example, see Hovey [9]), but it is in fact classically equivalent. With classical logic, the condition (2) in Definition 2.8 is derivable from (3). To be constructively precise, see the following lemma:

Lemma 2.9. Let C be a category, and let $C_-, C_+ \subseteq C$ be wide subcategories. Then we have the following:

10 FINITE REEDY CATEGORIES AS LOCALIZATIONS OF FINITE DIRECT CATEGORIES

- Under the assumption (3) in Definition 2.8, the subcategory C₋ (resp. C₊) is isomorphic to the category obtained by freely adjoining identities to some semicategory if and only if any morphism in C₋ (resp. C₊) is either an identity or non-identity.
- (2) Under the assumption (1) in Definition 2.8, the dichotomy of morphisms into identities and non-identities in C₋ (resp. C₊) is equivalent to the decidability of the membership of a morphism in C₊ (resp. C₋).

Proof. Follows from Lemma 2.6 and the fact that $C_- \cap C_+$ consists only of identities.

Corollary 2.10. If a category C and a pair of wide subcategories $C_{-}, C_{+} \subseteq C$ satisfies (1) and (3) in Definition 2.8, then the followings are equivalent:

- (1) (C, C_{-}, C_{+}) is a Reedy category.
- (2) The membership properties of a morphism in C_{-} and in C_{+} are decidable.
- (3) Any morphism in C is either an identity or non-identity.
- (4) Any morphism in C_{-} or C_{+} is either an identity or non-identity.

Proof. Follows from the previous lemma.

The condition (3) in Definition 2.8 is equivalently stated as the existence of a *degree function*. A degree function of C, in the sense of Reedy categories, is by definition a function $F: \operatorname{Ob}(C) \to W$, where W is some well-ordered set, such that all the non-identities $x \to y$ in C_+ and all the non-identities $x \leftarrow y$ in C_- satisfy F(x) < F(y). Some literature, including Hovey [9], requires the equipment of a degree function, but its existence is enough for us.

With this definition of Reedy categories, $Mor(C_{-})$ and $Mor(C_{+})$ are decidable subsets of Mor(C). We also see that C_{-} is an inverse category, and C_{+} a direct category, as can be proven classically.

In later sections, we will frequently draw a commutative diagram in a Reedy category. Within a diagram in a fixed Reedy category $C = (C, C_{-}, C_{+})$, we depict morphisms in C_{-} and C_{+} with \rightarrow and \rightarrow , respectively, to maintain the conciseness of these diagrams. Note that \rightarrow and \rightarrow will not necessarily stand for epimorphisms and monomorphisms when we consider a diagram in general Reedy categories, deviating from the general convention.

We shall cite a useful lemma concerning Reedy categories:

Lemma 2.11 (Lemma 3.9 from [5]). All idempotents in a Reedy category are split. For more precision, let (C, C_{-}, C_{+}) be a Reedy category and suppose that a morphism $\alpha: c \to c$ satisfies $\alpha \circ \alpha = \alpha$. Then there exists a unique pair of $\sigma \in \text{Mor } C_{-}$ and $\delta \in \text{Mor } C_{+}$ with $\delta \circ \sigma = \alpha$ and $\sigma \circ \delta = \text{id}$.

Proof. Uniqueness is clear. Let the following be the (C_{-}, C_{+}) -factorizations:

$$\alpha = \delta \circ \sigma, \quad \sigma \in \operatorname{Mor} C_{-}, \ \delta \in \operatorname{Mor} C_{+}; \tag{2.1}$$

$$\sigma \circ \delta = \delta' \circ \sigma', \sigma' \in \operatorname{Mor} C_{-}, \delta' \in \operatorname{Mor} C_{+}.$$
(2.2)

Since α is idempotent, we have:

$$\sigma \circ \delta = \alpha = \alpha \circ \alpha = \sigma \circ \delta \circ \sigma \circ \delta = \sigma \circ \sigma' \circ \delta' \circ \delta.$$

The uniqueness of (C_-, C_+) -factorization, the directness of C_+ , and the inverseness of C_- together imply $\sigma' = \delta' = id$, which reduces eq. (2.2) to $\sigma \circ \delta = id$. Combining this with eq. (2.1), we obtain the desired result.

As we declared in Section 1, we shall fix a Reedy category $C = (C, C_{-}, C_{+})$ in Section 3 and onward.

2.4. The simplex category. We will frequently use the simplex category Δ in this paper. Here we recall its definition and some of its properties. Remember we write **Poset** for the category of reflexively partially ordered sets (simply posets) and order-preserving maps between them.

Definition 2.12 (The simplex category). The category Δ_a , called the *augmented* simplex category, is the full subcategory of the **Poset** spanned by finite ordinals: the linearly ordered sets $[n] \coloneqq n + 1 = \{0 < 1 < \dots < n\}$ for integers $n \ge -1$. The simplex category Δ is the full subcategory of Δ_a spanned by inhabited finite ordinals: [n] for $n \ge 0$.

Definition 2.13 (The canonical Reedy structure). We shall write $(\Delta_a)_-$ and $(\Delta_a)_+$ for the wide subcategories of Δ_a consisting of surjections (or *degeneracies*) and injections (or *face maps*), respectively. The pair $((\Delta_a)_-, (\Delta_a)_+)$ is a Reedy structure on Δ_a , which is considered as the canonical one on Δ_a . The restrictions $\Delta_- := (\Delta_a)_- \cap \Delta$ and $\Delta_+ := (\Delta_a)_+ \cap \Delta$ form the canonical Reedy structure on Δ_a .

Notation 2.14. We shall employ the conventional notation for morphisms in Δ_a (and hence in Δ). If $0 \le k \le n$, the morphisms $\delta_k^n : [n-1] \rightarrow [n]$ in $(\Delta_a)_+$ and $\sigma_k^n : [n+1] \twoheadrightarrow [n]$ in $(\Delta_a)_-$ are given by:

$$\begin{split} \delta^n_k(i) &\coloneqq \begin{cases} i & \text{if } 0 \leq i < k; \\ i+1 & \text{if } k \leq i < n; \end{cases} \\ \sigma^n_k(i) &\coloneqq \begin{cases} i & \text{if } 0 \leq i \leq k; \\ i-1 & \text{if } k < i \leq n+1 \end{cases} \end{split}$$

Also, We add some non-conventional notation. Let $[n] \in \Delta_a$ and say we have the following subset:

$$S = \{s_0 < s_1 < s_2 < \ldots < s_{k-1}\} \subseteq [n].$$

Then we write $\iota_S^n = \iota_S \colon [k-1] \to [n]$ for the morphism in $(\Delta_a)_+$ defined by $\iota_S(i) = s_i$. The particular case where S is a singleton $S = \{s\}$ is also denoted by $\iota_S^n = \iota_S \coloneqq \iota_{\{s\}}^n \colon [0] \to [n]$.

2.5. Simplicial sets, quasicategories and $(\infty, 1)$ -categories. In this subsection, we partially review the theory of simplicial sets and quasicategories. The notable textbook references for this are Lurie's [16, 15]. Especially, [16] will be repeatedly referred to in Sections 8 and 9. The content of this subsection will only be used in Sections 8 and 9, so we will use classical logic here.

Definition 2.15 (Simplicial sets). A simplicial set is a functor $\Delta^{\text{op}} \to \text{Set}$. The category of simplicial sets, which is the functor category, is denoted by Set_{Δ} . The morphisms in the category are called simplicial maps. If $X \in \text{Set}_{\Delta}$, an *n*-simplex of X is an element of X_n , which stands for X evaluated at $[n] \in \Delta$. An *n*-simplex σ is degenerate if it is in the image of the map induced by some non-identity morphism in Δ_- ; it is non-degenerate otherwise. A simplicial set is finite if it has finitely many non-degenerate simplices across all dimensions.

Notation 2.16 (Simplices). We set $\Delta := \mathbf{y} : \mathbf{\Delta} \to \mathbf{Set}_{\mathbf{\Delta}}$ for the special case of the Yoneda embedding. We may extend this functor to $\mathbf{\Delta}_a$ by setting Δ to be:

$$\Delta_a \xrightarrow{\mathbf{y}} \operatorname{Set}_{\Delta_a} \xrightarrow{\operatorname{restrict}} \operatorname{Set}_{\Delta}.$$

In other words, $\Delta[-1] \coloneqq \emptyset$.

Notation 2.17 (degenerate edge). Let X be a simplicial set. If $x \in X_0$ is a vertex of X, its corresponding degenerate edge $(\sigma_0^0)^*(x) = X(\sigma_0^0)(x)$ is also denoted by id_x .

Notation 2.18 (Boundaries and horns). The simplicial sets $\partial \Delta[n] \subseteq \Delta[n]$, for $n \ge 0$, and $\Lambda_k[n] \subseteq \partial \Delta[n]$, for $0 \le k \le n$, are defined as follows:

$$\begin{split} \partial \Delta[n]_l &\coloneqq \{\alpha \colon [l] \to [n] \mid \alpha \text{ is not surjective} \} \,, \\ \Lambda_k[n]_l &\coloneqq \{\alpha \colon [l] \to [n] \mid \{0, \dots, l-1, l+1, \dots, n\} \not\subseteq \operatorname{Im} \alpha \} \,. \end{split}$$

Definition 2.19 (lifting property). Let C be a (possibly large) category and $f: A \to B, g: X \to Y$ be morphisms in C. We say that the morphism f has *left lifting property* with respect to g, or that g has *right lifting property* with respect to f, if for any morphism $u: A \to X$ and $v: B \to Y$ commutating the following outer square, there is a morphism that fits into the dotted line, commutating the two triangles:

$$\begin{array}{ccc} A & \stackrel{u}{\longrightarrow} X \\ f \downarrow & & \stackrel{\pi}{\searrow} \\ B & \stackrel{\pi}{\longrightarrow} Y \end{array}$$

Definition 2.20 (inner fibration, inner anodyne morphism, and quasicategories). Let $f: X \to Y$ be a morphism of simplicial sets. We say that f is an *inner fibration* if it has the right lifting property with respect to all horn inclusions $\Lambda_k[n] \hookrightarrow [n]$, for all pair of integers 0 < k < n. We say that a simplicial set X is a *quasicategory* if the canonical morphism $X \to *$ to the terminal object is an inner fibration. We say that a morphism $i: A \to B$ of simplicial sets is *inner anodyne* if it has the left lifting property with respect to all inner fibrations.

We refer the reader to [16, 15] for:

- equivalences (or isomorphisms) in a quasicategory;
- equivalences of quasicategories and its generalization categorical equivalences;
- joins of simplicial sets, denoted $X \star Y$.

Notation 2.21 (function complex). The exponential object in the category of simplicial sets will be denoted by $Fun(\bullet, \bullet)$, i.e.:

$$\operatorname{Fun}(X,Y)_n \coloneqq \operatorname{Hom}_{\operatorname{\mathbf{Set}}_{\Delta}}(\Delta[n] \times X,Y).$$

We now proceed to the definition of localization. In [16], the following simplicial subset is denoted by $\operatorname{Fun}(X[E^{-1}], Q)$. However, to avoid the misleading impression that it involves a simplicial set called $X[E^{-1}]$, we instead adopt a modified version of the notation used for the theory of marked simplicial sets in [15].

Notation 2.22 (complex of maps inverting some edges, cf. [15, Chapter 3]). Let $X \in \mathbf{Set}_{\Delta}$ be a simplicial set, $E \subseteq X_1$ a set of edges, and Q a quasicategory. We define a simplicial subset $\operatorname{Fun}((X, E), Q^{\natural}) \subseteq \operatorname{Fun}(X, Q)$. The set $\operatorname{Fun}((X, E), Q^{\natural})_0$ of vertices is defined to be the set of simplicial maps $X \to Q$ that send edges in E to equivalences in Q. The set $\operatorname{Fun}((X, E), Q^{\natural})_n$ of *n*-simplices is defined to be the set of all *n*-simplices $f \in \operatorname{Fun}(X, Q)_n$ whose vertices are in $\operatorname{Fun}((X, E), Q^{\natural})_0$.

This is the definition of localization that appears in our main theorem:

Definition 2.23 (localization [16, Definition 01MP]). Let $f: X \to Y$ be a simplicial map, and $W \subseteq X_1$ be a collection of edges. We say that f exhibits Y

as an $((\infty, 1)$ -)localization of X at, or with respect to, W, if for any quasicategory Q, the precomposition functor $f^* \colon \operatorname{Fun}(Y, Q) \to \operatorname{Fun}(X, Q)$ factors through $\operatorname{Fun}((X, W), Q^{\natural}) \subseteq \operatorname{Fun}(X, Q)$, and induces an equivalence of quasicategories

$$\operatorname{Fun}(Y,Q) \to \operatorname{Fun}((X,W),Q^{\natural}).$$

An $((\infty, 1)$ -)*localization map* is a simplicial map that exhibits the target as a localization of the source at some collection of edges.

Of course, this notion is a refinement of the 1-localization: according to [16, Remark 01MV], if f exhibits Y as an $(\infty, 1)$ -localization of X at $W \subseteq X_1$ and if h X and h Y denote the *homotopy category* of X and Y, respectively, then the induced functor h f: h $X \to$ h Y is a weak 1-localization of h X at the image of W under the canonical map $X \to N(h X)$.

To facilitate our argument, we introduce a stronger notion of localization:

Definition 2.24 (universal localization [16, Definition 02M0, Proposition 02M1]). Let $f: X \to Y$ be a simplicial map. Then f is said to be *universally localizing* or to be a *universal localization* if one, and hence both, of the following equivalent conditions hold:

- (1) Let $\varphi \colon \Delta[n] \to Y$ be any simplex of Y. Then the pullback $X \times_Y \Delta[n] \to \Delta[n]$ of f along φ is a localization map.
- (2) Let $\varphi: Z \to Y$ be any simplicial map from an arbitrary simplicial set. Then the pullback $X \times_Y Z \to Z$ of f along φ exhibits Z as an $(\infty, 1)$ -localization of $X \times_Y Z$ at the preimage of the degeneracy edges of Z.

We need some closure properties of localizations. The localizations and universal localizations are closed under filtered colimits [16, Propositions 01N6 and 02M9] and certain pushouts [16, Propositions 01N7 and 02MA] in $(\mathbf{Set}_{\Delta})^{[1]}$. Easier to verify is the closure under products:

Lemma 2.25 (closure under product of localizations, [16, Proposition 02LV]). Let K be a simplicial set. Assume that a simplicial map $f: X \to Y$ exhibits Y as a localization of X at $W \subseteq X_1$. Then the induced map $f \times id_K: X \times K \to Y \times K$ on products exhibits $Y \times K$ as a localization of $X \times K$ at:

$$W \times K_0 \coloneqq \{(e, \mathrm{id}_x) \mid e \in W, x \in K_0\} \subseteq (X \times K)_1.$$

Proof. See [16, Proposition 02LV]; this follows from the simple computation of mapping simplicial sets. \Box

Lemma 2.26 (closure under product of universal localizations). Let $f: X \to Y$ be a universally localizing simplicial map, and let K be a simplicial set. Then the induced map $f \times id_K: X \times K \to Y \times K$ on products is universally localizing.

Proof. If the left square in the following diagram is a pullback, then the whole rectangle is a pullback:

The proof of the closure under join is contributed by Maxime Ramzi as an answer [10] to my MathOverflow question:

Lemma 2.27 (closure under join of localizations, [10]). Let K be a simplicial set. Assume that a simplicial map $f: X \to Y$ exhibits Y as a localization of X at $W \subseteq X_1$. Then the induced map $f \star id_K \colon X \star K \to Y \star K$ on joins exhibits $Y \star K$ as a localization of $X \star K$ at:

$$W \subseteq X_1 = (X \star \emptyset)_1 \subseteq (X \star K)_1.$$

Proof. [16, Construction 01HN] constructs the following commutative cube of simplicial sets:

As [16, Proposition 01HP] states, the back and the front faces are *categorical* pushout squares in the sense of [16, Definition 01F7], a.k.a. homotopy pushout squares with respect to the Joyal model structure on \mathbf{Set}_{Δ} . Then [16, Proposition 01N7] implies that, in order to demonstrate our lemma, it suffices to check that the three maps that connect the pushout-defining spans are localization maps at appropriate sets of edges. We inspect the construction of the commutative cube to see that we need to show all of the following:

- $f \times id: X \times \partial \Delta[1] \times K \to Y \times \partial \Delta[1] \times K$ is a localization at $W \times (\partial \Delta[1] \times K)_0$;
- $f \times \mathrm{id} \colon X \times \Delta[1] \times K \to Y \times \Delta[1] \times K$ is a localization at $W \times (\Delta[1] \times K)_0$;
- $f \amalg \operatorname{id}: (X \times \{0\}) \amalg (\{1\} \times K) \to (Y \times \{0\}) \amalg (\{1\} \times K)$ is a localization at $W \times \{0\}$.

All of these are trivial or follow from Lemma 2.25.

Corollary 2.28 (closure under join of universal localizations). Let $f: X \to Y$ be a universally localizing simplicial map, and let K be a simplicial set. Then the induced map $f \star id_K: X \star K \to Y \star K$ on joins is universally localizing.

Proof. Let $\varphi \colon \Delta[n] \to Y \star K$ be any simplex of $Y \star K$. Then there are objects $[m_0], [m_1] \in \text{Ob} \, \Delta_a$ with $m_0 + m_1 + 1 = n$, and simplicial maps $\varphi_0 \colon \Delta[m_0] \to Y$ and $\varphi_1 \colon \Delta[m_1] \to K$ with:

$$\varphi = \varphi_0 \star \varphi_1 \colon \Delta[n] = \Delta[m_0] \star \Delta[m_1] \to Y \star K.$$

Now, note that the following diagram is a pullback of simplicial sets:

$$\begin{array}{ccc} (X \times_Y \Delta[m_0]) \star \Delta[m_1] \longrightarrow X \star K \\ & & & \downarrow \\ & & \downarrow \\ \Delta[m_0] \star \Delta[m_1] \xrightarrow{\varphi_0 \star \varphi_1} Y \star K \end{array}$$

Therefore, it suffices to show that the left vertical map is a localization map. This is a direct consequence of Lemma 2.27. $\hfill \Box$

We remember few more notations:

Notation 2.29 (nerve). Let C be a category. The *nerve* N(C) of C is the simplicial set defined by

$$N(C)_n \coloneqq \operatorname{Hom}_{\operatorname{Cat}}([n], C).$$

It is common to write just C for the nerve of C, but we will use N(C) to facilitate the notation for colimits.

Notation 2.30 (skeleton). Given a simplicial set X, the *n*-skeleton of X is denoted by $\operatorname{Sk}_n X$. To be specific, let $\Delta_{\leq n}$ denote the full subcategory of Δ spanned by $[0], \ldots, [n]$, and consider the restriction functor $\operatorname{tr}_n \colon \operatorname{Set}_{\Delta} \to \operatorname{Set}_{\Delta\leq n}$ and its left adjoint $\operatorname{sk}_n \colon \operatorname{Set}_{\Delta\leq n} \to \operatorname{Set}_{\Delta}$. Then we have $\operatorname{Sk}_n = \operatorname{sk}_n \circ \operatorname{tr}_n$.

To conclude this section, we list some references to lemmas that will be useful to prove that a simplicial map is a monomorphism or an inner anodyne map. We begin with the following definition, which is here merely for the formulation of Lemma 2.32:

Definition 2.31. Let C be a category, and $L, R \subseteq \text{Mor } C$ be classes of morphisms. We say that (L, R) is a *weak factorization system* on C if the following conditions are satisfied:

- for any morphism $f: x \to y$ in C, there exist an object $z \in Ob C$ and morphisms $l: x \to z$ in L and $r: z \to y$ in R such that $f = r \circ l$;
- $f \in Mor C$ is in L if and only if it has the right lifting property with respect to all morphisms in R;
- $f \in Mor C$ is in R if and only if it has the left lifting property with respect to all morphisms in L.

Notice that L may be monomorphisms or inner anodyne maps in \mathbf{Set}_{Δ} .

Lemma 2.32. Let (L, R) be a weak factorization system on a bicomplete category C. Let the following be a commutative diagram in C:



If the morphisms $a \to x$, $b \to y$, and $y \cup_b c \to z$ induced by the diagram are in L, then the morphism $a \cup_b c \to x \cup_y z$ between pushouts belongs to L.

Proof. We regard C as a model category by taking L as the class of cofibrations, R as the class of fibrations, and all morphisms as weak equivalences. Consider the Reedy category $S = \{0 \leftarrow 1 \rightarrow 2\}$, where the Reedy structure is shown by the shape of the arrows. As the commutative diagram in the statement may then be seen as a cofibration in the Reedy model category C^S , the result follows if we can show that colim: $C^S \rightarrow C$ is a left Quillen functor. Since Proposition 15.10.2 from Hirschhorn [8] demonstrates that S has fibrant constants in the sense of Definition 15.10.1 from the same book, we can apply Theorem 15.10.8 from [8] to conclude that colim is a left Quillen functor.

The following lemma is useful and I believe it is a well-known folklore, but I could not find a reference. Therefore I include a proof here. Note that the class I of morphisms can be the class of injections or the class of inner anodyne maps in \mathbf{Set}_{Δ} :

Lemma 2.33. Let C be a cocomplete (large) category. Let $I \subseteq \text{Mor } C$ be a class of morphisms, closed under transfinite compositions and pushouts. More specifically, assume that I satisfies the following conditions:

• Let $\alpha > 0$ be an ordinal and $F: \alpha \to C$ be a functor with $F(!_{\beta,\beta+1}) \in I$ for any β with $\beta + 1 < \alpha$. If F preserves all colimits of the diagrams $\beta \to \alpha$ for $\beta < \alpha$, then the canonical morphism $F(0) \to \operatorname{colim}_{\beta < \alpha} F(\beta)$ is in I. If f: x → y is a morphism in I and g: x → z is a morphism in C, then the pushout z → y ∪_x z of f along g is in I.

Let $F, G: \Delta \to C$ be functors. Let the colimit-preserving functors $\hat{F}, \hat{G}: \mathbf{Set}_{\Delta} \to C$ be defined by the left Kan extensions of F and G along Δ , as follows:

$$\hat{F}(X) \coloneqq \operatorname{colim}_{\Delta[n] \to X} F[n], \qquad \qquad \hat{G}(X) \coloneqq \operatorname{colim}_{\Delta[n] \to X} G[n].$$

Assume that a natural transformation $\theta: F \Rightarrow G$ is Reedy in I, i.e., for each $[n] \in \Delta$, the morphism $F[n] \cup_{\hat{F}(\partial \Delta[n])} \hat{G}(\partial \Delta[n]) \to G[n]$ induced by θ belongs to I. Then, for any monomorphism $i: A \hookrightarrow B$ in \mathbf{Set}_{Δ} , the map $\hat{F}(B) \cup_{\hat{F}(A)} \hat{G}(A) \to \hat{G}(B)$ induced by θ and i is in I. In particular, the constituent simplicial maps of the natural transformation $\hat{\theta}: \hat{F} \Rightarrow \hat{G}$ induced by θ all belong to I.

Proof. As is well known, the monomorphism $i: A \to B$ can be written as a transfinite composition of the pushouts of the maps of the form $\partial \Delta[n] \to \Delta[n]$. Let us take such a transfinite composition $X: \alpha + 1 \to \mathbf{Set}_{\Delta}$. To be more explicit: the functor $X: \alpha + 1 \to \mathbf{Set}_{\Delta}$ has the successor of an ordinal α as its domain, satisfies $X(0) = A, X(\alpha) = B$ and $X(!_{0,\alpha}) = i$, and preserves colimits of the diagrams $\beta \to \alpha \to \alpha + 1$ for $\beta \leq \alpha$. Also, for each $\beta < \alpha$, we have a pushout diagram in \mathbf{Set}_{Δ} of the following form:

$$\begin{array}{c} \partial \Delta[n_{\beta}] & \longrightarrow & \Delta[n_{\beta}] \\ \downarrow & \qquad \downarrow \\ X(\beta) & \stackrel{X(!)}{\longleftarrow} & X(\beta+1) \end{array}$$

Define a functor $Y: \alpha + 1 \to C$ by $Y(\beta) \coloneqq \hat{F}(B) \cup_{\hat{F}(X(\beta))} \hat{G}(X(\beta))$. Since \hat{F} and \hat{G} preserve colimits and X preserves colimits of the diagrams $\beta \hookrightarrow \alpha \hookrightarrow \alpha + 1$ for $\beta \leq \alpha$, we see that Y preserves colimits of such diagrams as well. We also see that $Y(!_{0,\alpha})$ is the morphism $\hat{F}(B) \cup_{\hat{F}(A)} \hat{G}(A) \to \hat{G}(B)$ in question. Therefore, it suffices to prove that $Y(!): Y(\beta) \to Y(\beta + 1)$ is in I for each $\beta < \alpha$, for it and our assumption on I together imply that $Y(!_{0,\alpha})$ is in I.

To see this remaining claim, let $\beta < \alpha$ and notice that there is the following pushout square in C:



Since the top vertical arrow is in I by assumption, the bottom arrow is also in I, as I is closed under pushouts. This completes the proof.

In combination with the previous lemma, the following lemmas are useful. The proof can be found in Section 14.3 from Riehl [18]:

Definition 2.34 ([18, Section 14.3]). Let C be a small category and $F: \Delta \to \mathbf{Set}^C$ be a functor. Then F is said to be *unaugmentable* if the pullback of the two morphisms $F(\delta_0^1), F(\delta_1^1): F[0] \Rightarrow F[1]$ is empty, i.e., the initial object.

Lemma 2.35 (A part of [18, Lemma 14.3.8]). Let C be a small category and $F: \Delta \to \mathbf{Set}^C$ be a functor. Then the unique natural transformation $\emptyset \Rightarrow F$ from the initial functor to F is a Reedy monomorphism if and only if F is unaugmentable.

Corollary 2.36. Let C be a small category and $F: \Delta \to \mathbf{Set}^C$ be a functor. Then the left Kan extension $\mathbf{Set}_{\Delta} \to \mathbf{Set}^C$ of F along the Yoneda embedding $\Delta: \Delta \to \mathbf{Set}_{\Delta}$ preserves monomorphism if and only if F is unaugmentable.

Lemma 2.37 (The rest of [18, Lemma 14.3.8]). Let C be a small category and $F, G: \Delta \to \mathbf{Set}^C$ be unaugmentable functors. Then a natural transformation $F \Rightarrow G$ is a Reedy monomorphism if and only if it is a pointwise monomorphism.

Corollary 2.38. Let C be a small category and $F, G: \Delta \to \mathbf{Set}^C$ be unaugmentable functors. Let $\theta: F \Rightarrow G$ be a natural transformation. Let $\hat{F}, \hat{G}: \mathbf{Set}_{\Delta} \to \mathbf{Set}^C$ denote the left Kan extensions of F and G along the Yoneda embedding $\Delta: \Delta \to \mathbf{Set}_{\Delta}$. Then the induced natural transformation $\hat{\theta}: \hat{F} \Rightarrow \hat{G}$ is a pointwise monomorphism if and only if θ is so.

3. Several categories related to a Reedy category

We remind the reader that we will fix a Reedy category C for the rest of the paper. In this section, we construct the category Down(C) from C. I, the author, appreciate Lumsdaine's helpful answer [11] to my MathOverflow question, which has reformulated the categories constructed in this section using Grothendieck construction and categorical nerves. This reformulation is explicitly included in Definition 3.1 immediately below and further discussed in Remark 3.4.

Our construction begins with the following category, $\int \mathbf{N}(C)$, which appears to be standard, although we could not locate a convenient reference.

Definition 3.1 $(\int \mathbf{N}(C))$. We shall denote by $\int \mathbf{N}(C)$ the total category of the Grothendieck construction of the categorical nerve

$$\mathbf{N}(C): \mathbf{\Delta}^{\mathrm{op}} \to \mathbf{Cat}; \ [n] \mapsto C^{[n]}$$

Explicitly, the category $\int \mathbf{N}(C)$ is described as follows:

- An object of $\int \mathbf{N}(C)$ is a pair ([n], X) of an object $[n] \in Ob \Delta$ and a functor $X : [n] \to C$.
- A morphism $([m], X) \to ([n], Y)$ is a pair (α, θ) of a morphism $\alpha \colon [m] \to [n]$ in Δ and a natural transformation $\theta \colon X \Rightarrow Y \circ \alpha$.
- The composition of

$$(\alpha, \theta) \colon ([l], X) \to ([m], Y), \text{ and}$$

 $(\beta, \phi) \colon ([m], Y) \to ([n], Z)$

is
$$(\beta \circ \alpha, (\phi \triangleright \alpha) \circ \theta)$$
.

In order to intuitively explain the category $\int \mathbf{N}(C)$, we compare it to the functor category $C^{[n]}$. Diagrammatically, the latter category consists of objects which are vertical chains of n composable morphisms in C, and morphisms which are horizontal commutative ladders consisting of n + 1 rungs connecting between those objects, as depicted in:



By contrast, the category $\int \mathbf{N}(C)$ has vertical chains of any finite length as objects, and skew ladders as morphisms, where by "skew" we mean that the rungs can be non-horizontal, as long as each object in the source chain connects to exactly one rung and the rungs do not cross midway, as shown in:



Since our actual interest is in two subcategories of $\int \mathbf{N}(C)$, we proceed to their definitions.

Definition 3.2 $(\int \mathbf{N}^{-,+}(C))$. The subcategory $\int \mathbf{N}^{-,+}(C)$ of $\int \mathbf{N}(C)$ is defined to consist of:

- the objects ([n], X) in which X can be considered as a functor $[n] \to C_-$, and
- the morphisms (α, θ) : $([m], X) \to ([n], Y)$ such that each constituent morphism

$$\theta_i \colon X(i) \to Y(\alpha(i))$$

lies in C_+ for $i \in [m] = \{0, 1, \dots, m\}.$

Definition 3.3 $(\int \mathbf{N}_{+}^{-,+}(C))$. We will write $\int \mathbf{N}_{+}^{-,+}(C)$ for the subcategory of $\int \mathbf{N}^{-,+}(C)$ specified by:

- its objects are the objects $([n], X) \in \text{Ob} \int \mathbf{N}^{-,+}(C)$ such that $X: [n] \to C_{-}$ is conservative, or equivalently reflects identities;
- its morphisms are the morphisms (α, θ) : $([m], X) \to ([n], Y)$ in $\int \mathbf{N}^{-,+}(C)$ such that α lies within Δ_+ .

Although we shall investigate $\int \mathbf{N}^{-,+}(C)$ and $\int \mathbf{N}^{-,+}_+(C)$ mainly using the explicit and combinatorial formulation as the categories of pairs, it might be helpful to justify the symbols having categorical intent:

Remark 3.4 (Lumsdaine [11]). The categories $\int \mathbf{N}^{-,+}(C)$ and $\int \mathbf{N}^{-,+}_+(C)$, as the symbols indicate, are Grothendieck constructions of functors

$$\mathbf{N}^{-,+}(C) \colon \mathbf{\Delta}^{\mathrm{op}} \to \mathbf{Cat}, \text{ and}$$

 $\mathbf{N}^{--,+}_+(C) \colon \mathbf{\Delta}^{\mathrm{op}}_+ \to \mathbf{Cat}.$

Here, the functor $\mathbf{N}^{-,+}(C)$ is a subfunctor of $\mathbf{N}(C)$, and $\mathbf{N}^{--,+}_+(C)$ is also a subfunctor of the restriction

$$\mathbf{\Delta}^{\mathrm{op}}_+ \longleftrightarrow \mathbf{\Delta}^{\mathrm{op}} \overset{\mathbf{N}^{-,+}(C)}{\longrightarrow} \mathbf{Cat}$$

We do not need the actual definitions of the two functors for the argument in this paper, but they should be clear from Definitions 3.2 and 3.3. As the two categories in question are Grothendieck constructions, they come equipped with Grothendieck fibrations

$$\int \mathbf{N}^{-,+}(C) \to \mathbf{\Delta}, \text{ and}$$
$$\int \mathbf{N}^{--,+}_+(C) \to \mathbf{\Delta}_+.$$

We will utilize these two functors later in this section, but we need not see them as a part of Grothendieck construction. They can be considered as the mere first projections, in accordance with the explicit constructions in Definitions 3.2 and 3.3.

The purpose of the next lemma is another remark we need to make. Although Definition 3.3 imposes a condition on the morphisms of $\int \mathbf{N}^{-,+}(C)$ to define those of $\int \mathbf{N}^{-,+}_+(C)$, it turns out that these conditions are always inherently satisfied; hence the condiction is logically redundant:

Lemma 3.5. The subcategory $\int \mathbf{N}_{+}^{-,+}(C) \subseteq \int \mathbf{N}^{-,+}(C)$ is full. Explicitly, if

$$([m], X), ([n], Y) \in \operatorname{Ob} \int \mathbf{N}_{+}^{--,+}(C) \subseteq \operatorname{Ob} \int \mathbf{N}^{-,+}(C)$$

are objects and (α, θ) : $([m], X) \to ([n], Y)$ is a morphism in $\int \mathbf{N}^{-,+}(C)$, then the morphism α : $[m] \to [n]$ in Δ also belongs to Δ_+ , qualifying (α, θ) as an arrow in $\int \mathbf{N}^{-,+}_+(C)$.

Proof. Set (α, θ) : $([m], X) \to ([n], Y)$ as assumed in the statement. It is sufficient to show α : $[m] \to [n]$ is injective. Assume that $0 \le i \le j \le m$ satisfy $\alpha(i) = \alpha(j)$. The naturality of θ implies the commutativity of the following diagram:

$$\begin{array}{ccc} X(i) & \xrightarrow{\theta_i} & Y(\alpha(i)) \\ X(!) & & & \\ X(j) & \xrightarrow{\theta_i} & Y(\alpha(j)) \end{array}$$

Here, $! = !_{ij}$ is the unique morphism $i \to j$ in [m]. In the diagram, the uniqueness of (C_-, C_+) -factorization applies and we obtain $X(!_{ij}) = id$. Since X is assumed to reflect identity by the definition of Ob $\int \mathbf{N}_+^{-,+}(C)$, we have i = j, as required. \Box

We proceed to the definitions of Down(C) and its inherently related category $\text{Down}_*(C)$. For that purpose, we first need to enrich the three previously defined categories over the category **Poset** of (reflexively) partially ordered sets, since our desired categories are obtained by changing the enriching base of the two **Poset**-categories $\int \mathbf{N}^{-,+}(C)$ and $\int \mathbf{N}^{--,+}_+(C)$.

Remember the canonical **Poset**-enrichment of Δ as a subcategory of the cartesian closed category **Poset**. Explicitly stated, for any parallel pair of morphisms $\alpha, \alpha' : [m] \to [n]$, we consider that $\alpha \leq \alpha'$ if and only if we have $\alpha(i) \leq \alpha'(i)$ for all $i \in [m] = \{0, 1, \ldots, m\}$. This enrichment induces the following **Poset**-enrichments:

Definition 3.6. Let $C = (C, C_{-}, C_{+})$ be as above. Let

$$(\alpha, \theta), (\alpha', \theta') \colon ([m], X) \to ([n], Y)$$

be a parallel pair of morphisms in any of the three categories $\int \mathbf{N}(C)$, $\int \mathbf{N}^{-,+}(C)$, and $\int \mathbf{N}^{-,+}_+(C)$. We say $(\alpha, \theta) \leq (\alpha', \theta')$ if and only if it holds that $\alpha \leq \alpha'$ and the following diagram of natural transformations is commutative:

$$X \xrightarrow[\theta]{\theta'} Y \circ \alpha \\ \downarrow Y \circ \alpha'$$

Here, the unique natural transformation $!: \alpha \Rightarrow \alpha'$ corresponds to the inequality $\alpha \leq \alpha'$ and $Y \triangleleft !$ stands for its left whiskering with Y. With this partial order, we regard the three categories as **Poset**-enriched categories, and hence as 2-categories.

Let $\pi_0: \mathbf{Poset} \to \mathbf{Set}$ denote the left adjoint functor of the discrete poset functor $\mathbf{Set} \to \mathbf{Poset}$. The functor π_0 sends a poset P to the quotient of the set P by the symmetric transitive closure of the partial order on P. Since this functor is strongly cartesian monoidal, the following Definition 3.8, of the category we want to construct in this section, makes sense.

Definition 3.7. Let

$$(\alpha, \theta), (\beta, \phi) \colon ([m], X) \to ([n], Y)$$

be a parallel pair of morphisms in any of the three categories $\int \mathbf{N}(C)$, $\int \mathbf{N}^{-,+}(C)$, and $\int \mathbf{N}^{-,+}_+(C)$. We say they are *equivalent* and write $(\alpha, \theta) \sim (\beta, \phi)$ if the two arrows are related by the symmetric transitive closure of the order on the hom-set $\operatorname{Hom}(([m], X), ([n], Y))$.

Definition 3.8 (Down(C)). Let $C = (C, C_{-}, C_{+})$ be a Reedy category. The category Down(C) is defined to be the **Set**-category obtained by changing the enriching base of $\int \mathbf{N}_{+}^{-,+}(C)$ by the functor $\pi_{0} \colon \mathbf{Poset} \to \mathbf{Set}$. To be more explicit, Down(C) has the same set of objects as $\int \mathbf{N}_{+}^{-,+}(C)$, and we declare the hom-sets of Down(C) to be the quotients of the original hom-sets by the equivalence we detailed above. Similarly, we define Down_{*}(C) to be the change of enriching base by π_{0} of $\int \mathbf{N}^{-,+}(C)$.

There are two paths of construction of the category $\operatorname{Down}(C)$ from $\int \mathbf{N}^{-,+}(C)$. As we have done, we may first take the full subcategory $\int \mathbf{N}_{+}^{-,+}(C) \subseteq \int \mathbf{N}^{-,+}(C)$, and then take the quotient in hom-sets to obtain $\operatorname{Down}(C)$. On the other hand, as will be seen in Remark 6.1, we can also regard $\operatorname{Down}(C)$ as a full subcategory of $\operatorname{Down}_{*}(C)$, which is obtained by taking quotients in hom-sets of $\int \mathbf{N}^{-,+}(C)$. The two intermediate categories $\operatorname{Down}_{*}(C)$ and $\int \mathbf{N}_{+}^{-,+}(C)$ serve different purposes in our investigation of $\operatorname{Down}(C)$. The quotient $\operatorname{Down}_{*}(C)$ is better in its categorical nature, and used in the proof that $\operatorname{Down}(C) \to C$ is a localization. The full subcategory $\int \mathbf{N}_{+}^{-,+}(C)$ is better in its combinatorial simplicity, and used in the proof that $\operatorname{Down}(C)$ is finite and direct.

Now we have defined the category Down(C); however, the combinatorial nature of the category Down(C) is still unclear. For a better understanding of this combinatorics, the next section will be devoted to the analysis of the order and the equivalence relation on the Hom-sets of $\int \mathbf{N}(C)$, $\int \mathbf{N}^{-,+}(C)$, and $\int \mathbf{N}^{+,+}_{+,-}(C)$.

4. The hom-posets of the categories of skew ladders

The symmetric transitive closure of a partial order, used in the definitions of Down(C) and $\text{Down}_*(C)$, is generally tricky to discuss and compute with. However, the simplicity of the order on the hom-sets on $\int \mathbf{N}_+^{--,+}(C)$, shown in this section, makes it manageable.

We will prepare some notations for the subsequent discussion, in which we will analyze the **Poset**-enrichment of the three categories $\int \mathbf{N}(C)$, $\int \mathbf{N}^{-,+}(C)$, and $\int \mathbf{N}^{-,+}_{+}(C)$. During the analysis, Γ will denote any of these three categories. If $p \in P$ is an element of a poset P, we shall write

$$[p,\infty]_P = [p,\infty] \coloneqq \{x \in P \mid p \le x\}$$

for the upward unbounded interval above p. If $p, q \in P$, then we set

$$[p,q]_P = [p,q] \coloneqq \{x \in P \mid p \le x \le q\}$$

to be the closed bounded interval between p and q. Also, let us remind ourselves of the notation from Definition 2.1: if $x \leq y \in P$ are elements in a poset, we write $! = !_{xy}: x \to y$ for the unique morphism in P seen as a category, as we already did in Definition 3.6 and in the proof of Lemma 3.5.

We can consider the three categories $\int \mathbf{N}(C)$, $\int \mathbf{N}^{-,+}(C)$, and $\int \mathbf{N}^{-,+}_+(C)$ as Grothendieck constructions, as stated in Definition 3.1 and Remark 3.4; thus they come equipped with Grothendieck fibrations

$$\int \mathbf{N}(C) \to \mathbf{\Delta},$$

$$\int \mathbf{N}^{-,+}(C) \to \mathbf{\Delta}, \text{ and }$$

$$\int \mathbf{N}^{--,+}_{+}(C) \to \mathbf{\Delta}_{+}.$$

They are the first projections in terms of the explicit construction in Definitions 3.1–3.3. By post-composing the inclusion $\Delta_+ \hookrightarrow \Delta$ as necessary, we get a functor pr: $\Gamma \to \Delta$.

In the following series of lemmas, we will analyze upward unbounded intervals of the hom-posets of Γ , by comparing such intervals with the intervals of the hom-posets of Δ . The functor pr will be central in this analysis.

Lemma 4.1. Let (α, θ) : $([m], X) \to ([n], Y)$ be a morphism in Γ . Then the canonical order-preserving map

pr:
$$[(\alpha, \theta), \infty|_{\operatorname{Hom}_{\Gamma}(([m], X), ([n], Y))} \to [\alpha, \infty|_{\operatorname{Hom}_{\Delta}([m], [n])},$$

induced by the functor pr above, is injective and reflects order.

Proof. Follows from the definition of the order.

Lemma 4.2. We consider the cases $\Gamma = \int \mathbf{N}^{-,+}(C)$ and $\Gamma = \int \mathbf{N}^{-,+}_+(C)$. Let

$$(\alpha, \theta) \colon ([m], X) \to ([n], Y)$$

be a morphism in Γ . Let $\beta \colon [m] \to [n]$ be a morphism of Δ with $\alpha \leq \beta$. Then the following conditions are equivalent.

(1) There exists a morphism in Γ of the form

$$(\beta, \phi) \colon ([m], X) \to ([n], Y)$$

satisfying $(\alpha, \theta) \leq (\beta, \phi)$.

(2) For each $0 \le i \le m$, the composite ϕ_i in the following commutative diagram belongs to C_+ :



Additionally, if the equivalent conditions hold, the morphism (β, ϕ) stated to exist in (1) is unique, and is specified by $\phi = (\phi_i)_{0 \le i \le m}$ using the notation from (2).

Proof. The implication $(1) \implies (2)$ and the uniqueness part is a direct consequence of the definition of the order on $\operatorname{Hom}_{\Gamma}(([m], X), ([n], Y))$. It just remains to demonstrate $(2) \implies (1)$.

Assume the condition (2). Remember Lemma 3.5; in order to construct a legitimate morphism (β, ϕ) , it suffices to prove that $\phi = (\phi_0, \phi_1, \dots, \phi_m)$ is a natural transformation. Let $0 \le i \le j \le m$. Consider the following diagram:



The commutativity of the two triangles follows from the definition of ϕ . The naturality of θ commutes the left small quadrilateral. The functoriality of Y ensures the commutativity of the right small quadrilateral. Combining them, we obtain the commutativity of the outer square, which is the naturality we desire.

Now we have ensured that

$$(\beta, \phi) \colon ([m], X) \to ([n], Y)$$

is a well-defined morphism in Γ , and the inequality $(\alpha, \theta) \leq (\beta, \phi)$ is already included in our assumption. This establishes (1), and completes the proof. \Box

Lemma 4.3. Let (α, θ) : $([m], X) \to ([n], Y)$ be a morphism in Γ . Then there is the largest element in the upward unbounded interval

$$[(\alpha,\theta),\infty[_{\operatorname{Hom}_{\Gamma}(([m],X),([n],Y))}.$$

Proof. For $\Gamma = \int \mathbf{N}(C)$, the obvious surjectivity of

pr:
$$[(\alpha, \theta), \infty[_{\operatorname{Hom}_{\{\mathbf{N}(C)}(([m], X), ([n], Y))} \to [\alpha, \infty[_{\operatorname{Hom}_{\Delta}([m], [n])}])$$

and Lemma 4.1 together imply that the unique inverse image of the maximum element of $\operatorname{Hom}_{\Delta}([m], [n])$ is the required largest element.

We proceed to the cases of $\Gamma = \int \mathbf{N}^{-,+}(C)$ and $\Gamma = \int \mathbf{N}^{-,+}_+(C)$. For each $i \in [m]$, define a natural number $\alpha'(i) \in [n] = \{0, 1, \dots, n\}$ as the biggest $\alpha(i) \leq k \leq n$ for which the following composition lies within C_+ :

$$X(i) \xrightarrow{\theta_i} Y(\alpha(i)) \xrightarrow{Y(!)} Y(k).$$

Remember that $Mor(C_+)$ is a decidable subset of Mor(C); therefore the maximum k does exist. This defines a map $\alpha' \colon [m] \to [n]$ of sets.

We need to verify that α' preserves order. Let $0 \leq i \leq j \leq m$. We wish to show $\alpha'(i) \leq \alpha'(j)$. If $\alpha'(i) < \alpha(j)$, there is nothing to prove; we consider the case $\alpha'(i) \geq \alpha(j)$. Consider the following commutative diagram:

$$\begin{array}{c|c} X(i) & \xrightarrow{\theta_i} Y(\alpha(i)) \\ X(!) & & \downarrow Y(!) \\ X(j) & \xrightarrow{\theta_j} Y(\alpha(j)) \\ \sigma & & \downarrow Y(!) \\ Z & \longrightarrow Y(\alpha'(i)) \end{array}$$

where $X(j) \to Z \to Y(\alpha'(i))$ is the (C_-, C_+) -factorization of the composite $X(j) \to Y(\alpha(j)) \to Y(\alpha'(i))$. By the definition of $\alpha'(i)$, the morphism $X(i) \to Y(\alpha'(i))$ in

the diagram belongs to C_+ ; therefore the uniqueness of (C_-, C_+) -factorization implies that $X(i) \twoheadrightarrow Z$ is an identity. It follows from the inverseness of C_- that $X(!_{ij}) = \sigma = \text{id.}$ Therefore $\alpha'(i) = \alpha'(j)$ by definition.

Now we have proved that α' is a morphism of Δ . By Lemma 4.2, there is a unique morphism in Γ of the form

$$(\alpha', \theta') \colon ([m], X) \to ([n], Y)$$

satisfying $(\alpha, \theta) \leq (\alpha', \theta')$. By the construction of α' and Lemma 4.2, we see that this is the maximum we desired.

Proposition 4.4. Let (α, θ) : $([m], X) \to ([n], Y)$ be a morphism in Γ . Then the canonical order-preserving map

pr:
$$|(\alpha, \theta), \infty|_{\operatorname{Hom}_{\Gamma}(([m], X), ([n], Y))} \to |\alpha, \infty|_{\operatorname{Hom}_{\Delta}([m], [n])}$$

has the largest element α' in its image. Furthermore, the map is an order isomorphism when restricted as

$$[(\alpha, \theta), \infty[_{\operatorname{Hom}_{\Gamma}(([m], X), ([n], Y))} \to [\alpha, \alpha']_{\operatorname{Hom}_{\Delta}([m], [n])}$$

Proof. Since pr in the statement preserves order, the map sends the maximum element $(\alpha', \theta') \in [(\alpha, \theta), \infty[_{\text{Hom}_{\Gamma}(([m], X), ([n], Y))})$, shown to exist in Lemma 4.3, to the biggest element α' in the image of the map. Thanks to Lemma 4.1, all that is left to prove is the surjectivity of

pr:
$$[(\alpha, \theta), \infty[_{\operatorname{Hom}_{\Gamma}(([m], X), ([n], Y))} \to [\alpha, \alpha']_{\operatorname{Hom}_{\Delta}([m], [n])},$$

which is obvious for $\Gamma = \int \mathbf{N}(C)$. We prove this for $\Gamma = \int \mathbf{N}^{-,+}(C)$ and $\Gamma = \int \mathbf{N}^{-,+}_+(C)$.

Let $\beta \colon [m] \to [n]$ in Δ satisfy $\alpha \leq \beta \leq \alpha'$. We would like to show that there is a morphism in Γ having the form

$$(\beta, \phi) \colon ([m], X) \to ([n], Y)$$

such that $(\alpha, \theta) \leq (\beta, \phi) \leq (\alpha', \theta')$. By Lemmas 4.1 and 4.2, it suffices to prove, for each $0 \leq i \leq m$, that the composite

$$X(i) \xrightarrow{\theta_i} Y(\alpha(i)) \xrightarrow{Y(!)} Y(\beta(i))$$

resides in C_+ . Examine the following commutative diagram, formed by successively taking (C_-, C_+) -factorization twice:

$$\begin{split} X(i) & \xrightarrow{\theta_i} Y(\alpha(i)) \\ \exists ! \downarrow & \qquad \downarrow Y(!) \\ \exists ! Z_\beta & \xrightarrow{\exists !} Y(\beta(i)) \\ \exists ! \downarrow & \qquad \downarrow Y(!) \\ \exists ! Z_{\alpha'} & \xrightarrow{\exists !} Y(\alpha'(i)) \end{split}$$

Since (α, θ) is a morphism in Γ , $X(i) \to Y(\alpha'(i))$ belongs to C_+ . From the uniqueness of (C_-, C_+) -factorization follows that $X(i) \twoheadrightarrow Z_{\alpha'}$ is an identity. The inverseness of C_- implies that $X(i) \twoheadrightarrow Z_{\beta}$ is an identity; therefore we obtain that $X(i) \to Y(\beta(i))$ is in C_+ .

Thanks to Proposition 4.4, we see that upward unbounded intervals of the homposets of Γ has a good structure: 24 FINITE REEDY CATEGORIES AS LOCALIZATIONS OF FINITE DIRECT CATEGORIES

Corollary 4.5. Let (α, θ) : $([m], X) \to ([n], Y)$ be a morphism in Γ . Then the upward unbounded interval $[(\alpha, \theta), \infty[_{\operatorname{Hom}_{\Gamma}(([m], X), ([n], Y))}]$ enjoys the structure of a bounded distributive lattice.

Proof. Let α' be as specified in Proposition 4.4. For $[\alpha, \alpha']_{\text{Hom}_{\Delta}([m], [n])}$ is a bounded distributive lattice, Proposition 4.4 directly verifies the corollary.

Now we are ready to simplify the definition of the equivalence relation on the hom-sets of Γ in a way that is easier to investigate.

Proposition 4.6. Let $(\alpha, \theta), (\alpha', \theta') \colon ([m], X) \to ([n], Y)$ be a parallel pair of morphisms in Γ . Then the following conditions are equivalent:

- (1) $(\alpha, \theta) \sim (\alpha', \theta')$.
- (2) There exists a common upper bound of (α, θ) and (α', θ') .
- (3) Let

$$(\tilde{\alpha}, \tilde{\theta}) \in [(\alpha, \theta), \infty[_{\operatorname{Hom}_{\Gamma}(([m], X), ([n], Y))}, and$$

 $(\tilde{\alpha}', \tilde{\theta}') \in [(\alpha', \theta'), \infty[_{\operatorname{Hom}_{\Gamma}(([m], X), ([n], Y))}$

be the largest elements, shown to exist in Lemma 4.3. Then they coincide: $(\tilde{\alpha}, \tilde{\theta}) = (\tilde{\alpha}', \tilde{\theta}').$

Proof. We verify each part of $(1) \implies (2) \implies (3) \implies (1)$ separately.

(1) \implies (2): By the definition of the equivalence, it suffices to show that the condition (2) is a symmetric transitive relation when seen as a relation between (α, θ) and (α', θ') . Symmetry is trivial. For transitivity, let

 $(\alpha_0, \theta_0), (\alpha_1, \theta_1), (\alpha_2, \theta_2) \colon ([m], X) \to ([n], Y)$

be a parallel triple of arrows in Γ . Assume that

$$(\beta_1, \phi_1), (\beta_2, \phi_2) \colon ([m], X) \to ([n], Y)$$

satisfy $(\alpha_i, \theta_i) \leq (\beta_j, \phi_j)$ for j = 1, 2 and i = j - 1, j. Since (α_1, θ_1) bounds both (β_1, ϕ_1) and (β_2, ϕ_2) from below, Corollary 4.5 allows us to take their join, i.e., their least upper bound, which we call (γ, ψ) . The transitivity of the order gives $(\alpha_i, \theta_i) \leq (\gamma, \psi)$ for i = 0, 2, which verifies the transitivity of the relation (2), as desired.

(2) \implies (3): If (β, ϕ) is a common upper bound of (α, θ) and (α', θ') , then the upward unbounded interval $[(\beta, \phi), \infty[$ is a common upper-closed subset of $[(\alpha, \theta), \infty[$ and $[(\alpha', \theta'), \infty[$. Since $[(\beta, \phi), \infty[$ is inhabited, $[(\beta, \phi), \infty[$ shares any largest element of $[(\alpha, \theta), \infty[$ or $[(\alpha', \theta'), \infty[$; this immediately implies (3).

(3)
$$\implies$$
 (1): We see $(\alpha, \theta) \sim (\tilde{\alpha}, \tilde{\theta}) = (\tilde{\alpha}', \tilde{\theta}') \sim (\alpha', \theta').$

As a direct consequence of the previous proposition, we get:

Corollary 4.7. If $([m], X), ([n], Y) \in Ob \Gamma$ are objects, every equivalence class in $\operatorname{Hom}_{\Gamma}(([m], X), ([n], Y))$ possesses a maximum with respect to the equipped order.

5. Reedy structure, directness and finiteness

In this section, we will construct a Reedy structure of $\int \mathbf{N}^{-,+}(C)$, and prove the directness of $\int \mathbf{N}^{-,+}_+(C)$ and Down(C).

Set $\Gamma \coloneqq \int \mathbf{N}^{-,+}(C)$ until Proposition 5.3, where we would like to establish a Reedy structure of Γ .

Definition 5.1 $(\int \mathbf{N}_{-}^{-,{\mathrm{id}}}(C), \int \mathbf{N}_{+}^{-,+}(C))$. Let wide subcategories

$$\Gamma_{-} = \int \mathbf{N}_{-}^{-,\{\mathrm{id}\}}(C), \text{ and}$$

$$\Gamma_{+} = \int \mathbf{N}_{+}^{-,+}(C)$$

of $\Gamma = \int \mathbf{N}^{-,+}(C)$ be defined as follows:

$$\operatorname{Mor} \Gamma_{-} := \left\{ (\sigma, \operatorname{id}_{X \circ \sigma}) \colon ([m], X \circ \sigma) \to ([n], X) \middle| \begin{array}{c} ([n], X) \in \operatorname{Ob} \Gamma, \\ \sigma \colon [m] \to [n] \text{ in } \mathbf{\Delta}_{-} \end{array} \right\}, \text{ and} \\ \operatorname{Mor} \Gamma_{+} := \left\{ (\delta, \theta) \colon ([m], X) \to ([n], Y) \text{ in } \Gamma \mid \delta \colon [m] \to [n] \text{ in } \mathbf{\Delta}_{+} \right\}.$$

Lemma 5.2. $\Gamma = \int \mathbf{N}^{-,+}(C)$ admits unique (Γ_{-},Γ_{+}) -factorization.

Proof. A (Γ_{-}, Γ_{+}) -factorization of (α, θ) is a commutative diagram of the form (5.1) having a morphism from Γ_{-} and then one from Γ_{+} in the bottom row:

We break down the condition that the diagram (5.1) is a (Γ_{-}, Γ_{+}) -factorization. The membership conditions $([l], X') \in \text{Ob} \Gamma$, $(\sigma, \text{id}_{X' \circ \sigma}) \in \text{Mor} \Gamma_{-}$, and $(\delta, \theta') \in \text{Mor} \Gamma_{+}$ are equivalent to the following set of conditions:

$$[l] \in \mathrm{Ob}\,\boldsymbol{\Delta},\tag{5.2}$$

$$\sigma \in \operatorname{Hom}_{\Delta_{-}}([m], [l]), \tag{5.3}$$

$$\delta \in \operatorname{Hom}_{\Delta_{+}}([l], [n]), \tag{5.4}$$

$$X': [l] \to C_{-}$$
 a functor, (5.5)

$$\theta' \colon X' \Rightarrow Y \circ \delta \colon [l] \to C \text{ a natural transformation},$$
 (5.6)

$$\theta'_i \in \operatorname{Hom}_{C_+}(X'(i), Y(\delta(i))), \, \forall i \in [l].$$
(5.7)

For the commutativity of the diagram (5.1), it is necessary and sufficient to have all of (5.8)–(5.10):

$$\alpha = \delta \circ \sigma \quad : [m] \to [n], \tag{5.8}$$

$$X = X' \circ \sigma \colon [m] \to C_{-},\tag{5.9}$$

$$\theta = \theta' \triangleright \sigma \ : X = X' \circ \sigma \Rightarrow Y \circ \alpha = Y \circ \delta \circ \sigma. \tag{5.10}$$

Since $(\Delta, \Delta_{-}, \Delta_{+})$ is a Reedy category, there exists a unique triple $([l], \sigma, \delta)$ satisfying (5.2)–(5.4) and (5.8). We fix $([l], \sigma, \delta)$ as such.

Now it suffices to establish the unique existence of a pair (X', θ') enjoying the conditions (5.5)–(5.7), (5.9) and (5.10). Remembering that every morphism in Δ_{-} is a split epimorphism in Δ and that the **Poset**-enriched category Δ is in fact enriched over finite bounded lattices, take the largest section $\epsilon : [l] \to [m]$ of σ . By precomposing ϵ on the both sides of eqs. (5.9) and (5.10), we get:

$$X \circ \epsilon = X' \colon [l] \to C_{-},\tag{5.11}$$

$$\theta \triangleright \epsilon = \theta' : X' \Rightarrow Y \circ \delta. \tag{5.12}$$

From these equations, the uniqueness of (X', θ') is immediate.

It just remains to show that the pair (X', θ') defined by eqs. (5.11) and (5.12) satisfies the conditions (5.5)–(5.7), (5.9) and (5.10), of which (5.5)–(5.7) are obvious. We wish to show eqs. (5.9) and (5.10). The maximality of ϵ yields the inequality

 $\operatorname{id}_{[m]} \leq \epsilon \circ \sigma \colon [m] \to [m]$; we have a natural transformation $! = !_{\operatorname{id},\epsilon \circ \sigma} \colon \operatorname{id}_{[m]} \Rightarrow \epsilon \circ \sigma$. Notice:

$$\alpha \triangleleft \mathbf{!}_{\mathrm{id},\epsilon\circ\sigma} = \mathbf{!} = \mathrm{id} \colon \alpha \Rightarrow \alpha \circ \epsilon \circ \sigma = \delta \circ \sigma \circ \epsilon \circ \sigma = \alpha.$$

$$(5.13)$$

The interchange law of $\theta: X \Rightarrow Y \circ \alpha$ and $!: id_{[m]} \Rightarrow \epsilon \circ \sigma$ may be written, combining eqs. (5.11)–(5.13), as the following commutative square of natural transformations:

$$\begin{array}{ccc} X & \xrightarrow{\theta} & Y \circ \alpha \\ X_{\triangleleft} \downarrow & & & \\ X' \circ \sigma & \xrightarrow{\theta' \triangleright \sigma} & Y \circ \alpha \end{array} \tag{5.14}$$

In the diagram (5.14), every constituent morphism of $X \triangleleft !: X \Rightarrow X' \circ \sigma$ belongs to C_- , and C_+ has all the component morphisms of natural transformations drawn horizontally in the diagram. Therefore the uniqueness of the (C_-, C_+) -factorization applies and we get eqs. (5.9) and (5.10), which concludes the proof.

Proposition 5.3. The pair of wide subcategories (Γ_{-}, Γ_{+}) in Definition 5.1 is a Reedy structure on $\Gamma = \int \mathbf{N}^{-,+}(C)$.

Proof. In light of Lemma 5.2, it just remains to show the conditions (2) and (3) from Definition 2.8. We first verify that the property of morphisms being an identity is decidable. For a morphism $(\alpha, \theta) : ([m], X) \to ([n], Y)$ to be identity, it is necessary and sufficient to have identities as the components $\alpha, \theta_0, \theta_1, \ldots, \theta_m$. Since we have dichotomy of each of these morphisms into identities and non-identies, we obtain the desired decidablity.

Now it suffices to establish the well-foundedness of $Ob \Gamma$. Since C is a Reedy category, C_+ is a direct category. That is, the relation $<_+$ defined by the following is a well-founded relation on $Ob C = Ob C_+$:

for
$$x, y \in \operatorname{Ob} C$$
, $x <_+ y :\iff \exists f \colon x \to y \text{ in } C_+, f \neq \operatorname{id}_x$.

Consider the following set:

$$(\operatorname{Ob} C)^{<\omega} = \prod_{n \in \mathbb{N}} (\operatorname{Ob} C)^n,$$

which consists of finitely long sequences $(x_0, x_1, \ldots, x_{n-1})$ of objects of C. For each pair of elements $\boldsymbol{x} = (x_0, x_1, \ldots, x_{n-1}), \boldsymbol{y} = (y_0, y_1, \ldots, y_{m-1}) \in (Ob C)^{<\omega}$, let us say $\boldsymbol{x} < \boldsymbol{y}$ if and only if we have either m < n or all of the followings hold:

- m = n,
- for some $0 \le i < m, x_i <_+ y_i$, and
- for all $0 \le i < m$, $x_i = y_i$ or $x_i <_+ y_i$.

The well-foundedness of this relation $\langle \text{ on } (\operatorname{Ob} C)^{\langle \omega \rangle}$ follows from those of \mathbb{N} and $\operatorname{Ob} C_+$. Now, consider the following map of sets:

$$F\colon \operatorname{Ob} \Gamma \to (\operatorname{Ob} C)^{<\omega}; \ ([n], \theta) \mapsto (\theta_0, \theta_1, \dots, \theta_n).$$

Now we can readily confirm that any non-identity $\boldsymbol{\alpha} \to \boldsymbol{\beta}$ in Γ_+ satisfies $F(\boldsymbol{\alpha}) < F(\boldsymbol{\beta})$, and that any non-identity $\boldsymbol{\alpha} \to \boldsymbol{\beta}$ in Γ_- satisfies $F(\boldsymbol{\alpha}) > F(\boldsymbol{\beta})$. This implies the well-foundedness $Ob \Gamma$ with respect to the relation specified in (3) from Definition 2.8.

Remark 5.4. Let $\Gamma = \int \mathbf{N}^{-,\text{all}}(C)$ be the full subcategory of $\int \mathbf{N}(C)$ spanned by those objects ([n], X) for which the constituent functor $X : [n] \to C$ factors through C_- . This category Γ admits a Reedy structure (Γ_-, Γ_+) , and the Reedy structure on $\int \mathbf{N}^{-,+}(C)$ in Proposition 5.3 is the restriction of this Reedy structure. Here, letting \pm denote either - or +, a morphism

$$(\alpha, \theta) \colon ([m], X) \to ([n], Y)$$

belongs to $\Gamma_{\pm} = \int \mathbf{N}_{\pm}^{-,\pm}(C)$ if and only if $\alpha \in \mathbf{\Delta}_{\pm}$ and $\theta_i \in C_{\pm}$ for every $i \in [m]$. The proof of this statement is similar to the proof in Lemma 5.2, but it calls for a slightly more careful construction of the Reedy factorization.

From the Reedy structure of $\int \mathbf{N}^{-,+}(C)$, the directness of $\int \mathbf{N}^{-,+}_+(C)$ may be deduced.

Corollary 5.5. The category $\int \mathbf{N}_{+}^{-,+}(C)$ is direct.

Proof. By Proposition 5.3, the cagegory $\int \mathbf{N}_{+}^{-,+}(C)$ is the direct part of the Reedy structure of $\int \mathbf{N}^{-,+}(C)$, and is therefore direct; the category $\int \mathbf{N}_{+}^{-,+}(C)$ in question is a (full) subcategory of $\int \mathbf{N}_{+}^{-,+}(C)$.

We now show the directness of Down(C).

Lemma 5.6. Let $([n], X) \in Ob \int \mathbf{N}^{-,+}(C)$. Then the following conditions are equivalent:

- (1) $([n], X) \in \text{Ob} \int \mathbf{N}_{+}^{--,+}(C).$
- (2) The element $\operatorname{id}_{([n],X)} \in \operatorname{Hom}_{\int \mathbf{N}^{-,+}(C)}(([n],X),([n],X))$ is maximal, i.e., for any $\boldsymbol{\alpha} \geq \operatorname{id}_{([n],X)}$: $([n],X) \to ([n],X)$, we have $\boldsymbol{\alpha} = \operatorname{id}_{([n],X)}$.
- (3) The element $\operatorname{id}_{([n],X)} \in \operatorname{Hom}_{\int \mathbf{N}^{-,+}(C)}(([n],X),([n],X))$ is minimal, i.e., for any $\boldsymbol{\alpha} \leq \operatorname{id}_{([n],X)}$: $([n],X) \to ([n],X)$, we have $\boldsymbol{\alpha} = \operatorname{id}_{([n],X)}$.
- (4) The element $\operatorname{id}_{([n],X)} \in \operatorname{Hom}_{\int \mathbf{N}^{-,+}(C)}(([n],X),([n],X))$ is the largest in its own equivalence class.
- (5) The equivalence class of $id_{([n],X)} \in Hom_{\int \mathbf{N}^{-,+}(C)}(([n],X),([n],X))$ is a singleton.

Proof. By the definition of the equivalence of parallel morphisms, (5) is equivalent to the conjuction of (2) and (3). The equivalence of (2) and (4) follows from Proposition 4.6. We show (1) \implies (2) \implies (3) \implies (1) part by part.

(1) \implies (2): Let $([n], X) \in \operatorname{Ob} \int \mathbf{N}_{+}^{-,+}(C)$, and let:

$$\boldsymbol{\alpha} = (\alpha, \theta) \ge \mathrm{id}_{([n], X)} \colon ([n], X) \to ([n], X).$$

Let $i \in [n]$. We have the following commutative diagram:

$$\begin{array}{c} X(i) = & X(i) \\ & \swarrow \\ \theta_i \\ & \downarrow \\ X(\alpha(i)) \end{array}$$

By the uniqueness of (C_-, C_+) -factorization, we get $X(i) = X(\alpha(i)), \ \theta_i = \mathrm{id}_{X(i)},$ and $X(!_{\alpha(i)i}) = \mathrm{id}_{X(i)} \colon X(\alpha(i)) \to X(i)$. From the last equation of the three and the assumption (1), we obtain $\alpha(i) = i$. Wrapping up, we get $\alpha = \mathrm{id}_{[n]}, \ \theta = \mathrm{id}_X,$ and hence $\boldsymbol{\alpha} = \mathrm{id}_{([n],X)}$, as required.

(2) \implies (3): Assume (2), and let $\boldsymbol{\alpha} = (\alpha, \theta) \leq \operatorname{id}_{([n],X)} : ([n], X) \rightarrow ([n], X)$. Define a morphism $\beta : [n] \rightarrow [n]$ in $\boldsymbol{\Delta}$ by $\beta(i) \coloneqq \max \alpha^{-1} \{\alpha(i)\}$ for $i \in [n]$. By definition, we see that $\beta \geq \operatorname{id}_{[n]}$. From this and Lemma 4.2 follows that β corresponds to a unique morphism $\boldsymbol{\beta} = (\beta, \phi) : ([n], X) \rightarrow ([n], X)$ in $\int \mathbf{N}^{-,+}(C)$ satisfying $\boldsymbol{\beta} \geq \operatorname{id}_{([n],X)}$, which, by our assumption (2), implies $\boldsymbol{\beta} = \operatorname{id}_{([n],X)}$. Therefore α is injective, and hence $\alpha = \operatorname{id}_{[n]}$. Combining this with $\boldsymbol{\alpha} \leq \operatorname{id}_{([n],X)}$, we obtain the desired equation $\boldsymbol{\alpha} = \operatorname{id}_{([n],X)}$.

(3) \implies (1): Suppose (3). Assume that $0 \le k \le l \le n$ satisfy:

$$X(!_{kl}) = \mathrm{id}_{X(k)} \colon X(k) \to X(k) = X(l).$$

It suffices to show k = l. By the inverseness of C_{-} , for each pair $k \leq i \leq j \leq l$, it holds that:

$$X(!_{ij}) = \mathrm{id}_{X(k)} \colon X(i) = X(k) \to X(k) = X(j).$$
(5.15)

Define $\alpha \colon [n] \to [n]$ in $\boldsymbol{\Delta}$ by

$$\alpha(i) \coloneqq \begin{cases} k & \text{if } k \le i \le l, \\ i & \text{if } 0 \le i < k \text{ or } l < i \le n \end{cases}$$

By eq. (5.15), we see $X = X \circ \alpha$; therefore $(\alpha, \mathrm{id}_X) : ([n], X) \to ([n], X)$ is a valid morphism in $\int \mathbf{N}^{-,+}(C)$. By eq. (5.15) again, we see that $(\alpha, \mathrm{id}_X) \leq \mathrm{id}_{([n],X)}$, and hence that $(\alpha, \mathrm{id}_X) = \mathrm{id}_{([n],X)}$ by the assumption (3). From this we obtain $l = \mathrm{id}_{[n]}(l) = \alpha(l) = k$, which concludes the proof.

Proposition 5.7. The category Down(C) is direct.

Proof. By Lemma 5.6, the equivalence class of any idenitity in $\int \mathbf{N}_{+}^{-,+}(C)$ is a singleton. Therefore the result immediately follows from Corollary 5.5.

We conclude this section by showing the finiteness of Down(C).

Lemma 5.8. If C is finite, then $\int \mathbf{N}_{+}^{-,+}(C)$ is finite.

Proof. Since C is finite, Mor C has decidable equality. Therefore, for any pair of objects $([m], X), ([n], Y) \in Ob \int \mathbf{N}(C)$, the hom-set $\operatorname{Hom}_{\int \mathbf{N}(C)}(([m], X), ([n], Y))$ is a decidable subset of the finite product $\operatorname{Hom}_{\Delta}([m], [n]) \times (\operatorname{Mor} C)^{m+1}$ of finite sets, and hence is finite. Assume that $([m], X), ([n], Y) \in Ob \int \mathbf{N}_{+}^{-,+}(C)$. The decidability of the subset $\operatorname{Mor} C_{+} \subseteq \operatorname{Mor} C$, which we have assumed as part of the definition of Reedy structures, implies the decidability of the following subset, and hence its finiteness:

 $\operatorname{Hom}_{\int \mathbf{N}_{-}^{-,+}(C)}(([m], X), ([n], Y)) \subseteq \operatorname{Hom}_{\int \mathbf{N}(C)}(([m], X), ([n], Y)).$

Thus it suffices to show that the set of objects of $\int \mathbf{N}_{+}^{--,+}(C)$ is finite. If the number of objects of C is N, then any object ([n], X) of $\int \mathbf{N}_{+}^{--,+}(C)$ satisfies n < N. Since the set of functors $[n] \to C_{-}$, which X is a member of, is finite, the object set in question may be seen as the disjoint union of a finite family of finite sets, and is therefore finite.

Corollary 5.9. If C is finite, then Down(C) is finite.

Proof. The category Down(C) is obtained by taking quotients of the hom-sets of $\int \mathbf{N}_{+}^{-,+}(C)$, which is finite by Lemma 5.8. Therefore it suffices to check the equivalence relations on the hom-sets producing the quotient is decidable. For this, see Proposition 4.6 and remember that the category C has decidable equality on morphisms.

6. Functors between the constructed categories

In this section, we examine canonical functors between the categories defined in Section 3. Here we list such functors:

Remark 6.1. By the construction of the five categories constructed in Section 3, we have the following commutative diagram of functors:

The two horizontal arrows in the upper row are inclusions of subcategories. The vertical arrows are the canonical projections: to be pedantic on categorical formalism, it is induced by the canonical natural transformation

forget
$$\Rightarrow \pi_0 \colon \mathbf{Poset} \to \mathbf{Set}$$
.

The lower horizontal arrow $\text{Down}(C) \hookrightarrow \text{Down}_*(C)$ is induced by the inclusion $\int \mathbf{N}_+^{-,+}(C) \hookrightarrow \int \mathbf{N}^{-,+}(C)$, which, by Definition 3.6 and Lemma 3.5, is a **Poset**-fully faithful **Poset**-enriched functor, meaning that the functor is order isomorphisms in hom-posets. Therefore, the $\text{Down}(C) \hookrightarrow \text{Down}_*(C)$ may be considered as the inclusion of a full subcategory.

The main contents of this section are the following two results:

- The inclusion $\text{Down}(C) \hookrightarrow \text{Down}_*(C)$ of a full subcategory is a categorical equivalence (Lemma 6.2).
- The canonical surjective functor $\int \mathbf{N}^{-,+}(C) \twoheadrightarrow \text{Down}_*(C)$ is a strict 1-localization functor (Proposition 6.3).

Lemma 6.2. The inclusion $\text{Down}(C) \hookrightarrow \text{Down}_*(C)$ is a categorical equivalence.

Proof. Since the inclusion is fully faithful, it suffices to explicitly construct, for each $([m], X) \in Ob(Down_*(C))$, a pair of an object $([n], Y) \in Ob(Down(C))$ and an isomorphism $([n], Y) \xrightarrow{\sim} ([m], X)$ in $Down_*(C)$. Let $([m], X) \in Ob(Down_*(C))$ and consider its identity morphism in $\int \mathbf{N}^{-,+}(C)$:

$$id_{([m],X)} = (id_{[m]}, id_X) \colon ([m], X) \to ([m], X).$$

Let $\alpha: ([m], X) \to ([m], X)$ be the largest element of the equivalence class of $\mathrm{id}_{([m],X)}$, which is shown to exist in Corollary 4.7. Since $\mathrm{id}_{([m],X)} \leq \alpha$, we have $\alpha \leq \alpha \circ \alpha$. The maximality of α and $\alpha \sim \alpha \circ \alpha$ together give $\alpha \geq \alpha \circ \alpha$; we therefore get $\alpha = \alpha \circ \alpha$. By Lemma 2.11 and Proposition 5.3, the idempotent α has a unique splitting of the following form:



Since $\alpha \sim id$, we have constructed an isomorphism $([m], X) \cong ([n], Y)$ in $\text{Down}_*(C)$. It only remains to show that

$$([n], Y) \in \operatorname{Ob}(\operatorname{Down}(C)) = \operatorname{Ob} \int \mathbf{N}_{+}^{--,+}(C).$$

Let $\boldsymbol{\alpha}': ([n], Y) \to ([n], Y)$ be the largest element of the equivalence class of the identity $\mathrm{id}_{([n],Y)} \in \mathrm{Hom}_{\int \mathbf{N}^{-,+}(C)}(([n],Y), ([n],Y))$. By Lemma 5.6, it suffices to prove $\boldsymbol{\alpha}' = \mathrm{id}_{([n],Y)}$. By repeating the argument above, we may take $\boldsymbol{\sigma}'$ and $\boldsymbol{\delta}'$ with the following commutative diagram:



Since $\operatorname{id}_{([n],Y)} \leq \alpha'$, we have:

$$\boldsymbol{\alpha} = \boldsymbol{\delta} \circ \operatorname{id}_{([n],Y)} \circ \boldsymbol{\sigma} \leq \boldsymbol{\delta} \circ \boldsymbol{\alpha'} \circ \boldsymbol{\sigma} = \boldsymbol{\delta} \circ \boldsymbol{\delta'} \circ \boldsymbol{\sigma'} \circ \boldsymbol{\sigma}.$$

From this and the maximality of α , we can derive:

$$\delta \circ \sigma = \alpha = \delta \circ \delta' \circ \sigma' \circ \sigma.$$

Proposition 5.3 implies ([l], Z) = ([n], Y), $\boldsymbol{\delta} = \boldsymbol{\delta} \circ \boldsymbol{\delta}'$ and $\boldsymbol{\sigma} = \boldsymbol{\sigma}' \circ \boldsymbol{\sigma}$, which require $\boldsymbol{\delta}' = \boldsymbol{\sigma}' = \mathrm{id}_{([n],Y)}$. Hence $\boldsymbol{\alpha}' = \mathrm{id}_{([n],Y)}$, as desired.

Next, we consider the quotient functor $\int \mathbf{N}^{-,+}(C) \to \operatorname{Down}_*(C)$:

30 FINITE REEDY CATEGORIES AS LOCALIZATIONS OF FINITE DIRECT CATEGORIES

Proposition 6.3. The quotient functor $q: \int \mathbf{N}^{-,+}(C) \to \text{Down}_*(C)$ exhibits the category $\text{Down}_*(C)$ as the strict 1-localization of $\int \mathbf{N}^{-,+}(C)$ at $\int \mathbf{N}^{-,\{\text{id}\}}(C)$ in Definition 5.1.

Proof. We first show that the functor q sends every morphism in $\int \mathbf{N}_{-}^{-,\{\mathrm{id}\}}(C)$ to an isomorphism in $\mathrm{Down}_{*}(C)$. Let $(\sigma, \mathrm{id}_{X \circ \sigma}) \colon ([m], X \circ \sigma) \to ([n], X)$ be a morphism in $\int \mathbf{N}_{-}^{-,\{\mathrm{id}\}}(C)$. Using $\sigma \in \mathrm{Mor} \Delta_{-}$, which follows from the definition of the wide subcategory $\int \mathbf{N}_{-}^{-,\{\mathrm{id}\}}(C)$, take the largest section $\delta \colon [n] \to [m]$ of $\sigma \colon [m] \to [n]$. We easily calculate to see that $(\delta, \mathrm{id}_X) \colon ([n], X) = ([n], X \circ \sigma \circ \delta) \to ([m], X \circ \sigma)$ is a legitimate morphism in $\int \mathbf{N}^{-,+}(C)$ and is a section of $(\sigma, \mathrm{id}_{X \circ \sigma})$. In addition, we have:

$$\mathrm{id}_{([m],X\circ\sigma)} = (\mathrm{id}_{[m]},\mathrm{id}_{X\circ\sigma}) \leq (\delta\circ\sigma,\mathrm{id}_{X\circ\sigma}) = (\delta,\mathrm{id}_X)\circ(\sigma,\mathrm{id}_{X\circ\sigma}).$$

Thus, $q(\sigma, \mathrm{id}_{X \circ \sigma})$ and $q(\delta, \mathrm{id}_X)$ are mutual inverses in $\mathrm{Down}_*(C)$, as desired.

Next, we address the conditional unique factorization through q. Assume that a functor $F: \int \mathbf{N}_{-}^{-,\{\mathrm{id}\}}(C) \to D$ sends every morphism in $\int \mathbf{N}_{-}^{-,\{\mathrm{id}\}}(C)$ to an isomorphism in D. Since q is a quotient functor, it suffices to prove that F respects the equivalence relations on hom-sets: explicitly, $F(\alpha, \theta) = F(\alpha', \theta')$ for any parallel pair of morphisms $(\alpha, \theta) \ge (\alpha', \theta'): ([m], X) \to ([n], Y)$. Take any such parallel pair. To facilitate the proof, we construct a sequence of morphisms:

$$(\alpha, \theta) = (\alpha^0, \theta^0) \ge (\alpha^1, \theta^1) \ge \dots \ge (\alpha^m, \theta^m) \ge (\alpha^{m+1}, \theta^{m+1}) = (\alpha', \theta')$$
$$: ([m], X) \to ([n], Y).$$

Specifically, for each $0 \le k \le m+1$, the k-th entry of the sequence is given by:

$$\alpha^{k}(i) \coloneqq \begin{cases} \alpha'(i) & \text{if } 0 \leq i < k, \text{ and} \\ \alpha(i) & \text{if } k \leq i \leq m; \end{cases}$$
$$\left(\theta^{k}\right)_{i} \coloneqq \begin{cases} \theta'_{i} & \text{if } 0 \leq i < k, \text{ and} \\ \theta_{i} & \text{if } k \leq i \leq m. \end{cases}$$

Now, we need only demonstrate $F(\alpha^k, \theta^k) = F(\alpha^{k+1}, \theta^{k+1})$ for every $0 \le k \le m$. Let the morphisms $\sigma_k^m \colon [m+1] \to [m]$ and $\delta_k^{m+1}, \delta_{k+1}^{m+1} \colon [m] \to [m+1]$ in Δ be those from the usual convention. By assumption, the functor F sends the morphism

$$(\sigma_k^m, \mathrm{id}_{X \circ \sigma_k^m}) \colon ([m+1], X \circ \sigma_k^m) \to ([m], X)$$

to an isomorphism. Since the two morphisms

$$(\delta_k^{m+1}, \mathrm{id}_X), (\delta_{k+1}^{m+1}, \mathrm{id}_X) \colon ([m], X) \to ([m+1], X \circ \sigma_k^m)$$

are both sections of $(\sigma_k^m, \operatorname{id}_{X \circ \sigma_k^m})$, their images by F coincide. Define the morphism $(\tilde{\alpha}^k, \tilde{\theta}^k) \colon ([m+1], X \circ \sigma_k^m) \to ([n], Y)$ by:

$$\begin{split} \tilde{\alpha}^k(i) &\coloneqq \begin{cases} \alpha'(i) & \text{if } 0 \leq i \leq k, \text{ and} \\ \alpha(i-1) & \text{if } k < i \leq m+1; \end{cases} \\ (\tilde{\theta}^k)_i &\coloneqq \begin{cases} \theta'_i & \text{if } 0 \leq i \leq k, \text{ and} \\ \theta_{i-1} & \text{if } k < i \leq m+1. \end{cases} \end{split}$$

We see $(\tilde{\alpha}^k, \tilde{\theta}^k) \circ (\delta_l^{m+1}, \operatorname{id}_X) = (\alpha^l, \theta^l)$ for l = k, k+1; therefore we get the following, as required:

$$F(\alpha^{k}, \theta^{k})$$

$$= F(\tilde{\alpha}^{k}, \tilde{\theta}^{k}) \circ F(\delta_{k}^{m+1}, \mathrm{id}_{X})$$

$$= F(\tilde{\alpha}^{k}, \tilde{\theta}^{k}) \circ F(\delta_{k+1}^{m+1}, \mathrm{id}_{X})$$

$$= F(\alpha^{k+1}, \theta^{k+1}).$$

 \square

Now, we add an obvious corollary to the above proposition: we can consider both Down(C) and $\text{Down}_*(C)$ as the localization of $\int \mathbf{N}^{-,+}(C)$ at $\int \mathbf{N}^{-,\{\text{id}\}}(C)$:

Corollary 6.4. Write $q: \int \mathbf{N}^{-,+}(C) \to \text{Down}_*(C)$ for the canonical projection, and let $r: \text{Down}_*(C) \to \text{Down}(C)$ stand for the quasi-inverse of the inclusion $\text{Down}(C) \hookrightarrow \text{Down}_*(C)$, constructed in Lemma 6.2. Then the pairs $(\text{Down}_*(C), q)$ and $(\text{Down}(C), r \circ q)$ are both weak 1-localizations of $\int \mathbf{N}^{-,+}(C)$ at $\int \mathbf{N}^{-,\{\text{id}\}}(C)$.

Proof. Follows from Proposition 6.3 and Remark 2.3.

7. The proof of 1-localization

Now in this section, we compare Down(C) and C. We first construct the functor that compares the two categories:

Definition 7.1 (The last component functor). The following describes three functors $\int \mathbf{N}(C) \to C$, $\int \mathbf{N}^{-,+}(C) \to C$, and $\int \mathbf{N}^{-,+}_{+}(C) \to C$, which will all be denoted commonly by last:

$$\begin{split} \mathfrak{last}([n],X) &\coloneqq X(n) & \text{for } ([n],X) \text{ an object of the domain;} \\ \mathfrak{last}(\alpha,\theta) &\coloneqq Y(!_{\alpha(m),n}) \circ \theta_m & \text{for } (\alpha,\theta) \colon ([m],X) \to ([n],Y). \end{split}$$

These functors respect the equivalence relations on the hom-sets of the domain categories, so that they induce the functors $\text{Down}(C) \to C$ and $\text{Down}_*(C) \to C$. These two will also be denoted by \mathfrak{last} , abusing the notation. When we need to distinguish the five functors above, we write $\mathfrak{last}_{\Gamma} \colon \Gamma \to C$.

Our purpose is to prove that last presents C as a localization of Down(C). Therefore we will also need the class of weak equivalences:

Definition 7.2 (last-weak equivalence). Let Γ denote one of the five categories $\int \mathbf{N}(C)$, $\int \mathbf{N}^{-,+}(C)$, $\int \mathbf{N}^{-,+}_+(C)$, $\operatorname{Down}_*(C)$, and $\operatorname{Down}(C)$. A morphism in Γ is said to be a last-weak equivalence (in Γ) if last: $\Gamma \to C$ maps it to some identity in C. We will write:

$$W^{\mathfrak{last}} = W_{\Gamma}^{\mathfrak{last}} \coloneqq \{ \boldsymbol{\alpha} \in \operatorname{Mor} \Gamma \mid \mathfrak{last}(\boldsymbol{\alpha}) = \operatorname{id} \} \subseteq \operatorname{Mor} \Gamma$$

for the collection of *last*-weak equivalences.

The main theorem of this section is the following one:

Theorem 7.3. Let Γ be one of the four categories $\int \mathbf{N}(C)$, $\int \mathbf{N}^{-,+}(C)$, $\operatorname{Down}(C)$, and $\operatorname{Down}_*(C)$. The category C and the functor $\operatorname{last}_{\Gamma}$ in Definition 7.1 constitute a weak 1-localization at last -weak equivalences.

Notice the *four* in the statement of the theorem; we have one counterexample:

Remark 7.4. The functor $\text{last}: \int \mathbf{N}_{+}^{--,+}(C) \to C$ is not in general a 1-localization. For a simple counterexample, define C to be the poset $\{0 \leq 2 \leq 1\}$ equipped with the Reedy structure (C_{-}, C_{+}) determined by $g \coloneqq \mathbf{!}_{21} \in C_{-}, f \coloneqq \mathbf{!}_{02} \in C_{+}$, and $g \circ f = \mathbf{!}_{01} \in C_{+}$. Let C' be the category obtained by freely adjoining $h: 0 \to 1$ to C. In diagram:



Let incl: $C \hookrightarrow C'$ be the inclusion. Let us define a functor $F: \int \mathbf{N}_{+}^{-,+}(C) \to C'$. On objects, we set:

$$F([n], X) \coloneqq X(n) = \operatorname{incl}(\mathfrak{last}([n], X)).$$

Let (α, θ) : $([m], X) \to ([n], Y)$ be a morphism in $\int \mathbf{N}_{+}^{-,+}(C)$. Then we set:

$$F(\alpha, \theta) \coloneqq \begin{cases} h & \text{if } m = 0, \ X(0) = 0, \ Y(\alpha(0)) = 1, \text{ and } \theta_0 = g \circ f; \\ \text{incl}(\mathfrak{last}(\alpha, \theta)) & \text{otherwise.} \end{cases}$$

Then F is a well-defined functor and sends all the last-weak equivalences to isomorphisms. However it does not factor through last even up to natural isomorphism; indeed, F maps the parallel pair of morphisms

$$(\delta_1^1, f), (\delta_0^1, g \circ f) \colon ([0], 0) \to ([1], 2 \xrightarrow{g} 1)$$

to an unequal parallel pair $g \circ f$ and h, while \mathfrak{last} sends them to an equal morphism $g \circ f$. This establishes a counterexample: $\mathfrak{last}: \int \mathbf{N}_{+}^{-,+}(C) \to C$ is not a 1-localization.

The rest of this section is devoted to the proof of Theorem 7.3. The demonstration is broken down to several lemmas, which will be wrapped up in page 37. We begin with the most trivial two of these auxiliary propositions:

Lemma 7.5. The five functors denoted by last in Definition 7.1 send last-weak equivalences to isomorphisms. In fact, last-weak equivalences are exactly the morphisms mapped to isomorphisms by last.

Proof. The first statement follows from the definition of last-weak equivalences. The second follows from the assumption that C is a Reedy category.

Lemma 7.6. The functors last and the families W^{last} are compatible with the following strictly commutative diagram of functors. Explicitly, for any arrow $F: \Gamma \to \Gamma'$ depicted in the diagram, we have $\text{last}_{\Gamma'} \circ F = \text{last}_{\Gamma}$ and $F^{-1}(W^{\text{last}}_{\Gamma'}) = W^{\text{last}}_{\Gamma}$.

$$\int \mathbf{N}_{-}^{-,+}(C) \longleftrightarrow \int \mathbf{N}^{-,+}(C) \longleftrightarrow \int \mathbf{N}(C)$$

$$\begin{array}{c} \text{quotient} \\ \text{quotient} \\ \text{Down}(C) \longleftrightarrow \text{Down}_{*}(C) \end{array} \tag{7.1}$$

Proof. By definition.

To facilitate a clearer presentation, we commence with the relatively simple case of $\Gamma = \int \mathbf{N}(C)$ in Theorem 7.3:

Lemma 7.7. The functor last: $\int \mathbf{N}(C) \to C$ exhibits C as a weak 1-localization of $\int \mathbf{N}(C)$ at last-weak equivalences.

Proof. The functor in question admits the following right adjoint:

$$i: C \to \int \mathbf{N}(C);$$

$$Ob C = Ob C^{[0]} \ni x \mapsto ([0], x);$$

$$(f: x \to y) \mapsto (\mathrm{id}_{[0]}, f).$$

Since this right adjoint is fully faithful, we may consider C as a reflective subcategory of $\int \mathbf{N}(C)$, and last is its reflector. Now the claim follows from Lemma 7.5. \Box

The other cases are a little more intricate; therefore we prove them part by part. The next lemma shows the weak uniqueness property of the factorization through \mathfrak{last}_{Γ} , which is a part of the definition of weak 1-localization in Definition 2.2. It holds even for $\Gamma = \int \mathbf{N}_{+}^{-,+}(C)$:

Lemma 7.8. Let $last = last_{\Gamma} \colon \Gamma \to C$ be any of the five functors defined in Definition 7.1. Let $F, G \colon C \to D$ be functors. If:

$$\epsilon \colon F \circ \mathfrak{last} \Rightarrow G \circ \mathfrak{last} \colon \Gamma \to D$$

is a natural transformation, there is a unique natural transformation $\tilde{\epsilon} \colon F \Rightarrow G$ satisfying $\tilde{\epsilon} \triangleright \mathsf{last} = \epsilon$.

Proof. The equation $\tilde{\epsilon} \triangleright \mathfrak{last} = \epsilon$ implies:

$$\tilde{\epsilon}_x = \tilde{\epsilon}_{\mathfrak{last}([0],x)} = \epsilon_{([0],x)} \colon F(x) \to G(x)$$
(7.2)

for any $x \in \operatorname{Ob} C = \operatorname{Ob} C^{[0]}$. This shows uniqueness. It suffices to prove that (7.2) defines a natural transformation $\tilde{\epsilon} \colon F \Rightarrow G$ and to check $\tilde{\epsilon} \triangleright \mathfrak{last} = \epsilon$.

We shall first show:

$$\tilde{\epsilon}_{\mathfrak{last}([n],X)} = \epsilon_{([n],X)} \colon F(\mathfrak{last}([n],X)) \to G(\mathfrak{last}([n],X)) \tag{7.3}$$

for any $([n], X) \in Ob \Gamma$. Once we show the naturality of $\tilde{\epsilon}$, (7.3) should automatically prove $\tilde{\epsilon} \triangleright \mathsf{last} = \epsilon$. Let $([n], X) \in Ob \Gamma$ be arbitrary. Remember our notation; the morphism $\iota_n \colon [0] \to [n]$ in Δ is the one with $\iota_n(0) = n$. We apply the naturality of ϵ to $(\iota_n, \mathrm{id}_{X(n)}) \colon ([0], X(n)) \to ([n], X)$, which belongs to any choice of Γ , to get the following commutative diagram:

Since the vertical arrows reduce to $id_{F(X(n))}$ and $id_{G(X(n))}$, the horizontal arrows must be equal. Hence the desired equality (7.3) follows from the defining equation (7.2).

It just remains to confirm the naturality. Since C is a Reedy category, it suffices to show the commutativity of the naturality diagram:

$$F(x) \xrightarrow{\epsilon_x} G(x)$$

$$F(f) \downarrow \qquad \qquad \downarrow^{G(f)} \qquad (7.4)$$

$$F(y) \xrightarrow{\epsilon_y} G(y)$$

in the cases $f \in \operatorname{Mor} C_+$ and $f \in \operatorname{Mor} C_-$.

If $f \in \text{Mor } C_+$, the diagram (7.4) coincides with the following commutative diagram, which is the naturality of ϵ for $(\text{id}_{[0]}, f): ([0], x) \to ([0], y):$

$$\begin{split} F(\mathfrak{last}([0],x)) & \xrightarrow{\epsilon_{([0],x)}} G(\mathfrak{last}([0],x)) \\ F(\mathfrak{last}(\mathrm{id}_{[0]},f)) & \downarrow G(\mathfrak{last}(\mathrm{id}_{[0]},f)) \\ F(\mathfrak{last}([0],y)) & \xrightarrow{\epsilon_{([0],y)}} G(\mathfrak{last}([0],y)) \end{split}$$

We discuss the remaining case: $f \in \operatorname{Mor} C_-$. Remember the property of f being an identity is decidable. If f is some identity, the desired naturality follows from the case $f \in \operatorname{Mor} C_+$. Otherwise, we use (7.3) to see that the desired commutative diagram (7.4) is exactly the naturality of ϵ for $(\delta_1^1, \operatorname{id}_x): ([0], x) \to ([1], x \xrightarrow{f} y)$, which is:

$$\begin{array}{c} F(\mathfrak{last}([0], x)) \xrightarrow{\epsilon_{([0], x)}} G(\mathfrak{last}([0], x)) \\ F(\mathfrak{last}(\delta_1^1, \mathrm{id}_x)) \downarrow & \qquad \qquad \downarrow G(\mathfrak{last}(\delta_1^1, \mathrm{id}_x)) \\ F(\mathfrak{last}([1], f)) \xrightarrow{\epsilon_{([1], f)}} G(\mathfrak{last}([1], f)) \end{array}$$

This now concludes the proof.

Now we demonstrate the remaining condition of 1-localization: the conditional existence of the factorization through $\operatorname{last}_{\Gamma}$ for $\Gamma = \operatorname{Down}_*(C)$. The cases $\Gamma = \int \mathbf{N}^{-,+}(C)$ and $\Gamma = \operatorname{Down}(C)$ are equivalent to this case, so their treatment is postponed. We split out a small lemma from the proof to make it easier to read:

Lemma 7.9. Let Γ stand for any of $\int \mathbf{N}(C)$, $\int \mathbf{N}^{-,+}(C)$ and $\int \mathbf{N}^{-,+}_+(C)$. Then any morphism (α, θ) : $([l], X) \to ([n], Z)$ decomposes into the composition $(\alpha, \theta) = (\gamma, \psi) \circ (\beta, \phi)$ of a (not necessarily unique) pair of morphisms

$$(\beta, \phi) \colon ([l], X) \to ([m], Y),$$

$$(\gamma, \psi) \colon ([m], Y) \to ([n], Z)$$

satisfying $\beta(l) = m$, $Y(m) = Z(\gamma(m))$, and $\psi_m = id_{Y(m)} = id_{Z(\gamma(m))}$.

Proof. Set $m \coloneqq \alpha(l)$. There is a factorization that fits into the following commutative diagram, given by $\alpha'(i) = \alpha(i)$:

Here, $\iota_{\leq m} = \iota_{\{0,1,\ldots,m\}} \colon [m] \rightarrowtail [n]$ is specified by $\iota_{\leq m}(i) = i$. Now,

$$(\beta, \phi) \coloneqq (\alpha', \theta) \colon ([l], X) \to ([m], Y \circ \iota_{\leq m}),$$

$$\gamma, \psi) \coloneqq (\iota_{\leq m}, \operatorname{id}_{Y \circ \iota_{\leq m}}) \colon ([m], Y \circ \iota_{\leq m}) \to ([n], Y)$$

meets the requirements.

Now we proceed to the actual proof of the desired existence.

Proposition 7.10. Let D be a category. Suppose that a functor $F: \text{Down}_*(C) \to D$ maps every last-weak equivalence to an isomorphism. Then there exists a functor $\tilde{F}: C \to D$ and a natural isomorphism $\theta: \tilde{F} \circ \text{last} \cong F$.

Proof. For the sake of readability of the construction of the functor \tilde{F} , we begin by specifying its two restrictions $\tilde{F}_+: C_+ \to D$ and $\tilde{F}_-: C_- \to D$. For any object $x \in \operatorname{Ob} C = \operatorname{Ob} C^{[0]}$, we set $\tilde{F}(x) = \tilde{F}_+(x) = \tilde{F}_-(x) \coloneqq F([0], x)$. If $d: x \to y$ be

a morphism in C_+ , then we define $\tilde{F}_+(d) \coloneqq F(\mathrm{id}_{[0]}, d)$, through the identification $C = C^{[0]}$. This obviously makes \tilde{F}_+ a functor.

Now consider a morphism $s: x \twoheadrightarrow y$ in C_- . The object $(x \twoheadrightarrow y) \in Ob(C_-)^{[1]}$ constitutes an object $([1], s) \in Ob \operatorname{Down}_*(C)$. We define $\tilde{F}_-(s)$ to be the unique morphism that commutates the following diagram:

We need to show the functoriality of \tilde{F}_- . Since putting x = y, $s = \mathrm{id}_x$, and $\tilde{F}_-(\mathrm{id}_x) = \mathrm{id}_{\tilde{F}_-(x)}$ commutes the diagram (7.5), we see that $\tilde{F}_-(\mathrm{id}_x) = \mathrm{id}_{\tilde{F}_-(x)}$. We prove that \tilde{F}_- preserves composition. Let $X = (x \xrightarrow{s} y \xrightarrow{t} z) \in \mathrm{Ob}(C_-)^{[2]}$ be an arbitrary composable pair of morphisms. We have the following commutative diagram in $\mathrm{Down}_*(C)$ (in fact in $\int \mathbf{N}^{-,+}(C)$):

$$([0], x) \xrightarrow{(\delta_{1}^{1}, \operatorname{id}_{x})} ([1], t \circ s) \xleftarrow{(\delta_{0}^{1}, \operatorname{id}_{z})} ([0], z)$$

$$(\delta_{1}^{1}, \operatorname{id}_{x}) \downarrow (\iota_{0}, \operatorname{id}_{x}) \xrightarrow{((\delta_{1}^{2}, \operatorname{id}_{tos}))} ([2], X) \xleftarrow{(\iota_{2}, \operatorname{id}_{z})} ([1], t)$$

$$([1], s) \xrightarrow{(\delta_{2}^{2}, \operatorname{id}_{s})} ([2], X) \xleftarrow{(\delta_{0}^{2}, \operatorname{id}_{t})} ([1], t) \qquad (7.6)$$

$$(\delta_{0}^{1}, \operatorname{id}_{y}) \uparrow ([0], y) \xrightarrow{(\delta_{1}^{1}, \operatorname{id}_{y})} ([0], y)$$

The functor F sends this to the commutative diagram below, which demonstrates $\tilde{F}_{-}(t) \circ \tilde{F}_{-}(s) = \tilde{F}_{-}(t \circ s)$, as required:

$$\tilde{F}_{-}(x) \xrightarrow{F_{-}(t \circ s)} \tilde{F}_{-}(z) \xrightarrow{F_{-}(t \circ s)}$$

Thus we obtain that \tilde{F}_{-} is a functor.

Now we set $\tilde{F}(d \circ s) \coloneqq \tilde{F}_+(d) \circ \tilde{F}_-(s)$ for any composable pair $x \xrightarrow{s} y \xrightarrow{d} z$ in C. The unique factorization of the Reedy structure of C ensures that this gives a well-defined family of functions $\operatorname{Hom}_C(x, z) \to \operatorname{Hom}_D(x, z)$ for $x, z \in \operatorname{Ob} C$.

We wish to verify the functoriality of \tilde{F} . Consider any commutative square of the following form:

The diagram (7.8) represents a natural transformation $\phi_{d'd} \colon s \Rightarrow s' \colon [1] \to C$, which comprises a morphism $(\mathrm{id}_{[1]}, \phi_{d'd}) \colon ([1], s) \to ([1], s')$ in $\int \mathbf{N}^{-,+}(C)$.

Now, notice that the desired functoriality follows from the outer square of the diagram (7.9) below. However, in the following (7.9), the commutativity of each inner small trapezoid is definitionally guaranteed, and that of each small triangle is obtained by simple calculation:

$$\tilde{F}(w) \xrightarrow{F_{+}(d')} \tilde{F}(x) \xrightarrow{F([0],w)} F([0],w) \xrightarrow{F(\mathrm{id}_{[0]},d')} F([0],x) \xrightarrow{F(\delta_{1}^{1},\mathrm{id}_{w})} F([0],x) \xrightarrow{F(\delta_{1}^{1},\mathrm{id}_{w})} \xrightarrow{F(\delta_{1}^{1},\mathrm{id}_{w})} F([1],s') \xrightarrow{F(\delta_{0}^{1},\mathrm{id}_{w})} \xrightarrow{F(\delta_{0}^{1},\mathrm{id}_{w})} F([1],s') \xrightarrow{F(\delta_{0}^{1},\mathrm{id}_{w})} \xrightarrow{F([0],w)} F([0],z) \xrightarrow{F(\mathrm{id}_{[0]},d)} F([0],z) \xrightarrow{\tilde{F}_{+}(d)} \tilde{F}(z)$$
(7.9)

Hence the outer square in (7.9) is commutative, which completes the proof of the functoriality of \tilde{F} .

Now we wish to define $\theta \colon \tilde{F} \circ \mathfrak{last} \xrightarrow{\sim} F$ by

$$\theta_{([n],X)} \coloneqq F(\iota_n, \mathrm{id}_{X \circ \iota_n}) \colon \tilde{F}(\mathfrak{last}([n], X)) = F([0], X(n)) \xrightarrow{\sim} F([n], X).$$

Since $(\iota_n, \mathrm{id}_{X \circ \iota_n})$ is a last-weak equivalence, the morphism above is an isomorphism. It just remains to demonstrate the naturality of θ : for any α : $([m], X) \to ([n], Y)$, we shall check the commutativity of the following diagram:

Let (α, ϕ) : $([m], X) \to ([n], Y)$ in $\int \mathbf{N}^{-,+}(C)$ be any representative of the equivalence class $\boldsymbol{\alpha} \in \text{Mor Down}_*(C)$. By Lemma 7.9, we may confine our consideration into two cases: the one where $\alpha(m) = n$ and the one where $X(m) = Y(\alpha(m))$ and $\phi_m = \text{id}_{X(m)}$.

If $\alpha(m) = n$, then $\mathfrak{last}(\alpha) = \phi_m \in \operatorname{Mor} C_+$; hence by definition we have:

$$\tilde{F}(\mathfrak{last}(\alpha)) = \tilde{F}_+(\phi_m) = F(\mathrm{id}_{[0]}, \phi_m) \colon F([0], X(m)) \to F([0], Y(n)).$$

Now send the following commutative diagram under F to obtain the commutativity of (7.10):

Finally, we consider the remaining case: $X(m) = Y(\alpha(m))$ and $\phi_m = \operatorname{id}_{X(m)}$. Let us write $s \coloneqq Y(!_{\alpha(m),n}) \colon Y(\alpha(m)) \twoheadrightarrow Y(n)$. The premise of the case implies that $\operatorname{last}(\alpha) = s$ and that $\tilde{F}(\operatorname{last}(\alpha)) = \tilde{F}_{-}(s)$. Let $\beta \colon [1] \to [n]$ in Δ be given by $\beta(0) = \alpha(m)$ and $\beta(1) = n$. By the defining diagram (7.5), the commutativity of (7.10) follows from the image by F of the following commutative diagram:

Now we have constructed the both that have been required.

Now we wrap up the lemmas above to achieve our goal of this section:

Proof of Theorem 7.3. The case $\Gamma = \int \mathbf{N}(C)$ is addressed in Lemma 7.7. The case $\Gamma = \text{Down}_*(C)$ follows from Lemmas 7.5 and 7.8 and Proposition 7.10. By Lemmas 6.2 and 7.6, the cases $\Gamma = \text{Down}_*(C)$ and $\Gamma = \text{Down}(C)$ are equivalent.

The remaining case is $\Gamma = \int \mathbf{N}^{-,+}(C)$. In Proposition 6.3, note that:

$$\operatorname{Mor} \int \mathbf{N}_{-}^{-, {\operatorname{id}}}(C) \subseteq W_{\int \mathbf{N}^{-, +}(C)}^{\mathfrak{last}}.$$

Therefore, by Lemma 7.6, the case in question is again equivalent to the case $\Gamma = \text{Down}_*(C)$. This concludes the proof.

8. Shapes of $(\infty, 1)$ -diagrams used in the proof of $(\infty, 1)$ -localization

In the preceding discussion, we worked on finite or constructive foundation. In this and the next section, we will work on a sufficiently strong classical set theoretic foundation, like ZFC.

In Theorem 7.3 from the previous section, we proved that the functor $\operatorname{last}_{\Gamma}$ is a 1-localization functor for $\Gamma = \int \mathbf{N}(C)$, $\int \mathbf{N}^{-,+}(C)$, $\operatorname{Down}_*(C)$, $\operatorname{Down}(C)$. This section presents the preparatory technical arguments needed for the next Section 9, where we prove that $\operatorname{last}_{\Gamma}$ is an $(\infty, 1)$ -localization map for the same four instances of Γ . To be more precise, the generalities discussed here will affect the proof for three of those four: $\Gamma = \int \mathbf{N}^{-,+}(C)$, $\operatorname{Down}_*(C)$, $\operatorname{Down}(C)$. The remaining case, $\Gamma = \int \mathbf{N}(C)$, simply follows from the argument of a reflexive subcategory.

More concretely speaking, in this section, we shall construct the following six endofunctors on the category \mathbf{Set}_{Δ} of simplicial sets, and study certain natural transformations between them:

- Dcp and DcpI (Definition 8.1);
- ESd (Definition 8.3) and ESdI (Definition 8.5);
- ESd' and ESdI' (Definition 8.6).

The transformations fit into the following commutative diagram (see Lemma 8.13):



As we will see in Sections 8.3–8.6, the maps $\text{Dcp} X \to \text{ESd}' X$ and $\text{DcpI} X \to \text{ESdI}' X$ are localization maps for any X.

These functors are used to describe the shapes of the $(\infty, 1)$ -diagrams that appear in the proof that $last_{\Gamma}$ is an $(\infty, 1)$ -localization. Specifically, in Lemma 9.3 from the next section, we will construct the following simplicial maps:

- $\operatorname{Dcp} \operatorname{N}(C) \to \operatorname{N}(\Gamma);$
- $\operatorname{DcpI} N(\Gamma) \to N(\Gamma);$
- $\operatorname{ESd}' \operatorname{N}(C) \to \operatorname{N}(C);$
- $\operatorname{ESdI'} \operatorname{N}(\Gamma) \to \operatorname{N}(C).$

Suppose that a quasi-category-valued simplicial map $N(\Gamma) \to Q$ sends last-weak equivalences to equivalences. Then the general properties of the transformations investigated here will allow us to factor $Dcp N(C) \to N(\Gamma) \to Q$ through ESd' N(C)and $DcpI N(\Gamma) \to N(\Gamma) \to Q$ through ESdI' $N(\Gamma)$. This will give $N(C) \to Q$ and $N(\Gamma) \times \Delta[1] \to Q$, which shows the factorization property of a localization map.

This section is organized as follows: in Section 8.1, we will define the six endofunctors on \mathbf{Set}_{Δ} mentioned above. In Section 8.2, we will construct the natural transformations between them, as shown in the diagram above. Sections 8.3–8.5 will be devoted to the investigation of specific properties of these transformations:

- in Section 8.3, we will show that $\text{Dcp } X \to \text{ESd } X$ and $\text{DcpI } X \to \text{ESdI } X$ are universal localization maps;
- in Section 8.4, we will prove that $\operatorname{ESd} X \to \operatorname{ESd}' X$ is an inner anodyne map;
- in Section 8.5, we will demonstrate that $\operatorname{ESdI} X \cup_{\operatorname{ESd} X} \operatorname{ESd}' X \to \operatorname{ESdI}' X$ is an inner anodyne map.

Section 8.6 will conclude this Section 8 by introducing new notation and restating the results. It will facilitate the reference from the next section.

8.1. Some endofunctors on Set_{Δ} . We shall begin by defining two simplicial sets that elaborate to higher dimensions the commutative diagrams used in the proof of Proposition 7.10.

Definition 8.1. For any object $[n] \in \Delta$, we define the category $\text{Dcp}_C[n]$ as the full subcategory of the product category $[n] \times (\Delta/[n])$ spanned by the objects $(x, \alpha: [k] \to [n])$ such that $x \leq \alpha(0)$. Also, we define the category $\text{DcpI}_C[n]$ by the following formulae:

 $\operatorname{Ob}(\operatorname{DcpI}_{C}[n])\coloneqq\operatorname{Ob}(\operatorname{Dcp}_{C}[n])\amalg[n]=(\{0\}\times\operatorname{Ob}\operatorname{Dcp}_{C}[n])\cup(\{1\}\times[n]);$

 $\operatorname{Hom}_{\operatorname{DcpI}_{C}[n]}((0,p),(0,q)) \coloneqq \operatorname{Hom}_{\operatorname{Dcp}_{C}[n]}(p,q);$

 $\operatorname{Hom}_{\operatorname{DcpI}_C[n]}((0,(x,\alpha\colon [k]\to [n])),(1,y))\coloneqq\operatorname{Hom}_{[n]}(\alpha(k),y);$

$$\operatorname{Hom}_{\operatorname{DcpI}_{C}[n]}((1,x),(0,q)) \coloneqq \emptyset;$$

$$\operatorname{Hom}_{\operatorname{DcpI}_{C}[n]}((1,x),(1,y)) := \operatorname{Hom}_{[n]}(x,y).$$

Here, the composition of morphisms is clear, for every composition has its value in a singleton set or is of the form $(0, p) \rightarrow (0, q) \rightarrow (0, r)$. In an evident way, the categories $\text{Dcp}_C[n]$ and $\text{DcpI}_C[n]$ assemble into cosimplicial categories

$$\operatorname{Dcp}_C, \operatorname{DcpI}_C \colon \Delta \to \operatorname{Cat}.$$

We shall denote by Dcp, DcpI: $\mathbf{Set}_{\Delta} \to \mathbf{Set}_{\Delta}$, respectively, the left Kan extensions of $\mathbb{N} \circ \mathrm{Dcp}_{C}$, $\mathbb{N} \circ \mathrm{DcpI}_{C}$: $\Delta \to \mathbf{Set}_{\Delta}$ along the Yoneda embedding $\Delta : \Delta \to \mathbf{Set}_{\Delta}$.

Remark 8.2. Explicitly, for any simplicial set X, we have

$$\operatorname{Dcp} X = \int^{[n] \in \mathbf{\Delta}} N(\operatorname{Dcp}_{C}[n]) \times X_{n} = \operatorname{colim}_{\Delta[n] \to X} N(\operatorname{Dcp}_{C}[n]);$$

$$\operatorname{DcpI} X = \int^{[n] \in \mathbf{\Delta}} N(\operatorname{DcpI}_{C}[n]) \times X_{n} = \operatorname{colim}_{\Delta[n] \to X} N(\operatorname{DcpI}_{C}[n]).$$

In Proposition 7.10, we have constructed a functor $F: C \to D$ and a natural isomorphism $\theta: \tilde{F} \circ \mathfrak{last} \cong F$ from a functor $F: \operatorname{Down}_*(C) \to D$ that sends every \mathfrak{last} -weak equivalence to an isomorphism. The simplicial set $\operatorname{Dcp}(\mathcal{N}(C))$ captures the shape of the diagrams used for the construction of \tilde{F} , namely (7.7) and (7.9). The construction of θ , which used the commutative diagrams (7.11) and (7.12), is covered by the simplicial set $\operatorname{DcpI}(\mathcal{N}(\operatorname{Down}_*(C)))$.

For a good control of compositions and inverses in quasi-categories, we need a few more endofunctors of \mathbf{Set}_{Δ} . We go on to define them.

Definition 8.3. For the purpose of this definition, let $F: \Delta \to \Delta$ denote the functor $F[n] = [n] \star [n] = [2n + 1]$. We shall define the endofunctor ESd: $\mathbf{Set}_{\Delta} \to \mathbf{Set}_{\Delta}$ as the precomposition of $F^{\mathrm{op}}: \Delta^{\mathrm{op}} \to \Delta^{\mathrm{op}}$: for any simplicial set X, we have $\mathrm{ESd}(X)_n = X_{2n+1}$.

Remark 8.4. The endofunctor ESd: $\operatorname{Set}_{\Delta} \to \operatorname{Set}_{\Delta}$ is not new: it is the composition of the two well-known functors. Let $\operatorname{Set}_{\Delta \times \Delta}$ denote the category of bisimplicial sets: set-valued presheaves on the product category $\Delta \times \Delta$. The *total décalage functor* Dec: $\operatorname{Set}_{\Delta} \to \operatorname{Set}_{\Delta \times \Delta}$ and the *diagonal functor* diag: $\operatorname{Set}_{\Delta \times \Delta} \to \operatorname{Set}_{\Delta}$ are given by the precomposition of the *join* \star : $\Delta \times \Delta \to \Delta$ and the *diagonal inclusion* $\Delta \to \Delta \times \Delta$. It is easy to see that these two functors compose to produce our ESd.

Since the left and the right Kan extensions of F^{op} gives the left and the right adjoints of ESd, note that ESd preserves limits and colimits, as should also be clear from the formula $\text{ESd}(X)_n = X_{2n+1}$.

Definition 8.5. We define the cosimplicial simplicial set $\mathrm{ESdI}_{\Delta} : \Delta \to \operatorname{Set}_{\Delta}$ by

$$\mathrm{ESdI}_{\mathbf{\Delta}}[n]_k \coloneqq \prod_{\substack{I \sqcup J = [k] \\ I < J}} \mathrm{Hom}_{\mathbf{Poset}}(I \star I \star J, [n]).$$

Here, the disjoint union II is taken over all the partitions $[k] = I \sqcup J$ satisfying i < j for any $i \in I$ and $j \in J$. We define the endofunctor ESdI: $\mathbf{Set}_{\Delta} \to \mathbf{Set}_{\Delta}$ as the left Kan extension of ESdI_{Δ} along the Yoneda embedding $\Delta : \Delta \to \mathbf{Set}_{\Delta}$.

With this definition, the set of n-simplices of ESdI X can be expressed as the following disjoint union:

$$(\operatorname{ESdI} X)_n \cong \prod_{i=0}^{n+1} X_{n+i}.$$

Definition 8.6. Let us define the cosimplicial posets $\text{ESd}'_P, \text{ESdI}'_P: \Delta \to \text{Poset}$ by:

$$\operatorname{ESd}_{P}^{\prime}[n] \coloneqq [n]^{[1]};$$

$$\operatorname{ESdI}_{P}^{\prime}[n] \coloneqq (\{0\} \times [n]^{[1]}) \cup (\{1\} \times [n]).$$

Here, the latter set is regarded as a poset by pulling back the order through the following injective map, whose codomain is endowed with the product order:

$$\operatorname{ESdI}_{P}^{\prime}[n] \to [1] \times [n]^{[1]};$$

(0, f) \mapsto (0, f);
(1, x) \mapsto (1, const_x).

We define the endofunctors $\mathrm{ESd}', \mathrm{ESdI}' : \mathbf{Set}_{\Delta} \to \mathbf{Set}_{\Delta}$ as the left Kan extensions along the Yoneda embedding $\Delta : \Delta \to \mathbf{Set}_{\Delta}$ of the post-compositions of ESd'_P and ESdI'_P with the embedding $\mathbf{Poset} \hookrightarrow \mathbf{Cat} \hookrightarrow \mathbf{Set}_{\Delta}$. Let us explain the shapes and the intended meanings of the simplicial sets we have defined, emphasizing lower dimensions. We begin with Dcp. Since we would like to show that $N(\mathfrak{last}): N(\mathrm{Down}(C)) \to N(C)$ is a localization map, we need to construct a map $N(C) \to Q$, given a quasi-category-valued simplicial map $N(\mathrm{Down}(C)) \to Q$ that sends \mathfrak{last} -weak equivalences to equivalences. This endofunctor Dcp is designed to make $\mathrm{Dcp}(N(C))$ capture how simplices in the nerve N(C) of our fixed Reedy category C should be mapped to $N(\mathrm{Down}(C))$, considering \mathfrak{last} -weak equivalences as invertibles. Let us say we have a 2-simplex in N(C): a commutative triangle $X_0 \to X_1 \to X_2$ in C. Assume that the following is the Reedy factorization of the triangle:



Let us write:

$$S \coloneqq \left(X_0 \xrightarrow{s_2} Y_2 \xrightarrow{s} Y_1 \right)$$

for the length-2 composable chain in C, representing an object in $\text{Down}_*(C)$. The factorized triangle $X_0 \to X_1 \to X_2$ should be mapped to the following in $\text{Down}_*(C)$:



Here, the labels for objects and morphisms are abbreviated for simplicity. Also note that this diagram includes the diagrams (7.7) and (7.9) from the proof of Proposition 7.10.

The diagram (8.1) is, itself, not a single 2-simplex in $N(\text{Down}_*(C))$. However, since all arrows labeled with \sim are *last*-weak equivalences, if we consider these arrows as invertible, we can compute the "composition" of this diagram to yield a valid 2-simplex.

In order to describe the shape of the concrete diagram (8.1), we consider the following 2-dimensional simplicial set, which is an incomplete prototype of $\text{Dcp }\Delta[2]$:



Here, ~ on an edge indicates that the corresponding morphism in (8.1) is a lastweak equivalence, meaning it preserves the left coordinate and the maximum of the right coordinate. If we were to formulate the exact shape of the diagram as $Dcp \Delta[2]$, we would have the following:

$${}^{"}\mathrm{Dcp}_{P}[n] {}^{"} \stackrel{!}{\coloneqq} \{(x,\alpha) \mid x \in [n], \emptyset \neq \alpha \subseteq [n], x \leq \min \alpha \} \text{ poset}; \\ {}^{"}\mathrm{Dcp}\,\Delta[n] {}^{"} \stackrel{?}{\coloneqq} \mathrm{N}(\mathrm{Dcp}_{P}[n]).$$

However, for a better interaction with degeneracies, we would like to consider as α general [n]-valued morphisms in Δ , not just the [n]-valued injectives; hence the construction in Definition 8.1. This makes the simplicial set Dcp Δ [2] higher dimensional than 2, and too visually complex to be depicted here, but we can more easily manipulate it in our proofs.

In each vertex $(x, \alpha) \in \operatorname{Ob}(\operatorname{Dcp}_C[n])$ of $\operatorname{Dcp}\Delta[n]$, its right coordinate α parametrizes the length of a chain in C_- , which is an object in $\operatorname{Down}_*(C)$, and the left coordinate x controls the Reedy factorization from which the chain is obtained. More rigourously put, if $X_0 \to X_1 \to \cdots \to X_n$ is an *n*-simplex in $\operatorname{N}(C)$, then the vertex $(x, \alpha : [k] \to [n])$ in $\operatorname{Dcp}(\Delta[n])$ corresponds to the chain $Z_0 \twoheadrightarrow \cdots \twoheadrightarrow Z_k$ below in C_- , representing an object in $\operatorname{Down}_*(C)$:

This glues together to form a simplicial map $Dcp(N(C)) \rightarrow N(Down_*(C))$, as shall be demonstrated in Lemma 9.3.

In order to compose simplices forming diagrams like (8.1), we need to properly treat the reverse-direction simplices. For that purpose, we simply collapse the arrows with ~ and simplices containing them into degeneracies. If we do so in $Dcp(\Delta[2])$, then we obtain $ESd(\Delta[2])$, which has the following shape:



Here, a vertex (x, α) in $\text{Dcp}(\Delta[2])$ is collapsed into the vertex $(x, \max \alpha)$. This collapse works as intended. This generalizes to higher dimensions and glues together to form a natural simplicial map $\text{Dcp} X \to \text{ESd} X$ for any simplicial set X. This natural map is a localization map, as we will show in Proposition 8.19.

We are left only with forward-direction simplices in ESd, but the composition of simplices in $\operatorname{ESd}(\Delta[n])$ is not specified in the simplicial set: for example, note that the non-degenerate simplices in $\operatorname{ESd}(\Delta[2])$ are the vertices, the arrows, and the four small triangle explicitly depicted in the diagram (8.2). Therefore, even if we have a map with domain $\operatorname{ESd}(N(C))$, the values of simplices in N(C) is not determined. We need an extension that contains the "composition" of these simplices. This is the job ESd' does. The simplicial set $\operatorname{ESd}'(\Delta[n]) = N([n]^{[1]})$ is obtained by considering a poset that is obtained by all lacking compositions in $\operatorname{ESd}(\Delta[n])$. This makes simplicial set $\operatorname{ESd}'(\Delta[n])$ a quasi-category for each n. Most desirably, for each X, we have natural inclusions $X \hookrightarrow \operatorname{ESd}' X$ and $\operatorname{ESd} X \hookrightarrow \operatorname{ESd}' X$, the latter of which is, as shown in Proposition 8.23, inner anodyne.

The three endofunctors Dcp, ESd, and ESd' are used to describe the shapes of the diagram used to construct a factorizing functor $N(C) \rightarrow Q$ from a functor $N(Down(C)) \rightarrow Q$. The construction of the natural transformation that ensures the functor is a factorization, requires the other three endofunctors: DcpI, ESdI, and ESdI'. These endofunctors with "I" evaluated at a simplicial set X serve as a cylinder that connects the value at X of the version without "I" and the original simplicial set X. For example, ESdI(Δ [1]) has the following shape:



To conclude this subsection, we shall check an important property of the functors we have defined:

Lemma 8.7. The functors Dcp, DcpI, ESd, ESdI, ESd', and ESdI' preserve monomorphisms.

Proof. By Corollary 2.36, the claim follows from the construction of the functors in question. \Box

8.2. Some natural transformations between the endofunctors. We shall now construct some natural transformations between the endofunctors we have defined. They should fit into the following commutative diagram (see Lemma 8.13):

Here, the numbers stand for the definition where the natural transformation is constructed.

Definition 8.8. Let $X \in \mathbf{Set}_{\Delta}$ be a simplicial set. Below we shall construct surjective simplicial maps $\operatorname{Dcp} X \twoheadrightarrow \operatorname{ESd} X$ and $\operatorname{DcpI} X \twoheadrightarrow \operatorname{ESdI} X$, which are natural in X. We shall consider these natural maps as the canonical connecting maps between those simplicial sets.

Construction. Since the functors in question are all colimit-preserving and colimits preserve surjections, we may confine our attention to the case where X is a representable simplicial set $\Delta[n]$. We first construct $\text{Dcp}\,\Delta[n] \to \text{ESd}\,\Delta[n]$. Let $\varphi: [n] \to \text{Dcp}_C[n]$ be a k-simplex in $\text{Dcp}\,\Delta[n] = N(\text{Dcp}_C[n])$. Write:

$(x_i, \alpha_i \colon [m_i] \to [n]) \coloneqq \varphi(i)$	for $0 \le i \le k$;
$(!_{x_i,x_j},\beta_{i,j}:\alpha_i\to\alpha_j)\coloneqq\varphi(!_{i,j})$	for $0 \le i \le j \le k$.

Since $\beta_{i,j}$ above is a morphism in $\mathbf{\Delta}/[n]$, we deduce:

$$x_k \leq \alpha_k(0) \leq \alpha_k(\beta_{0,k}(0)) = \alpha_0(0) \leq \alpha_0(m_0);$$

$$\alpha_i(m_i) = \alpha_j(\beta_{i,j}(m_i)) \le \alpha_j(m_j) \quad \text{for } 1 \le i \le j \le k.$$

Therefore, the following chain of inequalities holds:

 $x_0 \leq x_1 \leq \cdots \leq x_k \leq \alpha_0(m_0) \leq \alpha_1(m_1) \leq \cdots \leq \alpha_k(m_k).$

The above (2k + 2)-chain in [n] represents a k-simplex in ESd $\Delta[n]$. This defines a simplicial map $\operatorname{Dcp}\Delta[n] \to \operatorname{ESd}\Delta[n]$, natural in [n], which is surjective by construction.

Next, we construct $\text{DcpI}\Delta[n] \twoheadrightarrow \text{ESdI}\Delta[n]$. Let $\varphi \colon [k] \to \text{DcpI}_C[n]$ be a k-simplex in $\text{DcpI}\Delta[n] = N(\text{DcpI}_C[n])$. We wish to assign to φ an element of

$$\operatorname{ESdI}_{\boldsymbol{\Delta}}[n]_{k} = \coprod_{\substack{I \sqcup J = [k] \\ I < J}} \operatorname{Hom}_{\operatorname{\mathbf{Poset}}}(I \star I \star J, [n]).$$

Let the subposet $I = \{0, 1, ..., l-1\}$ and $J = \{l, l+1, ..., k\}$ of [k], respectively, be the inverse images of $\{0\} \times \text{Ob} \operatorname{Dcp}_C[n]$ and $\{1\} \times [n]$ under φ . Write:

$$\begin{aligned} (0,(x_i,\alpha_i\colon [m_i]\to [n])) &\coloneqq \varphi(i) \quad \text{for } i\in I; \\ (1,y_i) &\coloneqq \varphi(i) \quad \text{for } i\in J. \end{aligned}$$

From the same reasoning as above, we deduce that the following chain of inequalities holds, yielding an order-preserving map $I \star I \star J \to [n]$:

$$x_0 \le x_1 \le \dots \le x_{l-1}$$
$$\le \max \alpha_0 \le \max \alpha_1 \le \dots \le \max \alpha_{l-1}$$
$$\le y_l \le y_{l+1} \le \dots \le y_k.$$

This constitutes an element of $\operatorname{ESdI}\Delta[n]_k = \operatorname{ESdI}\Delta[n]_k$ above, and tying these corresponding elements together gives a simplicial map $\operatorname{DcpI}\Delta[n] \to \operatorname{ESdI}\Delta[n]$, naturally in [n], which is surjective by construction.

Definition 8.9. Let $X \in \mathbf{Set}_{\Delta}$ be a simplicial set. In the subsequent Construction, we shall give injective simplicial maps $\mathrm{ESd} X \hookrightarrow \mathrm{ESd}' X$ and $\mathrm{ESdI} X \hookrightarrow \mathrm{ESdI}' X$, which are natural in X. We shall think of these natural transformations as the canonical ones between the involved simplicial sets.

Construction. Again, we may confine our attention to the case where X is a representable simplicial set $\Delta[n]$. We first construct $\operatorname{ESd}\Delta[n] \hookrightarrow \operatorname{ESd}'\Delta[n]$. For any $[k] \in \Delta$, consider the following order-preserving bijection (which is not an isomorphism of posets):

$$u_{[k]}: [1] \times [k] \to [k] \star [k] = [2k+1]; \ (i,x) \mapsto i * (k+1) + x.$$

This is natural in [k]. Let an order-preserving map $f: [k] \star [k] \to [n]$ represent a k-simplex in ESd $\Delta[n]$. The composite $f \circ u_{[k]}: [1] \times [k] \to [n]$ has the adjunct map $[k] \to [n]^{[1]}$, which is an element of ESd' $\Delta[n]_k = \operatorname{N}(\operatorname{ESd}'_P[n])_k$. By corresponding this adjunct to the k-simplex f, we obtain a simplicial map ESd $\Delta[n] \hookrightarrow \operatorname{ESd}' \Delta[n]$, naturally in [n]. This family of maps induces a natural transformation ESd $\Rightarrow \operatorname{ESd}'$. By Corollary 2.38, the injectivity of the induced transformation follows from that of each ESd $\Delta[n] \hookrightarrow \operatorname{ESd}' \Delta[n]$, which is clear.

Next, we construct $\operatorname{ESdI}\Delta[n] \hookrightarrow \operatorname{ESdI'}\Delta[n]$. In order to arbitrarily take a ksimplex in $\operatorname{ESdI}\Delta[n]$, let $[k] = I \sqcup J$ be a partition with I < J, and let $f: I \star I \star J \to$ [n] be a map. By applying the above argument to $f|_{I\star I}: I \star I \to [n]$, we may obtain a map $I \to [n]^{[1]}$. Bundling this with the map $f|_J: J \to [n]$ gives a map

$$[k] = I \sqcup J = I \star J \to \{0\} \times [n]^{[1]} \cup \{1\} \times [n] = \mathrm{ESdI}'_P[n].$$

This gives a k-simplex in ESdI' $\Delta[n] = N(\text{ESdI}'_P[n])$. We have thus constructed a simplicial map ESdI $\Delta[n] \hookrightarrow \text{ESdI}' \Delta[n]$, naturally in [n].

The injectivity of ESdI $X \to \text{ESdI}' X$ for any simplicial set X may be shown by Corollary 2.38.

Definition 8.10. For a simplicial set $X \in \mathbf{Set}_{\Delta}$, we will demonstrate the construction of canonical injective simplicial maps $X \hookrightarrow \mathrm{ESd}' X$ and $X \times \Delta[1] \hookrightarrow \mathrm{ESdI}' X$, which are natural in X. These constructions will be referred to as the canonical connecting maps between these simplicial sets.

Construction. For the third time, we may consider only the case where X is a representable simplicial set $\Delta[n]$. The map $\Delta[n] \hookrightarrow \operatorname{ESd}' \Delta[n]$ may be obtained by sending the constant inclusion

const:
$$[n] \hookrightarrow [n]^{[1]}; x \mapsto \text{const}_x = (x, x),$$

which is natural in [n], with the nerve functor N: Poset \rightarrow Cat \rightarrow Set_{Δ}.

In order to construct the map $\Delta[n] \times \Delta[1] \hookrightarrow \operatorname{ESdI}' \Delta[n]$, condider the following inclusion of posets:

E a 1

 \square

$$\begin{split} & [n] \times [1] \hookrightarrow \mathrm{ESdI}'_P[n] = \{0\} \times [n]^{[1]} \cup \{1\} \times [n]; \\ & (x,0) \mapsto (0, \mathrm{const}_x), \\ & (x,1) \mapsto (1,x). \end{split}$$

This is natural in [n]. We may then apply the nerve functor to this inclusion to obtain the desired map.

The injectivity of these maps follows from Corollary 2.38, as before.

Definition 8.11. Given a simplicial set $X \in \mathbf{Set}_{\Delta}$, we aim to construct canonical injective simplicial maps $\operatorname{Dcp} X \hookrightarrow \operatorname{DcpI} X$, $\operatorname{ESd} X \hookrightarrow \operatorname{ESdI} X$, and $\operatorname{ESd}' X \hookrightarrow \operatorname{ESdI}' X$. These constructions, detailed below, are natural in X and will serve as the canonical natural transformation between the involved functors.

Construction. For the fourth time, we may confine our attention to the case where X is a representable simplicial set $\Delta[n]$. $\operatorname{DcpI}_C[n]$ and $\operatorname{ESdI}'_P[n]$ are defined in such a way that there is an obvious inclusions $\operatorname{Dcp}_C[n] \hookrightarrow \operatorname{DcpI}_C[n]$ and $\operatorname{ESd}'_P[n] \hookrightarrow \operatorname{ESdI}'_P[n]$, which induce the desired simplicial maps $\operatorname{Dcp}\Delta[n] \hookrightarrow \operatorname{DcpI}\Delta[n]$ and $\operatorname{ESd}'\Delta[n] \hookrightarrow \operatorname{ESdI}'\Delta[n]$. This leaves us with the construction of $\operatorname{ESd}\Delta[n] \hookrightarrow \operatorname{ESdI}\Delta[n]$. Unwinding the definition:

$$\operatorname{ESd}\Delta[n]_{k} \cong \operatorname{Hom}_{\operatorname{Poset}}([k] \star [k], [n]);$$
$$\operatorname{ESdI}\Delta[n]_{k} = \coprod_{\substack{I \sqcup J = [k]\\ I < I}} \operatorname{Hom}_{\operatorname{Poset}}(I \star I \star J, [n])$$

Therefore, the set $\operatorname{ESd} \Delta[n]_k$ embeds isomorphically to the component of $\operatorname{ESdI} \Delta[n]_k$ indexed by the partition I = [k] and $J = \emptyset$. This gives the desired simplicial map.

The naturality of these inclusions is clear from the definitions. The injectivity of these maps may be shown by using Corollary 2.38. \Box

Definition 8.12. Let X be a simplicial set. The injective simplicial maps $X \hookrightarrow$ DcpI X and $X \hookrightarrow$ ESdI X, natural in X and constructed below, will be considered as the canonical ones between the involved simplicial sets.

Construction. For the fifth time, we may confine our attention to the case where X is a representable simplicial set $\Delta[n]$. The first of the desired maps:

$$\mathbf{N}[n] = \Delta[n] \hookrightarrow \mathrm{DcpI}\,\Delta[n] = N(\mathrm{DcpI}_C[n])$$

may be obtained by applying the nerve functor to the inclusion of the full subcategory:

$$[n] = \{1\} \times [n] \hookrightarrow \mathrm{DcpI}_C[n]; \ x \mapsto (1, x).$$

Note that this is natural in [n].

For the construction of the other map $\Delta[n] \hookrightarrow \operatorname{ESdI} \Delta[n]$, remember:

$$\begin{split} \Delta[n]_k &= \operatorname{Hom}_{\operatorname{\textbf{Poset}}}([k], [n]);\\ \operatorname{ESdI}\Delta[n]_k &= \operatorname{ESdI}_{\Delta}[n]_k = \coprod_{\substack{I \sqcup J = [k]\\ I < J}} \operatorname{Hom}_{\operatorname{\textbf{Poset}}}(I \star I \star J, [n]). \end{split}$$

Therefore,

$$\operatorname{Hom}_{\operatorname{\textbf{Poset}}}([k], [n]) \cong \operatorname{Hom}_{\operatorname{\textbf{Poset}}}(\emptyset \star \emptyset \star [k], [n]) \hookrightarrow \coprod_{\substack{I \sqcup J = [k] \\ I < J}} \operatorname{Hom}_{\operatorname{\textbf{Poset}}}(I \star I \star J, [n])$$

gives a map between the sets of k-simplices. Since this is natural in $[k] \in \Delta^{\text{op}}$ and $[n] \in \Delta$, we obtain what we want.

To show the injectivity of these maps, it suffices to remember Corollary 2.38. \Box

Finally, we remember the diagram that we presented at the beginning of this subsection:

Lemma 8.13. The diagram (8.3) at the beginning of this subsection, which we shall reproduce here for the reader's convenience, commutes:

$$\begin{array}{c|c} \operatorname{Dcp} X \times \{0\} & \xrightarrow{8.8} \operatorname{ESd} X \times \{0\} & \xrightarrow{8.9} \operatorname{ESd}' X \times \{0\} & \xleftarrow{8.10} X \times \{0\} \\ & s.11 & s.11 & s.11 & \| \\ & \operatorname{DcpI} X & \xrightarrow{8.8} & \operatorname{ESdI} X & \xrightarrow{8.9} \operatorname{ESdI'} X & X \times \{0\} \\ & s.12 & s.12 & & & \\ & X \times \{1\} & \xrightarrow{8.10} & & & \\ & X \times \{1\} & \xrightarrow{8.10} & X \times \{1\} & \xrightarrow{8.10} & X \times \Delta[1] \end{array}$$

Remember that the numbers stand for the definition where the natural transformation is constructed.

Proof. By construction.

8.3. Properties of the transformations 1: two universal localizations. In Definition 8.8, we have constructed the canonical natural transformations $Dcp \Rightarrow ESd$ and $DcpI \Rightarrow ESdI$. The purpose of this section is to show that the following three classes of natural maps are universal localizations in the sense of [16, Definition 02M0]:

- the canonical map $\operatorname{Dcp} X \twoheadrightarrow \operatorname{ESd} X$ for each simplicial set X (Proposition 8.19);
- the canonical map $\operatorname{DcpI} X \twoheadrightarrow \operatorname{ESdI} X$ for each simplicial set X (Proposition 8.21);
- the map $\operatorname{Dcpl} X \cup_{\operatorname{Dcp} X} \operatorname{Dcp} Y \twoheadrightarrow \operatorname{ESdl} X \cup_{\operatorname{ESd} X} \operatorname{ESd} Y$ for any simplicial map $X \to Y$, induced by the canonical transformations (Corollary 8.22).

Using Proposition 02M9 and Proposition 02MA from [16], we can show that these three types of maps are universal localizations by decomposing both the domain and codomain into smaller parts, verifying universal localization on each part, and then assembling the results via colimits. Lemma 8.15 and Corollary 8.16 presented below provide the fundamental building blocks of these proofs.

Notation 8.14. Let $[n] \in Ob \Delta_a$. For the purpose of this subsection, we use the following symbol for a functor:

$$\lambda_{[n]} \colon \mathbf{\Delta}/[n] \to [n]; \; (\alpha \colon [m] \to [n]) \mapsto \alpha(m).$$

Here, note that $\Delta/[n]$ may be extended to the case n = -1 by regarding it as a full subcategory of $\Delta_a/[n]$, i.e., $\Delta/[-1] = \emptyset$. Note that this is natural in $[n] \in \Delta$. We shall also write

$$\lambda_{[n]} \coloneqq \mathrm{N}(\lambda_{[n]}) \colon \mathrm{N}(\mathbf{\Delta}/[n]) \to \Delta[n]$$

Remember our notation: $\Delta[-1] = N[-1] = \emptyset$.

Lemma 8.15. For any $[n] \in \Delta_a$, the functor $\lambda_{[n]} \colon N(\Delta/[n]) \to \Delta[n]$ in Notation 8.14 is a universal localization in the sense of [16, Definition 02M0].

Proof. According to [16, Proposition 04JT], to show our claim, it suffices to construct a section $u: \Delta[n] \to \mathcal{N}(\mathbf{\Delta}/[n])$ to $\lambda_{[n]}$, and to check that the composition $u \circ \lambda_{[n]}$ and the identity $\mathrm{id}_{\Delta[n]}$ belong to the same connected component of the mapping space $\mathrm{Fun}_{\Delta[n]}(\lambda_{[n]}, \lambda_{[n]})$ over $\Delta[n]$ (see [16, Construction 01AB] for the notation). In terms of 1-categories, it is enough to construct a functor $u: [n] \to \mathbf{\Delta}/[n]$ strictly satisfying $\lambda_{[n]} \circ u = \mathrm{id}_{[n]}$, and a natural transformation $\theta: \mathrm{id}_{\mathbf{\Delta}/[n]} \Rightarrow u \circ \lambda_{[n]}$ satisfying $\lambda_{[n]} \triangleleft \theta = \mathrm{id}_{\lambda_{[n]}}$.

satisfying $\lambda_{[n]} \triangleleft \theta = \mathrm{id}_{\lambda_{[n]}}$. We define u as follows. For each object $x \in [n]$, we set $u(x) \coloneqq \iota_{[x]}^n$. Remember, for any non-empty subset $S = \{x_0 < \cdots < x_k\} \subseteq [n]$, the injective order-preserving map $\iota_S^n \colon [k] \to [n]$ has been defined by $\iota_S^n(i) = x_i$, and the morphism $\iota_{[x]}^n \colon [x] \to [n]$ in Δ gives an object of $\Delta/[n]$. If $x \leq y$ in [n], the morphism

$$u(!_{x,y}): u(x) = \iota^n_{[x]} \to \iota^n_{[y]} = u(y)$$

is given by $u(!_{x,y}) \coloneqq \iota^y_{[x]} \colon [x] \to [y]$, which evidently satisfies the condition for a morphism in Δ to be a morphism in $\Delta/[n]$. This defines a functor $u \colon [n] \to \Delta/[n]$, and we can easily check:

$$(\lambda_{[n]} \circ u)(x) = \iota_{[x]}^n(x) = x = \mathrm{id}_{[n]}(x).$$

We proceed to construct the natural transformation θ . For each object in $\Delta/[n]$, i.e., a morphism $\alpha \colon [m] \to [n]$ in Δ , we need to define a morphism

$$\theta_{\alpha} \colon \alpha = \mathrm{id}_{\Delta/[n]}(\alpha) \to u(\lambda_{[n]}(\alpha)) = \iota^{n}_{[\alpha(m)]}$$

in $\Delta/[n]$. This is given by the morphism $[m] \to [\alpha(m)]$; $x \mapsto \alpha(x)$ in Δ , which is indeed a morphism $\alpha \to \iota^n_{[\alpha(m)]}$ in $\Delta/[n]$. The naturality of the family θ_α in α may be simply checked. Indeed, let the following commutative triangle in Δ present an arbitrary morphism β in $\Delta/[n]$:



Then, for any $x \in [m]$, we compute:

$$(\theta_{\alpha'} \circ \mathrm{id}_{[m]}(\beta))(x) = \alpha'(\beta(x)) = \alpha(x) = \iota_{[\alpha(m)]}^{\alpha'(m')}(\alpha(x)) = ((u \circ \lambda_{[n]})(\beta) \circ \theta_{\alpha})(x),$$

which is the desired naturality. Since [n] is a poset, the equation $\lambda_{[n]} \triangleleft \theta = \mathrm{id}_{\lambda_{[n]}}$ trivially holds. This completes the proof.

Corollary 8.16. Let $[m], [n] \in \Delta_a$ be objects of the augmented simplex category. Using the notation from Notation 8.14, consider the following simplicial map:

 $\mathrm{id}_{\Delta[m]} \times \lambda_{[n]} \colon \Delta[m] \times \mathrm{N}(\mathbf{\Delta}/[n]) \to \Delta[m] \times \Delta[n].$

Then this map is a universal localization in the sense of [16, Definition 02M0].

Proof. Since the product preserves universal localizations (Lemma 2.26), this follows from Lemma 8.15. $\hfill\square$

Building on the minimal components of the domain and codomain of our maps, treated in Lemma 8.15 and Corollary 8.16, we now proceed to slightly larger parts:

Lemma 8.17. Let $[n] \in \Delta_a$ be an object of the augmented simplex category. Then the canonical map $\text{Dcp }\Delta[n] \to \text{ESd }\Delta[n]$ from Definition 8.8 is a universal localization in the sense of [16, Definition 02M0].

Proof. The case n = -1 is trivial, so we may assume $n \ge 0$.

For each $-1 \leq l \leq n$, let $C_{\leq l}^n$ and C_l^n denote the full subcategories of $\text{Dcp}_C[n]$ determined by:

$$Ob(C_{\leq l}^{n}) \coloneqq \{(x, \alpha) \in Ob(Dcp_{C}[n]) \mid x \leq l\};\$$
$$Ob(C_{l}^{n}) \coloneqq \{(x, \alpha) \in Ob(Dcp_{C}[n]) \mid x \leq l \leq \alpha(0)\}$$

We set $F_{\leq l} := \mathcal{N}(C^n_{\leq l}), F_l := \mathcal{N}(C^n_l) \subseteq \operatorname{Dcp} \Delta[n]$. Then we have the following filtration of simplicial sets:

$$\emptyset = F_{\leq -1} \subseteq F_{\leq 0} \subseteq F_{\leq 1} \subseteq \cdots \subseteq F_{\leq n} = \operatorname{Dcp} \Delta[n].$$

We shall also consider the following simplicial subsets of ESd $\Delta[n]$: for each $-1 \leq l \leq n$:

$$\begin{aligned} (G_{\leq l})_k &\coloneqq \{f \colon [2k+1] = [k] \star [k] \to [n] \mid f(k) \leq l\}; \\ (G_l)_k &\coloneqq \{f \colon [2k+1] = [k] \star [k] \to [n] \mid f(k) \leq l \leq f(k+1)\}. \end{aligned}$$

Then we again obtain the following filtration of simplicial sets:

$$\emptyset = G_{\leq -1} \subseteq G_{\leq 0} \subseteq G_{\leq 1} \subseteq \cdots \subseteq G_{\leq n} = \operatorname{ESd} \Delta[n].$$

Notice that the canonical map $\operatorname{Dcp} \Delta[n] \to \operatorname{ESd} \Delta[n]$ restricts to $F_{\leq l} \to G_{\leq l}$ and $F_l \to G_l$ for each $-1 \leq l \leq m$.

By induction, in the increasing order of l, we shall show that the map $F_{\leq l} \to G_{\leq l}$ is a universal localization, which will imply our desired lemma. The base case l = -1 is trivial. Let $0 \leq l \leq m$ and suppose that $F_{\leq l-1} \to G_{\leq l-1}$ is a universal localization; we need to show that $F_{\leq l} \to G_{\leq l}$ is a universal localization. Consider the following commutative cube, consisting of inclusions of simplicial subsets (\hookrightarrow) and the restrictions of the canonical map $\text{Dcp } \Delta[n] \to \text{ESd } \Delta[n] (\to)$:



In this cube, the front and the back faces are pushout squares by construction, and the hooked arrows are injective. Therefore, by [16, Proposition 02MA], in order to show that $F_{\leq l} \to G_{\leq l}$ is a universal localization, it suffices to check that the three maps $F_{l-1} \cap F_l \to G_{l-1} \cap G_l$, $F_l \to G_l$, and $F_{\leq l-1} \to G_{\leq l-1}$ are universal localizations. Of those three, the third map $F_{\leq l-1} \to G_{\leq l-1}$ is a universal localization by the induction hypothesis. Now, using the notation from Notation 8.14, we observe that the first map $F_{l-1} \cap F_l \to G_{l-1} \cap G_l$ is isomorphic to:

$$\mathrm{id}_{\Delta[l-1]} \times \lambda_{[n-l]} \colon \Delta[l-1] \times \mathrm{N}(\mathbf{\Delta}/[n-l]) \to \Delta[l-1] \times \Delta[n-l],$$

and that the second map $F_l \to G_l$ is isomorphic to the map:

$$\mathrm{id}_{\Delta[l]} \times \lambda_{[n-l]} \colon \Delta[l] \times \mathrm{N}(\mathbf{\Delta}/[n-l]) \to \Delta[l] \times \Delta[n-l].$$

48 FINITE REEDY CATEGORIES AS LOCALIZATIONS OF FINITE DIRECT CATEGORIES

These two maps are universal localizations by Corollary 8.16. Thus we have proved the sufficient condition for the map $F_{\leq l} \to G_{\leq l}$ to be a universal localization. This completes the induction, and we have obtained the desired claim.

Corollary 8.18. Let $[m], [n] \in \Delta$ be objects. Then the following join of the canonical simplicial map from Definition 8.8 and the identity map is a universal localization in the sense of [16, Definition 02M0]:

$$\operatorname{Dcp} \Delta[m] \star \Delta[n] \to \operatorname{ESd} \Delta[m] \star \Delta[n].$$

Proof. By using the fact that the join $\star X$ of a simplicial set X is preserves universal localizations (Corollary 2.28), this follows from Lemma 8.17.

Now, we are ready to prove one of the main results of this section:

Proposition 8.19. Let $X \in \mathbf{Set}_{\Delta}$ be a simplicial set. Then the canonical simplicial map $\operatorname{Dcp} X \to \operatorname{ESd} X$ in Definition 8.8 is a universal localization in the sense of [16, Definition 02M0].

Proof. As [16, Proposition 02M9] states, the class of universal localizations is closed under filtered colimits in $\mathbf{Set}_{\Delta}^{[1]}$. Therefore, we may assume that X is a finite simplicial set. We shall show the claim by induction on the dimension and the number of the highest-dimensional non-degenerate simplices of X. If X is empty, the claim is trivial. If X is non-empty, let $\sigma: \Delta[n] \to X$ be one of the highestdimensional non-degenerate simplices of X. There is the following pushout square, with X' a subcomplex of X having fewer dimension-n non-degenerate simplices:



By sending this square under Dcp and ESd, we obtain the following commutative cube:



In this cube, the front and the back faces are pushout squares by construction, and the hooked arrows are injective. The maps $\text{Dcp}\,\partial\Delta[n] \to \text{ESd}\,\partial\Delta[n]$ and $\text{Dcp}\,X' \to \text{ESd}\,X'$ are universal localizations by the induction hypothesis. The map $\text{Dcp}\,\Delta[n] \to \text{ESd}\,\Delta[n]$ is a universal localization by Lemma 8.17. Therefore, by [16, Proposition 02MA], the map $\text{Dcp}\,X \to \text{ESd}\,X$ is a universal localization. This completes the induction, and we have obtained the desired claim. \Box

The following lemma is the largest component of the remaining main universal localization result:

Lemma 8.20. Let $[n] \in \Delta$ be a simplex. Then the canonical simplicial map $DcpI \Delta[n] \rightarrow ESdI \Delta[n]$ is a universal localization in the sense of [16, Definition 02M0].

Proof. For each $-1 \leq l \leq n$, let $C_{\leq l}^n$ and C_l^n denote the full subcategories of $\text{DcpI}_C[n]$ determined by:

$$\operatorname{Ob}(C^n_{\leq l}) \coloneqq (\{0\} \times \{(x, \alpha) \in \operatorname{Ob}(\operatorname{Dcp}_C[n]) \mid \max \alpha \leq l\}) \cup (\{1\} \times [n]);$$

 $\operatorname{Ob}(C_l^n)\coloneqq \left(\{0\}\times\{(x,\alpha)\in\operatorname{Ob}(\operatorname{Dcp}_C[n])\mid \max\alpha\leq l\}\right)\cup \left\{(1,x)\mid l\leq x\leq n\right\}.$

We set $F_{\leq l} \coloneqq \mathcal{N}(C^n_{\leq l}), F_l \coloneqq \mathcal{N}(C^n_l) \subseteq \operatorname{DcpI}\Delta[n]$. Then we have the following filtration of simplicial sets:

$$\{1\} \times \Delta[n] \cong F_{\leq -1} \subseteq F_{\leq 0} \subseteq F_{\leq 1} \subseteq \cdots \subseteq F_{\leq n} = \operatorname{DepI} \Delta[n].$$

We also need a filtration of the codomain. For notational ease, regard:

$$\operatorname{ESdI}\Delta[n]_k = \operatorname{ESdI}_{\Delta}[n]_k = \coprod_{\substack{I \sqcup J = [k] \\ I < J}} \operatorname{Hom}_{\operatorname{\textbf{Poset}}}(I \star I \star J, [n]),$$

as the set of triplets of the form (I, J, f), where $I \sqcup J = [k]$, I < J is a partition, and $f: I \star I \star J \to [n]$ is an order-preserving function. For each $-1 \leq l \leq n$, we define the following simplicial subsets of ESdI $\Delta[n]$:

$$(G_{\leq l})_k \coloneqq \{ (I, J, f) \mid \max f \mid_{I \star I} \leq l \}; (G_l)_k \coloneqq \{ (I, J, f) \mid \max f \mid_{I \star I} \leq l \leq \min f \mid_J \}.$$

Here, we set $\max f|_{I \star I} = -1$ if $I = \emptyset$, and $\min f|_J = n$ if $J = \emptyset$. Then we obtain the following filtration of simplicial sets:

$$\Delta[n] \cong G_{\leq -1} \subseteq G_{\leq 0} \subseteq G_{\leq 1} \subseteq \dots \subseteq G_{\leq n} = \operatorname{ESdI} \Delta[n].$$

Notice that the canonical map $\operatorname{DcpI}\Delta[n] \to \operatorname{ESdI}\Delta[n]$ restricts to $F_{\leq l} \to G_{\leq l}$ and $F_l \to G_l$ for each $-1 \leq l \leq n$.

Now, we repeat an inductive argument similar to that in the proof of Lemma 8.17. We shall show that $F_{\leq l} \to G_{\leq l}$ is a universal localization for each $-1 \leq l \leq n$, which is enough for the establishment of the lemma. For the base case l = -1, the map $F_{\leq -1} \to G_{\leq -1}$ is the identity map, which is a universal localization.

For an inductive case, let $0 \leq l \leq m$ and suppose that $F_{\leq l-1} \to G_{\leq l-1}$ is a universal localization; we need to show that $F_{\leq l} \to G_{\leq l}$ is a universal localization. We again wish to use [16, Proposition 02MA]. Consider the following commutative cube, where \hookrightarrow denotes inclusions of simplicial subsets, and \to denotes restrictions of the canonical map DcpI $\Delta[n] \to \text{ESdI} \Delta[n]$:



Since the front and the back faces are pushout squares by construction, and the hooked arrows are injective, [16, Proposition 02MA] applies: to show that $F_{\leq l} \rightarrow G_{\leq l}$ is a universal localization, it suffices to check that the three maps $F_{l-1} \cap F_l \rightarrow G_{l-1} \cap G_l$, $F_l \rightarrow G_l$, and $F_{\leq l-1} \rightarrow G_{\leq l-1}$ are universal localizations.

The first map $F_{l-1} \cap F_l \to G_{l-1} \cap G_l$ is isomorphic to:

$$\operatorname{Dcp} \Delta[l-1] \star \Delta[n-l] \to \operatorname{ESd} \Delta[l-1] \star \Delta[n-l].$$

The second map $F_l \to G_l$ is isomorphic to:

$$\operatorname{Dcp}\Delta[l] \star \Delta[n-l] \to \operatorname{ESd}\Delta[l] \star \Delta[n-l]$$

50 FINITE REEDY CATEGORIES AS LOCALIZATIONS OF FINITE DIRECT CATEGORIES

Therefore these two maps are universal localizations by Corollary 8.18. The third map $F_{\leq l-1} \rightarrow G_{\leq l-1}$ is a universal localization by the induction hypothesis; thus we have shown that $F_{\leq l} \rightarrow G_{\leq l}$ is a universal localization. This completes the induction, and we have obtained the desired lemma.

Now, it only remains to prove the main results of this subsection:

Proposition 8.21. Let $X \in \mathbf{Set}_{\Delta}$ be a simplicial set. Then the canonical simplicial map $\operatorname{DcpI} X \to \operatorname{ESdI} X$ in Definition 8.8 is a universal localization in the sense of [16, Definition 02M0].

Proof. By virtue of Lemma 8.20, the proof goes exactly in the same way as that of Proposition 8.19. $\hfill \Box$

Corollary 8.22. Let $f: X \to Y$ be a simplicial map. Consider the following morphism of spans of simplicial sets:



Then the following induced map of pushouts is a universal localization in the sense of [16, Definition 02M0]:

 $(\operatorname{Dcp} Y) \cup_{\operatorname{Dcp} X} (\operatorname{DcpI} X) \to (\operatorname{ESd} Y) \cup_{\operatorname{ESd} X} (\operatorname{ESdI} X).$

Proof. By Propositions 8.19 and 8.21, the claim follows from [16, Proposition 02MA]. \Box

8.4. **Properties of the transformations 2: an inner anodyne map.** The goal of this subsection is the following:

Proposition 8.23. Let $X \in \mathbf{Set}_{\Delta}$ be a simplicial set. Then the canonical simplicial map $\mathrm{ESd} X \to \mathrm{ESd}' X$ in Definition 8.9 is inner anodyne.

For the proof of this proposition, the following lemma is crucial:

Lemma 8.24. Let $[n] \in \Delta$ be a simplex. Consider the following commutative square:



Here, the vertical maps are from Definition 8.9, and the horizontal maps are induced by the canonical inclusions. Then the corresponding simplicial map

 $(\operatorname{ESd}\Delta[n]) \cup_{\operatorname{ESd}\partial\Delta[n]} (\operatorname{ESd}'\partial\Delta[n]) \to \operatorname{ESd}'\Delta[n]$

out of the pushout is inner anodyne.

With this lemma, the proof of Proposition 8.23 is straightforward.

Proof of Proposition 8.23 from Lemma 8.24. Lemma 8.24 says that the canonical natural transformation $\text{ESd} \circ \Delta \Rightarrow \text{ESd}' \circ \Delta \colon \Delta \to \text{Set}_{\Delta}$ is Reedy inner anodyne. Therefore from Lemma 2.33 follows that the map in question, obtained by the left Kan extension of this natural transformation along the Yoneda embedding, is inner anodyne. The rest of this subsection is devoted to the proof of Lemma 8.24. The proof of the lemma is combinatorial and straightforward: since inner horn inclusion $\Lambda_k[n] \hookrightarrow \Delta[n]$ adds two non-degenerate simplices in pair, we just need to add nondegenerate simplices two-by-two in an appropriate order. However, making such pairs (called *horn pairs*) gracefully takes a bit of care and a long argument. To enhance readability, we shall break the proof down into several other lemmas. The concluding proof of Lemma 8.24 is given in page 54, after Lemma 8.32. We shall first introduce some notation and terminology used for the rest of this subsection:

Notation 8.25. Here we list the terminology and notation used in the proof of Lemma 8.24. The notation is used only throughout this subsection.

- We fix a simplex $[n] \in \Delta$.
- We shall write $f: X \hookrightarrow Y$ for the map in focus. Specifically, f is the canonical injective map between:

$$X := (\operatorname{ESd} \Delta[n]) \cup_{\operatorname{ESd} \partial \Delta[n]} (\operatorname{ESd}' \partial \Delta[n]);$$
$$Y := \operatorname{ESd}' \Delta[n].$$

- We define an *outsider* to be a simplex of Y that is not in the image of f. If an outsider is a k-simplex, we shall say that it is a k-outsider.
- We shall denote the set of *non-degenerate* outsiders by \mathcal{O} . The set of non-degenerate k-outsiders is denoted by \mathcal{O}_k .
- If σ is a k-simplex of Y, it is represented by an order-preserving map $[k] \rightarrow [n]^{[1]}$. We shall denote the adjunct of this map by $u_{\sigma} : [k] \times [1] \rightarrow [n]$.
- A horn pair is an ordered pair (σ, τ) of non-degenerate outsiders σ and τ that satisfy the following conditions for some $0 < i < k := \dim \sigma$:
 - (1) $u_{\sigma}(i-1,0) = u_{\sigma}(i,0);$
 - (2) $u_{\sigma}(i-1,1) < u_{\sigma}(i,1) = u_{\sigma}(k,0);$
 - (3) the simplex τ is the *i*-th facet of σ .
- The horn position of a horn pair (σ, τ) is the integer *i* in the definition above. Note that the integer *i* is unique, since we have $u_{\sigma}(i-1,1) < u_{\sigma}(i,1) = u_{\sigma}(k,0)$ and u_{σ} is order-preserving.
- The anticipated horn position of a non-degenerate k-outsider σ is the number of $0 \le j \le k$ such that $u_{\sigma}(j, 1) < u_{\sigma}(k, 0)$.
- We denote the set of horn pairs by \mathcal{H} .
- If (σ, τ) is a horn pair, we shall say that σ is the *(horn) core* of the pair, and τ is the *(horn) periphery* of the pair. We shall also simply say that σ is a *horn core* to mean that, for some τ , the pair (σ, τ) is a horn pair. A *horn periphery* is defined similarly.

In order to prove our desired Lemma 8.24, it suffices to show that non-degenerate outliers are divided into mutually disjoint horn pairs, and well-order these horn pairs appropriately. We shall first see the equivalent conditions for a simplex to be a non-degenerate outsider:

Lemma 8.26. Let σ be any k-simplex in Y. Then we have the following:

- (1) The simplex σ is non-degenerate if and only if, for each $0 \le i < k$, we have $(u_{\sigma}(i,0), u_{\sigma}(i,1)) < (u_{\sigma}(i+1,0), u_{\sigma}(i+1,1))$ with respect to the product order.
- (2) The simplex σ does not belong to the injective image of ESd' $\partial \Delta[n]$ if and only if u_{σ} is surjective.
- (3) The simplex σ does not belong to the injective image of $\operatorname{ESd}\Delta[n]$ if and only if $u_{\sigma}(k,0) > u_{\sigma}(0,1)$.
- (4) The simplex σ is an outsider if and only if $u_{\sigma}(k,0) > u_{\sigma}(0,1)$ and u_{σ} is surjective.

Proof. By direct calculation.

A number of lemmas below are needed to show that non-degenerate outsiders are divided into horn pairs:

Lemma 8.27. Let $\sigma \in \mathcal{O}_k$ be any non-degenerate k-outsider. Let $0 \leq i \leq k+1$ be the anticipated horn position of σ . Then we have the following:

- (1) It holds that $0 < i \le k$.
- (2) If σ belongs to a horn pair $p \in \mathcal{H}$, either as its core or its periphery, then the horn position of p is i.

Proof. (1): Since u_{σ} is order-preserving, we have $i \leq k$. The other inequality i > 0 follows from the assumption that σ is an outsider; see Lemma 8.26 (4).

(2): If σ is the core of p, the claim is immediate from the definition of a horn pair and the order preservation of u_{σ} . Assume that p is of the form (τ, σ) , and let us write i for the horn position of p. Since σ is the i-th face of τ , we have, for each $0 \le j < i$:

$$u_{\sigma}(j,1) = u_{\tau}(j,1) \le u_{\tau}(i-1,1) < u_{\tau}(i,1) = u_{\tau}(k+1,0) = u_{\sigma}(k,0).$$

Here we used i < k + 1. We also have, for each $i \leq j \leq k$:

$$u_{\sigma}(j,1) = u_{\tau}(j+1,1) \ge u_{\tau}(i,1) = u_{\tau}(k+1,0) = u_{\sigma}(k,0).$$

The claim should be clear from these inequalities.

Lemma 8.28. A horn core is not a horn periphery.

Proof. For contradition, suppose that there were to be two horn pairs (σ_0, σ_1) and (σ_1, σ_2) . Let 0 < i < k be the horn position of the second pair, where $k = \dim \sigma_1$. By applying Lemma 8.27 (2) to σ_1 , we see that *i* is also the horn position of the first pair (σ_0, σ_1) . We may now deduce:

$$u_{\sigma_0}(i,0) = u_{\sigma_0}(i-1,0) = u_{\sigma_1}(i-1,0) = u_{\sigma_1}(i,0) = u_{\sigma_0}(i+1,0);$$

$$u_{\sigma_0}(i,1) = u_{\sigma_0}(k+1,0) = u_{\sigma_1}(k,0) = u_{\sigma_1}(i,1) = u_{\sigma_0}(i+1,1).$$

These equations and Lemma 8.26 (1) imply that σ_0 is a degenerate simplex, which contradicts the definition of a horn pair.

Lemma 8.29. Let σ be a non-degenerate k-outsider, and let $0 < i \leq k$ be the anticipated horn position of σ ; see Lemma 8.27 (1). If we have $u_{\sigma}(i-1,0) = u_{\sigma}(i,0)$ and $u_{\sigma}(i,1) = u_{\sigma}(k,0)$, then σ is the core of a unique horn pair.

Proof. Consider the *i*-th face τ of σ . By the definition of a horn position and Lemma 8.27 (2), the uniqueness is clear: if σ is the core of a horn pair, its periphery can only be τ .

It remains to demonstrate that $p = (\sigma, \tau)$ is a horn pair, for which it suffices to show that i < k, that $u_{\sigma}(i-1,1) < u_{\sigma}(i,1)$, and that $\tau \in \mathcal{O}$. Of these three the second is immediate from the definition of i. The first claim i < k follows from the fact that u_{σ} is order-preserving: if we were to have i = k, then the following contradictory inequalities would hold:

$$u_{\sigma}(i-1,1) < u_{\sigma}(i,1) = u_{\sigma}(k,0) = u_{\sigma}(i,0) = u_{\sigma}(i-1,0) \le u_{\sigma}(i-1,1).$$

We shall now check that τ is a non-degenerate outsider. The non-degeneracy of τ is trivial, since τ is a face of the non-degenerate σ and Y is the nerve of a poset. To show that τ is an (k-1)-outsider, we remember Lemma 8.26 (4). Since 0 < i < k, we have the following inequality:

$$u_{\tau}(k-1,0) = u_{\sigma}(k,0) > u_{\sigma}(0,1) = u_{\tau}(0,1).$$

Also, u_{τ} is surjective, since we have:

$$\operatorname{Im} u_{\tau} = \operatorname{Im} \left(\left. u_{\sigma} \right|_{\left([k] \times [1] \right) \setminus \left(\{i\} \times [1] \right)} \right) = \operatorname{Im} u_{\sigma} = [n].$$

Here, the second equality follows from $u_{\sigma}(i-1,0) = u_{\sigma}(i,0)$ and $u_{\sigma}(i,1) = u_{\sigma}(k,0)$. Therefore, τ is a non-degenerate (k-1)-outsider, and the claim has now been proven.

Lemma 8.30. We apply Notation 8.25 here. Let τ be a non-degenerate k-outsider, and let $0 < i \le k$ be the anticipated horn position of τ . Assume that we have either $u_{\tau}(i-1,0) < u_{\tau}(i,0)$ or $u_{\tau}(i,1) > u_{\tau}(k,0)$. Then τ is the periphery of a unique horn pair.

Proof. We begin with the uniqueness. Let σ be any non-degenerate (k+1)-outsider such that $p = (\sigma, \tau)$ is a horn pair. The horn position of (σ, τ) is equal to *i* by Lemma 8.27 (2). Since τ is the *i*-th face of σ , we have:

$$u_{\sigma}(j,l) = u_{\tau}(j,l)$$
 for $0 \le j < i, 0 \le l \le 1;$ (8.4)

$$u_{\sigma}(j,l) = u_{\tau}(j-1,l) \text{ for } i < j \le k+1, 0 \le l \le 1.$$
 (8.5)

By the definition of a horn pair, we also have:

$$u_{\sigma}(i,0) = u_{\sigma}(i-1,0) = u_{\tau}(i-1,0); \qquad (8.6)$$

$$u_{\sigma}(i,1) = u_{\sigma}(k+1,0) = u_{\tau}(k,0). \tag{8.7}$$

Equations (8.4)–(8.7) together uniquely determine the map u_{σ} , and hence the simplex σ . This is the desired uniqueness.

We proceed to the existence of the horn pair. By the definition of the anticipated horn position, we have $u_{\tau}(i,1) \geq u_{\tau}(k,0)$. Therefore, eqs. (8.4)–(8.7) well-define an order-preserving map $u_{\sigma}: [k+1] \times [1] \to [n]$, and hence a (k+1)-simplex σ of Y. It suffices to show that $p = (\sigma, \tau)$ is a horn pair. By the definition of i and σ , we see that $u_{\sigma}(i-1,0) = u_{\sigma}(i,0)$; that $u_{\sigma}(i-1,1) < u_{\sigma}(i,1) = u_{\sigma}(k+1,0)$; and that τ is the *i*-th face of σ .

The only thing left to check is that σ is a non-degenerate outsider. We should be reminded of (1) and (4) from Lemma 8.26. The surjectivity of u_{σ} is immediate from that of u_{τ} . The inequality $u_{\sigma}(0,1) < u_{\sigma}(k+1,0)$ is also clear from the corresponding inequality $u_{\tau}(0,1) < u_{\tau}(k,0)$. Since τ is non-degenerate and we have $u_{\sigma}(i-1,1) < u_{\sigma}(i,1)$, the only non-trivial inequality needed for the nondegeneracy of σ is $(u_{\sigma}(i,0), u_{\sigma}(i,1)) < (u_{\sigma}(i+1,0), u_{\sigma}(i+1,1))$ with respect to the product order. There are two cases to consider: $u_{\tau}(i-1,0) < u_{\tau}(i,0)$ and $u_{\tau}(i,1) > u_{\tau}(k,0)$. In the former case, we compute:

$$u_{\sigma}(i,0) = u_{\tau}(i-1,0) < u_{\tau}(i,0) = u_{\sigma}(i+1,0).$$

In the latter case, we deduce:

$$u_{\sigma}(i,1) = u_{\tau}(k,0) < u_{\tau}(i,1) = u_{\sigma}(i+1,1).$$

Corollary 8.31. Every non-degenerate outsider is contained in a unique horn pair, either as its core or its periphery.

Either way, σ is a non-degenerate outsider, and the claim has been proven.

Proof. This is a direct consequence of Lemmas 8.28–8.30. If σ is a non-degenerate outsider and $0 < i \leq k$ is its anticipated horn position, we only need to perform a case analysis on whether or not we have both $u_{\sigma}(i-1,0) = u_{\sigma}(i,0)$ and $u_{\sigma}(i,1) = u_{\sigma}(k,0)$.

The final lemma before the proof of Lemma 8.24 is the following; we shall define a well-order on the set of horn pairs:

Lemma 8.32. There is an irreflexive well-order \prec on the finite set \mathcal{H} of horn pairs that satisfies the following property (\star) :

- (*) Let σ be a non-degenerate outsider, and τ be a proper face of σ of any dimension. Let p be the horn pair containing σ (see Corollary 8.31). Then exactly one of the following holds:
 - (1) we have $p = (\sigma, \tau)$;
 - (2) the simplex τ is contained in a horn pair $q \in \mathcal{H}$ such that $q \prec p$;
 - (3) the simplex τ is in the image of f.

Proof. As Y is finite, it is automatic that \mathcal{H} is finite. Note that the required property (\star) is preserved under extension of the partial order \prec . Since any order may be linearly extended and any finite linear order is well-ordered, it suffices to define an irreflexive *partial* order \prec on \mathcal{H} that satisfies the property (\star) .

Consider the set $\mathbb{N}^3 = \mathbb{N}$ of ordered triples of natural numbers, and equip this set with the (irreflexive) lexicographic order <, where each coordinate \mathbb{N} is ordered in the usual way. We define a map $\phi: \mathcal{H} \to \mathbb{N}^3$ by sending each horn pair $p = (\sigma, \tau)$ of horn position *i* to the triple (dim σ , u_{σ} (dim σ , 0), *i*). We define the irreflexive partial order \prec on \mathcal{H} by pulling back the lexicographic order < on \mathbb{N}^3 under ϕ .

We shall now show that \prec satisfies the property (*). Let σ be a non-degenerate outsider, and τ be a proper face of σ . We first remind the reader that Y is nerve of a poset, so that τ , being a face of non-degenerate σ , is also non-degenerate. Also, notice that the conditions (1)–(3) in the statement are mutually exclusive; we only need to show that σ and τ enjoy at least one of the three conditions (1)–(3).

If τ is in the image of f, then there is nothing to prove; hence we may assume that τ is an outsider. Let p and q be the horn pairs containing σ and τ , respectively. Write $p = (\sigma_0, \sigma_1)$ and $q = (\tau_0, \tau_1)$. If either $\sigma = \sigma_1$ or $\tau = \tau_0$ holds, then dim $\tau < \dim \sigma$, dim $\sigma_0 = \dim \sigma_1 + 1$, and dim $\tau_0 = \dim \tau_1 + 1$ imply that dim $\tau_0 < \dim \sigma_0$, and hence that $q \prec p$. Therefore, we may assume that $\sigma = \sigma_0$, i.e., σ is the core of p, and that $\tau = \tau_1$, i.e., τ is the periphery of q.

Let $k = \dim \sigma$, and let 0 < i < k be the horn position of p. If $\dim \tau < k - 1$, then we have $\dim \tau_0 < \dim \sigma_0$, and hence $q \prec p$. Therefore, we may assume that $\dim \tau = k - 1$, which gives $\dim \tau_0 = k = \dim \sigma_0$. Say that the τ is the *j*-th facet of σ . Since τ is assumed to be a horn periphery, by Lemmas 8.28 and 8.29, we only have three cases to consider: j = i - 1, j = i, and j = k. If j = i, then we have $p = q = (\sigma, \tau)$, and we are done.

Next, consider the case j = i - 1. In this case, we have:

$$u_{\tau_0}(k,0) = u_{\tau}(k-1,0) = u_{\sigma}(k,0).$$

The horn position of q, equal to the anticipated horn position i - 1 of τ , is smaller than i. Therefore we have $q \prec p$.

Finally, consider the case j = k. In order for τ to be a periphery, $u_{\tau_0}(k,0) = u_{\tau}(k-1,0) = u_{\sigma}(k-1,0)$ must be strictly less than $u_{\sigma}(k,0)$. In this case, we have $q \prec p$ by definition.

We have analyzed all possible cases, and the claim has been proven.

Now we have paired the simplices out of the image of f so that they may be added via inner horn fillings, so we may now proceed to the proof of the lemma in question.

Proof of Lemma 8.24. We employ Notation 8.25 throughout this proof. We shall show that the map $f: X \hookrightarrow Y$ is inner anodyne by decomposing f into the composition of a finite sequence of inner horn fillings. Here, an inner horn filling stands for the pushout along any simplicial map of any inner horn inclusion: $\Lambda_l[k] \hookrightarrow [k]$ with 0 < l < k.

Let \prec be an irreflexive well-order on the finite set \mathcal{H} that has been asserted to exist in Lemma 8.32. Write \mathcal{H} as

$$\mathcal{H} = \{p_0 \prec p_1 \prec \cdots \prec p_{m-1}\}.$$

Write $p_i = (\sigma_i, \tau_i)$. For any $0 \le i \le m$, we define $X_i \subseteq Y$ as the smallest subcomplex of Y that contains the image of f and the simplices σ_j for j < i. Then there is the following filtration:

$$\operatorname{Im} f = X_0 \subseteq X_1 \subseteq \dots \subseteq X_{m-1} \subseteq X_m$$

Notice that $X_m = Y$, because every outsider is a degeneracy of a non-degenerate outsider, and X_m contains all non-degenerate outsiders.

It suffices to show that the inclusion $X_{j-1} \hookrightarrow X_j$ is an inner horn filling for every $0 < j \leq m$. Write $p_{j-1} = (\sigma_{j-1}, \tau_{j-1})$. Let $k = \dim \sigma_{j-1}$, and let *i* be the horn position of p_{j-1} . Then, by the property of the well-order \prec that Lemma 8.32 guarantees, we have the following pushout square:

$$\begin{array}{ccc} \Lambda_i[k] & \longleftrightarrow & \Delta[k] \\ & & & & & \int \sigma_{j-1} \\ X_{j-1} & \longleftrightarrow & X_j \end{array}$$

This is exactly what we wanted to show.

8.5. Properties of the transformations 3: another inner anodyne map. There is another natural transformation that we need to show are pointwise inner anodyne:

Proposition 8.33. Let $X \in \mathbf{Set}_{\Delta}$ be a simplicial set. Consider the following commutative diagram:



Here, the vertical maps are from Definition 8.11, and the horizontal maps are from Definition 8.9. Then the corresponding simplicial map

$$(\operatorname{ESd}' X) \cup_{\operatorname{ESd} X} (\operatorname{ESdI} X) \to \operatorname{ESdI}' X$$

out of the pushout is inner anodyne.

Before providing the proof of this proposition, we state a simple corollary:

Corollary 8.34. Let $f: X \to Y$ be any map of simplicial sets. Consider the following morphism of spans of simplicial sets:

$$\begin{array}{cccc} \operatorname{ESd} Y & \xleftarrow{f} & \operatorname{ESd} X & \longrightarrow & \operatorname{ESdI} X \\ & & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ \operatorname{ESd}' Y & \xleftarrow{f} & & \operatorname{ESd}' X & \longleftrightarrow & \operatorname{ESdI}' X \end{array}$$

Then, the following induced map of pushouts is inner anodyne:

 $(\operatorname{ESd} Y) \cup_{\operatorname{ESd} X} (\operatorname{ESdI} X) \to (\operatorname{ESd}' Y) \cup_{\operatorname{ESd}' X} (\operatorname{ESdI}' X).$

Proof assuming Proposition 8.33. The previous Proposition 8.33 states that the morphism of the spans in question is Reedy inner anodyne, which immediately implies the claim. $\hfill \Box$

The proof of Proposition 8.33 will be conducted similarly to that of Proposition 8.23; it will be simple with the presence of the following lemma:

Lemma 8.35. Let $[n] \in \Delta$ be a simplex. Consider the following commutative cube:



Let X denote the colimit of the cube except $Y \coloneqq \text{ESdI}' \Delta[n]$. Then the canonical simplicial map $X \to Y$ induced by the cube is inner anodyne.

Proof of Proposition 8.33 from Lemma 8.35. By Lemma 2.33, it suffices to prove that the natural transformation

$$(\operatorname{ESd}'\Delta[n]) \cup_{\operatorname{ESd}\Delta[n]} (\operatorname{ESdI}\Delta[n]) \to \operatorname{ESdI}'\Delta[n]$$

in $[n] \in \Delta$ is Reedy inner anodyne, which is proved in Lemma 8.35.

The rest of this section will be devoted to the proof of Lemma 8.35. As with the previous subsection, we shall first establish some preliminary lemmas. The concluding proof will appear in page 60 after Lemma 8.41. We again start with some notations:

Notation 8.36. The notations listed here will be used solely for the proof of Lemma 8.35. This applies to the rest of this subsection.

- We use some notations from Notation 8.25 with varying values of n. Since Notation 8.25 fixes n, we need to rename some terminologies and notations for each case where n = m:
 - A horn pair shall be called a *m*-horn pair. The set of all *m*-horn pairs will be denoted by \mathcal{H}^m .
 - An outsider shall be called a (m, \bullet) -outsider; a k-outsider a (m, k)outsider. The set of all non-degenerate (m, k)-outsiders will be denoted
 by \mathcal{O}_k^m ; the set of all non-degenerate (m, \bullet) -outsiders will be denoted
 by \mathcal{O}^m .
 - The notation $u_{\sigma} \colon [k] \times [1] \to [m]$, where σ is a simplex of ESd' $\Delta[m]$, will be used and does not need to be renamed.
 - The terms *horn position*, *core*, and *periphery* of an *m*-horn pair will be used as is.
 - Other notations and terminologies will not be used; in particular, X, Y, and f are reserved for definitions of this Notation's own.
- Apart from m, which varies for the use of Notation 8.25, we shall fix a simplex $[n] \in \Delta$.
- The map in question, which we aim to show is inner anodyne, will be denoted by $f: X \to Y$.
- A k-simplex σ of $Y \cong \mathcal{N}(\mathrm{ESdI}'_P[n])$ may be considered as a map of posets

 $\sigma \colon [k] \to \operatorname{ESdI}_P'[n] = (\{0\} \times [n]^{[1]}) \cup (\{1\} \times [n]),$

the order of whose codomain is given in Definition 8.6. Let $[k'] \subseteq [k]$ in $\Delta_a \subseteq \mathbf{Poset}$ be the preimage of $\{0\} \times [n]^{[1]}$ under σ ; let $-1 \leq n' \leq n$ be the smallest integer such that $\sigma([k']) \subseteq \{0\} \times [n']^{[1]}$. Then:

- The integer $-1 \le k' \le k$ is called the *switch position* of σ ; the integer $-1 \le n' \le n$ is called the *switch value* of σ .

- The zeroth part of σ is the map

$$\sigma_0: [k'] \to \{0\} \times [n']^{[1]} \cong [n']^{[1]},$$

where the left arrow is the restriction of σ . Note that, if $k' \ge 0$, then $n' \ge 0$ and we may consider σ_0 as a k'-simplex of ESd' $\Delta[n']$.

- The first part of σ is the map

$$\sigma_1: [k] \setminus [k'] \to \{1\} \times \{n', n'+1, \dots, n\} \cong \{n', n'+1, \dots, n\},\$$

where the left arrow is the restriction of σ .

- We say that σ has a non-empty zeroth part if $k' \ge 0$; has an empty zeroth part if k' = -1; has a non-empty first part if k' < k; has an empty first part if k' = k.
- We say that σ has a straight first part if σ_1 is an order embedding, and its image either is the full codomain $\{n', n'+1, \ldots, n\}$ or misses only the smallest number: $\{n'+1, n'+2, \ldots, n\}$.
- An *I-outsider* is a simplex of Y that is not in the image of the map f. A k-*I-outsider* is an *I*-outsider of dimension k.
- The set of all non-degenerate k-I-outsiders will be denoted by $\mathcal{O}_k^{\mathrm{I}}$; that of all non-degenerate I-outsiders by \mathcal{O}^{I} .
- An *I-horn pair* is a pair (σ, τ) of simplices of Y satisfying the followings:
 - The simplex τ is some facet of σ .
 - The simplex σ has a non-empty and straight first part.
 - The switch value n' of σ and τ are equal and non-negative, implying that the two simplices have non-empty zeroth parts.
 - Let σ_0 and τ_0 denote the zeroth parts of σ and τ , respectively. Then (σ_0, τ_0) is an *n'*-horn pair.
- The set of all I-horn pairs will be denoted by \mathcal{H}^{I} .
- If $p = (\sigma, \tau)$ is an I-horn pair, then σ is called the *core* of p, and τ the *periphery* of p.

Except that we separately prove that f is injective, the proof of Lemma 8.35 will be conducted in a similar manner to that of Lemma 8.24: we show that every I-outsider is contained in a unique I-horn pair, and construct a well-order on the set of I-horn pairs to determin the order in which the horns are to be filled.

Lemma 8.37. We invoke Notation 8.36. The map $f: X \to Y$ is injective.

Proof. Consider the commutative cube (8.8) in the statement of Lemma 8.35. The desired injectivity of f follows from the injectivity of all arrows in the cube and the fact that every face of the cube is a pullback square.

Lemma 8.38. Let σ be a k-simplex of Y. Let $\sigma_0: [k'] \to [n']^{[1]}$ and $\sigma_1: [k] \setminus [k'] \to \{n', \ldots, n\}$ be the zeroth and first parts, respectively, of σ . Then we have the following:

- (1) The simplex σ is non-degenerate if and only if both σ_0 and σ_1 are injective.
- (2) Given that σ has a non-empty zeroth part, the zeroth part σ_0 is injective if and only if it is a non-degenerate simplex in ESd' $\Delta[n']$.
- (3) The simplex σ is not in the injective image of ESd' $\Delta[n]$ if and only if it has a non-empty first part.
- (4) The simplex σ is not in the injective image of ESdI $\Delta[n]$ if and only if it has a non-empty zeroth part and the zeroth part σ_0 satisfies $u_{\sigma_0}(0,1) < u_{\sigma_0}(k',0)$.
- (5) The simplex σ is not in the injective image of ESdI' $\partial \Delta[n]$ if and only if the adjunct $u_{\sigma_0}: [k'] \times [1] \rightarrow [n']$ of the zeroth part is surjective and the image of σ_1 includes $\{n'+1, n'+2, \ldots, n\}$.

58 FINITE REEDY CATEGORIES AS LOCALIZATIONS OF FINITE DIRECT CATEGORIES

(6) The simplex σ is a non-degenerate I-outsider if and only if it has a nonempty and straight first part and a non-empty zeroth part σ_0 that is a nondegenerate (n', k')-outsider.

Proof. The first five claims follows from constrution. The last claim is a direct consequence of the first five: see Lemma 8.26. \Box

Lemma 8.39. Let (σ, τ) be an I-horn pair. Let k be the dimension of σ ; n' be the switch value of σ and τ ; $0 \le k' < k$ be the switch position of σ ; σ_i and τ_i be the *i*-th part of σ and τ , respectively, for i = 0, 1. Then we have the following:

- (1) The switch position of τ is $k' 1 \ge 0$.
- (2) The simplex τ is the *i*-th facet of σ for some unique $0 \le i \le k$, which satisfies 0 < i < k'.
- (3) Let q denote the unique order isomorphism

$$q\colon [k-1]\setminus [k'-1]\to [k]\setminus [k']; \quad i\mapsto i+1.$$

Then the following triangle is commutative:

$$\begin{matrix} [k-1] \setminus [k'-1] & \xrightarrow{\sim} & [k] \setminus [k'] \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\ & & & & & & \\$$

- (4) The simplex τ has a non-empty and straight first part.
- (5) The simplices σ and τ are non-degenerate I-outsiders.

Proof. (1): By definition, we have $(\sigma_0, \tau_0) \in \mathcal{H}^{n'}$; therefore, τ_0 must be a facet of σ_0 , which proves the claim.

(2): By inspecting each facet of σ , we see that *i* must be the horn position of (σ_0, τ_0) .

(3): Follows from (2).

(4): Follows from (3) and the fact that σ satisfies the same property.

(5): We apply Lemma 8.38 (6) to σ and τ . Since (σ_0, τ_0) is an n'-horn pair, we see that σ and τ has non-empty zeroth parts, that $\sigma_0 \in \mathcal{O}_{k'}^{n'}$, and that $\tau_0 \in \mathcal{O}_{k'-1}^{n'}$. By (4) and the definition of I-horn pairs, σ and τ has non-empty and straight first parts. The claim follows from these observations.

Lemma 8.40. Every non-degenerate I-outsider belongs to a unique I-horn pair.

Proof. Let $\sigma \in \mathcal{O}_k^{\mathrm{I}}$ be a non-degenerate k-I-outsider. By Lemma 8.38 (6), σ has a non-empty and straight first part and a non-empty zeroth part σ_0 that is a non-degenerate (n', k')-outsider, which implies $k \leq 2$ and 0 < k' < k. Let p be the n'-horn pair containing σ_0 . We have two cases to consider: that where σ_0 is the core of p, and that where σ_0 is the periphery of p.

For the first case, let us assume that $p = (\sigma_0, \tau_0)$ for some τ_0 . Define an orderpreserving map $\tau : [k-1] \to \text{ESdI}'_P[n]$ by:

$$\tau(i) \coloneqq \begin{cases} (0, \tau_0(i)) & \text{if } 0 \le i < k'; \\ (1, \sigma(i+1)) & \text{if } k' \le i < k. \end{cases}$$

We consider τ as a (k-1)-simplex of $\text{ESdI}'_{P}[n]$. Now, we may easily check that (σ, τ) is an I-horn pair containing σ : consult the definitions of I-horn pairs and n'-horn pairs.

For the second case, let us assume that $p = (\tau_0, \sigma_0)$ for some τ_0 . Define an order-preserving map $\tau : [k+1] \to \operatorname{ESdI}_P'[n]$ by:

$$\tau(i) \coloneqq \begin{cases} (0, \tau_0(i)) & \text{if } 0 \le i \le k' + 1; \\ (1, \sigma(i)) & \text{if } k' + 1 < i \le k + 1. \end{cases}$$

We consider τ as a (k + 1)-simplex of $\operatorname{ESdI}_P'[n]$. Now, by the definitions of I-horn pairs and n'-horn pairs, it is easy to see that (τ, σ) is an I-horn pair containing σ .

We have shown that every non-degenerate I-outsider belongs to an I-horn pair. It only remains to show that the I-horn pair is unique. Assume that σ is contained in I-horn pairs p, p'. Let τ and τ' be the other consituent of p and p', respectively. Write $\sigma_i, \tau_i, \tau'_i$ for the *i*-th part of σ, τ, τ' , respectively, for i = 0, 1. By definition, σ_0 and τ_0 , with an appropriate order, form an n'-horn pair p_0 , and σ_0 and τ'_0 form an n'-horn pair p'_0 . By Corollary 8.31, we have either:

$$(\sigma_0, \tau_0) = p_0 = p'_0 = (\sigma_0, \tau'_0), \text{ or}$$

 $(\tau_0, \sigma_0) = p_0 = p'_0 = (\tau'_0, \sigma_0).$

Either way we get $\tau_0 = \tau'_0$. This also implies that σ is either the core of both p and p', or the periphery of the both. Therefore, in order to conclude the proof, it suffices to show that $\tau_1 = \tau'_1$, which is a direct consequence of Lemma 8.39 (3). \Box

Lemma 8.41. We employ Notation 8.36. There is an irreflexive well-order \prec^{I} on the finite set \mathcal{H}^{I} of I-horn pairs that satisfies the following property (\star) :

- (*) Let σ be a non-degenerate I-outsider, and τ be a proper face of σ . Let p be the I-horn pair containing σ (see Lemma 8.40). Then exactly one of the following holds:
 - (1) $p = (\sigma, \tau);$
 - (2) the simplex τ is contained in a horn pair $q \in \mathcal{H}^{\mathrm{I}}$ such that $q \prec^{\mathrm{I}} p$;
 - (3) the simplex τ is in the image of f.

Proof. The finiteness of \mathcal{H}^{I} follows from the finiteness of Y.

In order to define an irreflexive well-order \prec^{I} on \mathcal{H}^{I} , we need some auxiliary orders on other sets. For each $0 \leq m \leq n$, take an irreflexive well-order \prec^m on \mathcal{H}^m guaranteed to exist by Lemma 8.32. Consider the following disjoint union of sets:

$$\mathcal{H}^{ullet} \coloneqq \bigcup_{0 \le m \le n} \mathcal{H}^m = \bigsqcup_{0 \le m \le n} \mathcal{H}^m$$

Equip this with an irreflexive well-order \prec^{\bullet} by considering this union as the join or the ordinal sum $H^0 \star H^1 \star \cdots \star H^n$. That is, if $p \in \mathcal{H}^m$ and $q \in \mathcal{H}^{m'}$, then $p \prec^{\bullet} q$ precisely if either m < m' or m = m' and $p \prec^m q$. Consider the irreflexively well-ordered set $P := \mathbb{N} \times \mathcal{H}^{\bullet}$, where the order is the lexicographic order.

We define a map $\psi: \mathcal{H}^{\mathrm{I}} \to P$ as follows: for each I-horn pair $p = (\sigma, \tau)$ with the zeroth parts of the components forming an n'-horn pair p_0 , we set $\psi(p) :=$ $(\dim \sigma, p_0)$. We define the order \prec^{I} on \mathcal{H}^{I} by $p \prec^{\mathrm{I}} q$ if and only if $\psi(p) \prec^{\bullet} \psi(q)$. By Lemmas 8.38–8.40, We see that ψ is injective; hence \prec^{I} is an irreflexive well-order.

We now show that \prec^{I} satisfies (*). Let σ be a non-degenerate I-outsider, and τ be a proper face of σ . Since Y is nerve of a poset, the simplex τ is non-degenerate. By applying the same argument as in the proof of Lemma 8.32 to Lemmas 8.39 and 8.40, we see that the conditions (1)–(3) are mutually exclusive, and that the only non-trivial case to consider is when σ is the core of some I-horn pair $p = (\sigma, \sigma')$ and τ is a facet of σ and the periphery of another I-horn pair $q = (\tau', \tau)$.

Let k' be the switch position of σ and m be the switch value of σ . Say that τ is the *i*-th facet of σ . By the definition of I-horn pairs and Lemma 8.28, in order that τ is the periphery of q, we must have $0 \le i \le k'$. If the switch value m' of τ

is not equal to m, then we have i = k' and m' < m; therefore, $q \prec^{\mathrm{I}} p$. We consider the case m' = m. Let $p_0 = (\sigma_0, \sigma'_0)$ and $q_0 = (\tau'_0, \tau_0)$ be the *m*-horn pairs formed by the zeroth parts of the components of p and q, respectively. Since τ_0 is a facet of σ_0 , the defining property of \prec^m from Lemma 8.32 applies and we have exactly one of $p_0 = (\sigma_0, \tau_0) = q_0$ or $q_0 \prec^m p_0$. In the former case, we have $p = (\sigma, \tau) = q$; in the latter, we have $q \prec^{\mathrm{I}} p$.

Now we conclude this subsection by proving Lemma 8.35:

Proof of Lemma 8.35. By Lemma 8.37, it suffices to prove that the inclusion of the image of f into Y is inner anodyne. By virtue of Lemmas 8.40 and 8.41, the rest of the proof goes exactly the same as that of Lemma 8.24 in page 54.

8.6. Main lemma for $(\infty, 1)$ -localization. The purpose of this subsection is Corollary 8.44, which is a summarized form of the results of this section. For the sake of conciseness, we begin with a new notation:

Definition 8.42. Let $f: X \to Y$ be any map of simplicial sets. We shall define the simplicial sets DcpC f and ESdC' f (C for "(mapping) cylinder") as those that fills in the following pushout squares:

$$\begin{array}{cccc} \operatorname{Dcp} X & \longrightarrow & \operatorname{DcpI} X & & \operatorname{ESd}' X & \longrightarrow & \operatorname{ESdI}' X \\ \operatorname{Dcp}(f) & & \downarrow & & \operatorname{and} & & \operatorname{ESd}'(f) \\ & & & & & \\ \operatorname{Dcp} Y & \longmapsto & \operatorname{DcpC} f & & & \operatorname{ESd}' Y & \longrightarrow & \operatorname{ESdC}' f \end{array}$$

Note that there is a canonical map $\operatorname{DcpC} f \to \operatorname{ESdC'} f$. We shall write $W_f^C \subseteq (\operatorname{DcpC} f)_1$ for the set of edges that are mapped to degenerate edges in $\operatorname{ESdC'} f$.

Proposition 8.43. Let $f: X \to Y$ be any map of simplicial sets. Then the canonical map DcpC $f \to \text{ESdC}' f$ exhibits ESdC' f as the localization of DcpC f at W_f^C .

Proof. Follows from Corollaries 8.22 and 8.34; note that an edge in ESdC' f is degenerate precisely if it is the image of a degenerate edge in ESd $Y \cup_{\text{ESd } X} \text{ESdI } X$, for the map between these two simplicial sets is a monomorphism and an isomorphism on the set of vertices. See Remark 01MX and Proposition 02M1 in [16] for more details.

Corollary 8.44. Let $u: X \to Y$ be any map of simplicial sets, and $A \subseteq B$ be an inclusion of simplicial sets. Set $K := B \times \text{DcpC} u$ and $L := B \times \text{ESdC}' u$. Consider the following unions of simplicial subsets:

$$\begin{aligned} K' &:= (A \times \operatorname{DcpC} u) \cup (B \times (X \times \{1\})) \subseteq K, \\ L' &:= (A \times \operatorname{ESdC}' u) \cup (B \times (X \times \{1\})) \subseteq L. \end{aligned}$$

Let $Q \in \mathbf{Set}_{\Delta}$ be a quasi-category, and assume that there is a following commutative square of simplicial sets:

$$\begin{array}{cccc}
K' & \longrightarrow & L' \\
& & \downarrow g \\
K & & \longrightarrow & Q
\end{array}$$
(8.9)

Here, unlabeled edges represent canonical maps (see Definition 8.42 for $K' \to L'$), and the labeled morphisms f and g are arbitrary. Suppose that f sends the edges in the following subset to equivalences in Q:

$$W_K \coloneqq \left\{ (\mathrm{id}_b, e) \mid b \in B_0, e \in W_u^C \right\} \subseteq B_1 \times (\mathrm{DcpC}\, u)_1 = K_1$$

Then there exists an extension $h: L \to Q$ of g, such that the composite $K \to L \xrightarrow{n} Q$ is naturally equivalent to f relatively to K'.

Proof. By Proposition 8.43 and some simple computation, we see that the canonical $K \to L$ exhibits L as the localization of K at W_K . Moreover, we claim that its restriction $K' \to L'$ also exhibits L' as the localization of K' at:

$$W_{K'} \coloneqq W_K \cap K' = (A \times W_u^C) \cup \{ \mathrm{id}_p \mid p \in (B \times (X \times \{1\}))_0 \}.$$

To see the claim, note that the subcomplexes K' and L' are the following pushouts, and that $K' \to L'$ is induced by the obvious transformation of the pushout-defining spans:

$$K' = (A \times \operatorname{DcpC} u) \cup_{A \times (X \times \{1\})} (B \times (X \times \{1\})),$$

$$L' = (A \times \operatorname{ESdC}' u) \cup_{A \times (X \times \{1\})} (B \times (X \times \{1\})).$$

Each of the three consituent maps of the transformation, say $U \to V$, exhibits V as the localization of U at $W_K \cap U$. Since the two arrows in each of the two spans are inclusions, we derive from [16, Proposition 01N7] that the localization property is inherited by the pushout, as we have claimed.

Apply Fun(•, Q) to the canonical commutative square (8.10) below to get another one. The latter square, by the definition of localizations, admits the following factorization (8.11) through subcomplexes, where the arrows with ~ are equivalences of quasi-categories:

From $L' \subseteq L$, $K' \subseteq K$, and $W_{K'} \subseteq W_K$, we see that the vertical maps in (8.11) are isofibrations of quasi-categories. To detail the derivation, for the left and the right vertical arrows, you may use [16, Corollary 01F3], $L' \subseteq L$, and $K' \subseteq K$. Next, we derive from [16, Remark 01MN] that $\operatorname{Fun}((K', W_{K'}), Q^{\natural}) \hookrightarrow \operatorname{Fun}(K', Q)$ is an isofibration of quasi-categories. Now note that the outer rectangle in the following commutative diagram is a pullback:

Since the lower horizontal composite is an isofibration of quasi-categories, [16, Corollary 01H4] implies that $\operatorname{Fun}((K, W_K), Q^{\natural}) \to \operatorname{Fun}((K', W_{K'}), Q^{\natural})$ is an isofibration of quasi-categories.

We have assumed in (8.9) that $g \in \operatorname{Fun}(L', Q)$ goes to $f|_{K'} \in \operatorname{Fun}((K', W_{K'}), Q^{\natural})$ in the diagram (8.11). Let us write F_g for the fiber of $\operatorname{Fun}(L, Q) \to \operatorname{Fun}(L', Q)$ over g, and F_f for the fiber of $\operatorname{Fun}((K, W_K), Q^{\natural}) \to \operatorname{Fun}((K', W_{K'}), Q^{\natural})$ over $f|_{K'}$. It suffices to show that the restriction $F_g \to F_f$ of the middle vertical map in (8.11) is an equivalence of quasi-categories, for the map h we seek is a vertex of F_g that maps to an object equivalent to f in F_f . Consider the following 1-categorical pullback diagrams in \mathbf{Set}_{Δ} that define F_g and F_f :

$$\begin{array}{cccc} F_g & & & & & F_f & & & F_m((K, W_K), Q^{\natural}) \\ & & & \downarrow & & & \downarrow & & \downarrow \\ \Delta[0] & & & & & \Delta[0] & & & \Delta[0] & & & Fun((K', W_{K'}), Q^{\natural}) \end{array}$$

In these squares, the vertices are all quasi-categories, and as we have seen, the right vertical arrows are both isofibrations of quasi-categories. Therefore these pullbacks are in fact homotopy pullback squares with respect to the Joyal model structure on \mathbf{Set}_{Δ} , which are equivariant under categorical equivalences, which shows our sufficient claim for our corollary.

9. The proof of $(\infty, 1)$ -localization

Now, we are finally ready to show that $\operatorname{last}_{\Gamma}$ is a localization map, where Γ is one of the four categories $\int \mathbf{N}(C)$, $\int \mathbf{N}^{-,+}(C)$, $\operatorname{Down}_*(C)$, and $\operatorname{Down}(C)$. We begin with the easiest case, $\Gamma = \int \mathbf{N}(C)$.

Proposition 9.1. The simplicial map $N(\mathfrak{last}): N(\int \mathbf{N}(C)) \to N(C)$ exhibits the domain $N(\int \mathbf{N}(C))$ as the localization of N(C) at \mathfrak{last} -weak equivalences.

Proof. This follows from the proof of Lemma 7.7, which shows that $last: \int \mathbf{N}(C) \rightarrow C$ is a reflective localization.

We now treat the remaining cases, which are of our main interest.

Notation 9.2. In the following discussion, we set S = N(C), where C is the Reedy category that was fixed throughout this paper, and $T = N(\Gamma)$, where Γ is any of the three categories $\int \mathbf{N}^{-,+}(C)$, $\text{Down}_*(C)$, and Down(C). For the purpose of aesthetics, We shall abbreviate as $\lambda = \text{last}$ the functor $N(\text{last}): T \to S$ of quasicategories corresponding to $\text{last}: \Gamma \to C$ from Definition 7.1.

We begin with defining the $(\infty, 1)$ -diagrams used in the proof of our (infty, 1)-localization theorem.

Lemma 9.3. There are simplicial maps $D: \operatorname{Dcp} S \to T, D^I: \operatorname{Dcp} T \to T, E: \operatorname{ESd}' S \to S, and E^I: \operatorname{ESdI}' T \to S$ that satisfy the following properties:

(1) The following diagram commutes:



- (2) The maps D^I and E are retractions of the canonical injections.
- (3) We have the following commutative square:



- (4) The map D: Dcp S → T sends the inverse image of degenerate edges under Dcp S → ESd S to last-weak equivalences in T.
- (5) Let $e \in (\text{DcpI} T)_1$ be an edge. If the image of e under $\text{DcpI} T \twoheadrightarrow \text{ESdI} T$ is either degenerate or in the subcomplex $\text{ESdI}(\text{Sk}_0 T) \subseteq \text{ESdI} T$, then the map D^I sends e to a last-weak equivalence in T.

Proof. Let $\Phi: C^{[1]} \to C^{[2]}$ denote the functorial factorization that corresponds to the Reedy factorization of C. That is: Φ is a strict section of the composition functor $(\delta_1^2)^*: C^{[2]} \to C^{[1]}$; for each $u \in \operatorname{Ob} C^{[1]} = \operatorname{Mor} C$, we have $(\delta_2^2)^*(\Phi(u)) \in C_-$ and $(\delta_0^2)^*(\Phi(u)) \in C_+$. Write $Q := (\iota_1)^* \circ \Phi: C^{[1]} \to C$ for the "midpoint" functor of the factorization Φ . The natural transformation that corresponds to $(\delta_0^2)^* \circ \Phi: C^{[1]} \to C^{[1]}$ shall be denoted by $\eta: Q \Rightarrow \operatorname{cod} = (\iota_2^2)^*: C^{[1]} \to C$: the "second arrow." If $(f,g): u \to v$ is a morphism in $C^{[1]}$, then Q and η fit into the following commutative diagram:



Note that if $f \in C_-$, then $Q(f,g) \in C_-$, and if $g \in C_+$, then $Q(f,g) \in C_+$.

We shall first construct E. With the following definitions in mind:

$$\operatorname{ESd}' S = \operatorname{colim}_{\Delta[n] \to S} \operatorname{N}(\operatorname{ESd}'_{P}[n]), \quad S = \operatorname{N}(C),$$

we see that it suffices to compatibly define E_{φ} : $\mathrm{ESd}'_{P}[n] \to C$ for each *n*-simplex $\varphi: [n] \to C$ of $S = \mathrm{N}(C)$. Therefore we set E_{φ} as the composite of:

$$\operatorname{ESd}_P'[n] = [n]^{[1]} \xrightarrow{\varphi_*} C^{[1]} \xrightarrow{Q} C,$$

which is compatible with the structure maps of the colimit-defining diagram for $\operatorname{ESd}' S$. Combining, we obtain a simplicial map $E \colon \operatorname{ESd}' S \to S$.

The construction of E^I is similar; for each *n*-simplex $\varphi \colon [n] \to T$ of $T = \mathcal{N}(\Gamma)$, we set E^I_{φ} as the composite of:

$$\mathrm{ESdI}'_{P}[n] \, \hookrightarrow \, [1] \times [n]^{[1]} \twoheadrightarrow [n]^{[1]} \xrightarrow{\varphi_{*}} \Gamma^{[1]} \xrightarrow{\mathfrak{last}_{*}} C^{[1]} \xrightarrow{Q} C.$$

Here, the first inclusion is mentioned in the definition of ESdI'_P itself in Definition 8.6. The construction is natural in φ , so bundles together to yield a simplicial map E^I : ESdI' $T \to S$.

The construction of D requires a little harder work. Remember $T = \mathcal{N}(\Gamma)$; we may focus on the case $\Gamma = \int \mathbf{N}^{-,+}(C)$, for the cases $\Gamma = \text{Down}_*(C)$ and $\Gamma = \text{Down}(C)$ are treated through the post-composition of the quotient functor $\int \mathbf{N}^{-,+}(C) \to \text{Down}_*(C)$ and the quasi-inverse of the equivalence $\text{Down}(C) \hookrightarrow$ $\text{Down}_*(C)$.

Let $\varphi \colon [n] \to C$ be an *n*-simplex of $S = \mathcal{N}(C)$. Similarly to above, we wish to compatibly define a functor $D_{\varphi} \colon \mathrm{Dcp}_{C}[n] \to \Gamma$. Let $(x, \alpha \colon [m] \to [n]) \in \mathrm{Ob}(\mathrm{Dcp}_{C}[n])$. We set $D_{\varphi}(x, \alpha) \coloneqq ([m], Q_{(x,\alpha)}^{\varphi})$, where $Q_{(x,\alpha)}^{\varphi} \colon [m] \to C_{-}$ is the following functor:

$$[m] \ni i \longmapsto Q(\varphi(!_{x,\alpha(i)})) \in \operatorname{Ob} C;$$
$$(!_{i,j}: i \leq j) \longmapsto Q(\operatorname{id}_{\varphi(x)}, \varphi(!_{\alpha(i),\alpha(j)})).$$

Let the following be an arbitrary morphism in $Dcp_C[n]$:

$$(!_{x,x'},\beta)\colon (x,\alpha\colon [m]\to [n])\to (x',\alpha'\colon [m']\to [n]).$$

Remember, since $\beta : \alpha \to \alpha'$ is a morphism in $\Delta/[n]$, it is a morphism $\beta : [m] \to [m']$ in Δ . We shall set:

$$D_{\varphi}(!_{x,x'},\beta) \coloneqq (\beta, \theta^{\varphi x x' \alpha}) \colon ([m], Q_{(x,\alpha)}^{\varphi}) \to ([m'], Q_{(x',\alpha')}^{\varphi}).$$

Here, the natural transformation

$$Q^{\varphi xx'\alpha} \colon Q^{\varphi}_{(x,\alpha)} \Rightarrow Q^{\varphi}_{(x',\alpha)} = Q^{\varphi}_{(x',\alpha')} \circ \beta \colon [m] \to C$$

is defined by, for each $i \in [m]$:

$$\theta_i^{\varphi x x' \alpha} \coloneqq Q(\varphi(!_{x,x'}), \mathrm{id}_{\varphi(\alpha(i))}) \colon Q(\varphi(!_{x,\alpha(i)})) \to Q(\varphi(!_{x',\alpha(i)})) = Q(\varphi(!_{x',\alpha'(\beta(i))})),$$

which belongs to C_+ by the property of Q. The naturality of $\theta^{\varphi xx'\alpha}$ follows from the structure of the domain $C^{[1]}$ of Q and the functoriality of Q; thus we have declared the mapping D_{φ} on objects and morphisms. The functoriality of D_{φ} follows from that of Q and φ , so we obtain a simplicial map $N(D_{\varphi})$: $\text{Dcp }\Delta[n] \to T$. Since the naturality of $N(D_{\varphi})$ in φ may be straightforwardly proven using the definition of the functor $\text{Dcp}_C \colon \Delta \to \mathbf{Cat}$, we obtain a map $D \colon \text{Dcp } S \to T$ of simplicial sets.

We finally construct D^I : DcpI $T \to T$. We begin with the case $\Gamma = \int \mathbf{N}^{-,+}(C)$. Given an *n*-simplex $\varphi \colon [n] \to \Gamma$ of $T = \mathcal{N}(\Gamma)$, we need to compatibly define a functor D_{φ}^I : DcpI_C $[n] \to \Gamma$. In this construction of D_{φ}^I , we will use the following symbols:

$$([m_x], X_x) \coloneqq \varphi(x) \in \operatorname{Ob} \Gamma \qquad \text{for } x \in [n]; \\ (\beta_{xy}, \theta^{xy}) \coloneqq \varphi(!_{xy}) \colon \varphi(x) \to \varphi(y) \quad \text{for } x \leq y.$$

First remember that $\text{DcpI}_C[n]$ has $\{0\} \times (\text{Dcp}_C[n])$ and $\{1\} \times [n]$ as disjoint full subcategories. We set:

$$\begin{aligned} D_{\varphi}^{I} \Big|_{\{0\} \times (\mathrm{Dcp}_{C}[n])} &\coloneqq D_{\lambda(\varphi)} = D_{\mathfrak{lasto}\varphi}, \\ D_{\varphi}^{I} \Big|_{\{1\} \times [n]} &\coloneqq \varphi. \end{aligned}$$

Since every object of $\text{DcpI}_C[n]$ belongs to one of these two full subcategories, it only remains to define how D_{φ}^I acts on morphisms that connect objects in the distinct subcategories.

By the definition of $\text{DcpI}_C[n]$, we only need to consider the unique morphism $!_{(0,(x,\alpha)),(1,y)}: (0,(x,\alpha)) \to (1,y)$, where $(x,\alpha:[m] \to [n]) \in \text{Ob}(\text{Dcp}_C[n])$ and $y \in [n]$ are objects satisfying $\alpha(m) \leq y$. We wish to construct

$$D_{\varphi}^{I}(!_{(0,(x,\alpha)),(1,y)}) = (\beta,\theta) \colon ([m], Q_{(x,\alpha)}^{\lambda(\varphi)}) \to \varphi(y) = ([m_y], X_y).$$

We would like to define $\beta \colon [m] \to [m_y]$ by, for each $i \in [m]$:

$$\beta(i) \coloneqq \beta_{\alpha(i),y}(m_{\alpha(i)}) \in [m_y]$$

This is indeed a morphism in Δ , as we have, for each pair $i \leq j$ in [m]:

$$\beta(i) = \beta_{\alpha(i),y}(m_{\alpha(i)}) = \beta_{\alpha(j),y}(\beta_{\alpha(i),\alpha(j)}(m_{\alpha(i)})) \le \beta_{\alpha(j),y}(m_{\alpha(j)}) = \beta(j).$$

We next have to define a natural transformation $\theta: Q_{(x,\alpha)}^{\lambda(\varphi)} \Rightarrow X_y \circ \beta$. Let $i \in [m]$. We have the following morphisms in C_+ :

$$\begin{split} \eta_{\mathrm{last}(\varphi(\mathbf{I}_{x,\alpha(i)}))} \colon Q_{(x,\alpha)}^{\lambda(\varphi)}(i) &= Q(\mathrm{last}(\varphi(\mathbf{I}_{x,\alpha(i)}))) \rightarrowtail \mathrm{last}(\varphi(\alpha(i))), \\ \theta_{m_{\alpha(i)}}^{\alpha(i)y} \colon \ \mathrm{last}(\varphi(\alpha(i))) &= X_{\alpha(i)}(m_{\alpha(i)}) \rightarrowtail X_y(\beta_{\alpha(i)y}(m_{\alpha(i)})) = X_y(\beta(i)) \end{split}$$

We set θ_i as the composite of these two morphisms, which are both natural in *i*.

Now we have defined D_{φ}^{I} on objects and morphisms, and it follows from the naturality of η and the functoriality of φ that D_{φ}^{I} is a functor. The naturality in

 φ of the simplicial map $\mathcal{N}(D_{\varphi}^{I})$: DcpI $\Delta[n] \to T$ is straightforward to prove, so we obtain a simplicial map D^{I} : DcpI $T \to T$.

Now we need to consider the cases $\Gamma = \text{Down}_*(C)$ and $\Gamma = \text{Down}(C)$, but we may obtain D^I from the following commutative diagram:

$$\begin{array}{ccc} \operatorname{DcpI}\operatorname{N}(\int \mathbf{N}^{-,+}(C)) & \longrightarrow & \operatorname{DcpI}\operatorname{N}(\operatorname{Down}_*(C)) & \longleftrightarrow & \operatorname{DcpI}\operatorname{N}(\operatorname{Down}(C)) \\ & & & & \downarrow^{\exists^{1}}D_{I} & & \downarrow^{\exists^{1}}D_{I} \\ & & & & & \bigwedge(\int \mathbf{N}^{-,+}(C)) & \longrightarrow & \operatorname{N}(\operatorname{Down}_*(C)) & \longrightarrow & \operatorname{N}(\operatorname{Down}(C)) \end{array}$$

Here, the horizontal arrows are the canonical ones: two in the left are induced by the quotient functor $\int \mathbf{N}^{-,+}(C) \rightarrow \text{Down}_*(C)$, the other two in the right comes from the equivalence $\text{Down}(C) \hookrightarrow \text{Down}_*(C)$ of categories in Lemma 6.2.

The required properties are now easy to verify, and we have obtained the desired lemma. $\hfill \Box$

We can now prove our theorem in the form of a lifting property, as follows:

Lemma 9.4. Let $A \subseteq B$ be a simplicial set and a simplicial subset, and let Q be a quasi-category. Assume that we have the following (strictly) commutative square of simplicial sets, where f and g are arbitrary:

$$\begin{array}{ccc} A \times T & \xrightarrow{\operatorname{id}_A \times \lambda = \operatorname{id}_A \times \mathfrak{last}} & A \times S \\ & & & \downarrow g \\ B \times T & & & \downarrow g \\ & & & & f \end{array}$$

Suppose that f takes any edge of the form $(id_b, t) \in B_1 \times T_1 = (B \times T)_1$, with $id_b \in B_1$ a degenerate edge and $t \in T_1$ a last-weak equivalence, to an equivalence in Q. Then, there exists a strict extension $h: B \times S \to Q$ of g such that its precomposition $h \circ (id_B \times \lambda): B \times T \to Q$ and f are naturally equivalent relative to $A \times T$.

Proof. Take D, D^I , E, and E^I as in Lemma 9.3. The universality of pushouts and Lemma 9.3 (1) allow us construct \tilde{D} : DcpC $\lambda \to T$ and \tilde{E} : ESdC' $\lambda \to S$ from these four maps. In terms of \tilde{D} and \tilde{E} , Lemma 8.13 and the properties (1)–(3) from Lemma 9.3 can be wrapped up in the following commutative diagram:



We see that the following maps \tilde{f} and \tilde{g} are well-defined and satisfy the constraint of Corollary 8.44, using our assumption, the diagram (9.1), and the properties (4) and (5) from Lemma 9.3:

$$\begin{split} \tilde{f} &: B \times \operatorname{DcpC} \lambda \xrightarrow{\operatorname{id}_B \times D} B \times T \xrightarrow{f} Q; \\ \hat{g} &: A \times \operatorname{ESdC}' \lambda \xrightarrow{\operatorname{id}_A \times \tilde{E}} A \times S \xrightarrow{g} Q; \\ \tilde{g} &:= \hat{g} \cup f : (A \times \operatorname{ESdC}' \lambda) \cup (B \times (T \times \{1\})) \to Q. \end{split}$$

. . *~*

Therefore, the map \tilde{g} can be extended to $h: B \times \text{ESdC}' \lambda \to Q$ whose precomposition with the canonical $B \times \text{DcpC} \lambda \to B \times \text{ESdC}' \lambda$ is homotopic to \tilde{f} relative to $A \times \text{DcpC} \lambda$.

Let us define $h: B \times S \to Q$ as the following composite:

$$B \times (S \times \{0\}) \hookrightarrow B \times ((S \times \{0\}) \cup_{T \times \{0\}} (T \times \Delta[1])) \hookrightarrow B \times \text{ESdC'} \lambda \xrightarrow{h} Q.$$

If we write $u \subseteq v$ to mean a simplicial map u is a restriction of v, we have:

$$g\subseteq \hat{g}\subseteq \tilde{g}\subseteq h\supseteq h,$$

where the first \subseteq is due to the commutativity of the rightmost triangle in the diagram (9.1). Since dom $g = A \times S \times \{0\} \subseteq B \times S \times \{0\} = \text{dom } h$, we see $g \subseteq h$. Now, define $H: (B \times T) \times \Delta[1] \to Q$ as the following composite:

$$B \times (T \times \Delta[1]) \longrightarrow B \times ((S \times \{0\}) \cup_{T \times \{0\}} (T \times \Delta[1])) \hookrightarrow B \times \text{ESdC}' \lambda \xrightarrow{h} Q.$$

By definition, we see $H|_{(B \times T) \times \{0\}} = h \circ (\mathrm{id}_B \times \lambda)$. We also have $H|_{(B \times T) \times \{1\}} = f$, because $f \subseteq \tilde{g} \subseteq \tilde{h} \supseteq H|_{(B \times T) \times \{1\}}$. As $\hat{g} \subseteq \tilde{g} \subseteq \tilde{h}$, the map H restricts to the $A \times T \times \Delta[1] \to Q$ in the following commutative diagram:

$$\begin{array}{c|c} A \times (T \times \Delta[1]) \longrightarrow A \times ((S \times \{0\}) \cup_{T \times \{0\}} (T \times \Delta[1])) \hookrightarrow A \times \operatorname{ESdC'} \lambda \xrightarrow{g} Q \\ & & & & \\ \operatorname{proj} & & & & \\ & & & & \\ A \times T \xrightarrow{\qquad} A \times (S \cup_T T) \xrightarrow{\qquad} A \times S \xrightarrow{g} Q \end{array}$$

Here we have used the commutativity of the rightmost triangle in the diagram (9.1) and the definition of \hat{g} . Thus we see that H is a simplicial homotopy, or a natural transformation, from $h \circ (\mathrm{id}_B \times \lambda)$ to f relative to $A \times T$.

It only remains to demonstrate that the natural transformation H is a natural equivalence. We wish to prove, for each vertex $p \in (B \times T)_0$, that the edge

$$e_p \coloneqq p \times \mathrm{id}_{\Delta[1]} \colon \Delta[1] = \Delta[0] \times \Delta[1] \to (B \times T) \times \Delta[1]$$

in $(B \times T) \times \Delta[1]$ is sent by H to an equivalence in Q. If we write $\lambda_0: \operatorname{Sk}_0 T \to \operatorname{Sk}_0 S$ for the restriction of **last**, then the image of e_p in $B \times \operatorname{ESdC'} \lambda$ lies within the subcomplex $B \times \operatorname{ESdC'} \lambda_0 \subseteq B \times \operatorname{ESdC'} \lambda$. By simple inspection we see that the canoncial map from $\operatorname{DcpC} \lambda_0$ surjects onto $\operatorname{ESdC'} \lambda_0$; therefore it reduces to show that the following composite sends every edge in the domain to an equivalence in Q:

$$\operatorname{Sk}_0 B \times \operatorname{DcpC} \lambda_0 \longrightarrow B \times \operatorname{DcpC} \lambda \longrightarrow B \times \operatorname{ESdC}' \lambda \xrightarrow{h} Q.$$

The restriction \tilde{f}_0 of \tilde{f} to $\operatorname{Sk}_0 B \times \operatorname{DcpC} \lambda_0$ is naturally equivalent to the composite above, by the construction of \tilde{h} . By Lemma 9.3 (5) and our assumption on f, we see that \tilde{f}_0 sends every edge to an equivalence in Q; hence the same holds for the composite above, as desired.

Now, our theorem is now simple to prove.

Theorem 9.5. The functor $\lambda = \text{last}: T \to S$ exhibits S as the localization of T at the last-weak equivalences.

Proof. Follows from the previous Lemma 9.4; see [16, Proposition 01MS]. \Box

REFERENCES

10. Conclusion: proof of the main theorem

In this short section, we shall summarize the results of this paper into the theorem stated in Section 1.

Theorem 10.1 (Restatement of Theorem 1.1). Let C be a small Reedy category. Then there is a concrete construction of a direct category Down(C), a set $W^{\text{last}} \subseteq$ Mor Down(C) of morphisms, and a functor last: $\text{Down}(C) \to C$ that exhibits C as the localization of Down(C) at W^{last} . Furthermore, if C is finite, then Down(C)is also finite.

Proof. The category Down(C) is constructed in Definition 3.8; the functor **last** is constructed in Definition 7.1; and the set W^{last} is constructed in Definition 7.2. The directness of Down(C) is proven in Proposition 5.7. It is shown that (Down(C), last) is a 1-localization of C at W^{last} in Theorem 7.3 constructively; and it is demonstrated that the pair is $(\infty, 1)$ -categorical localization in Theorem 9.5. Finally, the finiteness of Down(C) is proven in Corollary 5.9.

References

- Peter Aczel. "The Type Theoretic Interpretation of Constructive Set Theory". In: Logic Colloquium '77. Ed. by A. MacIntyre, L. Pacholski, and J. Paris. North-Holland, 1978, pp. 55–66.
- [2] Steve Awodey. Category Theory. 2nd. USA: Oxford University Press, Inc., 2010. ISBN: 0-19-923718-2.
- C. Barwick and D. M. Kan. A Thomason-like Quillen equivalence between quasi-categories and relative categories. 2011. arXiv: 1101.0772 [math.AT]. URL: https://arxiv.org/abs/1101.0772.
- C. Barwick and D. M. Kan. Relative categories: Another model for the homotopy theory of homotopy theories. 2011. arXiv: 1011.1691 [math.AT]. URL: https://arxiv.org/abs/1011.1691.
- Julia Bergner and Charles Rezk. "Reedy categories and the Θ-construction". In: Mathematische Zeitschrift 274 (Oct. 2011), pp. 499–514. DOI: 10.1007/ s00209-012-1082-0.
- [6] Francis Borceux. Handbook of Categorical Algebra. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 1994.
- [7] P. Gabriel and M. Zisman. Calculus of Fractions and Homotopy Theory. Calculus of Fractions and Homotopy Theory. Springer-Verlag, 1967. ISBN: 978-0-387-03777-6.
- P.S. Hirschhorn. Model Categories and Their Localizations. Mathematical surveys and monographs. American Mathematical Society, 2003. ISBN: 978-0-8218-4917-0.
- [9] M. Hovey. *Model Categories*. Mathematical surveys and monographs. American Mathematical Society, 2007. ISBN: 978-0-8218-4361-1.
- [10] Maxime Ramzi. Does join of simplicial sets preserve (∞, 1)-categorical localizations? MathOverflow answer. Dec. 27, 2024. URL: https://mathoverflo w.net/q/484893 (visited on 02/06/2025).
- [11] Peter LeFanu Lumsdaine. What is the name for the construction of this poset related to coherence of degeneracies of the simplex category? MathOverflow answer. June 4, 2022. URL: https://mathoverflow.net/q/423963 (visited on 02/06/2025).
- [12] Mike Shulman. In the constructive theory of direct categories, is it decidable whether an arbitrary morphism is an identity or not? MathOverflow answer. Sept. 18, 2023. URL: https://mathoverflow.net/q/454848 (visited on 02/06/2025).

REFERENCES

- [13] Nicolai Kraus and Christian Sattler. Space-Valued Diagrams, Type-Theoretically (Extended Abstract). 2017. arXiv: 1704.04543 [math.LO]. URL: https: //arxiv.org/abs/1704.04543.
- [14] Robert S. Lubarsky. "IKP and Friends". In: The Journal of Symbolic Logic 67.4 (2002), pp. 1295-1322. ISSN: 00224812. URL: http://www.jstor.org/ stable/3648574 (visited on 02/06/2025).
- [15] J. Lurie. *Higher Topos Theory*. Academic Search Complete. Princeton University Press, 2009. ISBN: 978-0-691-14048-3.
- [16] Jacob Lurie. *Kerodon*. 2024. URL: https://kerodon.net.
- [17] Saunders MacLane. Categories for the Working Mathematician. Graduate Texts in Mathematics, Vol. 5. New York: Springer-Verlag, 1971, pp. ix+262.
- [18] Emily Riehl. *Categorical Homotopy Theory*. New Mathematical Monographs. Cambridge University Press, 2014.
- [19] Michael Shulman. Reedy categories and their generalizations. 2015. arXiv: 1507.01065 [math.AT]. URL: https://arxiv.org/abs/1507.01065.
- [20] Michael Shulman. "Univalence for inverse diagrams and homotopy canonicity". In: *Mathematical Structures in Computer Science* 25.5 (Nov. 2014), pp. 1203–1277. ISSN: 1469-8072. DOI: 10.1017/s0960129514000565.
- [21] Michael A. Shulman. Set theory for category theory. 2008. arXiv: 0810.1279 [math.CT]. URL: https://arxiv.org/abs/0810.1279.
- [22] The Univalent Foundations Program. Homotopy Type Theory: Univalent Foundations of Mathematics. Institute for Advanced Study, 2013. URL: https: //homotopytypetheory.org/book.

68