On representations of the crystallization of the quantized function algebra $C(SU_q(n+1))$

Saikat Das and Ayan Dey Indian Statistical Institute, Delhi, India

Abstract

The crystallization $C(SU_0(n+1))$ of the q-family of C^* -algebras $C(SU_q(n+1))$ was defined by Giri & Pal for all $n \geq 2$ and they observed that any irreducible representation at the q=0 level is the norm limit of irreducible representations of $C(SU_q(n+1))$ as $q \to 0^+$. In this article, we will generalize this to any non-degenerate representation (respectively faithful) of $C(SU_0(n+1))$. As a consequence, one can alternatively realize $C(SU_0(n+1))$ as the C^* -algebra generated by the limit operators of faithful representations of $C(SU_q(n+1))$.

AMS Subject Classification No.: 20G42, 46L67, 58B32

Keywords. Quantum groups, q-deformation, quantized function algebras, representations.

1 Introduction

The crystallization of the C^* -algebra $C(K_q)$ was introduced by Giri & Pal in [4] for type A_n compact Lie groups K = SU(n+1) and subsequently by Matassa & Yuncken in [8] for all simple simply connected compact Lie groups K. These two definitions of crystallization are slightly different but it is expected that the two crystallizations give the same C^* -algebra. However, we will only work with the definition of crystallization given in [4] where the authors had observed that for every irreducible representation of $C(K_q)$, limits of the canonical generating elements, scaled appropriately, obey the same set of relations as $q \to 0^+$ and used those relations to define the crystallized algebra. Throughout this article, we will denote $C(SU_q(n+1))$ by $A_q(n)$ for any $q \in [0, 1)$. Furthermore, to make the notation more convenient, we may sometimes omit n and write A_q for $A_q(n)$. In [5] Giri & Pal studied irreducible representations of the crystallized algebra $A_0(n)$ in great detail. In particular, they found all the irreducible representations of the crystallized algebra and proved that every irreducible representation of $A_0(n)$ is the norm limit of irreducible representations of $A_q(n)$ with appropriately scaled generators. The goal of this article is to extend this result for any arbitrary non-degenerate faithful representation of $A_0(n)$ and, conversely, any non-degenerate faithful representation of $A_q(n)$ gives rise to a non-degenerate faithful representation of $A_0(n)$ in a natural way (see Theorem 4.15).

Now we will say a few words about the content of this paper. In section 2, the representation theory of $A_0(n)$ is recalled very briefly, and some of the important results of [5] are listed. Section 3 is devoted to understanding the topology and Borel structure of the spectrum of $A_q(n)$. The crucial observation is that the spectrums \hat{A}_0 and \hat{A}_q are Borel isomorphic via a specific map and a necessary condition for factorization of irreducibles is described in terms of Bruhut ordering of S_{n+1} . The initial part of section 4 is all about decomposing any arbitrary non-degenerate representation of $A_q(n)$ into multiple copies of a direct integral of irreducibles using a result from [3] and, finally we then come to the main results of this paper that any non-degenerate representation (respectively faithful) of $A_0(n)$ on a separable Hilbert space \mathcal{H} is the norm limit of non-degenerate representations (respectively faithful) of $A_q(n)$ on the same Hilbert space \mathcal{H} with appropriately scaled generators, and the converse. The two crucial steps of the proof of faithfulness for different values of $q \in [0,1)$ are identifying the minimal faithful part of a faithful representation and capturing faithfulness in terms of support of the underlying positive Borel measure of direct integrals. As a consequence, we will then be able to realize the crystallized algebra $A_0(n)$ as the C^* -algebra generated by the limit operators of faithful representations of $A_q(n)$ with suitably scaled generators (see Corollary 4.16).

2 Irreducible representations of $A_0(n)$

For any $q \in (0,1)$, $A_q(n)$ is a well-studied universal C^* -algebra with $(n+1)^2$ many generators $z_{i,j}(q)$ satisfying certain relations (see [4]). Given two representations π_1 and π_2 of $A_q(n)$, their convolution product is defined by the formula $\pi_1 * \pi_2 := (\pi_1 \otimes \pi_2) \circ \Delta$ where $\Delta : A_q \to A_q \otimes A_q$ is the comultiplication map on A_q associated to the quantum group structure of A_q . \mathcal{S}_{n+1} denotes the permutation group on n+1 letters and the collection $\{s_i = (i,i+1) : 1 \leq i \leq n\}$ of Coxeter generators generates the group \mathcal{S}_{n+1} . We use the symbol S for the left shift/backward shift operator on $l^2(\mathbb{N})$ and, q^N is the operator on $l^2(\mathbb{N})$ that acts on the standard basis $\{e_n\}_{n\geq 0}$ by, $e_n \to q^n e_n$ for all $n \in \mathbb{N} \cup \{0\}$. Then for any $s_r \in \mathcal{S}_{n+1}$ one can define,

$$\pi_{s_r}^{(q)}(z_{i,j}(q)) := \begin{cases} S\sqrt{1 - q^{2N}} & i = j = r \\ \sqrt{1 - q^{2N}}S^* & i = j = r + 1 \\ -q^{N+1} & i = r, j = r + 1 \\ q^N & i = r + 1, j = r \\ \delta_{i,j}I & \text{otherwise.} \end{cases}$$

For $a, b \in \mathbb{N}$ with $a \leq b$, let us denote by $s_{[a,b]}$ the product $s_b s_{b-1} \cdots s_a$ in \mathcal{S}_{n+1} . Let $1 \leq k \leq n$, $1 \leq b_k \leq b_{k-1} \leq \cdots \leq b_1 \leq n$, and $1 \leq a_i \leq b_i$ for $1 \leq i \leq k$. It follows from the strong exchange condition and the deletion condition in the characterization of the Coxeter system (see [1]) that $w = s_{[a_k,b_k]} s_{[a_{k-1},b_{k-1}]} \cdots s_{[a_1,b_1]}$ is a reduced word in \mathcal{S}_{n+1} . For any reduced word w in this form we define,

$$\pi_w^{(q)} := \pi_{s_{[a_k,b_k]}}^{(q)} * \pi_{s_{[a_{k-1},b_{k-1}]}}^{(q)} * \dots * \pi_{s_{[a_1,b_1]}}^{(q)}.$$

On the other hand, for any $t \in (S^1)^n$ we have all one-dimensional irreducibles of $A_q(n)$ defined as follows,

$$\chi_t^{(q)}(z_{i,j}(q)) := \begin{cases} t_1 \delta_{i,j} & i = 1\\ \overline{t_n} \delta_{i,j} & i = n+1\\ \overline{t_{i-1}} t_i \delta_{i,j} & \text{otherwise} \end{cases}$$

Theorem 2.1 ([7]). Any irreducible representation of $A_q(n)$ is of the form $\pi_{t,w}^{(q)} := \chi_t^{(q)} * \pi_w^{(q)}$ for some $t \in (S^1)^n$ and reduced word $w \in \mathcal{S}_{n+1}$. Furthermore, $\pi_{t,w}^{(q)}$'s are all mutually inequivalent.

The definition of $A_0(n)$ is given in [4] with generators $\{z_{k,j}(0): 1 \leq k, j \leq n+1\}$ satisfying certain relations. Giri & Pal observed that the following limits exist in the norm,

$$\pi_{t,w}^{(0)}(z_{i,j}(0)) := \begin{cases} \lim_{q \to 0^+} \pi_{t,w}^{(q)}(z_{i,j}(q)) & \text{if } i \ge j \\ \lim_{q \to 0^+} (-q)^{i-j} \pi_{t,w}^{(q)}(z_{i,j}(q)) & \text{if } i < j \end{cases}$$

for $t \in (S^1)^n$ and any reduced word $w \in \mathcal{S}_{n+1}$ in the above form. In particular, for $s_r \in \mathcal{S}_{n+1}$ we have

$$\pi_{s_r}^{(0)}(z_{i,j}(0)) := \begin{cases} S & i = j = r \\ S^* & i = j = r + 1 \\ P_0 & i = r, j = r + 1 \\ P_0 & i = r + 1, j = r \\ \delta_{i,j}I & \text{otherwise} \end{cases}$$

where $P_0 = I - S^*S$ is the orthogonal projection onto the subspace of $l^2(\mathbb{N})$ generated by e_0 .

Remark 2.2. Recall that even though the elements $z_{i,j}(q)$ belong to different C^* -algebras $A_q(n)$ for different values of q, they are images of the generating elements $z_{i,j}(t)$ of the space $\mathcal{O}_t(SU(n+1))$ under the specialization maps θ_q (see Section 2, [4] for details; there $z_{i,j}(q)$ and $z_{i,j}(t)$ are denoted by $u_{i,j}(q)$ and $u_{i,j}(t)$ respectively).

Theorem 2.3 ([5]). 1. $\pi_{t,w}^{(0)}$'s are all mutually inequivalent irreducible representations of the crystallized algebra $A_0(n)$.

2. $A_0(n)$ is a separable C^* -algebra of type I i.e. the image of any irreducible representation on a Hilbert space \mathcal{H} contains $K(\mathcal{H})$.

Altogether we have the following important and desirable property of the crystallized algebra.

Theorem 2.4 ([5]). Denote by $z_{k,j}(q)$ the generators of $A_q(n)$ for $q \in [0,1)$. Let $\pi^{(0)}$ be an irreducible representation of $A_0(n)$ on a seperable Hilbert space \mathcal{H} . Then there exists irreducible representations $\pi^{(q)}$ of A_q on the same Hilbert space \mathcal{H} such that

$$\pi^{(0)}(z_{k,j}(0)) = \lim_{q \to 0+} \pi^{(q)} \left((-q)^{\min\{k-j,0\}} z_{k,j}(q) \right), \qquad k,j \in \{1,2,\dots,n+1\}.$$

Although the authors have found all the irreducible representations of A_0 in [5], they did not prove that the definition of these representations is independent of the choice of the reduced form taken for a reduced word $w \in \mathcal{S}_{n+1}$. To prove this we now make use of the Coxeter structure of \mathcal{S}_{n+1} , in particular the Bruhat ordering \leq on \mathcal{S}_{n+1} . For further details on Bruhat ordering, we refer to [1]. In the proof of the next proposition, we will write $\pi_{n,w}^{(q)}$ for the representation $\pi_w^{(q)}$ of $A_n(q)$ for avoiding notational confusion and these notations won't be used anywhere else.

Proposition 2.5. For any two reduced expressions of w_1 and w_2 of a permutation w, $\pi_{w_1}^{(0)}$ and $\pi_{w_2}^{(0)}$ are unitarily equivalent. In particular, the collection of irreducible representations of $A_0(n)$ is indexed by the elements of $(S^1)^n \times S_{n+1}$.

Proof. As observed in Theorem 3.3.1 of [1], given two reduced forms of an element of S_{n+1} one can reach from one reduced form to another by performing a finite sequence of braid moves i.e. $s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}$ and $s_i s_j = s_j s_i$ for $|i-j| \geq 2, 1 \leq i, j \leq n$. Keeping this in mind we proceed to prove the proposition first for the algebra $A_0(2)$. Define a *-automorphism ϕ of $A_0(2)$ by sending $z_{i,j}(0) \to z_{4-i,4-j}^*(0)$ for $1 \leq i, j \leq 3$. Then one can directly verify $\pi_{2,s_1s_2s_1}^{(0)} = \pi_{2,s_2s_1s_2}^{(0)} \circ \phi$. Therefore, $\pi_{2,s_2s_1s_2}^{(0)}$ is irreducible and hence using Theorem 2.3, $\pi_{2,s_2s_1s_2}^{(0)}$ is equivalent to $\pi_{2,s_1s_2s_1}^{(0)}$. Let U be an intertwining unitary between these two representations, that is $U^*\pi_{2,s_2s_1s_2}^{(0)}U = \pi_{2,s_1s_2s_1}^{(0)}$. The proof for $A_0(n)$ is done in two steps. For $1 \leq i, j \leq n$ with $|i-j| \geq 2$, $\pi_{n,s_is_j}^{(0)} \cong \pi_{n,s_js_i}^{(0)}$ via the flip operator on $l^2(\mathbb{N})^{\otimes 2}$. For the other braid move i.e. $s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}$, observe that one can write $\pi_{n,s_is_{i+1}s_i}^{(0)} = \pi_{2,s_1s_2s_1}^{(0)} \circ \phi_1$ and $\pi_{n,s_{i+1}s_is_{i+1}}^{(0)} = \pi_{2,s_2s_1s_2}^{(0)} \circ \phi_1$ where $\phi_1 : A_0(n) \to A_0(2)$ is a *-homomorphism that sends $z_{k,l} \to z_{k-i+1,l-i+1}$ if $k,l \in \{i,i+1,i+2\}$ and $\delta_{k,l}I$ on other generators. Therefore, the same unitary U intertwines $\pi_{n,s_is_{i+1}s_i}^{(0)}$ and $\pi_{n,s_{i+1}s_is_{i+1}}^{(0)}$ and $\pi_{n,s_{i+1}s_is_{i+1}}^{(0)}$ and $\pi_{n,s_{i+1}s_is_{i+1}}^{(0)}$ is an $\pi_{n,s_is_{i+1}s_i}^{(0)}$ and $\pi_{n,s_is_{i+1}s_i}^{(0)}$ and $\pi_{n,s_is_{i+1}s_i}^{(0)}$ if I is an I denoted by the same unitary I intertwines I intertwines I and I intertwines I intertwines

For $q \neq 0$, the above result is already known but observe that the proof given above does not work for this case.

So, from now on, we will freely use the notation w for any reduced word irrespective of the choice of reduced form.

3 Topology and Borel structure on \hat{A}_0

Let A be a separable C^* -algebra and J be an ideal of A. J is called primitive if J is the kernel of some nonzero irreducible representation of A. Let Prim(A) be the collection of all primitive ideals of A. We will work with the Jacobson topology on Prim(A) whose closed sets can be characterized as follows. $S \subseteq Prim(A)$ is closed if and only if there is a subset M of A such that S is exactly the collection of all primitive ideals containing M. Let \hat{A} be the collection of all equivalence classes of non-zero irreducible representations of A. The topology τ on \hat{A} is then

defined as pullback of the Jacobson topology via the canonical map from $\hat{A} \to Prim(A)$. This topology can be very pathological because it is not even Hausdorff in general (see Corollary 3.5). However, if we take A to be a separable type I C^* -algebra then $(\hat{A}, \mathcal{B}_{\tau})$ is a standard Borel space (see 4.6.1, [3]) where \mathcal{B}_{τ} is the Borel sigma-algebra generated by pullback of the Jacobson topology on \hat{A} . Furthermore, the canonical map $\hat{A} \to Prim(A)$ becomes a homeomorphism (see 3.1.6, 9.1, [3]).

For any $q \in [0,1)$ we will denote the collection of equivalence classes of all one-dimensional representations and the collection of all equivalence classes of infinite-dimensional irreducible representations on $l^2(\mathbb{N})$ by $\hat{A}_{q,1}$ and $\hat{A}_{q,\infty}$ respectively. Let $w \neq 1$ be any reduced word. Denote $\hat{A}_{q,w} = \{[V_w^* \pi_{t,w}^{(q)}(.)V_w] : t \in (S^1)^n\} \subseteq \hat{A}_q$ where $V_w : l^2(\mathbb{N}) \to l^2(\mathbb{N})^{\otimes l(w)}$ is a fixed unitary and l(w) stands for the length of the reduced word. Also, note that since we are considering the class of irreducibles, the choice of unitary V_w does not matter. Then using Theorems 2.1 and 2.3 one can write

$$\hat{A}_q = \hat{A}_{q,1} \bigsqcup \hat{A}_{q,\infty} = \hat{A}_{q,1} \bigsqcup \left(\bigsqcup_{w \neq 1} \hat{A}_{q,w} \right).$$

One of the goals of this section is to show that the above decomposition is into Borel subsets. Hence, we have the following decomposition

$$\int_{\hat{A}_q}^{\bigoplus} F \cong \int_{\hat{A}_{q,1}}^{\bigoplus} F \bigoplus \left(\bigoplus_{w \neq 1} \int_{\hat{A}_{q,w}}^{\bigoplus} F \right).$$

We will use this decomposition with a suitable integrand F later. Let $w \neq 1$ be a fixed reduced word in S_{n+1} . We write $Irr_{\infty}(A_q)$ for the collection of all non-zero infinite-dimensional irreducible representations of A_q on $l^2(\mathbb{N})$ and the set $Irr_{V_w,w}(A_q) = \{V_w^*\pi_{t,w}^{(q)}(.)V_w : t \in (S^1)^n\} \subseteq Irr_{\infty}(A_q)$.

Lemma 1. For $q \in [0,1)$ and a reduced word $w \neq 1$ we have

- 1. $Irr_{\infty}(A_q)$ with the topology of strong pointwise convergence is a polish space.
- 2. The natural map $\rho: Irr_{\infty}(A_q) \to \hat{A}_{q,\infty}(\subset \hat{A}_q)$ is open, continuous and onto. Hence, it is a Borel map from a standard space $Irr_{\infty}(A_q)$ to a standard space \hat{A}_q .
- 3. $\rho(Irr_{V_w,w}(A_q)) = \hat{A}_{q,w}$ and ρ restricted to $Irr_{V_w,w}(A_q)$ is injective.

Proof. 1 and 2 are obvious from results 3.7.4 and 3.5.8 in [3]. The equality $\rho(Irr_{V_w,w}(A_q)) = \hat{A}_{q,w}$ follows immediately once we realise $l^2(\mathbb{N})$ and $l^2(\mathbb{N})^{\otimes l(w)}$ are isomorphic Hilbert spaces and injectivity of the restriction of ρ is a consequence of Theorems 2.1 and 2.3.

Lemma 2. Let $w \neq 1$ be a fixed reduced word. Then the map $f: Irr_{V_w,w}(A_q) \to (S^1)^n$ defined by

$$V_w^* \pi_{t,w}^{(q)}(.) V_w \to t.$$

is a homeomorphism.

Proof. Surjectivity is obvious and injectivity again follows from Theorems 2.1 and 2.3. Now we prove f is continuous. Let $\{V_w^*\pi_{t_\alpha,w}^{(q)}(.)V_w\}_\alpha$ be a net in $Irr_{V_w,w}(A_q)$ converging to $V_w^*\pi_{t,w}^{(q)}(.)V_w$ under the topology of strong pointwise convergence. That means for the generator $z_{1,1}(q) \in A_q$ we have

$$|t_{\alpha}^{1} - t^{1}| \|V_{w}^{*}\pi_{w}^{(q)}(z_{1,1}(q))V_{w}(\eta)\| \to 0$$

where $\eta \in l^2(\mathbb{N})$. The above convergence forces us to conclude $t^1_{\alpha} \to t^1$ in S^1 . Similarly, using other diagonal generators $z_{i,i}(q)$, one can show $t^i_{\alpha} \to t^i$ for all $i = 2, 3 \cdots n$. To prove the continuity of the inverse let us take a net $t_{\alpha} \to t$ in $(S^1)^n$. Then the same proof given above shows that

$$\left\| V_w^* \pi_{t_\alpha, w}^{(q)}(a) V_w(\eta) - V_w^* \pi_{t, w}^{(q)}(a) V_w(\eta) \right\| \to 0$$

for $a=z_{k,l}(q)$ or $z_{k,l}^*(q)$ and $\eta\in l^2(\mathbb{N})$. Hence, the above convergence is also true for any $a\in A_q(n)$.

Therefore, $Irr_{V_w,w}(A_q)$ is a compact subset of a polish space $Irr_{\infty}(A_q)$. So, $Irr_{V_w,w}(A_q)$ is itself closed and hence a Borel subset of $Irr_{\infty}(A_q)$.

Proposition 3.1. $\hat{A}_{q,w}$ is a Borel subset of \hat{A}_q and restriction of ρ on $Irr_{V_w,w}(A_q)$ is a Borel isomorphism between $Irr_{V_w,w}(A_q)$ and $\hat{A}_{q,w}$. Furthermore, $\hat{A}_{q,w}$ is Borel isomorphic to $(S^1)^n$ for all $q \in [0,1)$.

Proof. It follows immediately from Corollary 15.2 of [6] or Appendix B21 of [3].

Using the previous results, now we can safely write that for $q \in (0,1)$, the map $T_q^{-1}: \hat{A}_{q,\infty} \to \hat{A}_{0,\infty}$ defined by, $[V_w^* \pi_{t,w}^{(q)}(.) V_w] \to [V_w^* \pi_{t,w}^{(0)}(.) V_w]$ is a Borel isomorphism. It further extends to a Borel isomorphism between \hat{A}_q and \hat{A}_0 naturally. We record this observation in the next theorem.

Theorem 3.2. For any $q \in (0,1)$ the map $T_q : \hat{A}_0 \to \hat{A}_q$ is a Borel isomorphism

Remark 3.3. It is important to note that \hat{A}_0 and \hat{A}_q are always Borel isomorphic as both of them are standard Borel spaces with the same cardinality. But that does not imply the map $T_q^{-1}:\hat{A}_q\to\hat{A}_0$ is a Borel isomorphism. We need this specific map T_q^{-1} to be our required isomorphism for the main result. Although \hat{A}_q is Borel isomorphic to $\bigsqcup_{(n+1)!}(S^1)^n$ for any $q\in[0,1)$, they can never be homeomorphic as pullback of the Jacobson topology on \hat{A}_q is not Hausdorff (see Corollary 3.5).

Theorem 3.4. Let u and w be two permutations in S_{n+1} with $u \leq w$. Then $\pi_{t,u}^{(q)}$ factorizes through $\pi_{t,w}^{(q)}$ for any $t \in (S^1)^n$ and $q \in [0,1)$.

Proof. We will denote the Toeplitz algebra by \mathscr{T} . Now using the subword property of the Bruhat ordering on S_{n+1} (Theorem 2.2.2, [1]) one can obtain a reduced expression for u by deleting some Coxeter generators in a reduced word expression for w. Therefore, one can define a *-homomorphism $ev: \mathscr{T}^{l(w)} \to \mathscr{T}^{l(u)}$ by sending $S \to 1$ for the deleted components in the reduced expression for w and all other remaining components $S \to S$. Now, together with Proposition 2.5, it is easy to see that $ev \circ \pi_{t,w}^{(q)} = \pi_{t,u}^{(q)}$.

We will call the factorization map defined in the proof of the last theorem factorization by evaluations. One should also note that $\pi_{t,u}^{(q)}$ factorizes through $\pi_{t,w}^{(q)}$ if and only if $Ker \pi_{t,w}^{(q)} \subseteq Ker \pi_{t,u}^{(q)}$ for any $t \in (S^1)^n$.

Corollary 3.5. For any $q \in [0,1)$, the topology on \hat{A}_q is not Hausdorff.

Proof. Choose a $t \in (S^1)^n$ and a reduced word $w_2 \in \mathcal{S}_{n+1}$. If w_1 is a Bruhat subword of w_2 , then there is a C^* -homomorphism $\phi: \mathcal{F}^{l(w_2)} \to \mathcal{F}^{l(w_1)}$ such that $\phi \circ \pi_{t,w_2}^{(q)} = \pi_{t,w_1}^{(q)}$. That means $Ker \pi_{t,\omega_2}^{(q)} \subseteq Ker \pi_{t,\omega_1}^{(q)}$. Therefore, from the description of closed sets of the Jacobson topology on $Prim(A_q)$ it follows that the set $\{Ker \pi_{t,\omega_2}^{(q)}\}$ is not closed. Hence, its pullback $\{[V_{w_2}^* \pi_{t,\omega_2}^{(q)}(.)V_{w_2}]\} \subseteq \hat{A}_q$ is not closed.

We will use the notation $w_L = s_1(s_2s_1)(s_3s_2s_1)\cdots(s_ns_{n-1}\cdots s_2s_1)$ for the longest word in S_{n+1} . Any reduced word w is a Bruhat subword of w_L .

Corollary 3.6. For all $q \in [0,1)$, \hat{A}_{q,w_L} is dense in \hat{A}_q .

Proof. Since any reduced word is a Bruhat subword of the longest word w_L , $\bigcap_t Ker \pi_{t,w_L}^{(q)} = \{0\}$. This follows from the fact that no non-zero element can lie in the kernel of every irreducible representation of a C^* -algebra. Therefore, any element $[\pi] \in \hat{A}_q$ is weakly contained in \hat{A}_{q,w_L} that is $\bigcap_t Ker \pi_{t,w_L}^{(q)} \subseteq Ker \pi$. Now, the density of \hat{A}_{q,w_L} follows from Theorem 3.4.10 of [3]. \square

4 Main results

Our goal of this section is to extend Theorem 2.4 for any non-degenerate (respectively faithful) representations of $A_0(n)$ and describe $A_0(n)$ as the C^* -algebra generated by the limit operators of faithful representations of $A_q(n)$. We need a decomposition theorem for representations involving disintegration to prove these results, which we will discuss first. We refer to chapter 8 of [3] for more details. Let Z be a Borel space and \mathcal{H}_n be the standard n-dimensional Hilbert space. By A we denote a separable C^* -algebra of type I. \hat{A} stands for its spectrum and \hat{A}_n stands for the collection of equivalence classes of irreducible representations of A on \mathcal{H}_n .

Definition 4.1. A constant field Γ_n of Hilbert spaces $\tau \to \mathcal{H}(\tau)$ defined by \mathcal{H}_n is a field with the following properties,

- $\mathcal{H}(\tau) = \mathcal{H}_n$ for all $\tau \in \mathbb{Z}$.
- $\Gamma_n = \{ \tau \to x(\tau) : \tau \to \langle x(\tau) | a \rangle \text{ is Borel for all } a \in \mathcal{H}_n \}.$

Then for every positive Borel measure μ on Z one has $\Gamma_n \subseteq \Gamma_n^{\mu}$ where

$$\Gamma_n^{\mu} = \{ \tau \to x(\tau) : \tau \to \langle x(\tau) | a \rangle \text{ is } \mu\text{-measurable for all } a \in \mathcal{H}_n \}.$$

Let n be a cardinal and $n \leq \aleph_0$. There is a unique Borel field of Hilbert spaces $\Gamma_c : \tau \to \mathcal{H}(\tau)$ on \hat{A} such that Γ reduces on \hat{A}_n to the constant field defined by \mathcal{H}_n . For any positive Borel measure μ on \hat{A} , Γ_c is endowed with a μ -measurable field structure that is $\Gamma_c \subseteq \Gamma_c^{\mu}$ where

$$\Gamma_c^{\mu} = \{ \tau \to x(\tau) : (\tau \to x(\tau))|_{\hat{A}_n} \in \Gamma_n^{\mu} \}.$$

We will frequently talk about representations of A on the Hilbert space $\int_{\hat{A}}^{\bigoplus} \mathcal{H}(\tau)d\mu$ defined by the field $((\mathcal{H}(\tau))_{\tau \in \hat{A}}, \Gamma_c^{\mu})$.

Theorem 4.2 (8.6.2, [3]). 1. There is a field $\tau \to \psi(\tau)$ of irreducible representations of A (defined on \hat{A}) in the Hilbert spaces $\mathcal{H}(\tau)$ such that $\psi(\tau) \in \tau$ and this field is μ -measurable for any positive measure μ on \hat{A} .

2. If there is another field of irreducible representations $\tau \to \psi_1(\tau)$ with the same properties as above, then for every positive Borel measure μ on \hat{A} , we have

$$\int_{\hat{A}}^{\bigoplus} \psi(\tau) d\mu \cong \int_{\hat{A}}^{\bigoplus} \psi_1(\tau) d\mu.$$

This equivalence class of representations is denoted by $\int_{\hat{A}}^{\bigoplus} \tau d\mu$.

Theorem 4.3 (8.6.6, [3]). Let A be a separable type I C^* -algebra and $\pi: A \to B(\mathcal{H})$ be a non-degenerate representation of A on a separable Hilbert space \mathcal{H} . Then there are mutually singular positive Borel measures $\mu_1, \mu_2, ..., \mu_{\infty}$ on \hat{A} such that

$$\pi \cong \int_{\hat{A}}^{\bigoplus} \tau d\mu_1(\tau) \bigoplus 2 \int_{\hat{A}}^{\bigoplus} \tau d\mu_2(\tau) \bigoplus \cdots \bigoplus \aleph_0 \int_{\hat{A}}^{\bigoplus} \tau d\mu_\infty(\tau).$$

The field of irreducible representations that appeared in part 1 of Theorem 4.2 is completely existential. Therefore we will directly construct a field of irreducible representations $\tau_q \to \psi_q(\tau_q)$ of A_q which satisfies all the properties of part 1 of Theorem 4.2 and hence by part 2 of Theorem 4.2 we can conclude that

$$\int_{\hat{A}_q}^{\bigoplus} \tau_q d\mu(\tau_q) = \left[\int_{\hat{A}_q}^{\bigoplus} \psi_q(\tau_q) d\mu(\tau_q) \right].$$

Let $\mathcal{H}_1, \mathcal{H}_2, \dots \mathcal{H}_{\infty}$ be standard Hilbert spaces of dimension $1, 2, \dots \aleph_0$. For each cardinal $n \leq \aleph_0$, the collection of all representations of A on \mathcal{H}_n is denoted by $Rep_n(A)$. The Borel structure on $Rep_n(A)$ is defined by the topology of strong pointwise convergence. One can equip $Rep(A) = \bigcup_{n \leq \aleph_0} Rep_n(A)$ with the sum Borel structure and Irr(A) is then a Borel subset of Rep(A) because for each n, $Irr_n(A)$ is a Borel subset of $Rep_n(A)$ (3.8, [3]).

Let (Z, \mathcal{B}, μ) be a Borel space and $\tau \to \mathcal{H}(\tau)$ be a μ -measurable field of Hilbert spaces defined on Z.

Proposition 4.4 (8.1.8, [3]). Suppose that Z is the union of mutually disjoint Borel sets $Z_1, Z_2, ... Z_{\infty}$ and that $\tau \to \mathcal{H}(\tau)$ reduces on Z_n to the constant field $\tau \to \mathcal{H}_n$. Let $\tau \to \pi(\tau)$ be a field of representations of A on the Hilbert spaces $\mathcal{H}(\tau)$. Then $\tau \to \pi(\tau)$ is μ -measurable if and only if $\tau \to \pi(\tau)$ is equal μ -almost everywhere to a Borel mapping of Z into Rep(A).

For each $q \in [0,1)$, A_q is a separable C^* -algebra of type I (see [2], [5]). Define a field of irreducible representations $\tau_q \to \psi_q(\tau_q)$ on \hat{A}_q as follows,

$$\psi_q(\tau_q) = \begin{cases} V_w^* \pi_{t,w}^{(q)}(.) V_w & \tau_q = [V_w^* \pi_{t,w}^{(q)}(.) V_w] \\ \chi_t^{(q)} & \tau_q = [\chi_t^{(q)}] \end{cases}$$

Corollary 4.5. Let $q \in [0, 1)$. Then $\tau_q \to \psi_q(\tau_q)$ is a μ -measurable field of representations for any positive Borel measure μ on \hat{A}_q .

Proof. Restrict $\tau_q \to \psi_q(\tau_q)$ to $\hat{A}_{q,w}$. Then by Proposition 3.1 this restricted map is a Borel map from $\hat{A}_{q,w}$ into $Irr_{V_w,w}(A_q) \subseteq Irr(A_q) \subseteq Rep(A_q)$). Since $\hat{A}_{q,1}$ and $\hat{A}_{q,w}$ for $w \neq 1$ are all disjoint Borel subsets of \hat{A}_q , it is easy to show that $\tau_q \to \psi_q(\tau_q)$ is a Borel mapping from \hat{A}_q into $Rep(A_q)$. Hence, the result follows if we take $Z = \hat{A}_q, Z_1 = \hat{A}_{q,1}, Z_\infty = \hat{A}_{q,\infty}$ and the empty set to be the null set appearing in Proposition 4.4.

Remark 4.6. Therefore, from part 2 of Theorem 4.2, it follows that

$$\int_{\hat{A}_q}^{\bigoplus} \tau_q d\mu(\tau_q) = \left[\int_{\hat{A}_q}^{\bigoplus} \psi_q(\tau_q) d\mu(\tau_q) \right].$$

for any positive Borel measure μ on \hat{A}_q and for all $q \in [0,1)$. The key observation here is that the unitary V_w 's used to define this field of irreducibles is independent of $q \in [0,1)$ and plays a crucial role in proving the main result.

Now we will extend Theorem 2.4 for any non-degenerate representation of the crystallized algebra.

Theorem 4.7. Let $\pi^{(0)}$ be a non-degenerate representation of A_0 on a seperable Hilbert space \mathcal{H} . Then there exist non-degenerate representations $\pi^{(q)}$ of A_q on the same Hilbert space \mathcal{H} such that

$$\pi^{(0)}(z_{k,j}(0)) = \lim_{q \to 0+} \pi^{(q)} \left((-q)^{\min\{k-j,0\}} z_{k,j}(q) \right), \qquad k,j \in \{1,2,\dots,n+1\}.$$

Proof. Let $\mu_1, \mu_2, ... \mu_{\infty}$ be mutually singular positive measures on \hat{A}_0 corresponding to the following decomposition of $\pi^{(0)}$ (Theorem 4.3)

$$\int_{\hat{A}_0}^{\bigoplus} \psi_0(\tau_0) d\mu_1 \bigoplus 2 \int_{\hat{A}_0}^{\bigoplus} \psi_0(\tau_0) d\mu_2 \bigoplus \cdots \bigoplus \aleph_0 \int_{\hat{A}_0}^{\bigoplus} \psi_0(\tau_0) d\mu_{\infty}.$$

For $q \in (0,1)$, define a representation $\pi^{(q)}$ of A_q as follows,

$$\pi^{(q)} := \int_{\hat{A}_0}^{\bigoplus} \psi_q(T_q(\tau_0)) d\mu_1 \bigoplus 2 \int_{\hat{A}_0}^{\bigoplus} \psi_q(T_q(\tau_0)) d\mu_2 \bigoplus \cdots \bigoplus \aleph_0 \int_{\hat{A}_0}^{\bigoplus} \psi_q(T_q(\tau_0)) d\mu_\infty$$

where $T_q: \hat{A}_0 \to \hat{A}_q$ is the Borel isomorphism of Theorem 3.2 discussed earlier. $\pi^{(q)}$ is well defined that follows from Theorem 3.2 and Corollary 4.5. Let $z_{k,j}(0)$ be a generator of A_0 . For any $\varepsilon > 0$, there is a natural number $m \in \mathbb{N}$ such that whenever $q < \frac{1}{m}$ we have,

$$\left\| \pi_{t,w}^{(q)}(c_{k,j}(q)z_{k,j}(q)) - \pi_{t,w}^{(0)}(z_{k,j}(0)) \right\| < \varepsilon$$

for all $t \in (S^1)^n$ and for all reduced words w and $c_{k,j}(q) = (-q)^{\min\{k-j,0\}}$. Note that we have chosen the natural number m independent of $t \in (S^1)^n$ and reduced word w. This follows from the way $\pi_{t,w}^{(q)}$ is defined for all $q \in [0,1)$. For any positive measure μ_i as above, we can write

$$\int_{\hat{A}_0}^{\bigoplus} \psi_0(\tau_0) d\mu_i \cong \int_{\hat{A}_{0,1}}^{\bigoplus} \psi_0(\tau_0) d\mu_i \bigoplus \left(\bigoplus_{w \neq 1} \int_{\hat{A}_{0,w}}^{\bigoplus} \psi_0(\tau_0) d\mu_i \right).$$

Similarly, we can break $\int_{\hat{A}_0}^{\bigoplus} \psi_q(T_q(\tau_0)) d\mu_i$ using Proposition 3.1. It is easy to see that

$$\left\| \int_{\hat{A}_{0,1}}^{\bigoplus} \psi_q(T_q(\tau_0))(c_{k,j}(q)z_{k,j}(q))d\mu_i - \int_{\hat{A}_{0,1}}^{\bigoplus} \psi_0(\tau_0)(z_{k,j}(0))d\mu_i \right\| = 0.$$

Let $\tau_0 = [V_w^* \pi_{t,w}^{(0)}(.) V_w] \to x(\tau_0)$ be a unit vector field i.e., $\int_{\hat{A}_{0,w}} \|x(\tau_0)\|^2 d\mu_i = 1$. Then

$$\begin{split}
& \left\| \left[\int_{\hat{A}_{0,w}}^{\bigoplus} \psi_{q}(T_{q}(\tau_{0}))(c_{k,j}(q)z_{k,j}(q))d\mu_{i} - \int_{\hat{A}_{0,w}}^{\bigoplus} \psi_{0}(\tau_{0})(z_{k,j}(0))d\mu_{i} \right] (x(\tau_{0})) \right\|^{2} \\
&= \int_{\hat{A}_{0,w}} \left\| V_{w}^{*} \left[\pi_{t,w}^{(q)}(c_{k,j}(q)z_{k,j}(q)) - \pi_{t,w}^{(0)}(z_{k,j}(0)) \right] V_{w}x(\tau_{0}) \right\|^{2} d\mu_{i} \\
&< \varepsilon^{2}. \int_{\hat{A}_{0,w}} \|x(\tau_{0})\|^{2} d\mu_{i} \\
&< \varepsilon^{2}.
\end{split}$$

That means for any reduced word $w \neq 1$,

$$\left\| \int_{\hat{A}_{0,w}}^{\bigoplus} \psi_q(T_q(\tau_0))(c_{k,j}(q)z_{k,j}(q))d\mu_i - \int_{\hat{A}_{0,w}}^{\bigoplus} \psi_0(\tau_0)(z_{k,j}(0))d\mu_i \right\| < \varepsilon$$

whenever $q < \frac{1}{m}$. Hence, we can write

$$\left\| \int_{\hat{A}_0}^{\bigoplus} \psi_q(T_q(\tau_0))(c_{k,j}(q)z_{k,j}(q))d\mu_i - \int_{\hat{A}_0}^{\bigoplus} \psi_0(\tau_0)(z_{k,j}(0))d\mu_i \right\| < \varepsilon.$$

Since the above inequality is true for all positive measures μ_i 's appearing in the decomposition of π_0 , therefore for any $q < \frac{1}{m}$ we have

$$\|\pi^{(q)}(c_{k,j}(q)z_{k,j}(q)) - \pi^{(0)}(z_{k,j}(0))\| \le \varepsilon.$$

Non-degeneracy of $\pi^{(q)}$ follows from the fact that irreducible representations are non-degenerate and a direct integral of non-degenerate representations is again non-degenerate (8.1.4, [3]). This concludes the proof of the theorem.

Our next goal is to start with a faithful representation $\pi^{(0)}$ and prove that we can choose $\pi^{(q)}$'s to be faithful. For this, we need some further results. The very first step is to observe that \hat{A}_{0,w_L} is an open dense subset of \hat{A}_0 . For any $1 \leq k \leq n$ we will write w_k for the reduced word $(s_1)(s_2s_1)\cdots(s_{k-1}s_{k-2}\cdots s_1)(s_ks_{k-1}\cdots s_2)(s_{k+1}s_k\cdots s_2s_1)\cdots(s_ns_{n-1}\cdots s_2s_1)$. That is the generator s_1 is only missing at the k-th bracket.

Proposition 4.8.

- 1. $(z_{n+1,1})(z_{n,1}z_{n+1,2})(z_{n-1,1}z_{n,2}z_{n+1,3})\cdots(z_{2,1}z_{3,2}z_{4,3}\cdots z_{n+1,n}) \in Ker \,\pi_{t,w}^{(0)} \text{ for all } t \in (S^1)^n$ and $w \neq w_L$. We will denote this element by a.
- 2. $\pi_{w_L}^{(0)}(a)$ is a non-zero operator which is elementary tensor product of P_0 . Hence \hat{A}_{0,w_L} is an open dense subset of \hat{A}_0 .
- 3. $\pi_{w_L}^{(0)}$ applied to any bracket element of a is again a positive operator consisting of elementary tensor product of P_0 and I.

Proof. 1. From the relations of $A_n(0)$ (see [4]) it follows that $z_{1,n+1}^* = (z_{2,1}z_{3,2}z_{4,3}\cdots z_{n+1,n}) \in Ker \pi_{t,w_1}^{(0)}$ with $w_1 = (s_2s_1)(s_3s_2s_1)\cdots(s_ns_{n-1}\cdots s_1)$. Similarly, one can prove by direct computation that $z_{3,1}z_{4,2}\cdots z_{n+1,n}\in Ker \pi_{t,w_2}^{(0)}$ with $w_2 = (s_1)(s_2)(s_3s_2s_1)\cdots(s_ns_{n-1}\cdots s_1)$ so on and finally $z_{n+1,1}\in Ker \pi_{t,w_n}^{(0)}$ with $w_n = (s_1)(s_2s_1)(s_3s_2s_1)\cdots(s_ns_{n-1}\cdots s_2)$. Since $\{w_1,w_2,\cdots w_n\}$ exhausts the list of all maximal Bruhat subwords of w_L , the rest of the proof follows from Theorem 3.4. The proof of both part 2 and part 3 follows from direct calculations. To prove $\hat{A}_{0.w_L}$ is open observe that its complement in \hat{A}_0 is closed which follows easily from the description of closed sets under the Jacobson topology.

Let $\pi^{(0)}$ be a non-degenerate representation of A_0 on a separable Hilbert space. For the sake of simplicity, let us further assume $\pi^{(0)}$ is just a single direct integral of irreducibles over \hat{A}_0 with respect to some non-negative Borel measure μ . The general case follows similarly with slight modifications. The next theorem helps us to filter out the minimal faithful part of a faithful representation of A_0 .

Theorem 4.9. $\pi^{(0)}$ is faithful if and only if $Q_L := \int_{\hat{A}_{0,w_L}}^{\bigoplus} V_{w_L}^* \pi_{t,w_L}^{(0)}(.) V_{w_L} d\mu$ is faithful.

Proof. Suppose that $\pi^{(0)}$ is faithful. Then using part 1 of Proposition 4.8 it follows that $\mu(\hat{A}_{0,w_L}) > 0$. Consider an element z such that $Q_L(z) = 0$. For showing z = 0, it suffices to prove that for any $s \in (S^1)^n$, $\pi^{(0)}_{s,w_L}(z)(e_{m_1} \otimes e_{m_2} \otimes \cdots \otimes e_{m_{n(n+1)/2}}) = 0$. Since A_0 is of type I therefore we have an element $y \in A_0$ such that $\pi^{(0)}_{s,w_L}(y) = (S^*)^{m_1} P_0 \otimes \cdots \otimes (S^*)^{m_{n(n+1)/2}} P_0$. Now, observe that $zya \in Ker \pi^{(0)}$. Since $\pi^{(0)}$ is faithful, therefore one has zya = 0. Now use part 2 of the Proposition 4.8 to conclude $\pi^{(0)}_{s,w_L}(z)(e_{m_1} \otimes e_{m_2} \otimes \cdots \otimes e_{m_{n(n+1)/2}}) = 0$. Converse is trivial.

The next proposition captures the faithfulness of a direct integral of irreducible representations in terms of support of the underlying positive Borel measure.

Proposition 4.10. Let A be any separable type I C^* -algebra and \hat{A} be the spectrum of A. Then any non-degenerate representation $\pi := \int_{\hat{A}}^{\bigoplus} \psi(\tau) d\mu$ (see Theorem 4.2) is faithful if and only if $supp(\mu) = \hat{A}$.

Proof. Assume π is faithful. Suppose there is an irreducible representation $\rho \notin \operatorname{supp}(\mu)$. As $\operatorname{supp}(\mu)$ is a closed subset of \hat{A} , using the definition of Jacobson topology one can find a non-zero element $x \in A$ such that $\sigma(x) = 0$ for all $\sigma \in \operatorname{supp}(\mu)$ and $\rho(x) \neq 0$. But this implies $\pi(x) = 0$ which contradicts the faithfulness of π .

Conversely, let us assume μ has full support but π is not faithful. Then there is a non-zero element $x \in A$ such that $\pi(x) = 0$. Consider the following subset $U := \{\tau \in \hat{A} : \psi(\tau)(x) \neq 0\}$ of \hat{A} . U is non-empty because an irreducible representation sends x to a non-zero operator as x is itself non-zero. Also observe that U is open in \hat{A} and $\mu(U) = 0$. But this contradicts our assumption μ has full support.

Corollary 4.11. Any non-degenerate representation π with decomposition as in Theorem 4.3 is faithful if and only if $supp(\sum_i \mu_i) = \hat{A}$.

Proof. Note that for each i, $\operatorname{supp}(\mu_i) \subseteq \operatorname{supp}(\sum_j \mu_j)$ and the rest of the proof follows similarly as above.

Currently it is not clear to us whether $T_q: \hat{A}_0 \to \hat{A}_q$ is homeomorphism, but its restriction to the dense open subset \hat{A}_{0,w_L} is indeed a homeomorphism from $\hat{A}_{0,w_L} \to \hat{A}_{q,w_L}$. We now proceed to prove this. It is already known that the natural map from $(S^1)^n \to \hat{A}_{q,w_L}$ is a continuous bijection for $q \in [0,1)$ (see Lemma 1). Therefore, our required homeomorphism can be achieved if we can prove \hat{A}_{0,w_L} is Hausdorff because homeomorphism will then automatically follow as for $q \neq 0$, \hat{A}_{q,w_L} is homeomorphic with $(S^1)^n$ (Theorem 4.1, [9]). The map $\rho: Irr_{\infty}(A_0) \to \hat{A}_{0,\infty}$ is open continuous onto map (see Lemma 1). Observe that the following set $Irr_{w_L,\infty}(A_0) :=$

 $\{W^*\pi_{t,w_L}^{(0)}(.)W: t\in (S^1)^n, W: l^2(\mathbb{N})\to l^2(\mathbb{N})^{\otimes l(w_L)} \text{ is unitary }\}=\rho^{-1}(\hat{A}_{0,w_L}) \text{ is a saturated open set. Therefore, } \rho|_{res}: Irr_{w_L,\infty}(A_0)\to \hat{A}_{0,w_L} \text{ is again an open quotient map defined on a Hausdorff space.}$

Lemma 3. Let R be an equivalence relation on a Hausdorff topological space X and $p: X \to X/R$ be an open quotient map. Then X/R is Hausdorff if the relation R is a closed relation that is $Q := \{(x,y) : xRy\} \subseteq X \times X$ is closed.

Proof. Since the map $p: X \to X/R$ is open therefore, the map $p \times p: X \times X \to (X/R) \times (X/R)$ defined by $p \times p(x,y) = (p(x),p(y))$ is also open. Now one can easily verify that

$$p \times p((X \times X) \setminus Q) = (X/R \times X/R) \setminus D_{X/R}$$

where $D_{X/R} := \{(p(x), p(x)) : x \in X\}$. As Q is closed, $D_{X/R}$ is also closed. That implies X/R is Hausdorff.

We will apply this result for $\rho|_{res}$. The equivalence relation will be closed in our case due to the following proposition. In the proof of the next proposition, we will use the notation $t = (t^{(1)}, t^{(2)}, \dots, t^{(n)})$ for any $t \in (S^1)^n$.

Proposition 4.12. Let U_m and U be unitaries from $l^2(\mathbb{N}) \to l^2(\mathbb{N})^{\otimes n(n+1)/2}$ such that for any $z \in A_0(n)$ and $x, y \in l^2(\mathbb{N})$ we have $\lim_{m \to \infty} \langle x | U_m^* \pi_{t_m, w_L}^{(0)}(z) U_m(y) \rangle = \langle x | U^* \pi_{t, w_L}^{(0)}(z) U(y) \rangle$. Then $\lim_{m \to \infty} t_m = t$.

Proof. Using the elements $z_{1,n+1}$ and $z_{n+1,1}$ one can show the first and last component of the sequence t_m converges to the first and last component of t respectively. For the convergence of other components of t_m , we need part 3 of the Proposition 4.8 particularly the positivity of the operators. Choosing x=y and bracket elements of a suitably one can conclude that all other components of t_m converge to components of t. For instance, let us focus on the convergence $t_m^{(n-1)} \to t^{(n-1)}$. Choose x=y appropriately so that

$$\overline{t^{(n-1)}}\langle x|U^*\pi_{w_L}^{(0)}(z_{n,1}z_{n+1,2})U(x)\rangle = \langle x|U^*\pi_{t,w_L}^{(0)}(z_{n,1}z_{n+1,2})U(x)\rangle \neq 0.$$

Therefore for all but finitely many m,

$$\overline{t_m^{(n-1)}} \langle x | U_m^* \pi_{w_L}^{(0)}(z_{n,1} z_{n+1,2}) U_m(x) \rangle = \langle x | U_m^* \pi_{t_m,w_L}^{(0)}(z_{n,1} z_{n+1,2}) U_m(x) \rangle \neq 0.$$

Also by part 3 of Proposition 4.8 we have

$$\langle x|U_m^*\pi_{w_L}^{(0)}(z_{n,1}z_{n+1,2})U_m(x)\rangle > 0, \ \langle x|U^*\pi_{w_L}^{(0)}(z_{n,1}z_{n+1,2})U(x)\rangle > 0$$

as $\pi_{w_L}^{(0)}(z_{n,1}z_{n+1,2})$ is a positive operator consisting of elementary tensor of P_0 and I. Therefore, we have

$$\begin{split} \lim_{m \to \infty} \langle x | U_m^* \pi_{w_L}^{(0)}(z_{n,1} z_{n+1,2}) U_m(x) \rangle &= \lim_{m \to \infty} \left| \langle x | U_m^* \pi_{t_m,w_L}^{(0)}(z_{n,1} z_{n+1,2}) U_m(x) \rangle \right| \\ &= \left| \langle x | U^* \pi_{t,w_L}^{(0)}(z_{n,1} z_{n+1,2}) U(x) \rangle \right| \\ &= \langle x | U^* \pi_{w_L}^{(0)}(z_{n,1} z_{n+1,2}) U(x) \rangle. \end{split}$$

Now it is evident that $t_m^{(n-1)} \to t^{(n-1)}$.

Therefore, the relative topology on \hat{A}_{0,w_L} is Hausdorff despite \hat{A}_0 being non-Hausdorff.

We now proceed to prove that $\int_{\hat{A}_0}^{\bigoplus} V_{w_L}^* \pi_{t,w_L}^{(q)}(.) V_{w_L} d\mu = \int_{\hat{A}_0}^{\bigoplus} \psi_q(T_q(\tau_0)) d\nu = \pi^{(q)} \text{ is faithful}$ if $\int_{\hat{A}_{t}}^{\bigoplus} V_{w_{L}}^{*} \pi_{t,w_{L}}^{(0)}(.) V_{w_{L}} d\mu = \int_{\hat{A}_{t}}^{\bigoplus} \psi_{0}(\tau_{0}) d\nu = \pi^{(0)}$ is faithful where ν is a new Borel measure on \hat{A}_0 which is equal to μ on $\hat{A}_{0,\underline{w}_L}$ and 0 otherwise. Using the standard change of variable formula one can write $\pi^{(q)} \cong \int_{\hat{A}}^{\bigoplus} \psi_q(\tau_q) dT_q \nu$. So, for proving the faithfulness of $\pi^{(q)}$, it is enough to show that $supp(T_q\nu) \stackrel{\cdot}{=} \hat{A}_q$ (see Proposition 4.10). As support is a closed set, it further reduces to prove $\hat{A}_{q,w_L} \subseteq supp(T_q\nu)$ (see Corollary 3.6).

Proposition 4.13. $\pi^{(q)}$ is a faithful if and only if $\pi^{(0)}$ is faithful.

Proof. Assume that $\operatorname{supp}(\nu) = \hat{A}_0$ but there is an element $x \in \hat{A}_{q,w_L}$ and x is outside $\operatorname{supp}(T_q \nu)$. That means we can find a non-empty open subset $V \subseteq \hat{A}_q$, $x \in V$ and $T_q \nu(V) = 0$. Hence $x \in \mathcal{A}_q$ $V \cap \hat{A}_{q,w_L}$ is a non-empty open subset of \hat{A}_{q,w_L} with $T_q \nu(V \cap \hat{A}_{q,w_L}) = 0$. That implies $\nu(T_q^{-1}(V \cap \hat{A}_{q,w_L})) = 0$. $\hat{A}_{q,w_L})) = 0$. But $T_q^{-1}(V \cap \hat{A}_{q,w_L})$ is non-empty open in \hat{A}_{0,w_L} (using the homeomorphism). By part 2 of the Proposition 4.8, \hat{A}_{0,w_L} is open in \hat{A}_0 . So, $T_q^{-1}(V \cap \hat{A}_{q,w_L})$ is a non-empty open subset of \hat{A}_0 with $\nu(T_q^{-1}(V \cap \hat{A}_{q,w_L})) = 0$ which contradicts our assumption.

Converse follows along the same lines.

Now we are ready to formulate our concluding results.

Theorem 4.14. Let $\pi^{(0)}$ be a non-degenerate faithful representation of A_0 on a separable Hilbert space \mathcal{H} . Then there exist non-degenerate faithful representations $\pi^{(q)}$ of A_q on the same Hilbert space H such that

$$\pi^{(0)}(z_{k,j}(0)) = \lim_{q \to 0+} \pi^{(q)} \left((-q)^{\min\{k-j,0\}} z_{k,j}(q) \right), \qquad k,j \in \{1,2,\dots,n+1\}.$$

Proof. Choose $\pi^{(q)}$ as given in the proof of the Theorem 4.7. Use Theorem 4.9 to filter out the minimal faithful part of $\pi^{(0)}$. Faithfulness of $\pi^{(q)}$ now follows from the Propositions 4.10 and 4.13. The next theorem establishes the converse that any non-degenerate faithful representation of $A_q(n)$ for $q \neq 0$ gives rise to a non-degenerate faithful representation of the crystallized algebra in a natural way.

Theorem 4.15. For any $q_0 \in (0,1)$, let $\pi^{(q_0)}$ be a non-degenerate faithful representation of A_{q_0} on a seperable Hilbert space \mathcal{H} . Then there exist non-degenerate faithful representations $\pi^{(q)}$ of A_q on the same Hilbert space \mathcal{H} for all $q \in [0, q_0)$ such that

$$\pi^{(0)}(z_{k,j}(0)) = \lim_{q \to 0+} \pi^{(q)} \left((-q)^{\min\{k-j,0\}} z_{k,j}(q) \right), \qquad k,j \in \{1,2,\dots,n+1\}.$$

Proof. To prove this, we need a $q \neq 0$ version of Proposition 4.9, that is, $\pi^{(q)} \cong \int_{\hat{A}_q}^{\bigoplus} \psi_q(\tau_q) d\mu$

is faithful if and only if $Q_L:=\int_{\hat{A}_{q,w_L}}^{\bigoplus}V_{w_L}^*\pi_{t,w_L}^{(q)}(.)V_{w_L}d\mu$ is faithful. It can similarly be proved by exploiting the fact that \hat{A}_{q,w_L} is an open dense subset of \hat{A}_q for $q\neq 0$ (Theorem 4.1, [9]) and $A_q(n)$ is a type I C^* -algebra. Let $\pi^{(q)}$ be faithful and $z\in A_q$ be such that $Q_L(z)=0$. We have to prove that z=0. It again suffices to show that for any $s\in (S^1)^n$, $\pi_{s,w_L}^{(q)}(z)=0$. Now using the description of closed subsets of \hat{A}_q we can find a non-zero element $a\in \bigcap_{[\pi]\in \hat{A}_q\setminus \hat{A}_{q,w_L}} Ker\,\pi$ but $\pi_{s,w_L}^{(q)}(a)=K\neq 0$. Let K_1 be a any compact operator on $l^2(\mathbb{N})^{\otimes l(w_L)}$. Since A_q is of type I, there exists an element $z_{K_1}\in A_q$ such that $\pi_{s,w_L}^{(q)}(z_{K_1})=K_1$. Now observe that $zz_{K_1}a\in Ker\pi^{(q)}$. So we have $zz_{K_1}a=0$. This again implies $\pi_{s,w_L}^{(q)}(z)K_1(K)=0$ where K_1 can be choosen to be any arbitrary compact operator. Hence it follows that $\pi_{s,w_L}^{(q)}(z)=0$.

The rest of the argument follows similarly by performing the steps mentioned in the proof of Theorem 4.14 at $q \neq 0$ level, using the canonical homeomorphism between \hat{A}_{0,w_L} and \hat{A}_{q,w_L} and Proposition 4.10.

Corollary 4.16. Given any $q_0 \in (0,1)$ and a non-degenerate faithful representation $\pi^{(q_0)}$ of $C(SU_{q_0}(n+1))$ on a separable Hilbert space \mathcal{H} , there exist non-degenerate faithful representations $\pi^{(q)}$ of $C(SU_q(n+1))$ on the same Hilbert space \mathcal{H} for all $q \in (0,q_0)$ such that the following limits exist in norm

$$\lim_{q \to 0+} \pi^{(q)} \left((-q)^{\min\{k-j,0\}} z_{k,j}(q) \right) \in B(\mathcal{H}), \qquad k, j \in \{1, 2, \dots, n+1\}$$

and the C^* -subalgebra of $B(\mathcal{H})$ generated by these limit operators is isomorphic to the crystallized algebra $C(SU_0(n+1))$.

Remark 4.17. In particular, one can choose $\pi^{(q)}$'s to be the Soibelman representations of $C(SU_q(n+1))$ for $q \neq 0$, that is the crystallized algebra $C(SU_0(n+1))$ is isomorphic to the C^* -algebra generated by the limit operators of the Soibelman representation of $C(SU_q(n+1))$ as observed in Theorem 8.4 of [5].

Acknowledgement- We are greatly indebted to Prof. Arup K. Pal for his valuable suggestions and constant support time and again. We would also like to thank the Indian Statistical Institute for supporting us both in the form of a Ph.D. fellowship.

References

- [1] A. Bjorner and F. Brenti. Combinatorics of Coxeter Groups. Graduate texts in Mathematics 231, 2005, Springer.
- [2] K. Bragiel. The Twisted SU(N) Group. On the C^* -Algebra $C(S_{\mu}U(N))$. Letters in Mathematical Physics 20: 251-257, 1990.
- [3] Jacques Dixmier. C*-algebras. Volume 15, 1977, North Holland publishing company.
- [4] Manabendra Giri and Arup Kumar Pal. Quantized function algebras at q=0: type A_n case. Proc. Indian Acad. Sci. (Math. Sci.) 134, 30 (2024).
- [5] Manabendra Giri and Arup Kumar Pal. Irreducible representations of the crystallization of the quantized function algebras $C(SU_q(n+1))$. J. Noncommut. Geom. Doi 10.4171/JNCG/608.
- [6] Alexander S Kechris. Classical Descriptive Set theory. 1994, Springer.
- [7] L. I. Korogodski and Y. S. Soibelman. Algebras of functions on quantum groups. Part I. Math. Surveys Monogr. 56, American Mathematical Society, Providence, RI, 1998.
- [8] Marco Matassa and Robert Yuncken. Crystal limits of compact semisimple quantum groups as higher-rank graph algebras. J. Reine Angew. Math., 802: 173–221, 2023.
- [9] S. Neshveyev and L. Tuset. Quantized algebras of functions on homogeneous spaces with Poisson stabilizers. Commun. Math. Phys. 312(1), 223–250 (2012)

Saikat Das

Theoretical Statistics and Mathematics Unit, Indian Statistical Institute, Delhi, India; saikat20r@isid.ac.in, saikat1811@gmail.com

Ayan Dey

Theoretical Statistics and Mathematics Unit, Indian Statistical Institute, Delhi, India; ayan22r@isid.ac.in, studentayandey@gmail.com