HOMOTOPY TYPES OF SMALL SEMIGROUPS

DENNIS SWEENEY

ABSTRACT. We algorithmically compute integral Eilenberg-MacLane homology of all semigroups of order at most 8 and present some particular semigroups with notable classifying spaces, refuting conjectures of Nico. Along the way, we give an alternative topological proof of the fact that if a finite semigroup Shas a left-simple or right-simple minimal ideal K(S), then the classifying space BS is homotopy equivalent to the classifying space B(GS) of the group completion. We also describe an algorithm for computing the group completion GS of a finite semigroup S using asymptotically fewer than $|S|^2$ semigroup operations. Finally, we show that the set of homotopy types of classifying spaces of finite monoids is closed under suspension.

CONTENTS

1. Introduction	1
2. Classifying Spaces	3
3. McDuff's Result	4
4. First Properties	5
5. The Structure of the Minimal Ideal	6
6. Group Completions	8
7. K-thin Semigroups have the Classifying Space of a Group	10
8. Discussion from Homology Computations	13
9. Joins and Suspensions	18
References	20
Appendix A. Computing Homology as Tor	22
Appendix B. Tables of Homology	28

1. INTRODUCTION

The classifying space BG of a discrete group G is an Eilenberg-MacLane space of type K(G, 1). However, McDuff [McD79] showed that the classifying space BS of a monoid S has no such homotopical restriction: BS can attain any connected weak homotopy type. Later, Fiedorowicz [Fie84] conjectured that any homotopy type of a simply connected finite simplicial complex is attainable as the classifying space of some finite monoid. While this conjecture remains open, we will give some partial results on the classifying spaces of finite semigroups and monoids. The results are summarized as follows.

For any finite semigroup or monoid S with minimal two-sided ideal K(S):

- If K(S) is left-simple or right-simple then we have a homotopy equivalence $BS \simeq BGS$, where GS is the group completion of S (Theorem 7.3). This is implied by a cohomological result of [AR67] or [Nic69], but here we give a proof that directly describes the homotopies defining the equivalence, without using Tor or Ext functors (Remark 7.6).
 - This holds when S is left-commutative, inverse, or left-cancellative.
 - This also holds when S satisfies some equation $x_{\lambda_1} \cdots x_{\lambda_{\ell}} = x_{\lambda'_1} \cdots x_{\lambda'_{\ell'}}$ in which the leftmost variable on the two sides of the equation is not the same, such as xy = yx for commutativity or xya = yxa for leftcommutativity (Theorem 7.9).
 - This condition is satisfied by 99.99999999990% of the semigroups of order 10, up to isomorphism and anti-isomorphism (Figure 2).
- A Rees matrix semigroup structure for K(S) can be computed using O(|S|) semigroup operations, and so we can very quickly determine when the first bullet holds (Algorithm 5.1).
- We present Algorithm 6.1 to compute GS in $O(|S|\log(|S|)\alpha(|S|))$ time, using many fewer than $|S|^2$ semigroup operations. Here α is the inverse Ackermann function.
- We present Algorithm A.1 for computing the homology of *BS*. Several integral homology groups are listed for all semigroups of order at most 8 (Appendix B).

We compile a list of finite semigroups to serve as examples, especially those with homology notably different from that of any finite group:

- The 5-element semigroup S of Example 8.1 has no left or right zero element but has contractible BS.
- The 6-element semigroup S of Example 8.2 has $BS \simeq \mathbb{S}^3$.
- The 5-element semigroup S of Example 8.5 has $H_i(BS; \mathbb{Z}) \cong \mathbb{Z}$ for all $i \ge 2$, which disproves the conjectures of the introduction to [Nic69] that homology in high dimensions is determined by subgroups. However, some similar weakened statements may apply to bands or regular semigroups (Questions 8.4 and 8.6).
- It is known that the rectangular band semigroup Rect_b^a satisfies $B \operatorname{Rect}_b^a \simeq \bigvee_{k=1}^{(a-1)(b-1)} \mathbb{S}^2$ (Proposition 3.6).
- The 9-element semigroup S of Example 8.7 has $BS \simeq \Sigma^2(\mathbb{R}P^2)$, but no subgroup of order 2.
- The 12-element semigroup S of Example 8.9 has $BS \simeq \Sigma(\mathbb{R}P^2)$.
- The semigroup S of Example 8.10 has $H_6(BS;\mathbb{Z}) \cong \mathbb{Z}^9 \times C_{1494640}$, but only 10 elements. Here and throughout, C_k denotes the cyclic group of order k.
- The 6-element semigroup S of Example A.5 has $H_i(BS;\mathbb{Z}) = \mathbb{Z}^{2^{i-2}}$ for $i \ge 2$.

Finally, we show that for any monoid S and any $n \ge 1$, there is a monoid S' with (1+n)|S| elements satisfying $BS' \simeq \bigvee_{i=1}^{n-1} \Sigma(BS)$ (Corollary 9.4).

- Taking n = 2, this implies that the set of homotopy types of finite monoids (or semigroups) is closed under suspension.
- This also gives examples of finite monoids with infinitely generated cohomology rings (Remark 9.5).




FIGURE 1. An illustration of the low-dimensional cells in BS

Acknowledgements. The author would like to thank Sanjeevi Krishnan for many helpful conversations and suggestions.

2. Classifying Spaces

This paper investigates the nerve construction: we build a topological space based on the algebraic data of a (discrete) semigroup or monoid, and investigate the correspondence between the algebra of the semigroup and the the algebraic topology of the space. There are also purely-algebraic definitions of monoid homology as in [CL24], but we will emphasize topological definitions.

Definition 2.1. The classifying space $BS = B_{Sgrp}S$ of a semigroup S will mean the realization of the Δ -set consisting of an n-simplex for each n-tuple of elements of S (see Figure 1), with face maps $d_i^n : S^n \to S^{n-1}$ given by

$$d_i^n(x_1, \dots, x_n) = \begin{cases} (x_2, \dots, x_n) & i = 0\\ (x_1, \dots, x_{i-1}, x_i x_{i+1}, x_{i+2}, \dots, x_n) & 0 < i < n\\ (x_1, \dots, x_{n-1}) & i = n. \end{cases}$$

This is essentially the definition given in [Hat02]. One can can distinguish the combinatorial data $\mathcal{N}_{Sgrp}S$ of this Δ -set (usually called the *nerve*) and its realization $B_{Sgrp}S = |\mathcal{N}_{Sgrp}S|$ as a topological space.

The definition for monoids incorporates the data of the unit:

Definition 2.2. The classifying space $BS = B_{\text{Mon}}S$ of a monoid S is the realization of the simplicial set with the same n-cells and face maps as $B_{\text{Sgrp}}S$, but additionally equipped with degeneracy maps $s_i^n \colon S^n \to S^{n+1}$ given by

$$s_i^n(x_1, \dots, x_n) = (x_1, \dots, x_i, 1_S, x_{i+1}, \dots, x_n) \text{ for } 0 \le i \le n.$$

As above, one can distinguish between the nerve $\mathcal{N}_{\mathsf{Mon}}S$ and its realization $B_{\mathsf{Mon}}S$.

Fortunately, equipping the degeneracy maps does not change the homotopy type: for any monoid S, $B_{Sgrp}S \simeq B_{Mon}S$, since $B_{Sgrp}S$ and $B_{Mon}S$ are the thick realization $\|\cdot\|$ and the standard realization $|\cdot|$ of the same simplicial set:

$$B_{\mathsf{Sgrp}}S = |\mathcal{N}_{\mathsf{Sgrp}}S| = ||\mathcal{N}_{\mathsf{Mon}}S|| \simeq |\mathcal{N}_{\mathsf{Mon}}S| = B_{\mathsf{Mon}}S.$$

The equivalence $\|\cdot\| \simeq |\cdot|$ is show in Appendix A of [Seg74]. This shows that the forgetful functor Mon \rightarrow Sgrp preserves the homotopy type of the classifying space.

The forgetful functor has a left adjoint $(\cdot)^{1}$: Sgrp \rightarrow Mon that also preserves the classifying space:

Definition 2.3. For any semigroup S, write $S^{1} := S \sqcup \{1\}$ for the monoid formed by freely adjoining a unit to S, even if S is already a monoid.

Since the nondegenerate cells of $B_{Mon}S^{\mathbb{1}}$ are exactly the cells of $B_{Sgrp}S$, these are homeomorphic. Alternately, one could simply define $B_{Sgrp}S \coloneqq B_{Mon}S^{\mathbb{1}}$. Thanks to these equivalences, we can generally write BS to mean $B_{Mon}S$ or $B_{Sgrp}S$ without confusion.

Remark 2.4. When G is a group, maps $X \to BG$ classify principle G-bundles over nice spaces X. This theory can be extended to monoids and categories—see [Wei05] for example. Another variant of this idea is presented in [FH10]. We do not discuss either any further here.

3. McDuff's Result

The central theorem about classifying spaces of discrete monoids and semigroups was proved by Dusa McDuff in 1979:

Theorem 3.1 (Theorem 1 in [McD79]). Every path-connected space has the same weak homotopy type as the classifying space BS of some (discrete) monoid S.

To prove this, McDuff approximates the space by a simplicial complex \mathcal{K} , then constructs a small category with an object for each vertex of \mathcal{K} , an invertible morphism for each edge of \mathcal{K} , and an idempotent for each simplex of dimension ≥ 2 in \mathcal{K} , essentially subject to relations of the form $\tau \sigma = \sigma$ for any face τ of a simplex σ . She then shows by induction that the classifying space of this category is equivalent to \mathcal{K} , and since all objects in the category are isomorphic, the skeleton of the category is a monoid. This monoid is typically infinite.

Remark 3.2. The corresponding result for categories is easier: every simplicial complex is homotopy equivalent to the classifying space of some category because the nerve of the poset of faces in a simplicial complex is exactly the barycentric subdivision of the simplicial complex.

Remark 3.3. The Kan-Thurston Theorem [KT76] gives that every path-connected weak homotopy type is *homology equivalent* to the classifying space of some group, i.e., groups are as homologically expressive as possible. However, since BG is a K(G, 1), the class of homotopy types arising from groups is severely restricted. Theorem 3.1 thus states that broadening the available objects from groups to monoids removes this homotopical restriction.

Fiedorowicz gives in [Fie84] an alternative proof of Theorem 3.1, and he suggests that the following question likely has a positive answer:

Question 3.4. Is every finite simply connected simplicial complex homotopy equivalent to the classifying space of some finite monoid?

The simple-connectedness restriction (or some other restriction on $\pi_1(BS)$) is required because a finite monoid or semigroup S cannot have infinite $\pi_1(BS)$ — $\pi_1(BS)$ is the group completion GS (Lemma 6.2), and the group completion GS of a finite semigroup is finite. For example, the circle \mathbb{S}^1 is not homotopy equivalent to the classifying space of some finite monoid or semigroup. On the other hand, some important simply connected homotopy types can already be constructed from finite semigroups without too many elements:

Definition 3.5. For $a, b \in \mathbb{N}$, an *a*-by-*b* rectangular band is a semigroup

$$\operatorname{Rect}_{b}^{a} \coloneqq \{x_{ij} : i \in \{1, \dots, a\}, j \in \{1, \dots, b\}\}$$

with operation $x_{ij}x_{k\ell} \coloneqq x_{i\ell}$.

Proposition 3.6. $B(\operatorname{Rect}_{b}^{a}) \simeq \bigvee_{k=1}^{(a-1)(b-1)} \mathbb{S}^{2}$, the wedge of (a-1)(b-1) copies of the 2-sphere.

A proof will follow in Section 8. Fiedorowicz [Fie84; Fie02] describes the case of $B(\text{Rect}_2^2) \simeq \mathbb{S}^2$, using the 4-element semigroup Rect_2^2 . A further generalization appears in [Ste24].

4. First Properties

We list some basic properties of the functors $B: \mathsf{Mon} \to \mathsf{Top} \text{ and } B: \mathsf{Sgrp} \to \mathsf{Top}$.

Proposition 4.1. For a monoid S, the monoid S^{1} (that has an additional unit adjoined) satisfies $BS \simeq B(S^{1})$, and the same holds for adjoining a unit to a semigroup.

Proof. For monoids S

$$B_{\mathsf{Mon}}(S^{\mathbb{1}}) = B_{\mathsf{Sgrp}}(S) \simeq B_{\mathsf{Mon}}(S),$$

and for semigroups S,

$$B_{\mathsf{Sgrp}}(S^{\mathbb{1}}) \simeq B_{\mathsf{Mon}}(S^{\mathbb{1}}) = B_{\mathsf{Sgrp}}(S).$$

Remark 4.2. Since we can adjoin identities as above arbitrarily many times, the order of a finite semigroup or monoid does not carry essential information about the homotopy type of its classifying space, which is in contrast to the situation for groups: multiplication by the order of a finite group G annihilates $H^n(G;\mathbb{Z})$ ([Bro82], Corollary 10.2).

Proposition 4.3. If S and T are monoids, then $B(S \times T) \cong BS \times BT$.

Proof. We have an isomorphism of simplicial sets:

 $((s_1,t_1),\ldots,(s_n,t_n))\leftrightarrow((s_1,\ldots,s_n),(t_1,\ldots,t_n))).$

Geometric realization of simplicial sets preserves products so long as S and T are countable (see Theorem A.6 in [Hat02]). Otherwise, we must agree that products of spaces are implicitly taken in a convenient subcategory of spaces.

Warning 4.4. B does not preserve products of semigroups. For instance, using Proposition 3.6, we have:

 $B(\operatorname{Rect}_2^1 \times \operatorname{Rect}_1^2) = B(\operatorname{Rect}_2^2) \simeq \mathbb{S}^2 \not\simeq * = * \times * = B(\operatorname{Rect}_2^1) \times B(\operatorname{Rect}_2^2).$

However, based on Propositions 4.1 and 4.3, for semigroups S and T we do have

 $B(S) \times B(T) \cong B(S^{\mathbb{1}}) \times B(T^{\mathbb{1}}) \simeq B(S^{\mathbb{1}} \times T^{\mathbb{1}}).$

Units are essential to form the product space as a nerve: a k-cell in B(S) can be paired with the 0-cell in B(T) to form one of the k-cells in $B(S) \times B(T)$, but $B(S \times T)$ has no way to represent this pairing if T has no unit. Next we see that B also respects pointed coproducts, but we won't use this, as free products of nontrivial monoids or semigroups are always infinite.

Proposition 4.5. If S and T are monoids or semigroups, $B(S * T) \simeq BS \lor BT$.

Proof. The result for monoids follows from Lemma 3 of [McD79], where the inclusions $1 \to S$ and $1 \to T$ are right-free, as S and T are unions of disjoint cosets $x1 = \{x\}$. To extend to semigroups S and T, we have $(S *_{\mathsf{Sgrp}} T)^{\mathbb{1}} \cong S^{\mathbb{1}} *_{\mathsf{Mon}} T^{\mathbb{1}}$, so

 $B(S\ast_{\mathsf{Sgrp}} T)\cong B\bigl((S\ast_{\mathsf{Sgrp}} T)^{\mathbb{1}}\bigr)\cong B\bigl(S^{\mathbb{1}}\ast_{\mathsf{Mon}} T^{\mathbb{1}}\bigr)\simeq BS^{\mathbb{1}}\vee BT^{\mathbb{1}}\cong BS\vee BT.\quad \Box$

Proposition 4.6. If S is a semigroup or monoid with a left zero element z (i.e. zx = z for all $x \in S$), then BS is contractible.

Proof. If S is only a semigroup, then z remains a left zero in S^{1} because z1 = z, so without loss of generality, assume S is a monoid. Viewing S as a category with one object, we can show $BS \simeq *$ by providing natural transformations as homotopies. The composition $* \to S \to *$ is the identity functor on the nose, and the composition $S \to * \to S$ is the constant functor const₁: $S \to S$ where const₁(x) = 1 for $x \in S = S(*,*)$. There is a natural transformation η : id_S \Rightarrow const₁ with unique component $\eta_* = z$ and naturality square

$$\begin{array}{c} * \xrightarrow{\eta_* = z} * \\ \text{id} x = x \downarrow & \downarrow \\ * \xrightarrow{\eta_* = z} * \end{array}$$

which commutes because zx = z = 1z, so η provides the nullhomotopy of id_{BS}. \Box

For any semigroup or monoid S, note that by symmetry there is a homeomorphism $BS \cong B(S^{\text{op}})$, where S^{op} is the opposite semigroup, i.e., the same underlying set as S but with operation $(x, y) \mapsto yx$. Hence, we have a dual statement to the previous proposition: we also know BS is contractible if S has a right zero element.

Remark 4.7. Proposition 4.6 implies that the classifying spaces of the majority of semigroups are contractible: [KRS76] shows that for labelled semigroups on the set $\{1, \ldots, n\}$, the fraction of associative multiplication tables having a zero element (and having a stronger property called 3-*nilpotence*) approaches 100% as n grows. It is believed by semigroup theorists [DM12] that the same holds for semigroups of order n up to isomorphism, but this has not been proved. It was verified in [Dis+12] that 99.9942% of semigroups of order 10, up to isomorphism and anti-isomorphism, are 3-nilpotent and thus have a zero element.

5. The Structure of the Minimal Ideal

While not as rigidly structured as groups, semigroups nonetheless admit structural study via their ideals and *Green's Relations*. A classical text is [CP61], and helpful introductions include [How95] and the first chapter of [HS12]. Of particular interest for us will be the study of the minimal ideal of a finite semigroup.

For a semigroup S, a subset $A \subseteq S$ is called a *left ideal* of S if $SA \subseteq A$, a *right ideal* if $AS \subseteq A$, and an *ideal* if both $SA \subseteq A$ and $AS \subseteq A$. Equivalently, A is an ideal if $S^{\mathbb{1}}AS^{\mathbb{1}} \subseteq A$ or equivalently $S^{\mathbb{1}}AS^{\mathbb{1}} = A$. Some authors require ideals to be nonempty. Note that the intersection of two nonempty ideals $A \ni a$ and $A' \ni a'$ is nonempty: $A \cap A' \ni aa'$. We can thus define:

Definition 5.1. For a nonempty finite semigroup S, the minimal ideal K(S) is the (nonempty) intersection of all nonempty ideals of S.

Proposition 5.2. For a finite semigroup $S = \{x_1, \ldots, x_n\}$ with elements listed in any order, the product $k \coloneqq x_1 \cdots x_n$ of all elements of S belongs to K(S). In fact $K(S) = SkS = S^{\mathbb{1}}kS^{\mathbb{1}}$.

Proof. Each nonempty ideal A contains some x_i , so

 $k = (x_1 \cdots x_{i-1}) x_i (x_{i+1} \cdots x_n) \in S^{\mathbb{1}} A S^{\mathbb{1}} \subseteq A,$

and so $k \in \bigcap_A A = K(S)$. Because $k \in K(S)$, we know $S^{\mathbb{1}}kS^{\mathbb{1}}$ and SkS are nonempty ideals and subsets of K(S), so they are equal to K(S) by minimality. \Box

Definition 5.3. The *Rees matrix semigroup* $\mathcal{M}(H; I, J; \llbracket \rrbracket)$ is a semigroup defined on the set $I \times H \times J$, where I and J are sets and H is a group, and where $\llbracket \rrbracket : J \times I \to$ H is any function. The multiplication on $\mathcal{M}(H; I, J; \llbracket \rrbracket)$ is defined by

$$(i, h, j)(i', h', j') = (i, h[[j, i']]h', j')$$

The table [[]] of H elements is called the *sandwich matrix*.

A Rees matrix semigroup $\mathcal{M}(1; I, J; \llbracket])$ with trivial H is a rectangular band $\operatorname{Rect}_{|J|}^{|I|}$. More generally, if $\llbracket i, j \rrbracket = 1$ for all i, j, then $\mathcal{M}(H; I, J; \llbracket])$ is a direct product $H \times \operatorname{Rect}_{|J|}^{|I|}$. A Rees matrix semigroup with a nontrivial sandwich matrix can be thought of as a "twisted product" of a group and a rectangular band.

Rees matrix semigroups are *simple*—they have no proper nonempty ideals. It happens that all finite simple semigroups take this form, which is implied by the next theorem. The statement below follows Theorems 1.62 and 1.63 of [HS12], but the essential content was first proved by Suschkewitsch in [Sus28], considered the first paper properly belonging to abstract semigroup theory (see the history in [Hol09]). Later, Rees [Ree40] generalized the structure theorem to certain infinite semigroups, but we concern ourselves with the finite case.

Theorem 5.4 (The Structure Theorem). For any finite nonempty semigroup S, the minimal ideal K = K(S) is isomorphic as a semigroup to $\mathcal{M}(H; I, J; [\![]\!])$ for some finite sets I and J, some finite group H, and some sandwich matrix $[\![]\!]$. Furthermore:

- The sandwich matrix can be taken to be *normalized* to some $i_0 \in I$ and $j_0 \in J$ by assuming that $[\![j_0, i]\!] = e_H$ for all $i \in I$ and $[\![j, i_0]\!] = e_H$ for all $j \in J$.
- The minimal right ideals of S are exactly the subsets of K corresponding to $\{i\} \times H \times J$ for each *i*, and likewise minimal left ideals of S take the form $I \times H \times \{j\}$.
- The maximal subgroups of K are the sets $\{i\} \times H \times \{j\}$, which are all isomorphic to H.

To construct the isomorphism for Theorem 5.4, we can choose an arbitrary $k \in K$, take $I = \{i \in S^{\mathbb{1}}k : ii = i\}$, $J = \{j \in kS^{\mathbb{1}} : jj = j\}$, $H = kS^{\mathbb{1}}k$, and [[j,i]] = ji. Each element of K has a unique representation as (i, h, j) := ihj where $i \in I$, $h \in H$, and $j \in J$, and (i, h, j)(i', h', j') = ihji'h'j' = (i, h[[j,i']]h', j'). The identity e_H is the unique idempotent of H, and $i_0 = j_0 = e_H$. See [HS12] for a detailed proof. All of this is explicitly constructive, so we will call this Algorithm 5.1. Algorithm 5.1: Computing the structure of K(S) in O(|S|) time.


 $\begin{array}{l} \textbf{Input: A nonempty finite semigroup } S = \{x_1, \ldots, x_{|S|}\} \text{ with operation } \ast \\ \textbf{Output: Subsets } I, H, J \subseteq S \text{ so that } K(S) = IHJ \cong \mathcal{M}(H; I, J; \ast) \\ \textbf{Function getMinIdeal}(S, \ast): \\ k \coloneqq x_1 \ast x_2 \ast \cdots \ast x_{|S|}; \quad /\ast \text{ Find } k \in K(S) \text{ as in Proposition 5.2 } \ast / \\ I \coloneqq \{x \ast k : x \in S \text{ with } (x \ast k) \ast (x \ast k) = x \ast k\}; \\ J \coloneqq \{k \ast x : x \in S \text{ with } (k \ast x) \ast (k \ast x) = k \ast x\}; \\ H \coloneqq \{k \ast x \ast k : x \in S\}; \\ \textbf{return } I, H, J; \end{aligned}$

6. GROUP COMPLETIONS

One application of the Structure Theorem is the computation of the group completion of a semigroup or monoid.

Definition 6.1. The group completion functor $G: \mathsf{Sgrp} \to \mathsf{Grp}$ is the left adjoint to the forgetful functor $\mathsf{Grp} \to \mathsf{Sgrp}$.

For a semigroup S, the group completion semigroup homomorphism $S \to GS$ is characterized by the universal property that for any group Γ , any semigroup homomorphism $S \to \Gamma$ factors as a composition $S \to GS \to \Gamma$. As such, GS is alternately referred to as the maximal homomorphic group image of S as in [HM03], or as the enveloping group of S as in [MS76]. The group completion of a semigroup with a certain semigroup presentation is given by reinterpreting the same generators and relations as a group presentation. The group completion of a monoid is similar, and the following commutes:



The group GS is significant to the topology of BS:

Lemma 6.2. For any monoid or semigroup S, we have $\pi_1(BS) \cong GS$.

The above is Lemma 1 in [McD79] and is proved by checking the universal property. The following is proved in [Lar72], Lemme II.3; see also [HM03].

Theorem 6.3. Any finite nonempty semigroup S has GS = H/N, where

- $K(S) \cong \mathcal{M}(H; I, J; \llbracket \rrbracket),$
- $\llbracket \rrbracket$ is normalized to some $j_0 \in J$ and $i_0 \in I$, and
- $N \leq H$ is the normal closure of $\{ [j, i] : i \in I, j \in J \}$.

The mapping $S \to H/N$ is $x \mapsto [(i_0, e, j_0)x(i_0, e, j_0)]_N$. Note that the group completion of a finite semigroup depends only on its minimal ideal: GS = G(K(S)). Further note that the map $S \to GS$ is surjective when S is finite.

Assuming S is a semigroup with a computational representation that allows efficient iteration, constant-time multiplication, and constant-time checks for equality of elements, Algorithm 6.1 shows that one can compute the group completion GSin close to linear time in |S|. Notably, we don't need to access all $|S|^2$ products in S, nor do we need to explicitly produce the $|GS|^2$ products of the result—we can compute $GS = (R, *_R)$ by finding a set $R \subset S$ of representatives of the elements of GS and then multiply representatives in constant time as $r_1 *_R r_2 \coloneqq \rho(r_1 r_2)$.

Algorithm 6.1 is a direct implementation of Theorem 6.3. It computes a set $(JI)^H$ of generators for the normal closure of JI in $O(|I||J||H|) = O(|K(S)|) \subseteq O(|S|)$ time, then generates the normal closure $N = \langle (JI)^H \rangle$. Finding $(JI)^H$ requires computing inverses, which is possible in linear time using Algorithm 6.2 (where each loop computes some n inverses in O(n) time¹). Generating $N = \langle (IJ)^H \rangle$ is efficient using Algorithm 6.3, which makes use of a disjoint-set forest data structure that represents equivalence relations in such a way that new equivalences can be added in amortized $O(\alpha(|S|))$ time, where α is the extremely slow-growing inverse Ackermann function—see [CLRS], Chapter 21.3. The outer loop in Algorithm 6.3 maintains the invariant that after the first n elements $\{x_1, \ldots, x_n\}$ of X have been processed, the equivalence relation has partitioned H into cosets of $\langle x_1, \ldots, x_n \rangle$. Each time the body of the if-statement runs, the cosets grow by a factor of at least 2, so there can be at most $\log_2(|H|)$ executions of this body, each of which takes $O(|H|\alpha(|H|))$ time.

An alternative method to compute GS would be to find generators for S and compute a presentation as in [FP97], then reinterpret this as a group presentation, but computing with general group presentations is not straightforward. Another alternative would be to begin by searching for x, y satisfying xy = x or similar relations, but even this already requires $|S|^2$ semigroup operations. Note that $\Omega(|S|)$ operations are necessary in the worst case for any algorithm because for any group Γ , the two semigroups Γ^{1} and Γ^{0} (with an adjoined zero) have $G\Gamma^{1} = \Gamma$ but $G\Gamma^{0} = 0$, while sharing the subgroup Γ —every element must therefore be accessed in general.

Algorithm 6.1:	Computing	GS in O	$ S \log($	$ S)\alpha(S)$)) time.

Input: A nonempty finite semigroup $S = \{x_1, \ldots, x_{|S|}\}$ with operation ***Output:** A set $R \subset S$ of representatives for GS and a mapping $\rho: S \to R$ to the assigned representative Function groupCompletion(S, *): $I, H, J \coloneqq \texttt{getMinIdeal}(S, *);$ /* Use Algorithm 5.1 */ $e, \sigma \coloneqq \texttt{getIdentityAndInverses}(H, *);$ /* Use Algorithm 6.2 */ $(JI)^H \coloneqq \{h * j * i * \sigma(h) : i \in I, j \in J, h \in H\};$ $N \coloneqq \texttt{getGeneratedSubgroup}(H, *, e, (JI)^H);$ /* Use Algorithm 6.3 */ $H_{\text{remaining}} \coloneqq H; \ R \coloneqq \emptyset; \ \rho = \text{emptyMapping()};$ for each $h \in H$ do if $h \in H_{\text{remaining}}$ then $R \coloneqq R \cup \{h\};$ for each $n \in N$ do $H_{\text{remaining}} \coloneqq H_{\text{remaining}} \setminus \{h * n\};$ $\rho(h*n) \coloneqq h;$ foreach $x \in S$ do $\rho(x) \coloneqq \rho(e * x * e);$ return $R, \rho;$

 $^{^1\}mathrm{Thanks}$ to Alex Li for the idea of a linear-time version Algorithm 6.2

Algorithm 6.2: Finding the identity and inverses of a group O(|H|) time.

Input: A finite group H with operation * Output: The identity $e \in H$ and a mapping $\sigma: H \to H$ where $\sigma(h) = h^{-1}$ Function getIdentityAndInverses(H, *): $\{e\} := \{h \in H : h * h = h\};$ $\sigma := emptyMapping();$ $\sigma(e) := e;$ foreach $h \in H$ do C := [e, h]; // 0-indexed list while C.last is not in the domain of σ do [C.append(C.last * h);for $\ell \in \{1, \dots, C.length - 2\}$ do $[\sigma(C[\ell]) := \sigma(C.last) * C[C.length - \ell];$ return $e, \sigma;$

Algorithm 6.3: Finding a generated subgroup in $O(|H| \log |H| \alpha(|H|))$ time.

Input: A finite group H with operation * and identity e, and a subset $X \subseteq H$ Output: The generated subgroup $\langle X \rangle \leq H$ Function getGeneratedSubgroup(H, *, e, X): u := InitializeDisjointSetForest(<math>H); /* See [CLRS], Chapter 21.3 */ foreach $x \in X$ do if not u.sameComponent(e, x) then foreach $h \in H$ do u.mergeComponents(h, h * x); return { $x \in X : u.sameComponent(x, e)$ };

7. K-THIN SEMIGROUPS HAVE THE CLASSIFYING SPACE OF A GROUP

As pointed out by [AR67] and [Nic69], certain isomorphism types for K(S) give strong conditions on homological properties of S.

Definition 7.1. Call a finite nonempty semigroup S K-thin if its minimal ideal K(S) has a normalized Rees matrix semigroup structure $\mathcal{M}(H; I, J; [])$ in which |I| = 1 or |J| = 1. Equivalently, S is K-thin when K(S) is left-simple or right-simple.

The smallest nonempty semigroup that is not K-thin is Rect_2^2 . Algorithm 5.1 can be used to determine whether S is K-thin in O(|S|) time. If S is K-thin, then by the normalization of [[]], we know the subgroup $N \leq H$ of Theorem 6.3 is trivial, and so the maximal subgroups H of K(S) are then isomorphic to GS.

Proposition 7.2. A finite nonempty semigroup S has a left zero or a right zero if and only if S is K-thin and GS is trivial.

Proof. If $z \in S$ is a left zero then $zS^{1} = \{z\}$ is necessarily a minimal right ideal, which by Theorem 5.4 must have the form $\{z\} = \{i\} \times H \times J$, so H and J are singletons and GS = H = 1. Conversely, if $K(S) \cong \mathcal{M}(1; 1, J; [])$ then any singleton $\{(1, 1, j)\} \subseteq K(S)$ is a minimal right ideal, so (1, 1, j)x = (1, 1, j) for all

 $x \in S$, and so (1,1,j) is a left zero. The symmetric argument applies for right zeros.

The following is thus a generalization of Proposition 4.6:

Theorem 7.3. If S is K-thin then $BS \simeq BGS$.

The above theorem and required lemma below are the topological versions of, and follow from, results of [AR67] and [Nic69]; see Remark 7.6.

Lemma 7.4. If S is a semigroup with some idempotent e, and the subsemigroup eSe satisfies either eSe = Se or eSe = eS, then $BS \simeq B(eSe)$.

Proof. By replacing the operation on S with $(x, y) \mapsto yx$ as necessary, assume without loss of generality that eSe = eS. Write T := eSe, and note that e is an identity for T, so exe = ex for all $x \in S$. Define $r: S \to T$ by r(x) = ex. The map r is a semigroup homomorphism because exey = exy. Write $\iota: T \to$ S for the inclusion, so then $r \circ \iota = \operatorname{id}_T$. We will show that ι and r define a homotopy equivalence. Since $B_{\mathsf{Sgrp}}S \cong B_{\mathsf{Mon}}S^{\mathbb{1}}$ is a natural homeomorphism, it suffices to show that $r^{\mathbb{1}}: S^{\mathbb{1}} \to T^{\mathbb{1}}$ and $\iota^{\mathbb{1}}: T^{\mathbb{1}} \to S^{\mathbb{1}}$ define a homotopy equivalence $Br^{\mathbb{1}}: BS^{\mathbb{1}} \to BT^{\mathbb{1}}$ and $B\iota^{\mathbb{1}}: BT^{\mathbb{1}} \to BS^{\mathbb{1}}$.

Interpreting $T^{\mathbb{1}}$ and $S^{\mathbb{1}}$ as categories with a single object, we can provide natural transformations as homotopies. the first composition is $r^{\mathbb{1}} \circ \iota^{\mathbb{1}} = (r \circ \iota)^{\mathbb{1}} = \mathrm{id}_{T}^{\mathbb{1}} = \mathrm{id}_{T}^{\mathbb{1}}$ on the nose. For the second, define a natural transformation $\eta \colon \mathrm{id}_{S^{\mathbb{1}}} \Rightarrow \iota^{\mathbb{1}} \circ r^{\mathbb{1}}$ by taking unique component $\eta_* = e$. The naturality square for each $x \in S^{\mathbb{1}}$ is:

$$\begin{array}{ccc} & \ast & \stackrel{\eta_*=e}{\longrightarrow} & \ast \\ \mathrm{id}_{S^1} x=x & & \downarrow & \downarrow_{\iota^1 r^1 x} \\ & \ast & \stackrel{\eta_*=e}{\longrightarrow} & \ast \end{array}$$

This commutes because e1 = 1e for x = 1 and ex = exe for $x \in S$.

Proof of Theorem 7.3. Without loss of generality assume K(S) has a normalized Rees matrix semigroup structure with $J = \{j_0\}$. Then the Structure Theorem implies that we have a minimal right ideal

$$(i_0, e, j_0)S = \{i_0\} \times H \times J = \{i_0\} \times H \times \{j_0\},\$$

so $(i_0, e, j_0)S(i_0, e, j_0) = (i_0, e, j_0)S$. Since this maximal subgroup of K(S) is isomorphic to H, the lemma gives that $BS \simeq BH$. Finally, the subgroup $N \trianglelefteq H$ is trivial by the normalization of [[]], so GS = H/N = H.

Remark 7.5. The proof of Theorem 7.3 extends to infinite semigroups S so long as S has a minimal ideal isomorphic to a Rees matrix semigroup. Various hypotheses could guarantee this—for example, [HS12] gives the Structure Theorem for when S has a minimal ideal containing an idempotent.

Remark 7.6. The following purely-algebraic version of Lemma 7.4 is proved as Corollary 3 in [AR67], and is also proved in [Nic69]: for any semigroup S with some idempotent e and left ideal Se = eSe, for any S-module D we have

$$H^n(S,D) = H^n(Se,D) = H^n(Se,eD),$$

where $H^n(S, D) := \operatorname{Ext}_{\mathbb{Z}S}^n(\mathbb{Z}, D)$. If D is a GS-module, then by an analogous argument to Theorem A.1, we have $H^n(S, D) = H^n(BS; D)$, but the algebraic version allows S-modules that are not GS-modules. In fact, we can deduce Lemma 7.4

from the above by arguing that if Se = eSe then GS = G(eSe) and therefore for any local coefficient system (i.e. GS-module) D we have

$$H^n(BS; D) \cong H^n(S, D) \cong H^n(eSe, D) \cong H^n(B(eSe); D),$$

and so $r: S \rightarrow eSe$ induces isomorphisms on (co)homology with all local coefficient systems. [AR67] and [Nic69] also deduce a corresponding S-module strengthening of Theorem 7.3. However, our proofs of Lemma 7.4 and Theorem 7.3 have the advantage of explicitly providing the homotopies and do not require any mention of Tor or Ext functors.

There are many properties that imply that a finite semigroup S is K-thin, so then BS will be a K(GS, 1). For example:

Proposition 7.7. Finite commutative semigroups are K-thin.

Proof. Taking an arbitrary pair of elements of the minimum ideal K, we have

$$(i, h[[j, i']]h', j') = (i, h, j)(i', h', j') = (i', h', j')(i, h, j) = (i', h'[[j', i]]h, j),$$

so $i = i'$ and $j = j'$, and so $|I| = |J| = 1.$

It is already known for general commutative semigroups S that $BS \simeq BGS$ (see [Fie84], Remark 4.3), but the above is a fast proof for the finite case that easily generalizes:

Proposition 7.8. Finite left-commutative semigroups (i.e. those semigroups satisfying xya = yxa for all x, y, a) are K-thin.

Proof. Extending the above and taking a third element $a = (i'', h'', j'') \in K$, we see that

$$(i, h[[j, i']], j')(i'', h'', j'') = (i', h'[[j', i]]h, j)(i'', h'', j''),$$

so multiplying out gives $i = i'$, so $|I| = 1$.

Theorem 7.9. Generalizing the previous two propositions, suppose some finite semigroup S satisfies some equation $x_{\lambda_1} \cdots x_{\lambda_\ell} = x_{\lambda'_1} \cdots x_{\lambda'_{\ell'}}$ over some set of variables $\{x_1, x_2, x_3, \ldots\}$ (for example xy = yx defining commutativity or xya = yxa defining left-commutativity) for all substitutions of elements of S into the variables. If $\lambda_1 \neq \lambda'_1$ then S is K-thin. Dually, if $\lambda_\ell \neq \lambda'_{\ell'}$ then S is K-thin.

Proof. Substituting $x_{\lambda_1} \coloneqq (i, e, j)$ and $x_{\lambda'_1} \coloneqq (i', e, j')$, the product $x_{\lambda_1} \cdots x_{\lambda_{\ell}}$ evaluates to some element of $\{i\} \times H \times J$, while the product $x_{\lambda'_1} \cdots x_{\lambda'_{\ell'}}$ evaluates to an element of $\{i'\} \times H \times J$. Because the products are equal, i = i', so |I| = 1. The dual proof applies for the dual statement.

Proposition 7.10. Finite left-cancellative semigroups are K-thin.

Proof. For $i, i' \in I$ we have $(i_0, e, j_0)(i, e, j_0) = (i_0, e, j_0) = (i_0, e, j_0)(i', e, j_0)$, so we can cancel to see that $(i, e, j_0) = (i', e, j_0)$, and so i = i'.

Proposition 7.11. Finite *inverse semigroups* (see [CP61], Section 1.9) are K-thin.

Proof. By Corollary 2.19 in [CP61], each Green's \mathcal{L} -class and each Green's \mathcal{R} -class of an inverse semigroup must have exactly one idempotent. The minimal left ideals $I \times H \times \{j\}$ are Green's \mathcal{L} -classes, but each $(i, [j, i]^{-1}, j)$ is an idempotent, so we must have |I| = 1 (and likewise |J| = 1).

Order	Number not K -thin	Percentage K -thin
1-3	0	100%
4	1	99.21%
5	4	99.66%
6	48	99.70%
7	614	99.927%
8	10865	99.99941%
9	229613	99.999999567%
10	6257785	99.999999999950%

FIGURE 2. The numbers of semigroups (up to (anti-)isomorphism) of each order that are not K-thin based on an exhaustive search, and the percentage of all semigroups that are K-thin, from dividing by the totals in [DM24; Dis+12] and subtracting from 1.

The prevalence of semigroups in Appendix B whose classifying space has the same homology as some group, such as $[H_1(C_k), H_2(C_k), \ldots] = [C_k, 0, C_k, 0, C_k, 0, \ldots]$, is largely explained by Theorem 7.3. Exhaustive search for non-K-thin semigroups using the Minion solver [GJM06], splitting cases based on the isomorphism type of K(S) and using the symmetry-breaking techniques of [Dis10], gave the counts in Figure 2, which shows that the vast majority of known semigroups are K-thin.

8. DISCUSSION FROM HOMOLOGY COMPUTATIONS

Appendix B lists some homology groups for the classifying space of each semigroup with at most 8 elements. The present section describes some results learned during these computations and extensions to more elements. We will freely use projective resolutions of the form described in Appendix A to compute $H_*(BS;\mathbb{Z}) =$ $\operatorname{Tor}_*^{\mathbb{Z}S^1}(\mathbb{Z},\mathbb{Z})$.

Example 8.1. The converse of Proposition 4.6 does not hold: consider the 5element semigroup $S = \{x_{11}, x_{12}, x_{21}, x_{22}, q\}$ with operation defined by $x_{ij}x_{i'j'} = x_{ij'}$ and $x_{ij}q = x_{i1}$ and $qx_{ij} = x_{ij}$ and qq = q. The classifying space BS is contractible, but S does not have a zero element. This is the unique (up to isomorphism or anti-isomorphism) smallest such semigroup.

Proof. Take the following projective resolution of the form described in Appendix A:

$$\mathbb{Z} \stackrel{\varepsilon}{\leftarrow} \mathbb{Z}S^{\mathbb{1}}x_{11} \stackrel{\cdot (x_{21} - x_{11})}{\leftarrow} \mathbb{Z}S^{\mathbb{1}} \stackrel{\cdot 1}{\leftarrow} \mathbb{Z}S^{\mathbb{1}}q \leftarrow 0$$

Removing the augmentation and tensoring with \mathbb{Z} gives $\mathbb{Z} \stackrel{0}{\leftarrow} \mathbb{Z} \stackrel{1}{\leftarrow} \mathbb{Z} \leftarrow 0$, which has the homology of a point. The group completion GS is trivial, so BS is a simply connected CW complex with trivial homology.

Example 8.2. The semigroup $S = \{x_{11}, x_{12}, x_{21}, x_{22}, q, r\}$ with multiplication defined by $x_{ij}x_{i'j'} = x_{ij'}$ and $x_{ij}q = x_{ij}r = x_{i1}$ and $qx_{ij} = rx_{ij} = x_{ij}$ and qq = qr = q and rr = rq = r (extending Example 8.1) has $BS \simeq \mathbb{S}^3$.

Proof. Take the following projective resolution of the form described in Appendix A:

$$\mathbb{Z} \stackrel{\varepsilon}{\leftarrow} \mathbb{Z}S^{\mathbb{1}}x_{11} \stackrel{\cdot (x_{21}-x_{11})}{\longleftarrow} \mathbb{Z}S^{\mathbb{1}}q \stackrel{(1 \quad r-q)}{\longleftarrow} \frac{\mathbb{Z}S^{\mathbb{1}}x_{11}}{\mathbb{Z}S^{\mathbb{1}}} \stackrel{\begin{pmatrix} 0 & 0\\ 1 & 1 \end{pmatrix}}{\bigoplus} \frac{\mathbb{Z}S^{\mathbb{1}}x_{12}}{\mathbb{Z}S^{\mathbb{1}}q} \stackrel{(0)}{\leftarrow} 0$$

The matrices represent maps φ as constructed in Lemma A.3. Removing the augmentation and tensoring with Z gives

$$\mathbb{Z} \stackrel{0}{\leftarrow} \mathbb{Z} \stackrel{(1 \quad 0)}{\longleftarrow} \mathbb{Z}^2 \stackrel{(0 \quad 0)}{\longleftarrow} \mathbb{Z}^2 \leftarrow 0,$$

which has the homology of \mathbb{S}^3 . Since GS is trivial, BS is simply connected, so by the homotopy-uniqueness of Moore spaces (see [Hat02], Example 4.34), we in fact have a homotopy equivalence.

Remark 8.3. Much is known about the (co)homology of (the classifying spaces of) finite groups. For example, Corollary 10.2 of [Bro82] gives that multiplication by the order |G| of a finite group annihilates the positive-dimensional cohomology—and therefore homology—of its classifying space. In particular $H_i(BG; \mathbb{Z})$ is a finite abelian group for all $i \ge 1$. Furthermore, a theorem of Swan [Swa60] gives that the classifying space of a nontrivial finite group has infinitely many nontrivial homology groups.

The appearance of $H_3(BS) = \mathbb{Z}$ in Example 8.2 showed a deviation from the above patterns that hold for group homology. Another example is given by rectangular bands:

Proof of Proposition 3.6. The following proof generalizes the argument in [Fie02]. We first show that the homology of $B \operatorname{Rect}_b^a$ is the homology of $\bigvee_{i=1}^{(a-1)(b-1)} \mathbb{S}^2$. Take the projective resolution

$$\mathbb{Z} \stackrel{\varepsilon}{\leftarrow} \mathbb{Z}S^{\mathbb{I}}x_{11} \stackrel{(x_{21} - x_{11} \cdots x_{a1} - x_{11})}{\underbrace{(1 \cdots 1}_{i=2} \cdots \underbrace{(1 \cdots 1}_{i=2} \cdots \underbrace{(1 \cdots 1}_{j=1} \cdots \underbrace{(1 \cdots 1}_{i=2} \cdots \underbrace{(1 \cdots 1}_{j=1} \cdots \underbrace{(1 \cdots 1}_{$$

The matrices represent maps φ as constructed in Lemma A.3, where the (a-1)-by-b matrix of of zeros and ones represents adding together the *j*-summands within each *i*-summand. Removing the augmentation and tensoring with \mathbb{Z} gives

$$\mathbb{Z} \stackrel{0}{\leftarrow} \mathbb{Z}^{a-1} \xleftarrow{\begin{pmatrix} 1 & \cdots & 1 & & \\ & & 1 & \cdots & 1 & \\ & & & & \ddots \end{pmatrix}}{\longleftarrow} \mathbb{Z}^{(a-1)b} \leftarrow 0,$$

which has the homology of $\bigvee \mathbb{S}^2$. Since GS is trivial, BS is simply connected, so by the uniqueness of Moore spaces we have a homotopy equivalence.

Proposition 3.6 is greatly generalized by [Ste24], where $H_i(B\mathcal{M}(G; I, J; [[]))$ is computed for any Rees matrix semigroup.

It appears that Example 8.2 and Proposition 3.6 can be generalized to find finite S with $BS \simeq \bigvee_{k=1}^{\ell} \mathbb{S}^n$ for arbitrary $n \ge 2$ and $\ell \ge 1$ (see also Corollary 9.4). Adjoining units and taking products of these provides examples of free abelian integral homology groups, nonzero only in finitely many dimensions. Generalizing from this, a semigroup in which all elements are idempotent is called a *band*, and it was verified that all bands of order at most 11 have classifying spaces with free abelian homology groups that vanish in sufficiently high dimensions.

Question 8.4. Do the following hold for every finite band S?

- (a) $H_i(BS;\mathbb{Z})$ is free abelian for all *i*.
- (b) $H_i(BS;\mathbb{Z})$ vanishes for all sufficiently large *i*.
- (c) If S is a monoid then S has a finite projective resolution, i.e., S has homological type **FP**.

Certain cases of Question 8.4(b) are answered by [Nic69], particularly when the Green's \mathcal{J} -classes of S are linearly ordered.

The introduction of [Nic69] conjectures that homology is determined in high enough dimensions by the subgroups of S, and in particular that $H_i(BS; \mathbb{Z})$ vanishes for large i if S has no nontrivial subgroups. The following is a counterexample:

Example 8.5. Let S be the 5-element semigroup on the set $\{x_{11}, x_{12}, x_{21}, x_{22}, y_{11}\}$ with $x_{ij}x_{i'j'} \coloneqq x_{ij'}$ and $y_{11}x_{ij} = x_{1j}$ and $x_{ij}y_{11} = x_{i1}$ and $y_{11}y_{11} = x_{11}$. Then $H_1(BS) = 0$, but $H_i(BS) = \mathbb{Z}$ for all $i \ge 2$.

Proof. Take the projective resolution

$$\mathbb{Z} \stackrel{\varepsilon}{\leftarrow} \mathbb{Z}S^{\mathbb{1}}x_{11} \stackrel{\cdot (x_{21}-x_{11})}{\leftarrow} \mathbb{Z}S^{\mathbb{1}} \stackrel{(y_{11} \quad x_{12})}{\leftarrow} \mathbb{Z}S^{\mathbb{1}} \oplus \mathbb{Z}S^{\mathbb{1}}x_{12}$$

$$\stackrel{(y_{11}-x_{11} \quad x_{12}-x_{11})}{\leftarrow} \stackrel{(y_{11} \quad x_{12})}{\leftarrow} \mathbb{Z}S^{\mathbb{1}} \oplus \mathbb{Z}S^{\mathbb{1}}x_{12}$$

where the last two modules and maps alternate indefinitely. Removing the augmentation and tensoring with \mathbb{Z} results in

$$\mathbb{Z} \stackrel{0}{\leftarrow} \mathbb{Z} \stackrel{(1 \quad 1)}{\longleftarrow} \mathbb{Z}^2 \stackrel{0}{\leftarrow} \mathbb{Z}^2 \stackrel{(1 \quad 1)}{\longleftarrow} \mathbb{Z}^2 \stackrel{0}{\leftarrow} \mathbb{Z}^2 \stackrel{(1 \quad 1)}{\longleftarrow} \mathbb{Z}^2 \stackrel$$

which has the claimed homology.

The semigroup in Example 8.5 is not *regular* because not every element $y \in S$ has some $x \in S$ satisfying yxy = y—the element y_{11} in particular is not regular because it is not the result of any multiplication. It seems that non-regularity might be necessary to have infinitely many \mathbb{Z} s:

Question 8.6. Does every finite regular semigroup S have $H_i(BS;\mathbb{Z})$ finite for all sufficiently large i?

	0	1	2	3	4	5	6	7	8
0	0	1	0 0	1	0	0	0	0	0
1	0	1	0	1	0	0	0	1	1
2	2	3	2	3	2	2	2	2	2
3	2	3	2	3	2	2	2	3	3
4	0	1	0	1	0	4	4	0	0
5	0	1	2	3	0	5	5	0	0
6	0	1	0 2 2 0 2 2 2 2	3	0	6	6	0	0
7	0	1	0	1	4	4	0	$\overline{7}$	8
8	0	1		1	4	0	4	7	8

FIGURE 3. The multiplication table for a semigroup S where BS is a Moore space of type $M(C_2, 3)$.

This question is answered affirmatively in certain cases by [Nic69].

Turning from the free part of integral homology to the torsion parts and looking at the data in Appendix B, one sees that for semigroups S with at most 8 elements, if $H_i(BS;\mathbb{Z})$ has nontrivial torsion for some *i* then this is the case for more than one *i*, likely infinitely many as is the case for groups (Remark 8.3). This fails for a certain semigroup with 9 elements:

Example 8.7. Let S be a semigroup on the 9 elements $\{x_0, \ldots, x_8\}$ where the operation $x_i x_j = x_k$ is recorded by an integer k in entry j of row i in the table in Figure 3. Then $H_3(BS) = C_2$ and $\tilde{H}_i(BS) = 0$ for $i \neq 3$. In fact, $BS \simeq \Sigma^2(\mathbb{RP}^2)$, the twofold suspension of the real projective plane.

Proof. Take the projective resolution

$$\mathbb{Z} \stackrel{\varepsilon}{\leftarrow} \mathbb{Z}S^{1}x_{0} \stackrel{\leftarrow (x_{2}-x_{0})}{\leftarrow} \mathbb{Z}S^{1}x_{5} \stackrel{\leftarrow (x_{4} - x_{6} - x_{5})}{\leftarrow} \mathbb{Z}S^{1}x_{7} \\ \xrightarrow{\begin{pmatrix} 0 & 0 & x_{7} - x_{0} & 0 \\ 1 & 1 & x_{7} & x_{7} + x_{8} \end{pmatrix}}_{\leftarrow} \mathbb{Z}S^{1}x_{5} \\ \xrightarrow{\oplus} \mathbb{Z}S^{1}x_{7} \\ \oplus \mathbb{Z}S^{1}x_{7} \\ \oplus \mathbb{Z}S^{1}x_{7} \\ \oplus \mathbb{Z}S^{1}x_{1} \\ \leftarrow \\ \xrightarrow{\begin{pmatrix} -1 & -2 & 0 \\ 0 & 0 & -2 \\ 1 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix}}_{\leftarrow} \mathbb{Z}S^{1}x_{0} \\ \xrightarrow{\oplus} \mathbb{Z}S^{1}x_{1} \\ \oplus \mathbb{Z}S^{1}x_{1} \\ \oplus \mathbb{Z}S^{1}x_{1} \\ \leftarrow \\ \end{bmatrix}$$

Tensoring with \mathbbm{Z} and taking homology gives

$$H_{3}(BS) = \frac{\ker(\mathbb{Z}^{4} \to \mathbb{Z}^{2})}{\operatorname{im}(\mathbb{Z}^{3} \to \mathbb{Z}^{4})} = \frac{\mathbb{Z}\left\langle (-1, 0, 1, 0), (-1, 1, 0, 0), (-2, 0, 0, 1) \right\rangle}{\mathbb{Z}\left\langle (-1, 0, 1, 0), (-2, 0, 0, 1), (0, -2, 0, 1) \right\rangle} \\ = \frac{\mathbb{Z}\left\langle (-1, 0, 1, 0), (-1, 1, 0, 0), (-2, 0, 0, 1) \right\rangle}{\mathbb{Z}\left\langle (-1, 0, 1, 0), (-2, 2, 0, 0), (-2, 0, 0, 1) \right\rangle} \cong C_{2},$$

and so BS has the desired homology. The group completion GS is trivial because the minimal ideal $K(S) = \{x_0, x_1, x_2, x_3\}$ has trivial group completion, so BS is simply connected, and so BS is a Moore space of type $M(C_2, 3)$. The result follows from the uniqueness of Moore spaces.

	0	1	2	3	4	5	6	$\overline{7}$	8	9	10	11
0	0	0	0	0	4	4	4	4	0	0	4	4
1	1	1	1	1	5	5	5	5	1	1	5	5
2	2	2	2	2	6	6	6	6	2	2	6	6
3	3	3	3	3	7	7	7	7	3	3	$\overline{7}$	7
4	0	0	0	0	4	4	4	4	4	4	0	0
5	1	1	1	1	5	5	5	5	5	5	1	1
6	2	2	2	2	6	6	6	6	6	6	2	2
$\overline{7}$	3	3	3	3	7	7	7	7	7	7	3	3
8	0	3	2	3	4	7	6	7	8	9	10	11
9	0	2	2	3	4	6	6	7	8	9	10	11
10	0	2	3	2	4	6	7	6	10	11	8	9
11	0	3	3	2	4	7	7	6	10	11	8	9

FIGURE 4. The multiplication table for a semigroup S where BS is a Moore space of type $M(C_2, 2)$.

Remark 8.8. For finite groups G, as implied by Remark 8.3, if $H_*(BG)$ has nontrivial *p*-torsion for some prime p, then p divides |G|, so G has a subgroup of order p. On the other hand, the semigroup S in Example 8.7 has 2-torsion in $H_3(BS)$, but S has no sub-group of order 2. In fact, Green's \mathcal{H} relation on S is trivial, so S has no nontrivial subgroups at all.

An $M(C_2, 2)$ is also possible, though this example has subgroups $\{x_8, x_{10}\}$ and $\{x_9, x_{11}\}$ of order 2:

Example 8.9. Let S be the semigroup on the 12 elements $\{x_0, \ldots, x_{11}\}$ where the operation $x_i x_j = x_k$ is recorded by an integer k in entry j of row i of the table in Figure 4. Then $BS \simeq \Sigma(\mathbb{RP}^2)$, the suspension of the real projective plane.

Proof. Take the projective resolution:

$$\mathbb{Z} \stackrel{\varepsilon}{\leftarrow} \mathbb{Z}S^{\mathbb{1}}x_0 \stackrel{\cdot(x_1-x_0)}{\leftarrow} \mathbb{Z}S^{\mathbb{1}} \stackrel{(x_0 \quad x_4 \quad x_{11}-x_8)}{\leftarrow} \mathbb{Z}S^{\mathbb{1}}x_0 \\ \stackrel{(x_0 \quad x_0)}{\leftarrow} \mathbb{Z}S^{\mathbb{1}}x_8 \\ \stackrel{(x_0 \quad -x_0)}{\leftarrow} \stackrel{(x_4 \quad x_0)}{\leftarrow} \mathbb{Z}S^{\mathbb{1}}x_0 \\ \stackrel{(x_0 \quad -x_0)}{\leftarrow} \mathbb{Z}S^{\mathbb{1}}x$$

Tensoring with \mathbb{Z} gives

$$\mathbb{Z} \xleftarrow{0}{\leftarrow} \mathbb{Z} \xleftarrow{\begin{pmatrix} 1 & 1 & 0 \end{pmatrix}}{\leftarrow} \mathbb{Z}^3 \xleftarrow{\begin{pmatrix} -1 & 1 \\ 1 & -1 \\ 1 & 1 \end{pmatrix}} \mathbb{Z}^2 \leftarrow 0,$$

which has the stated homology. The group completion GS is trivial because the minimal ideal $K(S) = \{x_0, \ldots, x_7\}$ has trivial group completion, so BS is simply connected, and so BS is a Moore space of type $M(C_2, 2)$. The result follows from the uniqueness of Moore spaces.

•	0	1			4	5	6	7	8	9
0	0	1	0	1	0	0	0	0	0	0
1	0	1	0	1	0	0	0	0	0	0
2			2	3	2	2	2	2	2	2
3	2	3	2	3	2	2	2	2	2	2
4	0	1	0	1	0	0	0	0	0	0
5	0	1	0	1	0	0	0	0	0	0
6	0	1	0	1	0	0	0	0	0	0
$\overline{7}$	0	1	0	1	0	0	0	4	4	5
8	0	1	0	1	0	0	0	6	5	6
9	0	1	0	1	0	0	0	0	6	4

FIGURE 5. The multiplication table for a semigroup S with $H_6(BS;\mathbb{Z}) \cong \mathbb{Z}^9 \times C_{1494640}$.

The failure of multiplication by |S| to annihilate $H_*(BS; \mathbb{Z})$ is even more extreme: we can find torsion of order $1494640 = 157 \cdot 17 \cdot 7 \cdot 5 \cdot 2^4$ from a semigroup of order only 10:

Example 8.10. The semigroup S defined by Figure 5 has first several integral homology groups $H_i = H_i(BS; \mathbb{Z})$ as follows:

 $[H_1, H_2, H_3, H_4, H_5, H_6, H_7] = [0, \mathbb{Z}, \mathbb{Z}^3, \mathbb{Z}^6, \mathbb{Z}^9, \mathbb{Z}^9 \times C_{1494640}, \mathbb{Z}^{27} \times C_{17}].$

The computer calculation following Algorithm ${\rm A.1}$ is omitted.

9. Joins and Suspensions

In the table of 6-element semigroups in Appendix B, one can notice that the entry $[H_1, H_2, H_3, \ldots] = [0, C_2, 0, C_2, 0, C_2, \ldots]$ describes a semigroup S with 6 elements consistent with $BS \simeq \Sigma(BC_2) \cong \Sigma(\mathbb{R}P^{\infty})$, where Σ is suspension. This equivalence in fact holds, and we will now show that the set of homotopy types of classifying spaces of finite monoids (or semigroups) is closed under suspension.

Definition 9.1. For a monoid S and a set Y, define a monoid $J^Y(S)$ on the (1 + |Y|)|S| symbols $S \sqcup \{yx : y \in Y, x \in S\}$, with operation "." defined by

For $y \in Y$, identify $y \coloneqq y \in J^Y(S)$, so $Y \subseteq J^Y(S)$ is a subsemigroup with yy' = y.

Note that $J^{Y}(S)$ is the union of the monoid S with a |Y|-by-|S| rectangular band YS, where S acts as the identity on the left of YS and acts as multiplication in S on the right of YS. This section will prove the following:

Theorem 9.2. For a monoid S and set Y, the monoid $J^Y(S)$ has classifying space $B(J^Y(S)) \simeq BS * Y^{\text{discrete}}$, the topological join of BS with the discrete set Y.

From this, the following corollaries are immediate:

Corollary 9.3. For nonempty Y and arbitrary $y_0 \in Y$,

$$B(J^{Y}(S)) \simeq \bigvee_{\substack{y \in Y \setminus \{y_0\}\\18}} \Sigma(BS).$$

Proof. Working topologically from the result of the theorem,

$$BS * Y^{\text{discrete}} \simeq \Sigma \Big(BS \wedge Y^{\text{discrete}} \Big) \simeq \Sigma \left(\bigvee_{y \in Y \setminus \{y_0\}} BS \right) \cong \bigvee_{y \in Y \setminus \{y_0\}} \Sigma (BS). \quad \Box$$

Corollary 9.4. $B(J^{\{1,2\}}(S)) \simeq \Sigma(BS)$, so the set of homotopy types of classifying spaces of finite monoids is closed under suspension.

Remark 9.5. Finite groups G have finitely generated cohomology rings $H^*(BG)$ [Eve61]. On the other hand, taking $S = J^{\{1,2\}}(C_2)$, we know BC_2 and therefore BS have nonzero cohomology in infinitely many dimensions, while the cup products on BS are trivial. Thus the cohomology ring $H^*(BS)$ is not finitely generated.

To prove Theorem 9.2, we use the collapsing scheme defined in [Bro92] for the nerve of a monoid with a complete rewriting system. See [BO93] for more information on string rewriting systems.

Proposition 9.6. $J^Y(S)$ is a monoid with a complete rewriting system on the alphabet $S \sqcup Y$ with rewriting rules

$xx' \to x''$	for each $xx' = x''$ in S ,
$xy \rightarrow y$	for each $x \in S, y \in Y$, and
$yy' \to y$	for each $y, y' \in Y$.

Proof. There are no infinite chains of reductions because each rule reduces the length of a string. The system is confluent because the critical pairs resolve:

- $(xx')y \to x''y \to y$ agrees with $x(x'y) \to xy \to y$.
- $(xy)y' \to yy' \to y$ agrees with $x(yy') \to xy \to y$.
- Critical pairs only in S or only in Y certainly resolve.

The irreducible words take the form x, y, or yx for any $x \in S$ or $y \in Y$, and the rewriting rules give exactly the operation on $J^Y(S)$.

Lemma 9.7. There is a homotopy equivalence $B(J^Y(S)) \to Q$, where Q is the subcomplex of $B(J^Y(S))$ consisting of cells of the form $(x_1, \ldots, x_k, y_1, \ldots, y_\ell)$ for $k, \ell \ge 0$.

Proof. Using the complete rewriting system from Proposition 9.6, The collapsing scheme of [Bro92] defines a homotopy equivalence from $B(J^Y(S))$ onto a certain quotient complex with a cell for each *essential* cell of $B(J^Y(S))$. Brown's essential cells are tuples (w_1, \ldots, w_n) of irreducible words over the alphabet, with the first entry w_1 a single letter, the concatenation $w_i w_{i+1}$ of any pair of adjacent entries reducible, and no proper prefix of any $w_i w_{i+1}$ reducible.

For our system, the essential cells cannot include (\ldots, x, yx', \ldots) and cannot include $(\ldots, y, y'x', \ldots)$ because xy and yy' are reducible proper prefixes of the concatenations. Thus, because these longer irreducible words cannot follow a single-letter word, all entries of our essential cells are single letters. Because any yx is irreducible, our essential cells cannot include (\ldots, y, x, \ldots) either, so we are left with only the stated cells of Q.

Note that for arbitrary complete rewriting systems, the quotient complex is a more general CW complex. However, because our essential cells form a subcomplex of the simplicial set $B(J^Y(S))$, we can take Q to be the sub-simplicial-set. \Box

Recall that for simplicial sets X, X', regarded as augmented with a unique (-1)-cell \emptyset , the join has simplices

$$(X * X')^n \coloneqq \bigsqcup_{\substack{(a+1)+(b+1)=(n+1)\\a,b \ge -1}} X^a \times X'^b$$

written as $\sigma^a * \tau^b \in X^a \times X'^b$, with face maps

$$d_i^{a+b+1}(\sigma^a * \tau^b) \coloneqq \begin{cases} d_i^a \sigma^a * \tau^b & 0 \leqslant i \leqslant a \\ \sigma^a * d_{i-(a+1)}^b \tau^b & a+1 \leqslant i \leqslant a+b+1, \end{cases}$$

and degeneracy maps defined similarly.

Proof of Theorem 9.2. Using the simplicial set Q from Lemma 9.7, we show the following zigzag of equivalences:

$$BS * Y^{\text{discrete}} \xleftarrow[(1)]{} BS * \bigsqcup_{y \in Y} BY \xrightarrow[(2)]{} \frac{BS * \bigsqcup_{y \in Y} BY}{() * \bigsqcup_{y \in Y} BY} \cong \frac{Q}{(3)} \xrightarrow[(4)]{} Q \xleftarrow[(5)]{} B(J^Y(S)).$$

Here, $Y \subseteq J^Y(S)$ has operation yy' = y, so BY is contractible by Lemma 4.6, and so (4) is an equivalence. Joins respect equivalences, so (1) is an equivalence. A join with a single point () is a cone, and hence is contractible, so (2) is an equivalence. The map (5) is an equivalence by the lemma, so it remains to give the isomorphism (3) of simplicial sets. The correspondence (3) on nondegenerate cells is

the unique 0-cell \longleftrightarrow the unique 0-cell

$$[(x_1, \dots, x_k) * \emptyset] \longleftrightarrow [(x_1, \dots, x_k)] \qquad \text{for } k \ge 1$$

 $[(x_1,\ldots,x_k)*\iota_{y_0}(y_1,\ldots,y_\ell)]\longleftrightarrow [(x_1,\ldots,x_k,y_0,y_1,\ldots,y_\ell)] \quad \text{for } k \ge 1, \ell \ge 0.$

For the last case, the face map d_k is respected because $x_k y_0 = y_0$, and the face map d_{k+1} is respected because $y_0 y_1 = y_0$.

References

- [AR67] William W. Adams and Marc A. Rieffel. "Adjoint functors and derived functors with an application to the cohomology of semigroups". In: *Jour*nal of Algebra 7 (1967), pp. 25–34. DOI: 10.1016/0021-8693(67)90065-8.
- [BO93] Ronald V. Book and Friedrich Otto. "String-Rewriting Systems". In: Monographs in Computer Science (1993). DOI: 10.1007/978-1-4613-9771-7.
- [Bro82] Kenneth S. Brown. *Cohomology of Groups*. Graduate Texts in Mathematics. Springer New York, NY, 1982.
- [Bro92] Kenneth S. Brown. "The Geometry of Rewriting Systems: A Proof of the Anick-Groves-Squier Theorem". In: Algorithms and Classification in Combinatorial Group Theory. Springer New York, 1992, pp. 137–163.
 ISBN: 978-1-4613-9730-4. DOI: 10.1007/978-1-4613-9730-4_6.
- [CL24] Antonio M. Cegarra and Jonathan Leech. *The Cohomology of Monoids*. Springer Cham, 2024. DOI: 10.1007/978-3-031-50258-3.
- [CP61] A.H. Clifford and G.B. Preston. The Algebraic Theory of Semigroups, Volume I. Mathematical Surveys and Monographs. American Mathematical Society, 1961. ISBN: 9780821802717.

- [CLRS] Thomas H. Cormen et al. Introduction to Algorithms, Third Edition. The MIT Press, 2009. ISBN: 0262033844.
- [DM24] A. Distler and J. Mitchell. Smallsemi, A library of small semigroups, Version 0.7.1. GAP package. Aug. 2024. URL: https://gap-packages. github.io/smallsemi/.
- [Dis10] Andreas Distler. "Classification and enumeration of finite semigroups". In: Ph.D. Thesis. University of St Andrews Research Repository, 2010.
- [DM12] Andreas Distler and James D. Mitchell. "The number of nilpotent semigroups of degree 3". In: *The Electronic Journal of Combinatorics* 19 (2012). DOI: 10.37236/2441.
- [Dis+12] Andreas Distler et al. "The Semigroups of Order 10". In: Principles and Practice of Constraint Programming. Ed. by Michela Milano. Berlin, Heidelberg: Springer Berlin Heidelberg, 2012, pp. 883–899. ISBN: 978-3-642-33558-7.
- [Eve61] Leonard Evens. "The Cohomology Ring of a Finite Group". In: Transactions of the American Mathematical Society 101.2 (1961), pp. 224-239.
 ISSN: 00029947. URL: http://www.jstor.org/stable/1993372 (visited on 01/28/2025).
- [Fie84] Zbigniew Fiedorowicz. "Classifying Spaces of Topological Monoids and Categories". In: American Journal of Mathematics 106 (1984), p. 301.
- [Fie02] Zbigniew Fiedorowicz. "A counterexample to a group completion conjecture of J C Moore". In: Algebraic & Geometric Topology 2 (2002), pp. 33–35.
- [FP97] Véronique Froidure and Jean-Eric Pin. "Algorithms for computing finite semigroups". In: Foundations of Computational Mathematics. Springer Berlin Heidelberg, 1997, pp. 112–126. ISBN: 978-3-642-60539-0.
- [FH10] Jonathon Funk and Pieter Hofstra. "Topos theoretic aspects of semigroup actions". In: vol. 24. 6. 2010, pp. 117–147.
- [GAP] GAP—Groups, Algorithms, and Programming. The GAP Group. URL: https://www.gap-system.org.
- [GJM06] Ian P. Gent, Chris Jefferson, and Ian Miguel. "MINION: A Fast, Scalable, Constraint Solver". In: Proceedings of the 2006 Conference on ECAI 2006: 17th European Conference on Artificial Intelligence August 29 - September 1, 2006, Riva Del Garda, Italy. NLD: IOS Press, 2006, pp. 98–102. ISBN: 1586036424.
- [Hat02] Allen Hatcher. Algebraic topology. Cambridge Univ. Press, 2002.
- [HS12] Neil Hindman and Dona Strauss. Algebra in the Stone-Cech Compactification. De Gruyter, 2012. ISBN: 9783110258356. DOI: doi:10.1515/ 9783110258356.
- [HM03] Göran Högnäs and Arunava Mukherjea. "Maximal Homomorphic Group Image and Convergence of Convolution Sequences on a Semigroup". In: *Journal of Theoretical Probability* 16 (4 2003), pp. 847–854. DOI: 10. 1023/B:J0TP.0000011996.44195.2b.
- [Hol09] Christopher Hollings. "The Early Development of the Algebraic Theory of Semigroups". In: Archive for History of Exact Sciences 63.5 (2009), pp. 497-536. ISSN: 00039519, 14320657. URL: http://www.jstor.org/ stable/41134321.

- [How95] John M. Howie. Fundamentals of Semigroup Theory. Oxford University Press Inc., New York, 1995.
- [KT76] D.M. Kan and W.P. Thurston. "Every connected space has the homology of a K(π ,1)". In: *Topology* 15.3 (1976), pp. 253–258. ISSN: 0040-9383. DOI: 10.1016/0040-9383(76)90040-9.
- [KRS76] Daniel J. Kleitman, Bruce R. Rothschild, and Joel H. Spencer. "The Number of Semigroups of Order n". In: Proceedings of the American Mathematical Society 55.1 (1976), pp. 227–232. DOI: 10.2307/2041879.
- [Lar72] Jean Larisse. "Marches au hasard sur les demi-groupes discrets, II". In: Annales de l'institut Henri Poincaré, Section B: Calcul des probabilités et statistiques 8.2 (1972), pp. 127–173.
- [McD79] Dusa McDuff. "On the classifying spaces of discrete monoids". In: Topology 18.4 (1979), pp. 313–320. ISSN: 0040-9383. DOI: 10.1016/0040-9383(79)90022-3.
- [MS76] Dusa McDuff and Graeme Segal. "Homology fibrations and the 'groupcompletion' theorem". In: *Invent Math* 31 (1976), pp. 279–284. DOI: 10.1007/BF01403148.
- [Nic69] William R Nico. "On the cohomology of finite semigroups". In: Journal of Algebra 11.4 (1969), pp. 598–612. ISSN: 0021-8693. DOI: 10.1016/0021-8693(69)90093-3.
- [Ree40] D. Rees. "On semi-groups". In: Mathematical Proceedings of the Cambridge Philosophical Society 36.4 (1940), pp. 387–400. DOI: 10.1017/ S0305004100017436.
- [Seg74] Graeme Segal. "Categories and cohomology theories". In: *Topology* 13.3 (1974), pp. 293–312. ISSN: 0040-9383. DOI: 10.1016/0040-9383(74) 90022-6.
- [Ste24] Benjamin Steinberg. The homology of completely simple semigroups. 2024. arXiv: 2405.06594 [math.GR].
- [Sus28] Anton Suschkewitsch. "Über die endlichen Gruppen ohne das Gesetz der eindeutigen Umkehrbarkeit". In: *Mathematische Annalen* 99 (1928), pp. 30–50. DOI: doi:10.1007/BF01459084.
- [Swa60] Richard G. Swan. "The Nontrivality of the Restriction Map in the Cohomology of Groups". In: Proceedings of the American Mathematical Society 11.6 (1960), pp. 885–887. ISSN: 00029939, 10886826. DOI: 10. 2307/2034431.
- [Wei05] Michael Weiss. "What does the Classifying Space of a Category Classify?" In: *Homology, Homotopy and Applications* 7 (2005), pp. 185–195.

APPENDIX A. COMPUTING HOMOLOGY AS TOR

To do computations, we make use of the following fact:

Theorem A.1. For a monoid S with group completion $G = GS = \pi_1(BS)$, and for any G-module D, we have

$$H_i(BS; D) \cong \operatorname{Tor}_i^{\mathbb{Z}S}(\mathbb{Z}, D),$$

The left side is homology with local coefficients—the trivial *G*-module $D = \mathbb{Z}$ gives integral homology of *BS*. On the right side *D* is regarded as an *S*-module via the group completion map $p: S \to G$.

This fact is already understood, by [Fie02] for example, but we elaborate here for completeness. The expression $\operatorname{Tor}_{i}^{\mathbb{Z}S}(\mathbb{Z};D)$ (for general *S*-modules *D*) is called the Eilenberg-MacLane homology of *S*—see [CL24] for various generalizations.

Proof. We make use of the natural generalization of Definition 2.2 for categories, where *n*-simplices are *n*-tuples of composable morphisms. Let ES be the nerve of the category in which the objects are elements of S and the morphisms are $x_0 \xrightarrow{x_1} x_1 x_0$, with the natural composition and identities. Then ES is contractible because there is a natural transformation η : const₁ \Rightarrow id defined by $\eta_x = x$; naturality is

$$\operatorname{const}_{1} x_{1} = 1 \bigcup_{1} \begin{array}{c} & \stackrel{\eta_{x_{0}}}{\longrightarrow} & x_{0} \\ & \downarrow & \downarrow \operatorname{id} x_{1} = x_{1} \\ & 1 \xrightarrow{\eta_{x_{1}x_{0}}} & x_{1}x_{0} \end{array}$$

Recall that EG is the universal cover of BG, but ES is not generally the universal cover of BS because BS may not have contractible universal cover. To construct the universal cover, take a pullback:

$$\begin{array}{ccc} \tilde{BS} & & & EG \\ \downarrow & & \downarrow \\ BS & \xrightarrow{Bp} & BG \end{array}$$

The pullback simplicial set \tilde{BS} has as its *n*-simplices length-*n* chains

$$g_0 \xrightarrow{x_1} p(x_1)g_0 \xrightarrow{x_2} \cdots \xrightarrow{x_n} p(x_n \cdots x_2 x_1)g_0,$$

with the natural boundaries and degeneracies, where $x_i \in S$ and $g_0 \in G$. We now show the following:

$$H_*(BS;D) \cong H\left(C_*\left(\tilde{BS}\right) \underset{\mathbb{Z}G}{\otimes} D\right) \cong H\left(C_*(ES) \underset{\mathbb{Z}S}{\otimes} D\right) \cong \operatorname{Tor}_*^{\mathbb{Z}S}(\mathbb{Z},D).$$

Here C_* means the simplicial chain complex, and H means homology. An element $\gamma \in G$ acts on $C_*(\tilde{BS})$ by replacing g_0 with $g_0\gamma$. The first isomorphism above is a definition of local coefficients. For the third isomorphism, the complex $C_*(ES)$ is a resolution of \mathbb{Z} because ES is contractible, and $C_*(ES)$ is free as a right $\mathbb{Z}S$ -module because $C_n(ES) \cong \bigoplus_{x_1,\ldots,x_n} \mathbb{Z}S$. The middle isomorphism is due to the fact that we can write any pure tensor of $C_n(\tilde{BS}) \otimes_G D$ as

$$\left(g_0 \xrightarrow{x_1} \cdots \xrightarrow{x_n} p(x_n \cdots x_1)g_0\right) \otimes_G d = \left(1 \xrightarrow{x_1} \cdots \xrightarrow{x_n} p(x_n \cdots x_1)\right) \otimes_G g_0 d,$$

and we can write any pure tensor of $C_*(ES) \otimes_S D$ as

$$\left(x_0 \xrightarrow{x_1} \cdots \xrightarrow{x_n} x_n \cdots x_1 x_0\right) \otimes_S d = \left(1 \xrightarrow{x_1} \cdots \xrightarrow{x_n} x_n \cdots x_1\right) \otimes_S p(x_0) d,$$

so we have a bijection between the pure tensors.

Computing Tor will involve finding projective resolutions of \mathbb{Z} , so it will be convenient to have a collection of projective left $\mathbb{Z}S$ modules to choose from.

Proposition A.2. If S is a monoid with some idempotent e, then $\mathbb{Z}Se$ is a projective left $\mathbb{Z}S$ module.

Proof. Since e is an idempotent of the ring $\mathbb{Z}S$, we can decompose $\mathbb{Z}S = \mathbb{Z}Se \oplus \mathbb{Z}S(1-e)$ to see $\mathbb{Z}Se$ as a direct summand of the free module $\mathbb{Z}S$. \Box

It will also be helpful to have a collection of $\mathbb{Z}S$ -linear maps, and to understand their interaction with the functor $\mathbb{Z} \bigotimes_{\mathbb{Z}S} -$ to compute Tor.

Lemma A.3. Let S be a monoid. Let $e_1, \ldots, e_n, f_1, \ldots, f_m \in S$ be idempotents. For $(i, j) \in \{1, \ldots, m\} \times \{1, \ldots, n\}$, suppose $a_j^i \in \mathbb{Z}S$, and suppose that $e_j a_j^i \in \mathbb{Z}Sf_i$ for all i, j. Define the map

$$\varphi \colon \bigoplus_{j=1}^n \mathbb{Z}Se_j \to \bigoplus_{i=1}^m \mathbb{Z}Sf_i$$

by $\pi_i(\varphi(\iota_j(x))) = xa_j^i$, where π_i is projection to summand i and ι_j is inclusion of summand j. Then φ is left $\mathbb{Z}S$ -linear. By applying the functor $\mathbb{Z} \otimes -$, we can identify the domain of φ with \mathbb{Z}^n , the codomain of φ with \mathbb{Z}^m , and the tensor product $\mathrm{id}_{\mathbb{Z}} \otimes \varphi$ with the map $\overline{\varphi} \colon \mathbb{Z}^n \to \mathbb{Z}^m$ defined by $\pi_i(\overline{\varphi}(\iota_j(x))) = x\varepsilon(a_j^i)$, where $\varepsilon \colon \mathbb{Z}S \to \mathbb{Z}$ is augmentation.

Proof. The $\mathbb{Z}S$ -linearity of φ follows from the associativity of S. Tensor products distribute over direct sums, so

$$\mathbb{Z} \underset{\mathbb{Z}S}{\otimes} \bigoplus_{i=1}^{n} \mathbb{Z}Se_{j} \cong \bigoplus_{i=1}^{n} \mathbb{Z} \underset{\mathbb{Z}S}{\otimes} \mathbb{Z}Se_{j} \cong \bigoplus_{i=1}^{n} \mathbb{Z},$$

where the first isomorphism is $1 \otimes \iota_i(x) \mapsto \iota_i(1 \otimes x)$, and the second isomorphism is $\iota_i(1 \otimes x) \mapsto \iota_i(1)$, which is an isomorphism because for $s \in Se_j$ we have $1 \otimes s = 1 \otimes se_j = 1 \otimes e_j$. By functoriality, we then know $\operatorname{id}_{\mathbb{Z}} \otimes (\pi_i \varphi \iota_j) = \pi_i \overline{\varphi} \iota_j$. so it remains to show that $\operatorname{id}_{\mathbb{Z}} \otimes \rho_{a_j^i} \cong \rho_{\varepsilon(a_j^i)}$, where ρ means right-multiplication. Indeed, any $1 \otimes x \in \mathbb{Z} \otimes \mathbb{Z}Se_j$ is sent to $1 \otimes xa_j^i = 1 \otimes xa_j^i f_i = \varepsilon(xa_j^i) \otimes f_i$, and so under the identifications to \mathbb{Z}^n and \mathbb{Z}^m , the map is multiplication by $\varepsilon(a_j^i)$. \Box

Algorithm A.1 computes $H_i(BS; \mathbb{Z})$ by finding a projective resolution with maps of the form in Lemma A.3, with several optimizations: (1) we apply Theorem 7.3 to reduce to group homology whenever possible, (2) we cache and re-use the modules and maps of a resolution to allow representation of an infinite resolution by a finite graph with cycles, and (3) we attempt to fork the resolution at each module into smaller resolutions to maximize the potential for re-use.

This sort of computation via projective resolutions was almost always faster at computing $H_i(BS)$ for finite S than any equivalent computations using complete rewriting systems and collapsing schemes as in [Bro92].

Example A.4. When computing $H_i(BC_k;\mathbb{Z})$, where $C_k = \langle a | a^k = 1 \rangle$, Algorithm A.1 finds the usual resolution:

$$\mathbb{Z} \stackrel{\varepsilon}{\leftarrow} \mathbb{Z}C_k \stackrel{a-1}{\longleftarrow} \mathbb{Z}C_k \stackrel{N}{\leftarrow} \mathbb{Z}C_k \stackrel{a-1}{\longleftarrow} \mathbb{Z}C_k \stackrel{N}{\leftarrow} \mathbb{Z}C_k \leftarrow \cdots$$

Here $N = 1 + a + \cdots + a^{k-1}$. Re-using parts of this infinite resolution allows it to be represented as a graph with a cycle:

$$\mathbb{Z} \stackrel{\varepsilon}{\leftarrow} \mathbb{Z}C_k \stackrel{a-1}{\longleftarrow} \mathbb{Z}C_k \stackrel{N}{\leftarrow} \mathbb{Z}C_k \longleftarrow$$

Algorithm A.1: Computing $H_*(BS;\mathbb{Z})$: pre-processing and setup

Input: A nonempty finite semigroup S with operation *, and some $m \in \mathbb{N}$ **Output:** A list $[H_1(BS;\mathbb{Z}),\ldots,H_m(BS;\mathbb{Z})]$ of finitely generated abelian groups Function getHomology(S, *, m): $I, H, J \coloneqq \texttt{getMinIdeal}(S, *);$ /* Use Algorithm 5.1 */ if |I| = 1 or |J| = 1 then /* Detect K-thin semigroups */ // Short circuit: identify pre-computed small groups H. if |H| = 1 then return [0, 0, 0, 0, ...];else if $|H| \in \{2, 3, 5, 7\}$ then return $[C_{|H|}, 0, C_{|H|}, 0, \dots];$ else if |H| = 4 then if $|\{h * h : h \in H\}| = 1$ then return $[C_2^2, C_2, C_2^3, C_2^2, C_2^4, C_2^3, \dots];$ else return $[C_4, 0, C_4, 0, ...];$ else return getHomologyWithProjectiveResolution(*H*, *, *m*); else if some idempotent $e \in S$ has eSe = eS or eSe = Se then // Apply Lemma 7.4 if |eSe| < |S| then return getHomology(eSe, *, m); else return getHomologyWithProjectiveResolution(S, *, m); else return getHomologyWithProjectiveResolution($S \sqcup \{1\}, *^{\mathbb{1}}, m$); Function getHomologyWithProjectiveResolution(S, *, m): // Here we assume (S, *) is a monoid. e := an idempotent of S with smallest |Se|; $C_0 \coloneqq \mathbb{Z}Se; //$ the module $\mathbb{Z}Se$ represented as a list [e] $Z_0 \coloneqq \ker(\varepsilon \colon C_0 \to \mathbb{Z}); // a \mathbb{Z}$ -basis for this kernel $C_1, \partial_1 \coloneqq \texttt{getCoverByMapping}(S, *, C_0, Z_0);$ /* See Algorithm A.2 */ $root := ResolutionNode(S, *, C_1, C_0, \partial_1);$ /* See Algorithm A.3 */ nodeCache := emptyMapping(); nodeCache($[C_1, C_0, \partial_1]$) := root; return [root.homologyWithShift(*i*, nodeCache) for $0 \le i \le m-1$]

Algorithm A.2: Computing $H_*(BS; \mathbb{Z})$: covering submodules

Input: A monoid (S, *), a $\mathbb{Z}S$ -module $C = \mathbb{Z}Sf_1 \oplus \cdots \oplus \mathbb{Z}Sf_n$, and a submodule A of C

Output: A $\mathbb{Z}S$ -module $C' = \mathbb{Z}Se_1 \oplus \cdots \oplus \mathbb{Z}Se_m$ with a map $\partial : C' \to C$ such that $\partial(C') = A$

Function getCoverByMapping(S, *, C, A):

for v in a \mathbb{Z} -basis of A do

X := a mutable submodule of \mathbb{Z}^{ℓ} where $\ell = |Sf_1| + \dots + |Sf_k|$;

 $C' \coloneqq$ zero module; $\partial \coloneqq$ a k-by-0 matrix with entries in $\mathbb{Z}S$;

// Traversing A in a strategic order here can make C^\prime small

if $v \notin X$ then

Find an idempotent $e \in S$ with ev = v and |Se| as small as possible; Add a direct summand $\mathbb{Z}Se$ to C';

Append v as a column of ∂ ;

foreach $s \in S$ do X.addVector(sv);

return $C', \partial;$

Algorithm A.3: Computing $H_*(BS;\mathbb{Z})$: the node class

Class ResolutionNode: **Instance Data** A finite monoid (S, *); A list $E = [e_1, \ldots, e_n]$ of idempotents of S representing $C_i = \bigoplus_i \mathbb{Z}Se_i$; A list $F = [f_1, \ldots, f_m]$ of idempotents of S representing $C_{i-1} = \bigoplus_i \mathbb{Z}Sf_j$; A matrix in $(\mathbb{Z}S)^{m \times n}$ representing a map $\partial_i : C_i \to C_{i-1}$ as in Lemma A.3; A partition $E = E_1 \sqcup \cdots \sqcup E_k$ representing some $C_i = C_{i,1} \oplus \cdots \oplus C_{i,k}$; A list children of ResolutionNodes; A mapping hCache from integers to finitely generated abelian groups; Method this.makeChildren(nodeCache): // Split this covering problem into smaller problems $Z \coloneqq \ker(\partial_i \colon C_i \to C_{i-1}); // a \mathbb{Z}$ -basis for this kernel Partition $E = E_1 \sqcup \cdots \sqcup E_k$ according to which e_i are linked by occurring as nonzero entries in some common basis element in Z. Write $C_{i,j}$ for the $\mathbb{Z}S$ -module $C_{i,j} \coloneqq \bigoplus_{e \in E_j} \mathbb{Z}Se;$ $Z_1,\ldots,Z_k\coloneqq [],\ldots,[];$ foreach $v \in Z$ do Find E_j corresponding to the nonzero indices of v; Add to the corresponding Z_i a representation of v, eliminating the zero entries of v from the complement of E_j ; children := [];for $j \in \{1, ..., k\}$ do $C_{i+1,j}, \partial_{i+1,j} \coloneqq \texttt{getCoverByMapping}(S, *, C_{i,j}, Z_j);$ if $[C_{i+1,j}, C_{i,j}, \partial_{i+1,j}]$ is in nodeCache then child := nodeCache([$C_{i+1,j}, C_{i,j}, \partial_{i+1,j}$]); else child := ResolutionNode($S, *, C_{i+1,j}, C_{i,j}, \partial_{i+1,j}$); nodeCache([$C_{i+1,j}, C_{i,j}, \partial_{i+1,j}$]) := child; children.append(child); Method this.computeHomology(): $M_{\text{out}} \coloneqq \text{augmentation}(\partial_i); // \text{See Lemma A.3}$ images := [augmentation($\partial_{i+1,j}$) for $j \in \{1, \ldots, k\}$]; // from children // Embed each column of the individual image matrices // and collect together into one matrix: $M_{\mathrm{in}} \coloneqq \bigoplus_{i=1}^k \mathrm{images}; // \mathrm{a map} \ \mathbb{Z} \otimes \bigoplus_i C_{i+1,j} \to \mathbb{Z} \otimes C_i$ // Use standard algorithms for \mathbb{Z} -matrices: return homologyFromMatrices(M_{out} , M_{in}); Method this.homologyWithShift(shift, nodeCache): if this.hCache(shift) is defined then return this.hCache(shift); if this.children is not computed then this.makeChildren(nodeCache); if shift = 0 then $H_i \coloneqq$ this.computeHomology(); else $H_i := \bigoplus_{i=1}^k \text{children}[j]$.homologyWithShift(shift - 1, nodeCache); $hCache(shift) := H_i;$ return H_i ;

This is exactly the structure created by Algorithm A.1 for C_k .

Example A.5. Let $S = \{x_{11}, x_{12}, x_{21}, x_{22}, y_{11}, z_{11}\}$ with operation defined by $a_{ij}b_{k\ell} = x_{i\ell}$ for $a, b \in \{x, y, z\}$, extending Example 8.5. Algorithm A.1 finds the following projective resolution:

$$\mathbb{Z} \xleftarrow{\varepsilon} \mathbb{Z}S^{1}x_{11} \xleftarrow{\partial_{1}} \mathbb{Z}S^{1} \xleftarrow{\partial_{2}} \mathbb{Z}S^{1}x_{11} \xleftarrow{0} \mathbb{Z}S^{1}x_{1} \xleftarrow{0} \mathbb{Z}S^{1$$

The three bold square brackets above represent the **ResolutionNodes** in Algorithm A.1. The maps here, as matrices of right-multipliers as in Lemma A.3 are:

$$\begin{aligned} \partial_1 &= \begin{pmatrix} x_{21} - x_{11} \end{pmatrix}; \quad \partial_2 &= \begin{pmatrix} x_{12} & x_{11} & y_{11} & z_{11} \end{pmatrix}; \\ \partial_3 &= \begin{pmatrix} x_{11} & x_{11} & x_{11} & x_{11} & x_{11} & x_{11} & x_{11} \\ -x_{21} & 0 & -x_{11} & -y_{11} & -z_{11} & 0 & 0 & 0 \\ 0 & -x_{21} & 0 & 0 & 0 & -x_{11} & -y_{11} & -z_{11} \end{pmatrix}. \end{aligned}$$

The kernels ker ∂_2 and ker ∂_3 split along the rows displayed in each node: ker $\partial_2 = 0 \oplus \operatorname{im} \partial_3$ and ker $\partial_3 = 0 \oplus 0 \oplus \operatorname{im} \partial_3 \oplus \operatorname{im} \partial_3$. The looped arrows indicate that we could repeat two new copies of the last node as its own children, and similarly extend those two children, and so on, with the overall size of the entire direct sum C_i growing exponentially in *i*. However, using **nodeCache** allowed this to be represented using only the finite data above. After tensoring with \mathbb{Z} , this doubling structure gives

$$H_i(BS;\mathbb{Z}) = \begin{cases} 0 & i=1\\ \mathbb{Z}^{2^{i-2}} & i \ge 2 \end{cases}$$

Indeed, an implementation of Algorithm A.1 computed $H_{10000}(BS;\mathbb{Z}) = \mathbb{Z}^{2^{9998}}$ in about one second.

APPENDIX B. TABLES OF HOMOLOGY

The smallsemi package [DM24] for the GAP system [GAP] lists all semigroups up to order 8, up to isomorphism and anti-isomorphism. Using the techniques from Appendix A, we can compute homology groups $H_i(BS)$ for all of their classifying spaces. Some results are displayed in the subsections that follow.

B.1. Semigroups of order at most 6. The tables below list the integral homology for the classifying spaces of each semigroup of order at most 6. Each row lists six homology groups, along with the number of semigroups, up to isomorphism and anti-isomorphism, having the given order and the given homology groups for its classifying space. Note that by Proposition 4.1, each row in the table for order n is also included in the table for order n + 1. A "." denotes the trivial group.

oru	er 1	:				
H_1	H_2	H_3	H_4	H_5	H_6	#
•	•	•	•	•	•	1

Order 3:

H_1	H_2	H_3	H_4	H_5	H_6	#
•	•	•	•	•	•	14
C_2	•	C_2	•	C_2	•	3
C_3	•	C_3	•	C_3	•	1

Ord	ler 2	:						
H_1	H_2	H_3	H_4	H_5	H_6	#		
•	•	•	•	•	•	3		
C_2	•	C_2	•	C_2	•	1		
Order 4:								
H_1	H_{2}	H_{2}	H_{4}	$H_{\rm E}$	H_{c}	#		

H_1	H_2	H_3	H_4	H_5	H_6	#
•	•			•		102
C_2		C_2		C_2		18
$\tilde{C_3}$		α		$\tilde{C_3}$		3
•	\mathbb{Z}	•				1
C_{2}^{2}	C_2	C_2^3	C_{2}^{2}	C_2^4	C_2^3	1
$\tilde{C_4}$		$\tilde{C_4}$		$\tilde{C_4}$		1

Order 6:

Order 5:

Orc	ier ə	:				
H_1	H_2	H_3	H_4	H_5	H_6	#
•	•	•	•	•	•	996
C_2	•	C_2	•	C_2	•	136
C_3	•	C_3	•	C_3	•	17
C_4	•	C_4	•	C_4	•	4
C_2^2	C_2	C_2^3	C_2^2	C_2^4	C_2^3	3
•	\mathbb{Z}	•	•	•	•	2
•	\mathbb{Z}	\mathbb{Z}	\mathbb{Z}	\mathbb{Z}	\mathbb{Z}	1
C_5	•	C_5	•	C_5	•	1

H_1	H_2	H_3	H_4	H_5	H_6	#
•	•	•	•	•	•	14334
C_2	•	C_2	•	C_2	•	1420
C_3	•	C_3	•	C_3	•	139
C_4		C_4		C_4		25
C_{2}^{2}	C_2	C_2^3	C_2^2	C_2^4	C_2^3	18
.	\mathbb{Z}	•				17
	\mathbb{Z}	\mathbb{Z}	\mathbb{Z}	\mathbb{Z}	\mathbb{Z}	6
C_5	•	C_5		C_5	•	3
	\mathbb{Z}	\mathbb{Z}^{2}	\mathbb{Z}^4	$\mathbb{Z}^{\tilde{8}}$	\mathbb{Z}^{16}	3
•	•	\mathbb{Z}	•	•	•	2
•	\mathbb{Z}	C_2		C_2		2
•	C_2	•	C_2	•	C_2	1
	$\mathbb{Z}^{\overline{2}}$					1
C_2	•	C_6	•	C_2	•	1
C_6	•	C_6	•	C_6	•	1

ables.								
H_1	H_2	H_3	H_4	H_5	H_6	H_7	H_8	#
•	•		•	•		•	•	811817
C_2		C_2		C_2		C_2		21982
C_3		C_3		C_3		C_3		1484
C_4	•	C_4	•	C_4	•	C_4	•	222
	\mathbb{Z}	•	•	•	•		•	163
C_{2}^{2}	C_2	C_2^3	C_2^2	C_2^4	C_2^3	C_2^5	C_2^4	147
	\mathbb{Z}	\mathbb{Z}	\mathbb{Z}	\mathbb{Z}	\mathbb{Z}	\mathbb{Z}	\mathbb{Z}	72
	\mathbb{Z}	\mathbb{Z}^2	\mathbb{Z}^4	\mathbb{Z}^8	\mathbb{Z}^{16}	\mathbb{Z}^{32}	\mathbb{Z}^{64}	18
C_5	•	C_5		C_5	•	C_5		18
	•	\mathbb{Z}	•		•			16
	•	\mathbb{Z}	\mathbb{Z}	\mathbb{Z}	\mathbb{Z}	\mathbb{Z}	\mathbb{Z}	14
	\mathbb{Z}	C_2	•	C_2	•	C_2		12
	\mathbb{Z}	•	\mathbb{Z}		\mathbb{Z}		\mathbb{Z}	9
	\mathbb{Z}	\mathbb{Z}^2	\mathbb{Z}^3	\mathbb{Z}^4	\mathbb{Z}^5	\mathbb{Z}^6	\mathbb{Z}^7	9
	•	\mathbb{Z}	•	\mathbb{Z}	•	\mathbb{Z}		7
	C_2	•	C_2		C_2		C_2	6
C_6	•	C_6	•	C_6	•	C_6	•	5
•	•	\mathbb{Z}^2	•	•	•	•	•	4
	\mathbb{Z}	\mathbb{Z}^3	\mathbb{Z}^9	\mathbb{Z}^{27}	\mathbb{Z}^{81}	\mathbb{Z}^{243}	\mathbb{Z}^{729}	4
C_2	•	C_6	•	C_2	•	C_6	•	4
	\mathbb{Z}^2	•	•	•	•	•	•	2
	•	C_2	•	C_2	•	C_2	•	1
	•	\mathbb{Z}	C_2		C_2		C_2	1
•	\mathbb{Z}	C_3	•	C_3	•	C_3	•	1
•	\mathbb{Z}	$\mathbb{Z} \times C_2$	$\mathbb{Z} \times C_2^2$			$\mathbb{Z} \times C_2^{21}$		1
•	\mathbb{Z}^2	\mathbb{Z}^2	\mathbb{Z}^2	\mathbb{Z}^2	\mathbb{Z}^2	\mathbb{Z}^2	\mathbb{Z}^2	1
C_7	·	C_7	•	C_7	•	C_7	•	1

B.2. Semigroups of order 7. The table below lists eight integral homology groups for the classifying spaces of each semigroup of order 7, formatted as in the previous tables.

$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	H_1	H_2	H_3	H_4	H_5	H_6	H_7	H_8	#
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		•	•	•	•	•		•	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	1		C_2		C_2		C_2		
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$									
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$									
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$							•		
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	C_2^2		C_2^3	C_2^2	C_2^4	C_2^3	C_2^5	C_2^4	
$\begin{array}{cccccccccccccccccccccccccccccccccccc$									
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$						•		•	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		7.		\mathbb{Z}^4	\mathbb{Z}^8	\mathbb{Z}^{16}	Z^{32}	Z^{64}	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$									
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	C-								
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		77.					C_{5}		
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$				778	$\frac{C_{2}}{\pi^{21}}$	755	7^{144}	7/377	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$			<i>1</i> ,2		<i>1</i> 2				
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	•		•		•		•		
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	•	\mathbb{C}_2			772				
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		•						<i>Ш</i>	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		•		772		775		7713	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	· ·								
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		L							
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	•	٠	L		"		⊿-∘		
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	•		773		· 7727		· rz243	r#729	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	•		Ш°						
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	· ·	Z		Ľ	Ľ	<u>"</u>	Ш°	Ш°	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		•		•	•	•	•	•	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	C_6		C_6	•	C_6		C_6		
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	•	Z	•		•		•		
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	· ·	•							
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	•	•		\mathbb{Z}^2		\mathbb{Z}^4		\mathbb{Z}^{6}	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	C_2	•	C_6	•	C_2	•		•	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	· ·		·	•	•	•	•	•	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	· ·			•		•		•	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	•		$\mathbb{Z} \times C_2$		$\mathbb{Z} \times C_2^5$	$\mathbb{Z} \times C_2^{10}$	$\mathbb{Z} \times C_2^{21}$	$\mathbb{Z} \times C_2^{42}$	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	· ·	\mathbb{Z}	•	\mathbb{Z}	•	•	•	•	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	· ·	•		•		•	\mathbb{Z}^{8}	•	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	· ·	•		C_2	•	C_2	•	C_2	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	· ·	\mathbb{Z}		\mathbb{Z}^2		\mathbb{Z}^{8}		\mathbb{Z}^{32}	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	· ·	•		•	C_2	•	C_2	•	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	· ·	\mathbb{Z}							
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$.	•			$\mathbb{Z} \times C_2$	$\mathbb{Z} \times C_2^2$			8
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	·	•	C_2	C_2	C_2^2	C_2^2	C_{2}^{3}	C_{2}^{3}	8
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$			\mathbb{Z}		•	•	•		7
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$			C_3	•	C_3	•	C_3		
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		\mathbb{Z}^2		\mathbb{Z}^2	\mathbb{Z}^2	\mathbb{Z}^2	\mathbb{Z}^2	\mathbb{Z}^2	
$\begin{array}{cccccccccccccccccccccccccccccccccccc$		•		•	•	•	•	•	6
$\begin{array}{cccccccccccccccccccccccccccccccccccc$.	\mathbb{Z}^2			\mathbb{Z}^{16}	\mathbb{Z}^{32}	\mathbb{Z}^{64}	\mathbb{Z}^{128}	
$C_7 \cdot C_7 \cdot C_7 \cdot C_7 \cdot C_7 \cdot C_7$.	•	\mathbb{Z}		\mathbb{Z}	C_2^3		C_2^7	4
\mathbb{Z} \mathbb{Z}^2 \mathbb{Z}^3 \mathbb{Z}^5 \mathbb{Z}^9 \mathbb{Z}^{15} \mathbb{Z}^{25} 3	C_7	•			C_7	•	C_{7}		
0	.	\mathbb{Z}	\mathbb{Z}^2	\mathbb{Z}^3	\mathbb{Z}^5	\mathbb{Z}^9	\mathbb{Z}^{15}	\mathbb{Z}^{25}	3

B.3. Semigroups of order 8. The table below lists eight integral homology groups for the classifying spaces of each semigroup of order 8, formatted as in the previous tables. The table continues on the next page.

.	\mathbb{Z}	$\mathbb{Z}^2 \times C_2$	$\mathbb{Z}^4 imes C_2^4$		$\mathbb{Z}^{16} \times C_2^{50}$		$\mathbb{Z}^{64} \times C_2^{504}$	3
	\mathbb{Z}	C_2	\mathbb{Z}	C_2^3	$\mathbb{Z} \times C_2$	C_2^7	$\mathbb{Z} \times C_2^5$	3
	\mathbb{Z}	•	\mathbb{Z}	C_2	\mathbb{Z}	C_{2}^{3}	$\mathbb{Z} \times C_2$	3
	C_2	C_2	C_2^2	C_2^2	C_2^3	C_2^3	C_2^4	3
.		\mathbb{Z}	$\mathbb{Z} \times C_2$	$\mathbb{Z} \times C_2^2$	$\mathbb{Z} \times C_2^5$	$\mathbb{Z} \times C_2^{10}$	$\mathbb{Z} \times C_2^{21}$	3
.	\mathbb{Z}	\mathbb{Z}^2	\mathbb{Z}^3	\mathbb{Z}^6	\mathbb{Z}^{11}	\mathbb{Z}^{19}	\mathbb{Z}^{36}	2
	\mathbb{Z}	\mathbb{Z}^2	\mathbb{Z}^3	\mathbb{Z}^5	\mathbb{Z}^9	\mathbb{Z}^{16}	\mathbb{Z}^{28}	2
	\mathbb{Z}	$\mathbb{Z} \times C_2$	$\mathbb{Z} \times C_2$	$\mathbb{Z} \times C_2^2$	$\mathbb{Z}\times C_2^2$	$\mathbb{Z} \times C_2^3$	$\mathbb{Z} imes C_2^3$	2
	\mathbb{Z}	C_4	•	C_4	•	C_4		2
	\mathbb{Z}	C_2^2	C_2	C_{2}^{3}	C_2^2	C_2^4	C_2^3	2
	•	$\mathbb{Z}^{\overline{2}}$	C_2^2	•	C_2^2		C_2^2	2
•	•	C_2^2	•	C_2^2		C_2^2	•	2
C_8	•	C_8	•	C_8		C_8		1
C_2	\mathbb{Z}	C_2	•	C_2		C_2	•	1
$C_2 \times C_4$	C_2	$C_2^2 \times C_4$	C_2^2	$C_2^3 \times C_4$	C_{2}^{3}	$C_2^4 \times C_4$	C_2^4	1
$\begin{array}{c} C_2^2 \\ C_2^2 \end{array}$	C_2	$C_2^2 \times C_4$	C_{2}^{2}	$\begin{array}{c} C_2^4 \\ C_2^2 \end{array}$	C_{2}^{3}	$C_2^4 \times C_4$	C_2^4	1
C_2^2	•	C_8	•	C_2^2		C_8	•	1
C_2^3	C_2^3	C_2^7	C_{2}^{8}	C_{2}^{13}	C_{2}^{15}	C_{2}^{21}	C_{2}^{24}	1
•	\mathbb{Z}	\mathbb{Z}^2	\mathbb{Z}^2	\mathbb{Z}^4	\mathbb{Z}^8	\mathbb{Z}^8	\mathbb{Z}^{16}	1
	\mathbb{Z}	$\mathbb{Z} \times C_3$	$\mathbb{Z} \times C_3^2$	$\mathbb{Z} \times C_3^5$	$\mathbb{Z}\times C_3^{10}$	$\mathbb{Z}\times C_3^{21}$	$\mathbb{Z}\times C_3^{42}$	1
· ·	\mathbb{Z}^2	C_2^2	•	C_2^2	•	C_2^2	•	1
· ·	\mathbb{Z}^3	•	•		•	•	•	1
· ·	C_2^2	•	C_2^2		C_2^2	•	C_2^2	1
•	•	\mathbb{Z}	C_3	•	C_3	•	C_3	1