HYBRID SUBCONVEXITY BOUNDS FOR TWISTS OF $GL_2 \times GL_2$ *L*-FUNCTIONS

CHENCHEN SHAO AND HUIMIN ZHANG

ABSTRACT. In this paper, we prove hybrid subconvexity bounds for $GL_2 \times GL_2$ Rankin– Selberg *L*-functions twisted by a primitive Dirichlet character χ modulo a prime power, in the *t* and depth aspects.

1. INTRODUCTION

The subconvexity problem for general *L*-functions is one of the central questions in analytic number theory and it has many applications in number theory. Let f and g be holomorphic or Maass cusp forms for $SL_2(\mathbb{Z})$ with normalized Fourier coefficients $\lambda_f(n)$ and $\lambda_g(n)$ respectively. Let χ be a primitive Dirichlet character of prime power conductor p^{κ} with p prime. The $GL_2 \times GL_2 \times GL_1$ Rankin–Selberg *L*-function is defined by

$$L(s, f \times g \times \chi) = L(2s, \chi^2) \sum_{n=1}^{\infty} \frac{\lambda_f(n)\lambda_g(n)\chi(n)}{n^s},$$

for $\operatorname{Re}(s) > 1$. And this *L*-function has an analytic continuation to the whole complex plane. The subconvexity problem for $\operatorname{GL}_2 \times \operatorname{GL}_2 \times \operatorname{GL}_1$ *L*-functions is to obtain a bound of the form

$$L\left(\frac{1}{2}+it, f \times g \times \chi\right) \ll (p^{\kappa}(|t|+1))^{1-\delta+\varepsilon}$$

with $0 \leq \delta \leq 1$.

The convexity bounds correspond to the bounds with $\delta = 0$, which follow from the functional equation and the Phragmén–Lindelöf convexity principle. The Lindelöf hypothesis is the statement with $\delta = 1$.

The literature on the subconvexity problem is rich and the first result can be traced back over 100 years ago. For the GL_1 case, Weyl [24] proved a subconvexity bound for Riemann zeta function. Burgess [6] proved a subconvexity bound for Dirichlet *L*-functions. For GL_2 case, see Michel–Venkatesh [21] and the references therein. Many progress on GL_3 *L*-functions has been made in the past ten years (see [3], [17], [19], [20], [11], etc.).

Subconvexity for $GL_2 \times GL_2$ *L*-functions have been solved by many number theorists. For the *t*-aspect, see Bernstein–Reznikov [2], Acharya–Sharma–Singh [1], Blomer–Jana–Nelson [4], Huang–Sun–Zhang [10]. For the depth aspect, see Sun [22].

²⁰¹⁰ Mathematics Subject Classification. 11F66.

Key words and phrases. Hybrid subconvexity, twists, Delta method, $GL_2 \times GL_2$ Rankin–Selberg L-functions.

This work is in part supported by the National Key R&D Program of China (No. 2021YFA1000700). H. Zhang is also supported by NSFC (No. 12031008) and China Scholarship Council (No. 202206220071).

In this paper, we extend the techniques to prove hybrid subconvexity bounds for twists of $GL_2 \times GL_2$ *L*-functions, which generalizes the best known bounds using delta method in the depth aspect and in the *t*-aspect simultaneously.

Theorem 1.1. Let f and g be holomorphic or Maass cusp forms for $SL_2(\mathbb{Z})$ and let χ be a primitive Dirichlet character of prime power conductor p^{κ} with p an odd prime and $\kappa > 12$. The following hybrid subconvexity bound holds:

$$L\left(\frac{1}{2}+it, f \times g \times \chi\right) \ll_{\varepsilon} p^{\frac{3}{4}} (p^{\kappa})^{1-\frac{1}{16}+\varepsilon} (|t|+1)^{1-\frac{1}{10}+\varepsilon}.$$

Acknowledgements. The authors would like to thank Professors Valentin Blomer, Bingrong Huang and Lilu Zhao for many helpful comments and encouragement. The authors also thank Dr. Xinchen Miao for pointing out some references. A large part of this research is finished when Huimin Zhang is visiting the University of Bonn. She is grateful to the China Scholarship Council for supporting her studies at the University of Bonn, and also thanks Mathematical Institute of the University of Bonn for their hospitality.

Notation. Throughout the paper, the letters ε and A denote arbitrarily small and large positive constants, respectively, not necessarily the same at different occurrences. We use $A \simeq B$ to mean that $c_1B \leq |A| \leq c_2B$ for some positive constants c_1 and c_2 . The symbol $\ll_{a,b,c}$ denotes that the implied constant depends at most on a, b and c. As usual, $e(x) = e^{2\pi i x}$ and $q \sim R$ means $R < q \leq 2R$.

2. Outline of the proof

We give a brief sketch of the proof of Theorem 1.1. In section 4.1, after using the approximate functional equation, we are left with estimating the sum

$$S(N) = \sum_{n \sim N} \lambda_f(n) \lambda_g(n) n^{-it} \chi(n)$$

with $N \ll p^{2\kappa}t^2$. In section 4.2, we apply the conductor lowering mechanism for the depth aspect, and apply a version of the delta method of Duke, Friedlander and Iwaniec to decouple the oscillations. For simplicity we consider the generic case, i.e. $N = p^{2\kappa}t^2$ and $q \sim Q = \sqrt{N/p^{\lambda}K}$ with (q, p) = 1, then we essentially have

$$S(N) \approx \int_{u \sim 1} \frac{1}{Qp^{\lambda}} \sum_{\substack{q \sim Q \\ (q,p)=1}} \frac{1}{q} \sum_{a \pmod{qp^{\lambda}}} \sum_{n \sim N} \lambda_g(n) e\left(-\frac{na}{qp^{\lambda}}\right) e\left(-\frac{nu}{Qqp^{\lambda}}\right)$$
$$\cdot \sum_{m \sim N} \lambda_f(m) m^{-it} \chi(m) e\left(\frac{ma}{qp^{\lambda}}\right) e\left(\frac{mu}{Qqp^{\lambda}}\right) du$$

Trivially we have $S(N) \ll N^2$. Note that we don't need the conductor lowering trick for the *t*-aspect as [1], in fact, the *u*-integral above plays the same role.

Next we apply the GL_2 Voronoi summation formula on the sums over n and m in section 5. For the *n*-sum, we roughly arrive at

$$S_1 = \sum_{n \sim N} \lambda_g(n) e\left(-\frac{na}{qp^{\lambda}}\right) e\left(-\frac{nu}{Qqp^{\lambda}}\right) \approx \frac{N}{qp^{\lambda}} \sum_{n=1}^{\infty} \lambda_g(n) e\left(\frac{\overline{a}n}{qp^{\lambda}}\right) \Phi_1\left(\frac{nN}{q^2p^{2\lambda}}\right),$$

for certain weight function Φ_1 . The conductor is $K^2 Q^2 p^{2\lambda}$, so the dual length becomes $n \simeq \frac{K^2 Q^2 p^{2\lambda}}{N} = K p^{\lambda}$. By Lemma 5.1, we can trivially estimate the dual sum as $\frac{N}{Qp^{\lambda}} \cdot K p^{\lambda} \cdot (\frac{K p^{\lambda} N}{Q^2 p^{2\lambda}})^{-1/2}$. So we save $\frac{N^{1/2}}{p^{\lambda/2} K^{1/2}}$ from *n*-sum. By the stationary phase method, we save $K^{1/2}$ from the *u*-integral. And for the *m*-sum, we get

$$S_2 \approx \frac{N}{qp^{\kappa}} \sum_m \lambda_f(m) e\left(\frac{-\overline{a^*}m}{qp^{\kappa}}\right) \Phi_2\left(\frac{mN}{q^2p^{2\kappa}}\right),$$

for certain weight function Φ_2 . The conductor is $t^2 Q^2 p^{2\kappa}$, and hence the dual length becomes $m \simeq \frac{t^2 Q^2 p^{2\kappa}}{N} = \frac{t^2}{K} p^{2\kappa-\lambda}$. By Lemma 3.5, we can trivially estimate the dual sum as $\frac{N}{qp^{\kappa}} \cdot (\frac{N}{q^2 p^{2\kappa}})^{-1/4} \cdot (\frac{t^2 Q^2 p^{2\kappa}}{N})^{3/4} \cdot t^{-1/2}$. So we save $\frac{N^{1/2} p^{\lambda/2} K^{1/2}}{p_{\kappa}^{\kappa} t}$ from *m*-sum.

Because of a Ramanujan sum appearing in the character sum and assuming square-root cancellation for the remaining part of the character sum, until now we have saved

$$\frac{N^{1/2}}{p^{\lambda/2}K^{1/2}} \times K^{1/2} \times \frac{N^{1/2}p^{\lambda/2}K^{1/2}}{p^{\kappa}t} \times Qp^{\lambda/2} \sim N.$$

So we are on the convex bound and need to save a little more.

After an application of the Cauchy–Schwarz inequality and the Poisson summation formula to the sum over m in Section 6, we arrive at

$$S(N) \ll \frac{Q}{p^{(\kappa+\lambda)/2}} \sum_{\substack{q \sim R \\ (q,p)=1}} \left(\sum_{\substack{n_1 \asymp \frac{N}{Q^2} \\ n_2 \equiv n_1 \mod q}} |\lambda_g(n_1)|^2 \sum_{\substack{n_2 \asymp \frac{N}{Q^2} \\ n_2 \equiv n_1 \mod q}} \frac{R^2 p^{\kappa} t^2}{Nq} \sum_{m \in \mathbb{Z}} \mathfrak{C}(m, n_1, n_2) \cdot \mathfrak{I}(m, n_1, n_2) \right)^{1/2}$$

for certain character sum $\mathfrak{C}(m, n_1, n_2)$ in (6.4) and integral transform $\mathfrak{I}(m, n_1, n_2)$ in (6.5). In total, we save $\left(\frac{p^{\lambda}K}{Q}\right)^{1/2} = \frac{p^{3\lambda/4-\kappa/2}K^{3/4}}{t^{1/2}}$ from the zero frequency and save

$$\left(\frac{p^{\kappa}t^2p^{\kappa-\lambda}}{QK^2}\right)^{1/2} \times \left(\frac{p^{\kappa+\lambda}}{p^{\frac{5\kappa}{2}-(\kappa-\lambda)}}\right)^{1/2} \times K^{1/4} = p^{\frac{\kappa-\lambda}{4}} \frac{t^{1/2}}{K^{1/2}}$$

from the non-zero frequencies. So the optimal choice for λ and K are given by $\lambda = 3\kappa/4$ and $K = t^{4/5}$. In total, we have saved $N \times p^{\kappa/16} t^{1/10}$. It follows that

$$S(N) \ll N^{1/2} p^{\frac{3}{4}} (p^{\kappa})^{1-\frac{1}{16}+\varepsilon} (|t|+1)^{1-\frac{1}{10}+\varepsilon}.$$

3. Preliminaries

3.1. The delta method. Define $\delta : \mathbb{Z} \to \{0, 1\}$ with $\delta(0) = 1$ and $\delta(n) = 0$ for $n \neq 0$. We will use a version of the delta method of Duke, Friedlander and Iwaniec (see [13, Chapter 20]) which states that for any $n \in \mathbb{Z}$ and $Q \in \mathbb{R}^+$, we have

$$\delta(n) = \frac{1}{Q} \sum_{q \le Q} \frac{1}{q} \sum_{a \mod q}^{\star} e\left(\frac{na}{q}\right) \int_{\mathbb{R}} g(q, u) e\left(\frac{nu}{qQ}\right) du$$
(3.1)

where the \star on the sum indicates that the sum over *a* is restricted to (a, q) = 1. The function *g* has the following properties (see (20.158) and (20.159) of [13] and [9, Lemma 15])

$$g(q,u) \ll |u|^{-A}, \quad g(q,u) = 1 + h(q,u) \text{ with } h(q,u) = O\left(\frac{Q}{q}\left(\frac{q}{Q} + |u|\right)^{A}\right)$$
(3.2)

for any A > 1 and

$$u^{j}\frac{\partial^{j}}{\partial u^{j}}g(q,u) \ll (\log Q) \min\left\{\frac{Q}{q}, \frac{1}{|u|}\right\}, \qquad j \ge 1.$$
(3.3)

In particular the first property in (3.2) implies that the effective range of the integration in (3.1) is $[-Q^{\varepsilon}, Q^{\varepsilon}]$.

Sun [22] shows a new variant of delta method, which can reduce Q to a smaller value by an idea of Munshi [19].

Lemma 3.1. [22, Lemma 1] Suppose $\lambda \geq 1$. Let $U_0 \in \mathcal{C}^{\infty}_c(-4, 4)$ be a smooth positive function with $U_0(x) = 1$ if $x \in [-2, 2]$ and satisfying $U_0^{(j)}(x) \ll_j 1$. Then

$$\begin{split} \delta(n) &= \sum_{k=0}^{\lambda} \frac{1}{Q} \sum_{\substack{q \leq Q \\ (q,p)=1}} \frac{1}{q p^{\lambda}} \sum_{\substack{a \bmod q p^{\lambda-k}}} e\left(\frac{na}{q p^{\lambda-k}}\right) \int_{\mathbb{R}} U_0\left(\frac{u}{N^{\varepsilon}}\right) g(q,u) e\left(\frac{nu}{Q q p^{\lambda}}\right) \mathrm{d}u \\ &+ \sum_{r=1}^{\lfloor \log Q/\log p \rfloor} \frac{1}{Q} \sum_{\substack{q \leq Q/p^r \\ (q,p)=1}} \frac{1}{q p^{\lambda+r}} \sum_{\substack{a \bmod q p^{\lambda+r}}} e\left(\frac{na}{q p^{\lambda+r}}\right) \\ &\int_{\mathbb{R}} U_0\left(\frac{u}{N^{\varepsilon}}\right) g(p^r q, u) e\left(\frac{nu}{Q q p^{\lambda+r}}\right) \mathrm{d}u + O_A(N^{-A}) \end{split}$$

for any A > 0.

Proof. See [22, Lemma 1].

First we recall some basic results on automorphic forms for GL_2 (see e.g. [8]).

3.2. Holomorphic cusp forms for GL₂. Let f be a holomorphic cusp form of weight k_f for $SL_2(\mathbb{Z})$ with Fourier expansion

$$f(z) = \sum_{n=1}^{\infty} \lambda_f(n) n^{(k_f - 1)/2} e(nz)$$

for Im z > 0, normalized such that $\lambda_f(1) = 1$. By the Ramanujan–Petersson conjecture proved by Deligne [7], we have $\lambda_f(n) \ll \tau(n) \ll n^{\varepsilon}$ with $\tau(n)$ being the divisor function.

For $\varphi(x) \in \mathcal{C}_c(0,\infty)$, we set

$$\Phi(x) = 2\pi i^{k_f} \int_0^\infty \varphi(y) J_{k_f-1}(4\pi\sqrt{xy}) \mathrm{d}y, \qquad (3.4)$$

where J_{k_f-1} is the usual J-Bessel function of order $k_f - 1$. We have the following Voronoi summation formula (see [16, Theorem A.4]).

Lemma 3.2. Let $q \in \mathbb{N}$ and $a \in \mathbb{Z}$ be such that (a, q) = 1. For N > 0, we have

$$\sum_{n=1}^{\infty} \lambda_f(n) e\left(\frac{an}{q}\right) \varphi\left(\frac{n}{N}\right) = \frac{N}{q} \sum_{n=1}^{\infty} \lambda_f(n) e\left(-\frac{\overline{a}n}{q}\right) \Phi\left(\frac{nN}{q^2}\right),$$

where \overline{a} denotes the multiplicative inverse of a modulo q.

The function $\Phi(x)$ has the following asymptotic expansion when $x \gg 1$ (see [18, Lemma 3.2]).

Lemma 3.3. For any fixed integer $J \ge 1$ and $x \gg 1$, we have

$$\Phi(x) = x^{-1/4} \int_0^\infty \varphi(y) y^{-1/4} \sum_{j=0}^J \frac{c_j e(2\sqrt{xy}) + d_j e(-2\sqrt{xy})}{(xy)^{j/2}} dy + O_{k_f,J} \left(x^{-J/2 - 3/4} \right) +$$

where c_i and d_i are constants depending on k_f .

3.3. Maass cusp forms for GL_2 . Let f be a Hecke–Maass cusp form for $SL_2(\mathbb{Z})$ with Laplace eigenvalue $1/4 + t_f^2$. Then f has a Fourier expansion

$$f(z) = \sqrt{y} \sum_{n \neq 0} \lambda_f(n) K_{it_f}(2\pi |n|y) e(nx),$$

where K_{it_f} is the modified Bessel function of the third kind. The Fourier coefficients satisfy

$$\lambda_f(n) \ll n^{\theta + \varepsilon},\tag{3.5}$$

where, here and throughout the paper, θ denotes the exponent towards the Ramanujan conjecture for GL₂ Maass forms. The Ramanujan conjecture states that $\theta = 0$ and the current record due to Kim and Sarnak [14] is $\theta = 7/64$. It is well-known that

$$\sum_{n \le N} |\lambda_f(n)|^2 \ll_f N \tag{3.6}$$

by the Rankin–Selberg theory.

For $\varphi(x) \in \mathcal{C}_c^{\infty}(0,\infty)$, we define the integral transforms

$$\Phi^{+}(x) = \frac{-\pi}{\sin(\pi i t_{f})} \int_{0}^{\infty} \varphi(y) \left(J_{2it_{f}}(4\pi\sqrt{xy}) - J_{-2it_{f}}(4\pi\sqrt{xy}) \right) dy,$$

$$\Phi^{-}(x) = 4\varepsilon_{f} \cosh(\pi t_{f}) \int_{0}^{\infty} \varphi(y) K_{2it_{f}}(4\pi\sqrt{xy}) dy,$$

(3.7)

where ε_f is an eigenvalue under the reflection operator. We have the following Voronoi summation formula (see [16, Theorem A.4]).

Lemma 3.4. Let $q \in \mathbb{N}$ and $a \in \mathbb{Z}$ be such that (a, q) = 1. For N > 0, we have

$$\sum_{n=1}^{\infty} \lambda_f(n) e\left(\frac{an}{q}\right) \varphi\left(\frac{n}{N}\right) = \frac{N}{q} \sum_{\pm} \sum_{n=1}^{\infty} \lambda_f(n) e\left(\mp \frac{\overline{a}n}{q}\right) \Phi^{\pm}\left(\frac{nN}{q^2}\right),$$

where \overline{a} denotes the multiplicative inverse of a modulo q.

For $x \gg 1$, we have (see (3.8) in [18])

$$\Phi^{-}(x) \ll_{t_f,A} x^{-A}.$$
(3.8)

For $\Phi^+(x)$ and $x \gg 1$, we have a similar asymptotic formula as for $\Phi(x)$ in the holomorphic case (see [18, Lemma 3.4]).

Lemma 3.5. For any fixed integer $J \ge 1$ and $x \gg 1$, we have

$$\Phi^{+}(x) = x^{-1/4} \int_{0}^{\infty} \varphi(y) y^{-1/4} \sum_{j=0}^{J} \frac{c_{j} e(2\sqrt{xy}) + d_{j} e(-2\sqrt{xy})}{(xy)^{j/2}} dy + O_{t_{f,J}} \left(x^{-J/2 - 3/4} \right),$$

where c_i and d_i are some constants depending on t_f .

Remark 3.6. For $x \gg N^{\varepsilon}$, we can choose J sufficiently large so that the contribution from the O-terms in Lemmas 3.3 and 3.5 is negligible. For the main terms we only need to analyze the leading term j = 0, as the analysis of the remaining lower order terms is the same and their contribution is smaller compared to that of the leading term.

Notice that Lemma 3.5 are only valid for $x \gg N^{\varepsilon}$. So we also need the facts which state that, for $y > 0, k \ge 0$ and $\operatorname{Re} \nu = 0$, one has (see [16, Lemma C.2])

$$y^{k} J_{\nu}^{(k)}(y) \ll_{k,\nu} \frac{1}{(1+y)^{1/2}},$$

$$y^{k} K_{\nu}^{(k)}(y) \ll_{k,\nu} \frac{e^{-y} (1+|\log y|)}{(1+y)^{1/2}}.$$
(3.9)

3.4. Estimates for exponential integrals. Let

$$I = \int_{\mathbb{R}} w(y) e^{ih(y)} dy.$$

Firstly, we have the following estimates for exponential integrals (see [5, Lemma 8.1]).

Lemma 3.7. Let w(x) be a smooth function supported on [a, b] and h(x) be a real smooth function on [a, b]. Suppose that there are parameters Q, U, Y, Z, R > 0 such that

$$h^{(i)}(x) \ll_i Y/Q^i, \qquad w^{(j)}(x) \ll_j Z/U^j,$$

for $i \geq 2$ and $j \geq 0$, and

$$|h'(x)| \ge R.$$

Then for any $A \ge 0$ we have

$$I \ll_A (b-a)Z\left(\frac{Y}{R^2Q^2} + \frac{1}{RQ} + \frac{1}{RU}\right)^A.$$

Next, we need the following evaluation for exponential integrals which are Lemma 8.1 and Proposition 8.2 of [5] in the language of inert functions (see [15, Lemma 3.1]).

Let \mathcal{F} be an index set, $Y : \mathcal{F} \to \mathbb{R}_{\geq 1}$ and under this map $T \mapsto Y_T$ be a function of $T \in \mathcal{F}$. A family $\{w_T\}_{T \in \mathcal{F}}$ of smooth functions supported on a product of dyadic intervals in $\mathbb{R}^d_{>0}$ is called Y-inert if for each $j = (j_1, \ldots, j_d) \in \mathbb{Z}^d_{>0}$ we have

$$C(j_1, \dots, j_d) = \sup_{T \in \mathcal{F}} \sup_{(y_1, \dots, y_d) \in \mathbb{R}^d_{>0}} Y_T^{-j_1 - \dots - j_d} \left| y_1^{j_1} \cdots y_d^{j_d} w_T^{(j_1, \dots, j_d)}(y_1, \dots, y_d) \right| < \infty$$

Lemma 3.8. Suppose that $w = w_T(y)$ is a family of Y-inert functions, with compact support on [Z, 2Z], so that $w^{(j)}(y) \ll (Z/Y)^{-j}$. Also suppose that h is smooth and satisfies $h^{(j)}(y) \ll$ H/Z^j for some $H/X^2 \ge R \ge 1$ and all y in the support of w.

(1) If $|h'(y)| \gg H/Z$ for all y in the support of w, then $I \ll_A ZR^{-A}$ for A arbitrarily large.

(2) If $h''(y) \gg H/Z^2$ for all y in the support of w, and there exists $y_0 \in \mathbb{R}$ such that $h'(y_0) = 0$ (note y_0 is necessarily unique), then

$$I = \frac{e^{ih(y_0)}}{\sqrt{h''(y_0)}} F(y_0) + O_A(ZR^{-A}), \qquad (3.10)$$

where $F(y_0)$ is an Y-inert function (depending on A) supported on $y_0 \simeq Z$.

Finally, we also need the rth derivative test with $r \ge 2$ (see [12, Lemma 5.1.3, Lemma 5.1.4]).

Lemma 3.9. Let h(x) be real and r times differentiable on the open interval (a, b) with $h^{(r)}(x) \gg \lambda_0 > 0$ on (a, b). Let w(x) be real on [a, b] and let V_0 be its total variation on [a, b] plus the maximum modulus of w(x) on [a, b]. Then

$$I \ll \frac{V_0}{{\lambda_0}^{1/r}}.$$

4. Setting

Now we start to prove Theorem 1.1.

4.1. Approximate functional equation. By the approximate functional equation [13, Theorem 5.3], we have

$$L\left(\frac{1}{2} + it, f \times g \times \chi\right) \ll (p^{\kappa}t)^{\varepsilon} \sup_{N \ll (p^{\kappa}t)^{2+\varepsilon}} \frac{S(N)}{N^{1/2}} + (p^{\kappa}t)^{-A}$$
(4.1)

for A > 0. Here

$$S(N) = \sum_{n=1}^{\infty} \lambda_f(n) \lambda_g(n) n^{-it} \chi(n) W\left(\frac{n}{N}\right)$$

with W some fixed smooth functions supported in [1, 2] and satisfying $W^{(j)}(x) \ll_j 1$. Note that by the Cauchy–Schwarz inequality and the Rankin–Selberg estimate (3.6), we have the trivial bound $S(N) \ll_{f,g,\varepsilon} N$. Therefore our goal is to obtain any extra savings on this trivial bound at this point.

4.2. Applying DFI's delta method. We write

$$S(N) = \sum_{n=1}^{\infty} \lambda_g(n) V\left(\frac{n}{N}\right) \sum_{m=1}^{\infty} \lambda_f(m) m^{-it} \chi(m) W\left(\frac{m}{N}\right) \delta(m-n),$$

where V is a smooth function supported in [1/2, 5/2], V(x) = 1 for $x \in [1, 2]$ and $V^{(j)}(x) \ll_j 1$ for any $j \in \mathbb{N}$. Applying Lemma 3.1 with $Q = \sqrt{N/Kp^{\lambda}}$, we express S(N) as

$$\begin{split} &\sum_{n=1}^{\infty} \lambda_g(n) V\left(\frac{n}{N}\right) \sum_{m=1}^{\infty} \lambda_f(m) m^{-it} \chi(m) W\left(\frac{m}{N}\right) \\ &\times \left(\sum_{k=0}^{\lambda} \frac{1}{Q} \sum_{\substack{q \leq Q \\ (q,p)=1}} \frac{1}{q p^{\lambda}} \sum_{\substack{a \bmod q p^{\lambda-k}}} e\left(\frac{a(m-n)}{q p^{\lambda-k}}\right) \right) \\ &\int_{\mathbb{R}} U_0\left(\frac{u}{N^{\varepsilon}}\right) g(q,u) e\left(\frac{(m-n)u}{Q q p^{\lambda}}\right) du \\ &+ \sum_{r=1}^{\lfloor \log Q/\log p \rfloor} \frac{1}{Q} \sum_{\substack{q \leq Q/p^r \\ (q,p)=1}} \frac{1}{q p^{\lambda+r}} \sum_{\substack{a \bmod q p^{\lambda+r}}} e\left(\frac{a(m-n)}{q p^{\lambda+r}}\right) \\ &\int_{\mathbb{R}} U_0\left(\frac{u}{N^{\varepsilon}}\right) g(p^r q,u) e\left(\frac{(m-n)u}{Q q p^{\lambda+r}}\right) du \right) + O_A(N^{-A}) \end{split}$$

for any A > 0. Note that λ and K are parameters to be determined later with $\lambda \in \mathbb{N}$ $(2 \leq \lambda < \kappa)$ and 0 < K < t.

In the following we will only treat the terms with k = 0 and r = 0, because the terms with $k \neq 0$ or $r \neq 0$ are lower order terms that can be handled by the same way. Inserting a smooth partition of unity for the *u*-integral and a dyadic partition for the *q*-sum, we have

$$S(N) \ll N^{\varepsilon} \sup_{X \ll N^{\varepsilon}} \sup_{1 \ll R \ll Q} |S^{\flat}(N)| + N^{-A}$$

$$(4.2)$$

for any A > 0, where

$$S^{\flat}(N) = \sum_{n=1}^{\infty} \lambda_g(n) V\left(\frac{n}{N}\right) \sum_{m=1}^{\infty} \lambda_f(m) m^{-it} \chi(m) W\left(\frac{m}{N}\right) \\ \times \frac{1}{Q} \sum_{\substack{q \sim R \\ (q,p)=1}} \frac{1}{q p^{\lambda}} \sum_{\substack{a \bmod q p^{\lambda}}} e\left(\frac{a(m-n)}{q p^{\lambda}}\right) \int_{\mathbb{R}} U\left(\frac{\pm u}{X}\right) g(q,u) e\left(\frac{(m-n)u}{Q q p^{\lambda}}\right) du.$$
(4.3)

Here U is a fixed compactly supported 1-inert function with supp $U \subset [1, 2]$.

Rearranging the order of the sums and integrals in (4.3) we get

$$S^{\flat}(N) = \frac{1}{Q} \sum_{\substack{q \sim R \\ (q,p)=1}} \frac{1}{qp^{\lambda}} \sum_{a \bmod qp^{\lambda}} \int_{\mathbb{R}} U\left(\frac{\pm u}{X}\right) g(q, u) \sum_{n=1}^{\infty} \lambda_{g}(n) e\left(-\frac{na}{qp^{\lambda}}\right) e\left(-\frac{nu}{Qqp^{\lambda}}\right) V\left(\frac{n}{N}\right)$$
$$\times \sum_{m=1}^{\infty} \lambda_{f}(m) m^{-it} \chi(m) e\left(\frac{ma}{qp^{\lambda}}\right) e\left(\frac{mu}{Qqp^{\lambda}}\right) W\left(\frac{m}{N}\right) du.$$
(4.4)

Without loss of generality, we only consider the contribution from u > 0 and t > 0 (the contribution from u < 0 or t < 0 can be estimated similarly). By abuse of notation, we still write the contribution from u > 0 and t > 0 as $S^{\flat}(N)$.

5. VORONOI SUMMATION FORMULAS

In this section, we transform the m and n sums in (4.4) by Voronoi summation formulas. We first apply the GL₂ Voronoi summation formula to the n-sum

$$S_1 := \sum_{n=1}^{\infty} \lambda_g(n) e\left(-\frac{na}{qp^{\lambda}}\right) e\left(-\frac{nu}{Qqp^{\lambda}}\right) V\left(\frac{n}{N}\right).$$

Applying Lemma 3.4 with $\varphi_1(x) = e\left(-\frac{Nux}{Qqp^{\lambda}}\right)V(x)$, we obtain that

$$S_1 = \frac{N}{qp^{\lambda}} \sum_{\pm} \sum_{n=1}^{\infty} \lambda_g(n) e\left(\pm \frac{\overline{a}n}{qp^{\lambda}}\right) \Phi_1^{\pm}\left(\frac{nN}{q^2 p^{2\lambda}}\right)$$
(5.1)

where if g is holomorphic, $\Phi_1^+(y) = \Phi_1(y)$ with $\Phi_1(y)$ given by (3.4) and $\Phi_1^-(y) = 0$, while for g a Hecke–Maass cusp form, $\Phi_1^{\pm}(y)$ are given by (3.7).

Lemma 5.1. (1) For $y = \frac{nN}{q^2 p^{2\lambda}} \gg N^{\varepsilon}$, we have $\Phi_1^-(y) \ll N^{-A}$. For $y = \frac{nN}{q^2 p^{2\lambda}} \gg N^{\varepsilon}$ and $n \asymp \frac{Nu^2}{Q^2} \asymp \frac{NX^2}{Q^2}$, we have $\Phi_1^+(y) = y^{-\frac{1}{2}} V_1\left(y \frac{Q^2 q^2 p^{2\lambda}}{N^2 u^2}\right) e\left(y \frac{Q q p^{\lambda}}{N u}\right) + O(N^{-A}).$ (2) For $y = \frac{nN}{q^2 p^{2\lambda}} \ll N^{\varepsilon}$ and $\frac{NX}{p^{\lambda} RQ} \gg N^{\varepsilon}$, we have $\Phi_1^{\pm}(y) \ll N^{-A}$. (3) For $y = \frac{nN}{q^2 p^{2\lambda}} \ll N^{\varepsilon}$ and $\frac{NX}{p^{\lambda} RQ} \ll N^{\varepsilon}$, we have $\Phi_1^{\pm}(y) \ll N^{\varepsilon}$.

Proof. For $y = \frac{nN}{q^2 p^{2\lambda}} \gg N^{\varepsilon}$, we know that $\Phi_1^-(y) \ll N^{-A}$ by (3.8), and

$$\Phi_1^+(y) = y^{-\frac{1}{4}} \int_0^\infty V(\xi) \xi^{-\frac{1}{4}} e\left(-\frac{Nu}{Qqp^{\lambda}} \xi + 2\sigma_1 \sqrt{y\xi}\right) d\xi + O(N^{-A})$$
(5.2)

with $\sigma_1 = \pm 1$. For $\Phi_1^+(y)$, it is negligibly small by repeated integration by parts unless $\sqrt{y} \approx \frac{Nu}{Qqp^{\lambda}}$ and $\operatorname{sgn} \sigma_1 = +$, in which case we apply stationary phase method to the ξ -integral in (5.2). Let

$$h_1(\xi) = -\frac{Nu}{Qqp^{\lambda}}\xi + 2\sqrt{y\xi}.$$

Then we have

$$h_1'(\xi) = -\frac{Nu}{Qqp^{\lambda}} + \sqrt{y}\xi^{-1/2}, \quad h_1''(\xi) = -\frac{1}{2}\sqrt{y}\xi^{-3/2} \asymp \sqrt{y}, \quad h_1^{(j)}(\xi) \ll_j \sqrt{y}, \ j \ge 3.$$

Note that the stationary point is $\xi_{01} = y \frac{Q^2 q^2 p^{2\lambda}}{N^2 u^2}$ and we have $h_1(\xi_{01}) = y \frac{Q q p^{\lambda}}{N u}$. Let $w_1(\xi) = V(\xi) \xi^{-1/4}$. Then we have $w_1^{(j)}(\xi) \ll_j 1$ and $\operatorname{supp} w_1 \subset [1/4, 4]$. By Lemma 3.8(2), we obtain that

$$\Phi_1^+(y) = y^{-\frac{1}{2}} V_1\left(y \frac{Q^2 q^2 p^{2\lambda}}{N^2 u^2}\right) e\left(y \frac{Q q p^{\lambda}}{N u}\right) + O(N^{-A}),$$
(5.3)

where V_1 is an 1-inert function (depending on A) supported on $\xi_{01} \approx 1$. Hence, case (1) is proved.

For $y = \frac{nN}{q^2p^{2\lambda}} \ll N^{\varepsilon}$, by (3.7) we may regard $\Phi_1^{\pm}(y)$ as

$$\Phi_1^{\pm}(y) = \int_0^\infty W(\xi) e\left(-\frac{Nu\xi}{Qqp^{\lambda}}\right) J_g^{\pm}(y\xi) \mathrm{d}\xi, \qquad (5.4)$$

where

$$J_g^+(y) = \frac{-\pi}{\sin(\pi i t_g)} \left(J_{2it_g}(4\pi\sqrt{y}) - J_{-2it_g}(4\pi\sqrt{y}) \right),$$

and

$$J_g^-(y) = 4\varepsilon_g \cosh(\pi t_g) K_{2it_g}(4\pi\sqrt{y})$$

Then, by partial integration and (3.9), $\Phi_1^{\pm}(y)$ is negligible unless $\frac{Nu}{Qqp^{\lambda}} \ll N^{\varepsilon}$, in which case $\Phi_1^{\pm}(y) \ll N^{\varepsilon}$. Therefore, case (2) and (3) are proved.

According to Lemma 5.1, we first consider the estimations for the case $\frac{nN}{q^2p^{2\lambda}} \simeq \left(\frac{NX}{RQp^{\lambda}}\right)^2 \gg N^{\varepsilon}$. For the complementary range $\frac{nN}{q^2p^{2\lambda}} \ll N^{\varepsilon}$ and $\frac{NX}{p^{\lambda}RQ} \ll N^{\varepsilon}$, we will analyse the contribution of the case in §10.

Plugging (5.3) into (5.1), we find that the *n*-sum is asymptotically equal to

$$N^{\frac{1}{2}} \sum_{n} \frac{\lambda_g(n)}{n^{1/2}} e\left(\frac{\overline{a}n}{qp^{\lambda}}\right) V_1\left(\frac{nQ^2}{Nu^2}\right) e\left(\frac{nQ}{qp^{\lambda}u}\right),$$

and we have $n \asymp \frac{Nu^2}{Q^2} \asymp \frac{NX^2}{Q^2}$. Hence

$$S^{\flat}(N) = \frac{N^{\frac{1}{2}}}{Q} \sum_{\substack{q \sim R \\ (q,p)=1}} \frac{1}{qp^{\lambda}} \sum_{a \bmod qp^{\lambda}} \sum_{n \asymp \frac{NX^{2}}{Q^{2}}} \frac{\lambda_{g}(n)}{n^{1/2}} e\left(\frac{\overline{a}n}{qp^{\lambda}}\right) \sum_{m} \lambda_{f}(m)\chi(m)m^{-it}e\left(\frac{am}{qp^{\lambda}}\right) V\left(\frac{m}{N}\right)$$
$$\cdot \int_{\mathbb{R}} U\left(\frac{u}{X}\right) g(q,u)V_{1}\left(\frac{nQ^{2}}{Nu^{2}}\right) e\left(\frac{nQ}{qp^{\lambda}u} + \frac{mu}{Qqp^{\lambda}}\right) \mathrm{d}u + O(N^{-A}).$$

Making a change of variable $u = X\xi$, we have

$$I(X) = X \int_{\mathbb{R}} U(\xi)g(q, X\xi)V_1\left(\frac{nQ^2}{NX^2\xi^2}\right)e\left(\frac{nQ}{p^{\lambda}qX\xi} + \frac{mX\xi}{Qqp^{\lambda}}\right)d\xi.$$
 (5.5)

Next, we apply stationary phase method to (5.5). Let

$$h_2(\xi) = \frac{nQ}{p^\lambda q X \xi} + \frac{mX\xi}{Qqp^\lambda}.$$

Then we have

$$h_2'(\xi) = -\frac{nQ}{p^\lambda q X \xi^2} + \frac{mX}{Qqp^\lambda}, \quad h_2''(\xi) = \frac{2nQ}{p^\lambda q X \xi^3} \asymp \frac{NX}{p^\lambda RQ}$$

Let $h'_{2}(\xi_{02}) = 0$. Then we have the stationary point $\xi_{02} = \frac{n^{1/2}Q}{m^{1/2}X}$ and $h_{2}(\xi_{02}) = 2\frac{m^{1/2}n^{1/2}}{qp^{\lambda}}$. By Lemma 3.8(2), we obtain that

$$I(X) = X\left(\frac{nQ}{p^{\lambda}qX}\right)^{-1/2} V_2\left(\frac{nQ^2}{mX^2}\right) e\left(2\frac{m^{1/2}n^{1/2}}{qp^{\lambda}}\right) + O(N^{-A}),$$
(5.6)

where V_2 is an 1-inert function (depending on A) supported on $\xi_{02} \simeq 1$. Hence

$$S^{\flat}(N) = \frac{X^{\frac{3}{2}}N^{\frac{1}{2}}}{Q^{\frac{3}{2}}} \sum_{\substack{q \sim R \\ (q,p)=1}} \frac{1}{(qp^{\lambda})^{1/2}} \sum_{a \bmod qp^{\lambda}} \sum_{n \asymp \frac{NX^{2}}{Q^{2}}} \frac{\lambda_{g}(n)}{n} e\left(\frac{\overline{a}n}{qp^{\lambda}}\right)$$
$$\cdot \sum_{m} \lambda_{f}(m)\chi(m)m^{-it}e\left(\frac{am}{qp^{\lambda}}\right) V\left(\frac{m}{N}\right) V_{2}\left(\frac{nQ^{2}}{mX^{2}}\right) e\left(2\frac{m^{1/2}n^{1/2}}{qp^{\lambda}}\right) + O(N^{-A}).$$
(5.7)

Let

$$S_2 := \sum_m \lambda_f(m) \chi(m) m^{-it} e\left(\frac{am}{qp^{\lambda}}\right) e\left(2\frac{m^{1/2}n^{1/2}}{qp^{\lambda}}\right) V\left(\frac{m}{N}\right) V_2\left(\frac{nQ^2}{mX^2}\right).$$

By the Fourier expansion of χ in terms of additive characters, we have

$$S_2 = \frac{1}{\tau(\overline{\chi})} \sum_{c \bmod p^{\kappa}} \overline{\chi}(c) \sum_m \lambda_f(m) e\left(\frac{(ap^{\kappa-\lambda} + cq)m}{qp^{\kappa}}\right) m^{-it} e\left(2\frac{m^{1/2}n^{1/2}}{qp^{\lambda}}\right) V\left(\frac{m}{N}\right) V_2\left(\frac{nQ^2}{mX^2}\right)$$

Note that $(ap^{\kappa-\lambda} + cq, q) = 1$ and $(ap^{\kappa-\lambda} + cq, p) = (cq, p) = 1$. Thus $(ap^{\kappa-\lambda} + cq, qp^{\kappa}) = 1$. Denote $a^* = ap^{\kappa-\lambda} + cq$.

Next, we apply the GL_2 Voronoi summation formula in Lemma 3.4 with

$$\varphi_2(x) = x^{-it} e\left(2\frac{N^{1/2}n^{1/2}}{qp^{\lambda}}x^{1/2}\right)V(x)V_2\left(\frac{nQ^2}{NX^2x}\right)$$

and obtain that

$$S_2 = \frac{1}{\tau(\overline{\chi})} \frac{N^{1-it}}{qp^{\kappa}} \sum_{c \bmod p^{\kappa}} \overline{\chi}(c) \sum_m \lambda_f(m) e\left(\frac{-\overline{a^*}m}{qp^{\kappa}}\right) \Phi_2^{\pm}\left(\frac{mN}{q^2 p^{2\kappa}}\right),$$

where $\Phi_2^{\pm}(y)$ are given by (3.7).

Lemma 5.2. If $y = \frac{mN}{q^2 p^{2\kappa}} \ll N^{\varepsilon}$, then $\Phi_2^{\pm}(y)$ is negligible unless $t \simeq \frac{N^{1/2} n^{1/2}}{q p^{\lambda}} \simeq \frac{NX}{p^{\lambda} RQ}$. *Proof.* For $\frac{mN}{q^2 p^{2\kappa}} \ll N^{\varepsilon}$, by (3.7) we may regard $\Phi_2^{\pm}(y)$ as

$$\Phi_2^{\pm}(y) = \int_0^\infty V(\xi) V_2\left(\frac{nQ^2}{NX^2\xi}\right) e\left(-\frac{t\log\xi}{2\pi} + 2\frac{N^{1/2}n^{1/2}}{qp^{\lambda}}\xi^{1/2}\right) J_f^{\pm}(y\xi) \mathrm{d}\xi, \qquad (5.8)$$

where

$$J_f^+(y) = \frac{-\pi}{\sin(\pi i t_f)} \left(J_{2it_f}(4\pi\sqrt{y}) - J_{-2it_f}(4\pi\sqrt{y}) \right),$$

and

$$J_f^-(y) = 4\varepsilon_f \cosh(\pi t_f) K_{2it_f}(4\pi \sqrt{y}).$$

Then, by partial integration together with (3.9), $\Phi_2^{\pm}(y)$ is negligible unless $t \approx \frac{N^{1/2}n^{1/2}}{qp^{\lambda}} \approx \frac{NX}{p^{\lambda}RQ}$.

We will analyse the contribution of the case $\frac{mN}{q^2p^{2\kappa}} \ll N^{\varepsilon}$ to S(N) in §10. In this section we consider the estimations for the case $\frac{mN}{q^2p^{2\kappa}} \gg N^{\varepsilon}$.

For $y = \frac{mN}{q^2 p^{2\kappa}} \gg N^{\varepsilon}$, in view of (3.8), $\Phi_2^-(y)$ is negligible. By Lemma 3.5, we have $\Phi_2^+(y) = y^{-\frac{1}{4}} \int_0^\infty V(\xi) V_2\left(\frac{nQ^2}{NX^2\xi}\right) \xi^{-\frac{1}{4}} e\left(-\frac{t\log\xi}{2\pi} + 2\frac{N^{1/2}n^{1/2}}{qp^{\lambda}}\xi^{1/2} + 2\sigma_2\sqrt{y\xi}\right) d\xi + O(N^{-A})$ (5.9)

with $\sigma_2 = \pm 1$. Making a change of variable $\xi^{1/2} \to \xi$ and taking $y = \frac{mN}{q^2 p^{2\kappa}}$, we obtain

$$\Phi_2^+ \left(\frac{mN}{q^2 p^{2\kappa}}\right) = 2\left(\frac{mN}{q^2 p^{2\kappa}}\right)^{-\frac{1}{4}} \int_0^\infty V_3(\xi) e\left(-\frac{t\log\xi}{\pi} + 2\frac{N^{1/2}n^{1/2}}{qp^{\lambda}}\xi + 2\sigma_2\frac{\sqrt{mN}}{qp^{\kappa}}\xi\right) d\xi + O(N^{-A}),$$

where $V_3(\xi) = V(\xi^2) V_2\left(\frac{nQ^2}{NX^2\xi^2}\right) \xi^{\frac{1}{2}}$ is an 1-inert function.

By repeated integration by parts, one can truncate the *m*-sum at $m \ll \max\{\frac{t^2 R^2 p^{2\kappa}}{N}, \frac{N X^2 p^{2\kappa-2\lambda}}{Q^2}\}$. Hence, it is sufficient to estimate

$$\frac{N^{5/4-it}X^{3/2}}{Q^{3/2}\tau(\overline{\chi})p^{(\kappa+\lambda)/2}} \sum_{\substack{q\sim R\\(q,p)=1}} \frac{1}{q} \sum_{\substack{n\asymp \frac{NX^2}{Q^2}}} \frac{\lambda_g(n)}{n} \sum_{\substack{m\ll\max\{\frac{t^2R^2p^{2\kappa}}{N}, \frac{NX^2p^{2\kappa-2\lambda}}{Q^2}\}}} \frac{\lambda_f(m)}{m^{1/4}} \sum_{\substack{a \bmod qp^{\lambda}}} \left(\frac{\overline{a}n}{qp^{\lambda}}\right) \\
\cdot \sum_{c \bmod p^{\kappa}} \overline{\chi}(c)e\left(\frac{-\overline{a^*}m}{qp^{\kappa}}\right) \int_0^\infty V_3(\xi)e\left(-\frac{t\log\xi}{\pi} + 2\frac{N^{1/2}n^{1/2}}{qp^{\lambda}}\xi + 2\sigma_2\frac{N^{1/2}m^{1/2}}{qp^{\kappa}}\xi\right) d\xi.$$
(5.10)

Note that

$$e\left(\frac{-\overline{a^*}m}{qp^{\kappa}}\right) = e\left(\frac{-(\overline{ap^{\kappa-\lambda}+cq})m}{qp^{\kappa}}\right) = e\left(\frac{-(\overline{ap^{\kappa-\lambda}+cq})m\bar{q}}{p^{\kappa}}\right)e\left(\frac{-\overline{ap^{2\kappa-\lambda}}m}{q}\right)$$

The character sum in (5.10) is

$$\sum_{a \mod qp^{\lambda}}^{\star} \left(\frac{\overline{a}n}{qp^{\lambda}}\right) \sum_{c \mod p^{\kappa}} \overline{\chi}(c) e\left(\frac{-\overline{a^{*}m}}{qp^{\kappa}}\right)$$
$$= \sum_{c \mod p^{\kappa}} \overline{\chi}(c) \sum_{b \mod p^{\lambda}}^{\star} e\left(\frac{-(\overline{ap^{\kappa-\lambda}+cq})m\bar{q}}{p^{\kappa}} + \frac{n\bar{q}\bar{b}}{p^{\lambda}}\right) \sum_{n \equiv mp^{\lambda} \frac{d|q}{p^{2\kappa-\lambda}} \mod d} d\mu\left(\frac{q}{d}\right).$$
(5.11)

Plugging (5.11) into (5.10), we are led to estimate

$$\frac{N^{5/4-it}X^{3/2}}{Q^{3/2}\tau(\overline{\chi})p^{(\kappa+\lambda)/2}} \sum_{\substack{q\sim R\\(q,p)=1}} \frac{1}{q} \sum_{d|q} d\mu\left(\frac{q}{d}\right) \sum_{\substack{n\asymp \frac{NX^2}{Q^2}}} \frac{\lambda_g(n)}{n} \\
\times \sum_{\substack{m\ll \max\{\frac{t^2R^2p^{2\kappa}}{N}, \frac{NX^2p^{2\kappa-2\lambda}}{Q^2}\}\\n\equiv mp^{\lambda}\overline{p^{2\kappa-\lambda}} \mod d}} \frac{\lambda_f(m)}{m^{1/4}} \mathfrak{C}^*(m,n,q)I(m,n,q)$$
(5.12)

with

$$\mathfrak{C}^*(m,n,q) := \sum_{c \bmod p^{\kappa}} \overline{\chi}(c) \sum_{b \bmod p^{\lambda}} e\left(\frac{-(\overline{ap^{\kappa-\lambda} + cq})m\overline{q}}{p^{\kappa}} + \frac{n\overline{q}\overline{b}}{p^{\lambda}}\right)$$

and

$$I(m,n,q) := \int_0^\infty V_3(\xi) e\left(-\frac{t\log\xi}{\pi} + 2\frac{N^{1/2}n^{1/2}}{qp^{\lambda}}\xi + 2\sigma_2\frac{N^{1/2}m^{1/2}}{qp^{\kappa}}\xi\right) \mathrm{d}\xi.$$

6. Applying Cauchy–Schwarz inequality and Poisson summation formula

Applying the Cauchy–Schwarz inequality to the m-sum in (5.12) and using the Rankin–Selberg estimate in (3.6), we get

$$S^{\flat}(N) \ll \sup_{\substack{M_{1} \ll \max\{\frac{t^{2}R^{2}p^{2\kappa}}{N}, \frac{NX^{2}p^{2\kappa-2\lambda}}{Q^{2}}\}}} \frac{N^{5/4}X^{3/2}}{Q^{3/2}p^{(2\kappa+\lambda)/2}} \sum_{\substack{q \sim R \\ (q,p)=1}} \frac{1}{q} \sum_{d|q} d$$

$$\cdot \sum_{m \sim M_{1}} \frac{|\lambda_{f}(m)|}{m^{1/4}} \left| \sum_{\substack{n \simeq \frac{NX^{2}}{Q^{2}}\\ n \equiv mp^{\lambda}p^{2\kappa-\lambda} \bmod d}} \frac{\lambda_{g}(n)}{n} \mathfrak{C}^{*}(m,n,q)I(m,n,q) \right| \qquad (6.1)$$

$$\ll \sup_{M_{1} \ll \max\{\frac{t^{2}R^{2}p^{2\kappa}}{N}, \frac{NX^{2}p^{2\kappa-2\lambda}}{Q^{2}}\}} \frac{N^{5/4}X^{3/2}M_{1}^{1/4}}{Q^{3/2}p^{(2\kappa+\lambda)/2}} \sum_{\substack{q \sim R \\ (q,p)=1}} \frac{1}{q} \sum_{d|q} d \cdot \Sigma(d,q)^{1/2},$$

where

$$\Sigma(d,q) := \sum_{m} W_1\left(\frac{m}{M_1}\right) \left| \sum_{\substack{n \leq \frac{NX^2}{Q^2}\\n \equiv mp^{\lambda} \overline{p^{2\kappa-\lambda}} \bmod d}} \frac{\lambda_g(n)}{n} \mathfrak{C}^*(m,n,q) I(m,n,q) \right|^2$$

with W_1 supported on [1,2] and satisfying $W_1^{(j)}(x) \ll_j 1$. Expanding the absolute square and switching the order of summations, we obtain

$$\Sigma(d,q) = \sum_{\substack{n_1 \asymp \frac{NX^2}{Q^2}}} \frac{\lambda_g(n_1)}{n_1} \sum_{\substack{n_2 \asymp \frac{NX^2}{Q^2} \\ n_2 \equiv n_1 \bmod d}} \frac{\lambda_g(n_2)}{n_2} \sum_{\substack{m \ge 1 \\ m \equiv n_1 \overline{p^\lambda} p^{2\kappa - \lambda} \bmod d}} W_1\left(\frac{m}{M_1}\right) \\ \cdot \mathfrak{C}^*(m,n_1,q) \overline{\mathfrak{C}^*(m,n_2,q)} I(m,n_1,q) \overline{I(m,n_2,q)}.$$

Note that $\lambda_g(n_1)\lambda_g(n_2) \ll |\lambda_g(n_1)|^2 + |\lambda_g(n_2)|^2$. Hence we have

$$\Sigma(d,q) \ll \frac{Q^4}{N^2 X^4} \sum_{n_1 \asymp \frac{NX^2}{Q^2}} |\lambda_g(n_1)|^2 \sum_{\substack{n_2 \asymp \frac{NX^2}{Q^2} \\ n_2 \equiv n_1 \bmod d}} |\mathcal{K}|,$$
(6.2)

where

$$\mathcal{K} = \sum_{\substack{m \ge 1 \\ m \equiv n_1 \overline{p^{\lambda}} p^{2\kappa - \lambda} \mod d}} W_1\left(\frac{m}{M_1}\right) \mathfrak{C}^*(m, n_1, q) \overline{\mathfrak{C}^*(m, n_2, q)} I(m, n_1, q) \overline{I(m, n_2, q)}.$$

Applying the Poisson summation formula with modulo dp^{κ} , we get

$$\mathcal{K} = \frac{M_1}{dp^{\kappa}} \sum_{m \in \mathbb{Z}} \mathfrak{C}(m, n_1, n_2) \cdot \mathfrak{I}(m, n_1, n_2), \tag{6.3}$$

where

$$\mathfrak{C}(m,n_1,n_2) = \sum_{\gamma \bmod p^{\kappa}} \mathfrak{C}^*(\gamma,n_1,q) \overline{\mathfrak{C}^*(\gamma,n_2,q)} e\left(\frac{m\left(n_1 p^{3\kappa-\lambda} \overline{p^{\kappa+\lambda}} + \gamma d\bar{d}\right)}{dp^{\kappa}}\right)$$
(6.4)

and

$$\mathfrak{I}(m,n_1,n_2) := \int_{\mathbb{R}} W_1(z) I(M_1z,n_1,q) \overline{I(M_1z,n_2,q)} e\left(-\frac{mM_1}{p^{\kappa}d}z\right) \mathrm{d}z \tag{6.5}$$

with

$$I(M_1z, n, q) = \int_0^\infty V_3(\xi) e\left(-\frac{t\log\xi}{\pi} + 2\frac{N^{1/2}n^{1/2}}{qp^{\lambda}}\xi + 2\sigma_2\frac{N^{1/2}M_1^{1/2}z^{1/2}}{qp^{\kappa}}\xi\right) d\xi.$$
(6.6)

For the character sum, Sun [22, Lemma 4] has the following estimation:

Lemma 6.1. Assume $(2\kappa+1)/3 \leq \lambda \leq 3\kappa/4$. The character sum vanishes unless $(n_1n_2, p) = 1$ and $p^{\kappa-\lambda}|m$. Let $m = m'p^{\kappa-\lambda}$, $\lambda = 2\alpha + \delta_1$ and $\kappa = 2\beta + \delta_2$ with $\delta_1, \delta_2 = 0$ or $1, \alpha \geq 1$ and $\beta \geq 1$.

(1) If
$$p^{\beta-\kappa+\lambda}|m'$$
, then $n_1 \equiv n_2 \pmod{p^{\beta-\kappa+\lambda}}$ and
 $\mathfrak{C}(m, n_1, n_2) \ll p^{2\kappa+\lambda+\delta_1}$.
(2) If $p^{\ell}||m'|$ with $\ell < \beta - \kappa + \lambda$, then $p^{\ell}||n_1 - n_2|$ and
 $\mathfrak{C}(m, n_1, n_2) \ll p^{5\kappa/2+2\ell+\delta_1+\delta_2/2}$.

(3) For m = 0, we have

$$\mathfrak{C}(m, n_1, n_2) = p^{2\kappa} \sum_{d \mid (n_1 - n_2, p^{\lambda})} d\mu \left(p^{\lambda}/d \right).$$

Next, we deal with $\Im(m, n_1, n_2)$.

6.1. The large modulo case. Assume $N^{\varepsilon} \ll \frac{NX}{p^{\lambda}RQ} \ll t^{1-\varepsilon}$. Then we have

$$\frac{N^{1/2}n^{1/2}}{qp^{\lambda}} \asymp \frac{NX}{p^{\lambda}RQ} \ll t^{1-\varepsilon}.$$

We first consider $I(M_1z, n, q)$. For $I(M_1z, n, q)$, it is negligibly small by repeated integration by parts unless $t \approx \frac{N^{1/2}M_1^{1/2}z^{1/2}}{qp^{\kappa}}$ (i.e. $M_1 \approx \frac{t^2R^2p^{2\kappa}}{N}$) and $\operatorname{sgn} \sigma_2 = +$, in which case we apply the stationary phase method to the ξ -integral. Let $h_3(\xi) = -\frac{t\log\xi}{\pi} + 2\frac{N^{1/2}n^{1/2}}{qp^{\lambda}}\xi + 2\frac{N^{1/2}M_1^{1/2}z^{1/2}}{qp^{\kappa}}\xi$. Then we have

$$h_3'(\xi) = -\frac{t}{\pi\xi} + 2\frac{N^{1/2}n^{1/2}}{qp^{\lambda}} + 2\frac{N^{1/2}M_1^{1/2}z^{1/2}}{qp^{\kappa}}, \quad h_3''(\xi) = \frac{t}{\pi\xi^2} \asymp t.$$

The stationary point is

$$\xi_{03} = \frac{t}{2\pi \left(\frac{N^{1/2}M_1^{1/2}z^{1/2}}{qp^{\kappa}} + \frac{N^{1/2}n^{1/2}}{qp^{\lambda}}\right)},$$

and

$$h_3(\xi_{03}) = \frac{t}{\pi} \log \frac{2e\pi(\frac{N^{1/2}M_1^{1/2}z^{1/2}}{qp^{\kappa}} + \frac{N^{1/2}n^{1/2}}{qp^{\lambda}})}{t}$$

By Lemma 3.8(2), we have

$$I(M_1z, n, q) = t^{-\frac{1}{2}} V_4 \left(\frac{\frac{N^{1/2} M_1^{1/2} z^{1/2}}{qp^{\kappa}} + \frac{N^{1/2} n^{1/2}}{qp^{\lambda}}}{t/2\pi} \right) e \left(\frac{t}{\pi} \log \frac{2e\pi \left(\frac{N^{1/2} M_1^{1/2} z^{1/2}}{qp^{\kappa}} + \frac{N^{1/2} n^{1/2}}{qp^{\lambda}}\right)}{t} \right),$$
(6.7)

where V_4 is an 1-inert function supported on $\xi_{03} \simeq 1$. The oscillation in the above exponential function is of size t, which is quite large. Next, we will reduce the oscillation. This is one of the key observations in our paper. Let

$$h(M_1z, n, q) := \frac{t}{\pi} \log \frac{2e\pi \left(\frac{N^{1/2}M_1^{1/2}z^{1/2}}{qp^{\kappa}} + \frac{N^{1/2}n^{1/2}}{qp^{\lambda}}\right)}{t}$$

$$= \frac{t}{\pi} \log \frac{2e\pi}{t} + \frac{t}{\pi} \log \frac{N^{1/2}M_1^{1/2}z^{1/2}}{qp^{\kappa}} + \frac{t}{\pi} \log \left(1 + \frac{p^{\kappa - \lambda}n^{1/2}}{(M_1z)^{1/2}}\right).$$
(6.8)

Since

$$p^{2\kappa-2\lambda}n \asymp p^{2\kappa-2\lambda}\frac{NX^2}{Q^2} \ll p^{2\kappa-2\lambda}\frac{R^2p^{2\lambda}t^2}{N}t^{-2\varepsilon} \asymp M_1t^{-2\varepsilon},$$

by the Taylor expansion we get

$$\log\left(1 + \frac{p^{\kappa - \lambda} n^{1/2}}{(M_1 z)^{1/2}}\right) = \frac{p^{\kappa - \lambda} n^{1/2}}{(M_1 z)^{1/2}} + \frac{p^{2\kappa - 2\lambda} n}{2M_1 z} + \sum_{j \ge 3} c_j \left(\frac{p^{\kappa - \lambda} n^{1/2}}{(M_1 z)^{1/2}}\right)^j.$$
(6.9)

By (6.8) and (6.9), we have

$$h(M_1z, n, q) = h_1(M_1z, q) + h_2(M_1z, n, q),$$
(6.10)

where

$$h_1(M_1z,q) = \frac{t}{\pi} \log \frac{2e\pi}{t} + \frac{t}{\pi} \log \frac{N^{1/2} M_1^{1/2} z^{1/2}}{q p^{\kappa}}$$

and

$$h_2(M_1z, n, q) = \frac{t}{\pi} \left(\frac{p^{\kappa - \lambda} n^{1/2}}{(M_1 z)^{1/2}} + \frac{p^{2\kappa - 2\lambda} n}{2M_1 z} + \sum_{j \ge 3} c_j \left(\frac{p^{\kappa - \lambda} n^{1/2}}{(M_1 z)^{1/2}} \right)^j \right).$$

Plugging (6.7) and (6.10) into (6.5), we get

$$\Im(m, n_1, n_2) = t^{-1} \int_{\mathbb{R}} W_2(z) e\left(h_2\Big(M_1 z, n_1, q\Big) - h_2\Big(M_1 z, n_2, q\Big) - \frac{mM_1}{p^{\kappa} d}z\right) dz, \quad (6.11)$$

where W_2 is an 1-inert function.

Then we deal with z-integral.

Lemma 6.2. We have

(1) If $m \gg \frac{N^2 X d}{R^3 Q t^2 p^{\kappa+\lambda}} N^{\varepsilon}$, then $\Im(m, n_1, n_2) \ll N^{-A}$. (2) If $\frac{dN}{R^2 p^{\kappa} t^2} N^{\varepsilon} \ll m \ll \frac{N^2 X d}{R^3 Q t^2 p^{\kappa+\lambda}} N^{\varepsilon}$, then $\Im(m, n_1, n_2) \ll N^{-A}$ unless $|n_1 - n_2| \frac{Q}{p^{\lambda} R X} \asymp \frac{|m|R^2 p^{\kappa} t^2}{N d}$ in which case we have

$$\Im(m, n_1, n_2) \ll \left(\frac{dN}{|m|R^2 p^{\kappa} t^4}\right)^{1/2}.$$

(3) If
$$m \ll \frac{dN}{R^2 p^{\kappa} t^2} N^{\varepsilon}$$
, then $\Im(m, n_1, n_2) \ll N^{-A}$ unless
 $n^{\lambda} R X$

$$|n_1 - n_2| \ll \frac{p^* R X}{Q} N^{\varepsilon},$$

in which case we have $\Im(m, n_1, n_2) \ll t^{-1}$.

Proof. (1). Let

$$\rho(z) = h_2 \Big(M_1 z, n_1, q \Big) - h_2 \Big(M_1 z, n_2, q \Big) - \frac{m M_1}{p^{\kappa} d} z$$

Note that

$$h_2\Big(M_1z, n_1, q\Big) - h_2\Big(M_1z, n_2, q\Big)$$

= $\frac{t}{\pi} \left(\frac{p^{\kappa-\lambda}}{(M_1z)^{1/2}} \left(n_1^{1/2} - n_2^{1/2}\right) + \frac{p^{2\kappa-2\lambda}}{2M_1z} \left(n_1 - n_2\right) + \sum_{j\geq 3} c_j \left(\frac{p^{\kappa-\lambda}}{(M_1z)^{1/2}}\right)^j \left(n_1^{j/2} - n_2^{j/2}\right)\right).$

For $n_1 \simeq n_2 \simeq N X^2 / Q^2$, we have

$$\frac{\partial^k}{\partial z^k} \left(h_2 \Big(M_1 z, n_1, q \Big) - h_2 \Big(M_1 z, n_2, q \Big) \right) \ll_k \frac{NX}{p^\lambda RQ}, \quad k \ge 1$$

By repeated integration by parts, we have $\Im(m, n_1, n_2)$ is negligibly small unless $m \ll$ $\frac{\frac{N^2 X d}{R^3 Q t^2 p^{\kappa+\lambda}} N^{\varepsilon}}{(2). \text{ Note that}}$

$$\left(\frac{p^{\kappa-\lambda}}{(M_1z)^{1/2}}\right)^j \left(n_1^{j/2} - n_2^{j/2}\right) = \frac{p^{\kappa-\lambda}}{(M_1z)^{1/2}} \left(n_1^{1/2} - n_2^{1/2}\right) \left(\frac{p^{\kappa-\lambda}}{(M_1z)^{1/2}}\right)^{j-1} \sum_{i+\ell=j-1} n_1^{i/2} n_2^{\ell/2} \\ \approx \frac{p^{\kappa-\lambda}}{(M_1z)^{1/2}} \left(n_1^{1/2} - n_2^{1/2}\right) \left(\frac{N^{1/2}}{tRp^{\lambda}}\right)^{j-1} \left(\frac{N^{1/2}X}{Q}\right)^{j-1} \\ \ll \frac{p^{\kappa-\lambda}}{(M_1z)^{1/2}} \left(n_1^{1/2} - n_2^{1/2}\right) t^{-(j-1)\varepsilon},$$

for $j \geq 2$. We have

$$\frac{\partial^k}{\partial z^k} \left(h_2 \Big(M_1 z, n_1, q \Big) - h_2 \Big(M_1 z, n_2, q \Big) \right) \asymp_k |n_1^{1/2} - n_2^{1/2}| \frac{N^{1/2}}{p^\lambda R} \asymp |n_1 - n_2| \frac{Q}{p^\lambda R X},$$

for any $k \ge 1$.

By repeated integration by parts, we have $\Im(m, n_1, n_2)$ is negligibly small unless $|n_1 - n_2| \frac{Q}{p^{\lambda}RX} \approx \frac{|m|R^2 p^{\kappa} t^2}{Nd}$. Note that

$$\rho''(z) \asymp |n_1 - n_2| \frac{Q}{p^{\lambda} R X} \asymp \frac{|m| R^2 p^{\kappa} t^2}{N d}$$

By Lemma 3.9 with r = 2, we have

$$\Im(m, n_1, n_2) \ll t^{-1} \left(\frac{|m|R^2 p^{\kappa} t^2}{Nd}\right)^{-1/2} = \left(\frac{dN}{|m|R^2 p^{\kappa} t^4}\right)^{1/2}.$$

(3). For $m \ll \frac{dN}{R^2 p^{\kappa} t^2} N^{\varepsilon}$, by repeated integration by parts, $\Im(m, n_1, n_2)$ is negligibly small unless $|n_1 - n_2| \ll \frac{p_{\lambda RX}}{Q} N^{\varepsilon}$, in which case $\Im(m, n_1, n_2) \ll t^{-1}$. 6.2. The small modulo case. Assume that $\frac{NX}{p^{\lambda}RQ} \gg t^{1-\varepsilon}$. Then we have $R \ll \frac{NX}{p^{\lambda}Qt^{1-\varepsilon}}$. Recall that

$$\Im(m, n_1, n_2) = \int_{\mathbb{R}} W_1(z) I(M_1 z, n_1, q) \overline{I(M_1 z, n_2, q)} e\left(-\frac{mM_1}{p^{\kappa} d}z\right) dz$$

with

$$I(M_1z, n, q) = \int_0^\infty V_3(\xi) e\left(-\frac{t\log\xi}{\pi} + 2\frac{N^{1/2}n^{1/2}}{qp^{\lambda}}\xi + 2\sigma_2\frac{N^{1/2}M_1^{1/2}z^{1/2}}{qp^{\kappa}}\xi\right)d\xi.$$

Lemma 6.3. (1) For $m \gg \frac{N^{1/2}d}{M_1^{1/2}q} N^{\varepsilon}$, we have $\Im(m, n_1, n_2) \ll N^{-A}$. (2) For $\frac{p^{\kappa}d}{M_1} N^{\varepsilon} \ll m \ll \frac{N^{1/2}d}{M_1^{1/2}q} N^{\varepsilon}$, we have $\Im(m, n_1, n_2) \ll \frac{1}{t} \left(\frac{mNM_1^2}{q^2p^{3\kappa}t^2d}\right)^{-1/3}$. (3) For $m \ll \frac{p^{\kappa}d}{M_1} N^{\varepsilon}$, we have $\Im(m, n_1, n_2) \ll \frac{1}{t}$.

Proof. (1). Plugging (6.6) into (6.5) and interchanging the order of integrals, we express $\Im(m, n_1, n_2)$ as

$$\int_{0}^{\infty} \int_{0}^{\infty} V_{3}(\xi_{1}) \overline{V_{3}(\xi_{2})} e\left(-\frac{t \log \xi_{1}}{\pi} + \frac{t \log \xi_{2}}{\pi} + 2\frac{N^{1/2} n_{1}^{1/2}}{q p^{\lambda}} \xi_{1} - \frac{N^{1/2} n_{2}^{1/2}}{q p^{\lambda}} \xi_{2}\right) \\ \cdot \left(\int_{\mathbb{R}} W_{1}(z) e\left(2\sigma_{2} \frac{N^{1/2} (M_{1}z)^{1/2}}{q p^{\kappa}} \xi_{1} - 2\sigma_{2} \frac{N^{1/2} (M_{1}z)^{1/2}}{q p^{\kappa}} \xi_{2} - \frac{m M_{1}}{p^{\kappa} d} z\right) dz\right) d\xi_{1} d\xi_{2}.$$
(6.12)

We first deal with the z-integral in (6.12). Let

$$\rho_2(z) = 2\sigma_2 \frac{N^{1/2} \left(M_1 z\right)^{1/2}}{q p^{\kappa}} \xi_1 - 2\sigma_2 \frac{N^{1/2} \left(M_1 z\right)^{1/2}}{q p^{\kappa}} \xi_2 - \frac{m M_1}{p^{\kappa} d} z$$

Note that

$$\frac{\partial^k}{\partial z^k} \left(2\sigma_2 \frac{N^{1/2} \left(M_1 z\right)^{1/2}}{q p^{\kappa}} \xi_1 - 2\sigma_2 \frac{N^{1/2} \left(M_1 z\right)^{1/2}}{q p^{\kappa}} \xi_2 \right) \ll_k \frac{N^{1/2} \left(M_1\right)^{1/2}}{q p^{\kappa}}, \quad k \ge 1.$$

By repeated integration by parts, we have the z-integral is negligibly small unless $m\ll \frac{N^{1/2}d}{M_1^{1/2}q}N^{\varepsilon}.$

(2). Now we consider the case $m \ll \frac{N^{1/2}d}{M_1^{1/2}q}N^{\varepsilon}$. For i = 1, 2, we have

$$I(M_1z, n_i, q) = \int_0^\infty V_3(\xi_i) e\left(-\frac{t\log\xi_i}{\pi} + 2\frac{N^{1/2}n_i^{1/2}}{qp^{\lambda}}\xi_i + 2\sigma_2\frac{N^{1/2}(M_1z)^{1/2}}{qp^{\kappa}}\xi_i\right) \mathrm{d}\xi_i.$$

For $I(M_1z, n_i, q)$, it is negligibly small by repeated integration by parts unless $t \simeq \frac{N^{1/2} n_i^{1/2}}{qp^{\lambda}} + \sigma_2 \frac{N^{1/2} (M_1z)^{1/2}}{qp^{\kappa}}$, in which case we apply the stationary phase method to the ξ_i -integral. Let $h_4(\xi_i) = -\frac{t\log\xi_i}{\pi} + 2\frac{N^{1/2} n_i^{1/2}}{qp^{\lambda}} \xi_i + 2\sigma_2 \frac{N^{1/2} (M_1z)^{1/2}}{qp^{\kappa}} \xi_i$. Then we have

$$h_4'(\xi_i) = -\frac{t}{\pi\xi_i} + 2\frac{N^{1/2}n_i^{1/2}}{qp^{\lambda}} + 2\sigma_2\frac{N^{1/2}(M_1z)^{1/2}}{qp^{\kappa}}, \quad h_4''(\xi_i) = \frac{t}{\pi\xi_i^2} \asymp t.$$

Then we have the stationary point

$$\xi_{i0} = \frac{t}{2\pi \left(\frac{N^{1/2} n_i^{1/2}}{q p^{\lambda}} + \sigma_2 \frac{N^{1/2} (M_1 z)^{1/2}}{q p^{\kappa}}\right)} \approx 1,$$

and

$$h_4(\xi_{i0}) = \frac{t}{\pi} \log \frac{2e\pi \left(\frac{N^{1/2} n_i^{1/2}}{qp^{\lambda}} + \sigma_2 \frac{N^{1/2} (M_1 z)^{1/2}}{qp^{\kappa}}\right)}{t}.$$

Hence, we have

$$I(M_1z, n_i, q) = t^{-\frac{1}{2}} V_{1i} \left(\frac{\frac{N^{1/2} n_i^{1/2}}{qp^{\lambda}} + \sigma_2 \frac{N^{1/2} (M_1z)^{1/2}}{qp^{\kappa}}}{t/2\pi} \right) e \left(\frac{t}{\pi} \log \frac{2e\pi \left(\frac{N^{1/2} n_i^{1/2}}{qp^{\lambda}} + \sigma_2 \frac{N^{1/2} (M_1z)^{1/2}}{qp^{\kappa}} \right)}{t} \right)$$
(6.13)

where V_{1i} are 1-inert functions supported on $\xi_{i0} \simeq 1$. We insert (6.13) into (6.5), make a change of variable $z \to z^2$ and write $a(n, z) := \frac{N^{1/2}n^{1/2}}{qp^{\lambda}} + \sigma_2 \frac{N^{1/2}M_1^{1/2}z}{qp^{\kappa}}$, then we obtain

$$\Im(m, n_1, n_2) = \frac{2}{t} \int_{\mathbb{R}} z W_1(z^2) V_{11}\left(\frac{a(n_1, z)}{t/2\pi}\right) \overline{V_{12}\left(\frac{a(n_2, z)}{t/2\pi}\right)} e\left(\rho_3(z)\right) \mathrm{d}z, \tag{6.14}$$

where

$$\rho_3(z) = \frac{t}{\pi} \log a(n_1, z) - \frac{t}{\pi} \log a(n_2, z) - \frac{mM_1}{p^{\kappa} d} z^2$$

Note that

$$\frac{\partial}{\partial z} \left(\frac{t}{\pi} \log a(n_1, z) - \frac{t}{\pi} \log a(n_2, z) \right) = \frac{t}{\pi} \frac{\sigma_2 N^{1/2} M_1^{1/2}}{q p^{\kappa}} \left(\frac{1}{a(n_1, z)} - \frac{1}{a(n_2, z)} \right)$$

For $\frac{p^{\kappa}d}{M_1}N^{\varepsilon} \ll m \ll \frac{N^{1/2}d}{M_1^{1/2}q}N^{\varepsilon}$, by repeated integration by parts, $\Im(m, n_1, n_2)$ is negligible small unless

$$\frac{tN^{1/2}M_1^{1/2}}{qp^{\kappa}} \left(\frac{1}{a(n_1,z)} - \frac{1}{a(n_2,z)}\right) \asymp \frac{mM_1}{p^{\kappa}d},\tag{6.15}$$

in which case we consider the third derivative of $\rho_3(z)$

$$\frac{\partial^3}{\partial z^3} \rho_3(z) = \frac{2t}{\pi} \left(\frac{\sigma_2 N^{1/2} M_1^{1/2}}{q p^{\kappa}} \right)^3 \left(\frac{1}{a(n_1, z)^3} - \frac{1}{a(n_2, z)^3} \right).$$
(6.16)

Note that $\frac{1}{a^2} + \frac{1}{ab} + \frac{1}{b^2} \gg \min(\frac{1}{a^2}, \frac{1}{b^2})$ for any $a, b \neq 0$. Then we have

$$\frac{1}{a(n_1,z)^3} - \frac{1}{a(n_2,z)^3} = \left(\frac{1}{a(n_1,z)} - \frac{1}{a(n_2,z)}\right) \left(\frac{1}{a(n_1,z)^2} + \frac{1}{a(n_1,z)a(n_2,z)} + \frac{1}{a(n_2,z)^2}\right) \\ \gg \left(\frac{1}{a(n_1,z)} - \frac{1}{a(n_2,z)}\right) \frac{1}{t^2}.$$
(6.17)

Therefore, by (6.15), (6.16) and (6.17) we obtain

$$\frac{\partial^3}{\partial z^3} \rho_3(z) \gg t \left(\frac{N^{1/2} M_1^{1/2}}{q p^{\kappa}}\right)^3 \left(\frac{1}{a(n_1, z)} - \frac{1}{a(n_2, z)}\right) \frac{1}{t^2} \asymp \frac{m M_1}{p^{\kappa} d} \left(\frac{N^{1/2} M_1^{1/2}}{q p^{\kappa} t}\right)^2 = \frac{m N M_1^2}{q^2 p^{3\kappa} t^2 d}$$

By Lemma 3.9 with r = 3, we have

$$z$$
-integral $\ll \left(\frac{mNM_1^2}{q^2p^{3\kappa}t^2d}\right)^{-1/3}$

Hence,

$$\Im(m, n_1, n_2) \ll \frac{1}{t} \left(\frac{m N M_1^2}{q^2 p^{3\kappa} t^2 d} \right)^{-1/3}$$

(3). For $m \ll \frac{p^{\kappa}d}{M_1}N^{\varepsilon}$, we estimate (6.14) trivially and obtain $\Im(m, n_1, n_2) \ll \frac{1}{t}$.

7. The zero frequency

In this section, we deal with the terms with m = 0. The character sum with m = 0 is

$$\mathfrak{C}(0,n_1,n_2) = p^{2\kappa} \sum_{d \mid (n_1-n_2,p^{\lambda})} d\mu \left(p^{\lambda}/d \right).$$

The contributions of the zero frequency to $\Sigma(d,q)$ and $S^{\flat}(N)$ are denoted by $\Sigma_0(d,q)$ and $S_0^{\flat}(N)$, respectively.

7.1. Large modulo case: $N^{\varepsilon} \ll \frac{NX}{p^{\lambda}RQ} \ll t^{1-\varepsilon}$. In this case, we have $|n_1 - n_2| \ll \frac{p^{\lambda}RX}{Q}N^{\varepsilon}$, $\Im(0, n_1, n_2) \ll \frac{1}{t}$. The contributions of this part to $\Sigma(d, q)$ and $S^{\flat}(N)$ are denoted by $\Sigma_{01}(d, q)$ and $S_{01}^{\flat}(N)$, respectively. By (6.2), (6.3) and $M_1 \simeq \frac{t^2R^2p^{2\kappa}}{N}$, we get

$$\Sigma_{01}(d,q) \ll \frac{Q^4}{N^2 X^4} \frac{R^2 p^{\kappa} t^2}{Nd} \sum_{n_1 \asymp \frac{NX^2}{Q^2}} |\lambda_g(n_1)|^2 \left(\sum_{\substack{n_2 \asymp \frac{NX^2}{Q^2} \\ n_2 \equiv n_1 \bmod dp^{\lambda} \\ |n_1 - n_2| \ll \frac{p^{\lambda} RX}{Q} N^{\varepsilon}}} \frac{p^{2\kappa + \lambda}}{n_2 \equiv n_1 \bmod dp^{\lambda - 1}} \right)$$
$$\ll \frac{R^2 p^{\kappa} t^2}{Nd} \frac{Q^2}{NX^2} N^{\varepsilon} \left(1 + \frac{p^{\lambda} RX}{Qdp^{\lambda}} \right) \frac{p^{2\kappa + \lambda}}{t}}{t}$$
$$\ll \frac{R^2 Q^2 t N^{\varepsilon}}{N^2 X^2 d} p^{3\kappa + \lambda} + \frac{R^3 Q t N^{\varepsilon}}{N^2 X d^2} p^{3\kappa + \lambda}.$$
(7.1)

By (6.1), (7.1) and $N \ll (p^{\kappa}t)^{2+\varepsilon}$, we obtain

$$\begin{split} S^{\flat}_{01}(N) \ll & \frac{NX^{3/2}R^{1/2}t^{1/2}}{Q^{3/2}p^{(\kappa+\lambda)/2}} \sum_{\substack{q \sim R \\ (q,p)=1}} \frac{1}{q} \sum_{d|q} d\left(\frac{R^2Q^2tN^{\varepsilon}}{N^2X^2d} p^{3\kappa+\lambda} + \frac{R^3QtN^{\varepsilon}}{N^2Xd^2} p^{3\kappa+\lambda}\right)^{1/2} \\ \ll & N^{1/2+\varepsilon} \left(p^{3\kappa/2-3\lambda/4}\frac{t^{3/2}}{K^{3/4}} + p^{\kappa-\lambda/2}\frac{t}{K^{1/2}}\right). \end{split}$$

7.2. Small modulo case: $\frac{NX}{p^{\lambda}RQ} \gg t^{1-\varepsilon}$. We have $\Im(0, n_1, n_2) \ll \frac{1}{t}$, $M_1 \ll \frac{NX^2p^{2\kappa-2\lambda}}{Q^2}$ and $R \ll \frac{NX}{p^{\lambda}Qt^{1-\varepsilon}}$. The contributions of this part to $\Sigma(d, q)$ and $S^{\flat}(N)$ are denoted by $\Sigma_{02}(d, q)$ and $S^{\flat}_{02}(N)$, respectively. By (6.2) and (6.3), we get

$$\Sigma_{02}(d,q) \ll \frac{Q^4}{N^2 X^4} \frac{M_1}{p^{\kappa} d} \sum_{n_1 \asymp \frac{NX^2}{Q^2}} |\lambda_g(n_1)|^2 \left(\sum_{\substack{n_2 \asymp \frac{NX^2}{Q^2} \\ n_2 \equiv n_1 \bmod dp^{\lambda}}} \frac{p^{2\kappa+\lambda}}{t} + \sum_{\substack{n_2 \asymp \frac{NX^2}{Q^2} \\ n_2 \equiv n_1 \bmod dp^{\lambda}}} \frac{p^{2\kappa+\lambda-1}}{t} \right)$$
$$\ll \frac{M_1}{p^{\kappa} d} \frac{Q^2}{NX^2} \left(1 + \frac{NX^2}{Q^2 dp^{\lambda}} \right) \frac{p^{2\kappa+\lambda}}{t}$$
$$\ll \frac{Q^2 M_1 N^{\varepsilon}}{NX^2 dt} p^{\kappa+\lambda} + \frac{M_1 N^{\varepsilon}}{d^2 t} p^{\kappa}.$$
(7.2)

By (6.1), (7.2) and $N \ll (p^{\kappa}t)^{2+\varepsilon}$, we obtain

$$\begin{split} S_{02}^{\flat}(N) &= \sup_{M_1 \ll \frac{NX^2 p^{2\kappa - 2\lambda}}{Q^2}} \frac{N^{5/4} X^{3/2} M_1^{1/4}}{Q^{3/2} p^{(2\kappa + \lambda)/2}} \sum_{\substack{q \sim R \\ (q,p) = 1}} \frac{1}{q} \sum_{d|q} d \cdot \Sigma_0^{1/2} \\ &\ll \sup_{M_1 \ll \frac{NX^2 p^{2\kappa - 2\lambda}}{Q^2}} \frac{N^{5/4} X^{3/2} M_1^{1/4} N^{\varepsilon}}{Q^{3/2} p^{(2\kappa + \lambda)/2}} \sum_{\substack{q \sim R \\ (q,p) = 1}} \frac{1}{q} \sum_{d|q} d \left(\frac{Q^2 M_1 N^{\varepsilon}}{NX^2 dt} p^{\kappa + \lambda} + \frac{M_1 N^{\varepsilon}}{d^2 t} p^{\kappa} \right)^{1/2} \\ &\ll N^{1/2 + \varepsilon} \left(p^{3\kappa/2 - 3\lambda/4} \frac{K^{5/4}}{t^{1/2}} + p^{\kappa - \lambda/2} \frac{K^{3/2}}{t^{1/2}} \right). \end{split}$$

8. The non-zero frequencies

In this section, we deal with the terms with $m \neq 0$.

8.1. Large modulo case:
$$N^{\varepsilon} \ll \frac{NX}{p^{\lambda}RQ} \ll t^{1-\varepsilon}$$
.

8.1.1. $m \ll \frac{dN}{R^2 p^{\kappa} t^2} N^{\varepsilon}$. For $m \ll \frac{dN}{R^2 p^{\kappa} t^2} N^{\varepsilon}$ and $m \neq 0$, we have $|n_1 - n_2| \ll \frac{p^{\lambda} RX}{Q} N^{\varepsilon}$ and $\Im(m, n_1, n_2) \ll t^{-1}$. The contributions of this part to $\Sigma(d, q)$ and $S^{\flat}(N)$ are denoted by

 $\Sigma_{11}(d,q)$ and $S_{11}^{\flat}(N)$, respectively. By (6.2), (6.3) and Lemma 6.1(1)(2), we get

$$\begin{split} \Sigma_{11}(d,q) \ll \frac{Q^4}{N^2 X^4} \frac{R^2 p^{\kappa} t^2}{Nd} \sum_{n_1 \asymp \frac{NX^2}{Q^2}} |\lambda_g(n_1)|^2 \cdot \left(\sum_{\substack{n_2 \asymp \frac{NX^2}{Q^2} \\ n_2 \equiv n_1 \bmod dp^{\beta-\kappa+\lambda} p^{\kappa-\lambda+\beta-\kappa+\lambda} \mid m}} \sum_{\substack{m \ll \frac{dN}{R^2 p^{\kappa} t^2} N^{\varepsilon} \\ n_2 \equiv n_1 \bmod dp^{\beta-\kappa+\lambda} p^{\kappa-\lambda+\beta-\kappa+\lambda} \mid m}} \right) \\ &+ \sum_{0 \le \ell \le \lambda-\kappa/2} \sum_{\substack{n_2 \asymp \frac{NX^2}{Q^2} \\ n_2 \equiv n_1 \bmod dp^{\ell}, p^{\ell} \mid n_1 - n_2}} \sum_{\substack{m \ll \frac{dN}{R^2 p^{\kappa} t^2} N^{\varepsilon} \\ p^{\kappa-\lambda+\ell} \mid m}} \frac{p^{5\kappa/2+2\ell+\delta_1+\delta_2/2}}{t} \right) \\ \ll \frac{R^2 Q^2 t^2 p^{\kappa}}{N^2 X^2 d} N^{\varepsilon} \left(\left(1 + \frac{p^{\lambda} RX}{Q dp^{\beta-\kappa+\lambda}}\right) \frac{N d \cdot p^{2\kappa+\lambda+\delta_1}}{R^2 p^{\kappa} t^2 p^{\beta} \cdot t} + \frac{p^{\lambda} RX}{Q dp^{\ell}} \frac{N d \cdot p^{5\kappa/2+2\ell+\delta_1+\delta_2/2}}{R^2 p^{\kappa-\lambda+\ell} \cdot t} \right) \\ \ll \frac{Q^2 N^{\varepsilon}}{NX^2 t} p^{3\kappa/2+\lambda+\delta_1+\delta_2/2} + \frac{QRN^{\varepsilon}}{NX dt} p^{2\kappa+\lambda+\delta_1+\delta_2} + \frac{QRN^{\varepsilon}}{NX dt} p^{3\kappa/2+2\lambda+\delta_1+\delta_2/2}. \end{split}$$

$$\tag{8.1}$$

By (6.1), (8.1) and $N \ll (p^{\kappa}t)^{2+\varepsilon}$, we obtain

$$S_{11}^{\flat}(N) = \frac{NX^{3/2}R^{1/2}t^{1/2}}{Q^{3/2}p^{(\kappa+\lambda)/2}} \sum_{\substack{q \sim R \\ (q,p)=1}} \frac{1}{q} \sum_{d|q} d \cdot \Sigma_{11}(d,q)^{1/2} \\ \ll N^{1/2+\varepsilon} \left(p^{5\kappa/4-\lambda/2+3/4} \frac{t}{K^{1/2}} + p^{\kappa-\lambda/4+1} \frac{t^{1/2}}{K^{1/2}} + p^{3\kappa/4+\lambda/4+3/4} \frac{t^{1/2}}{K^{1/4}} \right).$$

8.1.2. $\frac{dN}{R^2 p^{\kappa} t^2} N^{\varepsilon} \ll m \ll \frac{N^2 X d}{R^3 Q t^2 p^{\kappa+\lambda}} N^{\varepsilon}$. For $m \sim M$ with $\frac{dN}{R^2 p^{\kappa} t^2} N^{\varepsilon} \ll M \ll \frac{N^2 X d}{R^3 Q t^2 p^{\kappa+\lambda}} N^{\varepsilon}$, we have $|n_1 - n_2| \asymp \frac{MR^3 X t^2 p^{\kappa+\lambda}}{NQd}$ and $\mathfrak{I}(m, n_1, n_2) \ll \left(\frac{dN}{MR^2 p^{\kappa} t^4}\right)^{1/2}$. The contributions of this part to $\Sigma(d, q)$ and $S^{\flat}(N)$ are denoted by $\Sigma_{12}(d, q)$ and $S^{\flat}_{12}(N)$, respectively.

$$\begin{split} \Sigma_{12}(d,q) \ll & \frac{Q^4}{N^2 X^4} \frac{R^2 p^{\kappa} t^2}{Nd} \sum_{n_1 \asymp \frac{NX^2}{Q^2}} |\lambda_g(n_1)|^2 \bigg(\sum_{\substack{n_2 \asymp NX^2 \\ n_2 \equiv n_1 \bmod dp^{\beta-\kappa+\lambda} \\ |n_1-n_2| \ll \frac{MR^3 X t^2 p^{\kappa+\lambda}}{NQd}}} \sum_{\substack{p^{\kappa-\lambda+\beta-\kappa+\lambda}|m}} \sum_{\substack{p^{\kappa-\lambda+\beta} | n^{\kappa/2} \\ M^{1/2} R p^{\kappa/2} t^2}} + \sum_{\substack{0 \le \ell \le \lambda - \kappa/2 \\ n_2 \equiv n_1 \bmod dp^{\ell}, \ p^{\ell} | n_1-n_2|}} \sum_{\substack{m \approx M \\ p^{\kappa-\lambda+\ell}|m}} \sum_{\substack{p^{\kappa-\lambda+\ell} | m}} \frac{p^{5\kappa/2+2\ell+\delta_1+\delta_2/2} d^{1/2} N^{1/2}}{M^{1/2} R p^{\kappa/2} t^2}} \bigg) \\ \ll \frac{RQ^2 p^{\kappa/2} N^{\varepsilon}}{M^{1/2} N^{3/2} X^2 d^{1/2}} \bigg(\bigg(1 + \frac{MR^3 X t^2 p^{\kappa+\lambda}}{NQ d^2 p^{\beta-\kappa+\lambda}} \bigg) \frac{Mp^{2\kappa+\lambda+\delta_1}}{p^{\beta}} + \sup_{\ell} \frac{MR^3 X t^2 p^{\kappa+\lambda}}{NQ d^2 p^{\ell}} \frac{Mp^{5\kappa/2+2\ell+3/2}}{p^{\kappa-\lambda+\ell}} \bigg) \\ \ll \frac{Q^{3/2} p^{3\kappa/2+\lambda/2+3/2} N^{\varepsilon}}{R^{1/2} X^{3/2} N^{1/2} t} + \frac{N^{1/2} X^{1/2} p^{2\kappa-\lambda/2+1} N^{\varepsilon}}{R^{1/2} Q^{1/2} dt} + \frac{N^{1/2} X^{1/2} p^{3\kappa/2+\lambda/2+3/2} N^{\varepsilon}}{R^{1/2} Q^{1/2} dt}. \end{split}$$

$$(8.2)$$

By (6.1), (8.2) and $N \ll (p^{\kappa}t)^{2+\varepsilon}$, we obtain

$$S_{12}^{\flat}(N) = \frac{NX^{3/2}R^{1/2}t^{1/2}}{Q^{3/2}p^{(\kappa+\lambda)/2}} \sum_{\substack{q \sim R \\ (q,p)=1}} \frac{1}{q} \sum_{d|q} d \cdot \Sigma_{12}(d,q)^{1/2} \\ \ll N^{1/2+\varepsilon} \left(p^{5\kappa/4-\lambda/2+3/4} \frac{t}{K^{1/4}} + p^{\kappa-\lambda/4+1}t^{1/2}K^{1/2} + p^{3\kappa/4+\lambda/4+3/4}t^{1/2}K^{1/2} \right).$$

8.2. Small modulo case: $\frac{NX}{p^{\lambda}RQ} \gg t^{1-\varepsilon}$.

8.2.1. $m \ll \frac{p^{\kappa_d}}{M_1} N^{\varepsilon}$. For $m \ll \frac{p^{\kappa_d}}{M_1} N^{\varepsilon}$ and $m \neq 0$, we have $\Im(m, n_1, n_2) \ll \frac{1}{t}$. The contributions of this part to $\Sigma(d, q)$ and $S^{\flat}(N)$ are denoted by $\Sigma_{13}(d, q)$ and $S^{\flat}_{13}(N)$, respectively. By (6.2), (6.3) and Lemma 6.1(1)(2), we get

$$\begin{split} \Sigma_{13}(d,q) \ll \frac{Q^4}{N^2 X^4} \frac{M_1}{p^{\kappa} d} \sum_{n_1 \asymp \frac{NX^2}{Q^2}} |\lambda_g(n_1)|^2 \bigg(\sum_{\substack{n_2 \asymp \frac{NX^2}{Q^2} \\ n_2 \equiv n_1 \bmod dp^{\beta-\kappa+\lambda} p^{\kappa-\lambda+\beta-\kappa+\lambda} | m}} \sum_{\substack{m \ll \frac{p^{\kappa} d}{M_1} N^{\varepsilon} \\ n_2 \equiv n_1 \bmod dp^{\beta-\kappa+\lambda} p^{\kappa-\lambda+\beta-\kappa+\lambda} | m}} + \sum_{\substack{0 \le \ell \le \lambda - \kappa/2 \\ n_2 \equiv n_1 \bmod dp^{\ell} p^{\kappa-\lambda+\ell} | m}} \sum_{\substack{m \ll \frac{p^{\kappa} d}{M_1} N^{\varepsilon} \\ n_2 \equiv n_1 \bmod dp^{\ell} p^{\kappa-\lambda+\ell} | m}} p^{5\kappa/2+2\ell+\delta_1+\delta_2/2} \frac{1}{t}} \bigg) \\ \ll \frac{Q^2}{NX^2} \frac{M_1}{p^{\kappa} d} \bigg(\bigg(1 + \frac{NX^2}{Q^2 dp^{\beta-\kappa+\lambda}} \bigg) \frac{\frac{p^{\kappa} d}{M_1} N^{\varepsilon}}{p^{\beta}} \frac{p^{2\kappa+\lambda+\delta_1}}{t} + \sup_{\ell} \frac{NX^2}{Q^2 dp^{\ell}} \frac{\frac{p^{\kappa} d}{M_1} N^{\varepsilon}}{p^{\kappa-\lambda+\ell}} \frac{p^{5\kappa/2+2\ell+3/2}}{t} \bigg) \\ \ll \frac{Q^2 p^{3\kappa/2+\lambda+3/2} N^{\varepsilon}}{NX^2 t} + \frac{p^{2\kappa+2} N^{\varepsilon}}{dt} + \frac{p^{3\kappa/2+\lambda+3/2} N^{\varepsilon}}{dt}. \end{split}$$

$$(8.3)$$

By (6.1), (8.3) and $N \ll (p^{\kappa}t)^{2+\varepsilon}$, we obtain

$$S_{13}^{\flat}(N) = \sup_{M_1 \ll \frac{NX^2 p^{2\kappa - 2\lambda}}{Q^2}} \frac{N^{5/4} X^{3/2} M_1^{1/4}}{Q^{3/2} p^{(2\kappa + \lambda)/2}} \sum_{\substack{q \sim R \\ (q,p) = 1}} \frac{1}{q} \sum_{d|q} d \cdot \Sigma_{13}^{1/2}$$
$$\ll N^{1/2 + \varepsilon} \left(p^{5\kappa/4 - \lambda/2 + 3/4} \frac{K}{t^{1/2}} + p^{\kappa - \lambda/4 + 1} \frac{K^{5/4}}{t^{1/2}} + p^{3\kappa/4 + \lambda/4 + 3/4} \frac{K^{5/4}}{t^{1/2}} \right).$$

8.2.2. $\frac{p^{\kappa}d}{M_1}N^{\varepsilon} \ll m \ll \frac{N^{1/2}d}{M_1^{1/2}q}N^{\varepsilon}$. For $\frac{p^{\kappa}d}{M_1}N^{\varepsilon} \ll m \ll \frac{N^{1/2}d}{M_1^{1/2}q}N^{\varepsilon}$, we have $\Im(m, n_1, n_2) \ll \frac{1}{t}\left(\frac{mNM_1^2}{q^2p^{3\kappa}t^2d}\right)^{-1/3}$. The contributions of this part to $\Sigma(d,q)$ and $S^{\flat}(N)$ are denoted by $\Sigma_{14}(d,q)$

and $S_{14}^{\flat}(N)$, respectively. By (6.2), (6.3) and Lemma 6.1(1)(2), we get

$$\begin{split} \Sigma_{14}(d,q) \ll & \frac{Q^4}{N^2 X^4} \frac{M_1}{p^{\kappa} d} \sum_{n_1 \asymp \frac{NX^2}{Q^2}} |\lambda_g(n_1)|^2 \Big(\sum_{\substack{n_2 \asymp \frac{NX^2}{Q^2} \\ n_2 \equiv n_1 \bmod dp^{\beta-\kappa+\lambda}}} \sum_{\substack{m \ll \frac{N^{1/2} d}{M_1^{1/2} q} N^{\varepsilon} \\ p^{\kappa-\lambda+\beta-\kappa+\lambda} \mid m}} p^{2\kappa+\lambda+\delta_1} \frac{1}{t} \left(\frac{mNM_1^2}{q^2 p^{3\kappa} t^2 d} \right)^{-1/3}}{t} \\ &+ \sum_{0 \le \ell \le \lambda-\kappa/2} \sum_{\substack{n_2 \asymp \frac{NX^2}{Q^2} \\ n_2 \equiv n_1 \bmod dp^{\ell} \\ p^{\ell} \mid n_1 - n_2}} \sum_{\substack{m \ll \frac{N^{1/2} d}{M_1^{1/2} q} N^{\varepsilon} \\ p^{\kappa-\lambda+\ell} \mid m}} p^{5\kappa/2+2\ell+\delta_1+\delta_2/2} \frac{1}{t} \left(\frac{mNM_1^2}{q^2 p^{3\kappa} t^2 d} \right)^{-1/3}}{t} \Big) \\ \ll \frac{Q^2 M_1^{1/3} q^{2/3}}{N^{4/3} X^2 d^{2/3} t^{1/3}} \left(\left(1 + \frac{NX^2}{Q^2 d p^{\beta-\kappa+\lambda}} \right) \frac{\left(\frac{N^{1/2} d}{M_1^{1/2} q} N^{\varepsilon} \right)^{2/3} p^{2\kappa+\lambda+\delta_1}}{p^{\beta}} \\ &+ \sup_{\ell} \frac{NX^2}{Q^2 d p^{\ell}} \frac{\left(\frac{N^{1/2} d}{M_1^{1/2} q} N^{\varepsilon} \right)^{2/3} p^{5\kappa/2+2\ell+3/2}}{p^{\kappa-\lambda+\ell}} \right) \\ \ll \frac{Q^2 p^{3\kappa/2+\lambda+3/2} N^{\varepsilon}}{NX^2 t^{1/3}} + \frac{p^{2\kappa+2} N^{\varepsilon}}{dt^{1/3}} + \frac{p^{3\kappa/2+\lambda+3/2} N^{\varepsilon}}}{dt^{1/3}}. \end{split}$$

$$(8.4)$$

By (6.1), (8.4) and $N \ll (p^{\kappa}t)^{2+\varepsilon}$, we obtain

$$S_{14}^{\flat}(N) = \sup_{M_1 \ll \frac{NX^2 p^{2\kappa - 2\lambda}}{Q^2}} \frac{N^{5/4} X^{3/2} M_1^{1/4}}{Q^{3/2} p^{(2\kappa + \lambda)/2}} \sum_{\substack{q \sim R \\ (q,p) = 1}} \frac{1}{q} \sum_{d|q} d \cdot \Sigma_{14}^{1/2}$$
$$\ll N^{1/2 + \varepsilon} \left(p^{5\kappa/4 - \lambda/2 + 3/4} \frac{K}{t^{1/6}} + p^{\kappa - \lambda/4 + 1} \frac{K^{5/4}}{t^{1/6}} + p^{3\kappa/4 + \lambda/4 + 3/4} \frac{K^{5/4}}{t^{1/6}} \right).$$

9. CONCLUSION

By (4.2) and the upper bounds of $S_{01}^{\flat}(N)$, $S_{02}^{\flat}(N)$, $S_{11}^{\flat}(N)$, $S_{12}^{\flat}(N)$, $S_{13}^{\flat}(N)$, $S_{14}^{\flat}(N)$, $S_{b0}^{\flat}(N)$, $S_{b1}^{\flat}(N)$, $S_{c0}^{\flat}(N)$, $S_{c1}^{\flat}(N)$, $S_{c1}^{\flat}(N)$, the best choice of parameters is $K = t^{\frac{4}{5}}$ and $\lambda = [\frac{3}{4}\kappa]$, where [x] denotes the largest integer which does not exceed x. Therefore, we have shown the following bound

$$S(N) \ll N^{1/2} p^{3/4} (p^{\kappa})^{1 - \frac{1}{16} + \varepsilon} t^{1 - \frac{1}{10} + \varepsilon}.$$
(9.1)

Substituting the estimate in (9.1) for S(N) into (4.1), we obtain

$$L\left(\frac{1}{2}+it, f \times g \times \chi\right) \ll_{\varepsilon} p^{\frac{3}{4}} (p^{\kappa})^{1-\frac{1}{16}+\varepsilon} t^{1-\frac{1}{10}+\varepsilon}.$$

This completes the proof of Theorem 1.1.

10. The complementary cases

In this section we focus on the complementary cases whose estimations can be bounded by that of the main case. And we give the brief proofs for the complementary cases. Next, we consider the following three cases:

case
$$a \quad \frac{nN}{q^2 p^{2\lambda}} \ll N^{\varepsilon}, \quad \frac{NX}{p^{\lambda} RQ} \ll N^{\varepsilon} \text{ and } \frac{mN}{q^2 p^{2\kappa}} \ll N^{\varepsilon},$$

case $b \quad \frac{nN}{q^2 p^{2\lambda}} \asymp \left(\frac{NX}{RQp^{\lambda}}\right)^2 \gg N^{\varepsilon} \text{ and } \frac{mN}{q^2 p^{2\kappa}} \ll N^{\varepsilon},$
case $c \quad \frac{nN}{q^2 p^{2\lambda}} \ll N^{\varepsilon}, \quad \frac{NX}{p^{\lambda} RQ} \ll N^{\varepsilon} \text{ and } \frac{mN}{q^2 p^{2\kappa}} \gg N^{\varepsilon}.$

It is easy to see that the contribution from case a to S(N) is negligibly small by repeated integration by parts for (5.8).

10.1. Case b: $\frac{nN}{q^2p^{2\lambda}} \approx \left(\frac{NX}{RQp^{\lambda}}\right)^2 \gg N^{\varepsilon}$, $\frac{mN}{q^2p^{2\kappa}} \ll N^{\varepsilon}$. We first use the same method as in §5 to *n*-sum and *u*-integral. Then we use the GL₂ Voronoi summation formula to *m*-sum and obtain that

$$S_{b}^{\flat}(N) = \frac{X^{\frac{3}{2}}N^{\frac{1}{2}}}{Q^{\frac{3}{2}}} \sum_{\substack{q \sim R \\ (q,p)=1}} \frac{1}{(qp^{\lambda})^{1/2}} \sum_{a \bmod qp^{\lambda}} \sum_{n \asymp \frac{NX^{2}}{Q^{2}}} \frac{\lambda_{g}(n)}{n} e\left(\frac{\overline{a}n}{qp^{\lambda}}\right)$$

$$\cdot \frac{1}{\tau(\overline{\chi})} \frac{N^{1-it}}{qp^{\kappa}} \sum_{c \bmod p^{\kappa}} \overline{\chi}(c) \sum_{m} \lambda_{f}(m) e\left(-\frac{\overline{a^{*}}m}{qp^{\kappa}}\right) \Phi_{2}^{\pm}\left(\frac{mN}{q^{2}p^{2\kappa}}\right) + O(N^{-A}).$$
(10.1)

By Lemma 5.2, $\Phi_2^{\pm}(y)$ is negligible small unless $t \simeq \frac{N^{1/2}n^{1/2}}{qp^{\lambda}} \simeq \frac{NX}{p^{\lambda}RQ}$, in which case

$$\Phi_2^{\pm}(y) = \int_0^\infty V(\xi) V_2\left(\frac{nQ^2}{NX^2\xi}\right) e\left(-\frac{t\log\xi}{2\pi} + 2\frac{N^{1/2}n^{1/2}}{qp^{\lambda}}\xi^{1/2}\right) J_f^{\pm}(y\xi) \mathrm{d}\xi, \tag{10.2}$$

with $y^k J_f^{\pm(k)}(y) \ll_{k,f} N^{\varepsilon}$.

Inserting (10.2) into (10.1) and rearranging the order of summations and integrals, we arrive at

$$S_b^{\flat}(N) = \frac{N^{3/2 - it} X^{3/2}}{\tau(\overline{\chi}) Q^{3/2} p^{(2\kappa+\lambda)/2}} \sum_{\substack{q \sim R \\ (q,p)=1}} \frac{1}{q^{\frac{3}{2}}} \sum_{d|q} d\mu\left(\frac{q}{d}\right) \sum_{\substack{n \asymp \frac{NX^2}{Q^2}}} \frac{\lambda_g(n)}{n}$$
$$\cdot \sum_{\substack{m \ll \frac{q^2 p^{2\kappa} N^{\varepsilon}}{N} \\ n \equiv mp^{\lambda} \overline{p^{2\kappa-\lambda}} \bmod d}} \lambda_f(m) \mathfrak{C}^*(m,n,q) \Phi_2^{\pm}\left(\frac{mN}{q^2 p^{2\kappa}}\right)$$

with

$$\mathfrak{C}^*(m,n,q) = \sum_{c \bmod p^{\kappa}} \overline{\chi}(c) \sum_{b \bmod p^{\lambda}} e\left(\frac{-(\overline{ap^{\kappa-\lambda} + cq})m\bar{q}}{p^{\kappa}} + \frac{n\bar{q}\bar{b}}{p^{\lambda}}\right).$$

Applying the Cauchy–Schwarz inequality and the Poisson summation formula to m-sum, we have

$$S_b^{\flat}(N) \ll \frac{NX^{3/2}}{Q^{3/2} p^{(\kappa+\lambda)/2}} \sum_{\substack{q \sim R \\ (q,p)=1}} \frac{1}{q^{\frac{1}{2}}} \sum_{d|q} d \cdot \Sigma_b(d,q)^{1/2}, \tag{10.3}$$

where

$$\Sigma_{b}(d,q) \ll \frac{Q^{4}R^{2}p^{\kappa}}{N^{3}X^{4}d} \sum_{\substack{n_{1} \asymp \frac{NX^{2}}{Q^{2}}\\ n_{2} \equiv n_{1} \bmod d}} |\lambda_{g}(n_{1})|^{2} \sum_{\substack{n_{2} \asymp \frac{NX^{2}}{Q^{2}}\\ n_{2} \equiv n_{1} \bmod d}} \sum_{m \in \mathbb{Z}} |\mathfrak{C}(m,n_{1},n_{2}) \cdot \mathfrak{I}_{b}(m,n_{1},n_{2})|,$$

with

$$\mathfrak{C}(m,n_1,n_2) = \sum_{\gamma \bmod p^{\kappa}} \mathfrak{C}^*(\gamma,n_1,q) \overline{\mathfrak{C}^*(\gamma,n_2,q)} e\left(\frac{m\left(n_1 p^{3\kappa-\lambda} \overline{p^{\kappa+\lambda}} + \gamma d\overline{d}\right)}{dp^{\kappa}}\right)$$

and

$$\mathfrak{I}_{b}(m,n_{1},n_{2}) := \int_{\mathbb{R}} V\left(\xi\right) \Phi_{2}^{\pm}\left(\xi\right) \overline{\Phi_{2}^{\pm}\left(\xi\right)} e\left(-\frac{mR^{2}p^{\kappa}}{Nd}\xi\right) \mathrm{d}\xi$$

Lemma 10.1. (1) If $m \gg \frac{N^{1+\varepsilon}d}{R^2p^{\kappa}}$, we have $\Im(m, n_1, n_2) \ll N^{-A}$. (2) If $m \ll \frac{N^{1+\varepsilon}d}{R^2p^{\kappa}}$, we have $\Im(m, n_1, n_2) \ll 1$.

Proof. The first result can be done by applying integration by parts with respect to the ξ -integral. For $m \ll \frac{N^{1+\epsilon}d}{R^2p^{\kappa}}$, we have the trivial bound $\Im(m, n_1, n_2) \ll 1$.

The contribution from the zero frequency to $\Sigma_b(d,q)$ is

$$\Sigma_{b0}(d,q) \ll \frac{Q^4 R^2 p^{\kappa}}{N^3 X^4 d} \sum_{\substack{n_1 \asymp \frac{NX^2}{Q^2}}} |\lambda_g(n_1)|^2 \left(\sum_{\substack{n_2 \asymp \frac{NX^2}{Q^2} \\ n_2 \equiv n_1 \bmod dp^{\lambda}}} p^{2\kappa+\lambda} + \sum_{\substack{n_2 \asymp \frac{NX^2}{Q^2} \\ n_2 \equiv n_1 \bmod dp^{\lambda}}} p^{2\kappa+\lambda-1} \right).$$
(10.4)

Plugging (10.4) into (10.3), we get

$$S_{b0}^{\flat}(N) \ll N^{1/2+\varepsilon} \left(p^{3\kappa/2 - 3\lambda/4} \frac{K^{5/4}}{t^{3/2}} + p^{\kappa - \lambda/2} \frac{K^{3/2}}{t^{3/2}} \right).$$

The contribution from the non-zero frequencies to $\Sigma_b(d,q)$ is

$$\Sigma_{b1}(d,q) \ll \frac{Q^4 R^2 p^{\kappa}}{N^3 X^4 d} \sum_{n_1 \asymp \frac{NX^2}{Q^2}} |\lambda_g(n_1)|^2 \bigg(\sum_{\substack{n_2 \asymp \frac{NX^2}{Q^2} \\ n_2 \equiv n_1 \bmod dp^{\beta-\kappa+\lambda} p^{\kappa-\lambda+\beta-\kappa+\lambda} | m}} \sum_{\substack{m \ll \frac{dN^{1+\varepsilon}}{R^2 p^{\kappa}} \\ p^{\kappa-\lambda+\beta-\kappa+\lambda} | m}} p^{2\kappa+\lambda+\delta_1} + \sum_{\substack{0 \le \ell \le \lambda-\kappa/2 \\ n_2 \equiv n_1 \bmod dp^{\ell}}} \sum_{\substack{m \ll \frac{dN^{1+\varepsilon}}{R^2 p^{\kappa}} \\ p^{\kappa-\lambda+\ell} | m}} p^{5\kappa/2+2\ell+\delta_1+\delta_2/2}} \bigg),$$
(10.5)

Plugging (10.5) into (10.3), we get

$$S_{b1}^{\flat}(N) \ll N^{1/2+\varepsilon} \left(p^{5\kappa/4 - \lambda/2 + 3/4} \frac{K}{t^{1/2}} + p^{\kappa - \lambda/4 + 1} \frac{K^{5/4}}{t^{1/2}} + p^{3\kappa/4 + \lambda/4 + 3/4} \frac{K^{5/4}}{t^{1/2}} \right).$$

10.2. Case c: $\frac{nN}{q^2p^{2\lambda}} \ll N^{\varepsilon}$, $\frac{NX}{p^{\lambda}RQ} \ll N^{\varepsilon}$, $\frac{mN}{q^2p^{2\kappa}} \gg N^{\varepsilon}$. Applying Lemma 3.4 to *m*-sum in (4.4) and by (3.8), we are led to estimate

$$\frac{1}{\tau(\overline{\chi})} \frac{N^{1-it}}{qp^{\kappa}} \sum_{c \bmod p^{\kappa}} \overline{\chi}(c) \sum_{m} \lambda_f(m) e\left(-\frac{\overline{a^*}m}{qp^{\kappa}}\right) \Phi_2^+\left(\frac{mN}{q^2 p^{2\kappa}}\right)$$

with

$$\Phi_2^+(y) = y^{-\frac{1}{4}} \int_0^\infty V(\xi) \xi^{-\frac{1}{4}} e\left(-\frac{t\log\xi}{2\pi} + \frac{Nu}{Qqp^{\lambda}} \xi \pm 2\sqrt{y\xi}\right) \mathrm{d}\xi + O(N^{-A}).$$
(10.6)

By repeated integration by parts, $\Phi_2^+(y)$ is negligibly small unless $\frac{\sqrt{mN}}{Rp^{\kappa}} \approx t$, in which case by making a change of variable $\xi \to \xi^2$ and applying the stationary phase method we have

$$\Phi_2^+(y) = 2y^{-\frac{1}{4}}t^{-1/2}e\left(-\frac{t}{\pi}\log\frac{Rtp^{\kappa}}{2\pi e\sqrt{mN}}\right)W\left(\frac{mN}{R^2p^{2\kappa}t^2}\right) + O(N^{-A}).$$

Then we apply the GL_2 Voronoi summation formula to n-sum and making a change of variable $u \to Xu$, we have

$$S_{c}^{\flat}(N) \ll \frac{XN^{7/4+\varepsilon}}{p^{2\lambda+\kappa}Q\sqrt{t}} \sum_{\substack{q\sim R\\(q,p)=1}} \frac{1}{q^{5/2}} \sum_{a \bmod qp^{\lambda} c \bmod p^{\kappa}} \overline{\chi}(c) \sum_{\eta=\pm 1} \sum_{n} \lambda_{g}(n) e\left(\eta \frac{\overline{a}n}{qp^{\lambda}}\right) \sum_{m} \frac{\lambda_{f}(m)}{m^{1/4}} e\left(\frac{-\overline{a^{*}}m}{qp^{\kappa}}\right) + \int_{\mathbb{R}} W_{q}(u) g(q, Xu) \Phi_{1}^{\operatorname{sgn}\eta}\left(\frac{nN}{q^{2}p^{2\lambda}}\right) e\left(-\frac{t}{\pi}\log\frac{Rtp^{\kappa}}{2\pi e\sqrt{mN}}\right) W\left(\frac{mN}{R^{2}p^{2\kappa}t^{2}}\right) du.$$

$$(10.7)$$

By Lemma 5.1(3),

$$\Phi_1^{\operatorname{sgn}\eta}(y) = \int_0^\infty W(\xi) e\left(-\frac{NXu\xi}{Qqp^\lambda}\right) J_g^{\operatorname{sgn}\eta}(y\xi) \mathrm{d}\xi,\tag{10.8}$$

with $y^k J_g^{\operatorname{sgn} \eta(k)}(y) \ll_{k,g} N^{\varepsilon}$. Applying the Cauchy–Schwarz inequality, we get

$$S_{c}^{\flat}(N) \ll \frac{XN^{3/2}R^{1/2}}{p^{2\lambda + \kappa/2}Q} \sum_{\substack{q \sim R \\ (q,p)=1}} \frac{1}{q^{5/2}} \sum_{d|q} d \cdot \Sigma_{c}(d,q)^{1/2},$$
(10.9)

where

$$\Sigma_{c}(d,q) := \sum_{m} W_{2}\left(\frac{m}{\frac{R^{2}p^{2\kappa}t^{2}}{N}}\right) \bigg| \sum_{\substack{n \ll \frac{q^{2}p^{2\lambda}N^{\varepsilon}}{N} \\ n \equiv mp^{\lambda}\overline{p^{2\kappa-\lambda}} \bmod d}} \lambda_{g}(n)\mathfrak{C}^{*}(m,n,q)J(m,n)\bigg|^{2},$$

with

$$\mathfrak{C}^*(m,n,q) = \sum_{c \bmod p^{\kappa}} \overline{\chi}(c) \sum_{b \bmod p^{\lambda}} e\left(\frac{-\overline{a^*}m\overline{q}}{p^{\kappa}} + \frac{\eta n\overline{q}\overline{b}}{p^{\lambda}}\right)$$

and

$$J(m,n) = \int_{\mathbb{R}} W_q(u) g(q, Xu) e\left(-\frac{t}{\pi} \log \frac{Rtp^{\kappa}}{2\pi e\sqrt{mN}}\right) \Phi_1^{\operatorname{sgn}\eta}\left(\frac{nN}{q^2 p^{2\lambda}}\right) W\left(\frac{mN}{R^2 p^{2\kappa} t^2}\right) \mathrm{d}u$$

We apply the Poisson summation formula on m with modulo dp^{κ} and get

$$\Sigma_c(d,q) \ll \frac{R^2 p^{\kappa} t^2}{Nd} \sum_{\substack{n_1 \ll \frac{q^2 p^{2\lambda_N \varepsilon}}{N}}} |\lambda_g(n_1)|^2 \sum_{\substack{n_2 \ll \frac{q^2 p^{2\lambda_N \varepsilon}}{N} \\ n_2 \equiv n_1 \mod d}} \sum_{\substack{m \in \mathbb{Z}}} |\mathfrak{C}(m,n_1,n_2)\mathfrak{I}_c(m,n_1,n_2)|,$$

where

$$\mathfrak{C}(m,n_1,n_2) = \sum_{\gamma \bmod p^{\kappa}} \mathfrak{C}^*(\gamma,n_1,q) \overline{\mathfrak{C}^*(\gamma,n_2,q)} e\left(\frac{m\left(n_1 p^{3\kappa-\lambda} \overline{p^{\kappa+\lambda}} + \gamma d\overline{d}\right)}{dp^{\kappa}}\right)$$

and

$$\mathfrak{I}_{c}(m,n_{1},n_{2}) = \int_{\mathbb{R}} W_{2}\left(\xi\right) J\left(\frac{R^{2}p^{2\kappa}t^{2}\xi}{N},n_{1}\right) \overline{J\left(\frac{R^{2}p^{2\kappa}t^{2}\xi}{N},n_{2}\right)} e\left(-\frac{mR^{2}p^{\kappa}t^{2}}{Nd}\xi\right) \mathrm{d}\xi. \quad (10.10)$$

Lemma 10.2. (1) If $m \gg \frac{N^{1+\varepsilon}d}{R^2p^{\kappa}t^2}$, we have $\mathfrak{I}_c(m, n_1, n_2) \ll N^{-A}$. (2) If $m \ll \frac{N^{1+\varepsilon}d}{R^2p^{\kappa}t^2}$, we have $\mathfrak{I}_c(m, n_1, n_2) \ll 1$.

Proof. The first result can be done by applying integration by parts with respect to the ξ -integral. For $m \ll \frac{N^{1+\varepsilon_d}}{R^2 p^{\kappa_t 2}}$, we have the trivial bound $\Im_c(m, n_1, n_2) \ll 1$.

The contribution from the zero frequency to $\Sigma_c(d,q)$ is

$$\Sigma_{c0}(d,q) \ll \frac{R^2 p^{\kappa} t^2}{Nd} \sum_{\substack{n_1 \ll \frac{q^2 p^{2\lambda_N \varepsilon}}{N}}} |\lambda_g(n_1)|^2 \left(\sum_{\substack{n_2 \ll \frac{q^2 p^{2\lambda_N \varepsilon}}{N}} p^{2\kappa+\lambda} + \sum_{\substack{n_2 \ll \frac{q^2 p^{2\lambda_N \varepsilon}}{N}} n_2 \equiv n_1 \mod dp^{\lambda-1}}} p^{2\kappa+\lambda-1} \right).$$
(10.11)

Plugging (10.11) into (10.9), we get

$$S_{c0}^{\flat}(N) \ll N^{1/2+\varepsilon} \left(p^{3\kappa/2-3\lambda/4} \frac{t^{3/2}}{K^{5/4}} + p^{\kappa-\lambda/2} \frac{t}{K^{3/2}} \right).$$

The contribution from the non-zero frequencies to $\Sigma_c(d,q)$ is

$$\Sigma_{c1}(d,q) \ll \frac{R^2 p^{\kappa} t^2}{Nd} \sum_{n_1 \ll \frac{q^2 p^{2\lambda_N \varepsilon}}{N}} |\lambda_g(n_1)|^2 \bigg(\sum_{\substack{n_2 \ll \frac{q^2 p^{2\lambda_N \varepsilon}}{N} \\ n_2 \equiv n_1 \bmod dp^{\beta-\kappa+\lambda}}} \sum_{\substack{m \ll \frac{N^{1+\varepsilon}d}{R^2 p^{\kappa+2}} \\ p^{\kappa-\lambda+\beta-\kappa+\lambda} \mid m}} p^{2\kappa+\lambda+\delta_1} + \sum_{\substack{0 \le \ell \le \lambda-\kappa/2}} \sum_{\substack{n_2 \ll \frac{q^2 p^{2\lambda_N \varepsilon}}{N} \\ n_2 \equiv n_1 \bmod dp^\ell}} \sum_{\substack{m \ll \frac{N^{1+\varepsilon}d}{R^2 p^{\kappa+2}} \\ p^{\kappa-\lambda+\ell} \mid m}} p^{5\kappa/2+2\ell+\delta_1+\delta_2/2}} \bigg).$$

$$(10.12)$$

Plugging (10.12) into (10.9), we get

$$S_{c1}^{\flat}(N) \ll N^{1/2+\varepsilon} \left(p^{5\kappa/4 - \lambda/2 + 3/4} \frac{t}{K} + p^{\kappa - \lambda/4 + 1} \frac{t^{1/2}}{K^{5/4}} + p^{3\kappa/4 + \lambda/4 + 3/4} \frac{t^{1/2}}{K^{5/4}} \right).$$

References

- R. Acharya, P. Sharma, S. Singh, t-aspect subconvexity for GL(2)×GL(2) L-function. J. Number Theory 240 (2022), 296–324.
- [2] J. Bernstein and A. Reznikov. Subconvexity bounds for triple L-functions and representation theory. Ann. of Math. (2) 172 (2010), no. 3, 1679–1718.
- [3] V. Blomer. Subconvexity for twisted L-functions on GL(3). Amer. J. Math. 134 (2012), no. 5, 1385–1421.
- [4] V. Blomer, S. Jana, and P. Nelson. The Weyl bound for triple product L-functions. Duke Math. J. 172 (2023), no. 6, 1173–1234.
- [5] V. Blomer, R. Khan, and M. Young. Distribution of mass of holomorphic cusp forms. *Duke Math. J.* 162 (2013), no. 14, 2609–2644.
- [6] D.A. Burgess. On character sums and L-series. II. Proc. London Math. Soc. (3) 13 (1963), no. 14, 524–536.
- [7] P. Deligne. La conjecture de Weil. I. Inst. Hautes Études Sci. Publ. Math. (1974), no. 43, 273–307.
- [8] D. Goldfeld. Automorphic forms and L-functions for the group $GL(n, \mathbf{R})$. With an appendix by Kevin A. Broughan. Cambridge Stud. Adv. Math. 99, Cambridge University Press, Cambridge, 2006.
- [9] B. Huang. On the Rankin-Selberg problem. Math. Ann. 381 (2021), no. 3-4, 1217–1251.
- [10] B. Huang, Q. Sun and H. Zhang. Analytic twists of GL₂ × GL₂ automorphic forms. Math. Nachr. 296 (2023), no. 6, 2366–2394.
- [11] B. Huang and Z. Xu. Hybrid subconvexity bounds for twists of GL(3) × GL(2) L-functions. Algebra & Number Theory 17 (2023), no. 10, 1715–1752.
- [12] M.N. Huxley. Area, lattice points, and exponential sums. London Mathematical Society Monographs. New Series, vol. 13, The Clarendon Press, Oxford University Press, New York, 1996.
- [13] H. Iwaniec and E. Kowalski. Analytic number theory. American Mathematical Society Colloquium Publications, Vol 53. American Mathematical Society, Providence, RI, 2004.
- [14] H. Kim and P. Sarnak. Appendix 2 in Functoriality for the exterior square of GL₄ and the symmetric fourth of GL₂. J. Amer. Math. Soc. 16 (2003), no. 1, 139–183.
- [15] E. Kıral, I. Petrow and M. Young. Oscillatory integrals with uniformity in parameters. J. Théor. Nombres Bordeaux 31 (2019), no. 1, 145–159.
- [16] E. Kowalski, Ph. Michel and J. VanderKam. Rankin–Selberg L-functions in the level aspect. Duke Math. J. 114 (2002), no. 1, 123–191.
- [17] X. Li. Bounds for GL(3) × GL(2) L-functions and GL(3) L-functions. Ann. of Math. (2) 173 (2011), no. 1, 301–336.
- [18] Y. Lin and Q. Sun. Analytic twists of GL₃×GL₂ automorphic forms. Int. Math. Res. Not. IMRN (2021), no. 19, 15143–15208.
- [19] R. Munshi. The circle method and bounds for L-functions, II: Subconvexity for twists of GL(3) L-functions. Amer. J. Math. 137 (2015), no. 3, 791–812.
- [20] R. Munshi. The circle method and bounds for L-functions-III: t-aspect subconvexity for GL(3) L-functions. J. Amer. Math. Soc. 28 (2015), no. 4, 913–938.
- [21] P. Michel and A. Venkatesh. The subconvexity problem for GL₂. Publ. Math. Inst. Hautes Etudes Sci. (2010), no. 111, 171–271.
- [22] Q. Sun. Bounds for GL₂ × GL₂ L-functions in the depth aspect. To appear in Manuscripta Math. https://doi.org/10.1007/s00229-023-01515-1
- [23] E.C. Titchmarsh. The theory of the Riemann zeta-function. Edited and with a preface by D. R. Heath-Brown, 2nd edn., pp.x+412. The Clarendon Press, Oxford University Press, New York, 1986.
- [24] H. Weyl. Zur Abschätzung von $\zeta(1+ti)$ Math. Z. 10 (1921), 88–101.

SCHOOL OF MATHEMATICS, SHANDONG UNIVERSITY, JINAN, SHANDONG 250100, CHINA *Email address:* scc_sdu@mail.sdu.edu.cn

DATA SCIENCE INSTITUTE AND SCHOOL OF MATHEMATICS, SHANDONG UNIVERSITY, JINAN, SHAN-DONG 250100, CHINA

Email address: hmzhang@mail.sdu.edu.cn