RESOLVENT BOUNDS FOR REPULSIVE POTENTIALS

ANDRÉS LARRAÍN-HUBACH, YULONG LI, JACOB SHAPIRO, AND JOSEPH TILLER

ABSTRACT. We prove limiting absorption resolvent bounds for the semiclassical Schrödinger operator with a repulsive potential in dimension $n \ge 3$, which may have a singularity at the origin. As an application, we obtain time decay for the weighted energy of the solution to the associated wave equation with a short range repulsive potential and compactly supported initial data.

1. INTRODUCTION AND STATEMENT OF RESULTS

The goal of this paper is to establish limiting absportion resolvent bounds for the semiclassical Schrödinger operator with a repulsive potential in dimensions three and higher. The dimension one case was studied in [ChDa21, Section 2]. As an application, we obtain time decay of a weighted energy for the solution to the associated wave equation with a short range repulsive potential and compactly supported initial data.

To fix some notation, let $\Delta := \sum_{j=1}^{n} \partial_j^2 \leq 0$ be the Laplacian on \mathbb{R}^n , $n \geq 3$. We use $(r, \theta) = (|x|, x/|x|) \in (0, \infty) \times \mathbb{S}^{n-1}$ for polar coordinates on $\mathbb{R}^n \setminus \{0\}$. Put $B(0, r_0) := \{x \in \mathbb{R}^n : |x| < r_0\}$. For a function u defined on a subset of \mathbb{R}^n , we write $u(r, \theta) := u(r\theta)$, and denote the radial derivative by $u' := \partial_r u$. If $E \subseteq \mathbb{R}^n$ is a Borel set, $\mathbf{1}_E$ stands for its indicator function.

Our Schrödinger operator takes the form

$$P(h) := -h^2 \Delta + V(x) : L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n), \qquad x \in \mathbb{R}^n,$$
(1.1)

where h > 0 is the semiclassical parameter. The conditions we place on the potential $V : \mathbb{R}^n \to \mathbb{R}$ are as follows. We suppose

$$V \ge 0, \tag{1.2}$$

$$rV\mathbf{1}_{B(0,1)} \in L^{\infty}(\mathbb{R}^n), \tag{1.3}$$

$$V\mathbf{1}_{\mathbb{R}^n \setminus B(0,1)} \in L^{\infty}(\mathbb{R}^n), \tag{1.4}$$

for each
$$\theta \in \mathbb{S}^{n-1}$$
, $(0,\infty) \ni r \mapsto V(r,\theta) := V(r\theta)$ has bounded variation. (1.5)

Recall that function f of locally bounded variation on an interval $I \subseteq \mathbb{R}$ has distributional derivative equal to a locally finite Borel measure, which we denote by df; it satisfies

$$\int_{(a,b]} df = f^R(b) - f^R(a),$$
(1.6)

for any interval (a, b] contained in the interior of I, where $f^R(x) := \lim_{\delta \to 0^+} f(x + \delta)$.

The last condition we impose on V is that there exists $C_V > 0$ so that for all $\theta \in \mathbb{S}^{n-1}$,

$$dV(\cdot,\theta) \le -C_V(r+1)^{-1}V(\cdot,\theta)dr.$$
(1.7)

in the sense of Borel measures on $(0, \infty)$.

A prototype potential satisfying the above conditions is

$$V(r,\theta) = g(\theta) \left(\mathbf{1}_{B(0,1)} r^{-1} + 2^{-1} \mathbf{1}_{\mathbb{R}^n \setminus B(0,1)} r^{-\delta} \right)$$

for some $\delta > 0$ and g a bounded function on \mathbb{S}^{n-1} . Note that if $V \in C^1(\mathbb{R}^n; [0, \infty))$, (1.7) implies each $V(\cdot, \theta)$ is repulsive in sense of classical mechanics, i.e., that $V(r, \theta) > 0$ implies $V'(r, \theta) < 0$. The condition (1.3) allows for potentials with an r^{-1} -singularity at the origin, most notably the repulsive Coulomb $(V(x) = r^{-1})$ and Yukawa $(V(x) = e^{-r}r^{-1})$ potentials [Yu]. Moreover, (1.3) and (1.4) imply (1.1) is self-adjoint with respect to the domain $H^2(\mathbb{R}^n)$ [Ne64, Theorem 8]. We utilize $H^2(\mathbb{R}^n)$ in a density argument in Appendix B.

Our main results are the following weighted limiting absorption resolvent bounds for P.

Theorem 1.1. Suppose $n \ge 3$ and V, satisfies (1.2) through (1.7). Define P(h) by (1.1), and equip it with the domain $H^2(\mathbb{R}^n)$. For all $s, s_1, s_2 > 1/2$ with $s_1 + s_2 > 2$, there is C > 0 such that for all $z \in \mathbb{C} \setminus [0, \infty)$ and h > 0,

$$\|(r+1)^{-s}(P(h)-z)^{-1}(r+1)^{-s}\|_{L^2(\mathbb{R}^n)\to L^2(\mathbb{R}^n)} \le \frac{C}{h|z|^{1/2}},$$
(1.8)

$$\|(r+1)^{-s_1}(P(h)-z)^{-1}(r+1)^{-s_2}\|_{L^2(\mathbb{R}^n)\to L^2(\mathbb{R}^n)} \le \frac{C}{h^2}.$$
(1.9)

Christiansen and Datchev [ChDa21, Section 2] obtained (1.8) and (1.9) for bounded, repulsive potentials on the half-line. Thus the novelty of our work is that it extends these bounds to higher dimensions for repulsive potentials which may have a singularity at r = 0.

Remark 1.2. In Appendix C, we recall how for the case V = 0 and n = 3, the conditions on s, s_1 and s_2 in Theorem 1.1, as well as the *h*- and *z*-dependencies of the right sides of (1.8) and (1.9), are nearly optimal in a suitable sense.

We prove Theorem 1.1 by means of the so-called spherical energy method, which is a popular strategy for obtaining resolvent estimates (see, e.g., [CaVo02, Da14, GaSh22]). It relies on separation of variables and the well known identity

$$r^{\frac{n-1}{2}}(-\Delta)r^{-\frac{n-1}{2}} = -\partial_r^2 + r^{-2}\Lambda,$$
(1.10)

where

$$\Lambda := -\Delta_{\mathbb{S}^{n-1}} + \frac{(n-1)(n-3)}{4},\tag{1.11}$$

and $\Delta_{\mathbb{S}^{n-1}}$ denotes the negative Laplace-Beltrami operator on \mathbb{S}^{n-1} . We use in a crucial way that $\Lambda \geq 0$ on $L^2(\mathbb{S}^{n-1})$, see (2.5). This is why our approach does not cover the case n = 2 where the effective potential $-1/(4r^2)$ has a strong negative singularity as $r \to 0$. We expect that repulsive potentials in dimension two can be treated by adapting the Mellin transform methods used in [DGS23, Ob24].

As an application of (1.9), we prove weighted energy decay for the solution to the wave equation

$$\begin{cases} (\partial_t^2 - \Delta + V(x))u(t, x) = 0 & (t, x) \in \mathbb{R} \times \mathbb{R}^n, \ n \ge 3, \\ u(0, x) = u_0(x), \ \partial_t u(0, x) = u_1(x) & x \in \mathbb{R}^n, \end{cases}$$
(1.12)

where $u_0 \in H^1(\mathbb{R}^n)$ and $u_1 \in L^2(\mathbb{R}^n)$ have compact support. The potential V again obeys (1.2) through (1.7), but also the extra short range condition

$$|\mathbf{1}_{\mathbb{R}^n \setminus B(0,1)}V| \le C(r+1)^{-\delta},$$
 (1.13)

for some C > 0 and $\delta > 0$ such that

$$\delta > \begin{cases} \frac{1}{2} + \frac{n+3}{4} & n \neq 8, \\ \frac{1}{2} + 3 & n = 8. \end{cases}$$
(1.14)

Since $P := P(1) = -\Delta + V$ is self-adjoint (and nonnegative) on $L^2(\mathbb{R}^n)$, we may use the spectral theorem for self-adjoint operators to represent the solution to (1.12) by

$$u(t,\cdot) = \cos(t\sqrt{P})u_0 + \frac{\sin(t\sqrt{P})}{\sqrt{P}}u_1.$$
(1.15)

For s > 0 fixed, define the weighted energy of the solution u to (1.12) to be

$$E_s[u](t) = E_s(t) := \int_{\mathbb{R}^n} \langle x \rangle^{-2s} (|\partial_t u(t,x)|^2 + |\nabla u(t,x)|^2 + |u(t,x)|^2) dx.$$

Set also

$$E(0) := \|\nabla u_0\|_{L^2}^2 + \|u_1\|_{L^2}^2$$

Theorem 1.3. Suppose V satisfies (1.2) through (1.7) as well as (1.13) and (1.14). Let $u_0 \in H^1(\mathbb{R}^n)$ and $u_1 \in L^2(\mathbb{R}^n)$ have compact support. For each s such that

$$s > \begin{cases} \frac{n+3}{4} & n \neq 8, \\ 3 & n = 8, \end{cases}$$
(1.16)

there exists $C_s > 0$ depending on s but independent of t, u_0 , and u_1 so that

$$E_s(t) \le C_s \langle t \rangle^{-2} E(0), \tag{1.17}$$

where $\langle t \rangle := (1 + t^2)^{1/2}$.

Remark 1.4. Since $u(-t, \cdot) = \cos(t\sqrt{P})u_0 + (\sin(t\sqrt{P})/\sqrt{P})(-u_1)$, to prove (1.17) it suffices to suppose $t \ge 0$.

For smooth, nonnegative potentials of compact support, the local energy

$$E_{r_0}(t) := \int_{B(0,r_0)} |\partial_t u(t,x)|^2 + |\nabla u(t,x)|^2 + |u(t,x)|^2 dx, \qquad r_0 > 0,$$

obeys

$$E_{r_0}(t) = \begin{cases} O(e^{-ct}) \text{ for some } c > 0 & n \ge 3 \text{ odd,} \\ O(t^{-2n}) & n \ge 4 \text{ even.} \end{cases}$$
(1.18)

Indeed, Vainberg showed (1.18) for compactly supported perturbations of the Laplacian satisfying the so called Generalized Huygens Principle (defined in [Vo04c]). From a propagation of singularities result of Melrose and Sjöstrand [MeSj78, MeSj82], the Generalized Huygens Principle holds for a large class of the so called nontrapping perturbations of Laplacian, which includes smooth, nonnegative potentials compact support. The study of energy decay for nontrapping perturbations has a long history, going back to the works of Lax, Morawetz, and Phillips [LMP63, Mo66, Mo75].

On the other hand, bounds similar to (1.17) were obtained in previous works for various classes of short range potentials. In [Za04], Zappacosta considered potentials $V \in C^1(\mathbb{R}^3; (0, \infty))$ with $\partial_x^{\alpha} V = O(\langle x \rangle^{-\delta - |\alpha|})$ for all $0 \leq |\alpha| \leq 1$ and some $\delta > 2$. For the each $\chi \in C_0^{\infty}(\mathbb{R}^3)$, the bound $\|\chi \sqrt{V}(\sin(t\sqrt{P})/\sqrt{P})\sqrt{V}\chi\|_{L^2 \to L^2}^2 = O(t^{-2})$ was obtained. Vodev extended [Za04] in [Vo04a] by obtaining $E_{r_0}(t) = O(t^{-2})$ in dimension $n \geq 3$, where $V \in C^1(\mathbb{R}^n; [0, \infty))$ obeys

$$\partial_x^{\alpha} V = O(\langle x \rangle^{-\delta_0}) \quad \text{for all } 0 \le |\alpha| \le 1 \text{ and some } \delta_0 > (n+1)/2, \text{ and} \quad (1.19)$$

$$2V + r\partial_r V \le C\langle x \rangle^{-\delta}$$
 for some $C > 0$ and some $\delta > 1$. (1.20)

We note that if $V \in C^1(\mathbb{R}^n; [0, \infty))$ satisfies (1.7) with $C_V \geq 2$, then (1.20) holds. Vodev also obtained weighted energy decay for a class of long range, nontrapping perturbations of the Laplacian that includes perturbation by a nonnnegative long range potential, provided the initial conditions are spectrally localized away from [0, a] for a > 0 sufficiently large [Vo04b].

The proof of Theorem 1.3 proceeds as in [Vo04a, Section 3], with some modifications to address the possible singular behavior of V at the origin. It suffices to establish

$$t^2 E_s(t) \le C t^2 \int_t^\infty E_s(\tau) d\tau \le C E(0), \qquad t \ge 1.$$
(1.21)

For this, we use Duhamel's formula and the Fourier transform, with t dual to λ , to represent the Fourier transform of u in terms of $(P - \lambda^2)^{-1}$, see (4.5). Since the initial data are compactly

supported, finite speed of propagation holds, and we may freely introduce a cutoff function η into this expression. So Plancherel's theorem gives a route to control $E_s(t)$, thanks to the bound on $\langle x \rangle^{-s} (P - \lambda^2)^{-1} \eta$ that comes from (1.9), see also Lemma 3.1. However, due the factor of t^2 in (1.21), and since multiplication by t is dual to differentiation with respect to λ , it is necessary to control $\langle x \rangle^{-s} \frac{d}{d\lambda} (P - \lambda^2)^{-1} \eta$ too. We obtain bounds on this derivative under the additional conditions (1.13), (1.14), and (1.16), see Lemma 3.4.

Another energy studied is the quantity

$$E_K^{(1)}[u](t) := \int_K |\partial_t u(t,x)|^2 + |\nabla u(t,x)|^2 + V(x)|u(t,x)|^2 dx,$$

where K is suitable subset of \mathbb{R}^n . In [Vo04a, Theorem 1.1], Vodev considered $K = B(0, \gamma_0 t) \subseteq \mathbb{R}^n$ for $n \geq 3$ and suitable $0 < \gamma_0 < 1$. He showed $E_K^{(1)}[u](t) = O(t^{-1})$ provided (1.19) and (1.20) hold. In [Ik23, Theorems 1.1 and 1.2], Ikehata considered exterior subdomains Ω of \mathbb{R}^n , $n \geq 2$, that exclude the origin. For K a compact subset of Ω , he showed $E_K^{(1)}[u](t) = O(t^{-1})$ provided V is nonnegative, C^1 , and obeys $x \cdot \nabla V + 2V \leq 0$. It seems $O(t^{-1})$ -decay was first obtained first by Morawetz [Mo61] for V = 0 in the exterior of a three-dimensional star-shaped obstacle (and later improved to exponential decay in [LMP63]).

As alluded to above, there is a large body literature on wave decay for higher order perturbations. See [Bu98, Vo99, Bu02, Bo11, CaVo04, Sh18, ChIk20] for more historical background.

The rest of the paper is organized as follows. In Section 2, we prove Theorem 1.1. In Section 3, we apply Theorem 1.1 to prove norm bounds for $\langle x \rangle^{-s} (P - \lambda^2)^{-1} \langle x \rangle^{-s}$ and its λ -derivative. In Section 4 we prove Theorem 1.3. Finally, we include several appendices of technical results that assist with the proofs of earlier sections.

ACKNOWLEDGEMENTS: We thank Kiril Datchev and Georgi Vodev for helpful discussions. J. S. and A. L-H. gratefully acknowledge support from NSF DMS-2204322. J. S. was also supported by a University of Dayton Research Council Seed Grant.

2. Proof of Theorem 1.1

In this section, we prove Theorem 1.1. Throughout this section, we take P(h) as in (1.1), and assume the potential V satisfies (1.2) through (1.7).

By (1.10),

$$P^{\pm}(h) := r^{\frac{n-1}{2}} \left(P(h) - E \pm i\varepsilon \right) r^{-\frac{n-1}{2}} = -h^2 \partial_r^2 + h^2 r^{-2} \Lambda + V - E \pm i\varepsilon.$$
(2.1)

Let $u \in r^{(n-1)/2} C_0^{\infty}(\mathbb{R}^n)$. Define a spherical energy functional F[u](r),

$$F(r) = F[u](r) := \|hu'(r, \cdot)\|^2 - \langle (h^2 r^{-2} \Lambda + V(r, \cdot) - E)u(r, \cdot), u(r, \cdot) \rangle,$$
(2.2)

where $\|\cdot\|$ and $\langle\cdot,\cdot\rangle$ denote the norm and inner product on $L^2(\mathbb{S}^{n-1}_{\theta})$. For a weight w(r) which is absolutely continuous, nonnegative, and increasing, one computes (wF)' in the sense of distributions

on $(0,\infty)$:

$$(wF)' = wF' + w'F$$

$$= w(-2\operatorname{Re}\langle (-h^{2}\partial_{r}^{2} + h^{2}r^{-2}\Lambda + V - E)u, u'\rangle$$

$$+ 2h^{2}r^{-3}\langle\Lambda u, u\rangle - \int_{\mathbb{S}^{n-1}} |u(\theta, r)|^{2}dV(r, \theta)d\theta\rangle$$

$$+ w'(||hu'||^{2} - \langle h^{2}r^{-2}\Lambda u, u\rangle + \langle (E - V)u, u\rangle$$

$$= -2w\operatorname{Re}\langle P^{\pm}(h)u, u'\rangle \mp 2\varepsilon w\operatorname{Im}\langle u, u'\rangle + w'||hu'||^{2}$$

$$+ (2wr^{-1} - w')\langle h^{2}r^{-2}\Lambda u, u\rangle + Ew'||u||^{2}$$

$$- \int_{\mathbb{S}^{n-1}} |u(\theta, r)|^{2}(w(r)dV(r, \theta) + w'(r)V(r, \theta))d\theta.$$
(2.3)

First we show (1.8). Since increasing s decreases the left side of (1.8), without loss of generality we may take $0 < \delta := 2s - 1 < 1$. We will show the last line of (2.3) can be made to have a suitable lower bound, using

$$w(r) := 1 - \frac{C_V}{C_V + \delta} (1+r)^{-\delta}.$$
(2.4)

For such w, we clearly have

$$w'(r) = \frac{\delta C_V}{C_V + \delta} (r+1)^{-1-\delta},$$

Therefore, on the one hand

$$2wr^{-1} - w' = 2r^{-1}\left(1 - \frac{C_V}{C_V + \delta}(r+1)^{-\delta}\left[1 - \frac{\delta}{2}\frac{r}{r+1}\right]\right) \ge 0,$$

since $\delta < 1$. On the other hand, in the sense of measures of Borel measures on $(0, \infty)$, using (1.7)

$$wdV + w'V = \frac{\delta C_V V}{(C_V + \delta)(r+1)^{1+\delta}} + wdV \le \frac{C_V V}{1+r}((r+1)^{-\delta} - 1) \le 0$$

Thus, the last two estimates and (2.3) imply, in the sense of distributions on $(0, \infty)$,

$$(wF)' \ge -2w \operatorname{Re} \langle P^{\pm}(h)u, u' \rangle \mp 2\varepsilon w \operatorname{Im} \langle u, u' \rangle + w' ||hu'||^2 + Ew' ||u||^2.$$
(2.5)

Integrating (2.5) with respect to dr from $r = r_0 > 0$ to $r = \infty$ implies (because u is compactly supported)

$$\int_{r_0}^{\infty} Ew' \|u\|^2 + w' \|hu'\|^2 dr + w(r_0) F(r_0) \le \int_{r_0}^{\infty} 2w \operatorname{Re} \langle P^{\pm}(h)u, u' \rangle \pm 2\varepsilon w \operatorname{Im} \langle u, u' \rangle dr.$$
(2.6)

Since $u = r^{(n-1)/2}v$ for some $v \in C_0^{\infty}(\mathbb{R}^n)$, we recognize that

$$F(r_0) = \|hu'(r_0, \cdot)\|^2 + r_0^{n-3} \langle h^2 \Delta_{\mathbb{S}^{n-1}} v(r_0, \cdot), v(r_0, \cdot) \rangle + (Er_0^{n-1} - h^2 4^{-1} (n-1)(n-3)r_0^{n-3}) \|v(r_0, \cdot)\|^2 + r_0^{n-1} \langle V(r_0, \cdot)v(r_0, \cdot), v(r_0, \cdot) \rangle.$$

$$(2.7)$$

We rewrite the term in (2.7) involving $\Delta_{\mathbb{S}^{n-1}}$ using the well known formula for the Laplacian in spherical coordinates:

$$r^{-2}\Delta_{\mathbb{S}^{n-1}} = \Delta - \partial_r^2 - (n-1)r^{-1}\partial_r.$$

Therefore,

We can express the differential operators ∂_r and ∂_r^2 with respect to the Euclidean coordinate system,

$$\partial_r = r^{-1} \sum_{j=1}^n x_j \partial_{x_j}, \qquad \partial_r^2 = r^{-2} \sum_{k=1}^n x_k \sum_{j=1}^n x_j \partial_{x_k} \partial_{x_j}.$$
(2.9)

Thus by (2.8) and (2.9), all terms in (2.7) tend to zero as $r_0 \to 0$, except for possibly $||hu'(r, \cdot)||^2$ in dimension three, which in that case tends to $|v(0)|^2 \int_{\mathbb{S}^{n-1}} d\theta$. We conclude

$$\lim_{r_0 \to 0} w(r_0) F(r_0) = w(0) F(0) = \begin{cases} \omega_{n-1} w(0) |v(0)|^2 & n = 3, \\ 0 & n \ge 4, \end{cases}$$

where ω_{n-1} is the (n-1)-dimensional volume of \mathbb{S}^{n-1} .

Thus in view of (2.6) and $0 < w \le 1$,

$$\int_{0}^{\infty} Ew' \|u\|^{2} + w' \|hu'\|^{2} dr
\leq 2 \Big(\int_{0}^{\infty} \frac{1}{h^{2}w'} \|P^{\pm}(h)u\|^{2} dr \Big)^{1/2} \Big(\int_{0}^{\infty} w' \|hu'\|^{2} dr \Big)^{1/2}
+ \frac{2\varepsilon}{h} \Big(\int_{0}^{\infty} \|u\|^{2} dr \Big)^{1/2} \Big(\int_{0}^{\infty} \|hu'\|^{2} dr \Big)^{1/2}.$$
(2.10)

We now estimate,

$$\begin{split} \int_{0}^{\infty} \|hu'\|^{2} dr &= \operatorname{Re} \int_{0}^{\infty} \langle u, -h^{2}u'' \rangle dr \\ &= \operatorname{Re} \left(\int_{0}^{\infty} \langle u, P^{\pm}(h)u \rangle dr + \int_{0}^{\infty} \langle u, (E - V - h^{2}r^{-2}\Lambda)u \rangle dr \mp i\varepsilon \int_{0}^{\infty} \|u\|^{2} dr \right) \\ &= \operatorname{Re} \int_{0}^{\infty} \langle u, P^{\pm}(h)u \rangle dr + \int_{0}^{\infty} \langle u, (E - V - h^{2}r^{-2}\Lambda)u \rangle dr \\ &\leq \left(\int_{0}^{\infty} \frac{1}{w'} \|P^{\pm}(h)u\|^{2} dr \right)^{1/2} \left(\int_{0}^{\infty} w' \|u\|^{2} dr \right)^{1/2} + E \int_{0}^{\infty} \|u\|^{2} dr, \end{split}$$

and

$$\begin{split} \varepsilon \int_0^\infty \|u\|^2 dr &= \varepsilon \|v\|_{L^2}^2 \\ &= |\operatorname{Im}\langle (P(h) - E \pm i\varepsilon)v, v\rangle_{L^2}| \\ &= |\operatorname{Im} \int_0^\infty \langle P^{\pm}(h)u, u\rangle dr| \\ &\leq \big(\int_0^\infty \frac{1}{w'} \|P^{\pm}(h)u\|^2 dr\big)^{1/2} \big(\int_0^\infty w' \|u\|^2 dr\big)^{1/2} \end{split}$$

Combining these gives

$$\frac{\varepsilon^2}{h^2} \int_0^\infty \|u\|^2 dr \cdot \int_0^\infty \|hu'\|^2 dr \le (E+\varepsilon) \int_0^\infty \frac{1}{h^2 w'} \|P^{\pm}(h)u\|^2 dr \cdot \int_0^\infty w' \|u\|^2 dr$$
in this into (2.10) yields

Pluggin

$$\int_{0}^{\infty} Ew' \|u\|^{2} + w' \|hu'\|^{2} dr \leq 2 \Big(\int_{0}^{\infty} \frac{1}{h^{2}w'} \|P^{\pm}(h)u\|^{2} dr \Big)^{1/2} \\ \cdot \Big(\Big(\int_{0}^{\infty} w' \|hu'\|^{2} dr \Big)^{1/2} + (E+\varepsilon)^{1/2} \Big(\int_{0}^{\infty} w' \|u\|^{2} dr \Big)^{1/2} \Big).$$

$$(2.11)$$

Completing the square in (2.11), we find

$$(E^{1/2} (\int_0^\infty w' \|u\|^2 dr)^{1/2} - \frac{(E+\varepsilon)^{1/2}}{E^{1/2}} (\int_0^\infty \frac{1}{h^2 w'} \|P^{\pm}(h)u\|^2 dr)^{1/2})^2 + ((\int_0^\infty w' \|hu'\|^2 dr)^{1/2} - (\int_0^\infty \frac{1}{h^2 w'} \|P^{\pm}(h)u\|^2 dr)^{1/2})^2$$

$$\leq \frac{2E+\varepsilon}{E} \int_0^\infty \frac{1}{h^2 w'} \|P^{\pm}(h)u\|^2 dr.$$

$$(2.12)$$

Dropping the second term on the left side of (2.12) implies

$$E^{1/2} \Big(\int_0^\infty w' \|u\|^2 dr \Big)^{1/2} \le \Big(\frac{(E+\varepsilon)^{1/2}}{E^{1/2}} + \frac{(2E+\varepsilon)^{1/2}}{E^{1/2}} \Big) \Big(\int_0^\infty \frac{1}{h^2 w'} \|P^{\pm}(h)u\|^2 dr \Big)^{1/2}.$$
(2.13)

Next, consider the sector $\{z \in \mathbb{C} : |\operatorname{Im} z| < \alpha \operatorname{Re} z\}$ for $0 < \alpha < 1$. Since $w' = C_V \delta(C_V + \delta)^{-1} (r+1)^{-1-\delta}$ and $\delta = 2s - 1$, from (2.13), we get for all h > 0, $E \pm i\varepsilon \in \{z \in \mathbb{C} : |\operatorname{Im} z| < \alpha \operatorname{Re} z\}$, and $u \in r^{n-1/2} C_0^{\infty}(\mathbb{R}^n)$,

$$|(E \pm i\varepsilon)^{1/2}| \Big(\int_0^\infty (r+1)^{-2s} ||u||^2 dr \Big)^{1/2} \le h^{-1} (1+\alpha^2)^{1/4} \Big(\frac{1}{\delta} + \frac{1}{C_V} \Big) ((1+\alpha)^{1/2} + (2+\alpha)^{1/2}) \Big(\int_0^\infty (r+1)^{2s} ||P^{\pm}(h)u||^2 dr \Big)^{1/2},$$
(2.14)

Here, our branch of the complex square root is chosen so that $\operatorname{Im}(E \pm i\varepsilon)^{1/2} > 0$, and we used that $|(E \pm i\varepsilon)|^{1/2} = (E^2 + \varepsilon^2)^{1/4} \le E^{1/2}(1 + \alpha^2)^{1/4}$ for $E \pm i\varepsilon \in \{z \in \mathbb{C} : |\operatorname{Im} z| < \alpha \operatorname{Re} z\}.$

Since $u \in r^{(n-1)/2}C_0^{\infty}(\mathbb{R}^n)$, a standard density argument, which we review in Appendix B, shows that (2.14) implies

$$\begin{aligned} \|z^{1/2}(r+1)^{-s}(P(h)-z)^{-1}(r+1)^{-s}\|_{L^2 \to L^2} \\ &\leq h^{-1}(1+\alpha^2)^{1/4} \left(\frac{1}{\delta} + \frac{1}{C_V}\right)((1+\alpha)^{1/2} + (2+\alpha)^{1/2}), \end{aligned}$$
(2.15)

on $\{z \in \mathbb{C} : |\operatorname{Im} z| < \alpha \operatorname{Re} z\}$ and for any $0 < \alpha < 1$. To extend this bound to all $z \in \mathbb{C} \setminus [0, \infty)$, we use the Phragmén-Lindelöf principle [EM] in the following way. For $u, v \in L^2(\mathbb{R}^n)$, put

$$U(z) := z^{1/2} \langle (r+1)^{-s} (P(h) - z)^{-1} (r+1)^{-s} u, v \rangle_{L^2}.$$

Then U(z) in analytic in $\Omega_{\alpha} := \{z \in \mathbb{C} : \alpha \operatorname{Re} z < |\operatorname{Im} z|\}$. By (2.15), on $\partial \Omega_{\alpha} \setminus \{0\}$ we have

$$|U(z)| \le h^{-1}(1+\alpha^2)^{1/4} \left(\frac{1}{\delta} + \frac{1}{C_V}\right) \left((1+\alpha)^{1/2} + (2+\alpha)^{1/2}\right) \|u\|_{L^2} \|v\|_{L^2}.$$
 (2.16)

On the other hand, in Ω_{α} , we have the standard bound

$$|U(z)| \le \frac{|z|^{1/2} ||u||_{L^2} ||v||_{L^2}}{\operatorname{dist}(z, [0, \infty))} = \begin{cases} \frac{||u||_{L^2} ||v||_{L^2}}{|z|^{1/2}} & \operatorname{Re} z < 0, \\ \frac{|z|^{1/2} ||u||_{L^2} ||v||_{L^2}}{|\operatorname{Im} z|} & \operatorname{Re} z \ge 0, \ z \in \Omega_{\alpha}, \end{cases}$$
(2.17)

where we used

$$\frac{1}{\operatorname{dist}(z, [0, \infty))} = \frac{1}{\inf_{r \ge 0} ((\operatorname{Re} z - r)^2 + (\operatorname{Im} z)^2)^{1/2}} = \begin{cases} \frac{1}{|z|} & \operatorname{Re} z < 0, \\ \frac{1}{|\operatorname{Im} z|} & \operatorname{Re} z \ge 0, \ z \in \Omega_{\alpha}. \end{cases}$$

Finally, define, $g(z) = e^{i(z^{-1})^{1/2}}$, where our branch of the square root is as above. In Ω_{α} , $|g(z)| \leq e^{-c|z|^{-1/2}}$ for some 0 < c < 1 depending on α . This with (2.17) says that for any $\sigma > 0$,

$$\limsup_{z \to 0, \, z \in \Omega_{\alpha}} |g(z)|^{\sigma} |U(z)| = 0.$$
(2.18)

Therefore, from (2.16) and (2.18), the Phragmén Lindelöf Theorem (Theorem A.1 in Appendix A) implies that (2.15) holds for all $z \in \Omega_{\alpha}$ too. Sending $\alpha \to 0^+$ completes the proof of (1.8).

To prove (1.9), start again at (2.12) and drop the first term on the left hand side. Still working on $\{z \in \mathbb{C} : |\operatorname{Im} z| < \alpha \operatorname{Re} z\}$, some manipulations give

$$\left(\int_0^\infty w' \|hu'\|^2 \, dr\right)^{1/2} \le (1 + \sqrt{2 + \alpha}) \left(\int_0^\infty \frac{1}{h^2 w'} \|P^{\pm}(h)u\|^2 \, dr\right)^{1/2}.$$
(2.19)

By integration by parts,

$$\begin{split} \int_0^\infty (r+1)^{-3-\delta} \|u\|^2 \, dr &= \frac{2}{2+\delta} \int_0^\infty (r+1)^{-2-\delta} \operatorname{Re}\langle u, u' \rangle \, dr \\ &\leq h^{-1} \left(\int_0^\infty (r+1)^{-1-\delta} \|hu'\|^2 \, dr \right)^{1/2} \left(\int_0^\infty (r+1)^{-3-\delta} \|u\|^2 \, dr \right)^{1/2}, \end{split}$$

which implies

$$\left(\int_0^\infty (r+1)^{-3-\delta} \|u\|^2 \, dr\right)^{1/2} \le h^{-1} \left(\int_0^\infty (r+1)^{-1-\delta} \|hu'\|^2 \, dr\right)^{1/2}.$$
(2.20)

From (2.19), (2.20) and $w' = C_V \delta (C_V + \delta)^{-1} (r+1)^{-1-\delta}$,

$$\left(\int_0^\infty (r+1)^{-3-\delta} \|u\|^2 \, dr\right)^{1/2} \le h^{-2} \left(\frac{1}{\delta} + \frac{1}{C_V}\right) (1 + \sqrt{2+\alpha}) \left(\int_0^\infty (r+1)^{1+\delta} \|P^{\pm}(h)u\|^2 \, dr\right)^{1/2}.$$

Using again the density argument in Appendix B, for $0 < \delta < 1$,

$$\|(1+r)^{-\frac{3+\delta}{2}}(P(h)-z)^{-1}(1+r)^{-\frac{1+\delta}{2}}\|_{L^2\to L^2} \le h^{-2}\left(\delta^{-1}+C_V^{-1}\right)(1+\sqrt{2+\alpha}),\tag{2.21}$$

on $\{z \in \mathbb{C} : |\operatorname{Im} z| < \alpha \operatorname{Re} z\}$. Then, as above, (2.17), the Phragmén Lindelöf Theorem, and sending $\alpha \to 0^+$, imply

$$\|(1+r)^{-\frac{3+\delta}{2}}(P(h)-z)^{-1}(1+r)^{-\frac{1+\delta}{2}}\|_{L^2\to L^2} \le h^{-2}\left(\delta^{-1}+C_V^{-1}\right)(1+\sqrt{2}), \qquad z\in\mathbb{C}\setminus[0,\infty).$$

Since the norm of an operator and its adjoint coincide,

$$\|(1+r)^{-\frac{1+\delta}{2}}(P(h)-z)^{-1}(1+r)^{-\frac{3+\delta}{2}}\|_{L^2\to L^2} \le h^{-2}\left(\delta^{-1}+C_V^{-1}\right)(1+\sqrt{2}), \qquad z\in\mathbb{C}\setminus[0,\infty).$$

The three lines lemma then says that for fixed $z \in \mathbb{C} \setminus [0, \infty)$, the analytic mapping

$$\lambda \mapsto (1+r)^{-\frac{3+\delta}{2}+\lambda} (P(h)-z)^{-1}(1+r)^{-\frac{1+\delta}{2}-\lambda}, \qquad 0 < \operatorname{Re} \lambda < 1,$$

(with values in the space of bounded operators $L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n))$ obeys

$$\|(1+r)^{-\frac{3+\delta}{2}+\theta}(P(h)-z)^{-1}(1+r)^{-\frac{1+\delta}{2}-\theta}\|_{L^2\to L^2} \le h^{-2}\left(\delta^{-1}+C_V^{-1}\right)(1+\sqrt{2}), \qquad \theta \in [0,1].$$
(2.22)

Having established (2.22), to finish, we need to see that we can choose δ and θ appropriately to arrive at (1.9). That is, we need to attain the more general weights characterized by $s_1, s_2 > 1/2$, $s_1+s_2 > 2$. However, because we have the restrictions $\delta \in (0, 1)$ and $\theta \in [0, 1]$, we first need to make reductions as follows. Since decreasing s_1 or s_2 in (1.9) increases the right side, it suffices to suppose $s_1, s_2 > 1/2, 2 < s_1+s_2 < 3$. Furthermore, by taking the adjoint, it is no restriction to have $s_1 \leq s_2$. If we write $s_1 = (1+2\delta_1)/2$ for some $\delta_1 > 0$, then we may replace s_2 by min $(s_2, (3+\delta_1)/2)$. Having made these reductions, (1.9) follows from (2.22) by setting $\delta = s_1+s_2-2 < 1, \theta = (s_2-s_1+1)/2 \leq (4-\delta_1)/4 < 1$.

3. Resolvent bounds for wave decay

In this section, we consider the operator $P = P(1) = -\Delta + V$, with P(h) as in (1.1); V obeys (1.2) through (1.7). As a consequence of Theorem 1.1, we prove several more resolvent bounds for P, which enable us in Section 4 to establish weighted energy decay for the solution to the wave equation (1.12). Throughout this section, C denotes a positive constant whose precise value may change, but is always independent λ , which plays the role of our spectral parameter.

Lemma 3.1. Fix $s_1, s_2 > 1/2$ with $s_1 + s_2 > 2$. There exist C > 0 so that for all $\lambda \in \mathbb{C}$ with $0 < |\operatorname{Im} \lambda| \le 1$, and for all multiindices α_1, α_2 with $|\alpha_1| + |\alpha_2| \le 2$,

$$\|\langle x\rangle^{-s_1}\partial_x^{\alpha_1}(P-\lambda^2)^{-1}\partial_x^{\alpha_2}\langle x\rangle^{-s_2}\|_{L^2(\mathbb{R}^n)\to L^2(\mathbb{R}^n)} \le C(1+|\operatorname{Re}\lambda|)^{|\alpha_1|+|\alpha_2|-1}.$$
(3.1)

Proof. Since $((P - \lambda^2)^{-1})^* = (P - (\overline{\lambda})^2)^{-1}$, to prove (3.1) is suffices to assume Im $\lambda > 0$. First, we treat the case $|\alpha_2| = 0$. Using (1.8) if $|\operatorname{Re} \lambda| > 1$ or (1.9) if $|\operatorname{Re} \lambda| \le 1$, we get

$$|\langle x \rangle^{-s_1} (P - \lambda^2)^{-1} \langle x \rangle^{-s_2} \|_{L^2 \to L^2} \le C (1 + |\operatorname{Re} \lambda|)^{-1}, \qquad 0 < \operatorname{Im} \lambda \le 1,$$
 (3.2)

Recall from standard elliptic theory that for all $f \in H^2(\mathbb{R}^n)$ and all $\gamma > 0$,

$$\|f\|_{H^2} \le C(\|f\|_{L^2} + \|\Delta f\|_{L^2}),$$

$$\|f\|_{H^1}^2 \le C\|f\|_{L^2}\|f\|_{H^2} \le C(\gamma^{-1}\|f\|_{L^2}^2 + \gamma\|\Delta f\|_{L^2}^2).$$

(3.3)

Therefore, for any $f \in L^2(\mathbb{R}^n)$,

$$\begin{aligned} \|\langle x \rangle^{-s_{1}} (P - \lambda^{2})^{-1} \langle x \rangle^{-s_{2}} f\|_{H^{2}(\mathbb{R}^{n})} \\ &\leq C(\|\langle x \rangle^{-s_{1}} (P - \lambda^{2})^{-1} \langle x \rangle^{-s_{2}} f\|_{L^{2}(\mathbb{R}^{n})} + \|(-\Delta) \langle x \rangle^{-s_{1}} (P - \lambda^{2})^{-1} \langle x \rangle^{-s_{2}} f\|_{L^{2}(\mathbb{R}^{n})}) \\ &\leq C(\|\langle x \rangle^{-s_{1}} (P - \lambda^{2})^{-1} \langle x \rangle^{-s_{2}} f\|_{H^{1}(\mathbb{R}^{n})} + \|\langle x \rangle^{-s_{1}} (-\Delta) (P - \lambda^{2})^{-1} \langle x \rangle^{-s_{2}} f\|_{L^{2}(\mathbb{R}^{n})}) \\ &\leq C(\gamma^{-1} + |\operatorname{Re} \lambda|^{2}) \|\langle x \rangle^{-s_{1}} (P - \lambda^{2})^{-1} \langle x \rangle^{-s_{2}} f\|_{L^{2}(\mathbb{R}^{n})} \\ &+ C\gamma \|\Delta \langle x \rangle^{-s_{1}} (P - \lambda^{2})^{-1} \langle x \rangle^{-s_{2}} f\|_{L^{2}(\mathbb{R}^{n})} \\ &+ C \|f\|_{L^{2}(\mathbb{R}^{n})}. \end{aligned}$$

Selecting γ sufficiently small depending on C, and applying (3.2) yields

$$\|\langle x \rangle^{-s_1} (P - \lambda^2)^{-1} \langle x \rangle^{-s_2} f\|_{H^2(\mathbb{R}^n)} \le C(1 + |\operatorname{Re} \lambda|) \|f\|_{L^2(\mathbb{R}^n)},$$
(3.4)

as desired. This confirms (3.1) for $|\alpha_1| = 2$. For $|\alpha_1| = 1$ (still with $|\alpha_2| = 0$), combine (3.2) and (3.4) via the second line of (3.3).

If $|\alpha_2| > 0$, let $f \in C_0^{\infty}(\mathbb{R}^n)$, and put $u = \langle x \rangle^{-s_1} (P - \lambda^2)^{-1} \langle x \rangle^{-s_2} \partial_x^{\alpha_2} f$. We need to show

$$||u||_{H^{|\alpha_2|}} \le C(1+|\operatorname{Re}\lambda|)^{|\alpha_1|+|\alpha_2|-1}||f||_{L^2}, \qquad H^0 := L^2(\mathbb{R}^n).$$

If $|\alpha_1| = 0$, we use self-adjointness and $||\langle x \rangle^{-s_2} (P - \lambda^2)^{-1} \langle x \rangle^{-s_1} f||_{H^1(\mathbb{R}^n)} \leq C ||f||_{L^2(\mathbb{R}^n)}$ to get

$$\begin{aligned} \|u\|_{L^{2}}^{2} &= \langle u, \langle x \rangle^{-s_{1}} (P - \lambda^{2})^{-1} \langle x \rangle^{-s_{2}} \partial_{x}^{\alpha_{2}} f \rangle_{L^{2}} \\ &\leq \|\partial_{x}^{\alpha_{2}} \langle x \rangle^{-s_{2}} (P - (\overline{\lambda})^{2})^{-1} \langle x \rangle^{-s_{1}} u\|_{L^{2}} \|f\|_{L^{2}} \\ &\leq C (1 + |\operatorname{Re} \lambda|)^{|\alpha_{2}| - 1} \|u\|_{L^{2}} \|f\|_{L^{2}}. \end{aligned}$$

If $|\alpha_1| = 1$, we recognize that $(P - \lambda^2)u = \langle x \rangle^{-s_1 - s_2} \partial_x^{\alpha_2} f + [-\Delta, \langle x \rangle^{-s_1}] \langle x \rangle^{s_1} u$. Then multiply by \overline{u} , integrate over \mathbb{R}^n , and integrate by parts as appropriate

$$\|\nabla u\|_{L^2}^2 = \int (\lambda^2 - V)|u|^2 - \int \partial_x^{\alpha_2} (\langle x \rangle^{-s_1 - s_2} \overline{u})f + \int \overline{u}[-\Delta, \langle x \rangle^{-s_1}] \langle x \rangle^{s_1} u.$$

Because both

$$[-\Delta, \langle x \rangle^{-s_1}] \langle x \rangle^{s_1} = (-\Delta \langle x \rangle^{-s_1}) \langle x \rangle^{s_1} - 2(\nabla \langle x \rangle^{-s_1}) \cdot \nabla \langle x \rangle^{s_1},$$

and $\partial_x^{\alpha_2} \langle x \rangle^{-s_1-s_2}$ are first order differential operators with bounded coefficients, we conclude, for all $\gamma > 0$,

$$\begin{aligned} \|\nabla u\|_{L^2}^2 &\leq C_{\gamma}((1+|\operatorname{Re}\lambda|)^2 \|u\|_{L^2}^2 + \|f\|_{L^2}^2) + \gamma \|\nabla u\|_{L^2}^2 \\ &\leq C_{\gamma}(1+|\operatorname{Re}\lambda|)^2 \|f\|_{L^2}^2 + \gamma \|\nabla u\|_{L^2}^2, \end{aligned}$$

for some $C_{\gamma} > 0$ depending on γ . Note also we used (1.3) and Lemma E.1 to estimate

$$\int V|u|^{2} \leq \int \mathbf{1}_{B(0,1)} V|u|^{2} + \int \mathbf{1}_{\mathbb{R}^{n} \setminus B(0,1)} V|u|^{2}$$
$$\leq C \int \mathbf{1}_{B(0,1)} |r^{-1}u||u| + C ||u||_{L^{2}}^{2}$$
$$\leq C(||u||_{L^{2}} ||\nabla u||_{L^{2}} + ||u||_{L^{2}}^{2}).$$

Fixing γ small enough, we absorb the second term on the right side into the left side, confirming (3.1) when $|\alpha_1| = |\alpha_2| = 1$.

Next, we prove an estimate for the derivative of the weighted resolvent, which requires an extra short range condition on the potential. As input we need the following bound for the weighted square of the free resolvent, which we prove in Appendix ??.

Lemma 3.2. Let $n \ge 3$, $j \in \{0, 1\}$, and suppose s satisfies (1.16). There exists C > 0 such that for all $\lambda \in \mathbb{C}$ with $0 < |\operatorname{Im} \lambda| \le 1$,

$$\|\lambda\langle x\rangle^{-s}\nabla^{j}(-\Delta-\lambda^{2})^{-2}\langle x\rangle^{-s}\|_{L^{2}(\mathbb{R}^{n})\to L^{2}(\mathbb{R}^{n})} \leq C(1+|\lambda|)^{j-1}.$$
(3.5)

Remark 3.3. In [Vo04a], the estimate (3.5) is stated to hold in any dimension $n \geq 3$ provided s > 3/2. However, our proof of Lemma 3.2 in dimension $n \geq 4$ needs s larger if (3.5) is to hold uniformly as $|\lambda| \to 0$. In our approach, we use the integral kernel of $\lambda \langle x \rangle^{-s} (-\Delta - \lambda^2)^{-2} \langle x \rangle^{-s}$ to assess L^2 -boundedness as $|\lambda| \to 0$. The kernel is given in terms of the Macdonald function [DLMF, 10.27.4, 10.27.5] of order n/2 - 2, along with other factors. We are able to conclude boundedness on $L^2(\mathbb{R}^n)$ for s as in (1.16).

Lemma 3.4. Let $n \ge 3$ and s as in (1.16). Assume V obeys (1.2) through (1.7) as well as (1.13) and (1.14). There exists C > 0 so that for all $\lambda \in \mathbb{C}$ with $0 < |\operatorname{Im} \lambda| \le 1$, and all $j_1, j_2 \in \{0, 1\}$ with $j_1 + j_2 \le 1$,

$$\left\|\frac{d}{d\lambda}\langle x\rangle^{-s}\lambda^{j_1}\nabla^{j_2}(P-\lambda^2)^{-1}\langle x\rangle^{-s}\right\|_{L^2(\mathbb{R}^n)\to L^2(\mathbb{R}^n)} \le C.$$
(3.6)

Proof. Without loss of generality, we take s sufficiently close to (n+3)/4 when $n \neq 8$, or sufficiently close to 3 when n = 8, so that by (1.14) we may fix s' > 1/2 so that $s + s' < \delta$.

We begin from the resolvent identity

$$(P - \lambda^2)^{-1} \langle x \rangle^{-s} (1 + K(\lambda)) = R_0(\lambda) \langle x \rangle^{-s},$$
(3.7)

where $K(\lambda) := V(x) \langle x \rangle^{s+s'} \langle x \rangle^{-s'} R_0(\lambda) \langle x \rangle^{-s}$ and $R_0(\lambda) := (-\Delta - \lambda^2)^{-1}$. It is well known that $\langle x \rangle^{-s'} R_0(\lambda) \langle x \rangle^{-s} : L^2(\mathbb{R}^n) \to H^2(\mathbb{R}^n)$ has a continuous extension from

It is well known that $\langle x \rangle^{-s} R_0(\lambda) \langle x \rangle^{-s} : L^2(\mathbb{R}^n) \to H^2(\mathbb{R}^n)$ has a continuous extension from either half-plane ($\pm \operatorname{Im} \lambda > 0$) to \mathbb{R} [GiMo74, Proposition 2.4]. Let us denote this extension by $R_{0,s',s}^{\pm}(\lambda)$ and put $K^{\pm}(\lambda) = V(x) \langle x \rangle^{s+s'} R_{0,s',s}^{\pm}(\lambda)$

We now show that $K^{\pm}(\lambda)$ is a compact operator $L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n)$. To see this, observe that we may write $K^{\pm}(\lambda)$ as the sum

$$K^{\pm}(\lambda) = (\chi \langle x \rangle^{s+s'} V) R^{\pm}_{0,s',s}(\lambda) + ((1-\chi)V \langle x \rangle^{\delta}) \langle x \rangle^{s+s'-\delta} R^{\pm}_{0,s',s}(\lambda)$$

where $\chi \in C_0^{\infty}(\mathbb{R}^n; [0, 1])$ is identically one near the origin in \mathbb{R}^n and supported in B(0, 1). The second operator on the right side is compact by [DyZw19, Theorem B.4]). The first operator on the right side is compact as follows: it may be viewed as the composition of bounded

 $R_{0,s',s}^{\pm}(\lambda): L^2(\mathbb{R}^n) \to H^2(\mathbb{R}^n)$ followed by multiplication by $\chi\langle x \rangle^{s+s'} V$. Due to (1.3) and Lemma E.1, we have $\|\chi\langle x \rangle^{s+s'} V u\|_{L^2(\mathbb{R}^n)} \leq C \|u\|_{H^1(B(0,1))}$ for some C > 0 and all $u \in H^2(\mathbb{R}^n)$. By the Kondrachov embedding theorem the inclusion $H^2(B(0,1)) \to H^1(B(0,1))$ is compact. So compactness of $(\chi\langle x \rangle^{s+s'} V) R_{0,s',s}^{\pm}(\lambda)$ holds as desired.

We claim further that $1 + K^{\pm}(\lambda)$ is invertible $L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n)$ for all λ with $\pm \operatorname{Im} \lambda \geq 0$. By compactness of $K^{\pm}(\lambda)$ and the Fredholm alternative [ReSi80, Theorem VI.14], we have that $1 + K^{\pm}(\lambda)$ is invertible if we can show $(1 + K^{\pm}(\lambda))g = 0$ implies g = 0. To this end, put $u := \langle x \rangle^{s'} R^{\pm}_{0,s',s}(\lambda)g$, which belongs to $\langle x \rangle^{s'} H^2(\mathbb{R}^n)$. If we can show u = 0, then in fact g = 0. This is because $(-\Delta - \lambda^2)\langle x \rangle^{-s'}u = \langle x \rangle^{-s}g$ in the distributional sense. Now let us show u = 0. If $\lambda^2 \in \mathbb{C} \setminus [0, \infty)$ (so that $K^{\pm}(\lambda) = K(\lambda)$), this follows immediately from

Now let us show u = 0. If $\lambda^2 \in \mathbb{C} \setminus [0, \infty)$ (so that $K^{\pm}(\lambda) = K(\lambda)$), this follows immediately from $(P - \lambda^2)u = \langle x \rangle^{-s}g + VR_0(\lambda)\langle x \rangle^{-s}g = \langle x \rangle^{-s}(1 + K(\lambda))g = 0$. If $\lambda^2 \in [0, \infty)$, the idea is the same, but we incorporate a limiting step. Set $u_{\pm,\varepsilon} = (-\Delta - \lambda^2 \pm i\varepsilon)^{-1}\langle x \rangle^{-s}g$. We have that [GiMo74, Proposition 2.4] implies that $\langle x \rangle^{-s'}u_{\pm,\varepsilon}$ converges to $\langle x \rangle^{-s'}u$ in $H^2(\mathbb{R}^n)$ as $\varepsilon \to 0^+$. We also have that

$$u_{\pm,\varepsilon} = (-\Delta - \lambda^2 \pm i\varepsilon)^{-1} \langle x \rangle^{-s} g$$

= $(P - \lambda^2 \pm i\varepsilon)^{-1} (P - \lambda^2 \pm i\varepsilon) (-\Delta - \lambda^2 \pm i\varepsilon)^{-1} \langle x \rangle^{-s} g$
= $(P - \lambda^2 \pm i\varepsilon)^{-1} \langle x \rangle^{-s} (I + V \langle x \rangle^s (-\Delta - \lambda^2 \pm i\varepsilon)^{-1} \langle x \rangle^{-s}) g$

Therefore, by (1.9),

$$\begin{aligned} \|\langle x \rangle^{-s_1} u\|_{L^2} &= \lim_{\varepsilon \to 0^+} \|\langle x \rangle^{-s_1} u_{\pm,\varepsilon} \|_{L^2} \\ &\leq C \lim_{\varepsilon \to 0^+} \|(I + V \langle x \rangle^s (-\Delta - \lambda^2 \pm i\varepsilon)^{-1} \langle x \rangle^{-s}) g\|_{L^2} \\ &= \|(1 + K^{\pm}(\lambda))g\|_{L^2} = 0. \end{aligned}$$

Thus we have demonstrated that $I+K^{\pm}(\lambda)$ is invertible for $\pm \operatorname{Im} \lambda \geq 0$. As $\lambda \to \infty$, $||K(\lambda)||_{L^2 \to L^2} \to 0$ thanks to (3.1), hence we can compute $(I + K^{\pm}(\lambda))^{-1}$ by a Neumann series, thanks to (3.1). Therefore

$$\|(I + K^{\pm}(\lambda))^{-1}\|_{L^2 \to L^2} \le C.$$
(3.8)

Now for $0 < |\operatorname{Im} \lambda| \le 1$ take the derivative of (3.7) with respect to λ ,

$$\left(\frac{d}{d\lambda}\langle x\rangle^{-s}\lambda^{j_1}\nabla^{j_2}(P-\lambda^2)^{-1}\langle x\rangle^{-s}\right)(I+K(\lambda)) = \frac{d}{d\lambda}\langle x\rangle^{-s}\lambda^{j_1}\nabla^{j_2}R_0(\lambda)\langle x\rangle^{-s} - 2\langle x\rangle^{-s}\lambda^{j_1}\nabla^{j_2}(P-\lambda^2)^{-1}\langle x\rangle^{-s'}V\langle x\rangle^{s+s'}\lambda\langle x\rangle^{-s}(-\Delta-\lambda^2)^{-2}\langle x\rangle^{-s},$$
(3.9)

where we used

$$\frac{d}{d\lambda}\langle x\rangle^{-s'}\nabla^j R_0(\lambda)\langle x\rangle^{-s} = 2\lambda\langle x\rangle^{-s'}\nabla^j (-\Delta - \lambda^2)^{-2}\langle x\rangle^{-s}, \qquad j \in \{0, 1\}.$$
(3.10)

The operator norm $L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n)$ of the term in the second line of (3.9) is bounded above by a constant due to (3.5) and (3.10). As for the third line, $\|\langle x \rangle^{-s} \lambda^{j_1} \nabla^{j_2} (P - \lambda^2)^{-1} \langle x \rangle^{-s'} \|_{L^2 \to L^2} \leq C$ by (3.1). Moreover

$$\begin{aligned} \|V\langle x\rangle^{s+s'}\lambda\langle x\rangle^{-s}(-\Delta-\lambda^2)^{-2}\langle x\rangle^{-s}\|_{L^2\to L^2} \\ &\leq C\|\lambda\langle x\rangle^{-s}(-\Delta-\lambda^2)^{-2}\langle x\rangle^{-s}\|_{H^1\to L^2} \end{aligned}$$

since multiplication by $V\langle x \rangle^{s+s'}$ is a bounded operator $H^1(\mathbb{R}^n) \to L^2(\mathbb{R}^n)$ (see (1.3) and Lemma E.1). Finally, because $\|\lambda\langle x \rangle^{-s}(-\Delta - \lambda^2)^{-2}\langle x \rangle^{-s}\|_{H^1 \to L^2} \leq C$ by (3.5), the proof of (3.6) is complete.

4. Proof of Theorem 1.3

In this section we prove Theorem 1.3 by combining the resolvent bounds of the previous section with an argument appearing in [Vo04a, Section 3]. As before we use the notation $P = -\Delta + V$: $L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n), n \ge 3$, where V obeys (1.2) through (1.7) along with (1.13) and (1.14).

In several steps below, we use that for all $0 \leq \alpha \leq 1$, there exists C > 0 so that for any $f \in H^1(\mathbb{R}^n)$,

$$\|V^{\alpha}f\|_{L^{2}}^{2} \leq C(\|\nabla f\|_{L^{2}}^{2} + \|f\|_{L^{2}}^{2}) \leq C\|\nabla f\|_{L^{2}}^{2}.$$
(4.1)

The first inequality follows from (1.3), (1.13) and Lemma E.1, while the second follows from the Poincaré inequality (as we work in dimension $n \ge 3$).

Given s > 0 and u as in (1.15) solving the wave equation (1.12), with compactly supported initial conditions $u(0, x) = u_0(x) \in H^1(\mathbb{R}^n)$, $\partial_t u(0, x) = u_1(x) \in L^2(\mathbb{R}^n)$, define

$$E_s(t) := \int_{\mathbb{R}^n} \langle x \rangle^{-2s} (|\partial_t u(t,x)|^2 + |\nabla u(t,x)|^2 + |u(t,x)|^2) dx,$$
$$E(0) := \|\nabla u_0\|_{L^2}^2 + \|u\|_{L^2}^2.$$

Lemma 4.1. If s > 1/2 and V satisfies (1.2) through (1.7), there exists C > 0 so that

$$\int_0^\infty E_s(\tau)d\tau \le CE(0). \tag{4.2}$$

If in addition s satisfies (1.16) and V (1.13) and (1.14), there exists C > 0 so that for $t \ge 1$,

$$\int_{t}^{\infty} E_s(\tau) d\tau \le C t^{-2} E(0).$$
(4.3)

Proof. Choose $\phi \in C^{\infty}(\mathbb{R}), \phi \ge 0, \phi(t) = 0$ near $(-\infty, 1/2], \phi(t) = 1$ near $[1, \infty)$. Since $(\partial_t^2 + P)u = 0$, where $P = -\Delta + V$, It holds that

$$(\partial_t^2 + P)\phi u = (\phi'' + 2\phi'\partial_t)u := v(t).$$
(4.4)

Thus, by Duhamel's formula for the solution to an inhomogeneous wave equation with zero initial conditions,

$$\phi u(t) = \int_0^t \frac{\sin(t-\tau)\sqrt{P}}{\sqrt{P}} v(\tau) d\tau.$$

On the other hand,

$$(P - (\lambda - i\varepsilon)^2)^{-1} = \int_0^\infty e^{-it(\lambda - i\varepsilon)} \frac{\sin(t\sqrt{P})}{\sqrt{P}} dt, \qquad \varepsilon > 0.$$

It follows from the last two identities that the Fourier transform ϕu of ϕu satisfies

$$\widehat{\phi u}(\lambda - i\varepsilon) := \int_{-\infty}^{\infty} e^{-it(\lambda - i\varepsilon)} \phi(t) u(\cdot, t) dt = (P - (\lambda - i\varepsilon)^2)^{-1} \widehat{v}(\lambda - i\varepsilon).$$
(4.5)

By finite propagation speed for the wave equation, $\operatorname{supp}_x v(t)$, and thus also $\operatorname{supp}_x \hat{v}(\lambda)$, is contained in a compact set independent of t. Choose $\eta \in C_0^{\infty}(\mathbb{R}^n)$ such that $\eta = 1$ near $\operatorname{supp}_x v(t)$ for all $t \in \mathbb{R}$. By (4.5),

$$\langle x \rangle^{-s} \widehat{\phi u}(\lambda - i\varepsilon) = \langle x \rangle^{-s} (P - (\lambda - i\varepsilon)^2)^{-1} \eta \widehat{v}(\lambda - i\varepsilon),$$

$$\langle x \rangle^{-s} \widehat{\partial_t(\phi u)}(\lambda - i\varepsilon) = \langle x \rangle^{-s} (\lambda - i\varepsilon) (P - (\lambda - i\varepsilon)^2)^{-1} \eta \widehat{v}(\lambda - i\varepsilon),$$

$$\langle x \rangle^{-s} \nabla \widehat{\phi u}(\lambda - i\varepsilon) = \langle x \rangle^{-s} (\lambda - i\varepsilon) \nabla (P - (\lambda - i\varepsilon)^2)^{-1} \eta \widehat{v}(\lambda - i\varepsilon).$$

Therefore, by (3.1), for s > 1/2 and V obeying (1.2) through (1.7), there is C > 0 independent of λ and ε , so that for all $\lambda \in \mathbb{R}$, $0 < \varepsilon \leq 1$, we have

$$\begin{aligned} \left\| \frac{d^{k}}{d\lambda^{k}} \langle x \rangle^{-s} \widehat{\partial_{t}(\phi u)}(\lambda - i\varepsilon) \right\|_{L^{2}} + \left\| \frac{d^{k}}{d\lambda^{k}} \langle x \rangle^{-s} \nabla \widehat{\phi u}(\lambda - i\varepsilon) \right\|_{L^{2}} \\ + \left\| \frac{d^{k}}{d\lambda^{k}} \langle x \rangle^{-s} \widehat{\phi u}(\lambda - i\varepsilon) \right\|_{L^{2}} \le C \| \widehat{v}(\lambda - i\varepsilon) \|_{L^{2}} + Ck \| \widehat{tv}(\lambda - i\varepsilon) \|_{L^{2}}. \end{aligned}$$

$$(4.6)$$

for k = 0. If in addition we suppose s satisfies (1.16) and V satisfies (1.13) and (1.14), then by (3.6), (4.6) holds for $k \in \{0, 1\}$. Note when k = 1 we used the product rule and the identity $\frac{d}{d\lambda}\hat{v}(\lambda - i\varepsilon) = -i\hat{t}\hat{v}(\lambda - i\varepsilon)$. Next, by (4.6) and Plancherel's theorem, there exist C_1 , C_2 , $C_2 \in C > 0$ independent of ε so that

Next, by (4.6) and Finicherer's theorem, there exist
$$C_1, C_2, C_3, C > 0$$
 independent of ε so that

$$\int_{-\infty}^{\infty} (\|\langle x \rangle^{-s} \partial_t(\phi u)\|_{L^2}^2 + \|\langle x \rangle^{-s} \nabla(\phi u)\|_{L^2}^2 + \|\langle x \rangle^{-s} \phi u\|_{L^2}^2) e^{-2\varepsilon t} dt$$

$$= C_1 \int_{-\infty}^{\infty} (\|\langle x \rangle^{-s} \widehat{\partial_t(\phi u)}(\lambda - i\varepsilon)\|_{L^2}^2 + \|\langle x \rangle^{-s} \nabla \widehat{\phi u}(\lambda - i\varepsilon)\|_{L^2}^2 + \|\langle x \rangle^{-s} \widehat{\phi u}(\lambda - i\varepsilon)\|_{L^2}^2) d\lambda$$

$$\leq C_2 \int_{-\infty}^{\infty} \|\hat{v}(\lambda - i\varepsilon)\|_{L^2}^2 d\lambda = C_3 \int_{-\infty}^{\infty} \|v(t)\|_{L^2}^2 e^{-2\varepsilon t} dt \leq C \sup_{t \in \mathbb{R}} \|v(t)\|^2.$$

$$(4.7)$$

The last constant C is independent of ε because v(t) has compact support in t, see (4.4). The proof of (4.2) is completed by sending $\varepsilon \to 0$ in (4.7) and observing

$$\|v(t)\|_{L^{2}} \leq C(\|u_{0}\|_{L^{2}} + \|\sqrt{P}u_{0}\|_{L^{2}} + \|u_{1}\|_{L^{2}})$$

$$\leq C(\|\nabla u_{0}\|_{L^{2}} + \|u_{1}\|_{L^{2}}) = C\sqrt{E(0)}.$$
(4.8)

Between lines one and two of (4.8), we used that for any $f \in H^2(\mathbb{R}^n)$ (and thus any $f \in H^1(\mathbb{R}^n)$, since $H^2(\mathbb{R}^n)$ is dense in $H^1(\mathbb{R}^n)$),

$$\|\sqrt{P}f\|_{L^2}^2 = \langle f, Pf \rangle_{L^2} = \|\nabla f\|_{L^2}^2 + \|\sqrt{V}f\|_{L^2}^2 \le C \|\nabla f\|_{L^2}^2,$$

with the second inequality due to (4.1).

To prove (4.3), we again use Plancherel's theorem with (4.6), so that for all $0 < \varepsilon \leq 1$ and $T \geq 1$,

$$T^{2} \int_{T}^{\infty} (\|\langle x \rangle^{-s} \partial_{t}(\phi u)\|_{L^{2}}^{2} + \|\langle x \rangle^{-s} \nabla(\phi u)\|_{L^{2}}^{2} + \|\langle x \rangle^{-s} \phi u\|_{L^{2}}^{2}) e^{-2\varepsilon t} dt$$

$$\leq \int_{-\infty}^{\infty} (\|\langle x \rangle^{-s} t \partial_{t}(\phi u)\|_{L^{2}}^{2} + \|\langle x \rangle^{-s} t \nabla(\phi u)\|_{L^{2}}^{2} + \|\langle x \rangle^{-s} t \phi u\|_{L^{2}}^{2}) e^{-2\varepsilon t} dt$$

$$= C_{1} \int_{-\infty}^{\infty} (\|\frac{d}{d\lambda} \langle x \rangle^{-s} \widehat{\partial_{t}(\phi u)}(\lambda - i\varepsilon)\|_{L^{2}}^{2} + \|\frac{d}{d\lambda} \langle x \rangle^{-s} \nabla \widehat{\phi u}(\lambda - i\varepsilon)\|_{L^{2}}^{2}$$

$$+ \|\frac{d}{d\lambda} \langle x \rangle^{-s} \widehat{\phi u}(\lambda - i\varepsilon)\|_{L^{2}}^{2} + \|\widehat{t} v(\lambda - i\varepsilon)\|_{L^{2}}^{2}) d\lambda$$

$$\leq C_{2} \int_{-\infty}^{\infty} \|\widehat{v}(\lambda - i\varepsilon)\|_{L^{2}}^{2} + \|\widehat{t} v(\lambda - i\varepsilon)\|_{L^{2}}^{2}) d\lambda$$

$$= C_{3} \int_{-\infty}^{\infty} (\|v(t)\|_{L^{2}}^{2} + \|tv(t)\|_{L^{2}}^{2}) e^{-2\varepsilon t} dt \leq C \sup_{t \in \mathbb{R}} \|v(t)\|^{2} \leq C E(0).$$
(4.9)

Once again sending $\varepsilon \to 0^+$ concludes the proof of (4.3).

The local energy decay (1.17) follows from (4.3) and

Lemma 4.2. If s > 1/2 and V satisfies (1.2) through (1.7), there exists C > 0 so that for all $t \ge 1$,

$$E_s(t) \le C \int_t^\infty E_s(\tau) d\tau.$$
(4.10)

Proof. The strategy is the same as that of [Vo04a, Lemma 3.2]. Computing $\frac{d}{dt}E_s(t)$, one finds

$$\frac{d}{dt}E_{s}(t) = -2\operatorname{Re}\int_{\mathbb{R}^{n}}\partial_{r}u(t,x)\overline{\partial_{t}u(t,x)}\partial_{r}\langle x\rangle^{-2s}dx + 2\operatorname{Re}\int_{\mathbb{R}^{n}}(-Vu(t,x)\overline{\partial_{t}u(t,x)} + u(t,x)\overline{\partial_{t}u(t,x)})\langle x\rangle^{-2s}dx.$$
(4.11)

By (4.1),

$$\begin{aligned} \|V\langle x\rangle^{-s}u(t,x)\|_{L^2} &\leq C \|\nabla\langle x\rangle^{-s}u(t,x)\|_{L^2} \\ &\leq C \|\langle x\rangle^{-s}\nabla u(t,x)\|_{L^2} + C \|\langle x\rangle^{-s}u(t,x)\|_{L^2}. \end{aligned}$$

for C > 0 independent of t, and whose precise value may change between lines. Thus we can bound the right side of (4.11) from above by Cauchy-Schwarz,

$$\frac{d}{dt}E_{s}(t) \leq C \|\langle x \rangle^{-s} \partial_{r} u(t,x)\|_{L^{2}} \|\langle x \rangle^{-s} \partial_{t} u(t,x)\|_{L^{2}} + C \|V\langle x \rangle^{-s} u(t,x)\|_{L^{2}} \|\langle x \rangle^{-s} \partial_{t} u(t,x)\|_{L^{2}}
+ C \|\langle x \rangle^{-s} u(t,x)\|_{L^{2}} \|\langle x \rangle^{-s} \partial_{t} u(t,x)\|_{L^{2}} \leq C E_{s}(t).$$

We then have, for all $T > t \ge 1$,

$$E_s(t) \le E_s(T) + C_s \int_t^T E_s(\tau) d\tau.$$
(4.12)

From (4.2), we also have a sequence $T_j \to \infty$ so that $\lim_{T_j\to\infty} E_s(T_j) = 0$. So setting $T = T_j$ in (4.12) and sending $T_j \to \infty$ completes the proof of (4.10).

Appendix A. Phragmén-Lindelöf Theorem

In this appendix we recall the Phragmén Lindelöf Theorem. Let f(z) be a holomorphic function in a domain D of the complex plane with boundary Γ . We say that f(z) does not exceed a number $M \ge 0$ in modulus at a boundary point $\zeta \in \Gamma$ if $\limsup_{z \to \zeta, z \in D} |f(z)| \le M$.

Theorem A.1 (Phragmén Lindelöf Theorem [EM]). Suppose $E \subseteq \Gamma$, and f analytic on D does not exceed M in modulus at any point of $\Gamma \setminus E$. Suppose also there is a function g(z) with the following properties:

(1) g(z) is analytic in D,

- (2) |g(z)| < 1 in D,
- (3) $g(z) \neq 0$ in D,

(4) For every $\sigma > 0$, the function $|g(z)|^{\sigma}|f(z)|$ does not exceed M is modulus at any $\zeta \in E$. Under these conditions, $|f(z)| \leq M$ everywhere in D.

Appendix B. Density argument: proof of (2.15) and (2.21)

In this appendix, we prove (2.15) and (2.21) as a consequence of

Lemma B.1. Fix $h, s_1 > 0, 0 < s_2 < 1$, and $z \in \mathbb{C} \setminus [0, \infty)$. Let P(h) be as in (1.1) with $V : \mathbb{R}^n \to \mathbb{R}$ obeying (1.3) and (1.4) (so that P(h) is self-adjoint with respect to the domain $H^2(\mathbb{R}^n)$). Suppose there exists C > 0 so that for all $v \in C_0^{\infty}(\mathbb{R}^n)$,

$$\|\langle x \rangle^{-s_1} v\|_{L^2}^2 \le C \|\langle x \rangle^{s_2} (P(h) - z) v\|_{L^2}^2.$$
(B.1)

Then

$$\|\langle x \rangle^{-s_1} (P(h) - z)^{-1} \langle x \rangle^{-s_2} \|_{L^2 \to L^2} \le C.$$
(B.2)

Proof. The operator

$$[P(h), \langle x \rangle^{s_2}] \langle x \rangle^{-s_2} = \left(-h^2 (\Delta \langle x \rangle^{s_2}) - 2h^2 (\nabla \langle x \rangle^{s_2}) \cdot \nabla \right) \langle x \rangle^{-s_2}$$

is bounded $H^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n)$. So, for $v \in H^2(\mathbb{R}^n)$ such that $\langle x \rangle^{s_2} v \in H^2(\mathbb{R}^n)$,

$$\begin{aligned} \|\langle x \rangle^{s_2} (P(h) - z) v\|_{L^2} &\leq \|(P(h) - z) \langle x \rangle^{s_2} v\|_{L^2} + \|[P(h), \langle x \rangle^{s_2}] \langle x \rangle^{-s_2} \langle x \rangle^{s_2} v\|_{L^2} \\ &\leq C_{z,h} \|\langle x \rangle^{s_2} v\|_{H^2}, \end{aligned}$$
(B.3)

for some constant $C_{z,h} > 0$ depending on z and h.

Given $f \in L^2(\mathbb{R}^n)$, the function $u = \langle x \rangle^{s_2} (P(h) - z)^{-1} \langle x \rangle^{-s_2} f \in H^2(\mathbb{R}^n)$ because

$$u = (P(h) - z)^{-1}(f + w), \qquad w = [P(h), \langle x \rangle^{s_2}]u,$$

with $[P(h), \langle x \rangle^{s_2}]$ being bounded $L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n)$ since $s_2 < 1$.

Now, choose a sequence $v_k \in C_0^{\infty}$ such that $v_k \to \langle x \rangle^{s_2} (P(h) - z)^{-1} \langle x \rangle^{-s_2} f$ in $H^2(\mathbb{R}^n)$. Define $\tilde{v}_k := \langle x \rangle^{-s_2} v_k$. Then, as $k \to \infty$,

$$\begin{aligned} \|\langle x \rangle^{-s_1} \tilde{v}_k - \langle x \rangle^{-s_1} (P(h) - z)^{-1} \langle x \rangle^{-s_2} f \|_{L^2} \\ &\leq \|v_k - \langle x \rangle^{s_2} (P(h) - z)^{-1} \langle x \rangle^{-s_2} f \|_{H^2} \to 0. \end{aligned}$$

Also, applying (B.3),

$$\|\langle x \rangle^{s_2} (P(h) - z) \tilde{v}_k - f\|_{L^2} \le C_{z,h} \|v_k - \langle x \rangle^{s_2} (P(h) - E \pm i\varepsilon)^{-1} \langle x \rangle^{-s_2} f\|_{H^2} \to 0.$$

Thus (B.2) follows by replacing v by \tilde{v}_k in (B.1) and sending $k \to \infty$.

Appendix C. Justification of Remark 1.2

In the setting of Theorem 1.1, consider the case of V = 0 and n = 3. In that scenario the integral kernel of $(P(h) - z)^{-1} = (-h^2\Delta - z)^{-1}$ with $z \in \mathbb{C} \setminus [0, \infty)$ is given by

$$R_0(x, y, z) := h^{-2} \frac{e^{i\frac{\sqrt{z}}{h}|x-y|}}{4\pi|x-y|}, \qquad \text{Im } \sqrt{z} > 0.$$

We recall why having a bound like (1.8) on $\langle \cdot \rangle^{-s_1} (-h^2 \Delta - z)^{-1} \langle \cdot \rangle^{-s_2} : L^2(\mathbb{R}^3) \to L^2(\mathbb{R}^3)$ requires $s_1, s_2 > 1/2$.

Since the norm of an operator and its adjoint coincide, it suffices to show $s_1 > 1/2$ is necessary. Use \sqrt{z} of the form $\sqrt{z} = E + i\varepsilon$ for E > 0 fixed and $\varepsilon > 0$ tending to zero. Then, as calculated in the proof of [DyZw19, Theorem 3.5], for $f \in C_0^{\infty}(\mathbb{R}^3)$,

$$\langle x \rangle^{-s_1} \int_{\mathbb{R}^3} R_0(x, y, z) f(y) dy$$

$$= h^{-2} \frac{\langle x \rangle^{-s_1}}{4\pi |x|} e^{\frac{i}{h} (E+i\varepsilon)|x|} (\hat{f}(\frac{E}{h} \frac{x}{|x|}) + o(1)) + O(|x|^{-2}), \quad \text{as } \varepsilon \to 0^+ \text{ and } |x| \to \infty.$$

$$(C.1)$$

If f is chosen so that $|\hat{f}| > c$ for some c > 0 on $\{|x| = E/h\}$, (C.1) and $s_1 \le 1/2$ imply $\|\langle x \rangle^{-s_1} \int_{\mathbb{R}^3} R_0(x, y, z) f(y) dy\|_{L^2} \to \infty$ as $\varepsilon \to 0^+$.

Next, supposing $s_1, s_2 > 1/2$, we show why a bound like (1.9) on $\langle \cdot \rangle^{-s_1} (-h^2 \Delta - z)^{-1} \langle \cdot \rangle^{-s_2} : L^2(\mathbb{R}^3) \to L^2(\mathbb{R}^3)$ requires additionally that $s_1 + s_2 \geq 2$, which is nearly the condition we impose

for (1.9). Using $\sqrt{z} = i\varepsilon$ for $\varepsilon > 0$ tending to zero, and $f_{\eta}(y) = \langle y \rangle^{-\eta - \frac{3}{2}}$, $\eta > 0$, we see as in [BoHa10, Proof of Remark 2] that

$$\begin{split} \langle x \rangle^{-s_1} \int_{\mathbb{R}^3} &R_0(x,y,z) \langle y \rangle^{-s_2} f(y) \\ &= h^{-2} \langle x \rangle^{-s_1} \int_{\mathbb{R}^3} \frac{e^{-\frac{\varepsilon}{h}|x-y|}}{4\pi |x-y|} \langle y \rangle^{-s_2} f(y) dy \\ &\gtrsim h^{-2} e^{-\frac{3\varepsilon}{2h}|x|} \langle x \rangle^{-s_1-1} \int_{|y| \le \frac{|x|}{2}} \langle y \rangle^{-s_2-\eta-\frac{3}{2}} dy \gtrsim h^{-2} e^{-\frac{3\varepsilon}{2h}|x|} \langle x \rangle^{-s_1-s_2-\eta+\frac{1}{2}}, \end{split}$$

where the implicit constants indicated by \gtrsim are independent of ε and η . First sending $\varepsilon \to 0^+$ gives $s_1 + s_2 \ge 2 - \eta$, but since $\eta > 0$ is arbitrary, we in turn get $s_1 + s_2 \ge 2$.

To see that the $O(|z|^{-\frac{1}{2}}h^{-1})$ -dependence of the right side of (1.8) is optimal, consider the function $u = e^{i\frac{\sqrt{z}}{h}x_1}\chi$ for nontrivial $\chi \in C_0^{\infty}(\mathbb{R}^3; [0, 1])$. We have

$$\langle x \rangle^s (-h^2 \Delta - z) u = -i \sqrt{z} h \langle x \rangle^s \partial_{x_1} \chi - h^2 e^{i \frac{\sqrt{z}}{h} x_1} \langle x \rangle^s \Delta \chi =: f,$$

whence $\langle \cdot \rangle^{-s} (-h^2 \Delta - z)^{-1} \langle \cdot \rangle^{-s} f = \langle \cdot \rangle^{-s} u$ and thus, as $h \to 0$,

$$\frac{\|\langle \cdot \rangle^{-s} (-h^2 \Delta - z)^{-1} \langle \cdot \rangle^{-s} f\|_{L^2}}{\|f\|_{L^2}} = \frac{\|\langle \cdot \rangle^{-s} u\|_{L^2}}{\|f\|_{L^2}} \gtrsim |z|^{-\frac{1}{2}} h^{-1}.$$

Finally, we argue why the $O(h^{-2})$ -dependence of the right side of (1.9) is sharp. As noted before [GiMo74, Proposition 2.4] gives that $R_{0,s_1,s_2}(\lambda) = \langle \cdot \rangle^{-s_1} (-h^2 \Delta - \lambda^2)^{-1} \langle \cdot \rangle^{-s_2}$ ($s_1, s_2 > 1/2, s_1 + s_2 > 2$), has extends continuously from $\text{Im } \lambda > 0$ to \mathbb{R} in the space of bounded opeartors $\mathbb{R}^3 \to \mathbb{R}^3$. In this case, we have,

$$h^{-2} \|\langle x \rangle^{-s_1} \int_{\mathbb{R}^3} \frac{1}{4\pi |x-y|} \langle y \rangle^{-s_2} dy \|_{L^2 \to L^2} = \lim_{\varepsilon \to 0} \|R_{0,s_1,s_2}(i\varepsilon)\|_{L^2 \to L^2}$$

Appendix D. Proof of Lemma 3.2

In this appendix we prove Lemma 3.2. The proof proceeds in two steps. First we treat the case $|\lambda| \ge 1$, followed by $|\lambda| \le 1$.

Proof of lemma 3.2. Initially we take j = 0 in (3.5), so may assume without loss of generality that $\operatorname{Im} \lambda > 0$. We treat the j = 1 case at the end of the proof. Observe that $\frac{d}{d\lambda} \langle x \rangle^{-s} (-\Delta - \lambda^2)^{-1} \langle x \rangle^{-s} = 2\lambda \langle x \rangle^{-s} (-\Delta - \lambda^2)^{-2} \langle x \rangle^{-s}$, so we can bound the $L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n)$ norm of either quantity. If $|\lambda| > 1$, begin from

$$-\lambda^{2} \langle x \rangle^{-s} (-\Delta - \lambda^{2})^{-2} \langle x \rangle^{-s} = -\frac{1}{2} \langle x \rangle^{-s} (-\Delta - \lambda^{2})^{-1} (-2\Delta) (-\Delta - \lambda^{2})^{-1} \langle x \rangle^{-s} + \langle x \rangle^{-s} (-\Delta - \lambda^{2})^{-1} \langle x \rangle^{-s}.$$
(D.1)

By (3.1), the $L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n)$ norm of the second line of (D.1) is bounded by $C(1+|\operatorname{Re} \lambda|)^{-1}$. So it suffices to investigate the term on the right side of the first line of (D.1). For notational brevity, put $R_0(\lambda) := (-\Delta - \lambda^2)^{-1}$. We show that for all $f \in C_0^{\infty}(\mathbb{R}^n)$,

$$\langle x \rangle^{-s} R_0(\lambda) (-2\Delta) R_0(\lambda) \langle x \rangle^{-s} f = -\langle x \rangle^{-s} R_0(\lambda) \partial_r (r \langle x \rangle^{-s} f) + \langle x \rangle^{-s} R_0(\lambda) \langle x \rangle^{-s} f + \langle x \rangle^{-s} r \partial_r R_0(\lambda) \langle x \rangle^{-s} f$$
 (D.2)

Since s > 3/2 by (1.16), the $L^2(\mathbb{R}^n)$ -norm of the right side of (D.2) is bounded by $C||f||_{L^2}$, thanks to (3.1). So it remains to show (D.2).

Recall the well known formula for the Laplacian in polar coordinates,

$$\Delta = \partial_r^2 + (n-1)r^{-1}\partial_r + r^{-2}\Delta_{\mathbb{S}^{n-1}},$$

which implies the commutator identity

$$[r\partial_r, \Delta] := r\partial_r(\Delta) - \Delta(r\partial_r) = -2\Delta.$$
(D.3)

Fix $f \in C_0^{\infty}(\mathbb{R}^n)$, $g \in L^2(\mathbb{R}^n)$, and put $u := R_0(\lambda) \langle x \rangle^{-s} f \in H^2(\mathbb{R}^n)$. Let $\{u_k\}_{k=1}^{\infty} \subseteq C_0^{\infty}(\mathbb{R}^n)$ be a sequence converging to u in $H^2(\mathbb{R}^n)$. Starting from the left side of (D.2) and applying (D.3),

$$\langle g, \langle x \rangle^{-s} R_0(\lambda)(-2\Delta) R_0(\lambda) \langle x \rangle^{-s} f \rangle_{L^2} = \lim_{k \to \infty} \langle g, \langle x \rangle^{-s} R_0(\lambda) [r \partial_r, \Delta] u_k \rangle_{L^2} = \langle g, \langle x \rangle^{-s} r \partial_r u \rangle_{L^2} - \lim_{k \to \infty} \langle g, \langle x \rangle^{-s} R_0(\lambda) r \partial_r (-\Delta - \lambda^2) u_k \rangle_{L^2}.$$
(D.4)

The purpose of the following calculations is to show that the last line of (D.4) equals $-\langle g, \langle x \rangle^{-s} R_0(\lambda) r \partial_r \langle x \rangle^{-s} f \rangle_{L^2}$. First, for any $v \in L^2(\mathbb{R}^n)$, $rR_0(\lambda) \langle x \rangle^{-1} v \in H^1(\mathbb{R}^n)$. This holds because, if we put $w := \langle x \rangle R_0(\lambda) \langle x \rangle^{-1} v$, then $rR_0(\lambda) \langle x \rangle^{-1} v = r \langle x \rangle^{-1} w$ and

$$(-\Delta - \lambda^2)w = [-\Delta, \langle x \rangle] R_0(\lambda) \langle x \rangle^{-1} v + v \implies w = R_0(\lambda) ([-\Delta, \langle x \rangle] R_0(\lambda) \langle x \rangle^{-1} v + v) \in H^2(\mathbb{R}).$$

Furthermore, for any $w, v \in C_0^{\infty}(\mathbb{R}^n)$,

$$\langle w, \partial_r v \rangle_{L^2} = \langle \partial_r^* w, v \rangle_{L^2} := (1-n) \langle r^{-1} w, v \rangle_{L^2} - \langle \partial_r w, v \rangle_{L^2}$$

Therefore, by the density of $C_0^{\infty}(\mathbb{R}^n)$ in $H^1(\mathbb{R}^n)$, and setting $\tilde{u}_k = (-\Delta - \lambda)u_k$, we get

$$\lim_{k \to \infty} \langle g, \langle x \rangle^{-s} R_0(\lambda)(r\partial_r) \tilde{u}_k \rangle_{L^2} = \lim_{k \to \infty} \langle (\partial_r)^* r R_0(\overline{\lambda}) \langle x \rangle^{-s} g, \tilde{u}_k \rangle_{L^2} = \langle (\partial_r)^* r R_0(\overline{\lambda}) \langle x \rangle^{-s} g, \langle x \rangle^{-s} f \rangle_{L^2} = \langle g, \langle x \rangle^{-s} R_0(\lambda) r \partial_r \langle x \rangle^{-s} f \rangle_{L^2}.$$
(D.5)

as desired. Taken together, (D.4) and (D.5) confirm (D.2).

Now we turn to the case $|\lambda| \leq 1$, and utilize the integral kernel of the free resolvent, which is given by [JeNe01, Section 3],

$$(-\Delta - \lambda^2)^{-1}(|x - y|) = \frac{1}{2\pi} \left(\frac{-i\lambda}{2\pi |x - y|} \right)^{\frac{n}{2} - 1} K_{\frac{n}{2} - 1}(-i\lambda |x - y|), \quad \text{Im } \lambda > 0, \quad (D.6)$$

where $K_{\nu}(z)$ is the Macdonald function of order ν [DLMF, 10.27.4, 10.27.5]. Now, if n = 3, then the integral kernel of $\langle x \rangle^{-s} \frac{d}{d\lambda} (-\Delta - \lambda^2)^{-1} \langle x \rangle^{-s}$ is given by $i(4\pi)^{-1} \langle x \rangle^{-s} e^{i\lambda|x-y|} \langle y \rangle^{-s}$, which has Hilbert-Schmidt norm bounded uniformly in $|\lambda| \leq 1$ provided s > 3/2. Moving on to $n \geq 4$, by [DLMF, 10.29.2],

$$\frac{d}{d\lambda} \left(\frac{-i\lambda}{2\pi|x-y|} \right)^{\frac{n}{2}-1} K_{\frac{n}{2}-1}(-i\lambda|x-y|) = \frac{-(-i)^{\frac{n}{2}}\lambda^{\frac{n}{2}-1}}{(2\pi)^{\frac{n}{2}-1}|x-y|^{\frac{n}{2}-2}} K_{\frac{n}{2}-2}(-i\lambda|x-y|).$$
(D.7)

The Macdonald function satisfies [DLMF, 10.25.3, 10.39.2, 10.30.2]

$$|K_{\nu}(z)| \leq \begin{cases} C|z|^{-\nu} & 0 < |z| \le 1, \, \nu > 0, \\ C|\ln|z|| & 0 < |z| \le 1, \, \nu = 0, \\ C|z|^{-1/2} & |z| \ge 1, \, \operatorname{Re} z \ge 0, \end{cases}$$
(D.8)

for C > 0 a constant independent of z. Therefore, for C > 0 independent of λ ,

$$\left| \frac{\lambda^{\frac{n}{2}-1} \langle x \rangle^{-s} \langle y \rangle^{-s}}{|x-y|^{\frac{n}{2}-2}} K_{\frac{n}{2}-2}(-i\lambda|x-y|) \right| \\
\leq \begin{cases} C|\lambda| \langle x \rangle^{-s} \langle y \rangle^{-s} |\ln(|\lambda||x-y|)| \mathbf{1}_{\{|\lambda||x-y|\leq 1\}} + C\frac{|\lambda|^{\frac{n}{2}-\frac{3}{2}} \langle x \rangle^{-s} \langle y \rangle^{-s}}{|x-y|^{\frac{n}{2}-\frac{3}{2}}} \mathbf{1}_{\{|\lambda||x-y|>1\}} & n = 4, \\ C\frac{|\lambda| \langle x \rangle^{-s} \langle y \rangle^{-s}}{|x-y|^{n-4}} \mathbf{1}_{\{|\lambda||x-y|\leq 1\}} + C\frac{|\lambda|^{\frac{n}{2}-\frac{3}{2}} \langle x \rangle^{-s} \langle y \rangle^{-s}}{|x-y|^{\frac{n}{2}-\frac{3}{2}}} \mathbf{1}_{\{|\lambda||x-y|>1\}} & n > 4. \end{cases}$$
(D.9)

As preparation for the conclusions we draw in the next paragraph, we observe that the first term in line three of (D.9) has the bound

$$\frac{|\lambda|\langle x\rangle^{-s}\langle y\rangle^{-s}}{|x-y|^{n-4}}\mathbf{1}_{\{|\lambda||x-y|\leq 1\}} = \frac{|\lambda||x-y|^{\alpha}\langle x\rangle^{-s}\langle y\rangle^{-s}}{|x-y|^{n-4+\alpha}}\mathbf{1}_{\{|\lambda||x-y|\leq 1\}} \le \frac{\langle x\rangle^{-s}\langle y\rangle^{-s}}{|x-y|^{n-4+\alpha}}\mathbf{1}_{\{|\lambda||x-y|\leq 1\}}.$$
(D.10)

for any $0 < \alpha \leq 1$.

In what follows we make repeated use of Lemma E.3. In (D.9), the second term in line two and the second term in line three are are uniformly bounded in Hilbert-Schmidt norm for $|\lambda| \leq 1$, provided s > (n+3)/4. This also holds for

the first term in line two if s > 3/2,

the first term in line three if s > 3/2 and n = 5 (by (D.10) with $\alpha = 1$),

the first term in line three if s > 3/2 and n = 6 (by (D.10) with α so that $s > 2 - (\alpha/2)$, and

the first term in line three if s > 7/4 and n = 7 (by (D.10) with α so that $s > 2 - (\alpha/2)$).

Finally, if n > 8, the first term in line three is uniformly bounded $L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n)$ for $|\lambda| \leq 1$ provided s > 3. This is due to the Schur test, see Lemma E.2.

We finish by resolving the j = 1 case for (3.5). By (3.5) in the j = 0 case, and by (3.3), we need to show $\|\lambda\langle x\rangle^{-s}(-\Delta - \lambda^2)^{-2}\langle x\rangle^{-s}f\|_{H^2} \leq O(1 + |\lambda|)\|f\|_{L^2}$. According to (E.1) below,

$$\begin{aligned} \|\lambda\langle x\rangle^{-s}(-\Delta-\lambda^2)^{-2}\langle x\rangle^{-s}f\|_{H^2} \\ &\leq C\|\lambda\langle x\rangle^{-s}(-\Delta-\lambda^2)^{-2}\langle x\rangle^{-s}f\|_{L^2} + C\|\lambda\langle x\rangle^{-s}(-\Delta)(-\Delta-\lambda^2)^{-2}\langle x\rangle^{-s}f\|_{L^2} \\ &= C\|f\|_{L^2} + C\|\lambda\langle x\rangle^{-s}(-\Delta)(-\Delta-\lambda^2)^{-2}\langle x\rangle^{-s}f\|_{L^2}. \end{aligned}$$

Then use

$$\langle x \rangle^{-s} (-\Delta) (-\Delta - \lambda^2)^{-2} \langle x \rangle^{-s} f = \langle x \rangle^{-s} (-\Delta - \lambda^2)^{-1} \langle x \rangle^{-s} f + \lambda^2 \langle x \rangle^{-s} (-\Delta - \lambda^2)^{-2} \langle x \rangle^{-s} f,$$

which in combination with (3.1), as well as (3.5) in the j = 0 case yields

$$|\lambda \langle x \rangle^{-s} (-\Delta) (-\Delta - \lambda^2)^{-2} \langle x \rangle^{-s} f \|_{L^2} \le C(1 + |\lambda|) \| f_{L^2},$$

completing the proof.

APPENDIX E. USEFUL LEMMAS

Lemma E.1 ([Fa67, Proposition]). Let $n \ge 3$. Then,

$$||r^{-1}u||_{L^2}^2 \le \left(\frac{2}{n-2}\right)^2 ||\nabla u||_{L^2}^2, \qquad u \in H^1(\mathbb{R}^n).$$

Lemma E.2 (Schur's test [DyZw19, Section A.5]). Suppose that K(x, y) is measurable on $\mathbb{R}^n \times \mathbb{R}^n$ and

$$\sup_{x} \int |K(x,y)| dy, \sup_{y} \int |K(x,y)| dy \le C.$$

Then the linear operator

$$Tf(x) = \int K(x,y)f(y)dy$$

obeys the estimate

$$||Tf||_{L^2} \le C ||f||_{L^2}$$

Lemma E.3 ([Pe24]). The necessary and sufficient conditions for

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \langle x \rangle^{-s} \langle y \rangle^{-t} |x-y|^{-p} dx dy < \infty,$$

are

$$s+p>n, \quad t+p>n, \quad s+p+t>2n, \quad p$$

Lemma E.4. Suppose $T: L^2(\mathbb{R}^n) \to H^2(\mathbb{R}^n)$ is a bounded operator. For any s > 0, there exists C > 0 so that

$$\|\langle x \rangle^{-s} T\|_{L^2 \to H^2} \le C(\|\langle x \rangle^{-s} T\|_{L^2 \to L^2} + \|\langle x \rangle^{-s} \Delta T\|_{L^2 \to L^2}).$$
(E.1)

Proof. Let $f \in L^2(\mathbb{R}^n)$ and put u = Tf. By the first line of (3.3), there exists C > 0, whose precise value may change from line to line, so that

$$\|\langle x \rangle^{-s} \tilde{u}\|_{H^2} \le C \|\langle x \rangle^{-s} \tilde{u}\|_{L^2} + C \|\Delta \langle x \rangle^{-s} \tilde{u}\|_{L^2}, \qquad \tilde{u} \in H^2(\mathbb{R}^n).$$
(E.2)

Then use the second line of (3.3),

$$\begin{split} \|\Delta\langle x\rangle^{-s}u\|_{L^{2}} &\leq \|[\Delta,\langle x\rangle^{-s}]u\|_{L^{2}} + \|\langle x\rangle^{-s}\Delta u\|_{L^{2}} \\ &\leq C\|\langle x\rangle^{-s}u\|_{H^{1}} + \|\langle x\rangle^{-s}\Delta u\|_{L^{2}} \\ &\leq C\gamma^{-1}\|\langle x\rangle^{-s}u\|_{L^{2}} + C\gamma\|\Delta\langle x\rangle^{-s}u\|_{L^{2}}) + \|\langle x\rangle^{-s}\Delta u\|_{L^{2}}, \qquad \gamma > 0. \end{split}$$

Fixing γ small enough yields,

$$\|\Delta \langle x \rangle^{-s} u\|_{L^2} \le C(\|\langle x \rangle^{-s} u\|_{L^2} + \|\langle x \rangle^{-s} \Delta u\|_{L^2}),$$

which in combination with (E.2) implies (E.1).

References

- [BoHa10] J.-F. Bony and D. Häfner. Low frequency resolvent estimates for long range perturbations of the Euclidean Laplacian. Math. Res. Lett. 17(2) (2010), 303–308 16
- [Bo11] J. Bouclet. Low frequency estimates and local energy decay for asymptotically Euclidean Laplacians. Comm. Partial Differential Equations, 36(7) (2011), 1239–1286 4
- [Bu98] N. Burq. Décroissance de l'énergie locale de l'équation des ondes pour le problème extérieur et absence de résonance au voisinage du réel. Acta Math. 180(1) (1998), 1–29 4
- [Bu02] N. Burq, Lower bounds for shape resonances widths of long range Schrödinger operators. Amer. J. Math., 124:4 (2002), pp. 677–735. 4
- [CaVo02] F. Cardoso and G. Vodev. Uniform Estimates of the Resolvent of the Laplace-Beltrami Operator on Infinite Volume Riemannian Manifolds. II. Ann. Henri Poincaré 4(3) (2002), 673–691 2
- [CaV004] F. Cardoso and G. Vodev. High Frequency Resolvent Estimates and Energy Decay of Solutions to the Wave Equation. Canad. Math. Bull., (4)47 (2004), 504–514 4
- [ChIk20] R. Charao and R. Ikehata. A note on decay rates of the local energy for wave equations with Lipschiz wavespeeds. J. Math. Anal. Appl. 483(2) (2020), 1–14 4
- [ChDa21] T. Christiansen and K. Datchev. Resolvent estimates on asymptotically cylindrical manifolds and on the half line. Ann. Sci. Éc. Norm. Supér. 54(4) (2021), 1051–1088 1, 2
- [Da14] K. Datchev. Quantitative limiting absorption principle in the semiclassical limit. Geom. Func. Anal. 24(3) (2014), 740–747 2
- [DGS23] K. Datchev, J. Galkowksi, and J. Shapiro. Semiclassical resolvent bounds for compactly supported radial potentials. J. Funct. Anal. 284(7), paper no. 109835 (2023), 28pp. 2
- [DLMF] NIST Digital Library of Mathematical Functions. http://dlmf.nist.gov/, Release 1.1.0 of 2020-12-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds. 10, 17

- [DyZw19] S. Dyatlov and M. Zworski. Mathematical Theory of Scattering Resonances. Graduate Studies in Mathematics 200. American Mathematical Society, Providence, RI (2019) 10, 15, 18
- [EM] Phragmén-Lindelöf theorem. Encyclopedia of Mathematics.
- URL: http://encyclopediaofmath.org/index.php?title=Phragm%C3%A9n-Lindel%C3%B6f_theorem&oldid=481767, 14
- [Fa67] W. Faris. The product formula for semigroups defined by Friedrichs extensions. Pacific J. Math. 22(1) (1967), 47–70 18
- [DyZw19] S. Dyatlov and M. Zworski. Mathematical Theory of Scattering Resonances. Graduate Studies in Mathematics 200. American Mathematical Society, Providence, RI (2019) 10, 15, 18
- [GaSh22] J. Galkowski and J. Shapiro. Semiclassical resolvent bounds for long range Lipschitz potentials. Int. Math. Res. Not. IMRN 2022(18) (2022), 14134–14150 2
- [GiMo74] J. Ginibre and M. Moulin. Hilbert sapce approach to the quantum mechanical three-body problem. Ann. Inst. H. Poincaré Sect. A (N.S.) 21(2) (1974), 97–145. 10, 11, 16
- [Ik23] R. Ikehata. A note on local energy decay results for wave equations with a potential. Asymptot. Anal. 134(1–2) (2023), 281–295 4
- [JeNe01] A. Jensen and G. Nenciu. A Unified approach to resolvent expansions at thresholds. Rev. Math. Phys. 13(6) (2001), 717–754 17
- [MeSj82] R. B. Melrose and J. Sjöstrand. Singularities of boundary value problems. II Comm. Pure Appl. Math. 35(2) (1982), 129–168 3
- [MeSj78] R. B. Melrose and J. Sjöstrand. Singularities of boundary value problems. II Comm. Pure Appl. Math. 31(5) (1978), 593–617 3
- [LMP63] P. D. Lax, C. S. Morawetz, and R. S. Phillips. Exponential decay of solutions of the wave equation in the exterior of a star-shaped obstacle. Comm. Pure Appl. Math. 16(4) (1962), 477–486 3, 4
- [Mo61] C. S. Morawetz. The decay of solutions of the exterior initial-boundary value problem for the wave equation. Comm. Pure Appl. Math. 14(3) (1961), 561–568 4
- [Mo66] C. S. Morawetz. Exponential decay of solutions of the wave equation. Comm. Pure Appl. Math. 19(4) (1966), 439–444 3
- [Mo75] C. S. Morawetz. Decay for solutions of the exterior problem for the wave equation. Comm. Pure Appl. Math. 28(2) (1975), 229–264–444 3
- [Ne64] E. Nelson. Feynman integrals and the Schrödinger equations. J. Mathematical Phys. 5(3) (1964), 332–343. 2
- [NiZe92] H. Niessen and A. Zettl. Singular Sturm-Liouville Problems: The Friedrichs Extension and Comparison of Eigenvalues. Proc. London Math. Soc. 64(3) (1992), 545–578
- [Ob24] D. Obovu. Resolvent bounds for Lipschitz potentials in dimension two and higher with singularities at the origin. J. Spectr. Theory 14(1) (2024) 163–183 2
- [Pe24] F. Petrov. Show that the kernel $|x y|^{-1}$ on $\mathbb{R}^3 \times \mathbb{R}^3$ is Hilbert Schmidt with respect to a weighted L^2 space, Math Overflow, https://mathoverflow.net/q/478823 19
- [ReSi80] Michael Reed and Barry Simon. Methods of Modern Mathematical Physics I: Functional Analysis, Revised and Enlarged Edition. Academic Press, San Diego (1980) 11
- [ReSi75] M. Reed and B. Simon. Methods of modern mathematical physics II. Fourier Analysis. Academic Press, San Diego, CA (1975)
- [Sh18] J. Shapiro. Local energy decay for Lipschitz wavespeeds. Comm. Partial Differential Equations 43 (5) (2018), 839–858 4
- [Va75] B. R. Vainberg. On the short wave asymptotic behaviour of solutions of stationary problems and the asymptotic behaviour as $t \to \infty$ of solutions of non-stationary problems. *Russ. Math. Surv.* 30(2) (1975), 1–58
- [Vo99] G. Vodev. On the uniform decay of the local energy. Serdica Math. J. 25(3) 1999, 191–206 4
- [Vo04a] G. Vodev. Local energy decay of solutions to the wave equation for short range potentials. Asymptot. Anal. 37(2) (2004), 175–187 3, 4, 10, 12, 14
- [Vo04b] G. Vodev. Local energy decay of solutions to the wave equation for nontrapping metrics. Ark. Mat. 42(2) (2004), 379–397 3
- [Vo04c] G. Vodev. Local energy decay of solutions to the wave equation for nontrapping metrics. Mat. Contemp. 26 (2004), 129–133 3
- [Yu] Yukawa potential. Wikipedia, The Free Encyclopedia,
- https://en.wikipedia.org/w/index.php?title=Yukawa_potential&oldid=1156529154 2
- [Ya10] D. R. Yafaev. Mathematical Scattering Theory, Analytic Theory. Mathematical Surveys and Monographs 158. American Mathematical Society, Providence, RI (2010)
- [Za04] S. Zappacosta. Resolvent estimates and applications to wave equations with potential. Asymptot. Anal. 38(3–4) (2004), 221–239 3

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DAYTON, DAYTON, OH 45469-2316, USA *Email address*: alarrainhubach1@udayton.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DAYTON, DAYTON, OH 45469-2316, USA *Email address*: yli004@udayton.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DAYTON, DAYTON, OH 45469-2316, USA *Email address*: jshapiro1@udayton.edu

UNIVERSITY OF DAYTON RESEARCH INSTITUTE, DAYTON, OH 45469-2316, USA *Email address*: jtiller1@udayton.edu