On the Convergence and Stability of Upside-Down Reinforcement Learning, Goal-Conditioned Supervised Learning, and Online Decision Transformers

Miroslav Štrupl ¹	MIROSLAV.STRUPL@IDSIA.CH
Oleg Szehr ¹	OLEG.SZEHR@IDSIA.CH
Francesco Faccio ^{1,2}	FRANCESCO.FACCIO@IDSIA.CH
Dylan R. Ashley ^{1,2}	DYLAN.ASHLEY@IDSIA.CH
Rupesh Kumar Srivastava	RUPSPACE@GMAIL.COM
Jürgen Schmidhuber ^{1,2,3}	JUERGEN.SCHMIDHUBER@KAUST.EDU.SA

¹ Dalle Molle Institute for Artificial Intelligence (IDSIA) - USI/SUPSI, Lugano, Switzerland

² Center of Excellence for Generative AI, King Abdullah University of Science and Technology, Thuwal. Saudi Arabia

³ NNAISENSE, Lugano, Switzerland

Abstract

This article provides a rigorous analysis of convergence and stability of Episodic Upside-Down Reinforcement Learning, Goal-Conditioned Supervised Learning and Online Decision Transformers. These algorithms performed competitively across various benchmarks, from games to robotic tasks, but their theoretical understanding is limited to specific environmental conditions. This work initiates a theoretical foundation for algorithms that build on the broad paradigm of approaching reinforcement learning through supervised learning or sequence modeling. At the core of this investigation lies the analysis of conditions on the underlying environment, under which the algorithms can identify optimal solutions. We also assess whether emerging solutions remain stable in situations where the environment is subject to tiny levels of noise. Specifically, we study the continuity and asymptotic convergence of command-conditioned policies, values and the goal-reaching objective depending on the transition kernel of the underlying Markov Decision Process. We demonstrate that near-optimal behavior is achieved if the transition kernel is located in a sufficiently small neighborhood of a deterministic kernel. The mentioned quantities are continuous (with respect to a specific topology) at deterministic kernels, both asymptotically and after a finite number of learning cycles. The developed methods allow us to present the first explicit estimates on the convergence and stability of policies and values in terms of the underlying transition kernels. On the theoretical side we introduce a number of new concepts to reinforcement learning, like working in segment spaces, studying continuity in quotient topologies and the application of the fixed-point theory of dynamical systems. The theoretical study is accompanied by a detailed investigation of example environments and numerical experiments.

Keywords: Reinforcement Learning, Upside-down reinforcement learning, Goal-Conditioned Supervised Learning, Online Decision Transformers, Convergence and Stability

 $[\]textcircled{O}2022$ Miroslav Štrupl, Oleg Szehr, Francesco Faccio, Dylan R. Ashley, Rupesh Kumar Srivastava, and Jürgen Schmidhuber.

Contents

1	Intr 1.1	oduction Outline	$\frac{4}{7}$	
2	Bac	kground	8	
	2.1	Markov Decision Process	8	
	2.2	Markov Decision Processes of Command Extension Type	9	
	2.3	Trajectory and Segment Distributions of CEs	11	
	2.4	The eUDRL Algorithm	13	
	2.5	The Relation of eUDRL to the ODT Recursion	14	
3	The	e eUDRL Recursion, Reward-Weighted Regression and Practical Im-		
	pler	nentations	16	
	3.1	The eUDRL Recursion in Specific Segment Spaces	16	
	3.2	The Relation of eUDRL to Reward-Weighted Regression	19	
	3.3	The Role of Prioritized Replay	19	
4	4 The Continuity of eUDRL in the Transition Kernel and Stability Prop			
	erti	es	21	
	4.1	Compatible Families and the Stability of Supports	21	
	4.2	On the Stability of Supports in Segment Sub-Spaces	24	
5	Con	vergence to Optimal Policies for Deterministic Kernels	26	
	5.1	Optimal Actions for Deterministic Transition Kernels	26	
	5.2	Convergence of eUDRL in the Space of all Segments	27	
	5.3	Convergence of eUDRL in Segment Sub-Spaces	28	
6	The	The Continuity of eUDRL at Deterministic Transition Kernels for a Finite		
	Nur	mber of Iterations	29	
	6.1	Continuity of Values and Segment Distribution Marginals	31	
	6.2	Relative Continuity of eUDRL Policies and Continuity of Related Values	32	
	6.3	Extending the Continuity Results to Other Segment Sub-Spaces	37	
	6.4	Continuity of the Goal-Reaching Objective for a Finite Number of Iterations	39	
7	The	e Continuity of eUDRL at Deterministic Transition Kernels as the		
	Nur	mber of Iterations Approaches Infinity	42	
	7.1	Asymptotic Continuity for Special Initial Distributions	42	
		7.1.1 Preliminary lemmata	43	
		7.1.2 The main theorem	48	
		7.1.3 Extending the continuity results to other segment sub-spaces	53	
		7.1.4 Estimating the location of accumulation points	54	
	7.2	Asymptotic Continuity Assuming Uniqueness of the Optimal Policy	59	
		7.2.1 Preliminary lemma	61	
		7.2.2 The main theorem	63	
		7.2.3 Extending the continuity results to other segment sub-spaces	67	

	7.2.4 Estimating the location of accumulation points	67
8	Regularized Recursion	72
9	Related Work	78
	9.1 Reward-Weighted Regression	78
	9.2 Goal-conditioned Reinforcement Learning Without Relabeling	78
	9.3 UDRL	79
	9.4 Transformers	82
10	Conclusion	83
	10.1 Investigation of Stability at a Finite Number of Iterations	83
	10.2 Convergence and Investigation of Stability at an Infinite Number of Iterations	s 83
	10.3 Further Results	84
A	The Segment Distribution and its Factorization	90
в	Motivating Examples	93
\mathbf{C}	Continuity at Interior Kernels for a Finite Number of Iterations	101
D	Regularized Recursion — Lemmas and Proofs	104
	D.1 Preliminary Lemmata	104
	D.2 The Main Theorem	109
	D.3 Extending the Continuity Results to Other Segment Sub-Spaces	113
	D.4 Estimating the Location of Accumulation Points	113

1 Introduction

Reinforcement Learning (RL) algorithms are designed to learn policies that choose optimal actions while interacting with an environment. The environment does not reveal optimal actions, but only provides higher rewards for taking better actions. This is in direct contrast with Supervised Learning (SL), where the correct output for each input is available for learning. Nevertheless, a series of algorithms have been proposed that attempt to solve reinforcement learning tasks using purely supervised learning techniques. Upside-Down Reinforcement Learning (Schmidhuber, 5 Dec 2019; Srivastava et al., 2019) (UDRL) inverts the traditional RL process by mapping desired returns/goals to actions, treating action prediction as a supervised learning problem. Goal-conditioned Supervised Learning (Ghosh et al., 2021) (GCSL) utilizes goal information to guide the model's learning process, and Online Decision Transformers (Zheng et al., 2022) (ODT) leverage transformer architectures to model entire trajectories, treating past states, actions, and rewards as sequences to predict optimal actions. Experiments have shown that in addition to being strikingly simple and scalable due to their dependence on SL, these algorithms can produce good results on several RL benchmarks (such as Vizdoom (Kempka et al., 2016), robotic manipulation (Ahn et al., 2020) and locomotion (Fu et al., 2020)). Their theoretical understanding, however, is limited to heuristics and the study of restrictive special cases. Through a rigorous analysis of convergence and stability, this work initiates the development of a theoretical foundation for algorithms that build on the broad idea of approaching RL via SL or sequence modeling. Two questions guide our investigation: 1) What can be said about the convergence of UDRL, GCSL and ODT assuming that an explicit model (transition kernel) for the underlying Markovian environment is given? What behavior can be expected of typical objects of interest, such as policies, state and action-values in the limit of infinite resources. 2) How stable are these quantities under perturbation or in the presence of errors in the environment model? Guarantees that ensure that algorithms reliably identify optimal solutions and remain stable under varying conditions are foundational for their practical deployment in real-world systems (Bertsekas and Tsitsiklis, 1996; Nocedal and Wright, 2006).

Stepping back to establish some basic background, it is notable that UDRL, GCSL, and ODT are very similar algorithms. Although architectural details vary, at their core stand common ideas about the acquisition of information by the learning agent. They all focus on directly predicting actions based on reward signals from trajectories, rather than learning value functions. The key shared ingredient is the interpretation of rewards, observations and planning horizons as task-defining inputs from which a command is computed for the learning agent. The agent's rule of action (the *policy*) is then updated through SL, to map previous trajectory observations and commands into actions, completing the learning process. More formally, suppose a number of trajectory samples have been collected by a learning agent that follows a certain rule of action π_{old} . Given a trajectory segment that starts with a state-action pair (s, a), has a length of h, and where the goal g is a quantity that is computed from features of the segment (such as the sequences of states and rewards), one could reason that the action a is useful for achieving g from s in h steps. It then appears natural to interpret (h, g) as a command for the agent, fitting the new rule of action π_{new} to the distribution a|s, h, g using SL,

$$\pi_{new} = \operatorname{argmax}_{\pi} \mathbb{E}[\operatorname{loss}(\pi(a|s, h, g))],$$

where loss is an appropriate loss function and the expectation is computed over all segments in the trajectory sample. Learning proceeds iteratively by replacing π_{old} with π_{new} , sampling new trajectories from π_{old} , computing the achieved horizons and goals (h, g), and finally using this information to update π_{new} .

Our analysis of the UDRL, GCSL, and ODT algorithms is conducted within the overarching framework of episodic UDRL (eUDRL), which is characterized by a specific structure of the goal. Specifically eUDRL assumes that a goal map ρ is applied to a segment's terminal state to evaluate whether the goal has been reached $g = \rho(s)$. GCSL can be viewed as a slightly restricted version of eUDRL, as it focuses solely on state-reaching tasks and operates with a fixed horizon. Decision Transformer (Chen et al., 2021) (DT) essentially corresponds to one iteration of eUDRL aimed at offline RL, and ODT can be seen as a form of eUDRL with entropy regularization; see the background section 2 for details. Unlike standard RL policies which define an agent's action probabilities based on a given state, the "policies" π_{old} and π_{new} also condition these probabilities on the command (h, g). This leads to the formalism of Command Extensions (cf. definition 1), a special class of Markov Decision Processes, where the command is included as part of the state, $\bar{s} = (s, h, g)$. On the technical side this article develops the mathematics of Command Extensions, which provides a solid foundation for exploring RL through SL.

eUDRL's learning process only requires a single SL step, with no need for value function or policy proxies. While simplicity and high experimental performance coincide remarkably in the eUDRL algorithm, it is evident only in deterministic environments that eUDRL identifies the optimal policy. This limitation was acknowledged already in the original work (Schmidhuber, 5 Dec 2019), where an alternative approach was proposed for nondeterministic environments (e.g., by operating with expected returns rather than just actual returns). In practice, eUDRL's simplicity has led to its adoption in non-deterministic environments as well. In part this was based on the observation that many non-deterministic environments exhibit only minor non-determinism, i.e. they can be viewed as perturbed deterministic environments. This approach was explored in the case of eUDRL in the article (Srivastava et al., 2019), as well as concurrently in the case of GCSL in (Ghosh et al., 2021), demonstrating the algorithm's practical utility in various domains, including several MuJoCo tasks. However, these articles did not provide solid convergence guarantees. While Ghosh et al. (2021) showed that GCSL optimizes a lower bound on the goal-reaching objective, the guarantees regarding its tightness are limited. In fact, eUDRL's goal-reaching objective can be sensitive to perturbations of the transition kernel even in near-deterministic environments. We will discuss this behavior in several instances throughout the article illustrating it with specific examples and computations. Consider figure 1 for an illustration of the mentioned discontinuity of the UDRL-generated goal-reaching objective in a specific environment (see section 6 for details). The figure shows the values of the goal-reaching objective (denoted by J_{α}) along two continuous one-parameter families of environment transition kernels, where the families are parametrized by α and intersect at a parameter value of $\alpha = 0$. In the case of figure 1a the one-parameter families intersect at a deterministic environment. In the case of figure 1b the families intersect a specific non-deterministic environment. While figure 1a suggests a continuous behavior, the goal-reaching objective appears to be discontinuous in figure 1b. Despite eUDRL's remarkable characteristics, the algorithm's stability is clearly a concern.



(a) Determinist. Kernel - Goal-R. Objective

(b) Non-Deter. Kernel - Goal-R. Objective

Figure 1: Illustration of discontinuity of eUDRL-generated goal-reaching objective along two continuous one-parameter rays (items A and C) of environments (transition kernels). Horizontal axes show the value of the ray-parameter α ; the point of intersection is $\alpha = 0$. The exact value of the respective quantities at $\alpha = 0$ is represented by a horizontal line (item B).

The instability of eUDRL (particularly UDRL, GCSL, and DT) in non-deterministic environments has been highlighted by Štrupl et al. (2022a) and Paster et al. (2022). The former analyzes the principles behind this instability, while the latter addresses issues with DT and aims for a fully functional variation of the algorithm. Since ODT can be seen as a variation of DT with online fine-tuning, the observed instability is also present in ODT. Subsequent studies, such as (Yang et al., 2023), have proposed various improvements to UDRL, GCSL, and DT. Although Brandfonbrener et al. (2022) provide an in-depth analysis of the first eUDRL iteration in the context of offline RL, a comprehensive treatment of the behavior of UDRL, GCSL and ODT at a finite number of iterations and in the asymptotic limit remains an open problem.

This article does not propose another approach to ensure that eUDRL works in nondeterministic environments, but seeks to understand its original design and justify its empirical success, as seen in (Srivastava et al., 2019) and (Ghosh et al., 2021). We aim to determine if iterating eUDRL beyond the first step leads to convergence to a desired asymptotic behavior, focusing on rigorous convergence guarantees and asymptotic error bounds. To extend convergence guarantees for deterministic transition kernels to nearly-optimal behavior for kernels close to deterministic ones we employ a topological approach. This approach includes mathematical tools that are new to the area of RL, introducing notions like continuity in quotient topologies that employ the weak convergence of measures relative to compact sets (the so-called *relative continuity*). We prove the relative continuity of policies and the continuity of the goal-reaching objectives at deterministic kernels after any finite

iteration. Despite its significance, this topic is largely unexplored, partly due to its complexity. As we will show, eUDRL policies become discontinuous from the second iteration onward, particularly at the boundary of the set of transition kernels (including deterministic kernels). This suggests that the first iteration is as effective as any finite iteration but it does not clarify whether repeated iterations reduce the error to optimal behavior. By establishing bounds for the eUDRL recursion under specific conditions, we demonstrate the relative continuity of accumulation point sets of eUDRL-generated policies and prove the continuity of related quantities like the goal-reaching objective, along with useful bounds and q-linear convergence rates. These conditions outline two important special cases: The first involves a condition on the support of the initial distribution and the second is characterized by the uniqueness of the optimal policy in a deterministic environment, see section 7 for details. While the general discussion of continuity and stability of eUDRL remains an open problem, we have found that the theory developed in this article is sufficient to address the regularized eUDRL recursion. The regularized eUDRL recursion closely approximates the ODT, which utilizes entropy regularization. Our analysis of the regularized eUDRL recursion, formulated as a convex combination between the updated policy with the uniform distribution, akin to ϵ -greedy policies, is presented in full generality, without restricting it to any special cases. We restrict all discussions to finite (discrete) environments, ignoring issues of function approximation and limited sample size.

1.1 Outline

Section 2 provides the necessary background for understanding the article.

Section 3 describes the eUDRL recursion in specific segment subspaces and discusses the connection to the algorithm Reward-weighted Regression.

Section 4 describes sets of states where eUDRL is stable under environment perturbations across all iterations, which proves instrumental for the discussion of continuity of eUDRL-generated quantities in the subsequent sections.

Section 5 proves that eUDRL converges to optimal policies in case of deterministic environments.

Section 6 investigates the continuity of eUDRL-generated quantities at a finite number of iteration for deterministic environments.

Section 7 investigates the continuity of sets of accumulation points of eUDRL-generated quantities for deterministic environments.

Section 8 investigates the continuity of sets of accumulation points of quantities generated by regularized eUDRL in full generality.

Section 9 discusses related work and section 10 concludes the presentation.

To make this article accessible to a wider audience several appendices are included.

Appendix A contains details about the construction of segment distribution.

Appendix B contains examples from the main text, worked out in full detail.

Appendix C investigates the continuity of eUDRL-generated quantities at interior points (of the set of all transition kernels) for a finite number of iterations.

Appendix D contains all details on lemmas and proofs about the regularized eUDRL recursion of section 8.

2 Background

This section provides the necessary background for understanding the article. It offers an orientation on the placement of our work within the "theoretical landscape", introducing fundamental theoretical concepts, particularly focusing on Markov Decision Processes of Command Extension type and distributions over segment space. Furthermore, this section outlines how our work fits within the existing literature, describing the integration of ODTs within the eUDRL framework. The innovative developments presented in this article will be introduced in the following sections.

We will be dealing exclusively with finite, real random variables. They are defined as maps $X : \Omega \to \mathbb{R}$ with a measure space $(\Omega, \mathcal{F}, \mathbb{P})$, where Ω is a finite set, \mathcal{F} is a σ -algebra over Ω and \mathbb{P} denotes a probability measure.

2.1 Markov Decision Process

Markov decision processes (MDPs) are a mathematical framework for sequential decision problems in uncertain dynamic environments (Puterman, 2014). Formally an MDP is a five-tuple $\mathcal{M} = (\mathcal{S}, \mathcal{A}, \lambda, \mu, R)$, with a set \mathcal{S} of admissible states, a set of possible actions \mathcal{A} , a transition probability kernel λ that defines the probability of entering a new state given that an action is taken in a given state, a distribution μ over initial states, and a (random) reward R that is granted through this transition. In MDPs an agent interacts iteratively with an environment over a sequence of time steps t (where we add the subscript t to random variables to emphasise that they belong to a specific point of time). Let the random variables $S_t : \Omega \to \mathcal{S}$ describe the state of the MDP and the random variables $A_t: \Omega \to \mathcal{A}$ describe the actions chosen by the agent. Beginning with an initial distribution over states $\mu(s) = \mathbb{P}(S_0 = s)$ at each step of the MDP the agent observes the current state $s \in \mathcal{S}$ and takes a respective action $a \in \mathcal{A}$ according a policy $\pi(a|s) = \mathbb{P}(A_t)$ $a|S_t = s$, subsequently the environment transitions to s' according to the probability $\lambda(s'|s,a) = \mathbb{P}(S_{t+1} = s' \mid S_t = s, A_t = a)$. A transition kernel λ is deterministic if for each $s, a \in \mathcal{S} \times \mathcal{A}$ the distribution $\lambda(\cdot | s, a)$ is deterministic, i.e. if there exists a state $s'_{s,a}$ such that $\lambda(s'_{s,a}|s,a) = 1$. Similarly a policy π is deterministic if for each $s \in S$ the distribution $\pi(\cdot|s)$ is deterministic. Consecutive sequences of state-action transitions are commonly called trajectories of the MDP. In this article we will always assume deterministic rewards, i.e., the reward is given by $R_{t+1}(S_{t+1}, S_t, A_t) = r(S_{t+1}, S_t, A_t)$ with a deterministic function r. The return $G_t = \sum_{k \in \mathbb{N}_0} R_{t+k+1}$ is the accumulated reward beginning with time t over the entire MDP episode, where we do not discount future rewards. The performance of an agent following a policy π can be measured by means of the state value function $V^{\pi}(s) = \mathbb{E}[G_t \mid S_t = s; \pi]$ and the action value function $Q^{\pi}(s, a) = \mathbb{E}[G_t \mid S_t = s, A_t = a; \pi]$. There exist a unique optimal state value function $V^* = \max_{\pi} V^{\pi}$ and a unique optimal action value function $Q^* = \max_{\pi} Q^{\pi}$, where the maximization is taken over the set of all policies. A policy π^* for which $V^{\pi^*} = V^*$ and consequently also $Q^{\pi^*} = Q^*$ is called optimal. In what follows we will work with special types of MDPs called *Command Extensions*. In this article we will reserve the symbols π , V^{π} , Q^{π} for the respective quantities for MDPs of Command Extension type.

2.2 Markov Decision Processes of Command Extension Type

The aim of the training process in eUDRL is to achieve that the agent becomes better at executing commands. In eUDRL the command is provided to the agent in the form of a "goal" and a "horizon". The goal says which state the agent is supposed to achieve and the horizon says when this state should be achieved. A goal map $\rho : S \to G$ will be employed to evaluate if the command's goal has been reached at the command's horizon, i.e. if $\rho(s)$ equals to the command's specified goal for the final state s. The codomain \mathcal{G} of the goal map is the set of all possible goals, weather they are reached in one specific trajectory or not. Notice that since ρ is defined on the entire state space \mathcal{S} , every state s at the horizon corresponds to a valid goal $\rho(s)$. However, this goal map allows one to study prototypical eUDRL tasks within a unified formalism: The state reaching task is covered by $\mathcal{G} = S, \rho = \mathrm{id}_{\mathcal{S}}$ (the identity map on \mathcal{S}), while goals related to return of the original MDP \mathcal{M} can be covered by extending the state by a component that accumulates rewards and defining ρ to be the projection on this component.

Apart from the MDP state the eUDRL agent takes an additional command input. Let the random variable $G_t : \Omega \to \mathcal{G}$ describe the goal of the eUDRL agent and let $H_t : \Omega \to \overline{\mathbb{N}}_0 = \{0, 1, ..., N\}, N \geq 1$, be a random variable that describes the remaining horizon. The eUDRL agent can be viewed as an ordinary agent operating in an MDP whose state space is extended by the command. In this context one has to provide an initial distribution over goals and horizons $\mathbb{P}(H_0 = h, G_0 = g \mid S_0 = s)$ and with each transition of the extended MDP the remaining horizon decreases by 1 until a horizon of 0 is reached. At this stage the extended MDP enters an absorbing state (see below for a definition), from which no further evolution occurs. A reward is granted if the specified goal is reached when the horizon turns to 0. In summary we have the following definition of command extensions:

Definition 1 (Command Extension) A Command Extension (CE) of an MDP of the form $\mathcal{M} = (\mathcal{S}, \mathcal{A}, \lambda, \mu, r)$ is an MDP of the form $\overline{\mathcal{M}} = (\overline{\mathcal{S}}, \mathcal{A}, \overline{\lambda}, \overline{\mu}, \overline{r}, \rho)$, where:

- A command is a pair $(g,h) \in \mathcal{G} \times \overline{\mathbb{N}}_0$, where $\mathcal{G} \subset \mathbb{Z}^{n_G}$ and $\overline{\mathbb{N}}_0 = \{0, 1, ..., N\}$. $N \ge 1$ stands for the maximum horizon and \mathcal{G} for the goal set of the CE. The goal map $\rho: S \to \mathcal{G}$ is used to evaluate if a goal $g \in \mathcal{G}$ has been reached.
- The extended state space is $\bar{S} = S \times \bar{\mathbb{N}}_0 \times G$ and the extended state is a triple $\bar{s} = (s, h, g) \in \bar{S}$ composed of an original MDP state s and the command (g, h).
- The initial distribution of *M* is given by a product of a distribution of commands and the initial distribution of *M*

$$\bar{\mu}(\bar{s}) = \mathbb{P}(H_0 = h, G_0 = g \mid S_0 = s)\mu(s).$$

• The transition kernel $\bar{\lambda}$ is defined for all $(s, h, g) \in \bar{S}$, all $s' \in S$ and $a \in A$ by

$$\begin{split} \bar{\lambda}\Big((s',h-1,g)\mid(s,h,g),a\Big) &= \lambda(s'\mid s,a) & \text{if } h > 0, \\ \bar{\lambda}\Big((s',h,g)\mid(s,h,g),a\Big) &= \delta_{ss'} & \text{if } h = 0. \end{split}$$

The reward function r̄ is defined for all s̄' = (s', h', g') ∈ S̄, s̄ = (s, h, g) ∈ S̄ and all a ∈ A by

$$\bar{r}((s',h',g'),(s,h,g),a) = \begin{cases} \mathbf{1}_{\{\rho(s')=g\}} & \text{if } h = 1, h' = 0, g' = g, \\ 0 & \text{otherwise.} \end{cases}$$

We will commonly refer to \mathcal{M} as the original MDP and to $\overline{\mathcal{M}}$ as the extended MDP. Similarly, s will be the original state and \overline{s} will be the extended state.

Our subsequent investigation almost exclusively revolves around the policies of extended MDPs rather than those of the underlying original MDPs. For brevity we reserve the notation π (rather than $\bar{\pi}$) for the policies of CE agents. The reward is defined in a way as to guide the agent towards achieving the intended command. The choice of a binary reward function allows one to interpret the expected reward values as probabilities. In summary the computation of state and action-value functions yields that for all $(s, h, g) \in \bar{S}$, h > 1, and all actions $a \in A$

$$Q^{\pi}((s,h,g),a) = \mathbb{P}\Big(\rho(S_{t+h}) = g \Big| S_t = s, H_t = h, G_t = g, A_t = a; \pi\Big),$$

$$V^{\pi}(s,h,g) = \mathbb{P}\Big(\rho(S_{t+h}) = g \Big| S_t = s, H_t = h, G_t = g; \pi\Big).$$
(2.1)

With these state and action-value functions, CEs could be interpreted as ordinary finitehorizon RL problems. One could then choose from a variety of algorithms that are guaranteed to converge to an optimal policy, see e.g. the monograph by Sutton and Barto (2018). However, this work is dedicated to the study of the eUDRL algorithm, which is not just an ordinary RL algorithm applied to an MDP of CE type. Instead eUDRL is defined through a specific iterative procedure on CEs (that involves trajectory sampling, supervised learning and policy updates), see section 2.4 for a detailed description, which (despite of certain convenient properties) is not always guaranteed to converge to the optimum. In the study of eUDRL it will be convenient to employ the *goal-reaching objective* of Ghosh et al. (2021), which is derived from the CE value function and can be written as

$$J^{\pi} = \sum_{\bar{s}\in\bar{\mathcal{S}}} \mu(\bar{s}) V^{\pi}(\bar{s}).$$
(2.2)

There exist a unique optimal goal-reaching objective $J^* = \max_{\pi} J^{\pi}$, where maximization is taken over the set of all policies. It is easy to see that for any optimal policy π^* it holds that $J^{\pi^*} = J^*$.

It will be convenient to introduce a separate notation of absorbing states of the CE. Recall that a state $s \in S$ is called *absorbing* for \mathcal{M} if once the absorbing state is reached the underlying process always remains in this state, i.e. for all states s, s' and actions a the kernel equals the Kronecker delta $\lambda(s'|sa) = \delta_{ss'}$.

Remark 2 (Absorbing State of CE) Let $\overline{\mathcal{M}} = (\overline{\mathcal{S}}, \mathcal{A}, \overline{\lambda}, \overline{\mu}, \overline{r}, \rho)$ be a CE. A state of the form $(s, g, h) \in \overline{\mathcal{S}}$ is an absorbing state if and only if h = 0. A state that is not absorbing is called transient. We denote by $\overline{\mathcal{S}}_A \subset \overline{\mathcal{S}}$ and $\overline{\mathcal{S}}_T \subset \overline{\mathcal{S}}$, respectively, the sets of absorbing and transient states.

Initializing a CE into an absorbing state results in no learnable information because the CE will remain in this state, irrespective of the chosen actions. For this reason we will focus our investigation on such CEs that are not initialized into an absorbing state. In other words we will assume that the initial distribution of the studied CE is such that it assigns 0 probability to absorbing states,

$$\operatorname{supp} \bar{\mu} \subset \bar{\mathcal{S}}_T. \tag{2.3}$$

Such CEs are called *non-degenerate*. Hereafter we will only study non-degenerate CEs, loosely referring to them as simply as CEs.

2.3 Trajectory and Segment Distributions of CEs

A key role in the eUDRL algorithm is played by segments of sampled CE trajectories. A CE trajectory contains a sequence of consecutive transitions comprised of extended states and actions. In full generality CE trajectories could be represented by infinite sequences of the form

$$\tau = ((s_0, h_0, g_0), a_0, (s_1, h_0 - 1, g_0), a_1, \dots, (s_{h_0 - 1}, 1, g_0), a_{h_0 - 1}, (s_{h_0}, 0, g_0), \dots).$$

However, once the maximum horizon of the CE is reached, the CE enters an absorbing state, from which no further evolution occurs. For this reason it is sufficient to represent trajectories by finite sequences of a maximum length N. Let $\text{Traj} \subset (\bar{S} \times \mathcal{A})^N \times \bar{S}$ denote the subset of finite trajectories of length N whose transitions satisfy the requirements of definition 1 on the dynamics of horizons and goals. In what follows we will represent trajectories by

$$\tau = ((s_0, h_0, g_0), a_0, (s_1, h_0 - 1, g_0), a_1, \dots \\ \dots, (s_{l(\tau)-1}, 1, g_0), a_{l(\tau)-1}, (s_{l(\tau)}, 0, g_0), \dots (s_N, 0, g_0)) \in \text{Traj.}$$

The quantity $l(\tau)$ stands for the number of transitions until an absorbing state is entered for the first time; it is equal to the initial horizon of τ . Assuming finite state and action spaces Traj is measurable and we denote by $\mathcal{T} : \Omega \to \text{Traj}$ a random variable with components $\mathcal{T} =$ $((S_0, H_0, G_0), A_0, \ldots, (S_N, H_N, G_N))$ whose outcomes are CE trajectories. The probability of $\tau \in \text{Traj}$ is given by

$$\mathbb{P}(\mathcal{T}=\tau;\pi) = \left(\prod_{t=1}^{l(\tau)} \lambda(s_t | a_{t-1}, s_{t-1})\right) \cdot \left(\prod_{t=0}^{l(\tau)-1} \pi(a_t | \bar{s}_t)\right) \cdot \bar{\mu}(\bar{s}_0).$$

Each segment contains a chunk of consecutive state-action transitions from a given trajectory and also the segment's initial horizon, goal and length. We will represent segments by tuples of the form

$$\sigma = (l(\sigma), s_0^{\sigma}, h_0^{\sigma}, g_0^{\sigma}, a_0^{\sigma}, s_1^{\sigma}, a_1^{\sigma}, \dots, s_{l(\sigma)}),$$

where $a_0^{\sigma}, \ldots, a_{l(\sigma)-1}^{\sigma}$ are the chosen actions, $s_0^{\sigma}, \ldots, s_{l(\sigma)}^{\sigma}$ are the reached states of the original MDP and the subscript *t* indicates the time of the respective quantity *within the segment* (i.e. irrespective of the actual time when the state or action occurred in the original MDP's

trajectory). The quantity $l(\sigma)$ stands for the length of the segment (i.e. the number of actions executed during the course of the segment) and h_0^{σ} and g_0^{σ} denote the segment's initial horizon and goal. In our notation we leave out the horizons and goals for all extended states except from the first one because they are already determined by the CE's horizon and goal dynamics. We write Seg for the set of all tuples of this form. Assuming finite state and action spaces Seg is measurable and we denote by $\Sigma : \Omega \to \text{Seg a random variable with components } \Sigma = (l(\Sigma), S_0^{\Sigma}, H_0^{\Sigma}, G_0^{\Sigma}, A_0^{\Sigma}, S_1^{\Sigma}, A_1^{\Sigma}, \dots, S_{l(\Sigma)}^{\Sigma})$ whose outcomes are segments of CE trajectories. We provide all formal details about the construction of the distribution of Σ in appendix A. Here we only summarize that for all $\sigma \in \text{Seg holds}$

$$\mathbb{P}(\Sigma = \sigma; \pi) = c^{-1} \sum_{t \le N - l(\sigma)} \mathbb{P}(S_t = s_0^{\sigma}, H_t = h_0^{\sigma}, G_t = g_0^{\sigma}, A_t = a_0^{\sigma}, \dots, S_{t+l(\sigma)} = s_{l(\sigma)}^{\sigma}; \pi),$$

where c is the normalization constant

$$c = \sum_{\sigma \in \text{Seg } t \le N - l(\sigma)} \mathbb{P}(S_t = s_0^{\sigma}, H_t = h_0^{\sigma}, G_t = g_0^{\sigma}, A_t = a_0^{\sigma}, \dots, S_{t+l(\sigma)} = s_{l(\sigma)}^{\sigma}; \pi).$$

It is shown in the appendix that the following bounds hold for the normalization constant:

$$0 < c \le \frac{N(N+1)}{2}.$$
(2.4)

We will also demonstrate the following "Markovianity properties" for the distribution of Σ in the appendix. For every segment of length k and any $i \leq k$ we have that

$$\mathbb{P}\left(A_{i}^{\Sigma} = a_{i} \middle| l(\Sigma) = k, S_{0}^{\Sigma} = s_{0}, H_{0}^{\Sigma} = h_{0}, G_{0}^{\Sigma} = g_{0}, A_{0}^{\Sigma} = a_{0}, \dots, S_{i}^{\Sigma} = s_{i}; \pi_{n}\right)$$

$$= \pi_{n}(a_{i} | s_{i}, h - i, g),$$
(2.5)

$$\mathbb{P}\left(S_{i}^{\Sigma}=s_{i}\Big|l(\Sigma)=k, S_{0}^{\Sigma}=s_{0}, H_{0}^{\Sigma}=h_{0}, G_{0}^{\Sigma}=g_{0}, A_{0}^{\Sigma}=a_{0}, \dots, S_{i-1}^{\Sigma}=s_{i-1}, A_{i-1}^{\Sigma}=a_{i-1}; \pi_{n}\right)$$
$$=\mathbb{P}(S_{i}^{\Sigma}=s_{i}|S_{i-1}^{\Sigma}=s_{i-1}, A_{i-1}^{\Sigma}=a_{i-1})=\lambda(s_{i}|s_{i-1}, a_{i-1}).$$
(2.6)

Restriction to Trailing Segments: Sometimes it is useful restrict the analysis to specific subsets of segments in Seg. For instance it will useful to consider so called "trailing segments". A segment $\sigma = (l(\sigma), s_0^{\sigma}, h_0^{\sigma}, g_0^{\sigma}, a_0^{\sigma}, \dots, s_{l(\sigma)}^{\sigma})$ is trailing if it is aligned with the end of a trajectory, i.e., if $l(\sigma) = h_0^{\sigma}$. The subspace of Seg consisting of all trailing segments will be denoted by Seg^{trail} := { $\sigma \in \text{Seg} \mid l(\sigma) = h_0^{\sigma}$ }. ODT and also sometimes eUDRL operates on trailing segments. In the case of eUDRL this restriction was motivated by speeding up the learning process reflecting the episodic nature of the problem.

Restriction to Diagonal Segments: A segment that satisfies $g_0^{\sigma} = \rho(s_{l(\sigma)}^{\sigma})$ is called a goal-diagonal segment. Such segments are characterized by the fact that the initial goal of the segment is actually achieved at the end of the segment. A segment which is trailing and goal-diagonal will be called diagonal. The subspace of Seg which consists of all diagonal segments will be denoted Seg^{diag} := { $\sigma \in \text{Seg}|l(\sigma) = h_0^{\sigma}, g_0^{\sigma} = \rho(s_{l(\sigma)}^{\sigma})$ }. We will consider these segments when discussing the relation to the Reward-Weighted Regression (Peters and Schaal, 2007), see section 3.2.

2.4 The eUDRL Algorithm

Suppose that a CE $\overline{\mathcal{M}}$ is given. Starting with an initial policy π_0 eUDRL generates a sequence of policies $(\pi_n)_{n\geq 0}$ in an iterative process. Each eUDRL iteration comprises the following steps. First, a batch of trajectories is generated from $\overline{\mathcal{M}}$ following the current policy π_n . Second, segments σ of trajectories are sampled from the batch according to the distribution $d_{\Sigma}^{\pi_n}$ of Σ . Finally, a subsequent policy π_{n+1} is fitted using supervised learning to the sampled trajectory segments. In practice, a class of parameterized policies will usually be assumed and π_{n+1} will be one of the policies that minimize log-likelihood loss within this class. Here, we investigate what eUDRL can achieve in principle. We assume no specific representation of policies and we suppose that π_{n+1} is computed using cross-entropy,

$$\pi_{n+1} = \operatorname{argmax}_{\pi} \mathop{\mathbb{E}}_{\sigma \sim d_{\Sigma}^{\pi_n}} \log \left(\pi \left(a_0^{\sigma} \middle| s_0^{\sigma}, l(\sigma), \rho(s_{l(\sigma)}^{\sigma}) \right) \right),$$
(2.7)

which reflects infinite sample size or complete knowledge about the distribution $d_{\Sigma}^{\pi_n}$. In other words the new policy π_{n+1} is fitted to the conditional probability¹

$$\pi_{n+1}(a \mid (s, h, g)) = \mathbb{P}\left(A_0^{\Sigma} = a \mid S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_{l(\sigma)}^{\Sigma}) = g; \pi_n\right).$$
(2.8)

The choice of the conditional in equation (2.7) can be motivated by a comparison with the Hindsight Experience Replay (HER) algorithm of Andrychowicz et al. (2017). The way eUDRL uses segment samples is similar to that of HER in that not only trajectories that achieve the intended goals are used for learning but also trajectories that do not achieve the intended goals, together with their effective outcomes. Following a similar line of reasoning, if a sample $\sigma \sim d_{\Sigma}^{\pi_n}$ is observed, one might assume that the first action a_0^{σ} is a good choice for executing the realized command $(l(\sigma), \rho(s_{l(\sigma)}^{\sigma}))$, irrespective of the actual command $(h_0^{\sigma}, g_0^{\sigma})$ chosen for this segment.

Many algorithms that follow the paradigm of "reinforcement learning via supervised learning" including the practical eUDRL implementations of Srivastava et al. (2019), Goal-Conditioned Supervised Learning (GCSL) of Ghosh et al. (2021) and Online Decision Transformer (ODT) of Zheng et al. (2022) fit into the framework of this article. It should be said that our assumption of infinite sample size implies a simplification as compared to the eUDRL variations discussed by Srivastava et al. (2019) or Ghosh et al. (2021), where we omit the replay buffer (since it is not required). The algorithm in Srivastava et al. (2019) uses a trajectory organized replay buffer prioritized by return, where goals are related to the trajectory returns. Further initial commands are sampled according to a best quantile of the distribution of returns in the replay buffer. This means that the initial distribution is changing over time, while here we assume that it is fixed. The GCSL algorithm of Ghosh et al. (2021) can be viewed as a simplified version of Srivastava et al. (2019) in that GCSL executes pure state-reaching tasks. GCSL omits the horizon component from commands assuming a fixed horizon for its tasks. Similarly to Srivastava et al. (2019), GCSL employs a replay buffer, but without any prioritization. The relation between eUDRL and ODT is described in a separate section, section 2.5, below.

^{1.} Notice that π_{n+1} is defined by (2.8) if and only if $\mathbb{P}(S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_{l(\sigma)}^{\Sigma}) = g; \pi_n) > 0$, otherwise we leave it undefined for now. Later, it will be convenient to choose wlog. $\pi_{n+1} = 1/|\mathcal{A}|$ if $\mathbb{P}(S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_{l(\sigma)}^{\Sigma}) = g; \pi_n) = 0$.

A convenient feature of the eUDRL iteration is that it is based solely on supervised learning as opposed to algorithms that employ Bellman-type updates like, e.g., in value iteration or actor-critic algorithms. It was hypothesized that if eUDRL is close to SL it could be easier used for training large scale networks which may require many stabilization tricks with traditional RL algorithms. While the motivation for the eUDRL iteration is to improve successive policies through updates, it was shown by Štrupl et al. (2022a) that the described updates neither guarantee monotonic policy improvements nor the convergence to an optimal policy in stochastic episodic environments (in sense of CE value functions). It is interesting that when introducing eUDRL, Schmidhuber (5 Dec 2019) reserved the described iteration for deterministic environments, where, as we will discuss in the main body of this article, the algorithm converges to an optimum. On the other hand, there are also successful reports about eUDRL's practical implementations, see work by Srivastava et al. (2019) and Ghosh et al. (2021), on certain stochastic environments.

2.5 The Relation of eUDRL to the ODT Recursion

The idea of combining the successful transformer architecture (see (Schmidhuber, 1992), (Vaswani et al., 2017), (Schlag et al., 2021)) with UDRL led to the concept of Online Decision Transformers (ODTs) that interpret RL as a sequence modeling problem. The sequences in question consist of tuples (s'_t, a'_t, g'_t) of states, actions and so-called "returns-to-go" of the MDP to be solved by ODT. The return-to-go g'_t is computed once a trajectory is completed. It is given by the rewards accumulated from the time t until the end of the trajectory.

In ODT the transformer model receives sequence of states and returns-to-go to predict the subsequent action. To approximate the distribution of a'_t at each time step t the transformer (with context length K) receives sequences of length min $\{t, K\}$ of preceding states $s'_{-K,t} := s'_{\min\{0,t-K\}:t}$ and returns-to-go $g'_{-K,t} := g'_{\min\{0,t-K\}:t}$. Similar to UDRL this distribution is employed as a policy that is conditioned on returns-to-go. In contrast to the original transformer architecture, which uses positional encoding, ODT uses time embeddings. Furthermore, ODT operates exclusively on trailing segments. As compared to its predecessor, the Decision Transformer (DT), ODT introduces a series of improvements that allow for online fine-tuning. First, ODT uses stochastic policies (while DT solely allowed for deterministic policies) and, second, these policies are trained using the maximum likelihood criterion (while DT relied on mean squares). These features are present in similar form also in the eUDRL algorithm. Furthermore, ODT introduced entropy regularization, which motivates the investigation of regularization techniques also for eUDRL. In section 8 we will consider eUDRL with policy regularization using a convex combination with a uniform distribution, which brings our investigation close to the ODT algorithm.

Like the eUDRL implementations of Srivastava et al. (2019) and Ghosh et al. (2021), ODT, too, needs some simplifications in order to conform to the CE framework. As before we omit replay buffers and we assume a fixed horizon, which entails that CE initializes the horizon to N. This is to ensure a one-to-one correspondence between the remaining horizon h and current time t via t = N - h. Consequently, after ODT performs the time embedding this difference becomes obsolete. With these simplifications ODTs match goal-reaching scenarios in the CE formalism in situations, where the reward is granted in the end of the

episode. The correspondence is as follows: First the states s'_t of the MDP to be solved by ODT are grouped together into sequences of past states $s_t = s'_{-K,t}$. The original MDP underlying the CE then operates on the grouped states s_t . The codomain of the goal map ρ is chosen as the set $\mathcal{G} = \{$ "all possible rewards" $\}$, where the rewards depend only on the terminal states s_N . Policy updates are executed using maximum likelihood matching to the conditional $a_t | s_t, l(\sigma), \rho(s_N)$. Using the ODT assumptions of trailing segments, fixed horizon $l(\sigma) = h_t$ and the fact that $\rho(s_N)$ computes rewards corresponding to s'_N this is equivalent to matching $a_t | s'_{-K,t}, h_t, \rho(s_N)$. The general scenario can be phrased in the language of CEs only partially by the described accumulation of rewards as part of the state. We assume the state is given by $s_t = (s'_{-Kt}, z_{-Kt})$ and z_t corresponds to the accumulated reward of the underlying MDP. The goal map is chosen to be $\rho(s_t) = z_t$. In this case the policy updates are executed by maximum likelihood matching to the conditional $a_t|(s'_{-K,t}, z_{-K,t}), h_t, z_N,$ where returns-to-go can be recovered from $z_N - z_{-K,t}$. This does not fit perfectly the CE framework since apart from the desired returns-to-go $z_N - z_{-K,t}$ we also condition on $(z_N, z_{-K,t})$. In principle this might introduce some noise into the learning process. However, this poses no problem as we assume no limitation on the number of available samples. The formulation of ODTs in the CE framework has no drawback as the effect of noise becomes negligible in this way.

3 The eUDRL Recursion, Reward-Weighted Regression and Practical Implementations

The standard eUDRL recursion is carried out on the entire segment space. In this section, we will show recursion formulas corresponding to the subspaces of trailing and diagonal segments. This will allow us to connect eUDRL to the Reward-Weighted Regression (RWR) algorithm, see (Dayan and Hinton, 1997) and (Peters and Schaal, 2007). The known convergence of RWR (Štrupl et al., 2022b) will be an intuitive guidance for our proofs in later sections.

3.1 The eUDRL Recursion in Specific Segment Spaces

The following lemma specifies how the eUDRL recursion can be written when only segments from a specific subset of Seg are used for fitting in the recursion (2.7), see the discussion of trailing and diagonal segments above. Recall that $\sigma = (l(\sigma), s_0^{\sigma}, h_0^{\sigma}, g_0^{\sigma}, a_0^{\sigma}, \dots, s_{l(\sigma)}^{\sigma}) \in \text{Seg}$ is trailing if $l(\sigma) = h_0^{\sigma}$ and it is diagonal if it is trailing and moreover $\rho(s_{l(\sigma)}^{\sigma}) = g_0^{\sigma}$. We will often use the proportional \propto sign instead of equality in the lemma (and also in its proof) in order to save some space by leaving out bulky normalizing factors in policy distributions (which can by anyway easily recovered). As a consequence here the "proportionality" refers to the action dimension only.

Lemma 3 Consider the recursive policy updates in eUDRL described by equation (2.7).

1. Suppose the recursion is carried out on the entire set Seg. Then for all states $(s, h, g) \in \overline{S}_T$ and all actions $a \in \mathcal{A}$ it holds that

$$\pi_{n+1}(a \mid s, h, g) = \mathbb{P}(A_0^{\Sigma} = a \mid S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g; \pi_n)$$

$$\propto \sum_{\substack{h' \ge h, g' \in \mathcal{G}}} \mathbb{P}(\rho(S_h) = g \mid A_0 = a, H_0 = h', G_0 = g', S_0 = s; \pi_n) \quad (3.1)$$

$$\cdot \pi_n(a \mid s, h', g') \mathbb{P}(H_0^{\Sigma} = h', G_0^{\Sigma} = g' \mid S_0^{\Sigma} = s, l(\Sigma) = h; \pi_n).$$

2. Suppose the recursion is carried out on the set Seg^{trail}. Then for all states $(s, h, g) \in \overline{S}_T$ and all actions $a \in \mathcal{A}$ it holds that

$$\pi_{n+1}^{\text{trail}}(a \mid s, h, g) = \mathbb{P}(A_0^{\Sigma} = a \mid S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g, l(\Sigma) = H_0^{\Sigma}; \pi_n^{\text{trail}})$$

$$\propto \sum_{g' \in \mathcal{G}} \mathbb{P}(\rho(S_h) = g \mid A_0 = a, H_0 = h, G_0 = g', S_0 = s; \pi_n^{\text{trail}})$$

$$\cdot \pi_n^{\text{trail}}(a \mid s, h, g') \mathbb{P}(H_0^{\Sigma} = h, G_0^{\Sigma} = g' \mid S_0^{\Sigma} = s, l(\Sigma) = h; \pi_n^{\text{trail}}).$$
(3.2)

3. Suppose the recursion is carried out on the set Seg^{diag}. Then for all states $(s, h, g) \in \overline{S_T}$ and all actions $a \in \mathcal{A}$ it holds that

$$\pi_{n+1}^{\text{diag}}(a \mid s, h, g) = \mathbb{P}(A_0^{\Sigma} = a \mid S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g, \rho(S_h^{\Sigma}) = G_0^{\Sigma}, l(\Sigma) = H_0^{\Sigma}; \pi_n^{\text{diag}}) \\ \propto \mathbb{P}(\rho(S_h) = g \mid A_0 = a, H_0 = h, G_0 = g, S_0 = s; \pi_n^{\text{diag}}) \\ \cdot \pi_n^{\text{diag}}(a \mid s, h, g) \mathbb{P}(H_0^{\Sigma} = h, G_0^{\Sigma} = g \mid S_0^{\Sigma} = s, l(\Sigma) = h; \pi_n^{\text{diag}}).$$
(3.3)

Moreover, the policy can be written in terms of the Q-function as

$$\begin{aligned} \pi_{n+1}^{\text{diag}}(a \mid s, h, g) &\propto Q^{\pi_n^{\text{diag}}}((s, h, g), a) \pi_n^{\text{diag}}(a \mid s, h, g) \\ & \cdot \mathbb{P}(H_0^{\Sigma} = h, G_0^{\Sigma} = g \mid S_0^{\Sigma} = s, l(\Sigma) = h; \pi_n^{\text{diag}}), \\ \pi_{n+1}^{\text{diag}}(a \mid s, h, g) &\propto Q^{\pi_n^{\text{diag}}}((s, h, g), a) \pi_n^{\text{diag}}(a \mid s, h, g). \end{aligned}$$

Notice that the recursion formulas in Seg, Seg^{trail} and Seg^{diag} bear a basic similarity in that while $\pi_{n+1}^{\text{diag/trail}}(a \mid s, h, g)$ depends on the conditional probabilities $\mathbb{P}(H_0^{\Sigma} = h, G_0^{\Sigma} = g \mid S_0^{\Sigma} = s, l(\Sigma) = h; \pi_n)$ it does not depend on the conditions $l(\Sigma) = H_0^{\Sigma}$ and $\rho(S_{l(\Sigma)}^{\Sigma}) = G_0^{\Sigma}$ that define Seg^{diag} and Seg^{trail}. The recursion formulas are different only through the sets over which the summations are taken. Later this property will allow us to derive common bounds to all three recursions (i.e. bounding π_{n+1} in terms of π_n) resulting in simpler proofs.

Proof Lemma 3, Point 1. The equality is just (2.8). We start by applying Bayes' rule, where we assume $\mathbb{P}(S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g; \pi_n) > 0$. Subsequently we marginalize over H_0^{Σ} and G_0^{Σ} provided that $H_0^{\Sigma} \ge l(\Sigma) = h$ (by definition a segment is always contained in a trajectory) and we apply the product rule. We obtain

$$\begin{split} \pi_{n+1}(a \mid s, h, g) &= \mathbb{P}(A_0^{\Sigma} = a \mid S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g; \pi_n) \\ &\propto \mathbb{P}(\rho(S_h^{\Sigma}) = g, A_0^{\Sigma} = a \mid S_0^{\Sigma} = s, l(\Sigma) = h; \pi_n) \\ &= \sum_{h' \ge h, g' \in \mathcal{G}} \mathbb{P}(\rho(S_h^{\Sigma}) = g, A_0^{\Sigma} = a, H_0^{\Sigma} = h', G_0^{\Sigma} = g' \mid S_0^{\Sigma} = s, l(\Sigma) = h; \pi_n) \\ &= \sum_{h' \ge h, g' \in \mathcal{G}} \mathbb{P}(\rho(S_h^{\Sigma}) = g \mid A_0^{\Sigma} = a, H_0^{\Sigma} = h', G_0^{\Sigma} = g', S_0^{\Sigma} = s, l(\Sigma) = h; \pi_n) \\ &\quad \cdot \mathbb{P}(A_0^{\Sigma} = a \mid H_0^{\Sigma} = h', G_0^{\Sigma} = g', S_0^{\Sigma} = s, l(\Sigma) = h; \pi_n) \\ &\quad \cdot \mathbb{P}(H_0^{\Sigma} = h', G_0^{\Sigma} = g' \mid S_0^{\Sigma} = s, l(\Sigma) = h; \pi_n) \\ &= \sum_{h' \ge h, g' \in \mathcal{G}} \mathbb{P}(\rho(S_h) = g \mid A_0 = a, H_0 = h', G_0^{\Sigma} = g', S_0 = s; \pi_n) \\ &\quad \cdot \pi_n(a \mid s, h', g') \mathbb{P}(H_0^{\Sigma} = h', G_0^{\Sigma} = g' \mid S_0^{\Sigma} = s, l(\Sigma) = h; \pi_n), \end{split}$$

where in the last equality we applied the segment distribution properties (2.5) and (2.6).

Lemma 3, Point 2. We have that

$$\begin{split} \pi_{n+1}^{\text{trail}}(a \mid s, h, g) &= \mathbb{P}(A_0^{\Sigma} = a \mid S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g, l(\Sigma) = H_0^{\Sigma}; \pi_n^{\text{trail}}) \\ &= \mathbb{P}(A_0^{\Sigma} = a, H_0^{\Sigma} = h \mid S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g, l(\Sigma) = H_0^{\Sigma}; \pi_n^{\text{trail}}). \end{split}$$

The first equality is a consequence of (2.7), where the condition $l(\Sigma) = H_0^{\Sigma}$ is added to restrict the equation to Seg^{trail}. For the second equality we made use of the implication $(l(\Sigma) = h) \land (l(\Sigma) = H_0^{\Sigma}) \Longrightarrow H_0^{\Sigma} = h$. Reasoning as above we apply Bayes' rule, where we assume $\mathbb{P}(S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g, l(\Sigma) = H_0^{\Sigma}; \pi_n^{\text{trail}}) > 0$. Subsequently we marginalize over H_0^{Σ} and G_0^{Σ} , apply the product rule and make use of the implication $(l(\Sigma) = h) \land (H_0^{\Sigma} = h) \Longrightarrow l(\Sigma) = H_0^{\Sigma}$ to eliminate the event $l(\Sigma) = H_0^{\Sigma}$. We obtain

$$\begin{split} \pi_{n+1}^{\text{trail}}(a \mid s, h, g) &\propto \mathbb{P}(\rho(S_h^{\Sigma}) = g, A_0^{\Sigma} = a, H_0^{\Sigma} = h \mid S_0^{\Sigma} = s, l(\Sigma) = h, l(\Sigma) = H_0^{\Sigma}; \pi_n^{\text{trail}}) \\ &= \sum_{g' \in \mathcal{G}} \mathbb{P}(\rho(S_h^{\Sigma}) = g, A_0^{\Sigma} = a, H_0^{\Sigma} = h, G_0^{\Sigma} = g' \mid S_0^{\Sigma} = s, l(\Sigma) = h, l(\Sigma) = H_0^{\Sigma}; \pi_n^{\text{trail}}) \\ &= \sum_{g' \in \mathcal{G}} \mathbb{P}(\rho(S_h^{\Sigma}) = g \mid A_0^{\Sigma} = a, H_0^{\Sigma} = h, G_0^{\Sigma} = g', S_0^{\Sigma} = s, l(\Sigma) = h, l(\Sigma) = H_0^{\Sigma}; \pi_n^{\text{trail}}) \\ &\quad \cdot \mathbb{P}(A_0^{\Sigma} = a \mid H_0^{\Sigma} = h, G_0^{\Sigma} = g', S_0^{\Sigma} = s, l(\Sigma) = h, l(\Sigma) = H_0^{\Sigma}; \pi_n^{\text{trail}}) \\ &\quad \cdot \mathbb{P}(H_0^{\Sigma} = h, G_0^{\Sigma} = g', l(\Sigma) = H_0^{\Sigma} \mid S_0^{\Sigma} = s, l(\Sigma) = h; \pi_n^{\text{trail}}) \\ &= \sum_{g' \in \mathcal{G}} \mathbb{P}(\rho(S_h) = g \mid A_0 = a, H_0 = h, G_0 = g', S_0 = s; \pi_n^{\text{trail}}) \\ &\quad \cdot \pi_n^{\text{trail}}(a \mid s, h, g') \mathbb{P}(H_0^{\Sigma} = h, G_0^{\Sigma} = g' \mid S_0^{\Sigma} = s, l(\Sigma) = h; \pi_n^{\text{trail}}). \end{split}$$

Lemma 3, Point 3. We have that

$$\begin{split} \pi_{n+1}^{\text{diag}}(a \mid s, h, g) \\ &= \mathbb{P}(A_0^{\Sigma} = a \mid S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g, \rho(S_h^{\Sigma}) = G_0^{\Sigma}, l(\Sigma) = H_0^{\Sigma}; \pi_n^{\text{diag}}) \\ &= \mathbb{P}(A_0^{\Sigma} = a, H_0^{\Sigma} = h, G_0^{\Sigma} = g \mid S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g, \rho(S_h^{\Sigma}) = G_0^{\Sigma}, l(\Sigma) = H_0^{\Sigma}; \pi_n^{\text{diag}}) \end{split}$$

The first equality is a consequence of (2.7), where the conditions $l(\Sigma) = H_0^{\Sigma}$ and $\rho(S_h^{\Sigma}) = G_0^{\Sigma}$ are added to restrict the equation to $\operatorname{Seg^{diag}}$. For the second equality we made use of the implications $(l(\Sigma) = h) \wedge (l(\Sigma) = H_0^{\Sigma}) \Longrightarrow H_0^{\Sigma} = h$ and $(\rho(S_h^{\Sigma}) = g) \wedge (\rho(S_h^{\Sigma}) = G_0^{\Sigma}) \Longrightarrow G_0^{\Sigma} = g$. Reasoning as above we apply Bayes' rule, where we assume $\mathbb{P}(S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g, \rho(S_h^{\Sigma}) = G_0^{\Sigma}, l(\Sigma) = H_0^{\Sigma}; \pi_n^{\operatorname{diag}}) > 0$. Subsequently we marginalize over H_0^{Σ} and G_0^{Σ} , apply the product rule and make use of the implications $(\rho(S_h^{\Sigma}) = g) \wedge (G_0^{\Sigma} = g) \Longrightarrow \rho(S_h^{\Sigma}) = G_0^{\Sigma}$ and $(l(\Sigma) = h) \wedge (H_0^{\Sigma} = h) \Longrightarrow l(\Sigma) = H_0^{\Sigma}$ to eliminate the events $\rho(S_h^{\Sigma}) = G_0^{\Sigma}$ and $l(\Sigma) = H_0^{\Sigma}$. We obtain

$$\begin{split} \pi_{n+1}^{\mathrm{diag}}(a \mid s, h, g) \\ \propto \mathbb{P}(\rho(S_h^{\Sigma}) = g, \rho(S_h^{\Sigma}) = G_0^{\Sigma}, A_0^{\Sigma} = a, H_0^{\Sigma} = h, G_0^{\Sigma} = g \mid S_0^{\Sigma} = s, l(\Sigma) = h, l(\Sigma) = H_0^{\Sigma}; \pi_n^{\mathrm{diag}}) \\ = \mathbb{P}(\rho(S_h^{\Sigma}) = g \mid A_0^{\Sigma} = a, H_0^{\Sigma} = h, G_0^{\Sigma} = g, S_0^{\Sigma} = s, l(\Sigma) = h, l(\Sigma) = H_0^{\Sigma}; \pi_n^{\mathrm{diag}}) \\ \cdot \mathbb{P}(A_0^{\Sigma} = a \mid H_0^{\Sigma} = h, G_0^{\Sigma} = g, S_0^{\Sigma} = s, l(\Sigma) = h, l(\Sigma) = H_0^{\Sigma}; \pi_n^{\mathrm{diag}}) \\ \cdot \mathbb{P}(H_0^{\Sigma} = h, G_0^{\Sigma} = g, l(\Sigma) = H_0^{\Sigma} \mid S_0^{\Sigma} = s, l(\Sigma) = h; \pi_n^{\mathrm{diag}}). \end{split}$$

Finally, making use of the expression (2.1) for the action-value functions we obtain

$$\begin{aligned} \pi_{n+1}^{\text{diag}}(a \mid s, h, g) \propto Q^{\pi_n^{\text{diag}}}((s, h, g), a) \pi_n^{\text{diag}}(a \mid s, h, g) \mathbb{P}(H_0^{\Sigma} = h, G_0^{\Sigma} = g | S_0^{\Sigma} = s, l(\Sigma) = h; \pi_n^{\text{diag}}) \\ \propto Q^{\pi_n^{\text{diag}}}((s, h, g), a) \pi_n^{\text{diag}}(a \mid s, h, g). \end{aligned}$$

3.2 The Relation of eUDRL to Reward-Weighted Regression

Reward-Weighted Regression (RWR, see (Dayan and Hinton, 1997), (Peters and Schaal, 2007), (Peng et al., 2019)) is an RL algorithm that, similarly to eUDRL, relies on a recursive sequence of policy updates. As in eUDRL a batch of trajectories is generated following the current policy π_n and the subsequent policy π_{n+1} is fitted using supervised learning on the sampled trajectories but the contribution of actions is weighted by trajectory returns. In other words RWR goes beyond vanilla imitation learning on sampled trajectories in that it weights the relevance of trajectories. RWR is characterized by update formulas of the form $\pi_{n+1}(a \mid s) \propto Q(s, a)\pi_n(a \mid s)$, see Štrupl et al. (2022b) for details. In the context of CE this translates to the following update rule

$$\pi_{n+1}(a \mid s, h, g) \propto Q^{\pi_n}((s, h, g), a) \pi_n(a \mid s, h, g).$$

In view of Lemma 3, point 3. this corresponds to the update rule of eUDRL in the set Seg^{diag} . In other words in this special situation eUDRL becomes RWR. Notice, however, that contrary to general eUDRL the RWR recursion always converges² to optimal policies Štrupl et al. (2022b). On the other hand, operating on Seg^{diag} instead of Seg, RWR uses a far smaller set of samples, which leads to a slower overall learning process. We will not rely on the convergence of RWR in our continuity proofs, but the intuition that "focusing solely on diagonal segments is sufficient" will guide our discussion.

3.3 The Role of Prioritized Replay

As an application of the described correspondence of eUDRL and RWR, we study the role of prioritized replay in eUDRL. The eUDRL implementation of Srivastava et al. (2019) employs a replay buffer, where initial goals or returns-to-go are sampled from such trajectories in the buffer that correspond to a high quantile of returns. The following examples illustrate that prioritization by return can both increase and decrease the performance of the learning system.

Performance deterioration through Prioritized replay: Consider an environment that is characterized by the following features:

i) High returns are hard to reach: Even for optimal sequences of action (in the sense of the CE) the probability of obtaining a high return is small.

ii) Moderate returns are easy to reach.

iii) Attempting to reach a high return entails sequences of actions that can lead to 'dead ends', i.e. low returns.

In this setup it might occur that the preference of the buffer for choosing high returns might lead to small expected returns. Choosing trajectories with high returns from the replay buffer might thus obfuscate the exploration of moderate returns that might have a higher expected value.

^{2.} Although Štrupl et al. (2022b) assumes positive rewards, while rewards are non-negative in the article at hand, a generalization of Štrupl et al. (2022b) is straightforward.

Performance improvement through Prioritized replay: Consider an environment that is characterized by the following features:

- i) There are two possible returns $\{0, 1\}$.
- *ii)* There are only trailing segments, i.e. segments are aligned with the end of the trajectory. In this setup the return-prioritized buffer is filled over time with trajectories of return

1. As a consequence sampling results in the same initial goals or returns-to-go of 1 and the buffer will be filled with trajectories with initial goal or return-to-go of 1 and realized return of 1. Together with the assumption that segments are trailing, it follows that the algorithm operates on diagonal segments. This effectively transforms the algorithm to RWR, which is proven to converge.

4 The Continuity of eUDRL in the Transition Kernel and Stability Properties

We investigate the continuity of various eUDRL-generated quantities like policies, associated values and the goal-reaching objective under small changes of the transition kernel. The established continuity results will be used to generalize the known optimality of eUDRL when kernels are deterministic (see section 5 below) to the case of near optimality of eUDRL (i.e. optimality up to a fixed error) when kernels are nearly deterministic (i.e. the kernels are located in a neighborhood of a deterministic kernel).

4.1 Compatible Families and the Stability of Supports

Let \mathcal{M} be a given MDP and let $\overline{\mathcal{M}}$ denote its CE. Given a state and an action, one can view the transition kernel $\lambda(\cdot | s, a)$ as a vector in a *probability simplex* ΔS , i.e., one can write $\lambda \in (\Delta S)^{S \times \mathcal{A}} \subset \mathbb{R}^{S \times S \times \mathcal{A}}$. We study how properties of \mathcal{M} and $\overline{\mathcal{M}}$, such as eUDRLgenerated policies and values depend on λ . For our purpose it is sufficient to study the sensitivity with respect to changes of λ (and $\overline{\lambda}$, respectively), where other components of the MDP and its CE remain fixed. To this end we define the notion of *compatible families* of MDPs.

Definition 4 (Compatible families of MDPs) Let λ_0 be a given transition kernel, and let $\mathcal{M} = (\mathcal{S}, \mathcal{A}, \lambda_0, \mu, r)$ be a respective MDP with CE $\overline{\mathcal{M}} = (\overline{\mathcal{S}}, \mathcal{A}, \overline{\lambda_0}, \overline{\mu}, \overline{r}, \rho)$. Write $\{\mathcal{M}_{\lambda} = (\mathcal{S}, \mathcal{A}, \lambda, \mu, r) | \lambda \in (\Delta \mathcal{S})^{\mathcal{S} \times \mathcal{A}} \}$ for the family of MDPs, whose kernel belongs to the same product of simplexes $(\Delta \mathcal{S})^{\mathcal{S} \times \mathcal{A}}$ as λ_0 . Similarly, write $\{\overline{\mathcal{M}}_{\lambda} = (\overline{\mathcal{S}}, \mathcal{A}, \overline{\lambda}, \overline{\mu}, \overline{r}, \rho) | \lambda \in (\Delta \mathcal{S})^{\mathcal{S} \times \mathcal{A}} \}$ for the family of the respective CEs. We will refer to any pair of MDP families resulting from the above construction as compatible families.

Hereafter we will reserve the subscript λ to refer to quantities that stem from the MDPs \mathcal{M}_{λ} or $\bar{\mathcal{M}}_{\lambda}$ of given compatible families. We will occasionally add the subscript λ to emphasize that a certain quantity (such as a family of policies) is associated with a family of MDPs.

The following definition summarizes some notions that will play an important role in our proofs.

Definition 5 Let $\{\mathcal{M}_{\lambda}|\lambda \in (\Delta S)^{S \times A}\}$ and $\{\bar{\mathcal{M}}_{\lambda}|\lambda \in (\Delta S)^{S \times A}\}$ be compatible families.

1. (Numerator and denominator in the eUDRL recursion) For all $a \in \mathcal{A}$ and all $(s, h, g) \in \overline{S}_T$ let $\operatorname{num}_{\lambda,\pi}(a, s, h, g)$ (numerator) and $\operatorname{den}_{\lambda,\pi}(s, h, g)$ (denominator) be defined by

$$\operatorname{num}_{\lambda,\pi}(a,s,h,g) = \mathbb{P}_{\lambda}(A_0^{\Sigma} = a, S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g; \pi),$$
$$\operatorname{den}_{\lambda,\pi}(s,h,g) = \mathbb{P}_{\lambda}(S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g; \pi).$$

2. (The CE state visitation distribution) The CE state visitation distribution for transition kernel λ and policy π is defined for all $\bar{s} \in \bar{S}$ by³

$$\nu_{\lambda,\pi}(\bar{s}) = \frac{\sum_{t < N} \mathbb{P}_{\lambda}(\bar{S}_t = \bar{s}, \bar{S}_t \in \bar{S}_T)}{\sum_{\bar{s} \in \bar{S}} \sum_{t < N} \mathbb{P}_{\lambda}(\bar{S}_t = \bar{s}, \bar{S}_t \in \bar{S}_T)}$$

^{3.} Here we can refer to Sutton and Barto (2018) equations (9.2) and (9.3) with addition that one has to restrict the computation only to transient states.

A state is called visited under λ and π if and only if $\nu_{\lambda,\pi}(\bar{s}) > 0$.

3. (The set of critical states) Let $\lambda_0 \in (\Delta S)^{S \times A}$ be a deterministic kernel and $\pi_0 > 0$ a policy. The set of critical states (for eUDRL learning) is

$$\mathcal{S}_{\lambda_0} = \operatorname{supp} \operatorname{den}_{\lambda_0, \pi_0} \cap \operatorname{supp} \nu_{\lambda_0, \pi_0}.$$

Notice that the particular choice of $\pi_0 > 0$ does not matter.

Using the numerator and denominator defined above the eUDRL recursion (2.8) can be written (for any $\pi_0 \in (\Delta \mathcal{A})^S$) as

$$\pi_{n+1}(a|s,h,g) = \frac{\operatorname{num}_{\lambda,\pi_n}(a,s,h,g)}{\operatorname{den}_{\lambda,\pi_n}(s,h,g)} \quad \text{for } (s,h,g) \in \operatorname{supp den}_{\lambda,\pi_n}, \tag{4.1}$$

where we set⁴ $\pi_{n+1}(a|s, h, g) = 1/|\mathcal{A}|$ outside the support supp den_{λ,π_n}. The state visitation distribution is restricted to the transient states because this is the set of states where the learned policy affects the evolution of the CE. The set of all visited states equals to the support of $\nu_{\lambda,\pi}$. The support can be written shortly as

$$\operatorname{supp} \nu_{\lambda,\pi} = \{ \bar{s} \in \bar{S}_T ; (\exists t < N) : \mathbb{P}_{\lambda}(\bar{S}_t = \bar{s}; \pi) > 0 \}.$$

While it is evident from the recursion equation that π_n depends on λ for n > 1, we assume that the initial π_0 is constant through the family of MDPs. Notice that the definition of "critical states" \bar{S}_{λ_0} depends on the specified family through the set $(\Delta S)^{S \times A}$. The definition of \bar{S}_{λ_0} is motivated by the fact that the non-trivial behavior of our continuity discussion occurs on these sets of states. In fact the chosen intersection of states ensures the following criteria:

i) the command can actually be satisfied (through $\mathbb{P}_{\lambda_0}(S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g; \pi_0) > 0$ on supp den_{λ_0, π_0}) and

ii) critical states have a non-zero visitation probability (on supp ν_{λ_0,π_0}).

In view of these points, inspecting formula (2.2) yields that given a fixed kernel λ_0 , the states that are outside \bar{S}_{λ_0} do not contribute to the goal-reaching objective. We will discuss rigorously in section 6 that policies and values in other states are not relevant for establishing the continuity of the goal-reaching objective.

The following lemma reveals some nice stability properties of S_{λ_0} that will be useful later. In particular, we demonstrate that \bar{S}_{λ_0} is stable under small perturbations of the transitions kernel λ within a fixed compatible family and also through the course of the eUDRL iteration process. For a $\delta > 0$ we write

$$U_{\delta}(\lambda_0) = \left\{ \lambda \mid \max_{(s,a) \in \mathcal{S} \times \mathcal{A}} \|\lambda(\cdot|s,a) - \lambda_0(\cdot|s,a)\|_1 < \delta \right\}$$

for the δ -neighborhood in composite max-1 norm.

^{4.} It is worth noting that our continuity proofs will not depend on the concrete way of defining π_{n+1} for $(s, h, g) \notin \operatorname{supp} \operatorname{den}_{\lambda, \pi_n}$. The discontinuities that we will present cannot be removed by an alternative definition. See the discussion below example 1 in the appendix B for details.

Lemma 6 (Stability of \bar{S}_{λ_0}) Let $\{\mathcal{M}_{\lambda}|\lambda \in (\Delta S)^{S \times A}\}$ and $\{\bar{\mathcal{M}}_{\lambda}|\lambda \in (\Delta S)^{S \times A}\}$ be compatible families. Let $(\pi_{n,\lambda})_{n\geq 0}$ be a sequence of policies generated by the eUDRL iteration given a transition kernel $\lambda \in (\Delta S)^{S \times A}$ and an initial policy π_0 (that does not depend on λ). Fix a deterministic transition kernel $\lambda_0 \in (\Delta S)^{S \times A}$ then for all initial conditions $\pi_0 > 0$ it holds:

- 1. For all $n \geq 0$ and all $\lambda \in U_2(\lambda_0)$ we have that $\operatorname{supp num}_{\lambda_0,\pi_0} \cap (\mathcal{A} \times \operatorname{supp} \nu_{\lambda_0,\pi_0}) \subset \operatorname{supp num}_{\lambda,\pi_n} \cap (\mathcal{A} \times \operatorname{supp} \nu_{\lambda,\pi_n})$, where the inclusion becomes equality for $\lambda = \lambda_0$.
- 2. For all $n \ge 0$ and all $\lambda \in U_2(\lambda_0)$ we have that $\bar{S}_{\lambda_0} \subset \operatorname{supp} \operatorname{den}_{\lambda,\pi_n} \cap \operatorname{supp} \nu_{\lambda,\pi_n}$, where the inclusion becomes an equality for $\lambda = \lambda_0$.
- 3. For all $n \ge 0$ and all $\lambda \in U_2(\lambda_0)$ we have that

$$\mathbb{P}_{\lambda}(S_0^{\Sigma}=s, l(\Sigma)=h, \rho(S_h^{\Sigma})=g, H_0^{\Sigma}=h, G_0^{\Sigma}=g; \pi_n)>0$$

for all $(s, h, g) \in \overline{S}_{\lambda_0}$.

- 4. For all $(a, s, h, g) \in \operatorname{supp num}_{\lambda_0, \pi_0} \cap (\mathcal{A} \times \operatorname{supp} \nu_{\lambda_0, \pi_0}), h > 1$ there exists an $s' \in \mathcal{S}$ so that $\lambda_0(s'|s, a) > 0$ and $(s', h 1, g) \in \overline{S}_{\lambda_0}$.
- 5. For all $(a, s, h, g) \notin \operatorname{supp num}_{\lambda_0, \pi_0}$ with $(s, h, g) \in \operatorname{supp den}_{\lambda_0, \pi_0}$ we have for any policy π that $Q_{\lambda_0}^{\pi}((s, h, g), a) = 0$.

It is straightforward to extend the lemma for arbitrary $\lambda_0 \in (\Delta S)^{S \times A}$, but we will need the result only for deterministic λ_0 .

Proof Assuming that λ_0 is deterministic, the following inclusion holds:

$$\forall \lambda \in U_2(\lambda_0)) : \operatorname{supp} \lambda_0 \subset \operatorname{supp} \lambda. \tag{4.2}$$

This means that any state that is reachable through an environment transition by the kernel λ_0 is also reachable by λ . Since λ_0 is deterministic there exists a unique s' such that $1 = \lambda_0(s'|s, a)$. It suffices to show that $\lambda(s'|s, a)$ is non-zero. We have

$$2 > \|\lambda(\cdot|s,a) - \lambda_0(\cdot|s,a)\|_1 = |\lambda(s'|s,a) - \lambda_0(s'|s,a)| + \sum_{s'' \neq s'} |\lambda(s''|s,a) - \lambda_0(s''|s,a)|$$
$$= 1 - \lambda(s'|s,a) + \sum_{s'' \neq s'} \lambda(s''|s,a) = 2(1 - \lambda(s'|s,a))$$

which is equivalent to $\lambda(s'|s, a) > 0$.

1. Assuming $(a, s, h, g) \in \text{supp num}_{\lambda_0, \pi_0} \cap (\mathcal{A} \times \text{supp } \nu_{\lambda_0, \pi_0})$, it follows that there exists a trajectory $\tau' = ((s'_0, H', g'), a'_0, (s'_1, H' - 1, g'), a'_1, \ldots)$ with positive probability $\mathbb{P}_{\lambda_0}(\mathcal{T} = \tau'; \pi_0) > 0$ with the following property: for a certain $t' \leq N-h$ we have $s'_{t'} = s, a'_{t'} = a, H' - t' \geq h$ and $\rho(s_{t'+h}) = g$. We claim that there exists a trajectory $\tau = ((s_0, H, g), a_0, (s_1, H - 1, g), \ldots)$ and $t \leq N$ with positive probability $\mathbb{P}_{\lambda_0}(\mathcal{T} = \tau; \pi_0) > 0$ with the property that $s_t = s, a_t = a, h = H - t$ and $\rho(s_{t+h}) = g$. The existence of a beginning segment of a trajectory (so-called prefix) $((s_0, H, g), a_0, \ldots, (s_t, H - t, g))$ that fits τ at t and has positive probability (under λ_0, π_0) follows from $(s_t, H - t, g) = (s, h, g) \in \overline{S}_{\lambda_0} \subset \text{supp } \nu_{\lambda_0, \pi_0}$. For the terminal segment of the trajectory (the so-called suffix) we take the suffix of τ' (starting at t') and adjust the horizon and goal components so that they coincide with those of τ at t. The entire trajectory will still have positive probability (under λ_0, π_0), i.e., $\mathbb{P}_{\lambda_0}(A_t = a_t, \bar{S}_{t+1} = (s_{t+1}, H - t + 1, g), A_{t+1} = a_{t+1}, \dots |\bar{S}_t = (s_t, H - t, g); \pi_0) > 0$, since we kept original MDP state components and we can use transitions from $\lambda_0(s'_{t'+i+1}|s'_{t'+i}, a'_{t'+i}), 0 \le i \le h - 1$ and $\pi_0 > 0$. This proves the existence of τ .

Making use of the inclusion (4.2) we have $\mathbb{P}_{\lambda}(\mathcal{T} = \tau; \pi_0) > 0$ for all $\lambda \in U_{\delta}(\lambda_0)$. Assuming $\lambda \in U_{\delta}(\lambda_0)$ fixed, we prove by induction that

$$\mathbb{P}_{\lambda}(\mathcal{T}=\tau;\pi_n) > 0. \tag{4.3}$$

The statement for n = 0 is established already. Assume that it holds for $n \ge 0$. Because the probability of τ is positive (under λ, π_n) we get $\forall i : \mathbb{P}_{\lambda}(A_0^{\Sigma} = a_i, S_0^{\Sigma} = s_i, l(\sigma) =$ $H - i, \rho(S_{H-i}^{\Sigma}) = g; \pi_n) > 0$. It follows that $\pi_{n+1}(a_i|s_i, H - i, g) = \mathbb{P}_{\lambda}(A_0^{\Sigma} = a_i|S_0^{\Sigma} =$ $s_i, l(\sigma) = H - i, \rho(S_{H-i}^{\Sigma}) = g; \pi_n) > 0$ for all i < H. Since the transition kernel λ and the initial distribution μ remain unchanged $\mathbb{P}_{\lambda}(\mathcal{T} = \tau; \pi_{n+1}) > 0$ follows.

From (4.3) follows that $\operatorname{num}_{\lambda,\pi_n}(a,s,h,g) > 0$ and $\nu_{\lambda,\pi_n}(s,h,g) > 0$, also we see that $\mathbb{P}_{\lambda}(A_0^{\Sigma} = a, S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g, H_0^{\Sigma} = h, G_0^{\Sigma} = g; \pi_n) > 0$. This finishes the proof of the inclusion. To establish equality in case of $\lambda = \lambda_0$ it suffices to show supp $\operatorname{num}_{\lambda_0,\pi_n} \subset \operatorname{supp} \operatorname{num}_{\lambda_0,\pi_0}$ and $\operatorname{supp} \nu_{\lambda_0,\pi_n} \subset \operatorname{supp} \nu_{\lambda_0,\pi_0}$. But this follows since it is assumed that π_0 has maximal support.

2. This follows immediately from point 1. and the fact that for any λ we have

 $(\forall n \ge 0, \forall (s, h, g) \in \bar{S}_T) : (s, h, g) \in \operatorname{supp} \operatorname{den}_{\lambda, \pi_n} \iff (\exists a \in \mathcal{A} : (a, s, h, g) \in \operatorname{supp} \operatorname{num}_{\lambda, \pi_n}).$

3. The fact that $\mathbb{P}_{\lambda}(S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g; H_0^{\Sigma} = h, G_0^{\Sigma} = g, \pi_n) > 0$ follows from inequality (4.3).

4. This follows immediately from existence of the trajectory τ constructed in the proof of point 1.

5. For the sake of contradiction assume $Q_{\lambda_0}^{\pi}((s,h,g),a) > 0$ for some π . By the maximality of the support of π_0 it follows that $Q_{\lambda_0}^{\pi_0}((s,h,g),a) > 0$. From $(s,h,g) \in$ supp den $_{\lambda_0,\pi_0}$ and the equivalence used in the proof of point 2. above we deduce the existence of a trajectory τ' with positive probability (under λ_0, π_0) (as in point 1. above). We will need just its prefix $\mathbb{P}_{\lambda_0}(S_0 = s'_0, H_0 = H', G_0 = g', \ldots S_{t'} = s, H_{t'} = h', G_{t'} = g'; \pi_0) > 0$, $h' \geq h$. Since $0 < Q_{\lambda_0}^{\pi_0}((s,h,g),a) = \mathbb{P}_{\lambda_0}(\rho(S_h) = g|S_0 = s, H_0 = h, G_0 = g, A_0 = a; \pi_0)$ it follows that $\mathbb{P}_{\lambda_0}(\rho(S_{t'+h}) = g|S_{t'} = s, H_{t'} = h', G_{t'} = g', A_{t'} = a; \pi_0) > 0$, where we used that support of π_0 neglects horizon and goal components. Similarly it holds $\pi_0(a|s,h',g') > 0$. Putting this together we conclude that there exists a trajectory $((s'_0, H', g'), \ldots, (s, h', g'), a, \ldots (s_{t'+h}, h' - h, g'), \ldots)$ with $\rho(s_{t'+h}) = g$ with positive probability (under λ_0, π_0) which demonstrates that $\operatorname{num}_{\lambda_0,\pi_0}(a, s, h, g) > 0$ and contradicts our assumption.

4.2 On the Stability of Supports in Segment Sub-Spaces

The discussion of stability of sets of critical states in compatible families presented in section 4.1 translate mutatis mutandis to a discussion of the eUDRL recursion on segment sub-spaces Seg^{trail} and Seg^{diag}. Motivated by the theoretical description of eUDRL-type iterations in algorithms like ODT or RWR we briefly outline the respective stability properties in this section. Let $\operatorname{num}_{\lambda,\pi}^{\operatorname{trail/diag}}$ and $\operatorname{den}_{\lambda,\pi}^{\operatorname{trail/diag}}$ denote the numerator and denominator of the eUDRL recursion (compare definition 4.1). The quantities are modified by introducing conditions that define the respective segment sub-spaces. For example, for all $a \in \mathcal{A}$ and all $(s, h, g) \in \overline{S}_T$ we set

$$\begin{split} \operatorname{num}_{\lambda,\pi}^{\operatorname{diag}}(a,s,h,g) &:= \mathbb{P}_{\lambda}(A_{0}^{\Sigma} = a, S_{0}^{\Sigma} = s, l(\Sigma) = h, S_{h}^{\Sigma} = g | l(\Sigma) = H_{0}^{\Sigma}, \rho(S_{l(\Sigma)}^{\Sigma}) = G_{0}^{\Sigma}; \pi), \\ \operatorname{den}_{\lambda,\pi}^{\operatorname{diag}}(s,h,g) &:= \mathbb{P}_{\lambda}(S_{0}^{\Sigma} = s, l(\Sigma) = h, S_{h}^{\Sigma} = g | l(\Sigma) = H_{0}^{\Sigma}, \rho(S_{l(\Sigma)}^{\Sigma}) = G_{0}^{\Sigma}; \pi). \end{split}$$

The eUDRL recursion on Seg^{trail} and Seg^{diag} has the form as equation (4.1) with the renaming $\pi_n \to \pi_n^{\text{diag/trail}}$, num \to num^{diag/trail} and den \to den^{diag/trail}.

Lemma 7 (Stability of \bar{S}_{λ_0} in trailing and diagonal sub-spaces) In the setting of Lemma 6 consider restrictions of the eUDRL recursion to the sets Seg^{diag} and Seg^{diag}. Then the set of critical states \bar{S}_{λ_0} remains unchanged and the conclusion of Lemma 6 remain valid under the renaming $\pi_n \to \pi_n^{\text{diag/trail}}$, num \to num^{diag/trail} and den \to den^{diag/trail}.

Proof We follow the proof of Lemma 6 and only highlight the differences. Consider any $(a, s, h, g) \in \operatorname{supp} \operatorname{num}_{\lambda_0, \pi_0}^{\operatorname{diag/trail}} \cap (\mathcal{A} \times \operatorname{supp} \nu_{\lambda_0, \pi_0})$ that is contained in a trajectory τ' . We construct a trajectory τ with positive probability that contains (a, s, h, g) in a diagonal segment. As the diagonal/trailing segment is also contained in Seg, we can take the same approach as before. As a byproduct of the construction of τ we conclude that

$$\operatorname{supp\,num}_{\lambda_0,\pi_0} \cap (\mathcal{A} \times \operatorname{supp} \nu_{\lambda_0,\pi_0}) = \operatorname{supp\,num}_{\lambda_0,\pi_0}^{\operatorname{diag/trail}} \cap (\mathcal{A} \times \operatorname{supp} \nu_{\lambda_0,\pi_0}), \qquad (4.4)$$

and consequently

$$\bar{\mathcal{S}}_{\lambda_0} = \operatorname{supp} \operatorname{den}_{\lambda_0, \pi_0} \cap \operatorname{supp} \nu_{\lambda_0, \pi_0} = \operatorname{supp} \operatorname{den}_{\lambda_0, \pi_0}^{\operatorname{diag/trail}} \cap \operatorname{supp} \nu_{\lambda_0, \pi_0}.$$
(4.5)

In other words the set \bar{S}_{λ_0} and its state-action variant $\operatorname{num}_{\lambda_0,\pi_0} \cap (\mathcal{A} \times \operatorname{supp} \nu_{\lambda_0,\pi_0})$ remain unchanged. To prove that $\mathbb{P}_{\lambda}(\mathcal{T} = \tau; \pi_n^{\operatorname{diag/trail}}) > 0$ it suffice to realize that the segments used to prove $\pi_{n+1}(a_i|s_i, H-i, g) > 0$ were diagonal, i.e. $\pi_{n+1}^{\operatorname{diag/trail}}(a_i|s_i, H-i, g) > 0$. The rest of point 1. and points 2.,3.,4. follow similarly. In the proof of point 5. the fact that we began with a diagonal/trailing segment is employed to demonstrate that $\operatorname{num}_{\lambda_0,\pi_0}^{\operatorname{diag/trail}}(a, s, h, g) > 0$ and to obtain a contradiction.

5 Convergence to Optimal Policies for Deterministic Kernels

While the eUDRL recursion converges to optimal policies for deterministic environments, it can result in sub-optimal behavior in the presence of stochasticity. This fact is apparent already in the early paper by Schmidhuber (5 Dec 2019), where the eUDRL recursion in stochastic environments is replaced by an algorithm that operates on expected rather than actual returns. The fact that GCSL converges to an optimal policy is mentioned by Ghosh et al. (2021). A related fact is also discussed by Brandfonbrener et al. (2022), focusing on the first UDRL iteration (offline RL) in return-reaching tasks, where it is proved that intended returns coincide with expected returns in deterministic environments. The fact that GCSL converges to optimal policies is rather straightforward such that a full proof is usually omitted. We provide a complete proof in this article as we will later build on this result and it is convenient to introduce our notation. We begin by characterizing the set of optimal actions for CEs in the case of a deterministic environment in section 5.1. Subsequently, in sections 5.2 and 5.3 we show that this set corresponds to the support of the policies emerging in the course of the eUDRL recursion. As a consequence the support of π_n is constant for $n \ge 1$, which demonstrates that eUDRL-generated policies are optimal in deterministic environments for $n \ge 1$.

5.1 Optimal Actions for Deterministic Transition Kernels

The goal of this section is to identify the sets of optimal actions for CEs arising form MDPs with a given deterministic transition kernel. We begin by defining the notion of an optimal action.

Definition 8 (Optimal actions) Let $\mathcal{M} = (\mathcal{S}, \mathcal{A}, \lambda, \mu, r)$ be an MDP. An action $a \in \mathcal{A}$ is called optimal in the state $s \in \mathcal{S}$ if and only if there exists a policy π with $\pi(a|s) > 0$ and π reaches maximal (optimal) value in s. We write $\mathcal{O}(s)$ for the set of optimal actions at s.

Notice that, since CE is a particular MDP, this definition includes optimal actions in the context of CEs. Accordingly, in the context of CEs, any action that maximizes the probability of achieving the intended command is optimal.

Lemma 9 (The set of optimal actions of CEs for deterministic kernels) Let λ_0 be a given deterministic transition kernel and let $\mathcal{M} = (\mathcal{S}, \mathcal{A}, \lambda_0, \mu, r)$ be a respective MDP with CE $\overline{\mathcal{M}} = (\overline{\mathcal{S}}, \mathcal{A}, \overline{\lambda_0}, \overline{\mu}, \overline{r}, \rho)$ and consider a policy $\pi_0 > 0$. For any state $\overline{s} \in \operatorname{supp} \operatorname{den}_{\lambda_0, \pi_0}(\cdot, \overline{s})$ the set of optimal actions is $\mathcal{O}(\overline{s}) = \operatorname{supp} \operatorname{num}_{\lambda_0, \pi_0}(\cdot, \overline{s})$.

Lemma 9 identifies the set of optimal actions for deterministic kernels for general CEbased algorithm but, it has no reference to eUDRL specifically. Notice that the occurring quantities are related to the segment space Seg, which can be constructed without resorting to eUDRL. In our subsequent discussion of continuity we will investigate the impact of changing λ around some fixed deterministic kernel λ_0 . We will drop the index $\mathcal{O}_{\lambda_0}(\bar{s})$ in our notation, writing $\mathcal{O}(\bar{s})$ to denote optimal actions at fixed λ_0 (we will not investigate $\mathcal{O}_{\lambda}(\bar{s})$ in situations that the kernel varies).

Proof We begin with the reverse implication. Fix a state $\bar{s} \in \text{supp den}_{\lambda_0,\pi_0}$ and an action $a \in \mathcal{O}(\bar{s})$. Since $\text{num}_{\lambda_0,\pi_0}(a,\bar{s}) > 0$ there exists a trajectory $\tau' = ((s_0, H', g'), a_0, (s_1, H' - C_0))$

1, g'), a_1, \ldots) with positive probability under π_0 with $s_t = s, a_t = a, \rho(s_{t+h}) = g, t + h \leq H'$. Moreover $\lambda_0(s_{i+1}|s_i, a_i) = 1$ for i < H', since λ_0 is deterministic. Since $Q((s_{t+h-1}, 1, g), a_{t+h-1}) = \sum_{s' \in \rho^{-1}(\{g\})} \lambda_0(s'|s_{t+h-1}, a_{t+h-1}) = \lambda_0(s_{t+h}|s_{t+h-1}, a_{t+h-1}) = 1$ is the maximal value, a_{t+h-1} is optimal in $(s_{t+h-1}, 1, g)$. Therefore, $\pi^*(a_{t+h-1}|s_{t+h-1}, 1, g) = 1$ is an optimal policy in this state. Furthermore,

$$V^{\pi^*}(s_{t+h-1}, 1, g) = \sum_{a \in \mathcal{A}} Q((s_{t+h-1}, 1, g), a)\pi^*(a|s_{t+h-1}, 1, g)$$
$$= Q((s_{t+h-1}, 1, g), a_{t+h-1})\pi^*(a_{t+h-1}|s_{t+h-1}, 1, g) = 1,$$

and

$$Q^{\pi^*}(s_{t+h-2}, 2, g, a_{t+h-2}) = \sum_{s' \in \mathcal{S}} V^{\pi^*}(s', 1, g)\lambda_0(s'|s_{t+h-2}, a_{t+h-2})$$
$$= V^{\pi^*}(s_{t+h-1}, 1, g)\lambda_0(s_{t+h-1}|s_{t+h-2}, a_{t+h-2}) = 1$$

which again achieves the maximal possible value. Therefore, a_{t+h-2} is optimal in $(s_{t+h-2}, 2, g)$. Repeating this procedure h times yields that $Q^{\pi^*}(s_t, h, g, a_t) = 1$, a_t is an optimal action in (s_t, h, g) , and the value is $V^{\pi^*}(s_t, h, g) = 1$.

For the forward implication, let $(s, h, g) \in \text{supp den}_{\lambda_0, \pi_0}$ and suppose a is optimal in (s, h, g). Following the same reasoning as in the proof of the reverse implication presented above, we can observe that (s, h, g) achieves the maximal value of 1. Since a is optimal, there exists a policy π^* that reaches the goal g in h steps from the state s following the action a. Due to the fact that $\pi_0 > 0$, it follows that π_0 also reaches g in h steps from s following a, i.e., $\mathbb{P}_{\lambda_0}(\rho(S_h) = g, A_0 = a|S_0 = s, H_0 = h, G_0 = g; \pi_0) > 0$. Since $(s, h, g) \in \text{supp den}_{\lambda, \pi_0}$, we can similarly as in proof of point 1. of lemma 6 find $\tau' = (s'_0, H', g'), \ldots, (s, h', g'), \ldots$) with $h' \geq h$ with positive probability under λ_0 and π_0 . It follows that $\mathbb{P}_{\lambda_0}(\rho(S_h) = g, A_0 = a|S_0 = s, H_0 = h', G_0 = g'; \pi_0) > 0$. Combining this with the suitable prefix of τ' we conclude the existence of a trajectory that demonstrates that $\max_{\lambda_0,\pi_0}(a, s, h, g) > 0$.

5.2 Convergence of eUDRL in the Space of all Segments

In the following lemma, we utilize the notion of optimal actions to demonstrate that eUDRLgenerated policies $(\pi_n)_{n>1}$ are optimal if the transition kernel is deterministic.

Lemma 10 (Optimality of eUDRL policies for deterministic transition kernels) Let λ_0 be a given deterministic transition kernel and let $\mathcal{M} = (S, \mathcal{A}, \lambda_0, \mu, r)$ be a respective MDP with $CE \ \bar{\mathcal{M}} = (\bar{S}, \mathcal{A}, \bar{\lambda_0}, \bar{\mu}, \bar{r}, \rho)$. Assume $\pi_0 > 0$ and let $(\pi_n)_{n\geq 0}$ be the policy sequence generated by the eUDRL iteration given π_0 and λ_0 . Then it holds:

- 1. For all $n \ge 1$ and all $\bar{s} \in \bar{S}_{\lambda_0}$ the support of $\pi_{n+1}(\cdot|\bar{s})$ is identical to the set of optimal actions $\mathcal{O}(\bar{s})$. In particular, the policy π_n is an optimal policy on \bar{S}_{λ_0} .
- 2. For all $n \geq 1$ and all $\bar{s} \in \bar{S}_{\lambda_0}$ the value is maximal, i.e.,

$$V^{\pi_n}(\bar{s}) = 1$$

3. For all $n \geq 1$, all $\bar{s} \in \bar{S}_{\lambda_0}$, and all $a \in A$, the action-value function is given by

$$Q^{\pi_n}(\bar{s}, a) = \begin{cases} 1 & if a \in \mathcal{O}(\bar{s}), \\ 0 & otherwise. \end{cases}$$

Proof 1. Let $n \geq 0$, $(s, h, g) \in \overline{S}_{\lambda_0}(= \operatorname{supp} \operatorname{den}_{\lambda_0, \pi_n} \cap \operatorname{supp} \nu_{\lambda_0, \pi_n})$. We have $a \in \mathcal{O}(s, h, g) \iff \operatorname{num}_{\lambda_0, \pi_0}(a, s, h, g) > 0$. By Lemma 6, point 1., this is equivalent to $\operatorname{num}_{\lambda_0, \pi_n}(a, s, h, g) > 0$ and it follows

$$\pi_{n+1}(a|s,h,g) = \frac{\operatorname{num}_{\lambda_0,\pi_n}(a,s,h,g)}{\operatorname{den}_{\lambda_0,\pi_n}(s,h,g)} > 0 \iff a \in \mathcal{O}(s,h,g).$$

Thus the support of $\pi_{n+1}(\cdot|s, h, g)$ is identical to the set of optimal actions.

2. Let $n \ge 1$, $(s, h, g) \in \overline{S}_{\lambda_0} \subset \text{supp den}_{\lambda_0, \pi_0}$. As in the proof of lemma 9, $V^*(s, h, g) = 1$ and from point 1. above, we know that π_n is optimal on \overline{S}_{λ_0} . Therefore $V^{\pi_n}(s, h, g) = 1$.

3. Let $n \geq 1$, $(s,h,g) \in \bar{S}_{\lambda_0}$. First assume $a \in \mathcal{O}(s,h,g)$. By lemma 6 point 4. there exists $s' \in S$ so that $\lambda_0(s'|s,a) = 1$ and $(s',h-1,g) \in \bar{S}_{\lambda_0}$. By point 2. above, $V^{\pi_n}(s',h-1,g) = 1$ and so $Q^{\pi_n}((s,h,g),a) = \sum_{s'' \in S} V^{\pi_n}(s'',h-1,g)\lambda(s''|s,a) = 1$. If $a \notin \mathcal{O}(s,h,g)$ the assertion follows from lemma 6 point 5.

5.3 Convergence of eUDRL in Segment Sub-Spaces

In the context of CEs, the set of optimal actions $\mathcal{O}(\bar{s})$, $\bar{s} \in \text{supp} \operatorname{den}_{\lambda_0,\pi_0}$ is linked to the distribution on the segment space Seg through the quantities $\operatorname{num}_{\lambda_0,\pi_0}$ and $\operatorname{den}_{\lambda_0,\pi_0}$, cf. lemma 9. However, from (4.4) we conclude that for all $\bar{s} \in \bar{S}_{\lambda_0}$ it holds that

$$\mathcal{O}(\bar{s}) = \operatorname{supp\,num}_{\lambda_0, \pi_0}^{\operatorname{diag/trail}}(\cdot, \bar{s}), \tag{5.1}$$

such that $\operatorname{supp} \operatorname{num}_{\lambda_0,\pi_0}^{\operatorname{diag/trail}}(\cdot, \bar{s})$ remains the set of optimal actions even when the algorithm operates on $\operatorname{Seg}^{\operatorname{diag/trail}}$ as, e.g., in the case of RWR. The variants of lemma 10 for the mentioned segment sub-spaces remain valid mutatis mutandis.

Lemma 11 (Optimality of eUDRL policies for deterministic kernels in Seg^{diag/trail}) In the setting of Lemma 10 consider restrictions of the eUDRL recursion to the sets Seg^{diag/trail}. The conclusions of the lemma remain valid under the renaming $\pi_n \to \pi_n^{\text{diag/trail}}$ (cf. (3.3), (3.2)).

Proof The proof follows the same lines as the proof of lemma 10, except that we apply lemma 7 (instead of lemma 6) and equations (4.5), (4.4).

6 The Continuity of eUDRL at Deterministic Transition Kernels for a Finite Number of Iterations



Figure 2: Illustration of discontinuity of eUDRL-generated quantities at the boundary of $(\Delta S)^{S \times A}$. The figures present values of policy and goal-reaching objective along two continuous one-parameter rays in $(\Delta S)^{S \times A}$, see items A and C in the legend. Horizontal axes show the value of the ray-parameter α ; the boundary is reached at $\alpha = 0$. The exact value of the respective quantities at the boundary is represented by a horizontal line, see item B.

We begin our discussion by observing that eUDRL-generated quantities, like the policies $\pi_{n,\lambda}$ and the respective goal-reaching objective $J_{\lambda}^{\pi_{n,\lambda}}$, in general do not need to be continuous (functions of the transition kernel) at all points $\lambda \in (\Delta S)^{S \times A}$. Kernels located at the boundary⁵ of $(\Delta S)^{S \times A}$ are particularly problematic. Figure 2 illustrates this by showing the presence of continuous and discontinuous behavior of the policy and the goal-reaching objective near two specific kernels (where one is deterministic⁶ and the other is non-deterministic) at the boundary of $(\Delta S)^{S \times A}$. All details of the construction of the MDPs underlying figure 2 can be found in the appendix B, examples 1 and 2. Since eU-DRL converges to the optimum already for $n \ge 1$ in case of a deterministic kernel, we use n=2 for our illustration. Specifically, figure 2 shows a comparison of the policy and the goal-reaching objective along two distinct, continuous rays of kernels in $(\Delta S)^{S \times A}$, where each ray is parametrized by a parameter α . The rays intersect at the boundary when $\alpha = 0$. Subfigure 2b suggests that the goal-reaching objective $J_{\lambda}^{\pi_{2,\lambda}}$ is continuous at the boundary (as $\alpha = 0+$) if the kernel is deterministic, but subfigure 2d suggest that the goal-reaching objective can exhibit a non-removable discontinuity at the boundary in general. In view of the behavior in subfigure 2d the continuity of $J_{\lambda}^{\pi_{2,\lambda}}$ at a deterministic point (a product of ΔS vertices, which also lies on the boundary) might come as a surprise and, as it turns out, is non-trivial to prove (see below). Notice, however, that eUDRL-generated policies are discontinuous even in the case of a deterministic kernel (see subfigures 2a, 2c). This can be seen as a consequence of the non-uniqueness of policies near optimality. This fact will play a major role in our proofs and cause a number of technical complications. One might wonder how $\pi_{n,\lambda}$ and $J_{\lambda}^{\pi_{n,\lambda}}$ behave in the interior of $(\Delta S)^{S \times A}$, even though this plays a minor role for our discussion of convergence of eUDRL. We leave the proof of continuity of eUDRLgenerated $\pi_{n,\lambda}$ and $J_{\lambda}^{\pi_{n,\lambda}}$ strictly in the interior of $(\Delta S)^{S \times A}$ to appendix C. We also include a discussion describing the causes leading to discontinuities of eUDRL-generated quantities at the end of appendix C.

The remaining part of this section is structured as follows. Section 6.1 introduces the discussion of continuity along given compatible families of CEs. In section 6.2, we define a notion of relative limit and relative continuity and demonstrate the relative continuity of eUDRL-generated policies and related values at deterministic kernels for finite number of iterations. We extend the results to other segment subspaces in section 6.3. We conclude by showing the continuity of the goal-reaching objective at deterministic kernels for finite number of iterations and discussing the error to optimality in section 6.4.

Since we always deal with finite-dimensional vector spaces, all norms are equivalent and we will often omit a norm specification from our results. Similarly, we will use norms interchangeably in some of our continuity-proofs.

^{5.} In this article we always interpret the terms boundary/interior in the elementary way, viewing $(\Delta S)^{S \times A}$ as continuously embedded into \mathbb{R}^m , where $m = |\mathcal{A}||\mathcal{S}|(|\mathcal{S}|-1)$. The term boundary/interior then refers to the elementary geometric understanding of the boundary/interior of a simplex.

^{6.} Notice that deterministic kernels are located on the boundary of the simplex in general. This is because an arbitrary small perturbation that makes the zero-entries of a deterministic kernel negative will leave the simplex of probabilities.

6.1 Continuity of Values and Segment Distribution Marginals

While eUDRL-generated policies and goal-reaching objectives are not generally continuous functions of the transition kernel, this section shows that state-value functions, action-value functions and the segment distribution of marginals are indeed continuous (functions of the transition kernel). The difference is that here we interpret (see below) the state/action-values and segment-distribution marginals as functions of two variables λ, π , whereas we interpret d them as a function solely of λ in the discussion of figure 2. In other words, in our discussion of figure 2 the discontinuity appears because the value function is concatenated with a λ -dependent policy.

Lemma 12 (Continuity in compatible families) Let $\{\mathcal{M}_{\lambda}|\lambda \in (\Delta S)^{S \times A}\}$ and $\{\bar{\mathcal{M}}_{\lambda}|\lambda \in (\Delta S)^{S \times A}\}$ be compatible families.

- 1. (Continuity of value) Fix $\bar{s} \in \bar{S}_T$ and $a \in A$. Then the values $V^{\pi}_{\lambda}(\bar{s})$ and action-values $Q^{\pi}_{\lambda}(\bar{s}, a)$ are continuous when viewed as a function of (λ, π) . The optimal values $V^*_{\lambda}(\bar{s})$ and action values $Q^*_{\lambda}(\bar{s}, a)$ are continuous as function of λ .
- 2. (Upper bound on \mathbb{P}_{λ}) Fix $\bar{s} \in \bar{S}_T$ and $a \in \mathcal{A}$. Then for all $h' \geq h$ and $g' \in \mathcal{G}$ it holds that

$$\mathbb{P}_{\lambda}(\rho(S_h) = g | S_0 = s, H_0 = h', G_0 = g', A_0 = a; \pi) \le Q_{\lambda}^*((s, h, g), a).$$

- 3. (Continuity at value zero) Let $\pi_0 > 0$ be a given policy and let $\lambda_0 \in (\Delta S)^{S \times A}$ be a given transition kernel. Let $\lambda \mapsto \pi_{\lambda}$ denote a policy that depends on a transition kernel. Then for all $\bar{s}^* \in \bar{S}_T$ with $V_{\lambda_0}^{\pi_0}(\bar{s}^*) = 0$ it holds that:
 - $\lambda \mapsto V_{\lambda}^{\pi_{\lambda}}(\bar{s}^*)$ is continuous at λ_0 and
 - $V_{\lambda_0}^{\pi_{\lambda_0}}(\bar{s}^*) = 0.$
- 4. (Continuity of marginals) Segment distribution marginals are continuous in the transition kernel and the policy. The normalization constant of segment distribution $c(\lambda, \pi)$ and $c(\lambda, \pi)^{-1}$ are continuous.

Proof 1. Values $V_{\lambda}^{\pi}(\bar{s})$ and action-values $Q_{\lambda}^{\pi}(\bar{s}, a)$ and jointly continuous in λ and π because the recursion relation for the value function only contains a finite number of multiplication and summation operations involving λ and π . The continuity of $Q_{\lambda}^{*}(\bar{s}, a)$ follows from $Q_{\lambda}^{*}(\bar{s}, a) = \max_{\pi} Q_{\lambda}^{\pi}(\bar{s}, a)$ and similarly for $V_{\lambda}^{*}(\bar{s})$.

2. If π^* is a policy that achieves the highest probability of executing the command "reach goal g in exactly h steps", then

$$\mathbb{P}_{\lambda}(\rho(S_h) = g | S_0 = s, H_0 = h', G_0 = g', A_0 = a; \pi)$$

$$\leq \mathbb{P}_{\lambda}(\rho(S_h) = g | S_0 = s, H_0 = h, G_0 = g, A_0 = a; \pi^*) = Q_{\lambda}^*((s, h, g), a).$$

3. There exists $\epsilon > 0$ with $\pi_{\lambda} \leq 1 \leq \epsilon^{-1}\pi_0$. We claim that for all $\bar{s} = (s, h, g) \in \bar{S}_T$ it holds that

$$V_{\lambda}^{\pi_{\lambda}}(\bar{s}) \le V_{\lambda}^{\pi_{0}}(\bar{s})\epsilon^{-h}.$$

This is shown by induction. For h = 1 we get for all $\bar{s} = (s, 1, g) \in \bar{S}_T$ that

$$V_{\lambda}^{\pi_{\lambda}}(\bar{s}) = \sum_{a \in \mathcal{A}} Q_{\lambda}((s, 1, g), a) \pi_{\lambda}(a|s, 1, g) \le \sum_{a \in \mathcal{A}} Q_{\lambda}((s, 1, g), a) \epsilon^{-1} \pi_{0}(a|s, 1, g) = \frac{1}{\epsilon} V_{\lambda}^{\pi_{0}}(s, 1, g)$$

If the claim holds for $h \ge 1$ then, for all $\bar{s} = (s, h + 1, g) \in \bar{S}_T$, it also holds for h + 1:

$$\begin{aligned} V_{\lambda}^{\pi_{\lambda}}(\bar{s}) &= \sum_{a \in \mathcal{A}} (\sum_{s' \in \mathcal{S}} V_{\lambda}^{\pi_{\lambda}}((s',h,g))\lambda(s'|s,a))\pi_{\lambda}(a|s,1,g) \\ &\leq \sum_{a \in \mathcal{A}} (\sum_{s' \in \mathcal{S}} \epsilon^{-h} V_{\lambda}^{\pi_{0}}(s',h,g)\lambda(s'|s,a))\epsilon^{-1}\pi_{0}(a|s,1,g) = \epsilon^{-h-1} V_{\lambda}^{\pi_{0}}(s,h+1,g). \end{aligned}$$

Since $\lambda \mapsto V_{\lambda}^{\pi_0}$ is continuous by point 1. we conclude that

$$0 \le V_{\lambda}^{\pi_{\lambda}}(\bar{s}^*) \le \epsilon^{-h} V_{\lambda}^{\pi_0}(\bar{s}^*) \to \epsilon^{-h} V_{\lambda_0}^{\pi_0}(\bar{s}^*) = 0 \text{ as } \lambda \to \lambda_0.$$

4. Since $\mathbb{P}_{\lambda}(\mathcal{T} = \tau; \pi)$ is polynomial in the components of λ and π , it is continuous in (λ, π) . Since the overall number of trajectories is finite the normalizing term $c(\lambda, \pi)$ is continuous in (λ, π) as a finite sum of marginals $\mathbb{P}_{\lambda}(\mathcal{T} = \tau; \pi)$. Further, $c(\lambda, \pi) > 0$ due to the condition (2.3) on the initial distribution $\bar{\mu}$. Therefore, the segment probabilities $\mathbb{P}_{\lambda}(\Sigma = \sigma; \pi)$ are continuous in (λ, π) . The continuity of the marginals of $\mathbb{P}_{\lambda}(\Sigma = \sigma; \pi)$ follows since they are composed by finite sums of continuous functions $\mathbb{P}_{\lambda}(\Sigma = \sigma; \pi)$. The continuity of $c^{-1}(\lambda, \pi)$ follows from the continuity and positivity of $c(\lambda, \pi)$.

6.2 Relative Continuity of eUDRL Policies and Continuity of Related Values

Policies generated by eUDRL are not generally continuous in deterministic kernels (see figure 2c and example 2 in the appendix for details). As explained above discontinuities in policies can lead to discontinuous behavior of derived quantities, like the goal-reaching objective. Another key aspect is the non-uniqueness of optimal policies, which additionally complicates the analysis of continuity. However, a weaker notion of continuity still holds for eUDRL's policies, which will be useful in the study of continuity of emergent quantities. We will utilize the notion of relative weak limit of Štrupl et al. (2022b), except that in the finite setting of the article at hand weak convergence coincides with ordinary point-wise convergence and we drop the predicate "weak". For completeness we recall the underlying notion of convergence of probability distributions on finite quotient spaces, which we call relative convergence in this article. The standard notion of continuity then applies to maps of the form $f: Y \to \Delta X$ (e.g. policies) with respect to the quotient topology.

Definition 13 Let X be a finite set, let $F \subset X$ be a subset, let ΔX denote the probability simplex over X.

1. We say that a sequence of probabilities $(P_k)_{k\geq 0}$, $P_k \in \Delta X$ converges relatively to F, $P_k \xrightarrow{F} P$, if and only if it converges over the quotient X/F, i.e.

$$(\forall x \in X \setminus F) : P_k(x) \to P(x) \quad and \quad \sum_{x \in F} P_k(x) \to \sum_{x \in F} P(x).$$

2. Let (Y,d) be a metric space. We say that $f: Y \to \Delta X$ is continuous at $y' \in Y$ relative to F if and only if for every sequence $(y_k)_{k\geq 0}$, $y_k \in Y$ that converges to y' the sequence $(f(y_k))_{k\geq 0}$ converges to f(y') relatively to F. In this case we will write:

$$f(y) \xrightarrow{F} f(y')$$
, as $y \to y'$.

Remark 14 Recall that the above continuity of f at y' relative to F can be written equivalently using limit of a function, i.e.

$$(\forall x \in X \setminus F) : [f(y)](x) \to [f(y')](x) \quad and \quad \sum_{x \in F} [f(y)](x) \to \sum_{x \in F} [f(y')](x) \quad as \quad y \to y'.$$

Here the functions $y \mapsto [f(y)](x)$, $x \in X \setminus F$ and $y \mapsto \sum_{x \in F} [f(y)](x)$. Furthermore, the sequential approach used in point 2 of the definition is equivalent to the ϵ - δ formalism, since we are working in metric spaces.

In what follows we will demonstrate that the mapping $\lambda \mapsto \pi_{n,\lambda}$ has a relative limit that at deterministic kernels equals to optimal policies. Thus $\pi_{n,\lambda}$ are relatively continuous at deterministic kernels. Before we will state a simple lemma which will be needed during the proof.

Lemma 15 Assume the same situation as in definition 13, where f is continuous at y' relative to F. Moreover, consider a second function $h: Y \to \mathbb{R}^X$ that is continuous at $y' \in Y$ and assume h(y') is constant on F. Then the following holds as $y \to y'$:

$$\sum_{x \in X} [f(y)](x) \cdot [h(y)](x) \to \sum_{x \in X} [f(y')](x) \cdot [h(y')](x).$$

Proof We employ the defining properties of convergence relative to F. Expanding the left hand side yields

$$\sum_{x \in X} [f(y)](x)[h(y)](x) = \sum_{x \in F} [f(y)](x)[h(y)](x) + \sum_{x \in X \setminus F} [f(y)](x)[h(y)](x).$$

Since convergence relative to F behaves like point-wise convergence on $X \setminus F$ we have

$$\sum_{x \in X \setminus F} [f(y)](x) \cdot [h(y)](x) \to \sum_{x \in X \setminus F} [f(y')](x) \cdot [h(y')](x).$$

On the other hand, let $(y_k)_{k\geq 0}$, $y_k \in Y$ be a sequence that converges to y'. Since h(y') = c is constant on F, choosing k_0 large enough, we have for all $k > k_0$ that $|h(y_k) - c| \leq \varepsilon$. It follows for any $k > k_0$ that

$$\left| \sum_{x \in F} [f(y_k)](x)[h(y_k)](x) - \sum_{x \in F} [f(y')](x)[h(y')](x) \right| \le c \left| \sum_{x \in F} [f(y_k)](x) - \sum_{x \in F} [f(y')](x) \right| + |X| \cdot \varepsilon,$$

where |X| stands for the finite number of elements in X. Making use of $f(y) \xrightarrow{F} f(y')$ the right hand side converges to $\epsilon |X|$. Since we can choose ϵ arbitrary small the left-hand side

of the inequality has to converge to 0. This completes the proof.

The following theorem is the main result of this section. It demonstrates that eUDRLgenerated policies converge relatively to their sets of optimal actions. The result is stated for the eUDRL recursion on Seg space (3.1), but it can be modified mutatis mutandis for recursions on trailing or diagonal segments.

Theorem 16 (Relative continuity of eUDRL policies and continuity of value at deterministic kernels) Let $\{\mathcal{M}_{\lambda} : \lambda \in (\Delta S)^{S \times A}\}$ and $\{\bar{\mathcal{M}}_{\lambda} : \lambda \in (\Delta S)^{S \times A}\}$ be compatible families of MDPs. Let $\lambda_0 \in (\Delta S)^{S \times A}$ be a deterministic kernel and let $(\pi_{n,\lambda})_{n\geq 0}$ with $\pi_0 > 0$ be a eUDRL-generated sequence of policies.

- 1. For all $\bar{s} \in \bar{S}_{\lambda_0}$ the policy $\pi_{n,\lambda}(\cdot|\bar{s})$ is continuous relative to $\mathcal{O}(\bar{s})$ at λ_0 .
- 2. For all $\bar{s} \in \bar{S}_{\lambda_0}$ and for all $a \in A$, the values $Q_{\lambda}^{\pi_n}(\bar{s}, a)$, $V_{\lambda}^{\pi_n}(\bar{s})$ are continuous functions of λ at λ_0 .
- 3. There exists a sequence of neighborhoods $(U_{\delta_n}(\lambda_0))_{n\geq 0}$ with $U_{\delta_{n+1}}(\lambda_0) \subset U_{\delta_n}(\lambda_0)$ such that for all $n \geq 0$ it holds:
 - $\pi_{n,\lambda}(a|\bar{s})$ is separated from 0 for all $(a, \bar{s}, \lambda) \in (\operatorname{supp num}_{\lambda_0, \pi_0} \cap (\mathcal{A} \times \operatorname{supp} \nu_{\lambda_0, \pi_0})) \times U_{\delta_n}(\lambda_0)$, and
 - the probabilities $\mathbb{P}_{\lambda}(H^{\Sigma} = h, G_0^{\Sigma} = g | S_0^{\Sigma} = s, l(\Sigma) = h; \pi_n)$ are separated from 0 for all $((s, h, g), \lambda) \in \bar{S}_{\lambda_0} \times U_{\delta_n}(\lambda_0)$.

Proof The proof proceeds by induction on the index n of the eUDRL iteration.

Base case (n=0): Since π_0 is positive, constant in λ and state and action spaces are finite it follows that π_0 is continuous, relatively continuous, and separated from 0. Since π_0 is continuous in λ , and values are continuous in (λ, π_0) by lemma 12 point 1, values are continuous in λ at λ_0 (after composition with π_0). Choose $2 > \delta_0 > 0$ such that $\|\lambda - \lambda_0\|_{\max} < \frac{1}{2}$ for $\lambda \in \overline{U_{\delta_0}(\lambda_0)}$. From lemma 12 point 4. it follows that the segment distribution marginal $\mathbb{P}_{\lambda}(H_0^{\Sigma} = h, G_0^{\Sigma} = g, S_0^{\Sigma} = s, l(\Sigma) = h; \pi_0)$ is continuous. Since $(s, h, g) \in \overline{S}_{\lambda_0}$, it follows from lemma 6 point 3. that the marginal is positive. In summary, the marginal is a positive and continuous function on the compact set $\overline{U_{\delta_0}(\lambda_0)}$, which implies that it has a positive minimum.

Induction: Assume all three statements hold for n. We aim to prove them for n+1. By point 3. for all $\lambda \in \overline{U_{\delta_n}(\lambda_0)}$ we have that lemma 6 applies. Consequently choosing a fixed $(s, h, g) \in \overline{S}_{\lambda_0} \subset \operatorname{supp} \operatorname{den}_{\lambda, \pi_n} \cap \operatorname{supp} \nu_{\lambda, \pi_n}$ we can use lemma 3 point 1 to get

$$\pi_{n+1}(a|s,h,g) = \frac{u(a)}{\sum_{a\in\mathcal{A}} u(a)} \text{ with}$$

$$u(a) := \sum_{N\geq h'\geq h, g'\in\mathcal{G}} \mathbb{P}_{\lambda}(\rho(S_h^{\Sigma}) = g|A_0^{\Sigma} = a, S_0^{\Sigma}, H_0^{\Sigma} = h', G_0^{\Sigma} = g', l(\Sigma) = h; \pi_n)$$

$$\cdot \pi_n(a|s,h',g') \mathbb{P}_{\lambda}(H_0^{\Sigma} = h', G_0^{\Sigma} = g'|S_0^{\Sigma}, l(\Sigma) = h; \pi_n). \quad (6.1)$$

To demonstrate point 1. we begin by providing lower and upper estimates on u. Using lemma 12 point 2. we have and upper bound

$$u(a) \le \sum_{N \ge h' \ge h, g' \in \mathcal{G}} Q_{\lambda}^{*}((s, h, g), a) \cdot 1 \cdot \mathbb{P}_{\lambda}(H_{0}^{\Sigma} = h', G_{0}^{\Sigma} = g'|S_{0}^{\Sigma}, l(\Sigma) = h; \pi_{n}) \le Q_{\lambda}^{*}((s, h, g), a)$$
(6.2)

and the lower bound

$$u(a) \geq Q_{\lambda}^{\pi_{n}}((s,h,g),a)\pi_{n}(a|s,h,g)\mathbb{P}_{\lambda}(H_{0}^{\Sigma}=h,G_{0}^{\Sigma}=g|S_{0}^{\Sigma},l(\Sigma)=h;\pi_{n})$$

$$\geq \begin{cases} Q_{\lambda}^{\pi_{n}}((s,h,g),a)\epsilon \text{ for } a \in \mathcal{O}(s,h,g), \\ 0 \text{ otherwise}, \end{cases}$$
(6.3)

where the existence of $\epsilon > 0$ follows by point 3. of the theorem by the induction assumption. Using these bounds we now show that $\pi_{n+1}(a|s,h,g) \xrightarrow{\mathcal{O}(s,h,g)} \pi_{\lambda_0}^*$ as $\lambda \to \lambda_0$. For this it is sufficient to show that $\pi_{n+1}(a|s,h,g) \to 0$ for all $a \notin \mathcal{O}(s,h,g)$ because $\pi_{n+1}(\mathcal{O}(s,h,g)|s,h,g) \to 1$ follows automatically. We show that $u(a) \to 0, a \notin \mathcal{O}(s,h,g)$ and that the denominator $\sum_{a \in \mathcal{A}} u(a)$ is separated from 0. First, $u(a) \to 0, a \notin \mathcal{O}(s,h,g)$ since the upper bound $Q_{\lambda}^*((s,h,g),a) \to Q_{\lambda_0}^*((s,h,g),a) = 0$ for $a \notin \mathcal{O}(s,h,g)$ by the continuity of $Q_{\lambda}^*((s,h,g),a)$. By assumption $Q_{\lambda}^{\pi_n}((s,h,g),a)$ is continuous and we can choose $\delta_n \geq \delta_{n+1} > 0$ such that $Q_{\lambda}^{\pi_n}((s,h,g),a) \geq \frac{1}{2}$ for $a, (s,h,g), \lambda \in (\text{supp num}_{\lambda_0,\pi_0} \cap (\mathcal{A} \times \text{supp } \nu_{\lambda_0,\pi_0})) \times U_{\delta_{n+1}}(\lambda_0)$. Finally, using the lower bound we conclude that

$$\sum_{a \in \mathcal{A}} u(a) \geq \epsilon \sum_{a \in \mathcal{O}(s,h,g)} Q_{\lambda}^{\pi_n}((s,h,g),a) \geq \epsilon \frac{1}{2} > 0.$$

Thus $\pi_{n+1}(a|s, h, g)$ is relatively continuous in λ_0 . Using both bounds on u we show that $\pi_{n+1}(a|s, h, g)$ is also separated form 0 for all $(s, h, g), a, \lambda \in (\text{supp num}_{\lambda_0, \pi_0} \cap (\mathcal{A} \times \text{supp } \nu_{\lambda_0, \pi_0})) \times U_{\delta_{n+1}}(\lambda_0),$

$$\pi_{n+1}(a|s,h,g) = \frac{u(a)}{\sum_{a \in \mathcal{A}} u(a)} \ge \frac{Q_{\lambda}^{\pi_n}((s,h,g),a)\epsilon}{\sum_{a \in \mathcal{A}} Q_{\lambda}^*((s,h,g),a)} \ge \frac{\frac{1}{2}\epsilon}{\sum_{a \in \mathcal{A}} 1} \ge \frac{1}{2|\mathcal{A}|}\epsilon > 0$$

To demonstrate the continuity of state/action-values we will employ an additional induction argument (for fixed n) over the remaining horizon $N \ge h \ge 1$.

Sub-induction base case (h = 1): For h = 1, $(s, 1, g) \in \overline{S}_T$, $a \in \mathcal{A}$ the action-value function

$$Q_{\lambda}^{\pi_{n+1}}((s,1,g),a) = \sum_{s' \in \rho^{-1}(\{g\})} \lambda(s'|s,a)$$

is continuous in λ .

Sub-induction: Fix the remaining horizon $N \ge h' \ge 1$. Assume that for all $(s, h, g) \in \overline{S}_{\lambda_0}$, h = h', $a \in \mathcal{A}$ the action values $Q_{\lambda}^{\pi_{n+1}}((s, h, g), a)$ are continuous in λ at λ_0 . It remains to prove continuity of $V_{\lambda}^{\pi_{n+1}}(s, h, g)$ for all $(s, h, g) \in \overline{S}_{\lambda_0}$, h = h' and of $Q_{\lambda}^{\pi_{n+1}}((s, h+1, g), a)$ for all $(s, h + 1, g) \in \overline{S}_{\lambda_0}$, h = h', $a \in \mathcal{A}$ at λ_0 . From lemma 15 and the relative continuity

of π_{n+1} we obtain

$$\lim_{\lambda \to \lambda_0} V_{\lambda}^{\pi_{n+1}}(s,h,g) = \lim_{\lambda \to \lambda_0} \sum_{a \in \mathcal{A}} Q_{\lambda}^{\pi_{n+1}}((s,h,g),a)\pi_{n+1}(a|s,h,g)$$
$$= \sum_{a \in \mathcal{O}(s,h,g)} \pi_{\lambda_0}^*(a|s,h,g) = 1 = V_{\lambda_0}^*(s,h,g).$$

Thus the continuity $V_{\lambda}^{\pi_{n+1}}(s,h,g)$ in λ_0 follows. Now for all $(s,h+1,g) \in \bar{S}_{\lambda_0}$, h = h', $a \in \mathcal{A}$ the limit of the action-value function can be computed as

$$\lim_{\lambda \to \lambda_0} Q_{\lambda}^{\pi_{n+1,\lambda}}((s,h+1,g),a) = \lim_{\lambda \to \lambda_0} V_{\lambda}^{\pi_{n+1,\lambda}}(s',h,g)\lambda(s'|s,a),$$

where we used that λ_0 is deterministic, chose s' so that $\lambda_0(s'|s, a) = 1$ and made use of the fact that $\lambda(s''|s, a) \to 0$ as $\lambda \to \lambda_0$ for $s'' \neq s'$. It remains to show the continuity of $V_{\lambda}^{\pi_{n+1,\lambda}}(s', h, g)$ and that $V_{\lambda_0}^{\pi_{n+1,\lambda_0}}(s', h, g) = Q_{\lambda_0}^{\pi_{n+1,\lambda_0}}((s, h+1, g), a) = Q_{\lambda_0}^*((s, h+1, g), a)$. First assume $a \in \mathcal{O}(s, h+1, g)$ then form lemma 6 point 4. it holds that $(s', h, g) \in \bar{S}_{\lambda_0}$. But we have just proved $V_{\lambda}^{\pi_{n+1,\lambda}}(s', h, g) \to V_{\lambda_0}^{\pi_{n+1,\lambda_0}}(s', h, g) = 1, \lambda \to \lambda_0$ for such states. Finally assume $a \notin \mathcal{O}(s, h+1, g)$ then from lemma 6 point 5. we conclude that $Q_{\lambda_0}^{\pi_{n+1,\lambda_0}}((s, h+1, g), a) = 0 = V_{\lambda_0}^{\pi_{n+1,\lambda_0}}((s', h, g)) \cdot 1$. Then lemma 12 point 3. implies

$$V_{\lambda}^{\pi_{n+1,\lambda}}((s',h,g)) \to V_{\lambda_0}^{\pi_{n+1,\lambda_0}}((s',h,g)) = 0 \text{ for } \lambda \to \lambda_0.$$

This completes the sub-induction.

It remains to show that the probabilities $\mathbb{P}_{\lambda}(H_0^{\Sigma} = h, G_0^{\Sigma} = g \mid S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g; \pi_{n+1})$ remain separated from 0 on a neighborhood of λ_0 . Notice that the segment distribution marginal $\mathbb{P}_{\lambda}(H_0^{\Sigma} = h, G_0^{\Sigma} = g, S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g; \pi_{n+1}) > 0$ is continuous in $(\lambda, \pi_{n+1,\lambda})$ (viewed as independent variables) but $\pi_{n+1,\lambda}$ is not generally continuous in λ at λ_0 (but only relatively continuous). Hence, it would not be sufficient to argue that "a positive continuous function on a compact must have a positive minimum" and use lemma 6 point 3. and lemma 12 point 4. This motivates the following, more elaborate, approach.

From lemma 6 points 2., 3. we have that $(s, h, g) \in \overline{S}_{\lambda_0} = \operatorname{supp} \operatorname{den}_{\lambda_0, \pi_{n+1}} \cap \operatorname{supp} \nu_{\lambda_0, \pi_{n+1}}$ and that

$$\mathbb{P}_{\lambda_0}(S_0^{\Sigma} = s, H_0^{\Sigma} = h, G_0^{\Sigma} = g, l(\Sigma) = h; \pi_{n+1}) > 0.$$

It follows that there exists a segment σ that makes the above marginal of the segment distribution positive. In turn there must be a trajectory τ with positive probability. We end up with the following inequalities

$$\mathbb{P}_{\lambda_0}(S_0^{\Sigma} = s, H_0^{\Sigma} = h, G_0^{\Sigma} = g, l(\Sigma) = h; \pi_{n+1}) \ge \mathbb{P}_{\lambda_0}(\Sigma = \sigma; \pi_{n+1}) \ge \frac{\mathbb{P}_{\lambda_0}(\mathcal{T} = \tau; \pi_{n+1})}{c_{\lambda_0}}.$$

The normalizing factor of the segment distribution c_{λ} can be bounded above $c_{\lambda} \leq N(N + 1)/2$ (see equation (2.4)). Since the transition probabilities in $\mathbb{P}_{\lambda_0}(\mathcal{T} = \tau; \pi_{n+1})$ are 1 as λ_0 is deterministic and $\|\lambda - \lambda_0\|_{\max} < \frac{1}{2}$ on $U_{\delta_{n+1}}(\lambda_0) \subset U_{\delta_0}(\lambda_0)$ we bound them on $U_{\delta_{n+1}}(\lambda_0)$
by $\frac{1}{2}$ from below. Similarly, since $\pi_{n+1,\lambda_0} = \pi^*$ chooses actions from the optimal action set $\mathcal{O}(\cdot)$, where $\pi_n > 0$ is separated from 0 on $U_{\delta_{n+1}}(\lambda_0)$, we conclude

$$\mathbb{P}_{\lambda}(S_{0}^{\Sigma} = s, H_{0}^{\Sigma} = h, G_{0}^{\Sigma} = g, l(\Sigma) = h; \pi_{n+1}) \geq \frac{\mathbb{P}_{\lambda}(\mathcal{T} = \tau; \pi_{n+1})}{c_{\lambda}} = \frac{\bar{\mu}(s_{0}, h_{0}, g) \prod_{t=0}^{N-1} \lambda(s_{t+1}|s_{t}, a_{t}) \pi_{n+1}(a_{t}|s_{t}, h_{0} - t, g)}{c_{\lambda}} \geq \underbrace{\frac{2\bar{\mu}(s_{0}, h_{0}, g)(\frac{1}{2})^{N}(\epsilon')^{N}}{N(N+1)}}_{C(s, h, g):=} > 0,$$

where $\epsilon' > 0$ denotes the respective lower bound. This implies that the probabilities $\mathbb{P}_{\lambda}(H_0^{\Sigma} = h, G_0^{\Sigma} = g \mid S_0^{\Sigma} = s, l(\Sigma) = h; \pi_{n+1})$ are well-defined and bounded from below by C(s, h, g). Since the set \bar{S}_{λ_0} is finite we can choose a common positive lower bound as the minimum of all positive lower bounds.

As an illustration of theorem 16 and of the behavior of the continuity of eUDRL-generated policies, figure 3 shows plots of the quantity $\min_{\bar{s}\in\bar{S}_{\lambda_0}}\pi_n(\mathcal{O}(\bar{s})|s)$ in case of a random walk on \mathbb{Z}_3 . All details of the construction of the MDP underlying figure 3 can be found in the appendix B, example 3. The figure shows how $\min_{\bar{s}\in\bar{S}_{\lambda_0}}\pi_n(\mathcal{O}(\bar{s})|s)$ depends on n for varying distances from a deterministic kernel (as measured by $\delta = ||\lambda - \lambda_0||_1$) and varying initial conditions. The initial conditions are given by the starting policy for eUDRL and are reflected by the offset of the respective graph on the vertical axis (at n = 0). Different initial conditions are depicted using different colors. The dependency $\min_{\bar{s}\in\bar{S}_{\lambda_0}}\pi_n(\mathcal{O}(\bar{s})|s)$ on δ is illustrated by different graphs, where the increase of $\min_{\bar{s}\in\bar{S}_{\lambda_0}}\pi_n(\mathcal{O}(\bar{s})|s)$ as δ decreases is consistent with the established continuity of policy π_n in λ . It is also visible that for small values of δ the eUDRL iteration leads to an increase of $\min_{\bar{s}\in\bar{S}_{\lambda_0}}\pi_n(\mathcal{O}(\bar{s})|s)$, whereas for large values of δ the eUDRL iteration leads to a deterioration of performance. The dependency of eUDRL's overall performance on the initial conditions at fixed δ can be estimated from the graphs by comparing the asymptotic behavior when n is increasing.

6.3 Extending the Continuity Results to Other Segment Sub-Spaces

To cover algorithms like ODT, RWR (and some specific variants of eUDRL) a modification of theorem 16 is needed that restricts the recursion (cf. section 2.4 and following) to Seg^{trail} or Seg^{diag} subspaces.

Theorem 17 (Relative continuity of eUDRL policies and continuity of values in deterministic kernels) Theorem 16 remains valid under the renaming $\pi_n \to \pi_n^{\text{diag/trail}}$.

Notice that the renaming applies to policies only. It ignores the sets \bar{S}_{λ_0} and $\sup \operatorname{num}_{\lambda_0,\pi_0} \cap (\mathcal{A} \times \sup \nu_{\lambda_0,\pi_0})$, which is justified by the equations (4.4) and (4.5). There is also no need to investigate trailing/diagonal variants of the probability in theorem 16, point 3 (by conditioning on the events $H_0^{\Sigma} = l(\Sigma)$, $G_0^{\Sigma} = \rho(S_h^{\Sigma})$). The probability to bound remains the same for all segment sub-spaces. This is a consequence of lemma 3, particularly equation (6.1), which permits rewriting the eUDRL recursion for all segment sub-spaces exclusively using terms that appear in the recursion rewrite for Seg. More broadly, lemma 3 simplifies proofs in Seg^{diag/trail} considerably since it allows us to restrict to quantities without the conditioning on $H_0^{\Sigma} = l(\Sigma)$, $G_0^{\Sigma} = \rho(S_h^{\Sigma})$ in all three versions of the proof.



Figure 3: Illustration of the behavior of $\min_{\bar{s}\in \bar{S}_{\lambda_0}} \pi_n(\mathcal{O}(\bar{s})|s)$ when varying the distance to a deterministic kernel and varying the initial policy for a random walk on \mathbb{Z}^3 . The legend shows the distance δ of the respective graph to the deterministic kernel. Each initial condition is depicted by a different color and plotted for various distances δ . One particular initial condition is highlighted in black.

Proof We follow the proof of theorem 16 and only highlight differences. First, we employ lemma 11 for Seg^{diag/trail} instead of lemma 10, which we employed for Seg. Second, as explained above, there is no need to bound diagonal/trailing variants of the probability in point 3. Third, the fact that we restricted the proof to $\bar{s} \in \bar{S}_{\lambda_0}$ (in (6.1)) implies that not only $\pi_{n+1,\lambda}(\cdot|\bar{s})$ is defined but also $\pi_{n+1,\lambda}^{\text{trail}}(\cdot|\bar{s})$ and $\pi_{n+1,\lambda}^{\text{diag}}(\cdot|\bar{s})$ are defined, see (4.5) and lemma 7 point 2. For the recursion in Seg^{trail} or Seg^{diag} we have to modify u (compare (6.1) and (3.2), (3.3)) choosing

$$\begin{split} u^{\text{trail}}(\pi_n^{\text{trail}}) &:= \sum_{g' \in \mathcal{G}} \mathbb{P}_{\lambda}(\rho(S_h^{\Sigma}) = g | A_0^{\Sigma} = a, S_0^{\Sigma}, H_0^{\Sigma} = h, G_0^{\Sigma} = g', l(\Sigma) = h; \pi_n^{\text{trail}}) \\ & \quad \cdot \pi_n^{\text{trail}}(a | s, h, g') \mathbb{P}_{\lambda}(H_0^{\Sigma} = h, G_0^{\Sigma} = g' | S_0^{\Sigma}, l(\Sigma) = h; \pi_n^{\text{trail}}), \\ u^{\text{diag}}(\pi_n^{\text{diag}}) &:= \mathbb{P}_{\lambda}(\rho(S_h^{\Sigma}) = g | A_0^{\Sigma} = a, S_0^{\Sigma}, H_0^{\Sigma} = h, G_0^{\Sigma} = g, l(\Sigma) = h; \pi_n^{\text{diag}}) \\ & \quad \cdot \pi_n^{\text{diag}}(a | s, h, g) \mathbb{P}_{\lambda}(H_0^{\Sigma} = h, G_0^{\Sigma} = g | S_0^{\Sigma}, l(\Sigma) = h; \pi_n^{\text{diag}}). \end{split}$$

Notice that given a policy π_n it holds that

$$u^{\operatorname{diag}}(\pi_n) \le u^{\operatorname{trail}}(\pi_n) \le u(\pi_n).$$

Since the introduced upper bound (6.2) bounds $u(\pi_n)$ it also applies for $u^{\text{diag}}(\pi_n)$ and $u^{\text{trail}}(\pi_n)$. Similarly, since all non-diagonal terms were lower bounded by zero, the introduced lower bound (6.3) bounds $u^{\text{diag}}(\pi_n)$ and consequently it bounds also $u^{\text{trail}}(\pi_n)$. Thus the upper and lower estimates that are used in the proof of theorem 16 apply also for $u^{\text{trail}}(\pi_n)$ and $u^{\text{diag}}(\pi_n)$. Thus apart from the change of the definition of u the proof remains unchanged.

We will apply this method to derive universal bounds by lower bounding u^{diag} and upper bounding u also in later proofs when studying the eUDRL-type recursions.

6.4 Continuity of the Goal-Reaching Objective for a Finite Number of Iterations

Corollary 18 (Continuity of the goal-reaching objective at deterministic points) Let $(\pi_{n,\lambda})_{n\geq 0}$ with $\pi_0 > 0$ be a sequence of eUDRL-generated policies. Let $\lambda_0 \in (\Delta S)^{S \times A}$ be a deterministic kernel. Then the goal-reaching objective, viewed as a function of λ , $J_{\lambda}^{\pi_{n,\lambda}} = \sum_{\bar{s}\in\bar{S}_T}\bar{\mu}(\bar{s})V_{\lambda}^{\pi_{n,\lambda}}$ is continuous at λ_0 for all $n\geq 0$.

Proof The goal-reaching objective can be written as follows

$$J_{\lambda}^{\pi_{n,\lambda}} = \sum_{\bar{s} \in \operatorname{supp} \bar{\mu}} \bar{\mu}(\bar{s}) V_{\lambda}^{\pi_{n,\lambda}} = \sum_{\bar{s} \in \operatorname{supp} \bar{\mu} \setminus \bar{\mathcal{S}}_{\lambda_0}} \bar{\mu}(\bar{s}) V_{\lambda}^{\pi_{n,\lambda}} + \sum_{\bar{s} \in \operatorname{supp} \bar{\mu} \cap \bar{\mathcal{S}}_{\lambda_0}} \bar{\mu}(\bar{s}) V_{\lambda}^{\pi_{n,\lambda}}$$

where the continuity of the second summand follows from theorem 16, point 2. For the first summand we prove that for all $\bar{s} \in \operatorname{supp} \bar{\mu} \setminus \bar{S}_{\lambda_0} = \operatorname{supp} \bar{\mu} \setminus \operatorname{supp} \operatorname{den}_{\lambda_0,\pi_0}$ it holds that $V_{\lambda_0}^{\pi_0}(\bar{s}) = 0$. Continuity then follows from lemma 12, point 3. For the sake of contradiction assume that $V_{\lambda_0}^{\pi_0}(\bar{s}) > 0$ for a certain $\bar{s} = (s, h, g) \in \operatorname{supp} \bar{\mu} \setminus \operatorname{supp} \operatorname{den}_{\lambda_0,\pi_0}$ (recall that $\operatorname{supp} \bar{\mu} \subset \bar{S}_T$). This means that there exists a trajectory $\tau = \bar{s}_0, a_0, \ldots, \bar{s}_h$ with $\bar{s} = (s, h, g) = \bar{s}_0, \rho(s_h) = g$ and $\mathbb{P}_{\lambda_0}(A_0 = a_0, \ldots, \rho(S_h) = g | \bar{S}_0 = \bar{s}; \pi_0) > 0$. Since $\bar{\mu}(\bar{s}) > 0$ we conclude $\mathbb{P}_{\lambda_0}(\mathcal{T} = \tau; \pi_0) > 0$ but this contradicts $\bar{s} \notin \operatorname{supp} \operatorname{den}_{\lambda_0,\pi_0}$.

We have shown that the goal-reaching objective derived from eUDRL-generated policies is continuous for any iteration $n \geq 0$. This implies that if an environment possesses a transition kernel that is close to a deterministic kernel, then the eUDRL generated policy $\pi_{n,\lambda}$, $n \geq 1$ (with initial condition $\pi_0 > 0$) has a goal-reaching objective $J_{\lambda}^{\pi_{n,\lambda}}$ close to the optimal goal-reaching objective $J_{\lambda}^{\pi^*}$. We summarize this observation in a corollary.

Corollary 19 (Near-optimal behavior near deterministic kernels) Let λ_0 be a deterministic transition kernel and let $(\pi_{n,\lambda})_{n\geq 0}$ with $\pi_0 > 0$ be a eUDRL-generated sequence of policies. Then for any fixed n and all $\epsilon > 0$ there exists $\delta > 0$ such that if $\lambda \in U_{\delta}(\lambda_0)$ then

$$|J_{\lambda}^{\pi_{n,\lambda}} - J_{\lambda}^{\pi_{\lambda}^*}| < \epsilon.$$

Proof Due to continuity of the optimal values in λ (see lemma 12), we can choose $\delta_1 > 0$ such that

$$|J_{\lambda}^{\pi_{\lambda}^*} - J_{\lambda_0}^{\pi_{\lambda_0}^*}| < \frac{\epsilon}{2}.$$

By lemma 10 the eUDRL-generated policies are optimal for deterministic kernels when $n \geq 1$. By the continuity of $J_{\lambda}^{\pi_{n,\lambda}}$ at λ_0 (see lemma 18), we can chose $\delta_2 > 0$ such that for all $\lambda \in U_{\delta_2}(\lambda_0)$ it holds that

$$\frac{\epsilon}{2} > |J_{\lambda}^{\pi_{n,\lambda}} - J_{\lambda_0}^{\pi_{n,\lambda_0}}| = |J_{\lambda}^{\pi_{n,\lambda}} - J_{\lambda_0}^{\pi_{\lambda_0}^*}|.$$

By the triangle inequality and choosing $\delta := \min\{\delta_1, \delta_2\}$ we obtain $|J_{\lambda}^{\pi_{n,\lambda}} - J_{\lambda_0}^{\pi_{\lambda_0}^*}| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$.

This says that $J_{\lambda}^{\pi_{n,\lambda}}$ has a small error to optimality in a small neighborhood around a deterministic transition kernel for all finite $n \geq 1$. However, the lemma provides no information on how for a given level of error the size of the neighborhood δ (for given fixed ϵ) depends on the iteration. As shown in (Štrupl et al., 2022a) by explicit counterexamples there exist non-deterministic environments, where eUDRL-generated policies remain constant for $n \geq 1$, such that there is no improvement at all. The same article shows that even if initialized with the optimal policy a priori, eUDRL might produce non-optimal policies such that there is no monotonic improvement.

An illustration of the behavior of the goal-reaching objective for eUDRL-generated policies in case of a random walk on \mathbb{Z}_3 is shown in the figure 4. All details of the construction of the MDP underlying figure 4 can be found in the appendix B, example 3. Figure 4a shows the dependency of $J_{\lambda}^{\pi_{n,\lambda}}$ on the iteration for varying distances from a deterministic kernel (as measured by $\delta = \|\lambda - \lambda_0\|_1$) and varying initial conditions. The initial conditions are given by the starting policy for eUDRL and are reflected by the offset of the respective graph on the vertical axis (at n = 0). Different initial conditions are depicted using different colors. The dependency $J_{\lambda}^{\pi_{n,\lambda}}$ on δ is illustrated by different graphs, where the increase of values $J_{\lambda}^{\pi_{n,\lambda}}$ as δ decreases is consistent with the established continuity of $J_{\lambda}^{\pi_{n,\lambda}}$ in λ . It is also visible that for small values of δ the eUDRL iteration leads to an increase of $J_{\lambda}^{\pi_{n,\lambda}}$, whereas for large values of δ the eUDRL iteration leads to a deterioration of performance. The dependency of eUDRL's overall performance on the initial conditions at fixed δ can be estimated from the graphs by comparing the asymptotic behavior when n is increasing. Figure 4b shows a closer resolution of one of the graphs of figure 4a for $\delta = 0.25$. It is visible that two distinct regimes are present in the performance of eUDRL in this example, with a sharp increase for small values of n and a slow subsequent deterioration. Furthermore, considering the asymptotic behavior present in figure 4 for different values of δ it appears that the set of accumulation points of the sequences of values of the goal-reaching objective also exhibits a form of continuity in λ . The proofs of continuity properties of the sets of accumulation points of eUDRL-generated policies and the associated goal-reaching objectives is the content of the coming section.



Figure 4: Illustration of the continuity of the goal-reaching objective for varying distance to a deterministic kernel and varying initial policy for a random walk on \mathbb{Z}^3 . The legend shows the distance δ of the respective graph to the deterministic kernel. Each initial condition is depicted by a different color and plotted for various distances δ . One particular initial condition is highlighted in black in (a). One particular example is highlighted in closer zoom in (b).

7 The Continuity of eUDRL at Deterministic Transition Kernels as the Number of Iterations Approaches Infinity

The main objective of this section is to investigate the relative continuity of eUDRLgenerated policies, the associated values and goal-reaching objective as the number of iterations approaches infinity. The rigorous asymptotic analysis of eUDRL-type recursion schemes is open in the literature. A major hurdle is that it is unknown if, in general, eUDRL converges to a limiting policy. Since all policies form a compact simplex we will focus on the set of accumulation points of eUDRL-generated policies instead. For this we will introduce a notion of "relative continuity of sets of accumulation points of policies". Although, we do not provide a complete analysis of the continuity of asymptotic polices and values, we contribute significantly to the progress of the theory of "RL as SL" algorithms providing the first rigorous account of the topic. The study of continuity of eUDRL addresses the algorithm's asymptotic stability, i.e. the question whether small changes within the environment entail a small change in the overall learning performance. Observe that the largest values of the goal-reaching objective in figure 4 seem to be reached roughly at the first to third iteration. Increasing the number of iterations further results in an effective deterioration of performance. This raises concerns on whether the eUDRL recursion scheme makes sense at all. Thinking along this line one might ask for guarantees on the behavior of the accumulation points of eUDRL-generated policies and derived quantities. Since optimal values are always continuous (see lemma 12 point 1.) and eUDRL-values are optimal for deterministic kernels (both in case of a finite number of iterations and also in the limit) a discontinuity of eUDRL-values in λ_0 would imply a discontinuous error between eUDRL-values and optimal values in the limit (in the sense that $\|\lambda - \lambda_0\|_1 \leq \delta$ but the values have a positive separation). Thus the set of accumulation points would show a discontinuous behavior. Since this deficiency does not occur at a finite number of iterations (see Theorem 16), this would imply that eUDRL is asymptotically unstable, making the effort to achieve any limiting behavior via eUDRL a questionable pursuit.

As we are working in a metric space (rather than an abstract topological space), establishing a proof of continuity is tantamount to the derivation of error bounds on eUDRLgenerated quantities. Given a bound δ on the distance $\|\lambda - \lambda_0\|_1$ one could determine an upper bound ϵ on the error of, e.g., the goal-reaching objective. Building on this observation, the estimates derived during our exploration of continuity can help determine whether it is worthwhile to continue with the eUDRL iteration to achieve a specific level of error. This observation extends mutatis mutandis to GCSL, ODT and other "RL as SL" approaches.

We have no estimates on the change of the visitation distribution on the set of optimal policies (in contrast to, e.g., the action value function which is constant) to our disposal, which constitutes a major technical hurdle. We address this by restricting our discussion to the following special cases: (1) the case when the support of the initial distribution includes \bar{S}_{λ_0} , which will be discussed in section 7.1, and (2) the case when an optimal policy is unique on \bar{S}_{λ_0} , which will be discussed in section 7.2.

7.1 Asymptotic Continuity for Special Initial Distributions

The main result of this section is to demonstrate that the set of accumulation points of the sequence of eUDRL-generated policies is relatively continuous at a deterministic kernel λ_0 if

 $\bar{S}_{\lambda_0} \subset \operatorname{supp} \bar{\mu}$ (see theorem 25). Similar to the preceding sections, the core idea that enables our asymptotic continuity analysis is to derive bounds for the eUDRL policy recursion. This is achieved by bounding various individual terms occurring in the recursion, like Q-values (see lemmas 20 and 21) and visitation probabilities (lemma 22). However, the analysis must be carried out more carefully here as the individual estimates for the numerator and denominator of the policy recursion turn out to be insufficient. This leads us to develop a joint analysis of numerator and denominator during the course of the proof of the main result of this section, theorem 25. Unsurprisingly, bounding the policy recursion via bounds to the respective individual terms can be achieved through monotonicity bounds for the underlying rational function (see remark 23), which plays a key role in our discussion and will be used repeatedly. This finally leads to an estimation of the eUDRL recursion through a dynamical system (which we introduce in lemma 24). This dynamical system possesses a unique fixed point that bounds the accumulation points of eUDRL's policy recursion and enjoys desirable properties as $\lambda \to \lambda_0$.

We derive some preliminary lemmata in section 7.1.1. The main theorem is contained in section 7.1.2. In section 7.1.3 we show how it can be extended to the Seg^{diag/trail} segment spaces. As a corollary, we study the continuity of accumulation point sets also for other quantities like values and the goal-reaching objective in section 7.1.4. This yields upper bounds on the errors of these quantities as a bonus. Somewhat surprisingly, these error bounds turn out to be independent of the initial condition $\pi_0 > 0$. Furthermore, we establish a preliminary result about the convergence rate of the policy error. We conclude with a few examples that illustrate the obtained bounds.

7.1.1 Preliminary Lemmata

Bounding optimal Values: Lemma 12 point 3. asserts that eUDRL values are continuous near 0. We begin by showing a stronger version of the lemma, which makes the continuity explicit by bounding the discrepancy in values given a discrepancy δ in the transition kernels.

Lemma 20 (Continuity at value zero) Let $\{\mathcal{M}_{\lambda}|\lambda \in (\Delta S)^{S \times A}\}$ and $\{\overline{\mathcal{M}}_{\lambda}|\lambda \in (\Delta S)^{S \times A}\}$ be compatible families. Let λ_0 be a deterministic kernel. Choose $\delta \in (0, 2)$ and assume $\lambda \in U_{\delta}(\lambda_0) = \{\lambda | \max_{s,a \in S \times A} ||\lambda(\cdot|s,a) - \lambda_0(\cdot|s,a)||_1 < \delta\}$. Then for any policy π , state $\bar{s} = (s, h, g) \in \bar{S}_T$ and action $a \in \mathcal{A}$ the following holds:

1. Suppose
$$V_{\lambda_0}^*(\bar{s}) = 0$$
 then $V_{\lambda}^{\pi}(\bar{s}) \leq \frac{\delta h}{2} \leq \frac{\delta N}{2}$.

2. Suppose
$$Q_{\lambda_0}^*(\bar{s}, a) = 0$$
 then $Q_{\lambda}^{\pi}(\bar{s}, a) \leq \frac{\delta h}{2} \leq \frac{\delta N}{2}$.

Proof We begin by showing that point 1. implies point 2. Let s^* be the unique state for which $\lambda_0(s^*|s, a) = 1$, let $\bar{s}^* = (s^*, h - 1, g)$. Notice that $V^*_{\lambda_0}(\bar{s}^*) = 0$ (for $h \ge 2$) since the

opposite would contradict the assumption $Q_{\lambda_0}^*(\bar{s}, a) = 0$.

$$\begin{aligned} Q_{\lambda}^{\pi}(\bar{s},a) &= \mathbb{P}_{\lambda}(\rho(S_{h}) = g | \bar{S}_{0} = \bar{s}, A_{0} = a; \pi) = \sum_{s' \in \mathcal{S}} \mathbb{P}_{\lambda}(\rho(S_{h}) = g, \bar{S}_{1} = \bar{s}' | \bar{S}_{0} = \bar{s}, A_{0} = a; \pi) \\ &= \sum_{s' \in \mathcal{S}} \mathbb{P}_{\lambda}(\rho(S_{h}) = g | \bar{S}_{1} = \bar{s}'; \pi) \mathbb{P}_{\lambda}(\bar{S}_{1} = \bar{s}' | \bar{S}_{0} = \bar{s}, A_{0} = a; \pi) \\ &= \mathbb{P}_{\lambda}(\rho(S_{h}) = g | \bar{S}_{1} = \bar{s}^{*}; \pi) \lambda(s^{*} | s, a) + \sum_{s' \in \mathcal{S}, s' \neq s^{*}} \mathbb{P}_{\lambda}(\rho(S_{h}) = g | \bar{S}_{1} = \bar{s}'; \pi) \lambda(s' | s, a), \end{aligned}$$

where $\bar{s}' = (s', h - 1, g)$. If h = 1 the first summand is 0 and the second summand can be bounded by $\sum_{s' \in S, s' \neq s^*} \lambda(s'|s, a) \leq \frac{\delta}{2}$. For $h \geq 2$ we get

$$Q_{\lambda}^{\pi}(\bar{s},a) \leq V_{\lambda}^{\pi}(\bar{s}^*) + \sum_{s' \in \mathcal{S}, s' \neq s^*} \lambda(s'|s,a) \leq \frac{\delta(h-1)}{2} + \frac{\delta}{2}.$$

We show point 1. Recall that $V_{\lambda}^{\pi}(\bar{s}) = \mathbb{P}_{\lambda}(\rho(S_h) = g | \bar{S}_0 = \bar{s})$. Reaching g from s in h steps has a 0 probability under λ_0 because $V_{\lambda_0}^*(\bar{s}) = 0$. Thus we decompose the probability by restricting to the event

 $\Lambda_k = \{ \text{The first transition with 0 probability according to } \lambda_0 \text{ occurred at step } k \},\$

i.e.

$$\mathbb{P}_{\lambda}(\rho(S_h) = g | \bar{S}_0 = \bar{s}) = \sum_{k=1}^h \mathbb{P}_{\lambda}(\rho(S_h) = g, \Lambda_k | \bar{S}_0 = \bar{s}).$$

We bound the individual summands by

$$\begin{aligned} \mathbb{P}_{\lambda}(\rho(S_{h}) &= g, \Lambda_{k} | \bar{S}_{0} = \bar{s}) \\ &= \sum_{\substack{\bar{s}' = (s', h - (k-1), g) \in \bar{S}_{T} \\ \bar{s}'' = (s'', h - k, g) \in \bar{S}_{T}}} \mathbb{P}_{\lambda}(\bar{S}_{k-1} = \bar{s}' | \bar{S}_{0} = \bar{s}) \mathbb{P}_{\lambda}(\bar{S}_{k} = \bar{s}'', \Lambda_{k} | \bar{S}_{k-1} = \bar{s}') \mathbb{P}_{\lambda}(\rho(S_{h}) = g | \bar{S}_{k} = \bar{s}'') \\ &\leq \sum_{\substack{\bar{s}' = (s', h - (k-1), g) \in \bar{S}_{T}}} \sum_{\substack{\bar{s}'' = (s'', h - k, g) \in \bar{S}_{T}}} \mathbb{P}_{\lambda}(\bar{S}_{k-1} = \bar{s}' | \bar{S}_{0} = \bar{s}) \mathbb{P}_{\lambda}(\bar{S}_{k} = \bar{s}'', \Lambda_{k} | \bar{S}_{k-1} = \bar{s}') \\ &= \sum_{\substack{\bar{s}' = (s', h - (k-1), g) \in \bar{S}_{T}}} \mathbb{P}_{\lambda}(\bar{S}_{k-1} = \bar{s}' | \bar{S}_{0} = \bar{s}) \sum_{a \in \mathcal{A}} \sum_{\substack{\bar{s}'' = (s'', h - k, g) \in \bar{S}_{T}}} \lambda(s'', \Lambda_{k} | s', a') \pi(a' | \bar{s}') \\ &\leq \sum_{\substack{\bar{s}' = (s', h - (k-1), g) \in \bar{S}_{T}}} \mathbb{P}_{\lambda}(\bar{S}_{k-1} = \bar{s}' | \bar{S}_{0} = \bar{s}) \sum_{a \in \mathcal{A}} \frac{\delta}{2} \pi(a' | \bar{s}') \leq \frac{\delta}{2}. \end{aligned}$$

Continuity of action-values in quotient topology: The continuity of Q_{λ}^{π} as a function of (π, λ) was proven in lemma 12 point 1. The following lemma establishes a weaker variant that only requires the relative continuity of policies (theorem 16). It establishes the continuity of Q_{λ}^{π} in a factor topology. It will also be useful for the proof of the main theorem below to show this form of continuity for a fixed remaining horizon. This is done in the second part of the lemma, where an explicit estimate is provided that implies the continuity of Q_{λ}^{π} by induction over the remaining horizon.

Lemma 21 (Continuity of action-values in quotient topology) Let $\{\mathcal{M}_{\lambda}|\lambda \in (\Delta S)^{S \times A}\}$ and $\{\overline{\mathcal{M}}_{\lambda}|\lambda \in (\Delta S)^{S \times A}\}$ be compatible families. Let λ_0 be a deterministic kernel. For all $\epsilon > 0$ there exists $\delta > 0$ such that if $\lambda \in U_{\delta}(\lambda_0)$ and if the policy π satisfies $2(1 - \pi(\mathcal{O}(\bar{s})|\bar{s})) < \delta$ for all $\bar{s} \in \overline{S}_{\lambda_0}$ then $|Q_{\lambda}^{\pi}(\bar{s}, \cdot) - Q_{\lambda_0}^{*}(\bar{s}, \cdot)| \leq \epsilon$ for all $\bar{s} \in \overline{S}_{\lambda_0}$. The statement can be made explicit by making use of the following recursive estimate: For all $\bar{s} \in \overline{S}_{\lambda_0}$ with $h \geq 2$ and all $a \in \mathcal{O}(\bar{s})$ it holds that

$$\begin{aligned} |Q_{\lambda}^{\pi}(\bar{s},a) - Q_{\lambda_{0}}^{*}(\bar{s},a)| &\leq \|\lambda(\cdot|s,a) - \lambda_{0}(\cdot|s,a)\|_{1} \\ &+ \max_{\bar{s}' = (s',h-1,g) \in \bar{\mathcal{S}}_{\lambda_{0}}} 2(1 - \pi(\mathcal{O}(\bar{s}')|\bar{s}')) \\ &+ \max_{\bar{s}' = (s',h-1,g) \in \bar{\mathcal{S}}_{\lambda_{0}}} |Q_{\lambda}^{\pi}(\bar{s}',a') - Q_{\lambda_{0}}^{*}(\bar{s}',a')|. \end{aligned}$$

We establish the estimate for the distance of Q_{λ}^{π} values in the lemma for all $h \geq 2$ by induction over the remaining horizon h. The inequality implies continuity at any fixed remaining horizon $h \geq 2$. As Q_{λ}^{π} is continuous for h = 1 the overall continuity of Q_{λ}^{π} in the quotient topology follows.

Proof Base case (h = 1): In this case $Q_{\lambda}^{\pi}((s, 1, g), a) = \sum_{s' \in \rho^{-1}(\{g\})} \lambda(s'|s, a)$ is continuous in λ and independent of π .

Induction: Now assume the estimate in the lemma holds for h-1 and let π^* be an optimal policy for λ_0 . Without loss of generality $\pi^*(a|\bar{s}') \geq \pi(a|\bar{s}')$ for all $a \in \mathcal{O}(\bar{s}')$, $\bar{s}' \in \bar{S}_{\lambda_0}$. For all $\bar{s} = (s, h, g) \in \bar{S}_{\lambda_0}$ and $a \in \mathcal{O}(\bar{s})$ it holds that

$$\begin{split} |Q_{\lambda}^{\pi}(\bar{s},a) - Q_{\lambda_{0}}^{*}(\bar{s},a))| &= \left| \sum_{s'\in\bar{S}} \lambda(s'|s,a) \sum_{a'\in\mathcal{A}} \pi(a'|s',h-1,g) Q_{\lambda}^{\pi}((s',h-1,g),a') \right| \\ &- \sum_{s'\in\bar{S}} \lambda_{0}(s'|s,a) \sum_{a'\in\mathcal{A}} \pi^{*}(a'|s',h-1,g) Q_{\lambda_{0}}^{*}((s',h-1,g),a') \right| \\ &\leq \sum_{s'\in\bar{S}} |\lambda(s'|s,a) - \lambda_{0}(s'|s,a)| \sum_{a'\in\mathcal{A}} \pi(a'|s',h-1,g) Q_{\lambda}^{\pi}((s',h-1,g),a') \\ &+ \sum_{s'\in\bar{S}} \lambda_{0}(s'|s,a) \sum_{a'\in\mathcal{A}} |\pi(a'|s',h-1,g) - \pi^{*}(a'|s',h-1,g)| Q_{\lambda}^{\pi}((s',h-1,g),a') \\ &+ \sum_{s'\in\bar{S}} \lambda_{0}(s'|s,a) \sum_{a'\in\mathcal{A}} \pi^{*}(a'|s',h-1,g) |Q_{\lambda}^{\pi}((s',h-1,g),a') - Q_{\lambda_{0}}^{*}((s',h-1,g),a')| \\ &\leq \|\lambda(\cdot|s,a) - \lambda_{0}(\cdot|s,a)\|_{1} + 2(1 - \pi(\mathcal{O}(\bar{s}'')|\bar{s}'') + \max_{a\in\mathcal{O}(\bar{s}'')} |Q_{\lambda}^{\pi}(\bar{s}'',a') - Q_{\lambda_{0}}^{*}(\bar{s}'',a')| \quad (7.1) \\ &\leq \delta + \delta + \|Q_{\lambda}^{\pi}(\bar{s}'',\cdot) - Q_{\lambda_{0}}^{*}(\bar{s}'',\cdot)\|_{\max} \end{split}$$

where we made use of the fact that if $\lambda_0(\cdot|s, a)$ is deterministic then there is exactly one $s'', \bar{s}'' = (s'', h - 1, g)$ with $\lambda_0(s''|s, a) = 1$ and 0 otherwise. The assumption $||\lambda(s'|s, a) - \lambda_0(s'|s, a)||_1 \leq \delta$ is used in the last line. Further by lemma 6 point 4. $\bar{s}'' \in \bar{\mathcal{S}}_{\lambda_0}$ and it follows that $\sum_{a' \in \mathcal{A}} |\pi(a'|\bar{s}'') - \pi^*(a'|\bar{s}'')| = 2 \sum_{a' \in \mathcal{O}(\bar{s}'')} \pi^*(a'|\bar{s}'') - \pi(a'|\bar{s}'') = 2(1 - \pi(\mathcal{O}(\bar{s}'')|\bar{s}'') < \delta$ since $Q_{\lambda}^{\pi}((s', h - 1, g), a') \leq 1$. Further we used that $\pi^*(a'|\bar{s}'')$ is zero for $a' \notin \mathcal{O}(\bar{s}'')$. It follows from the induction assumption that $|Q_{\lambda}^{\pi}((\bar{s}''), a') - Q_{\lambda_0}^*((\bar{s}''), a')| \to 0$ for $\delta \to 0$ and the right goes to 0. To complete the induction we prove continuity also in the remaining case $a \notin \mathcal{O}(\bar{s})$. For all $\bar{s} = (s, h, g) \in \bar{\mathcal{S}}_{\lambda_0}$ and $a \notin \mathcal{O}(\bar{s})$ it holds

$$|Q_{\lambda}^{\pi}(\bar{s},a) - Q_{\lambda_0}^{*}(\bar{s},a))| = Q_{\lambda}^{\pi}(\bar{s},a) \le \frac{\delta N}{2} \to 0, \ \delta \to 0,$$

where we used $Q_{\lambda_0}^*(\bar{s}, a) = 0$ by lemma 6 point 5. and lemma 20. This completes the induction.

Visitation probabilities are separated from 0: The assumption $\bar{S}_{\lambda_0} \subset \operatorname{supp} \bar{\mu}$ allows one to bound state visitation probabilities in eUDRL recursion rewrites (compare lemma 3).

Lemma 22 (Lower bound on visitation probabilities) Let $\{\mathcal{M}_{\lambda}|\lambda \in (\Delta S)^{S \times A}\}$ and $\{\overline{\mathcal{M}}_{\lambda}|\lambda \in (\Delta S)^{S \times A}\}$ be compatible families. Let λ_0 be a deterministic kernel. Assume that $\overline{S}_{\lambda_0} \subset \operatorname{supp} \overline{\mu}$, where $\overline{\mu}$ denotes the CE's initial distribution. Then for all $\overline{s} = (s, h, g) \in \overline{S}_{\lambda_0}$, all transition kernels λ and all policies π it holds that

$$\mathbb{P}_{\lambda}(H_0^{\Sigma}=h,G_0^{\Sigma}=g|S_0^{\Sigma}=s,l(\Sigma)=h;\pi)>\alpha,$$

where

$$\alpha = \frac{2}{N(N+1)} \min_{\bar{s}\in\bar{\mathcal{S}}_{\lambda_0}} \bar{\mu}(\bar{s}) > 0.$$

Proof Write $v(s, h, g) := \mathbb{P}_{\lambda}(H_0^{\Sigma} = h, G_0^{\Sigma} = g, S_0^{\Sigma} = s, l(\Sigma) = h; \pi)$. By the definition of the segment distribution we obtain that

$$\begin{aligned} v(s,h,g) &= \sum_{\sigma:l(\sigma)=h,\bar{s}_0^{\sigma}=(s,h,g)} c_{\lambda,\pi_n}^{-1} \sum_{t< N-l(\sigma)} \mathbb{P}_{\lambda}(\bar{S}_{t+l(\sigma)} = \bar{s}_{l(\sigma)}^{\sigma}, \dots, \bar{S}_t = \bar{s}_0^{\sigma}; \pi) \\ &\geq \sum_{\sigma:l(\sigma)=h,\bar{s}_0^{\sigma}=(s,h,g)} c_{\lambda,\pi_n}^{-1} \mathbb{P}_{\lambda}(\bar{S}_h = \bar{s}_h^{\sigma}, \dots, \bar{S}_0 = \bar{s}_0^{\sigma}; \pi) \\ &= \sum_{\sigma:l(\sigma)=h,\bar{s}_0^{\sigma}=(s,h,g)} c_{\lambda,\pi_n}^{-1} \mathbb{P}_{\lambda}(\bar{S}_h = \bar{s}_h^{\sigma}, \dots, \bar{S}_1 = \bar{s}_1^{\sigma} | \bar{S}_0 = \bar{s}_0^{\sigma}; \pi) \bar{\mu}(\bar{s}_0^{\sigma}) \\ &= c_{\lambda,\pi}^{-1} \bar{\mu}(\bar{s}_0^{\sigma}). \end{aligned}$$

In the last equation we made use of the fact that by summing over all possible segments with the starting condition (s, h, g) we are actually summing over all h - 1 tuples of extended states compatible with the CE framework, leaving out only tuples whose probability is 0. The latter are tuples where the remaining horizon does not decrease by 1 or where the goal is not constant. By equation (2.4) we have $c_{\lambda,\pi}^{-1} \geq \frac{2}{N(N+1)}$. We choose $\alpha = \frac{2}{N(N+1)} \min_{\bar{s} \in \bar{S}_{\lambda_0}} \bar{\mu}(\bar{s}) > 0$ and conclude for all $(s, h, g) \in \bar{S}_{\lambda_0}$ that

$$\mathbb{P}_{\lambda}(H_0^{\Sigma} = h, G_0^{\Sigma} = g | S_0^{\Sigma} = s, l(\Sigma) = h; \pi) = \frac{v(s, h, g)}{\mathbb{P}_{\lambda}(S_0^{\Sigma} = s, l(\Sigma) = h; \pi)} \ge \frac{v(s, h, g)}{1}$$
$$\ge c_{\lambda, \pi}^{-1} \bar{\mu}(s, h, g) \ge \frac{2}{N(N+1)} \min_{\bar{s} \in \bar{\mathcal{S}}_{\lambda_0}} \bar{\mu}(\bar{s}) = \alpha.$$

Method for bounding the eUDRL recursion: Our analysis of eUDRL's convergence draws on the properties of dynamical systems induced by certain rational functions. Specifically, we proceed describing the eUDRL recursion in terms of a multi-dimensional rational function $F : D \to \mathbb{R}$ on a domain $D = \prod_{k < K} [y_k, 1], y = (y_1, ..., y_{K-1}) \in [0, 1]^K$. The recursive application of this function leads to a set of accumulation points that, in turn, is bounded by studying the fixed points of a simplified dynamical system that emerges through the iterative application of a scalar rational function.

Remark 23 (Monotonicity) The function $g : [0,1] \to \mathbb{R}$ with $g(x) := \frac{ax+b}{ax+b+c}$, with $a, b \ge 0$, c > 0 is non-decreasing.

Bounding of F proceeds in K steps, where in each step k < K the following estimates are used: 1. For all $z \in D$ the function $x_k \mapsto F(z_0, z_1, \ldots, z_{k-1}, x_k, z_{k+1}, \ldots, z_{K-1})$ has the a form specified in the remark. 2. In conclusion for all $z \in D$ it holds that $F(y_0, y_1, \ldots, y_{k-1}, z_k, z_{k+1}, \ldots, z_{K-1}) \geq F(y_0, y_1, \ldots, y_{k-1}, y_k, z_{k+1}, \ldots, z_{K-1})$. Applying this reasoning along all coordinates leads to the estimate $F(z) \geq F(y)$ for all $z \in D$. F(y), in turn, is fully described by a scalar rational function. The following lemma summarizes the features of a dynamical system that arises through the iterative application of this function. Given a function $f: X \to X$ on a set X write $f^{\circ n}: X \to X$ for the *n*-fold composition of f with itself, i.e. $f^{\circ n}(x) = (f \circ f \circ \ldots \circ f)(x)$.

Lemma 24 (f-lemma) Let $\gamma \in (0,1)$ and $f_{\gamma} : (0,1] \to (0,1]$ be defined as $f_{\gamma}(x) = \frac{x}{x+\gamma}$. Then the following assertions hold.

- 1. f_{γ} is strictly monotonically increasing.
- 2. f_{γ} has a unique fixed point $x^*(\gamma) = 1 \gamma$ and

$$\begin{aligned} &f_{\gamma}(x) > x \quad for \ x < x^{*}(\gamma), \\ &f_{\gamma}(x) < x \quad for \ x > x^{*}(\gamma). \end{aligned}$$

- 3. For all $x \in (0,1]$ the iterated application of f_{γ} converges point-wise, $f_{\gamma}^{\circ n}(x) \xrightarrow{n \to \infty} x^*(\gamma)$.
- 4. Assume a sequence $(y_n)_{n\geq 0}$, $y_n \in (0,1]$ such that for all $n \geq 0$ we have $y_{n+1} \geq f_{\gamma}(y_n)$. Then $y_n \geq f_{\gamma}^{\circ n}(y_0)$ and $\liminf_n y_n \geq x^*(\gamma)$.



Figure 5: Illustration of the dynamical system induced by iterative application of f_{γ} . The figure shows the dependence of $f_{\gamma}^{\circ n}(x)$ given initial condition x on iteration n.

Figure 5 illustrates the dynamical system induced by iterative application of f_{γ} .

Proof 1. The derivative of f is positive.

2. Solving the equation $f_{\gamma}(x) = x$ shows that $x^* = 1 - \gamma$ is a unique fixed point. The remaining statements follow by plugging f into the respective inequalities.

3. Write $I_1 = (0, x^*]$ and $I_2 = [x^*, 1]$ and $f_1 = f \upharpoonright_{I_1}, f_2 = f \upharpoonright_{I_2}$. f is increasing so that $f(I_1) \leq f(x^*) = x^*$ and thus $f(I_1) \subset I_1$, i.e. $f_1 : I_1 \to I_1$. Similarly, $f_2 : I_2 \to I_2$. Fix $x \in I_1$ then the sequence $f^{\circ n}(x) = f_1^{\circ n}(x)$ is bounded from above by x^* . The sequence is also increasing by point 2. Thus the sequence converges to a limit in I_1 . Since f_1 is continuous on I_1 , the limit has to be a fixed point. Since x^* is the only fixed point we conclude $f_1^{\circ n}(x) \xrightarrow{n \to \infty} x^*$. Similarly, for $x \in I_2$ we get $f^{\circ n}(x) = f_2^{\circ n}(x) \xrightarrow{n \to \infty} x^*$.

4. We proceed by induction. For n = 0 we have $y_0 \ge f^{\circ 0}(y_0) = y_0$. Assume that $y_n \ge f^{\circ n}(y_0)$. Since f is increasing $f(y_n) \ge f(f^{\circ n}(y_0))$. Making use of $y_{n+1} \ge f(y_n)$ yields $y_{n+1} \ge f(y_n) \ge f(f^{\circ n}(y_0))$, which concludes the induction. $\liminf_n y_n \ge x^*$ then follows directly from point 3.

7.1.2 The main theorem

We describe the continuity of eUDRL-generated policies when the number of iterations approaches infinity. We have already discussed that the set of accumulation points is not empty. We proved in theorem 10 that for every finite iteration the eUDRL-generated policy is optimal if the transition kernel λ_0 is deterministic. Since for all states $\bar{s} \in \bar{S}_{\lambda_0}$ and actions $a \in \mathcal{A} \setminus \mathcal{O}(\bar{s})$ and $n \geq 1$ it holds that $\pi_n(a|\bar{s}) = 0$ it also follows that $\pi_n(a|\bar{s}) \to 0$ as $n \to \infty$. As a consequence the set of accumulation points of the sequence of policies for a deterministic kernel is a subset of $\{\pi_{\lambda_0}^* \mid \pi_{\lambda_0}^* \text{ is an optimal policy for } \lambda_0\}$.

Theorem 25 (Relative continuity of eUDRL limit policies in deterministic kernels – the case $\bar{S}_{\lambda_0} \subset \operatorname{supp} \bar{\mu}$) Let $\{\mathcal{M}_{\lambda} | \lambda \in (\Delta S)^{S \times \mathcal{A}}\}$ and $\{\bar{\mathcal{M}}_{\lambda} | \lambda \in (\Delta S)^{S \times \mathcal{A}}\}$ be compatible families. Let $\lambda_0 \in (\Delta S)^{S \times \mathcal{A}}$ be a deterministic kernel and let $(\pi_{n,\lambda})_{n\geq 0}$ be a sequence of eUDRL-generated policies with initial condition $\pi_0 > 0$ and transition kernel λ . Assume $\bar{S}_{\lambda_0} \subset \operatorname{supp} \bar{\mu}$. Then for all $\pi_0 > 0$ the following holds:

1. Let $\mathcal{L}(\pi_0, \lambda)$ denote the set of accumulation points of $(\pi_{n,\lambda})_{n\geq 0}$. Then any function $u: (\pi_0, \lambda) \mapsto u(\pi_0, \lambda) \in \mathcal{L}(\pi_0, \lambda)$ is relatively continuous in λ at λ_0 on $\bar{\mathcal{S}}_{\lambda_0}$, i.e. for all $\bar{s} \in \bar{\mathcal{S}}_{\lambda_0}$

$$[u(\pi_0,\lambda)](\cdot|\bar{s}) \xrightarrow{\mathcal{O}(\bar{s})} \pi^*_{\lambda_0}(\cdot|\bar{s}) \quad \text{as} \quad \lambda \to \lambda_0.$$

2. Let $\alpha > 0$ be chosen as in lemma 22 and let $\beta, \tilde{\beta} \in (0, 1)$ be such that $\gamma = \frac{\beta}{(1-\beta)\alpha} < 1$ and let

$$U_{\delta}(\lambda_0) = \{\lambda | \max_{(s,a) \in \mathcal{S} \times \mathcal{A}} \|\lambda(\cdot|s,a) - \lambda_0(\cdot|s,a)\|_1 < \delta\}.$$

There exists $\delta > 0$ so that for all $\lambda \in U_{\delta}(\lambda_0)$ and all $\bar{s} \in \bar{S}_{\lambda_0}$ it holds that

$$\liminf_{n} \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \ge x^*(\gamma) \quad \text{and} \quad x^*(\gamma) \to 1 \quad \text{as} \quad (\beta, \tilde{\beta}) \to 0,$$

where $x^*(\gamma) = 1 - \gamma$ (cf. lemma 24). As a consequence for all $\bar{s} \in \bar{S}_{\lambda_0}$ it holds that

$$\liminf_{n} \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \to 1 \quad \text{as} \quad \lambda \to \lambda_0.$$

At the core of the proof stands the analysis of dynamical systems and convergence induced by the iterative application of the rational function $f_{\gamma}(x) = \frac{x}{x+\gamma}$, see lemma 24. Indeed we will show that for all horizons $N \ge h \ge 1$, all initial conditions $\pi_0 > 0$ and all $\beta, \tilde{\beta} \in (0, 1)$ satisfying $\frac{\tilde{\beta}}{(1-\beta)\alpha} = \gamma < 1$ there exists n_0 and $\delta > 0$ such that for all $\lambda \in U_{\delta}(\lambda_0)$ and all $n > n_1 \ge n_0$ and all $\bar{s} \in \bar{S}_{\lambda_0}$ with remaining horizon h it holds that

$$\pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \ge f_{\gamma}^{\circ(n-n_1)}(\pi_{n_1,\lambda}(\mathcal{O}(\bar{s})|\bar{s})).$$

$$(7.2)$$

To apply lemma 24 we need to ensure the positivity of the argument of f_{γ} . By theorem 16, point 3. we have that $\pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) > 0$ for all $n \geq 0$ for all $\lambda \in U_{\delta_n}(\lambda_0)$ for suitable $\delta_n > 0$. However, we will require a slightly stronger statement that also ensures that $\pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) > 0$ for all $n \geq 0, \lambda \in U_2(\lambda_0), \bar{s} \in \bar{S}_{\lambda_0}$ that we show in the course of the main theorem, see equation (7.5). Then by lemma 24 the right hand side converges to $x^*(\gamma) = 1 - \gamma$, which implies point 2. of the theorem, see the proof for details. Furthermore point 1. is an immediate consequence of point 2. We include point 1. to showcase the relation to the notion of "continuity of the set of accumulation points relative to the set of optimal actions" that we mentioned earlier. Point 1. says that any function u that maps (π_0, λ) to an accumulation point is relatively continuous at λ_0 . Since the set of values of all such functions u at (π_0, λ) is exactly $\mathcal{L}(\pi_0, \lambda)$ one can interpret 1. as the relative continuity of the set $\mathcal{L}(\pi_0, \lambda)$. **Proof** It is obvious that point 2. implies point 1. since $\pi_{\lambda_0}^*(\mathcal{O}(\bar{s})|\bar{s}) = 1$ on $\bar{\mathcal{S}}_{\lambda_0}$. Second, we prove that equation (7.2) implies point 2. Fix $\pi_0 > 0$ and choose $\epsilon > 0$. Since $x^*(\gamma) \to 1$ as $\gamma \to 0$ (cf. lemma 24) and $\gamma \to 0$ as $(\beta, \tilde{\beta}) \to 0$ we can pick $\beta, \tilde{\beta} > 0$ so that $x^*(\gamma) > 1 - \epsilon$. Since equation (7.2) holds for every h there are $\delta(h) > 0$ and $n_0(h)$ so that for all $n > n_1 \ge n_0(h), \lambda \in U_{\delta(h)}(\lambda_0)$ and all $\bar{s} \in \bar{\mathcal{S}}_{\lambda_0}$ the equation remains valid. Let us fix $n_1 = \max_h n_0(h)$ and $\delta = \min_h \delta(h)$ and $\eta = \min_{\bar{s} \in \bar{\mathcal{S}}_{\lambda_0}} \pi_{n_1,\lambda}(\mathcal{O}(\bar{s})|\bar{s})$. By lemma 24 $f_{\gamma}^{\circ(n-n_1)}$ is increasing and it follows that $\pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \ge f_{\gamma}^{\circ(n-n_1)}(\eta)$. Since $f_{\gamma}^{\circ n}(\eta) \to x^*(\gamma)$ as $n \to \infty$ we obtain $\liminf_n \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \ge x^*(\gamma) > 1 - \epsilon$. As $\epsilon > 0$ is arbitrary we can always find $\delta > 0$ so that the above holds. We conclude that as $\lambda \to \lambda_0$

$$\liminf_{n} \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \to 1$$

It remains to prove equation (7.2). Fixing $2 > \delta > 0$ lemma 6 point 2. asserts that $\bar{S}_{\lambda_0} \subset \operatorname{supp} \operatorname{den}_{\lambda,\pi_n} \cap \operatorname{supp} \nu_{\lambda,\pi_n}$ for all $n \ge 0, \pi_0 > 0, \lambda \in U_{\delta}(\lambda_0)$. The policy $\pi_{n+1,\lambda}(\cdot|\bar{s})$ is then well-defined by

$$\pi_{n+1,\lambda}(\cdot|\bar{s}) = \frac{\operatorname{num}_{\lambda,\pi_n}(\cdot|\bar{s})}{\operatorname{den}_{\lambda,\pi_n}(\bar{s})}$$
(7.3)

for all $\bar{s} \in \bar{S}_{\lambda_0}$. Using the notation

$$\begin{aligned} Q_{\lambda}^{\pi_n,g,h}((s,h',g'),a) &:= \mathbb{P}_{\lambda}(\rho(S_h^{\Sigma}) = g | A_0^{\Sigma} = a, S_0^{\Sigma} = s, H_0^{\Sigma} = h', G_0^{\Sigma} = g'; \pi_n), \\ v_{\lambda,\pi_n}(h',g'|s,h) &:= \mathbb{P}_{\lambda}(\rho(H_0^{\Sigma}) = h', G_0^{\Sigma} = g' | S_0^{\Sigma} = s, l(\Sigma) = h; \pi_n) \end{aligned}$$

and writing $\bar{s} = (s, h, g) \in \bar{S}_{\lambda_0}$ lemma 3 point 1. allows one to rewrite the policy as

$$\pi_{n+1,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) = \frac{\sum_{a \in \mathcal{O}(\bar{s}), h' \ge h, g' \in \mathcal{G}} Q_{\lambda}^{\pi_n, g, h}((s, h', g'), a) \pi_{n,\lambda}(a|s, h', g') v_{\lambda, \pi_n}(h', g'|s, h)}{\sum_{a \in \mathcal{A}, h' \ge h, g' \in \mathcal{G}} Q_{\lambda}^{\pi_n, g, h}((s, h', g'), a) \pi_{n,\lambda}(a|s, h', g') v_{\lambda, \pi_n}(h', g'|s, h)}.$$
(7.4)

We show some estimates that will be used later in the proof.

Bounding state visitation: By lemma 22 and using $\bar{s} \in \bar{S}_{\lambda_0} \subset \text{supp } \bar{\mu}$ there exists $\alpha > 0$ so that

$$v_{\lambda,\pi_n}(h,g|s,h) > \alpha.$$

Separation of $\pi_{n,\lambda}$ from 0 on optimal actions: We will show that for each $n \ge 0$ there exist $\eta > 0$ such that for all $\bar{s} \in \bar{S}_{\lambda_0}$, $a \in \mathcal{O}(\bar{s})$, $\lambda \in U_{\delta}(\lambda_0)$ it holds that

$$\pi_{n,\lambda}(a \mid \bar{s}) > \eta > 0. \tag{7.5}$$

The proof proceeds by induction. The base case n = 0 follows since $\pi_0 > 0$ does not depend on λ . For the induction step assume $\pi_{n,\lambda}(a \mid \bar{s}) > \eta' > 0$ for all (\bar{s}, a, λ) above. Using (7.4) we get $\pi_{n,\lambda}(a \mid \bar{s}) \ge Q_{\lambda}^{\pi_n}(\bar{s}, a)\pi_n(a \mid \bar{s})v_{\lambda,\pi_n}(h, g \mid s, h)$. Because $\bar{s} = (s, h, g) \in \bar{S}_{\lambda_0}$ and $a \in \mathcal{O}(\bar{s})$ we can fix a sequence $\bar{s}_0, a_0, \ldots, \bar{s}_h$ with $\bar{s}_0 = \bar{s}, a_0 = a, \rho(s_h) = g$ evidencing that the probability $Q_{\lambda_0}^*(\bar{s}, a)$ is positive. We deduce that, for all $i = 0, \ldots, h - 1$, all \bar{s}_{i+1} belongs to \bar{S}_{λ_0} and all actions a_i are optimal and $\lambda_0(s_{i+1} \mid s_i, a_i) = 1$. Thus we can estimate $\lambda(s_{i+1} \mid s_i, a_i) > 1 - \frac{\delta}{2}$ (since $\lambda_0(s_{i+1} \mid s_i, a_i) - \lambda(s_{i+1} \mid s_i, a_i) < \frac{\delta}{2}$) and $\pi_{n,\lambda}(a_i \mid \bar{s}) > \eta'$. This gives us the desired lower bound

$$\pi_{n+1,\lambda}(a \mid \bar{s}) \ge \left(\prod_{i=0}^{h-1} \lambda(s_{i+1} \mid s_i, a_i) \pi_{n,\lambda}(a_i \mid s_i)\right) \cdot \alpha \ge (1 - \frac{\delta}{2})^h (\eta')^h \alpha > 0.$$

This completes proof of (7.5) by induction.

Bounding $Q_{\lambda}^{\pi_n,g,h}$: By lemma 12 and lemma 10 we have that for any policy π for all $\bar{s} = (s,h,g)$

$$Q_{\lambda}^{\pi,g,h}((s,h',g'),a) \le Q_{\lambda}^*((s,h,g),a) \to Q_{\lambda_0}^*((s,h,g),a), \ \lambda \to \lambda_0.$$

$$(7.6)$$

The proof proceeds by induction on the remaining horizon h. For the rest of the proof π_0 is assumed fixed. Base case (h = 1): We estimate $\pi_{n+1,\lambda}(\mathcal{O}(\bar{s})|\bar{s})$ by showing lower estimates on the recursion (7.4). By lemma 21 we have that if $\bar{s} = (s, 1, g) \in \bar{S}_{\lambda_0}$ and $a \in \mathcal{O}(\bar{s})$ then

$$|Q_{\lambda}^{\pi_{n}}(\bar{s},a) - Q_{\lambda_{0}}^{*}(\bar{s},a)| = |Q_{\lambda}^{\pi_{n}}(\bar{s},a) - 1| \to 0, \ \lambda \to \lambda_{0}.$$
(7.7)

Since h = 1 $Q_{\lambda}^{\pi_n}(\bar{s}, a)$ does not depend on π_n . Fix $\beta, \tilde{\beta} \in (0, 1)$ so that $1 > \gamma = \frac{\tilde{\beta}}{(1-\beta)\alpha}$. By (7.6) and (7.7) we can fix $2 > \delta > 0$ so that for all $\lambda \in U_{\delta}(\lambda_0)$ and $\bar{s} = (s, 1, g) \in \bar{S}_{\lambda_0}$ it holds that

$$\beta > Q_{\lambda}^{*}(\bar{s}, a), \text{ for } a \notin \mathcal{O}(\bar{s}), 1 - \beta < Q_{\lambda}^{\pi_{n}}(\bar{s}, a), \text{ for } a \in \mathcal{O}(\bar{s}).$$

$$(7.8)$$

We proceed by bounding $\pi_{n+1,\lambda}(\mathcal{O}(\bar{s})|\bar{s})$ in (7.4) from below. For the denominator we find using (7.8) and (7.6) that

$$\begin{split} \sum_{a \in \mathcal{A}, h' \geq 1, g' \in \mathcal{G}} Q_{\lambda}^{\pi_n, g, 1}((s, h', g'), a) \pi_{n, \lambda}(a|s, h', g') v_{\lambda, \pi_n}(h', g'|s, 1) \\ & \leq \sum_{a \in \mathcal{O}(\bar{s}), h' \geq 1, g' \in \mathcal{G}} Q_{\lambda}^{\pi_n, g, 1}((s, h', g'), a) \pi_{n, \lambda}(a|s, h', g') v_{\lambda, \pi_n}(h', g'|s, 1) + \tilde{\beta}. \end{split}$$

Applying the monotonicity remark 23 (M) to exploit the bounds in (7.8) and $v_{\lambda,\pi_n}(h,g|s,h) > \alpha$ yields that $(\forall n \ge 0, \forall \lambda \in U_{\delta}(\lambda_0), \forall \bar{s} = (s, 1, g) \in \bar{S}_{\lambda_0})$

$$\pi_{n+1,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \geq^{(M)} \frac{\sum\limits_{a\in\mathcal{O}(\bar{s})} (1-\beta)\pi_{n,\lambda}(a|s,1,g)v_{\lambda,\pi_n}(1,g|s,1)}{\sum\limits_{a\in\mathcal{O}(\bar{s})} (1-\beta)\pi_{n,\lambda}(a|s,1,g)v_{\lambda,\pi_n}(1,g|s,1)+\tilde{\beta}}$$

$$\geq^{(M)} \frac{\sum\limits_{a\in\mathcal{O}(\bar{s})} (1-\beta)\pi_{n,\lambda}(a|s,1,g)\alpha}{\sum\limits_{a\in\mathcal{O}(\bar{s})} (1-\beta)\pi_{n,\lambda}(a|s,1,g)\alpha+\tilde{\beta}}$$

$$= \frac{(1-\beta)\alpha\pi_{n,\lambda}(\mathcal{O}(\bar{s})|s,1,g)}{(1-\beta)\alpha\pi_{n,\lambda}(\mathcal{O}(\bar{s})|s,1,g)+\tilde{\beta}} = f_{\gamma}(\pi_{n,\lambda}(\mathcal{O}(\bar{s})|s,1,g)). \quad (7.9)$$

Induction: Fix h > 1 and assume that (7.2) holds for all h' < h. Fix $\beta, \tilde{\beta} \in (0, 1)$ so that $1 > \gamma = \frac{\tilde{\beta}}{(1-\beta)\alpha}$. By lemma 21 there exists δ' such that if the following two conditions are met

$$\lambda \in U_{\delta'}(\lambda_0),$$
$$(\forall \bar{s}' = (s', h', g') \in \bar{\mathcal{S}}_{\lambda_0}, h' < h): \ 1 - \pi(\mathcal{O}(\bar{s}')|\bar{s}') < \frac{\delta'}{2}$$

then it holds that

$$(\forall \bar{s} = (s, h, g) \in \bar{\mathcal{S}}_{\lambda_0}, \forall a \in \mathcal{O}(\bar{s})), \lambda \in U_{\delta'}(\lambda_0) : |1 - Q_{\lambda}^{\pi}(\bar{s}, a)| < \beta.$$

The first condition is met by the choice $0 < \delta < \delta' < 2$. To meet the second condition we will use the induction assumption choosing $\beta', \tilde{\beta}' > 0$ such that $\gamma' = \frac{\tilde{\beta}'}{(1-\beta')\alpha} < 1$ and $x^*(\gamma') > 1 - \delta'/2$. Applying the induction assumption requires a restriction on δ . From the induction assumption and equation (7.5) it follows that there exist n'_0 and $\eta > 0$ such that for all $n > n'_0$, $\lambda \in U_{\delta}(\lambda_0)$ holds $\pi_{n,\lambda}(\mathcal{O}(\bar{s}')|\bar{s}') > f_{\gamma'}^{\circ n-n'_0}(\eta)$. As $f_{\gamma'}^{\circ n-n'_0}(\eta) \to x^*(\gamma')$ there exists n_0 such that $\pi_{n,\lambda}(\mathcal{O}(\bar{s}')|\bar{s}') > 1 - \delta'/2$ for $n > n_0$, $\lambda \in U_{\delta}(\lambda_0)$. Making use of (7.6) we can assume that $\delta > 0$ is chosen such that

$$(\forall \lambda \in U_{\delta}(\lambda_0), \forall \bar{s} \in \bar{\mathcal{S}}_{\lambda_0}, a \notin \mathcal{O}(\bar{s})) : Q_{\lambda}^*(\bar{s}, a) < \beta,$$

We proceed by bounding $\pi_{n+1,\lambda}(\mathcal{O}(\bar{s})|\bar{s})$ in (7.4) from below. Assuming that $n \geq n_0$ we can utilize the obtained estimates as we already did for h = 1. As a consequence (for all $\bar{s} \in \bar{S}_{\lambda_0}$ with remaining horizon h and $\lambda \in U_{\delta}(\lambda_0)$) it holds

$$\pi_{n+1,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \geq \frac{\sum_{a \in \mathcal{O}(\bar{s}), h' \geq h, g' \in \mathcal{G}} Q_{\lambda}^{\pi_n, h, g}((s, h', g'), a) \pi_{n,\lambda}(a|s, h', g') v_{\lambda, \pi_n}(h', g'|s, h)}{\sum_{a \in \mathcal{A}, h' \geq h, g' \in \mathcal{G}} Q_{\lambda}^{\pi_n, h, g}((s, h', g'), a) \pi_{n,\lambda}(a|s, h', g') v_{\lambda, \pi_n}(h', g'|s, h)} \\ \geq \frac{\sum_{a \in \mathcal{O}(\bar{s})} (1 - \beta) \pi_{n,\lambda}(a|s, h, g) \alpha}{\sum_{a \in \mathcal{O}(\bar{s})} (1 - \beta) \pi_{n,\lambda}(a|s, h, g) \alpha + \tilde{\beta}} = \frac{(1 - \beta) \alpha \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s})}{(1 - \beta) \alpha \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) + \tilde{\beta}} \\ = f_{\gamma}(\pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s})),$$

which competes the proof.

An illustration of the accumulation points of $\min_{\bar{s}\in\bar{S}_{\lambda_0}} \pi_n(\mathcal{O}(\bar{s})|s)$ and the lower estimate provided by theorem 25, point 2. is given in Figure 6 in the case of a 2-armed bandit model. All details of the construction of the MDP underlying figure 6 can be found in the appendix B, example 4. Figure 6a shows how the lower bound derived in the theorem and the accumulation points of eUDRL-generated policies behave as functions of the distance to a deterministic kernel. Figure 6b shows how eUDRL approaches those accumulation points as a function of both the iteration n and also the initial policy (which is visible at n = 0). Remark that while the theorem does specify the behavior of γ as $\lambda \to \lambda_0$, it does not provide quantitative information on the specific dependency of γ on δ . The derivation of



(a) Dependency on distance to determin. kernel.

(b) Dependency on iteration.

Figure 6: Illustration of estimates of theorem 25, point 2. Plots show the behavior of $\min_{\bar{s}\in\bar{S}_{\lambda_0}} \pi_n(\mathcal{O}(\bar{s})|s)$ for a 2-armed bandit. Plot (a) shows the dependency on the distance δ to a deterministic kernel, where orange points depict accumulation points (approximated by values at n = 100) and the estimate is depicted in blue (cf. corollary 27). To illustrate the approach of eUDRL towards the accumulation points plot (b) shows the dependency on the iteration n for varying distances to a deterministic kernel (highlighted by different colors) and varying initial policy.

such estimates is the content of section 7.1.4 below. The graph for $x^*(\gamma)$ in Figure 6a is computed using the estimates of corollary 27 (which provides the quantitative details for theorem 25). Regrettably, the bounds derived from the corollary tend to become quite loose as the number of states and the maximum horizon increase. This limitation is an artifact of the bounding of visitation terms in the eUDRL recursion, which was handled somewhat crudely in lemma 22. We explore this issue in greater detail in section 7.1.4. However, the primary aim of section 7.1 was to establish the (relative) continuity of accumulation points of eUDRL-generated policies and the associated goal-reaching objective. The bounds are rather an unexpected byproduct of this work. We emphasize that despite this limitation the proof of continuity and the bounds themselves have not been previously investigated or mentioned in the literature.

7.1.3 Extending the continuity results to other segment sub-spaces

As before we need to modify theorem 25 to cover algorithms like ODT, RWR (and some specific variants of eUDRL) restricting the recursion to Seg^{trail} or Seg^{diag} subspaces.

Theorem 26 (Relative continuity of accumulation points of eUDRL-generated policies at deterministic kernels – the case $\bar{S}_{\lambda_0} \subset \operatorname{supp} \bar{\mu}$) Theorem 25 remains valid under the renaming $\pi_n \to \pi_n^{\text{diag/trail}}$.

The proof of theorem 26 follows along similar lines as that of theorem 25. The differences are rather small and resemble those described in section 6.3. First, to assert that eUDRL is optimal at deterministic points one applies lemma 11 instead of lemma 10. Second, to prove that that $\pi_{n+1}^{\text{diag/trail}}$ is well-defined through equation (7.3) one uses equation (4.5) and lemma 7, point 2. instead of lemma 6. Third, the application of point 1. of lemma 3 in equation (7.4) has to be replaced with points 2. or 3. of the lemma respectively. To bound the denominator of equation 7.4 one proceeds along the same line that leads to equation (7.9), yielding the exactly same result as in (7.9). The same reasoning applies to the induction step. Finally, using lemma 3, which allows one to rewrite the eUDRL recursion using only Seg-space quantities, we can show that there are no further differences even for $\pi_{n+1}^{\text{diag/trail}}$. The same approach applies also to the forthcoming corollary of theorem 25, which follows theorem 25 but provides explicit estimates for eUDRL-generated quantities.

7.1.4 Estimating the location of accumulation points

As noted before an important motivation for this work has been to determine whether or not continuing the eUDRL iteration from a certain state is expected to improve the policy. This can be achieved through explicit estimates for eUDRL-generated quantities and their distance to optimality in terms of the distance of transition kernels. The following corollary can be proved along the same lines as theorems 25 and 26 taking care of quantitative estimates explicitly. Recall that a convergent sequence $(y_n)_{n\geq 0}$ with limit y is said to be q-linearly convergent with rate γ if and only if

$$\lim_{n \to \infty} \left| \frac{y_{n+1} - y}{y_n - y} \right| = \gamma \in (0, 1).$$

Corollary 27 (Estimating the location of accumulation points – the case $\bar{S}_{\lambda_0} \subset \operatorname{supp} \bar{\mu}$) Under the conditions of theorem 25 assume that $1 > \delta > 0$ and set, as before, $\alpha = \frac{2}{N(N-1)} \min_{\bar{s} \in \bar{S}_{\lambda_0}} \bar{\mu}(\bar{s})$. Define the quantities $\tilde{\beta} = \frac{N\delta}{2}$, $x^*(\gamma) = 1 - \gamma$ (cf. lemma 24) and for a horizon h, $1 \leq h \leq N$,

$$\beta_h = \begin{cases} \max\{\delta, \tilde{\beta}\}, \text{ if } h = 1, \\ \delta + \kappa_{h-1} + \beta_{h-1}, \text{ if } h \ge 2, \end{cases}$$
$$\gamma_h = \frac{\tilde{\beta}}{(1 - \beta_h)\alpha}, \\ \kappa_h = 2(1 - x^*(\gamma_h)).$$

Further assume $\beta_h, \gamma_h \in (0,1)$ and notice that $\beta_h, \gamma_h, \kappa_h$ are increasing in h and that $\beta_h, \gamma_h, \kappa_h \to 0$ as $\delta \to 0$. Then the following assertions hold for all $\lambda \in U_{\delta}(\lambda_0)$ and for all $\pi_0 > 0$:

- 1. $\limsup_{n} \max_{\bar{s} \in \bar{\mathcal{S}}_{\lambda_0}} 2(1 \pi_n(\mathcal{O}(\bar{s})|\bar{s})) \le \kappa_N,$
- 2. $\limsup_{n} \max_{(\bar{s},a)\in\bar{\mathcal{S}}_{\lambda_{0}}\times\mathcal{A}} |Q_{\lambda}^{\pi_{n}}(\bar{s},a) Q_{\lambda_{0}}^{*}(\bar{s},a)| \leq \beta_{N},$

- 3. $\limsup_{n} \max_{\bar{s} \in \bar{\mathcal{S}}_{\lambda_0}} |V_{\lambda}^{\pi_n}(\bar{s}) V_{\lambda_0}^*(\bar{s})| \le \beta_N + \kappa_N,$
- 4. $\limsup_{n} |J_{\lambda}^{\pi_{n}} J_{\lambda_{0}}^{*}| \leq \frac{N\delta}{2} + \beta_{N} + \kappa_{N}$, and
- 5. for all $\epsilon > 0$ there exist a sequence of λ_0 -optimal policies $(\pi^*_{\lambda_0,n})_{n\geq 0}$ and n_0 so that for all $n \geq n_0$ and for all $\bar{s} \in \bar{S}_{\lambda_0}$ it holds that

$$\|\pi_{n,\lambda}(\cdot|\bar{s}) - \pi^*_{\lambda_0,n}(\cdot|\bar{s})\|_1 \le 2(1 - f^{\circ(n-n_0)}_{\gamma'}(x_0)),$$

where $x_0 = \min_{\bar{s} \in \bar{S}_{\lambda_0}} \pi_{n_0,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) > 0, \ \beta' = \beta_N + \epsilon$, where ϵ is chosen such that $\gamma' = \frac{\tilde{\beta}}{(1-\beta')\alpha} \in (0,1).$

The terms on the right hand side of 1.-4. converge to 0 as $\delta \to 0$. In 5. the sequence $(y_n)_{n\geq 0}$ defined by $y_n = 2(1 - f_{\gamma'}^{\circ(n-n_0)}(x_0))$ converges q-linearly to the limit $y = 2(1 - x_{\gamma'}^*)$ at a convergence rate of γ' which tends to 0 as $\delta \to 0$.

The bounds in the corollary are solely dependent on α and δ , where α is determined by the initial distribution and δ represents the distance to a deterministic kernel. Notably, there is no dependence on the initial policy $\pi_0 > 0$. This feature allows the estimates to be applied even when no information about the initial policy is available. However, this comes at the cost of producing relatively loose estimates. While the initial policy does not affect the asymptotic estimates, it can affect convergence time, compare figure 6b and figure 5 and consider the effect of x_0 in the point 5. All estimates converge to 0 as $\delta \to 0$, which implies the continuity of the respective quantities at deterministic kernels (relative continuity in the case of the policy). This can be interpreted as a form of continuity of the set of accumulation points (see the remark below). It will also sometimes be more convenient to work with $\pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s})$ rather than $2(1 - \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}))$. We summarize this in a remark.

Remark 28 The estimate in point 1. of corollary 27 is equivalent to the following estimate $\liminf_{\bar{s}\in\bar{S}_{\lambda_0}} \pi_n(\mathcal{O}(\bar{s})|\bar{s}) \geq x^*(\gamma_N)$, cf. figure 6, 7. The estimate in point 5. of the corollary is equivalent to $\pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \geq f_{\gamma'}^{\circ(n-n_0)}(x_0)$.

Remark 29 The estimates in the corollary can be stated also in terms of accumulation points of eUDRL-generated quantities. For example point 1. is equivalent to (for all $\lambda \in U_{\delta}(\lambda_0)$ and $\pi_0 > 0$)

$$\sup_{\pi \text{ accumulation point of } (\pi_{n,\lambda})} \max_{\bar{s} \in \bar{\mathcal{S}}_{\lambda_0}} 2(1 - \pi(\mathcal{O}(\bar{s})|\bar{s})) \leq \kappa_N$$

Proof The choice of $1 > \delta > 0$ and the form of $U_{\delta}(\lambda_0)$ is compliant with lemma 6. By this we mean that for all δ (with $U_{\delta}(\lambda_0)$ defined as above) the points 1., 2., and 3. of the lemma 6 hold. Similarly, the choice of α is compliant with lemma 22. The restriction on γ_h : $1 > \gamma_h > 0$ comes from lemma 24, which ensures that we can use them safely with $f_{\gamma}, x^*(\gamma)$ as defined in this lemma. 1. The result for the limits of $\beta_h, \kappa_h, \gamma_h, \tilde{\beta}$ follows from their definitions and the fact that $x^*(\gamma)$ is a decreasing function in γ with a limit 1 for $\gamma \to 0$. Further, that β_h, κ_h are increasing in h follows again from their definition and the previously stated properties of the function x^* .

During the proof we will also utilize that for all $\lambda \in U_{\delta}(\lambda_0), \bar{s} \in \bar{S}_{\lambda_0}, a \notin \mathcal{O}(\bar{s})$ holds

$$Q_{\lambda}^*(\bar{s}, a) \le \beta \le \beta_h$$

where $N \ge h \ge 1$. The first inequality follows from lemma 20. The inequality $\tilde{\beta} < \beta_1$ follows from the definition of β_1 . The rest follows from the previously stated increasing property of β_h . The result about $Q^*_{\lambda}(\bar{s}, a)$ is useful for ensuring that all action value function-related quantities $Q^{\pi_n,h,g}_{\lambda}((s, h', g'), a)$ for non-optimal actions a are bounded by β_h .

the claim: First we aim to prove the following claim:

$$\begin{aligned} (\forall N \ge h \ge 1, \forall \pi_0 > 0, \forall \lambda \in U_{\delta}(\lambda_0)) : \limsup_{\substack{n \\ a \in \mathcal{A}}} \max_{\substack{\bar{s} = (s, h', g) \in \bar{\mathcal{S}}_{\lambda_0}, h' = h, \\ a \in \mathcal{A}}} |Q_{\lambda}^{\pi_n}(\bar{s}, a) - Q_{\lambda_0}^*(\bar{s}, a)| \le \beta_h, \\ \limsup_{\substack{n \\ \bar{s} = (s, h', g) \in \bar{\mathcal{S}}_{\lambda_0}, h' = h}} \max_{\substack{2(1 - \pi_n(\mathcal{O}(\bar{s})|\bar{s})) \le \kappa_h.}} \end{aligned}$$

We will proceed by induction by the remaining horizon h.

the claim, base case: Assume h = 1 and $(s, 1, g) \in \overline{S}_{\lambda_0}$. It holds

$$\begin{aligned} |Q_{\lambda}^{\pi_{n}}((s,1,g),a) - Q_{\lambda_{0}}^{*}((s,1,g),a)| &= \left| \sum_{s' \in \rho^{-1}(\{g\})} \lambda(s'|s,a) - \sum_{s' \in \rho^{-1}(\{g\})} \lambda_{0}(s'|s,a) \right| \\ &\leq \sum_{s' \in \rho^{-1}(\{g\})} |\lambda(s'|s,a) - \lambda_{0}(s'|s,a)| \leq \|\lambda(\cdot|s,a) - \lambda_{0}(\cdot|s,a)\|_{1} \end{aligned}$$

which gives us, for all $\pi_0 > 0$, $\lambda \in U_{\delta}(\lambda_0)$ and $n \ge 0$,

$$\max_{\substack{\bar{s}=(s,1,g)\in\bar{\mathcal{S}}_{\lambda_0},\\a\in\mathcal{A}}} |Q_{\lambda}^{\pi_n}((s,1,g),a) - Q_{\lambda_0}^*((s,1,g),a)| \leq \max_{\substack{\bar{s}=(s,1,g)\in\bar{\mathcal{S}}_{\lambda_0},\\a\in\mathcal{A}}} \|\lambda(\cdot|s,a) - \lambda_0(\cdot|s,a)\|_1$$
$$\leq \delta \leq \beta_1,$$

from which the first inequality in the claim follows. For the second inequality, we will proceed like in (7.9) bounding $v_{\lambda,\pi_n}(h,g|s,h)$ terms by α and action value function-related terms using $\beta_1, \tilde{\beta}$. Notice that for h = 1 the lim \sup_n does not really matter (see the last equation: β_1 bounds the error for all n). This gives us (for all $n \ge 0$, $\pi_0 > 0$, $\lambda \in U_{\delta}(\lambda_0)$, $\bar{s} = (s, 1, g) \in \bar{S}_{\lambda_0}$)

$$\pi_{n+1,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \ge f_{\gamma_1}(\pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s})).$$

Using point 4. of lemma 24 then leads to

$$\liminf_{x \to \infty} \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \ge x^*(\gamma_1).$$

After rearranging, we get (for all $\pi_0 > 0, \lambda \in U_{\delta}(\lambda_0)$)

$$\limsup_{n} \max_{\bar{s}=(s,1,g)\in\bar{\mathcal{S}}_{\lambda_0}} 2(1-\pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s})) \le 2(1-x^*(\gamma_1)) = \kappa_1.$$

the claim, the induction step $h \ge 2$: Now assume that the claim holds for h-1. We aim to prove it for h. For the first inequality of the claim we will use the the recursive estimate from the lemma 21 (after substituting π with π_n and extending maximization in the last term from $a' \in \mathcal{O}(\bar{s}')$ to $a' \in \mathcal{A}$):

$$\begin{aligned} (\forall \bar{s} = (s, h, g) \in \mathcal{S}_{\lambda_0}, h \ge 2, \forall a \in \mathcal{O}(\bar{s}), \forall n \ge 0) : \\ |Q_{\lambda}^{\pi_n}(\bar{s}, a) - Q_{\lambda_0}^*(\bar{s}, a)| \le \|\lambda(\cdot|s, a) - \lambda_0(\cdot|s, a)\|_1 + \max_{\bar{s}' = (s', h-1, g) \in \bar{\mathcal{S}}_{\lambda_0}} 2(1 - \pi_n(\mathcal{O}(\bar{s}')|\bar{s}')). \\ &+ \max_{\bar{s}' = (s', h-1, g) \in \bar{\mathcal{S}}_{\lambda_0}, a' \in \mathcal{A}} |Q_{\lambda}^{\pi_n}(\bar{s}', a') - Q_{\lambda_0}^*(\bar{s}', a')|. \end{aligned}$$

Now we bound $\|\lambda(\cdot|s,a) - \lambda_0(\cdot|s,a)\|_1$ by δ and apply $\limsup_n(\cdot)$ on both sides of the inequality leading to

$$\limsup_{n} |Q_{\lambda}^{\pi_{n}}(\bar{s}, a) - Q_{\lambda_{0}}^{*}(\bar{s}, a)| \leq \delta + \limsup_{n} \max_{\bar{s}' = (s', h-1, g) \in \bar{\mathcal{S}}_{\lambda_{0}}} 2(1 - \pi_{n}(\mathcal{O}(\bar{s}')|\bar{s}')) \\ + \limsup_{n} \max_{\bar{s}' = (s', h-1, g) \in \bar{\mathcal{S}}_{\lambda_{0}}, a' \in \mathcal{A}} |Q_{\lambda}^{\pi_{n}}(\bar{s}', a') - Q_{\lambda_{0}}^{*}(\bar{s}', a')| \leq \delta + \kappa_{h-1} + \beta_{h-1} = \beta_{h},$$

where we applied the induction assumption. Further for all $\bar{s} = (s, h, g) \in \bar{S}_{\lambda_0}$ and $a \notin \mathcal{O}(\bar{s})$ holds $|Q_{\lambda}^{\pi_n}(\bar{s}, a) - Q_{\lambda_0}^*(\bar{s}, a)| = Q_{\lambda}^{\pi_n}(\bar{s}, a) \leq Q_{\lambda}^*(\bar{s}, a) \leq \tilde{\beta} \leq \beta_1 \leq \beta_h$. After applying \limsup_n we get

$$\limsup_{n} |Q_{\lambda}^{\pi_n}(\bar{s}, a) - Q_{\lambda_0}^*(\bar{s}, a)| \le \beta_h.$$

Putting both results together and maximizing over \bar{s}, a on the left-hand side (it is possible since the right-hand side is independent of \bar{s}, a and \bar{S} and A are finite) we obtain the first inequality of the claim.

Now we aim to prove the second inequality of the claim. We proceed similarly as in the case h = 1, the difference is that now \limsup_n in the previously proved inequality with β_h is important. To be able to bound the errors of the action value function-related terms for the horizon h we have to enlarge β_h ; define $\beta' = \beta_h + \epsilon$, with $\epsilon > 0$ being a fixed arbitrary value so that $1 > \gamma' := \frac{\tilde{\beta}}{(1-\beta')\alpha}$; and fix the π_n sequence such that $\pi_0 > 0$, $\lambda \in U_{\delta}(\lambda_0)$. Then we can find n_0 such that for all $n \ge n_0$ it holds that (note that \bar{S}_{λ_0} is finite)

$$\max_{\bar{s}=(s,h',g)\in\bar{\mathcal{S}}_{\lambda_0},h'=h,a\in\mathcal{A}}|Q_{\lambda}^{\pi_n}(\bar{s},a)-Q_{\lambda_0}^*(\bar{s},a)|\leq\beta'.$$

Then we can continue as before and get for all $n \ge n_0$, and all $\bar{s} = (s, h', g) \in \bar{S}_{\lambda_0}, h' = h$

$$\pi_{n+1,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) = f_{\gamma'}(\pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s})).$$

Using point 4. of lemma 24 then leads again to (for all $\bar{s} = (s, h', g) \in \bar{S}_{\lambda_0}, h' = h$)

$$\liminf_{n} \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \ge x^*(\gamma')$$

Since we could choose $\epsilon > 0$ arbitrarily (and $x^*(\gamma)$ is continuous in γ and $\gamma = \frac{\tilde{\beta}}{(1-\beta)\alpha}$ is continuous in β) we get (for all $\bar{s} = (s, h', g) \in \bar{S}_{\lambda_0}, h' = h$)

$$\liminf_{n} \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \ge x^*(\gamma_h),$$

and since the right side is independent of the fixed sequence we get

$$(\forall \pi_0 > 0, \forall \lambda \in U_{\delta}(\lambda_0), \forall \bar{s} = (s, h', g) \in \bar{\mathcal{S}}_{\lambda_0}, h' = h) : \liminf_{n} \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \ge x^*(\gamma_h).$$

From this follows

$$(\forall \pi_0 > 0, \forall \lambda \in U_{\delta}(\lambda_0)): \limsup_{n} \max_{\bar{s} = (s,h',g) \in \bar{\mathcal{S}}_{\lambda_0}, h' = h} 2(1 - \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \le 2(1 - x^*(\gamma_h)) = \kappa_h.$$

This completes the proof of the claim from which also follows both points 1. and 2. 3. For all $\lambda \in U_{\delta}(\lambda_0), \pi_0 > 0, n \ge 0, \bar{s} \in \bar{S}_{\lambda_0}$ it holds that

$$\begin{aligned} |V_{\lambda}^{\pi_{n}}(\bar{s}) - V_{\lambda_{0}}^{*}(\bar{s})| &\leq |\sum_{a \in \mathcal{A}} Q_{\lambda}^{\pi_{n}}(\bar{s}, a) \pi_{n,\lambda}(a|\bar{s}) - \sum_{a \in \mathcal{A}} Q_{\lambda_{0}}^{*}(\bar{s}, a) \pi_{n,\lambda}(a|\bar{s})| \\ &+ |\sum_{a \in \mathcal{A}} Q_{\lambda_{0}}^{*}(\bar{s}, a) \pi_{n,\lambda}(a|\bar{s}) - \sum_{a \in \mathcal{A}} Q_{\lambda_{0}}^{*}(\bar{s}, a) \pi_{\lambda_{0}}^{*}(\pi_{n,\lambda})(a|\bar{s})| \\ &\leq \max_{\bar{s}, a \in \bar{\mathcal{S}}_{\lambda_{0}} \times \mathcal{A}} |Q_{\lambda}^{\pi_{n}}(\bar{s}, a) - Q_{\lambda_{0}}^{*}(\bar{s}, a)| + \max_{\bar{s} \in \bar{\mathcal{S}}_{\lambda_{0}}} 2(1 - \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s})). \end{aligned}$$

Since the right side of the inequality is not dependent on \bar{s} we can also maximize over \bar{s} on the left side and the apply $\limsup_{n}(\cdot)$, which completes the proof of this point.

4. For all $\lambda \in U_{\delta}(\lambda_0), \pi_0 > 0, n \ge 0, \bar{s} \in \bar{S}_{\lambda_0}$ it holds that

$$\begin{split} |J_{\lambda}^{\pi_n} - J_{\lambda_0}^*| &\leq \sum_{\bar{s} \in \text{supp}\,\bar{\mu}} \bar{\mu}(\bar{s}) |V_{\lambda}^{\pi_n}(\bar{s}) - V_{\lambda_0}^*(\bar{s})| \\ &= \sum_{\bar{s} \in \text{supp}\,\bar{\mu} \setminus \bar{\mathcal{S}}_{\lambda_0}} \bar{\mu}(\bar{s}) V_{\lambda}^{\pi_n}(\bar{s}) + \sum_{\bar{s} \in \bar{\mathcal{S}}_{\lambda_0}} \bar{\mu}(\bar{s}) |V_{\lambda}^{\pi_n}(\bar{s}) - V_{\lambda_0}^*(\bar{s})|. \end{split}$$

Because of $\operatorname{supp} \bar{\mu} \subset \operatorname{supp} \nu_{\lambda_0,\pi_0}$ it holds that $\operatorname{supp} \bar{\mu} \setminus \bar{S}_{\lambda_0} = \operatorname{supp} \bar{\mu} \setminus \operatorname{supp} \operatorname{den}_{\lambda_0,\pi_0}$. Since $V_{\lambda_0}^*(\bar{s}) = 0$ on $\operatorname{supp} \bar{\mu} \setminus \operatorname{supp} \operatorname{den}_{\lambda_0,\pi_0}$ (similar to corollary 18), we can use lemma 20 to bound $V_{\lambda_0}^{\pi_n}(\bar{s})$ by $\frac{N\delta}{2}$. The result follows by applying $\limsup \operatorname{supp}(\cdot)$ and then using point 3. above.

5. We choose the sequence $(\pi_{\lambda_0,n}^*)_{n\geq 0}$ so that $\pi_{\lambda_0,n}^*(a \mid \bar{s}) \geq \pi_{n,\lambda}(a \mid \bar{s})$ for all $\bar{s} \in \bar{S}_{\lambda_0}$ and all $a \in \mathcal{O}(\bar{s})$, which implies $\|\pi_{n,\lambda}(\cdot|\bar{s}) - \pi_{\lambda_0,n}^*(\cdot|\bar{s})\|_1 = 2(1 - \pi_{n_0,\lambda}(\mathcal{O}(\bar{s})|\bar{s}))$. The first statement follows the same logic as the second statement of the claim and point 4. of lemma 24. The only difference is that we are now bounding for all the remaining horizons. Notice that the β' we choose works for all the horizons because $\beta' > \beta_N \ge \beta_h$, $h \le N$. The correctness of the choice of x_0 follows from the fact that $f_{\gamma'}$ is increasing. The statement about $y_n \to y$ follows also from point 4. of lemma 24. Finally, for the q-linear convergence we have (let us denote $x_n := f_{\gamma'}^{\circ(n-n_0)}(x_0)$)

$$\lim_{n \to \infty} \frac{y_{n+1} - y_L}{y_n - y_L} = \lim_{n \to \infty} \frac{2(1 - f_{\gamma'}(x_n)) - 2(1 - x^*(\gamma'))}{2(1 - x_n) - 2(1 - x^*(\gamma'))} = \lim_{n \to \infty} \frac{f_{\gamma'}(x_n) - x^*(\gamma')}{x_n - x^*(\gamma')}$$
$$= \overset{Heine}{\lim_{x \to x^*(\gamma')} \frac{f_{\gamma'}(x) - x^*(\gamma')}{x - x^*(\gamma')}}{\lim_{x \to x^*(\gamma')} f_{\gamma'}(x)} = \lim_{x \to x^*(\gamma')} (\frac{x}{x + \gamma'})' = \frac{\gamma'}{(x^*(\gamma') + \gamma')^2}$$
$$= \gamma' < 1,$$

where we used Heine theorem, L'Hospital rule and definition of $x^*(\gamma')$ from lemma 24.

Given a deviation of δ from a deterministic kernel, corollary 27 can be used to bound the deviation in the accumulation points of, e.g., the goal-reaching objective (point 4.), or the policy (points 1. and 5.). An illustration of the accumulation points of $\min_{\bar{s}\in\bar{S}_{\lambda_0}}\pi_n(\mathcal{O}(\bar{s})|s)$ and the estimates in corollary 27 is provided in figure 6 in the case of a 2-armed bandit model (see section 7.1.2 and the description there) and in figure 7 in case of a simple grid world model. All details of the construction of the MDP underlying figure 7 can be found in the appendix B, example 5. Figure 7a shows how the bound in the corollary, point 5. (blue line) and the accumulation points of eUDRL-generated policies (orange marks) behave as functions of the distance δ to a deterministic kernel. Figure 7b shows how eUDRL approaches those accumulation points as a function of the iteration n. Varying initial policies are visible at n = 0 and varying levels of δ are highlighted in different colors. Figure 7c contains the same data as Figure 7b but it is organized to reveal the dependency on δ . Notice that initial conditions uniformly fill the plot's area. Figure 7d shows a map of the grid world. Similar to the example presented in figure 6, it is apparent that the bound in figure 7a (blue line) originates from continuity: it becomes tight as δ approaches 0 and becomes loose as δ increases. Regrettably, the bound deteriorates quickly as δ increases above ~ 10⁻⁵, which limits the applicability in a practical setting. The origin of this behavior might be seen in lemma 22, which uses a uniform bound α to estimate the visitation probabilities ignoring the specific dynamics of the environment. It remains a question for forthcoming research to develop methods that provide practical estimates that hold on intervals of δ , N, number of states, etc. of practical interest.

7.2 Asymptotic Continuity Assuming Uniqueness of the Optimal Policy

The main result of this section is to demonstrate the continuity of accumulation points of the sequence of eUDRL-generated policies under the assumption that the optimal policy is unique and deterministic. This assumption can be concisely expressed as follows: for a deterministic kernel λ_0 the set of optimal actions has exactly one element $|\mathcal{O}(\bar{s})| = 1$ on $\bar{\mathcal{S}}_{\lambda_0}$ (compare definitions 8 and 5 point 3). Recall also that restricting to \bar{S}_{λ_0} does not result in a loss of generality as the values (and potential discontinuity) of a policy outside \bar{S}_{λ_0} have no impact on the goal-reaching objective. The assumption $|\mathcal{O}(\bar{s})| = 1$ entails that the factorization with respect to the set of optimal actions becomes trivial and the continuity relative to optimal actions reduces to ordinary continuity. In this situation theorem 16 implies that eUDRL-generated policies are continuous at λ_0 . Working with continuous policies, as opposed to relatively continuous ones, leads to a substantial simplification compared to the discussion in the section 7.1. The uniqueness condition on the optimal policy is commonly encountered in publications on the analysis of convergence and stability of RL algorithms. But in the context of eUDRL it is not straightforward to see how this condition could be employed without loss of generality. Similar to the preceding sections, the core idea that enables our asymptotic continuity analysis is to derive bounds for the eUDRL policy recursion through monotonicity bounds for repeated application of rational functions (see remark 23). This leads to the study of a dynamical system (see lemma 30), whose fixed-point properties reveal information about the accumulation points of eUDRL's policy recursion as $\lambda \to \lambda_0$.



(c) Dependency on distance to determin. kernel.



Figure 7: Illustration of estimates of corollary 27 on the behavior of $\min_{\bar{s}\in\bar{S}_{\lambda_0}} \pi_n(\mathcal{O}(\bar{s})|s)$ for the simple grid world example shown in plot (d). Plot (a) shows the dependency of accumulation points (depicted in orange, approximated by values at n = 100) and the estimate in the corollary (blue). To illustrate the approach of eUDRL towards the accumulation points, plot (b) shows the dependency on the iteration n for varying distances to a deterministic kernel highlighted by different colors and varying initial policy. Plot (c) contains the same information as (b) but it is organized to reveal the dependency on δ , where varying numbers of iteration are highlighted by different colors. Plot (d) shows the map of the grid world. A wall is depicted in gray, the set \bar{S}_{λ_0} in yellow, and the goal in red. Arrows depict optimal actions associated with the specific state and goal. The structure and content of this section roughly align with section 7.1. The discussion is often shorter due to the assumption of a unique optimal policy but our analysis is based on the study of a different rational function, which requires an additional lemma about the dynamical system properties. We derive the preliminary lemma in section 7.2.1. The main theorem is contained in section 7.1.2. The result is generalized to other segment subspaces in section 7.2.3. As a corollary of the main result, we provide explicit estimates on the location of accumulation point sets in section 7.2.4. This yields upper bounds on the errors of policies, values and goal-reaching objective. We conclude the discussion with examples.

7.2.1 Preliminary Lemma

Our analysis of eUDRL's convergence in case of a unique optimal policy draws on the properties of a dynamical system induced by a certain rational function. The emergence of the dynamical system and its relation to the eUDRL policy recursion are described in detail in section 7.1.1. Here we only analyze the relevant dynamical system in case of a unique optimal policy.

Lemma 30 (*h*-lemma) Let $N \ge 1$ be a natural number, let $b_0 = \frac{1}{2N} (\frac{2N-1}{2N})^{2N-1}$ and let $b \in (0, b_0)$. Let $h_b : [0, 1] \to [0, 1]$ be defined as

$$h_b(x) = \frac{x^{2N}}{x^{2N} + b}.$$

Then the following assertions hold.

- 1. $h_b \upharpoonright_{(0,1]}$ is increasing, h_b is non-decreasing.
- 2. h_b has exactly three fixed points 0, $x_l(b)$ and $x_u(b)$, where $0 < x_l(b) < x_u(b) < 1$. It holds that

$$h_b(x) > x$$
, for $x_l(b) < x < x_u(b)$,
 $h_b(x) < x$, for $0 < x < x_l(b)$, or $x_u(b) < x \le 1$.

- 3. $h_b((x_l(b), 1]) \subset (x_l(b), 1]$ and $h_b^{\circ n}(x) \to x_u(b)$ for all $x \in (x_l(b), 1]$.
- 4. Let $(y_n)_{n\geq 0}$, $y_n \in [0,1]$ be a sequence with $y_0 > x_l(b)$ and $y_{n+1} \ge h_b(y_n)$ for all $n \ge 0$. Then $y_n \ge h_b^{\circ n}(y_0)$ and $\liminf_n y_n \ge x_u(b)$.
- 5. There exist unique, continuous, strictly monotonic extensions $\bar{x}_l : [0, b_0] \rightarrow [0, \frac{2N-1}{2N}]$, $\bar{x}_u : [0, b_0] \rightarrow [\frac{2N-1}{2N}, 1]$, where \bar{x}_l is increasing and \bar{x}_u is decreasing, so that $\bar{x}_l(b) = x_l(b)$ and $\bar{x}_u(b) = x_u(b)$ on $(0, b_0)$. It follows that $x_l(b) \rightarrow 0$, $x_u(b) \rightarrow 1$ as $b \rightarrow 0$ and $x_l(b) \rightarrow \frac{2N-1}{2N}$, $x_u(b) \rightarrow \frac{2N-1}{2N}$ as $b \rightarrow b_0$. Since \bar{x}_l is just an extension of x_l we will denote it by the same symbol x_l . Similarly, we denote \bar{x}_u by x_u .
- 6. For $0 < b < b_0$ the values $x_u(b)$ and $x_l(b)$ can be computed by iterative application of the strict contractions $g_l : [0, \frac{2N-1}{2N}] \to [0, \frac{2N-1}{2N}], g_l(x) = (\frac{b}{1-x})^{\frac{1}{2N-1}}$ and $g_u : [\frac{2N-1}{2N}, 1] \to [\frac{2N-1}{2N}, 1], g_u(x) = 1 - \frac{b}{x^{2N-1}}.$

Proof 1. The derivative of h is non-negative. It is positive if x > 0.

2. 0 is a solution of h(x) = x and therefore it is a fixed point. If x > 0 then $h(x) > x \iff u(x) := x^{2N-1}(1-x) > b$ and $h(x) = x \iff u(x) = b$. Define u(0) := 0, such that u(x) = 0 implies $x \in \{0, 1\}$. Computing the derivative one can verify that u has two stationary points x = 0 and $x = \frac{2N-1}{2N}$ and is increasing on $[0, \frac{2N-1}{2N}]$ and decreasing on $[\frac{2N-1}{2N}, 1]$. u has a local maximum at $x = \frac{2N-1}{2N}$ with $u(\frac{2N-1}{2N}) = \frac{1}{2N}(\frac{2N-1}{2N})^{2N-1}$. Making use of the assumptions $0 < b < \frac{1}{2N}(\frac{2N-1}{2N})^{2N-1}$ the intermediate value theorem implies that u(x) = b has exactly two solutions on [0, 1]. The first solution, x_l , lies in $(0, \frac{2N-1}{2N})$. Similarly, the second solution, x_u , lies in $(\frac{2N-1}{2N}, 1)$. The rest of the statement follows since u(x) > b on (x_l, x_u) and u(x) < b on $[0, x_l) \cup (x_u, 1]$.

3. Denote $h_1 = h \upharpoonright_{I_1}, h_2 = h \upharpoonright_{I_2}$, where $I_1 = (x_l, x_u], I_2 = [x_u, 1]$. Since h is increasing $x_l = h_1(x_l) < h_1(I_1) \le h_1(x_u) = x_u$ and thus $h_1(I_1) \subset I_1$, i.e., $h_1 : I_1 \to I_1$. Similarly $h_2 : I_2 \to I_2$. We show that if $x \in (x_l, 1]$ then $h^{\circ n}(x) \to x_u$. If $x \in I_1$ then $h^{\circ n}(x) = h_1^{\circ n}(x)$ is an increasing and bounded sequence that has a limit. Since h is continuous this limit is a fixed point. Since there are two fixed points in $\overline{I_1}$ and x_l cannot be a limit for any $x \in I_1$ (since $h_1^{\circ n}(x) \ge x > x_l$) it follows $h_1^{\circ n}(x) \to x_u$. Similarly in the remaining case $h_2^{\circ n}(x) \to x_u$.

4. The proof follows the same line as in point 4. of lemma 24.

5. Consider the increasing, continuous function $u_1 = u \upharpoonright_{[0,\frac{2N-1}{2N}]}$ and set $\bar{x}_l = u_1^{-1}$, which is also increasing. For all $b \in (0, b_0) u_1^{-1}(b) \in [0, \frac{2N-1}{2N}]$ is the unique solution of the equation $u_1(x) = b$ in this interval, so that $\bar{x}_l(b) = x_l(b)$. Since $u_1(0) = 0$ and $u_1(\frac{2N-1}{2N}) = b_0$ it follows $\bar{x}_l(0) = 0$ and $\bar{x}_l(b_0) = \frac{2N-1}{2N}$. Since $u_1 : [0, \frac{2N-1}{2N}] \to [0, b_0]$ is continuous, defined on interval and increasing it is an open map. It follows that u_1 is a homeomorphism and \bar{x}_l is continuous. The discussion of \bar{x}_u follows along the same line by consider the decreasing, continuous function $u_2 = u \upharpoonright_{[\frac{2N-1}{2N}, 1]}$ and setting $\bar{x}_u = u_2^{-1}$.

6. Now assume that $b \in (0, b_0)$. We aim to prove that g_l is a strict contraction on a complete metric space with the fixed point $x_l(b)$. In order to prove $g_l(I_l) \subset I_l$ assume $x \in I_l$. Since $x \leq \frac{2N-1}{2N} \iff 1-x \geq \frac{1}{2N}$ it follows $g_l(x) = (\frac{b}{1-x})^{\frac{1}{2N-1}} < (\frac{b_0}{\frac{1}{2N}})^{\frac{1}{2N-1}} = \frac{2N-1}{2N}$. Further, $g_l(x) = (\frac{b}{1-x})^{\frac{1}{2N-1}} \geq 0$. Combining those points yields $g_l(x) \in I_l$. Completeness follows since I_l is a compact metric space. Note that for all $x \in I_l$ it holds that $\frac{dg_l}{dx}(x) = \frac{b^{\frac{1}{2N-1}}}{2N-1}(\frac{1}{1-x})^{1+\frac{1}{2N-1}} < \frac{b_0^{\frac{1}{2N-1}}}{2N-1}(2N)^{1+\frac{1}{2N-1}} = 1$. Since $0 \leq \frac{dg_l}{dx} < 1$ on I_l and I_l is compact and $\frac{dg_l}{dx}$ is continuous there exists L < 1 so that $|\frac{dg_l}{dx}| < L$. By Lagrange's mean value theorem it follows that for all $x, x' \in I_l$ it holds that $|g_l(x) - g_l(x')| < L|x - x'|$. This proves that g_l is a strict contraction on the complete metric space I_l . From Banach's fixed point theorem it follows that g_l has exactly one fixed point and the iteration $g_l^{\circ n}(x_0)$ converges to this fixed point for $n \to \infty$ for all $x_0 \in I_l$. It remains to show that $x_l(b)$ is the fixed point of g_l . Since $x_l(b)$ is the solution of $u_1(x) = b$ we have $b = u_1(x_l(b)) = x_l^{2N-1}(b)(1-x_l(b))$. Since b > 0 this is equivalent to $x_l(b) = (\frac{b}{1-x_l(b)})^{\frac{1}{2N-1}} = g_l(x_l(b))$.

The claim about g_u can be proved in similar way. Assume that $b \in (0, b_0)$ and $x \in I_u = [\frac{2N-1}{2N}, 1]$. Since $g_u(x) = 1 - \frac{b}{x^{2N-1}} > 1 - \frac{b_0}{(\frac{2N-1}{2N})^{2N-1}} = 1 - \frac{1}{2N} = \frac{2N-1}{2N}$ and $g_u(x) = 1 - \frac{b}{x^{2N-1}} < 1 - \frac{0}{1^{2N-1}} = 1$ we get $g_u(x) \in I_u$. Further for all $x \in I_u$ it



Figure 8: Illustration of the dynamical system induced by iterative application of h_b . The figure shows the dependence of $h_b^{\circ n}(x)$ given initial condition x on iteration n. The parameter b is chosen as $b = \frac{b_0}{2}$ for N = 2.

holds that $0 \leq \frac{dg_u}{dx}(x) = \frac{b(2N-1)}{x^{2N}} < \frac{b_0(2N-1)}{(\frac{2N-1}{2N})^{2N}} = 1$ for all $x \in I_u$. We conclude (similarly as in case of g_l) that g_u is a strict contraction on a complete metric space. Since $b = u_2(u_2^{-1}(b)) = u_2(x_u(b)) = x_u^{2N}(b)(1-x_u(b))$ we get $x_u(b) = 1 - \frac{b}{x_u^{2N}(b)} = g_u(x_u(b))$, i.e., $x_u(b)$ is fixed point of g_u .

Figure 8 illustrates the dynamical system induced by iterative application of h_b .

7.2.2 The main theorem

The following result is a counterpart of theorem 25 point 2. Instead of assuming $S_{\lambda_0} \subset \sup p\bar{\mu}$ (as in theorem 25) the following theorem applies in a situation where $|\mathcal{O}(\bar{s})| = 1$ on \bar{S}_{λ_0} . Together with lemma 10 the theorem demonstrates the continuity of the sets of accumulation points of UDRL generated policies. Since the reasoning is the same as in point 1. of theorem 25 we skip the analogue of theorem 25 point 1. and focus solely on point 2.

Theorem 31 (Continuity of eUDRL limit policies at deterministic points – the case $|\mathcal{O}(\bar{s})| = 1$ on \bar{S}_{λ_0}) Let $\{\mathcal{M}_{\lambda}|\lambda \in (\Delta S)^{S \times A}\}$ and $\{\bar{\mathcal{M}}_{\lambda}|\lambda \in (\Delta S)^{S \times A}\}$ be compatible families. Let $\lambda_0 \in (\Delta S)^{S \times A}$ be a deterministic kernel and let $(\pi_{n,\lambda})_{n\geq 0}$ be a sequence of eUDRL-generated policies with initial condition $\pi_0 > 0$ and transition kernel λ . Assume that $|\mathcal{O}(\bar{s})| = 1$ on \bar{S}_{λ_0} . Then for all $\pi_0 > 0$ there exists $\delta \in (0, 2)$ so that for all $\lambda \in U_{\delta}(\lambda_0)$ with

$$U_{\delta}(\lambda_0) = \{\lambda | \max_{(s,a) \in \mathcal{S} \times \mathcal{A}} \|\lambda(\cdot|s,a) - \lambda_0(\cdot|s,a)\|_1 < \delta \}$$

and all $\bar{s} \in \bar{S}_{\lambda_0}$ it holds that

$$\liminf_{n} \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \ge x_u(b) \quad and \quad x_u(b) \to 1 \quad as \quad \delta \to 0,$$

where

$$b = \frac{\delta N^2 (N+1)}{4(1-\frac{\delta}{2})^{2N} \min_{\bar{s}' \in \operatorname{supp} \bar{\mu}} \bar{\mu}(\bar{s}')}$$

and x_u is defined in lemma 30. It follows that

$$\liminf_{n \to \infty} \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \to 1 \quad as \quad \lambda \to \lambda_0.$$

In fact we will prove the slightly stronger assertion that there exists $\delta \in (0, 2)$ such that for all $\lambda \in U_{\delta}(\lambda_0)$ and $\bar{s} \in \bar{S}_{\lambda_0}$ it holds that

$$\min_{\bar{s}\in\bar{\mathcal{S}}_{\lambda_0}}\pi_{n+1,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \ge h_b(\min_{\bar{s}\in\bar{\mathcal{S}}_{\lambda_0}}\pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}))$$
(7.10)

where the function h_b and quantities $x_u(b)$, $x_l(b)$ are defined in lemma 30.

Proof We utilize the notation introduced in the proof of theorem 25. Assume $\pi_0 > 0$ is arbitrary and fixed. Fix arbitrary $\delta \in (0,2)$ and assume $\lambda \in U_{\delta}(\lambda_0)$. We show some estimates that will be used later in the proof.

Bounding state visitation: Assume $\bar{s} = (s, h, g) \in \bar{S}_{\lambda_0}$. By lemma 6 point 3. and lemma 10 (optimality at deterministic points) we have that for $n \ge 1$ it holds that

$$\begin{split} \mathbb{P}_{\lambda_0}(S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g, H_0^{\Sigma} = h, G_0^{\Sigma} = g; \pi_{\lambda_0}^*) \\ = \mathbb{P}_{\lambda_0}(S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g, H_0^{\Sigma} = h, G_0^{\Sigma} = g; \pi_n) > 0. \end{split}$$

It follows that there exists t such that $\mathbb{P}_{\lambda_0}(\bar{S}_t = \bar{s}; \pi^*_{\lambda_0}) > 0$. Thus there exists a prefix $\bar{s}_0, a_0, \ldots, \bar{s}_t = \bar{s}$ with $\mathbb{P}_{\lambda_0}(\bar{S}_t = \bar{s}, \ldots, a_0, \bar{s}_0; \pi^*_{\lambda_0}) > 0$ and it follows that $\bar{s}_i \in \bar{S}_{\lambda_0}$ for $i = 0, \ldots, t-1$. Assuming λ_0 and $\pi^*_{\lambda_0}$ are deterministic $\lambda_0(s_{i+1}|s_i, a_i) = 1$ and $\pi^*_{\lambda_0}(a_i|\bar{s}_i) = 1$ for the optimal action a_i . Noting that $\lambda(s_{i+1}|s_i, a_i) \geq 1 - \frac{\delta}{2}$ and $\pi(a_i|\bar{s}_i) = \pi(\mathcal{O}(\bar{s}_i)|\bar{s}_i) \geq \min_{\bar{s}' \in \bar{S}_{\lambda_0}} \pi(\mathcal{O}(\bar{s}')|\bar{s}')$ leads to the estimate

$$\mathbb{P}_{\lambda}(S_{t} = \bar{s}; \pi) \geq \mathbb{P}_{\lambda}(S_{t} = \bar{s}, \dots, a_{0}, \bar{s}_{0}; \pi)$$

$$= \bar{\mu}(\bar{s}_{0})\pi(a_{0}|\bar{s}_{0})\lambda(s_{1}|s_{0}, a_{0})\dots\lambda(s|s_{t-1}, a_{t-1})$$

$$\geq (1 - \frac{\delta}{2})^{N} \min_{\bar{s}' \in \operatorname{supp} \bar{\mu}} \bar{\mu}(\bar{s}') \left(\min_{\bar{s}' \in \bar{\mathcal{S}}_{\lambda_{0}}} \pi(\mathcal{O}(\bar{s}')|\bar{s}')\right)^{N}.$$

For the state visitation probability $v_{\lambda,\pi}(h,g|s,h)$ this yields that

$$\begin{aligned} v_{\lambda,\pi}(h,g|s,h) &\geq \mathbb{P}_{\lambda}(H_{0}^{\Sigma}=h,G_{0}^{\Sigma}=g,S_{0}^{\Sigma}=s,l(\Sigma)=h;\pi) \\ &= \sum_{\substack{\sigma:l(\sigma)=h,\\\bar{s}_{0}^{\sigma}=\bar{s}}} c^{-1} \sum_{\substack{t'\leq N-l(\sigma)\\\bar{s}_{0}^{\sigma}=\bar{s}}} \mathbb{P}_{\lambda}(\bar{S}_{t}=\bar{s}_{0}^{\sigma},\ldots,\bar{S}_{t+l(\sigma)}=\bar{s}_{l(\sigma)}^{\sigma};\pi) \\ &\geq \sum_{\substack{\sigma:l(\sigma)=h,\\\bar{s}_{0}^{\sigma}=\bar{s}}} c^{-1} \mathbb{P}_{\lambda}(\bar{S}_{t}=\bar{s}_{0}^{\sigma},\ldots,\bar{S}_{t+l(\sigma)}=\bar{s}_{l(\sigma)}^{\sigma};\pi) \\ &= c^{-1} \mathbb{P}_{\lambda}(\bar{S}_{t}=\bar{s};\pi) \sum_{\substack{\sigma:l(\sigma)=h,\\\bar{s}_{0}^{\sigma}=\bar{s}}} \mathbb{P}_{\lambda}(\bar{A}_{t}=a_{0}^{\sigma},\ldots,\bar{S}_{t+l(\sigma)}=\bar{s}_{l(\sigma)}^{\sigma}|\bar{S}_{t}=\bar{s}_{0}^{\sigma};\pi) \\ &\geq c^{-1} \mathbb{P}_{\lambda}(\bar{S}_{t}=\bar{s};\pi) \\ &\geq \frac{2}{N(N+1)} \min_{\bar{s}'\in \mathrm{supp}\,\bar{\mu}}\bar{\mu}(\bar{s}')(\min_{\bar{s}'\in\bar{S}_{\lambda_{0}}}\pi(\mathcal{O}(\bar{s}')|\bar{s}'))^{N}(1-\frac{\delta}{2})^{N}, \end{aligned}$$
(7.11)

where we used the fact that $\sum_{\sigma:l(\sigma)=h,\bar{s}_0^{\sigma}=\bar{s}} \mathbb{P}_{\lambda}(\bar{A}_t = a_0^{\sigma}, \dots, \bar{S}_{t+l(\sigma)} = \bar{s}_{l(\sigma)}^{\sigma} | \bar{S}_t = \bar{s}_0^{\sigma}; \pi) = 1.$ Bounding $Q_{\lambda}^{\pi_n,g,h}$: If $\bar{s} \in \bar{S}_{\lambda_0}$ and $a \in \mathcal{A} \setminus \mathcal{O}(\bar{s})$ then by lemma 20

$$Q_{\lambda}^{\pi_n,g,h}((s,h',g'),a) \le Q_{\lambda}^*(\bar{s},a) \le \frac{\delta N}{2}.$$
(7.12)

Bounding Q_{λ}^{π} for optimal actions: Let $\bar{s} \in \bar{S}_{\lambda_0}$ and $a \in \mathcal{O}(\bar{s})$. If h = 1 then $\bar{s} = (s, 1, g)$. As λ_0 is deterministic there exists s'' so that $\lambda_0(s''|s, a) = 1$. Notice that since $\bar{s} \in \bar{S}_{\lambda_0}$ and a is optimal it holds that $\rho(s'') = g$. Consequently $Q_{\lambda}^{\pi}((s, 1, g), a) \ge \lambda(s''|s, a) \ge 1 - \frac{\delta}{2}$. For $h = 2 \ \bar{s} = (s, 2, g)$ there exist s'' so that $(s'', 1, g) \in \bar{S}_{\lambda_0}$ and $\lambda_0(s''|s, a) = 1$ by lemma 6 point 4. It follows for $a \in \mathcal{O}(\bar{s})$ that

$$\begin{aligned} Q_{\lambda}^{\pi}((s,2,g),a) &= \sum_{s'\in\bar{S}_{T}} \lambda(s'|s,a) \sum_{a'\in\mathcal{A}} \pi(a'|(s',1,g)) Q_{\lambda}^{\pi}(s',1,g,a') \\ &\geq \lambda(s''|s,a) \sum_{a'\in\mathcal{O}(\bar{s})} \pi(a'|(s'',1,g)) Q_{\lambda}^{\pi}(s'',1,g,a') \\ &\geq (1-\frac{\delta}{2})^{2} \pi(\mathcal{O}((s'',1,g))|(s'',1,g)) \geq (1-\frac{\delta}{2})^{2} \min_{\bar{s}'\in\bar{S}_{\lambda_{0}}} \pi(\mathcal{O}(\bar{s}')|\bar{s}'). \end{aligned}$$

Reasoning inductively we obtain for horizon h

$$Q_{\lambda}^{\pi}((s,h,g),a) \ge (1-\frac{\delta}{2})^{N} \left(\min_{\bar{s}' \in \bar{\mathcal{S}}_{\lambda_0}} \pi(\mathcal{O}(\bar{s}')|\bar{s}') \right)^{N-1}.$$
(7.13)

Recursive bounding of policies on optimal states: We begin by estimating $\pi(\mathcal{O}(\bar{s})|\bar{s})$, $\bar{s} = (s, h, g) \in \bar{S}_{\lambda_0}$ similarly to the proof of theorem 25. We use lemma 6 point 2. and lemma 3 point 1. to get $(\forall n \ge 0, \forall \lambda \in U_{\delta}(\lambda_0))$ that

$$\begin{aligned} \pi_{n+1,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) &= \frac{\sum\limits_{a \in \mathcal{O}(\bar{s}), h' \ge h, g \in \mathcal{G}} Q_{\lambda}^{\pi_{n},g,h}((s,h',g'),a)\pi_{n,\lambda}(a|s,h',g')v_{\lambda,\pi_{n}}(h',g'|s,h)}{\sum\limits_{a \in \mathcal{A}, h' \ge h, g \in \mathcal{G}} Q_{\lambda}^{\pi_{n},g,h}((s,h',g'),a)\pi_{n,\lambda}(a|s,h',g')v_{\lambda,\pi_{n}}(h',g'|s,h)} \\ &\ge \frac{\sum\limits_{a \in \mathcal{O}(\bar{s}), h' \ge h, g \in \mathcal{G}} Q_{\lambda}^{\pi_{n},g,h}((s,h',g'),a)\pi_{n,\lambda}(a|s,h',g')v_{\lambda,\pi_{n}}(h',g'|s,h)}{\sum\limits_{a \in \mathcal{O}(\bar{s}), h' \ge h, g \in \mathcal{G}} Q_{\lambda}^{\pi_{n},g,h}((s,h',g'),a)\pi_{n,\lambda}(a|s,h',g')v_{\lambda,\pi_{n}}(h',g'|s,h) + \frac{\delta N}{2}}, \end{aligned}$$

where we made use of the estimate (7.12) and lemma 12 point 2. For $(h', g') \neq (h, g)$ we bound $Q_{\lambda}^{\pi_n,g,h}((s,h',g'),a)$ from below by zero (as they are probabilities) and use the monotonicity remark 23 to obtain

$$\pi_{n+1,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \geq \frac{\sum_{a \in \mathcal{O}(\bar{s})} Q_{\lambda}^{\pi_n}((s,h,g),a) \pi_{n,\lambda}(a|s,h,g) v_{\lambda,\pi_n}(h,g|s,h)}{\sum_{a \in \mathcal{O}(\bar{s})} Q_{\lambda}^{\pi_n}((s,h,g),a) \pi_{n,\lambda}(a|s,h,g) v_{\lambda,\pi_n}(h,g|s,h) + \frac{\delta N}{2}}$$

Employing remark 23 a second time and the estimates (7.13) and (7.11) gives then

$$\pi_{n+1,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \geq \frac{\frac{2}{N(N+1)}(\min_{\bar{s}\in \text{supp }\bar{\mu}}\bar{\mu}(\bar{s}))(1-\frac{\delta}{2})^{2N}(\min_{\bar{s}'\in\bar{\mathcal{S}}_{\lambda_0}}\pi_{n,\lambda}(\mathcal{O}(\bar{s}')|\bar{s}'))^{2N-1}\pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s})}{\frac{2}{N(N+1)}(\min_{\bar{s}\in \text{supp }\bar{\mu}}\bar{\mu}(\bar{s}))(1-\frac{\delta}{2})^{2N}(\min_{\bar{s}'\in\bar{\mathcal{S}}_{\lambda_0}}\pi_{n,\lambda}(\mathcal{O}(\bar{s}')|\bar{s}'))^{2N-1}\pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s})+\frac{\delta N}{2}}$$

Finally, bounding $\pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s})$ from below by its minimum over $\bar{\mathcal{S}}_{\lambda_0}$ we obtain $(\forall n \ge 0, \forall \lambda \in U_{\delta}(\lambda_0), \forall \bar{s} = (s, h, g) \in \bar{S}(\lambda_0))$:

$$\begin{aligned} \pi_{n+1,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) &\geq \frac{\frac{2}{N(N+1)} (\min_{\bar{s}\in \mathrm{supp}\,\bar{\mu}}\,\bar{\mu}(\bar{s}))(1-\frac{\delta}{2})^{2N} (\min_{\bar{s}'\in\bar{\mathcal{S}}_{\lambda_0}}\,\pi_{n,\lambda}(\mathcal{O}(\bar{s}')|\bar{s}'))^{2N}}{\frac{2}{N(N+1)} (\min_{\bar{s}\in \mathrm{supp}\,\bar{\mu}}\,\bar{\mu}(\bar{s}))(1-\frac{\delta}{2})^{2N} (\min_{\bar{s}'\in\bar{\mathcal{S}}_{\lambda_0}}\,\pi_{n,\lambda}(\mathcal{O}(\bar{s}')|\bar{s}'))^{2N}} \\ &= \frac{(\min_{\bar{s}'\in\bar{\mathcal{S}}_{\lambda_0}}\,\pi_{n,\lambda}(\mathcal{O}(\bar{s}')|\bar{s}'))^{2N}}{(\min_{\bar{s}'\in\bar{\mathcal{S}}_{\lambda_0}}\,\pi_{n,\lambda}(\mathcal{O}(\bar{s}')|\bar{s}'))^{2N}+b} = h_b(\min_{\bar{s}'\in\bar{\mathcal{S}}_{\lambda_0}}\,\pi_{n,\lambda}(\mathcal{O}(\bar{s}')|\bar{s}')), \end{aligned}$$

where

$$b = \frac{\frac{\delta N}{2}}{\frac{2}{N(N+1)} (\min_{\bar{s} \in \operatorname{supp} \bar{\mu}} \bar{\mu}(\bar{s})) (1 - \frac{\delta}{2})^{2N}}.$$

Since the left hand side of the inequality does no longer depend on \bar{s} one can replace it by the minimum over \bar{s} , which proves the inequality (7.10). Since $b \to 0$ as $\delta \to 0$ we can restrict $\delta \in (0,2)$ to ensure that $b < \frac{1}{2N} (\frac{2N-1}{2N})^{2N-1}$.

Without loss of generality we can assume that $\delta \in (0, 2)$ is also chosen so that $\pi_0(\mathcal{O}(\bar{s})|\bar{s}) > x_l(b)$ for all $\bar{s} \in \bar{S}_{\lambda_0}$. This is possible since $x_l(b) \to 0$ for $\delta \to 0$ (since $b \to 0$ as $\delta \to 0$) and $\pi_0(\mathcal{O}(\bar{s})|\bar{s}) > 0$ for all $\bar{s} \in \bar{S}_{\lambda_0}$. Having established $\min_{\bar{s} \in \bar{S}_{\lambda_0}} \pi_0(\mathcal{O}(\bar{s})|\bar{s}) > x_l(b)$ we can employ lemma 30 point 4. to inequality (7.10) to find for all $\lambda \in U_{\delta}(\lambda_0)$ and $\bar{s} \in \bar{S}_{\lambda_0}$ that

$$\liminf_{n} \min_{\bar{s} \in \bar{\mathcal{S}}_{\lambda_0}} \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \ge x_u(b)$$

where $x_u(b) \to 1, b \to 0$. Since $b \to 0, \delta \to 0$ we also have that $x_u(b) \to 1$.

7.2.3 Extending the continuity results to other segment sub-spaces

As before we need to modify theorem 31 to cover algorithms like ODT, RWR (and some specific variants of eUDRL) restricting the recursion to Seg^{trail} or Seg^{diag} subspaces.

Theorem 32 (Relative continuity of accumulation points of eUDRL-generated policies at deterministic kernels – the case $|\mathcal{O}(\bar{s})| = 1$ on \bar{S}_{λ_0}) Theorem 31 remains valid under the renaming $\pi_n \to \pi_n^{\text{diag/trail}}$.

The proof of theorem 32 follows along similar lines as that of theorem 31. The differences are rather small and resemble those described in section 7.1.3.

7.2.4 Estimating the location of accumulation points

We conclude with a corollary that provides explicit bounds on the accumulation points of eUDRL-generated policies, values and goal-reaching objectives. See section 7.1.4 for a motivation.

Corollary 33 (Estimating the location of accumulation points – the case $|\mathcal{O}(\bar{s})| = 1$ on \bar{S}_{λ_0}) Under the conditions of theorem 31 let $\delta_0 \in (0,2)$ be the unique solution of the equation (cf. theorem 31 and lemma 30)

$$b(\delta) = b_0, \text{ where}$$

$$b(\delta) = \frac{\delta N^2 (N+1)}{4(1-\frac{\delta}{2})^{2N} \min_{\bar{s} \in \text{supp } \bar{\mu}} \bar{\mu}(\bar{s})}$$

$$b_0 = \frac{1}{2N} \left(\frac{2N-1}{2N}\right)^{2N-1}.$$

Then there exist two continuous, strictly monotonic functions $x_l : [0, \delta_0] \to [0, \frac{2N-1}{2N}]$, $x_u : [0, \delta_0] \to [\frac{2N-1}{2N}, 1]$, where x_l is increasing and x_u decreasing with $x_l(0) = 0$, $x_u(0) = 1$ with the following property: For any fixed $\delta \in (0, \delta_0)$ suppose that the policy π_0 satisfies $\pi_0 > 0$ and $\pi_0(\mathcal{O}(\bar{s})|\bar{s}) > x_l(\delta)$ for all $\bar{s} \in \bar{S}_{\lambda_0}$ then for all $\lambda \in U_{\delta}(\lambda_0)$ and all $\bar{s} \in \bar{S}_{\lambda_0}$ the following conclusions hold:

- 1. $\liminf_n \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s}) \geq x_u(\delta) \to 1, \delta \to 0$ or equivalently (in form of error to an optimal policy) $\limsup_n (1 \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s})) \leq 1 x_u(\delta)$
- 2. $\limsup_n |V_{\lambda}^{\pi_n}(\bar{s}) V_{\lambda_0}^*(\bar{s})| \le 1 (1 \frac{\delta}{2})^N x_u^N(\delta)$
- 3. $\limsup_n |J_{\lambda}^{\pi_n} J_{\lambda_0}^*| \leq \frac{N\delta}{2} + (1 (1 \frac{\delta}{2})^N x_u^N(\delta))$
- 4. (q-linear convergence) There exists a sequence of λ_0 -optimal policies $(\pi^*_{\lambda_0,n})_{n\geq 0}$ such that for all n it holds

$$\|\pi_{n,\lambda}(\cdot|\bar{s}) - \pi^*_{\lambda_0,n}(\cdot|\bar{s})\|_1 \le 2(1 - h_b^{\circ n}(x_0)),$$

where $x_0 = \min_{\bar{s} \in \bar{S}_{\lambda_0}} \pi_0(\mathcal{O}(\bar{s})|\bar{s}).$

The terms on the right hand side of 1.-3. converge to 0 as $\delta \to 0$. In 4. the sequence $(y_n)_{n\geq 0}$ defined by $y_n = 2(1-h_b^{\circ n}(x_0))$ converges q-linearly to the limit $y = 2(1-x_u(\delta))$ at a convergence rate of $\frac{2Nbx_u^{2N-1}}{(x_u^{2N}+b)^2}$ which tends to 0 as $\delta \to 0$.

Recall that the functions x_l , x_u were defined in lemma 30 as functions of b, which, in turn, is a function of δ . We slightly abuse notation in corollary 33 writing $x_l(\delta)$ for $x_l(b(\delta))$ and $x_u(\delta)$ for $x_l(b(\delta))$.

Proof First we have to show the uniqueness of δ_0 , which is defined as a solution to the equation

$$b(\delta) = b_0, \quad \delta \in [0, 2).$$

For convenience we restrict the domain of $b(\delta)$ to be [0,2), so that the restricted function is continuous. Since the first derivative of b is positive

$$\frac{db}{d\delta} = \frac{N^2(N+1)}{4\min_{\bar{s}\in \mathrm{supp}\,\bar{\mu}}\bar{\mu}(\bar{s})} \frac{(1-\frac{\delta}{2})+N\delta}{(1-\frac{\delta}{2})^{2N+1}} > 0 \iff 1+\delta(N-\frac{1}{2}) > 0,$$

there can be at most one solution to the equation. Since b(0) = 0, $b(\frac{2}{N+1}) > b_0$ and making use of the mentioned continuity, there exists at least one solution to the equation. This proves the existence of a unique solution δ_0 .

Here, the symbols x_l, x_u refer to compositions $x_l := \bar{x}_l \circ b$, $x_u := \bar{x}_u \circ b$, where \bar{x}_l, \bar{x}_u were defined in point 5. of lemma 30. Since b is strictly increasing, continuous, and defined on the interval all claimed properties of the compound functions follow from 30 point 5.

1. This point is just the theorem 31. Similarly to the proof of the theorem 31 the condition $\pi_0(\mathcal{O}(\bar{s})|\bar{s}) > x_l(\delta)$ comes from lemma 30 point 4.

2. Assume $\bar{s} \in \bar{\mathcal{S}}_{\lambda_0}$. From (7.13) it follows

$$V_{\lambda}^{\pi_n}(\bar{s}) \ge \sum_{a \in \mathcal{O}(\bar{s})} Q_{\lambda}^{\pi_n}(\bar{s}, a) \pi_{n,\lambda}(a|\bar{s}) \ge (1 - \frac{\delta}{2})^N (\min_{\bar{s}' \in \bar{\mathcal{S}}_{\lambda_0}} \pi(\mathcal{O}(\bar{s}')|\bar{s}'))^N$$

Now we apply point 1.

$$\liminf_{n} V_{\lambda}^{\pi_n}(\bar{s}) \ge (1 - \frac{\delta}{2})^N (\min_{\bar{s}' \in \bar{\mathcal{S}}_{\lambda_0}} \liminf_{n} \pi(\mathcal{O}(\bar{s}')|\bar{s}'))^N \ge (1 - \frac{\delta}{2})^N x_u^N$$

Finally, we conclude that

$$\limsup_{n} |V_{\lambda_0}^*(\bar{s}) - V_{\lambda}^{\pi_n}(\bar{s})| = 1 - \liminf_{n} V_{\lambda}^{\pi_n}(\bar{s}) \le 1 - (1 - \frac{\delta}{2})^N x_u^N.$$

3. The proof follows similarly as for corollary 27 point 4. except we utilize the previous point

$$\begin{split} |J_{\lambda_0}^* - J_{\lambda}^{\pi_n}| &\leq \sum_{\bar{s} \in \operatorname{supp} \bar{\mu} \setminus \bar{S}_{\lambda_0}} \bar{\mu}(\bar{s}) |V_{\lambda}^{\pi_n}(\bar{s})| + \sum_{\bar{s} \in \operatorname{supp} \bar{\mu} \cap \bar{S}_{\lambda_0}} \bar{\mu}(\bar{s}) |V_{\lambda_0}^* - V_{\lambda}^{\pi_n}(\bar{s})| \\ &\leq \frac{N\delta}{2} + \sum_{\bar{s} \in \operatorname{supp} \bar{\mu} \cap \bar{S}_{\lambda_0}} \bar{\mu}(\bar{s}) |V_{\lambda_0}^* - V_{\lambda}^{\pi_n}|, \\ \lim\sup_n |J_{\lambda_0}^* - J_{\lambda}^{\pi_n}| &\leq \frac{N\delta}{2} + 1 - (1 - \frac{\delta}{2})^N x_u^N(\delta). \end{split}$$

4. The sequence $(\pi^*_{\lambda_0,n})_{n\geq 0}$ is chosen in the same way as in the corollary 27. The first inequality is just the theorem 31 and lemma 30. The the statement $y_n \to y$, $n \to \infty$ is lemma 30 point 3.

Consider a sequence $x_n = h_b^{\circ n}(x_0), n \ge 0$. The last statement about the convergence rate follows by using Heine theorem and L'Hospital rule

$$\lim_{n \to \infty} \frac{y_{n+1} - y}{y_n - y} = \lim_{n \to \infty} \frac{x_u - h_b(x_n)}{x_u - x_n}$$
$$=^{Heine} \lim_{x \to x_u} \frac{x_u - h_b(x)}{x_u - x}$$
$$=^{L'Hospital} \lim_{x \to x_u} h'_b(x) = \lim_{x \to x_u} \frac{2Nbx^{2N-1}}{(x^{2N} + b)^2} = \frac{2Nbx_u^{2N-1}}{(x^{2N}_u + b)^2}.$$

Since $x_u \to 1$, $\delta \to 0$ and $b \to 0$, $\delta \to 0$ the the rate $\frac{2Nbx_u^{2N-1}}{(x_u^{2N}+b)^2} \to 0$ for $\delta \to 0$.

Similar to corollary 27, we can employ corollary 33 to bound the deviation in the accumulation points of eUDRL-generated quantities given a deviation of δ from a deterministic kernel. In the case of corollary 27 we already presented the estimates for the accumulation points of $\min_{\bar{s}\in \bar{S}_{\lambda_0}} \pi_n(\mathcal{O}(\bar{s})|s)$ for the simple bandit example 4 in figure 6 (see also section 7.1.2 and the explanation there). In fact both conditions $\bar{S}_{\lambda_0} \subset \text{supp }\bar{\mu}$ and $(\forall \bar{s} \in \bar{S}_{\lambda_0} : |\mathcal{O}(\bar{s})| = 1)$ are simultaneously met by the bandit example, which allows us to compare the estimates from the corollaries 27 and 33. A comparison is provided in figure 9, which shows how the estimates from the corollaries (blue x^* , orange x_u , green x_l lines) and the accumulation points of eUDRL-generated policies (orange marks) behave as functions of the distance δ to a deterministic kernel. Recall that according to corollary 33 $x_l(\delta)$ is an unstable fixed point of the underlying dynamical system. If the initial condition on the policy is such that it is larger than this fixed point (cf. corollary) then the accumulation points of the eUDRL recursion are bounded from below by the fixed point $x_u(\delta)$. In other words the $x_l(\delta)$ graph separates the plane in two regions, where the lower bound by $x_u(\delta)$ is guaranteed in the upper region.

As a second example we illustrate how the estimates of corollary 33 apply to an ODT architecture trained to identify the optimal policy in a simple grid world (that has a unique optimal policy on \bar{S}_{λ_0}), see figure 10. As outlined in section 2 ODT can be interpreted as a special instance of algorithms that fit within CE framework, where eUDRL's recursion operates on the space of trailing segments Seg^{trail} rather than Seg. All details of the construction of the MDP underlying figure 10 and the details of the ODT algorithm are provided in appendix B, example 6. Figure 10a shows how the bound in the corollary (orange and green lines) and the accumulation points of ODT-generated policies (orange marks) behave as functions of the distance δ to a deterministic kernel. Figure 10b shows how ODT approaches those accumulation points as a function of the iteration n. Varying initial policies are visible at n = 0 and different levels of δ are highlighted in different colors. Figure 10d shows a map of the grid world. Given the structure of the grid world it is visible that the uniqueness condition on the optimal policy is restrictive in the sense that domains that satisfying the condition ($\forall \bar{s} \in \bar{S}_{\lambda_0} : |\mathcal{O}(\bar{s})| = 1$) seem to occur rarely.



Figure 9: Comparison of estimates of corollaries 27, 33 on the behavior of $\min_{\bar{s}\in\bar{S}_{\lambda_0}}\pi_n(\mathcal{O}(\bar{s})|s)$ in case of a simple bandit example. The plot shows the δ -dependency of accumulation points (depicted in orange, approximated by values at n = 100) and the estimates from corollary 27 (blue) and corollary 33 (orange, green).



(c) Dependency on distance to determin. kernel.



Figure 10: Illustration of the estimates of corollary 33 on the behavior of $\min_{\bar{s}\in\bar{S}_{\lambda_0}}\pi_n(\mathcal{O}(\bar{s})|s)$ for ODT training in a simple grid world shown in plot (d). Plot (a) shows the dependency of accumulation points (depicted in orange, approximated by values at n = 100) and the estimate in the corollary given (orange and green). To illustrate how ODT approaches towards the accumulation points, plot (b) shows the dependency on the iteration n for varying distances to a deterministic kernel highlighted by different colors and varying initial policy. Plot (c) contains the same information as (b) but it is organized to reveal the dependency on δ , where varying numbers of iteration are highlighted by different colors. Plot (d) shows the map of the grid world. A wall is depicted in gray, the set \bar{S}_{λ_0} in yellow, and the goal in red. Arrows depict optimal actions associated with the specific state and goal.

8 Regularized Recursion

This section examines regularization in eUDRL, which is motivated by the following two facts. First, the standard ODT architecture utilizes entropy regularization, whereas we have modeled ODT without regularization so far. Second, on the mathematical side, regularization often leads to significant simplification when discussing continuity. For instance it allows for fully general continuity proofs in situations where it guarantees uniform visitation of all states $\bar{s} \in \bar{S}_{\lambda_0}$ over all iterations of eUDRL (equivalent to a lower bound on the state visitation distribution). As a case study we study a simple form of regularization, which, akin to ϵ -greedy exploration, involves a convex combination of an algorithm's current policy and a uniform policy. More precisely, choosing $\epsilon \in (0, 1)$ we extend the eUDRL recursion of formula (4.1) to an ϵ -regularized recursion (ϵ -eUDRL) by

$$\pi_{n+1,\epsilon}(a|s,h,g) = (1-\epsilon) \frac{\operatorname{num}_{\lambda,\pi_{n,\epsilon}}(a,s,h,g)}{\operatorname{den}_{\lambda,\pi_{n,\epsilon}}(s,h,g)} + \epsilon \frac{1}{|\mathcal{A}|} \quad \text{for } (s,h,g) \in \operatorname{supp den}_{\lambda,\pi_{n,\epsilon}},$$

where $\pi_{n+1,\epsilon}(a|s, h, g) = 1/|\mathcal{A}|$ outside the support supp den_{$\lambda,\pi_{n,\epsilon}$}. The sequence of policies generated by this recursion automatically satisfies $\pi_{n,\epsilon} > 0$, which ensures that the relevant state visitation probabilities are strictly positive. As compared to the discussion of continuity in the preceding sections this simplification avoids the complicated maintenance of support terms in the course of proofs (as policy-supports are always maximal). It should be said that the price to pay for this simplification is that ϵ -eUDRL does not converge to optimal policies for $\epsilon > 0$.

Below, we will present versions of the results previously encountered, now with regularization applied. Many of the results presented in sections 4, 5 and 7 remain valid up to small modification. For brevity and to avoid repetition we decided to leave the precise statement of the ϵ -eUDRL versions of the involved lemmas to appendix D focusing only on the main finding in this section. From a conceptual perspective lemma 6 (*Stability of* \bar{S}_{λ_0}), lemma 10 (*Optimality of eUDRL policies for deterministic transition kernels*), lemma 21 (*Continuity of action-values in quotient topology*), lemma 22 (*Lower bound on visitation probabilities*), lemma 24 (*f-lemma*) played key roles in the derivations of our main continuity results in sections 6 and 7. We recapitulate their content highlighting the specific modifications required for ϵ -eUDRL. Lemma 40 (ϵ -eUDRL version of lemma 6) addresses the behavior of supports of ϵ -eUDRL-generated quantities along compatible families of MDPs. Lemma 41 (ϵ -eUDRL version of lemma 10) describes the behavior of ϵ -eUDRL at deterministic transition kernels. Writing λ_0 for a deterministic kernel, the set

$$\Pi_{\lambda_0,\epsilon}^* = \left\{ (1-\epsilon)\pi_{\lambda_0}^* + \frac{\epsilon}{|\mathcal{A}|} \mid \pi_{\lambda_0}^* \text{ is an optimal policy for } \lambda_0 \text{ on } \bar{\mathcal{S}}_{\lambda_0} \right\}$$

will take the role of the set of optimal policies. Lemma 41 asserts that the accumulation points of ϵ -eUDRL are contained in $\Pi^*_{\lambda_0,\epsilon}$. To demonstrate the relative continuity of ϵ eUDRL policies at an infinite number of iterations, thus, the set $\Pi^*_{\lambda_0,\epsilon}$ should constitute the relative limit. The lemma also establishes estimates on value functions corresponding to policies in $\Pi^*_{\lambda_0,\epsilon}$ that are important for the proof of the main theorem. Unlike the optimal action-value function $Q^*_{\lambda_0}$ for eUDRL, in the context of ϵ -eUDRL there are many possible action-value functions $Q^{\pi^*}_{\lambda_0}$ depending on the choice of $\pi^*_{\lambda_0,\epsilon} \in \Pi^*_{\lambda_0,\epsilon}$. $Q^{\pi^*_{\epsilon}}_{\lambda_0}$ are also not
constant on $\mathcal{O}(\bar{s})$ in general, and, consequently, do not represent an equivalence class under the quotient map relative to $\mathcal{O}(\bar{s})$. Lemma 44 (ϵ -eUDRL version of lemma 21) discusses a variation designed to address this point. A lower bound $\alpha(\delta, \epsilon)$ on the state visitation probabilities is provided in lemma 42 (ϵ -eUDRL version of the lemma 22). As before the main continuity results follow by studying a dynamical system to bound the ϵ -eUDRL policy recursion. The dynamical system properties emerging from an iterative application of a map $z_{\gamma,\epsilon,M}$ (ϵ -eUDRL version of f_{γ}) are discussed in lemma 45 (ϵ -eUDRL version of lemma 24). As it turns out this dynamical system has a unique fixed point $x^*(\gamma, \epsilon, M)$. The theorem shows the relative continuity of the set of accumulation points of ϵ -eUDRLgenerated policies in λ at λ_0 as n tends to infinity. Along the lines outlined in section 7.1.2 we conclude that the set of accumulation points of the sequence of ϵ -eUDRL-generated policies at $\lambda = \lambda_0$ is nonempty and contained in $\Pi^*_{\lambda_0,\epsilon}$.

Theorem 34 (Relative continuity of ϵ -eUDRL limit policies in deterministic kernels) Let $\{\mathcal{M}_{\lambda} : \lambda \in (\Delta S)^{S \times A}\}$ and $\{\overline{\mathcal{M}}_{\lambda} : \lambda \in (\Delta S)^{S \times A}\}$ be compatible families. Let λ_0 be a deterministic kernel and let $(\pi_{n,\lambda,\epsilon})_{n\geq 0}$ be a sequence of ϵ -eUDRL-generated policies with initial condition $\pi_0 > 0$, transition kernel λ and regularization parameter $\epsilon \in (0,1)$. Let $\epsilon_0 \in (0,1)$ be a fixed regularization parameter. Then for all $\pi_0 > 0$ the following statements hold:

1. Let $\mathcal{L}(\pi_0, \lambda, \epsilon)$ denote the set of accumulation points of $(\pi_{n,\lambda,\epsilon})_{n\geq 0}$. Then any function $u : (\pi_0, \lambda, \epsilon) \to u(\pi_0, \lambda, \epsilon) \in \mathcal{L}(\pi_0, \lambda, \epsilon)$ is relatively continuous in λ, ϵ at point λ_0, ϵ_0 on $\bar{\mathcal{S}}_{\lambda_0}$, *i.e.* for all $\bar{s} \in \bar{\mathcal{S}}_{\lambda_0}$

$$[u(\pi_0,\lambda,\epsilon_0)](\cdot|\bar{s}) \xrightarrow{\mathcal{O}(\bar{s})} \pi^*_{\lambda_0,\epsilon_0}(\cdot|\bar{s}) \quad as \quad \lambda \to \lambda_0.$$

2. Let $\alpha(\delta, \epsilon)$ is chosen as in lemma 42 and let $\beta, \tilde{\beta}, \epsilon \in (0, 1)$ be such that $1 > \gamma + \epsilon$, where $\gamma = \frac{\tilde{\beta}}{((1-\epsilon)^N - \beta)\alpha(\delta,\epsilon)}$ and let

$$U_{\delta}(\lambda_0) = \{\lambda \mid \max_{(s,a) \in \mathcal{S} \times \mathcal{A}} \|\lambda(\cdot \mid s, a) - \lambda_0(\cdot \mid s, a)\|_1 < \delta\}.$$

There exists $\delta > 0$ so that for all $\lambda \in U_{\delta}(\lambda_0)$ and all $\bar{s} \in \bar{S}_{\lambda_0}$ it holds

$$\liminf_{n} \pi_{n,\lambda,\epsilon}(\mathcal{O}(\bar{s})|\bar{s}) \ge x^*(\gamma,\epsilon,|\mathcal{O}(\bar{s})|)$$

and

$$x^*(\gamma,\epsilon,|\mathcal{O}(\bar{s})|) \to 1 - \epsilon_0 \left(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}\right) \quad as \quad (\beta,\tilde{\beta},\alpha,\epsilon) \to (0,0,\alpha(0,\epsilon_0),\epsilon_0),$$

where $x^*(\gamma, \epsilon, |\mathcal{O}(\bar{s})|)$ is given in lemma 45. Consequently for all $\bar{s} \in \bar{S}_{\lambda_0}$ it holds that

$$\liminf_{n} \pi_{n,\lambda,\epsilon}(\mathcal{O}(\bar{s})|\bar{s}) \to 1 - \epsilon_0 \left(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}\right) \quad as \quad (\lambda,\epsilon) \to (\lambda_0,\epsilon_0).$$

The theorem is proved in appendix D. Along the lines of section 7.1.3, we can extend the result also to the segment spaces Seg^{diag} and $\text{Seg}^{\text{trail}}$.

Theorem 35 (Relative continuity of accumulation points of ϵ -eUDRL-generated policies at deterministic kernels) Theorem 34 remains valid under the renaming $\pi_{n,\epsilon} \to \pi_{n,\epsilon}^{\text{diag/trail}}$.

Details are provided in appendix D. The estimates given in theorem 34 can also be made entirely explicit along the lines of section 7.1.4, providing estimates for ϵ -eUDRL-generated quantities as a functions of the distance δ from the deterministic kernel λ_0 .

Corollary 36 (Estimating the location of accumulation points – ϵ -eUDRL) Under the conditions of theorem 34 assume that $\delta \in (0, 1)$ and set

$$\alpha = \frac{2}{N(N-1)} \left(\min_{\bar{s} \in \bar{\mathcal{S}}_{\lambda_0}} \bar{\mu}(\bar{s}) \right) \left(\frac{\epsilon}{|\mathcal{A}|} \right)^N \left(1 - \frac{\delta}{2} \right)^N$$

Define the quantities $\tilde{\beta} = \frac{N\delta}{2}$, $x^*(\gamma, \epsilon, M) = \frac{\hat{x}^* + \sqrt{(\hat{x}^*)^2 + \frac{4\gamma\epsilon M}{|\mathcal{A}|}}}{2}$ where $\hat{x}^* = 1 - \epsilon(1 - \frac{M}{\mathcal{A}}) - \gamma$ (cf. lemma 45) and for horizon $h, 1 \le h \le N$,

$$\beta_{h} = \begin{cases} \max\{\delta, \tilde{\beta}\}, \text{ if } h = 1, \\ \delta + \kappa_{h-1} + \beta_{h-1}, \text{ if } h \ge 2, \end{cases}$$
$$\gamma_{h} = \frac{\tilde{\beta}}{((1-\epsilon)^{N} - \beta_{h})\alpha},$$
$$\kappa_{h} = \max_{\bar{s} = (s,h',g) \in \bar{\mathcal{S}}_{\lambda_{0}}, h' = h} 2\left(1 - \epsilon\left(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}\right) - x^{*}(\gamma_{h}, \epsilon, \mathcal{O}(\bar{s}))\right).$$

Further assume $\beta_h, \gamma_h \in (0, 1), 1 > \gamma_h + \epsilon$ and notice that $\beta_h, \kappa_h, \gamma_h$, are increasing in h and that $\beta_h, \kappa_h, \gamma_h \to 0$ as $\delta \to 0$. Then the following assertions hold for all $\lambda \in U_{\delta}(\lambda_0)$ and for all $\pi_0 > 0$:

1.
$$\lim_{n} \sup_{\bar{s} \in \bar{\mathcal{S}}_{\lambda_{0}}} 2\left(1 - \epsilon \left(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}\right) - \pi_{n,\epsilon}(\mathcal{O}(\bar{s})|\bar{s})\right) \leq \kappa_{N},$$

2.
$$\left(\exists (\pi_{n,\epsilon}^{*}), \pi_{n,\epsilon}^{*} \in \Pi_{\lambda_{0},\epsilon}^{*}\right) : \lim_{n} \sup_{\bar{s},a \in \bar{\mathcal{S}}_{\lambda_{0}} \times \mathcal{A}} \max_{\lambda_{0},\epsilon} |Q_{\lambda}^{\pi_{n,\epsilon}}(\bar{s},a) - Q_{\lambda_{0}}^{\pi_{n,\epsilon}^{*}}(\bar{s},a)| \leq \beta_{N},$$

3.
$$\left(\exists (\pi_{n,\epsilon}^{*}), \pi_{n,\epsilon}^{*} \in \Pi_{\lambda_{0},\epsilon}^{*}\right) : \limsup_{n} \max_{\bar{s} \in \bar{\mathcal{S}}_{\lambda_{0}}} |V_{\lambda}^{\pi_{n,\epsilon}}(\bar{s}) - V_{\lambda_{0}}^{\pi_{n,\epsilon}^{*}}(\bar{s})| \leq \beta_{N} + \kappa_{N},$$

4.
$$(\exists (\pi_{n,\epsilon}^*), \pi_{n,\epsilon}^* \in \Pi_{\lambda_0,\epsilon}^*)$$
: $\limsup_{n} |J_{\lambda}^{\pi_{n,\epsilon}} - J_{\lambda_0}^{\pi_{n,\epsilon}^*}| \le \frac{N\delta}{2} + \beta_N + \kappa_N,$

5. for all $\epsilon' > 0$ there exists n_0 and $(\pi_{n,\epsilon}^*)_{n \ge 0}$, $\pi_{n,\epsilon}^* \in \Pi_{\lambda_0,\epsilon}^*$ so that for all $n \ge n_0$ and for all $\bar{s} \in \bar{S}_{\lambda_0}$ it holds that

$$\|\pi_{n,\epsilon}(\cdot|\bar{s}) - \pi_{n,\epsilon}^*(\cdot|\bar{s})\|_1 \le 2\left(1 - \epsilon\left(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}\right) - z_{\gamma',\epsilon,|\mathcal{O}(\bar{s})|}^{\circ(n-n_0)}(x_0)\right)$$

where $x_0 = \frac{\epsilon |\mathcal{O}(\bar{s})|}{|\mathcal{A}|}, \ \beta' > \beta_N + \epsilon'$, where ϵ' is chosen such that $1 > \gamma' + \epsilon'$ with $\gamma' = \frac{\tilde{\beta}}{((1-\epsilon)^N - \beta')\alpha} > 0.$



(a) Accumulation point estimates - ϵ -eUDRL. (b) Acc. point estimates - ϵ -eUDRL & eUDRL.

Figure 11: Illustration of estimates of corollary 36 for ϵ -eUDRL. Plots show the behavior of $\min_{\bar{s}\in\bar{S}_{\lambda_0}} \pi_n(\mathcal{O}(\bar{s})|s)$ and estimates for a 2-armed bandit for several values of the regularization parameter. Plot (a) showcases the dependency on the distance δ to a deterministic kernel, where dotted graphs depict accumulation points (approximated by values at n = 100) and the estimates $\min_M x^*(\gamma, \epsilon, M)$ provided in the corollary are depicted by solid graphs. For comparison, plot (b) shows the estimates provided in the corollary 36 together with the estimates $x^*(\gamma_N)$ and $x_u(\delta)$ from corollaries 27 and 33.

The terms on the right hand side of 1.-4. converge to 0 as $\delta \to 0$. In 5. the sequence $(y_n)_{n\geq 0}$ defined by $y_n = 2(1-\epsilon(1-\frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|})-z^{\circ(n-n_0)}_{\gamma',\epsilon,|\mathcal{O}(\bar{s})|}(x_0))$ converges q-linearly to the limit $y = 2(1-\epsilon(1-\frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|})-x^*(\gamma',\epsilon,|\mathcal{O}(\bar{s})|))$ at convergence rate $\frac{(1-\epsilon)\gamma'}{(x^*(\gamma',\epsilon,|\mathcal{O}(\bar{s})|)+\gamma')^2}$ which tends to 0 as $\delta \to 0$.

The result is proved in appendix D. Similarly to remark 29, points 1.-4. of the corollary can be formulated in terms of accumulation points. An illustration of the accumulation points of $\min_{\bar{s}\in\bar{S}_{\lambda_0}} \pi_{n,\lambda,\epsilon}(\mathcal{O}(\bar{s})|\bar{s})$ and the lower estimate provided by corollary 36 is given in figure 11 in the case of a 2-armed bandit model. All details of the construction of the MDP underlying figure 11 can be found in the appendix B, example 4. Figure 11a shows the δ -dependency of accumulation points of ϵ -eUDRL together with the estimates on accumulation points $\min_{\bar{s}\in\bar{S}_{\lambda_0}} x^*(\gamma,\epsilon,|\mathcal{O}(\bar{s})|)$ for several values of regularization parameter. For reference, we also plot the estimates for accumulation points for eUDRL in figure 11b although they are not directly comparable to those of ϵ -eUDRL. Figure 12 illustrates the derived estimates in case of a regularized ODT recursion applied to a grid world. This example is characterized by a non-deterministic optimal policy, cf. 12b, such that the x_u based bound for ODT without regularization cannot be used. All details of the construction of the MDP underlying figure 12 can be found in the appendix B, example 7.





(a) Accumulation point estimates - reg. ODT

(b) Map of grid world.

Figure 12: Illustration of estimates of corollary 36 for a regularized ODT recursion. Plots show the behavior of accumulation points of $\min_{\bar{s}\in\bar{S}_{\lambda_0}}\pi_n(\mathcal{O}(\bar{s})|s)$ and respective estimates for a grid world example. Plot (a) shows accumulation points of $\min_{\bar{s}\in\bar{S}_{\lambda_0}}\pi_{n,\lambda,\epsilon}(\mathcal{O}(\bar{s})|\bar{s})$ of the ODT recursion with uniform regularization, $\epsilon = 0.1$, together with the lower bound $\min_M x^*(\gamma, \epsilon, M)$. Plot (b) shows the map of the grid world. A wall is depicted in gray, the set \bar{S}_{λ_0} in yellow, and the goal in red. Arrows depict optimal actions associated with the specific state and goal. We conclude this section with a brief summary. Adding regularization allowed us to carry out the discussion of continuity in full generality. This covers domains like, e.g., example 7, which does not fall into the special cases of section 7. On the other hand the more regularization is applied, the further the asymptotic policy must deviate from an optimal policy. Notice that policies from $\Pi^*_{\lambda_0,\epsilon}$, $\bar{s} \in \bar{S}_{\lambda_0}$, are only assigning a mass of $1 - \epsilon \left(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}\right)$ to $\mathcal{O}(\bar{s})$ while an optimal policy is assigning a mass of 1 to $\mathcal{O}(\bar{s})$, cf. the discussion after the lemma 44. On the other hand a smaller value of ϵ entails a smaller lower the bound on the state visitation terms $\alpha(\delta, \epsilon)$. This, in turn, affects γ in that a smaller distance to a deterministic kernel δ is needed to recover asymptotic behavior for $\delta \to 0$, cf. corollary 36 and lemma 45 for the behavior of respective quantities. The trade-off in the choice of ϵ is apparent the comparison of $x^*(\gamma, \epsilon, M)$ bounds for several values of ϵ in figure 11.

9 Related Work

In this section, we start by discussing key early works that looked at the underlying theme of using iterated supervised learning to solve reinforcement learning problems in section 9.1. We then look at works associated with the more specific goal-conditioned reinforcement learning in section 9.2. After, we tackle the most relevant works to this article—those directly relating to upside-down reinforcement learning—in section 9.3. We end this section by discussing critical work associated with the transformer architecture used by decision transformers and online decision transformers in section 9.4.

9.1 Reward-Weighted Regression

The idea of using *iterated supervised learning to solve reinforcement learning problems* was previously investigated by Peters and Schaal (2007) in the form of Reward-Weighted Regression (RWR), which itself built on the Expectation-Maximization framework for RL by Dayan and Hinton (1997). However, the setting explored in their works is very limited; the work of Dayan and Hinton (1997) considered only a bandit scenario (only one state and a finite number of actions) and Peters and Schaal (2007) restricted their work to immediate reward problems. The extension of RWR to consider a full episodic setting (considering not only immediate rewards but rather full returns) was done by Wierstra et al. (2008) and Kober and Peters (2011). Efficient off-policy schemes in the context of RWR were later discussed by Hachiya et al. (2009, 2011). Finally, the use of deep neural network approximators was introduced to RWR by Peng et al. (2019).

Theoretical studies of RWR are more sparse, with the monotonicity of RWR proved by Dayan and Hinton (1997) and Peters and Schaal (2007) in their respective settings. Both of the aforementioned monotonicity proofs build on the Expectation-Maximization paradigm. More than a decade after the work on monotonicity, RWR's convergence to a global optimum was proved by Štrupl et al. (2022b) for compact state and action spaces (assuming an infinite number of samples and no function approximation).

RWR remains particularly relevant for this work because eUDRL on Seg^{diag} coincides with RWR (as discussed in section 3). This link motivated our approach to the continuity proofs of eUDRL generated quantities at deterministic kernels.

9.2 Goal-conditioned Reinforcement Learning Without Relabeling

To our knowledge, the first work that could be considered goal-conditional RL was published by Schmidhuber and Huber (1991, 1990) in the context of learning selective attention. In this work, there are extra goal-defining input patterns that encode various tasks so that the RL machine knows which task to execute next. The rewards are granted at the end of each task. The need to propagate learning signals through a non-differentiable environment is elevated by a separate environment model network. Essentially, it is model-based RL with sparse rewards, which is equivalent to SL except samples are collected online. However, note that, in contrast to UDRL, this work and also all other works in this paragraph are missing segment/trajectory relabeling. The work by Schmidhuber and Huber (1991) also contributes to a body of literature on attention which provides grounds for today transformer architectures. The options framework of Sutton et al. (1999) could also be considered as goal-conditioned RL (an option determines a policy). Providing goals to a policy network provides possibility of generalization to unseen goals when representing the policy by deep network (similarly for value functions). Building on the work of Sutton et al. (2011), the work of Schaul et al. (2015) proposes convenient approximators of goalconditioned value functions. For an example of a more recent work on goal-conditioned RL, we can cite the paper by Faccio et al. (2023) which introduced goal-conditioned policy generators for deep policy networks.

9.3 UDRL

UDRL (Schmidhuber, 5 Dec 2019) extends the *iterated SL* idea by learning commandconditioned policies using trajectory segments relabeled with the achieved goals/returns. This was motivated by the desire to obtain generalization across goals/returns (when representing the UDRL policy by a deep neural network), and by the desire for more efficient data usage (there are more segments in a given trajectory than there are states). Trajectory relabeling appeared already before in Hindsight Experience Replay (HER) by Andrychowicz et al. (2017) and we relate to this work in more detail below.

Schmidhuber (5 Dec 2019) designed the specific algorithm that we denote here as eUDRL only for deterministic environments (though, some speculation was done by Schmidhuber (5 Dec 2019) on variants for stochastic environments).

However, it is clear that there are many useful stochastic environments which are close to deterministic. With this motivation (a return-reaching variant of) eUDRL was demonstrated by Srivastava et al. (2019) to solve successfully many non-deterministic environments (e.g. MuJoCo tasks) featuring transition kernels exhibiting only small non-determinism. This was afterwards followed by Ghosh et al. (2021) in context of GCSL (essentially eU-DRL restricted to fixed horizon and state reaching task) with further impressive benchmark performance. Further concurrently there appeared also work of Kumar et al. (2019) which is again essentially eUDRL.

The idea of combining eUDRL with transformer architectures Schmidhuber (1992); Vaswani et al. (2017); Schlag et al. (2021) as policy approximator was successfully demonstrated in the Decision Transformer (DT) architecture proposed by Chen et al. (2021). Chen et al. (2021), however, only looked at the offline RL setting (equivalent to a single eUDRL iteration). Shortly after the DT architecture was introduced, a generalized version of it was proposed by Furuta et al. (2021), which led to a non-trivial performance improvement.

Štrupl et al. (2022a) and Paster et al. (2022) described the problems and causes of eUDRL convergence issues in environments with stochastic transition dynamics. While the work of Štrupl et al. (2022a) rewrote the recursion in eUDRL to identify the convergence issues, the work of Paster et al. (2022) directly proposed a solution to this issue in the context of DTs.

Several later works were proposed to fix the issues with eUDRL stochastic environments, such as the work by Yang et al. (2023) in context of DT and the work of Faccio et al. (2023) in context of eUDRL.

More relevant to this work than much of the above is some of the theoretical work that has been done with regard to eUDRL and related algorithms. Here, the most related practical and theoretical works are those of Brandfonbrener et al. (2022), Ghosh et al. (2021), Kumar et al. (2019), Andrychowicz et al. (2017), and Paster et al. (2022). We provide a comprehensive treatment of each of these works below.

Brandfonbrener et al. (2022) discusses eUDRL (and other related algorithms) in context of offline RL, i.e., it investigates just one iteration of eUDRL aiming for sample bounds. In contrast, we are interested in developing continuity results and bounds for both any finite number of iterations and for the asymptotic case where there is an infinite sample size. So, there is an overlap between this work and that of Brandfonbrener et al. (2022) in some trivial results (namely, the continuity of expected return (goal-reaching objective in our case) for the first iteration as shown by theorem 1 of Brandfonbrener et al. (2022)). However, note that the continuity for the first iteration is trivial to obtain, as the initial policies do not depend on λ . The main difficulty comes from policies being functions of λ . This leads to a discontinuity of policies in λ at λ_0 (a deterministic kernel) in \bar{S}_{λ_0} for the second and later iterations.⁷ To overcome this difficulty, one has to utilize some weaker notion of continuity. Here, we used the notion of relative continuity (see the induction step in our proof of theorem 16). Developing asymptotic results is yet another level of complexity (see sections 7 and 8).

Ghosh et al. (2021) introduces the following lower bound on the goal-reaching objective (see theorem 3.1 of Ghosh et al. (2021)):

$$J^{\pi} \ge J_{\text{GCSL}}(\pi) - 4T(T-1)\alpha^2 + C,$$

where J^{π} denotes the goal-reaching objective for a policy π , $J_{\text{GCSL}}(\pi)$ denotes a GCSL objective for the policy π (the same as eUDRL objective (2.7) for trailing segments), Tdenotes a fixed horizon, $\alpha' := \max_{\bar{s}} D_{\text{TV}}(\pi(\cdot|\bar{s})|\pi_{\text{old}}(\cdot|\bar{s}))$ denotes total variation distance of the policy π from policy π_{old} which was used to collect trajectories, and C is a constant on π . There are (at least) two problems when one tries to adopt this result in our work to, for example, assess the continuity of the goal-reaching objective (or the δ -dependent error bounds for the goal-reaching objective).

The first of the aforementioned problems is that to use the bound in a meaningful way, one has to minimize α' . This could be done by assuming that a sequence of GCSL (eUDRL) policies has a limit (then $\alpha' \to 0$). This is, however, difficult to prove. To maintain a certain level of rigor, we carefully avoid the limit assumption and instead investigate all the possible accumulation points. From this perspective, this result is not useful as, for a finite number of iteration, α' could be large. Nor is it useful for the asymptotic case as we could not afford the limit assumption.

The second of the aforementioned problems is that the dependence on δ (the distance of the transition kernel λ to a deterministic kernel λ_0) is not considered, as, in order to use the bound (e.g. for assessing continuity in λ at λ_0), one has to determine the dependence on δ for all three terms on the right side. This seems to be more challenging than determining this dependence for J^{π} itself (as we do here) and would most likely still be much less precise (see the point above).

Kumar et al. (2019) introduced an algorithm for learning return-conditioned policies that is essentially eUDRL on Seg_{trail} with fixed horizon. However, while the authors do

^{7.} We refer the reader to the discussion at the end of appendix C which justifies this statement.

include some theoretical discussion on the properties of eUDRL, this discussion does not touch on the distance of an MDP transition kernel to a deterministic kernel (the role played by deterministic kernels is not investigated at all). Therefore, with regards to the continuity of the transition kernel we are particularly interested in here, this work has little to offer.

Andrychowicz et al. (2017) introduced Hindsight Experience Replay (HER), the popular extension of experience replay that has be applied to many off-policy RL algorithms, such as DQN and DDPG. HER is closely related to eUDRL in some aspects. Like eUDRL, HER deals with goals, a goal map, and goal dependent policies (note that a dependence on the time step/remaining horizon could be accounted for by including it into the state representation). HER is formulated for a fixed horizon with discounting (no discounting case is included). The principle of the HER extension relies in populating the replay buffer not only with the trajectory that the agent actually encountered, but also with its "relabeled" version, where the original goals are replaced with the actually achieved final state and the corresponding rewards. HER then uses an off-policy algorithm to learn value functions and a policy using data in the replay buffer.

The main distinguishing characteristics of eUDRL with respect to HER are that (1) HER uses not just the relabeled trajectories but also the original trajectories, i.e., with the original intended goal; and, (2) HER does not fit the next policy directly to some action conditional (at least when DQN or DDPG is used as the offline algorithm as in the original HER paper), e.g., DQN aims to rather learn the critic and derive the policy as an epsilon-greedy policy using the critic. These are the main problems when one wants to connect HER and eUDRL recursion. The second problem could be elevated by using a convenient RL algorithm (e.g. RWR) after which, because the substantial part of the data is not relabeled, we are left with some crossover between RWR and eUDRL recursions. This observation could already hint some properties of an HER extension of RWR. Note that some of the problems HER shares with eUDRL were already reported, analyzed, and/or fixed in the literature. We refer the interested reader to the works of Lanka and Wu (2018) and Schramm et al. (2023).

Paster et al. (2022) describes the causes behind DT divergence in stochastic environments and proposes a fix. To resolve the issues with convergence in stochastic domains, instead of conditioning on return, Paster et al. (2022) propose to condition on a statistics $I(\tau)$ (where τ stands for a trajectory) which is independent of the stochasticity of the environment. In their work, I is represented by NN and learned using an adversarial scheme. Trajectories are then clustered according to I values with I values subsequently mapped to the average return of their corresponding cluster. The result is then that by conditioning the policy on I values, the corresponding average return can be reached consistently, i.e., in expectation. This is in contrast to when conditioning on a statistics which is generally dependent on the environment's stochasticity, such as the trajectory return (or some general abstract goals like in eUDRL). For a detailed description of what it means to be "independence of the environment stochasticity" please refer to the original article by Paster et al. (2022).

9.4 Transformers

Transformers with unnormalized linearized self attention (also known as Unnormalized Linear Transformers, or ULTRAs), appeared at least as early as the 1990s in works by Schmidhuber (1992, 1991) under name fast-weight programmers (Schlag et al., 2021). The goal of FWPs was to obtain a more storage-efficient alternative to recurrent networks by processing each input with both a slow and a fast network, where the output of the slow network provided changes to the weights of the fast network. The attention terminology was introduced shortly after by Schmidhuber (1993). The modern transformer architecture was introduced by Vaswani et al. (2017) and scaled quadratically in input size. More recently, variants of the Vaswani et al. (2017) architecture have reverted to using linearized attention (e.g., see the work by Katharopoulos et al. (2020) and Choromanski et al. (2020)), as this results in a linear scaling.

10 Conclusion

Our contribution lies in being the first rigorous treatise on the convergence and stability of some key "reinforcement learning through supervised learning" approaches. Specifically, our study focuses on both MDPs of the CE type (on the side of their mathematical formulation) and on eUDRL (on the side of the algorithms investigated). Taken together, these frameworks offer sufficient generality to accommodate common "RL through SL" training schemes, including GCSL, ODT and RWR. In particular, we have shown how GCSL, ODT, and RWR can be expressed in the framework of CEs and interpreted as derivatives of eUDRL. This is achieved by restricting the technical analysis to specific segment spaces Seg^{trail} and Seg^{diag}. In essence, the training iteration of GCSL, ODT and RWR can be understood as instances of the eUDRL training iteration on Seg, Seg^{trail} or Seg^{diag}, for a conveniently chosen CE. To accommodate such training schemes, we expanded our analysis of convergence and stability to the segment spaces Seg^{trail} and Seg^{diag}, in addition to proving results for Seg. Our findings can roughly be categorized into three clusters: (1) stability analysis of training schemes at a finite number of iterations, (2) investigation of the asymptotic properties of training schemes, and (3) results that complement the discussion of algorithms targeting a broader and more complete mathematical picture.

10.1 Investigation of Stability at a Finite Number of Iterations

In this work, we have demonstrated the instability of eUDRL at the boundary of the space of transition kernels. Specifically, we have provided examples of environments that are characterized by non-deterministic kernels located on the boundary, where even a tiny perturbation of the kernel can result in a significant, arbitrary shift in the value or in the goal-reaching objective generated by the training schemes. Examples 1 and 2, representing non-deterministic or deterministic transition kernels, respectively, illustrate the presence of non-removable discontinuities in eUDRL-generated policies at the boundary of the transition kernel space for iterations n > 2. In contrast, within the interior of the kernel space, both policies and values remain continuous. However, at deterministic kernels, while values are continuous, policies might have non-removable discontinuities. To address this, we introduced the concept of relative continuity, which corresponds to a form of continuity with respect to a quotient topology. Using this concept, we showed that eUDRL policies are relatively continuous at deterministic kernels for any finite number of iterations. This entails the continuity of the goal-reaching objective and the stability of eUDRL-type training schemes at deterministic kernels at any finite number of iterations. Together with the convergence of eUDRL to optimality at deterministic kernels, this implies near-optimality of the goal-reaching objective at transition kernels that are close to deterministic.

10.2 Convergence and Investigation of Stability at an Infinite Number of Iterations

In this work, we investigated the asymptotic behavior of eUDRL-generated quantities as the number of iterations tends to infinity. We established the relative continuity of the sets of accumulation points of eUDRL policies and the continuity of the associated goal-reaching objectives at deterministic kernels, specifically for two important special cases. These cases are characterized by additional assumptions that facilitate the analysis of continuity.

- 1. The support of the CE's initial distribution contains the set of "critical states" $S_{\lambda_0} \subset \sup p\bar{\mu}$. In simple terms, \bar{S}_{λ_0} consists of the states that are essential to prove the continuity of the goal-reaching objective at a given deterministic kernel λ_0 . States that cannot be reached by the CE, in particular, are excluded from \bar{S}_{λ_0} .
- 2. The optimal policy is unique on \bar{S}_{λ_0} , which means that for all states $(\forall \bar{s} \in \bar{S}_{\lambda_0}) : |\mathcal{O}(\bar{s})| = 1$. This condition alleviates the complexity of studying relative continuity, significantly simplifying the analysis of stability and convergence.

Thus, both conditions entail near-optimal behavior of the goal-reaching objective at neardeterministic kernels in the asymptotic limit, as is also the case at any finite number of iterations. Making the established estimates fully explicit, we derived estimates that quantify the error from optimality based on the distance of the transition kernel from a deterministic kernel. We also derived a bound on the eUDRL policy error and assessed its q-linear convergence rate. Although we believe that the outlined conditions encompass a wide range of practical scenarios, a fully general discussion of the relative continuity of accumulation point sets for eUDRL-generated policies at deterministic kernels remains an open problem. Motivated by the widespread use of regularization in "RL through SL" training schemes (e.g., in ODT), we investigated ϵ -greedy regularization of the policy iteration of eUDRL. We established the relative continuity of the set of accumulation points of ϵ -eUDRL-generated policies in full generality. As before, this entails the continuity of the accumulation point sets of the corresponding goal-reaching objective. We also showed that the goal-reaching objective exhibits near-optimal behavior at near-deterministic kernels in the asymptotic limit for ϵ -eUDRL. Akin to the discussion without regularization, we provide estimates that quantify the error from optimality given the distance of the transition kernel from a deterministic kernel for ϵ -eUDRL. Notice, however, that the asymptotic analysis is conducted in full generality without relying on the special assumptions used in previous cases.

10.3 Further Results

We established the continuity of eUDRL-generated policies and goal-reaching objectives at any finite number of iterations for transition kernels located within the interior of the space of all kernels. Unlike deterministic kernels, this does not immediately entail convergence to near-optimality. We include this result to complete the mathematical picture and to stipulate further work. We also included a proof of the optimality of eUDRL at deterministic kernels: While this result may be considered as implicitly understood within the literature, we decided to write it up for completeness. The mathematical investigation is supplemented by a range of worked-out examples that might prove useful for the continued development of this research area. Specifically, we illustrated our results through examples involving a 2-armed bandit, a random walk on \mathbb{Z}_3 and a 3×3 grid world domain. The code which was used for computing the examples and generating the associated figures is available at https://github.com/struplm/eUDRL-GCSL-ODT-Convergence-public

Acknowledgments and Disclosure of Funding

This work was supported by the European Research Council (ERC, Advanced Grant Number 742870), the Swiss National Supercomputing Centre (CSCS, Project s1090), and by the Swiss National Science Foundation (Grant Number 200021_192356, Project NEUSYM). We also thank both the NVIDIA Corporation for donating a DGX-1 as part of the Pioneers of AI Research Award and IBM for donating a Minsky machine.

References

- Michael Ahn, Henry Zhu, Kristian Hartikainen, Hugo Ponte, Abhishek Gupta, Sergey Levine, and Vikash Kumar. Robel: Robotics benchmarks for learning with low-cost robots. In *Conference on robot learning*, pages 1300–1313. PMLR, 2020.
- Marcin Andrychowicz, Filip Wolski, Alex Ray, Jonas Schneider, Rachel Fong, Peter Welinder, Bob McGrew, Josh Tobin, OpenAI Pieter Abbeel, and Wojciech Zaremba. Hindsight experience replay. In I. Guyon, U. Von Luxburg, S. Bengio, H. Wallach, R. Fergus, S. Vishwanathan, and R. Garnett, editors, Advances in Neural Information Processing Systems, volume 30. Curran Associates, Inc., 2017.
- Dimitri P. Bertsekas and John N. Tsitsiklis. Neuro-Dynamic Programming. Athena Scientific, 1996.
- David Brandfonbrener, Alberto Bietti, Jacob Buckman, Romain Laroche, and Joan Bruna. When does return-conditioned supervised learning work for offline reinforcement learning? Advances in Neural Information Processing Systems, 35:1542–1553, 2022.
- Lili Chen, Kevin Lu, Aravind Rajeswaran, Kimin Lee, Aditya Grover, Misha Laskin, Pieter Abbeel, Aravind Srinivas, and Igor Mordatch. Decision transformer: Reinforcement learning via sequence modeling. Advances in neural information processing systems, 34:15084– 15097, 2021.
- Krzysztof Choromanski, Valerii Likhosherstov, David Dohan, Xingyou Song, Andreea Gane, Tamas Sarlos, Peter Hawkins, Jared Davis, Afroz Mohiuddin, Lukasz Kaiser, et al. Rethinking attention with performers. arXiv preprint arXiv:2009.14794, 2020.
- Peter Dayan and Geoffrey E. Hinton. Using expectation-maximization for reinforcement learning. Neural Comput., 9(2):271-278, 1997. doi: 10.1162/neco.1997.9.2.271. URL https://doi.org/10.1162/neco.1997.9.2.271.
- Francesco Faccio, Vincent Herrmann, Aditya A. Ramesh, Louis Kirsch, and Jürgen Schmidhuber. Goal-conditioned generators of deep policies. In Brian Williams, Yiling Chen, and Jennifer Neville, editors, Thirty-Seventh AAAI Conference on Artificial Intelligence, AAAI 2023, Thirty-Fifth Conference on Innovative Applications of Artificial Intelligence, IAAI 2023, Thirteenth Symposium on Educational Advances in Artificial Intelligence, EAAI 2023, Washington, DC, USA, February 7-14, 2023, pages 7503-7511. AAAI Press, 2023. doi: 10.1609/AAAI.V37I6.25912. URL https://doi.org/10.1609/aaai.v37i6.25912.
- Justin Fu, Aviral Kumar, Ofir Nachum, George Tucker, and Sergey Levine. D4rl: Datasets for deep data-driven reinforcement learning. arXiv preprint arXiv:2004.07219, 2020.
- Hiroki Furuta, Yutaka Matsuo, and Shixiang Shane Gu. Generalized decision transformer for offline hindsight information matching. arXiv preprint arXiv:2111.10364, 2021.
- Dibya Ghosh, Abhishek Gupta, Ashwin Reddy, Justin Fu, Coline Devin, Benjamin Eysenbach, and Sergey Levine. Learning to reach goals via iterated supervised learning. In International Conference on Learning Representations, 2021.

- Hirotaka Hachiya, Jan Peters, and Masashi Sugiyama. Efficient sample reuse in embased policy search. In Wray L. Buntine, Marko Grobelnik, Dunja Mladenic, and John Shawe-Taylor, editors, Machine Learning and Knowledge Discovery in Databases, European Conference, ECML PKDD 2009, Bled, Slovenia, September 7-11, 2009, Proceedings, Part I, volume 5781 of Lecture Notes in Computer Science, pages 469–484. Springer, 2009. doi: 10.1007/978-3-642-04180-8_48. URL https://doi.org/10.1007/ 978-3-642-04180-8_48.
- Hirotaka Hachiya, Jan Peters, and Masashi Sugiyama. Reward-weighted regression with sample reuse for direct policy search in reinforcement learning. *Neural Comput.*, 23(11): 2798–2832, 2011. doi: 10.1162/NECO_a_00199. URL https://doi.org/10.1162/NECO_a_00199.
- Angelos Katharopoulos, Apoorv Vyas, Nikolaos Pappas, and François Fleuret. Transformers are rnns: Fast autoregressive transformers with linear attention. In *International conference on machine learning*, pages 5156–5165. PMLR, 2020.
- Michał Kempka, Marek Wydmuch, Grzegorz Runc, Jakub Toczek, and Wojciech Jaśkowski. Vizdoom: A doom-based ai research platform for visual reinforcement learning. In 2016 IEEE conference on computational intelligence and games (CIG), pages 1–8. IEEE, 2016.
- Jens Kober and Jan Peters. Policy search for motor primitives in robotics. *Mach. Learn.*, 84 (1-2):171-203, 2011. doi: 10.1007/s10994-010-5223-6. URL https://doi.org/10.1007/s10994-010-5223-6.
- Aviral Kumar, Xue Bin Peng, and Sergey Levine. Reward-conditioned policies. arXiv preprint arXiv:1912.13465, 2019.
- Sameera Lanka and Tianfu Wu. Archer: Aggressive rewards to counter bias in hindsight experience replay, 2018. URL https://arxiv.org/abs/1809.02070.
- Jorge Nocedal and Stephen J. Wright. Numerical Optimization. Springer Series in Operations Research, 2006.
- Keiran Paster, Sheila McIlraith, and Jimmy Ba. You can't count on luck: Why decision transformers and rvs fail in stochastic environments. Advances in neural information processing systems, 35:38966–38979, 2022.
- Xue Bin Peng, Aviral Kumar, Grace Zhang, and Sergey Levine. Advantage-weighted regression: Simple and scalable off-policy reinforcement learning, 2019.
- Jan Peters and Stefan Schaal. Reinforcement learning by reward-weighted regression for operational space control. In Zoubin Ghahramani, editor, Machine Learning, Proceedings of the Twenty-Fourth International Conference (ICML 2007), Corvallis, Oregon, USA, June 20-24, 2007, volume 227 of ACM International Conference Proceeding Series, pages 745–750. ACM, 2007. doi: 10.1145/1273496.1273590. URL https://doi.org/10.1145/ 1273496.1273590.
- Martin L Puterman. Markov Decision Processes: Discrete Stochastic Dynamic Programming. John Wiley & Sons, 2014.

- Tom Schaul, Daniel Horgan, Karol Gregor, and David Silver. Universal value function approximators. In *International conference on machine learning*, pages 1312–1320. PMLR, 2015.
- Imanol Schlag, Kazuki Irie, and Jürgen Schmidhuber. Linear transformers are secretly fast weight programmers. In *International Conference on Machine Learning*, pages 9355–9366. PMLR, 2021.
- J. Schmidhuber. Learning to control fast-weight memories: An alternative to recurrent nets. Technical Report FKI-128-90, Institut f
 ür Informatik, Technische Universit
 ät M
 ünchen, 1991.
- J. Schmidhuber and R. Huber. Learning to generate focus trajectories for attentive vision. Technical Report FKI-128-90, Institut f
 ür Informatik, Technische Universit
 ät M
 ünchen, 1990.
- J. Schmidhuber and R. Huber. Learning to generate artificial fovea trajectories for target detection. *International Journal of Neural Systems*, 2(1 & 2):135–141, 1991. (Based on TR FKI-128-90, TUM, 1990).
- Jürgen Schmidhuber. Learning to control fast-weight memories: An alternative to dynamic recurrent networks. *Neural Computation*, 4(1):131–139, 1992.
- Jürgen Schmidhuber. Reducing the ratio between learning complexity and number of time varying variables in fully recurrent nets. In ICANN'93: Proceedings of the International Conference on Artificial Neural Networks Amsterdam, pages 460–463. Springer, 1993.
- Jürgen Schmidhuber. Reinforcement learning upside down: Don't predict rewards-just map them to actions. *Preprint arXiv:1912.02875*, 5 Dec 2019.
- Liam Schramm, Yunfu Deng, Edgar Granados, and Abdeslam Boularias. Usher: Unbiased sampling for hindsight experience replay. In *Conference on Robot Learning*, pages 2073– 2082. PMLR, 2023.
- Rupesh Kumar Srivastava, Pranav Shyam, Filipe Mutz, Wojciech Jaskowski, and Jürgen Schmidhuber. Training agents with upside-down reinforcement learning. NNAISENSE Technical Report 201911-02, arXiv:1912.02877, 2019. NeurIPS 2019 Deep RL workshop.
- Richard S. Sutton and Andrew G. Barto. Reinforcement Learning: An Introduction. A Bradford Book, USA, 2018. ISBN 0262039249, 9780262039246.
- Richard S. Sutton, Doina Precup, and Satinder P. Singh. Between mdps and semi-mdps: A framework for temporal abstraction in reinforcement learning. Artif. Intell., 112(1-2): 181–211, 1999. doi: 10.1016/S0004-3702(99)00052-1. URL https://doi.org/10.1016/ S0004-3702(99)00052-1.
- Richard S Sutton, Joseph Modayil, Michael Delp, Thomas Degris, Patrick M Pilarski, Adam White, and Doina Precup. Horde: A scalable real-time architecture for learning knowledge from unsupervised sensorimotor interaction. In *The 10th International Conference on Autonomous Agents and Multiagent Systems-Volume 2*, pages 761–768, 2011.

- Ashish Vaswani, Noam Shazeer, Niki Parmar, Jakob Uszkoreit, Llion Jones, Aidan N. Gomez, Lukasz Kaiser, and Illia Polosukhin. Attention is all you need. Advances in Neural Information Processing Systems, 2017.
- Daan Wierstra, Tom Schaul, Jan Peters, and Jürgen Schmidhuber. Episodic reinforcement learning by logistic reward-weighted regression. In Vera Kurková, Roman Neruda, and Jan Koutník, editors, Artificial Neural Networks - ICANN 2008, 18th International Conference, Prague, Czech Republic, September 3-6, 2008, Proceedings, Part I, volume 5163 of Lecture Notes in Computer Science, pages 407–416. Springer, 2008. doi: 10.1007/ 978-3-540-87536-9_42. URL https://doi.org/10.1007/978-3-540-87536-9_42.
- Sherry Yang, Dale Schuurmans, Pieter Abbeel, and Ofir Nachum. Dichotomy of control: Separating what you can control from what you cannot. In *The Eleventh International Conference on Learning Representations*, 2023.
- Qinqing Zheng, Amy Zhang, and Aditya Grover. Online decision transformer. In Kamalika Chaudhuri, Stefanie Jegelka, Le Song, Csaba Szepesvari, Gang Niu, and Sivan Sabato, editors, Proceedings of the 39th International Conference on Machine Learning, volume 162 of Proceedings of Machine Learning Research, pages 27042–27059. PMLR, 17–23 Jul 2022. URL https://proceedings.mlr.press/v162/zheng22c.html.
- Miroslav Štrupl, Francesco Faccio, Dylan R. Ashley, Jürgen Schmidhuber, and Rupesh Kumar Srivastava. Upside-down reinforcement learning can diverge in stochastic environments with episodic resets, 2022a.
- Miroslav Štrupl, Francesco Faccio, Dylan R. Ashley, Rupesh Kumar Srivastava, and Jürgen Schmidhuber. Reward-weighted regression converges to a global optimum. *Proceedings* of the AAAI Conference on Artificial Intelligence, 36(8):8361-8369, Jun. 2022b. doi: 10. 1609/aaai.v36i8.20811. URL https://ojs.aaai.org/index.php/AAAI/article/view/ 20811.

Appendix A. The Segment Distribution and its Factorization

Since we assume a deterministic reward function, rewards are completely determined by state-action-state transitions and can be omitted from segments or trajectories. Once a CE has reached an absorbing state in $\bar{\mathcal{S}}_A$, the policy becomes irrelevant as all actions lead to the same absorbing state. For this reason, it is sufficient to restrict the discussion to all policies π that deterministically select a unique action a_A in all absorbing states $\bar{\mathcal{S}}_A$, i.e., $\forall \bar{s} \in \mathcal{S}_A : \pi(a_A | \bar{s}) = 1$. In this section, we assume that all trajectories are generated by using one fixed policy π . The length $l(\cdot)$ of a trajectory is defined as the number of transitions until an absorbing state is entered for the first time. Regarding trajectories, we can consider just prefixes of length N because the CE MDP bounds the remaining horizon component by N. Further, we can restrict ourselves to the subspace $\operatorname{Traj} \subset (\bar{\mathcal{S}} \times \mathcal{A})^N \times \bar{\mathcal{S}}$ allowed by remaining horizon/goal dynamics of CE, i.e., for a trajectory $\tau = (\bar{s}_0, a_0, \dots, \bar{s}_N) \in \text{Traj}$, $\bar{s}_t = (s_t, h_t, g_t), \ 0 \le t \le N$, the remaining horizon h_t decreases by 1 from its initial value $h_0 = l(\tau)$ till 0 when entering to an absorbing state. The goal $g_t = g_0$ remains unchanged. Thus τ is fully determined by the initial horizon $h_0 = l(\tau)$, the initial goal g_0 , the states $s_0, \ldots, s_{l(\tau)}$, and actions $a_0, \ldots, a_{l(\tau)-1}$, i.e., $\tau = ((s_0, h_0, g_0), a_0, (s_1, h_0 - g_0), a_0, (s_1, h_0 - g_0))$ $(1, g_0), a_1, \ldots, (s_{l(\tau)}, 0, g_0), a_A, \ldots, (s_{l(\tau)}, 0, g_0))$. The probability of τ is given as:

$$\mathbb{P}(\mathcal{T}=\tau;\pi) = \left(\prod_{t=1}^{l(\tau)} \lambda(s_t | a_{t-1}, s_{t-1})\right) \times \left(\prod_{t=0}^{l(\tau)-1} \pi(a_t | \bar{s}_t)\right) \times \bar{\mu}(\bar{s}_0),$$

where $\mathcal{T}: \Omega \to \text{Traj}, \mathcal{T} = ((S_0, H_0, G_0), A_0, \dots, (S_N, H_N, G_N))$ is the trajectory random variable map.

Segment Distribution: Segments are assumed to be continuous chunks of trajectories in Traj, so they respect the CE horizon/goal dynamics. We assume them to be always contained within the length of a trajectory. Notice that eUDRL is learning actions just in a state which happens to be the first state of a segment. Since learning actions in absorbing states is not meaningful, we will assume the first state of a segment to be transient (i.e., $\in \bar{S}_T)^8$. Thus, segments σ are fully determined by the segment length, denoted by $l(\sigma)$ (the number of transitions), by the remaining horizon and goal at the segment beginning $h_0^{\sigma}, g_0^{\sigma}$, and by $l(\sigma) + 1$ states and $l(\sigma)$ actions. Without loss of generality we will identify σ with such a tuple $\sigma = (l(\sigma), s_0^{\sigma}, h_0^{\sigma}, g_0^{\sigma}, a_0^{\sigma}, s_1^{\sigma}, a_1^{\sigma}, \dots, s_{l(\sigma)}^{\sigma})$). The space of all such tuples will be denoted Seg. Formally, we will assume that there exists a random variable map $\Sigma : \Omega \to \text{Seg}, \Sigma = (l(\Sigma), S_0^{\Sigma}, H_0^{\Sigma}, G_0^{\Sigma}, A_0^{\Sigma}, S_1^{\Sigma}, A_1^{\Sigma}, \dots, S_{l(\Sigma)}^{\Sigma})$ with distribution d_{Σ}^{π} given below.

We construct the segment distribution d_{Σ}^{π} in a similar way as the state visitation distribution—summing across appropriate trajectory distribution marginals. This causes the result to be un-normalized (among other restrictions, e.g., on the first state of the

^{8.} One could likewise additionally assume the original MDP component of this state to be transient for the similar reason.

segment), therefore we have to include a normalization constant c. $(\forall \sigma \in \text{Seg})$:

$$\begin{aligned} (d_{\Sigma}^{\pi_{n}}(\sigma) :=) \mathbb{P}(\Sigma = \sigma; \pi) &= \\ &= c^{-1} \sum_{t \le N - l(\sigma)} \mathbb{P}(S_{t} = s_{0}^{\sigma}, H_{t} = h_{0}^{\sigma}, G_{t} = g_{0}^{\sigma}, A_{t} = a_{0}^{\sigma}, \dots, S_{t+l(\sigma)} = s_{l(\sigma)}^{\sigma}; \pi) \\ &= c^{-1} \sum_{t \le N - l(\sigma)} \left(\prod_{i=1}^{l(\sigma)} \lambda(s_{i}^{\sigma} | a_{i-1}^{\sigma}, s_{i-1}^{\sigma}) \right) \cdot \left(\prod_{i=0}^{l(\sigma)-1} \pi(a_{i}^{\sigma} | \bar{s}_{i}^{\sigma}) \right) \cdot \mathbb{P}(S_{t} = s_{0}^{\sigma}, H_{t} = h_{0}^{\sigma}, G_{t} = g_{0}^{\sigma}; \pi) \\ &= c^{-1} \left(\prod_{i=1}^{l(\sigma)} \lambda(s_{i}^{\sigma} | a_{i-1}^{\sigma}, s_{i-1}^{\sigma}) \right) \cdot \left(\prod_{i=0}^{l(\sigma)-1} \pi(a_{i}^{\sigma} | \bar{s}_{i}^{\sigma}) \right) \cdot \sum_{t \le N - l(\sigma)} \mathbb{P}(S_{t} = s_{0}^{\sigma}, H_{t} = h_{0}^{\sigma}, G_{t} = g_{0}^{\sigma}; \pi) \end{aligned}$$

where

$$c := \sum_{\sigma \in \text{Seg}} \sum_{t \le N - l(\sigma)} \mathbb{P}(S_t = s_0^{\sigma}, H_t = h_0^{\sigma}, G_t = g_0^{\sigma}, A_t = a_0^{\sigma}, \dots, S_{t+l(\sigma)} = s_{l(\sigma)}^{\sigma}; \pi) > 0.$$

Since we constrained $N \geq 1$, $\operatorname{supp} \bar{\mu} \subset \bar{S}_T$ (which prevents degenerate cases), there must be a non-zero probability assigned to some segments leading to the full summation in the definition of c being positive. This means that c^{-1} and the segment distribution are defined correctly. Given the factorized form of the segment distribution we can conclude the following (through computing marginals and calculating conditional probability ratios):

$$\mathbb{P}(l(\Sigma) = k, S_0^{\Sigma} = s_0, H_0^{\Sigma} = h_0, G_0^{\Sigma} = g; \pi) = c^{-1} \sum_{t \le N-k} \mathbb{P}(S_t = s_0, H_t = h_0, G_t = g; \pi)$$

$$\mathbb{P}(A_0^{\Sigma} = a_0, S_1^{\Sigma} = s_1, A_1^{\Sigma} = a_1, \dots, S_k^{\Sigma} = s_k | S_0^{\Sigma} = s_0, H_0^{\Sigma} = h_0, G_0^{\Sigma} = g_0, l(\Sigma) = k; \pi)$$
$$= \left(\prod_{i=1}^k \lambda(s_i | a_{i-1}, s_{i-1})\right) \times \prod_{i=0}^{k-1} \pi(a_i | \bar{s}_i)$$

and $(\forall 0 \leq i \leq k)$:

$$\mathbb{P}(A_{i}^{\Sigma} = a_{i}|S_{i}^{\Sigma} = s_{i}, \dots, S_{0}^{\Sigma} = s_{0}, H_{0}^{\Sigma} = h_{0}, G_{0}^{\Sigma} = g, l(\Sigma) = k; \pi)$$

$$= \pi(a_{i}|s_{i}, h_{0} - i, g)$$

$$= \mathbb{P}(A_{i}^{\Sigma} = a_{i}|S_{i}^{\Sigma} = s_{i}, H_{0}^{\Sigma} = h_{0}, G_{0}^{\Sigma} = g; \pi), \qquad (2.5)$$

$$\mathbb{P}(S_{i}^{\Sigma} = s_{i}|S_{i-1}^{\Sigma} = s_{i-1}, A_{i-1}^{\Sigma} = a_{i-1}, \dots, S_{0}^{\Sigma} = s_{0}, H_{0}^{\Sigma} = h_{0}, G_{0}^{\Sigma} = g_{0}, l(\Sigma) = k; \pi)$$

$$= \lambda(s_{i}|s_{i-1}, a_{i-1})$$

$$= \mathbb{P}(S_{i}^{\Sigma} = s_{i}|S_{i-1}^{\Sigma} = s_{i-1}, A_{i-1}^{\Sigma} = a_{i-1})$$
(2.6)

After defining $H_i^{\Sigma} := H_0^{\Sigma} - i, G_i^{\Sigma} := G_0^{\Sigma}$, we can write $\mathbb{P}(A_i^{\Sigma} = a_i | S_i^{\Sigma} = s_i, H_i^{\Sigma} = h_i, G_i^{\Sigma} = g_i; \pi) = \pi(a_i | s_i, h_i, g_i).$

Bounding c: For further investigation it is useful to bound the normalizing constant c. We can write

$$0 < c = \sum_{\sigma \in \text{Seg}} \sum_{t \le N - l(\sigma)} \mathbb{P}(S_t = s_0^{\sigma}, H_t = h_0^{\sigma}, G_t = g_0^{\sigma}, A_t = a_0^{\sigma}, \dots, S_{t+l(\sigma)} = s_{l(\sigma)}^{\sigma}; \pi)$$

$$= \sum_{k=1}^N \sum_{t \le N - k} \sum_{\sigma: l(\sigma) = k} \mathbb{P}(S_t = s_0^{\sigma}, H_t = h_0^{\sigma}, G_t = g_0^{\sigma}, A_t = a_0^{\sigma}, \dots, S_{t+l(\sigma)} = s_{l(\sigma)}^{\sigma}; \pi)$$

$$\leq \sum_{k=1}^N \sum_{t \le N - k} 1 = \frac{N(N+1)}{2}.$$
(2.4)

Appendix B. Motivating Examples

We begin this section by demonstrating discontinuities of eUDRL generated quantities in the transition kernel given specific example of compatible families of MDPs. In order to demonstrate a discontinuity at a specific point $\lambda_0 \in (\Delta S)^{S \times A}$ we take limits with respect to two distinct rays meeting at λ_0 and showing that they disagree. Apart of providing plots depicting how the value of investigated quantity depends on the ray parameter we also provide exact computation of the limits. During the computation we will use the form of eUDRL recursion introduced by Štrupl et al. (2022a) ($\forall a \in \mathcal{A}, (s, h, g) \in \text{supp den}_{\lambda, \pi_n}$):

$$\pi_{n+1}(a|s,h,g) = \mathbb{P}_{\lambda}(A_0^{\Sigma} = a|S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_{l(\Sigma)}^{\Sigma}) = g; \pi_n)$$

$$= \frac{Q_A^{\pi_n,g}(s,h,a)\pi_{A,n}(a|s,h)}{\sum_{a \in \mathcal{A}} Q_A^{\pi_n,g}(s,h,a)\pi_{A,n}(a|s,h)},$$
(B.1)

where Q_A denotes an "average" Q-value and $\pi_{A,n}$ denotes an "average" policy

$$Q_{A}^{\pi_{n},g}(s,h,a) = \mathbb{P}_{\lambda}(\rho(S_{l(\Sigma)}^{\Sigma})) = g|A_{0}^{\Sigma} = a, S_{0}^{\Sigma} = s, l(\Sigma) = h; \pi_{n})$$

$$\pi_{A,n}(a|s,h) = \sum_{h' \ge h, g' \in \mathcal{G}} \pi_{n}(a|h',g',s) \mathbb{P}_{\lambda}(H_{0}^{\Sigma} = h', G_{0}^{\Sigma} = g'|S_{0}^{\Sigma} = s, l(\Sigma) = h; \pi_{n}).$$
(B.2)

The derivation of the above eUDRL recursion rewrite uses the same techniques as the proof of lemma 3.

At points λ lying on a boundary of $(\Delta S)^{S \times A}$ the continuity can break. Therefore, we cannot prove continuity on the whole boundary. This is illustrated in the following counter-example.

Example 1 (non-removable discontinuity of goal reaching objective at a boundary point) Consider an MDP $\mathcal{M} = (S, \mathcal{A}, \lambda, \mu, r)$ with three states $S = \{0, 1, 2\}$ and three actions $\mathcal{A} = \{0, 1, 2\}$. The state 0 is the only initial state. We consider its CE \overline{M} with maximum horizon N = 1, $\mathcal{G} := S$, $\rho := id_S$. The CE's initial distribution fixes the initial remaining horizon at $H_0 = 1$ and initial goals are distributed as $\mathbb{P}(G_0 = 0) = \mathbb{P}(G_0 = 2) = \frac{1}{2}$ (so $\mathbb{P}(G_0 = 1) = 0$). It follows that we are left with only two transient states (0, 1, 0), $(0, 1, 2) \in S \times \mathbb{N}_0 \times \mathcal{G}$ differing just by the goal component which are actually visited. Thus it suffices to provide policies and values only for these states. Since we have N = 1 and the only initial state $S_0 = 0$ it suffice to give the transition kernel only from this state. We split the computation into three parts and label them by letters A, B and C for later reference.

A First we define the parametric transition kernel (with parameter $\alpha \in [0,1]$)

$$\begin{array}{c|c|c} \lambda_{\alpha}(g|a) & g=0 & g=1 & g=2\\ \hline a=0 & 1-\alpha & \frac{\alpha}{4} & \frac{3\alpha}{4}\\ a=1 & \frac{3\alpha}{4} & 1-\alpha & \frac{\alpha}{4}\\ a=2 & \frac{1}{2} & \frac{1}{2} & 0 \end{array}$$

Which we identify with the matrix

$$\begin{bmatrix} \lambda_{\alpha}(g|a) \end{bmatrix} = \begin{pmatrix} 1-\alpha & \frac{\alpha}{4} & \frac{3\alpha}{4} \\ \frac{3\alpha}{4} & 1-\alpha & \frac{\alpha}{4} \\ \frac{1}{2} & \frac{1}{2} & 0 \end{pmatrix} \xrightarrow{\alpha \to 0+} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ \frac{1}{2} & \frac{1}{2} & 0 \end{pmatrix} = \begin{bmatrix} \lambda_{0+}(g|a) \end{bmatrix}.$$

We will be interested how values (policies) behave at and around the point $\alpha = 0$. For computing the eUDRL policies we will use the recursion (B.1). We will note argument names inside kernels and value functions (in case they were substituted by actual values) so it is not necessary to always look for how we fixed individual argument positions.⁹ Using (B.2) we get $Q_A^{\pi_n,g}(s=0, h=1, a) = \lambda(g|a, s=0)$. Since the condition $\operatorname{supp} \overline{\mu} \subset \overline{S}_T$ and fixed horizon 1 the segments coincide with trajectories. Therefore, the distribution of the first extended state of a segment is the same as the initial distribution $\overline{\mu}$: $\mathbb{P}(H_0^{\Sigma} = 1, G_0^{\Sigma} =$ $g'|S_0^{\Sigma} = 0, l(\Sigma) = 1; \pi_n) = \mathbb{P}(G_0 = g')$. Thus we get the following relation for the "average" policy: $\pi_{A,n}(a|s=0, h=1) = \sum_{g' \in \mathcal{G}} \pi_n(a|s=0, h'=1, g')\mathbb{P}(G_0 = g')$. The recursion (B.1) can then be rewitten

$$egin{aligned} \pi_{n+1}(a|s=0,h=1,g) \propto Q_A^{\pi_n,g}(s=0,h=1,a)\pi_{A,n}(a|s=0,h=1)\ &=\lambda(g|a,s=0)\pi_{A,n}(a|s=0,h=1). \end{aligned}$$

Assuming we start with uniform initial condition π_0 we obtain also uniform $\pi_{A,0} = \frac{1}{3}$ leaving $\pi_1(a|s=0, h=1, g) \propto \lambda(g|a, s=0) \frac{1}{3} \propto \lambda(g|a, s=0)$. Thus we get

$$\begin{bmatrix} \pi_{1,\alpha}(a|g) \end{bmatrix} = \begin{pmatrix} \frac{1-\alpha}{1-\alpha+\frac{3\alpha}{4}+\frac{1}{2}} & \frac{\alpha}{4}+1-\alpha+\frac{1}{2} & \frac{3}{4} \\ \frac{3\alpha}{4}+\frac{1}{2} & \frac{1-\alpha}{4}+1-\alpha+\frac{1}{2} & \frac{1}{4} \\ \frac{1}{2} & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2}-\alpha+\frac{3\alpha}{4}+\frac{1}{2} & \frac{\alpha}{4}+1-\alpha+\frac{1}{2} & 0 \end{pmatrix} \xrightarrow{\alpha \to 0+} \begin{pmatrix} \frac{2}{3} & 0 & \frac{3}{4} \\ 0 & \frac{2}{3} & \frac{1}{4} \\ \frac{1}{3} & \frac{1}{3} & 0 \end{pmatrix} = [\pi_{1,0+}(a|g)].$$

This gives us

$$\begin{bmatrix} \pi_{A,1,0+} \end{bmatrix} = \begin{bmatrix} \pi_{1,0+}(a|g) \end{bmatrix} \begin{bmatrix} \mathbb{P}(G_0 = g) \end{bmatrix} = \begin{bmatrix} \pi_{1,0+}(a|g) \end{bmatrix} \begin{pmatrix} \frac{1}{2} \\ 0 \\ \frac{1}{2} \end{pmatrix} = \begin{pmatrix} \frac{17}{24} \\ \frac{1}{8} \\ \frac{1}{6} \end{pmatrix}$$

We deduce from $\pi_2(a|s=0, h=1, g) \propto \lambda(g|a, s=0)\pi_{A,1}(a)$ that

....

$$\lim_{\alpha \to 0+} \pi_2(a|s=0, h=1, g) = \frac{\lim_{\alpha \to 0+} \lambda(g|a, s=0) \cdot \lim_{\alpha \to 0+} \pi_{A,1}(a)}{\sum_{a \in \mathcal{A}} \lim_{\alpha \to 0+} \lambda(g|a, s=0) \cdot \lim_{\alpha \to 0+} \pi_{A,1}(a)}$$

provided all limits exists and division makes sense. Fortunately we will be interested just in the column for g = 0 and g = 1

$$[\pi_{2,0+}(a|g=0)] = \begin{pmatrix} \frac{\frac{1}{24}1}{\frac{17}{24}1 + \frac{1}{8}0 + \frac{1}{6}\frac{1}{2}} \\ \frac{\frac{1}{8}0}{\frac{1}{24}1 + \frac{1}{8}0 + \frac{1}{6}\frac{1}{2}} \\ \frac{\frac{1}{6}\frac{1}{2}}{\frac{1}{7}\frac{1}{24}1 + \frac{1}{8}0 + \frac{1}{6}\frac{1}{2}} \end{pmatrix} = \begin{pmatrix} \frac{17}{\frac{19}{24}} \\ 0 \\ \frac{1}{\frac{19}{24}} \\ \frac{19}{24} \end{pmatrix} = \begin{pmatrix} \frac{17}{19} \\ 0 \\ \frac{2}{19} \end{pmatrix}, \quad [\pi_{2,0+}(a|g=1)] = \begin{pmatrix} 0 \\ \frac{3}{5} \\ \frac{2}{5} \\ \frac{2}{5} \end{pmatrix}.$$

Finally using

$$V_{0+}^{\pi_2}(s=0,h=1,g) = \sum_{a\in\mathcal{A}} \lambda_{0+}(g|s=0,a)\pi_{2,0+}(a|g)$$

^{9.} This is only to improve readability of the example by helping the reader to keep track of substitutions. We use different color (gray) for these extra notes.

and

$$J_{\lambda,\pi_2,0+} = \sum_{g \in \mathcal{G}} V_{0+}^{\pi_2}(s=0, h=1, g) \bar{\mu}(s=0, h=1, g) = \sum_{g \in \mathcal{G}} V_{0+}^{\pi_2}(s=0, h=1, g) \mathbb{P}(G_0 = g)$$

the values $V_{0+}^{\pi_2}(s=0, h=1, g=0) = \frac{18}{19}$, $V_{0+}^{\pi_2}(s=0, h=1, g=2) = 0$ and the goal reaching objective $J_{\lambda,\pi_2,0+} = \frac{9}{19}$ can be computed.

B Now we aim to re-compute everything for $\alpha = 0$. Since λ is continuous in λ we get $\lambda_0 = \lambda_{0+}$. Further since $\lambda_0(g=2|\cdot) = 0$ we get $\pi_{1,0}(\cdot|g=2) = \frac{1}{\mathcal{A}} = \frac{1}{3}$. Thus

$$\begin{bmatrix} \pi_{1,0}(a|g) \end{bmatrix} = \begin{pmatrix} \frac{2}{3} & 0 & \frac{1}{3} \\ 0 & \frac{2}{3} & \frac{1}{3} \\ \frac{1}{3} & \frac{1}{3} & \frac{1}{3} \end{pmatrix}, \quad \begin{bmatrix} \pi_{A,1,0} \end{bmatrix} = \begin{bmatrix} \pi_{1,0}(a|g) \end{bmatrix} \begin{pmatrix} \frac{1}{2} \\ 0 \\ \frac{1}{2} \end{pmatrix} = \begin{pmatrix} \frac{1}{2} \\ \frac{1}{6} \\ \frac{1}{3} \end{pmatrix}$$

$$\begin{bmatrix} \pi_{2,0}(a|g=0) \end{bmatrix} = \begin{bmatrix} \frac{\lambda_0(a|g=0))\pi_{A,1,0}(a)}{\sum_{a \in \mathcal{A}} \lambda_0(a|g=0))\pi_{A,1,0}(a)} \end{bmatrix} = \begin{pmatrix} \frac{\frac{1}{2}1}{\frac{1}{2}1 + \frac{1}{6}0 + \frac{1}{3}\frac{1}{2}} \\ \frac{\frac{1}{6}0}{\frac{1}{2}1 + \frac{1}{6}0 + \frac{1}{3}\frac{1}{2}} \\ \frac{1}{2}1 + \frac{1}{6}0 + \frac{1}{3}\frac{1}{2}} \end{pmatrix} = \begin{pmatrix} \frac{3}{4} \\ 0 \\ \frac{1}{4} \end{pmatrix},$$

similarly

$$[\pi_{2,0}(a|g=1)] = \begin{pmatrix} 0\\ \frac{1}{2}\\ \frac{1}{2} \end{pmatrix}$$

Finally we get

$$V_0^{\pi_2}(s=0,h=1,g=0) = \frac{7}{8}, \quad V_0^{\pi_2}(s=0,h=1,g=2) = 0, \quad J_{\lambda,\pi_2,0} = \frac{7}{16},$$

which means that we have a discontinuity at $\alpha = 0$: a boundary point.

C Moreover, this discontinuity could not be removed as can be seen by computing the limit with respect to another ray, e.g.,

$$\begin{bmatrix} \lambda_{\alpha}'(g|a) \end{bmatrix} = \begin{pmatrix} 1-\alpha & \frac{3\alpha}{4} & \frac{\alpha}{4} \\ \frac{\alpha}{4} & 1-\alpha & \frac{3\alpha}{4} \\ \frac{1}{2} & \frac{1}{2} & 0 \end{pmatrix}.$$

Notice that $\lambda'_{0+} = \lambda'_0 = \lambda_{0+} = \lambda_0$. Following the same procedure as in part A we obtain the following results:

$$\begin{bmatrix} \pi'_{1,0+}(a|g) \end{bmatrix} = \begin{pmatrix} \frac{3}{2} & 0 & \frac{1}{4} \\ 0 & \frac{2}{3} & \frac{3}{4} \\ \frac{1}{3} & \frac{1}{3} & 0 \end{pmatrix}, \quad \begin{bmatrix} \pi'_{A,1,0+}(a) \end{bmatrix} = \begin{pmatrix} \frac{11}{24} \\ \frac{3}{8} \\ \frac{1}{6} \end{pmatrix},$$

$$\begin{bmatrix} \pi'_{2,0+}(a|g=0) \end{bmatrix} = \begin{pmatrix} \frac{11}{13} \\ 0 \\ \frac{2}{13} \end{pmatrix}, \quad \begin{bmatrix} \pi'_{2,0+}(a|g=1) \end{bmatrix} = \begin{pmatrix} 0 \\ \frac{9}{11} \\ \frac{2}{11} \end{pmatrix},$$

$$V_{0+}^{'\pi_{2}}(g=0) = \frac{12}{13}, \quad V_{0+}^{'\pi_{2}}(g=2) = 0, \quad J_{0+}^{'\pi_{2}} = \frac{6}{13}.$$

Discussing alternative definitions of eUDRL recursion outside of $\operatorname{supp} \operatorname{den}_{\lambda,\pi_n}$: Since the computation of the limits in A and C does not depend on the way we defined the eUDRL recursion outside of $\operatorname{supp} \operatorname{den}_{\lambda,\pi_n}$ in (4.1), the presented non-removable discontinuities of π_2 and J^{π_2} at λ_0 cannot vanish by employing any alternative definition of the recursion outside of $\operatorname{supp} \operatorname{den}_{\lambda,\pi_n}$. This means there is no suitable alternative definition of eUDRL recursion (outside of $\operatorname{supp} \operatorname{den}_{\lambda,\pi_n}$) which would remove the present discontinuity, e.g. in the goal-reaching objective. Therefore, the concrete definition of the recursion in (4.1) outside of $\operatorname{supp} \operatorname{den}_{\lambda,\pi_n}$ does not matter much from this point of view (the discontinuities will always be there). The same reasoning applies also for discontinuities presented in the next example.

The next example illustrates non-removable discontinuity of eUDRL generated policies (for $n \geq 2$) at deterministic points. However, while we will see the limits with respect to different rays disagree, they are all optimal policies. This could motivate us to prove relative continuity (some weaker notion of continuity) for these policies, as was successfully achieved in theorem 16.

Example 2 (non-removable discontinuity of policy at deterministic point) Let the MDP \mathcal{M} be the same as in example 1 except that we vary the parametric transition kernels to illustrate the behavior around (and at) the specific deterministic transition kernel. Further, we change the distribution of the initial goals to the uniform distribution, i.e., $\mathbb{P}(G_0 = 0) = \mathbb{P}(G_0 = 1) = \mathbb{P}(G_0 = 2) = \frac{1}{3}$. Since the computation proceeds as in the example 1 we just introduce the new kernels and summarize the results.

A

$$\begin{bmatrix} \lambda_{\alpha}(g|a) \end{bmatrix} = \begin{pmatrix} 1-\alpha & \alpha & 0 \\ 0 & 1-\alpha & \alpha \\ \alpha & 1-\alpha & 0 \end{pmatrix} \xrightarrow{\alpha \to 0+} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} = \begin{bmatrix} \lambda_{0+}(g|a) \end{bmatrix} (= \begin{bmatrix} \lambda_0(g|a) \end{bmatrix}).$$

Notice continuity at $\alpha = 0$ ($\lambda_{0+} = \lambda_0$). The results follow:

$$\begin{split} [\pi_{1,0+}(a|g)] &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{2} & 1 \\ 0 & \frac{1}{2} & 0 \end{pmatrix}, \quad [\pi_{A,1,0+}(a)] = \begin{pmatrix} \frac{1}{3} \\ \frac{1}{2} \\ \frac{1}{6} \end{pmatrix}, \\ [\pi_{2,0+}(a|g=0)] &= \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad [\pi_{2,0+}(a|g=1)] = \begin{pmatrix} 0 \\ \frac{3}{4} \\ \frac{1}{4} \end{pmatrix}, \\ V_{0+}^{\pi_2}(g=0) &= 1, \quad V_{0+}^{\pi_2}(g=1) = 1, \quad V_{0+}^{\pi_2}(g=2) = 0, \quad J_{0+}^{\pi_2} = \frac{2}{3}. \end{split}$$

 \boldsymbol{B}

$$\begin{aligned} \pi_{1,0}(a|g)] &= \begin{pmatrix} 1 & 0 & \frac{1}{3} \\ 0 & \frac{1}{2} & \frac{1}{3} \\ 0 & \frac{1}{2} & \frac{1}{3} \end{pmatrix}, \quad [\pi_{A,1,0}(a)] = \begin{pmatrix} \frac{4}{9} \\ \frac{5}{18} \\ \frac{5}{18} \end{pmatrix}, \quad [\pi_{2,0}(a|g)] = \begin{pmatrix} 1 & 0 & \frac{1}{3} \\ 0 & \frac{1}{2} & \frac{1}{3} \\ 0 & \frac{1}{2} & \frac{1}{3} \end{pmatrix}, \\ V_0^{\pi_2}(g=0) &= 1, \quad V_0^{\pi_2}(g=1) = 1, \quad V_0^{\pi_2}(g=2) = 0, \quad J_0^{\pi_2} = \frac{2}{3}. \end{aligned}$$

 \boldsymbol{C}

$$\begin{bmatrix} \lambda'_{1,\alpha}(a|g) \end{bmatrix} := \begin{pmatrix} 1-\alpha & 0 & \alpha \\ \alpha & 1-\alpha & 0 \\ 0 & 1-\alpha & \alpha \end{pmatrix}.$$

Notice that $\lambda'_{1,0+} = \lambda'_{1,0} = \lambda_{1,0} = \lambda_{1,0+}$.

$$\begin{bmatrix} \pi'_{1,0+}(a|g) \end{bmatrix} = \begin{pmatrix} 1 & 0 & \frac{1}{2} \\ 0 & \frac{1}{2} & 0 \\ 0 & \frac{1}{2} & \frac{1}{2} \end{pmatrix}, \quad \begin{bmatrix} \pi'_{A,1,0+}(a) \end{bmatrix} = \begin{pmatrix} \frac{1}{2} \\ \frac{1}{6} \\ \frac{1}{3} \end{pmatrix}, \quad \begin{bmatrix} \pi'_{2,0+}(a|g) \end{bmatrix} = \begin{pmatrix} 1 & 0 & \frac{3}{5} \\ 0 & \frac{1}{3} & 0 \\ 0 & \frac{2}{3} & \frac{2}{5} \end{pmatrix},$$

$$V_{0+}^{'\pi_{2}}(g=0) = 1, \quad V_{0+}^{'\pi_{2}}(g=1) = 1, \quad V_{0+}^{'\pi_{2}}(g=2) = 0, \quad J_{0+}^{'\pi_{2}} = \frac{2}{3}.$$

The dependence of the goal reaching objective and the specific π_2 component on α is depicted in figure 2 (for both examples 1 and 2). While policies are clearly discontinuous at $\alpha = 0$ in both examples, we see continuity of a goal-reaching objective (on A and C rays) at the deterministic kernel (example 2). This hints at a possible continuity of goal reaching objectives and relative continuity of policies at deterministic points (kernels), as was later proved in this paper. Unfortunately, we cannot hope for the similar (general) result in a case of non-deterministic boundary points (kernels) which was clearly demonstrated in example 1 and figures 2c and 2d. In addition to the examples above, we include a discussion describing the causes that lead to discontinuities of eUDRL-generated quantities at the end of section C.

In the following example, we want to illustrate (relative) continuity of eUDRL generated policies and goal reaching objectives for a finite number of iterations, which was proved in section 6.

Example 3 (example featuring random walk on \mathbb{Z}_3) In this example, we consider an MDP $\mathcal{M} = (\mathcal{S}, \mathcal{A}, \lambda, \mu, r)$ with state space $\mathcal{S} = \mathbb{Z}_3$ and action space $\mathcal{A} = \{0, 1\}$. The transition kernel depends on the parameter $\alpha \in [0, 1]$ that regulates the stochasticity. When action 0 is selected, the MDP stays in the current state $s \in \mathcal{S}$ with probability $1 - \alpha$ and transits to the state s + 1 with probability α . When action 1 is selected, the MDP stays in the current state $s \in \mathcal{S}$ with probability $1 - \alpha$, i.e., the states $w \in \mathbb{S}$ with probability α and transits to the state s + 1 with probability α and transits to the state s + 1 with probability $1 - \alpha$, i.e., the states we are transiting stay the same; just the probabilities are swapped. For both actions, the transition kernels are essentially different random walks on $\mathcal{S} = \mathbb{Z}_3$. We consider \mathcal{M} 's CE $\overline{\mathcal{M}}$ with remaining horizon N = 8, $\mathcal{G} := \mathcal{S}$, $\rho := \mathrm{id}_{\mathcal{S}}$. The CE's initial distribution is a uniform distribution over all extended states.

The figures 3 and 4 show the dependence of $\min_{\bar{s}\in\bar{S}_{\lambda_0}} \pi_{n,\lambda}(\mathcal{O}(\bar{s}|\bar{s}))$ and the goal reaching objective respectively on iteration n. Continuity in the transition kernel is hinted at by these figures as the plots for different $\delta := \|\lambda - \lambda_0\|_1$ approach the plot for $\delta = 0$ (i.e., λ_0 plot) smoothly as $\delta \to 0$. Plots in this particular example are interesting as they exhibit significant oscillations in the first few iterations. A more detailed plot for the oscillating goal reaching objective (for one specific λ and one specific initial condition) is shown in 4b. Here we see that the best results are obtained for either iteration 2 or 3. Furthermore, the performance of the goal reaching objective deteriorates when eUDRL is continued. The general non-monotonicity of a goal reaching objective could raise the



(a) Dependency on iteration. (b) Depen

(b) Dependency on distance to determin. kernel.

Figure 13: Illustration of behavior of $\min_{\bar{s}\in\bar{S}_{\lambda_0}} \pi_n(\mathcal{O}(\bar{s})|s)$ for eUDRL recursion in a example 4. Plot (a) shows the dependency on the iteration n for varying distances to a deterministic kernel highlighted by different colors and varying initial policy. Plot (b) contains the same information as (a) but it is organized to reveal the dependency on δ , where varying numbers of iteration are highlighted by different colors.

question of whether or not it makes sense to continue eUDRL iterations in order to reach some "optimal" behavior. These questions motivate our theoretical investigation of the asymptotic properties of eUDRL iteration in section 7.

In order to compare bounds derived for all special cases from section 7, we include a simple bandit example below (which complies with both of the conditions $(\forall \bar{s} \in \bar{S}_{\lambda_0})$: $|\mathcal{O}(\bar{s})| = 1$ and $\bar{S}_{\lambda_0} \subset \operatorname{supp} \bar{\mu}$).

Example 4 (a simple bandit) We consider an MDP $\mathcal{M} = (S, \mathcal{A}, \lambda, \mu, r)$ with a state space $S = \{0, 1\}$ and an action space $\mathcal{A} = \{0, 1\}$. The transition kernel depends on parameter $\alpha \in [0, 1]$ which regulates stochasticity. When action 0 is selected, the MDP stays in the current state $s \in S$ with probability $1 - \alpha$ and transitions to the other state with probability α . When action 1 is selected, the MDP stays in the current state $s \in S$ with probability $1 - \alpha$ and transitions to the other state with probability α and transitions to the other state with probability $1 - \alpha$. We consider \mathcal{M} 's $CE \overline{\mathcal{M}}$ with remaining horizon N = 1, $\mathcal{G} := S$, $\rho := id_S$. The CE's initial distribution is a uniform distribution over states with the original MDP state component s = 0. Since the initial horizon can be just 1, there are just two initial states in the CE which differ in the goal component: (0, 1, 0) and (0, 1, 1).

For detailed plots concerning the bandit example, see figures 6, 9, 11 and 13.

Next, we include an example featuring a tiny grid world domain. In this example, we wanted to demonstrate x^* bound on eUDRL policy accumulation points introduced in

section 7. In order to satisfy the assumption $S_{\lambda_0} \subset \operatorname{supp} \bar{\mu}$ required by this bound, we chosen the uniform initial distribution for the associated CE.

Example 5 (a tiny grid world domain) The domain is described by a 3 by 3 map (see figure 7d). The gray color depicts a wall. Other positions correspond to the MDP states. Thus, we have a domain $\mathcal{M} = (S, \mathcal{A}, \lambda, \mu, r)$ with 8 states (|S| = 8) and 4 actions $(\mathcal{A} = \{$ "right", "left", "up", "down" $\}$). The transition kernel is parameterized by the parameter $\alpha \in [0, 1]$, regulating the distance from a deterministic kernel (for $\alpha = 0$ the kernel is deterministic and we denote it λ_0). In case of $\alpha = 0$, the kernel transits as indicated by the name of an action if possible. If it is not possible (one would move out of the grid or into a wall), the MDP stays in its current state. In case of non-deterministic kernels ($\alpha > 0$), the exact kernel depends on "available states" which is union (over all actions) of all states the deterministic to s' and "available states" $\{s'\} \neq \emptyset$, the kernel transits to s' with probability α is uniformly distributed between all states in "available states" $\{s'\} = \emptyset$, then the kernel transits to s' with probability 1. We consider \mathcal{M} 's $CE \ mathcal{M}$ with the remaining horizon N = 4, $\mathcal{G} := S$, $\rho := id_S$. The CE's initial distribution is the uniform distribution over all CE states.

The bunch of example accumulation points, together with the lower bound $x^*(\gamma_N)$ on the kernel distance $\delta := \|\lambda - \lambda_0\|_1$, is plotted in the figure 7a. The map of the grid world domain is shown in figure 7d.

Further, we give an example which features the x_u bound on the eUDRL policy accumulation points introduced in section 7 together with ODT recursion. In order to do this, we followed the approach explained in the background section which assumes fixed horizon scenario with the reward at the end. This approach employs first a state space transformation (which consists of forming a K-tuples of states) and then builds CE on top of the new state space. The fixed horizon and the optimal policy uniqueness (just on \bar{S}_{λ_0}) requirements are handled by a convenient choice of the CE initial distribution and the goal map. Finally, we utilized the fact that ODT recursion becomes just eUDRL recursion (on the described CE with transformed states) on the restricted segment space Seg^{trail}.

Example 6 (a tiny grid world domain with ODT recursion) The underlying MDP is the same as in the previous example 5 with just the initial distribution having restricted support (we will comment on it more precisely when detailing the associated CE). Then we form a new MDP by considering K-tuples of the underlying MDP states with K = 3 and extend the kernel accordingly. Over this new MDP, we form the CE. The CE's maximal horizon was set to N = 4 and the goal space was defined $\mathcal{G} := \{0,1\}$ with the goal map mapping K-tuples finishing with position (0,2) to 1 and others to 0. We pick just one initial state for this CE corresponding to position (2,2) with a remaining horizon 4 and goal 1 (see the grid world map on 10d). This setting ensured the fixed horizon requirement (horizon was fixed to 4 by the choice of initial state) and the requirement for the uniqueness of the optimal policy on \bar{S}_{λ_0} . Now, note that the position (2,2) and the goal 1 (the position (0,2)) of length N = 4 going through these states. Therefore, \bar{S}_{λ_0} spans over K-tuples finishing

with the remaining positions. Moreover, since there is only one trajectory connecting (2,2)with (0,2) in 4 steps, there is only one remaining horizon available for each position in \bar{S}_{λ_0} . This leads to the uniqueness of the optimal policy on \bar{S}_{λ_0} . So we can employ the x_u bound for this domain.

The example's accumulation points of $\min_{\bar{s}\in\bar{S}_{\lambda_0}} \pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s})$ of ODT recursion together with the lower bound $x_u(\delta)$ are shown in figure 10a.

Finally, we give an example of a grid word which does not fall in the special cases investigated in section 7. On this example we will showcase ODT recursion with regularization (using uniform distribution). This recursion (in the specific setting of the example) can be understood as a special case of ϵ -eUDRL which was investigated in section 8.

Example 7 (a tiny grid world domain with ODT recursion allowing for a non-deterministic optimal policy on \bar{S}_{λ_0}) This example is exactly the same as the previous example 6 except that we change the initial state position to (2,0). This change allows for non-determinism of an optimal policy on \bar{S}_{λ_0} (see the map in the figure 12b). Thus, both special conditions we have introduced for bounding the accumulation points of eUDRL (ODT without regularization) recursion fail. However, we can still bound regularized eUDRL (ODT) recursions.

The example accumulation points of $\min_{\bar{s}\in\bar{\mathcal{S}}_{\lambda_0}}\pi_{n,\lambda}(\mathcal{O}(\bar{s})|\bar{s})$ of the ODT recursion together with the ϵ -eUDRL based lower bound $\min_M x^*(\gamma_N, \epsilon, \delta)$ are plotted in figure 12a.

Appendix C. Continuity at Interior Kernels for a Finite Number of Iterations

In this section, we aim to prove continuity of eUDRL-generated policies and related quantities at the interior points of the set of all transition kernels $(\Delta S)^{S \times A}$. Note that an interior point $\lambda \in ((\Delta S)^{S \times A})^{\circ} = ((\Delta S^{\circ})^{S \times A})$ can be equivalently characterized as a transition kernel satisfying $\lambda > 0$. The following lemma, which could be understood as the interior point counterpart of the lemma 6, discusses the stability properties of the set supp den_{λ,π} subject to changes in the transition kernel $\lambda > 0$ and the policy π .

Lemma 37 (support properties) Let $\{\mathcal{M}_{\lambda} : \lambda \in (\Delta S)^{S \times A}\}$ and $\{\overline{\mathcal{M}}_{\lambda} : \lambda \in (\Delta S)^{S \times A}\}$ be compatible families of MDPs. Let $\lambda, \lambda' \in (\Delta S)^{S \times A}$ be transition kernels with $\lambda, \lambda' > 0$ and $\pi, \pi' \in (\Delta A)^S$ be policies. Then it holds that

 $\operatorname{supp} \operatorname{den}_{\lambda,\pi} = \operatorname{supp} \operatorname{den}_{\lambda',\pi'}.$

Proof Assume $\bar{s} = (s, h, g) \in \text{supp den}_{\lambda,\pi}$. This means that there exist a trajectory τ and $0 \leq t \leq N$ with $\mathbb{P}_{\lambda,\pi}(\mathcal{T} = \tau) > 0$, $l(\tau) \geq t + h$, $s_t^{\tau} = s$, $\rho(s_{t+h}^{\tau}) = g$. Let us construct a new trajectory τ' from τ by altering actions so that $a_{t'}^{\tau'} \in \text{supp } \pi'(\cdot|\bar{s}_{t'}^{\tau})$ for all $0 \leq t' \leq N$. Since $\lambda' > 0$, we also get $\mathbb{P}_{\lambda',\pi'}(\mathcal{T} = \tau') > 0$ causing $\bar{s} \in \text{supp den}_{\lambda',\pi'}$.

Lemma 37 asserts that the set supp $\text{den}_{\lambda,\pi}$ is constant over the interior of the set of all transition kernels and all policies. This is in striking difference with the behavior of supp $\text{den}_{\lambda,\pi}$ on the neighborhood of a deterministic kernel (or generally a kernel located at the boundary of the set of all transition kernels), as discussed in lemma 6 and example 2 (or example 1 for boundary points), where on every neighborhood the supp $\text{den}_{\lambda,\pi}$ could change abruptly. In the following lemma we will exploit this property to deliver a much simpler continuity proof than in the case of a deterministic kernel (cf. theorem 16).

Lemma 38 (Continuity of eUDRL policies and values at interior points) Let $\{\mathcal{M}_{\lambda} : \lambda \in (\Delta S)^{S \times \mathcal{A}}\}$ and $\{\overline{\mathcal{M}}_{\lambda} : \lambda \in (\Delta S)^{S \times \mathcal{A}}\}$ be compatible families of MDPs. Let $(\pi_{n,\lambda}), \pi_0 \in (\Delta \mathcal{A})^S$ be eUDRL generated sequence of policies. Let $\lambda_0 > 0$ be a transition kernel, i.e., λ_0 is an interior point $\lambda_0 \in ((\Delta S)^{S \times \mathcal{A}})^\circ$. Then for all $n \geq 0$ it holds that:

- 1. For all $\bar{s} = (s, h, g) \in \bar{S}_T$ the policy $\pi_{n+1,\lambda}(\cdot|\bar{s})$ is continuous in λ at λ_0 .
- 2. For all $\bar{s} = (s, h, g) \in \bar{S}_T$ the values $V_{\lambda}^{\pi_n}(\bar{s})$ and $Q_{\lambda}^{\pi_n}(\bar{s}, \cdot)$ are continuous in λ at the point λ_0 . In addition, the goal reaching objective $J_{\lambda}^{\pi_n}$ is continuous in λ at the point λ_0 . (Here we view $V_{\lambda}^{\pi_n}$, $Q_{\lambda}^{\pi_n}$, and $J_{\lambda}^{\pi_n}$ as functions of a single parameter λ resulting from composition with $\pi_{n,\lambda}$.)

Proof Since the interior $((\Delta S)^{S \times A})^{\circ}$ is an open set, we can fix $\delta > 0$ so that $U_{\delta}(\lambda_0)$ is contained in the interior.

1. The proof proceeds by induction on n.

Base case n = 0: π_0 is continuous in λ as it is constant in λ .

Induction: Assume the statement holds for $n \ge 0$, we aim to prove it for n+1. Let us fix $\bar{s} \in \bar{S}_T$. First, assume $\bar{s} \notin \text{supp den}_{\lambda_0,\pi_n}$ then from lemma 37 it holds that $\bar{s} \notin \text{supp den}_{\lambda,\pi_n}$

for all $\lambda \in U_{\delta}(\lambda_0)$. This in turn means that, as defined by (4.1), $\pi_{n+1,\lambda}(\cdot|\bar{s}) = \frac{1}{|\mathcal{A}|}$ for all $\lambda \in U_{\delta}(\lambda_0)$, and therefore $\pi_{n+1,\lambda}(\cdot|\bar{s})$ is continuous in λ at point λ_0 . Finally, assume $\bar{s} \in \operatorname{supp} \operatorname{den}_{\lambda_0,\pi_n}$. Then it holds that $\bar{s} \in \operatorname{supp} \operatorname{den}_{\lambda,\pi_n}$ for all $\lambda \in U_{\delta}(\lambda_0)$. This means that $\pi_{n+1,\lambda}(\cdot|\bar{s})$ is defined as

$$\pi_{n+1,\lambda}(a|\bar{s}) = \frac{\operatorname{num}_{\lambda,\pi_n}(\bar{s},a)}{\operatorname{den}_{\lambda,\pi_n}(\bar{s})}$$

for all $\lambda \in U_{\delta}(\lambda_0)$. From point 4 of lemma 12, both $\operatorname{num}_{\lambda,\pi_n}(\bar{s}, \cdot)$ and $\operatorname{den}_{\lambda,\pi_n}(\bar{s})$ are continuous in (λ, π_n) as segment distribution marginals. Further, from the induction assumption, $\pi_{n,\lambda}(\cdot|\bar{s}')$ is continuous in λ at λ_0 for all $\bar{s}' \in \bar{S}_T$. Therefore both $\operatorname{num}_{\lambda,\pi_n}(\bar{s}, \cdot)$ and $\operatorname{den}_{\lambda,\pi_n}(\bar{s})$, when compounded with $\pi_{n,\lambda}$, are continuous in λ at λ_0 . This together with $\operatorname{den}_{\lambda_0,\pi_n}(\bar{s}) > 0$ implies that $\pi_{n+1,\lambda}(a|\bar{s})$ is continuous in λ at λ_0 .

2. Assume n > 0. First, we prove the statement about values. Let us fix $\bar{s} \in \bar{S}_T$. The values $V_{\lambda}^{\pi_n}(\bar{s})$ and $Q_{\lambda}^{\pi_n}(\bar{s}, a)$ are both continuous in (λ, π_n) from point 1 of lemma 12. Since $\pi_{n,\lambda}$ are continuous in λ at λ_0 from point 1. both $V_{\lambda}^{\pi_n}(\bar{s})$ and $Q_{\lambda}^{\pi_n}(\bar{s}, a)$, when compounded with $\pi_{n,\lambda}$, are continuous in λ at λ_0 . Finally, the continuity of the goal-reaching objective

$$J_{\lambda}^{\pi_n} = \sum_{\bar{s}\in\bar{S}_T} V_{\lambda}^{\pi_n}(\bar{s})\bar{\mu}(\bar{s})$$

in λ at λ_0 follows from the continuity of $V_{\lambda}^{\pi_n}$.

When and why discontinuities arise at boundary points: Now, when we introduced the proof of continuity of eUDRL-generated quantities at the interior points (of $\Delta S^{S \times A}$ the set of all transition kernels), we are at the right position to investigate when this type of proof would break if we replace an interior point with a boundary point. In order to discuss general boundary points (not only deterministic ones) we will need the following two generalizations¹⁰. First, we need to generalize the notion of \bar{S}_{λ_0} for general transition kernels $\lambda_0 \in \Delta S^{S \times A}$ (allowing also for non-deterministic λ_0) which is trivial. Second, we will need to generalize lemma 6 to a general kernel λ_0 .

Lemma 39 (Stability of \bar{S}_{λ_0}) The conclusions of lemma 6 remain valid under following changes: We replace "a deterministic transition kernel λ_0 " with "a transition kernel λ_0 ". We replace quantification in points 1., 2. and 3. "for all $n \ge 0$ and all $\lambda \in U_2(\lambda_0)$ " with "There exists $2 > \delta > 0$ such that for all $n \ge 0$ and all $\lambda \in U_{\delta}(\lambda_0)$ ".

Proof The proof remains the same as in the original lemma 6 except that we have to find a convenient $\delta > 0$ for points 1., 2. and 3., i.e., we have to find $\delta > 0$ so that the following variation of (4.2) holds

$$(\forall \lambda \in U_{\delta}(\lambda_0)) : \operatorname{supp} \lambda_0 \subset \operatorname{supp} \lambda.$$

But this is just a consequence of the continuity of the identity map $\lambda \mapsto \lambda$.

^{10.} Note that these can be omitted if one is interested solely in deterministic points.

Now we will follow the proof of lemma 38 in the first two iterations of eUDRL while assuming λ_0 to be a boundary point and see when it fails. First, we fix $\delta > 0$ so that conclusions of points 1., 2. and 3. of lemma 39 hold.

Iteration n = 0: Here, π_0 is assumed constant in λ and therefore continuous in λ at λ_0 . Similarly, from point 1. of lemma 12, the values $V_{\lambda}^{\pi_0}(\bar{s})$ and $Q_{\lambda}^{\pi_0}(\bar{s}, a)$ (after composing with π_0) are continuous in λ at λ_0 for all $\bar{s} \in \bar{S}_T$. Finally, the goal-reaching objective $J_{\lambda}^{\pi_0}$ (cf. (2.2)) is continuous in λ at λ_0 due to the described continuity of values $V_{\lambda}^{\pi_0}(\bar{s})$.

Iteration n = 1: Let us fix $\bar{s} \in \bar{S}_T$. First assume $\bar{s} \notin \text{supp} \det_{\lambda_0,\pi_0}$. For any neighborhood $U_{\delta'}(\lambda_0)$, $0 < \delta' < \delta$ we can find an interior point $\lambda' \in U_{\delta'}(\lambda_0)$, i.e., we can find a sequence of interior points $(\lambda'_k)_{k\geq 0}$ converging to λ_0 . From lemma 37, only one of the following cases can happen: either $\bar{s} \in \text{supp} \det_{\lambda'_k,\pi_0}$ for all k or $\bar{s} \notin \text{supp} \det_{\lambda'_k,\pi_0}$ for all k. If the first case is true, $\pi_{1,\lambda}(\cdot|\bar{s})$ can be discontinuous at λ_0 due to discontinuity of supp \det_{λ,π_0} at λ_0 , as evidenced by the sequence $(\lambda'_k)_{k\geq 0}$ (the definition of $\pi_{1,\lambda}(\cdot|\bar{s})$, cf. 4.1, changes abruptly from num/den ratio to $1/|\mathcal{A}|$). This behavior is illustrated in example 1 (of non-deterministic boundary point λ_0) in the appendix (see $\pi_{1,0+}, \pi_{1,0}, \pi'_{1,0+}$ in the third column (g = 2)), and also in example 2 (of deterministic λ_0) in the appendix (see again the third column).

In case $\bar{s} \in \operatorname{supp} \operatorname{den}_{\lambda_0,\pi_0}$, it holds that $\bar{s} \in \operatorname{supp} \operatorname{den}_{\lambda_0,\pi_0} \subset \operatorname{supp} \operatorname{den}_{\lambda,\pi_0}$ for all $\lambda \in U_{\delta}(\lambda_0)$ by point 2. of the lemma 39. This means that $\pi_{1,\lambda}(\cdot|\bar{s})$ is defined on the whole $U_{\delta}(\lambda_0)$, using the ratio $\operatorname{num}_{\lambda,\pi_0}(\bar{s},a)/\operatorname{den}_{\lambda,\pi_0}(\bar{s})$. Following the reasoning in point 1. of lemma 38, utilizing the continuity of π_0 , point 4. of the lemma 12, and $\operatorname{den}_{\lambda_0,\pi_0}(\bar{s}) > 0$, we conclude that $\pi_{1,\lambda}(\cdot|\bar{s})$ is continuous in λ at λ_0 .

Iteration n = 2: Apart from the discontinuities of $\pi_{2,\lambda}(\cdot|\bar{s})$ (in λ at λ_0) which can arise due to $\bar{s} \notin \text{supp den}_{\lambda_0,\pi_1}$, in the same way as was already described for n = 1, there can be, additionally, also discontinuities for states in the set \bar{S}_{λ_0} . In order to discuss these, let us fix $\bar{s} \in \bar{S}_{\lambda_0}$. Since $\bar{S}_{\lambda_0} \subset \text{supp den}_{\lambda,\pi_1} \cap \text{supp } \nu_{\lambda,\pi_1}$ (by point 2. of lemma 39) the policy $\pi_{2,\lambda}(\cdot|\bar{s})$ is defined by the ratio

$$\pi_{2,\lambda}(\cdot|\bar{s}) = \frac{\operatorname{num}_{\lambda,\pi_1}(\bar{s},\cdot)}{\operatorname{den}_{\lambda,\pi_1}(\bar{s})}$$

for all $\lambda \in U_{\delta}(\lambda_0)$. However, since $\pi_{1,\lambda}$ can be already discontinuous for some $\bar{s}' \in \bar{S}_T \setminus$ supp den $_{\lambda_0,\pi_0}$ (as we have described in paragraph for n = 1), this discontinuity can propagate to $\operatorname{num}_{\lambda,\pi_1}(\bar{s}, \cdot)$, den $_{\lambda,\pi_1}(\bar{s})$ causing a discontinuity of $\pi_{2,\lambda}(\cdot|\bar{s})$ defined via the above ratio.

This behavior is illustrated in example 1 in the appendix (see $\pi_{2,0+}$, $\pi_{2,0}$, $\pi'_{2,0+}$ first column (g = 0)), and also in example 2 in the appendix (see the second column and figure 2a).

Summary: To sum up, policies can become discontinuous already at iteration n = 1, but this occurs outside of \bar{S}_{λ_0} . At iteration n = 2, we start also observing discontinuities on \bar{S}_{λ_0} , which is more problematic. While at deterministic kernels the policies on \bar{S}_{λ_0} are relatively continuous, causing the goal-reaching objective to be continuous (cf. theorem 16 and corollary 18). In non-deterministic boundary points this is not the case and policy discontinuities (for $n \geq 2$) propagates through values to goal-reaching objective (cf. figure 2d).

Appendix D. Regularized Recursion — Lemmas and Proofs

Here we introduce and prove the various lemmas and theorems used in section 8.

D.1 Preliminary Lemmata

The following lemma is the ϵ -eUDRL version of lemma 6.

Lemma 40 (ϵ -eUDRL version of support stability) Let $\{\mathcal{M}_{\lambda} : \lambda \in (\Delta S)^{S \times A}\}$ and $\{\overline{\mathcal{M}}_{\lambda} : \lambda \in (\Delta S)^{S \times A}\}$ be compatible families. Let $(\pi_{n,\lambda,\epsilon})_{n\geq 0}$ be a sequence of policies generated by the ϵ -eUDRL iteration given a transition kernel $\lambda \in (\Delta S)^{S \times A}$ and an initial condition π_0 (that does not depend on λ) and a regularization parameter $1 > \epsilon > 0$. Fix a deterministic transition kernel $\lambda_0 \in \Delta S^{S \times A}$. Then for all initial conditions $\pi_0 > 0$ it holds:

- 1. For all $n \geq 0$ and all $\lambda \in U_{\delta}(\lambda_0)$ we have that $\operatorname{supp num}_{\lambda_0,\pi_0} \cap (\mathcal{A} \times \operatorname{supp} \nu_{\lambda_0,\pi_0}) \subset \operatorname{supp num}_{\lambda,\pi_{n,\epsilon}} \cap (\mathcal{A} \times \operatorname{supp} \nu_{\lambda,\pi_{n,\epsilon}})$, where the inclusion becomes equality for $\lambda = \lambda_0$.
- 2. For all $n \geq 0$ and all $\lambda \in U_{\delta}(\lambda_0)$ we have that $\bar{S}_{\lambda_0} \subset \operatorname{supp} \operatorname{den}_{\lambda,\pi_{n,\epsilon}} \cap \operatorname{supp} \nu_{\lambda,\pi_{n,\epsilon}}$, where the inclusion becomes an equality for $\lambda = \lambda_0$.
- 3. For all $n \ge 0$ and all $\lambda \in U_{\delta}(\lambda_0)$ we have that $\mathbb{P}_{\lambda}(S_0^{\Sigma} = s, l(\Sigma) = h, \rho(S_h^{\Sigma}) = g, H_0^{\Sigma} = h, G_0^{\Sigma} = g; \pi_{n,\epsilon}) > 0$ for all $(s, h, g) \in \bar{S}_{\lambda_0}$.

The points 4. and 5. of the original lemma 6 hold also for ϵ -eUDRL.

Proof The equation (4.2) is proved exactly the same as in the proof of the original lemma 6.

1. There is an easier alternative proof for this point which we present now. Since $\pi_{n,\epsilon} > 0$ for $n \ge 0$ (because of the regularization and the assumption that $\pi_0 > 0$) we trivially have $\operatorname{supp} \pi_{n,\epsilon} = \operatorname{supp} \pi_0$. Now fix $\lambda \in U_2(\lambda_0)$. From $\operatorname{supp} \lambda_0 \subset \operatorname{supp} \lambda$ (cf. equation (4.2)) and just stated fact about policy supports it follows:

$$\operatorname{supp} \nu_{\lambda_0,\pi_0} \subset \operatorname{supp} \nu_{\lambda,\pi_{n,\epsilon}}, \quad \operatorname{supp} \operatorname{num}_{\lambda_0,\pi_0} \subset \operatorname{supp} \operatorname{num}_{\lambda,\pi_{n,\epsilon}}. \tag{D.1}$$

Now we take the first inclusion and perform the cartesian product with \mathcal{A} and intersection with supp $\operatorname{num}_{\lambda_0,\pi_0}$. Further we take the second inclusion and perform intersection with $\operatorname{supp} \nu_{\lambda,\pi_{n,\epsilon}} \times \mathcal{A}$. We obtain the following chain of inclusions:

$$(\operatorname{supp}\nu_{\lambda_0,\pi_0}\times\mathcal{A})\cap\operatorname{supp}\operatorname{num}_{\lambda_0,\pi_0}\subset(\operatorname{supp}\nu_{\lambda,\pi_{n,\epsilon}}\times\mathcal{A})\cap\operatorname{supp}\operatorname{num}_{\lambda_0,\pi_0}\subset(\operatorname{supp}\nu_{\lambda,\pi_{n,\epsilon}}\times\mathcal{A})\cap\operatorname{supp}\operatorname{num}_{\lambda,\pi_{n,\epsilon}},$$

which leaves the result. The equality is proved exactly the same as in the original proof.

2. The proof is unchanged.

3. The proof can be simplified but otherwise repeated with minimal changes as the new version of (4.3) trivially holds (since $\pi_{n,\epsilon} > 0$ for all $n \ge 0$).

The remaining proof of 4. and 5. is unchanged.

The following lemma is the ϵ -eUDRL version of the lemma 10.

Lemma 41 (optimality of ϵ -eUDRL policies for deterministic transition kernel) Let λ_0 be a deterministic transition kernel and let $\mathcal{M} = (\mathcal{S}, \mathcal{A}, \lambda_0, \mu, r)$ be a respective MDP with CE $\overline{\mathcal{M}} = (\overline{\mathcal{S}}, \mathcal{A}, \overline{\lambda}_0, \overline{\mu}, \overline{r}, \rho)$. Assume $\pi_0 > 0$ and let $(\pi_{n,\epsilon})_{n\geq 0}$ be the policy sequence generated by the ϵ -eUDRL iteration given π_0, λ_0 and regularisation parameter ϵ . Then it holds:

1. For all $n \ge 0$ the policy $\pi_{n+1,\epsilon}$ has the form (on \bar{S}_{λ_0})

$$\pi_{n+1,\epsilon} = (1-\epsilon)\pi_{n+1}^* + \frac{\epsilon}{|\mathcal{A}|},$$

where

$$\pi_{n+1}^* = \frac{\operatorname{num}_{\lambda_0, \pi_{n,\epsilon}}(a, s, h, g)}{\operatorname{den}_{\lambda_0, \pi_{n,\epsilon}}(s, h, g)}$$

is an optimal policy on \bar{S}_{λ_0} .

2. For all $n \ge 1$ and all $\bar{s} = (s, h, g) \in \bar{S}_{\lambda_0}$ it holds $V^{\pi_{n,\epsilon}}(\bar{s}) \ge (1-\epsilon)^h \ge (1-\epsilon)^N.$

3. For all $n \ge 1$ and all $\bar{s} = (s, h, g) \in \bar{S}_{\lambda_0}$ it holds

$$Q^{\pi_{n,\epsilon}}(\bar{s}, a) \ge (1-\epsilon)^{h-1} \ge (1-\epsilon)^N \quad \text{for } a \in \mathcal{O}(\bar{s}),$$

= 0 otherwise.

Proof 1. The equations is just a rewrite of the ϵ -eUDRL recursion. The easiest way to prove optimality of π_n^* is to realize that π_n^* is defined similarly as π_n (in the original lemma) except that it uses ϵ -version of "num/den"-ratio. Since the proof of point 1. (in the original lemma) is essentially just a statement about the supports and these do not change (due to points 1. and 2. of lemma 6 and points 1. and 2. of lemma 40), i.e.,

$$(\operatorname{supp} \nu_{\lambda_0,\pi_n} \times \mathcal{A}) \cap \operatorname{supp} \operatorname{num}_{\lambda_0,\pi_n} = (\operatorname{supp} \nu_{\lambda_0,\pi_0} \times \mathcal{A}) \cap \operatorname{supp} \operatorname{num}_{\lambda_0,\pi_0} = (\operatorname{supp} \nu_{\lambda_0,\pi_{n,\epsilon}} \times \mathcal{A}) \cap \operatorname{supp} \operatorname{num}_{\lambda_0,\pi_{n,\epsilon}}, \operatorname{supp} \operatorname{den}_{\lambda_0,\pi_n} \cap \operatorname{supp} \nu_{\lambda_0,\pi_n} = \bar{\mathcal{S}}_{\lambda_0} = \operatorname{supp} \operatorname{den}_{\lambda_0,\pi_{n,\epsilon}} \cap \operatorname{supp} \nu_{\lambda_0,\pi_{n,\epsilon}},$$

the π_n^* has to be optimal too.

2. & 3. The proof follows by induction on the remaining horizon h. For h = 1 the Q-values are independent of a policy and therefore are optimal, i.e., for all $\bar{s} = (s, 1, g) \in \bar{S}_{\lambda_0}$, it holds that $Q^{\pi_{n,\epsilon}}((s, 1, g), a) = 1$ for $a \in \mathcal{O}(s, 1, g)$ and $Q^{\pi_{n,\epsilon}}((s, 1, g), a) = 0$ otherwise. Now assume that 2. and 3. holds for a fixed horizon h. We obtain (for all $\bar{s} = (s, h, g) \in \bar{S}_{\lambda_0}$)

$$\begin{aligned} V^{\pi_{n,\epsilon}}(\bar{s}) &= \sum_{a \in \mathcal{A}} \pi_{n,\epsilon}(a|\bar{s})Q^{\pi_{n,\epsilon}}(\bar{s},a) \\ &\geq \sum_{a \in \mathcal{O}(\bar{s})} \pi_{n,\epsilon}(a|\bar{s})(1-\epsilon)^{h-1} = (1-\epsilon)^{h-1}\pi_{n,\epsilon}(\mathcal{O}(\bar{s})|\bar{s}) \\ &= (1-\epsilon)^{h-1}((1-\epsilon)\pi_n^*(\mathcal{O}(\bar{s})|\bar{s}) + \frac{\epsilon|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}) = (1-\epsilon)^{h-1}(1-\epsilon(1-\frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|})) \\ &\geq (1-\epsilon)^h, \end{aligned}$$

where we used the induction assumption and the point 1. Further for all $\bar{s} = (s, h+1, g) \in \bar{S}_{\lambda_0}$ it holds

$$\begin{aligned} Q^{\pi_{n,\epsilon}}(\bar{s},a) &= \sum_{s' \in \mathcal{S}} \lambda_0(s'|s,a) V^{\pi_{n,\epsilon}}(s',h,g) = \lambda_0(s''|s,a) V^{\pi_{n,\epsilon}}(s'',h,g) \ge (1-\epsilon)^h \quad \text{for } a \in \mathcal{O}(\bar{s}) \\ &= 0 \quad \text{otherwise}, \end{aligned}$$

where for $a \in \mathcal{O}(\bar{s})$ we used that the kernel is deterministic and therefore there exists a $s'' \in S$ such that $\lambda_0(s''|s, a) = 1$ and, further, that from point 4. of lemma 40, $(s'', h, g) \in \bar{S}_{\lambda_0}$. The statement for $a \notin \mathcal{O}(\bar{s})$ is just point 5. of lemma 40. This completes the induction.

The class of policies $\Pi_{\lambda_0,\epsilon}^* := \{(1-\epsilon)\pi_{\lambda_0}^* + \frac{\epsilon}{|\mathcal{A}|} \mid \pi_{\lambda_0}^*$ is an optimal policy for λ_0 on $\bar{S}_{\lambda_0}\}$ which was discussed in the previous lemma will be of great importance in the following text because (given $\bar{s} \in \bar{S}_{\lambda_0}$) it constitutes the limit relative to $\mathcal{O}(\bar{s})$ we would like to achieve when proving relative continuity of ϵ -eUDRL policies in λ at λ_0 . Unlike the optimal action-value function $Q_{\lambda_0}^*$, which is constant on $\mathcal{O}(\bar{s})$ (and therefore factors through a quotient map relative to $\mathcal{O}(\bar{s})$), there are many $Q_{\lambda_0}^{\pi_{\epsilon}^*}$ depending on $\pi_{\lambda_0,\epsilon}^* \in \Pi_{\lambda_0,\epsilon}^*$ and which are generally non-constant on $\mathcal{O}(\bar{s})$. Note that during the proof of points 2. and 3. of the above lemma we exclusively used policy properties following from $\pi_{n,\epsilon} \in \Pi_{\lambda_0,\epsilon}^*$ for n > 0. Therefore, the points 2. and 3. describe lower bounds on the values $Q_{\lambda_0}^{\pi_{\epsilon}^*}$ and $V_{\lambda_0}^{\pi_{\epsilon}^*}$, respectively, where $\pi_{\lambda_0,\epsilon}^* \in \Pi_{\lambda_0,\epsilon}^*$.

The following lemma is the ϵ -eUDRL version of the lemma 22.

Lemma 42 (ϵ -eUDRL version of the lower bound on visitation probabilities) Let { \mathcal{M}_{λ} : $\lambda \in (\Delta S)^{S \times \mathcal{A}}$ } and { $\overline{\mathcal{M}}_{\lambda} : \lambda \in (\Delta S)^{S \times \mathcal{A}}$ } be compatible families. Let λ_0 be a deterministic kernel. Let $(\pi_{n,\lambda,\epsilon})_{n\geq 0}$ denotes ϵ -eUDRL generated sequence given the initial condition $\pi_0 > 0$, the transition kernel λ , and the regularization parameter ϵ . Then for all n > 0 all $\overline{s} = (s, h, g) \in \overline{S}_{\lambda_0}$ and all $\lambda \in U_2(\lambda_0)$ it holds

$$\mathbb{P}_{\lambda}(H_0^{\Sigma} = h, G_0^{\Sigma} = g | S_0^{\Sigma} = s, l(\Sigma) = h; \pi_{n,\epsilon}) \ge \alpha(\delta, \epsilon),$$

where

$$\alpha(\delta,\epsilon) = \frac{2}{N(N+1)} (\min_{\bar{s}' \in \operatorname{supp} \bar{\mu}} \bar{\mu}(\bar{s}')) (\frac{\epsilon}{|\mathcal{A}|})^N (1 - \frac{\delta}{2})^N > 0.$$

Notice that $\alpha(\delta, \epsilon) \to \alpha(0, \epsilon_0)$ as $(\delta, \epsilon) \to (0, \epsilon_0)$, where we assume an $\epsilon_0 \ge 0$, and where

$$\alpha(0,\epsilon_0) = \frac{2}{N(N+1)} (\min_{\bar{s}' \in \operatorname{supp} \bar{\mu}} \bar{\mu}(\bar{s}')) \frac{1}{|\mathcal{A}|^N} \epsilon_0^N > 0$$

for $\epsilon_0 > 0$.

Proof We can bound the ϵ -eUDRL policy

$$(\forall n > 0, \bar{s} \in \bar{\mathcal{S}}_{\lambda_0}) : \pi_{n,\lambda,\epsilon}(\cdot|\bar{s}) \ge \frac{\epsilon}{|\mathcal{A}|} =: \alpha' > 0.$$

Further, we continue similarly as in theorem 31 "Bounding the visitation term" except we utilize the above α' bound for lower bounding the policy terms. In short, from $\bar{s} \in \bar{S}_{\lambda_0}$ we have $\bar{s} \in \text{supp } \nu_{\lambda_0,\pi_0}$ meaning there exists a prefix $\bar{s}_0, a_0, \ldots, \bar{s}_t = \bar{s}$ with positive probability $\mathbb{P}_{\lambda_0}(\bar{S}_t = \bar{s}, \ldots, A_0 = a_0, \bar{S}_0 = \bar{s}_0; \pi_0) > 0$. Rewriting this probability as a product of λ_0 terms and π_0 terms then replacing the λ_0 with λ and π_0 with $\pi_{n,\lambda,\epsilon}$ doing exactly the same bounding of λ terms as in theorem 31 and bounding $\pi_{n,\lambda,\epsilon}$ using the α' bound above we obtain

$$\mathbb{P}_{\lambda}(\bar{S}_t = \bar{s}; \pi_{n,\lambda,\epsilon}) \ge \left(\min_{\bar{s}' \in \operatorname{supp} \bar{\mu}} \bar{\mu}(\bar{s}')\right) (\alpha')^N (1 - \frac{\delta}{2})^N$$

the rest of the proof is the same as in theorem 31 leading to the α bound above.

Note that the bound could be easily extended to $n \ge 0$ by putting

$$\alpha' := \min\{\min_{\bar{s}' \in \bar{\mathcal{S}}_{\lambda_0}, a \in \mathcal{A}} \pi_0(a|\bar{s}'), \frac{\epsilon}{|\mathcal{A}|}\}.$$

Although we will suffice with the above simpler version.

We will call a policy $\pi \epsilon$ -regular if and only if, for all $\bar{s} \in \bar{S}_{\lambda_0}$ and $a \in A$, it holds that $\pi(a|\bar{s}) > \frac{\epsilon}{|A|}$. The following remark describes equivalent statements about the distance of an ϵ -regular policy to $\Pi^*_{\lambda_0,\epsilon}$.

Remark 43 (distance of ϵ -regular policy to $\Pi^*_{\lambda_0,\epsilon}$) Assume $\delta > 0$ and π_{ϵ} is an ϵ -regular policy, then the following statements are equivalent:

(a) $2(1 - \epsilon(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}) - \pi_{\epsilon}(\mathcal{O}(\bar{s}')|\bar{s}')) < \delta$

$$(\mathbf{b}) \ (\exists \pi_{\epsilon}^* \in \Pi_{\lambda_0, \epsilon}^*) : \quad \|\pi_{\epsilon}^* - \pi_{\epsilon}\|_1 < \delta \quad \land \quad ((\forall \bar{s} \in \bar{\mathcal{S}}_{\lambda_0}, \forall a \in \mathcal{O}(\bar{s})) : \pi_{\epsilon}^*(a|\bar{s}) \ge \pi_{\epsilon}(a|\bar{s}))$$

(c)
$$(\exists \tilde{\pi}^*_{\epsilon} \in \Pi^*_{\lambda_0, \epsilon}) : \|\tilde{\pi}^*_{\epsilon} - \pi_{\epsilon}\|_1 < \delta$$

Note that ϵ -eUDRL generated policies are ϵ -regular, as well as policies from $\Pi^*_{\lambda_0,\epsilon}$. A policy from $\Pi^*_{\lambda_0,\epsilon}$ can be characterized by being ϵ -regular and putting maximum mass on $\mathcal{O}(\bar{s})$ for all $\bar{s} \in \bar{S}_{\lambda_0}$, where this maximum is $1 - \epsilon (1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|})$.

Proof (of the remark 43) We begin by the proof of (c) \implies (b). Assume (c). Since π_{ϵ} might not be in $\Pi^*_{\lambda_0,\epsilon}$, we have $\pi_{\epsilon}(\mathcal{O}(\bar{s})|\bar{s}) \leq 1 - \epsilon(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}) \ (= \tilde{\pi}^*_{\epsilon}(\mathcal{O}(\bar{s})|\bar{s}))$. We can easily construct $\pi^*_{\epsilon} \in \Pi^*_{\lambda_0,\epsilon}$ satisfying the second condition in (b), e.g., by

$$\pi_{\epsilon}^{*}(a|\bar{s}) = \begin{cases} \pi_{\epsilon}(a|\bar{s}) + (1 - \epsilon(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}) - \pi_{\epsilon}(\mathcal{O}(\bar{s})|\bar{s})) / |\mathcal{O}(\bar{s})| & \text{for } a \in \mathcal{O}(\bar{s}) \\ \frac{\epsilon}{|\mathcal{A}|} & \text{otherwise.} \end{cases}$$

Since $\tilde{\pi}_{\epsilon}^*$ might not satisfy the second condition in (b), we have $\delta > \|\tilde{\pi}_{\epsilon}^* - \pi_{\epsilon}\|_1 \ge \|\pi_{\epsilon}^* - \pi_{\epsilon}\|_1$. This concludes the proof of (c) \implies (b).

The implication (b) \implies (a) is trivial:

$$\begin{split} \delta > \|\pi_{\epsilon}^* - \pi_{\epsilon}\|_1 &= \pi_{\epsilon}^*(\mathcal{O}(\bar{s})|\bar{s}) - \pi_{\epsilon}(\mathcal{O}(\bar{s})|\bar{s}) + \pi_{\epsilon}(\mathcal{A} \setminus \mathcal{O}(\bar{s})|\bar{s}) - \pi_{\epsilon}^*(\mathcal{A} \setminus \mathcal{O}(\bar{s})|\bar{s}) \\ &= 2(\pi_{\epsilon}^*(\mathcal{O}(\bar{s})|\bar{s}) - \pi_{\epsilon}(\mathcal{O}(\bar{s})|\bar{s})) \\ &= 2(1 - \epsilon(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}) - \pi_{\epsilon}(\mathcal{O}(\bar{s})|\bar{s}))), \end{split}$$

where we used $\pi_{\epsilon}^*(\mathcal{A}|\bar{s}) = \pi_{\epsilon}(\mathcal{A}|\bar{s}) = 1.$

The implication (a) \implies (c) is proved as follows. Assuming (a) we can construct $\tilde{\pi}_{\epsilon}^* \in \Pi_{\lambda_0,\epsilon}^*$ in the same way as we were constructing π_{ϵ}^* in the proof of (c) \implies (b).

In the following lemma, we will aim to show that when π_{ϵ} (we will consider just ϵ -regularized policies) is close to $\Pi^*_{\lambda_0,\epsilon}$ and λ is close to λ_0 , then $Q^{\pi_{\epsilon}}_{\lambda}$ is close to the set $\{Q^{\pi_{\epsilon}}_{\lambda_0} \mid \pi^*_{\lambda_0,\epsilon} \in \Pi^*_{\lambda_0,\epsilon}\}$. The following lemma is the ϵ -eUDRL version of the lemma 21.

Lemma 44 (ϵ -eUDRL version of the continuity of action-values in quotient topology) Let $\{\mathcal{M}_{\lambda} : \lambda \in (\Delta S)^{S \times A}\}$ and $\{\bar{\mathcal{M}}_{\lambda} : \lambda \in (\Delta S)^{S \times A}\}$ be compatible families. Let λ_0 be a deterministic kernel. For all $\epsilon' > 0$ there exists $\delta > 0$ such that if $\lambda \in U_{\delta}(\lambda_0)$ and the ϵ -regular policy π_{ϵ} satisfies, for all $\bar{s} = (s, h, g) \in \bar{S}_{\lambda_0}$,

$$2(1-\epsilon(1-\frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|})-\pi_{\epsilon}(\mathcal{O}(\bar{s})|\bar{s}))<\delta,$$

then $|Q_{\lambda}^{\pi_{\epsilon}}(\bar{s},\cdot) - Q_{\lambda_{0}}^{\pi_{\epsilon}^{*}}(\bar{s},\cdot)| < \epsilon'$ for all $\bar{s} = (s,h,g) \in \bar{S}_{\lambda_{0}}$ for some $\pi_{\lambda_{0},\epsilon}^{*} \in \Pi_{\lambda_{0},\epsilon}^{*}$. The statement can be made explicit by a recursive estimate: for all $\bar{s} = (s,h,g) \in \bar{S}_{\lambda_{0}}$, all $h \geq 2$, and all $a \in \mathcal{O}(\bar{s})$ it holds that

$$\begin{aligned} |Q_{\lambda}^{\pi_{\epsilon}}(\bar{s},a) - Q_{\lambda_{0}}^{\pi_{\epsilon}}(\bar{s},a)| &\leq \|\lambda(\cdot|s,a) - \lambda_{0}(\cdot|s,a)\|_{1} \\ &+ \max_{\bar{s}'=(s',h-1,g)\in\bar{\mathcal{S}}_{\lambda_{0}}} 2(1 - \epsilon(1 - \frac{|\mathcal{O}(\bar{s}')|}{|\mathcal{A}|}) - \pi_{\epsilon}(\mathcal{O}(\bar{s}')|\bar{s}')) \\ &+ \max_{\bar{s}'=(s',h-1,g)\in\bar{\mathcal{S}}_{\lambda_{0}},a'\in\mathcal{O}(\bar{s}')} |Q_{\lambda}^{\pi_{\epsilon}}(\bar{s}',a') - Q_{\lambda_{0}}^{\pi_{\epsilon}^{*}}(\bar{s}',a')|. \end{aligned}$$

Proof The proof follows as in lemma 21 by induction on the remaining horizon. In the induction step, we simply show that $|Q_{\lambda}^{\pi_{\epsilon}}(\bar{s},a) - Q_{\lambda_{0}}^{\pi_{\epsilon}^{*}}(\bar{s},a)| \to 0$ for $\delta \to 0$ by bounding it by δ and using the induction assumption. The trick is to use the implication (a) \implies (b) of the remark 43 to select a convenient π_{ϵ}^{*} for a given π_{ϵ} .

The following lemma is the ϵ -eUDRL version of lemma 24.

Lemma 45 Let $\gamma, \epsilon > 0, 1 > \gamma + \epsilon, |\mathcal{A}| \ge M > 0$ and $z_{\gamma,\epsilon,M} : [0,1] \rightarrow [0,1]$ be defined as

$$z_{\gamma,\epsilon,M}(x) = (1-\epsilon)\frac{x}{x+\gamma} + \epsilon \frac{M}{|\mathcal{A}|}$$

Then the following assertions hold

- 1. $z_{\gamma,\epsilon,M}$ is increasing.
- 2. $z_{\gamma,\epsilon,M}$ has a unique fixed point $x^*(\gamma,\epsilon,M)$

$$x^*(\gamma, \epsilon, M) = \frac{\hat{x}^* + \sqrt{(\hat{x}^*)^2 + \frac{4\gamma\epsilon M}{|\mathcal{A}|}}}{2},$$
where $\hat{x}^* = 1 - \epsilon(1 - \frac{M}{|\mathcal{A}|}) - \gamma$, $\hat{x}^* \leq x^*(\gamma, \epsilon, M)$ and $1 > x^*(\gamma, \epsilon, M) > 0$. Further, $x^*(\gamma, \epsilon, M) \to x^*(0, \epsilon_0, M) = 1 - \epsilon_0(1 - \frac{M}{|\mathcal{A}|})$ as $(\gamma, \epsilon) \to (0, \epsilon_0)$,

for any $1 > \epsilon_0 \ge 0$, and it holds that

$$\begin{split} &z_{\gamma,\epsilon,M}(x) > x \quad for \ x < x^*(\gamma,\epsilon,M), \\ &z_{\gamma,\epsilon,M}(x) < x \quad for \ x > x^*(\gamma,\epsilon,M). \end{split}$$

- 3. For all $x \in [0,1]$ the iterated application of $z_{\gamma,\epsilon,M}$ converges pointwise, $z_{\gamma,\epsilon,M}^{\circ n}(x) \xrightarrow{n \to \infty} x^*(\gamma,\epsilon,M)$.
- 4. Assume a sequence $(y_n), y_n \in [0,1]$ such that for all $n \ge 0$ we have $y_{n+1} \ge z_{\gamma,\epsilon,M}(y_n)$. Then $y_n \ge z_{\gamma,\epsilon,M}^{\circ n}(y_0)$ and $\liminf_n y_n \ge x^*(\gamma,\epsilon,M)$.

Proof 1. The derivative of z is positive.

2. We follow the same approach as in lemma 24 except the resulting equality/inequality is now quadratic (instead of linear). However there is just one root in [0, 1] (the other one is always negative) which is also the only fixed point $x^*(\gamma, \epsilon, M)$. From $\frac{4\gamma\epsilon M}{|\mathcal{A}|} > 0$ follows the inequality $\hat{x}^* \leq x^*$. To prove the bounds $1 > x^*(\gamma, \epsilon, M) > 0$ is trivial.

3. & 4. The rest follows as in lemma 24 except it is somewhat more simplified as there are no complications with the point x = 0.

The figure 14 illustrates the dynamical system induced by $z_{\gamma,\epsilon,M}$. The lemma shows some of its properties (e.g., the fixed point).

D.2 The Main Theorem

The following theorem is the ϵ -eUDRL version of theorem 25.

Theorem 34 (Relative continuity of ϵ -eUDRL limit policies in deterministic kernels) Let $\{\mathcal{M}_{\lambda} : \lambda \in (\Delta S)^{S \times A}\}$ and $\{\overline{\mathcal{M}}_{\lambda} : \lambda \in (\Delta S)^{S \times A}\}$ be compatible families. Let λ_0 be a deterministic kernel and let $(\pi_{n,\lambda,\epsilon})_{n\geq 0}$ be a sequence of ϵ -eUDRL-generated policies with initial condition $\pi_0 > 0$, transition kernel λ and regularization parameter $\epsilon \in (0,1)$. Let $\epsilon_0 \in (0,1)$ be a fixed regularization parameter. Then for all $\pi_0 > 0$ the following statements hold:

1. Let $\mathcal{L}(\pi_0, \lambda, \epsilon)$ denote the set of accumulation points of $(\pi_{n,\lambda,\epsilon})_{n\geq 0}$. Then any function $u : (\pi_0, \lambda, \epsilon) \to u(\pi_0, \lambda, \epsilon) \in \mathcal{L}(\pi_0, \lambda, \epsilon)$ is relatively continuous in λ, ϵ at point λ_0, ϵ_0 on $\bar{\mathcal{S}}_{\lambda_0}$, *i.e.* for all $\bar{s} \in \bar{\mathcal{S}}_{\lambda_0}$

$$[u(\pi_0,\lambda,\epsilon_0)](\cdot|\bar{s}) \xrightarrow{\mathcal{O}(\bar{s})} \pi^*_{\lambda_0,\epsilon_0}(\cdot|\bar{s}) \quad as \quad \lambda \to \lambda_0.$$

2. Let $\alpha(\delta, \epsilon)$ is chosen as in lemma 42 and let $\beta, \tilde{\beta}, \epsilon \in (0, 1)$ be such that $1 > \gamma + \epsilon$, where $\gamma = \frac{\tilde{\beta}}{((1-\epsilon)^N - \beta)\alpha(\delta,\epsilon)}$ and let $U_{\delta}(\lambda_0) = \{\lambda \mid \max_{(s,a)\in \mathcal{S}\times\mathcal{A}} \|\lambda(\cdot \mid s, a) - \lambda_0(\cdot \mid s, a)\|_1 < \delta\}.$



Figure 14: The map $z_{\gamma,\epsilon,M}$ as a dynamical system: the dependence of $z_{\gamma,\epsilon,M}^{\circ n}(x)$ (repeated application of $z_{\gamma,\epsilon,M}$ given initial condition x) on iteration n for various initial conditions x. The parameters are set with $|\mathcal{A}| = 4$, M = 1, and $\gamma = 0.4$. The regularization parameter is set to $\epsilon = 0.1$ for black trajectories and to $\epsilon = 0.3$ for green trajectories, respectively.

There exists $\delta > 0$ so that for all $\lambda \in U_{\delta}(\lambda_0)$ and all $\bar{s} \in \bar{S}_{\lambda_0}$ it holds

$$\liminf_{n} \pi_{n,\lambda,\epsilon}(\mathcal{O}(\bar{s})|\bar{s}) \ge x^*(\gamma,\epsilon,|\mathcal{O}(\bar{s})|)$$

and

$$x^*(\gamma,\epsilon,|\mathcal{O}(\bar{s})|) \to 1 - \epsilon_0 \left(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}\right) \quad as \quad (\beta,\tilde{\beta},\alpha,\epsilon) \to (0,0,\alpha(0,\epsilon_0),\epsilon_0),$$

where $x^*(\gamma, \epsilon, |\mathcal{O}(\bar{s})|)$ is given in lemma 45. Consequently for all $\bar{s} \in \bar{S}_{\lambda_0}$ it holds that

$$\liminf_{n} \pi_{n,\lambda,\epsilon}(\mathcal{O}(\bar{s})|\bar{s}) \to 1 - \epsilon_0 \left(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}\right) \quad as \quad (\lambda,\epsilon) \to (\lambda_0,\epsilon_0)$$

At the core of the proof is the analysis of dynamical systems and convergence induced by iterative application of the rational function $z_{\gamma,\epsilon,M}(x) = (1-\epsilon)\frac{x}{x+\gamma} + \epsilon \frac{M}{|\mathcal{A}|}$ (see lemma 45). Indeed, we will show that for all horizons $N \ge h \ge 1$ and all $\beta, \tilde{\beta}, \epsilon \in (0, 1)$ satisfying $\gamma + \epsilon < 1$ (with γ as in point 2. above), there exists n_0 and $\delta > 0$ such that, for all initial conditions $\pi_0 > 0$ all $\lambda \in U_{\delta}(\lambda_0), n > n_1 \ge n_0$, and all $\bar{s} \in \bar{S}_{\lambda_0}$ with remaining horizon h, it holds that

$$\pi_{n,\lambda,\epsilon}(\mathcal{O}(\bar{s})|\bar{s}) \ge z_{\gamma,\epsilon,|\mathcal{O}(\bar{s})|}^{\circ(n-n_1)}(\frac{\epsilon|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}).$$
(D.2)

Then, by lemma 45, the right-hand side converges to $x^*(\gamma, \epsilon, |\mathcal{O}(\bar{s})|)$, which implies point 2. of the theorem (see the proof for details). Furthermore, point 1. is a consequence of point 2. The argument for this is the same as in section 7.1.2.

Proof The proof is analogous to the proof of theorem 25 thus we will concentrate on discussing the small differences (of course, with the lemmas introduced in this appendix instead of their original versions).

Since the lower bound $\pi_{n,\epsilon}(\mathcal{O}(\bar{s})) \geq \frac{\epsilon |\mathcal{O}(\bar{s})|}{|\mathcal{A}|} \geq \frac{\epsilon}{|\mathcal{A}|} > 0$ (for all $\bar{s} \in \bar{S}_{\lambda_0}$) coming from ϵ -regularity of ϵ -eUDRL-generated policies is used in (D.2), we do not need to prove any analogies to (7.5).

In point 1., $(\beta, \beta, \alpha(\delta, \epsilon), \epsilon) \to (0, 0, \alpha(0, \epsilon_0), \epsilon_0)$ (where $\epsilon_0 > 0$ and $\alpha(0, \epsilon_0) > 0$ from lemma 45) implies $\gamma \to 0$ (continuity of γ at point $(0, 0, \alpha(0, \epsilon_0), \epsilon_0)$). Further, $(\gamma, \epsilon) \to (0, \epsilon_0)$ causes $x^* \to 1 - \epsilon_0(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|})$, according to lemma 45.

The proof proceeds by showing implications 2. \implies 1. and (D.2) \implies 2. similarly as in the original proof. Here we will focus on the proof of (D.2). Fixing $2 > \delta > 0$ and $\epsilon > 0$, lemma 40 asserts that $\bar{S}_{\lambda_0} \subset \operatorname{supp} \operatorname{den}_{\lambda,\pi_{n,\epsilon}} \cap \operatorname{supp} \nu_{\lambda,\pi_n}$ for all $n \ge 0, \pi_0 \ge 0, \lambda \in U_{\delta}(\lambda_0)$. The policy $\pi_{n+1,\lambda,\epsilon}$ is then well defined by

$$\pi_{n+1,\lambda,\epsilon}(\cdot|\bar{s}) = \frac{\operatorname{num}_{\lambda,\pi_{n,\epsilon}}(\cdot|\bar{s})}{\operatorname{den}_{\lambda,\pi_{n,\epsilon}}(\cdot|\bar{s})} + \frac{\epsilon}{|\mathcal{A}|}$$

for all $\bar{s} \in \bar{S}_{\lambda_0}$. Analogous to (7.4), we can rewrite the above equation similarly as in theorem 25 using $Q_{\lambda}^{\pi_{n,\epsilon},g,h}$ and $v_{\lambda,\pi_{n,\epsilon}}$. The proof proceeds by induction on the remaining horizon h.

Base case (h = 1): Now we fix $\beta, \tilde{\beta} \in (0, 1)$ arbitrary so that $1 > \gamma + \epsilon$, where $\gamma = \frac{\tilde{\beta}}{((1-\epsilon)^N - \beta)\alpha(\delta,\epsilon)}$. Note that we will be restricting (decreasing) $\delta > 0$ several times in the proof. Since $\alpha(\delta, \epsilon)$ is deceasing in δ , the decrease in δ will cause an increase in α , which further causes a decrease in γ leaving the constraint $1 > \gamma + \epsilon$ in place. According to point 1. of lemma 12 (continuity of Q_{λ}^*), point 5. of lemma 40, lemma 44, and lemma 41, we can fix $2 > \delta > 0$ so that for all $\lambda \in U_{\delta}(\lambda_0)$, all $\bar{s} = (s, 1, g) \in \bar{S}_{\lambda_0}$, and all $n \ge 1$ it holds that

$$\hat{\beta} > Q_{\lambda}^{*}(\bar{s}, a) \quad \text{for } a \notin \mathcal{O}(\bar{s}),$$
$$(1 - \epsilon)^{N} - \beta < Q_{\lambda}^{\pi_{n,\epsilon}}(\bar{s}, a) \quad \text{otherwise.}$$

Further, we lower bound the recursion for $\pi_{n+1,\lambda,\epsilon}(\mathcal{O}(\bar{s})|\bar{s})$ in exactly the same way as in theorem 25 using the above $\tilde{\beta}$ and β bounds, and also the α bound on the state visitation terms leaving $(\forall n \geq 0, \forall \pi_0 > 0, \forall \lambda \in U_{\delta}(\lambda_0), \forall \bar{s} = (s, 1, g) \in \bar{S}_{\lambda_0})$

$$\pi_{n+1,\lambda,\epsilon}(\mathcal{O}(\bar{s})|\bar{s}) \geq \frac{((1-\epsilon)^N - \beta)\alpha\pi_{n,\lambda,\epsilon}(\mathcal{O}(\bar{s})|\bar{s})}{((1-\epsilon)^N - \beta)\alpha\pi_{n,\lambda,\epsilon}(\mathcal{O}(\bar{s})|\bar{s}) + \tilde{\beta}} + \frac{\epsilon|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}$$
$$= z_{\gamma,\epsilon,\mathcal{O}(\bar{s})}(\pi_{n,\lambda,\epsilon}(\mathcal{O}(\bar{s})|\bar{s})). \tag{D.3}$$

Induction step: Now we aim to prove that the statement holds for a fixed h with h > 1, while working under the assumption that it holds for a smaller h (induction assumption). Fix $\beta, \tilde{\beta} \in (0, 1)$ arbitrary so that $1 > \gamma + \epsilon$, where $\gamma = \frac{\tilde{\beta}}{((1-\epsilon)^N - \beta)\alpha(\delta,\epsilon)}$. By lemma 44 there exists $\delta' > 0$ such that if the following two conditions are met

$$\lambda \in U_{\delta'}(\lambda_0),$$

$$(\forall \bar{s}' = (s', h', g') \in \bar{\mathcal{S}}_{\lambda_0}, h' < h): \quad 1 - \epsilon (1 - \frac{\mathcal{O}(\bar{s}')}{|\mathcal{A}|}) - \pi_{\epsilon}(\mathcal{O}(\bar{s}')|\bar{s}') < \frac{\delta'}{2}$$

then it holds that

$$(\forall \bar{s} = (s, h, g) \in \bar{\mathcal{S}}_{\lambda_0}, \forall a \in \mathcal{O}(\bar{s}), \forall \lambda \in U_{\delta'}(\lambda_0)): \quad Q_{\lambda}^{\pi_{\epsilon}}(\bar{s}, a) \ge (1 - \epsilon)^N - \beta$$

The first condition is met by the choice $0 < \delta < \delta' < 2$. To meet the second condition, we will use the induction assumption choosing $\beta', \tilde{\beta}' > 0$ such that $\gamma' + \epsilon < 1$ (where $\gamma' = \frac{\tilde{\beta}'}{((1-\epsilon)^N - \beta')\alpha(\delta,\epsilon)}$) and $x^*(\gamma', \epsilon, |\mathcal{O}(\bar{s}')|) > 1 - \epsilon(1 - \frac{|\mathcal{O}(\bar{s}')|}{|\mathcal{A}|}) - \delta'/2$ (this can be done since $x^*(\gamma', \epsilon, |\mathcal{O}(\bar{s}')|) \to 1 - \epsilon(1 - \frac{|\mathcal{O}(\bar{s}')|}{|\mathcal{A}|})$ for $(\beta', \tilde{\beta}') \to 0$). Applying the induction assumption requires a restriction on δ . From the induction assumption, it follows that there exists n'_0 such that, for all $n > n'_0, \lambda \in U_{\delta}(\lambda_0)$, it holds that $\pi_{n,\lambda,\epsilon}(\mathcal{O}(\bar{s}')|\bar{s}') > z^{n-n'_0}_{\gamma',\epsilon,|\mathcal{O}(\bar{s}')|}(\frac{\epsilon|\mathcal{O}(\bar{s}')|}{|\mathcal{A}|})$. As $z^{n-n'_0}_{\gamma',\epsilon,|\mathcal{O}(\bar{s}')|}(\frac{\epsilon|\mathcal{O}(\bar{s}')|}{|\mathcal{A}|}) \to x^*(\gamma',\epsilon,|\mathcal{O}(\bar{s}')|)$ there exists n_0 such that $\pi_{n,\lambda,\epsilon}(\mathcal{O}(\bar{s}')|\bar{s}') > 1 - \epsilon(1 - \frac{|\mathcal{O}(\bar{s}')|}{|\mathcal{A}|}) - \delta'/2$ for $n > n_0, \lambda \in U_{\delta}(\lambda_0)$.

In addition, by restricting $\delta > 0$ accordingly, we force the bound $Q_{\lambda}^*(\bar{s}, a) < \tilde{\beta}$ for all $\bar{s} \in \bar{S}_{\lambda_0}$ and all $a \notin \mathcal{O}(\bar{s})$. Finally, we can apply β , $\tilde{\beta}$ and state visitation bounds as we did before (for h = 1) to get the desired result.

D.3 Extending the Continuity Results to Other Segment Sub-Spaces

As before we need to modify theorem 34 to cover algorithms like ODT with regularization restricting the recursion to Seg^{diag} or Seg^{trail}. First, we will introduce Seg^{diag/trail} variants of lemmas 40 and 41. Note that equations (4.5), (4.4) and (5.1) depends only on π_0 and are therefore applicable also in Seg^{diag/trail}. Following the same reasoning as in section 4.2, one can extend the results in lemma 40 also to Seg^{diag/trail}.

Lemma 46 (ϵ -eUDRL version of support stability for Seg^{diag/trail}) The lemma 40 remains valid under renaming $\pi_{n,\epsilon} \to \pi_{n,\epsilon}^{\text{diag/trail}}$.

Similarly we can derive the variant of lemma 41 for Seg^{diag/trail}.

Lemma 47 (Optimality of ϵ -eUDRL policies for deterministic transition kernel in Seg^{diag/trail}) The lemma 41 remains valid under renaming $\pi_{n,\epsilon} \to \pi_{n,\epsilon}^{\text{diag/trail}}$.

The proof follows the same lines as in lemma 41 except in point 1. we use lemma 46 instead of lemma 40 and equations (4.5) and (4.4). Finally, we can state the $\text{Seg}^{\text{diag/trail}}$ version of the theorem 34.

Theorem 35 (Relative continuity of accumulation points of ϵ -eUDRL-generated policies at deterministic kernels) Theorem 34 remains valid under the renaming $\pi_{n,\epsilon} \to \pi_{n,\epsilon}^{\text{diag/trail}}$.

The proof follows similarly as the proof of theorem 34 except of small differences which resembles those described already in section 7.1.3. First, to assert that ϵ -eUDRL-generated policies are in $\Pi^*_{\lambda_0,\epsilon}$ at deterministic points one applies lemma 47 instead of lemma 41. Second, to prove that $\pi^{\text{diag/trail}}_{n+1,\epsilon}$ is well defined through equation (D.2), one uses equation (4.5) and point 2. of lemma 46 instead of lemma 40. Third, the application of point 1. of lemma 3 has to be replaced with points 2. and 3. of the lemma respectively. To bound the ϵ eUDRL recursion, one proceeds along the same lines yielding the exactly the same result as in (D.3). The same reasoning applies to the induction step. Finally, making use of lemma 3, which allows us to rewrite the ϵ -eUDRL recursion using only Seg-space-related quantities, shows that there are no further differences even for $\pi^{\text{diag/trail}}_{n+1,\epsilon}$. The same approach also applies to the forthcoming corollary of the theorem 34.

D.4 Estimating the Location of Accumulation Points

The following corollary is the ϵ -eUDRL version of the corollary 27.

Corollary 48 (Estimating the location of accumulation points – ϵ -eUDRL) Under the conditions of theorem 34 assume that $\delta \in (0, 1)$ and set

$$\alpha = \frac{2}{N(N-1)} \left(\min_{\bar{s} \in \bar{\mathcal{S}}_{\lambda_0}} \bar{\mu}(\bar{s}) \right) \left(\frac{\epsilon}{|\mathcal{A}|} \right)^N \left(1 - \frac{\delta}{2} \right)^N$$

Define the quantities $\tilde{\beta} = \frac{N\delta}{2}$, $x^*(\gamma, \epsilon, M) = \frac{\hat{x}^* + \sqrt{(\hat{x}^*)^2 + \frac{4\gamma\epsilon M}{|\mathcal{A}|}}}{2}$ where $\hat{x}^* = 1 - \epsilon(1 - \frac{M}{\mathcal{A}}) - \gamma$ (cf. lemma 45) and for horizon $h, 1 \le h \le N$,

$$\beta_{h} = \begin{cases} \max\{\delta, \tilde{\beta}\}, \text{ if } h = 1, \\ \delta + \kappa_{h-1} + \beta_{h-1}, \text{ if } h \ge 2, \end{cases}$$
$$\gamma_{h} = \frac{\tilde{\beta}}{((1-\epsilon)^{N} - \beta_{h})\alpha},$$
$$\kappa_{h} = \max_{\bar{s} = (s,h',g) \in \bar{\mathcal{S}}_{\lambda_{0}}, h' = h} 2\left(1 - \epsilon\left(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}\right) - x^{*}(\gamma_{h}, \epsilon, \mathcal{O}(\bar{s}))\right).$$

Further assume $\beta_h, \gamma_h \in (0, 1), 1 > \gamma_h + \epsilon$ and notice that $\beta_h, \kappa_h, \gamma_h$, are increasing in h and that $\beta_h, \kappa_h, \gamma_h \to 0$ as $\delta \to 0$. Then the following assertions hold for all $\lambda \in U_{\delta}(\lambda_0)$ and for all $\pi_0 > 0$:

1.
$$\lim_{n} \sup_{\bar{s} \in \bar{\mathcal{S}}_{\lambda_{0}}} 2\left(1 - \epsilon \left(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}\right) - \pi_{n,\epsilon}(\mathcal{O}(\bar{s})|\bar{s})\right) \le \kappa_{N},$$

2.
$$(\exists (\pi_{n,\epsilon}^{*}), \pi_{n,\epsilon}^{*} \in \Pi_{\lambda_{0},\epsilon}^{*}): \lim_{n} \sup_{\bar{s},a \in \bar{\mathcal{S}}_{\lambda_{0}} \times \mathcal{A}} \max_{\lambda_{0},\epsilon} |Q_{\lambda_{0}}^{\pi_{n,\epsilon}}(\bar{s},a) - Q_{\lambda_{0}}^{\pi_{n,\epsilon}^{*}}(\bar{s},a)| \le \beta_{N},$$

3.
$$(\exists (\pi_{n,\epsilon}^*), \pi_{n,\epsilon}^* \in \Pi_{\lambda_0,\epsilon}^*)$$
: $\limsup_{n} \max_{\bar{s} \in \bar{\mathcal{S}}_{\lambda_0}} |V_{\lambda}^{\pi_{n,\epsilon}}(\bar{s}) - V_{\lambda_0}^{\pi_{n,\epsilon}^*}(\bar{s})| \le \beta_N + \kappa_N,$

$$4. \ (\exists (\pi_{n,\epsilon}^*), \pi_{n,\epsilon}^* \in \Pi_{\lambda_0,\epsilon}^*): \ \limsup_{n} |J_{\lambda}^{\pi_{n,\epsilon}} - J_{\lambda_0}^{\pi_{n,\epsilon}^*}| \le \frac{N\delta}{2} + \beta_N + \kappa_N,$$

5. for all $\epsilon' > 0$ there exists n_0 and $(\pi^*_{n,\epsilon})_{n \ge 0}$, $\pi^*_{n,\epsilon} \in \Pi^*_{\lambda_0,\epsilon}$ so that for all $n \ge n_0$ and for all $\bar{s} \in \bar{S}_{\lambda_0}$ it holds that

$$\|\pi_{n,\epsilon}(\cdot|\bar{s}) - \pi^*_{n,\epsilon}(\cdot|\bar{s})\|_1 \le 2\left(1 - \epsilon\left(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}\right) - z^{\circ(n-n_0)}_{\gamma',\epsilon,|\mathcal{O}(\bar{s})|}(x_0)\right)$$

where $x_0 = \frac{\epsilon |\mathcal{O}(\bar{s})|}{|\mathcal{A}|}, \ \beta' > \beta_N + \epsilon'$, where ϵ' is chosen such that $1 > \gamma' + \epsilon'$ with $\gamma' = \frac{\tilde{\beta}}{((1-\epsilon)^N - \beta')\alpha} > 0.$

The terms on the right hand side of 1.-4. converge to 0 as $\delta \to 0$. In 5. the sequence $(y_n)_{n\geq 0}$ defined by $y_n = 2(1 - \epsilon(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}) - z_{\gamma',\epsilon,|\mathcal{O}(\bar{s})|}^{\circ(n-n_0)}(x_0))$ converges q-linearly to the limit $y = 2(1 - \epsilon(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}) - x^*(\gamma',\epsilon,|\mathcal{O}(\bar{s})|))$ at convergence rate $\frac{(1-\epsilon)\gamma'}{(x^*(\gamma',\epsilon,|\mathcal{O}(\bar{s})|)+\gamma')^2}$ which tends to 0 as $\delta \to 0$.

Proof The proof follows the same as the proof of corollary 27 except for slight variations in the corollary statement which has to be addressed. However, the main arguments are the same and we will not repeat them here. Again we rather try to concentrate on differences. Of course, one has to use lemmas introduced in this appendix instead of their original versions. The definition of α comes from lemma 42. We decided to drop δ , ϵ arguments for convenience because in this corollary proof there are no any frequent readjustments of δ in opposite to the previous theorem.

The claim has now a sightly different form $(\forall N \ge h \ge 1, \forall \pi_0 > 0, \forall \lambda \in U_{\delta}(\lambda_0))$:

$$(\exists (\pi_{n,\lambda_{0},\epsilon}^{*}), \pi_{n,\lambda_{0},\epsilon}^{*} \in \Pi_{\lambda_{0},\epsilon}^{*}) : \limsup_{n} \max_{\bar{s}=(s,h',g)\in\bar{S}_{\lambda_{0}},h'=h,a\in\mathcal{A}} |Q_{\lambda}^{\pi_{n,\epsilon}}(\bar{s},a) - Q_{\lambda_{0}}^{\pi_{n,\epsilon}^{*}}(\bar{s},a)| \leq \beta_{h},$$
$$\limsup_{n} \max_{\bar{s}=(s,h',g)\in\bar{S}_{\lambda_{0}},h'=h} 2(1-\epsilon(1-\frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}) - \pi_{n,\epsilon}(\mathcal{O}(\bar{s})|\bar{s})) \leq \nu_{h}$$

Notice that even before we were using convenient $\pi_{\lambda_0}^*$ to a policy π in the original version of lemma 44. The difference is that now we have to mention the chosen sequence $(\pi_{n,\lambda_0,\epsilon}^*)$ explicitly in the claim statement, because $Q_{\lambda_0}^{\pi_{n,\epsilon}^*}$ depends on the particular choice of $\pi_{n,\epsilon}^* \in \Pi_{\lambda_0,\epsilon}^*$ (unlike the optimal value $Q_{\lambda_0}^*$).

Points 1. and 2. follows again directly from the claim. points 3., 4., and 5. are proved in exactly the same way as in the original corollary. We comment only on point 5. here. We chose the sequence $(\pi_{n,\epsilon}^*)_{n\geq 0}$ so that $\pi_{n,\epsilon}^*(a \mid \bar{s}) \geq \pi_{n,\epsilon}(a \mid \bar{s})$, for all $\bar{s} \in \bar{S}_{\lambda_0}$ and all $a \in \mathcal{O}(\bar{s})$, which implies $\|\pi_{n,\epsilon}(\cdot|\bar{s}) - \pi_{n,\epsilon}^*(\cdot|\bar{s})\|_1 = 2(1 - \epsilon(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}) - \pi_{n,\epsilon}(\mathcal{O}(\bar{s})|\bar{s}))$. We conclude with the computation of the convergence rate. After applying Heine and L'Hospital theorems (in exactly the same way as in the original proof) one is left with

$$\begin{split} \lim_{n \to \infty} \frac{y_{n+1} - y_L}{y_n - y_L} &= \lim_{x \to x^*(\gamma', \epsilon, |\mathcal{O}(\bar{s})|)} z'_{\gamma', \epsilon, |\mathcal{O}(\bar{s})|} \\ &= \lim_{x \to x^*(\gamma', \epsilon, |\mathcal{O}(\bar{s})|)} \frac{(1 - \epsilon)\gamma'}{(x + \gamma')^2} = \frac{(1 - \epsilon)\gamma'}{(x^*(\gamma', \epsilon, |\mathcal{O}(\bar{s})|) + \gamma')^2} \\ &\leq \frac{(1 - \epsilon)\gamma'}{(\hat{x}^*(\gamma', \epsilon, |\mathcal{O}(\bar{s})|) + \gamma')^2} = \frac{(1 - \epsilon)\gamma'}{(1 - \epsilon(1 - \frac{|\mathcal{O}(\bar{s})|}{|\mathcal{A}|}))^2} \\ &\leq \frac{\gamma'}{(1 - \epsilon)} = \gamma' + \frac{\epsilon\gamma'}{(1 - \epsilon)} \leq \gamma' + \frac{\epsilon\gamma'}{\gamma'} = \gamma' + \epsilon < 1. \end{split}$$